

Robust Formulation of Wick’s theorem for Computing Matrix Elements Between Hartree–Fock–Bogoliubov Wavefunctions

Guo P. Chen^{1, a)} and Gustavo E. Scuseria^{1, 2}

¹⁾*Department of Chemistry, Rice University, Houston, TX 77005-1892*

²⁾*Department of Physics and Astronomy, Rice University, Houston, TX 77005-1892*

(Dated: April 26, 2023)

Numerical difficulties associated with computing matrix elements of operators between Hartree–Fock–Bogoliubov (HFB) wavefunctions have plagued the development of HFB-based many-body theories for decades. The nonorthogonal variant of Wick’s theorem leads to a zero-over-zero form in the limit of vanishing HFB overlap. This numerically ill-posed situation appears frequently in many theories. In this paper, we present a robust formulation of Wick’s theorem that stays well-behaved regardless of whether the HFB states are orthogonal or not. Our formula explicitly eliminates self-interaction, which otherwise causes additional numerical challenges. A computationally efficient version of our formalism enables robust symmetry-projected HFB calculations with the same computational cost as mean-field theories. We treat even and odd number of particles on the same footing, and reduce to Hartree–Fock as a natural limit. As proof of concept, we present a numerically stable and accurate solution to a Jordan–Wigner-transformed Hamiltonian, whose singularities motivated the present work. Our robust formulation of Wick’s theorem is a most promising development for methods using Bogoliubov product states.

I. INTRODUCTION

The Hartree–Fock–Bogoliubov (HFB) wavefunction is a fermionic mean-field ansatz allowing particle number symmetry breaking.¹ It encompasses the Hartree–Fock (HF) wavefunction in the same sense that spin-unrestricted HF (UHF) reduces to spin-restricted HF (RHF) without spin S^2 symmetry breaking. HFB-based methods are not as conventional in quantum chemistry as in nuclear structure theory. This is a consequence of repulsive electronic interactions in the former, which results in number symmetry not breaking spontaneously in mean-field.²

Nevertheless, the emergence of variation-after-projection approaches to symmetry breaking and restoration^{3–6} has sparked growing interest in HFB and related anzätze in electronic structure theory.^{5, 7–24} Methods based on symmetry-projected HFB (PHFB) are particularly promising for describing strongly correlated systems, where traditional HF-based methods fail.^{5, 25–27}

Essential for the implementation of PHFB-based methods are matrix elements of many-body operators between nonorthogonal HFB states. Indeed, the PHFB formalism can be considered a general form of nonorthogonal configuration interaction (NOCI).^{28, 29} The computation of relevant matrix elements relies on the generalized Wick’s theorem with respect to nonorthogonal HFB states, also known as the nonorthogonal Wick’s theorem.^{30, 31}

However, when the overlap between right and left HFB states vanishes, the standard formula in the nonorthogonal Wick’s theorem becomes numerically ill-conditioned, and eventually ill-defined, causing large round-off errors. This problem occurs in PHFB calculations^{32, 33} and is ex-

acerbated by self-interaction^{34, 35} if exchange- or pairing-type contractions are neglected, *e.g.*, within the context of nuclear density functional theory.^{35–38}

Our recent work mapping spin systems to fermions using the Jordan–Wigner transformation^{39, 40} is another example where orthogonal HFB wavefunctions arise and lead to large numerical errors. This happens when fermionic on-site occupations are equal to 1/2, which is frequently encountered in frustrated phases of transformed spin systems.

The main objective of this paper is demonstrating that with proper algorithmic design, the nonorthogonal Wick’s theorem can be extended to the orthogonal limit, resulting in accurate and efficient computation of matrix elements between arbitrary HFB states. Related algorithms have been discussed for HF-based NOCI,^{41–43} and our formalism reduces to them in that limit. Remedies for the numerical problems encountered in the HFB case have been proposed for specific applications^{32, 33} or under certain restrictions.^{44, 45} However, a universal, low-scaling formula for robust computation of matrix elements between HFB states, as the one presented here, was lacking.

II. NONORTHOGONAL WICK’S THEOREM

We define unnormalized HFB wavefunctions, $|\Phi_0\rangle$ and $|\Phi_1\rangle$ as

$$|\Phi_j\rangle = \prod_{p=1}^M \beta_p^j |-\rangle \quad (1)$$

for $j = 0, 1$, where $|-\rangle$ denotes the physical vacuum, and M is the dimensionality of the one-particle (spin-orbital) basis. The Bogoliubov quasiparticle annihilation operators β_p^j are related to the fermionic particle operators c_q^\dagger

^{a)}guo.chen@rice.edu

and c_q through a unitary canonical transformation known as the *Bogoliubov transformation*:

$$\beta_p^j = \sum_{q=1}^M (U_{qp}^{j*} c_q + V_{qp}^{j*} c_q^\dagger). \quad (2)$$

To ensure nonvanishing $|\Phi_j\rangle$, we assume that V^j is nonsingular, *i.e.*, all of the canonical orbitals of $|\Phi_j\rangle$ obtained from the Bloch–Messiah decomposition⁴⁶ of U^j and V^j (*vide infra*) are at least infinitesimally occupied. We will eliminate this assumption and address the normalization of $|\Phi_j\rangle$ in Sec. V.

The nonorthogonal Wick’s theorem in its most general form⁴⁷ states that, for $\langle \Phi_0 | \Phi_1 \rangle \neq 0$,

$$\langle \Phi_0 | \gamma^1 \gamma^2 \dots \gamma^{2d} | \Phi_1 \rangle = \langle \Phi_0 | \Phi_1 \rangle \text{pf}(\Gamma), \quad (3)$$

where $\{\gamma^k\}_{1 \leq k \leq 2d}$ is a set of arbitrary fermionic operators, and Γ is an antisymmetric matrix whose strict upper triangular part is defined by *contractions* of the form

$$\Gamma_{kl} = \overline{\gamma^k \gamma^l} = \frac{\langle \Phi_0 | \gamma^k \gamma^l | \Phi_1 \rangle}{\langle \Phi_0 | \Phi_1 \rangle} \quad (4)$$

for $1 \leq k < l \leq 2d$. The Pfaffian $\text{pf}(\Gamma)$ automatically generates all possible full contractions of the fermionic operators with the correct signs, reflecting the fermionic anticommutation relations. For example, Equation (3)

reduces to the conventional statement of the nonorthogonal Wick’s theorem^{1,30,31} for the two-particle reduce transition matrix (2-RTM):^{48–50}

$$\mathcal{D}_{pq,p'q'}^2 = \langle \Phi_0 | c_p^\dagger c_q^\dagger c_{q'} c_{p'} | \Phi_1 \rangle \quad (5a)$$

$$= \langle \Phi_0 | \Phi_1 \rangle (c_p^\dagger c_q^\dagger c_{q'} c_{p'} + c_p^\dagger c_q^\dagger c_{q'} c_{p'} + c_p^\dagger c_q^\dagger c_{q'} c_{p'}) \quad (5b)$$

$$= \langle \Phi_0 | \Phi_1 \rangle (\rho_{p'p}^{01} \rho_{q'q}^{01} - \rho_{p'q}^{01} \rho_{q'p}^{01} + \kappa_{pq}^{10*} \kappa_{p'q'}^{01}), \quad (5c)$$

where

$$\rho_{pq}^{01} = \overline{c_p^\dagger c_q}, \quad \kappa_{pq}^{01} = \overline{c_q c_p}, \quad \kappa_{pq}^{10*} = \overline{c_p^\dagger c_q^\dagger}, \quad (6)$$

and we follow the convention that each crossing between contraction lines introduces a minus sign, which is implied by the properties of the Pfaffian. In general, we can write γ^k as a quasiparticle

$$\gamma^k = \sum_{q=1}^M (A_{qk}^* c_q + B_{qk}^* c_q^\dagger), \quad (7)$$

where A and B are $M \times 2d$ matrices. Note that for $k \neq l$, γ^k and γ^l may come from different Bogoliubov transformations, hence the superscript in our notation.

We see from Equation (4) that the nonorthogonal Wick’s theorem becomes ill-defined when $\langle \Phi_0 | \Phi_1 \rangle = 0$. In this case, we may fall back to the original Wick’s theorem with respect to the physical vacuum.^{51–53} It follows that⁵⁴

$$\langle \Phi_0 | \gamma^1 \gamma^2 \dots \gamma^{2d} | \Phi_1 \rangle = (-1)^{M(M-1)/2} \langle - | \beta_1^{0\dagger} \beta_2^{0\dagger} \dots \beta_M^{0\dagger} \gamma^1 \gamma^2 \dots \gamma^{2d} \beta_1^1 \beta_2^1 \dots \beta_M^1 | - \rangle \quad (8a)$$

$$= (-1)^{M(M-1)/2} \text{pf} \begin{pmatrix} V^{0T} U^0 & V^{0T} B^* & V^{0T} V^{1*} \\ -B^\dagger V^0 & \Gamma^{(-)} & A^\dagger V^{1*} \\ -V^{1\dagger} V^0 & -V^{1\dagger} A^* & U^{1\dagger} V^{1*} \end{pmatrix}, \quad (8b)$$

where $\Gamma^{(-)}$ is an antisymmetric matrix defined by

$$\Gamma_{kl}^{(-)} = \langle - | \gamma^k \gamma^l | - \rangle \quad (9)$$

for $1 \leq k < l \leq 2d$. Compared with Equation (3), Equation (8) is always well-defined, but the Pfaffian of a much larger matrix needs to be evaluated, greatly increasing the computational cost.

In the following, we sketch a proof of the nonorthogonal Wick’s theorem stated in Equation (3), introducing our notation along the way. Unlike the proof by induction presented in Ref. 47, our proof follows a more direct approach inspired by Ref. 55. We start by rewriting Equation (8) using the properties of the Pfaffian:

$$\langle \Phi_0 | \gamma^1 \gamma^2 \dots \gamma^{2d} | \Phi_1 \rangle = (-1)^{M(M-1)/2} \text{pf}(\mathcal{M}), \quad (10)$$

where

$$\mathcal{M} = \begin{pmatrix} \mathcal{S} & \mathcal{G} \\ -\mathcal{G}^T & \Gamma^{(-)} \end{pmatrix}, \quad (11)$$

$$\mathcal{S} = \begin{pmatrix} V^{0T} U^0 & V^{0T} V^{1*} \\ -V^{1\dagger} V^0 & U^{1\dagger} V^{1*} \end{pmatrix}, \quad \mathcal{G} = \mathcal{V}^T \mathcal{C}^* \quad (12)$$

and

$$\mathcal{V} = \begin{pmatrix} 0 & -V^{1*} \\ V^0 & 0 \end{pmatrix}, \quad \mathcal{C} = \begin{pmatrix} A \\ B \end{pmatrix}. \quad (13)$$

When $\langle \Phi_0 | \Phi_1 \rangle \neq 0$, the antisymmetric matrix \mathcal{S} is nonsingular since⁵⁴

$$\langle \Phi_0 | \Phi_1 \rangle = (-1)^{M(M-1)/2} \text{pf}(\mathcal{S}). \quad (14)$$

We therefore have the Pfaffian identity

$$\text{pf}(\mathcal{M}) = \text{pf}(\mathcal{S})\text{pf}(\mathcal{M}/\mathcal{S}), \quad (15)$$

where \mathcal{M}/\mathcal{S} denotes the Schur complement of the block \mathcal{S} of the supermatrix \mathcal{M} , *i.e.*,

$$\mathcal{M}/\mathcal{S} = \Gamma^{(-)} + \mathcal{G}^T \mathcal{S}^{-1} \mathcal{G}. \quad (16)$$

On the other hand, as shown in detail in the supplementary material, we find

$$\mathcal{K} = \mathcal{V} \mathcal{S}^{-1} \mathcal{V}^T = \begin{pmatrix} -\kappa^{01} & -\rho^{01} \\ \rho^{01 T} & \kappa^{10*} \end{pmatrix}, \quad (17)$$

and it follows that

$$\mathcal{M}/\mathcal{S} = \Gamma^{(-)} + \mathcal{C}^\dagger \mathcal{K} \mathcal{C}^* = \Gamma. \quad (18)$$

Inserting Equations (14) and (18) into Equations (10) and (15) completes the proof.

Without loss of generality, we hereafter assume that $\gamma^1 \gamma^2 \dots \gamma^{2d}$ are particle operators in normal order with respect to the physical vacuum. Thus, we have $\Gamma^{(-)} = 0_{2d \times 2d}$ and

$$\langle \Phi_0 | \gamma^1 \gamma^2 \dots \gamma^{2d} | \Phi_1 \rangle = (-1)^{M(M-1)/2} \text{pf}(\mathcal{S}) \text{pf}(\mathcal{G}^T \mathcal{S}^{-1} \mathcal{G}). \quad (19)$$

These particular matrix elements include those that arise in the d -particle or d th-order reduce transition matrix (d -RTM),^{48–50}

$$\mathcal{D}_{p_1 p_2 \dots p_d, q_1 q_2 \dots q_d}^d = \langle \Phi_0 | c_{p_1}^\dagger c_{p_2}^\dagger \dots c_{p_d}^\dagger c_{q_d} \dots c_{q_2} c_{q_1} | \Phi_1 \rangle. \quad (20)$$

Equation (19) is reminiscent of Löwdin's formula for d -RTM between HF wavefunctions,⁵⁶ for which singular value decomposition (SVD) can be used to extract the zeros and poles in order to evaluate the zero-overlap limit.^{41–43} We show in the next section that this idea can be generalized to the HFB case.

III. ROBUST WICK'S THEOREM

The antisymmetric matrix \mathcal{S} can be written in canonical form,^{57,58}

$$\mathcal{S} = \mathcal{Q} \bar{\mathcal{S}} \mathcal{Q}^T, \quad (21)$$

where \mathcal{Q} is unitary and

$$\bar{\mathcal{S}} = \begin{pmatrix} 0 & \text{diag}(s) \\ -\text{diag}(s) & 0 \end{pmatrix}. \quad (22)$$

As shown in supplementary material, the elements of the vector s are non-negative and they reduce to the singular values of the overlap matrix in the HF case. Moreover, it follows from Equation (14) that

$$\langle \Phi_0 | \Phi_1 \rangle = \zeta \prod_r s_r, \quad (23)$$

where $\zeta = \det(\mathcal{Q})$ is a complex phase factor. We observe that the poles of \mathcal{S}^{-1} get cancelled out by the zeros of the overlap, hinting at a more general and robust formula for computing matrix elements.

Define

$$\tilde{\mathcal{K}}^r = \begin{pmatrix} -\tilde{\kappa}^{01,r} & -\tilde{\rho}^{01,r} \\ (\tilde{\rho}^{01,r})^T & (\tilde{\kappa}^{10,r})^* \end{pmatrix} \quad (24)$$

where

$$\tilde{\rho}_{pq}^{01,r} = [L_{11}]_{pr} [L_{22}]_{qr} - [L_{12}]_{pr} [L_{21}]_{qr}, \quad (25a)$$

$$\tilde{\kappa}_{pq}^{01,r} = [L_{11}]_{pr} [L_{12}]_{qr} - [L_{12}]_{pr} [L_{11}]_{qr}, \quad (25b)$$

$$(\tilde{\kappa}_{pq}^{10,r})^* = -[L_{21}]_{pr} [L_{22}]_{qr} + [L_{22}]_{pr} [L_{21}]_{qr}, \quad (25c)$$

and

$$\mathcal{L} = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix} = \mathcal{V} \mathcal{Q}^* \quad (26)$$

It is straightforward to show

$$\Gamma = \sum_r s_r^{-1} \tilde{\Gamma}^r, \quad \tilde{\Gamma}^r = \mathcal{C}^\dagger \tilde{\mathcal{K}}^r \mathcal{C}^*. \quad (27)$$

We now propose a robust formulation of Wick's theorem for computing matrix elements between HFB wavefunctions,

$$\langle \Phi_0 | \gamma^1 \gamma^2 \dots \gamma^{2d} | \Phi_1 \rangle = \zeta \times \sum_{r_1 r_2 \dots r_d} \lambda_{r_1 r_2 \dots r_d} \tilde{\text{pf}}(\tilde{\Gamma}^{r_1}, \tilde{\Gamma}^{r_2}, \dots, \tilde{\Gamma}^{r_d}) \quad (28)$$

with

$$\lambda_{r_1 \dots r_d} = \begin{cases} \prod_{r \neq r_1, \dots, r_d} s_r & (r_1, \dots, r_d \text{ are distinct}) \\ 0 & (\text{otherwise}) \end{cases} \quad (29)$$

$$\tilde{\text{pf}}(\tilde{\Gamma}^{r_1}, \tilde{\Gamma}^{r_2}, \dots, \tilde{\Gamma}^{r_d}) = \frac{1}{2^d d!} \times \sum_{\sigma \in S_{2d}} \text{sgn}(\sigma) \tilde{\Gamma}_{\sigma(1)\sigma(2)}^{r_1} \tilde{\Gamma}_{\sigma(3)\sigma(4)}^{r_2} \dots \tilde{\Gamma}_{\sigma(2d-1)\sigma(2d)}^{r_d} \quad (30)$$

where σ enumerates permutations in the symmetric group S_{2d} , and $\text{sgn}(\sigma)$ denotes the signature of σ . Equation (30) defines the generalized Pfaffian, $\tilde{\text{pf}}$, for a sequence of antisymmetric matrices, which reduces to the conventional Pfaffian when all the antisymmetric matrices are the same.

The proof of the theorem is presented in supplementary material. We emphasize that the formula in Equation (28) remains well-behaved with vanishing $\langle \Phi_0 | \Phi_1 \rangle$ because the denominator of each contraction has been cancelled out by construction. Moreover, the cancellation of self-interaction terms is guaranteed by the properties

of the generalized Pfaffian, and it is further enforced numerically by setting $\lambda_{r_1 \dots r_d}$ to zero when its indices are not distinct. From the definition of $\lambda_{r_1 \dots r_d}$, we can also see that the matrix element vanishes if the number of zero elements in s is greater than d , consistent with the Slater–Condon rules between HF wavefunctions.^{48,59,60}

For 2-RTM, the robust Wick’s theorem implies

$$\mathcal{D}_{pq,p'q'}^2 = \langle \Phi_0 | \overline{c_p^\dagger c_q^\dagger c_{q'} c_{p'}} + \overline{c_p^\dagger c_q^\dagger c_{q'} c_{p'}} + \overline{c_p^\dagger c_q^\dagger c_{q'} c_{p'}} | \Phi_1 \rangle \quad (31a)$$

$$= \zeta \sum_{r_1 r_2} \lambda_{r_1 r_2} \left(\tilde{\rho}_{p'p}^{01,r_1} \tilde{\rho}_{q'q}^{01,r_2} - \tilde{\rho}_{p'q}^{01,r_1} \tilde{\rho}_{q'p}^{01,r_2} + (\tilde{\kappa}_{pq}^{10,r_1})^* \tilde{\kappa}_{p'q'}^{01,r_2} \right). \quad (31b)$$

Computing the full 2-RTM using Equation (31b) scales as $\mathcal{O}(M^5)$. In practice, however, we never construct the full 2-RTM explicitly. To compute the matrix element of a two-body operator, we directly contract its parameters (*e.g.*, two-electron integrals) with the factorized form of

the 2-RTM (Equation (31b)). For example, the tensor contraction between the two-electron integrals and the 2-RTM should scale as $\mathcal{O}(M^3)$ to $\mathcal{O}(M^4)$ after exploiting locality using standard techniques.^{61–63}

IV. LOW-SCALING VERSION

We may further reduce the scaling by enforcing the cancellation only of the smallest elements in s . We partition Γ into singular (S) and regular (R) parts according to

$$\Gamma = \Gamma^S + \Gamma^R = \sum_{r \in S} s_r^{-1} \tilde{\Gamma}^r + \Gamma^R, \quad (32)$$

where $S = \{s_r < \epsilon \mid 1 \leq r \leq M, \epsilon > 0\}$. The value of ϵ should be so chosen that Γ^R is well-conditioned while the size of S is small. Similarly, we define $\rho^{01,R}$, $\kappa^{01,R}$, and $\kappa^{10,R}$ as the regular parts of ρ^{01} , κ^{01} , and κ^{10} , respectively. We can now establish a low-scaling robust Wick’s theorem, which states that

$$\begin{aligned} \langle \Phi_0 | \gamma^1 \gamma^2 \dots \gamma^{2d} | \Phi_1 \rangle &= \zeta \lambda^R \left(\lambda^S \text{pf}(\Gamma^R) + \sum_{r_1 \in S} \lambda_{r_1}^S \left(\text{pf}(\tilde{\Gamma}^{r_1}, \Gamma^R, \dots, \Gamma^R) + \text{pf}(\Gamma^R, \tilde{\Gamma}^{r_1}, \dots, \Gamma^R) + \dots + \text{pf}(\Gamma^R, \Gamma^R, \dots, \tilde{\Gamma}^{r_1}) \right) \right. \\ &+ \sum_{r_1 r_2 \in S} \lambda_{r_1 r_2}^S \left(\text{pf}(\tilde{\Gamma}^{r_1}, \tilde{\Gamma}^{r_2}, \Gamma^R, \dots, \Gamma^R) + \text{pf}(\tilde{\Gamma}^{r_1}, \Gamma^R, \tilde{\Gamma}^{r_2}, \dots, \Gamma^R) + \text{pf}(\Gamma^R, \dots, \Gamma^R, \tilde{\Gamma}^{r_1}, \tilde{\Gamma}^{r_2}) \right) \\ &+ \dots + \left. \sum_{r_1 r_2 \dots r_d \in S} \lambda_{r_1 r_2 \dots r_d}^S \text{pf}(\tilde{\Gamma}^{r_1}, \tilde{\Gamma}^{r_2}, \dots, \tilde{\Gamma}^{r_d}) \right), \quad (33) \end{aligned}$$

where

$$\lambda^S = \prod_{r \in S} s_r, \quad \lambda^R = \prod_{r \notin S} s_r, \quad (34a)$$

$$\lambda_{r_1 \dots r_d}^S = \begin{cases} \prod_{S \ni r \neq r_1, \dots, r_d} s_r & (r_1, \dots, r_d \text{ are distinct}) \\ 0 & (\text{otherwise}) \end{cases} \quad (34b)$$

Specifically for 2-RTM, we have

$$\begin{aligned} \langle \Phi_0 | \overline{c_p^\dagger c_q^\dagger c_{q'} c_{p'}} | \Phi_1 \rangle &= \zeta \lambda^R \\ &\times \left(\lambda^S \rho_{pq}^{01,R} \rho_{p'q'}^{01,R} + \sum_{r_1 \in S} \lambda_{r_1}^S \tilde{\rho}_{pq}^{01,r_1} \rho_{p'q'}^{01,R} \right. \\ &+ \sum_{r_1 \in S} \lambda_{r_1}^S \rho_{pq}^{01,R} \tilde{\rho}_{p'q'}^{01,r_1} + \left. \sum_{r_1, r_2 \in S} \lambda_{r_1 r_2}^S \tilde{\rho}_{pq}^{01,r_1} \tilde{\rho}_{p'q'}^{01,r_2} \right) \quad (35) \end{aligned}$$

and similarly for $\langle \Phi_0 | \overline{c_p^\dagger c_q^\dagger c_{q'} c_{p'}} | \Phi_1 \rangle$ and

$\langle \Phi_0 | \overline{c_p^\dagger c_q^\dagger c_{q'} c_{p'}} | \Phi_1 \rangle$. Each term in the above expression stays factorized, facilitating low-scaling tensor contractions with the electron integrals. Besides, the indices r_1, r_2 now only run over a small subset of s elements. As a result, the computational cost of the matrix element of a two-body operator scales the same as that of HF, *i.e.*, $\mathcal{O}(M^2)$ to $\mathcal{O}(M^3)$, albeit with a larger prefactor. For d -RTM with $2 < d \ll M$, the reduction of computational scaling from Equation (28) to Equation (33) is even more significant.

We can now generalize the Slater–Condon rules to the HFB case using the low-scaling robust Wick’s theorem. Let μ be the number of s elements that are strictly zero. Loosely speaking, μ equals the smallest number of levels we need to block in both $|\Phi_0\rangle$ and $|\Phi_1\rangle$ in order to make them nonorthogonal. The blocking should be done in a biorthogonal basis that simultaneously brings both $|\Phi_0\rangle$ and $|\Phi_1\rangle$ to the canonical form of the Bardeen–Cooper–Schrieffer ansatz.^{44,45,64} This biorthogonal basis is generally nonunitary and does not always exist as noted in Ref. 45. It is readily seen from Equation (33) that the λ^S term vanishes for $\mu = 1$, the λ^S and $\lambda_{r_1}^S$ terms van-

ish for $\mu = 2$, and so forth. These results resemble the generalized Slater–Condon rules for HF transition matrix elements,^{43,65} in which case μ corresponds to the smallest number of biorthogonal occupied orbital pairs that need to be emptied in order to make the HF wavefunctions nonorthogonal. Here the biorthogonalization is guaranteed by Löwdin pairing and realized by SVD.^{56,66} We should point out that, as opposed to the generalized Slater–Condon rules, our Equation (33) also ensures numerical stability when some elements of s are small but not strictly zero.

V. ROBUST NORMALIZATION

So far we have considered unnormalized HFB wavefunctions $|\Phi_j\rangle$ for $j = 0, 1$ as defined in Equation (1). The corresponding normalized HFB wavefunction is⁵⁴

$$|\Phi'_j\rangle = \frac{\det(C^j)}{\prod_{p=1}^{m_P^j} v_p^j} |\Phi_j\rangle, \quad (36)$$

where C^j and v^j come from the Bloch–Messiah decomposition,⁴⁶

$$U^j = D^j \bar{U}^j C^j, \quad V^j = (D^j)^* \bar{V}^j C^j \quad (37)$$

with D^j and C^j being unitary and

$$\bar{U}^j = \begin{pmatrix} 0_{M_C^j \times M_C^j} & & & \\ & m_P^j & & \\ & -\bigoplus_{p=1} i\sigma_y u_p^j & & \\ & & & I_{M_V^j} \end{pmatrix}, \quad (38a)$$

$$\bar{V}^j = \begin{pmatrix} I_{M_C^j} & & & \\ & m_P^j & & \\ & \bigoplus_{p=1} I_2 v_p^j & & \\ & & & 0_{M_V^j \times M_V^j} \end{pmatrix}. \quad (38b)$$

Here I_k denotes the $k \times k$ identity matrix, and σ_y denotes the y -component of the 2×2 Pauli matrices; u_p^j, v_p^j are real and satisfy $(u_p^j)^2 + (v_p^j)^2 = 1$. Note that we use an unconventional definition of (\bar{U}^j, \bar{V}^j) in Equation (38), which is physically inconsequential but will ease the notation in what follows. The Bloch–Messiah decomposition implies that, in the canonical basis of $|\Phi_j\rangle$ with orbital coefficients D^j , we have M_C^j fully occupied core orbitals, $2m_P^j$ paired fractionally occupied orbitals, and M_V^j unoccupied or virtual orbitals, where $M_C^j + 2m_P^j + M_V^j = M$.

The normalization factor in Equation (36) is generally unbounded, which may introduce additional numerical challenges. To overcome this problem, we define a set of unnormalized quasiparticles by an unnormalized Bogoliubov transformation:

$$\beta_p^{j'} = \sum_{q=1}^M \left((U_{qp}^{j'})^* c_q + (V_{qp}^{j'})^* c_q^\dagger \right) \quad (39)$$

for $1 \leq p \leq M_C^j + 2m_P^j$, with

$$U^{j'} = D^{j'} \bar{U}^{j'} (\bar{V}^{j'})^{-\frac{1}{2}}, \quad V^{j'} = (D^{j'})^* (\bar{V}^{j'})^{\frac{1}{2}}, \quad (40)$$

where $D^{j'}$ is built from the first $M_C^j + 2m_P^j$ columns of D^j as suggested in Ref. 67, and

$$\bar{U}^{j'} = \begin{pmatrix} 0_{M_C^j \times M_C^j} & & & \\ & m_P^j & & \\ & -\bigoplus_{p=1} i\sigma_y u_p^j & & \\ & & & I_{M_V^j} \end{pmatrix}, \quad (41a)$$

$$\bar{V}^{j'} = \begin{pmatrix} I_{M_C^j} & & & \\ & m_P^j & & \\ & \bigoplus_{p=1} I_2 v_p^j & & \\ & & & 0_{M_V^j \times M_V^j} \end{pmatrix}. \quad (41b)$$

It is readily shown that the normalized HFB wavefunction $|\Phi'_j\rangle$ is a Bogoliubov product state of a set of unnormalized quasiparticles, *i.e.*,

$$|\Phi'_j\rangle = \prod_{p=1}^{M_C^j + 2m_P^j} \beta_p^{j'} |-\rangle. \quad (42)$$

Furthermore,

$$\mathcal{S}' = \begin{pmatrix} (V^{0'})^\top U^{0'} & (V^{0'})^\top (V^{1'})^* \\ - (V^{1'})^\dagger V^{0'} & (U^{1'})^\dagger (V^{1'})^* \end{pmatrix} \quad (43a)$$

$$= \begin{pmatrix} \bar{U}^{0'} & (V^{0'})^\top (V^{1'})^* \\ - (V^{1'})^\dagger V^{0'} & -\bar{U}^{1'} \end{pmatrix}, \quad (43b)$$

whose elements are bounded by 1 in absolute value. Therefore, we no longer have to deal with the problematic normalization factor in Equation (36) as long as we compute the overlap between normalized HFB wavefunctions using

$$\langle \Phi'_0 | \Phi'_1 \rangle = (-1)^{(M_C^0 + 2m_P^0)(M_C^0 + 2m_P^0 - 1)/2} \text{pf}(\mathcal{S}'). \quad (44)$$

This procedure for computing the overlap is similar to the one proposed in Ref. 68; however, we emphasize that our formalism treats even- and odd-particle systems on the same footing, since the number of core orbitals M_C^j is not restricted to be even. More importantly, Equation (42) suggests that the robust Wick's theorem and its low-scaling version, Equations (28) and (33), remain valid for the d -RTM between the normalized $|\Phi'_0\rangle$ and $|\Phi'_1\rangle$ after formally replacing (U^j, V^j) with $(U^{j'}, V^{j'})$, or equivalently replacing \mathcal{S} with \mathcal{S}' and \mathcal{V} with

$$\mathcal{V}' = \begin{pmatrix} 0 & - (V^{1'})^* \\ V^{0'} & 0 \end{pmatrix}. \quad (45)$$

It is easier to see this correspondence from Equation (19), whose derivation does not rely on the normalization of the Bogoliubov transformations.

Finally, we note in passing that, when $m_P^j = 0$, $|\Phi'_j\rangle$ reduces to a HF wavefunction, so the robust Wick's theorem can also be used to compute HF reduced transition matrices, with all pairing-type contractions removed due to vanishing κ^{01} and κ^{10} .




FIG. 1. Energy errors for the 8-site XXZ chain with $\langle S^z \rangle = 0$. Left panel: Errors with respect to the full configuration interaction (FCI) energy. Hartree–Fock (HF) results are plotted to show symmetry breaking. Right panel: Numerical errors in absolute value with respect to the Hartree–Fock–Bogoliubov (HFB) energy computed without Jordan–Wigner (JW) strings according to Equation (49).

VI. NUMERICAL EXAMPLE

We here present a prototypical example to demonstrate the power of our proposed robust Wick’s theorem. The Heisenberg XXZ Hamiltonian can be treated as a fermionic system after Jordan–Wigner (JW) transformation. The claim to fame of this mapping is that in 1D at $\Delta = 0$, a maximally degenerate point, the transformed Hamiltonian becomes a free fermion system readily solvable by mean-field HF theory. Here, Δ is the anisotropy parameter. We refer the reader to Ref. 40 for details of the transformed Hamiltonian. For $\Delta < 0$, number symmetry breaks spontaneously, necessitating HFB solutions. Matrix elements of the form

$$\langle \Phi | c_p^\dagger c_q | \Phi^{pq} \rangle = \langle \Phi | c_p^\dagger c_q \phi_p \phi_q | \Phi \rangle \quad (46)$$

need to be computed for $1 \leq p < q \leq M$, where the JW string

$$\phi_p = \prod_{q < p} (1 - 2n_q) = \exp(i\pi \sum_{q < p} n_q) \quad (47)$$

is a Thouless rotation acting on $|\Phi\rangle$ with $n_q = c_q^\dagger c_q$. The overlap

$$\langle \Phi | \Phi^{pq} \rangle = \langle \Phi | \phi_p \phi_q | \Phi \rangle = \langle \Phi | \prod_{r=p}^{q-1} (1 - 2n_r) | \Phi \rangle \quad (48)$$

vanishes when $\langle \Phi | n_r | \Phi \rangle = 1/2$ for $p \leq r < q$, a situation we observe across the critical $-1 \leq \Delta \leq 1$ region of the 1D phase diagram. On the other hand, because XXZ possess only nearest neighbor interactions, one can avoid dealing with JW strings all together, and analytically show that

$$\langle \Phi | c_p^\dagger c_{p+1} | \Phi^{p,p+1} \rangle = - \langle \Phi | c_p^\dagger c_{p+1} | \Phi \rangle. \quad (49)$$

The right hand side of Equation (49) is numerically well-posed because $\langle \Phi | \Phi \rangle = 1$, after applying the normalization procedure in Sec. V. Therefore, we have two analytically equivalent forms of evaluating the same matrix elements. Comparing numerical results from Equation (46), which generates orthogonal HFB states, and Equation(49), we can measure the round-off errors due to vanishing $\langle \Phi | \Phi^{pq} \rangle$ and verify that our robust Wick’s theorem eliminates them.

Consider a JW-transformed 8-site XXZ chain with open boundary conditions. We constrain $\langle \Phi | \sum_p n_p | \Phi \rangle = M/2$, which corresponds to $\langle S^z \rangle = 0$. We find that $\langle \Phi | \Phi^{p,p+1} \rangle = 0$ for all $1 \leq p < M$ at $-1.12 \leq \Delta \leq 0.98$. We observe equal site occupations ($\langle S_p^z \rangle = 0$ in the original spin representation) within this range of Δ values, consistent with our previous calculations using a spin antisymmetrized geminal power ansatz.⁶⁹ As shown in Fig. 1, the HFB energies obtained from Equation (46) using the nonorthogonal Wick’s theorem indeed suffer from severe numerical error in this regime. In contrast, the robust Wick’s theorem, yields energies on top of he exact results computed without JW strings using Equation (49).

VII. CONCLUDING REMARKS

It is frequently assumed in quantum mechanical calculations with nonorthogonal basis that Hamiltonian matrix elements are small whenever the corresponding overlap matrix elements are small. This assumption is seemingly consistent with the standard formulation of the nonorthogonal Wick’s theorem in Equation (3), where the matrix element is proportional to the overlap $\langle \Phi_0 | \Phi_1 \rangle$. However, as our analysis shows, Equation (3) is ill-defined when $\langle \Phi_0 | \Phi_1 \rangle = 0$, while its limit as $\langle \Phi_0 | \Phi_1 \rangle$

tends to 0 is always well-defined and can be nonzero in general, disproving the assumption above.

The idea of taking the limit before numerical evaluation is central to our robust formulation of Wick's theorem. The resulting formula for matrix elements between HFB wavefunctions is universally applicable and numerically stable. In addition to the zero-overlap limit, the limits of fully occupied and completely empty orbitals have been considered, which also allow us to treat even- and odd-particle systems on the same footing. Moreover, we have presented a robust normalization procedure compatible with the robust Wick's theorem. This is achieved by formally viewing a normalized HFB wavefunction as a Bogoliubov product state of unnormalized quasiparticles. Although the coefficients defining these unnormalized quasiparticles are unbounded in general, the \mathcal{S}' and \mathcal{V}' matrices entering the final expressions for the matrix elements remains bounded.

To summarize, our robust formulation of Wick's theorem for computing matrix elements between HFB states resolves formal and numerical limitations of pre-existing formulations. We expect it to pave the way for future developments of many-body HFB-based theories.

ACKNOWLEDGMENTS

This work was supported by the U.S. National Science Foundation under Grant No. CHE-2153820. G.E.S. is a Welch Foundation Chair (Grant No. C-0036). G.P.C. thanks Dr. Thomas M. Henderson for helpful discussions and useful comments.

- ¹P. Ring and P. Schuck, *The nuclear many-body problem* (Springer-Verlag, Berlin Heidelberg, 1980).
- ²E. H. Lieb, "Variational principle for many-fermion systems," *Phys. Rev. Lett* **46**, 457–459 (1981).
- ³J. A. Sheikh and P. Ring, "Symmetry-projected Hartree–Fock–Bogoliubov equations," *Nucl. Phys. A* **665**, 71–91 (2000).
- ⁴K. Schmid, "On the use of general symmetry-projected Hartree–Fock–Bogoliubov configurations in variational approaches to the nuclear many-body problem," *Prog. Part. Nucl. Phys.* **52**, 565–633 (2004).
- ⁵G. E. Scuseria, C. A. Jiménez-Hoyos, T. M. Henderson, K. Samanta, and J. K. Ellis, "Projected quasiparticle theory for molecular electronic structure," *J. Chem. Phys.* **135**, 124108 (2011).
- ⁶J. A. Sheikh, J. Dobaczewski, P. Ring, L. M. Robledo, and C. Yannouleas, "Symmetry restoration in mean-field approaches," *J. Phys. G: Nucl. Part. Phys.* **48**, 123001 (2021).
- ⁷K. Naftchi-Ardebili, N. W. Hau, and D. A. Mazziotti, "Rank restriction for the variational calculation of two-electron reduced density matrices of many-electron atoms and molecules," *Phys. Rev. A* **84**, 052506 (2011).
- ⁸E. Neuscamman, "Size consistency error in the antisymmetric geminal power wave function can be completely removed," *Phys. Rev. Lett.* **109**, 203001 (2012).
- ⁹E. Neuscamman, "The Jastrow antisymmetric geminal power in Hilbert space: Theory, benchmarking, and application to a novel transition state," *J. Chem. Phys.* **139**, 194105 (2013).
- ¹⁰C. A. Jiménez-Hoyos, R. Rodríguez-Guzmán, and G. E. Scuseria, " N -electron Slater determinants from nonunitary canonical transformations of fermion operators," *Phys. Rev. A* **86**, 052102 (2012).
- ¹¹W. Uemura, S. Kasamatsu, and O. Sugino, "Configuration interaction with antisymmetrized geminal powers," *Phys. Rev. A* **91**, 062504 (2015).
- ¹²T. Tsuchimochi, M. Welborn, and T. Van Voorhis, "Density matrix embedding in an antisymmetrized geminal power bath," *J. Chem. Phys.* **143**, 024107 (2015).
- ¹³M. Degroote, T. M. Henderson, J. Zhao, J. Dukelsky, and G. E. Scuseria, "Polynomial similarity transformation theory: A smooth interpolation between coupled cluster doubles and projected BCS applied to the reduced BCS hamiltonian," *Phys. Rev. B* **93**, 125124 (2016).
- ¹⁴H. Shi and S. Zhang, "Many-body computations by stochastic sampling in Hartree–Fock–Bogoliubov space," *Phys. Rev. B* **95**, 045144 (2017).
- ¹⁵P. A. Johnson, P. A. Limacher, T. D. Kim, M. Richer, R. A. Miranda-Quintana, F. Heidar-Zadeh, P. W. Ayers, P. Bultinck, S. De Baerdemacker, and D. Van Neck, "Strategies for extending geminal-based wavefunctions: Open shells and beyond," *Comput. Theor. Chem.* **1116**, 207–219 (2017).
- ¹⁶A. Mahajan and S. Sharma, "Symmetry-projected Jastrow mean-field wave function in variational Monte Carlo," *J. Phys. Chem. A* **123**, 3911–3921 (2019).
- ¹⁷A. Khamoshi, T. M. Henderson, and G. E. Scuseria, "Efficient evaluation of AGP reduced density matrices," *J. Chem. Phys.* **151**, 184103 (2019).
- ¹⁸T. M. Henderson and G. E. Scuseria, "Correlating the antisymmetrized geminal power wave function," *J. Chem. Phys.* **153**, 084111 (2020).
- ¹⁹R. Dutta, T. M. Henderson, and G. E. Scuseria, "Geminal replacement models based on AGP," *J. Chem. Theory Comput.* **16**, 6358–6367 (2020).
- ²⁰P. A. Johnson, C. E. Fecteau, F. Berthiaume, S. Cloutier, L. Carrier, M. Gratton, P. Bultinck, S. D. Baerdemacker, D. V. Neck, P. Limacher, and P. W. Ayers, "Richardson–Gaudin mean-field for strong correlation in quantum chemistry," *J. Chem. Phys.* **153**, 104110 (2020).
- ²¹A. Khamoshi, F. A. Evangelista, and G. E. Scuseria, "Correlating AGP on a quantum computer," *Quantum Sci. Technol.* **6**, 014004 (2021).
- ²²A. Khamoshi, G. P. Chen, T. M. Henderson, and G. E. Scuseria, "Exploring non-linear correlators on AGP," *J. Chem. Phys.* **154**, 074113 (2021).
- ²³R. Dutta, G. P. Chen, T. M. Henderson, and G. E. Scuseria, "Construction of linearly independent non-orthogonal AGP states," *J. Chem. Phys.* **154**, 114112 (2021).
- ²⁴A. Khamoshi, G. P. Chen, F. A. Evangelista, and G. E. Scuseria, "AGP-based unitary coupled cluster theory for quantum computers," *Quantum Sci. Technol.* **8**, 015006 (2023).
- ²⁵C. N. Yang, "Concept of off-diagonal long-range order and the quantum phases of liquid He and of superconductors," *Rev. Mod. Phys.* **34**, 694–704 (1962).
- ²⁶A. J. Coleman, "Structure of fermion density matrices. II. Antisymmetrized geminal powers," *J. Math. Phys.* **6**, 1425–1431 (1965).
- ²⁷L. M. Sager and D. A. Mazziotti, "Cooper-pair condensates with nonclassical long-range order on quantum devices," *Phys. Rev. Research* **4**, 013003 (2022).
- ²⁸D. L. Hill and J. A. Wheeler, "Nuclear constitution and the interpretation of fission phenomena," *Phys. Rev.* **89**, 1102–1145 (1953).
- ²⁹R. E. Peierls and J. Yoccoz, "The collective model of nuclear motion," *Proc. Phys. Soc. A* **70**, 381 (1957).
- ³⁰N. Onishi and S. Yoshida, "Generator coordinate method applied to nuclei in the transition region," *Nucl. Phys.* **80**, 367–376 (1966).
- ³¹R. Balian and E. Brezin, "Nonunitary Bogoliubov transformations and extension of Wick's theorem," *Nuovo Cimento B* **64**, 37 (1969).
- ³²N. Tajima, H. Flocard, P. Bonche, J. Dobaczewski, and P.-H. Heenen, "Generator coordinate kernels between zero- and two-

- quasiparticle BCS states,” Nucl. Phys. A **542**, 355–367 (1992).
- ³³M. Anguiano, J. Egido, and L. Robledo, “Particle number projection with effective forces,” Nucl. Phys. A **696**, 467–493 (2001).
- ³⁴J. P. Perdew and A. Zunger, “Self-interaction correction to density-functional approximations for many-electron systems,” Phys. Rev. B **23**, 5048–5079 (1981).
- ³⁵M. Bender, T. Duguet, and D. Lacroix, “Particle-number restoration within the energy density functional formalism,” Phys. Rev. C **79**, 044319 (2009).
- ³⁶J. Dobaczewski, M. V. Stoitsov, W. Nazarewicz, and P.-G. Reinhard, “Particle-number projection and the density functional theory,” Phys. Rev. C **76**, 054315 (2007).
- ³⁷D. Lacroix, T. Duguet, and M. Bender, “Configuration mixing within the energy density functional formalism: Removing spurious contributions from nondiagonal energy kernels,” Phys. Rev. C **79**, 044318 (2009).
- ³⁸T. Duguet, M. Bender, K. Bennaceur, D. Lacroix, and T. Lesinski, “Particle-number restoration within the energy density functional formalism: Nonviability of terms depending on noninteger powers of the density matrices,” Phys. Rev. C **79**, 044320 (2009).
- ³⁹P. Jordan and E. Wigner, “Über das paulische äquivalenzverbot,” Z. Phys. **47**, 631–651 (1928).
- ⁴⁰T. M. Henderson, G. P. Chen, and G. E. Scuseria, “Strong-weak duality via Jordan–Wigner transformation: Using fermionic methods for strongly correlated $su(2)$ spin systems,” J. Chem. Phys. **157**, 194114 (2022).
- ⁴¹H. Koch and E. Dalgaard, “Linear superposition of optimized non-orthogonal slater determinants for singlet states,” Chem. Phys. Lett. **212**, 193–200 (1993).
- ⁴²J. Rodriguez-Laguna, L. M. Robledo, and J. Dukelsky, “Efficient computation of matrix elements of generic slater determinants,” Phys. Rev. A **101**, 012105 (2020).
- ⁴³H. G. A. Burton, “Generalized nonorthogonal matrix elements: Unifying Wick’s theorem and the Slater–Condon rules,” J. Chem. Phys. **154**, 144109 (2021).
- ⁴⁴F. Döna, “Canonical form of transition matrix elements,” Phys. Rev. C **58**, 872–877 (1998).
- ⁴⁵J. Dobaczewski, “Generalization of the Bloch–Messiah–Zumino theorem,” Phys. Rev. C **62**, 017301 (2000).
- ⁴⁶C. Bloch and A. Messiah, “The canonical form of an antisymmetric tensor and its application to the theory of superconductivity,” Nucl. Phys. **39**, 95–106 (1962).
- ⁴⁷Q.-L. Hu, Z.-C. Gao, and Y. Chen, “Matrix elements of one-body and two-body operators between arbitrary HFB multi-quasiparticle states,” Phys. Lett. B **734**, 162–166 (2014).
- ⁴⁸P.-O. Löwdin, “Quantum theory of many-particle systems. I. Physical interpretations by means of density matrices, natural spin-orbitals, and convergence problems in the method of configurational interaction,” Phys. Rev. **97**, 1474–1489 (1955).
- ⁴⁹R. McWeeny, “Some recent advances in density matrix theory,” Rev. Mod. Phys. **32**, 335–369 (1960).
- ⁵⁰D. A. Mazziotti, “Comparison of contracted Schrödinger and coupled-cluster theories,” Phys. Rev. A **60**, 4396–4408 (1999).
- ⁵¹G. C. Wick, “The evaluation of the collision matrix,” Phys. Rev. **80**, 268–272 (1950).
- ⁵²M. Gaudin, “Une démonstration simplifiée du théorème de Wick en mécanique statistique,” Nucl. Phys. **15**, 89–91 (1960).
- ⁵³E. H. Lieb, “A theorem on Pfaffians,” J. Comb. Theory **5**, 313–319 (1968).
- ⁵⁴G. F. Bertsch and L. M. Robledo, “Symmetry restoration in Hartree–Fock–Bogoliubov based theories,” Phys. Rev. Lett. **108**, 042505 (2012).
- ⁵⁵T. Mizusaki and M. Oi, “A new formulation to calculate general HFB matrix elements through the Pfaffian,” Phys. Lett. B **715**, 219–224 (2012).
- ⁵⁶P.-O. Löwdin, “Quantum theory of many-particle systems. II. Study of the ordinary Hartree–Fock approximation,” Phys. Rev. **97**, 1490–1508 (1955).
- ⁵⁷L.-K. Hua, “On the theory of automorphic functions of a matrix variable I—Geometrical basis,” Am. J. Math. **66**, 470 (1944).
- ⁵⁸M. Wimmer, “Algorithm 923: Efficient numerical computation of the Pfaffian for dense and banded skew-symmetric matrices,” ACM Trans. Math. Softw. **38**, 1–17 (2012).
- ⁵⁹J. C. Slater, “The theory of complex spectra,” Phys. Rev. **34**, 1293–1322 (1929).
- ⁶⁰E. U. Condon, “The theory of complex spectra,” Phys. Rev. **36**, 1121–1133 (1930).
- ⁶¹M. Häser and R. Ahlrichs, “Improvements on the direct SCF method,” J. Comput. Chem. **10**, 104–111 (1989).
- ⁶²D. L. Strout and G. E. Scuseria, “A quantitative study of the scaling properties of the Hartree–Fock method,” J. Chem. Phys. **102**, 8448–8452 (1995).
- ⁶³M. Challacombe and E. Schwegler, “Linear scaling computation of the Fock matrix,” J. Chem. Phys. **106**, 5526–5536 (1997).
- ⁶⁴K. Burzyński and J. Dobaczewski, “Quadrupole-collective states in a large single- j shell,” Phys. Rev. C **51**, 1825–1841 (1995).
- ⁶⁵J. Verbeek and J. H. Van Lenthe, “The generalized Slater–Condon rules,” Int. J. Quantum Chem. **40**, 201–210 (1991).
- ⁶⁶A. T. Amos, G. G. Hall, and H. Jones, “Single determinant wave functions,” Proc. Phys. Soc. A **263**, 483–493 (1961).
- ⁶⁷L. M. Robledo, “Technical aspects of the evaluation of the overlap of Hartree–Fock–Bogoliubov wave functions,” Phys. Rev. C **84**, 014307 (2011).
- ⁶⁸B. G. Carlsson and J. Rotureau, “New and practical formulation for overlaps of Bogoliubov vacua,” Phys. Rev. Lett. **126**, 172501 (2021).
- ⁶⁹Z. Liu, F. Gao, G. P. Chen, T. M. Henderson, J. Dukelsky, and G. E. Scuseria, “Exploring spin AGP ansätze for strongly correlated spin systems,” (2023), arXiv:2303.04925.