Andrés J. Moreno and Julieth Saavedra

1st May 2023

#### Abstract

In this work, we approach the Laplacian coflow of a coclosed  $G_2$ -structure  $\varphi$  using the formulae for the irreducible  $G_2$ -decomposition of the Hodge Laplacian and the Lie derivative of the Hodge dual 4-form of  $\varphi$ . In terms of this decomposition, we characterise the conditions for a vector field as an infinitesimal symmetry of a coclosed  $G_2$ -structure, as well as the soliton condition for the Laplacian coflow. More specifically, we provide an easier proof for the absence of compact shrinking solitons of the Laplacian coflow. Moreover, we revisit the Laplacian coflow of coclosed  $G_2$ -structures on almost Abelian Lie groups addressed by Fino-Bagaglini [BF18]. However, our approach is based on the bracket flow point of view. Notably, by showing that the norm of the Lie bracket is strictly decreasing, we prove that we have long-time existence for any coclosed Laplacian coflow solution.

## Contents

1	Introduction	1
2	Preliminaries	5
	2.1 G <sub>2</sub> -structures and their torsion	5
	2.2 Properties of coclosed $G_2$ -structures	6
3	Laplacian coflow of $G_2$ -structures	7
4	Almost Abelian Lie groups revisited	12
	4.1 The bracket flow	14
	4.2 Algebraic solitons	18
A	Contraction of $G_2$ and $SU(3)$ -identities	21
References		21

# **1** Introduction

A G<sub>2</sub>-structure is defined by a positive 3-form  $\varphi$ , which, in turn, defines the metric g and the corresponding Hodge dual 4-form  $\psi := *\varphi$ . The main goal in G<sub>2</sub>-geometric is the study of *torsion free* G<sub>2</sub>-structures, i.e.  $\nabla \varphi = 0$ , which is equivalent to the *closed* d $\varphi = 0$  and the *coclosed* condition d $\psi = 0$  (e.g [FG82]). Using Ricci flow ideas, Bryant introduced the *Laplacian flow* of closed G<sub>2</sub>-structures [Bry06], which is an evolution of an initial closed G<sub>2</sub>-structure along its Hodge Laplacian, namely

$$\frac{\partial \varphi(t)}{\partial t} = \Delta_t \varphi(t), \quad \varphi(0) = \varphi.$$
(1.1)

The Laplacian flow is not parabolic, however, when the initial condition is closed, the flow (1.1) preserves the closed condition and it evolves as a Ricci-like flow on  $\Omega^3$ . It allows using DeTurck's trick and, then, the Laplacian flow becomes parabolic in the direction of closed forms. In [BX11], Bryant and Xu addressed this approach in order to prove the short time existence of (1.1).

Motivated by Bryant and Xu ideas on the Laplacian flow of closed  $G_2$ -structures, Karigiannis, McKay and Tsui introduced the *Laplacian coflow* of coclosed  $G_2$ -structures in [KMT12]. It means that, instead of considering the heat flow equation for  $\varphi$ , they deal with the flow:

$$\frac{\partial \psi(t)}{\partial t} = \Delta_t \psi(t), \quad \psi(0) = \psi.$$
(1.2)

Equally to the Laplacian flow, if the initial condition satisfies  $d\psi = 0$ , the flow (1.2) preserves the coclosed condition. On one side, the Laplacian coflow is interesting, because coclosed G<sub>2</sub>-structures exist in any (compact and non-compact) spin and orientable 7-manifold by a parametric *h*-principle (see [CN15]). Unfortunately, the analytic approach employed for the Laplacian flow does not apply in the case (1.2), since it is not parabolic in the direction of the coclosed forms. Hence, the short-time existence of the Laplacian coflow is still an open problem. Nevertheless, in [Gri13], Grigorian proposed a modification of (1.2) fixing the failure of the Laplacian coflow to be parabolic, specifically the *modified Laplacian coflow* of coclosed G<sub>2</sub>-structures is the evolution given by

$$\frac{\partial \psi}{\partial t} = \Delta_t \psi(t) + 2d((A - \operatorname{tr}_{g(t)} T(t))\varphi(t)), \quad \text{for} \quad A > 0.$$
(1.3)

However, the critical points of (1.3) are no longer torsion-free G<sub>2</sub>-structures. For instance, if  $\varphi$  is a *nearly parallel* G<sub>2</sub>-structure, i.e.  $d\varphi = 4\psi$ , it vanishes the left hand side of (1.3) for A = 5. So, despite the fact that the modified Laplacian coflow can be seen as a tool for improving the torsion of  $\varphi$ , it does not search only for the torsion-free ones.

Regardless of the absence of an analytical theory of the Laplacian coflow in the general setting, the flow (1.2) had received the attention of some authors for manifolds with either a symmetry or an additional geometrical structure. For instance:

Assuming short time existence and uniqueness of (1.2), in [KMT12], Karigiannis, McKay and Tsui studied soliton solutions on warped products of a circle or an interval with a compact 6-manifold N with SU(3)-structure  $(\omega, Re(\Omega))$ . Running the Laplacian coflow among cohomogeneity-one solutions, when  $(N, \omega, Re(\Omega))$  is a Calabi-Yau manifold, they proved that the unique soliton solutions on the warped product are the steady ones. In particular, in the compact case, the soliton solutions are given by translations and phase rotations of the standard torsion-free  $G_2$ -structure.

On Sasakian manifolds  $(M, \xi, \eta, \Phi)$  with a contact Calabi-Yau structure  $(\omega := d\eta, \text{Re}(\Upsilon))$  in [LSES22], Lotay, Sá Earp and Saavedra proved the existence of a family of G<sub>2</sub>-structures by solving the Laplacian coflow, choosing  $\varepsilon \in \mathbb{R}^*$  and initial data  $\varphi = \varepsilon \eta \wedge \omega + \text{Re}(\Upsilon)$ , which is coclosed and the solution exists in  $t \in (-\frac{1}{10\varepsilon^2}, \infty)$ . Hence, the solution of the Laplacian coflow is immortal with a finite singularity at  $t = -\frac{1}{10\varepsilon^2}$ . It was the first example of compact solution to the Laplacian coflow which had an infinite time type *IIB* singularity.

On 3-Sasakian manifolds, there exists two non-equivalent nearly parallel  $G_2$ -structures [FKMS97]. Moreover, using the natural SU(2)-action, there is a 4-parameter family of coclosed  $G_2$ -structures (up to sign), which contains the nearly parallel ones. Under a special Ansatz of this family of coclosed  $G_2$ -structures, Kennon and Lotay proved that any solution of the Laplacian coflow starting at a coclosed  $G_2$ -structure converges, after rescaling, to one of the nearly parallel  $G_2$ -structure in the same family of the initial data [KL23]. In particular, the nearly parallel  $G_2$ -structures are both stable within their families.

In the other hand, when M = G/H is a homogeneous space and the solutions of (1.2) are required to be *G*-invariant, the Laplacian coflow becomes an ordinary differential equation. Namely, let  $\mathfrak{g}$  and  $\mathfrak{h}$  be the Lie algebras of *G* and *H* respectively, and  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$  a *reductive decomposition* (i.e.  $\operatorname{Ad}(H)$ -invariant), any *G*-invariant solution of (1.2) on *M* is determined by an  $\operatorname{Ad}(K)$ -invariant 4-form  $\psi(t)$  on  $\mathfrak{m} \simeq T_o M$  (where  $o = 1_G H$ ). Then, since  $\Delta \psi$  is invariant by diffeomorphisms of *M*, the flow (1.2) restricted to *G*-invariant solutions is equivalent with:

$$\frac{d}{dt}\psi(t) = \Delta_{\psi(t)}\psi(t) \quad \text{for} \quad \psi(t) \in \left(\Lambda^4\mathfrak{m}^*\right)^{\operatorname{Ad}(H)}.$$
(1.4)

Hence, short time existence and uniqueness of (1.4) are followed by the well known ODE arguments, since the linear map  $\Delta$  on  $\Lambda^4 \mathfrak{m}^*$  is continuous. For instance, in [KL21], Kath and Lauret obtained expanding solitons and immortal solutions of the Laplacian coflow when M is the connected and simply connected Lie group with Lie algebra  $\mathfrak{a} \ltimes \mathbb{R}^4$ , where  $\mathfrak{a}$  is any maximal  $\mathbb{R}$ -split torus of  $\mathfrak{sl}(\mathbb{R}^4)$ . The latest have been obtained using the bracket flow approach (see [Lau16] for a deep exposition of this method). Conversely, using a direct method, Bagaglini, Fernández and Fino obtained explicit immortal solutions of (1.4) when M is the 7-dimensional Heisenberg group [BFF20]. In [BF18], Bagaglini and Fino gave explicit immortal solutions and solitons of the Laplacian coflow for a subclass of almost Abelian Lie groups.

In this work, we study the Laplacian coflow of invariant coclosed  $G_2$ -structures. Intending to proceed with this, in Section 2, we provide some preliminaries on coclosed  $G_2$ -structures to establish the notation that is going to be used for the rest of the paper. In Section 3, we recall the definition of the Laplacian coflow of coclosed

G<sub>2</sub>-structures and its soliton solutions. Specifically for the parameter  $\lambda \in \mathbb{R}$  and the vector field  $X \in \mathscr{X}(M)$ , such that  $\varphi$  satisfies the soliton equation (3.3). We characterise the soliton condition in terms of the full torsion tensor T of  $\varphi$  by

$$\operatorname{div} T = -\frac{1}{2} (\operatorname{Curl} X)^{\flat} + X \lrcorner T \quad \text{and} \quad -\operatorname{Ric} + \frac{1}{2} T \circ T + (\operatorname{tr} T) T = \frac{\lambda}{4} g + \frac{1}{2} \mathcal{L}_X g. \tag{1.5}$$

where  $T \circ T$  and divT are defined in (2.11), and (2.12), respectively. Ric denotes the Ricci curvature induced by  $\varphi$  and  $(\operatorname{Curl} X)_c = \nabla_a X_b \varphi^{ab}{}_c$  denotes the curl of X. As an application of (1.5), we give in Corollary 3.1 an alternative proof for the non-existence of compact shriking solitons of the Laplacian coflow. Finally, in Section 4, we address the Laplacian coflow of invariant coclosed G<sub>2</sub>-structures on almost Abelian Lie groups  $G_A$ , with Lie algebra  $\mathfrak{g}$  and Lie bracket  $A \in \mathfrak{gl}(\mathbb{R}^6)$ . Using the bracket flow, we write the Laplacian coflow (1.4) in terms of the Lie bracket as

$$\frac{d}{dt}A = -\left(\frac{1}{2}\operatorname{tr}(S_A)^2 + \frac{1}{4}(\operatorname{tr}JA)^2\right)A + \frac{1}{2}[A, [A, A^t]] + \frac{1}{2}[A, S_A \circ_6 S_A],\tag{1.6}$$

where  $S_A$  denotes the symmetric part of A, J is the canonical almost complex structure on  $\mathbb{R}^6$  and the product  $S_A \circ_6 S_A$  is defined in (4.6). Hereby, we are able to prove:

**Theorem 1.1.** Let  $\mathfrak{g}$  be an almost Abelian Lie algebra with Lie bracket  $A \in \mathfrak{gl}(\mathbb{R}^6)$  and coclosed (non-flat)  $G_2$ -structure  $\varphi$ . Then, the solution  $\varphi(t)$  of the Laplacian coflow starting at  $\varphi$  is immortal, i,e, it is defined for all  $t \in (\varepsilon_1, \infty)$ .

In spite of not obtaining explicit solutions of (1.4) as it has been done in [BF18] for a subclass of almost Abelian Lie algebras, the Theorem (1.1) generalise the result of long-time existence of solutions for any almost Abelian Lie algebra. Moreover, the ODE (1.6) allows to study the dynamical behavior of the 2-parameter family

$$A = \begin{bmatrix} B & 0 \\ 0 & -B^t \end{bmatrix} \quad \text{with} \quad B = \begin{bmatrix} 0 & x & 0 \\ y & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad x, y \in \mathbb{R}$$

showing in Proposition 4.8 that the Laplacian coflow is stable. To conclude, we study the invariant solitons of the Laplacian coflow in terms of the Lie brackets  $A \in \mathfrak{gl}(\mathbb{R}^6)$ , satisfying the time independent equation (see Theorem 4.3)

$$[A, A^{t}] + S_{A} \circ_{6} S_{A} = -\left(\operatorname{tr} S_{A}^{2} - \frac{1}{2}(\operatorname{tr} JA)^{2} + 2d\right)I_{6} + (D + D^{t})|_{\mathbf{R}^{6}},$$
(1.7)

where D is a derivation of  $(\mathfrak{g}, A)$  and

$$d = \frac{|[A, A^t]|^2 + \langle S_A \circ_6 S_A, [A, A^t] \rangle}{2|A|^2}$$

As an application of (1.7), firstly we prove if A is skew-symmetric, then  $(\mathfrak{g}, A, \varphi)$  defines an invariant soliton of the Laplacian coflow (see Corollary 4.1). Secondly, we prove that any invariant (non-flat) soliton on an almost Abelian Lie group is an expanding solition (Proposition 4.10). Finally, as far as we know it, we provide the first example of a semi-algebraic soliton of the Laplacian coflow, which is no algebraic (Example 4.1).

#### Note

Fino and Bagaglini [BF18] have substantial overlap with this paper. However, while a number of conclusions are similar, the point of view on the Laplacian coflow is different. In this paper, we use the bracket flow introduced by Lauret in [Lau16], while in [BF18], a more traditional geometric flow approach is used. Both approaches are valuable and complementary, since they provide different perspectives on the same phenomenon. Since we are studying the same flow in the same space, we want to emphasise that this paper has different techniques, and both papers will give a better understanding of the Laplacian coflow.

#### Notation

Let (M, g) be a smooth oriented Riemannian 7-manifold. We use the Einstein summation convention throughout. We compute in a local orthonormal frame, so all indices are subscripts and any repeated indices are summed over all values from 1 to 7. A differential k-form  $\alpha$  on M will be written as

$$\alpha = \frac{1}{k!} \alpha_{i_1 i_2 \cdots i_k} \, \mathrm{d} x^{i_1} \wedge \mathrm{d} x^{i_2} \wedge \cdots \wedge \mathrm{d} x^{i_k}$$

in local coordinates  $(x^1, \ldots, x^7)$ , where  $\alpha_{i_1 i_2 \cdots i_k}$  is completely skew-symmetric in its indices. With this convention, the interior product  $\partial_m \lrcorner \alpha$  of  $\alpha$  with a coordinate vector field  $\partial_m$  is the (k-1)-form

$$\partial_m \lrcorner \alpha = \frac{1}{(k-1)!} \alpha_{mi_1 i_2 \cdots i_{k-1}} \, \mathrm{d} x^{i_1} \wedge \mathrm{d} x^{i_2} \wedge \cdots \wedge \mathrm{d} x^{i_{k-1}}.$$

The metric g on a Riemannian manifold M induces a metric on k-forms, such that the inner product of  $\alpha$  and  $\beta$  is

$$g(\alpha,\beta) = \frac{1}{k!} \alpha_{i_1 \cdots i_k} \beta_{j_1 \dots j_k} g^{i_1 j_1} \dots g^{i_k j_k}.$$

The Levi-Civita connection associated to g is denoted by  $\nabla$ , and its Christoffel symbols by  $\Gamma_{ij}^k$ . We write  $\nabla_i$  for covariant differentiation in the  $\partial_i$  direction. If  $T_{i_1\cdots i_k}$  is a tensor of type (0, k), then  $\nabla_m T_{i_1\cdots i_k}$  always means  $(\nabla_m T)_{i_1\cdots i_k}$ . We write the exterior derivative  $d\alpha$  of a k-form  $\alpha$  as

$$\mathrm{d}\alpha = \frac{1}{k!} \left( \nabla_m \alpha_{i_1 \cdots i_k} \right) \mathrm{d}x^m \wedge \mathrm{d}x^{i_1} \cdots \wedge \mathrm{d}x^{i_k}$$

in terms of the covariant derivative. The metric g defines an isomorphism between TM and  $T^*M$  (raising and lowering indices.) If v is a vector field, then the metric dual 1-form  $v^{\flat}$  is defined by  $v^{\flat}(w) = g(v, w)$ . In coordinates,  $(\partial_i)^{\flat} = g_{ik} dx^k$ . Similarly, the 1-form  $\alpha$  has a metric dual vector field  $\alpha^{\sharp}$ , and  $(dx^i)^{\sharp} = g^{ik}\partial_k$ .

We use 'vol' to denote the volume form on M associated to the metric g and an orientation. The Hodge star operator \* taking k-forms to (7 - k)-forms is defined by

$$\alpha \wedge *\beta = g(\alpha, \beta)$$
vol

Our convention for labelling the Riemann curvature tensor is

$$R_{ijkm}\frac{\partial}{\partial x^m} = \left(\nabla_i \nabla_j - \nabla_j \nabla_i\right) \frac{\partial}{\partial x^k}$$

in terms of coordinate vector fields. With this convention, the Ricci tensor is  $R_{jk} = R_{ljkl}$  and the first Bianchi identity of the Riemann curvature tensor is:

$$R_{abmn} + R_{amnb} + R_{anbm} = 0. ag{1.8}$$

We use  $\Gamma(E)$  to denote the space of smooth sections of E. As special instances, we denote the following cases as:

- $\Omega^k := \Gamma \left( \Lambda^k \left( T^* M \right) \right)$  is the space of smooth k-forms on M;
- $S := \Gamma(S^2(T^*M))$  is the space of smooth symmetric 2-tensors on M.
- $\mathscr{X}(M) := \Gamma(TM)$  the space of vector fields.

With respect to the metric g on M, we use  $S_0$  to denote those sections h of S that are traceless. That is,  $S_0$  consists of those sections of S, such that  $\operatorname{Tr} h = g^{ij}h_{ij} = 0$  in local coordinates. Then  $S \simeq \Omega^0 \oplus S_0$ , where  $h \in S$  is decomposed as  $h = \frac{1}{7}(\operatorname{Tr} h)g + h_0$ . Then, we have  $\Gamma(T^*M \otimes TM) = \Omega^0 \oplus S_0 \oplus \Omega^2$ , where the splitting is pointwise orthogonal with respect to the metric on  $T^*M \otimes TM$  induced by g.

**Acknowledgements:** The authors would like to thank Jorge Lauret for introducing them to the idea of the bracket flow in this context, and also the Universidad Nacional de Cordoba for hosting that conversation in 2019. Also, we are grateful to Henrique Sá Earp for the meaningful discussions and advises. This work stems on the MATHAMSUD Regional Program 21-MATH-06 collaborations. AM was funded by the Sao Paulo Research Foundation (Fapesp) [2021/08026-5] and JPS was supported by the Coordination for the Improvement of Higher Education Personnel-Brazil (CAPES) [88887.648550/2021-00].

# 2 Preliminaries

In this section we collect some results related to G<sub>2</sub>-structures that will be needed in the present paper. Any result of this section can be found in [Kar07, Gri13, Bry06].

### **2.1** G<sub>2</sub>-structures and their torsion

A G<sub>2</sub>-structure on a 7-manifold M is given by a differential 3-form  $\varphi$  on M, which is pointwise isomorphic to the 3-form

$$\varphi_0 = e^{123} + e^{145} + e^{167} + e^{246} - e^{257} - e^{347} - e^{356} \in \Lambda^3(\mathbb{R}^7)^*,$$

where  $e^{ijk} = e^i \wedge e^j \wedge e^k$  and  $\{e^1, \ldots, e^7\}$  is the dual basis of the canonical basis of  $\mathbb{R}^7$ . The G<sub>2</sub>-structure  $\varphi$  determines a Riemannian metric  $g_{\varphi}$  and a volume form  $\operatorname{vol}_{\varphi}$  so that

$$6g_{\varphi}(X,Y)\mathrm{vol}_{\varphi} = (X \lrcorner \varphi) \land (Y \lrcorner \varphi) \land \varphi \quad \text{for} \quad X,Y \in \mathscr{X}(M).$$

In addition,  $\varphi$  induces a Hodge star operator  $*_{\varphi}$  and we denote its dual 4-form by  $\psi = *_{\varphi}\varphi$ . For simplicity, we will write  $g = g_{\varphi}$  and  $* = *_{\varphi}$ . A G<sub>2</sub>-structure gives rise to a decomposition of the space of differential k-forms  $\Omega^k$  on M into irreducible G<sub>2</sub>-submodules. For instance,

$$\Omega^2 = \Omega_7^2 \oplus \Omega_{14}^2 \quad \text{and} \quad \Omega^3 = \Omega_1^3 \oplus \Omega_7^3 \oplus \Omega_{27}^3, \tag{2.1}$$

where  $\Omega_l^k$  has (pointwise) dimension *l*. In [Bry06], R. Bryant defines an injective map  $i_{\varphi} : S^2 \to \Omega^3$ , given in local coordinates  $x^1, \ldots x^7$  by

$$i_{\varphi}(h) = \frac{1}{3!} i_{\varphi}(h)_{ijk} dx^{ijk} = \frac{1}{3!} (h_i^m \varphi_{mjk} + h_j^m \varphi_{imk} + h_k^m \varphi_{ijm}) dx^{ijk},$$
(2.2)

where  $h \in S^2$  is a symmetric 2-tensor field on M. Additionally, the map  $i_{\varphi}$  is surjective on  $\Omega_1^3 \oplus \Omega_{27}^3$  and its Hodge dual satisfies (e.g. [Kar07, Proposition 2.8])

$$* i_{\varphi}(h) = \frac{1}{4!} (\bar{h}_{i}^{m} \psi_{mjkl} + \bar{h}_{j}^{m} \psi_{imkl} + \bar{h}_{k}^{m} \psi_{ijml} + \bar{h}_{l}^{m} \psi_{ijkm}) dx^{ijkl} =: i_{\psi}(\bar{h}),$$
(2.3)

where  $\bar{h} = \frac{1}{4} \operatorname{tr}(h)g - h$ . In particular, for any trace-free symmetric 2-tensors  $h \in S_0^2$ , we have  $i_{\varphi}(h) \in \Omega_{27}^3$  and  $i_{\psi}(h) \in \Omega_{27}^4 = *(\Omega_{27}^3)$ . According with the G<sub>2</sub>-decomposition of  $\Omega^4$  and  $\Omega^5$ , the exterior derivative of  $\varphi$  and  $\psi$  are completely described in term of the *torsion forms*  $\tau_0 \in \Omega^0$ ,  $\tau_1 \in \Omega^1$ ,  $\tau_2 \in \Omega_{14}^2$  and  $\tau_3 \in \Omega_{27}^3$ , given by (see [Bry06, Proposition 1])

$$d\varphi = \tau_0\psi + 3\tau_1 \wedge \varphi + *\tau_3 \in \Omega_1^4 \oplus \Omega_7^4 \oplus \Omega_{27}^4 \quad \text{and} \quad d\psi = 4\tau_1 \wedge \psi + \tau_2 \wedge \varphi \in \Omega_7^5 \oplus \Omega_{14}^5.$$
(2.4)

Moreover, for the *full torsion tensor* T of  $\varphi$ , which is defined locally by (see [Kar07])

$$\nabla_i \varphi_{jkl} = T_i^m \psi_{mjkl},\tag{2.5}$$

we may use the torsion forms to relate them with T, by

$$T = \frac{\tau_0}{4}g - \tau_{27} - \tau_1^{\sharp} \Box \varphi - \frac{1}{2}\tau_2, \qquad (2.6)$$

where  $\tau_{27}$  is the trace-free symmetric 2-tensor satisfying  $\tau_3 = i_{\varphi}(\tau_{27})$  and  $\tau_1^{\sharp}$  denotes the unique vector field induced by the Riemannian metric g, (i.e.  $g(\tau_1^{\sharp}, X) = \tau_1(X)$  for any  $\in \mathscr{X}(M)$ ). In addition, from (2.5) for the 4-form  $\psi$ , we have

$$\nabla_m \psi_{ijkl} = -(T_{mi}\varphi_{jkl} - T_{mj}\varphi_{ikl} - T_{mk}\varphi_{jil} - T_{ml}\varphi_{jki}).$$
(2.7)

#### **2.2** Properties of coclosed G<sub>2</sub>-structures

A G<sub>2</sub>-structure  $\varphi$  is *coclosed* if it satisfies  $d\psi = 0$ , in terms of (2.4) the coclosed condition is equivalent with  $\tau_1 = 0$  and  $\tau_2 = 0$ . Hence, the full torsion tensor of a coclosed G<sub>2</sub>-structure simplifies to the symmetric 2-tensor

$$T = \frac{\tau_0}{4}g - \tau_{27} \in \mathcal{S}^2.$$
(2.8)

In addition,  $d\varphi \in \Omega_1^4 \oplus \Omega_{27}^4$  thus, by (2.2), (2.4) and (2.8), we have

$$d\varphi = *i_{\varphi} \left(\frac{1}{3}(\operatorname{tr} T)g - T\right).$$
(2.9)

The following proposition include some well known identities of coclosed  $G_2$ -structures given in [Gri13], obtained as consequence of a general formulae of the exterior derivative of a generic 3-form. Here, we give an alternative proof of those identities, using the called  $G_2$ -*Bianchi type identity* 

$$\nabla_i T_{jk} - \nabla_j T_{ik} = \left(\frac{1}{2}R_{ijmn} - T_{im}T_{jn}\right)\varphi_k^{mn},\tag{2.10}$$

where  $T_{ij}$  is the coordinate of (2.8) and  $R_{ijmn}$  denotes the Riemann curvature tensor. We remark that the identity (2.10) can be read as the infinitesimal version of the diffeomorphism invariance of T as a function  $\varphi$ . (see [Kar07, Section 4] for an extensive discussion in the G<sub>2</sub>-case and [FLMSE22] for any H-structure.). In the statement, for any  $h, k \in S^2$ , we denote the inner product  $\langle h, k \rangle$  and the circ product  $h \circ k \in S^2$  by

$$\langle h, k \rangle = h_{ij} k_{ab} g^{ia} g^{jb}$$
 and  $(h \circ k)_{ab} = \varphi_{amn} \varphi_{bpq} h^{mp} k^{nq}$ . (2.11)

As well as the divergence and the curl of h, given in coordinates by

$$\operatorname{div} h_a = \nabla_b h_a^b$$
 and  $\operatorname{Curl} h_{ab} = \nabla_m h_{an} \varphi_b^{mn}$ . (2.12)

**Proposition 2.1.** Let  $\varphi$  be a coclosed G<sub>2</sub>-structure with full torsion tensor *T*, then the divergence and the curl of *T* satisfy we have

$$\operatorname{div} T_a = \nabla_a \operatorname{tr} T \quad and \quad \operatorname{Curl} T_{ab} = \operatorname{Curl} T_{ba}. \tag{2.13}$$

In addition, the Ricci tensor and the scalar curvature are

$$Ric = -CurlT - T^{2} + (trT)T \quad and \quad R = (trT)^{2} - |T|^{2}.$$
(2.14)

*Proof.* Using (1.8) and the symmetries of  $R_{abmn}$  is easy to prove that

$$R_{abmn}\varphi^{bmn} = 0 \quad \text{and} \quad R_{amnp}\psi_b{}^{mnp} = 0.$$
(2.15)

Now, since T is symmetric, using (2.10) and (2.15) for the divergence T, we have

$$\operatorname{div} T_a = \nabla_b T_a^b = \nabla_a T_b^b + \left(\frac{1}{2}R_{bamn} - T_{am}T_{bn}\right)\varphi^{bmn} = \nabla_a \operatorname{tr} T,$$

and in addition, by (A.3) for the curl of T, we get

$$\begin{aligned} \operatorname{Curl} T_{ab} - \operatorname{Curl} T_{ba} = & \nabla_m T_{an} \varphi_b{}^{mn} - \nabla_m T_{bn} \varphi_a{}^{mn} \\ &= \left(\frac{1}{2} R_{mapq} - T_{mp} T_{aq}\right) \varphi_n{}^{pq} \varphi_b{}^{mn} - \left(\frac{1}{2} R_{mbpq} - T_{mp} T_{bq}\right) \varphi_n{}^{pq} \varphi_a{}^{mn} \\ &= \left(\frac{1}{2} R_{mapq} - T_{mp} T_{aq}\right) \left(g_b^p g^{qm} - g_b^q g^{pm} + \psi_b{}^{mpq}\right) \\ &- \left(\frac{1}{2} R_{mbpq} - T_{mp} T_{bq}\right) \left(g_a^p g^{qm} - g_a^q g^{pm} + \psi_a{}^{mpq}\right) \\ &= \frac{1}{2} R_{mabq} g^{mq} - \frac{1}{2} R_{mapb} g^{mp} - T_{am} T_b^m + \operatorname{tr}(T) T_{ab}\end{aligned}$$

$$-\frac{1}{2}R_{mbaq}g^{mq} + \frac{1}{2}R_{mbpa}g^{mp} + T_{bm}T_a^m - \operatorname{tr}(T)T_{ba}$$
$$= -\operatorname{Ric}_{ab} + \operatorname{Ric}_{ba} = 0.$$

The formula for Ric can be derived from the computation above and for the scalar curvature, it follow from the observation

$$\operatorname{Curl} T_{aa} = \nabla_m T_{an} \varphi_a{}^{mn} = 0.$$

Similar to [Bry06, Corollary 2] for the case of closed  $G_2$ -structures, we can characterise the Einstein metrics induced by a coclosed  $G_2$ -structure:

**Corollary 2.1.** A coclosed  $G_2$ -structure  $\varphi$  induces an Einstein metric if and only if the full torsion tensor satisfies

$$\mathbf{i}_{\varphi}(\operatorname{Curl} T) = \frac{3}{7} |T|^2 \varphi - (\operatorname{tr} T) \tau_3 - \mathbf{i}_{\varphi}(T^2).$$
(2.16)

*Proof.* The result follows by applying the map  $i_{\varphi}$  in (2.14).

*Remark* 2.1. Using the expression of the full torsion tensor in terms of the torsion forms (2.8), the equation (2.16) becomes

$$i_{\varphi}(\operatorname{Curl}\tau_{27}) = \frac{3}{7} |\tau_{27}|^2 \varphi - \frac{5\tau_0}{4} \tau_3 - i_{\varphi}(\tau_{27}^2).$$
(2.17)

It is well know that a metric induced by the nearly parallel G<sub>2</sub>-structure (i.e.  $\tau_3 = i_{\varphi}(\tau_{27}) = 0$ ) is Einstein. It is easy to check that (2.17) is satisfied trivially for a nearly G<sub>2</sub>-structure.

### **3** Laplacian coflow of G<sub>2</sub>-structures

In this section, we recall the definition of the Laplacian coflow and we also study soliton solutions and symmetries of coclosed G<sub>2</sub>-structure. For these we follow [KMT12, Gri13].

**Definition 3.1.** A time-dependent G<sub>2</sub>-structure  $\{\varphi(t)\}_{t \in (\varepsilon_1, \varepsilon_2)}$  on a 7-manifold M, satisfies the Laplacian coflow of coclosed G<sub>2</sub>-structures, if for any  $t \in (\varepsilon_1, \varepsilon_2)$  we have

$$\frac{\partial}{\partial t}\psi(t) = \Delta_t\psi(t) \quad \text{and} \quad \mathrm{d}\psi(t) = 0,$$
(3.1)

where  $\psi(t) = *_t \varphi(t)$  and  $\Delta_t = dd^{*t} + d^{*t} d$  is the Hodge Laplacian with respect to the metric  $g(t) = g_{\varphi(t)}$ .

As for many geometric flows, we are interested in considering self-similar solutions

$$\varphi(t) = \lambda(t)f(t)^*\varphi$$
 where  $\lambda(t) \in C^{\infty}(M)$  and  $f(t) \in \text{Diff}(M)$ , (3.2)

it means, solutions that evolves the initial data  $\varphi$  by diffeomorphisms and scalings, since these kind of solutions are expected to be related to singularities of the flow. In particular, self-similar solutions with initial condition  $\varphi$  are equivalent with a time independent equation of  $\psi = *\varphi$ , called the *soliton equation*, namely,  $\varphi$  is called a *soliton* for the Laplacian coflow (3.1), if  $\psi$  satisfies the soliton equation:

$$\Delta_{\psi}\psi = \mathcal{L}_{X}\psi + \lambda\psi \tag{3.3}$$

where  $\lambda \in \mathbb{R}$  and X is a complete vector field on M. Moreover, the soliton  $(\varphi, \lambda, X)$  is called *expanding, steady*, or *shrinking*, if  $\lambda > 0$ ,  $\lambda = 0$  or  $\lambda < 0$ , respectively.

The following lemma, decomposes the Hodge Laplacian of  $\psi$  according to the G<sub>2</sub>-irreducible decomposition of  $\Omega^2$ , it appeared originally in [Gri13, Proposition 4.6]. Here, we provide the computations in detail for the self-contained of the work, for it we follow the computation given in [LW17] for  $\Delta_{\varphi}\varphi$  in the closed case.

**Lemma 3.1.** Let  $\varphi$  be a coclosed  $G_2$ -structure on a manifold M with associated metric g. Then,

$$\begin{aligned} \Delta_{\psi}\psi &= \frac{2}{7}((\operatorname{tr} T)^{2} + |T|^{2})\psi \oplus (\operatorname{d}\operatorname{tr} T) \wedge \varphi \\ &\oplus *_{\varphi}\mathrm{i}_{\varphi}\Big(\operatorname{Ric} - \frac{1}{2}T \circ T - (\operatorname{tr} T)T + \frac{1}{14}\left((\operatorname{tr} T)^{2} + |T|^{2}\right)g\Big) \in \Omega_{1}^{4} \oplus \Omega_{7}^{4} \oplus \Omega_{27}^{4}. \end{aligned}$$

*Proof.* Since  $d\psi = 0$ , by (2.9) we have

$$\Delta_{\psi}\psi = \mathrm{dd}^{*}\psi = \mathrm{d} \ast \mathrm{d}\varphi = \mathrm{d}\beta \quad \text{where} \quad \beta := \mathrm{i}_{\varphi}(h) = \mathrm{i}_{\varphi}\left(\frac{1}{3}(\mathrm{tr}\,T)g - T\right) \in \Omega^{3}_{27}. \tag{3.4}$$

In local coordinates, we can write (3.4) as

$$\Delta_{\psi}\psi = \frac{1}{4!} (\Delta_{\psi}\psi)_{ijkl} dx^{ijkl},$$

where

$$(\Delta_{\psi}\psi)_{ijkl} = \nabla_i\beta_{jkl} - \nabla_j\beta_{ikl} + \nabla_k\beta_{ijl} - \nabla_l\beta_{ijk}.$$
(3.5)

We can decomposes  $\Delta_{\psi}\psi$  into irreducible summands as

$$\Delta_{\psi}\psi = a\psi + X^{\flat} \wedge \varphi + *\mathbf{i}_{\varphi}(s), \tag{3.6}$$

where a is a function, X a vector field and s is a trace-less symmetric 2-tensor. Now, we compute the expression of a, X and s in terms of the full torsion tensor of  $\varphi$ . For a, using (3.4), (3.5), (A.4) and (A.6), we have

$$\begin{split} a &= \frac{1}{7} \langle \Delta_{\psi}, \psi \rangle = \frac{1}{168} (\nabla_{i} \beta_{jkl} - \nabla_{j} \beta_{ikl} + \nabla_{k} \beta_{ijl} - \nabla_{l} \beta_{ijk}) \psi^{ijkl} \\ &= \frac{1}{42} \nabla_{i} (h_{j}^{m} \varphi_{mkl} + h_{k}^{m} \varphi_{jml} + h_{l}^{m} \varphi_{jkm}) \psi^{ijkl} \\ &= \frac{1}{14} (\nabla_{i} h_{j}^{m} \varphi_{mkl} + h_{j}^{m} T_{i}^{n} \psi_{nmkl}) \psi^{ijkl} \\ &= \frac{2}{7} (\nabla_{i} h_{j}^{m} \varphi_{m}^{ij} + \operatorname{tr} h \operatorname{tr} T - \langle h, T \rangle) \\ &= \frac{2}{7} \left( (\operatorname{tr} T)^{2} + |T|^{2} \right), \end{split}$$

where h is the symmetric 2-tensor given in (3.4). For the vector field X, we have

$$\langle \Delta_{\psi}\psi, e^m \wedge \varphi \rangle = *(X^{\flat} \wedge \varphi \wedge *(e^m \wedge \varphi)) = 4\langle X^{\flat}, e^m \rangle = 4X_n g^{nm}.$$

Thus, using (3.4), (A.3), (A.2), (A.4) and (2.13), we get

$$\begin{split} X_m &= \frac{1}{4} \langle \Delta_{\psi} \psi, e^n \wedge \varphi \rangle g_{mn} \\ &= \frac{1}{98} (\nabla_i \beta_{jkl} - \nabla_j \beta_{ikl} + \nabla_k \beta_{ijl} - \nabla_l \beta_{ijk}) (e^n \wedge \varphi)^{ijkl} g_{mn} \\ &= \frac{1}{4!} (\nabla_m \beta_{jkl} \varphi^{jkl} - 3 \nabla_j \beta_{mkl} \varphi^{jkl}) \\ &= \frac{1}{4!} (\nabla_m (\beta_{jkl} \varphi^{jkl}) - \beta_{jkl} \nabla_m \varphi^{jkl} - 3 \nabla_j (\beta_{mkl} \varphi^{jkl}) + 3 \beta_{mkl} \nabla_j \varphi^{jkl}) \\ &= \frac{1}{4!} \left( 3 \nabla_m (h_j^n \varphi_{njk} \varphi^{jkl}) - 3 h_j^n \varphi_{nkl} T_{mp} \psi^{pjkl} - 3 \nabla_j (h_m^n \varphi_{nkl} \varphi^{jkl} + 2 h_k^n \varphi_{mnl} \varphi^{jkl}) \right) \\ &= \frac{1}{8} \left( 6 \nabla_m h_j^n g_n^j - 4 h_j^n T_{mp} \varphi_n^{pj} - 6 \nabla_j h_m^j - 2 \nabla_m h_k^n g_n^k + 2 \nabla_n h_m^n \right) \\ &= \frac{1}{2} \left( \frac{4}{3} \nabla_m (\operatorname{tr} T) - \frac{1}{3} \nabla_m (\operatorname{tr} T) + \nabla_j T_m^j \right) = (\operatorname{div} T)_m. \end{split}$$

Finally, to find the symmetric 2-tensor s, we have:

$$(\Delta_{\psi}\psi)_{imnp}\psi_{j}^{mnp} + (\Delta_{\psi}\psi)_{jmnp}\psi_{i}^{mnp} = a(\psi_{imnp}\psi_{j}^{mnp} + \psi_{jmnp}\psi_{i}^{mnp}) + (*i_{\varphi}(s))_{imnp}\psi_{j}^{mnp} + (*i_{\varphi}(s))_{jmnp}\psi_{i}^{mnp},$$
(3.7)

Then, using (2.3),(A.6) and (A.7), we get

$$(*i_{\varphi}(s))_{imnp}\psi_j^{mnp} = -s_i^q\psi_{qmnp}\psi_j^{mnp} - 3s_m^q\psi_{iqnp}\psi_j^{mnp}$$

$$= -24s_i^q g_{qj} - 3s_m^q (4g_{ij}g_q^m - 4g_i^m g_{qj} + 2\psi_{iqj}^m) = -12s_{ij}$$

By symmetry, the right hand side of (3.7) becomes

$$(\Delta_{\psi}\psi)_{imnp}\psi_{j}^{mnp} + (\Delta_{\psi}\psi)_{jmnp}\psi_{i}^{mnp} = 24\left(2ag_{ij} - s_{ij}\right).$$
(3.8)

Now, using (3.5), (A.4), (A.4), (A.5) and (A.6), we have

$$\begin{split} (\Delta_{\psi}\psi)_{imnp}\psi_{j}^{mnp} =& (\nabla_{i}\beta_{mnp} - 3\nabla_{m}\beta_{inp})\psi_{j}^{mnp} \\ =& 3(\nabla_{i}h_{m}^{q}\varphi_{qnp} + h_{m}^{q}\nabla_{i}\varphi_{qnp})\psi_{j}^{mnp} - 3\nabla_{m}((h_{i}^{q}\varphi_{qnp} + 2h_{n}^{q}\varphi_{iqp})\psi_{j}^{mnp}) \\ & + 3(h_{i}^{q}\varphi_{qnp} + 2h_{n}^{q}\varphi_{iqp})\nabla_{m}\psi_{j}^{mnp} \\ =& 3\left(4\nabla_{i}h_{m}^{q}\varphi_{qj}^{m} + h_{m}^{q}T_{i}^{l}(4g_{lj}g_{q}^{m} - 4g_{l}^{m}g_{qj} + 2\psi_{lqj}^{m}) - 4\nabla_{m}(h_{i}^{q}\varphi_{qj}^{m}) \right. \\ & + 2\nabla_{m}(h_{i}^{q}\varphi_{qj}^{m} - h_{nj}\varphi_{i}^{mn} - h_{n}^{m}\varphi_{ji}^{n} - \operatorname{tr}h\varphi_{ij}^{m}) \\ & + (h_{i}^{q}\varphi_{qnp} + 2h_{n}^{q}\varphi_{iqp})(-T_{mj}\varphi^{mnp} + \operatorname{tr}T\varphi_{j}^{np} - T_{m}^{n}\varphi_{j}^{mp} + T_{m}^{p}\varphi_{j}^{mn})) \\ =& 6\left(2(\operatorname{tr}hT_{ij} - T_{i}^{m}h_{mj}) - \nabla_{m}h_{i}^{q}\varphi_{qj}^{m} - \nabla_{m}h_{nj}\varphi_{i}^{mn} - \nabla_{m}(h_{n}^{m}\varphi_{ji}^{n} + \operatorname{tr}h\varphi_{ij}^{m}) \right. \\ & - 3h_{i}^{m}T_{mj} - \operatorname{tr}hT_{ij} + h_{i}^{m}T_{mj} + 3\operatorname{tr}Th_{ij} + \operatorname{tr}T\operatorname{tr}h_{gj} - \operatorname{tr}Th_{ij} - \operatorname{tr}Th_{ij} + h_{i}^{m}T_{mj} \right. \\ & - T_{m}^{n}h_{m}^{m}g_{ij} + T_{i}^{m}h_{mj} - (T \circ h)_{ij}) \\ =& 6\left(\operatorname{tr}hT_{ij} - T_{i}^{m}h_{mj} - (\operatorname{Curl}h)_{ij} - (\operatorname{Curl}h)_{ji} - \nabla_{m}(h_{n}^{m}\varphi_{ji}^{n} + \operatorname{tr}h\varphi_{ij}^{m}) \right. \\ & -h_{i}^{m}T_{mj} + \operatorname{tr}Th_{ij} + (\operatorname{tr}T\operatorname{tr}h - \langle T, h \rangle)g_{ij} - (T \circ h)_{ij}) \,. \end{split}$$

Thus, replacing  $h = \frac{1}{3}(\operatorname{tr} T)g - T$  in the above expression and using (2.13), the left hand side of (3.7) becomes

$$(\Delta_{\psi}\psi)_{imnp}\psi_{j}{}^{mnp} + (\Delta_{\psi}\psi)_{jmnp}\psi_{i}{}^{mnp} = 24\left(T_{i}^{m}T_{mj} + (\operatorname{Curl}T)_{ij} + \frac{1}{2}((\operatorname{tr}T)^{2} + |T|^{2})g_{ij} + \frac{1}{2}(T\circ T)_{ij}\right).$$

Finally, from (3.8), we obtain

$$s_{ij} = -(\operatorname{Curl}T)_{ij} - T_i^m T_{mj} - \frac{1}{2}(T \circ T)_{ij} + \frac{1}{14}((\operatorname{tr}T)^2 + |T|^2)g_{ij}$$

Similar to the Laplacian of  $\psi$ , we can compute the decomposition of the Lie derivative with respect to any vector field. We recall that the vector field X is called an *infinitesimal symmetry* of  $\psi$ , if  $\mathcal{L}_X \psi = 0$ . The next result was done in [DGK21] for the 3-form  $\varphi$ .

**Proposition 3.2.** Let  $\varphi$  be a coclosed G<sub>2</sub>-structure on  $M^7$ , with associated metric g, and let X be a vector field on M. Then, if  $\psi = *\varphi$ ,

$$\mathcal{L}_X \psi = \frac{4}{7} (\operatorname{div} X) \psi \oplus (-\frac{1}{2} \operatorname{Curl} X + X \lrcorner T)^\flat \land \varphi \oplus \ast i_\varphi \left(\frac{1}{7} (\operatorname{div} X)g - \frac{1}{2} (\mathcal{L}_X g)\right) \in \Omega_1^4 \oplus \Omega_7^4 \oplus \Omega_{27}^4.$$
(3.9)

In particular, X is an infinitesimal symmetry of  $\psi$  if and only if X is a Killing vector field of g and satisfies  $\operatorname{Curl}(X) = 2X \lrcorner T$ .

*Proof.* Since  $\varphi$  is coclosed, i.e.  $d\psi = 0$ , we have

$$\mathcal{L}_X \psi = \mathrm{d}(X \lrcorner \psi) + X \lrcorner \mathrm{d}\psi = \mathrm{d}(X \lrcorner \psi).$$

Let  $\alpha = X \lrcorner \psi$ , so that locally  $\alpha_{ijk} = X^l \psi_{lijk}$  and

$$(\mathcal{L}_X\psi)_{ijkl} = (\mathrm{d}\alpha)_{ijkl} = \nabla_i \alpha_{jkl} - \nabla_j \alpha_{ikl} + \nabla_k \alpha_{ijl} - \nabla_l \alpha_{ijk}.$$

Denoting by  $\pi_l^k: \Omega^k \to \Omega_l^k$  the orthogonal projections, we decompose  $\mathcal{L}_X \psi$  as

$$\mathcal{L}_X \psi = \pi_1^4(\mathcal{L}_X \psi) + \pi_7^4(\mathcal{L}_X \psi) + \pi_{27}^4(\mathcal{L}_X \psi) = a\psi + W^\flat \wedge \varphi + *i_\varphi(h), \tag{3.10}$$

where  $a \in \Omega^0$ , and h is a trace-free symmetric 2-tensor on M. We compute a as follows:

$$a = \frac{1}{7} \langle \mathcal{L}_X \psi, \psi \rangle = \frac{1}{168} (\nabla_i \alpha_{jkl} - \nabla_j \alpha_{ikl} + \nabla_k \alpha_{ijl} - \nabla_l \alpha_{ijk}) \psi^{ijkl}$$
  
$$= \frac{1}{42} \nabla_i \alpha_{jkl} \psi^{ijkl} = \frac{1}{42} \nabla_i (\alpha_{jkl} \psi^{ijkl}) - \frac{1}{42} \alpha_{jkl} \nabla_i \psi^{ijkl}$$
  
$$= \frac{24}{42} \nabla_i (X^m g_{mi}) - \frac{1}{42} X^m \psi_{mjkl} (\nabla_i \psi^{ijkl}) = \frac{4}{7} \nabla_i X_i = \frac{4}{7} \text{div} X,$$
  
(3.11)

where we used (A.4) and because T is symmetric. To compute  $W^{\flat}$ , note that

$$\langle *((*\mathcal{L}_X\psi)\wedge\varphi), e^m\rangle = 4\langle W^\flat, e^m\rangle,$$

thus

$$4W^m = \ast ((\ast \mathcal{L}_X \psi) \land \varphi \land e^m) = \langle \varphi \land e^m, \mathcal{L}_X \psi \rangle = \langle \varphi \land e^m, d\alpha \rangle$$

Therefore, we obtain

$$W^{m} = \frac{1}{4} \langle \varphi \wedge e^{m}, d\alpha \rangle = \frac{1}{4!} (\nabla^{i} \alpha^{jkm} - \nabla^{j} \alpha^{ikm} + \nabla^{k} \alpha^{ijm} - \nabla^{m} \alpha^{ijk}) \varphi_{ijk}$$

$$= \frac{1}{4!} (3\nabla^{i} \alpha^{jkm} \varphi_{ijk} - \nabla^{m} \alpha^{ijk} \varphi_{ijk})$$

$$= \frac{3}{4!} \nabla^{i} (\alpha^{jkm} \varphi_{ijk}) - \frac{3}{4!} \alpha^{jkm} \nabla^{i} \varphi_{ijk} - \frac{1}{4!} \nabla^{m} (\alpha^{ijk} \varphi_{ijk}) + \frac{1}{4!} \alpha^{ijk} \nabla^{m} \varphi_{ijk}$$

$$= \frac{3}{4!} \nabla^{i} (X_{l} \psi^{ljkm} \varphi_{ijk}) - \frac{3}{4!} X_{l} \psi^{ljkm} T_{n}^{i} \psi_{ijk}^{n} - \frac{1}{4!} \nabla^{m} (X_{l} \psi^{lijk} \varphi_{ijk}) + \frac{1}{4!} X_{l} \psi^{lijk} T_{n}^{m} \psi_{ijk}^{n}$$

$$= -\frac{1}{2} \nabla^{i} (X_{l} \varphi_{i}^{lm}) + X_{l} T^{ml} = -\frac{1}{2} (\nabla^{i} X_{l} \varphi_{i}^{lm} + X_{l} \nabla_{i} \varphi_{i}^{lm}) + X_{l} T^{ml}$$

$$= -\frac{1}{2} \operatorname{Curl} X^{m} - \frac{1}{2} X_{l} T_{i}^{n} \psi_{ni}^{lm} + (X \Box T)^{m} = -\frac{1}{2} (\operatorname{Curl} X)^{m} + (X \Box T)^{m}$$
(3.12)

Finally, to compute h, observe that

$$(\mathcal{L}_X \psi)_{imnp} \psi_j^{mnp} + (\mathcal{L}_X \psi)_{jmnp} \psi_i^{mnp} = a(\psi_{imnp} \psi_j^{mnp} + \psi_{jmnp} \psi_i^{mnp}) + (*i_{\varphi}(h))_{imnp} \psi_j^{mnp} + (*i_{\varphi}(h))_{jmnp} \psi_i^{mnp},$$
(3.13)

where

$$(*i_{\varphi}(h))_{imnp} = -(h_i^q \psi_{qmnp} + h_m^q \psi_{iqnp} + h_n^q \psi_{imqp} + h_p^q \psi_{imnq})$$

Using (A.6) and (A.7), we get

$$(*i_{\varphi}(h))_{imnp}\psi_{j}^{mnp} = -h_{i}^{q}\psi_{qmnp}\psi_{j}^{mnp} - 3h_{m}^{q}\psi_{iqnp}\psi_{j}^{mnp} = -24h_{i}^{q}g_{qj} - 3h_{m}^{q}(4g_{ij}g_{q}^{m} - 4g_{i}^{m}g_{qj} + 2\psi_{iqj}^{m}) = -12h_{ij}$$

By symmetry, the right hand side of (3.13) becomes

$$(\mathcal{L}_X\psi)_{imnp}\psi_j{}^{mnp} + (\mathcal{L}_X\psi)_{jmnp}\psi_i{}^{mnp} = 24\left(\frac{8}{7}(\operatorname{div} X)g_{ij} - h_{ij}\right)$$
(3.14)

For the left-hand side of (3.14), using the identities (A.5), (A.4), (A.6) and (A.7), we have:

$$\begin{aligned} (\mathcal{L}_{X}\psi)_{imnp}\psi_{j}{}^{mnp} = &\nabla_{i}\alpha_{mnp}\psi_{j}{}^{mnp} - 3\nabla_{m}\alpha_{inp}\psi_{j}{}^{mnp} \\ = &\nabla_{i}(\alpha_{mnp}\psi_{j}{}^{mnp}) - \alpha_{mnp}\nabla_{i}\psi_{j}{}^{mnp} - 3\nabla_{m}(\alpha_{inp}\psi_{j}{}^{mnp}) + 3\alpha_{inp}\nabla_{m}\psi_{j}{}^{mnp} \\ = &24\nabla_{i}X_{j} - 12T_{i}^{m}(X \lrcorner \varphi)_{mj} + 12(\operatorname{div} X)g_{ij} - 12\nabla_{i}X_{j} - 6(\nabla_{m}X \lrcorner \psi)_{ij}{}^{m} \\ &- 6\operatorname{tr}(T)(X \lrcorner \varphi)_{ij} + 6(X \lrcorner T)_{m}\varphi^{m}{}_{ij} + 6T_{i}^{m}(X \lrcorner \varphi)_{mj} - 6T_{j}^{m}(X \lrcorner \varphi)_{mi} \\ &- 12\operatorname{tr}(T)(X \lrcorner \varphi)_{ij} + 6T_{j}^{m}(X \lrcorner \varphi)_{mi} + 6T_{i}^{m}(X \lrcorner \varphi)_{mj} + 6(X \lrcorner T)_{m}\varphi^{m}{}_{ij} \\ = &12\nabla_{i}X_{j} + 12(\operatorname{div} X)g_{ij} - 6(\nabla_{m}X \lrcorner \psi)_{ij}{}^{m} - 18\operatorname{tr}(T)(X \lrcorner \varphi)_{ij} + 12(X \lrcorner T)_{m}\varphi^{m}{}_{ij} \end{aligned}$$

By symmetry, we get

$$(\mathcal{L}_X\psi)_{imnp}\psi_j^{mnp} + (\mathcal{L}_X\psi)_{jmnp}\psi_i^{mnp} = 12(\nabla_i X_j + \nabla_j X_i) + 24(\operatorname{div} X)g_{ij}$$

So, using (3.11), (3.14) and the above expressions, we obtain

$$\frac{1}{2}(\nabla_i X_j + \nabla_j X_i) + (\operatorname{div} X)g_{ij} = \frac{8}{7}(\operatorname{div} X)h_{ij} - h_{ij}$$

which, upon re-arranging, it gives

$$h_{ij} = \frac{1}{7} (\text{div}X) g_{ij} - \frac{1}{2} (\mathcal{L}_X g)_{ij}$$
(3.15)

Hence, substituting (3.11), (3.12) and (3.15) into (3.10) we obtain (3.9).

**Proposition 3.3.** Let  $\varphi$  be a coclosed G<sub>2</sub>-structure on M with associated metric g. If  $(\varphi, X, \lambda)$  is a soliton of the Laplacian coflow as in (3.3), then its full torsion tensor T satisfies

$$\operatorname{div} T = -\frac{1}{2} (\operatorname{Curl} X)^{\flat} + X \lrcorner T,$$
  
-Ric +  $\frac{1}{2} T \circ T + (\operatorname{tr} T) T = \frac{\lambda}{4} g + \frac{1}{2} \mathcal{L}_X g.$  (3.16)

*Proof.* Using (2.13), (2.3) and Lemma 3.1, we obtain

$$\Delta_{\psi}\psi = (\operatorname{div} T)^{\flat} \wedge \varphi + \mathrm{i}_{\psi}\Big( -\operatorname{Ric} + \frac{1}{2}T \circ T + (\operatorname{tr} T)T\Big).$$

On the other hand, by (2.3) and Proposition 3.2 we have

$$\lambda \psi + \mathcal{L}_X \psi = \left(-\frac{1}{2} \mathrm{Curl} X + X \lrcorner T\right)^{\flat} \land \varphi + \mathrm{i}_{\psi} \left(\frac{\lambda}{4}g + \frac{1}{2}(\mathcal{L}_X g)\right)$$

and thus we get (3.16).

- *Remark* 3.1. We notice that (3.16) coincides with the soliton equation for a general geometric flow given in [FLMSE22, Definition 1.52].
  - The tuple  $(g, X, \lambda)$  is called a *Ricci soliton* if it satisfies  $\text{Ric} = \lambda g + \mathcal{L}_X g$ . The second equation of (3.16) can be view as a perturbation of the Ricci soliton equation using the torsion tensor *T*. A similar remark was done by Lotay-Wei for the Laplacian flow [LW17], but in contrast, the first equation of (3.16) coincides with one of the equation of the isometric soliton condition of the harmonic flow of G<sub>2</sub>-structures [KLL20, Definition 2.16].
  - From the second equation of (3.16) is natural to ask for solitons of the Laplacian coflow, inducing Ricci solitons, aside from the nearly parallel case where  $\Delta \psi = \lambda^2 \psi$  and Ric  $= \frac{3}{8} \tau_0^2 g$ . For instance, in [MSE19] the authors obtain an example of a Laplacian coflow soliton inducing a Ricci soliton on a solvable Lie group.

Using (3.16), we can give an alternative proof for the non-existence of shrinking solitons in the compact case [KMT12, Proposition 4.3], and we extend this result to non-compact cases with X divergence free:

**Corollary 3.1.** *1. There are no compact shrinking solitons of the Laplacian coflow.* 

- 2. The only compact steady solitons of the Laplacian coflow are given by torsion free  $G_2$ -structures.
- 3. There do not exist steady (non-trivial i.e. X = 0) and shrinking solitons of the Laplacian coflow with  $\operatorname{div} X = 0$ .

*Proof.* Taking the trace on the second equation of (3.16), we obtain

$$\frac{1}{2}\left((\operatorname{tr} T)^2 + |T|^2\right) = \frac{7}{4}\lambda + \operatorname{div} X,\tag{3.17}$$

since  $\operatorname{tr}(T \circ T) = (\operatorname{tr} T)^2 - |T|^2$  and  $\operatorname{tr} \operatorname{Ric} = R$  (see (2.14)). If  $\operatorname{div} X = 0$  then  $\lambda \ge 0$ . When the manifold M is compact, we have

$$\lambda \operatorname{vol}(M) = \frac{2}{7} \int_M \left( (\operatorname{tr} T)^2 + |T|^2 \right) \operatorname{vol} \ge 0.$$

Hence,  $\lambda > 0$  or  $\lambda = 0$  if and only if T = 0.

### **4** Almost Abelian Lie groups revisited

We study in this section the Laplacian coflow and its solitons in a class of solvable Lie group, which has a codimension one Abelian ideal using the bracket flow as described in [Lau16].

Let G be a Lie group, it is called *almost Abelian* if its Lie algebra  $\mathfrak{g}$  admits an Abelian ideal  $\mathfrak{h}$  of codimension 1. For dim G = 7, any invariant G<sub>2</sub>-structure is completely determined by a G<sub>2</sub>-structure on  $\mathfrak{g}$ . Moreover, since G<sub>2</sub> acts transitively on the 6-sphere, thus, for any orthonormal basis  $\{e_1, \dots, e_7\}$ , we can suppose that  $e_7 \perp \mathfrak{h}$  and that the G<sub>2</sub>-structure has the form

$$\varphi = \omega \wedge e^7 + \rho^+ = e^{127} + e^{347} + e^{567} + e^{135} - e^{146} - e^{245} - e^{236}, \tag{4.1}$$

where  $\omega = e^{12} + e^{34} + e^{56}$  and  $\rho_+ = e^{135} - e^{146} - e^{245} - e^{236}$  are the canonical SU(3)-structure of  $\mathfrak{h} \cong \mathbb{R}^6$ . Additionally, the induced dual 4-form is

$$\psi := *\varphi = \frac{1}{2}\omega^2 + \rho_- \wedge e^7 = e^{1234} + e^{1256} + e^{3456} - e^{2467} + e^{2357} + e^{1457} + e^{1367}, \tag{4.2}$$

where  $\rho_{-} = J^* \rho_+$  and J is the canonical complex structure on  $\mathbb{R}^6$  defined by  $\omega := \langle J \cdot, \cdot \rangle$ . Moreover, the Lie bracket of  $\mathfrak{g}$  is encoded by  $A \in \mathfrak{gl}(\mathbb{R}^6)$  where  $A := \mathrm{ad}(e_7)|_{\mathfrak{h}}$ 

The transitive action of  $GL(\mathfrak{g})$  on the space of  $G_2$ -structures, defined by  $h \cdot \varphi := (h^{-1})^* \varphi$  (for  $h \in GL(\mathfrak{g})$ ), yields an infinitesimal representation of the alternating 3-form

$$\Lambda^{3}(\mathfrak{g})^{*} = \theta(\mathfrak{gl}(\mathfrak{g}))\varphi \tag{4.3}$$

where  $\theta : \mathfrak{gl}(\mathfrak{g}) \to \operatorname{End}(\Lambda^3 \mathfrak{g}^*)$  is defined by

$$\theta(B)\varphi := \frac{d}{dt}\Big|_{t=0} e^{tB} \cdot \varphi = -\varphi(B\cdot, \cdot, \cdot) - \varphi(\cdot, B\cdot, \cdot) - \varphi(\cdot, \cdot, B\cdot).$$
(4.4)

Since the orbit  $GL(\mathfrak{g})\psi$  is also open in  $\Lambda^4\mathfrak{g}^*$ , the relation (4.3) also holds for the 4-forms and  $\psi$ , namely  $\Lambda^4(\mathfrak{g})^* = \theta(\mathfrak{gl}(\mathfrak{g}))\psi$ . Coclosed G<sub>2</sub>-structures on almost Abelian Lie algebras are equivalent with the Lie bracket constrain  $A \in \mathfrak{sp}(\mathbb{R}^6)$  [Fre12], where

$$\mathfrak{sp}(\mathbb{R}^6) = \{ A \in \mathfrak{gl}(\mathbb{R}^6) : AJ + JA^t = 0 \quad \Leftrightarrow \quad \theta(A)\omega = 0 \}.$$

In particular, the non-vanishing torsion forms  $\tau_0$  and  $\tau_3$  can be described in terms of A:

**Proposition 4.1.** [Mor22, Prop. 3.2 & Cor 3.3] Let  $(\mathfrak{g}, A)$  be an almost Abelian Lie algebra with coclosed  $G_2$ -structure  $\varphi$ . Hence, the torsion forms of  $\varphi$  are

$$\tau_0 = \frac{2}{7} \operatorname{tr}(JA) \quad and \quad \tau_{27} = \left( \frac{\frac{1}{14} \operatorname{tr}(JA) \operatorname{I}_{6 \times 6} - \frac{1}{2}[J,A] \mid 0}{0 \mid -\frac{3}{7} \operatorname{tr}(JA)} \right).$$

And its full torsion tensor is

$$T = \left( \begin{array}{c|c} \frac{1}{2}[J,A] & 0\\ \hline 0 & \frac{1}{2}\operatorname{tr}(JA) \end{array} \right).$$
(4.5)

Moreover, we can describe the Hodge Laplacian  $\Delta \psi$  in function of  $A \in \mathfrak{sp}(\mathbb{R}^6)$ , hence according with Lemma 3.1, we first compute the tensor  $T \circ T$  given in (2.11):

**Lemma 4.2.** Let  $(\mathfrak{g}, A)$  be an almost Abelian Lie algebra with coclosed  $G_2$ -structure  $\varphi$ . Denote by  $S_A := \frac{1}{2}(A + A^t)$  the symmetric part of A, then we have

$$T \circ T = \left( \begin{array}{c|c} -\frac{1}{2} (\operatorname{tr} JA) [J, A] - S_A \circ_6 S_A & 0\\ \hline 0 & -\operatorname{tr} S_A^2 \end{array} \right), \tag{4.6}$$

where  $\circ_6$  is the product on  $\mathfrak{gl}(\mathbb{R}^6)$ , defined by  $(S_A \circ_6 S_A)_{ab} := (S_A)_{mn}(S_A)_{pq}\rho^+_{mpa}\rho^+_{nqb}$ .

*Proof.* We first compute the entry  $(T \circ T)_{77}$ , thus by (4.5) we have

$$(T \circ T)_{77} = T_{mn} T_{pq} \varphi_{mp7} \varphi_{nq7} = \frac{1}{4} [J, A]_{mn} [J, A]_{pq} \omega_{mp} \omega_{nq}$$
$$= \frac{1}{4} (J[J, A])_{np} ([J, A]J)_{np}$$
$$= \frac{1}{4} \langle J[J, A], [J, A]J \rangle = -\operatorname{tr} S_A^2,$$

for the last equality we used  $A = JA^t J$  (i.e.  $A \in \mathfrak{sp}(\mathbb{R}^6)$ ). Now, for  $i \neq 7$  and j = 7, we have

$$(T \circ T)_{i7} = T_{mn} T_{pq} \varphi_{mpi} \omega_{nq} = \frac{1}{4} [J, A]_{mn} [J, A]_{pq} \rho^{+}_{mpi} \omega_{nq}$$
$$= \frac{1}{4} ([J, A] J [J, A])_{mp} \rho^{+}_{mpi} = \frac{1}{4} ([J, A]^2)_{mn} J_{np} \rho^{+}_{pmi}$$
$$= -\frac{1}{4} ([J, A]^2)_{mn} \rho^{-}_{nmi} = 0.$$

For the above computation, we used that  $[J, A] \in \mathfrak{sp}(\mathbb{R}^6)$  is symmetric and (A.8). Finally, for  $i \neq 7$  and  $j \neq 7$  we have

$$(T \circ T)_{ij} = 2T_{mn}T_{77}\omega_{mi}\omega_{nj} + T_{mn}T_{pq}\rho_{mpi}^{+}\rho_{nqj}^{+}$$
  
$$= \frac{1}{2}(\operatorname{tr} JA)[J, A]_{mn}J_{mi}J_{nj} + \frac{1}{4}[J, A]_{mn}[J, A]_{pq}\rho_{mpi}^{+}\rho_{nqj}^{+}$$
  
$$= -\frac{1}{2}(\operatorname{tr} JA)(J[J, A]J)_{ij} + (JS_A)_{mn}(JS_A)_{pq}\rho_{mpi}^{+}\rho_{nqj}^{+}$$
  
$$= -\frac{1}{2}(\operatorname{tr} JA)([J, A])_{ij} + J_{mk}(S_A)_{kn}J_{pl}(S_A)_{lq}\rho_{mpi}^{+}\rho_{nqj}^{+}.$$

Once again, we used the identities (A.8) Finally, combining each case of i and j, we get the expression for  $T \circ T$ .

Now, for almost Abelian Lie algebras  $(\mathfrak{g}, A)$  the Ricci curvature is [Arr13, Eq (8)]:

$$\operatorname{Ric}_{A} = \left(\begin{array}{c|c} \frac{1}{2}[A, A^{t}] & 0\\ \hline 0 & -\operatorname{tr} S_{A}^{2} \end{array}\right) \quad \text{for} \quad A \in \mathfrak{sp}(\mathbb{R}^{6}),$$

$$(4.7)$$

and since tr(T) is constant, we have div T = 0. Therefore, we can write Lemma 3.1 in function of the Lie bracket: **Proposition 4.3.** Let  $(\mathfrak{g}, A)$  be an almost Abelian Lie algebra with coclosed G<sub>2</sub>-structure  $\varphi$ . Thus, the Hodge Laplacian of  $\psi$  is  $\Delta_A \psi = \theta(Q_A) \psi$  where

$$Q_A = \left(\begin{array}{c|c} Q_A^{\mathfrak{h}} & 0\\ \hline 0 & q_A \end{array}\right) = \left(\begin{array}{c|c} \frac{1}{2}[A, A^t] + \frac{1}{2}S_A \circ_6 S_A & 0\\ \hline 0 & -\frac{1}{2}\operatorname{tr}(S_A)^2 - \frac{1}{4}(\operatorname{tr} JA)^2 \end{array}\right)$$
(4.8)

In particular,  $Q_A \in \mathfrak{gl}(\mathfrak{g})$  is symmetric.

*Proof.* The result follows by applying equations (4.5), (4.7) and (4.6) into Lemma 3.1.

Notice that the closed condition on  $\psi$  implies that  $\Delta_{\psi}\psi = dd^*\psi$  is also closed. Similarly, it is interpreted as  $Q_A^{\mathfrak{h}} \in \mathfrak{sp}(\mathbb{R}^6)$  for  $A \in \mathfrak{sp}(\mathbb{R}^6)$ . Indeed:

**Lemma 4.4.** If  $A \in \mathfrak{sp}(\mathbb{R}^6)$  is symmetric then  $A \circ_6 A \in \mathfrak{sp}(\mathbb{R}^6)$ .

*Proof.* Notice that  $B := A \circ_6 A$  is symmetric, thus it is enough to prove the equality JBJ = B. Hence

$$(JBJ)_{ij} = J_{ik}(A \circ_6 A)_{kl}J_{lj} = J_{ik}A_{mn}A_{pq}\rho^+_{mpk}\rho^+_{nql}J_{lj}$$
  
$$= -(JAJ)_{mn}A_{pq}\rho^-_{mpi}\rho^-_{nqj} = -J_{mr}A_{rs}J_{sn}A_{pq}\rho^-_{mpi}\rho^-_{nqj}$$
  
$$= A_{rs}A_{pq}\rho^+_{rpi}\rho^+_{sqj} = B_{ij}$$

Later, we used the identities (A.8) time and again, as well as the symmetry of A.

### 4.1 The bracket flow

In this section we adapt the general approach of geometric flows of homogeneous geometric structures, proposed by J. Lauret, to the framework of the Laplacian coflow (3.1) on almost Abelian Lie algebras with coclosed G<sub>2</sub>-structures, for a broad exposition see [Lau16].

Let  $\{\varphi(t)\}_{t\in(\varepsilon_1,\varepsilon_2)}$  be a solution of the Laplacian coflow on  $(\mathfrak{g}, A)$  with initial condition  $\varphi(0) = \varphi_0$ . Since  $\varphi(t) \in \operatorname{GL}(\mathfrak{g})\varphi_0$ , we can write  $\varphi(t) = h(t)^*\varphi_0$  for  $h(t) \in \operatorname{GL}(\mathfrak{g})$  satisfying h(0) = I. Since  $*_{\varphi(t)} = (h^{-1})^* *_{\varphi_0} h^*$  (see [MSE19, Lemma 3.1]), we can write  $\psi(t) = h(t)^*\psi_0$  for  $\psi_0 = *_{\varphi_0}\varphi_0$  and by Proposition 4.3, we have

$$\Delta_A \psi(t) = \theta(Q_A(t))\psi(t),$$

hence, the Laplacian coflow is equivalent with

$$\frac{d}{dt}h(t) = -h(t)Q_A(t) \tag{4.9}$$

**Definition 4.1.** Let  $(G_1, \varphi_1)$  and  $(G_2, \varphi_2)$  be Lie groups with  $G_2$ -structure  $\varphi_i$  (for i = 1, 2). An isomorphism  $f : (G_1, \varphi_1) \to (G_2, \varphi_2)$  is called an *equivariant isomorphism* if it is a Lie group isomorphism such that  $\varphi_1 = f^* \varphi_2$  and this case,  $(G_1, \varphi_1)$  and  $(G_2, \varphi_2)$  are called *equivariant equivalent*.

Since  $\varphi(t) = *_t \psi(t)$  induces a SU(3)-structure on  $\mathfrak{h}$  for each t, we can write

$$h(t) = k(t) + a(t)e^7 \otimes e_7 \quad \text{where} \quad k(t) \in \mathrm{Gl}(\mathbb{R}^6) \quad \text{and} \quad a(t) \in \mathbb{R}^*.$$
(4.10)

Thus, we can defined the bracket  $A(t) = a(t)^{-1}k(t)Ak(t)^{-1}$  such that (4.10) becomes a Lie algebra isomorphism between  $(\mathfrak{g}, A, \varphi(t))$  and  $(\mathfrak{g}, A(t), \varphi)$  such that  $\varphi(t) = h(t)^* \varphi$ . Moreover, since  $\Delta_A \psi(t) = h(t)^* \Delta_{A(t)} \psi$  we get the relation  $Q_{A(t)} = h(t)Q_A(t)h(t)^{-1}$  and consequently, the equation (4.9) becomes an ODE on  $(\mathfrak{g}, A(t), \varphi)$ 

$$\frac{d}{dt}h(t) = -Q_{A(t)}h(t) \tag{4.11}$$

In particular, under the flow (4.11) the bracket A(t) evolves by:

$$\frac{d}{dt}A(t) = q_{A(t)}A(t) - [Q_{A(t)}^{\mathfrak{h}}, A(t)], \qquad (4.12)$$

where q(t) and  $Q_{A(t)}^{\mathfrak{h}}$  are defined in (4.8) for each  $t \in (\varepsilon_1, \varepsilon_2)$ . The ODE (4.12) is known as the *bracket flow* and it provides an equivalent analysis of the geometric flow of homogeneous geometric structures, varying the Lie bracket instead of the geometric structure:

**Theorem 4.1** ([Lau16, Theorem 5]). Let  $\{\varphi(t)\}_{t \in (\varepsilon_1, \varepsilon_2)}$  be a solution of the Laplacian coflow on  $(\mathfrak{g}, A)$  with initial condition  $\varphi(0) = \varphi_0$ . Then, there exist an equivariant isomorphism  $f(t) : (G_A, \varphi(t)) \to (G_{A(t)}, \varphi)$ , such that  $h(t) = df(t)_1$  solves either (4.9) or (4.11) for all  $t \in (\varepsilon_1, \varepsilon_2)$ . In addition, the solutions of (3.1) and (4.12) are

 $\varphi(t) = h(t)^* \varphi$  and  $A(t) = a(t)k(t)^{-1}Ak(t)$ ,

respectively, for  $t \in (\varepsilon_1, \varepsilon_2)$ .

Theorem 4.1 provides a useful tool for addressing long-time existence and regularity questions, since it shows that the Laplacian coflow and the bracket flow have the same maximal interval of solution. Hence, the bracket flow (4.12) is explicitly given by

**Proposition 4.5.** Let  $\mathcal{L} \simeq \mathfrak{gl}(\mathbb{R}^6)$  be the family of 7-dimensional almost Abelian Lie algebras. The subfamily  $\mathcal{L}_{coclosed} \simeq \mathfrak{sp}(\mathbb{R}^6) \subset \mathcal{L}$  of coclosed G<sub>2</sub>-structures is invariant under the bracket flow (4.12), which becomes equivalent to the following ODE for a one-parameter family of matrices  $A = A(t) \in \mathfrak{sp}(\mathbb{R}^6)$ :

$$\frac{d}{dt}A = -\left(\frac{1}{2}\operatorname{tr}(S_A)^2 + \frac{1}{4}(\operatorname{tr}JA)^2\right)A + \frac{1}{2}[A, [A, A^t]] + \frac{1}{2}[A, S_A \circ_6 S_A]$$
(4.13)

*Proof.* Notice that the velocity  $\dot{A}(t) = q_{A(t)}A + [A, Q_{A(t)}^{\mathfrak{h}}]$  lies in  $\mathfrak{sp}(\mathbb{R}^6)$ , since  $S_A \circ_6 S_A \in \mathfrak{sp}(\mathbb{R}^6)$  by Lemma 4.4, hence, the family  $\mathcal{L}_{coclosed} \subset \mathcal{L}$  is invariant under the bracket flow. Finally, replacing (4.8) into (4.12), we obtain (4.13).

**Proposition 4.6.** If A(t) is a bracket flow solution associated to the Laplacian coflow, then its norm evolves by

$$\frac{d}{dt}|A|^2 = -\left(|S_A|^2 + \frac{1}{2}(\operatorname{tr} JA)^2\right)|A|^2 - |[A, A^t]|^2 - \langle S_A \circ_6 S_A, [A, A^t]\rangle$$
(4.14)

*Proof.* From equation (4.13), we have

$$\frac{d}{dt}|A|^{2} = 2\langle \dot{A}, A \rangle = 2\operatorname{tr}(\dot{A}A^{t})$$

$$= -\left(\operatorname{tr}(S_{A})^{2} + \frac{1}{2}(\operatorname{tr}JA)^{2}\right)|A|^{2} + \operatorname{tr}([A, [A, A^{t}]]A^{t}) + \operatorname{tr}([A, S_{A} \circ_{6} S_{A}], A^{t})$$

$$= -\left(|S_{A}|^{2} + \frac{1}{2}(\operatorname{tr}JA)^{2}\right)|A|^{2} - |[A, A^{t}]| - \langle S_{A} \circ_{6} S_{A}, [A, A^{t}] \rangle$$

In order to proof long time existence solution for (4.13) we need the following identity.

**Lemma 4.7.** For the symmetric part  $S_A$  of the matrix  $A \in \mathfrak{sp}(\mathbb{R}^6)$ , we have

$$|S_A \circ_6 S_A|^2 = 4(|S_A|^2 |S_A|^2 - 2|S_A^2|^2 - \langle JS_A, S_A \rangle^2).$$

*Proof.* This identity follows by direct computations, using the contractions (A.8) and (A.11).

**Theorem 4.2.** The Laplacian coflow solution  $(\mathfrak{g}, A, \varphi(t))$  starting at any coclosed (non-flat) G<sub>2</sub>-structure is defined for all  $t \in (\varepsilon_1, \infty)$ .

*Proof.* Let  $\varphi(t)$  a solution of the Laplacian coflow defined for all  $t \in (\varepsilon_1, \varepsilon_2)$ , according to Theorem 4.1, we get that the solution  $A(t) \in \mathfrak{sp}(\mathbb{R}^6)$  of (4.13) is defined for all  $t \in (\varepsilon_1, \varepsilon_2)$ . Now, using the Cauchy-Schwarz and Peter-Paul inequalities (i.e.  $ab \leq \frac{a^2}{4} + b^2$  for  $a, b \geq 0$ ), we have

$$-\langle S_A \circ_6 S_A, [A, A^t] \rangle \leq |S_A \circ_6 S_A| |[A, A^t]| \\\leq \frac{|S_A \circ_6 S_A|^2}{4} + |[A, A^t]|^2 = |S_A|^2 |S_A|^2 - 2|S_A^2|^2 - \langle JS_A, S_A \rangle^2 + |[A, A^t]|^2.$$
(4.15)

Replacing the last inequality into equation (4.14), we have

$$\begin{aligned} \frac{d}{dt}|A|^2 &\leq -\left(|S_A|^2 + \frac{1}{2}(\operatorname{tr} JA)^2\right)|A|^2 - |[A, A^t]|^2 + |S_A|^2|S_A|^2 - 2|S_A^2|^2 - \langle JS_A, S_A \rangle^2 + |[A, A^t]|^2 \\ &= -|S_A|^2|S_A|^2 - \frac{1}{4}|S_A|^2|A - A^t|^2 - \frac{1}{2}(\operatorname{tr} JA)^2|A|^2 + |S_A|^2|S_A|^2 - 2|S_A^2|^2 - \langle JS_A, S_A \rangle^2 \\ &= -\frac{1}{4}|S_A|^2|A - A^t|^2 - \frac{1}{2}(\operatorname{tr} JA)^2|A|^2 - 2|S_A^2|^2 - \langle JS_A, S_A \rangle^2 \leq 0 \end{aligned}$$

Thus,  $|A|^2$  is non-increasing and non-negative, therefore A(t) is an inmortal solution, i.e. it is defined for all  $t \in (\varepsilon_1, \infty)$ . In particular,  $|A|^2$  is strictly decreasing unless  $(\mathfrak{g}, A(t), \varphi)$  is torsion free, that is

$$|A|^2 = 0 \quad \Leftrightarrow \quad A^t = -A \quad \text{and} \quad \operatorname{tr} JA = 0$$

and thus  $A(t) \equiv A_0 \in \mathfrak{sl}(\mathbb{C}^3) \cap \mathfrak{sp}(\mathbb{R}^6) = \mathfrak{su}(3)$  the bracket flow solution is constant.

*Remark* 4.1. In [BF18] Bagaglini and Fino address also the Laplacian coflow on almost Abelian Lie algebras, there the approach is different from ours, the authors find explicit solutions of the Laplacian coflow when  $A \in \mathfrak{sp}(\mathbb{R}^6)$  is normal. Notice that the above Corollary holds for any  $A \in \mathfrak{sp}(\mathbb{R}^6)$ .

**Proposition 4.8.** Let  $(\mathfrak{g}, A)$  be an almost Abelian algebra with the matrix A defined by

$$A = \begin{bmatrix} B & 0 \\ 0 & -B^t \end{bmatrix} \quad \text{with} \quad B = \begin{bmatrix} 0 & x & 0 \\ y & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{and} \quad x, y \in \mathbb{R}$$
(4.16)

Then, the bracket flow A(t) given by (4.13) is stable.



Figure 1: x-nulcline, y-nulcline, equilibrium points

*Proof.* The bracket flow (4.13) being A(t) the matrix given by (4.16) can be analized by the following nonlinear system  $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x})$  where  $\mathbf{f} : \mathbb{R}^2 \to \mathbb{R}^2$ ;  $\mathbf{x} \mapsto \mathbf{f} = (\mathbf{f_1}(\mathbf{x}), \mathbf{f_2}(\mathbf{x}))$  given by

$$\dot{x} = -2x(3x - y)(x + y)$$
 and  $\dot{y} = 2y(x - 3y)(x + y).$  (4.17)

A point  $\mathbf{x} \in \mathbb{R}^2$  is an *equilibrium point* if  $\mathbf{f}(\mathbf{x}) = \mathbf{0}$  which is given by the surface  $S = \{(x, y) \in \mathbb{R}^2 : x = -y\}$ . The x-nullclines (i.e,  $\mathbf{x} \in \mathbb{R}^2$  where  $\mathbf{f_1}(\mathbf{x}) = \mathbf{0}$ ) are the lines x = 0, x = -y and y = 3x and the y-nullclines (i.e,  $\mathbf{x} \in \mathbb{R}^2$  where  $\mathbf{f_2}(\mathbf{x}) = \mathbf{0}$ ) are the lines y = 0, x = -y and  $y = \frac{1}{3}x$ . The intersection of the x-nulcines and y-nulclines yield the equilibrium points. On the other hand, the lines y = 0 and x = 0 are invariants for the system (4.17). If we set y = 0 then we obtain  $\dot{x} = -6x^3$ . Therefore,  $\dot{x}$  is positive if x > 0 and negative if x < 0 which clearly shows the stability along the line y = 0.

To determine the trajectories, if  $\mathbf{x_0}$  is not a equilibrium point then, at least one of  $f_1(\mathbf{x_0})$  or  $f_2(\mathbf{x_0})$  is not zero. Let us suppose that  $f_1(\mathbf{x_0}) \neq \mathbf{0}$ . Then, there is an open neighborhood of  $\mathbf{x_0}$ , such that  $f_1(\mathbf{x_0}) \neq \mathbf{0}$ , so the orbit through  $\mathbf{x_0}$  can be defined as a solution of the nonautonomous scalar equation

$$\frac{dy}{dx} = -\frac{y(x-3y)}{x(3x-y)}.$$
(4.18)

This differential equation is homogeneous. Setting y = xv(x), we obtain

$$v + x\frac{dv}{dx} = -\frac{v(1-3v)}{3-v}.$$

That is,

$$x\frac{dv}{dx} = 4v\left(\frac{v-1}{3-v}\right).$$

The resulting ODE is separable, with solution  $x^{-4}v^{-3}(v-1)^2 = c$ . Reverting back to the original variables, the trajectories are level curves of

$$H(x(t), y(t)) = \frac{(y(t) - x(t))^2}{y(t)^3 x(t)^3}.$$

On the other hand, let

$$V(\mathbf{x}) = x^2 + y^2 + 2xy,$$

be a Lyapunov function. In fact,  $V(\mathbf{x}) = 0$  when  $\mathbf{x}$  is an equilibrium point for this system and  $V(x) = (x+y)^2 \ge 0$  if  $\mathbf{x}$  is not an equilibrium point. Computing  $\dot{V}(\mathbf{x})$ , we find

$$\dot{V}(\mathbf{x}) = -2(x+y)^2(6x^2 - 4xy + 6y^2),$$
(4.19)

where  $\dot{V}(x) = 0$  if x = -y and  $\dot{V}(x) \le 0$  otherwise. For any curve  $\gamma(r, \theta) = (r \cos \theta, r \sin \theta)$  with r > 0 and  $0 \le \theta \le 2\pi$ , we obtain

$$\dot{V}(\gamma(r,\theta)) = -2r^2(r\sin\theta + r\cos\theta)^2(6 - 2\sin(2\theta)) \le 0, \tag{4.20}$$

since  $|\sin(2\theta)| \le 1$  then we have  $6 - 2\sin(\theta) > 0$ . Therefore, the system is stable if  $\mathbf{x}_0$  is a equilibrium point.

**Proposition 4.9.** The scalar curvature R(t) of an invariant Laplacian coflow solution starting at a non-flat coclosed G<sub>2</sub>-structure is strictly increasing and satisfies the inequality

$$\frac{1}{-\frac{|t|}{2} + \frac{1}{R(0)}} \le R(t) \le 0 \quad \text{for} \quad \text{any} \quad t \in (\varepsilon_1, \infty),$$

$$(4.21)$$

In particular,  $|T|^2$  is strictly decreasing. and it converges to zero when  $t \to \infty$  as  $|A(t)|^2 \to 0$ .

*Proof.* From (4.7), we have  $R = -\operatorname{tr} S_A^2 = -\frac{1}{4}\operatorname{tr}(A + A^t)^2$ . Thus, using the bracket flow equation (4.13) we have

$$\begin{aligned} \frac{d}{dt} \operatorname{tr}(A+A^{t})^{2} &= 2\operatorname{tr}\left((A+A^{t})\frac{d}{dt}(A+A^{t})\right) \\ &= -\left(\operatorname{tr}S_{A}^{2} + \frac{1}{2}(\operatorname{tr}JA)^{2}\right)\operatorname{tr}(A+A^{t})^{2} + \operatorname{tr}\left((A+A^{t})[A-A^{t},[A,A^{t}]]\right) \\ &+ \operatorname{tr}\left((A+A^{t})[A-A^{t},S_{A}\circ_{6}S_{A}]\right) \\ &= -4\left(\operatorname{tr}S_{A}^{2} + \frac{1}{2}(\operatorname{tr}JA)^{2}\right)\operatorname{tr}S_{A}^{2} + \operatorname{tr}\left([A+A^{t},A-A^{t}][A,A^{t}]\right) \\ &+ \operatorname{tr}\left([A+A^{t},A-A^{t}]S_{A}\circ_{6}S_{A}\right) \\ &= -2\left(|S_{A}|^{2} + (\operatorname{tr}JA)^{2}\right)|S_{A}|^{2} - 2|[A,A^{t}]|^{2} - 2\langle[A,A^{t}]S_{A}\circ S_{A}\rangle \end{aligned}$$

Using the inequality (4.15), we obtain

$$\begin{aligned} \frac{d}{dt}\operatorname{tr}(A+A^{t})^{2} &\leq -2\left(2|S_{A}|^{2} + (\operatorname{tr}JA)^{2}\right)|S_{A}|^{2} + 2|S_{A}|^{4} - 4|S_{A}^{2}|^{2} - 2(\langle JS_{A}, S_{A}\rangle)^{2} \\ &\leq -2(\operatorname{tr}S_{A}^{2})^{2} = -\frac{1}{8}(\operatorname{tr}(A+A^{t})^{2})^{2}. \end{aligned}$$

For any  $t_1, t_2 \in (\varepsilon_1, \infty)$  satisfying  $t_1 \leq t_2$ , the last inequality implies

$$\frac{1}{R(t_2)} - \frac{1}{R(t_1)} \ge \frac{t_2 - t_1}{2}.$$

If  $t_1 = 0$  then we get

$$\frac{1}{-\frac{t_2}{2} + \frac{1}{R(0)}} \le R(t_2) < 0 \quad \text{any} \quad t_2 \in [0, \infty).$$

If  $t_2 = 0$  then we obtain

$$\frac{1}{\frac{t_1}{2} + \frac{1}{R(0)}} \le R(t_1) < 0 \quad \text{any} \quad t_1 \in (\varepsilon_1, 0].$$

Finally, by (2.13) and (4.5), the scalar curvature of a coclosed G<sub>2</sub>-structure is

$$R_A = -|T|^2 + (\operatorname{tr}(JA))^2.$$

Hence, using the Cauchy-Schwarz inequality, we have

$$|T|^{2} \leq -R(t) + |J|^{2}|A(t)|^{2} = -R(t) + 6|A(t)|^{2} \leq \frac{1}{\frac{|t|}{2} - \frac{1}{R(0)}} + 6|A(t)|^{2}.$$

Therefore,  $|T|^2$  is strictly decreasing, since  $|A(t)|^2$  is strictly decreasing as well and  $|T|^2$  goes to zero as  $|A(t)| \to 0$ .

### 4.2 Algebraic solitons

In this section, we characterise the invariant G<sub>2</sub>-structures on almost Abelian Lie algebras which are solitons of the Laplacian coflow, in terms of the Lie bracket  $A \in \mathfrak{sp}(\mathbb{R}^6)$ .

The self-similar solution (3.2) on the almost Abelian Lie group  $G_A$  is invariant if  $\lambda(t) \in \mathbb{R}^*$  and  $f(t) \in Aut(G_A)$ , then (3.2) corresponds with the solution

$$\psi(t) = \lambda(t)h(t)^*\psi \in \Lambda^4 \mathfrak{g}^* \quad \text{with} \quad \lambda(t) \in \mathbb{R}^* \quad \text{and} \quad h(t) \in \operatorname{Aut}(\mathfrak{g}, A), \tag{4.22}$$

with  $df(t)_1 = h(t)$  and then, the soliton equation (3.3) becomes  $\Delta_{\psi}\psi = \lambda\psi + \mathcal{L}_{X_D}\psi \in \Lambda^4\mathfrak{g}^*$  with  $\lambda \in \mathbb{R}$  and  $X_D := \frac{d}{dt}|_{t=0}h(t) =: -D \in \operatorname{Der}(\mathfrak{g}, A)$ . Using the representation (4.3), we have

$$\theta(Q_A)\psi = \Delta_{\psi}\psi = \lambda\psi + \mathcal{L}_{X_D}\psi$$
$$= \theta\left(-\frac{\lambda}{4}I_7\right)\psi + \frac{d}{dt}|_{t=0}h(t)^*\psi$$
$$= \theta\left(-\frac{\lambda}{4}I_7 + D\right)\psi.$$

By Proposition 4.3, the matrix  $Q_A$  is symmetric, hence, setting  $\lambda = -4c$  we say that  $\psi$  is a *semi-algebraic* soliton if

$$Q_A = cI_7 + \frac{1}{2}(D + D^t),$$

and  $\psi$  is an *algebraic soliton* if  $D^t \in \text{Der}(\mathfrak{g}, A)$ . Moreover, the self-similar solution (4.22) is given

$$\lambda(t) = (1 - 2ct)^2$$
 and  $h(t) = e^{-s(t)D}$  where  $s(t) = -\frac{1}{2c} \log |2ct - 1|$ 

(For c = 0 set s(t) = t). And the corresponding bracket solution of a semi-algebraic soliton is

$$A(t) = (1 - 2ct)^{-1/2} e^{s(t)E} A e^{-s(t)E} \quad \text{where} \quad E = \frac{1}{2} (D - D^t), \tag{4.23}$$

(e.g. [LL14, Remark 3.4] for the homogeneous Ricci soliton case). The next Theorem shows the (semi-) algebraic soliton equation in terms  $A \in \mathfrak{sp}(\mathbb{R}^6)$ .

**Theorem 4.3.** Let  $(\mathfrak{g}, A, \varphi)$  be an almost Abelian Lie algebra with coclosed  $G_2$ -structure:

(i)  $\psi$  is an algebraic soliton for the Laplacian coflow if and only if

$$[[A, A^{t}] + S_{A} \circ_{6} S_{A}, A] = \frac{|[A, A^{t}]|^{2} + \langle S_{A} \circ_{6} S_{A}, [A, A^{t}] \rangle}{|A|^{2}} A$$
(4.24)

In this case,  $D = Q_A - cI_7 \in \text{Der}(\mathfrak{g}, A)$  for

$$c = -\frac{1}{2} \Big( \operatorname{tr} S_A^2 + \frac{1}{2} (\operatorname{tr} JA)^2 + \frac{|[A, A^t]|^2}{|A|^2} + \frac{\langle S_A \circ_6 S_A, [A, A^t] \rangle}{|A|^2} \Big).$$
(4.25)

(ii)  $\psi$  is a semi-algebraic soliton if and only if

$$[A, A^{t}] + S_{A} \circ_{6} S_{A} = -\left(\operatorname{tr} S_{A}^{2} - \frac{1}{2}(\operatorname{tr} JA)^{2} + 2d\right)I_{6} + D_{1} + D_{1}^{t},$$
(4.26)

for some  $D_1 \in \mathfrak{gl}(\mathbb{R}^6)$  such that  $[D_1, A] = dA$ , where

$$d = \frac{|[A, A^t]|^2 + \langle S_A \circ_6 S_A, [A, A^t] \rangle}{2|A|^2}.$$

In this case  $Q_A = cI_7 + \frac{1}{2}(D+D^t)$  for

$$c = -\frac{1}{2} \left( \operatorname{tr} S_A^2 + \frac{1}{2} (\operatorname{tr} JA)^2 + \frac{|[A, A^t]|^2}{|A|^2} + \frac{\langle S_A \circ_6 S_A, [A, A^t] \rangle}{|A|^2} \right)$$
(4.27)

*Proof.* (i) Suppose that  $(\mathfrak{g}, A, \varphi)$  is an algebraic soliton i.e.  $Q_A = cI + D$  for  $c \in \mathbb{R}$  and  $D \in \text{Der}(\mathfrak{g}, A)$ . Then,

 $De_7 = de_7$  for some  $d \in \mathbb{R}$  and  $[Q_A^{\mathfrak{h}}, A] = [D|_{\mathfrak{h}}, A] = dA.$ 

Thus, by Proposition 4.3 we get

$$[[A, A^t], A] + [S_A \circ_6 S_A, A] = 2dA.$$

Taking the inner product between A and the above equation we obtain

$$d = \frac{|[A, A^t]|^2 + \langle S_A \circ_6 S_A, [A, A^t] \rangle}{2|A|^2}$$

The converse follows by taking  $D = Q_A - cI \in \text{Der}(\mathfrak{g}, A)$  and

$$c = q - d = -\frac{1}{2} \Big( \operatorname{tr} S_A^2 + \frac{1}{2} (\operatorname{tr} JA)^2 + \frac{|[A, A^t]|^2}{|A|^2} + \frac{\langle S_A \circ_6 S_A, [A, A^t] \rangle}{|A|^2} \Big).$$

(ii) Suppose that  $(\mathfrak{g}, A, \varphi)$  is a semi algebraic soliton, i.e.  $Q_A = cI_7 + \frac{1}{2}(D + D^t)$  for some  $c \in \mathbb{R}$  and  $D \in \text{Der}(\mathfrak{g}, A)$ . It implies the equations

$$Q_A^{\mathfrak{h}} = cI_6 + \frac{1}{2}(D_1 + D_1^t)$$
 and  $q = c + d$ 

where

$$De_7 = de_7$$
 for  $d \in \mathbb{R}$  and  $[D_1, A] = dA$  where  $D_1 = D|_{\mathfrak{h}}$ .

Since  $\langle [D_1, A], A \rangle = \langle A, [D_1^t, A] \rangle$ , by Proposition 4.3 we obtain (4.26). The converse follows immediately, and the formulae for *c* and *d* are obtained as in (i).

Using the condition (4.24) we describe a class of algebraic solitons.

**Corollary 4.1.** If  $A \in \mathfrak{sp}(\mathbb{R}^6)$  is skew-symmetric then  $(\mathfrak{g}, A, \varphi)$  is an algebraic soliton.

Using Lemma 4.7, we can prove the absence of shrinking (semi-) algebraic solitons for the Laplacian coflow on almost Abelian Lie algebras.

**Proposition 4.10.** If  $(\mathfrak{g}, A, \varphi)$  is a (semi-) algebraic soliton for the Laplacian coflow then it is expanding, and it is steady if it is torsion-free.

*Proof.* Using the inequality (4.15) in the equation (4.27), we have

$$2c \leq -\left(\operatorname{tr} S_{A}^{2} + \frac{1}{2} (\operatorname{tr} JA)^{2} + \frac{|[A, A^{t}]|^{2}}{|A|^{2}}\right) + \frac{1}{|A|^{2}} \left(|S_{A}|^{2}|S_{A}|^{2} - 2|S_{A}^{2}|^{2} - \langle JS_{A}, S_{A} \rangle^{2} + |[A, A^{t}]|^{2}\right)$$
$$\leq -\frac{1}{|A|^{2}} \left(|S_{A}|^{2} (|A|^{2} - |S_{A}|^{2}) + \frac{1}{2} (\operatorname{tr} JA)^{2}|A|^{2} + 2|S_{A}^{2}|^{2} + \langle JS_{A}, S_{A} \rangle^{2}\right) \leq 0.$$

If c = 0 then

$$\operatorname{tr} JA = 0 \quad \text{and} \quad S_A^2 = 0.$$

In particular  $S_A = 0$ , and thus A is skew-symmetric. And since  $A \in \mathfrak{sp}(\mathbb{R}^6)$  it implies that [J, A] = 0. Therefore, by equation (4.5) we get that the full torsion tensor T vanishes.

*Remark* 4.2. We remark that the previous proposition was prove in [BF18, Corollary 4.4] for the context of algebraic solitons and assuming that A is normal.

We conclude this section with an example of a semi-algebraic soliton which is not an algebraic one.

**Example 4.1.** Let  $(\mathfrak{g}, A, \varphi)$  be an almost Abelian Lie algebra with  $G_2$ -structure  $\varphi = \omega \wedge e^7 + \rho^+$ , where

$$\omega = e^{14} + e^{25} + e^{36}$$
 and  $\rho^+ = e^{123} - e^{156} + e^{246} - e^{345}$ 

and the Lie bracket is given by the 3-step nilpotent matrix

$$A = \left(\begin{array}{c|c} 0 & B \\ \hline C & 0 \end{array}\right) \in \mathfrak{sp}(\mathbb{R}^6), \quad B = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & \sqrt{2} & 0 \\ \sqrt{2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

We have that the matrix

$$D = \begin{pmatrix} D_1 & 0\\ 0 & d \end{pmatrix}, \quad D_1 = \begin{pmatrix} 2 & 0 & 0\\ 0 & 2 & 0\\ -\sqrt{2} & 0 & 4\\ \hline & & & \\ \hline & & & \\ 0 & 3 & 0\\ & & & & \\ 0 & 0 & 1 \end{pmatrix}, \quad d = 1,$$

satisfies the relation  $[D_1, A] = A$ , it means that  $D \in Der(\mathfrak{g}, A)$ . Now, for each term of (4.8), we obtain

$$[A, A^{t}] = \begin{pmatrix} -2 & 0 & 0 & | & & \\ 0 & -1 & 0 & & \\ 0 & 0 & 1 & & \\ \hline & & & 2 & 0 & 0 \\ & & & & 0 & 1 & 0 \\ & & & & 0 & 0 & -1 \end{pmatrix}, \quad S_{A} \circ_{6} S_{A} = \begin{pmatrix} 1 & 0 & -\sqrt{2} & & \\ 0 & 0 & 0 & & \\ -\sqrt{2} & 0 & 2 & & \\ \hline & & & & -1 & 0 & \sqrt{2} \\ & & & & 0 & 0 & \\ & & & & \sqrt{2} & 0 & -2 \end{pmatrix}$$

and

tr 
$$S_A^2 = 3$$
, tr  $JA = 0$ ,  $|[A, A^t]|^2 = 12$ ,  $|A|^2 = 6$  and  $\langle S_A \circ_6 S_A, [A, A^t] \rangle = 0$ .

Since the matrices A and D satisfy the equation (4.26), we have that  $(\mathfrak{g}, A, \varphi)$  is a semi-algebraic soliton with

$$Q_A = -\frac{5}{2}I + \frac{1}{2}(D + D^t).$$

Notice that  $[D_1^t, A] \neq A$ , so  $D^t \notin \text{Der}(\mathfrak{g}, A)$  thus  $(\mathfrak{g}, A, \varphi)$  is not an algebraic soliton. According to (4.23), the associated bracket flow solution is

$$A(t) = (1+5t)^{-1/2} e^{s(t)E} A e^{-s(t)E} = (1+5t)^{-1/2} \left( \cos \frac{s(t)}{\sqrt{2}} A + \sin \frac{s(t)}{\sqrt{2}} A^{\perp} \right)$$

where

$$E = \frac{1}{2}(D - D^{t}) = \frac{1}{\sqrt{2}} \begin{pmatrix} E_{1} & 0 \\ 0 & E_{1} \\ \hline & 0 \end{pmatrix}, \quad E_{1} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}$$

and

$$A^{\perp} = \begin{pmatrix} 0 & B' \\ \hline C' & 0 \end{pmatrix} \quad B' = \begin{pmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad C' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \sqrt{2} \\ 0 & \sqrt{2} & 0 \end{pmatrix}.$$

As in [Lau17, Example 5.28], we obtain that A(t)/|A(t)| runs on a circle and A(t) converges to zero rounding in a cone.

# A Contraction of $G_2$ and SU(3)-identities

Let  $\varphi$  be a G<sub>2</sub>-structure with Hodge dual 4-form  $\psi$  and induced SU(3)-structure  $(\omega, \rho^+ + i\rho^-) \in \Lambda^2(\mathbb{R}^6)^* \oplus \Lambda^3(\mathbb{C}^3)^*$ . From [Kar07, §A.3] and [BV07, §2.2], we gather the following contraction identities for G<sub>2</sub> and SU(3)-structures, respectively.

Contractions of  $\varphi$  with  $\varphi$ :

$$\varphi_{abc}\varphi^{abc} = 42,\tag{A.1}$$

$$\varphi_{abj}\varphi^{ab}_{\ k} = 6g_{jk},\tag{A.2}$$

$$\varphi_{apq}\varphi^a_{\ jk} = g_{pj}g_{qk} - g_{pk}g_{qj} + \psi_{pqjk}.\tag{A.3}$$

Contractions of  $\varphi$  with  $\psi$ :

$$\varphi_{ijk}\psi_a^{ijk} = 0,$$
  

$$\varphi_{ijq}\psi_{kl}^{ij} = 4\varphi_{qkl},$$
(A.4)

$$\varphi_{ipq}\psi^{i}_{jkl} = g_{pj}\varphi_{qkl} - g_{jq}\varphi_{pkl} + g_{pk}\varphi_{jql}$$

$$(A.5)$$

$$-g_{kq}\varphi_{jpl} + g_{pl}\varphi_{jkq} - g_{lq}\varphi_{jkp}. \tag{A.5}$$

Contractions of  $\psi$  with  $\psi$ :

$$\psi_{abcd}\psi^{ab}_{mn} = 4g_{cm}g_{dn} - 4g_{cn}g_{dm} + 2\psi_{abmn},\tag{A.6}$$

$$\psi_{abcd}\psi_m^{\ bcd} = 24g_{am},\tag{A.7}$$

$$\psi_{abcd}\psi^{abcd} = 168$$

. . .

Contractions of  $\omega$  with  $\omega$  and  $\rho^{\pm}$  with  $\omega$ :

$$\omega_{ip}\omega^{p}{}_{j} = -\delta_{ij}, \quad \rho^{+}_{iab}\omega^{ab} = 0, \quad \rho^{+}_{ijp}\omega^{p}{}_{k} = \rho^{-}_{ijk}, \quad \rho^{-}_{ijp}\omega^{p}{}_{k} = -\rho^{+}_{ijk}.$$
(A.8)

Contractions of  $\rho^{\pm}$  with  $\rho^{\pm}$ :

$$\rho_{ipq}^{+}\rho_{j}^{-pq} = 4\omega_{ij}, \quad \rho_{ipq}^{+}\rho_{j}^{+pq} = 4\delta_{ij} = \rho_{ipq}^{-}\rho_{j}^{-pq}, \tag{A.9}$$

$$\rho_{ijp}^{-}\rho_{kl}^{+p} = -\omega_{ik}\delta_{jl} + \omega_{jk}\delta_{il} + \omega_{il}\delta_{jk} - \omega_{jl}\delta_{ik}, \tag{A.10}$$

$$\rho_{ijp}^+ \rho_{kl}^{+p} = -\omega_{ik}\omega_{jl} + \omega_{il}\omega_{jk} + \delta_{ik}\delta_{jl} - \delta_{jk}\delta_{il} = \rho_{ijp}^- \rho_{kl}^{-p}.$$
(A.11)

## References

- [Arr13] R. Arroyo, The Ricci flow in a class of solvmanifolds, Differential Geom. Appl. 31 (2013), no. 4, 472–485. <sup>13</sup>
- [BF18] L. Bagaglini and A. Fino, The Laplacian coflow on almost-abelian Lie groups, Ann. Mat. Pura Appl. 197 (2018), no. 6, 1855–1873. ↑1, 2, 3, 15, 19
- [BFF20] L. Bagaglini, M. Fernández, and A. Fino, Laplacian coflow on the 7-dimensional Heisenberg group, Asian J. Math. 24 (2020), no. 2, 331–354. <sup>↑</sup>2
- [Bry06] R. L. Bryant, Some remarks on G<sub>2</sub>-structures, Proceedings of Gökova Geometry-Topology Conference 2005, 2006, pp. 75– 109. <sup>↑</sup>1, 5, 7
- [BV07] L. Bedulli and L. Vezzoni, The Ricci tensor of SU(3)-manifolds, J. Geom. Phys. 57 (2007), no. 4, 1125–1146. <sup>121</sup>
- [BX11] R. L. Bryant and Feng Xu, Laplacian flow for closed G<sub>2</sub>-structures: Short time behavior (2011). Available at arXiv:1101.2004.<sup>↑</sup>1
- [CN15] Diarmuid Crowley and Johannes Nordström, New invariants of G<sub>2</sub>-structures, Geom. Topol. 19 (2015), no. 5, 2949–2992. MR3416118 <sup>↑</sup>2
- [DGK21] S. Dwivedi, P. Gianniotis, and S. Karigiannis, A gradient flow of isometric G<sub>2</sub>-structures, J. Geom. Anal. 31 (2021), no. 2, 1855–1933. MR4215279 <sup>↑</sup>9
  - [FG82] M. Fernández and A. Gray, *Riemannian manifolds with structure group* G<sub>2</sub>, Ann. Mat. Pura Appl. **132** (1982), no. 1, 19–45 (1983). <sup>↑</sup>1

- [FKMS97] Th. Friedrich, I. Kath, A. Moroianu, and U. Semmelmann, On nearly parallel G<sub>2</sub>-structures, J. Geom. Phys. 23 (1997), no. 3-4, 259–286 (English). <sup>↑</sup>2
- [FLMSE22] D. Fadel, E. Loubeau, A. Moreno, and H. Sá Earp, Flows of geometric structures, arXiv preprint 2211.05197 (2022). <sup>+6</sup>, 11
  - [Fre12] M. Freibert, Cocalibrated structures on Lie algebras with a codimension one Abelian ideal, Ann. Glob. Anal. Geom 42 (2012), no. 4, 537–563. <sup>↑</sup>12
  - [Gri13] S. Grigorian, Short-time behaviour of a modified Laplacian coflow of G<sub>2</sub>-structures, Adv. Math **248** (2013), 378–415. <sup>↑</sup>2, 5, 6, 7
  - [Kar07] S. Karigiannis, Flows of G2-structures, Q. J. Math. 60 (2007), no. 4. 15, 6, 21
  - [KL21] I. Kath and J. Lauret, A new example of a compact ERP G<sub>2</sub>-structure, Bull. London Math. Soc. 53 (2021), no. 6, 1692–1710. ↑2
  - [KL23] A. Kennon and J. Lotay, *Geometric flows of* G<sub>2</sub>-structures on 3-sasakian 7-manifolds, J. Geom. Phys. **187** (2023), 104793. ↑2
  - [KLL20] S. Karigiannis, N. C. Leung, and J. D. Lotay (eds.), Lectures and Surveys on G<sub>2</sub>-manifolds and related topics, Fields Inst. Commun., vol. 84, Springer US, 2020. <sup>↑</sup>11
  - [KMT12] S. Karigiannis, B. McKay, and M-P. Tsui, Soliton solutions for the Laplacian coflow of some G<sub>2</sub>-structures with symmetry, Differ. Geom. Appl. 30 (2012), 318–333. <sup>↑</sup>1, 2, 7, 11
  - [Lau16] J. Lauret, Geometric flows and their solitons on homogeneous spaces, Rend. Sem. Mat. Univ. Politec. Torino 74 (2016), 55–93. ↑2, 3, 12, 14
  - [Lau17] \_\_\_\_\_, Laplacian flow of homogeneous G<sub>2</sub>-structures and its solitons, Proc. Lond. Math. Soc.. 114 (2017), 527–560. <sup>↑</sup>20
  - [LL14] R. Lafuente and J. Lauret, On homogeneous ricci solitons, Q. J. Math. 65 (2014), no. 2, 399–419. <sup>18</sup>
  - [LSES22] J. Lotay, H. Sá Earp, and J. Saavedra, Flows of G<sub>2</sub>-structures on contact calabi-yau 7-manifolds, Ann. Glob. Anal. Geom 62 (2022), no. 2, 367–389. <sup>↑</sup>2
    - [LW17] J. D. Lotay and Y. Wei, Laplacian Flow for Closed G<sub>2</sub>-structures: Shi-type Estimates, Uniqueness and Compactness, Geom. Funct. Anal. 27 (2017), 165–233. <sup>↑</sup>7, 11
  - [Mor22] A. J. Moreno, Harmonic G<sub>2</sub>-structures on almost Abelian Lie groups, arXiv preprint:2212.05168 (2022). <sup>↑</sup>12
  - [MSE19] A. Moreno and H. N. Sá Earp, Explicit soliton for the Laplacian co-flow on a solvmanifold, Proceedings of the satellite conference to ICM2018 on Differential Geometry, 2019. <sup>↑</sup>11, 14
  - IMECC, UNIVERSITY OF CAMPINAS, CAMPINAS SÃO PAULO, BRAZIL A. Moreno: andres.moreno@ime.unicamp.br
  - FEDERAL UNIVERSITY OF CEARÁ, FORTALEZA CEARÁ, BRAZIL J. Saavedra: julieth.p.saavedra@gmail.com