Non-Hermitian Floquet Topological Matter – A Review

Longwen Zhou^{1, 2, 3, *} and Da-Jian Zhang^{4, †}

¹College of Physics and Optoelectronic Engineering, Ocean University of China, Qingdao 266100, China

²Key Laboratory of Optics and Optoelectronics, Qingdao 266100, China

³Engineering Research Center of Advanced Marine Physical Instruments and Equipment,

Ministry of Education, Qingdao 266100, China

⁴Department of Physics, Shandong University, Jinan 250100, China

(Dated: 2023-08-01)

Abstract

The past few years have witnessed a surge of interest in non-Hermitian Floquet topological matters due to their exotic properties resulting from the interplay between driving fields and non-Hermiticity. The present review sums up our studies on non-Hermitian Floquet topological matters in one and two spatial dimensions. We first give a bird's-eye view of the literature for clarifying the physical significance of non-Hermitian Floquet systems. We then introduce, in a pedagogical manner, a number of useful tools tailored for the study of non-Hermitian Floquet systems and their topological properties. With the aid of these tools, we present typical examples of non-Hermitian Floquet topological insulators, superconductors, and quasicrystals, with a focus on their topological invariants, bulk-edge correspondences, non-Hermitian skin effects, dynamical properties, and localization transitions. We conclude this review by summarizing our main findings and presenting our vision of future directions.

 $^{^{\}ast}$ zhoulw
13@u.nus.edu

[†] zdj@sdu.edu.cn

CONTENTS

I. Introduction	3
II. Backgrounds	7
A. Floquet theorem	8
B. Floquet eigenvalue equation	10
C. Floquet effective Hamiltonian and high-frequency expansion	14
D. Adiabatic perturbation theory	20
E. Symmetry and topological characterization	24
F. Dynamical indicators	30
1. Dynamic winding number (DWN)	30
2. Mean chiral displacement (MCD)	32
G. Localization transition and mobility edge	34
III. Non-Hermitian Floquet phases of matter	38
A. Non-Hermitian Floquet exceptional topology	38
B. Non-Hermitian Floquet topological insulators	42
1. First-order topological phase	42
2. Second-order topological phase	49
3. qth-root topological phase	52
C. Non-Hermitian Floquet topological superconductors	56
D. Non-Hermitian Floquet quasicrystals	59
IV. Conclusion and outlook	64
Acknowledgments	65
References	65

I. INTRODUCTION

When a physical system is driven periodically in time, its properties could be drastically modified, leading to new phases and phenomena beyond the static limit [1-6]. One such example, which can be traced back to the early history of human civilization, is the Archimedes screw pump. Under periodic operations, the Archimedes screw could transfer water from low-lying rivers into high-lying irrigation ditches, rather than letting the water to follow its natural flowing tendency. Another example is the tide caused by the combined effects of the gravitational forces exerted by the Moon and its periodic orbiting around the Earth. In the quantum domain, rich nonequilibrium features have been identified in periodically driven systems over the past decades, such as the Rabi oscillation [7–9], stimulated Raman adiabatic passage [10–12], dynamical localization [13–15], Thouless pump [16–18], time crystal [19–21], Floquet topological phase [22–24] and integer quantum Hall effect from chaos [25-27] (for more information, see the reviews [28-60] and the references therein). Among all these discoveries, the Floquet topological matter stands out as one pivotal impetus in the study of periodically driven quantum systems. Over the past twenty years, it has attracted great attention in the context of quantum dynamics [33, 36, 37], quantum simulation [34, 35, 43], condensed matter physics [38, 39, 42], and so on.

The topological properties of a periodically driven system are mainly coded in its Floquet states, which are eigenstates of the evolution operator over a driving period. This one-period propagator is called the Floquet operator. For a system described by the Hamiltonian $\hat{H}(t) = \hat{H}(t+T)$, the Floquet operator can be expressed as $\hat{U} = \hat{\mathsf{T}} e^{-i \int_{t_0}^{t_0+T} \hat{H}(t) dt}$. Here t denotes time, T is the driving period, t_0 denotes the initial time of the period, $\hat{\mathsf{T}}$ performs the time ordering, and we have set the Planck constant $\hbar = 1$. Solving the eigenvalue equation $\hat{U}|\psi\rangle = e^{-iE}|\psi\rangle$ gives us the Floquet eigenstate $|\psi\rangle$ and the quasienergy E. The latter is a phase factor and defined modulus 2π . Its range $E \in [-\pi, \pi)$ is usually referred to as the Floquet quasienergy Brillouin zone. Moreover, the quasienergy is stroboscopically conserved and plays the role of "energy" in driven systems with only discrete-time translational symmetries. If the system also obeys the spatial translational symmetry, the quasienergy bands, which are thus called Floquet bands. Under appropriate conditions, these quasienergy bands could show nontrivial topological properties, which can be captured by topological invariants of the corresponding Floquet-Bloch states or the Floquet operator itself. Periodically driven quantum systems could thus support a new class of nonequilibrium phases, which is nowadays known as the Floquet topological matter.

Similar to static topological phases, Floquet topological phases could also possess symmetry-protected edge states under open boundary conditions and exhibit quantized dynamical or transport signals. However, three key features distinguish Floquet topological phases from their static counterparts. First, suitable driving fields could break the symmetry dynamically and open gaps around the touching points of static energy bands, yielding band inversions and topologically nontrivial bandstructures [22]. This aspect of driving can usually be taken into account by the Floquet effective Hamiltonians obtained from high-frequency expansions of the driving potential |61-64|. Second, as the quasienergy E is bounded from above $(E = \pi)$ and below $(E = -\pi)$, Floquet bands could meet with each other at $E = \pm \pi$ and develop nontrivial windings around the whole quasienergy Brillouin zone $E \in [-\pi, \pi)$. These spectral windings result in unique states of matter in periodically driven systems, such as Floquet semimetals with Floquet band holonomy [65–67], degenerated edge modes at $E = \pm \pi$ (anomalous Floquet π modes) [68–71] and anomalous chiral edge states in Floquet topological insulators [72–74], which have no counterparts in static systems. Third, the driving field could assist in the formation of long-ranged, and even spatially non-decaying coupling among different degrees of freedom in a lattice [24], leading to Floquet phases with large topological invariants, rich topological transitions, and a substantial number of topological edge states [75-81]. These phases and boundary states go beyond the description of any static tight-binding models in typical situations (i.e., with finite ranged or spatially decaying hopping amplitudes). The investigation of the above-mentioned features have not only led to new classification schemes of Floquet matter in theories [82–85] but also promoted experimental realizations of numerous Floquet topological phases in solid-state and quantum simulator setups [86–101], bringing new hope for applications in ultrafast electronics [38] and topological quantum computing [102–104].

Non-Hermitian physics deals with classical and open quantum systems subject to measurements, dissipation, gain and loss or nonreciprocal effects [105–108]. A non-Hermitian system can usually be modeled by a Hamiltonian \hat{H} that is not self-adjoint, i.e., $\hat{H} \neq \hat{H}^{\dagger}$. The spectrum of such a non-Hermitian Hamiltonian is complex in general and the resulting dynamics is non-unitary. In recent years, the study of topological phases in non-Hermitian




FIG. 1. Non-Hermitian Floquet topological matter: a schematic diagram to illustrate the concepts and definitions of several typical phases.

systems have attracted much attention from both theoretical and experimental sides (see Refs. [109–133] for reviews). The investigation of these nonequilibrium phases may help us deepening our understanding of topological matter in open systems and optimize the design of noise-resilient quantum devices.

The recent booming of non-Hermitian topological matter is driven by a couple of key concepts, including the parity and time-reversal (PT) symmetry, the exceptional point (EP), and the non-Hermitian skin effect (NHSE). The PT symmetry can appear in open systems with balanced gain and loss. A PT-symmetric, non-Hermitian Hamiltonian has a real spectrum in the PT unbroken regime [134, 135]. The PT-symmetry was thus viewed as a possible way to lift the restriction of Hermiticity for Hamiltonians in its early studies [108]. Recently, it was found that the topological and transport properties of a system may also change when it undergoes a PT-symmetry-breaking transition, yielding phases unique to non-Hermitian Hamiltonians [136–138]. Models with the PT symmetry thus become focuses in the study of non-Hermitian topological matter [111–113, 115, 116, 119, 125, 133]. The combination of PT and other symmetries further results in various classification schemes for non-Hermitian topological phases that go beyond their Hermitian counterparts [139–147]. The EP is a class of level degeneracy point unique to non-Hermitian operators. At an EP, the geometric and algebraic multiplicities of a non-Hermitian matrix are different, leading to the breakdown of its diagonalizability and the coalescence of its eigenvectors [105, 109, 110]. In recent years, various gapless band structures (nodal points, lines, loops, surfaces, knots etc.) formed by EPs were discovered, giving rise to rich non-Hermitian topological (semi-)metallic phases with intriguing transport properties [117, 119, 123, 124, 126, 148]. Besides, EPs were also found to play key roles in the topological energy transfer [149–151], high-precision sensing [152–155], topological lasers [156–158] and strongly correlated phases [123, 124]. The NHSE refers to the accumulation of bulk states around the edges of an open-boundary non-Hermitian lattice. It highlights the extreme sensitivity of non-Hermitian physics to the boundary condition of a system [121, 127, 132]. This phenomenon not only blurs the distinction between bulk and edge states in non-Hermitian models but also breaks the wellestablished bulk-boundary correspondence in Hermitian topological matter [159–163]. Over the past few years, a couple of theoretical frameworks have been introduced to characterize the NHSE and its related topological phenomena [164-174], which are accompanied by the experimental observations of NHSE in AMO systems, electrical circuits and metamaterials [175–181]. Entanglement transitions associated with the NHSE was also identified in a recent study [182]. Further progresses have been made in the study of non-Hermitian topological matter in randomly disordered [183–187], quasiperiodic [188–192] and many-body systems [193-207].

With all these developments, a natural follow-up is to consider the system in a more general situation, in which it is subject to both time-periodic drivings and non-Hermitian effects. Such non-Hermitian Floquet systems may possess exotic dynamical phenomena and topological phases with no static or Hermitian analogies. On the theoretical side, the investigation of driven non-Hermitian systems may lead to the discovery of new topological states and bring about the improvement of classification schemes for nonequilibrium phases of matter in general. On the practical side, the exploration of non-Hermitian Floquet matter is helpful to the design of new approaches for preparing or stabilizing topologically nontrivial states and controlling material properties. It also stimulates new ideas for realizing quantum devices and quantum computing protocols that are robust to perturbations caused by the environment. Though still at the early stage, many progresses have been made in the realization and characterization of non-Hermitian Floquet phases [208–253]. In this review, we limit our scope to the discussion of a number of typical topological phases we discovered in non-Hermitian Floquet systems [254–264]. In Sec. II, we give a pedagogical introduction to some key aspects of Floquet systems, including their dynamical and topological characterizations. In Sec. III, we present typical examples of non-Hermitian Floquet topological insulators, superconductors, quasicrystals and summarize their main physical properties, with a focus on the features that are unique to driven non-Hermitian systems. In Sec. IV, we conclude this review, briefly mention some relevant studies and discuss potential future work.

II. BACKGROUNDS

We start with a recap of the basic description of a non-Hermitian Floquet system. The Hamiltonian of such a system satisfies $\hat{H}(t) = \hat{H}(t+T)$, and there exists $t \in [0,T]$ such that $\hat{H}(t) \neq [\hat{H}(t)]^{\dagger}$. Here t denotes time and T is the driving period. The state of the system evolves according to the Schrödinger equation

$$i\frac{\partial}{\partial t}|\Psi(t)\rangle = \hat{H}(t)|\Psi(t)\rangle,\tag{1}$$

where we have set $\hbar = 1$. We first show that this equation can be solved by Floquet states even though $\hat{H}(t)$ is non-Hermitian. This is followed by different ways of obtaining the Floquet states in general situations, in the high-frequency regime, and in the adiabatic regime. We next discuss the symmetry, topological invariants, and dynamical characterizations of non-Hermitian Floquet states, with a focus on the types of physical systems explored in our previous studies. We conclude this section by presenting some tools for characterizing the spectrum properties and localization transitions in non-Hermitian Floquet disordered systems.

A. Floquet theorem

We sketch a proof of the Floquet theorem in this subsection [265]. It follows the proof of the Bloch theorem for waves in one-dimensional periodic lattices [266]. Applying the Fourier expansion to our time-periodic Hamiltonian $\hat{H}(t)$ in the time-frequency domain, we find

$$\hat{H}(t) = \sum_{n} \hat{H}(\omega_{n})e^{i\omega_{n}t}, \qquad \omega_{n} = n\omega = n\frac{2\pi}{T}, \qquad n \in \mathbb{Z}.$$
(2)

Here ω is the driving frequency. For an infinite system, the set of plane waves $\{|\psi_{\varepsilon}(t)\rangle = |\varphi(\mathbf{r})\rangle e^{-i\varepsilon t}\}$ can be chosen as a suitable basis. We can write down a matrix expression for $\hat{H}(t)$ in the orthonormal and complete basis $\{|\psi_{\varepsilon}(t)\rangle\}$. Acting $\hat{H}(t)$ on $|\psi_{\varepsilon}(t)\rangle$, we obtain

$$\hat{H}(t)|\psi_{\varepsilon}(t)\rangle = \sum_{n} \hat{H}(\omega_{n})|\varphi(\mathbf{r})\rangle e^{-i(\varepsilon-\omega_{n})t} = \sum_{n} \hat{H}(\omega_{n})|\psi_{\varepsilon-\omega_{n}}(t)\rangle.$$
(3)

For any given $n \in \mathbb{Z}$, the state $|\psi_{\varepsilon-\omega_n}(t)\rangle$ belongs to the subspace

$$S_{\varepsilon} = \{ |\psi_{\varepsilon}(t)\rangle, |\psi_{\varepsilon\pm\omega}(t)\rangle, |\psi_{\varepsilon\pm2\omega}(t)\rangle, ..., |\psi_{\varepsilon\pm n\omega}(t)\rangle, ... \}.$$
(4)

It is clear that any two subspaces S_{ε} and $S_{\varepsilon'}$ ($\varepsilon' \neq \varepsilon$) are decoupled under the action of $\hat{H}(t)$ if $\operatorname{Re}(\varepsilon - \varepsilon') \in (-\omega, \omega)$. Moreover, S_{ε} and $S_{\varepsilon'}$ are equivalent if there exists an $n \in \mathbb{Z}$ such that $\varepsilon - \varepsilon' = n\omega$. Therefore, we can study the dynamics of the system separately in each subspace S_{ε} for $\operatorname{Re}(\varepsilon) \in [-\pi/T, \pi/T)$, which is usually called the first quasienergy Brillouin zone (BZ). The quasienergy ε is thus a conserved quantity due to the discrete-time translational symmetry of $\hat{H}(t)$, similar to the conserved quasimomentum due to the discrete-space translational symmetry of a static Hamiltonian. In the subspace S_{ε} , a solution of the Schrödinger equation can be written as

$$|\Psi_{\varepsilon}(t)\rangle = \sum_{n} c(\varepsilon + \omega_{n})|\psi_{\varepsilon + \omega_{n}}(t)\rangle = e^{-i\varepsilon t} \sum_{n} c(\varepsilon + \omega_{n})|\varphi(\mathbf{r})\rangle e^{-i\omega_{n}t},$$
(5)

where $\{c(\varepsilon + \omega_n)\}$ are complex coefficients. It is clear that Eq. (5) possesses a time-periodic component

$$|u_{\varepsilon}(t)\rangle \equiv \sum_{n} c(\varepsilon + \omega_{n})|\varphi(\mathbf{r})\rangle e^{-i\omega_{n}t} = |u_{\varepsilon}(t+T)\rangle.$$
(6)

Therefore, we can express the general solution $|\Psi_{\varepsilon}(t)\rangle$ as the product of an oscillating phase factor $e^{-i\varepsilon t}$ and a time-periodic Floquet mode $|u_{\varepsilon}(t)\rangle$, i.e.,

$$|\Psi_{\varepsilon}(t)\rangle = e^{-i\varepsilon t}|u_{\varepsilon}(t)\rangle.$$
(7)

It also implies that

$$|\Psi_{\varepsilon}(t+T)\rangle = e^{-i\varepsilon T}|\Psi_{\varepsilon}(t)\rangle.$$
(8)

The latter equation indicates that the only change of the state $|\Psi_{\varepsilon}(t)\rangle$ after undergoing a oneperiod evolution is to pick up an exponential factor $e^{-i\varepsilon T}$. We refer to the set $\{|\Psi_{\varepsilon}(t)\rangle|\text{Re}(\varepsilon) \in [-\pi/T, \pi/T)\}$ as Floquet eigenstates of the system. They form an orthonormal and complete basis at each instant of time t.

Since the evolution from $|\Psi_{\varepsilon}(t)\rangle$ to $|\Psi_{\varepsilon}(t+T)\rangle$ is governed by the Schrödinger equation, we can also express Eq. (8) as

$$\hat{U}(t+T,t)|\Psi_{\varepsilon}(t)\rangle = e^{-i\varepsilon T}|\Psi_{\varepsilon}(t)\rangle,\tag{9}$$

where $\hat{U}(t+T,t) = \hat{\mathsf{T}}e^{-i\int_{t}^{t+T}\hat{H}(t')dt'}$ is nothing but the Floquet operator (evolution operator over one driving period) of the system. When we are only concerned with the stroboscopic dynamics, the initial time dependence of Eq. (9) is not important. In this case, we can set t = 0 in Eq. (9) and express the Floquet eigenvalue equation as

$$\hat{U}|\Psi_E\rangle = e^{-iE}|\Psi_E\rangle,\tag{10}$$

where $\hat{U} = \hat{\mathsf{T}} e^{-i \int_0^T \hat{H}(t) dt}$ and we have introduced $E = \varepsilon T$ as the dimensionless quasienergy, whose first BZ is given by $[-\pi, \pi)$. To sum up, we find that the solution of Eq. (1) with a time-periodic $\hat{H}(t)$ has the form of Eq. (7) or (8), where $|\Psi_{\varepsilon}(t)\rangle$ is an eigenstate of the Floquet operator of the system. For stroboscopic observations, all the Floquet eigenstates can thus be obtained by solving the eigenvalue equation (10) of Floquet operator \hat{U} . Due to the completeness of Floquet eigenstate basis $\{|\Psi_E\rangle\}$, we can expand an arbitrary initial state of the system as $|\Psi(0)\rangle = \sum_E c_E |\Psi_E\rangle$. The resulting state after an evolution over n driving periods is then given by

$$|\Psi(t)\rangle = \sum_{E} c_{E} e^{-inE} |\Psi_{E}\rangle.$$
(11)

Note that for a non-Hermitian $\hat{H}(t)$, \hat{U} is generally non-unitary and the quasienergy E may have a nonvanishing imaginary part. In this case, the real part of E still belongs to the range of $[-\pi,\pi)$ and our arguments leading to the general solution (11) can be satisfied.

B. Floquet eigenvalue equation

In the most general situations, we can solve Eq. (10) numerically by the split-operator method [3]. Dividing the evolution periodic T into N segments with a large enough N, we can express the Floquet operator \hat{U} approximately as

$$\hat{U} \simeq e^{-i\hat{H}((N-1)\Delta t)\Delta t} e^{-i\hat{H}((N-2)\Delta t)\Delta t} \cdots e^{-i\hat{H}(\Delta t)\Delta t} e^{-i\hat{H}(0)\Delta t},$$
(12)

where $\Delta t = T/N$. Over each small time interval Δt , $\hat{H}(t)$ is approximately time-independent and we can diagonalize it numerically at $t = \ell \Delta t$ as

$$\hat{H}(\ell \Delta t) = V_{\ell} D_{\ell} V_{\ell}^{-1}.$$
(13)

Each column of V_{ℓ} represents a right eigenvector of $\hat{H}(\ell \Delta t)$. The evolution operator over the one-time interval Δt then takes the form

$$e^{-i\hat{H}(\ell\Delta t)\Delta t} = V_{\ell}e^{-iD_{\ell}\Delta t}V_{\ell}^{-1}, \qquad \ell = 0, 1, ..., N - 1.$$
(14)

The multiplication of all the $V_{\ell}e^{-iD_{\ell}\Delta t}V_{\ell}^{-1}$ from right to left for $\ell = 0$ to N-1 yields Eq. (12), which further converges to the exact Floquet operator in the limit $N \to \infty$. We can thus numerically solve the Floquet eigenvalue equation by diagonalizing the approximated \hat{U} in Eq. (12). This approach works in principle for systems with any individual or multiple driving frequencies. But it may become time-consuming in practice for certain continuously or slowly driven systems.

When the driving field takes the form of periodic kicking or quenching, the series in

Eq. (12) can be greatly simplified and even obtained exactly. Here we give several examples. Consider a time-periodic Hamiltonian of the form

$$\hat{H}(t) = \hat{H}_0 + \sum_{\ell \in \mathbb{Z}} \delta(t/T - \ell) \hat{H}_1,$$
(15)

where $\delta(t/T - \ell)$ is the delta function peaked at $t = \ell T$, i.e., each integer multiple of the driving period. The dynamics over each driving period thus constitutes a free evolution part controlled by \hat{H}_0 and a delta kicking force controlled by \hat{H}_1 . The quantum kicked rotor is one representative example of such a system [28]. Focusing on the one-period evolution from $t = \ell T - 0^+$ to $t = (\ell + 1)T - 0^+$, we find the Floquet operator of $\hat{H}(t)$ to be

$$\hat{U} = e^{-i\int_{\ell T+0^+}^{(\ell+1)T-0^+} \hat{H}(t)dt} e^{-i\int_{\ell T-0^+}^{\ell T+0^+} \hat{H}(t)dt}
= e^{-i\int_{\ell T+0^+}^{(\ell+1)T-0^+} \hat{H}_0dt} e^{-i\int_{\ell T-0^+}^{\ell T+0^+} \delta(t/T-\ell)\hat{H}_1dt}
= e^{-i\hat{H}_0T} e^{-i\hat{H}_1T}.$$
(16)

Similarly, if there are two kicks separated by a time interval τ within each driving period, the Hamiltonian could take the form of

$$\hat{H}(t) = \hat{H}_0 + \sum_{\ell \in \mathbb{Z}} \delta(t/T - \tau/T - \ell) \hat{H}_2 + \sum_{\ell \in \mathbb{Z}} \delta(t/T - \ell) \hat{H}_1.$$
(17)

For the evolution from time $t = \ell T - 0^+$ to $t = (\ell + 1)T - 0^+$, the Floquet operator now takes the form of

$$\hat{U} = e^{-i\hat{H}_0(T-\tau)} e^{-i\hat{H}_2 T} e^{-i\hat{H}_0 \tau} e^{-i\hat{H}_1 T}.$$
(18)

One typical example of such a system is the double-kicked quantum rotor [24]. For a periodically quenched Hamiltonian in the form of

$$\hat{H}(t) = \begin{cases} \hat{H}_1 & t \in [\ell T, \ell T + T_1) \\ \hat{H}_2 & t \in [\ell T + T_1, \ell T + T_1 + T_2) \end{cases},$$
(19)

where $T = T_1 + T_2$, we can also directly obtain the corresponding Floquet operator from

 $t = \ell T - 0^+$ to $t = (\ell + 1)T - 0^+$ as

$$\hat{U} = e^{-i\int_{\ell T+T_1}^{\ell T+T-0^+} \hat{H}(t)dt} e^{-i\int_{\ell T-0^+}^{\ell T+T_1-0^+} \hat{H}(t)dt}
= e^{-i\int_{\ell T+T_1}^{\ell T+T-0^+} \hat{H}_2dt} e^{-i\int_{\ell T-0^+}^{\ell T+T_1-0^+} \hat{H}_1dt}
= e^{-i\hat{H}_2T_2} e^{-i\hat{H}_1T_1}.$$
(20)

The discrete-time quantum walk can be viewed as one example of such a periodically quenched system [31]. Time-periodic quenches are also frequently implemented in the study of discrete time crystals [19–21]. When $[\hat{H}_1, \hat{H}_2] \neq 0$, the quenches (or kicks) may effectively generate long-range coupling in the system according to the Baker-Campbell-Hausdorff formula, leading to Floquet phases with large topological invariants and many boundary states. This point will be explicitly demonstrated by the examples discussed in Sec. III.

For a continuously driven system, the solution of the Floquet eigenvalue equation may also be obtained approximately in terms of the frequency (Sambe) space formalism [267–269]. Inserting the Floquet state in Eq. (7) into the Schrödinger equation (1) and reorganizing the terms, we find

$$\left[\hat{H}(t) - i\frac{\partial}{\partial t}\right] |u_{\varepsilon}(t)\rangle = \varepsilon |u_{\varepsilon}(t)\rangle.$$
(21)

Using the Fourier expansion of $\hat{H}(t)$ and $|u_{\varepsilon}(t)\rangle = \sum_{n} e^{-i\omega_{n}t} |u_{\varepsilon}(\omega_{n})\rangle$, we further obtain

$$\sum_{m,n} \hat{H}(\omega_m) e^{i(\omega_m - \omega_n)t} |u_{\varepsilon}(\omega_n)\rangle - \sum_n \omega_n e^{-i\omega_n t} |u_{\varepsilon}(\omega_n)\rangle = \varepsilon \sum_n e^{-i\omega_n t} |u_{\varepsilon}(\omega_n)\rangle.$$
(22)

Multiplying $\frac{1}{T}e^{i\omega_{\ell}t}$ from the left on both sides of Eq. (22) and performing the integral over a driving period T, we arrive at

$$\sum_{m} (\hat{H}_{m-n} - \omega_m \delta_{m,n}) | u_{\varepsilon}(\omega_m) \rangle = \varepsilon | u_{\varepsilon}(\omega_n) \rangle, \qquad (23)$$

where

$$\hat{H}_{m-n} \equiv \hat{H}(\omega_{m-n}) = \frac{1}{T} \int_0^T e^{-i\omega_{m-n}t} \hat{H}(t) dt, \qquad (24)$$

and $\omega_{m-n} = (m-n)\omega = (m-n)\frac{2\pi}{T}$. Equation (23) is an infinite-dimensional matrix equation

of the form

$$\begin{bmatrix} \ddots & \ddots \\ \ddots & \hat{H}_{0} + 2\omega & \hat{H}_{1} & \hat{H}_{2} & \hat{H}_{3} & \hat{H}_{4} & \ddots \\ \ddots & \hat{H}_{-1} & \hat{H}_{0} + \omega & \hat{H}_{1} & \hat{H}_{2} & \hat{H}_{3} & \ddots \\ \ddots & \hat{H}_{-2} & \hat{H}_{-1} & \hat{H}_{0} & \hat{H}_{1} & \hat{H}_{2} & \ddots \\ \ddots & \hat{H}_{-3} & \hat{H}_{-2} & \hat{H}_{-1} & \hat{H}_{0} - \omega & \hat{H}_{1} & \ddots \\ \ddots & \hat{H}_{-4} & \hat{H}_{-3} & \hat{H}_{-2} & \hat{H}_{-1} & \hat{H}_{0} - 2\omega & \ddots \\ \ddots & \ddots \\ \end{bmatrix} \begin{bmatrix} \vdots \\ |u_{\varepsilon}(\omega_{-1})\rangle \\ |u_{\varepsilon}(\omega_{0})\rangle \\ |u_{\varepsilon}(\omega_{1})\rangle \\ |u_{\varepsilon}(\omega_{2})\rangle \\ \vdots \end{bmatrix} = \varepsilon \begin{bmatrix} \vdots \\ |u_{\varepsilon}(\omega_{-1})\rangle \\ |u_{\varepsilon}(\omega_{0})\rangle \\ |u_{\varepsilon}(\omega_{1})\rangle \\ |u_{\varepsilon}(\omega_{2})\rangle \\ \vdots \end{bmatrix} .$$
(25)

Note that each \hat{H}_{m-n} has the same Hilbert space dimension d as the original Hamiltonian $\hat{H}(t)$. As an example, for a harmonically driven system described by the Hamiltonian

$$\hat{H}(t) = \hat{H}_0 + \hat{V}e^{i\omega t} + \hat{W}e^{-i\omega t}, \qquad (26)$$

the above equation reduces to the following block tridiagonal form

$$\begin{bmatrix} \ddots & \ddots \\ \ddots & \hat{H}_{0} + 2\omega & \hat{V} & 0 & 0 & 0 & \ddots \\ \ddots & \hat{W} & \hat{H}_{0} + \omega & \hat{V} & 0 & 0 & \ddots \\ \ddots & 0 & \hat{W} & \hat{H}_{0} & \hat{V} & 0 & \ddots \\ \ddots & 0 & 0 & \hat{W} & \hat{H}_{0} - \omega & \hat{V} & \ddots \\ \ddots & 0 & 0 & 0 & \hat{W} & \hat{H}_{0} - 2\omega & \ddots \\ \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \end{bmatrix} \begin{bmatrix} \vdots \\ |u_{\varepsilon}(\omega_{-2})\rangle \\ |u_{\varepsilon}(\omega_{-1})\rangle \\ |u_{\varepsilon}(\omega_{0})\rangle \\ |u_{\varepsilon}(\omega_{0})\rangle \\ |u_{\varepsilon}(\omega_{1})\rangle \\ |u_{\varepsilon}(\omega_{2})\rangle \\ \vdots \end{bmatrix} = \varepsilon \begin{bmatrix} \vdots \\ |u_{\varepsilon}(\omega_{-2})\rangle \\ |u_{\varepsilon}(\omega_{-1})\rangle \\ |u_{\varepsilon}(\omega_{0})\rangle \\ |u_{\varepsilon}(\omega_{1})\rangle \\ |u_{\varepsilon}(\omega_{2})\rangle \\ \vdots \end{bmatrix} .$$
(27)

In practical calculations, one should truncate the infinite-dimensional matrix in Eq. (25) at a sufficiently high harmonics $N\omega$, leading to an $Nd \times Nd$ dimensional Floquet effective Hamiltonian, whose eigenvalue problem can be numerically solved. Assuming the characteristic energy scale of \hat{H}_{m-n} for all m - n to be Ω , we can take a smaller N to do the truncation for a larger ratio of ω/Ω , i.e., for a high-frequency driving field. Instead, for a resonantly or slowly driven system, more harmonics should be kept during the truncation. All the discussions presented in this subsection hold for both Hermitian and non-Hermitian Hamiltonians $\hat{H}(t)$.

C. Floquet effective Hamiltonian and high-frequency expansion

From the Floquet operator \hat{U} of a periodically driven system, one can formally define its Floquet effective Hamiltonian as

$$\hat{H}_{\text{eff}} = \frac{i}{T} \ln \hat{U} \Leftrightarrow \hat{U} = e^{-i\hat{H}_{\text{eff}}T}.$$
(28)

Taking into account the fact that the quasienergies are defined modulus $2\pi/T$, the \hat{H}_{eff} contains the same physical information as \hat{U} . Yet, it provides us with more room to treat the properties of Floquet systems in analogy with static Hamiltonian models. For a continuously driven system, the explicit form of \hat{H}_{eff} is usually involved. This can be inspected from Eq. (12), as the $\hat{H}(\ell \Delta t)$ and $\hat{H}(m\Delta t)$ do not commute for $\ell \neq m$ in general. When the frequency of the driving field is high enough, an approximate series expression for \hat{H}_{eff} can be obtained via high-frequency expansion methods [61–64]. Here we recap one such method in its full generality, which is applicable to both Hermitian and non-Hermitian systems.

We first assume that the time-periodic Hamiltonian $\hat{H}(t)$ can be decomposed into a static part \hat{H}_0 and a periodically modulated part $\hat{V}(t)$, i.e.,

$$\hat{H}(t) = \hat{H}_0 + \hat{V}(t), \qquad \hat{V}(t) = \hat{V}(t+T).$$
(29)

Here $T = 2\pi/\omega$ is the driving period with ω denoting the driving frequency. Next, we apply a similarity transformation to the evolved state $|\Psi(t)\rangle$ in the Schrödinger equation (1), yielding a rotated state

$$|\Phi(t)\rangle = \hat{\mathcal{U}}(t)|\Psi(t)\rangle = e^{i\hat{K}(t)}|\Psi(t)\rangle.$$
(30)

Here $\hat{K}(t)$ is sometimes called the kick operator. It encodes the information regarding the micromotion dynamics of the system. Plugging $|\Psi(t)\rangle = e^{-i\hat{K}(t)}|\Phi(t)\rangle$ into Eq. (1) leads to the transformed Schrödinger equation

$$i\frac{\partial}{\partial t}|\Phi(t)\rangle = \hat{H}_{\text{eff}}|\Phi(t)\rangle,$$
(31)

where

$$\hat{H}_{\text{eff}} = e^{i\hat{K}(t)}\hat{H}(t)e^{-i\hat{K}(t)} + i\frac{de^{i\hat{K}(t)}}{dt}e^{-i\hat{K}(t)}.$$
(32)

The aim of the high-frequency expansion method is to find a time-independent \hat{H}_{eff} by transferring all time-dependent terms into the kick operator $\hat{K}(t)$. When such a purpose is formally achieved, we can express the Floquet evolution of a system as

$$\hat{U}(t_1, t_0) |\Psi(t_0)\rangle = e^{-i\hat{K}(t_1)} e^{-i\hat{H}_{\text{eff}}(t_1 - t_0)} e^{i\hat{K}(t_0)} |\Psi(t_0)\rangle.$$
(33)

That is, the system is subject to an initial kick associated with the operator $\hat{K}(t_0)$, then evolved under the time-independent \hat{H}_{eff} , and finally subject to a second kick carried out by the operator $\hat{K}(t_1)$. There is no need to perform any time-ordered integral in the calculation of $\hat{U}(t_1, t_0)$.

Assuming the driving frequency ω to be large, we may expand \hat{H}_{eff} and $\hat{K}(t)$ into power series of $1/\omega$ as

$$\hat{H}_{\text{eff}} = \sum_{n=0}^{\infty} \frac{1}{\omega^n} \hat{H}_{\text{eff}}^{(n)}, \qquad \hat{K} = \sum_{n=1}^{\infty} \frac{1}{\omega^n} \hat{K}^{(n)}.$$
 (34)

In the meantime, we can apply Taylor expansions to the two terms in Eq. (32) and express \hat{H}_{eff} as

$$\hat{H}_{\text{eff}} = \sum_{n=0}^{\infty} \frac{i^n}{n!} \underbrace{\tilde{(K, [\hat{K}, ..., \hat{K}, [\hat{K}, \hat{H}]]]}}_{+\sum_{n=1}^{\infty} i \frac{i^n}{n!}} \underbrace{\tilde{(K, [\hat{K}, ..., \hat{K}, [\hat{K}, \partial_t \hat{K}]]]}}_{(35)}$$

Note that we have concealed the explicit time-dependence of $\hat{K}(t)$ and $\hat{K}^{(n)}(t)$ in the above two equations for brevity. To proceed, we impose two further requirements for $\hat{K}(t)$,

$$\hat{K}(t) = \hat{K}(t+T), \qquad \int_{0}^{T} \hat{K}(t)dt = 0.$$
 (36)

Inserting Eq. (34) and the Fourier expansion

$$\hat{H}(t) = \hat{H}_0 + \sum_{m \neq 0} \hat{V}_m e^{im\omega t}$$
(37)

into Eq. (35), we arrive at

$$\sum_{n=0}^{\infty} \frac{1}{\omega^{n}} \hat{H}_{\text{eff}}^{(n)}$$

$$= \sum_{n=0}^{\infty} \frac{i^{n}}{n!} [\overbrace{\sum_{p} \frac{1}{\omega^{p}} \hat{K}^{(p)}, [\sum_{q} \frac{1}{\omega^{q}} \hat{K}^{(q)}, ..., \sum_{r} \frac{1}{\omega^{r}} \hat{K}^{(r)}, [\sum_{s} \frac{1}{\omega^{s}} \hat{K}^{(s)}, \left(\hat{H}_{0} + \sum_{m \neq 0} \hat{V}_{m} e^{im\omega t}\right)]]]$$

$$+ \sum_{n=1}^{\infty} \frac{i^{n+1}}{n!} [\overbrace{\sum_{p} \frac{1}{\omega^{p}} \hat{K}^{(p)}, [\sum_{q} \frac{1}{\omega^{q}} \hat{K}^{(q)}, ..., \sum_{r} \frac{1}{\omega^{r}} \hat{K}^{(r)}, [\sum_{s} \frac{1}{\omega^{s}} \hat{K}^{(s)}, \partial_{t} \sum_{m} \frac{1}{\omega^{m}} \hat{K}^{(m)}]]]. \quad (38)$$

The high-frequency expansions of \hat{H}_{eff} and \hat{K} are obtained by equating both sides of Eq. (38) at each order of $1/\omega$.

We can now carry out the calculations order-by-order to find the first few terms in the series of \hat{H}_{eff} and \hat{K} . At the zeroth order, we could obtain

$$\hat{H}_{\rm eff}^{(0)} = \hat{H}_0 + \hat{V}(t) - \frac{1}{\omega} \partial_t \hat{K}^{(1)}$$
(39)

from Eq. (38). To ensure that $\hat{H}_{\rm eff}^{(0)}$ is time-independent, we must choose

$$\frac{1}{\omega}\partial_t \hat{K}^{(1)} = \hat{V}(t) = \sum_{m \neq 0} \hat{V}_m e^{im\omega t},\tag{40}$$

such that

$$\hat{K}^{(1)} = \sum_{m \neq 0} \frac{1}{im} \hat{V}_m e^{im\omega t}.$$
(41)

Therefore, up to the zeroth order of $1/\omega$, we have the following approximation for the effective Hamiltonian

$$\hat{H}_{\text{eff}} \approx \hat{H}_{\text{eff}}^{(0)} = \hat{H}_0. \tag{42}$$

Up to the first order of $1/\omega$, we have the following approximation for the kick operator

$$\hat{K}(t) \approx \frac{1}{\omega} \hat{K}^{(1)} = \frac{1}{\omega} \sum_{m \neq 0} \frac{\hat{V}_m}{im} e^{im\omega t}.$$
(43)

At the first order, we could find from Eq. (38) that

$$\hat{H}_{\text{eff}}^{(1)} = i \left[\hat{K}^{(1)}, \hat{H}_{0} + \hat{V}(t) \right] - \frac{1}{\omega} \partial_{t} \hat{K}^{(2)} - \frac{i}{2} \left[\hat{K}^{(1)}, \frac{1}{\omega} \partial_{t} \hat{K}^{(1)} \right]
= i \left[\hat{K}^{(1)}, \hat{H}_{0} + \hat{V}(t) \right] - \frac{1}{\omega} \partial_{t} \hat{K}^{(2)} - \frac{i}{2} \left[\hat{K}^{(1)}, \hat{V}(t) \right]
= i \left[\hat{K}^{(1)}, \hat{H}_{0} \right] + \frac{i}{2} \left[\hat{K}^{(1)}, \hat{V}(t) \right] - \frac{1}{\omega} \partial_{t} \hat{K}^{(2)}.$$
(44)

By definition, any non-vanishing elements of $i \left[\hat{K}^{(1)}, \hat{H}_0 \right]$ and $-\frac{1}{\omega} \partial_t \hat{K}^{(2)}$ are time-dependent. The only time-independent term that could contribute to $\hat{H}_{\text{eff}}^{(1)}$ comes from the term $\frac{i}{2} \left[\hat{K}^{(1)}, \hat{V}(t) \right]$, which can be written explicitly as

$$\frac{i}{2} \left[\hat{K}^{(1)}, \hat{V}(t) \right] = \frac{i}{2} \left[\sum_{m \neq 0} \frac{1}{im} \hat{V}_m e^{im\omega t}, \sum_{m' \neq 0} \hat{V}_{m'} e^{im'\omega t} \right]$$
$$= \sum_{m,m' \neq 0} \frac{e^{i(m+m')\omega t}}{2m} \left[\hat{V}_m, \hat{V}_{m'} \right].$$
(45)

The time-independent terms are those with m' = -m. Collecting these terms together, we find

$$\hat{H}_{\text{eff}}^{(1)} = \sum_{m \neq 0} \frac{\left[\hat{V}_m, \hat{V}_{-m}\right]}{2m} = \sum_{m \neq 0} \frac{\hat{V}_m \hat{V}_{-m}}{m}.$$
(46)

Therefore, up to the first order of $1/\omega$, we have the following approximation for the effective Hamiltonian

$$\hat{H}_{\text{eff}} \approx \hat{H}_{\text{eff}}^{(0)} + \frac{1}{\omega} \hat{H}_{\text{eff}}^{(1)} = \hat{H}_0 + \frac{1}{\omega} \sum_{m \neq 0} \frac{\hat{V}_m \hat{V}_{-m}}{m}.$$
(47)

The form of $\hat{K}^{(2)}$ is determined by the remaining time-dependent terms. Performing the integration over time directly, we obtain

$$\hat{K}^{(2)} = \sum_{m \neq 0} \frac{1}{m} \left[\hat{V}_m, \hat{H}_0 \right] \int \omega e^{im\omega t} dt + \sum_{m,m' \neq 0,-m} \frac{\left[\hat{V}_m, \hat{V}_{m'} \right]}{2m} \int \omega e^{i(m+m')\omega t} dt$$
$$= \sum_{m \neq 0} \frac{1}{im^2} \left[\hat{V}_m, \hat{H}_0 \right] e^{im\omega t} + \sum_{m,m' \neq 0,-m} \frac{\left[\hat{V}_m, \hat{V}_{m'} \right]}{2im(m+m')} e^{i(m+m')\omega t}.$$
(48)

Up to the second order of $1/\omega$, we can thus approximate the kick operator by

$$\hat{K}(t) \approx \frac{1}{\omega} \hat{K}^{(1)} + \frac{1}{\omega^2} \hat{K}^{(2)}
= \frac{1}{\omega} \sum_{m \neq 0} \frac{\hat{V}_m}{im} e^{im\omega t} + \frac{1}{\omega^2} \sum_{m \neq 0} \frac{1}{im^2} \left[\hat{V}_m, \hat{H}_0 \right] e^{im\omega t}
+ \frac{1}{\omega^2} \sum_{m,m' \neq 0,-m} \frac{\left[\hat{V}_m, \hat{V}_{m'} \right]}{2im(m+m')} e^{i(m+m')\omega t}.$$
(49)

We could continue these self-consistent calculations to obtain higher-order terms in the high-frequency expansion of \hat{H}_{eff} and $\hat{K}(t)$. For example, for the second-order component $\hat{H}_{\text{eff}}^{(2)}$, we have

$$\hat{H}_{\text{eff}}^{(2)} = \frac{-1}{2} [\hat{K}^{(1)}, [\hat{K}^{(1)}, \hat{H}_0 + \hat{V}(t)]] + i [\hat{K}^{(2)}, \hat{H}_0 + \hat{V}(t)] + \frac{-1}{\omega} \partial_t \hat{K}^{(3)} + \frac{-i}{2\omega} [\hat{K}^{(1)}, \partial_t \hat{K}^{(2)}] + \frac{-i}{2\omega} [\hat{K}^{(2)}, \partial_t \hat{K}^{(1)}] + \frac{1}{6\omega} [\hat{K}^{(1)}, [\hat{K}^{(1)}, \partial_t \hat{K}^{(1)}].$$
(50)

Dropping the time-dependent terms, we are left with

$$\hat{H}_{\text{eff}}^{(2)} = \frac{1}{2} \sum_{m \neq 0} \frac{[[\hat{V}_m, \hat{H}_0], \hat{V}_{-m}]}{m^2} + \frac{1}{3} \sum_{l, m \neq 0} \frac{[\hat{V}_l, [\hat{V}_m, \hat{V}_{-l-m}]]}{lm}.$$
(51)

Therefore, up to the second-order correction in $1/\omega$, the effective Hamiltonian \hat{H}_{eff} turns out to be

$$\hat{H}_{\text{eff}} \approx \hat{H}_{0} + \frac{1}{\omega} \sum_{m \neq 0} \frac{\hat{V}_{m} \hat{V}_{-m}}{m} + \frac{1}{2\omega^{2}} \sum_{m \neq 0} \frac{\left[[\hat{V}_{m}, \hat{H}_{0}], \hat{V}_{-m} \right]}{m^{2}} + \frac{1}{3\omega^{2}} \sum_{l,m \neq 0} \frac{\left[\hat{V}_{l}, [\hat{V}_{m}, \hat{V}_{-l-m}] \right]}{lm}.$$
(52)

This approximation holds for both Hermitian and non-Hermitian Floquet systems under high-frequency driving fields. Note that for stroboscopic evolution, we have $t_1 - t_0 = NT$ with $N \in \mathbb{Z}$ and $\hat{K}(t_1) = \hat{K}(t_0)$ in Eq. (33). In this case, the term $\hat{K}(t_0)$ in Eq. (33) describes an initial phase, which can be set to zero as the expansion of $\hat{K}(t)$ at every order of $1/\omega$ is only determined up to a constant [see Eqs. (41) and (48)]. Therefore, for stroboscopic dynamics, we can simply use the H_{eff} to capture most of the essential physics. In Sec. III, we will showcase the application of the high-frequency methods presented here to the Floquet engineering of non-Hermitian quasicrystals.

To be concrete, we illustrate the usage of \hat{H}_{eff} here with a simple example. Consider a harmonically driven two-level Hamiltonian of the form

$$\hat{H}(t) = (h_x + i\gamma/2)[\cos(\omega t)\sigma_x + \sin(\omega t)\sigma_y] + h_z\sigma_z,$$
(53)

where $h_x, h_z, \gamma \in \mathbb{R}$, and σ_{α} for $\alpha = x, y, z$ are Pauli matrices. It is not hard to verify that by choosing the kick operator as

$$e^{i\hat{K}(t)} = \begin{pmatrix} 1 & 0\\ 0 & e^{-i\omega t} \end{pmatrix},$$
(54)

the Floquet effective Hamiltonian in the rotating frame can be exactly obtained from Eq. (32), i.e.,

$$\hat{H}_{\text{eff}} = \frac{\omega}{2}\sigma_0 + (h_x + i\gamma/2)\sigma_x + (h_z - \omega/2)\sigma_z, \qquad (55)$$

where σ_0 denotes the 2 × 2 identity matrix. We see that the two levels of H(t) have the instantaneous eigenenergies

$$E_{\pm} = \pm \sqrt{h_x^2 + h_z^2 - \gamma^2/4 + ih_x\gamma},$$
(56)

which are time-independent and complex in general. They could meet with each other at a second-order EP with E = 0 when $h_x = 0$ and $h_z = \pm \gamma/2$. Meanwhile, \hat{H}_{eff} in Eq. (55) has two quasienergy levels defined by

$$\varepsilon_{\pm} = \frac{\omega}{2} \pm \sqrt{h_x^2 + \left(h_z - \frac{\omega}{2}\right)^2 - \gamma^2/4 + ih_x\gamma}.$$
(57)

They could meet with each other at a second-order Floquet EP with $\varepsilon = \omega/2$ when $h_x = 0$ and $h_z = \omega/2 \pm \gamma/2$. It is clear that the EP is shifted by the driving in both its energy and location in the parameter space. If we define εT as the dimensionless quasienergy E, the Floquet EP of our two-level system appears at the quasienergy $E = \pi$. This could lead to Floquet exceptional topology and anomalous π edge modes, which are unique to driven non-Hermitian systems. We will give an explicit example to demonstrate this point in Sec. III.

D. Adiabatic perturbation theory

We now consider a non-Hermitian system that is subject to slow-in-time and cyclic modulations. It can be viewed as the opposite limit of a high-frequency driven system. Introducing a re-scaled and dimensionless time variable s = t/T, we can express the Schrödinger equation (1) as

$$i\frac{\partial}{\partial s}|\Psi(s)\rangle = T\hat{H}(s)|\Psi(s)\rangle.$$
(58)

Our purpose is to find an expansion for $|\Psi(s)\rangle$ in the series of 1/T. It constitutes an adiabatic perturbation theory (APT) of the system [270], providing that different energy levels of $\hat{H}(s)$ are gapped for all $s \in [0, 1]$.

The Hamiltonian $\hat{H}(s)$ is periodic in the scaled time s with $\hat{H}(s) = \hat{H}(s+1)$. We denote its instantaneous right and left biorthonormal eigenvectors as $\{|n(s)\rangle\}$ and $\{|\tilde{n}(s)\rangle\}$ [271], such that

$$\hat{H}(s)|n(s)\rangle = E_n(s)|n(s)\rangle, \qquad \langle \tilde{n}(s)|\hat{H}(s) = \langle \tilde{n}(s)|E_n(s),$$
(59)

$$\langle \widetilde{m}(s)|n(s)\rangle = \delta_{mn}, \qquad \sum_{n} |n(s)\rangle \langle \widetilde{n}(s)| = 1.$$
 (60)

Here $E_n(s)$ is the instantaneous eigenenergy associated with $|n(s)\rangle$, which could be complex if $\hat{H}(s) \neq \hat{H}^{\dagger}(s)$. To proceed, we introduce an ansatz solution for the time-evolved state $|\Psi(s)\rangle$ as

$$|\Psi(s)\rangle = \sum_{n} e^{-i\Omega_n(s)} c_n(s) |n(s)\rangle, \tag{61}$$

where

$$\Omega_n(s) \equiv T \int_0^s E_n(s') ds'$$
(62)

describes a dynamical phase accumulated over the time interval [0, s]. At s = 0, we have $\Omega_n(0) = 0$ and

$$c_n(0) = \langle \widetilde{n}(0) | \Psi(0) \rangle \tag{63}$$

for the initial state $|\Psi(0)\rangle$. Plugging Eq. (61) into Eq. (58), we obtain

$$\sum_{n} e^{-i\Omega_n(s)} \dot{c}_n(s) |n(s)\rangle = -\sum_{n} e^{-i\Omega_n(s)} c_n(s) |\dot{n}(s)\rangle, \tag{64}$$

where $\dot{c}_n(s) = dc_n/ds$ and $|\dot{n}(s)\rangle = d|n(s)\rangle/ds$. Acting $\langle \widetilde{m}(s)|$ from the left on both sides of the above equation and using Eq. (60), we find

$$\dot{c}_m(s) = -\sum_{n \neq m} e^{i\Omega_{mn}(s)} c_n(s) \langle \widetilde{m}(s) | \dot{n}(s) \rangle, \tag{65}$$

where $\Omega_{mn}(s) \equiv \Omega_m(s) - \Omega_n(s)$. We have made the parallel transport gauge choice so that $\langle \tilde{n}(s) | \dot{n}(s) \rangle = 0$. Performing the integration over s and keeping the terms on the right-handside up to the first order of 1/T (i.e., up to the first order non-adiabatic correction), we arrive at

$$c_m(s) = c_m(0) + \frac{1}{T} \sum_{n \neq m} \left[\frac{i \langle \tilde{m}(s') | \dot{n}(s') \rangle}{\Delta_{mn}(s')} e^{i\Omega_{mn}(s')} \right] \Big|_{s'=0}^{s'=s} c_n(0),$$
(66)

where $\Delta_{mn}(s) = E_m(s) - E_n(s)$. To reach Eq. (66), we have assumed $\Omega_{mn}(s)$ to be real for all $m \neq n$ and $s \in [0, 1]$. This can be achieved if every instantaneous energy level possesses the same imaginary part $i\gamma(s)$. On the other hand, Eq. (66) is valid if the $\hat{H}(s)$ is PT-invariant at every s, such that $E_n(s) \in \mathbb{R}$ for all n. The approximation in Eq. (66) does not hold if $\Delta_{mn}(s) \notin \mathbb{R}$, which would cause the exponential amplification or decay of the amplitude $c_n(s)$. So the APT developed here is more restrictive in applications than its Hermitian counterparts. Inserting Eq. (66) into Eq. (61) results in the solution to Eq. (58) up to first order non-adiabatic corrections, i.e.,

$$|\Psi(s)\rangle = \sum_{m} e^{-i\Omega_{m}(s)} \left\{ c_{m}(0) + \frac{1}{T} \sum_{n \neq m} \left[\frac{i\langle \widetilde{m}(s') | \dot{n}(s') \rangle}{\Delta_{mn}(s')} e^{i\Omega_{mn}(s')} \right] \Big|_{s'=0}^{s'=s} c_{n}(0) \right\} |m(s)\rangle.$$
(67)

Since we have employed the formalism of biorthonormal eigenvectors, we may consider another ansatz solution for the left eigenvector

$$\langle \widetilde{\Psi}(s) | = \sum_{n} e^{i\Omega_n(s)} c_n^*(s) \langle \widetilde{n}(s) |, \qquad (68)$$

which obeys a conjugate Schrödinger equation

$$-i\frac{\partial}{\partial s}\langle \widetilde{\Psi}(s)| = \langle \widetilde{\Psi}(s)|\hat{H}.$$
(69)

Following the same reasoning in deriving Eq. (67), we find, up to the first order of 1/T, that

$$\langle \widetilde{\Psi}(s) | = \sum_{m} e^{+i\Omega_{m}(s)} \left\{ c_{m}^{*}(0) + \frac{1}{T} \sum_{n \neq m} \left[\frac{i \langle \dot{\widetilde{n}}(s') | m(s') \rangle}{\Delta_{nm}(s')} e^{i\Omega_{nm}(s')} \right]_{s'=0}^{s'=s} c_{n}^{*}(0) \right\} \langle \widetilde{m}(s) |.$$
(70)

Assuming that initially only the state with $m = \ell$ is occupied such that $c_m(0) = \delta_{m\ell}$, we can further simplify Eqs. (67) and (70) to

$$|\Psi_{\ell}(s)\rangle = e^{-i\Omega_{\ell}(s)} \left[|\ell(s)\rangle + \frac{1}{T} \sum_{m \neq \ell} \frac{i\langle \widetilde{m}(s) | \dot{\ell}(s) \rangle}{\Delta_{m\ell}(s)} |m(s)\rangle \right] - \frac{1}{T} \sum_{m \neq \ell} e^{-i\Omega_{m}(s)} \frac{i\langle \widetilde{m}(0) | \dot{\ell}(0) \rangle}{\Delta_{m\ell}(0)} |m(s)\rangle,$$
(71)

$$\langle \widetilde{\Psi}_{\ell}(s) | = e^{+i\Omega_{\ell}(s)} \left[\langle \widetilde{\ell}(s) | + \frac{1}{T} \sum_{m \neq \ell} \frac{i\langle \widetilde{\ell}(s) | m(s) \rangle}{\Delta_{\ell m}(s)} \langle \widetilde{m}(s) | \right] - \frac{1}{T} \sum_{m \neq \ell} e^{+i\Omega_{m}(s)} \frac{i\langle \widetilde{\ell}(0) | m(0) \rangle}{\Delta_{\ell m}(0)} \langle \widetilde{m}(s) |.$$

$$\tag{72}$$

For any observable \hat{O} , up to the correction of order 1/T, we can now express its biorthogonal average over the states $\{|\Psi_{\ell}(s)\rangle, |\widetilde{\Psi}_{\ell}(s)\rangle\}$ as

$$\langle \widetilde{\Psi}_{\ell}(s) | \hat{O} | \Psi_{\ell}(s) \rangle = \langle \widetilde{\ell}(s) | \hat{O} | \ell(s) \rangle + \frac{1}{T} \sum_{m \neq \ell} \frac{i \langle \widetilde{m}(s) | \dot{\ell}(s) \rangle}{\Delta_{m\ell}(s)} \langle \widetilde{\ell}(s) | \hat{O} | m(s) \rangle + \frac{1}{T} \sum_{m \neq \ell} \frac{i \langle \widetilde{\ell}(s) | m(s) \rangle}{\Delta_{\ell m}(s)} \langle \widetilde{m}(s) | \hat{O} | \ell(s) \rangle.$$
(73)

We notice that this average does not contain any time-oscillating phase factors.

We now illustrate this APT with an application in the study of dynamical topological phenomena. Let us consider noninteracting particles in a one-dimensional (1D) periodic lattice, whose onsite potential is also varied slowly and periodically in time. This is the typical situation encountered in the topological Thouless pump [16–18]. The group velocity of the particle can be expressed as $\hat{v} = \partial_k \hat{H}$, where k is the quasimomentum. At the initial time s = 0, we assume that the band ℓ is uniformly filled, and it is separated from the other bands at all k and s. The pumped number of particles over one adiabatic cycle due to this initially filled band is then given by

$$N_{\ell} = \int_{-\pi}^{\pi} \frac{dk}{2\pi} \int_{0}^{1} ds \langle \widetilde{\Psi}_{\ell}(s) | \partial_{k} \hat{H} | \Psi_{\ell}(s) \rangle.$$
(74)

With the aid of Eq. (59), it is not hard to identify that

$$\langle \tilde{\ell}(s) | \partial_k \hat{H} | \ell(s) \rangle = \partial_k E_\ell(k, s),$$
(75)

$$\frac{\langle \tilde{\ell}(s) | \partial_k \hat{H} | m(s) \rangle}{\Delta_{m\ell}(s)} = \langle \tilde{\ell}(s) | \partial_k | m(s) \rangle, \qquad m \neq \ell,$$
(76)

where $E_{\ell}(k, s)$ denotes the energy dispersion of the ℓ th adiabatic Bloch band. Plugging Eqs. (75), (76), and (73) into Eq. (74), we obtain

$$N_{\ell} = \int_{-\pi}^{\pi} \frac{dk}{2\pi} \int_{0}^{1} ds \partial_{k} E_{\ell}(k,s)$$

$$+ \sum_{m \neq \ell} \int_{-\pi}^{\pi} \frac{dk}{2\pi i} \int_{0}^{T} dt \langle \partial_{k} \widetilde{\ell}(k,t) | m(k,t) \rangle \langle \widetilde{m}(k,t) | \partial_{t} \ell(k,t) \rangle$$

$$- \sum_{m \neq \ell} \int_{-\pi}^{\pi} \frac{dk}{2\pi i} \int_{0}^{T} dt \langle \partial_{t} \widetilde{\ell}(k,t) | m(k,t) \rangle \langle \widetilde{m}(k,t) | \partial_{k} \ell(k,t) \rangle.$$

$$(77)$$

Noting that $E_{\ell}(k = -\pi, s) = E_{\ell}(k = \pi, s)$ and $\sum_{m} |m(k, t)\rangle \langle \widetilde{m}(k, t)| = 1$, we can simplify the above equation and arrive at the pumped number of particles over an adiabatic cycle

$$N_{\ell} = \int_{-\pi}^{\pi} \frac{dk}{2\pi i} \int_{0}^{T} dt \left[\langle \partial_{k} \widetilde{\ell}(k,t) | \partial_{t} \ell(k,t) \rangle - \langle \partial_{t} \widetilde{\ell}(k,t) | \partial_{k} \ell(k,t) \rangle \right].$$
(78)

Equation (78) describes nothing but the Chern number of the adiabatic Bloch band ℓ , whose energy dispersion is defined on a two dimensional (2D) torus $(k, t) \in [-\pi, \pi) \times [0, T)$. The conditions for Eq. (78) to hold are as follows. First, the Hamiltonian of the system should be quasi-Hermitian with a real spectrum (e.g., PT-invariant) in our considered parameter regime. Second, the band E_{ℓ} should be well-gapped from the other bands throughout the 2D torus $(k, t) \in [-\pi, \pi) \times [0, T)$, and \hbar/T should be much smaller than $|\Delta_{\ell m}|$ for all $m \neq \ell$ in order to guarantee the adiabatic condition. Third, the evolution of left and right vectors of the system should follow Eqs. (58) and (69). Note here that the expression of Chern number is not sensitive to the choice of biorthonormal basis. We will get the same N_{ℓ} after changing $\tilde{\ell} \to \ell$ or $\ell \to \tilde{\ell}$ in Eq. (78), as proved before for non-Hermitian Chern bands [139].

E. Symmetry and topological characterization

Over the past few years, rich symmetry classifications and topological invariants have been identified for non-Hermitian topological matter [139–147]. In this subsection, we mainly recap two symmetries together with their associated topological numbers. They are the most relevant ones for the characterization of non-Hermitian Floquet topological phases reviewed in this work.

We first discuss the PT-symmetry, which is associated with the operator \mathcal{PT} . Here \mathcal{P} denotes the parity operator and \mathcal{T} denotes the time-reversal operator. When the Hamiltonian of a non-Hermitian system \hat{H} respects the PT-symmetry, we have $[\mathcal{PT}, \hat{H}] = 0$. In this case, the system could have a real spectrum in the PT unbroken regime, where the eigenstates $|\psi\rangle$ and $\mathcal{PT}|\psi\rangle$ of \hat{H} are coincident up to a global phase. To see this, let us consider a non-degenerate eigenstate $|\psi\rangle$ of \hat{H} that satisfies the eigenvalue equation

$$\hat{H}|\psi\rangle = E|\psi\rangle. \tag{79}$$

The PT-symmetry of \hat{H} then implies that

$$\hat{H}(\mathcal{PT}|\psi\rangle) = E^*(\mathcal{PT}|\psi\rangle).$$
(80)

Therefore, $\mathcal{PT}|\psi\rangle$ is also an eigenstate of \hat{H} with the energy E^* . If $|\psi\rangle$ and \hat{H} share the same PT-symmetry, $|\psi\rangle$ should be the common eigenstate of \hat{H} and \mathcal{PT} . $\mathcal{PT}|\psi\rangle$ can thus only differ from $|\psi\rangle$ up to a global phase, which means that $E = E^* \in \mathbb{R}$. However, with the change of system parameters (e.g., the strengths of gain and loss), the PT-symmetry of $|\psi\rangle$ could be spontaneously broken and the spectrum of \hat{H} could switch from real to complex after undergoing a PT-symmetry breaking transition. A topological invariant, defined as [188]

$$w = \int_0^{2\pi} \frac{d\theta}{2\pi i} \partial_\theta \ln \det \left[\hat{H}(\theta) - E_0 \right], \tag{81}$$

might be employed to characterize such a real-to-complex spectral transition. Here E_0 is a base energy chosen appropriately on the complex plane. The parametrized Hamiltonian $\hat{H}(\theta) = \hat{H}(\theta + 2\pi)$, where θ can be viewed as the quasimomentum along an artificial dimension. We have also taken the periodic boundary condition (PBC) for \hat{H} before implementing its θ -parametrization. The w in Eq. (81) thus depicts a spectral winding number with respect to the base energy E_0 , i.e., it counts the number of times that the spectrum of $\hat{H}(\theta)$ winds around E_0 on the complex plane when the synthetic quasimomentum θ is varied over a cycle. When the spectrum of $\hat{H}(\theta)$ is real, we must have w = 0, as a spectral loop cannot be formed on the complex plane in this case. When the spectrum of $\hat{H}(\theta)$ is complex, w may take an integer-quantized value if $\hat{H}(\theta)$ possesses spectral loops around E_0 . A suitably chosen E_0 could then yield a nonzero w when the first spectral loop appears on the complex-E plane, thereby detecting the topological changes in the spectrum of \hat{H} across the PT-breaking transition.

For a Floquet system, if the time-periodic Hamiltonian $\hat{H}(t)$ possesses the PT-symmetry at each instant t, the resulting Floquet operator also has the PT-symmetry. In this case, we can define the spectral winding number w as in Eq. (81) for the stroboscopic Floquet effective Hamiltonian so as to capture the PT-breaking transition in the quasienergy spectrum. We will provide explicit examples for this usage in Subsec. III D, where we consider PT transitions, localization transitions, and topological transitions in non-Hermitian Floquet quasicrystals.

We next consider the chiral (or sublattice) symmetry, whose associated operator will be denoted by S. When the Hamiltonian \hat{H} of a system respects the chiral symmetry S, we have $S\hat{H}S = -\hat{H}$, where S is both Hermitian and unitary [272]. The implication of this symmetry on the spectrum of \hat{H} is as follows. Suppose that $|\psi\rangle$ is an eigenstate of a chiral-symmetric \hat{H} with the energy E, i.e., $\hat{H}|\psi\rangle = E|\psi\rangle$. We have

$$\hat{H}(\mathcal{S}|\psi\rangle) = -E(\mathcal{S}|\psi\rangle). \tag{82}$$

Therefore, $S|\psi\rangle$ is also an eigenstate of \hat{H} with the energy -E. The eigenstates of a chiralsymmetric \hat{H} should then come in pairs of $\{|\psi\rangle, S|\psi\rangle\}$ with the energies $\{E, -E\}$ that are symmetric with respect to E = 0. This further leads to a chiral-symmetry protected degeneracy for any eigenstate with E = 0. When the spectrum of \hat{H} is gapped at E = 0, we can group its energy levels into two clusters with $\operatorname{Re} E < 0$ and $\operatorname{Re} E > 0$. If these two clusters meet with each other at E = 0 and then separate with the change of certain system parameters, the system may undergo a phase transition. In one dimension, the change of band topology of the system before and after such a transition could be characterized by a winding number w_0 , defined as [272–274]

$$w_0 = \int_{-\pi}^{\pi} \frac{dk}{4\pi} \operatorname{Tr}[\mathcal{SQ}(k)i\partial_k \mathcal{Q}(k)].$$
(83)

Here the PBC has been assumed and $k \in [-\pi, \pi)$ denotes the quasimomentum. The signresolved projector $\mathcal{Q}(k)$ is obtained from the spectral decomposition of $\hat{H} = \sum_k |k\rangle H(k)\langle k|$ with $H(k) = \sum_n E_n(k)|n(k)\rangle\langle \tilde{n}(k)|$ at each k by attributing +1 (-1) to every energy band n with $\operatorname{Re} E_n > 0$ ($\operatorname{Re} E_n < 0$). $\mathcal{Q}(k)$ can thus be expressed as [261]

$$\mathcal{Q}(k) = \sum_{n} \operatorname{sgn}\{\operatorname{Re}[E_n(k)]\}|n(k)\rangle\langle \widetilde{n}(k)|.$$
(84)

Here $H(k)|n(k)\rangle = E_n(k)|n(k)\rangle$ and $\langle \tilde{n}(k)|H(k) = \langle \tilde{n}(k)|E_n(k)$. The set $\{|n(k)\rangle, |\tilde{n}(k)\rangle\}$ of basis satisfies the biorthonormal relations in Eq. (60). Under the PBC, the winding number w_0 could characterize the bulk topological properties of a 1D chiral-symmetric Hamiltonian \hat{H} , either Hermitian or non-Hermitian. It could further distinguish between different bulk topological insulating phases by showing a quantized jump at the transition point, where the two band clusters of \hat{H} meet with each other at E = 0. However, under the open boundary condition (OBC), due to the possible existence of NHSE, the w_0 defined in Eq. (83) may not be able to correctly predict the gap-closing points of the spectrum in the parameter space and determine the number of degenerate edge modes at E = 0 in different parameter regions. This non-Hermiticity induced breakdown of bulk-edge correspondence may be recovered by introducing a real-space counterpart of w_0 , defined as [168]

$$W_0 = -\frac{1}{L_{\rm B}} \operatorname{Tr}_{\rm B}(\mathcal{SQ}[\mathcal{Q}, \hat{N}]).$$
(85)

Here S is the chiral symmetry operator of \hat{H} , \hat{N} is the position operator in real space, and Q is the flat band projector defined as

$$Q = \sum_{|\psi_j\rangle \in \text{bulk}} \text{sgn}[\text{Re}(E_j)] |\psi_j\rangle \langle \widetilde{\psi}_j |, \qquad (86)$$

where the summation is now taken over all the bulk eigenstates $\{|\psi_j\rangle\}$ of \hat{H} under the OBC. The whole lattice of length $L = L_{\rm B} + 2L_{\rm E}$ is decomposed into three segments, with a bulk region of length $L_{\rm B}$ in the middle and two edge regions of the same length $L_{\rm E}$ at the left and right boundaries of the open chain. The trace $\text{Tr}_{\rm B}(\cdot)$ is only taken over the bulk region, which excludes all possible interruptions caused by the NHSE in the edge regions. The resulting W_0 was found to be able to faithfully capture the topological phase transitions and bulkedge correspondence in 1D, chiral symmetric non-Hermitian systems even in the presence of NHSE [168]. It was also suggested to be equivalent to the topological winding number defined through the generalized Brillouin zone of non-Hermitian systems. By definition, the winding number W_0 in Eq. (85) is also robust to perturbations induced by symmetrypreserved disorders and impurities, making it applicable to more general situations. In the clean, Hermitian, and thermodynamic limit, the W_0 in Eq. (85) can be further reduced to the w_0 in Eq. (83) [272–274].

For a non-Hermitian Floquet system, we can state its chiral symmetry as follows [254]. From Eq. (28), we can express the Floquet operator as $\hat{U} = e^{-i\hat{H}_{\text{eff}}}$, where we have set the driving period T = 1 for brevity. Viewing the \hat{H}_{eff} as a static Hamiltonian, we say that it respects the chiral symmetry if there exists a unitary and Hermitian operator \mathcal{S} such that $\hat{\mathcal{S}H}_{\text{eff}}\mathcal{S} = -\hat{H}_{\text{eff}}$. At the level of \hat{U} , the chiral symmetry then implies that

$$\mathcal{S}\hat{U}\mathcal{S} = \hat{U}^{-1}.\tag{87}$$

As in the case of static systems, the chiral symmetry of \hat{U} has a direct implication for the symmetry of its spectrum. If $|\Psi_E\rangle$ is an eigenstate of \hat{U} with the quasienergy E, i.e., $\hat{U}|\Psi_E\rangle = e^{-iE}|\Psi_E\rangle$, we immediately have

$$\hat{U}(\mathcal{S}|\Psi_E\rangle) = e^{-i(-E)}(\mathcal{S}|\Psi_E\rangle),\tag{88}$$

which means that $S|\Psi_E\rangle$ is also an eigenstate of \hat{U} with the quasienergy -E. The Floquet spectrum of \hat{U} is then symmetric with respect to both the quasienergies E = 0 and $E = \pi$. The latter is because -E and E are identified at the quasienergy π . When the spectrum is gapped at E = 0 and π , the quasienergy levels of \hat{U} could be grouped into two clusters. One of them has the quasienergy $\operatorname{Re}E \in (-\pi, 0)$ and the other one has $\operatorname{Re}E \in (0, \pi)$. They could meet with each other at either the quasienergy zero or π , leading to two possible phase transitions. This implies that a complete topological characterization of a chiral symmetric Floquet system should require at least two winding numbers, which is rather different from the case of static systems where a single winding number is sufficient. To identify these winding numbers, let us consider the example of a periodically kicked 1D system, whose Hamiltonian and Floquet operator take the forms of Eqs. (15) and (16). We also assume that the two parts of Hamiltonians \hat{H}_0 and \hat{H}_1 in Eq. (15) have the same chiral symmetry S. At the level of $\hat{U} = e^{-i\hat{H}_0}e^{-i\hat{H}_1}$ (assuming T = 1), it is not straightforward to identify the form of a chiral symmetry. However, we can apply similarity transformations to \hat{U} and express it in two symmetric time frames [69] as

$$\hat{U}_1 = e^{-\frac{i}{2}\hat{H}_1} e^{-i\hat{H}_0} e^{-\frac{i}{2}\hat{H}_1},\tag{89}$$

$$\hat{U}_2 = e^{-\frac{i}{2}\hat{H}_0}e^{-i\hat{H}_1}e^{-\frac{i}{2}\hat{H}_0}.$$
(90)

It is then clear that $S\hat{U}_{\alpha}S = \hat{U}_{\alpha}^{-1}$ for $\alpha = 1, 2$, that is, the Floquet operators \hat{U}_1 and \hat{U}_2 in the two symmetric time frames respect the same chiral symmetry S. We can thus introduce a winding number for each of them under the PBC as [254]

$$w_{\alpha} = \int_{-\pi}^{\pi} \frac{dk}{4\pi} \operatorname{Tr}[\mathcal{SQ}_{\alpha}(k)i\partial_{k}\mathcal{Q}_{\alpha}(k)], \qquad (91)$$

where $k \in [-\pi, \pi)$, and

$$\mathcal{Q}_{\alpha}(k) = \sum_{n} \operatorname{sgn}\{\operatorname{Re}[E_{n}(k)]\}|n_{\alpha}(k)\rangle\langle \widetilde{n}_{\alpha}(k)|.$$
(92)

The $E_n(k)$ in Eq. (92) now denotes the quasienergy of the Floquet eigenstate $|n_{\alpha}(k)\rangle$ of \hat{U}_{α} at the quasimomentum k under the PBC. Using the w_1 and w_2 , we can construct another pair of winding numbers w_0 and w_{π} , given by [254]

$$w_0 = \frac{w_1 + w_2}{2}, \qquad w_\pi = \frac{w_1 - w_2}{2}.$$
 (93)

In Subsecs. III B and III C, we will demonstrate with explicit examples that the w_0 (w_{π}) could correctly capture the bulk topological transitions of non-Hermitian Floquet bands

through the gap closing/reopening at the quasienergy E = 0 ($E = \pi$) in various chiral symmetric, non-Hermitian Floquet insulating and superconducting models. Furthermore, in the absence of NHSE, the w_0 and w_{π} could also capture the numbers of Floquet edge modes at zero and π quasienergies under the OBC, and thus are capable of describing the bulk-edge correspondence of the related models. In the presence of NHSE, we can retrieve the characterization of topological transitions and bulk-edge correspondence in chiral symmetric, non-Hermitian Floquet systems under the OBC through the open-bulk winding numbers, in analogy with Eq. (85). For the \hat{U}_1 and \hat{U}_2 in Eqs. (89) and (90), we can define a winding number for each of them under the OBC as [261]

$$W_{\alpha} = -\frac{1}{L_{\rm B}} \operatorname{Tr}_{\rm B}(\mathcal{SQ}_{\alpha}[\mathcal{Q}_{\alpha}, \hat{N}]), \qquad \alpha = 1, 2.$$
(94)

Here, the meanings of $L_{\rm B}$, ${\rm Tr}_{\rm B}$, \mathcal{S} and \hat{N} are the same as those in Eq. (85). The Floquet band projector in the time frame α is given by

$$\mathcal{Q}_{\alpha} = \sum_{|\psi_j^{\alpha}\rangle \in \text{bulk}} \text{sgn}[\text{Re}(E_j)] |\psi_j^{\alpha}\rangle \langle \widetilde{\psi}_j^{\alpha} |, \qquad (95)$$

where $|\psi_j^{\alpha}\rangle$ is the *j*th bulk eigenstate of \hat{U}_{α} ($\alpha = 1, 2$) with the quasienergy E_j under the OBC. The linear combinations of W_1 and W_2 lead to another pair of winding numbers [261]

$$W_0 = \frac{W_1 + W_2}{2}, \qquad W_\pi = \frac{W_1 - W_2}{2}.$$
 (96)

In Subsec. III B, we will illustrate that with the help of the winding numbers (w_0, w_π) and (W_0, W_π) , a dual topological characterization of the phase transitions, edge states, and bulk-edge correspondence can be established for 1D, chiral symmetric non-Hermitian Floquet systems under different boundary conditions, regardless of whether the NHSE is present or not [261]. Interestingly, the winding numbers (w_0, w_π) may both become halfinteger quantized due to the presence of Floquet EP in the bulk, thus revealing the presence of Floquet exceptional topology. Meanwhile, the open-bulk winding numbers (W_0, W_π) are always integer quantized.

F. Dynamical indicators

In this subsetion, we review two complementary dynamical probes in position and momentum spaces. Both of them can be used to characterize the topological properties of 1D non-Hermitian Floquet systems with chiral symmetry. These indicators allow us to extract the topological winding numbers of the system from its long-time stroboscopic dynamics [254–256]. The measurement of these indicators could thus provide evidences for the existence of non-Hermitian Floquet topological matter.

1. Dynamic winding number (DWN)

The DWN [275, 276], obtained from the long-time stroboscopic average of spin textures, could provide us with information about the bulk topological properties of non-Hermitian Floquet systems [256]. Let us consider a 1D, chiral symmetric non-Hermitian Floquet system with two quasienergy bands. Under the PBC, we can express its Floquet operator in the symmetric time frame α (= 1, 2) as $\hat{U}_{\alpha} = \sum_{k \in \text{BZ}} |k\rangle e^{-iH_{\alpha}(k)} \langle k|$. Here $H_{\alpha}(k)$ is the effective Hamiltonian in time frame α and $k \in [-\pi, \pi)$ is the quasimomentum. \hat{U}_{α} and $H_{\alpha}(k)$ share the same chiral symmetry S, i.e., $SH_{\alpha}(k)S = -H_{\alpha}(k)$. For a non-Hermitian system, the \hat{U}_{α} is generally not unitary and $H_{\alpha}(k)$ is also not Hermitian. The right and left biorthonormal eigenvectors $\{|n_{\alpha}(k)\rangle\}$ and $\{|\tilde{n}_{\alpha}(k)\rangle\}$ of $H_{\alpha}(k)$ satisfy the eigenvalue equations

$$H_{\alpha}(k)|n_{\alpha}(k)\rangle = E_{n}(k)|n_{\alpha}(k)\rangle, \qquad (97)$$

$$\langle \widetilde{n}_{\alpha}(k) | H_{\alpha}(k) = \langle \widetilde{n}_{\alpha}(k) | E_n(k).$$
(98)

Here $n = \pm$ are the indices of the two Floquet bands with the quasienergies $E_{\pm}(k) \equiv \pm E(k)$. The biorthonormal relationship requires

$$\langle \widetilde{n}_{\alpha}(k) | n'_{\alpha}(k) \rangle = \delta_{nn'}, \quad \sum_{n=\pm} |n_{\alpha}(k)\rangle \langle \widetilde{n}_{\alpha}(k)| = 1, \quad \alpha = 1, 2.$$
(99)

We consider the case in which the system is prepared in a general initial state $|\psi_{\alpha}(k,0)\rangle = \sum_{n=\pm} c_n(k) |n_{\alpha}(k)\rangle$. The corresponding initial state in the left Hilbert space reads $|\widetilde{\psi}_{\alpha}(k,0)\rangle = \sum_{n\pm} c_n(k) |\widetilde{n}_{\alpha}(k)\rangle$, such that initially $\sum_{n=\pm} |c_n(k)|^2 = 1$ at each k. After the stroboscopic

evolution over a number of ℓ driving periods, the right initial state becomes

$$|\psi_{\alpha}(k,\ell)\rangle = \sum_{n=\pm} c_n(k) e^{-i\ell E_n(k)} |n_{\alpha}(k)\rangle.$$
(100)

For the left initial state, we assume it to be evolved by a different effective Hamiltonian $\widetilde{H}_{\alpha}(k) = \sum_{\pm} E_n(k) |\widetilde{n}_{\alpha}(k)\rangle \langle n_{\alpha}(k)|$, so that after the evolution over ℓ driving periods it reaches the state

$$|\widetilde{\psi}_{\alpha}(k,\ell)\rangle = \sum_{n=\pm} c_n(k) e^{-i\ell E_n(k)} |\widetilde{n}_{\alpha}(k)\rangle.$$
(101)

Note that the dynamical equation of $|\tilde{\psi}_{\alpha}(k,0)\rangle$ we used here is different from that employed in our study of the APT in Subsec. II D.

The stroboscopic average of an observable \hat{O} over $|\tilde{\psi}_{\alpha}(k,t)\rangle$ at the time $t = \ell T$ is then given by

$$\langle \hat{O}(k,\ell) \rangle_{\alpha} = \frac{\langle \tilde{\psi}_{\alpha}(k,\ell) | \hat{O} | \psi_{\alpha}(k,\ell) \rangle}{\langle \tilde{\psi}_{\alpha}(k,\ell) | \psi_{\alpha}(k,\ell) \rangle}.$$
(102)

Without the loss of generality, we can consider the chiral symmetric $H_{\alpha}(k)$ to be in the form of

$$H_{\alpha}(k) = h_{\alpha x}(k)\sigma_x + h_{\alpha y}(k)\sigma_y.$$
(103)

Note that any two out of the three Pauli matrices $(\sigma_x, \sigma_y, \sigma_z)$ can be chosen to enter the $H_{\alpha}(k)$, and the rest Pauli matrix [e.g., σ_z for the $H_{\alpha}(k)$ in Eq. (103)] plays the role of the chiral symmetry operator S. By diagonalizing $H_{\alpha}(k)$, we obtain the biorthonormal eigenvectors as

$$|n_{\alpha}(k)\rangle = \frac{1}{\sqrt{2}E_n(k)} \begin{bmatrix} h_{\alpha x}(k) - ih_{\alpha y}(k) \\ E_n(k) \end{bmatrix}, \qquad |\widetilde{n}_{\alpha}(k)\rangle = \frac{1}{\sqrt{2}E_n^*(k)} \begin{bmatrix} h_{\alpha x}^*(k) - ih_{\alpha y}^*(k) \\ E_n^*(k) \end{bmatrix},$$
(104)

where $E_n(k) = n\sqrt{h_{\alpha x}^2(k) + h_{\alpha y}^2(k)}$ for $n = \pm$. We can now compute the stroboscopicaveraged spin textures in the long-time limit. For the $H_{\alpha}(k)$ in Eq. (103), this means that we need to find the multi-cycle averages of σ_x and σ_y . According to Eq. (102), they are given by

$$r_j^{\alpha}(k) \equiv \lim_{N \to \infty} \frac{1}{N} \sum_{\ell=1}^{N} \langle \sigma_j(k,\ell) \rangle_{\alpha}, \qquad (105)$$

where j = x, y and $\alpha = 1, 2$. N counts the total number of driving periods. From the averaged spin textures $[r_x^{\alpha}(k), r_y^{\alpha}(k)]$, we can define the dynamic winding angle as

$$\theta_{yx}^{\alpha}(k) \equiv \arctan\left[\frac{r_{y}^{\alpha}(k)}{r_{x}^{\alpha}(k)}\right].$$
(106)

The net winding number of $\theta_{yx}^{\alpha}(k)$ over a cycle in k-space defines the dynamic winding number (DWN) in the time frame α , i.e.,

$$\nu_{\alpha} = \int_{-\pi}^{\pi} \frac{dk}{2\pi} \partial_k \theta_{yx}^{\alpha}(k), \qquad \alpha = 1, 2.$$
(107)

In Ref. [256], it was proved with straightforward calculations that in the limit $N \to \infty$, the ν_{α} converges to the w_{α} in Eq. (91) if the initial condition satisfies $c_{\pm}(k) \neq 0$ at each k. Therefore, by preparing the initial state at different k under this condition and measuring the averaged spin textures over a long stroboscopic time, we can obtain the winding numbers (w_0, w_{π}) of a chiral symmetric 1D Floquet system (either Hermitian or non-Hermitian) through the following combinations of (ν_1, ν_2) in two symmetric time frames [256], i.e.,

$$w_0 = \frac{\nu_1 + \nu_2}{2}, \qquad w_\pi = \frac{\nu_1 - \nu_2}{2}.$$
 (108)

In Sec. III, we will illustrate the application of this dynamical-topological correspondence to non-Hermitian Floquet topological insulators in one dimension. We will see that both integer and half-integer quantized topological winding numbers can be extracted from the DWN.

2. Mean chiral displacement (MCD)

The MCD allows us to detect the winding numbers of a Floquet system from the longtime averaged chiral displacement of an initially localized wavepacket. It was first proposed as a means to probe the topological invariants of chiral symmetric topological insulators in one dimension [277–279]. Later, the MCD was used to obtain the winding numbers of Floquet systems [78, 81] and also generalized to 2D higher-order topological insulators [80]. Its applicability was demonstrated both theoretically and experimentally [277, 279].

For a chiral symmetric non-Hermitian system, the chiral displacement in the symmetric

time frame α (= 1, 2) can be defined as

$$C_{\alpha}(t) = \operatorname{Tr}[\rho_0(\hat{\widetilde{U}}^{\dagger}_{\alpha})^t (\hat{N} \otimes \mathcal{S}) \hat{U}^t_{\alpha}].$$
(109)

Here, $t = \ell T$ denotes the stroboscopic time, with T being the driving period. \hat{N} is the unit cell position operator and S is the chiral symmetry operator. The initial state ρ_0 can be chosen as a state localized in the middle of the lattice. For example, ρ_0 may take the form of $(|0\rangle\langle 0| \otimes \sigma_0)/2$ for a 1D bipartite lattice, where $|0\rangle$ is the eigenbasis of the central unit cell and σ_0 is the identity operator acting on the internal space of the two sublattices. Both the Floquet operator \hat{U}_{α} and its dual $\hat{\tilde{U}}_{\alpha}$ respect the chiral symmetry S. In the lattice representation, they can be expressed as

$$\hat{U}_{\alpha} = \sum_{j=1}^{L} e^{-iE_j} |\psi_j^{\alpha}\rangle \langle \widetilde{\psi}_j^{\alpha} |, \qquad \hat{\widetilde{U}}_{\alpha} = \sum_{j=1}^{L} e^{-iE_j} |\widetilde{\psi}_j^{\alpha}\rangle \langle \psi_j^{\alpha} |.$$
(110)

Here L denotes the total number of degrees of freedom of the lattice. $|\psi_j^{\alpha}\rangle$ and $\langle \tilde{\psi}_j^{\alpha}|$ denote the right and left eigenvectors of \hat{U}_{α} with the quasienergy E_j . They form a biorthonormal basis such that

$$\langle \widetilde{\psi}_{j}^{\alpha} | \psi_{m}^{\alpha} \rangle = \delta_{jm}, \qquad \sum_{j=1}^{L} | \psi_{j}^{\alpha} \rangle \langle \widetilde{\psi}_{j}^{\alpha} | = 1.$$
 (111)

Note that $\hat{\tilde{U}}_{\alpha}$ is not the Hermitian conjugate of \hat{U}_{α} in general.

We now consider the stroboscopic long-time average of $C_{\alpha}(t)$ in the time frames $\alpha = 1, 2$ for a 1D non-Hermitian Floquet system with chiral symmetry. Under the PBC, taking the initial state to be $\rho_0 = (|0\rangle\langle 0| \otimes \sigma_0)/2$ and performing the Fourier transformation from position to momentum representations, we find the $C_{\alpha}(t)$ in Eq. (109) to be [255]

$$C_{\alpha}(t) = \frac{1}{2} \int_{-\pi}^{\pi} \frac{dk}{2\pi} \operatorname{Tr}[\widetilde{U}_{\alpha}^{\dagger t}(k) \mathcal{S}i\partial_{k}U_{\alpha}^{t}(k)].$$
(112)

Here $k \in [-\pi, \pi)$ is the quasimomentum. $U_{\alpha}(k) = \langle k | \hat{U}_{\alpha} | k \rangle$ and $\tilde{U}_{\alpha}(k) = \langle k | \hat{\tilde{U}}_{\alpha} | k \rangle$ act on the internal degrees of freedom (spins and/or sublattices) of the system at a fixed k. Taking the long-time stroboscopic average and incorporating the normalization factor $\text{Tr}[\rho_0(\hat{\tilde{U}}_{\alpha}^{\dagger})^t \hat{U}_{\alpha}^t]$,

we find the expression of MCD as [255]

$$\overline{C}_{\alpha} = \lim_{\ell \to \infty} \frac{1}{\ell T} \sum_{t=T}^{\ell T} \int_{-\pi}^{\pi} \frac{dk}{2\pi} \frac{\operatorname{Tr}\left[\widetilde{U}_{\alpha}^{\dagger t}(k)\mathcal{S}i\partial_{k}U_{\alpha}^{t}(k)\right]}{\operatorname{Tr}\left[\widetilde{U}_{\alpha}^{\dagger t}(k)U_{\alpha}^{t}(k)\right]}.$$
(113)

For a given $H_{\alpha}(k)$ of the form in Eq. (103), we have $\mathcal{S} = \sigma_z$, $U_{\alpha}(k) = e^{-iH_{\alpha}(k)}$, and $\widetilde{U}_{\alpha}(k) = e^{-iE(k)H_{\alpha}^{\dagger}(k)/E^{*}(k)}$, where $E(k) = \frac{1}{2}\text{Tr}[H_{\alpha}^{2}(k)]$. One can then work out Eq. (113) explicitly and obtain [255]

$$\overline{C}_{\alpha} = \frac{w_{\alpha}}{2}, \qquad \alpha = 1, 2.$$
(114)

Here the w_{α} is nothing but the winding number of \hat{U}_{α} as defined in Eq. (91). Therefore, by measuring the MCDs $(\overline{C}_1, \overline{C}_2)$ in two symmetric time frames, we can determine the topological winding numbers (w_0, w_{π}) of a 1D chiral symmetric Floquet system through the relations [255]

$$w_0 = C_0 \equiv \overline{C}_1 + \overline{C}_2, \qquad w_\pi = C_\pi \equiv \overline{C}_1 - \overline{C}_2.$$
(115)

The MCD provides a real-space complementary to the DWN. In Sec. III, we will demonstrate the usage of MCD to dynamically probing the winding numbers of first- and second-order non-Hermitian Floquet topological insulators in one [255] and two [259] spatial dimensions.

G. Localization transition and mobility edge

In the last part of this subsection, we recap some tools that can be used to characterize the real-to-complex spectral transitions, localization transitions, and mobility edges in disordered non-Hermitian Floquet systems [262, 264]. Considering the relevance to this review, we focus on a 1D quadratic lattice model of the form

$$\hat{H}(t) = \sum_{\langle n,n' \rangle} [J_n(t)e^{\gamma_n} \hat{c}_n^{\dagger} \hat{c}_{n'} + J_n^*(t)e^{-\gamma_n} \hat{c}_{n'}^{\dagger} \hat{c}_n] + \sum_n V_n(t)\hat{c}_n^{\dagger} \hat{c}_n.$$
(116)

Here $\langle n, n' \rangle$ includes the lattice site indices n and n' with n' > n. $\hat{c}_n^{\dagger}(\hat{c}_n)$ creates (annihilates) a particle on the site n and the parameter $\gamma \in \mathbb{R}$. The Hamiltonian $\hat{H}(t)$ is non-Hermitian if $\gamma_n \neq 0$ for some n (nonreciprocal hopping) or $V_n(t) \neq V_n^*(t)$ (onsite gain and loss). $\hat{H}(t)$ is further time-periodic if $J_n(t) = J_n(t+T)$ and $V_n(t) = V_n(t+T)$ for all n, with T being the driving period. The disorder terms may be included within $V_n(t)$ (diagonal disorder) or $J_n(t)$ (off-diagonal disorder). As an example, for a 1D non-Hermitian quasicrystal (NHQC) with correlated onsite disorder, the $V_n(t)$ may take the form of $V_n(t) = V(t) \cos(2\pi\alpha n + i\beta)$. Here α is irrational and $\beta \in \mathbb{R}$, with $i\beta$ describing an imaginary phase shift in the superlattice potential V_n .

The Floquet operator of the system described by the $\hat{H}(t)$ in Eq. (116) takes the general form of $\hat{U} = \hat{\mathsf{T}} e^{-i \int_0^T \hat{H}(t) dt} = e^{-i \hat{H}_{\text{eff}} T}$. If $\hat{H}(t)$ respects the PT-symmetry, \hat{H}_{eff} could have the same symmetry and the quasienergy spectrum of \hat{U} could be real. With the increase of the non-Hermitian parameter of $\hat{H}(t)$, the Floquet spectrum of \hat{U} may undergo a PT-breaking transition, after which certain quasienergies of \hat{U} (obtained by solving $\hat{U}|\Psi\rangle = e^{-iE}|\Psi\rangle$) may acquire nonvanishing imaginary parts. To take into account such a spectral transition, we introduce the following quantities

$$\max |\mathrm{Im}E| \equiv \max_{j \in \{1,\dots,L\}} (|\mathrm{Im}E_j|), \tag{117}$$

$$\rho \equiv N(\text{Im}E \neq 0)/L. \tag{118}$$

Here E_j is the *j*th quasienergy eigenvalue of \hat{U} . *L* counts the Hilbert space dimension of \hat{U} , i.e., the total number of degrees of freedom of the lattice. $N(\text{Im}E \neq 0)$ denotes the number of quasienergy eigenvalues whose imaginary parts are nonzero. ρ thus describes the density of states with complex quasienergies in the system. In the PT-invariant phase, we would have max $|\text{Im}E| = \rho = 0$, implying that all the quasienergies of \hat{U} are real. In the PT-broken phase, we would have max |ImE| > 0 and $\rho \in (0, 1]$, meaning that there is a finite number of eigenstates of \hat{U} whose quasienergies are complex. Specially, we would have $\rho \simeq 1$ if almost all the Floquet eigenstates of \hat{U} possess complex quasienergies. Therefore, by locating the positions where both the max |ImE| and ρ start to deviate from zero in the parameter space, we can identify the PT transition points for the Floquet spectrum of \hat{U} . We will see that due to the interplay between periodic drivings and non-Hermitian effects, both PT-symmetry breaking and restoring transitions could be induced by tuning a single parameter in the Floquet NHQC.

The localization nature of the Floquet eigenstates of \hat{U} can be characterized by the statistics of its quasienergy levels and the inverse participation ratios (IPRs). Let us denote the normalized right eigenvectors of \hat{U} and their corresponding quasienergies as $\{|\psi_j\rangle|j =$

1, ..., L} and $\{E_j | j = 1, ..., L\}$. Along the real axis, the spacing between the *j*th and the (j-1)th quasienergies of \hat{U} is given by $\epsilon_j = \text{Re}E_j - \text{Re}E_{j-1}$, from which we obtain the ratio between two adjacent spacings of quasienergy levels as

$$g_j = \frac{\min(\epsilon_j, \epsilon_{j+1})}{\max(\epsilon_j, \epsilon_{j+1})}, \qquad j = 2, \dots, L-1.$$
(119)

Here the $\max(\epsilon_j, \epsilon_{j+1})$ and $\min(\epsilon_j, \epsilon_{j+1})$ are the maximum and minimum between the ϵ_j and ϵ_{j+1} , respectively. The statistical property of adjacent gap ratios can then be obtained by averaging over all g_j in the thermodynamic limit, i.e.,

$$\overline{g} = \lim_{L \to \infty} \frac{1}{L} \sum_{j} g_j.$$
(120)

We have $\overline{g} \to 0$ if all the bulk Floquet eigenstates of \hat{U} are extended. Comparatively, we expect \overline{g} to approach a constant $\overline{g}_{\max} > 0$ if all the bulk eigenstates of \hat{U} are localized. If we find $\overline{g} \in (0, \overline{g}_{\max})$, extended and localized eigenstates of \hat{U} should coexist and be separated by some mobility edges in the quasienergy spectrum. The \overline{g} can thus be utilized to distinguish between phases with different localization nature in 1D non-Hermitian Floquet systems from the perspective of level statistics. In the lattice representation, we can expand the right eigenvector $|\psi_j\rangle$ as $|\psi_j\rangle = \sum_{n=1}^L \psi_n^j |n\rangle$, where $\psi_n^j = \langle n | \psi_j \rangle$ and $\sum_n |\psi_n^j|^2 = 1$. The inverse and normalized participation ratios of $|\psi_j\rangle$ in the real space can then be defined as

$$IPR_j = \sum_{n=1}^{L} |\psi_n^j|^4, \qquad NPR_j = \frac{1}{L} IPR_j^{-1}.$$
 (121)

For a localized Floquet eigenstate $|\psi_j\rangle$, we have $\text{IPR}_j \to \lambda_j$ and $\text{NPR}_j \to 0$ in the thermodynamic limit, where the Lyapunov exponent λ_j of $|\psi_j\rangle$ could be a function of its quasienergy E_j . If $|\psi_j\rangle$ represents an extended state, we have $\text{IPR}_j \to 0$ and $\text{NPR}_j \to 1$. The global localization features of the Floquet system described by \hat{U} can then be extracted from the combined information of $\{\text{IPR}_j | j = 1, ..., L\}$ and $\{\text{NPR}_j | j = 1, ..., L\}$. For ease of usage, we introduce the following localization indicators

$$IPR_{\max} = \max_{j \in \{1,\dots,L\}} (IPR_j), \qquad (122)$$

$$IPR_{\min} = \min_{j \in \{1, \dots, L\}} (IPR_j), \qquad (123)$$

$$\zeta = \log_{10}(\text{IPR}_{\text{ave}} \cdot \text{NPR}_{\text{ave}}), \qquad (124)$$

where the IPR_{ave} = $\frac{1}{L} \sum_{j=1}^{L} IPR_j$ and NPR_{ave} = $\frac{1}{L} \sum_{j=1}^{L} NPR_j$ are the averages of IPR_j and NPR_j over all Floquet eigenstates. It is not hard to see that in the metallic phase where all Floquet eigenstates are extended, we have IPR_{max} $\rightarrow 0$, IPR_{min} $\rightarrow 0$ and $\zeta \sim -\log_{10}(L) \rightarrow -\infty$ in the thermodynamic limit $L \rightarrow \infty$. Instead, in the insulating phase with all Floquet eigenstates being localized, we have IPR_{max} > 0, IPR_{min} > 0 and $\zeta \rightarrow -\infty$. In the critical phase where extended and localized Floquet eigenstates are coexistent, we have IPR_{max} > 0, IPR_{min} $\rightarrow 0$ together with a finite ζ . Assembling the information obtained from IPR_{max}, IPR_{min}, and ζ thus allows us to distinguish the extended, localized, and critical mobility edge phases of a non-Hermitian Floquet system with disorder. These tools will be applied to our study of the Floquet NHQC in Subsec. III D.

We can also probe the transport nature of distinct non-Hermitian Floquet phases from the wave packet dynamics. Let us consider a generic and normalized initial state $|\Psi(0)\rangle$ in the lattice representation. After the stroboscopic evolution over a number of ℓ driving periods by \hat{U} , the final state turns out to be $|\Psi'(t = \ell T)\rangle = \hat{U}^{\ell}|\Psi(0)\rangle$. Since the Floquet operator \hat{U} is not unitary for a non-Hermitian $\hat{H}(t)$ in general, the norm of $|\Psi(0)\rangle$ cannot be preserved during the evolution. We can express the normalized state at $t = \ell T$ as $|\Psi(t)\rangle = |\Psi'(t)\rangle/\sqrt{\langle \Psi'(t)|\Psi'(t)\rangle}$. The real space expansion $|\Psi(t)\rangle = \sum_{n=1}^{L} \psi_n(t)|n\rangle$ then provides us with the probability amplitude $\psi_n(t) = \langle n|\Psi(t)\rangle$ of the normalized final state $|\Psi(t)\rangle$ on the lattice site n at time t. Using the collection of amplitudes $\{\psi_n(t)|n=1,...,L\}$, we can define the following dynamical quantities

$$\langle x(t) \rangle = \sum_{n=1}^{L} n |\psi_n(t)|^2, \qquad \langle x^2(t) \rangle = \sum_{n=1}^{L} n^2 |\psi_n(t)|^2,$$
(125)

$$\Delta x(t) = \sqrt{\langle x^2(t) \rangle - \langle x(t) \rangle^2}, \qquad v(t) = \frac{1}{t} \sqrt{\langle x^2(t) \rangle}, \tag{126}$$

where the stroboscopic time $t = \ell T$ and $\ell \in \mathbb{Z}$. It is clear that the $\langle x(t) \rangle$, $\Delta x(t)$, and v(t) describe the center, standard deviation, and spreading speed of the wavepacket $|\Psi(t)\rangle$ in the lattice representation, respectively. For simplicity, we usually choose the initial state $|\Psi(0)\rangle$ to be exponentially localized at a single site that is deep inside the bulk of the lattice. If the

Floquet system described by \hat{U} resides in a localized phase, we expect the $\langle x(t) \rangle$ and $\Delta x(t)$ to stay around their initial values, which means that the wavepacket almost does not move and spread. In this case, the speed v(t) should also tend to zero for a long-time evolution $(\ell \gg 1)$. If the system stays in an extended phase, we expect the increasing of $\Delta x(t)$ with time due to the spreading of the initial wavepacket. Meanwhile, the $\langle x(t) \rangle$ may or may not change with time, depending on whether the hopping amplitudes are symmetric. For a nearestneighbor, nonreciprocal hopping as in Eq. (116), we may have $\langle x(t) \rangle \propto t$ and $\Delta x(t) \propto \sqrt{t}$ for a metallic phase of the system. The v(t) should take a maximal possible value $v_{\max}(t)$ in this case. If the system is prepared in a critical mobility edge phase, we expect both the $\langle x(t) \rangle$ and $\Delta x(t)$ to show intervening behaviors, while the averaged spreading speed v(t)should satisfy $0 < v(t) < v_{\max}(t)$. Therefore, we can exploit the dynamical quantities $\langle x(t) \rangle$, $\Delta x(t)$, and v(t) to discriminate phases with different localization properties in disordered non-Hermitian Floquet systems. We will illustrate such an application for Floquet NHQC in Subsec. III D.

III. NON-HERMITIAN FLOQUET PHASES OF MATTER

This section collects and treats typical examples of non-Hermitian Floquet phases discovered in our previous work [254–264]. We will see that the interplay between periodic driving fields and gain/loss or nonreciprocal effects could induce rich phases and transitions in non-Hermitian Floquet systems.

A. Non-Hermitian Floquet exceptional topology

Let us start with a simple model, whose Floquet effective Hamiltonian and quasienergy dispersion are given by Eqs. (55) and (57). To be explicit, we choose

$$h_x = \sin k, \qquad h_z = \mu + \cos k, \tag{127}$$

where $k \in (-\pi, \pi]$ denotes the quasimomentum in the first Brillouin zone, and the system parameter $\mu \in \mathbb{R}$. The H_{eff} in Eq. (55) thus describes the Bloch Hamiltonian of a 1D two-band lattice model under the PBC in the rotating frame, where μ corresponds to the amplitude of onsite potential and the nearest-neighbor hopping amplitude has been chosen to be the unit of energy. From Eqs. (57) and (127), we find that the two quasienergy bands of H_{eff} could meet with each other at the quasienergy $\omega/2$ when

$$\mu = \begin{cases} \frac{\omega}{2} \pm \frac{\gamma}{2} - 1, & \text{for} \quad k = 0, \\ \frac{\omega}{2} \pm \frac{\gamma}{2} + 1, & \text{for} \quad k = \pi. \end{cases}$$
(128)

If μ satisfies one of the above two equalities, there will be a second-order Floquet EP at k = 0 or $k = \pi$ in the conventional Brillouin zone. Note that both the quasienergy of this Floquet EP and its location in the parameter space depend on the frequency ω , which high-lights the impact of the harmonic driving field on phase transitions in the system. Moreover, the appearance of an EP in k-space usually implies the breakdown of bulk-edge correspondence in conventional topological phases. This issue might be overcome by incorporating the formalism of generalized Brillouin zone (GBZ) and non-Bloch band theory [163, 165]. Following the standard recipe, we first make the substitutions $e^{ik} \rightarrow \beta$ and $e^{-ik} \rightarrow \beta^{-1}$, where the complex number $\beta \in \text{GBZ}$. The effective Hamiltonian in Eq. (55) now takes the form of

$$H_{\text{eff}}(\beta) = \frac{\omega}{2}\sigma_0 + \left(\frac{\beta - \beta^{-1}}{2i} + i\frac{\gamma}{2}\right)\sigma_x + \left(\mu + \frac{\beta + \beta^{-1}}{2} - \frac{\omega}{2}\right)\sigma_z,\tag{129}$$

with the quasienergy bands

$$\varepsilon_{\pm}(\beta) = \frac{\omega}{2} \pm \sqrt{\left(\frac{\beta - \beta^{-1}}{2i} + \frac{i\gamma}{2}\right)^2 + \left(\mu - \frac{\omega}{2} + \frac{\beta + \beta^{-1}}{2}\right)^2}.$$
(130)

We next focus on the non-constant part of $\varepsilon_{\pm}(\beta)$, whose square takes the form of

$$\epsilon_{\pm}^2(\beta) = \frac{a\beta^2 + b\beta + c}{\beta},\tag{131}$$

where

$$a = \mu - \frac{\omega}{2} + \frac{\gamma}{2}, \qquad b = 1 + \left(\mu - \frac{\omega}{2}\right)^2 - \frac{\gamma^2}{4}, \qquad c = \mu - \frac{\omega}{2} - \frac{\gamma}{2}.$$
 (132)

The $\varepsilon_{\pm}^2(\beta)$ is a Laurent polynomial of β . According to Ref. [165] (see also Ref. [280]), $\beta \in \text{GBZ}$ for the $H_{\text{eff}}(\beta)$ if $\epsilon_{\pm}^2(\beta) = \epsilon_{\pm}^2(\beta e^{i\theta})$, with θ being a phase factor. For Eq. (131), this means that $\beta^2 = c e^{-i\theta}/a$. The GBZ is thus a circle of the radius

$$r = |\beta| = \left|\frac{\mu - \omega/2 - \gamma/2}{\mu - \omega/2 + \gamma/2}\right|^{1/2}.$$
(133)

It is clear that the GBZ radius is controlled by both the driving field (through ω) and the non-Hermitian effect (through γ). In the Hermitian limit ($\gamma \rightarrow 0$), we have $r \rightarrow 1$ and the GBZ is reduced to the conventional BZ, as expected. The GBZ becomes ill defined if $\mu - \omega/2 = \pm \gamma/2$, where the radius r is zero or infinity. Finally, we can use the GBZ to determine the gap closing (phase transition) points of the system under the OBC. Setting the $\epsilon_{\pm}^2(\beta) = 0$ in Eq. (131), we find

$$\mu = \frac{\omega}{2} \pm \begin{cases} \sqrt{\gamma^2/4 + 1}, & |\gamma| \le 2, \\ \sqrt{\gamma^2/4 \pm 1}, & |\gamma| > 2. \end{cases}$$
(134)

These bulk gap-closing conditions are clearly different from those found in conventional BZ under the PBC in Eq. (128). Using the transfer matrix method [167], we can further obtain the parameter regions in which degenerate Floquet edge states appear at the quasienergy $\omega/2$ (anomalous Floquet π edge modes), i.e.,

$$\mu - \frac{\omega}{2} \in \begin{cases} \left(-\sqrt{\gamma^2/4 + 1}, \sqrt{\gamma^2/4 + 1}\right), & |\gamma| \le 2, \\ \left(-\sqrt{\gamma^2/4 + 1}, -\sqrt{\gamma^2/4 - 1}\right) \cup \left(\sqrt{\gamma^2/4 - 1}, \sqrt{\gamma^2/4 + 1}\right), & |\gamma| > 2. \end{cases}$$
(135)

Finally, we can characterize the different topological phases of the system by a non-Bloch winding number [168]

$$W = \int_{\text{GBZ}} \frac{d\beta}{4\pi i} \text{Tr}[\sigma_y \mathcal{Q}(\beta) \partial_\beta \mathcal{Q}(\beta)], \qquad (136)$$

where σ_y is the chiral symmetry operator of the shifted effective Hamiltonian $H_{\text{eff}}(\beta) - \frac{\omega}{2}\sigma_0$. The Q matrix is defined as $\mathcal{Q}(\beta) = |\psi_+(\beta)\rangle \langle \tilde{\psi}_+(\beta)| - |\psi_-(\beta)\rangle \langle \tilde{\psi}_-(\beta)|$. $|\psi_\pm(\beta)\rangle$ are the right eigenvectors of $H_{\text{eff}}(\beta)$ with the quasienergies $\varepsilon_{\pm}(\beta)$. $\langle \tilde{\psi}_{\pm}(\beta)|$ are the corresponding left eigenvectors. For our model, their explicit expressions are given by

$$|\psi_{\pm}(\beta)\rangle = \frac{1}{\sqrt{2(\varepsilon_{\pm} - \frac{\omega}{2})(\varepsilon_{\pm} - h_z)}} \begin{pmatrix} h_x + i\gamma/2\\ \varepsilon_{\pm} - h_z \end{pmatrix},$$
(137)




FIG. 2. Floquet spectra and winding numbers of the harmonically driven 1D non-Hermitian lattice model. The gray dots at each μ denote the real [in (a) and (b)] and imaginary [in (c) and (d)] parts of quasienergies obtained under the PBC in the conventional BZ. The blue dots at each μ denote the real [in (a) and (b)] and imaginary [in (c) and (d)] parts of quasienergies obtained under the OBC in the GBZ. The yellow circles denote the winding number W, which is equal to 1 (0) in the topological (trivial) phases with (without) anomalous Floquet edge modes at the quasienergies $\pm \pi$, which are highlighted by the red solid lines. Other system parameters are chosen as $(\omega, \gamma) = (2\pi, 1)$ $[(\omega, \gamma) = (2\pi, 3)]$ for the panels (a) and (c) [(b) and (d)].

$$\langle \widetilde{\psi}_{\pm}(\beta) | = \frac{1}{\sqrt{2(\varepsilon_{\pm} - \frac{\omega}{2})(\varepsilon_{\pm} - h_z)}} \left(h_x + i\gamma/2 \ \varepsilon_{\pm} - h_z \right).$$
(138)

In Fig. 2, we show the Floquet spectra of H_{eff} under different boundary conditions and the non-Bloch winding numbers of H_{eff} versus μ for some typical cases. We see that the Floquet spectra and the gap closing points at the quasienergies $E = \pm \pi$ could indeed be very different under the PBC and the OBC. Some high-order EPs can be identified from the bulk Floquet spectra under the OBC, which clearly signify the emergence of Floquet exceptional topology. Besides, the non-Bloch winding number W can correctly discriminate different topological phases and characterize topological phase transitions accompanied by quasienergy-band touchings under the OBC at $E = \pi$. We find W = 1 (W = 0) in the topologically nontrivial (trivial) phases with (without) anomalous Floquet π edge modes. The bulk-edge correspondence is then recovered. The gap-closing points and the parameter regions with Floquet π modes are found to be perfectly coincident with the predictions of Eqs. (134) and (135). These observations demonstrate the applicability of non-Bloch band theories to non-Hermitian Floquet systems. It is also straightforward to check that the system possesses Floquet NHSEs in a broad parameter regime under the OBC. The simple model introduced in this subsection thus allows us to get a bird's-eye view on the nontrivial topological phenomena that could be brought about by the interplay between Floquet driving fields and non-Hermitian effects. Further examples with more striking features will be reviewed in the following subsections.

B. Non-Hermitian Floquet topological insulators

In this subsection, we review three types of non-Hermitian Floquet topological insulators in one and two spatial dimensions. For all the cases, we uncover that the collaboration between driving and gain/loss or nonreciprocal effects could induce topological insulating states unique to non-Hermitian Floquet systems. They are featured by large topological invariants, many topological edge or corner modes, and separated by rich topological phase transitions. The bulk-edge (or bulk-corner) correspondence will also be established for each class of systems considered in this subsection.

1. First-order topological phase

We start with the characterization of 1D non-Hermitian Floquet topological insulators. One typical model that incorporates their rich topological properties is the following periodically quenched dimerized tight-binding lattice, whose time-dependent Hamiltonian reads [254]

$$\hat{H}(t) = \begin{cases} \hat{H}_1, & t \in [\ell T, \ell T + T/2), \\ \hat{H}_2, & t \in [\ell T + T/2, \ell T + T), \end{cases}$$
(139)

where

$$\hat{H}_1 = \sum_n (ir_y \hat{c}_{n+1}^{\dagger} \hat{c}_n + \text{H.c.} + 2i\gamma \hat{c}_n^{\dagger} \hat{c}_n) \otimes \sigma_y, \qquad (140)$$

$$\hat{H}_2 = \sum_n (r_x \hat{c}_n^{\dagger} \hat{c}_{n+1} + \mu \hat{c}_n^{\dagger} \hat{c}_n + \text{H.c.}) \otimes \sigma_x.$$
(141)




FIG. 3. Non-Hermitian Floquet topological insulator: (a) a schematic diagram of the lattice model, (b)–(e) quasienergy spectra on the complex plane under the OBC, (f)–(g) topological phase diagrams under the PBC, and (h)–(j) Floquet spectra and edge states under the OBC [254].

Here T is the driving period and we will set $\hbar = T = 1$ in our calculations. \hat{c}_n^{\dagger} (\hat{c}_n) creates (annihilates) a particle in the unit cell n of the lattice. σ_{α} , $\alpha = x, y, z$, are Pauli matrices acting on the two sublattice degrees of freedom A and B within each unit cell. The system parameters r_x, r_y, μ, γ are all real. The non-Hermitian effect is introduced by the nonreciprocal intracell coupling term $2i\gamma\sigma_y$ applied over the first half of each driving period [see Fig. 3(a) for an illustration of the model]. Experimentally, such a term might be realized by coupled-resonator optical waveguide with asymmetric internal scattering [281].

Considering the one-period evolution of the system from $t = \ell + 0^-$ to $t = \ell + 1 + 0^-$, the Floquet operator of $\hat{H}(t)$ takes the form of $\hat{U} = e^{-\frac{i}{2}\hat{H}_2}e^{-\frac{i}{2}\hat{H}_1}$. Its quasienergy spectrum and Floquet eigenstates are obtained by solving the eigenvalue equation $\hat{U}|\psi\rangle = e^{-iE}|\psi\rangle$. Under the OBC, we show the Floquet spectrum of \hat{U} on the complex quasienergy plane $\operatorname{Re} E - \operatorname{Im} E$ for some typical cases in Figs. 3(b)–(e) [with $(\mu, r_y, \gamma) = (0, \pi/2, \operatorname{arccosh}(\sqrt{2}))$, $r_x = \pi/2, \pi, 3\pi/2, 2\pi$, and the number of unit cells N = 150]. The numbers of degenerate Floquet edge modes at the quasienergies E = 0 and $E = \pi$ are denoted by n_0 and n_{π} in the corresponding figure captions. We observe one or multiple pairs of edge modes at both the center (E = 0) and boundary $(E = \pi)$ of the first quasienergy Brillouin zone. Interestingly, when the numbers of these edge modes are the same, i.e., $n_0 = n_{\pi}$, we find them to appear in different types of quasienergy gaps [see Figs. 3(c) and 3(e)]. For example, we find a pair of Floquet zero modes in the line gap at E = 0, while another pair of Floquet π modes are found in the point gap at $E = \pi$. This is rather different from the situation in Hermitian Floquet systems, where the quasienergy is real and there is no distinction between point and line gaps. It is also different from the cases encountered in non-Hermitian static systems with two bands, where topological edge modes can only appear in either a point gap or a line gap at E = 0. The Floquet spectra with hybrid (point plus line) quasienergy gaps are thus unique to non-Hermitian Floquet systems.

To understand the origin of the non-Hermitian Floquet edge modes at E = 0 and $E = \pi$, we study the bulk topological properties of the system. Under the PBC, the Floquet operator of the system reads

$$U(k) = e^{-i(\mu + r_x \cos k)\sigma_x} e^{-i(r_y \sin k + i\gamma)\sigma_y},$$
(142)

where $k \in [-\pi, \pi)$ is the quasimomentum. The quasienergy spectrum of U(k) is given by

$$\pm E(k) = \pm \arccos[\cos(\mu + r_x \cos k) \cos(r_y \sin k + i\gamma)], \qquad (143)$$

and the gap-closing conditions can be obtained analytically by setting $\cos[E(k)] = \pm 1$ [254]. Transforming U(k) to symmetric time frames, we obtain

$$U_1(k) = e^{-\frac{i}{2}(\mu + r_x \cos k)\sigma_x} e^{-i(r_y \sin k + i\gamma)\sigma_y} e^{-\frac{i}{2}(\mu + r_x \cos k)\sigma_x},$$
(144)

$$U_2(k) = e^{-\frac{i}{2}(r_y \sin k + i\gamma)\sigma_y} e^{-i(\mu + r_x \cos k)\sigma_x} e^{-\frac{i}{2}(r_y \sin k + i\gamma)\sigma_y}.$$
(145)

It is clear that both $U_1(k)$ and $U_2(k)$ respect the chiral (sublattice) symmetry $S = \sigma_z$. We can thus characterize the non-Hermitian Floquet topological phases of U(k) by the winding numbers (w_0, w_π) according to Eq. (93). In Figs. 3(f) and 3(g), we present the w_0 and w_π of U(k) versus (r_x, r_y) respectively, yielding the topological phase diagram of the system. We find various non-Hermitian Floquet insulating phases. They are characterized by large




FIG. 4. Dynamical characterization of non-Hermitian Floquet topological phases: (a) winding numbers (w_0, w_π) , MCDs (C_0, C_π) , and (b)–(e) dynamic winding numbers from stroboscopic time-averaged spin textures [256].

integer winding numbers and separated by a series of topological phase transitions with quasienergy level crossings at E = 0 [white solid lines in Fig. 3(f)] and $E = \pi$ [white dashed lines in Fig. 3(g)]. The winding numbers (w_0, w_π) could become arbitrarily large with the increase of system parameters (e.g., the intercell hopping amplitude r_x), which implies that Floquet phases with large topological invariants could indeed survive even with finite non-Hermitian effects [$\mu = 0$ and $\gamma = \operatorname{arccosh}(\sqrt{2})$ in Figs. 3(f)–(g)]. Moreover, these winding numbers correctly count the numbers of degenerate Floquet edge modes n_0 and n_{π} at E = 0and $E = \pi$ under the OBC. That is, we have the bulk-edge correspondence for our 1D chiral symmetric non-Hermitian Floquet topological insulator as [254]

$$n_0 = 2|w_0|, \qquad n_\pi = 2|w_\pi|.$$
 (146)

Representative examples of the Floquet spectra under the OBC are shown in Figs. 3(h)– (j) [with $\mu = 0$, $r_y = \pi/2$, $\gamma = \operatorname{arccosh}(\sqrt{2})$ and the number of unit cells N = 150], which provide verifications for the found bulk-edge correspondence in Eq. (146). Note in passing that the system does not show NHSE under the OBC for $\mu = 0$, even though the Floquet spectra possess point quasienergy gaps under the PBC. Meanwhile, many pairs of degenerate edge modes at the quasienergies E = 0 and $E = \pi$ are found in the topological phases with large winding numbers. As these topological edge modes could survive in the presence of non-Hermitian effects, they may provide more resources for the realization of robust quantum state transfer and quantum computing schemes in open systems.

The winding numbers of non-Hermitian Floquet topological insulators can be experimentally probed by measuring the stroboscopic time-averaged spin textures [254] or the MCDs [255], as introduced in Subsecs. II F 1 and II F 2. Following Ref. [256], we consider a periodically quenched non-Hermitian lattice model, whose Floquet operator in the momentum space reads $U(k) = e^{-iJ_2 \sin k\sigma_y} e^{-iJ_1 \cos k\sigma_x}$. Here $J_1 = u_1 + iv_1$, $J_2 = u_2 + iv_2$, and $u_1, u_2, v_1, v_2 \in \mathbb{R}$. The winding numbers (w_0, w_π) of U(k) can be obtained using the approach of symmetric time frames [see Eqs. (89)–(93)]. Following the recipes outlined in Subsecs. II F 1 and II F 2, we obtain the MCDs in Fig. 4(a) [with $v = v_1 = v_2$ and $(u_1, u_2) = (4.5\pi, 0.5\pi)$] and the dynamic winding angles of time-averaged spin textures in Figs. 4(b)–(e) [with $(u_1, u_2) = (4.5\pi, 0.5\pi)$ and $v = v_1 = v_2 = 0.2\pi, 0.35\pi, 0.6\pi, 0.9\pi$]. In all the cases, the obtained MCDs and dynamic winding numbers are coincident with the winding numbers (w_0, w_π) of Floquet operator U(k)even when there are finite non-Hermitian effects $(v \neq 0)$. The dynamic winding numbers and the MCDs thus provide us with two complementary approaches to dynamically characterize 1D non-Hermitian Floquet topological phases with chiral (sublattice) symmetry in momentum and in real spaces, respectively. They can be applied to detect non-Hermitian Floquet matter in different types of experimental setups [213, 222].

The models we considered above in this subsection do not possess the NHSE. It remains unclear whether the rich non-Hermitian Floquet topology could coexist with NHSEs, and how to characterize the topological bulk-edge correspondence in the presence of Floquet NHSE. To address this issue, we consider a periodically quenched non-Hermitian SSH model [261], whose k-space Hamiltonian under the PBC takes the form of

$$H(k,t) = \begin{cases} (\mu + J_1 \cos k + i\lambda \sin k)\sigma_x, & t \in [\ell T, \ell T + T/2), \\ (J_2 \sin k + i\lambda \cos k)\sigma_y, & t \in [\ell T + T/2, \ell T + T). \end{cases}$$
(147)

Here μ and $J_1/2 + J_2/2$ are the intracell and intercell hopping amplitudes. The non-Hermiticity is introduced by the asymmetric parts of hopping amplitudes $\pm i\lambda/2$ between the two sublattices. A sketch of the model is shown in Fig. 5. The Floquet operator associated with H(k,t) (for $\hbar = 1$ and T = 2) reads

$$U(k) = e^{-i(J_2 \sin k + i\lambda \cos k)\sigma_y} e^{-i(\mu + J_1 \cos k + i\lambda \sin k)\sigma_x}.$$
(148)




FIG. 5. Non-Hermitian Floquet topological insulator with NHSE: a schematic diagram of the lattice model, (a)–(b) topological phase diagrams, (c)–(f) static and dynamic winding angles, (g) quasienergy gap functions under the OBC, and (h)–(k) Floquet topological corner modes at zero, π quasienergies and skin-localized bulk modes. Under the OBC, the numbers of Floquet zero and π edge modes in each topological phase are shown below the horizontal axis of (g). System parameters are $(J_1, J_2, \mu, \lambda) = (\pi, 0.5\pi, 0.4\pi, 0.25)$ and $(J_1, J_2, \mu, \lambda) = (2\pi, 0.5\pi, 0.4\pi, 0.25)$ for (h, i) and (j, k) [261].

In symmetric time frames, it respects the chiral symmetry $S = \sigma_z$ [261]. We may thus use the winding numbers (w_0, w_π) to characterize the Floquet topological phases of U(k). The calculation of such winding numbers following Eqs. (89)–(93) leads to the topological phase diagrams in Fig. 5(a)–(b). Interestingly, we find that despite Floquet topological insulators with large integer winding numbers, there are also various phases with half-integer quantized topological invariants. These unique phases are absent in the Hermitian limit of the model. Their appearance is due to the Floquet exceptional topology induced by non-Hermitian effects in the system. The parameter regions which accommodate half-integer topological phases may further show Floquet NHSEs under the OBC. In Figs. 5(c)–(f), we compare the dynamic winding angles of U(k) with the winding patterns of its Floquet Hamiltonian vectors in different time frames [with $(J_1, J_2, \mu, \lambda) = (2.4\pi, 0.5\pi, 0.4\pi, 0.25)$] [261]. Their coincidence implies that we can use the dynamic winding numbers introduced in Subsec. II F 1 to probe the half-integer quantized topological invariants of 1D non-Hermitian Floquet topological insulators with chiral symmetry.

Under the OBC, the bulk Floquet spectra and the gap-closing points of quasienergies are found to be different from those under the PBC, as shown in Fig. 5(g) [with $(J_2, \mu, \lambda) =$ $(0.5\pi, 0.4\pi, 0.25)$] [261]. This means that the bulk winding numbers w_0 (diamonds) and w_π (pentagrams) of U(k) cannot be used to characterize the edge states and topological phase transitions under the OBC. To overcome this issue, we switch to the OBC winding numbers (ν_0, ν_π) [defined as the (W_0, W_π) in Eq. (96)]. These winding numbers are found to be integer quantized. They change their values only when the Floquet spectrum of the system closes its gap at E = 0 or $E = \pi$ under the OBC. Furthermore, in each gapped phase, the winding numbers (ν_0, ν_π) [circles and crosses in Fig. 5(g)] are related to the numbers of zero and π Floquet edge modes (n_0, n_π) through the bulk-edge correspondence $(n_0, n_\pi) = 2(|\nu_0|, |\nu_\pi|)$. This relation holds even with NHSEs in our system [261]. Therefore, our work provides a dual topological characterization of non-Hermitian Floquet topological insulators under different boundary conditions. In Figs. 5(h)-(k), we present examples of the skin-localized bulk states [in Figs. 5(h) and 5(j)] and two types of Floquet edge states [in Figs. 5(i) and 5(k)] in our system, which are also consistent with the theoretical predictions [261].

Putting together, we found different types of 1D non-Hermitian Floquet topological insulators in a series of studies [254–256, 258, 261]. All the models considered in these studies exhibit rich non-Hermitian Floquet phases with large topological invariants, many topological edge states, and unique topological transitions induced by the interplay between non-Hermitian effects and time-periodic driving fields. The bulk-edge correspondences and dynamic topological characterizations of these intriguing new phases are also established, leading to an explicit and all-round physical picture of 1D non-Hermitian Floquet topological matter. In the following subsections, we uncover the essential role of Floquet engineering in other types of non-Hermitian topological matter.

2. Second-order topological phase

We next consider the example of a non-Hermitian Floquet second-order topological insulator (SOTI) in two dimensions. An *n*th-order topological insulator in d ($d \ge n$) spatial dimensions possesses localized eigenmodes along its (d - n)-dimensional boundaries, which are topologically nontrivial and protected by the symmetries of the *d*-dimensional bulk. For example, an SOTI in two dimensions usually owns localized topological states around its zero-dimensional geometric corners. Assisted by time-periodic drivings, degenerate corner modes can further appear at different quasienergies in Floquet second-order topological phases [80, 282].

In this subsection, we focus on one typical model of non-Hermitian Floquet SOTI, which is obtained following the recipe of coupled-wire construction [80]. A schematic diagram of the static lattice model is shown in Fig. 6(a). The system is formed by a stack of SSH chains along the y-direction, with dimerized couplings (J_1, J_2) between adjacent chains. We consider a Floquet variant of the model by applying time-periodic quenches to the coupling and onsite parameters (J_1, J_2, μ) along the y-direction. Under the PBC along both x and y directions, the time-dependent Bloch Hamiltonian of the Floquet model takes the form of

$$H(k_x, k_y) = H_x(k_x) \otimes \tau_0 + \sigma_0 \otimes H_y(k_y, t), \tag{149}$$

where $(k_x, k_y) \in [-\pi, \pi) \times [-\pi, \pi)$ are the quasimomenta. σ_0 and τ_0 are two by two identity matrices. The Hamiltonians of the 1D subsystems $H_x(k_x)$ and $H_y(k_y, t)$ are explicitly given by

$$H_x(k_x) = [(J-\delta) + (J+\delta)\cos k_x]\sigma_x + (J+\delta)\sin k_x\sigma_y,$$
(150)

$$H_y(k_y, t) = \begin{cases} 2J_1 \cos k_y \tau_x & t \in [\ell T, \ell T + T/2) \\ 2(\mu + J_2 \sin k_y) \tau_z & t \in [\ell T + T/2, \ell T + T) \end{cases}.$$
 (151)

Here T is the driving period and $\ell \in \mathbb{Z}$. $\sigma_{x,y,z}$ and $\tau_{x,y,z}$ are Pauli matrices acting on the sublattice degrees of freedom in x and y directions, respectively. Setting $\hbar = T = 1$, $J = \delta = \Delta/2$, and choosing $\mu = u + iv$ $(u, v \in \mathbb{R})$, the Floquet operator of the system in **k**-space is found to be [259]

$$U(k_x, k_y) = e^{-i\Delta(\cos k_x \sigma_x + \sin k_x \sigma_y)} \otimes e^{-ih_z(k_y)\tau_z} e^{-ih_x(k_y)\tau_x},$$
(152)

where

$$h_x(k_y) = J_1 \cos k_y, \qquad h_z(k_y) = u + iv + J_2 \sin k_y.$$
 (153)

The non-Hermitian effect is brought about by the onsite gain and loss encoded in the term $iv\tau_z$.

In symmetric time frames, the system has the chiral (sublattice) symmetry $S = \sigma_z \otimes \tau_y$ [259]. One can thus characterize its Floquet second-order topological phases by integer winding numbers [259]

$$\nu_0 = \frac{\nu_1 + \nu_2}{2}, \qquad \nu_\pi = \frac{\nu_1 - \nu_2}{2}.$$
(154)

Here $\nu_{\alpha} = ww_{\alpha}$ for $\alpha = 1, 2$. w is the winding number of the static Hamiltonian $\Delta(\cos k_x \sigma_x + \sin k_x \sigma_y)$, which is always equal to one for the topological flat-band limit of the SSH model. (w_1, w_2) are the winding numbers of the subsystem Floquet operator $e^{-ih_z(k_y)\tau_z}e^{-ih_x(k_y)\tau_x}$, which can be defined via the recipe outlined in Subsec. II E. They both take integer-quantized values even with the non-Hermitian effects considered in our system [259].

In Fig. 6(b), we show the topological phase diagrams of $U(k_x, k_y)$ in the parameter space $J_1 - v$ [with $(\Delta, J_2, u) = (\pi/20, 3\pi, 0)$]. We find that both the winding numbers (ν_0, ν_π) can take large values, which indicates the presence of non-Hermitian Floquet SOTI phases with large topological invariants in our setting. Furthermore, with the increase of the gain and loss strength v, the system can undergo topological phase transitions and even enter non-Hermitian Floquet SOTI phases with larger winding numbers. These non-Hermiticity boosted topological signatures are also not expected in non-driven systems. Their appearance is thus due to the corporation between Floquet driving fields and non-Hermitian gain




FIG. 6. Non-Hermitian Floquet SOTI: (a) a schematic diagram of the static lattice model, (b)–(c) topological phase diagrams, (d)–(e) quasienergy gap functions under the OBC, (f)–(h) Floquet topological corner modes at the quasienergy zero, and (i)–(j) Floquet topological corner modes at the quasienergy π [259].

and loss.

Under the OBC along both x and y directions, localized Floquet eigenmodes with the quasienergies zero and π appear at the four corners of the square lattice. Some spatial profiles of these corner modes are shown in Figs. 6(f)–(j) [with $(\Delta, J_1, J_2, u, v) = (\pi/20, 2.5\pi, 3\pi, 0, 2)$]. There are twelve (eight) Floquet corner modes at the quasienergy zero (π) . More generally, their numbers (n_0, n_π) are related to the winding numbers (ν_0, ν_π) through the algebraic relations [259]

$$(n_0, n_\pi) = 4(|\nu_0|, |\nu_\pi|). \tag{155}$$

We thus established the bulk-corner correspondence for a class of non-Hermitian Floquet SOTI with chiral symmetry. With the growth of the non-Hermitian parameter v, topological phase transitions accompanied by the increase of bulk winding numbers (ν_0, ν_π) and corner mode numbers (n_0, n_π) are observed from the gap functions $(F_0, F_\pi) \equiv (\frac{1}{\pi}|\varepsilon|, \frac{1}{\pi}\sqrt{(|\text{Re}\varepsilon| - \pi)^2 + (\text{Im}\varepsilon)^2})$ under the OBC (here ε denotes the quasienergy), as showcased in Figs. 6(d)–(e). The topological phase transition points v_1 and v_2 under the OBC are precisely consistent with the gap closing points of the 2D bulk under the PBC, which can be found analytically [259]. Finally, even with non-Hermitian effects, the bulk winding numbers (ν_0, ν_π) can be dynamically probed by 2D generalizations of the MCD as introduced in Subsec. II F 2 (see Ref. [259] for more details).

Overall, we discovered rich non-Hermitian Floquet SOTI phases in a periodically quenched 2D lattice with balanced gain and loss. Different from the static case, each SOTI phase is now depicted by two topological invariants (ν_0, ν_π) . Many non-Hermitian Floquet SOTIs with large topological invariants and gain- or loss-induced topological phase transitions were identified. Under the OBC, the winding numbers (ν_0, ν_π) determine the numbers of symmetry-protected Floquet corner modes at zero and π quasienergies. The interplay between driving and dissipation thus results in a series of non-Hermitian Floquet SOTI phases with multiple zero and π corner modes, which may find applications in topological state preparations, information scrambling and quantum computation. The work of Ref. [259] offers one of the earliest findings of rich non-Hermitian Floquet topological phases beyond one spatial dimension. In the meantime, it introduces a generic scheme to construct non-Hermitian Floquet higher-order topological phases across different physical dimensions, which is expected to be applicable across insulating, superconducting and semi-metallic systems. Note in passing that some latter studies also considered the NHSEs and anomalous π modes in Floquet higher-order topological phases with somewhat different focuses [235, 238, 246].

3. qth-root topological phase

We now consider an exotic class of non-Hermitian Floquet topological phase, which could have symmetry-protected boundary states at the quasienergies $p\pi/q$, with $p, q \in \mathbb{Z}^{\pm}$ and $p/q \neq 0, 1$. A systematic recipe of obtaining these new phases is to take the nontrivial qthroot of a propagator \hat{U} that describes Floquet topological matter. The general procedure is developed in Ref. [263], which generalizes the previous schemes of taking 2^n th and 3^n th roots for static Hamiltonians [283–292] to Floquet systems. The basic idea is first outlined in Ref. [293], and then expanded by utilizing \mathbb{Z}_q generalizations of Pauli matrices as ancillary degrees of freedom before taking the *q*th-root of a Floquet operator in an enlarged Hilbert space. It is in parallel with Dirac's original idea of adding internal degrees of freedom for electrons before taking the square-root to get their relativistic wave equation [294].

To be explicit, we consider the construction of a cubic-root non-Hermitian Floquet topological insulator. An illustration of the scheme is given in Fig. 7. Let us consider a three-step periodically quenched parent system, whose time-periodic Hamiltonian takes the form of [263]

$$\hat{H}(t) = \begin{cases} \hat{H}_3, & t \in [\ell, \ell + 1/3), \\ \hat{H}_2, & t \in [\ell + 1/3, \ell + 2/3), \\ \hat{H}_1, & t \in [\ell + 2/3, \ell + 1), \end{cases}$$
(156)

where $\ell \in \mathbb{Z}$ and we have assumed $\hbar = T = 1$. The Floquet operator of the system is then given by $\hat{U} = e^{-\frac{i}{3}\hat{H}_1}e^{-\frac{i}{3}\hat{H}_2}e^{-\frac{i}{3}\hat{H}_3}$. Following the methodology of Ref. [263], the nontrivial cubic-root $\hat{U}_{1/3}$ of \hat{U} is found to be

$$\hat{U}_{1/3} = \begin{pmatrix} 0 & e^{-i\frac{\hat{H}_1}{3}} & 0\\ 0 & 0 & e^{-i\frac{\hat{H}_2}{3}}\\ e^{-i\frac{\hat{H}_3}{3}} & 0 & 0 \end{pmatrix}.$$
(157)

The cube of $\hat{U}_{1/3}$ then gives

$$\hat{U}_{1/3}^{3} = \begin{pmatrix} e^{-\frac{i}{3}\hat{H}_{1}}e^{-\frac{i}{3}\hat{H}_{2}}e^{-\frac{i}{3}\hat{H}_{3}} & 0 & 0\\ 0 & e^{-\frac{i}{3}\hat{H}_{2}}e^{-\frac{i}{3}\hat{H}_{3}}e^{-\frac{i}{3}\hat{H}_{1}} & 0\\ 0 & 0 & e^{-\frac{i}{3}\hat{H}_{3}}e^{-\frac{i}{3}\hat{H}_{1}}e^{-\frac{i}{3}\hat{H}_{2}} \end{pmatrix}.$$
(158)

It is clear that $\hat{U}_{1/3}^3$ contains three identical copies of \hat{U} , in the sense that its three diagonal blocks are all related by similarity transformations and thus sharing the same spectrum and topological properties with \hat{U} [263].

As an example, we choose the Floquet model introduced in Ref. [261] as the parent




FIG. 7. Cubic-root non-Hermitian Floquet topological insulator: a schematic diagram of taking the cubit-root for a Floquet operator, (a) quasienergy spectrum under the PBC (in grey) and OBC (in blue), (b) gap functions under the PBC (in grey) and OBC (in red and blue), (c) Floquet topological edge modes with $E = \pi/3, 2\pi/3$, and (d)–(f) spatial profiles of Floquet edge modes and bulk skin modes [263].

system, whose Hamiltonian can now be expressed as

$$\hat{H}(t) = \begin{cases} \frac{3}{4}\hat{H}_1 & t \in [\ell, \ell + 1/3) \\ \frac{3}{2}\hat{H}_2 & t \in [\ell + 1/3, \ell + 2/3) , \\ \frac{3}{4}\hat{H}_1 & t \in [\ell + 2/3, \ell + 1) \end{cases}$$
(159)

where

$$\hat{H}_1 = J_2 \sum_n (i\hat{c}_{n+1}^{\dagger}\hat{c}_n + \text{H.c.}) \otimes \sigma_y + i\lambda \sum_n (\hat{c}_n^{\dagger}\hat{c}_{n+1} + \text{H.c.}) \otimes \sigma_y,$$
(160)

$$\hat{H}_2 = \sum_n (\mu \hat{c}_n^{\dagger} \hat{c}_n + J_1 \hat{c}_n^{\dagger} \hat{c}_{n+1} + \text{H.c.}) \otimes \sigma_x + i\lambda \sum_n (i \hat{c}_{n+1}^{\dagger} \hat{c}_n + \text{H.c.}) \otimes \sigma_x.$$
(161)

The cubit-root Floquet operator $\hat{U}_{1/3}$ of the system then takes the form of Eq. (157). The non-Hermitian effect is introduced by the asymmetric hopping amplitude $i\lambda$. The Floquet quasienergy spectrum and gap functions of $\hat{U}_{1/3}$ versus J_1 under the PBC (grey dots) and OBC (blue dots, red and blue lines) are shown in Figs. 7(a)–(b) [with $(J_2, \mu, \lambda) =$ $(0.5\pi, 0.4\pi, 0.25)$ and the length of lattice L = 400]. Here the gap function

$$F_{\varepsilon} \equiv \sqrt{(\operatorname{Re}E - \varepsilon)^2 + (\operatorname{Im}E)^2}, \qquad (162)$$

and the blue (red) lines in the Fig. 7(b) denote $F_{\pi/3}$ ($F_{2\pi/3}$). We observe a series of phase transitions accompanied by quasienergy-gap closings at $E = p\pi/q$ for p = 0, 1, 2, 3 and q = 3. After these transitions, more and more degenerate Floquet edge modes emerge at the fractional quasienergies $E = \pi/3$ and $2\pi/3$, which are impossible in conventional Floquet topological insulators with two quasienergy bands in one dimension. These fractional-quasienergy edge modes are thus unique to cubit-root Floquet topological insulators, either Hermitian or non-Hermitian. A zoom in of their quasienergies for one exemplary case is shown in Fig. 7(c) [with $(J_1, J_2, \mu, \lambda) = (\pi, 0.5\pi, 0.4\pi, 0.25)$ and the length of lattice L = 400], and their spatial profiles are shown in Figs. 7(d) [for the $E = \pi/3$ eigenmodes] and 7(e) [for the $E = 2\pi/3$ eigenmodes]. Moreover, the numbers of these fractional-quasienergy edge modes $(n_{\pi/3}, n_{2\pi/3})$ are determined by the open-bulk winding numbers (ν_0, ν_π) [261] of the parent system $\hat{U} = e^{-i\frac{\hat{H}_1}{4}}e^{-i\frac{\hat{H}_2}{2}}e^{-i\frac{\hat{H}_1}{4}}$. More precisely, we have the following bulk-edge correspondence for our cubic-root non-Hermitian Floquet topological insulator $\hat{U}_{1/3}$, i.e., [263]

$$n_0 = n_{2\pi/3} = 2|\nu_0|, \qquad n_{\pi/3} = n_{\pi} = 2|\nu_{\pi}|, \tag{163}$$

where n_0 and n_{π} are the numbers of zero and π degenerate Floquet edge modes. Finally, we notice that the system also possesses NHSE under the OBC, as reflected by the profiles of its Floquet bulk states in Fig. 7(f) [with $(J_1, J_2, \mu, \lambda) = (\pi, 0.5\pi, 0.4\pi, 0.25)$ and the length of lattice L = 400]. In Figs. 7(a) and 7(b), we also notice the discrepancy between the gapclosing points of the system under the PBC and OBC. Nevertheless, the open-bulk winding numbers (ν_0, ν_{π}) correctly capture the bulk-edge correspondence of the system under the OBC even with NHSEs. Therefore, the work conducted in Ref. [263] also established a dual topological characterization for *q*th-root non-Hermitian Floquet topological insulators. The formalism developed in Ref. [263] is equally applicable to the construction of *q*th-root Floquet topological insulators, superconductors, and semimetals in other (non-)Hermitian systems and across different physical dimensions. The fractional quasienergy edge modes at $E = p\pi/q$ might also be employed to generate boundary time crystals with different temporal periodicity and topological properties [98]. They may also find applications in the Floquet quantum computing.

C. Non-Hermitian Floquet topological superconductors

Similar to the static case, non-Hermitian Floquet topological phases could also appear in superconducting systems. We review one such example in this subsection, which admits many Floquet Majorana zero and π edge modes even with non-Hermitian effects.

The model we are going to consider describes a Floquet Kitaev chain, whose superconducting pairing terms are subject to time-periodic kicks. The Hamiltonian of the model takes the form of

$$\hat{H}(t) = \frac{1}{2} \sum_{n} [J(\hat{c}_{n}^{\dagger} \hat{c}_{n+1} + \text{H.c.}) + \Delta \delta_{T}(t)(\hat{c}_{n} \hat{c}_{n+1} + \text{H.c.}) + \mu(2\hat{c}_{n}^{\dagger} \hat{c}_{n} - 1)], \quad (164)$$

where J is the nearest-neighbor hopping amplitude, Δ is the p-wave superconducting pairing amplitude, and μ is the chemical potential. $\delta_T(t) \equiv \sum_{\ell \in \mathbb{Z}} \delta(t/T - \ell)$ implements delta kickings periodically with the period T. The system is made non-Hermitian by setting $J = J_r + iJ_i$ or $\mu = \mu_r + i\mu_i$ with $J_i \neq 0$ or $\mu_i \neq 0$, respectively. Possible experimental realizations of the delta kickings and non-Hermitian terms in this model are discussed in Ref. [257].

Under the PBC and in symmetric time frames, the Floquet operator of H(t) respects the chiral (sublattice) symmetry $S = \sigma_x$. Its bulk topological phases can then be characterized by the winding numbers (w_0, w_π) [see Eqs. (89)–(93)], for which we choose

$$\hat{H}_0 = \frac{1}{2} \sum_n [J(\hat{c}_n^{\dagger} \hat{c}_{n+1} + \text{H.c.}) + \mu(2\hat{c}_n^{\dagger} \hat{c}_n - 1)], \qquad (165)$$

$$\hat{H}_1 = \frac{1}{2} \sum_n \Delta(\hat{c}_n \hat{c}_{n+1} + \text{H.c.}), \qquad (166)$$

and assume $\hbar = T = 1$. Typical topological phase diagrams of the system are shown in Figs. 8(a) [for $J \in \mathbb{R}$, $\mu \in \mathbb{C}$, $(\mu_r, \Delta) = (0.3\pi, 0.5\pi)$] and 8(b) [for $\mu \in \mathbb{R}$, $J \in \mathbb{C}$, $(\mu, \Delta) = (0.4\pi, 0.9\pi)$]. Different topological phases correspond to different uniformly colored regions and the winding numbers (w_0, w_π) of each phase are labeled therein. The




FIG. 8. **Periodically kicked non-Hermitian Kitaev chain**: topological phase diagrams under the PBC [in (a) and (b)], Floquet spectra under the OBC [in (c) and (d)] and quasienergy gap functions under the OBC [in (e) and (f)]. The length of lattice is L = 1000 for (c)–(f). In (e) and (f), the solid and dashed lines at $F_0 = 0$ and $F_{\pi} = 0$ denote non-Hermitian Floquet Majorana zero and π edge modes [257].

boundaries between different phases can be analytically determined from the gap-closing conditions of the system (see Ref. [257]). Two notable features are brought about by the interplay between periodic drivings and non-Hermitian effects. First, the system could undergo rich topological phase transitions and enter Floquet superconducting phases with arbitrarily large topological winding numbers in principle. These phases could support arbitrarily many zero and π Majorana edge modes in the thermodynamic limit, which might be employed to engineer boundary time crystals and realize Floquet quantum computing protocols. Second, Floquet superconducting phases with larger winding numbers and therefore stronger topological signatures may appear with the increase of the non-Hermitian parameter. Therefore, new topological states could be induced and stabilized solely by non-Hermitian effects in Floquet superconducting systems. This is not expected in the non-driven counterparts of

μ_i [in Fig. 8(e)]	(w_0, w_π)	(n_0, n_π)	J_r [in Fig. 8(f)]	(w_0, w_π)	(n_0, n_π)
$(0, y_1)$	(4, 4)	(8,8)	$(0, y_1)$	(0, 0)	(0, 0)
(y_1, y_2)	(3,4)	(6,8)	(y_1,y_2)	(1, 0)	(2,0)
(y_2,y_3)	(3,3)	(6, 6)	(y_2,y_3)	(1, 1)	(2,2)
(y_3, y_4)	(3,2)	(6,4)	(y_3,y_4)	(1, 2)	(2, 4)
(y_4, y_5)	(2,2)	(4, 4)	(y_4,y_5)	(2, 2)	(4, 4)
(y_5,y_6)	(1,2)	(2,4)	(y_5,y_6)	(3, 2)	(6, 4)
(y_6, y_7)	(1,1)	(2,2)	(y_6,y_7)	(3,3)	(6, 6)
(y_7, y_8)	(1,0)	(2,0)	(y_7,y_8)	(3, 4)	(6, 8)
$(y_8, 6.5)$	(0, 0)	(0, 0)	$(y_8, 4\pi)$	(4, 4)	(8, 8)

TABLE I. **Periodically kicked non-Hermitian Kitaev chain**: bulk topological winding numbers (w_0, w_π) and numbers of Floquet Majorana zero and π edge modes (n_0, n_π) in different topological phases. The numerical values of phase transition points $y_1 \sim y_8$ for μ_i can be analytically found and are approximately given by 0.64, 1.03, 1.57, 1.94, 2.60, 3.15, 4.41, and 6.11. The numerical values of phase transition points $y_1 \sim y_8$ for J_r/π can also be found analytically and are given approximately by 0.42, 0.63, 1.47, 1.68, 2.51, 2.72, 3.56, and 3.77 [257].

our model, where nontrivial topological properties are usually destroyed with the growth of non-Hermitian effects.

Under the OBC, the quasienergy spectrum and edge states of the system can be obtained by diagonalizing the Floquet operator $\hat{U} = e^{-i\hat{H}_0}e^{-i\hat{H}_1}$ in Majorana or BdG bases [257]. As examples, the real parts of Floquet spectra and gap functions (F_0, F_π) versus μ_i [with $(J, \Delta, \mu_r) = (4\pi, 0.5\pi, 0.3\pi)$] and J_r [with $(J_i, \Delta, \mu) = (1, 0.4\pi, 0.9\pi)$] are shown in Figs. 8(c), 8(e), and Figs. 8(d), 8(f), respectively. $y_1 \sim y_8$ refer to gap closing points of the Floquet spectrum under the PBC (see Table I). The gap functions are defined as

$$F_0 = \frac{1}{\pi}\sqrt{(\operatorname{Re}\varepsilon)^2 + (\operatorname{Im}\varepsilon)^2}, \qquad F_\pi = \frac{1}{\pi}\sqrt{(|\operatorname{Re}\varepsilon| - \pi)^2 + (\operatorname{Im}\varepsilon)^2}, \tag{167}$$

where ε includes all the quasienergies of \hat{U} under the OBC. The eigenmodes with $F_0 = 0$ and $F_{\pi} = 0$ thus have the quasienergies zero and π , respectively. In Table I, we list the numbers of Floquet Majorana edge modes (n_0, n_{π}) with the quasienergies $(0, \pi)$ and the bulk topological winding numbers (w_0, w_{π}) . A simple bulk-edge correspondence can be further found between these numbers, i.e.,

$$n_0 = 2|w_0|, \qquad n_\pi = 2|w_\pi|.$$
 (168)

These relations hold in other parameter regions of our model as well. We conclude that our non-Hermitian Floquet Kitaev chain could indeed possess many Majorana zero and π edge

modes due to its large winding numbers. The localization nature of these Majorana modes was also demonstrated in Ref. [257].

Therefore, our study in Ref. [257] established the topological characterization and bulkedge correspondence of 1D non-Hermitian Floquet topological superconductors that belong to an extended BDI symmetry class. The collaboration between time-periodic driving fields and non-Hermitian effects was found to produce rich Floquet superconducting phases with large topological winding numbers and many Majorana edge modes at two distinct quasienergies zero and π . These non-Hermitian Floquet Majorana modes might allow Floquet quantum computing schemes to be more robust to environmental-induced nonreciprocity, dissipation, and quasiparticle poisoning effects. The existence of many pairs of Floquet Majorana modes may also create stronger transport signals at the ends of the chain, making it easier to experimentally detect their topological properties in open-system settings [226]. In future studies, topics like non-Hermitian Floquet topological superconductors under different driving protocols, in other symmetry classes, with more complicated lattice effects (e.g., sublattice structures, long-range hoppings or disorder), in higher spatial dimensions and with NHSEs, deserve to be explored. Possible changes in topological classifications due to many-body effects in non-Hermitian Floquet superconductors are also awaited to be revealed.

D. Non-Hermitian Floquet quasicrystals

In this subsection, we go beyond the clean limit of driven non-Hermitian lattices and showcase that the interplay among correlated disorder, temporal driving, and non-Hermitian effects could yield Floquet quasicrystals with rich PT-symmetry breaking transitions, localization transitions, and topological phase transitions. We will consider systems under both high-frequency and near-resonant driving fields.

We start with the example of a 1D non-Hermitian quasicrystal under high-frequency harmonic driving forces. The time-periodic Hamiltonian of the model takes the form of [262]

$$\hat{H}(t) = \sum_{n} \{ J(e^{-\gamma} \hat{c}_{n}^{\dagger} \hat{c}_{n+1} + e^{\gamma} \hat{c}_{n+1}^{\dagger} \hat{c}_{n}) + [V \cos(2\pi\alpha n) - nK \cos(\omega t)] \hat{c}_{n}^{\dagger} \hat{c}_{n} \}.$$
(169)




FIG. 9. Floquet NHQC under high-frequency driving forces: (a) a schematic diagram of the lattice model, (b) PT transitions of the Floquet spectrum, (c) localization transitions, and (d) topological transitions [262].

An illustration of the model is given in Fig. 9(a). Here $\gamma \in \mathbb{R}$ and $Je^{-\gamma} (Je^{\gamma})$ describes the right-to-left (left-to-right) nearest-neighbor hopping amplitude. The hopping is nonreciprocal and thus $\hat{H}(t) \neq \hat{H}^{\dagger}(t)$ if $\gamma \neq 0$. \hat{c}_{n}^{\dagger} (\hat{c}_{n}) creates (annihilates) a particle on the lattice site n. $V \in \mathbb{R}$ is the amplitude of an onsite potential, and α is chosen to be irrational in order for the potential to be spatially quasiperiodic. $K \in \mathbb{R}$ is the driving amplitude and ω is the driving frequency. This model can be viewed as a Floquet and quasicrystal variant of the Hatano-Nelson model [183]. In the absence of the driving force and under the PBC, all the eigenstates of the system are extended (localized) with complex (real) eigenvalues if $|V| < |2J|e^{|\gamma|}$ ($|V| > |2J|e^{|\gamma|}$) [189]. The system could thus undergo a PT transition, a localization-delocalization transition, and also a topological transition accompanied by the quantized change of its spectral winding number [Eq. (81)] at $|V| = |2J|e^{|\gamma|}$ [189].

Transforming the Hamiltonian $\hat{H}(t)$ in Eq. (169) to the rotating frame and applying the method discussed in Subsec. II C, we can obtain the Floquet effective Hamiltonian of our system in the high-frequency limit [262], i.e.,

$$\hat{H}_{\rm F} = \sum_{n} \left[\mathcal{J}_0\left(\frac{K}{\omega}\right) J(e^{-\gamma} \hat{c}_n^{\dagger} \hat{c}_{n+1} + e^{\gamma} \hat{c}_{n+1}^{\dagger} \hat{c}_n) + V \cos(2\pi\alpha n) \hat{c}_n^{\dagger} \hat{c}_n \right].$$
(170)

Here $\mathcal{J}_0(K/\omega)$ is the Bessel function of first kind, which is a non-monotonous function of K/ω . Under the PBC, the Floquet NHQC described by $\hat{H}_{\rm F}$ could thus undergo multiple and reentrant PT, localization and topological transitions with the change of the ratio K/ω between the driving amplitude and driving frequency whenever [262]

$$|V| = |2J\mathcal{J}_0(K/\omega)|e^{|\gamma|}.$$
(171)

This is indeed the case, as demonstrated by the maximum of the imaginary parts of quasienergies, the minimum of IPRs, and the winding numbers in Figs. 9(b)–(d) [for $(J, V, \alpha) = (1, 1, \frac{\sqrt{5}-1}{2})$ and the length of lattice L = 610]. In the metallic phase, we have max |ImE| > 0, min(IPR) $\rightarrow 0$, and the winding number w = -1, which means that all the Floquet eigenstates of $\hat{H}_{\rm F}$ are extended with complex quasienergies. In the insulating phase, we have max |ImE| = 0, min(IPR) > 0, and the winding number w = 0, implying that all the Floquet eigenstates of $\hat{H}_{\rm F}$ are localized with real quasienergies. The boundaries between these phases [red dashed lines in Figs. 9(b)–(d)] are precisely depicted by Eq. (171). The Lyapunov exponent $\lambda = \ln |(Ve^{-|\gamma|})/[2J\mathcal{J}_0(K/\omega)]|$ is quasienergy-independent for all the Floquet eigenstates, which is positive (negative and thus ill-defined) in the localized (extended) phases [262]. Therefore, it is clear that the phases and transitions in the quasienystal Hatano-Nelson model can be significantly modified by Floquet driving fields even in the high-frequency limit. The periodic driving also provides us with a flexible knob to control and engineer different types of phase transitions in NHQCs, with further examples discussed in Ref. [262].

We move on to the example of an NHQC under near-resonant driving fields. In this case, the interplay between driving and non-Hermitian effects not only induces multiple and reentrant localization transitions but also generates critical mobility edge phases that are




FIG. 10. Floquet NHQC under time-periodic kickings: (a) a schematic diagram of the lattice model, (b)–(f) PT transitions of the Floquet spectrum, localization transitions of Floquet states and critical phases with mobility edges of the system under the PBC. The initial state $|\psi(0)\rangle = \sum_{n} \delta_{n0} |0\rangle$ for the calculation of spreading speed v = v(t) is chosen to be localized at the center of the lattice, and the time average in Eq. (126) is taken over t = 1000 driving periods [264].

absent in non-driven limits. A schematic diagram of our model is shown in Fig. 10(a). Its Hamiltonian takes the form of $\hat{H}(t) = \hat{K}$ for $t \in [\ell T, \ell T + T_1)$ and $\hat{H}(t) = \hat{V}$ for $t \in [\ell T + T_1, \ell T + T_1 + T_2)$, where $\ell \in \mathbb{Z}$ and the driving period $T = T_1 + T_2$. The system is piecewisely quenched between $\hat{K} = J \sum (e^{\gamma} \hat{c}_n^{\dagger} \hat{c}_{n+1} + e^{-\gamma} \hat{c}_{n+1}^{\dagger} \hat{c}_n)$ and $\hat{V} = V \sum_n \cos(2\pi\alpha n) \hat{c}_n^{\dagger} \hat{c}_n$ over each driving period. For an irrational α , we thus arrive at a periodically quenched, spatially quasiperiodic variant of the Hatano-Nelson model, whose Floquet operator is given by [264]

$$\hat{U} = e^{-i\mathsf{V}\sum_{n}\cos(2\pi\alpha n)\hat{c}_{n}^{\dagger}\hat{c}_{n}}e^{-i\mathsf{J}\sum(e^{\gamma}\hat{c}_{n}^{\dagger}\hat{c}_{n+1} + e^{-\gamma}\hat{c}_{n+1}^{\dagger}\hat{c}_{n})},$$
(172)

where $V = VT_2/\hbar$ and $J = JT_1/\hbar$. Solving the eigenvalue equation $\hat{U}|\psi\rangle = e^{-iE}|\psi\rangle$ and using the tools introduced in Subsec. II G, we could obtain the maximal imaginary parts of quasienergies max |ImE|, the density of states with complex eigenvalues ρ , the minimum of IPRs and the spreading velocity of an initially localized wavepacket v. A collection of these quantities versus the strength of quasiperiodic potential V is shown in Fig. 10(b) for a typical case [with $(J, \gamma, \alpha) = (\pi/6, 0.8, \frac{\sqrt{5}-1}{2})$ and the length of lattice L = 4181]. We observe that with the increase of V from V = 0, the system first undergoes a complex-toreal PT transition in its quasienergy spectrum, which is also accompanied by a localization transition of all its Floquet eigenstates from spatially extended to localized. Interestingly, with the further enhancement of the quasiperiodic potential V (thus with stronger correlated disorder), some localized Floquet states can again become extended with real eigenvalues. The system then enters a critical phase, in which extended and localized eigenstates coexist and are separated by mobility edges on the complex quasienergy plane (see also the Fig. 3 of Ref. [264]). The further increase of V leads to reentrant transitions between localized and critical mobility edge phases in the system. This is also reflected by the two-parameter phase diagrams in Figs. 10(c)–(f) [see Subsec. II C for the definitions of ρ, \bar{g} , IPR_{min}, and ζ]. Note that both the critical phases and the reentrant localization transitions are absent in the non-driven Hatano-Nelson quasicrystal. They are brought about by the nearest-resonant Floquet driving field. It induces long-range spatial couplings and quasienergy windings in the system, thus yielding the observed phenomena (see Ref. [264] for more detailed discussions).

In experiments, Floquet NHQCs might be realized by ultracold atoms in driven and quasiperiodic optical superlattices with particle losses [180, 223]. Static NHQCs have also been realized by non-unitary photonic quantum walks [191, 192]. Signatures of PT breaking transitions, localization transitions, mobility edges, and topological properties related to NHSEs can be extracted from dynamical observables of initially localized wavepackets [191, 192]. Since quantum walk models are intrinsically dynamical, they provide natural platforms to realize and detect the Floquet NHQCs reviewed in this subsection. Meanwhile, as the discrete-time quantum walk carries a synthetic spin-half degree of freedom, it may also be utilized to simulate other types of NHQCs, such as those with lattice dimerizations or non-Abelian potentials [295–297]. Topological Anderson insulators induced by uncorrelated disorder could also be explored in similar settings [187].

Overall, we find that both high-frequency and near-resonant driving fields could be used to trigger, control, and enrich the phases and transitions in NHQCs, and even create unique Floquet NHQC phases that are absent in the static limit [262, 264]. Non-Hermitian disordered systems thus provide further playgrounds for the exploration of new physics that are enabled by Floquet engineering and time-periodic driving fields.

IV. CONCLUSION AND OUTLOOK

In this review, we recapitulated some progress we made in the study of intriguing topological phases in non-Hermitian Floquet systems. After a general introduction about the backgrounds and motivation, we first formulated the theoretical frameworks underpinning our study. These include the Floquet theory applicable in general situations and some approximation schemes suitable for exploring fast and slowly driven systems. Efficient means of characterizing the symmetry, topology, dynamical, and localization nature of non-Hermitian Floquet systems were then introduced. Equipped with these tools, we discussed prototypical examples of non-Hermitian Floquet topological phases in insulating, superconducting, and quasicrystalline systems, where the PT transitions, topological transitions, bulk-edge/corner correspondences and localization transitions within these non-Hermitian Floquet phases of matter were systematically characterized. Therefore, the collection of our works [254–264] lays a solid foundation for the study of topological phases that could emerge due to the interplay between periodic drivings and non-Hermitian effects.

We note that some other schemes concerning the topological classification and bulk-edge correspondence of non-Hermitian Floquet matter were discussed in Refs. [224, 228, 230, 236, 245]. The EPs and gapless topological phases in non-Hermitian Floquet systems were also considered in Refs. [215, 231, 234, 237, 239, 244, 247]. From the perspective of quantum control, the periodic driving fields may provide efficient means to stabilize non-Hermitian systems and control the PT transitions therein [210]. Some interesting aspects regarding the quantum dynamics and anomalous diffusion were revealed in several chaotic non-Hermitian Floquet systems [227, 233, 242, 248]. Aspects of PT-symmetry in non-Hermitian Floquet phases were also investigated in Refs. [209, 216, 217, 221, 232]. On the experimental side, the quantum walks in cold atoms, photonic setups, and circuit QED provide useful frameworks to realize and detect non-Hermitian Floquet topological matter [191, 192, 208, 211–214, 222, 223, 241].

Even with all the progress mentioned above, our knowledge about non-Hermitian Floquet topological matter is still quite limited. This is especially the case for systems beyond one spatial dimension, with uncorrelated disorders, and with many-body interactions. Dynamical quantum phase transitions may exhibit unique critical signatures triggered by the non-Hermitian Floquet exceptional topology [298–302]. The entanglement and transport properties of non-Hermitian Floquet systems [303] may also deviate significantly from their Hermitian or static counterparts. The exact connection between the topological phases of non-Hermitian Floquet Hamiltonians and the Floquet master equation (or Floquet Liouvillian) of driven open systems is unclear. More experimental efforts deserve to be made in order to realize and observe non-Hermitian Floquet phases with large topological invariants, many topological edge states, and multiple topological phase transitions. All these facts indicate that the area of non-Hermitian Floquet matter is still in its infancy, and further substantial developments are eagerly needed in this area.

ACKNOWLEDGMENTS

This research was funded by the National Natural Science Foundation of China (Grants Nos. 12275260, 11905211 and 12275155), the Fundamental Research Funds for the Central Universities (Grant No. 202364008), the Young Talents Project of Ocean University of China, and the Qilu Young Scholar Program of Shandong University.

- T. Dittrich, P. Hänggi, G.-L. Ingold, B. Kramer, G. Schön and W. Zwerger, <u>Quantum</u> Transport and Dissipation (Wiley-VCH, Weinheim, 1998).
- [2] H.-J. Stöckmann, Quantum Chaos: An Introduction (Cambridge University Press, UK, 1999).
- [3] D. J. Tannor, <u>Introduction to Quantum Mechanics: A time-dependent perspective</u> (University Science Books, USA, 2007).
- [4] C. J. Joachain, N. J. Kylstra, and R. M. Potvliege, <u>Atoms in Intense Laser Fields</u> (Cambridge University Press, UK, 2011).
- [5] F. Haake, S. Gnutzmann, and M. Kuś, <u>Quantum Signatures of Chaos</u> (Springer, Switzerland, 2018).
- [6] L. Reichl, The Transition to Chaos (Springer, Switzerland, 2021).
- [7] I. I. Rabi, Space Quantization in a Gyrating Magnetic Field, Phys. Rev. 51, 652 (1937).
- [8] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, <u>Atom-Photon Interactions: Basic</u> Processes and Applications (Wiley, New York, 1992).

- [9] R. Loudon, The Quantum Theory of Light, 3rd ed. (Oxford Science, Oxford, 2000).
- [10] J. Oreg, F. T. Hioe, and J. H. Eberly, Adiabatic following in multilevel systems, Phys. Rev. A 29, 690 (1984).
- [11] U. Gaubatz, P. Rudecki, S. Schiemann, and K. Bergmann, Population transfer between molecular vibrational levels by stimulated Raman scattering with partially overlapping laser fields. A new concept and experimental results, J. Chem. Phys. 92, 5363 (1990).
- [12] N. V. Vitanov, A. A. Rangelov, B. W. Shore, and K. Bergmann, Stimulated Raman adiabatic passage in physics, chemistry, and beyond, Rev. Mod. Phys. 89, 015006 (2017).
- [13] G. Casati and J. Ford, editors, <u>Stochastic Behaviour in Classical and Quantum Hamiltonian</u> Systems (Springer, Berlin, 1979).
- [14] S. Fishman, D. R. Grempel, and R. E. Prang, Chaos, Quantum Recurrences, and Anderson Localization, Phys. Rev. Lett. 49, 509 (1982).
- [15] F. L. Moore, J. C. Robinson, C. Bharucha, P. E. Williams and M. G. Raizen, Observation of Dynamical Localization in Atomic Momentum Transfer: A New Testing Ground for Quantum Chaos, Phys. Rev. Lett. 73, 2974 (1994).
- [16] D. J. Thouless, Quantization of particle transport, Phys. Rev. B 27, 6083 (1983).
- [17] Q. Niu and D. J. Thouless, Quantised adiabatic charge transport in the presence of substrate disorder and many-body interaction, J. Phys. A Math. Gen. 17, 2453-2462 (1984).
- [18] R. Citro and M. Aidelsburger, Thouless pumping and topology, Nat. Rev. Phys. 5, 87-101 (2023).
- [19] V. Khemani, A. Lazarides, R. Moessner, and S. L. Sondhi, Phase Structure of Driven Quantum Systems, Phys. Rev. Lett. 116, 250401 (2016).
- [20] D. V. Else, B. Bauer, and C. Nayak, Floquet Time Crystals, Phys. Rev. Lett. 117, 090402 (2016).
- [21] N. Y. Yao, A. C. Potter, I.-D. Potirniche, and A. Vishwanath, Discrete Time Crystals: Rigidity, Criticality, and Realizations, Phys. Rev. Lett. **118**, 030401 (2017).
- [22] T. Oka and H. Aoki, Photovoltaic Hall effect in graphene. Phys. Rev. B 79, 081406(R) (2009).
- [23] N. H. Lindner, G. Refael, and V. Galitski, Floquet topological insulator in semiconductor quantum wells, Nat. Phys. 7, 490-495 (2011).
- [24] D. Y. H. Ho and J. Gong, Quantized Adiabatic Transport In Momentum Space, Phys. Rev. Lett. 109, 010601 (2012).

- [25] P. Leboeuf, J. Kurchan, M. Feingold, and D. P. Arovas, Topological aspects of quantum chaos, Chaos 2, 125 (1992).
- [26] Y. Chen and C. Tian, Planck's Quantum-Driven Integer Quantum Hall Effect in Chaos, Phys. Rev. Lett. 113, 216802 (2014).
- [27] C. Tian, Y. Chen, and J. Wang, Emergence of integer quantum Hall effect from chaos, Phys. Rev. B 93, 075403 (2016).
- [28] G. Casati and L. Molinari, "Quantum Chaos" with Time-Periodic Hamiltonians, Progress of Theoretical Physics Supplement 98, 287-322 (1989).
- [29] M. Grifoni and P. Hänggi, Driven quantum tunneling, Phys. Rep. 304, 229-354 (1998).
- [30] S. Kohler, J. Lehmann, and P. Hänggi, Driven quantum transport on the nanoscale, Phys. Rep. 406, 379-443 (2005).
- [31] T. Kitagawa, Topological phenomena in quantum walks: elementary introduction to the physics of topological phases, Quantum Inf. Process. 11, 1107-1148 (2012).
- [32] J. Cayssol, B. Dóra, F. Simon, and R. Moessner, Floquet topological insulators, Phys. Status Solidi RRL 7, 101-108 (2013).
- [33] M. Bukov, L. D'Alessio, and A. Polkovnikov, Universal high-frequency behavior of periodically driven systems: from dynamical stabilization to Floquet engineering, Adv. Phys. 64, 139-226 (2015).
- [34] M. Holthaus, Floquet engineering with quasienergy bands of periodically driven optical lattices, J. Phys. B: At. Mol. Opt. Phys. 49, 013001 (2016).
- [35] A. Eckardt, Colloquium: Atomic quantum gases in periodically driven optical lattices, Rev. Mod. Phys. 89, 011004 (2017).
- [36] P. Weinberg, M. Bukov, L. D'Alessio, A. Polkovnikov, S. Vajna, and M. Kolodrubetz, Adiabatic perturbation theory and geometry of periodically-driven systems, Phys. Rep. 688, 1-35 (2017).
- [37] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: a theoretical overview, J. Phys. B: At. Mol. Opt. Phys. 51, 112001 (2018).
- [38] T. Oka and S. Kitamura, Floquet Engineering of Quantum Materials, Annu. Rev. Condens. Matter Phys. 10, 387 (2019).

- [39] M. S. Rudner and N. H. Lindner, Band structure engineering and non-equilibrium dynamics in Floquet topological insulators, Nat. Rev. Phys. 2, 229-244 (2020).
- [40] M. S. Rudner and N. H. Lindner, The Floquet Engineer's Handbook, arXiv:2003.08252 (2020).
- [41] F. Harper, R. Roy, M. S. Rudner, and S. L. Sondhi, Topology and Broken Symmetry in Floquet Systems, Annu. Rev. Condens. Matter Phys. 11, 345 (2020).
- [42] A. de la Torre, D. M. Kennes, M. Claassen, S. Gerber, J. W. McIver, and M. A. Sentef, Colloquium: Nonthermal pathways to ultrafast control in quantum materials, Rev. Mod. Phys. 93, 041002 (2021).
- [43] C. Weitenberg and J. Simonet, Tailoring quantum gases by Floquet engineering, Nat. Phys. 17, 1342-1348 (2021).
- [44] S. Bandyopadhyay, S. Bhattacharjee and D. Sen, Driven quantum many-body systems and out-of-equilibrium topology, J. Phys.: Condens. Matter 33, 393001 (2021).
- [45] A. Sen, D. Sen and K. Sengupta, Analytic approaches to periodically driven closed quantum systems: methods and applications, J. Phys.: Condens. Matter 33, 443003 (2021).
- [46] M. Rodriguez-Vega, M. Vogl, and G. A. Fiete, Low-frequency and Moiré-Floquet engineering: A review, Annals of Physics 435, 168434 (2021).
- [47] J. Bloch, A. Cavalleri, V. Galitski, M. Hafezi, and A. Rubio, Strongly correlated electronphoton systems, Nature 606, 41-48 (2022).
- [48] W. Zhu, J. Gong, and R. W. Bomantara, Topological π modes and beyond, Science Bulletin
 67, 2145-2148 (2022).
- [49] E. Galiffi, R. Tirole, S. Yin, H. Li, S. Vezzoli, P. A. Huidobro, M. G. Silveirinha, R. Sapienza, A. Alù, and J. B. Pendry, Photonics of time-varying media, Advanced Photonics 4, 014002 (2022).
- [50] M. Rodriguez-Vega, M. G. Vergniory, and G. A. Fiete, Quantum materials out of equilibrium, Physics Today 75, 42 (2022).
- [51] S. Yin, E. Galiffi, and A. Alù, Floquet metamaterials, eLight 2, 8 (2022).
- [52] C. Bao, P. Tang, D. Sun, and S. Zhou, Light-induced emergent phenomena in 2D materials and topological materials, Nat. Rev. Phys. 4, 33-48 (2022).
- [53] A. Haldar and A. Das, Statistical mechanics of Floquet quantum matter: exact and emergent conservation laws, J. Phys.: Condens. Matter 34, 234001 (2022).

- [54] T. Mori, Floquet States in Open Quantum Systems, Annu. Rev. Condens. Matter Phys. 14, 35-56 (2023).
- [55] W. W. Ho, T. Mori, D. A. Abanin, and E. G. Dalla Torre, Quantum and classical Floquet prethermalization, https://doi.org/10.1016/j.aop.2023.169297 (2023).
- [56] N. Tsuji, Floquet States, arXiv:2301.12676 (2023).
- [57] K. Sacha and J. Zakrzewski, Time crystals: a review, Rep. Prog. Phys. 81, 016401 (2018).
- [58] V. Khemani, R. Moessner, and S. L. Sondhi, A Brief History of Time Crystals, arXiv:1910.10745 (2019).
- [59] D. V. Else, C. Monroe, C. Nayak, and N. Y. Yao, Discrete Time Crystals, Annu. Rev. Condens. Matter Phys. 11, 467 (2020).
- [60] L. Guo and P. Liang, Condensed matter physics in time crystals, New J. Phys. 22, 075003 (2020).
- [61] S. Rahav, I. Gilary, and S. Fishman, Effective Hamiltonians for Periodically Driven Systems, Phys. Rev. A 68, 013820 (2003).
- [62] N. Goldman and J. Dalibard, Periodically Driven Quantum Systems: Effective Hamiltonians and Engineered Gauge Fields, Phys. Rev. X 4, 031027 (2014).
- [63] A. Eckardt and E. Anisimovas, High-frequency approximation for periodically driven quantum systems from a Floquet-space perspective, New J. Phys. 17, 093039 (2015).
- [64] T. Mikami, S. Kitamura, K. Yasuda, N. Tsuji, T. Oka, and H. Aoki, Brillouin-Wigner theory for high-frequency expansion in periodically driven systems: Application to Floquet topological insulators, Phys. Rev. B 93, 144307 (2016).
- [65] L. Zhou, C. Chen, and J. Gong, Floquet semimetal with Floquet-band holonomy, Phys. Rev. B 94, 075443 (2016).
- [66] L. K. Upreti, C. Evain, S. Randoux, P. Suret, A. Amo, and P. Delplace, Topological Swing of Bloch Oscillations in Quantum Walks, Phys. Rev. Lett. 125, 186804 (2020).
- [67] A. F. Adiyatullin, L. K. Upreti, C. Lechevalier, C. Evain, F. Copie, P. Suret, S. Randoux, P. Delplace, and A. Amo, Topological Properties of Floquet Winding Bands in a Photonic Lattice, Phys. Rev. Lett. 130, 056901 (2023).
- [68] L. Jiang, T. Kitagawa, J. Alicea, A. R. Akhmerov, D. Pekker, G. Refael, J. I. Cirac, E. Demler, M. D. Lukin, and P. Zoller, Majorana Fermions in Equilibrium and in Driven Cold-Atom Quantum Wires, Phys. Rev. Lett. 106, 220402 (2011).

- [69] J. K. Asbóth and H. Obuse, Bulk-boundary correspondence for chiral symmetric quantum walks, Phys. Rev. B 88, 121406(R) (2013).
- [70] D. Y. H. Ho and J. Gong, Topological effects in chiral symmetric driven systems, Phys. Rev. B 90, 195419 (2014).
- [71] Q. Cheng, Y. Pan, H. Wang, C. Zhang, D. Yu, A. Gover, H. Zhang, T. Li, L. Zhou, and S. Zhu, Observation of Anomalous π Modes in Photonic Floquet Engineering, Phys. Rev. Lett. 122, 173901 (2019).
- [72] M. S. Rudner, N. H. Lindner, E. Berg, and M. Levin, Anomalous Edge States and the Bulk-Edge Correspondence for Periodically Driven Two-Dimensional Systems, Phys. Rev. X 3, 031005 (2013).
- [73] M. Lababidi, I. I. Satija, and E. Zhao, Counter-propagating Edge Modes and Topological Phases of a Kicked Quantum Hall System, Phys. Rev. Lett. **112**, 026805 (2014).
- [74] P. Titum, E. Berg, M. S. Rudner, G. Refael, and N. H. Lindner, Anomalous Floquet-Anderson Insulator as a Nonadiabatic Quantized Charge Pump, Phys. Rev. X 6, 021013 (2016).
- [75] Q.-J. Tong, J.-H. An, J. Gong, H.-G. Luo, and C. H. Oh, Generating many Majorana modes via periodic driving: A superconductor model, Phys. Rev. B 87, 201109(R) (2013).
- [76] L. Zhou, H. Wang, D. Y. H. Ho, and J. Gong, Aspects of Floquet bands and topological phase transitions in a continuously driven superlattice, Eur. Phys. J. B 87, 204 (2014).
- [77] T. Xiong, J. Gong, and J. An, Towards large-Chern-number topological phases by periodic quenching, Phys. Rev. B 93, 184306 (2016).
- [78] L. Zhou and J. Gong, Floquet topological phases in a spin-1/2 double kicked rotor, Phys. Rev. A 97, 063603 (2018).
- [79] L. Zhou and J. Gong, Recipe for creating an arbitrary number of Floquet chiral edge states, Phys. Rev. B 97, 245430 (2018).
- [80] R. W. Bomantara, L. Zhou, J. Pan, and J. Gong, Coupled-wire construction of static and Floquet second-order topological insulators, Phys. Rev. B 99, 045441 (2019).
- [81] L. Zhou and Q. Du, Floquet topological phases with fourfold-degenerate edge modes in a driven spin-1/2 Creutz ladder, Phys. Rev. A 101, 033607 (2020).
- [82] F. Nathan and M. S. Rudner, Topological singularities and the general classification of Floquet-Bloch systems, New J. Phys. 17, 125014 (2015).

- [83] A. C. Potter, T. Morimoto, and A. Vishwanath, Classification of Interacting Topological Floquet Phases in One Dimension, Phys. Rev. X 6, 041001 (2016).
- [84] R. Roy and F. Harper, Periodic table for Floquet topological insulators, Phys. Rev. B 96, 155118 (2017).
- [85] S. Yao, Z. Yan, and Z. Wang, Topological invariants of Floquet systems: General formulation, special properties, and Floquet topological defects, Phys. Rev. B 96, 195303 (2017).
- [86] T. Kitagawa, M. A. Broome, A. Fedrizzi, M. S. Rudner, E. Berg, I. Kassal, A. Aspuru-Guzik, E. Demler, and A. G. White, Observation of topologically protected bound states in photonic quantum walks, Nat. Commun. 3, 882 (2012).
- [87] M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, Photonic Floquet topological insulators, Nature (London) 496, 196 (2013).
- [88] Y. Wang, H. Steinberg, P. Jarillo-Herrero, and N. Gedik, Observation of Floquet-Bloch states on the surface of a topological insulator, Science 342, 453 (2013).
- [89] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, Experimental realization of the topological Haldane model with ultracold fermions, Nature (London) 515, 237 (2014).
- [90] W. Hu, J. C. Pillay, K. Wu, M. Pasek, P. P. Shum, and Y. D. Chong, Measurement of a Topological Edge Invariant in a Microwave Network, Phys. Rev. X 5, 011012 (2015).
- [91] L. J. Maczewsky, J. M. Zeuner, S. Nolte, and A. Szameit, Observation of photonic anomalous Floquet topological insulators, Nat. Commun. 8, 13756 (2017).
- [92] L. Asteria, D. T. Tran, T. Ozawa, M. Tarnowski, B. S. Rem, N. FlÀschner, K. Sengstock, N. Goldman, and C. Weitenberg, Measuring quantized circular dichroism in ultracold topological matter, Nat. Phys. 15, 449 (2019).
- [93] K. Wintersperger, C. Braun, F. N. Ünal, A. Eckardt, M. D. Liberto, N. Goldman, I. Bloch, and M. Aidelsburger, Realization of an anomalous Floquet topological system with ultracold atoms, Nat. Phys. 16, 1058 (2020).
- [94] J. W. McIver, B. Schulte, F.-U. Stein, T. Matsuyama, G. Jotzu, G. Meier, and A. Cavalleri, Light-induced anomalous Hall effect in graphene, Nat. Phys. 16, 38 (2020).
- [95] S. Afzal, T. J. Zimmerling, Y. Ren, D. Perron, and V. Van, Realization of Anomalous Floquet Insulators in Strongly Coupled Nanophotonic Lattices, Phys. Rev. Lett. 124, 253601 (2020).

- [96] B. Chen, S. Li, X. Hou, F. Ge, F. Zhou, P. Qian, F. Mei, S. Jia, N. Xu, and H. Shen, Digital quantum simulation of Floquet topological phases with a solid-state quantum simulator, Photon. Res. 9, 81 (2021).
- [97] K. Yang, S. Xu, L. Zhou, Z. Zhao, T. Xie, Z. Ding, W. Ma, J. Gong, F. Shi, and J. Du, Observation of Floquet topological phases with large Chern numbers, Phys. Rev. B 106, 184106 (2022).
- [98] Z. Cheng, R. W. Bomantara, H. Xue, W. Zhu, J. Gong, and B. Zhang, Observation of π/2 Modes in an Acoustic Floquet System, Phys. Rev. Lett. **129**, 254301 (2022).
- [99] W. Zhu, H. Xue, J. Gong, Y. Chong, and B. Zhang, Time-periodic corner states from Floquet higher-order topology, Nat. Commun. 13, 11 (2022).
- S. Zhou, C. Bao, B. Fan, H. Zhou, Q. Gao, H. Zhong, T. Lin, H. Liu, P. Yu, P. Tang, S. Meng,
 W. Duan, and S. Zhou, Pseudospin-selective Floquet band engineering in black phosphorus,
 Nature (London) 614, 75 (2023).
- [101] J.-Y. Zhang, C.-R. Yi, L. Zhang, R.-H. Jiao, K.-Y. Shi, H. Yuan, W. Zhang, X.-J. Liu, S. Chen, and J.-W. Pan, Tuning Anomalous Floquet Topological Bands with Ultracold Atoms, Phys. Rev. Lett. 130, 043201 (2023).
- [102] R. W. Bomantara and J. B. Gong, Simulation of non-Abelian braiding in Majorana time crystals, Phys. Rev. Lett. 120, 230405 (2018).
- [103] R. W. Bomantara and J. B. Gong, Quantum computation via Floquet topological edge modes, Phys. Rev. B 98, 165421 (2018).
- [104] R. W. Bomantara and J. Gong, Measurement-only quantum computation with Floquet Majorana corner modes, Phys. Rev. B 101, 085401 (2020).
- [105] T. Kato, <u>Perturbation Theory for Linear Operators</u> (Springer, Berlin, 1976).
- [106] P. Ring and P. Schuck, The Nuclear Many Body Problem (Springer, New York, 1980).
- [107] N. Moiseyev, <u>Non-Hermitian Quantum Mechanics</u> (Cambridge University Press, Cambridge, 2011).
- [108] C. M. Bender, <u>PT Symmetry in Quantum and Classical Physics</u> (World Scientific, Singapore, 2019).
- [109] M. V. Berry, Physics of Nonhermitian Degeneracies, Czechoslovak Journal of Physics 54, 1039-1047 (2004).
- [110] W. D. Heiss, The physics of exceptional points, J. Phys. A: Math. Theor. 45, 444016 (2012).

- [111] V. V. Konotop, J. Yang, and D. A. Zezyulin, Nonlinear waves in PT-symmetric systems, Rev. Mod. Phys. 88, 035002 (2016).
- [112] S. V. Suchkov, A. A. Sukhorukov, J. Huang, S. V. Dmitriev, C. Lee, and Y. S. Kivshar, Nonlinear switching and solitons in PT-symmetric photonic systems, Laser Photonics Rev. 10, 177-213 (2016).
- [113] H. Schomerus, From scattering theory to complex wave dynamics in non-Hermitian PTsymmetric resonators, Phil. Trans. R. Soc. A 371, 20120194 (2016).
- [114] V. M. Martinez Alvarez, J. E. Barrios Vargas, M. Berdakin, and L.E.F. Foa Torres, Topological states of non-Hermitian systems, Eur. Phys. J. Spec. Top. 227, 1295-1308 (2018).
- [115] R. El-Ganainy, K. G. Makris, M. Khajavikhan, Z. H. Musslimani, S. Rotter and D. N. Christodoulides, Non-Hermitian physics and PT symmetry, Nat. Phys. 14, 11-19 (2018).
- [116] H. Zhao and L. Feng, Parity-time symmetric photonics, Natl. Sci. Rev. 5, 183-199 (2018).
- [117] M.-A. Miri and A. Alù, Exceptional points in optics and photonics, Science **363**, 42 (2019).
- [118] A. Ghatak and T. Das, New topological invariants in non-Hermitian systems, J. Phys.: Condens. Matter **31**, 263001 (2019).
- [119] S. K. Özdemir, S. Rotter, F. Nori, and L. Yang, Parity-time symmetry and exceptional points in photonics, Nat. Mater. 18, 783-798 (2019).
- [120] Y. Ashida, Z. Gong and M. Ueda, Non-Hermitian physics, Adv. Phys. 69, 249-435 (2020).
- [121] K. Yokomizo and S. Murakami, Non-Bloch band theory and bulk-edge correspondence in non-Hermitian systems, Prog. Theor. Exp. Phys. 2020, 12A102 (2020).
- [122] H. Kondo, Y. Akagi, and H. Katsura, Non-Hermiticity and topological invariants of magnon Bogoliubov-de Gennes systems, Prog. Theor. Exp. Phys. 2020, 12A104 (2020).
- [123] T. Yoshida, R. Peters, N. Kawakami, and Y. Hatsugai, Exceptional band touching for strongly correlated systems in equilibrium, Prog. Theor. Exp. Phys. 2020, 12A109 (2020).
- [124] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, Exceptional topology of non-Hermitian systems, Rev. Mod. Phys. 93, 015005 (2021).
- [125] C. Coulais, R. Fleury, and J. van Wezel, Topology and broken Hermiticity, Nat. Phys. 17, 9-13 (2021).
- [126] K. Ding, C. Fang, and G. Ma, Non-Hermitian topology and exceptional-point geometries, Nat. Rev. Phys. 4, 745-760 (2022).

- [127] X. Zhanga, T. Zhanga, M.-H. Lu, and Y.-F. Chen, A review on non-Hermitian skin effect, Advances in Physics: X 7, 2109431 (2022).
- [128] H. M. Hurst and B. Flebus, Non-Hermitian physics in magnetic systems, J. Appl. Phys. 132, 220902 (2022).
- [129] A. Banerjee, R. Sarkar, S. Dey, and A. Narayan, Non-Hermitian Topological Phases: Principles and Prospects, J. Phys.: Condens. Matter 35, 333001 (2023).
- [130] N. Okuma and M. Sato, Non-Hermitian Topological Phenomena: A Review, Annu. Rev. Condens. Matter Phys. 14, 83-107 (2023).
- [131] Q. Yan, B. Zhao, R. Zhou, R. Ma, Q. Lyu, S. Chu, X. Hu and Q. Gong, Advances and applications on non-Hermitian topological photonics, Nanophotonics, https://doi.org/10.1515/nanoph-2022-0775 (2023).
- [132] R. Lin, T. Tai, L. Li, and C. H. Lee, Topological Non-Hermitian skin effect, Front. Phys. 18, 53605 (2023).
- [133] V. Medeny, L. Grunwald, and D. M. Kennes, PT-symmetric, non-Hermitian quantum manybody physics – a methodological perspective, arXiv:2303.05956 (2023).
- [134] C. M. Bender, D. C. Brody, and H. F. Jones, Complex Extension of Quantum Mechanics, Phys. Rev. Lett. 89, 270401 (2002).
- [135] D.-J. Zhang, Q.-h. Wang, and J. Gong, Time-dependent PT-symmetric quantum mechanics in generic non-Hermitian systems, Phys. Rev. A 100, 062121 (2019).
- [136] C. E. Rüter, K. G. Makris, R. El-Ganainy, D. N. Christodoulides, M. Segev, and D. Kip, Observation of parity-time symmetry in optics, Nat. Phys. 6, 192-195 (2010).
- [137] Y. Wu, W. Liu, J. Geng, X. Song, X. Ye, C.-K. Duan, X. Rong, and J. Du, Observation of parity-time symmetry breaking in a single-spin system, Science 364, 878-880 (2019).
- [138] S. Xia, D. Kaltsas, D. Song, I. Komis, J. Xu, A. Szameit, H. Buljan, K. G. Makris, Z. Chen, Nonlinear tuning of PT symmetry and non-Hermitian topological states, Science **372**, 72-76 (2021).
- [139] H. Shen, B. Zhen, and L. Fu, Topological Band Theory for Non-Hermitian Hamiltonians, Phys. Rev. Lett. 120, 146402 (2018).
- [140] Z. Gong, Y. Ashida, K. Kawabata, K. Takasan, S. Higashikawa, and M. Ueda, Topological Phases of Non-Hermitian Systems, Phys. Rev. X 8, 031079 (2018).

- [141] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, Symmetry and Topology in Non-Hermitian Physics, Phys. Rev. X 9, 041015 (2019).
- [142] H. Zhou and J. Y. Lee, Periodic table for topological bands with non-Hermitian symmetries, Phys. Rev. B 99, 235112 (2019).
- [143] C.-H. Liu, H. Jiang, and S. Chen, Topological classification of non-Hermitian systems with reflection symmetry, Phys. Rev. B 99, 125103 (2019).
- [144] C. C. Wojcik, X. Sun, T. Bzdušek, and S. Fan, Homotopy characterization of non-Hermitian Hamiltonians, Phys. Rev. B 101, 205417 (2020).
- [145] A. Altland, M. Fleischhauer, and S. Diehl, Symmetry Classes of Open Fermionic Quantum Matter, Phys. Rev. X 11, 021037 (2021).
- [146] A. M. García-García, L. Sá, and J. J. M. Verbaarschot, Symmetry Classification and Universality in Non-Hermitian Many-Body Quantum Chaos by the Sachdev-Ye-Kitaev Model, Phys. Rev. X 12, 021040 (2022).
- [147] C. C. Wojcik, K. Wang, A. Dutt, J. Zhong, and S. Fan, Eigenvalue topology of non-Hermitian band structures in two and three dimensions, Phys. Rev. B 106, L161401 (2022).
- [148] Y. Xu, S.-T. Wang, and L.-M. Duan, Weyl Exceptional Rings in a Three-Dimensional Dissipative Cold Atomic Gas, Phys. Rev. Lett. 118, 045701 (2017).
- [149] J. Doppler, A. A. Mailybaev, J. Bohm, U. Kuhl, A. Girschik, F. Libisch, T. J. Milburn, P. Rabl, N. Moiseyev, and S. Rotter, Dynamically encircling an exceptional point for asymmetric mode switching, Nature 537, 76-79 (2016).
- [150] H. Xu, D. Mason, L. Jiang, and J. G. E. Harris, Topological energy transfer in an optomechanical system with exceptional points, Nature 537, 80-83 (2016).
- [151] H. Nasari, G. Lopez-Galmiche, H. E. Lopez-Aviles, A. Schumer, A. U. Hassan, Q. Zhong, S. Rotter, P. LiKamWa, D. N. Christodoulides, and M. Khajavikhan, Observation of chiral state transfer without encircling an exceptional point, Nature 605, 256-261 (2022).
- [152] H. Hodaei, A. U. Hassan, S. Wittek, H. Garcia-Gracia1, R. El-Ganainy, D. N. Christodoulides, and M. Khajavikhan, Enhanced sensitivity at higher-order exceptional points, Nature 548, 187-191 (2017).
- [153] W. Chen, Ş. K. Özdemir, G. Zhao, and L. Yang, Exceptional points enhance sensing in an optical microcavity, Nature 548, 192-196 (2017).

- [154] M. P. Hokmabadi, A. Schumer, D. N. Christodoulides, and M. Khajavikhan, Non-Hermitian ring laser gyroscopes with enhanced Sagnac sensitivity, Nature 576, 70-74 (2019).
- [155] R. Kononchuk, J. Cai, F. Ellis, R. Thevamaran, T. Kottos, Exceptional-point-based accelerometers with enhanced signal-to-noise ratio, Nature 607, 697-702 (2022).
- [156] L. Feng, Z. J. Wong, R. M. Ma, Y. Wang, and X. Zhang, Single-mode laser by parity-time symmetry breaking, Science 346, 972 (2014).
- [157] H. Hodaei, M. A. Miri, M. Heinrich, D. N. Christodoulides, and M. Khajavikhan, Paritytime-symmetric microring lasers, Science 346, 975 (2014).
- [158] M. A. Bandres, S. Wittek, G. Harari, M. Parto, J. Ren, M. Segev, D. N. Christodoulides, and M. Khajavikhan, Topological insulator laser: Experiments, Science 359, 1231 (2018).
- [159] T. E. Lee, Anomalous Edge State in a Non-Hermitian Lattice, Phys. Rev. Lett. 116, 133903 (2016).
- [160] Y. Xiong, Why does bulk boundary correspondence fail in some non-hermitian topological models, J. Phys. Commun. 2, 35043 (2018).
- [161] V. M. M. Alvarez, J. E. B. Vargas, and L. E. F. Foa Torres, Non-Hermitian robust edge states in one dimension: Anomalous localization and eigenspace condensation at exceptional points, Phys. Rev. B 97, 121401(R) (2018).
- [162] F. K. Kunst, E. Edvardsson, J. C. Budich, and E. J. Bergholtz, Biorthogonal Bulk-Boundary Correspondence in Non-Hermitian Systems, Phys. Rev. Lett. **121**, 026808 (2018).
- [163] S. Yao and Z. Wang, Edge States and Topological Invariants of Non-Hermitian Systems, Phys. Rev. Lett. 121, 086803 (2018).
- [164] C. H. Lee and R. Thomale, Anatomy of skin modes and topology in non-Hermitian systems, Phys. Rev. B 99, 201103(R) (2019).
- [165] K. Yokomizo and S. Murakami, Non-Bloch Band Theory of Non-Hermitian Systems, Phys. Rev. Lett. 123, 066404 (2019).
- [166] F. Song, S. Yao, and Z. Wang, Non-Hermitian Skin Effect and Chiral Damping in Open Quantum Systems, Phys. Rev. Lett. 123, 170401 (2019).
- [167] F. K. Kunst and V. Dwivedi, Non-Hermitian systems and topology: A transfer-matrix perspective, Phys. Rev. B 99, 245116 (2019).
- [168] F. Song, S. Yao, and Z. Wang, Non-Hermitian Topological Invariants in Real Space, Phys. Rev. Lett. 123, 246801 (2019).

- [169] C. H. Lee, L. Li, and J. Gong, Hybrid Higher-Order Skin-Topological Modes in Nonreciprocal Systems, Phys. Rev. Lett. 123, 016805 (2019).
- [170] D. S. Borgnia, A. J. Kruchkov, and R.-J. Slager, Non-Hermitian Boundary Modes and Topology, Phys. Rev. Lett. 124, 056802 (2020).
- [171] N. Okuma, K. Kawabata, K. Shiozaki, and M. Sato, Topological Origin of Non-Hermitian Skin Effects, Phys. Rev. Lett. 124, 086801 (2020).
- [172] K. Zhang, Z. Yang, and C. Fang, Correspondence between Winding Numbers and Skin Modes in Non-Hermitian Systems, Phys. Rev. Lett. 125, 126402 (2020).
- [173] M. Lu, X. Zhang, and M. Franz, Magnetic Suppression of Non-Hermitian Skin Effects Phys. Rev. Lett. 127, 256402 (2021).
- [174] K. Zhang, Z. Yang, and C. Fang, Universal non-Hermitian skin effect in two and higher dimensions, Nat. Commun. 13, 2496 (2022).
- [175] T. Helbig, T. Hofmann, S. Imhof, M. Abdelghany, T. Kiessling, L. W. Molenkamp, C. H. Lee, A. Szameit, M. Greiter, and R. Thomale, Generalized bulk-boundary correspondence in non-Hermitian topolectrical circuits, Nat. Phys. 16, 747-750 (2020).
- [176] L. Xiao, T. Deng, K. Wang, G. Zhu, Z. Wang, W. Yi, and P. Xue, Non-Hermitian bulkboundary correspondence in quantum dynamics, Nat. Phys. 16, 761-766 (2020).
- [177] A. Ghatak, M. Brandenbourger, J. van Wezel, and C. Coulais, Observation of non-Hermitian topology and its bulk-edge correspondence in an active mechanical metamaterial, PNAS 117, 29561 (2020).
- [178] X. Zhang, Y. Tian, J. Jiang, M. Lu, and Y. Chen, Observation of higher-order non-Hermitian skin effect, Nat. Commun. 12, 5377 (2021).
- [179] D. Zou, T. Chen, W. He, J. Bao, C. H. Lee, H. Sun, and X. Zhang, Observation of hybrid higher-order skin-topological effect in non-Hermitian topolectrical circuits, Nat. Commun. 12, 7201 (2021).
- [180] Q. Liang, D. Xie, Z. Dong, H. Li, H. Li, B. Gadway, W. Yi, and B. Yan, Dynamic Signatures of Non-Hermitian Skin Effect and Topology in Ultracold Atoms, Phys. Rev. Lett. **129**, 070401 (2022).
- [181] Z. Gu, H. Gao, H. Xue, J. Li, Z. Su, and J. Zhu, Transient non-Hermitian skin effect, Nat. Commun. 13, 7668 (2022).

- [182] K. Kawabata, T. Numasawa, and S. Ryu, Entanglement Phase Transition Induced by the Non-Hermitian Skin Effect, Phys. Rev. X 13, 021007 (2023).
- [183] N. Hatano and D. R. Nelson, Localization Transitions in Non-Hermitian Quantum Mechanics, Phys. Rev. Lett. 77, 570 (1996).
- [184] J. Feinberg and A. Zee, Non-Hermitian localization and delocalization, Phys. Rev. E 59, 6433 (1999).
- [185] K. Kawabata and S. Ryu, Nonunitary Scaling Theory of Non-Hermitian Localization, Phys. Rev. Lett. 126, 166801 (2021).
- [186] X. Luo, T. Ohtsuki, and R. Shindou, Universality Classes of the Anderson Transitions Driven by Non-Hermitian Disorder, Phys. Rev. Lett. **126**, 090402 (2021).
- [187] Q. Lin, T. Li, L. Xiao, K. Wang, W. Yi, P. Xue, Observation of non-Hermitian topological Anderson insulator in quantum dynamics, Nat. Commun. 13, 3229 (2022).
- [188] S. Longhi, Topological Phase Transition in non-Hermitian Quasicrystals, Phys. Rev. Lett. 122, 237601 (2019).
- [189] H. Jiang, L.-J. Lang, C. Yang, S.-L. Zhu, and S. Chen, Interplay of non-Hermitian skin effects and Anderson localization in nonreciprocal quasiperiodic lattices, Phys. Rev. B 100, 054301 (2019).
- [190] Q.-B. Zeng, Y.-B. Yang, and Y. Xu, Topological phases in non-Hermitian Aubry-André-Harper models, Phys. Rev. B 101, 020201(R) (2020).
- [191] S. Weidemann, M. Kremer, S. Longhi, and A. Szamei, Topological triple phase transition in non-Hermitian Floquet quasicrystals, Nature 601, 354-359 (2022).
- [192] Q. Lin, T. Li, L. Xiao, K. Wang, W. Yi, and P. Xue, Topological Phase Transitions and Mobility Edges in Non-Hermitian Quasicrystals, Phys. Rev. Lett. **129**, 113601 (2022).
- [193] Y. Ashida, S. Furukawa, and M. Ueda, Parity-time-symmetric quantum critical phenomena, Nat. Commun. 8, 15791 (2017).
- [194] M. Nakagawa, N. Kawakami, and M. Ueda, Non-Hermitian Kondo Effect in Ultracold Alkaline-Earth Atoms, Phys. Rev. Lett. 121, 203001 (2018).
- [195] H. Shen and L. Fu, Quantum Oscillation from In-Gap States and a Non-Hermitian Landau Level Problem, Phys. Rev. Lett. 121, 026403 (2018).
- [196] R. Hamazaki, K. Kawabata, and M. Ueda, Non-Hermitian Many-Body Localization, Phys. Rev. Lett. 123, 090603 (2019).

- [197] D. J. Luitz and F. Piazza, Exceptional points and the topology of quantum many-body spectra, Phys. Rev. Research 1, 033051 (2019).
- [198] T. Yoshida, K. Kudo and Y. Hatsugai, Non-Hermitian fractional quantum Hall states, Sci. Rep. 9, 16895 (2019).
- [199] D.-W. Zhang, Y.-L. Chen, G.-Q. Zhang, L.-J. Lang, Z. Li, and S.-L. Zhu, Skin super-fluid, topological Mott insulators, and asymmetric dynamics in an interacting non-Hermitian Aubry-André-Harper model, Phys. Rev. B 101, 235150 (2020).
- [200] C.-X. Guo, X.-R. Wang, and S.-P. Kou, Non-Hermitian avalanche effect: Non-perturbative effect induced by local non-Hermitian perturbation on a Z₂ topological order, EPL **131**, 27002 (2020).
- [201] N. Okuma and M. Sato, Non-Hermitian Skin Effects in Hermitian Correlated or Disordered Systems: Quantities Sensitive or Insensitive to Boundary Effects and Pseudo-Quantum-Number, Phys. Rev. Lett. **126**, 176601 (2021).
- [202] W. Xi, Z.-H. Zhang, Z.-C. Gu, and W.-Q. Chen, Classification of topological phases in one dimensional interacting non-Hermitian systems and emergent unitarity, Sci. Bull. 66, 1731-1739 (2021).
- [203] K. Yang, S. C. Morampudi, and E. J. Bergholtz, Exceptional Spin Liquids from Couplings to the Environment, Phys. Rev. Lett. 126, 077201 (2021).
- [204] S.-B. Zhang, M. M. Denner, T. Bzdušek, M. A. Sentef, and T. Neupert, Symmetry breaking and spectral structure of the interacting Hatano-Nelson model, Phys. Rev. B 106, L121102 (2022).
- [205] K. Kawabata, K. Shiozaki, and S. Ryu, Many-body topology of non-Hermitian systems, Phys. Rev. B 105, 165137 (2022).
- [206] W. N. Faugno and T. Ozawa, Interaction-Induced Non-Hermitian Topological Phases from a Dynamical Gauge Field, Phys. Rev. Lett. **129**, 180401 (2022).
- [207] K. Suthar, Y.-C. Wang, Y.-P. Huang, H. H. Jen, and J.-S. You, Non-Hermitian many-body localization with open boundaries, Phys. Rev. B 106, 064208 (2022).
- [208] M. S. Rudner and L. S. Levitov, Topological Transition in a Non-Hermitian Quantum Walk, Phys. Rev. Lett. 102, 065703 (2009).
- [209] C. Yuce, PT symmetric Floquet topological phase, Eur. Phys. J. D 69, 184 (2015).

- [210] J. Gong and Q.-h. Wang, Stabilizing non-Hermitian systems by periodic driving, Phys. Rev. A 91, 042135 (2015).
- [211] Y. Huang, Z. Yin, and W. Yang, Realizing a topological transition in a non-Hermitian quantum walk with circuit QED, Phys. Rev. A 94, 022302 (2016).
- [212] T. Rakovszky, J. K. Asbóth, and A. Alberti, Detecting topological invariants in chiral symmetric insulators via losses, Phys. Rev. B 95, 201407(R) (2017).
- [213] X. Zhan, L. Xiao, Z. Bian, K. Wang, X. Qiu, B. C. Sanders, W. Yi, and P. Xue, Detecting Topological Invariants in Nonunitary Discrete-Time Quantum Walks, Phys. Rev. Lett. 119, 130501 (2017).
- [214] L. Xiao, X. Zhan, Z. H. Bian, K. Wang, X. Zhang, X. Wang, J. Li, K. Mochizuki, D. Kim, N. Kawakami, W. Yi, H. Obuse, B. C. Sanders and P. Xue, Observation of topological edge states in parity-time-symmetric quantum walks, Nat. Phys. 13, 1117-1123 (2017).
- [215] S. Longhi, Floquet exceptional points and chirality in non-Hermitian Hamiltonians, J. Phys. A: Math. Theor. 50, 505201 (2017).
- [216] M. Chitsazi, H. Li, F. M. Ellis, and T. Kottos, Experimental Realization of Floquet PT-Symmetric Systems, Phys. Rev. Lett. 119, 093901 (2017).
- [217] T. Chen, B. Wang, and X. Zhang, Characterization of topological phases and selection of topological interface modes in the parity-time-symmetric quantum walk, Phys. Rev. A 97, 052117 (2018).
- [218] R. d. J. León-Montiel, M. A. Quiroz-Juárez, J. L. Domínguez-Juárez, R. Quintero-Torres, J. L. Aragón, A. K. Harter, and Y. N. Joglekar, Observation of slowly decaying eigenmodes without exceptional points in Floquet dissipative synthetic circuits, Commun. Phys. 1, 88 (2018).
- [219] N. Wang, Z.-Q. Zhang, and C. T. Chan, Photonic Floquet media with a complex time-periodic permittivity, Phys. Rev. B 98, 085142 (2018).
- [220] T. T. Koutserimpas and R. Fleury, Nonreciprocal Gain in Non-Hermitian Time-Floquet Systems, Phys. Rev. Lett. 120, 087401 (2018).
- [221] Z. Turker, S. Tombuloglu, and C. Yuce, PT symmetric Floquet topological phase in SSH model, Phys. Lett. A 382, 2013-2016 (2018).
- [222] K. Wang, X. Qiu, L. Xiao, X. Zhan, Z. Bian, B. C. Sanders, W. Yi, and P. Xue, Observation of emergent momentum-time skyrmions in parity-time-symmetric non-unitary quench dynamics,

Nat. Commun. 10, 2293 (2019).

- [223] J. Li, A. K. Harter, J. Liu, L. d. Melo, Y. N. Joglekar and L. Luo, Observation of paritytime symmetry breaking transitions in a dissipative Floquet system of ultracold atoms, Nat. Commun. 10, 855 (2019).
- [224] B. Höckendorf, A. Alvermann, and H. Fehske, Non-Hermitian Boundary State Engineering in Anomalous Floquet Topological Insulators, Phys. Rev. Lett. **123**, 190403 (2019).
- [225] M. Li, X. Ni, M. Weiner, A. Alù, and A. B. Khanikaev, Topological phases and nonreciprocal edge states in non-Hermitian Floquet insulators, Phys. Rev. B 100, 045423 (2019).
- [226] M. van Caspel, S. E. T. Arze, and I. P. Castillo, Dynamical signatures of topological order in the driven-dissipative Kitaev chain, SciPost Phys. 6, 026 (2019).
- [227] W.-L. Zhao, J. Wang, X. Wang, and P. Tong, Directed momentum current induced by the PT-symmetric driving, Phys. Rev. E 99, 042201 (2019).
- [228] X. Zhang and J. Gong, Non-Hermitian Floquet topological phases: Exceptional points, coalescent edge modes, and the skin effect, Phys. Rev. B 101, 045415 (2020).
- [229] B. Höckendorf, A. Alvermann, and H. Fehske, Topological origin of quantized transport in non-Hermitian Floquet chains, Phys. Rev. Research 2, 023235 (2020).
- [230] H. Wu and J.-H. An, Floquet topological phases of non-Hermitian systems, Phys. Rev. B 102, 041119(R) (2020).
- [231] A. Banerjee and A. Narayan, Controlling exceptional points with light, Phys. Rev. B 102, 205423 (2020).
- [232] A. K. Harter and Y. N. Joglekar, Connecting active and passive PT-symmetric Floquet modulation models, Prog. Theor. Exp. Phys. 2020, 12A106 (2020).
- [233] S. Mudute-Ndumbe and E.-M. Graefe, A non-Hermitian PT-symmetric kicked top, New J. Phys. 22 103011 (2020).
- [234] P. He and Z. Huang, Floquet engineering and simulating exceptional rings with a quantum spin system, Phys. Rev. A 102, 062201 (2020).
- [235] H. Wu, B.-Q. Wang, and J.-H. An, Floquet second-order topological insulators in non-Hermitian systems, Phys. Rev. B 103, L041115 (2021).
- [236] Y. Cao, Y. Li, and X. Yang, Non-Hermitian bulk-boundary correspondence in a periodically driven system, Phys. Rev. B 103, 075126 (2021).

- [237] L. Ding, K. Shi, Q. Zhang, D. Shen, X. Zhang, and W. Zhang, Experimental Determination of PT-Symmetric Exceptional Points in a Single Trapped Ion, Phys. Rev. Lett. **126**, 083604 (2021).
- [238] S. Wu, W. Song, S. Gao, Y. Chen, S. Zhu, and T. Li, Floquet π mode engineering in non-Hermitian waveguide lattices, Phys. Rev. Research **3**, 023211 (2021).
- [239] D. Chowdhury, A. Banerjee, and A. Narayan, Light-driven Lifshitz transitions in non-Hermitian multi-Weyl semimetals, Phys. Rev. A 103, L051101 (2021).
- [240] V. M. Vyas and D. Roy, Topological aspects of periodically driven non-Hermitian Su-Schrieffer-Heeger model, Phys. Rev. B 103, 075441 (2021).
- [241] L. Xiao, T. Deng, K. Wang, Z. Wang, W. Yi, and P. Xue, Observation of non-Bloch paritytime symmetry and exceptional points, Phys. Rev. Lett. **126**, 230402 (2021).
- [242] W.-L. Zhao, Y. Hu, Z. Li, and Q. Wang, Super-exponential growth of out-of-time-ordered correlators, Phys. Rev. B 103, 184311 (2021).
- [243] V. Mittal, A. Raj, S. Dey, and S. K. Goyal, Persistence of topological phases in non-Hermitian quantum walks, Sci. Rep. 11, 10262 (2021).
- [244] H. Wu and J.-H. An, Non-Hermitian Weyl semimetal and its Floquet engineering, Phys. Rev. B 105, L121113 (2022).
- [245] C.-H. Liu, H. Hu, and S. Chen, Symmetry and topological classification of Floquet non-Hermitian systems, Phys. Rev. B 105, 214305 (2022).
- [246] W. Zhu and J. Gong, Hybrid skin-topological modes without asymmetric couplings, Phys. Rev. B 106, 035425 (2022).
- [247] D. Chowdhury, A. Banerjee, and A. Narayan, Exceptional hexagonal warping effect in multi-Weyl semimetals, Phys. Rev. B 105, 075133 (2022).
- [248] W.-L. Zhao, Quantization of out-of-time-ordered correlators in non-Hermitian chaotic systems, Phys. Rev. Research 4, 023004 (2022).
- [249] A. K. Ghosh and T. Nag, Non-Hermitian higher-order topological superconductors in two dimensions: Statics and dynamics, Phys. Rev. B 106, L140303 (2022).
- [250] T. Banerjee and K. Sengupta, Emergent conservation in the Floquet dynamics of integrable non-Hermitian models, Phys. Rev. B 107, 155117 (2023).
- [251] C.-H. Liu, H. Hu, S. Chen, and X.-J. Liu, Anomalous second-order skin modes in Floquet non-Hermitian systems, arXiv:2303.11259 (2023).

- [252] Y. Sun, X. Hou, T. Wan, F. Wang, S. Zhu, Z. Ruan, Z. Yang, Photonic Floquet skintopological effect, arXiv:2306.03705 (2023).
- [253] Y. Li, C. Lu, S. Zhang, Y.-C. Liu, Loss-induced Floquet non-Hermitian skin effect, arXiv:2306.04460 (2023).
- [254] L. Zhou and J. Gong, Non-Hermitian Floquet topological phases with arbitrarily many realquasienergy edge states, Phys. Rev. B 98, 205417 (2018).
- [255] L. Zhou and J. Pan, Non-Hermitian Floquet topological phases in the double-kicked rotor, Phys. Rev. A 100, 053608 (2019).
- [256] L. Zhou, Dynamical characterization of non-Hermitian Floquet topological phases in one dimension, Phys. Rev. B 100, 184314 (2019).
- [257] L. Zhou, Non-Hermitian Floquet topological superconductors with multiple Majorana edge modes, Phys. Rev. B 101, 014306 (2020).
- [258] L. Zhou, Non-Hermitian Floquet Phases with Even-Integer Topological Invariants in a Periodically Quenched Two-Leg Ladder, Entropy 22, 746 (2020).
- [259] J. Pan and L. Zhou, Non-Hermitian Floquet second order topological insulators in periodically quenched lattices, Phys. Rev. B 102, 094305 (2020).
- [260] W.-L. Zhao, L. Zhou, J. Liu, P. Tong, and K. Huang, Superexponential diffusion in nonlinear non-Hermitian systems, Phys. Rev. A 102, 062213 (2020).
- [261] L. Zhou, Y. Gu, and J. Gong, Dual topological characterization of non-Hermitian Floquet phases, Phys. Rev. B 103, L041404 (2021).
- [262] L. Zhou, Floquet engineering of topological localization transitions and mobility edges in one-dimensional non-Hermitian quasicrystals, Phys. Rev. Research 3, 033184 (2021).
- [263] L. Zhou, R. W. Bomantara, and S. Wu, qth-root non-Hermitian Floquet topological insulators, SciPost Phys. 13, 015 (2022).
- [264] L. Zhou and W. Han, Driving-induced multiple PT-symmetry breaking transitions and reentrant localization transitions in non-Hermitian Floquet quasicrystals, Phys. Rev. B 106, 054307 (2022).
- [265] L. Zhou, <u>Adiabatic Pumping in Floquet and Open Quantum Systems: An Interplay among</u> <u>Geometry, Topology and Quantum Coherence</u>, Ph.D. thesis, Department of Physics, National University of Singapore (2015).
- [266] G. Grosso and G. P. Parravicini, Solid State Physics (Academic Press, UK, 2014).

- [267] J. H. Shirley, Solution of the Schrödinger Equation with a Hamiltonian Periodic in Time, Phys. Rev. 138, B979 (1965).
- [268] Y. B. Zel'dovich, The quasienergy of a quantum-mechanical system subjected to a periodic action, Sov. Phys. JETP 24, 1006 (1967).
- [269] H. Sambe, Steady States and Quasienergies of a Quantum-Mechanical System in an Oscillating Field, Phys. Rev. A 7, 2203 (1973).
- [270] G. Rigolin, G. Ortiz, and V. H. Ponce, Beyond the quantum adiabatic approximation: Adiabatic perturbation theory, Phys. Rev. A 78, 052508 (2008).
- [271] D. C. Brody, Biorthogonal quantum mechanics, J. Phys. A: Math. Theor. 47, 035305 (2014).
- [272] S. Ryu, A. P. Schnyder, A. Furusaki and A. W. W. Ludwig, Topological insulators and superconductors: tenfold way and dimensional hierarchy, New J. Phys. 12, 065010 (2010).
- [273] I. Mondragon-Shem, T. L. Hughes, J. Song, and E. Prodan, Topological Criticality in the Chiral-Symmetric AIII Class at Strong Disorder, Phys. Rev. Lett. 113, 046802 (2014).
- [274] J. Song and E. Prodan, AIII and BDI topological systems at strong disorder, Phys. Rev. B 89, 224203 (2014).
- [275] L. Zhang, L. Zhang, S. Niu, and X. Liu, Dynamical classification of topological quantum phases, Sci. Bull. 63, 1385 (2018).
- [276] B. Zhu, Y. Ke, H. Zhong, and C. Lee, Dynamic winding number for exploring band topology, Phys. Rev. Research 2, 023043 (2020).
- [277] F. Cardano, A. D'Errico, A. Dauphin, M. Maffei, B. Piccirillo, C. de Lisio, G. D. Filippis, V. Cataudella, E. Santamato, L. Marrucci, M. Lewenstein, and P. Massignan, Detection of Zak phases and topological invariants in a chiral quantum walk of twisted photons, Nat. Commun. 8, 15516 (2017).
- [278] M. Maffei, A. Dauphin, F. Cardano, M. Lewenstein, and P. Massignan, Topological characterization of chiral models through their long time dynamics, New J. Phys. 20, 013023 (2018).
- [279] E. J. Meier, F. A. An, A. Dauphin, M. Maffei, P. Massignan, T. L. Hughes, and B. Gadway, Observation of the topological Anderson insulator in disordered atomic wires, Science 362, 929 (2018).
- [280] J. Bartlett and E. Zhao, Unravelling the edge spectra of non-Hermitian Chern insulators, Phys. Rev. B 107, 035101 (2023).

- [281] S. Malzard, C. Poli, and H. Schomerus, Topologically Protected Defect States in Open Photonic Systems with Non-Hermitian Charge-Conjugation and Parity-Time Symmetry, Phys. Rev. Lett. 115, 200402 (2015).
- [282] L. Zhou, Generating Many Majorana Corner Modes and Multiple Phase Transitions in Floquet Second-Order Topological Superconductors, Symmetry 14, 2546 (2022).
- [283] J. Arkinstall, M. H. Teimourpour, L. Feng, R. El-Ganainy and H. Schomerus, Topological tight-binding models from nontrivial square roots, Phys. Rev. B 95, 165109 (2017).
- [284] M. Kremer, I. Petrides, E. Meyer, M. Heinrich, O. Zilberberg and A. Szameit, A squareroot topological insulator with non-quantized indices realized with photonic Aharonov-Bohm cages, Nat. Commun. 11, 907 (2020).
- [285] L. Song, H. Yang, Y. Cao, and P. Yan, Realization of the square-root higher-order topological insulator in electric circuits, Nano Lett. 20, 7566 (2020).
- [286] M. Yan, X. Huang, L. Luo, J. Lu, W. Deng and Z. Liu, Acoustic square-root topological states, Phys. Rev. B 102, 180102 (2020).
- [287] T. Mizoguchi, Y. Kuno, and Y. Hatsugai, Square-root higher-order topological insulator on a decorated honeycomb lattice, Phys. Rev. A 102, 033527 (2020).
- [288] M. Ezawa, Systematic construction of square-root topological insulators and superconductors, Phys. Rev. Res. 2, 033397 (2020).
- [289] A. M. Marques, L. Madail and R. G. Dias, One-dimensional 2ⁿ-root topological insulators and superconductors, Phys. Rev. B 103, 235425 (2021).
- [290] A. M. Marques and R. G. Dias, 2ⁿ-root weak, Chern, and higher-order topological insulators, and 2n-root topological semimetals, Phys. Rev. B 104, 165410 (2021).
- [291] T. Yoshida, T. Mizoguchi, Y. Kuno and Y. Hatsugai, Square-root topological phase with time-reversal and particle-hole symmetry, Phys. Rev. B 103, 235130 (2021).
- [292] Z. Lin, S. Ke, X. Zhu and X. Li, Square-root non-Bloch topological insulators in non-Hermitian ring resonators, Opt. Express 29, 8462 (2021).
- [293] R. W. Bomantara, Square-root Floquet topological phases and time crystals, Phys. Rev. B 106, L060305 (2022).
- [294] P. A. M. Dirac, The quantum theory of the electron, Proc. R. Soc. London A 117, 610 (1928).
- [295] L. Zhou and W. Han, Non-Hermitian quasicrystal in dimerized lattices, Chin. Phys. B 30, 100308 (2021).

- [296] W. Han, and L. Zhou, Dimerization-induced mobility edges and multiple reentrant localization transitions in non-Hermitian quasicrystals, Phys. Rev. B 105, 054204 (2022).
- [297] L. Zhou, Non-Abelian generalization of non-Hermitian quasicrystal: PT-symmetry breaking, localization, entanglement and topological transitions, Phys. Rev. B 108, 014202 (2023).
- [298] L. Zhou, Q.-h. Wang, H. Wang, and J. Gong, Dynamical quantum phase transitions in non-Hermitian lattices, Phys. Rev. A 98, 022129 (2018).
- [299] L. Zhou and Q. Du, Non-Hermitian topological phases and dynamical quantum phase transitions: a generic connection, New J. Phys. 23, 063041 (2021).
- [300] K. Yang, L. Zhou, W. Ma, X. Kong, P. Wang, X. Qin, X. Rong, Y. Wang, F. Shi, J. Gong, and J. Du, Floquet dynamical quantum phase transitions, Phys. Rev. B 100, 085308 (2019).
- [301] L. Zhou and Q. Du, Floquet dynamical quantum phase transitions in periodically quenched systems, J. Phys.: Condens. Matter 33, 345403 (2021).
- [302] J. Naji, R. Jafari, L. Zhou, and A. Langari, Engineering Floquet dynamical quantum phase transitions, Phys. Rev. B 106, 094314 (2022).
- [303] L. Zhou, Entanglement spectrum and entropy in Floquet topological matter, Phys. Rev. Research 4, 043164 (2022).