

ORBIT RECOVERY FOR BAND-LIMITED FUNCTIONS

DAN EDIDIN, MATTHEW SATRIANO

ABSTRACT. We study the third moment for functions on arbitrary compact Lie groups. We use techniques of representation theory to generalize the notion of band-limited functions in classical Fourier theory to functions on the compact groups $SU(n)$, $SO(n)$, $Sp(n)$. We then prove that for generic band-limited functions the third moment or, its Fourier equivalent, the bispectrum determines the function up to translation by a single unitary matrix. Moreover, if $G = SU(n)$ or $G = SO(2n + 1)$ we prove that the third moment determines the G -orbit of a band-limited function. As a corollary we obtain a large class of finite-dimensional representations of these groups for which the third moment determines the orbit of a generic vector. When $G = SO(3)$ this gives a result relevant to cryo-EM which was our original motivation for studying this problem.

1. INTRODUCTION

Let G be a compact Lie group. The purpose of this paper is construct a class of finite-dimensional representations V of G for which the third moment can determine the orbit of a generic vector $f \in V$. As we explain, this work is motivated by several applications including multi-reference alignment (MRA), cryo-EM, and machine learning.

In its basic form, the multi-reference alignment (MRA) problem seeks to recover a signal $f \in V$ from noisy group translates of the signal

$$y_i = g_i \cdot f + \epsilon_i$$

where the g_i are randomly selected from a uniform distribution on G and the ϵ_i are taken from a Gaussian distribution $\mathcal{N}(0, \sigma^2 I)$ which is independent of the group element g_i . Without prior knowledge of the group elements, there is no way to distinguish f from $g \cdot f$ for any $g \in G$. Thus the MRA problem is one of orbit recovery. The MRA problem has been extensively studied in recent years, beginning with action of \mathbb{Z}_N on \mathbb{R}^N by cyclic shifts [3, 25, 8, 1, 5]. Other models include the dihedral group [10] and the rotation group $SO(2)$ acting on band-limited functions on \mathbb{R}^2 [4, 24, 19]. The case where $G = SO(3)$ is particularly important because of its connection to cryo-EM, a leading technique in molecular imaging. There, the measured data can be modelled as $y_i = T(g \cdot f) + \epsilon$ where f is the Coulomb potential of an unknown molecular structure and T is a tomographic projection [6].

In the low-noise regime the products $g_i g_j^{-1}$ can be estimated using the method of synchronization and then the signal can be approximated by averaging [26]. However, in the high-noise regime, as is the case for cryo-EM measurements of small molecules, there is no way to accurately estimate the relation between the unknown group elements [7, Proposition 2.1]. One common approach to this problem is to use the *method of moments*. In this case, it can be shown [25] that the moments of the unknown signal can be accurately approximated by the computing the corresponding moments of the experimental data. Thus, a crucial aspect of the MRA problem is understanding how to recover a signal $f \in V$ from its moments which are, by definition, G -invariant tensors in the signal.

In machine learning it is desirable to build neural networks whose architecture reflects the intrinsic structure of the data. When the data has natural symmetries under a group G then we want to build the network from G -equivariant functions. (Recall that if \mathbb{V} and \mathbb{W} are sets with a G action then a function $f: \mathbb{V} \rightarrow \mathbb{W}$ is G -equivariant if $f(gv) = gf(v)$ for all $g \in G$.) The basic model of an equivariant neural network [14, 22] is a sequence of maps

$$\mathbb{R}^{n_0} \xrightarrow{A_1} \mathbb{R}^{n_1} \xrightarrow{\sigma_{b_1}} \mathbb{R}^{n_2} \dots \xrightarrow{\sigma_{b_{k-1}}} \mathbb{R}^{n_{k-1}} \xrightarrow{A_k} \mathbb{R}^{n_k}$$

where each \mathbb{R}^{n_i} is a representation of G , the A_i are G -equivariant linear transformations and the σ_{b_i} are non-linear maps.

One difficulty with this model is that there may be relatively few G -equivariant linear maps of representations $A_i: \mathbb{R}^n \rightarrow \mathbb{R}^m$, so an equivariant neural network built this way may not be sufficiently expressive. Two common, and mathematically related, ideas are to use equivariant linear maps on tensors $(\mathbb{R}^n)^{\otimes k} \rightarrow (\mathbb{R}^m)^{\otimes \ell}$, or invariant polynomials of degree ℓ from $\mathbb{R}^n \rightarrow \mathbb{R}^m$ [21, 15, 12].

In both MRA and machine learning the use of tensors is theoretically desirable, but the cost of computing tensors grows exponentially in the degree. In addition, in MRA the sample complexity (the minimum number of observations required for accurate approximation) grows as σ^{2d} where d is the number of moments used and σ^2 is the variance of the noise. For this reason, we wish to identify representations for which moments of low degree separate generic orbits. Previous work of Bendory and the first author demonstrated that neither the first or second moment carries enough information to separate orbits in all but the simplest representations [9]. Thus an important problem is to understand and classify representations of compact groups for which the third moment can separate orbits. Previous work [2] showed that for any finite group the generic orbit in the regular representation can be recovered from the third moment using Jennrich's algorithm. More generally it is proved in [27] that if G is a positive dimensional compact group then the G -orbit of a function in the infinite dimensional representation $L^2(G)$ with non-singular Fourier coefficients can be recovered from its third moment.

In this paper we show that if G is one of the classical groups $SU(n), SO(n), Sp(n)$ then it is possible to use representation theory to define the notion of *band-limited function* which generalizes the notion of band-limited function in Fourier theory as well as previous definitions for $G = SO(3)$. Our main result can be stated as follows.

Theorem 1.1 (Informal). *(i) If G is one of the classical groups then the generic band-limited function can be recovered up to translation by a single unitary matrix from its third moment.*

(ii) The orbit of any generic band-limited function in $L^2(SU(n))$ or real-valued band-limited function in $L^2(SO(2n+1))$ can be recovered from its third moment.

Remark 1.2. As is the case in Fourier theory, for any band level, the vector space of band limited functions is finite dimensional, so our result gives a large class of finite-dimensional representation for which the third moment separates generic orbits.

The proof of Theorem 1.1 uses representation theory to generalize the well-known frequency marching result that states that a band-limited function on S^1 can be recovered from its bispectrum. When $G = SO(3)$ Kakarala [20] showed that the $SO(3)$ -orbit of a band-limited real valued function can be recovered up to reflection from the bispectrum, and our

result eliminates the reflection ambiguity. Indeed the idea of using techniques of representation theory to study the orbit recovery problem was motivated by Kakarala's earlier work for $\text{SO}(3)$.

As an application we prove (Corollary 4.2) that if we consider the finite dimensional approximation of $L^2(\mathbb{R}^3)$ as band-limited functions on R spherical shells, where the number of shells exceeds the band-limit then the $\text{SO}(3)$ orbit of a generic function can be recovered from the third moment. This is a standard assumption in the cryo-EM literature [4] and our result affirms many cases of a conjecture made in [2].

2. MOMENTS OF REPRESENTATIONS OF COMPACT GROUPS

Let V be a unitary representation of a compact group G . The d -th moment of V is the function $m_d: V \rightarrow \underbrace{V \otimes \dots \otimes V}_{d-1 \text{ times}} \otimes V^*$ defined by the formula

$$(2.1) \quad f \mapsto m_d(f) := \int_G \overbrace{g \cdot f \otimes \dots \otimes g \cdot f}^{d-1 \text{ times}} \otimes \overline{g \cdot f} dg$$

Note that the formula of (2.1) is invariant under translation by G so for any $f \in V$, $m_d(f)$ a G -invariant element of the tensor $V^{d-1} \otimes V^*$ or equivalently a G -invariant element of $\text{Hom}(V, V^d)$.

If V is a real representation then the d -th moment is an element of $\text{Sym}^d V$ and the collection of moments form a set of generators for the invariant ring $\mathbb{R}[V]^G$.

When V is identified with a space of functions $D \rightarrow \mathbb{C}$ where D is a domain on which G acts (for example we can take $D = G$ and consider V as a subspace of $L^2(G)$) then $m_d(f)$ is viewed as a function $D^d \rightarrow \mathbb{C}$ defined by the formula

$$(2.2) \quad m_d(f)(x_1, \dots, x_d) = \int_G (g \cdot f)(x_1) \dots (g \cdot f)(x_{d-1}) \overline{(g \cdot f)(x_d)} dg$$

where $g \cdot f: D \rightarrow \mathbb{C}$ is the function $(g \cdot f)(x) = f(g^{-1}x)$.

2.1. Moments and the decomposition into irreducibles. A general finite dimensional representation of a compact group can be decomposed as $V = \bigoplus_{\ell=1}^L V_\ell^{R_\ell}$ where the V_ℓ are distinct non-isomorphic irreducible representations of G of dimension N_ℓ . An element $f \in V$ has a unique G -invariant decomposition as a sum

$$(2.3) \quad f = \sum_{\ell=1}^L \sum_{i=1}^{R_\ell} f_\ell[i]$$

Where $f_\ell[i]$ is in the i -th copy of the irreducible representation V_ℓ . For fixed ℓ the vectors $f_\ell[1], \dots, f_\ell[R_\ell]$ determine a $N_\ell \times R_\ell$ matrix $A_\ell(f)$ which we sometimes refer to as the *coefficient matrix of f in V_ℓ* .

The d -th moment is a map $m_d: V \rightarrow (V^{\otimes(d-1)} \otimes V^*)^G$, which takes a vector $f \in V$ to invariant part of the tensor $f \otimes f \otimes \dots \otimes \bar{f}$. Identifying $V^{\otimes(d-1)} \otimes V^* = \text{Hom}(V, V^{\otimes(d-1)})$, the d -th moment decomposes according to the decomposition of V into irreducibles as a functions

$$m_d[V_{i_1}, \dots, V_{i_d}]: V \rightarrow \bigoplus_{i_1, \dots, i_d} \text{Hom}(W_{i_d}, W_{i_1} \otimes \dots \otimes W_{i_{d-1}})^G$$

where $W_\ell = V_\ell^{R_\ell}$. By Schur's lemma $m_d[V_{i_1}, \dots, V_{i_d}]$ is non-zero if and only if the irreducible representation V_{i_d} is a summand in the tensor product $W_{i_1} \otimes \dots \otimes W_{i_{d-1}}$.

The map $m_d[V_{i_1}, \dots, V_{i_d}]$ can be described explicitly as follows: If $f = \sum_{\ell=1}^L A_\ell(f)$ where $A_\ell(f) = (f_\ell[1], \dots, f_\ell[R_L]) \in W_\ell$, let $B_{i_d}(f)$ be the projection of

$$A_{i_1}(f) \otimes \dots \otimes A_{i_{d-1}}(f) \in W_{i_1} \otimes \dots \otimes W_{i_{d-1}}$$

to the isotypic component, T_{i_d} of $W_{i_1} \otimes \dots \otimes W_{i_{d-1}}$ corresponding to the irreducible representation V_{i_d} . Then $m_d[V_{i_1}, \dots, V_{i_d}](f)$ is the element of $\text{Hom}(W_{i_d}, T_{i_d})$ represented by the matrix $A_{i_d}(f)B_{i_d}(f)^*$.

In the case when $d = 2$ there is a simple description of the information determined by the second moment.

Proposition 2.1. [9, Theorem 2.3] *Let V be a finite dimensional representation of G . The second moment $m_2(f)$ determines, for each irreducible V_ℓ appearing in V , the product $A_\ell(f)A_\ell(f)^*$. In particular, if $A_\ell(f)$ is full rank, then the second moment determines $A_\ell(f)$ up to translation by an element of $U(V_\ell)$, or if $A_\ell(f)$ is real, and element of $O(V_\ell)$.*

The next result proves that if V contains a copy of the trivial representation then the third moment determines the first and second moments.

Proposition 2.2. *If the coefficient matrix, $A_0(f)$, of the trivial representation is non-zero then $m_1(f)$ and $m_2(f)$ are determined from $m_3(f)$.*

Proof. Since the trivial representation V_0 is one-dimensional the coefficient matrix $A_0(f) = (f_0[1], \dots, f_0[R_0])$ is just a row vector of length R_0 where R_0 is the multiplicity. The first moment is simply the projection of $V \rightarrow V^G = V_0^{R_0}$ so $m_1(f) = A_0(f)$. On the other hand, consider the component $m_3[V_0, V_0, V_0]$ of the third moment which is a G -invariant map $V \rightarrow W_0 \otimes W_0 \otimes W_0^*$. Since G acts trivially on W_0 , the entire tensor product $W_0 \otimes W_0$ is the V_0 isotypic component. Thus $m_3(f)[V_0, V_0, V_0] = A_0(f)(A_0(f) \otimes A_0(f))^*$ where we view $A_0(f) \otimes A_0(f)$ as a matrix of size $1 \times R_0^2$. Among the entries of this matrix are the products $f_0[j]\overline{f_0[j]}f_0[j] = \overline{f_0[j]}|f_0[j]|^2$. Writing $f_0[j] = r_j e^{i\theta}$ for $r_j > 0$ we see that $f_0[j]$ is determined by $\overline{f_0[j]}|f_0[j]|^2$.

In particular if $f_0[i] \neq 0$ is known, $m_2(f)$ is determined by $\frac{1}{|A_0(f)|} \int_G A_0[f] \otimes g \cdot f \otimes \overline{g \cdot f}$ which is a sum of components of $m_3(f)$. \square

The following bootstrap result is a generalization of [2, Proposition 4.15].

Proposition 2.3. *Let V be a representation of G with $V = V_1^{R_1} \oplus \dots \oplus V_L^{R_L}$ where V_1, \dots, V_L distinct irreducibles. Assume that for every $1 \leq k \leq L$, the generic vector f in the representation $V_1^{R_1} \oplus \widehat{V_k^{R_k}} \dots \oplus V_L^{R_L}$ has trivial stabilizer. (Here the notation $\widehat{V_k^{R_k}}$ means that the summand is omitted.)*

If the orbit of a generic vector $f \in V$ is determined from the d -th moment $m_d(f)$ and $W \supset V$ is another representation with the same irreducible components then the orbit of a generic vector $h \in W$ is determined by its the d -th moment $m_d(h)$.

Example 2.4. The hypothesis that W has the same irreducible components as V is necessary. Consider the case where $G = S^1$ and $V = V_0 \oplus V_1$, $W = V_0 \oplus V_1 \oplus V_3$ where the V_n is the one-dimensional representation of S^1 where S^1 acts with weight n ; i.e. $e^{i\theta} \cdot v = e^{ni\theta}v$. If $v = (v_0, v_1) \in V$ then the third moment determines v_0 and $|v_1|^2$ which determines the vector v up to multiplication by an element S^1 . On the other hand, the third moment of $w = (w_0, w_1, w_3) \in W$ determines, $w_0, |w_1|^2, |w_3|^2$ which is not sufficient to determine the vector w up to multiplication by an element of S^1 .

Proof. By induction on the multiplicities we may reduce to the case that

$$W = V_1^{R_1} \oplus \dots \oplus V_\ell^{R_\ell+1} \oplus \dots \oplus V_L^{R_L}$$

; i.e. all multiplicities of irreducibles in W are the same as in V except for the multiplicity of V_ℓ which is R_ℓ in V and $R_\ell + 1$ in W . By reordering the irreducibles we assume that $\ell = 1$

Suppose $h \in W$ has coefficient matrices, B_1, B_2, \dots, B_L where $B_1 = (f_1[1], \dots, f_1[R_1 + 1])$. For each $j = 1, \dots, R+1$ consider the G -invariant projection $\pi_j: W \rightarrow V$ which sends h to the vector f_j with coefficient matrices B_1^j, B_2, \dots, B_L where $B_1^j = (f_1[1], \dots, \widehat{f_1[j]}, \dots, f_1[R+1])$. By assumption, the G -orbit of f_j is determined from $m_2(f_j) = m_2(\pi_j h)$. In particular, if h' is another vector in W with $m_d(h) = m_d(h')$ and coefficient matrices, B'_1, \dots, B'_L then there exists $g_1, \dots, g_{R+1} \in G$ such that $g_j(B_1^j, \dots, B_L) = (B_1^{j'}, \dots, B'_L)$. To show that g_j 's are all equal note that for any j_1, j_2 the vector in the representation $V_2^{R_2} \oplus \dots \oplus V_L^{R_L}$ with coefficient matrices B_2, \dots, B_L is fixed by $g_{j_1} g_{j_2}^{-1}$ so by assumption on the representation we can conclude that $g_{j_1} = g_{j_2}$. Thus, there exists $g \in G$ such that for every j , $g(B_1^j, B_2, \dots, B'_L) = (B_1^{j'}, B'_2, \dots, B'_L)$. Since every row vector in B_1 (resp B'_1) is in some matrix B_1^j (resp. $B_1^{j'}$) it follows that $g(B_1, \dots, B_L) = (B'_1, \dots, B'_L)$. In other words, $h' = gh$ for some $g \in G$. \square

2.2. The regular representation and higher-order spectra. Here we take $V = L^2(G)$ to be the regular representation. In this case

$$m_d(f)(g_1, \dots, g_d) = \int_G f(g^{-1}g_1) f(gg_2) \dots f(g^{-1}g_{d-1}) \overline{f(g^{-1}g_d)} dg$$

Applying the change of variables $g = g^{-1}g_d$ we can rewrite

$$m_d(f)(g_1, \dots, g_d) = m_d(f)(g_d^{-1}g_1, \dots, g_d^{-1}g_{d-1}, 1).$$

Hence after replacing g with g^{-1} we may view the d -th moment of the regular representation as the function on G^{d-1} .

$$(2.4) \quad m_d(f)(g_1, \dots, g_{d-1}) = \int_G f^*(g) f(gg_1) \dots f(gg_{d-1}) dg$$

where $f^*(g) = \overline{f(g)}$.

If G is a compact group and $f \in L^2(G)$ then the *Fourier transform* (see [27] as well as [18, Chapter 8]) of f is a matrix-valued function $F(f)$ defined on the set representations of G by the formula

$$(2.5) \quad F(f)(V) = \int_G f(g) D_V(g)^* dg \in \text{End}(V)$$

where $D_V(g)$ the unitary linear transformation $V \rightarrow V$ defined by $v \mapsto gv$.

The matrix $F(f)(V)$ is called the *Fourier coefficient* of V . Later, we will implicitly choose a basis for each irreducible representation so we can view the Fourier coefficient as a matrix. As is the case for the classical Fourier transform, a function $f \in L^2(G)$ is uniquely determined by its Fourier coefficients $F(f)(V)$ where V runs through all irreducible representations of G [18, Theorem 31.5].

The $(d-1)$ -st higher-spectrum $a_d(f)$ is defined as the Fourier transform of the function $m_d(f) \in L^2(G^{d-1})$. Since every irreducible representation of G^{d-1} is of the form $V_1 \otimes \dots \otimes V_{d-1}$

we have that

(2.6)

$$a_d(f)(V_1 \otimes \dots \otimes V_{d-1}) = \int_{G^{d-1}} \left(\int_G f^*(g) f(gg_1) \dots f(gg_{d-1}) dg \right) D_{V_1}(g_1)^* \otimes \dots \otimes D_{V_{d-1}}(g_{d-1}^*) dg_1 \dots dg_{d-1}$$

Using the change of coordinates where replace g_i with gg_i and reversing the order of integration the right-hand side of (2.6) becomes

(2.7)

$$\begin{aligned} & \int_G \int_{G^{d-1}} f(g_1) \dots f(g_{d-1}) [D_{V_1}(g_1)^* \otimes \dots \otimes D_{V_{d-1}}(g_{d-1}^*)] dg_1 \dots dg_{d-1} [D_{V_1}(g) \otimes \dots \otimes D_{V_{d-1}}(g)] f^*(g) dg \\ &= [F(f)(V_1) \otimes \dots \otimes F(f)(V_{d-1})] \int_G f^*(g) [D_{V_1}(g) \otimes \dots \otimes D_{V_{d-1}}(g)] dg \\ &= [F(f)(V_1) \otimes \dots \otimes F(f)(V_{d-1})] [F(f)(V_1 \otimes \dots \otimes V_{d-1})]^* \end{aligned}$$

where the product is taken in the ring $\text{End}(V_1 \otimes \dots \otimes V_{d-1})$. Since the $(d-1)$ -st higher spectrum is the Fourier transform of the d -th moment and the Fourier transform is invertible the $(d-1)$ -st higher spectrum carries the same information as the d -th moment.

2.3. The bispectrum. When $d = 3$ the third moment $m_d(f)$ carries the same information as the bispectrum $a_2(f)$ whose value on a tensor $V \otimes W$ is

$$(2.8) \quad a_2(f)(V \otimes W) = [F(f)(V) \otimes F(f)(W)] [F(f)(V \otimes W)]^*$$

For every pair of irreducible representations V, W we choose an isomorphism

$$V \otimes W \simeq V_1 \oplus \dots \oplus V_r$$

with the V_i not necessarily distinct irreducibles. It follows that there are unitary matrices $C_{V,W}$ such that for all $f \in L^2(G)$

$$F(f)(V \otimes W) = C_{V,W} (F(f)(V_1) \oplus \dots \oplus F(f)(V_r)) C_{V,W}^*$$

where we identify $F(f)(V \otimes W)$ as a matrix with respect to a pre-chosen basis for $V \otimes W$. Thus we can rewrite the bispectrum as

$$(2.9) \quad a_2(f)(V \otimes W) = [F(f)(V) \otimes F(f)(W)] C_{V,W} [F(f)(V_1)^* \oplus \dots \oplus F(f)(V_r)^*] C_{V,W}^*$$

Lemma 2.5. *Let V and W be irreducible representations of G and let V_i be any irreducible appearing as a summand in $V \otimes W$. If $f \in L^2(G)$ is chosen such that the Fourier coefficients $F(f)(V)$ and $F(f)(W)$ are invertible, then the Fourier coefficient $F(f)(V_i)$ is determined by $F(f)(V)$, $F(f)(W)$ and the coefficient $a_2(f)(V \otimes W)$ of the bispectrum.*

Proof. Since

$$a_2(f)(V \otimes W) = [F(f)(V) \otimes F(f)(W)] [F(f)(V \otimes W)]^*$$

and $F(f)(V) \otimes F(f)(W)$ is invertible by hypothesis, we obtain

$$F(f)(V \otimes W)^* = [F(f)(V) \otimes F(f)(W)] a_2(f)(V \otimes W)^{-1}.$$

Using the decomposition

$$F(f)(V_1 \otimes V_2) = C_{V,W} [F(f)(V_1) \oplus \dots \oplus F(f)(V_r)] C_{V,W}^*$$

where V_1, \dots, V_r are irreducible yields the lemma. \square

Remark 2.6. The value of Lemma 2.5 is that it shows that if V, W are irreducibles and V_i is an irreducible summand appearing in $V \otimes W$ then $F(f)(V_i)$ is determined from $F(f)(V)$, $F(f)(W)$ and the bispectrum coefficient $a_2(f)(V \otimes W)$. We will use this observation repeatedly.

Proposition 2.7. [27, Theorem 5] *If the Fourier coefficients of $f \in L^2(G)$ are all non-singular then f is uniquely determined by its bispectrum*

Proof Sketch. We first observe that for every irreducible representation V , $a_2(f)(V \otimes \mathbf{1}) = F(f)(V)F(f)(V)^*$ where $\mathbf{1}$ denotes the trivial representation. Hence we know the matrices $F(f)(V)$ up to multiplication by some unknown unitary matrix $u(V) \in U(V)$. In particular if h is a function with the same bispectrum as f then $F(h)(V) = F(f)(V)u(V)$ for all irreducible representations V . The goal is to show that these unitary matrices are all of the form $D_V(g)$ for a fixed $g \in G$ and $D_V(g)$.

Since we know that the function h whose Fourier coefficient $F(h)(V) = F(f)(V)u(V)$ has the same bispectrum as f we see that

$$(2.10) \quad [F(f)(V) \otimes F(f)(W)][(F(f)(V \otimes W))]^* = \\ [F(f)(V)u(V) \otimes F(f)(W)u(W)][F(f)(V \otimes W)u(V \otimes W)]^*$$

Since we assume that Fourier coefficients are all invertible we can conclude that $u(V) \otimes u(W) = u(V \otimes W)$. Moreover, if $A: V \rightarrow W$ is an intertwining operator between representations; i.e. a G -invariant element of $\text{Hom}(V, W)$ then $Au(V) = u(W)A$.

These facts imply that $u(V) = D_V(g)$ for some fixed element $g \in G$, by Tannaka-Krein duality [18, Theorem 30.43]. \square

3. BANDING FUNCTIONS FOR SIMPLE COMPACT LIE GROUPS

For functions on S^1 the notion of band-limiting is well understood. We say that $f \in L^2(S^1)$ is b -band-limited if the Fourier coefficient $f_n = \int_{S^1} e^{in\theta} f(\theta) d\theta = 0$ for $|n| > b$ where the functions $\{e^{in\theta}\}_{n \in \mathbb{Z}}$ form an orthonormal basis for $L^2(S^1)$. For functions on S^2 there is also a corresponding notion of banding using the fact that any $f \in L^2(S^2)$ can be expanded in terms of spherical harmonics $\{Y_\ell^m(\phi, \theta)\}$ where $\ell \in \mathbb{N}$ and for each ℓ , m ranges from $-\ell$ to ℓ . In this context we say that $f = \sum_{\ell, m} a_{\ell, m} Y_\ell^m$ is L -band limited if $a_{\ell, m} = 0$ for all $\ell > L$ and all $m \in [-\ell, \ell]$. Band limited functions on S^2 can be understood in terms of the representation theory of $\text{SO}(3)$ as follows. The space of functions $L^2(S^2)$ decomposes as a representation of $\text{SO}(3)$ into an infinite sum of irreducibles $\oplus_{\ell \geq 0} V_\ell$ where V_ℓ is the $(2\ell + 1)$ -dimensional irreducible representation of $\text{SO}(3)$ spanned by the functions $\{Y_\ell^m\}_{m=-\ell, \dots, \ell}$. With this notation any $f \in V_\ell$ has *band* ℓ and the space of L -band-limited functions is the finite-dimensional representation $\oplus_{\ell=0}^L V_\ell$.

Using the theory of highest-weight vectors we show that the irreducible representations of a large class of Lie groups, including the classical groups $\text{SU}(n), \text{SO}(2n), \text{Sp}(n)$ can be banded. However, for groups of rank more than one, there will be more than one representation with a given band b . Nevertheless, every irreducible representation of a given band b appears as a summand in the tensor product of irreducible representations of lower band. As a consequence we can use a generalized frequency marching argument to show (Proposition 3.2) that if suitable Fourier coefficients are invertible then the Fourier coefficients of irreducible representations of band $b > 1$ can be recovered from the bispectrum and the Fourier coefficients of the irreducible representations of band one. For the classical groups

we can go further and prove in Theorem 3.4 that the irreducible representations of band one can be ordered in such a way that they are determined from the bispectrum and the Fourier coefficient of the defining representation.

3.0.1. *Fundamental representations and banding for simply connected groups.* We begin with the case that G is simply connected which is true for $G = \mathrm{SU}(n)$ or $G = \mathrm{Sp}(n)$. Any compact Lie group is the maximal compact subgroup of a corresponding complex algebraic group $G_{\mathbb{C}}$ [13, Propositions 8.3, 8.6]. When G is simply connected, the representations of G correspond to the representations of \mathfrak{g} where \mathfrak{g} is the Lie algebra of $G_{\mathbb{C}}$ [17, Theorem 3.7]. Since \mathfrak{g} is a semi-simple Lie algebra any representation decomposes into a sum of irreducible representations. Each irreducible representation decomposes as a sum of *weight spaces*. These are the common eigenspaces for the action of the Cartan subalgebra \mathfrak{h} which is the maximal abelian Lie subalgebra of \mathfrak{g} . The dimension of \mathfrak{h} is called the *rank* of \mathfrak{g} . The eigenvalues for the action of \mathfrak{h} on all representations of \mathfrak{g} generate a lattice in \mathfrak{h}^* called the weight lattice Λ_W . The eigenvalues for the action of \mathfrak{g} on itself are called *roots* and they generate the root lattice Λ_R in \mathfrak{h}^* . The root lattice Λ_R has finite index in the weight lattice Λ_W and Λ_R is the dual lattice to Λ_W with respect to a natural inner product on \mathfrak{h}^* . The set of roots Φ of \mathfrak{g} can be divided (by choice of a hyperplane in \mathfrak{h}^*) into positive and negative roots. The *positive simple roots* form basis for the root lattice characterized by the property that any positive root is a non-negative integral linear combination of the positive simple roots. A weight vector $\lambda \in \Lambda_W$ is *dominant* if its inner product with every positive simple root is non-negative. Any dominant weight vector is a non-negative integral linear span of the *fundamental weights* $\omega_1, \dots, \omega_n$ where n is the rank of the Lie algebra \mathfrak{g} .

Irreducible representations of a semi-simple Lie algebra are determined by their *highest weight vectors*, which is the unique dominant weight of the irreducible representation which maximizes the sum of the inner products with the fundamental weights. Moreover, any dominant weight $\lambda = a_1\omega_1 + \dots + a_n\omega_n$ with $a_i \in \mathbb{N}$ is the highest weight vector for a unique irreducible representation V_λ [16, p.205].

When G is simply connected we define the *band* of the representation V_λ with highest weight vector $\lambda = a_1\omega_1 + \dots + \omega_n$ to be $b = a_1 + \dots + a_n$.

3.0.2. *Type A_{n-1} - the group $\mathrm{SU}(n)$.* The compact group $\mathrm{SU}(n)$ is a compact form the algebraic group $\mathrm{SL}(n, \mathbb{C})$ and because $\mathrm{SU}(n)$ is simply connected representations of $\mathrm{SU}(n)$ bijectively correspond to representations of the complex Lie algebra \mathfrak{sl}_n which has type A_{n-1} for $n \geq 2$. The Lie algebra \mathfrak{sl}_n is the vector space of traceless $n \times n$ complex matrices and the Cartan subalgebra \mathfrak{h} is the subspace of diagonal traceless matrices.

The weight lattice is the lattice spanned by vectors L_1, \dots, L_{n-1}, L_n with $L_1 + \dots + L_n = 0$, where L_i is the function on \mathfrak{h} which reads the i -th entry along the diagonal. In this case the positive simple roots are

$$L_1 - L_2, \dots, L_{n-1} - L_n$$

and the fundamental weights are [16, p. 216]

$$\omega_i = \sum_{j \leq i} L_j$$

for $i < n$. The irreducible representation corresponding to ω_1 is the n -dimensional defining representation V ; i.e. the representation $\mathrm{SU}(n) \subset U(n)$. The other representations of band one (i.e. those associated to the fundamental representations) are the exterior powers

$V_k = \wedge^k V$ for $i = 1, \dots, n-1$. (Note that $\wedge^n V$ is the trivial representation which is consistent with the fact that $L_1 + \dots + L_n = 0$.)

3.0.3. Type C_n - the group $\mathrm{Sp}(n)$. The compact symplectic group $\mathrm{Sp}(n)$ is the intersection of the complex symplectic group $\mathrm{Sp}(2n, \mathbb{C})$ with the unitary group $U(2n, \mathbb{C})$. Since this group is simply connected the irreducible representations of $\mathrm{Sp}(2n)$ are the same as the irreducible representations of the complex Lie algebra \mathfrak{sp}_{2n} which has type C_n in the classification of Lie algebras. In this case the weight lattice is freely generated by vectors L_1, \dots, L_n . The positive simple roots are

$$L_1 - L_2, \dots, L_{n-1} - L_n, 2L_n$$

and the fundamental weights are

$$\omega_i = \sum_{j \leq i} L_j$$

for $i \leq n$ [16, Section 17.1]. In this case the irreducible representation V_k with highest weight vector ω_k is the kernel of the contraction map $\wedge^k V \rightarrow \wedge^{k-2} V$ [16, Theorem 17.5].

3.1. The non-simply connected groups $\mathrm{SO}(n)$. As discussed in [16, Section 23.1] a general simple Lie group G has a finite abelian fundamental group, so it is a quotient \tilde{G}/Z where \tilde{G} is simply connected and Z is a finite abelian group. Any irreducible representation of G is an irreducible representation of \tilde{G} but not every irreducible representation of \tilde{G} descends to an irreducible representation of G . The set of weights of representations of G forms a sublattice of the set of weights of representations of the universal cover \tilde{G} of finite index. In this case there need not be an analogue of fundamental weights. That is, we cannot guarantee that the weight lattice of G has a basis $\omega_1, \dots, \omega_n$ such that every highest weight vector can be written as non-negative integral linear combination of the ω_i . For the group $\mathrm{SO}(2n+1)$ it is possible to find such weights, and thus we can define the band of an irreducible representation as above. For the group $\mathrm{SO}(2n)$ the weight lattice does not have fundamental system of weights. Despite this, we are still able to define the band of an irreducible representation as we see below.

3.1.1. Type B_n - the group $\mathrm{SO}(2n+1, \mathbb{R})$. The compact group $\mathrm{SO}(2n+1, \mathbb{R})$ is a compact form of the complex group $\mathrm{SO}(2n+1, \mathbb{C})$ with Lie algebra \mathfrak{so}_{2n+1} which has type B_n . The root system of type B_n has weight space generated by L_1, \dots, L_n . The positive simple roots are [16, Section 19.4]

$$\alpha_1 = L_1 - L_2, \dots, \alpha_{n-1} = L_{n-1} - L_n, \alpha_n = L_n$$

and the fundamental weights are

$$\omega_i = \sum_{j \leq i} L_j$$

for $i < n$ and $\omega_n = \frac{1}{2} \sum_{j=1}^n L_j$.

Note that because $\mathrm{SO}(2n+1)$ is not simply connected not every irreducible representation of \mathfrak{so}_{2n+1} gives rise to an irreducible representation of $\mathrm{SO}(2n+1)$. The representations of the Lie algebra \mathfrak{so}_{2n+1} are in bijective correspondence with the representations of the simply connected spin group $\mathrm{Spin}(2n+1)$ and the irreducible representations of $\mathrm{SO}(2n+1)$ are exactly the representations with highest weight vectors $a_1 \omega_1 + \dots + a_{n-1} \omega_{n-1} + a_n \omega_n$ where a_n is required to be even [16, Proposition 23.13(iii)]. In particular we can take the vectors

$\omega_k = L_1 + \dots + L_k$ for $k < n$ and $\omega'_n = 2\omega_n = L_1 + \dots + L_n$ to be a set of fundamental weights for the Lie group $\mathrm{SO}(2n+1)$.

The representations V_1, \dots, V_n associated to the fundamental weights are the exterior powers $V_1 = \wedge^1 V, \dots, V_n = \wedge^n V$ where V is the defining representation of $\mathrm{SO}(2n+1)$. Note that V has dimension $2n+1$ [16, Theorem 19.14].

Example 3.1. The group $\mathrm{SU}(2)$ is the universal cover of $\mathrm{SO}(3)$. Since these groups have rank one, the weights are just integers. For $\mathrm{SU}(2)$ the fundamental weight is $\omega = 1$, while for $\mathrm{SO}(3)$ the fundamental weight is $\omega = 2$.

3.2. Type D_n - the group $\mathrm{SO}(2n, \mathbb{R})$. The group $\mathrm{SO}(2n, \mathbb{R})$ is the compact form of $\mathrm{SO}(2n, \mathbb{C})$ whose Lie algebra \mathfrak{so}_{2n} has type D_n . Its weight space is generated by L_1, \dots, L_n . The positive simple roots are

$$\alpha_1 = L_1 - L_2, \dots, \alpha_{n-1} = L_{n-1} - L_n, \alpha_n = L_{n-1} + L_n$$

and the fundamental weights are [16, Section 19.2]

$$\omega_i = \sum_{j \leq i} L_j$$

for $i < n-1$,

$$\omega_{n-1} = \frac{1}{2} \sum_{j=1}^n L_j, \quad \text{and} \quad \omega_n = \frac{1}{2}(L_1 + \dots + L_{n-1} - L_n).$$

Once again, $\mathrm{SO}(2n)$ is not simply connected so not every irreducible representation of \mathfrak{so}_{2n} gives rise to a representation of the Lie group $\mathrm{SO}(2n)$. The irreducible representations of $\mathrm{SO}(2n)$ are precisely those with highest weight vector $\sum_{i=1}^n a_i \omega_i$ where $a_{n-1} + a_n$ is even [16, Proposition 23.13(iii)]. In this case every highest weight vector can be expressed non-uniquely as non-negative linear combination of the weights $\omega_1, \dots, \omega_{n-2}$ and

$$\omega'_{n-1} = \omega_{n-1} + \omega_n = L_1 + \dots + L_{n-1},$$

$$\omega'_n = 2\omega_n = L_1 + \dots + L_n,$$

$$\omega'_{n+1} = 2\omega_n = L_1 + \dots + L_{n-1} - L_n.$$

If λ is a highest weight vector for an irreducible representation of $\mathrm{SO}(2n)$ and we write

$$\lambda = b_1 \omega_1 + \dots + b_{n-2} \omega_{n-2} + b_{n-1} \omega'_{n-1} + b_n \omega'_n$$

then, although the non-negative integers b_i are not unique, the sum $\sum_{i=1}^n b_i$ is independent of the choice of b_i 's. For example, the weight $2L_1 + \dots + 2L_{n-1}$ has band 2 since it can be expressed as $2\omega'_{n-1}$ or as $\omega'_n + \omega'_{n+1}$. Hence we can define the band of λ to be $\sum_{i=1}^n b_i$. In particular, the $n+1$ irreducible representations $\omega_1, \dots, \omega_{n-2}, \omega'_{n-1}, \omega'_n$ all have band one.

By [16, Remark page 289], if $k \leq n-2$ then the representation V_k with highest weight vector ω_k is the exterior product $\wedge^k V$. Likewise, the representation V_{n-1} with highest weight ω'_{n-1} is exterior power $\wedge^{n-1} V$. Finally the exterior power $\wedge^n V$ is the sum of the representations V_n and V_{n+1} which have highest weights ω'_n and ω'_{n+1} respectively.

3.3. Reduction to representations of band one.

Proposition 3.2. *Let G be a compact Lie group whose irreducible representations can be banded. If $f \in L^2(G)$ and W is an irreducible representation of band $b > 1$ then the Fourier coefficient $F(f)(W)$ is determined from the bispectrum coefficients $a_2(f)(W_1 \otimes W_{b-1})$ and the Fourier coefficients of $F(f)(W_1)$, $F(f)(W_{b-1})$ for some W_1 of band one and some W_{b-1} of band $b - 1$, provided that these coefficients are invertible.*

Proof. Since the band of W is b , there are weight vectors $\omega_1, \dots, \omega_r$ of band one such that the highest weight vector of W is $b_1\omega_1 + \dots + b_r\omega_r$ with $b_i \in \mathbb{N}$ and $\sum_i b_i = b$. Since $b > 1$ we know that one of the b_i s is positive. Take W_1 to be the irreducible representation with highest weight ω_i and W_{b-1} to be the irreducible representation with highest weight $b_1\omega_1 + \dots + (b_i - 1)\omega_i + \dots + b_r\omega_r$. Since weights are additive on tensor products, $b_1\omega_1 + \dots + b_r\omega_r$ is the highest weight in the tensor product $W_1 \otimes W_{b-1}$. Hence W is a summand in this tensor. Therefore by Lemma 2.5 we conclude that $F(f)(W)$ is determined. \square

Corollary 3.3. *With the hypotheses on G as above, if $f \in L^2(G)$ is a function which is band-limited at band $b > 1$ and the Fourier coefficients of all representations of band $0 \leq i \leq \lceil b/2 \rceil$ are invertible, then all Fourier coefficients can be determined from the Fourier coefficients of band one and the bispectrum.*

Proof. Let W be an irreducible representation with highest weight vector $\lambda = b_1\omega_1 + \dots + b_r\omega_r$ where the b_i are non-negative integers and the ω_i have band one. We can decompose the vector $(b_1, \dots, b_r) = (c_1, \dots, c_r) + (d_1, \dots, d_r)$ with b_i, c_j non-negative integers and $\sum c_i, \sum d_i \leq \lceil b/2 \rceil$. If W_1 is the irreducible representation with highest weight $c_1\omega_1 + \dots + c_r\omega_r$ and W_2 is the irreducible representation with highest weight $d_1\omega_1 + \dots + d_r\omega_r$, then W appears as a summand in the tensor product $W_1 \otimes W_2$. Thus, if $F(f)(W_1)$ and $F(f)(W_2)$ are invertible then $F(f)(W)$ can be determined from $A_2 f(W_1 \otimes W_2)$ and $F(f)(W_1)$, $F(f)(W_2)$ regardless of whether $F(f)(W)$ is invertible. By Proposition 3.2 and induction $F(f)(W_1)$, $F(f)(W_2)$ can be determined from the Fourier coefficients of band one. \square

3.4. Reduction to the defining representation. The previous propositions only required that the irreducible representations of our compact Lie group G be banded. We now state a result specific to the classical groups $SU(n)$, $SO(2n + 1)$, $Sp(n)$, $SO(2n)$.

Theorem 3.4. *Let G be one of the classical compact Lie groups $SU(n)$, $SO(n)$, $Sp(2n)$ with defining representation V_1 . If the Fourier coefficients of the irreducible representations of band one are all invertible, then the Fourier coefficient $F(f)(V_\ell)$ of any band-one representation V_ℓ is determined by $F(f)(V_1)$ and the bispectrum matrices $a_2(f)(V_1 \otimes V_k)$ with V_k of band-one and $k < \ell$.*

Proof. The proof is based on a case-by-case analysis, but the overall structure is the same in each case and makes use of a simple lemma.

Lemma 3.5. *Let V be a representation of a compact Lie group G . Then $\wedge^{k+1}V$ appears as a G -invariant summand in $V \otimes \wedge^k V$.*

Proof. The action of G on V defines a homomorphism $G \rightarrow U(V)$ where $U(V)$ is the group of unitary transformations of V . In particular, it suffices to prove the lemma when G is the unitary group $U(d)$ where $d = \dim V$. The statement then follows from the fact that the map $V \otimes \wedge^k V \rightarrow \wedge^{k+1} V$, defined by

$$v \otimes (v_1 \wedge \dots \wedge v_k) \mapsto v \wedge v_1 \dots \wedge v_k$$

is surjective and commutes with the respective actions of the unitary group $U(V)$ on $V \otimes \wedge^k V$ and $\wedge^{k+1} V$. \square

3.4.1. *Type A_{n-1} - the group $SU(n)$.* As noted above the representation V_k with highest weight $\omega_k = L_1 + \dots + L_k$ is $\wedge^k V_1$ where V_1 is the defining representation.

By Lemma 2.5 and induction it suffices to show that V_{k+1} is a summand in the tensor product $V_1 \otimes V_k$ which follows from Lemma 3.5.

3.4.2. *Type B_n - the group $SO(2n + 1)$.* Here we have a fundamental system of weights $\omega_1, \dots, \omega_{n-1}, \omega'_n$ and the corresponding irreducible representations are the exterior product $\wedge^k V_1$ for $1 \leq k \leq n$. Once again by Lemma 2.5 we just need to show that V_{k+1} is a summand in $V_1 \otimes V_k$ which follows from Lemma 3.5.

3.4.3. *Type C_n - the group $Sp(2n)$.* Unlike the case of $SU(n)$ and $SO(2n + 1)$ the irreducibles associated to the fundamental weights are not exterior powers of the defining representation V_1 . However, [16, Theorem 17.5] states that if $k > 1$ then V_k is the kernel of the contraction map $\wedge^k V_1 \rightarrow \wedge^{k-2} V_1$. Hence $\wedge^k V_1 = V_k \oplus V_{k-2} \oplus \dots \oplus V_0$ if k is even and $\wedge^k V_1 = V_k \oplus V_{k-2} \oplus \dots \oplus V_1$ if k is odd. Since the Fourier coefficient of the trivial representation $V_0 = \wedge^0 V_1$ is known from the bispectrum by Proposition 2.2, we are able to inductively determine the Fourier coefficient $F(f)(V_k)$ from the bispectrum and knowledge of $F(f)(V_1)$ as follows: Assume by induction that we have determined the Fourier coefficients $F(f)(V_\ell)$ for $\ell \leq k$. Since $\wedge^k(V_1) = \oplus_\ell V_{k-2\ell}$ we know the Fourier $F(f)(\wedge^k V_1)$ by induction. Since $\wedge^{k+1} V_1$ appears as a summand in $V_1 \otimes \wedge^k V_1$ and V_{k+1} is a summand in $\wedge^{k+1} V$ we see that V_{k+1} is a summand in $V_1 \otimes \wedge^k V$. Since the Fourier coefficients $F(f)(\wedge^k V_1)$ is known by induction, it follows from Lemma 2.5 that $F(f)(V_{k+1})$ is determined by $F(f)(V_1)$ and the bispectrum coefficient $a_2(f)(V_1 \otimes \wedge^k V_1)$.

3.4.4. *Type D_n - the case of $SO(2n)$.* In this case we have $n + 1$ weights of band one $\omega_1, \dots, \omega_{n-2}, \omega'_{n-1}, \omega'_n, \omega'_{n+1}$ and associated representations V_1, \dots, V_{n+1} . As noted above, $V_k = \wedge^k V_1$ for $1 \leq k \leq n - 1$. Hence the same argument used in the case of $SO(2n + 1)$ implies that the bispectrum and $F(f)(V_1)$ determine $F(f)(V_k)$ if $k \leq n - 1$. However we also know that V_n and V_{n+1} are summands in $\wedge^n V_1$ so we can determine them from $F(f)(V_{n-1}), F(f)(V_1)$ and the bispectrum coefficient $F(f)(V_1 \otimes V_{n-1})$. \square

4. SHARP RESULTS FOR $SO(2n + 1), SU(n)$

For the groups $G = SU(n)$ and $G = SO(2n + 1)$ we prove that the G -orbit of a generic band limited function in $L^2(G)$ is determined by its bispectrum.

Theorem 4.1. (i) *If $f \in L^2(SU(n))$ is band limited with band $b \geq 1$ and all Fourier coefficients of irreducible representations whose band is at most $\lceil b/2 \rceil$ are invertible then the $SU(n)$ orbit of f is determined by its bispectrum.*

(ii) *If $f \in L^2(SO(2n + 1))$ is real valued and band limited with band $b \geq 1$ and all Fourier coefficients of irreducible representations whose band is at most $\lceil b/2 \rceil$ are invertible then the $SO(2n + 1)$ orbit of f is determined by its bispectrum.*

Proof. By Theorem 3.4 we know that Fourier coefficients of all irreducibles are determined by the Fourier coefficient $F(f)(V)$ of the defining representation. Moreover, by Propositions 2.1 and 2.2, we also know $F(f)(V)F(f)(V)^*$ so we know $F(f)(V)$ up to translation by an element of $U(V)$ if f is complex valued and $O(V)$ if f is real-valued.

Suppose that $f' \in L^2(\text{SU}(n))$ has the same bispectrum as f . Then we know that $F(f')(V) = uF(f)(V)$ for some $u \in U(V) = U(n)$. Our goal is to show that $u \in \text{SU}(n)$. Any element of $U(n)$ can be factored as $u = (\text{diag } e^{i\theta})r$ where $r \in \text{SU}(n)$. Replace f with the function whose Fourier coefficient of V is $rF(V)$ and has the same bispectrum. (Such a function exists because the bispectrum is invariant under the action of $\text{SU}(n)$ on $L^2(\text{SU}(n))$.) By doing so, we may reduce to the case that $F(f')(V) = e^{i\theta}F(f)(V)$ and has the same bispectrum. Our goal is to show that $\text{diag } e^{i\theta} \in \text{SU}(n)$ or, equivalently, that $e^{i\theta}$ is a n -th root of unity.

Since the bispectra of f and f' are equal we see that

$$\begin{aligned} a_2(f)(V \otimes V) &= [F(f)(V) \otimes F(f)(V)]F(f)(V \otimes V)^* \\ &= [F(f')(V) \otimes F(f')(V)]F(f')(V \otimes V)^* \\ &= e^{2i\theta}[F(f)(V) \otimes F(f)(V)]F(f')(V \otimes V)^* \end{aligned}$$

where the last equality follows from the fact that $\text{diag } e^{i\theta} \otimes \text{diag } e^{i\theta}$ is $e^{2i\theta}$ times the identity operator on $V \otimes V$.

It follows that for any irreducible W appearing in $V \otimes V$, we have that $F(f')(W) = e^{2i\theta}F(f)(W)$. In particular if $V_2 = \wedge^2 V$ then $F(f')(V_2) = e^{2i\theta}F(f)(V)$. Continuing this way we see that for the fundamental representations $V = V_1, \dots, V_{n-1}$, $F(f')(V_k) = e^{ik\theta}F(f)(V_k)$. On the other hand we know that the bispectrum uniquely determines the Fourier coefficient of the trivial representation V_0 , so we must have that $F(f')(V_0) = F(f)(V_0)$. However, V_0 appears as a summand in $V \otimes V_{n-1}$ so by our previous argument we see that $F(f')(V_0) = e^{in\theta}F(f)(V_0)$. Therefore $e^{in\theta} = 1$ as desired.

The proof for $\text{SO}(2n+1)$ is similar to the proof for $\text{SU}(n)$ but requires a slightly more complicated representation-theoretic argument. If $f \in L^2(\text{SO}(2n+1))$ is real-valued and f' is another real valued function with the same bispectrum then we know that $F(f')(V) = oF(f)(V)$ where $o \in O(2n+1)$. Now any element in $O(2n+1)$ can be written as $\pm r$ where $r \in \text{SO}(2n+1)$. We will prove the result by showing that if $o \in O(2n+1) \setminus \text{SO}(2n+1)$ we obtain a contradiction. Assuming that $o \notin \text{SO}(2n+1)$ we can reduce to the case that $F(f')(V) = -F(f)(V)$ and f', f have the same bispectrum. The tensor product $V \otimes V$ contains the fundamental representation $V_2 = \wedge^2 V$ as a summand, so we see that $F(f')(V_2) = (-1)^2 F(f)(V)$. Continuing this way we see that $F(f')(V_k) = (-1)^k F(f)(V)$ for $1 \leq k \leq n$. Since $V_n = \wedge^n V$ we know by Lemma 3.5 that $V \otimes V_n$ contains a copy of $\wedge^{n+1} V$. Since $\dim V = 2n+1$, the exterior products $\wedge^n V$ and $\wedge^{n+1} V$ are dual representations. However, we also know that as $\text{SO}(2n+1)$ representations, $\wedge^k V$ is self-dual for $k \leq n$. Hence $V \otimes V_n$ contains a copy of V_n . This implies that $F(f')(V_n) = -F(f)(V_n)$, which is a contradiction. \square

4.1. $\text{SO}(3)$ and unprojected cryo-EM. Let $L^2(\mathbb{R}^3)$ be Hilbert space of complex valued L^2 functions on \mathbb{R}^3 . The action of $\text{SO}(3)$ on \mathbb{R}^3 induces a corresponding action on $L^2(\mathbb{R}^3)$, which we view as an infinite-dimensional representation of $\text{SO}(3)$. In cryo-EM we are interested in the action of $\text{SO}(3)$ on the subspace of $L^2(\mathbb{R}^3)$ corresponding to the Fourier transforms of real valued functions on \mathbb{R}^3 , representing the Coulomb potential of an unknown molecular structure.

Using spherical coordinates (ρ, θ, ϕ) , we consider a finite dimensional approximation of $L^2(\mathbb{R}^3)$ by discretizing $f(\rho, \theta, \phi)$ with R samples r_1, \dots, r_R , of the radial coordinates and band limiting the corresponding spherical functions $f(r_i, \theta, \phi)$. This is a standard assumption in the cryo-EM literature, see for example [4]. Mathematically, this means that we

approximate the infinite-dimensional representation $L^2(\mathbb{R}^3)$ with the finite dimensional representation $V = (\oplus_{\ell=0}^L V_\ell)^R$, where L is the band limit, and V_ℓ is the $(2\ell + 1)$ -dimensional irreducible representation of $\text{SO}(3)$, corresponding to harmonic polynomials of frequency ℓ . An orthonormal basis for V_ℓ is the set of spherical harmonic polynomials $\{Y_\ell^m(\theta, \phi)\}_{m=-\ell}^\ell$. We use the notation $Y_\ell^m[r]$ to consider the corresponding spherical harmonic as a basis vector for functions on the r -th spherical shell. The dimension of this representation is $R(L^2 + 2L + 1)$.

Viewing an element of V as a radially discretized function on \mathbb{R}^3 , we can view $f \in V$ as an R -tuple

$$f = (f[1], \dots, f[R]),$$

where $f[r] \in L^2(S^2)$ is an L -bandlimited function. Each $f[r]$ can be expanded in terms of the basis functions $Y_\ell^m(\theta, \varphi)$ as follows

$$(4.1) \quad f[r] = \sum_{\ell=0}^L \sum_{m=-\ell}^{\ell} A_\ell^m[r] Y_\ell^m.$$

Therefore, the problem of determining a structure reduces to determining the unknown coefficients $A_\ell^m[r]$ in (4.1).

Note that when f is the Fourier transform of a real valued function, the coefficients $A_\ell^m[r]$ are real for even ℓ and purely imaginary for odd ℓ [11].

For the case of $\text{SO}(3)$ we can combine our results and obtain the following corollary.

Corollary 4.2. *If $L \geq 3$ and any $R \geq L + 2$ the generic orbit in the $\text{SO}(3)$ representation $V = (\oplus_{\ell=0}^L V_\ell)^R$ can be recovered from the third moment.*

Proof. By Proposition 2.3 it suffices to prove the corollary when $R = L + 1$. Consider the $\text{SO}(3)$ -module

$$W = V_0 \oplus V_1^3 \oplus \dots \oplus V_{\lfloor L/2 \rfloor}^{2\lfloor L/2 \rfloor + 1} \oplus V_{\lfloor L/2 \rfloor + 1}^{L+2} \oplus \dots \oplus V_L^{L+2}.$$

Since $L+2 \geq \lfloor L/2 \rfloor + 1$, we can once more invoke Proposition 2.3 and prove the result for the representation W . We view W as an $\text{SO}(3)$ -submodule of the vector space of L -bandlimited functions in $L^2(\text{SO}(3))$ since the latter representation is isomorphic to $\oplus_{\ell=0}^L V_\ell^{2L+1}$. The generic element of W viewed as a submodule of $\oplus_{\ell=0}^L V_\ell^{2L+1}$ has invertible Fourier coefficients up to band $\lfloor L/2 \rfloor$. Therefore, by Corollary 3.3 if $f \in W$ then the Fourier coefficients $F(f)(V_\ell)$ are determined by the bispectrum and the single Fourier coefficient $F(f)(V_1)$ of band one. By Proposition 2.2 the third moment determines the second moment and by Proposition 2.1 the second moment determines $A_1(f)A_1(f)^T$ where $A_1(f)$ is the full-rank real 3×3 matrix $(-iA_1^m[r])_{-1 \leq m \leq 1, 1 \leq r \leq 3}$. (Note that the coefficients $A_1^m[r]$ are purely imaginary so we multiply by $-i$ to obtain a real valued matrix.) In particular the matrix $A_1(f)$ (which we can identify with the Fourier coefficient in a suitable basis) is determined up to multiplication by an element $O(3)$. By Theorem 4.1 $A_1(f)$ is determined up to rotation by an element of $\text{SO}(3)$. Hence the orbit of the generic vector $f \in W$ is determined from its third moment. \square

4.2. Discussion and further directions. In [2] it is conjectured that for any L the third moment separates generic orbits provided $R \geq 3$. This conjecture was verified for $1 \leq L \leq 15$ using techniques from computational commutative algebra and a computational algebra algorithm was presented for recovering the orbit using frequency marching. In a recent paper [23], Liu and Moitra view $\oplus_{\ell=0}^L V_\ell$ as the space of band-limited functions on S^2 and use the method of smoothed analysis to give a probabilistic algorithm for approximating an orbit in $\oplus_{\ell=0}^L V_\ell$. Note that Corollary 4.2 is weaker than the conjectured bounds of [2] in

that we require the multiplicities of the irreducible representations to grow with the band. An interesting direction for further work is to refine the the methods used here to determine whether the third moment carries enough information to separate orbits when the irreducible representations have constant multiplicity which is independent of the band limit.

Another interesting avenue of investigation is the case of finite groups. It is known [2] that the third moment separates generic orbits in any representation containing the regular representation. However, there are essentially no known non-trivial examples of smaller representations of finite groups where the third moment separates generic orbits.

ACKNOWLEDGMENT

We thank Tamir Bendory for helpful discussions. The work of D.E. was supported by the BSF grant no. 2020159 and NSF-DMS 2205626. The work of M.S. was partially supported by a Discovery Grant from the National Science and Engineering Research Council of Canada as well as a Mathematics Faculty Research Chair from the University of Waterloo.

REFERENCES

- [1] Emmanuel Abbe, Tamir Bendory, William Leeb, João M Pereira, Nir Sharon, and Amit Singer. Multireference alignment is easier with an aperiodic translation distribution. *IEEE Transactions on Information Theory*, 65(6):3565–3584, 2018.
- [2] Afonso S Bandeira, Ben Blum-Smith, Joe Kileel, Amelia Perry, Jonathan Weed, and Alexander S Wein. Estimation under group actions: recovering orbits from invariants. *arXiv preprint arXiv:1712.10163*, 2017.
- [3] Afonso S Bandeira, Moses Charikar, Amit Singer, and Andy Zhu. Multireference alignment using semi-definite programming. In *Proceedings of the 5th conference on Innovations in theoretical computer science*, pages 459–470, 2014.
- [4] Afonso S Bandeira, Yutong Chen, Roy R Lederman, and Amit Singer. Non-unique games over compact groups and orientation estimation in cryo-EM. *Inverse Problems*, 36(6):064002, 2020.
- [5] Afonso S Bandeira, Jonathan Niles-Weed, and Philippe Rigollet. Optimal rates of estimation for multi-reference alignment. *Mathematical Statistics and Learning*, 2(1):25–75, 2020.
- [6] Tamir Bendory, Alberto Bartesaghi, and Amit Singer. Single-particle cryo-electron microscopy: Mathematical theory, computational challenges, and opportunities. *IEEE signal processing magazine*, 37(2):58–76, 2020.
- [7] Tamir Bendory, Nicolas Boumal, William Leeb, Eitan Levin, and Amit Singer. Toward single particle reconstruction without particle picking: Breaking the detection limit. *arXiv preprint arXiv:1810.00226*, 2018.
- [8] Tamir Bendory, Nicolas Boumal, Chao Ma, Zhizhen Zhao, and Amit Singer. Bispectrum inversion with application to multireference alignment. *IEEE Transactions on Signal Processing*, 66(4):1037–1050, 2017.
- [9] Tamir Bendory and Dan Edidin. The sample complexity of sparse multi-reference alignment and single-particle cryo-electron microscopy. *arXiv preprint, arXiv:2210.15727*, 2022.
- [10] Tamir Bendory, Dan Edidin, William Leeb, and Nir Sharon. Dihedral multi-reference alignment. *IEEE Transactions on Information Theory*, 68(5):3489–3499, 2022.
- [11] Tejal Bhamre, Teng Zhang, and Amit Singer. Orthogonal matrix retrieval in cryo-electron microscopy. In *2015 IEEE 12th International Symposium on Biomedical Imaging (ISBI)*, pages 1048–1052. IEEE, 2015.
- [12] Ben Blum-Smith and Soledad Villar. Machine learning and invariant theory. *arXiv preprint arXiv:2209.14991*, 2023.
- [13] Theodor Bröcker and Tammo tom Dieck. *Representations of compact Lie groups*, volume 98 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. Translated from the German manuscript, Corrected reprint of the 1985 translation.

- [14] Taco S. Cohen and Max Welling. Group equivariant convolutional networks. *arXiv preprint arXiv:1602.07576*, 2016.
- [15] Nadav Dym and Haggai Maron. On the universality of rotation equivariant point cloud networks. *arXiv preprint, arXiv:2010.02449*, 2020.
- [16] William Fulton and Joe Harris. *Representation theory*, volume 129 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991. A first course, Readings in Mathematics.
- [17] Brian Hall. *Lie groups, Lie algebras, and representations*, volume 222 of *Graduate Texts in Mathematics*. Springer, Cham, second edition, 2015. An elementary introduction.
- [18] Edwin Hewitt and Kenneth A. Ross. *Abstract harmonic analysis. Vol. II: Structure and analysis for compact groups. Analysis on locally compact Abelian groups*. Die Grundlehren der mathematischen Wissenschaften, Band 152. Springer-Verlag, New York-Berlin, 1970.
- [19] Noam Janco and Tamir Bendory. An accelerated expectation-maximization algorithm for multi-reference alignment. *IEEE Transactions on Signal Processing*, 70:3237–3248, 2022.
- [20] Ramakrishna Kakarala. The bispectrum as a source of phase-sensitive invariants for Fourier descriptors: a group-theoretic approach. *J. Math. Imaging Vision*, 44(3):341–353, 2012.
- [21] Risi Kondor, Zhen Lin, and Shubhendu Trivedi. Clebsch-gordan nets: a fully fourier space spherical convolutional neural network. *arXiv preprint arXiv:1806.09231*, 2018.
- [22] Lek-Heng Lim and Bradley J. Nelson. What is an equivariant neural network? *arXiv preprint arXiv:2205.07362*, 2022.
- [23] Allen Liu and Ankur Moitra. Algorithms from invariants: Smoothed analysis of orbit recovery over $SO(3)$. *arXiv preprint arXiv:2106.02680*, 2021.
- [24] Chao Ma, Tamir Bendory, Nicolas Boumal, Fred Sigworth, and Amit Singer. Heterogeneous multireference alignment for images with application to 2D classification in single particle reconstruction. *IEEE Transactions on Image Processing*, 29:1699–1710, 2019.
- [25] Amelia Perry, Jonathan Weed, Afonso S Bandeira, Philippe Rigollet, and Amit Singer. The sample complexity of multireference alignment. *SIAM Journal on Mathematics of Data Science*, 1(3):497–517, 2019.
- [26] Amit Singer. Angular synchronization by eigenvectors and semidefinite programming. *Applied and Computational Harmonic Analysis*, 30(1):20–36, 2011.
- [27] Fethi Smach, Cedric Lemaître, Jean-Paul Gauthier, Johel Miteran, and Mohamed Atri. Generalized Fourier descriptors with applications to objects recognition in SVM context. *J. Math. Imaging Vision*, 30(1):43–71, 2008.

APPENDIX A. REPRESENTATION THEORY

A.1. Terminology on representations. Let G be a group. A representation of G is a homomorphism, $G \xrightarrow{\pi} \text{GL}(V)$, where V is a vector space over a field and $\text{GL}(V)$ is the group of invertible linear transformations $V \rightarrow V$. Given a representation of a group G , we can define an action of G on V by $g \cdot v = \pi(g)v$. Since $\pi(g)$ is a linear transformation, the action of G is necessarily linear, meaning that for any vectors v_1, v_2 and scalars $\lambda, \mu \in \mathbb{C}$ $g \cdot (\lambda v_1 + \mu v_2) = \lambda(g \cdot v_1) + \mu(g \cdot v_2)$. Conversely, given a linear action of G on a vector space V , we obtain a homomorphism $G \rightarrow \text{GL}(V)$, $g \mapsto T_g$, where $T_g: V \rightarrow V$ is the linear transformation $T_g(v) = (g \cdot v)$. Thus, giving a representation of G is equivalent to giving a linear action of G on a vector space V . Given this equivalence, we will follow standard terminology and refer to a vector space V with a linear action of G as a *representation of G* .

A representation V of G is *finite dimensional* if $\dim V < \infty$. In this case, a choice of basis for V identifies $\text{GL}(V) = \text{GL}(N)$, where $N = \dim V$. If V is a complex vector space with Hermitian inner product $\langle \cdot, \cdot \rangle$ on V , we say that a representation is *unitary* if for any two vectors $v_1, v_2 \in V$ $\langle g \cdot v_1, g \cdot v_2 \rangle = \langle v_1, v_2 \rangle$. Likewise if V is a real vector space with inner product $\langle \cdot, \cdot \rangle$ then we say that the representation is *orthogonal* if the action of G preserves the inner product. If we choose an orthonormal basis for V , then the representation of

G is unitary (resp. orthogonal) if and only if the image of G under the homomorphism $G \rightarrow \text{GL}(N)$ lies in the subgroup $U(N)$ (resp. $O(N)$) of unitary (resp. orthogonal) matrices.

A representation V of a group G is *irreducible* if V contains no non-zero proper G -invariant subspaces.

A.2. Representations of compact groups. Any compact group G has a G -invariant measure called a Haar measure. The Haar measure dg is typically normalized so that $\int_G dg = 1$. If V is a finite-dimensional representation of a compact group and (\cdot, \cdot) is any Hermitian inner product, then the inner product $\langle \cdot, \cdot \rangle$ defined by the formula $\langle v_1, v_2 \rangle = \int_G (g \cdot v_1, g \cdot v_2) dg$ is G -invariant. As a consequence we obtain the following fact.

Proposition A.1. *Every finite dimensional representation of a compact group is unitary.*

Using the invariant inner product we can then obtain the following decomposition theorem for finite dimensional representations of compact group.

Proposition A.2. *Any finite dimensional representation of a compact group decomposes into a direct sum of irreducible representations.*

If V is a representation, then $V^G = \{v \in V | g \cdot v = v\}$ is a subspace which is called the subspace of invariants.

A.3. Schur's Lemma. A key property of irreducible unitary representations is Schur's Lemma. Recall that a linear transformation Φ is G -invariant if $g \cdot \Phi v = \Phi g \cdot v$.

Lemma A.3. *Let $\Phi: V_1 \rightarrow V_2$ be a G -invariant linear transformation of finite dimensional irreducible representations of a group G (not necessarily compact). Then, Φ is either zero or an isomorphism. Moreover, if V is a finite dimensional irreducible unitary representation of a group G then any G -invariant linear transformation $\phi: V \rightarrow V$ is multiplication by a scalar.*

A.4. Dual, Hom and tensor products of representations. If V_1 and V_2 are representations of a group G , then the vector space $\text{Hom}(V_1, V_2)$ of linear transformations $V_1 \rightarrow V_2$ has a natural linear action of G given by the formula $(g \cdot A)(v_1) = g \cdot A(g^{-1}v_1)$. In particular, if V is a representation of G , then $V^* = \text{Hom}(V, \mathbb{C})$ has a natural action of G given by the formula $(g \cdot f)(v) = f(g^{-1}v)$.

A choice of inner product on V determines an identification of vector spaces $V = V^*$, given by the formula $v \mapsto \langle \cdot, v \rangle$. If V is a unitary representation of G then with the identification of $V = V^*$ the dual action of G on V is given by the formula $g \cdot_* v = \bar{g} \cdot v$. Likewise, if V_1 and V_2 are two representations then we can define an action of G on $V_1 \otimes V_2$ by the formula $g \cdot (v_1 \otimes v_2) = (g \cdot v_1) \otimes (g \cdot v_2)$.

Given two representations spaces V_1, V_2 there is an isomorphism of representations $V_1 \otimes V_2^* \rightarrow \text{Hom}(V_2, V_1)$ given by the formula $v_1 \otimes f_2 \mapsto \phi$, where the linear transform $\phi: V_2 \rightarrow V_1$ is defined by the formula $\phi(v_2) = f_2(v_2)v_1$. In particular, we can identify $V \otimes V^*$ with $\text{Hom}(V, V)$.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211
Email address: edidind@missouri.edu

UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO, ONTARIO, CANADA
 N2L 3G1
Email address: msatrian@uwaterloo.ca