AMBARZUMYAN-TYPE THEOREM FOR THE STURM-LIOUVILLE OPERATOR ON THE LASSO GRAPH

FENG WANG AND CHUAN-FU YANG

Abstract. We consider the Sturm-Liouville operator on the lasso graph with a segment and a loop joined at one point, which has arbitrary length. The Ambarzumyan's theorem for the operator is proved, which says that if the eigenvalues of the operator coincide with those of the zero potential, then the potential is zero.

1. introduction

Inverse spectral problems consist in recovering the coefficients of an operator from their spectral characteristics. The first inverse spectral result on Sturm-Liouville operators is given by Ambarzumyan [\[1\]](#page-8-0), which describes the following theorem:

If q is a continuous real-valued function on [0, 1], and $\{n^2\pi^2 : n = 0, 1, 2, ...\}$ is the set of eigenvalues of the boundary value problem

$$
-y'' + q(x)y = \lambda y, \ x \in (0,1), \quad y'(0) = y'(1) = 0,
$$

then $q(x) \equiv 0$ on [0, 1].

This theorem is called Ambarzumyan's theorem, and has been generalized in many directions (see [4-6, 9-16, 18-19] and other papers). Here we mention Ambarzumyantype theorems on star graphs [\[13,](#page-9-0) [14\]](#page-9-1), Ambarzumyan-type theorems on trees [\[4,](#page-8-1) [12\]](#page-9-2), and Ambarzumyan-type theorems on graphs with cycles [\[11,](#page-9-3) [15\]](#page-9-4).

Differential operators on graphs often appear in natural sciences and engineering (see [\[2,](#page-8-2) [3\]](#page-8-3) and the references therein). Such operators can be used to model the motion of quantum particles confined to certain low dimensional structures. In recent years, Ambarzumyan-type theorems on graphs have attracted the attention of many researchers. Some Ambarzumyan-type theorems for the Sturm-Liouville operator on graphs have been achieved in the literatures mentioned above and other works. Most of the works in this direction are devoted to the graphs with equal length edges. It is more difficult to study the Ambarzumyan-type theorems on graphs with unequal length edges. Nevertheless, it is worth noting that C. K. Law and E. Yanagida [\[12\]](#page-9-2) studied the Ambarzumyan-type theorems on trees with unequal length edges.

Date: June 2, 2023.

²⁰⁰⁰ Mathematics Subject Classification. 34A55; 34B24; 47E05.

Key words and phrases. Sturm-Liouville operator, Ambarzumyan-type theorem, Lasso graph.

Figure 1. Lasso graph G

In addition, we have also noticed that an Ambarzumian-type theorem for the Sturm-Liouville operator with a bounded real potential on arbitrary compact graphs is found by Davies [\[6\]](#page-8-4). In this paper, using the method different from [6], we obtain an Ambarzumiantype theorem for the Sturm-Liouville operator with a square integrable real potential on the lasso graph with arbitrary edge lengths. Our approach is based on the Hadamard's factorization theorem and the variational principle, which is simpler and can be applied to arbitrary compact graphs.

In this paper we consider the lasso graph G (see Figure [1\)](#page-1-0). The edge e_1 is a boundary edge of length l_1 , the edge e_2 is a loop of length l_2 . Each edge e_j is parameterized by the parameter $x_i \in [0, l_i]$. The value $x_1 = 0$ corresponds to the boundary vertex, and $x_1 = l_1$ corresponds to the internal vertex. For the loop e_2 , both ends $x_2 = 0$ and $x_2 = l_2$ correspond to the internal vertex.

A function y on G may be represented as a vector function $y = \{y_i\}_{i=1,2}$, where the function $y_i(x_j)$, $x_j \in [0, l_j]$, is defined on the edge e_j . Consider the following Sturm-Liouville equations on G:

$$
\varphi(y_j) := -y_j''(x_j) + q_j(x_j)y_j(x_j) = \lambda y_j(x_j), \quad x_j \in [0, l_j], \quad j = 1, 2,
$$
 (1.1)

where q_j , $j = 1, 2$, are real-valued functions from $L^2[0, l_j]$, λ is the spectral parameter, the functions $y_j, y'_j, j = 1, 2$, are absolutely continuous on $[0, l_j]$ and satisfy the following matching conditions in the internal vertex:

$$
y_1(l_1) = y_2(l_2) = y_2(0),
$$
 $y'_1(l_1) + y'_2(l_2) - y'_2(0) = 0.$ (1.2)

Matching conditions (1.2) are called the standard conditions. In electrical circuits, (1.2) expresses Kirchhoff's law; in elastic string network, it expresses the balance of tension, and so on.

Denote $q = \{q_j\}_{j=1,2}$ called the potential on G. Let us consider the boundary value problem $L(q)$ on G for equation [\(1.1\)](#page-1-2) with the standard matching conditions [\(1.2\)](#page-1-1) and with Neumann boundary condition in the boundary vertex:

$$
y_1'(0) = 0.\t(1.3)
$$

It is obvious that the spectrum $\sigma(L(q))$ of the boundary value problem $L(q)$ is a discrete real sequence, bounded from below, diverging to $+\infty$. Let $\sigma(L(q)) = {\lambda_n(q)}_{n=0}^{\infty}$, which can be arranged in an ascending order as (counting with their multiplicities)

$$
\lambda_0(q) \leq \lambda_1(q) \leq \cdots \leq \lambda_n(q) \leq \cdots \longrightarrow +\infty.
$$

The main result in this paper is as follows.

Theorem 1.1. If $\lambda_n(q) = \lambda_n(0)$ for all $n = 0, 1, 2 \cdots$, then $q_i(x_i) = 0$ a.e. on $[0, l_i]$, $j = 1, 2.$

2. Analysis of the characteristic function

In this section we analyze the characteristic function of the boundary value problem $L(q)$, which plays a key role in the proof of Theorem [1.1.](#page-2-0)

Let $C_j(x_j, \lambda)$, $S_j(x_j, \lambda)$, $j = 1, 2$, be the solutions of equation [\(1.1\)](#page-1-2) on the edge e_j with the initial conditions

$$
C_j(0,\lambda) = S'_j(0,\lambda) = 1,
$$
 $C'_j(0,\lambda) = S_j(0,\lambda) = 0.$

For each fixed $x_j \in [0, l_j]$, the functions $C_j^{(\nu)}$ $g_j^{(\nu)}(x_j,\lambda),\ S_j^{(\nu)}$ $j^{(\nu)}(x_j,\lambda), j=1,2, \nu=0,1$, are entire in λ of order $\frac{1}{2}$.

Then the solutions of equation (1.1) which satisfy condition (1.3) are represented as

$$
\begin{cases}\ny_1(x_1, \lambda) = A_1(\lambda)C_1(x_1, \lambda), \\
y_2(x_2, \lambda) = A_2(\lambda)C_2(x_2, \lambda) + B_2(\lambda)S_2(x_2, \lambda),\n\end{cases} (2.1)
$$

where $A_1(\lambda)$, $A_2(\lambda)$ and $B_2(\lambda)$ are only dependent on λ . Substituting [\(2.1\)](#page-2-1) into match-ing conditions [\(1.2\)](#page-1-1) we obtain a linear algebraic system s with respect to $A_1(\lambda)$, $A_2(\lambda)$ and $B_2(\lambda)$. The determinant $\Delta(\lambda)$ of s has the form

$$
\Delta(\lambda) = C_1(l_1, \lambda) \Big(C_2(l_2, \lambda) + S_2'(l_2, \lambda) - 2 \Big) + C_1'(l_1, \lambda) S_2(l_2, \lambda).
$$
 (2.2)

The function $\Delta(\lambda)$ is entire in λ of order $\frac{1}{2}$, and its zeros coincide with the eigenvalues of the boundary value problem $L(q)$. The function $\Delta(\lambda)$ is called the characteristic function for the boundary value problems $L(q)$.

Let $\lambda = \rho^2$, $\tau = Im\rho$, then it follows from [\[7\]](#page-8-5) that the following asymptotic formulas hold uniformly in $x_j \in [0, l_j]$:

$$
\begin{cases}\nS_j(x_j, \lambda) = \frac{\sin \rho x_j}{\rho} - \frac{\cos \rho x_j}{2\rho^2} \int_0^{x_j} q_j(t)dt + o\left(\frac{e^{|\tau|x_j|}}{\rho^2}\right), \\
S'_j(x_j, \lambda) = \cos \rho x_j + \frac{\sin \rho x_j}{2\rho} \int_0^{x_j} q_j(t)dt + o\left(\frac{e^{|\tau|x_j|}}{\rho}\right), \\
C_j(x_j, \lambda) = \cos \rho x_j + \frac{\sin \rho x_j}{2\rho} \int_0^{x_j} q_j(t)dt + o\left(\frac{e^{|\tau|x_j|}}{\rho}\right), \\
C'_j(x_j, \lambda) = -\rho \sin \rho x_j + \frac{\cos \rho x_j}{2} \int_0^{x_j} q_j(t)dt + o\left(e^{|\tau|x_j|}\right),\n\end{cases} (2.3)
$$

as $|\rho| \to \infty$. According to [\(2.2\)](#page-2-2) and [\(2.3\)](#page-2-3) we have

$$
\Delta(\lambda) = \Delta_0(\lambda) + \frac{2(\sin \rho l_1)(\cos \rho l_2 - 1) + (\cos \rho l_1)(\sin \rho l_2)}{2\rho} [q_1]
$$

$$
+ \frac{2(\cos \rho l_1)(\sin \rho l_2) + (\sin \rho l_1)(\cos \rho l_2)}{2\rho} [q_2]
$$

$$
+ o\left(\frac{e^{|\tau|(l_1 + l_2)}}{\rho}\right), \qquad |\rho| \to \infty, \tag{2.4}
$$

where

$$
\Delta_0(\lambda) = 2(\cos \rho l_1)(\cos \rho l_2 - 1) - (\sin \rho l_1)(\sin \rho l_2),
$$
\n(2.5)

$$
[q_j] = \int_0^{l_j} q_j(t)dt, \qquad j = 1, 2. \tag{2.6}
$$

The function $\Delta_0(\lambda)$ is entire in λ of order $\frac{1}{2}$, and the characteristic function for the boundary value problems $L(0)$ with zero potential.

Let us show that the specification of the spectrum $\sigma(L(q)) = {\lambda_n(q)}_{n=0}^{\infty}$ uniquely determines the characteristic function $\Delta(\lambda)$. To this end, we consider together with $L(q)$ the boundary value problems $L(\tilde{q})$ of the same form but with different $\tilde{q} = {\tilde{q}}_j\}_{j=1,2}$. We agree that if a certain symbol β denotes an object related to $L(q)$, then $\tilde{\beta}$ will denote the analogous object related to $L(\tilde{q})$.

Lemma 2.1. If $\lambda_n(q) = \lambda_n(\tilde{q})$ for all $n = 0, 1, 2 \cdots$, then $\Delta(\lambda) = \tilde{\Delta}(\lambda)$.

Proof. Denote

$$
\lambda_n^1(0) = \begin{cases} \lambda_n(0) & \text{if } \lambda_n(0) \neq 0, \\ 1 & \text{if } \lambda_n(0) = 0, \end{cases}
$$

where $\lambda_n(0)$, $n = 0, 1, 2 \cdots$, are eigenvalues of the boundary value problems $L(0)$. By Hadamard's factorization theorem, we have

$$
\Delta_0(\lambda) = C_0 \prod_{n=0}^{\infty} \frac{\lambda_n(0) - \lambda}{\lambda_n^1(0)},
$$
\n(2.7)

where

$$
C_0 = \frac{(-1)^m}{m!} \left(\frac{\partial^m}{\partial \lambda^m} \Delta_0(\lambda) \right) |_{\lambda = 0},
$$

and $m \geq 0$ is the multiplicity of the zero eigenvalue of $L(0)$. Note that the infinite product in [\(2.7\)](#page-3-0) is absolutely convergent (e.g. see Theorem B.2 in [\[8\]](#page-8-6)).

Denote

$$
\lambda_n^1(q) = \begin{cases} \lambda_n(q) & \text{if } \lambda_n(q) \neq 0, \\ 1 & \text{if } \lambda_n(q) = 0. \end{cases} \tag{2.8}
$$

Using Hadamard's factorization theorem again, we get

$$
\Delta(\lambda) = C \prod_{n=0}^{\infty} \frac{\lambda_n(q) - \lambda}{\lambda_n^1(q)},
$$
\n(2.9)

where $C \neq 0$ is a constant. Combining the equations [\(2.7\)](#page-3-0) and [\(2.9\)](#page-4-0) yields

$$
\frac{\Delta(\lambda)}{\Delta_0(\lambda)} = \frac{C}{C_0} \prod_{n=0}^{\infty} \frac{\lambda_n^1(0)}{\lambda_n^1(q)} \prod_{n=0}^{\infty} \left(1 + \frac{\lambda_n(q) - \lambda_n(0)}{\lambda_n(0) - \lambda} \right).
$$
\n(2.10)

Using properties of the characteristic functions and the eigenvalues one gets for large negative λ (see [\[17\]](#page-9-5), Sec. 2.3 and 2.4):

$$
\lim_{\lambda \to -\infty} \frac{\Delta(\lambda)}{\Delta_0(\lambda)} = 1.
$$

Consequently by [\(2.10\)](#page-4-1), we obtain

$$
C = C_0 \left[\lim_{\lambda \to -\infty} \prod_{n=0}^{\infty} \frac{\lambda_n^1(0)}{\lambda_n^1(q)} \prod_{n=0}^{\infty} \left(1 + \frac{\lambda_n(q) - \lambda_n(0)}{\lambda_n(0) - \lambda} \right) \right]^{-1}.
$$
 (2.11)

Similarly,

$$
\widetilde{\Delta}(\lambda) = \widetilde{C} \prod_{n=0}^{\infty} \frac{\lambda_n(\widetilde{q}) - \lambda}{\lambda_n^1(\widetilde{q})},
$$
\n(2.12)

where

$$
\widetilde{C} = C_0 \left[\lim_{\lambda \to -\infty} \prod_{n=0}^{\infty} \frac{\lambda_n^1(0)}{\lambda_n^1(\widetilde{q})} \prod_{n=0}^{\infty} \left(1 + \frac{\lambda_n(\widetilde{q}) - \lambda_n(0)}{\lambda_n(0) - \lambda} \right) \right]^{-1},
$$
\n(2.13)

and

$$
\lambda_n^1(\tilde{q}) = \begin{cases} \lambda_n(\tilde{q}) & \text{if } \lambda_n(\tilde{q}) \neq 0, \\ 1 & \text{if } \lambda_n(\tilde{q}) = 0. \end{cases} \tag{2.14}
$$

According to [\(2.8\)](#page-3-1), [\(2.11\)](#page-4-2), [\(2.13\)](#page-4-3) and [\(2.14\)](#page-4-4), together with assumptions of the lemma, we have $\lambda_n^1(q) = \lambda_n^1(\tilde{q})$ and $C = \tilde{C}$. Thus, it follows from [\(2.9\)](#page-4-0) and [\(2.12\)](#page-4-5) that $\Delta(\lambda) = \tilde{\lambda}(\lambda)$ $\widetilde{\Delta}(\lambda)$.

From Lemma [2.1,](#page-3-2) we can immediately get the following corollary.

Corollary 2.2. If $\lambda_n(q) = \lambda_n(0)$ for all $n = 0, 1, 2 \cdots$, then $\Delta(\lambda) = \Delta_0(\lambda)$.

6 F. WANG AND C. F. YANG

3. Proof of Theorem 1.1

This section is devoted to the proof of Theorem [1.1.](#page-2-0) The following lemma plays an important role in the proof of Theorem [1.1.](#page-2-0)

Lemma 3.1. If $\Delta(\lambda) = \Delta_0(\lambda)$, then $[q_1] = [q_2] = 0$.

Proof. (1) When $l_1 = l_2 = l$, from [\(2.4\)](#page-3-3), together with $\Delta(\lambda) = \Delta_0(\lambda)$, we have

$$
\frac{(\sin \rho l)(3\cos \rho l-2)}{2\rho}[q_1] + \frac{3(\sin \rho l)(\cos \rho l)}{2\rho}[q_2] = o\left(\frac{e^{2|\tau|l}}{\rho}\right), \quad |\rho| \to \infty.
$$

In the above estimate, taking $\rho = \frac{(n+\frac{1}{2})\pi}{l}$ $\frac{\frac{1}{2}\pi}{l}$ to get $[q_1] = 0$ and $\rho = \frac{2n\pi + \arccos \frac{2}{3}}{l}$ to get $[q_2] = 0.$

(2) When $l_1 \neq l_2$, it can be proved in two cases.

(i) Let $\frac{l_2}{l_1}$ be an irrational number. From [\(2.4\)](#page-3-3), together with $\Delta(\lambda) = \Delta_0(\lambda)$, we have

$$
\frac{2(\sin \rho l_1)(\cos \rho l_2 - 1) + (\cos \rho l_1)(\sin \rho l_2)}{2\rho} [q_1]
$$

$$
+ \frac{2(\cos \rho l_1)(\sin \rho l_2) + (\sin \rho l_1)(\cos \rho l_2)}{2\rho} [q_2]
$$

$$
= o\left(\frac{e^{|\tau|(l_1+l_2)}}{\rho}\right), \quad |\rho| \to \infty. \tag{3.1}
$$

Taking $\rho = \frac{2n\pi}{l_1}$ $\frac{n\pi}{l_1}$ in the estimate [\(3.1\)](#page-5-0), we obtain

$$
\sin\left(\frac{l_2}{l_1}2n\pi\right)[q_1] + 2\sin\left(\frac{l_2}{l_1}2n\pi\right)[q_2] = o(1), \qquad n \to \infty.
$$
 (3.2)

Note that

$$
\left(\sin\left(\frac{l_2}{l_1}2n\pi\right)\right)^{-1} = O(1), \qquad n \to \infty,
$$
\n(3.3)

since $\frac{l_2}{l_1}$ is an irrational number. Combining [\(3.2\)](#page-5-1) and [\(3.3\)](#page-5-2) yields

$$
[q_1] + 2[q_2] = o(1), \qquad n \to \infty,
$$

which follows

$$
[q_1] + 2[q_2] = 0. \t\t(3.4)
$$

Taking $\rho = \frac{(2n+1)\pi}{l_0}$ $\frac{1}{l_2}$ in the estimate [\(3.1\)](#page-5-0), we obtain $2\sin\left(\frac{l_1}{l}\right)$ $\frac{l_1}{l_2}(2n+1)\pi\bigg) [q_1] + \sin\left(\frac{l_1}{l_2}\right)$ $\frac{l_1}{l_2}(2n+1)\pi\bigg) [q_2] = o(1), \qquad n \to \infty.$ (3.5) Note that

$$
\left(\sin\left(\frac{l_1}{l_2}(2n+1)\pi\right)\right)^{-1} = O(1), \qquad n \to \infty,
$$
\n(3.6)

since $\frac{l_1}{l_2}$ is an irrational number. Combining [\(3.5\)](#page-5-3) and [\(3.6\)](#page-6-0) yields

$$
2[q_1] + [q_2] = o(1), \quad n \to \infty,
$$

which follows

$$
2[q_1] + [q_2] = 0. \t\t(3.7)
$$

It is easy to see that equations [\(3.4\)](#page-5-4) and [\(3.7\)](#page-6-1) imply $[q_1] = [q_2] = 0$.

(ii) Let $\frac{l_2}{l_1}$ be a rational number. Without losing generality, we assume $l_1 = k_1 l$, $l_2 = k_2 l$, where k_1, k_2 are two positive integers and relatively prime. From [\(2.4\)](#page-3-3), together with $\Delta(\lambda) = \Delta_0(\lambda)$, we have

$$
\frac{2(\sin \rho k_1 l)(\cos \rho k_2 l - 1) + (\cos \rho k_1 l)(\sin \rho k_2 l)}{2\rho} [q_1]
$$

$$
+\frac{2(\cos \rho k_1 l)(\sin \rho k_2 k) + (\sin \rho k_1 l)(\cos \rho k_2 l)}{2\rho} [q_2]
$$

$$
= o\left(\frac{e^{|\tau|(k_1+k_2)l}}{\rho}\right), \quad |\rho| \to \infty.
$$
(3.8)

When $k_1 = k_2$ then $[q_1] = [q_2] = 0$ is obvious from Case (1). The following is divided into cases $k_2 < k_1$ and $k_2 > k_1$.

a) Case $k_2 < k_1$. In the estimate [\(3.8\)](#page-6-2), taking $\rho = \frac{(2k_1n+1)\pi}{k_1l}$ $\frac{n+1}{k_1 l}$ and let $n \to \infty$, one can get

$$
\sin\left(\frac{k_2\pi}{k_1}\right)[q_1] + 2\sin\left(\frac{k_2\pi}{k_1}\right)[q_2] = 0,\tag{3.9}
$$

which follows

$$
[q_1] = -2[q_2],\tag{3.10}
$$

since $0 < \sin\left(\frac{k_2\pi}{k_1}\right)$ $\overline{k_1}$ $\left(3.8 \right)$, taking $\rho = \frac{(2k_1 n + \frac{1}{2})\pi}{k_1 l}$ $\frac{n+1}{k_1}$ and let $n \to \infty$, one can get

$$
2\left(\cos\left(\frac{k_2\pi}{2k_1}\right) - 1\right)[q_1] + \cos\left(\frac{k_2\pi}{2k_1}\right)[q_2] = 0.
$$
\n(3.11)

Substituting (3.10) into (3.11) , we have

$$
\left(4 - 3\cos\left(\frac{k_2\pi}{2k_1}\right)\right)[q_2] = 0.
$$

Since $4-3\cos\left(\frac{k_2\pi}{2k_1}\right)$ $\overline{2k_1}$ $\Big) \neq 0$, this yields $[q_2] = 0$. Thus $[q_1] = 0$ from [\(3.10\)](#page-6-3).

b) Case $k_2 > k_1$. In the estimate [\(3.8\)](#page-6-2), taking $\rho = \frac{(2k_2n+1)\pi}{k_2l}$ $\frac{n+1}{k_2 l}$ and let $n \to \infty$, one can get

$$
-4\sin\left(\frac{k_1\pi}{k_2}\right)[q_1] - \sin\left(\frac{k_1\pi}{k_2}\right)[q_2] = 0,
$$
\n(3.12)

which follows

$$
[q_2] = -4[q_1], \tag{3.13}
$$

since $0 < \sin\left(\frac{k_1 \pi}{k_2}\right)$ $\overline{k_{2}}$ $\left(3.8 \right)$, taking $\rho = \frac{(2k_2 n + \frac{1}{2})\pi}{k_2 k_1}$ $\frac{n+1}{k_2 l}$ and let $n \to \infty$, one can get

$$
\left(-2\sin\left(\frac{k_1\pi}{2k_2}\right) + \cos\left(\frac{k_1\pi}{2k_2}\right)\right)[q_1] + 2\cos\left(\frac{k_1\pi}{2k_2}\right)[q_2] = 0. \tag{3.14}
$$

Substituting (3.13) into (3.14) , we have

$$
\left(2\sin\left(\frac{k_1\pi}{2k_2}\right) + 7\cos\left(\frac{k_1\pi}{2k_2}\right)\right)[q_1] = 0.
$$

Since $2 \sin \left(\frac{k_1 \pi}{2 k_0} \right)$ $2k_2$ $+ 7 \cos \left(\frac{k_1 \pi}{2 k_0} \right)$ $2k_2$ $\Big) \neq 0$, this yields $[q_1] = 0$. Thus $[q_2] = 0$ from [\(3.13\)](#page-7-0). The proof of Lemma [3.1](#page-5-5) is complete. \Box

Let us introduce the Hilbert space $H := L^2(0, l_1) \oplus L^2(0, l_2)$ with the inner product

$$
(f,g) = \sum_{j=1}^{2} \int_0^{l_j} f_j(x) \overline{g_j(x)} dx,
$$

where $f = (f_1, f_2)^T \in H$, $g = (g_1, g_2)^T \in H$, and g^T denotes the transpose of the vector g. The domain of self-adjoint operator $L(q)$ is

$$
D(L(q)) = \left\{ y = (y_1, y_2)^T \in H \mid y_j \in AC(0, l_j), y'_j \in AC[0, l_j], j = 1, 2, satisfying (1.2) and (1.3), \varphi(y) := (\varphi(y_1), \varphi(y_2))^T \in H \right\},\
$$

where $AC[0, l_j]$ represents a set of all absolutely continuous functions on $[0, l_j]$.

Proof of Theorem [1.1](#page-2-0). It is readily verified that the operator $L(0)$ is non-negative and $0 \in \sigma(L(0))$, so zero is its smallest eigenvalue, namely $\lambda_0(0) = 0$.

Next we show that $y_0 = (1, 1)^T$ is an eigenfunction of $L(q)$ corresponding to the eigenvalue zero. By the variational principle, we obtain

$$
0 = \lambda_0(0) = \lambda_0(q) = \inf_{0 \neq y \in D(L(q))} \frac{(\varphi(y), y)}{(y, y)}
$$

$$
= \inf_{0 \neq y \in D(L(q))} \frac{\sum_{j=1}^{2} \left(- \int_{0}^{l_{j}} y''_{j}(x) \overline{y_{j}(x)} dx + \int_{0}^{l_{j}} q_{j}(x) |y_{j}(x)|^{2} dx \right)}{\sum_{j=1}^{2} \int_{0}^{l_{j}} |y_{j}(x)|^{2} dx}
$$

Now $y_0 \in D(L(q))$ is obvious, and so

$$
0 \le \frac{(\varphi(y_0), y_0)}{(y_0, y_0)} = \frac{\sum_{j=1}^2 \int_0^{l_j} q_j(x) dx}{\sum_{j=1}^2 l_j} = \frac{[q_1] + [q_2]}{l_1 + l_2},
$$

by Corollary [2.2](#page-4-6) and Lemma [3.1,](#page-5-5) the right hand side is exactly zero, which implies that the test function y_0 is an eigenfunction of $L(q)$ corresponding to the eigenvalue zero. Substituting y_0 , which is the eigenfunction of eigenvalue zero, into equations [\(1.1\)](#page-1-2), we obtain $q_j = 0$ in $L^2[0, l_j], j = 1, 2$. The proof is finished. \Box

Acknowledgments. This work was supported in part by the National Natural Science Foundation of China (11871031) and the National Natural Science Foundation of Jiang Su (BK20201303).

REFERENCES

- [1] V. A. Ambarzumyan, Über eine Frage der Eigenwerttheorie, $Z. Phys.$, 53 (1929), 690-695.
- [2] G. Berkolaiko, R. Carlson, S. Fulling and P. Kuchment, Quantum Graphs and Their Applications, Amer. Math. Soc., Providence, RI: Contemp. Math. 415 (2006).
- [3] G. Berkolaiko and P. Kuchment, Introduction to Quantum Graphs, Amer. Math. Soc., Providence, RI (2013).
- [4] R. Carlson and V. N. Pivovarchik, Ambarzumyan's theorem for the trees. *Electronic* J. Diff. Equa., 2007 (2007), 142, 1-9.
- [5] H. H. Chern and C. L. Shen, On the n-dimensional Ambarzumyan's theorem, In verse Problems, 13 (1997), 15-18.
- [6] E.B. Davies, An inverse spectral theorem, *J. Oper. Theory.*, **69** (1)(2013), 195-208.
- [7] G. Freiling and V. A. Yurko, Inverse Sturm-Liouville Problems and their Applications, Nova Science Publishers, New York, (2001).
- [8] F. Gesztesy and B. Simon, Inverse spectral analysis with partial information on the potential, II. the case of discrete spectrum, Trans. Amer. Math. Soc., 352, No.6 (1999), 2765-2787.
- [9] M. Horváth, On the stability in Ambarzumian theorems, *Inverse Problems*, **31** (2015), 025008, 9pp.
- [10] A. A. Kirac, On the Ambarzumyan's theorem for the quasi-periodic boundary conditions, Anal. Math. Phys., 6 (2016), 297-300.

.

10 F. WANG AND C. F. YANG

- [11] M. Kiss, Spectral determinants and Ambarzumian type theorem on graphs, Integr. Equ. Oper. Theory, 92 (2020), 24.
- [12] C. K. Law and E. Yanagida, A solution to an Ambarzumyan problem on trees, Kodai J. Math., 35 (2012), 358-373.
- [13] V. N. Pivovarchik, Ambarzumyan's theorem for for a Sturm-Liouville boundary value problem on a star-shaped graph, Funct. Anal. Appl., 39 (2005), 148-151.
- [14] C. F. Yang, Z. Y. Huang and X. P. Yang, Ambarzumyan-type theorems for the Sturm-Liouville equation on a graph, Rocky Mountain J. Math., 39 (2009), 1353- 1372.
- [15] C. F. Yang and X. C. Xu, Ambarzumyan-type theorems on graphs with loops and double edges, J. Math. Anal. Appl., 444 (2016), 1348-1358.
- [16] C. F. Yang, F. Wang and Z. Y. Huang, Ambarzumyan theorems for Dirac operators, Acta Mathematicae Applicatae Sinica-English Series, 37 (2) 2021, 287-298.
- [17] V. A. Yurko, Inverse problems for Sturm-Liouville operators on graphs wiyh a cycle, Operators and Matrices, 2(4) (2008), 543-553.
- [18] V. A. Yurko, On Ambarzumyan-type theorems, Appl. Math. Lett., 26 (2013), 506- 509.
- [19] R. Zhang and C. F. Yang, Ambarzumyan-type theorem for the impulsive Sturm-Liouville operator, J. Inv. Ill-Posed Problems, 29 (2021), 21-25.

SCHOOL OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF SCIENCE AND TECHNOLOGY, Nanjing, 210094, Jiangsu, China

 $Email \ address:$ wangfengmath@njust.edu.cn

Department of Mathematics, School of Mathematics and Statistics, Nanjing University of Science and Technology, Nanjing, 210094, Jiangsu, People's Republic of China $Email$ $address:$ chuanfuyang@njust.edu.cn