

From sparse to dense functional data in high dimensions: Revisiting phase transitions from a non-asymptotic perspective

Shaojun Guo¹, Dong Li², Xinghao Qiao³, and Yizhu Wang⁴

¹*Institute of Statistics and Big Data, Renmin University of China, P.R. China*

²*Center for Statistical Science, Tsinghua University, P.R. China*

³*Department of Statistics, London School of Economics and Political Science, U.K.*

⁴*Department of Mathematical Sciences, Tsinghua University, P.R. China*

Abstract

Nonparametric estimation of the mean and covariance functions is ubiquitous in functional data analysis and local linear smoothing techniques are most frequently used. Zhang and Wang (2016) explored different types of asymptotic properties of the estimation, which reveal interesting phase transition phenomena based on the relative order of the average sampling frequency per subject T to the number of subjects n , partitioning the data into three categories: “sparse”, “semi-dense” and “ultra-dense”. In an increasingly available high-dimensional scenario, where the number of functional variables p is large in relation to n , we revisit this open problem from a non-asymptotic perspective by deriving comprehensive concentration inequalities for the local linear smoothers. Besides being of interest by themselves, our non-asymptotic results lead to elementwise maximum rates of L_2 convergence and uniform convergence serving as a fundamentally important tool for further convergence analysis when p grows exponentially with n and possibly T . With the presence of extra $\log p$ terms to account for the high-dimensional effect, we then investigate the scaled phase transitions and the corresponding elementwise maximum rates from sparse to semi-dense to ultra-dense functional data in high dimensions. Finally, numerical studies are carried out to confirm our established theoretical properties.

Keywords: Concentration inequality; Covariance function; Elementwise maximum rate; L_2 and uniform convergence; Local linear smoother; Mean function.

1 Introduction

A fundamental issue in functional data analysis is the nonparametric estimation of the mean and covariance functions based on discretely sampled and noisy curves. The estimated quantities are not only of interest by themselves, but also serve as building blocks for dimension reduction and subsequent modelling of functional data, such as functional principal component analysis (Yao et al., 2005a; Li and Hsing, 2010) and functional linear regression (Yao et al., 2005b). Among candidate nonparametric smoothers, we focus on the most commonly-adopted local linear smoothing method due to its simplicity and attractive local and boundary correction properties.

In a typical functional data setting, we have n random curves, representing n subjects, observed with errors, at T_i randomly sampled time points for the i th subject. The sampling frequency T_i plays a pivotal role in the estimation, as it may affect the choice of the estimation procedure. The literature can be loosely divided into two categories. The first category corresponds to dense functional data, where T_i 's are larger than some order of n . A conventional approach to handle such data implements nonparametric smoothing to the observations from each subject to eliminate the noise, thus reconstructing each individual curve before subsequent analysis (Zhang and Chen, 2007). The second category, referred to as sparse functional data, accords with bounded T_i 's. Under such a scenario, the pre-smoothing step is no longer applicable, an alternative post-smoothing type approach considers pooling the data from all subjects to build strength across all observations (Yao et al., 2005a; Li and Hsing, 2010). More recently, Zhang and Wang (2016) provided a comprehensive analysis of phase transitions and the associated rates of convergence for three types of asymptotic properties: local asymptotic normality, L_2 convergence, and uniform convergence. They proposed to further partition dense functional data into new categories: “semi-dense” and “ultra-dense”, depending on whether the root- n rate is achieved with negligible asymptotic bias or not. However, these aforementioned asymptotic results are only suitable to handle univariate or low-dimensional multivariate functional data.

With recent advances in data collection technology, high-dimensional functional datasets become increasingly available. Examples include time-course gene expression data, and electroencephalography and functional magnetic resonance imaging data, where signals are

measured over time at a large number of regions of interest (Zhu et al., 2016; Li and Solea, 2018; Zapata et al., 2022; Fang et al., 2023). Those data can be represented as a p -vector of random functions $\mathbf{X}_i(\cdot) = \{X_{i1}(\cdot), \dots, X_{ip}(\cdot)\}^\top$ for $i = 1, \dots, n$ defined on a compact set \mathcal{U} , where the dimension p can be diverging with, or even larger than, the number of subjects n . In practice, each $X_{ij}(\cdot)$ is observed subject to error contamination at T_{ij} random time points; see (1) below. In this paper, we propose a unified local linear smoothing approach to estimate the mean and covariance functions for both sparsely and densely sampled high-dimensional functional data.

Within the framework of high-dimensional statistics, it is essential to conduct non-asymptotic analysis of the estimators by developing concentration inequalities under given performance metric, which can lead to probabilistic error bounds in elementwise maximum norm as a function of n , p and possibly T_{ij} 's under our setup. The literature has mainly focused on fully observed functional data, based on which concentration inequalities for the estimated covariance functions were established in Qiao et al. (2019) and Zapata et al. (2022). For the practical scenario where the curves are partially observed with errors, one has to deal with the second order U -statistics with complex dependence structure, posing a technically challenging task. Qiao et al. (2020) made the first attempt to derive some sub-optimal concentration results for the estimated covariance functions only under a restrictive finite-dimensional setting. Lee et al. (2022) and Fang et al. (2023) developed their methodologies by simply assuming that the elementwise maximum rates of the covariance estimators are not faster than the parametric counterpart without specifying the exact forms. Therefore, it remains an important open problem to investigate the non-asymptotic proprieties of both the mean and covariance estimators when handling partially observed functional data.

We present in this paper a systematic and unified non-asymptotic analysis of the local linear smoothers for the mean and covariance functions to accommodate both sparsely and densely sampled functional data in high dimensions. The new contribution of this paper is threefold.

- First, we develop generalized sub-Gaussian type concentration inequalities in both L_2 norm and supremum norm for each functional element of the mean and covariance estimators. Compare to the asymptotic results in Zhang and Wang (2016), our

non-asymptotic error bounds lead to the same rates of L_2 convergence and uniform convergence, and reveal the same phase transition phenomena depending on the relative order of the average sampling frequency per subject to $n^{1/4}$ for dense functional data.

- Second, we derive elementwise maximum rates of both L_2 and uniform convergence for the mean and covariance estimators, serving as a fundamental tool to provide theoretical guarantees for a set of high-dimensional functional data models, such as functional graphical models (Li and Solea, 2018; Qiao et al., 2019, 2020; Solea and Li, 2022; Zapata et al., 2022; Lee et al., 2022), functional covariance estimation (Fang et al., 2023), and functional additive regressions (Fan et al., 2014, 2015; Kong et al., 2016; Luo and Qi, 2017).
- Third, with the presence of additional $\log p$ terms to account for the high-dimensional effect in our established elementwise maximum rates, the scaled phase transitions for high-dimensional dense functional data occur based on the ratios of the average sampling frequency per subject to $n^{1/4}(\log p)^{-1/4}$. This leads to a further partition of dense functional data into categories of “semi-dense” and “ultra-dense”, depending on whether the parametric rate $(\log p)^{1/2}n^{-1/2}$ can be attained or not. With suitable choices of optimal bandwidths, we also present the optimal elementwise maximum rates from sparse to semi-dense to ultra-dense functional data, which correspondingly extend the optimal rates in Zhang and Wang (2016) to the high-dimensional setting.

Outline of the paper. In Section 2, we present the nonparametric smoothing approach to estimate the mean and covariance functions. In Section 3, we investigate the non-asymptotic and convergence properties of the proposed local linear smoothers and discuss the associated phase transition phenomena. The established theoretical results are validated through simulations in Section 4. All technical proofs are relegated to the Appendix.

Notation. We summarize here some notation to be used throughout the paper. For a positive integer q , we write $[q] = \{1, \dots, q\}$. For $x, y \in \mathbb{R}$, we write $x \vee y = \max(x, y)$ and $x \wedge y = \min(x, y)$. We use $I(\cdot)$ to denote an indicator function. Let $L_2(\mathcal{U})$ be a Hilbert space of square-integrable functions on a compact interval \mathcal{U} equipped with the inner product

$\langle f, g \rangle = \int f(u)g(u)du$ for $f(\cdot), g(\cdot) \in L_2(\mathcal{U})$ and the induced L_2 norm $\|\cdot\| = \langle \cdot, \cdot \rangle^{1/2}$. For any bivariate function $\Phi(\cdot, \cdot)$ in the tensor product space $L_2(\mathcal{U}) \otimes L_2(\mathcal{U})$, we also use Φ to denote the linear operator induced from the kernel function $\Phi(\cdot, \cdot)$, that is, for any $f(\cdot) \in L_2(\mathcal{U})$, $\Phi(f)(\cdot) = \int \Phi(\cdot, v)f(v)dv \in L_2(\mathcal{U})$, and denote its Hilbert–Schmidt norm by $\{\int \int \Phi(u, v)^2 dudv\}^{1/2}$. For two positive sequences $\{a_n\}$ and $\{b_n\}$, we write $a_n \lesssim b_n$ or $b_n \gtrsim a_n$ if there exist a positive constant c such that $\limsup_{n \rightarrow \infty} a_n/b_n \leq c$. We write $a_n \asymp b_n$ if and only if $a_n \lesssim b_n$ and $b_n \lesssim a_n$ hold simultaneously.

2 Methodology

Let $\mathbf{X}_i(\cdot) = \{X_{i1}(\cdot), \dots, X_{ip}(\cdot)\}^T$ for $i \in [n]$ be independently and identically distributed copies of $\mathbf{X}(\cdot)$ defined on \mathcal{U} with p -vector of mean functions $\boldsymbol{\mu}(\cdot) = \{\mu_1(\cdot), \dots, \mu_p(\cdot)\}^T = E\{\mathbf{X}(\cdot)\}$ and $(p \times p)$ -matrix of marginal- and cross-covariance functions $\boldsymbol{\Sigma}(u, v) = \{\Sigma_{jk}(u, v)\}_{p \times p}$ whose (j, k) -th entry $\Sigma_{jk}(u, v) = \text{cov}\{X_{ij}(u), X_{ik}(v)\}$. For any $i \in [n]$ and $j \in [p]$, $X_{ij}(\cdot)$ is not directly observable in practice. Instead it is observed, with random errors, at T_{ij} random time points, $U_{i1}, \dots, U_{iT_{ij}} \in \mathcal{U}$. Let Y_{ijt} be the observed value of $X_{ij}(U_{ijt})$ satisfying

$$Y_{ijt} = X_{ij}(U_{ijt}) + \varepsilon_{ijt}, \quad (1)$$

where the errors ε_{ijt} 's, independent of X_{ij} 's, are independently and identically distributed copies of ε_j with $E(\varepsilon_j) = 0$ and $\text{var}(\varepsilon_j) = \sigma_j^2 < \infty$.

Based on the observed data $\{(U_{ijt}, Y_{ijt}) : i \in [n], j \in [p], t \in [T_{ij}]\}$, we present a unified procedure to estimate the mean functions $\mu_j(\cdot)$'s and the marginal- and cross-covariance functions $\Sigma_{jk}(\cdot, \cdot)$'s for both sparsely and densely sampled functional data. In what follows, denote $K_h(\cdot) = h^{-1}K(\cdot/h)$ for a univariate kernel K with bandwidth $h > 0$. For each j , a local linear smoother is firstly applied to $\{(U_{ijt}, Y_{ijt}) : i \in [n], t \in [T_{ij}]\}$, and hence the estimated mean function is attained via $\hat{\mu}_j(u) = \hat{b}_0$, where

$$(\hat{b}_0, \hat{b}_1) = \arg \min_{b_0, b_1} \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \left\{ Y_{ijt} - b_0 - b_1(U_{ijt} - u) \right\}^2 K_{h_{\mu, j}}(U_{ijt} - u). \quad (2)$$

The weight v_{ij} is attached to each observation for the i th subject and the j th functional variable such that $\sum_{i=1}^n T_{ij}v_{ij} = 1$ (Zhang and Wang, 2016).

For each $i \in [n], j, k \in [p], t \in [T_{ij}]$ and $s \in [T_{ik}]$, once the mean functions are estimated, let $\Theta_{ijkts} = \{Y_{ijt} - \hat{\mu}_j(U_{ijt})\}\{Y_{iks} - \hat{\mu}_k(U_{iks})\}$ be the ‘‘raw covariance’’ between Y_{ijt} and

Y_{iks} . Observe that $\text{cov}(Y_{ijt}, Y_{iks}) = \Sigma_{jk}(U_{ijt}, U_{iks}) + \sigma_j^2 I(j = k)I(t = s)$. To estimate the marginal-covariance function $\Sigma_{jj}(\cdot, \cdot)$ for each j or the cross-covariance function $\Sigma_{jk}(\cdot, \cdot)$ for each $j \neq k$, we next employ local linear surface smoothers to the off-diagonals of the raw marginal-covariances $(\Theta_{ijjts})_{1 \leq t \neq s \leq T_{ij}}$ or to the raw cross-covariances $(\Theta_{ijkts})_{t \in [T_{ij}], s \in [T_{ik}]}$. Specifically, we minimize

$$\sum_{i=1}^n w_{ijk} \sum_{(t,s) \in \mathcal{T}} \left\{ \Theta_{ijkts} - \beta_0 - \beta_1(U_{ijt} - u) - \beta_2(U_{iks} - v) \right\}^2 K_{h_{\Sigma, jk}}(U_{ijt} - u) K_{h_{\Sigma, jk}}(U_{iks} - v). \quad (3)$$

with respect to $(\beta_0, \beta_1, \beta_2)$, where the set \mathcal{T} equals to $\{(t, s) : t \in [T_{ij}], s \in [T_{ij}], t \neq s\}$ if $j = k$ or $\{(t, s) : t \in [T_{ij}], s \in [T_{ik}]\}$ if $j \neq k$, and the weight w_{ijk} is assigned to each triplet (i, j, k) such that $\sum_{i=1}^n T_{ij} \{T_{ik} - I(j = k)\} w_{ijk} = 1$. See the weights to estimate marginal-covariance functions in [Zhang and Wang \(2016\)](#). The resulting marginal- or cross-covariance estimator is $\hat{\Sigma}_{jk}(u, v) = \hat{\beta}_0$. For ease of presentation, we assume that the mean functions $\mu_j(\cdot)$'s are known in advance when discussing the concentration and convergence results related to the covariance estimators $\hat{\Sigma}_{jk}(u, v)$'s in [Section 3](#) below. However, it is noteworthy that all our discussions remain valid even when $\mu_j(\cdot)$'s are unknown as long as a few additional technical assumptions are imposed.

Our estimation procedure allows general weighting schemes for $\{v_i\}_{i \in [n]}$, $\{w_{ijk}\}_{i \in [n], j, k \in [p]}$ such that two types of frequently-used schemes in existing literature are special cases of them. One type assigns the same weights to each observation ([Yao et al., 2005a](#)) with $v_{ij} = (\sum_{i=1}^n T_{ij})^{-1}$ and $w_{ijk} = [\sum_{i=1}^n T_{ij} \{T_{ik} - I(j = k)\}]^{-1}$, so a subject with a larger number of observations receives more weights in total. The other type assigns the same weights to each subject ([Li and Hsing, 2010](#)), thus leading to $v_{ij} = (nT_{ij})^{-1}$ and $w_{ijk} = [nT_{ij} \{T_{ik} - I(j = k)\}]^{-1}$.

Suppose that the estimated mean and covariance functions are evaluated at a grid of $R \times R$ locations over \mathcal{U}^2 . Under the high-dimensional setting, it is apparent that our proposed nonparametric smoothing approach suffers from high computational cost in kernel evaluations, particularly when estimating $p(p + 1)/2$ marginal- and cross-covariance functions. In a common practical scenario, where each $X_{ij}(\cdot)$ is observed at the same set of time points $U_1, \dots, U_{T_i} \in \mathcal{U}$ across $j \in [p]$, the number of kernel evaluations is reduced from $O(\sum_{i=1}^n \sum_{j=1}^p T_{ij} R)$ to $O(\sum_{i=1}^n T_i R)$. To further speed up the computation, we can adopt the linear binning technique ([Fan and Marron, 1994](#)) to approximate the mean and co-

variance estimation. This would largely reduced the number of kernel evaluations to $O(R)$ while requiring only $O(\sum_{i=1}^n T_i)$ additional operations. See the detailed implementation of binning in Fang et al. (2023). Our conducted numerical experiments show that such binned implementation offers substantially improved computational efficiency without sacrificing any estimation accuracy.

3 Theory

Before presenting the concentration and convergence results, we impose the following regularity assumptions.

Assumption 1 For each $i \in [n]$ and $j \in [p]$, $X_{ij}(\cdot)$ is a sub-Gaussian random process and ε_{ij} is a sub-Gaussian random variable, that is, there exists some positive constant c such that $E\{\exp(\langle x, X_{ij} - \mu_j \rangle)\} \leq \exp\{2^{-1}c^2\langle x, \Sigma_{jj}(x) \rangle\}$ for all $x(\cdot) \in L^2(\mathcal{U})$ and $E\{\exp(\varepsilon_{ij}z)\} \leq \exp(2^{-1}c^2\sigma_j^2z^2)$ for all $z \in \mathbb{R}$.

Assumption 2 For each $i \in [n]$ and $j \in [p]$, under the sparse design, $T_{ij} \leq T_0 < \infty$, and, under the dense design, $T_{ij} \rightarrow \infty$ and there exists some positive constant \bar{c} such that $\max_{i,j} T_{ij}(\min_{i,j} T_{ij})^{-1} \leq \bar{c}$.

Assumption 3 Under the dense design, there exists some positive constant c_0 such that $\max_{i,j} v_{ij}(\min_{i,j} v_{ij})^{-1} \leq c_0$ and $\max_{i,j,k} w_{ijk}(\min_{i,j,k} w_{ijk})^{-1} \leq c_0$.

Assumption 4 (i) Let $\{U_{ijt} : i \in [n], t \in [T_{ij}]\}$ be independently and identically distributed copies of a random variable U defined on \mathcal{U} . The density $f_U(\cdot)$ of U satisfies $0 < m_f \leq \inf_{\mathcal{U}} f_U(u) \leq \sup_{\mathcal{U}} f_U(u) \leq M_f < \infty$ for some positive constants m_f and M_f ; (ii) $\mathbf{X}(\cdot)$, U and $\{\varepsilon_j\}_{j \in [p]}$ are mutually independent.

Assumption 5 Let $B_{jk} = [(k-1)\tilde{T}_j^{-1}, k\tilde{T}_j^{-1}]$ for $k \in [\tilde{T}_j]$ with $\tilde{T}_j = \max_i T_{ij}$, there exists some constant $C > 0$ such that the cardinality $\#\{U_{ijt} : U_{ijt} \in B_{jk}, t \in [T_{ij}]\} \leq C$ for each $i \in [n], j \in [p]$ and $k \in [\tilde{T}_j]$.

Assumption 6 (i) $K(\cdot)$ is symmetric probability density function on $[-1, 1]$ with $\int u^2 K(u) < \infty$ and $\int K(u)^2 du < \infty$. (ii) $K(\cdot)$ is Lipschitz continuous: there exists some positive constant L such that $|K(u) - K(v)| \leq L|u - v|$ for any $u, v \in [-1, 1]$.

Assumption 7 (i) $\partial^2 \mu_j(u)/\partial u^2$ is uniformly bounded over $u \in \mathcal{U}$ and $j \in [p]$; (ii) $\partial^2 \Sigma_{jk}(u, v)/\partial u^2$, $\partial^2 \Sigma_{jk}(u, v)/\partial u \partial v$, and $\partial^2 \Sigma_{jk}(u, v)/\partial v^2$ are uniformly bounded over $(u, v) \in \mathcal{U}^2$ and $(j, k) \in [p]^2$.

The sub-Gaussianities in Assumption 1 for both Hilbert space-valued random elements $X_{ij}(\cdot)$'s and random errors ε_j 's together imply that the observations Y_{ijt} 's in (1) are sub-Gaussian, which plays a crucial role in deriving our subsequent concentration inequalities. The dense case in Assumption 2 corresponds to a common practical scenario, where the sampling frequencies T_{ij} 's are of the same order across $i \in [n]$ and $j \in [p]$. Under such a scenario, Assumption 3 is automatically satisfied by two frequently-used weighting schemes including “equal weight per observation” and “equal weight per subject”. Assumption 5 means that all observational time points are distributed in the sense of “uniformly on \mathcal{U} ”. This circumvents the possibility of an extreme case where a large number of time points are concentrated in some small areas, while leaving too few points in other regions. Assumptions 4, 6 and 7 are standard in the literature of local linear smoothing for functional data (Yao et al., 2005a; Zhang and Wang, 2016) adaptable to the multivariate setting.

Theorem 1 Suppose that Assumptions 1-6 hold. For each $j \in [p]$, let $\gamma_{n,T,h,j} = n(1 \wedge \bar{T}_{\mu,j} h_{\mu,j})$ with the corresponding average sampling frequency per subject $\bar{T}_{\mu,j} = n^{-1} \sum_{i=1}^n T_{ij}$, then there exist some positive constants c_1, c_2 (independent of $n, p, \bar{T}_{\mu,j}$'s) and arbitrarily small $\epsilon_1 > 0$ such that for any $\delta \in (0, 1]$,

$$\Pr(\|\hat{\mu}_j - \tilde{\mu}_j\| \geq \delta) \leq c_2 \exp(-c_1 \gamma_{n,T,h,j} \delta^2), \quad (4)$$

$$\Pr\left\{\sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \geq \delta\right\} \leq \frac{c_2 (n^{\epsilon_1} \gamma_{n,T,h,j})^{1/2}}{h_{\mu,j}} \exp(-c_1 \gamma_{n,T,h,j} \delta^2), \quad (5)$$

where $\tilde{\mu}_j(u)$ is a deterministic univariate function that converges to $\mu_j(u)$ as $h_{\mu,j} \rightarrow 0$. See (A.1) in Appendix A for the exact form of $\tilde{\mu}_j(u)$.

Theorem 2 Suppose that Assumptions 1-6 hold. For each $j, k \in [p]$, let $\nu_{n,T,h,jk} = n(1 \wedge \bar{T}_{\Sigma,jk}^2 h_{\Sigma,jk}^2)$ with the corresponding average sampling frequency per subject $\bar{T}_{\Sigma,jk} = [n^{-1} \sum_{i=1}^n T_{ij} \{T_{ik} - I(j=k)\}]^{1/2}$, then there exist some positive constants c_3, c_4 (independent of $n, p, \bar{T}_{\Sigma,jk}$'s) and arbitrarily small $\epsilon_2 > 0$ such that for any $\delta \in (0, 1]$,

$$\text{pr}\left(\|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_{\mathcal{S}} \geq \delta\right) \leq c_4 \exp\left(-c_3 \nu_{n,T,h,jk} \delta^2\right), \quad (6)$$

$$\text{pr}\left\{\sup_{(u,v) \in \mathcal{U}^2} \left|\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)\right| \geq \delta\right\} \leq \frac{c_4 n^{\epsilon_2} \nu_{n,T,h,jk}}{h_{\Sigma,jk}^2} \exp\left(-c_3 \nu_{n,T,h,jk} \delta^2\right), \quad (7)$$

where $\widetilde{\Sigma}_{jk}(u,v)$ is a deterministic bivariate function that converges to $\Sigma_{jk}(u,v)$ as $h_{\Sigma,jk} \rightarrow 0$. See (A.17) in Appendix A for the exact form of $\widetilde{\Sigma}_{jk}(u,v)$.

The concentration inequalities in Theorems 1 and 2 imply that $\widehat{\mu}_j$ and $\widehat{\Sigma}_{jk}$ are nicely concentrated around $\tilde{\mu}_j$ and $\widetilde{\Sigma}_{jk}$, respectively, in both L_2 norm and supremum norm with generalized sub-Gaussian type tail behaviors. It is worth mentioning that such L_2 and uniform concentration results are derived based on the local concentration inequalities of $\widehat{\mu}_j(u)$ and $\widehat{\Sigma}_{jk}(u,v)$ for fixed interior points $u, v \in \mathcal{U}$, which enjoy the same tail behaviors as (4) and (6). Besides being fundamental to derive elementwise maximum error bounds that are essential for further convergence analysis under high-dimensional scaling, these non-asymptotic results lead to the same rates of L_2 convergence and uniform convergence compared to those in Zhang and Wang (2016). Specifically, under extra Assumption 7, it holds that

$$\begin{aligned} \|\widehat{\mu}_j - \mu_j\| &= O_P\left\{n^{-1/2} + (n\bar{T}_{\mu,j}h_{\mu,j})^{-1/2} + h_{\mu,j}^2\right\}, \\ \sup_{u \in \mathcal{U}} |\widehat{\mu}_j(u) - \mu_j(u)| &= O_P\left[(\log n)^{1/2} n^{-1/2} \{1 + (\bar{T}_{\mu,j}h_{\mu,j})^{-1/2}\} + h_{\mu,j}^2\right], \\ \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} &= O_P\left\{n^{-1/2} + (n\bar{T}_{\Sigma,jk}^2 h_{\Sigma,jk}^2)^{-1/2} + h_{\Sigma,jk}^2\right\}, \\ \sup_{(u,v) \in \mathcal{U}^2} \left|\widehat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)\right| &= O_P\left[(\log n)^{1/2} n^{-1/2} \{1 + (\bar{T}_{\Sigma,jk}^2 h_{\Sigma,jk}^2)^{-1/2}\} + h_{\Sigma,jk}^2\right]. \end{aligned}$$

The above rates of convergence reveal interesting phase transition phenomena depending on the ratio of the average sampling frequency per subject $\bar{T}_{\mu,j}$ (or $\bar{T}_{\Sigma,jk}$) to $n^{1/4}$. In the following, we use different rates of L_2 convergence for $\widehat{\mu}_j$ and $\widehat{\Sigma}_{jk}$ to illustrate a systematic partition of functional data into three categories:

1. Under the sparse design, when $h_{\mu,j} \asymp n^{-1/5}$,

$$\|\widehat{\mu}_j - \mu_j\| = O_P(n^{-1/2} h_{\mu,j}^{-1/2} + h_{\mu,j}^2) = O_P(n^{-2/5});$$

when $h_{\Sigma,jk} \asymp n^{-1/6}$,

$$\|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P(n^{-1/2}h_{\Sigma,jk}^{-1} + h_{\Sigma,jk}^2) = O_P(n^{-1/3}).$$

2. Under the dense design, when $\bar{T}_{\mu,j}n^{-1/4} \rightarrow 0$ with $h_{\mu,j} \asymp (n\bar{T}_{\mu,j})^{-1/5}$,

$$\|\hat{\mu}_j - \mu_j\| = O_P(n^{-1/2}\bar{T}_{\mu,j}^{-1/2}h_{\mu,j}^{-1/2} + h_{\mu,j}^2) = O_P\{(n\bar{T}_{\mu,j})^{-2/5}\};$$

when $\bar{T}_{\Sigma,jk}n^{-1/4} \rightarrow 0$ with $h_{\Sigma,j} \asymp (n\bar{T}_{\Sigma,jk}^2)^{-1/6}$,

$$\|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P(n^{-1/2}\bar{T}_{\Sigma,jk}^{-1}h_{\Sigma,jk}^{-1} + h_{\Sigma,jk}^2) = O_P\{(n\bar{T}_{\Sigma,jk}^2)^{-1/3}\}.$$

3. Under the dense design, when $\bar{T}_{\mu,j}n^{-1/4} \rightarrow \tilde{c}$ (some positive constant) with $h_{\mu,j} \asymp n^{-1/4}$ or $\bar{T}_{\mu,j}n^{-1/4} \rightarrow \infty$ with $h_{\mu,j} = o(n^{-1/4})$ and $\bar{T}_{\mu,j}h_{\mu,j} \rightarrow \infty$,

$$\|\hat{\mu}_j - \mu_j\| = O_P(n^{-1/2});$$

when $\bar{T}_{\Sigma,jk}n^{-1/4} \rightarrow \tilde{c}$ with $h_{\Sigma,jk} \asymp n^{-1/4}$ or $\bar{T}_{\Sigma,jk}n^{-1/4} \rightarrow \infty$ with $h_{\Sigma,jk} = o(n^{-1/4})$ and $\bar{T}_{\Sigma,jk}h_{\Sigma,jk} \rightarrow \infty$,

$$\|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P(n^{-1/2}).$$

As $\bar{T}_{\mu,j}$ and $\bar{T}_{\Sigma,jk}$ grow very fast, case 3 results in the root- n rate complying with the parametric rate for fully observed functional data. As $\bar{T}_{\mu,j}$ and $\bar{T}_{\Sigma,jk}$ grow moderately fast, case 2 corresponds to the optimal minimax rates (Zhang and Wang, 2016), which are slower than root- n but faster than the counterparts for sparsely sampled functional data. Our established convergence rates in cases 1, 2 and 3 allow free choices of (j, k) , and are respectively consistent to those of the mean and covariance estimators under categories of “sparse”, “semi-dense” and “ultra-dense” functional data introduced in Zhang and Wang (2016).

Theorem 3 *Suppose that the assumptions in Theorem 1 and Assumption 7(i) hold, and $\log p(\min_j \gamma_{n,T,h,j})^{-1} \rightarrow 0$, $\max_j h_{\mu,j} \rightarrow 0$ as $n, p \rightarrow \infty$. It then holds that*

$$\max_{j \in [p]} \|\hat{\mu}_j - \mu_j\| = O_P \left\{ \left(\frac{\log p}{\min_j \gamma_{n,T,h,j}} \right)^{1/2} + \max_j h_{\mu,j}^2 \right\}, \quad (8)$$

and, if $\min_j h_{\mu,j} \asymp \{\log(p \vee n)/n\}^{\kappa_1}$ for some $\kappa_1 \in (0, 1/2]$,

$$\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)| = O_P \left[\left\{ \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\}^{1/2} + \max_j h_{\mu,j}^2 \right]. \quad (9)$$

Theorem 4 Suppose that the assumptions in Theorem 2 and Assumption 7(ii) hold and $\log p(\min_{j,k} \nu_{n,T,h,jk})^{-1} \rightarrow 0$, $\max_{j,k} h_{\Sigma,jk} \rightarrow 0$ as $n, p \rightarrow \infty$. It then holds that

$$\max_{j,k \in [p]} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P \left\{ \left(\frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right)^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right\}, \quad (10)$$

and, if $\min_{j,k} h_{\Sigma,jk} \asymp \{\log(p \vee n)/n\}^{\kappa_2}$ for some $\kappa_2 \in (0, 1/2]$,

$$\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)| = O_P \left[\left\{ \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\}^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right]. \quad (11)$$

We observe that the elementwise maximum rates of L_2 convergence and uniform convergence are governed by both dimensionality parameters $(n, p, \{\bar{T}_{\mu,j}\}_{j \in [p]}, \{\bar{T}_{\Sigma,jk}\}_{j,k \in [p]})$ and internal parameters $(\{h_{\mu,j}\}_{j \in [p]}, \{h_{\Sigma,jk}\}_{j,k \in [p]})$. Each convergence rate is composed of two terms reflecting our familiar variance-bias tradeoff in nonparametric statistics. It is easy to see that the variance terms are determined by the least frequently sampled and smoothed components, that is the smallest $\bar{T}_{\mu,j}$ (or $\bar{T}_{\Sigma,jk}$) and $h_{\mu,j}$ (or $h_{\Sigma,jk}$) across j, k , whereas the highest level of smoothness with the largest $h_{\mu,j}$ (or $h_{\Sigma,jk}$) controls the bias terms.

To facilitate further discussion, we consider the simplified setting where $\bar{T}_{\mu,j} \asymp \bar{T}_{\mu}$, $h_{\mu,j} \asymp h_{\mu}$ and $\bar{T}_{\Sigma,jk} \asymp \bar{T}_{\Sigma}$, $h_{\Sigma,jk} \asymp h_{\Sigma}$ for each j and k . Compared to cases 1–3 above, the corresponding elementwise maximum rates of convergence for $\{\hat{\mu}_j\}_{j \in [p]}$ (or $\{\widehat{\Sigma}_{jk}\}_{j,k \in [p]}$) in Theorem 3 (or Theorem 4) reveal scaled phase transitions for dense functional data depending on the relative order of \bar{T}_{μ} (or \bar{T}_{Σ}) to $n^{1/4}(\log p)^{-1/4}$ instead of $n^{1/4}$. In the following, we use elementwise maximum rates of L_2 convergence to illustrate the phase transition phenomena and the optimal estimation from sparse to dense functional data in high dimensions. In terms of the uniform convergence, the same phenomena occur as long as $p \gtrsim n$.

i. Under the sparse design, when $h_{\mu} \asymp (\log p)^{1/5} n^{-1/5}$,

$$\max_j \|\hat{\mu}_j - \mu_j\| = O_P \left\{ \left(\frac{\log p}{nh_{\mu}} \right)^{1/2} + h_{\mu}^2 \right\} = O_P \left\{ \left(\frac{\log p}{n} \right)^{2/5} \right\};$$

when $h_{\Sigma} \asymp (\log p)^{1/6} n^{-1/6}$,

$$\max_{j,k} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P \left\{ \left(\frac{\log p}{nh_{\Sigma}^2} \right)^{1/2} + h_{\Sigma}^2 \right\} = O_P \left\{ \left(\frac{\log p}{n} \right)^{1/3} \right\}.$$

ii. Under the dense design, when $\bar{T}_\mu(\log p)^{1/4}n^{-1/4} \rightarrow 0$ with $h_\mu \asymp (\log p)^{1/5}(n\bar{T}_\mu)^{-1/5}$,

$$\max_j \|\hat{\mu}_j - \mu_j\| = O_P \left\{ \left(\frac{\log p}{n\bar{T}_\mu h_\mu} \right)^{1/2} + h_\mu^2 \right\} = O_P \left\{ \left(\frac{\log p}{n\bar{T}_\mu} \right)^{2/5} \right\}; \quad (12)$$

when $\bar{T}_\Sigma(\log p)^{1/4}n^{-1/4} \rightarrow 0$ with $h_\Sigma \asymp (\log p)^{1/6}(n\bar{T}_\Sigma^2)^{-1/6}$,

$$\max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left(\frac{\log p}{n\bar{T}_\Sigma^2 h_\Sigma^2} \right)^{1/2} + h_\Sigma^2 \right\} = O_P \left\{ \left(\frac{\log p}{n\bar{T}_\Sigma^2} \right)^{1/3} \right\}. \quad (13)$$

iii. Under the dense design, when $\bar{T}_\mu(\log p)^{1/4}n^{-1/4} \rightarrow \tilde{c}$ with $h_\mu \asymp (\log p)^{1/4}n^{-1/4}$ or $\bar{T}_\mu(\log p)^{1/4}n^{-1/4} \rightarrow \infty$ with $h_\mu = o\{(\log p)^{1/4}n^{-1/4}\}$ and $\bar{T}_\mu h_\mu \rightarrow \infty$,

$$\max_j \|\hat{\mu}_j - \mu_j\| = O_P \left\{ \left(\frac{\log p}{n} \right)^{1/2} \right\};$$

when $\bar{T}_\Sigma(\log p)^{1/4}n^{-1/4} \rightarrow \tilde{c}$ with $h_\Sigma \asymp (\log p)^{1/4}n^{-1/4}$ or $\bar{T}_\Sigma(\log p)^{1/4}n^{-1/4} \rightarrow \infty$ with $h_\Sigma = o\{(\log p)^{1/4}n^{-1/4}\}$ and $\bar{T}_\Sigma h_\Sigma \rightarrow \infty$,

$$\max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left(\frac{\log p}{n} \right)^{1/2} \right\}.$$

In a similar spirit to the partitioned three categories for univariate functional data (see cases **1**, **2** and **3** above), we can also term the high-dimensional functional data in cases **i**, **ii**, and **iii** as “sparse”, “semi-dense” and “ultra-dense”, respectively. The main difference lies in the presence of additional $\log p$ terms to account for the high-dimensional effect. As \bar{T}_μ and \bar{T}_Σ grow at least in the order of $n^{1/4}(\log p)^{-1/4}$, the attained optimal rate $(\log p)^{1/2}n^{-1/2}$ is identical to that for the fully observed functional data (Zapata et al., 2022), presenting that the theory for high-dimensional ultra-dense functional data falls in the parametric paradigm. As \bar{T}_μ and \bar{T}_Σ diverge slower than $n^{1/4}(\log p)^{-1/4}$, if we let $h_\mu \asymp (\log p)^{1/5}(n\bar{T}_\mu)^{-1/5}$ and $h_\Sigma \asymp (\log p)^{1/6}(n\bar{T}_\Sigma^2)^{-1/6}$ to balance the corresponding variance and bias terms, the optimal rates for high-dimensional semi-dense functional data are respectively achieved in (12) and (13). These rates degenerate to the minimax rates in case **2** when p is fixed. With the choice of elementwise optimal bandwidths $h_\mu \asymp (n\bar{T}_\mu)^{-1/5}$ and $h_\Sigma \asymp (n\bar{T}_\Sigma^2)^{-1/6}$, we obtain $\max_j \|\hat{\mu}_j - \mu_j\| = O_P\{(\log p)^{1/2}(n\bar{T}_\mu)^{-2/5}\}$ and $\max_{j,k} \|\hat{\Sigma}_{jk} - \Sigma_{jk}\| = O_P\{(\log p)^{1/2}(n\bar{T}_\Sigma)^{-1/3}\}$, which are respectively slower than the optimal rates in (12) and (13). Such discussion applies analogously to the sparse functional setting; see cases **1** and **i**. Compared to the asymptotic results for cases **1**, **2** and **3** under a fixed p scenario,

the high-dimensionality in cases **i**, **ii** and **iii** leads to the scaled phase transitions, optimal selected bandwidths and corresponding optimal rates, each of which is up to a factor of $\log p$ at some polynomial order. It is worth mentioning that [Lee et al. \(2022\)](#) simply assumed the elementwise maximum rate $\max_{j,k} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} = O_P(\log pn^{-\tau})$, which was further reduced to $O_P\{(\log p)^{1/2}n^{-\tau}\}$ in [Fang et al. \(2023\)](#). The involved quantity $\tau \in (0, 1/2]$ reflects the average sampling frequency with larger values yielding denser observational points. We fundamentally improve their rates in the sense of precisely specifying the largest values of τ under cases with different sampling frequencies.

4 Simulations

In this section, we examine the finite-sample performance of the local linear smoothers for the mean and covariance function estimation in high dimensions.

We generalize the simulated example for univariate functional data in [Zhang and Wang \(2016\)](#) to the multivariate setting by generating

$$X_{ij}(u) = \mu_j(u) + \boldsymbol{\phi}(u)^{\text{T}} \boldsymbol{\theta}_{ij}, \quad i \in [n], j \in [p], u \in \mathcal{U} = [0, 1],$$

where the true mean function $\mu_j(u) = 1.5 \sin\{3\pi(u + 0.5)\} + 2u^3$, the basis function $\boldsymbol{\phi}(u) = \{\sqrt{2} \cos(2\pi u), \sqrt{2} \sin(2\pi u), \sqrt{2} \cos(4\pi u), \sqrt{2} \sin(4\pi u)\}^{\text{T}}$ and the basis coefficient vector $\boldsymbol{\theta}_i = (\boldsymbol{\theta}_{i1}^{\text{T}}, \dots, \boldsymbol{\theta}_{ip}^{\text{T}})^{\text{T}} \in \mathbb{R}^{4p}$ is sampled independently from a mean zero multivariate Gaussian distribution with block covariance matrix $\boldsymbol{\Lambda} \in \mathbb{R}^{4p \times 4p}$ whose (j, k) th block is given by $\boldsymbol{\Lambda}_{jk} = \rho^{|j-k|} \text{diag}\{2^{-2}, \dots, 5^{-2}\} \in \mathbb{R}^{4 \times 4}$ for $j, k \in [p]$. Hence the (j, k) th entry of the true covariance functions $\boldsymbol{\Sigma}(\cdot, \cdot) = \{\Sigma_{jk}(\cdot, \cdot)\}_{p \times p}$ is $\Sigma_{jk}(u, v) = \boldsymbol{\phi}(u)^{\text{T}} \boldsymbol{\Lambda}_{jk} \boldsymbol{\phi}(v)$. We then generate the observed values $Y_{ijt} = X_{ij}(U_{ijt}) + \varepsilon_{ijt}$ for $t = 1, \dots, T_{ij} = T$, where the time points U_{ijt} 's and errors ε_{ijt} 's are sampled independently from Uniform[0, 1] and $\mathcal{N}(0, 0.5^2)$, respectively.

We use the Epanechnikov kernel with bandwidth values varying on a dense grid. To evaluate the performance of $\widehat{\mu}_j(\cdot)$ for $j \in [p]$ and $\widehat{\Sigma}_{jk}(\cdot, \cdot)$ for $(j, k) \in [p]^2$ given specific bandwidth $h_{\mu,j}$ and $h_{\Sigma,jk}$, we define the corresponding mean integrated squared errors (MISE) as $\text{MISE}(\widehat{\mu}_j, h_{\mu,j}) = \int_{\mathcal{U}} \{\widehat{\mu}_j(u) - \mu_j(u)\}^2 du$ and $\text{MISE}(\widehat{\Sigma}_{jk}, h_{\Sigma,jk}) = \int_{\mathcal{U}} \int_{\mathcal{U}} \{\widehat{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)\}^2 dudv$. We first calculate the elementwise minimal MISEs for the mean and covariance estimators over the grids of candidate bandwidths in prespecified sets \mathcal{H}_{μ} and \mathcal{H}_{Σ} ,

respectively. We then compute their averages and maximums over $j \in [p]$ for the mean functions, that is,

$$\text{AveMISE}(\mu) = p^{-1} \sum_j \min_{h_{\mu,j}} \text{MISE}(\hat{\mu}_j, h_{\mu,j}), \quad \text{MaxMISE}(\mu) = \max_j \min_{h_{\mu,j}} \text{MISE}(\hat{\mu}_j, h_{\mu,j}),$$

and over $(j, k) \in [p]^2$ for the covariance functions, that is,

$$\text{AveMISE}(\Sigma) = p^{-2} \sum_j \sum_k \min_{h_{\Sigma,jk}} \text{MISE}(\hat{\Sigma}_{jk}, h_{\Sigma,jk}), \quad \text{MaxMISE}(\Sigma) = \max_{j,k} \min_{h_{\Sigma,jk}} \text{MISE}(\hat{\Sigma}_{jk}, h_{\Sigma,jk}).$$

We next use the example of estimating the mean functions to illustrate the rationale of the above measures. While $\text{AveMISE}(\mu)$ presents the averaged elementwise minimal MISEs across $j \in [p]$, some simple calculations in Appendix B show that the attainable quantity of the minimal elementwise maximum of MISEs, $\min_{(h_{\mu,1}, \dots, h_{\mu,p}) \in \mathcal{H}_\mu^p} \max_{j \in [p]} \text{MISE}(\hat{\mu}_j, h_{\mu,j})$, is equal to $\text{MaxMISE}(\mu)$.




Figure 1: Plots of the average MaxMISEs (black) and AveMISEs (red) against T with $p = 50$ (solid), 100 (dashed) and 150 (dotted) for the mean estimators (left) and the covariance estimators (right).

We consider settings of $n = 100$, $p = 50, 100, 150$, and $T = 5, 10, 20, 40, 60, 80, 100, 120, 140, 160$, varying from sparse to semi-dense to ultra-dense measurement schedules. We ran each simulation 100 times. Figure 1 plots the average AveMISEs and MaxMISEs as functions of the sampling frequency T for the estimated mean and covariance functions. A few apparent patterns are observable from Figure 1. First, both MaxMISEs and AveMISEs display a similar trend as T increases from 5 to 160, with a steep decline followed by a slight

decrease and then a period of stability. Such trend roughly corresponds to the three categories of “sparse”, “semi-dense” and “ultra-dense”, respectively. Second, while AveMISEs reflect the performance for univariate functional data, MaxMISEs gradually enlarges as p increases from 50 to 150, providing empirical evidence to support that the associated rates in high-dimensional settings all depend on $\log p$ -based multiplicative factors. Additionally, it is observable that the high-dimensionality causes the transition phase between semi-dense and ultra-dense functional data to slightly shift to the left. Third, compared to the results for the estimated mean functions, an increase in T for sparse and semi-dense functional data leads to an enhanced reduction in relative MISEs for the covariance estimators. These observations further validate our established theoretical results in Section 3.

A Technical proofs

The appendix contains proofs of all theorems. Throughout, we use c, c_1, c_2, \dots to denote generic positive finite constants that may be different in different uses.

A.1 Proof of Theorem 1

We organize the proof in four steps. First, we will define $\hat{\mu}(\cdot)$, $\tilde{\mu}(\cdot)$ and obtain the decomposition of $\hat{\mu}(\cdot) - \tilde{\mu}(\cdot)$. Second, we will prove the local concentration inequality for fixed interior point $u \in \mathcal{U}$. Third, we will prove the concentration inequality in L_2 norm. Finally, we will prove the concentration inequality in supremum norm.

A.1.1 Definition and decomposition

Without loss of generality, we let $h_{\mu,j} = h$ for $j \in [p]$ and denote $\mathbf{e}_0 = (1, 0)^\top$, $\tilde{\mathbf{U}}_{ijt} = \{1, (U_{ijt} - u)/h\}^\top$,

$$\begin{aligned}\hat{\mathbf{S}}_j(u) &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} \tilde{\mathbf{U}}_{ijt}^\top K_h(U_{ijt} - u), \\ \hat{\mathbf{R}}_j(u) &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} K_h(U_{ijt} - u) Y_{ijk}.\end{aligned}$$

A simple calculation yields $\hat{\mu}_j(u) = \mathbf{e}_0^\top \{\hat{\mathbf{S}}_j(u)\}^{-1} \hat{\mathbf{R}}_j(u)$. Let

$$\tilde{\mu}_j(u) = \mathbf{e}_0^\top [E\{\hat{\mathbf{S}}_j(u)\}]^{-1} E\{\hat{\mathbf{R}}_j(u)\}. \quad (\text{A.1})$$

We can decompose $\hat{\mu}_j(u) - \tilde{\mu}_j(u)$ as

$$\begin{aligned} \hat{\mu}_j(u) - \tilde{\mu}_j(u) &= \mathbf{e}_0^\top [E\{\hat{\mathbf{S}}_j(u)\}]^{-1} [\hat{\mathbf{R}}_j(u) - E\{\hat{\mathbf{R}}_j(u)\}] \\ &\quad - \mathbf{e}_0^\top \{\hat{\mathbf{S}}_j(u)\}^{-1} [\hat{\mathbf{S}}_j(u) - E\{\hat{\mathbf{S}}_j(u)\}] [E\{\hat{\mathbf{S}}_j(u)\}]^{-1} \hat{\mathbf{R}}_j(u), \end{aligned}$$

which then implies that

$$\begin{aligned} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| &\leq \|E\{\hat{\mathbf{S}}_j(u)\}\|_{\min}^{-1} \|\hat{\mathbf{R}}_j(u) - E\{\hat{\mathbf{R}}_j(u)\}\| \\ &\quad + \|\hat{\mathbf{S}}_j(u)\|_{\min}^{-1} \|E\{\hat{\mathbf{S}}_j(u)\}\|_{\min}^{-1} \|\hat{\mathbf{R}}_j(u)\| \|\hat{\mathbf{S}}_j(u) - E\{\hat{\mathbf{S}}_j(u)\}\|_{\text{F}}, \end{aligned} \quad (\text{A.2})$$

where, for any vector $\mathbf{b} = (b_1, \dots, b_p)^\top$, we write $\|\mathbf{b}\| = (\sum_i b_i^2)^{1/2}$ and, for any matrix $\mathbf{B} = (B_{ij})_{p \times q}$, we write $\|\mathbf{B}\|_{\min} = \{\lambda_{\min}(\mathbf{B}^\top \mathbf{B})\}^{1/2}$ and $\|\mathbf{B}\|_{\text{F}} = (\sum_{i,j} B_{ij}^2)^{1/2}$ to denote its Frobenius norm.

A.1.2 Local concentration inequality

We will firstly show that there exists some positive constant c such that for any $\delta > 0$ and $u \in \mathcal{U}$,

$$\text{pr} \left\{ \|\hat{\mathbf{S}}_j(u) - E\{\hat{\mathbf{S}}_j(u)\}\|_{\text{F}} \geq \delta \right\} \leq 8 \exp \left(- \frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta} \right). \quad (\text{A.3})$$

For $k, l = 1, 2$, let $\hat{S}_{jkl}(u)$ be the (k, l) th entry of $\hat{\mathbf{S}}_j(u)$. Under Assumptions 4 and 6, we obtain that for any integer $q = 2, 3, \dots$ and $s = 0, 1, 2$,

$$E \left\{ \left| \left(\frac{U_{ijt} - u}{h} \right)^s K_h(U_{ijt} - u) \right|^q \right\} \leq \int h^{-q} K^q \left(\frac{t-u}{h} \right) \left| \frac{t-u}{h} \right|^{sq} f_U(t) dt \leq ch^{1-q}. \quad (\text{A.4})$$

Note that Assumption 3 implies that the weights v_{ij} 's are of the same order $v_{ij} \asymp (n\bar{T}_{\mu,j})^{-1}$. By (A.4), it holds that $\sum_{i=1}^n \sum_{t=1}^{T_{ij}} E \left\{ \left| \left(\frac{U_{ijt} - u}{h} \right)^s K_h(U_{ijt} - u) \right|^2 \right\} \leq cn\bar{T}_{\mu,j}h^{-1}$, and

$$\sum_{i=1}^n \sum_{t=1}^{T_{ij}} E \left\{ \left| \left(\frac{U_{ijt} - u}{h} \right)^s K_h(U_{ijt} - u) \right|^q \right\} \leq 2^{-1} q! cn\bar{T}_{\mu,j} h^{-1} h^{2-q} \text{ for } q \geq 3.$$

By the Bernstein inequality (see Theorem 2.10 and Corollary 2.11 of Boucheron et al. (2014)), we obtain that there exists some positive constant c such that for any $\delta > 0$

$$\text{pr} \left\{ |\hat{S}_{jkl}(u) - E\{\hat{S}_{jkl}(u)\}| \geq \delta \right\} \leq 2 \exp \left(- \frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta} \right),$$

for $k, l = 1, 2$, which, by the union bound of probability, implies that (A.3) holds.

For $k = 1, 2$, let $\widehat{R}_{jk}(u)$ be the k th element of $\widehat{\mathbf{R}}_j(u)$. We will next show that there exists some positive constant c such that for any $\delta > 0$ and $u \in \mathcal{U}$,

$$\text{pr}\left\{|\widehat{R}_{jk}(u) - E\{\widehat{R}_{jk}(u)\}| \geq \delta\right\} \leq c \exp\left(-\frac{c\gamma_{n,T,h,j}\delta^2}{1+\delta}\right), \quad (\text{A.5})$$

where $\gamma_{n,T,h,j} = n(1 \wedge \bar{T}_{\mu,j}h)$. We only need to consider the case $k = 1$, while the case $k = 2$ can be demonstrated in a similar manner. Denote that $\xi_{ijt} = Y_{ijt} - \mu_j(U_{ijt})$, $\widehat{R}_{j3}(u) = \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)\mu_j(U_{ijt})$, and $\widehat{R}_{j4}(u) = \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)\xi_{ijt}$. Then $\widehat{R}_{j1}(u) - E\{\widehat{R}_{j1}(u)\}$ can be rewritten as

$$\widehat{R}_{j1}(u) - E\{\widehat{R}_{j1}(u)\} = \widehat{R}_{j3}(u) - E\{\widehat{R}_{j3}(u)\} + \widehat{R}_{j4}(u). \quad (\text{A.6})$$

Following the same procedure to prove (A.3) and using the Bernstein inequality, we can obtain that there exists some positive constant c such that for any $\delta > 0$ and $u \in \mathcal{U}$,

$$\text{pr}\left\{|\widehat{R}_{j3}(u) - E\{\widehat{R}_{j3}(u)\}| \geq \delta\right\} \leq 2 \exp\left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta}\right). \quad (\text{A.7})$$

Now we consider the tail behavior of $\widehat{R}_{j4}(u)$. Define the event $V_j = \{U_{ijt}, t \in [T_{ij}], i \in [n]\}$. It follows from sub-Gaussianities in Assumption 1 and (1) that $E\{\exp(\lambda\xi_{ijt})|V_j\} \leq \exp(\lambda^2c^2)$ for any $\lambda \in \mathbb{R}$. Rewrite $\widehat{R}_{j4}(u) = \sum_{i=1}^n v_{ij}\psi_{ij1}(u)$ with $\psi_{ij1}(u) = \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)\xi_{ijt}$. Note that for each i Assumption 3 implies that $v_{ij} \asymp (n\bar{T}_{\mu,j})^{-1}$. If $\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) > 0$, by Jensen's inequality, we have

$$\begin{aligned} & E\left[\exp\{\lambda v_{ij}\psi_{ij1}(u)\}|V_j\right] \\ & \leq \frac{1}{\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)} \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) E\left[\exp\left\{\lambda v_{ij}\xi_{ijt} \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)\right\}|V_j\right] \\ & \leq \exp\left[\lambda^2c^2(n\bar{T}_{\mu,j})^{-2}\left\{\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)\right\}^2\right]. \end{aligned}$$

It is obvious that the above inequality still holds even if $\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) = 0$. Assumption 5 implies that the number of nonzero terms in $\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u)$ has an upper bound $c(1 \vee \bar{T}_{\mu,j}h)$, which yields that $\sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \leq ch^{-1}(1 \vee \bar{T}_{\mu,j}h)$. Therefore, for any $\lambda \in \mathbb{R}$, we

obtain that

$$\begin{aligned} E \left[\exp \left\{ \lambda \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \right\} \middle| V_j \right] &\leq \exp \left[\lambda^2 c^2 \sum_{i=1}^n v_{ij}^2 \left\{ \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \right\}^2 \right] \\ &\leq \exp \left\{ \lambda^2 c^2 (n\bar{T}_{\mu,j}h)^{-2} h(1 \vee \bar{T}_{\mu,j}h) \sum_{i=1}^n \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \right\}. \end{aligned}$$

For any $\delta > 0$, define the event $\Omega_{j,1}(\delta) = \left\{ \sum_{i=1}^n \sum_{t=1}^{T_{ij}} K_h(U_{ijt} - u) \leq c(1 + \delta)n\bar{T}_{\mu,j} \right\}$. We have

$$\begin{aligned} E \left[\exp \left\{ \lambda \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \right\} \middle| \Omega_{j,1}(\delta) \right] &\leq \exp \left\{ \lambda^2 c^2 (1 + \delta) (n\bar{T}_{\mu,j}h)^{-2} n\bar{T}_{\mu,j}h(1 \vee \bar{T}_{\mu,j}h) \right\} \\ &= \exp \left\{ (1 + \delta) \lambda^2 c^2 (n\bar{T}_{\mu,j}h)^{-1} (1 \vee \bar{T}_{\mu,j}h) \right\}. \end{aligned}$$

As a consequence, we obtain that

$$\text{pr} \left\{ \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \geq \delta \middle| \Omega_{j,1}(\delta) \right\} \leq \exp \left\{ -\lambda\delta + (1 + \delta) \lambda^2 c^2 (n\bar{T}_{\mu,j}h)^{-1} (1 \vee \bar{T}_{\mu,j}h) \right\}. \quad (\text{A.8})$$

With the choice of $\lambda = \frac{n\bar{T}_{\mu,j}h\delta}{2(1+\delta)c^2(1 \vee \bar{T}_{\mu,j}h)}$, (A.8) degenerates to

$$\text{pr} \left\{ \sum_{i=1}^n v_{ij} \psi_{ij1}(u) \geq \delta \middle| \Omega_{j,1}(\delta) \right\} \leq \exp \left\{ -\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1 + \delta} \right\}. \quad (\text{A.9})$$

Note that $\sum_{i=1}^n \sum_{t=1}^{T_{ij}} E\{K_h(U_{ijt} - u)\} \leq cn\bar{T}_{\mu,j}$. By the Bernstein inequality, we obtain that there exists some positive constant c such that for any $\delta > 0$

$$\text{pr} \left(\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \left[K_h(U_{ijt} - u) - E\{K_h(U_{ijt} - u)\} \right] \geq n\bar{T}_{\mu,j}\delta \right) \leq \exp \left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1 + \delta} \right),$$

which implies that

$$1 - \text{pr}\{\Omega_{j,1}(\delta)\} \leq \exp \left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1 + \delta} \right). \quad (\text{A.10})$$

Combining (A.9) and (A.10), we obtain that there exists some constant $c > 0$ such that for any $\delta > 0$

$$\text{pr} \left\{ \widehat{R}_{j4}(u) \geq \delta \right\} \leq 2 \exp \left\{ -\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1 + \delta} \right\}.$$

and, consequently,

$$\text{pr} \left\{ |\widehat{R}_{j4}(u)| \geq \delta \right\} \leq 4 \exp \left\{ -\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1 + \delta} \right\}.$$

This together with (A.6) and (A.7) yields that

$$\Pr\left\{\left|\widehat{R}_{j1}(u) - E\{\widehat{R}_{j1}(u)\}\right| \geq \delta\right\} \leq 6 \exp\left\{-\frac{cn(1 \wedge \bar{T}_{\mu,j}h)\delta^2}{1+\delta}\right\}.$$

Define the event $\Omega_{j,2}(\delta) = \{\|\widehat{\mathbf{S}}_j(u) - E\{\widehat{\mathbf{S}}_j(u)\}\|_{\text{F}} \leq \delta/2\}$. Note that $E\{\widehat{\mathbf{S}}_j(u)\}$ is positive definite. On the event $\Omega_{j,2}(\delta)$ with $\delta \in (0, 1]$, we obtain that

$$\|\widehat{\mathbf{S}}_j(u)\|_{\min} \geq c(1 - \delta/2). \quad (\text{A.11})$$

By (A.3), we have

$$1 - \Pr\{\Omega_{j,2}(\delta)\} \leq 8 \exp\left(-\frac{cn\bar{T}_{\mu,j}h\delta^2}{1+\delta}\right). \quad (\text{A.12})$$

Define the event $\Omega_{j,3}(\delta) = \{\|\widehat{\mathbf{R}}_j(u) - E\{\widehat{\mathbf{R}}_j(u)\}\| \leq \delta\}$. Note that, under Assumption 3 with $v_{ij} \asymp (n\bar{T}_{\mu,j})^{-1}$, $\sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} E\{K_h(U_{ijt} - u)\} \leq c$ and $\mu_j(\cdot)$ is uniformly bounded over \mathcal{U} , hence $\|E\{\widehat{\mathbf{R}}_j(u)\}\|$ is uniformly bounded over \mathcal{U} . On the event $\Omega_{j,3}(\delta)$, we have

$$\|\widehat{\mathbf{R}}_j(u)\| \leq c(1 + \delta). \quad (\text{A.13})$$

On the event $\Omega_{j,2}(\delta) \cap \Omega_{j,3}(\delta)$ with $\delta \in (0, 1]$, it follows from (A.2), (A.11) and (A.13) that

$$|\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \leq c\delta + c(1 - \delta/2)^{-1}(1 + \delta)\delta \leq c_3\delta.$$

This together with concentration inequalities in (A.5) and (A.12) imply that there exist some positive universal constants c_1 and c_2 such that for any $\delta \in (0, 1]$ and $u \in \mathcal{U}$,

$$\Pr\left\{|\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \geq \delta\right\} \leq c_2 \exp(-c_1\gamma_{n,T,h,j}\delta^2),$$

which completes the proof of local concentration inequality for the mean estimator.

A.1.3 Concentration inequality in L_2 norm

We will derive the L_2 concentration inequality of $\|\widehat{\mu}_j - \widetilde{\mu}_j\|$. Let

$$\widetilde{\mathbf{S}}_j(u) = (n\bar{T}_j)^{-1} \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \widetilde{\mathbf{U}}_{ijt} \widetilde{\mathbf{U}}_{ijt}^{\text{T}} K_h(U_{ijt} - u).$$

Then we have that $\|\widehat{\mathbf{S}}_j(u)\|_{\min} \geq c\|\widetilde{\mathbf{S}}_j(u)\|_{\min}$. We now give a lower bound on $\|\widetilde{\mathbf{S}}_j(u)\|_{\min}$. Denote $W = \sup_{u \in \mathcal{U}} \|\widetilde{\mathbf{S}}_j(u) - E\{\widetilde{\mathbf{S}}_j(u)\}\|_{\text{F}}$. In an analogy to Lemma 13.5 of Boucheron et al. (2014), we can show that $E(W) \leq c(n\bar{T}_j h)^{-1/2}$. Moreover, it follows from the facts

$|((U_{ijt} - u)^a h^{-a} K_h(U_{ijt} - u)| - E\{|(U_{ijt} - u)^a h^{-a} K_h(U_{ijt} - u)|\}) = 0$, $E[(n\bar{T}_j)^{-2}\{(U_{ijt} - u)^a h^{-a} K_h(U_{ijt} - u)\}^2] \leq c(n\bar{T}_j)^{-2} h^{-1} \leq 1$ for $a = 0, 1, 2$, and Theorem 11.10 of [Boucheron et al. \(2014\)](#) that the variance of W is bounded by $c(n\bar{T}_j h)^{-1}$. Applying Theorem 12.5 of [Boucheron et al. \(2014\)](#) yields that there exists some positive constant c such that, for any $\delta > 0$,

$$\text{pr}\{W - E(W) > \delta\} \leq 4 \exp\left(-\frac{cn\bar{T}_j h \delta^2}{1 + \delta}\right). \quad (\text{A.14})$$

Define the event $\Omega_{j,4}(\delta) = \{\sup_{u \in \mathcal{U}} \|\tilde{\mathbf{S}}_j(u) - E\{\tilde{\mathbf{S}}_j(u)\}\|_{\mathbb{F}} \leq \delta/2\}$ with $\delta \in (0, 1]$. By (A.14), we obtain that there exists some constant $c > 0$ such that, for any $\delta \in (0, 1]$,

$$1 - \text{pr}\{\Omega_{j,4}(\delta)\} \leq 4 \exp(-cn\bar{T}_j h \delta^2).$$

On the event $\Omega_{j,4} = \Omega_{j,4}(\delta_1)$ with $c(n\bar{T}_j h)^{-1/2} < \delta_1 \leq 1$, $\|\hat{\mathbf{S}}_j(u)\|_{\min} \geq c\|\tilde{\mathbf{S}}_j(u)\|_{\min} \geq c(1 - \delta_1/2) \geq c/2$. Note that $E\{\hat{\mathbf{S}}_j(u)\}$ is positive definite and $\|E\{\hat{\mathbf{R}}_j(u)\}\|$ is uniformly bounded over \mathcal{U} . On the event $\Omega_{j,4}$, it thus follows from (A.2) and $\|\hat{\mathbf{R}}_j(u)\| \leq \|\hat{\mathbf{R}}_j(u) - E\{\hat{\mathbf{R}}_j(u)\}\| + \|E\{\hat{\mathbf{R}}_j(u)\}\|$ that

$$|\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \leq c\|\hat{\mathbf{R}}_j(u) - E\{\hat{\mathbf{R}}_j(u)\}\| + c\|\hat{\mathbf{S}}_j(u) - E\{\hat{\mathbf{S}}_j(u)\}\|_{\mathbb{F}}. \quad (\text{A.15})$$

Combining (A.15) with (A.3), (A.5) and applying the first part of Lemma 6 in [Guo and Qiao \(2023\)](#) yields that, for any $u \in \mathcal{U}$ and integer $q \geq 1$,

$$E\left\{|\hat{\mu}_j(u) - \tilde{\mu}_j(u)|^{2q} \mid \Omega_{j,4}\right\} \leq q! c \left(\frac{4}{c\gamma_{n,T,h,j}}\right)^q + (2q)! c \left(\frac{4}{c\gamma_{n,T,h,j}}\right)^{2q}.$$

Applying the second part of Lemma 6 in [Guo and Qiao \(2023\)](#), we can show that, for each $\delta \in (0, 1]$, (4) in Theorem 1 holds, which completes the proof of concentration inequality for the mean estimator in L_2 norm.

A.1.4 Concentration inequality in supremum norm

We will derive the uniform concentration bound of $\sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)|$. We partition the interval $\mathcal{U} = [0, 1]$ into N subintervals I_s for $s \in [N]$ of equal length. Let u_s be the center of I_s , then we have

$$\sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \leq \max_{s \in [N]} \left[|\hat{\mu}_j(u_s) - \tilde{\mu}_j(u_s)| + \left| \{\hat{\mu}_j(u_s) - \hat{\mu}_j(u)\} - \{\tilde{\mu}_j(u_s) - \tilde{\mu}_j(u)\} \right| \right].$$

We need to bound the second term. By some calculations, it suffices to bound $\left| \{\widehat{R}_{jk}(u) - \widehat{R}_{jk}(u_s)\} - [E\{\widehat{R}_{jk}(u)\} - E\{\widehat{R}_{jk}(u_s)\}] \right|$ and $\left| \{\widehat{S}_{jkl}(u) - \widehat{S}_{jkl}(u_s)\} - [E\{\widehat{S}_{jkl}(u)\} - E\{\widehat{S}_{jkl}(u_s)\}] \right|$ for $k, l = 1, 2$, which means that we need to bound $|\widehat{R}_{jk}(u) - \widehat{R}_{jk}(u_s)|$ and $|\widehat{S}_{jkl}(u) - \widehat{S}_{jkl}(u_s)|$. Let $u \in I_s$ and consider $|\widehat{R}_{j1}(u) - \widehat{R}_{j1}(u_s)|$ first. Define the event $\Omega_{R,j1} = \{|\sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} Y_{ijt}| \leq E(|\sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} Y_{ijt}|) + 1\}$. On this event, it follows from Assumption 6(ii) that

$$\begin{aligned} |\widehat{R}_{j1}(u) - \widehat{R}_{j1}(u_s)| &\leq \left| \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} Y_{ijt} \{K_h(U_{ijt} - u) - K_h(U_{ijt} - u_s)\} \right| \\ &\leq \frac{c|u - u_s|}{h} \left| \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} Y_{ijt} \right| \leq \frac{c}{Nh} \left\{ E \left(\left| \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} Y_{ijt} \right| \right) + 1 \right\} \leq \frac{c}{Nh}. \end{aligned}$$

Applying similar techniques as above, we can define events $\Omega_{R,jk}$ and $\Omega_{S,jkl}$ for $k, l = 1, 2$. On the intersection of these events, we can obtain that $|\widehat{R}_{jk}(u) - \widehat{R}_{jk}(u_s)| \leq c(Nh)^{-1}$ and $|\widehat{S}_{jkl}(u) - \widehat{S}_{jkl}(u_s)| \leq c(Nh)^{-1}$. Combing the above results, we have

$$\sup_{u \in \mathcal{U}} |\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \leq \max_{s \in [N]} |\widehat{\mu}_j(u_s) - \widetilde{\mu}_j(u_s)| + \frac{c}{Nh}.$$

By the Bernstein inequality, we have $\text{pr}(\Omega_{R,jk}^c) \leq c \exp(-cn) \leq c \exp(-c\gamma_{n,T,h,j})$ and $\text{pr}(\Omega_{S,jkl}^c) \leq c \exp(-cn) \leq c \exp(-c\gamma_{n,T,h,j})$ for $k, l = 1, 2$. It follows from the above results and the union bound of probability with the choice of $N = c(h\delta)^{-1}$ that there exist some positive constants c_1 and c_2 such that, for any $\delta \in (0, 1]$,

$$\text{pr} \left\{ \sup_{u \in \mathcal{U}} |\widehat{\mu}_j(u) - \widetilde{\mu}_j(u)| \geq \delta \right\} \leq \frac{c_2}{h\delta} \exp(-c_1\gamma_{n,T,h,j}\delta^2). \quad (\text{A.16})$$

Take arbitrarily small $\epsilon_1 > 0$. If $n^{\epsilon_1}\gamma_{n,T,h,j}\delta^2 \geq 1$, then the right side of (A.16) reduces to $c_2\{n^{\epsilon_1}\gamma_{n,T,h,j}\}^{1/2}h^{-1}\exp(-c_1\gamma_{n,T,h,j}\delta^2)$. If $n^{\epsilon_1}\gamma_{n,T,h,j}\delta^2 \leq 1$, we can choose c_2 and $n^{\epsilon_1} > c$ such that $c_2 \exp(-c_1c^{-1}) \geq 1$ and the same bound $c_1\{n^{\epsilon_1}\gamma_{n,T,h,j}\}^{1/2}h^{-1}\exp(-c_1\gamma_{n,T,h,j}\delta^2)$ can therefore still be used. Hence (5) in Theorem 1 holds, which completes the proof of concentration inequality for the mean estimator in supremum norm.

A.2 Proofs of Theorem 2

We organize the proof in four steps. First, we will define $\widehat{\Sigma}(\cdot, \cdot)$, $\widetilde{\Sigma}(\cdot, \cdot)$ and obtain the decomposition of $\widehat{\Sigma}(\cdot, \cdot) - \widetilde{\Sigma}(\cdot, \cdot)$. Second, we will prove the local concentration inequality

for fixed $(u, v) \in \mathcal{U}^2$. Third, we will prove the concentration inequality in Hilbert–Schmidt norm. Finally, we will prove the concentration inequality in supremum norm.

A.2.1 Definition and decomposition

Without loss of generality, let $h_{\Sigma, jk} = h$ for $(j, k) \in [p]^2$ and denote $\tilde{\mathbf{e}}_0 = (1, 0, 0)^\top$, $\tilde{\mathbf{U}}_{ijjts}(u, v) = \{1, (U_{ijt} - u)/h, (U_{iks} - v)/h\}^\top$. For $j = k$, let

$$\begin{aligned}\hat{\mathbf{\Xi}}_{jj}(u, v) &= \sum_{i=1}^n w_{ijj} \sum_{1 \leq t \neq s \leq T_{ij}} \tilde{\mathbf{U}}_{ijjts}(u, v) \tilde{\mathbf{U}}_{ijjts}^\top(u, v) K_h(U_{ijt} - u) K_h(U_{ijs} - v), \\ \hat{\mathbf{Z}}_{jj}(u, v) &= \sum_{i=1}^n w_{ijj} \sum_{1 \leq t \neq s \leq T_{ij}} \tilde{\mathbf{U}}_{ijjts}(u, v) \Theta_{ijjts} K_h(U_{ijt} - u) K_h(U_{ijs} - v).\end{aligned}$$

For $j \neq k$, let

$$\begin{aligned}\hat{\mathbf{\Xi}}_{jk}(u, v) &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijjts}(u, v) \tilde{\mathbf{U}}_{ijjts}^\top(u, v) K_h(U_{ijt} - u) K_h(U_{iks} - v), \\ \hat{\mathbf{Z}}_{jk}(u, v) &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijjts}(u, v) \Theta_{ijjts} K_h(U_{ijt} - u) K_h(U_{iks} - v).\end{aligned}$$

A simple calculation yields $\hat{\Sigma}_{jk}(u, v) = \tilde{\mathbf{e}}_0^\top \{\hat{\mathbf{\Xi}}_{jk}(u, v)\}^{-1} \hat{\mathbf{Z}}_{jk}(u, v)$. Let

$$\tilde{\Sigma}_{jk}(u, v) = \tilde{\mathbf{e}}_0^\top [E\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}]^{-1} E\{\hat{\mathbf{Z}}_{jk}(u, v)\}. \quad (\text{A.17})$$

We can decompose $\hat{\Sigma}_{jk}(u, v) - \tilde{\Sigma}_{jk}(u, v)$ as

$$\begin{aligned}\hat{\Sigma}_{jk}(u, v) - \tilde{\Sigma}_{jk}(u, v) &= \tilde{\mathbf{e}}_0^\top (\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}^{-1} - [E\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}]^{-1}) \hat{\mathbf{Z}}_{jk}(u, v) \\ &\quad + \tilde{\mathbf{e}}_0^\top [E\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}]^{-1} [\hat{\mathbf{Z}}_{jk}(u, v) - E\{\hat{\mathbf{Z}}_{jk}(u, v)\}],\end{aligned}$$

which further implies that

$$\begin{aligned}& |\hat{\Sigma}_{jk}(u, v) - \tilde{\Sigma}_{jk}(u, v)| \\ & \leq \|E\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}\|_{\min}^{-1} \|\hat{\mathbf{Z}}_{jk}(u, v) - E\{\hat{\mathbf{Z}}_{jk}(u, v)\}\| \\ & \quad + \|E\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}\|_{\min}^{-1} \|\hat{\mathbf{\Xi}}_{jk}(u, v)\|_{\min}^{-1} \|\hat{\mathbf{Z}}_{jk}(u, v)\| \|\hat{\mathbf{\Xi}}_{jk}(u, v) - E\{\hat{\mathbf{\Xi}}_{jk}(u, v)\}\|_{\text{F}}.\end{aligned} \quad (\text{A.18})$$

In the following, we will prove the concentration results for the case $j \neq k$, and the results for the case $j = k$ can be proved in a similar manner.

A.2.2 Local concentration inequality

We will firstly show that there exists some positive constant c such that for any $\delta > 0$ and $(u, v) \in \mathcal{U}^2$,

$$\text{pr} \left\{ \left\| \widehat{\Xi}_{jk}(u, v) - E\{\widehat{\Xi}_{jk}(u, v)\} \right\|_{\text{F}} \geq \delta \right\} \leq 18 \exp \left(- \frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right). \quad (\text{A.19})$$

For $m, l = 1, 2, 3$, let $\widehat{\Xi}_{jkm}(u, v)$ be the (m, l) th entry of $\widehat{\Xi}_{jk}(u, v)$. It follows from Assumptions 4 and 6 that for any integer $q = 2, 3, \dots$ and $s, s' = 0, 1, 2$,

$$\begin{aligned} & E \left\{ \left| \left(\frac{U_{ijt} - u}{h} \right)^s \frac{U_{ikt'} - v}{h} \right|^{s'} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \right\} \\ & \leq \int h^{-2q} K^q \left(\frac{t - u}{h} \right) K^q \left(\frac{t' - v}{h} \right) \left| \frac{t - u}{h} \right|^{sq} \left| \frac{t' - v}{h} \right|^{s'q} f_U(t) f_U(t') dt dt' \leq ch^{2-2q}. \end{aligned} \quad (\text{A.20})$$

Note that Assumption 3 implies that the weights are of the same order $w_{ijk} \asymp (n\bar{T}_{jk}^2)^{-1}$. By (A.20), $\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} E \left\{ \left| \left(\frac{U_{ijt} - u}{h} \right)^s \left(\frac{U_{ikt'} - v}{h} \right)^{s'} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \right|^2 \right\} \leq cn\bar{T}_{jk}^2 h^{-2}$,

$$\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} E \left\{ \left| \left(\frac{U_{ijt} - u}{h} \right)^s \left(\frac{U_{ikt'} - v}{h} \right)^{s'} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \right|^q \right\} \leq 2^{-1} q! cn\bar{T}_{jk}^2 h^{2-2q}$$

for $q \geq 3$. Applying the Bernstein inequality yields that there exists some positive constant c such that for any $\delta > 0$

$$\text{pr} \left\{ \left| \widehat{\Xi}_{jkm}(u, v) - E\{\widehat{\Xi}_{jkm}(u, v)\} \right| \geq \delta \right\} \leq 2 \exp \left(- \frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right),$$

for $m, l = 1, 2, 3$, which, by the union bound of probability, implies that (A.19) holds.

For $m = 1, 2, 3$, let $\widehat{Z}_{jkm}(u, v)$ be the m th element of $\widehat{\mathbf{Z}}_{jk}(u, v)$. We will next show that, there exists some positive constant c such that for any $\delta > 0$ and $(u, v) \in \mathcal{U}^2$,

$$\text{pr} \left\{ \left| \widehat{Z}_{jkm}(u, v) - E\{\widehat{Z}_{jkm}(u, v)\} \right| \geq \delta \right\} \leq c \exp \left(- \frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right), \quad (\text{A.21})$$

where $\nu_{n,T,h,jk} = n(1 \wedge \bar{T}_{jk}^2 h^2)$. We only need to consider the case $m = 1$, while the results for cases $m = 2, 3$ can be proved in a similar way. Denote that $\zeta_{ijks} = \{Y_{ijt} - \mu_j(U_{ijt})\} \{Y_{iks} - \mu_k(U_{iks})\} - \Sigma_{jk}(U_{ijt}, U_{iks})$, $\widehat{Z}_{jk4}(u, v) = \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{ikt'} - v) \Sigma_{jk}(U_{ijt}, U_{iks})$, and $\widehat{Z}_{jk5}(u, v) = \sum_{i=1}^n w_{ij} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) \zeta_{ijks}$. Then we rewrite $\widehat{Z}_{jk1}(u, v) - E\{\widehat{Z}_{jk1}(u, v)\}$ as

$$\widehat{Z}_{jk1}(u, v) - E\{\widehat{Z}_{jk1}(u, v)\} = \widehat{Z}_{jk4}(u, v) - E\{\widehat{Z}_{jk4}(u, v)\} + \widehat{Z}_{jk5}(u, v). \quad (\text{A.22})$$

Following the same procedure to prove (A.19) with the aid of the Bernstein inequality, we can obtain that there exists some positive constant c such that for any $\delta > 0$ and $(u, v) \in \mathcal{U}^2$,

$$\text{pr}\left\{\left|\widehat{Z}_{jk4}(u, v) - E\{\widehat{Z}_{jk4}(u, v)\}\right| \geq \delta\right\} \leq 2 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \quad (\text{A.23})$$

Now we consider the tail behavior of $\widehat{Z}_{jk5}(u, v)$. Define the event $\widetilde{V}_{jk} = \{(U_{ijt}, U_{iks}), t \in [T_{ij}], s \in [T_{ik}], i \in [n]\}$. The sub-Gaussianities under Assumption 1 implies that, conditional on the event \widetilde{V}_{jk} , $Y_{ijt} - \mu_j(U_{ijt})$ and $Y_{iks} - \mu_k(U_{iks})$ are sub-Gaussian random variables, then $\{Y_{ijt} - \mu_j(U_{ijt})\}\{Y_{iks} - \mu_k(U_{iks})\}$ is a sub-exponential random variable, and hence we have $E\{\exp(\lambda\zeta_{ijkt})|\widetilde{V}_{jk}\} \leq \exp\{(1-c\lambda)^{-1}c\lambda^2\}$ for any $\lambda \in (0, c^{-1})$. Rewrite $\widehat{Z}_{jk5}(u, v) = \sum_{i=1}^n w_{ijk}\phi_{ijk1}(u, v)$ with $\phi_{ijk1}(u, v) = \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v)\zeta_{ijkt}$. Note that for each i Assumption 3 implies that $w_{ijk} \asymp (n\bar{T}_{jk}^2)^{-1}$. If $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v) > 0$ holds, it follows from Jensen's inequality that

$$\begin{aligned} & E\left[\exp\{\lambda w_{ijk}\phi_{ijk1}(u, v)\}|\widetilde{V}_{jk}\right] \\ & \leq \frac{1}{\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v)} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \left(K_h(U_{ijt}-u)K_h(U_{iks}-v)\right) \\ & \quad \times E\left[\exp\left\{\lambda w_{ijk}\zeta_{ijkt} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v)\right\}|\widetilde{V}_{jk}\right] \\ & \leq \exp\left[\frac{c\lambda^2(n\bar{T}_{jk}^2)^{-2}\left\{\sum_{t,s} K_h(U_{ijt}-u)K_h(U_{iks}-v)\right\}^2}{1-c\lambda(n\bar{T}_{jk}^2)^{-1}\sum_{t,s} K_h(U_{ijt}-u)K_h(U_{iks}-v)}\right]. \end{aligned}$$

where $0 < \lambda(n\bar{T}_{jk}^2)^{-1}\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v) < c^{-1}$. It is obvious that the above inequality still holds even if $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v) = 0$. Assumption 5 implies that the number of nonzero terms in $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v)$ has an upper bound $c(1 \vee \bar{T}_{jk}^2 h^2)$, which yields that $\sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v) \leq ch^{-2}(1 \vee \bar{T}_{jk}^2 h^2)$. Therefore, for any λ satisfying $0 < \lambda(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2) < c^{-1}$ for some constant $c > 0$, we obtain that

$$\begin{aligned} & E\left[\exp\left\{\lambda \sum_{i=1}^n w_{ijk}\phi_{ijk1}(u, v)\right\}|\widetilde{V}_{jk}\right] \\ & \leq \exp\left\{\frac{c\lambda^2(n\bar{T}_{jk}^2)^{-2}h^{-2}(1 \vee \bar{T}_{jk}^2 h^2) \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt}-u)K_h(U_{iks}-v)}{1-c\lambda(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2)}\right\}. \end{aligned}$$

For any $\delta > 0$, define the event

$$\Lambda_{jk,1}(\delta) = \left\{ \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} K_h(U_{ijt} - u) K_h(U_{iks} - v) \leq c(1 + \delta) n \bar{T}_{jk}^2 \right\}.$$

We have

$$E \left[\exp \left\{ \lambda \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u, v) \right\} \middle| \Lambda_{jk,1}(\delta) \right] \leq \exp \left\{ \frac{c\lambda^2(1 + \delta)(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2)}{1 - c\lambda(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2)} \right\}.$$

Consequently, we obtain that

$$\text{pr} \left\{ \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u) \geq \delta \middle| \Lambda_{jk,1}(\delta) \right\} \leq \exp \left\{ -\lambda\delta + \frac{c\lambda^2(1 + \delta)(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2)}{1 - c\lambda(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2)} \right\}. \quad (\text{A.24})$$

With the choice of $\lambda = \frac{n\bar{T}_{jk}^2 h^2 \delta}{2c(1+\delta)(1 \vee \bar{T}_{jk}^2 h^2) + c\delta(1 \vee \bar{T}_{jk}^2 h^2)}$, (A.24) reduces to

$$\text{pr} \left\{ \sum_{i=1}^n w_{ijk} \phi_{ijk1}(u, v) \geq \delta \middle| \Lambda_{jk,1}(\delta) \right\} \leq \exp \left\{ -\frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right\}, \quad (\text{A.25})$$

where the constant c is chosen to satisfy $c\lambda(n\bar{T}_{jk}^2 h^2)^{-1}(1 \vee \bar{T}_{jk}^2 h^2) \leq 1/2$. Note that

$$\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} E \{ K_h(U_{ijt} - u) K_h(U_{iks} - v) \} \leq cn\bar{T}_{jk}^2.$$

By the Bernstein inequality, we obtain that there exists some positive constant c such that for any $\delta > 0$

$$\begin{aligned} & \text{pr} \left(\sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \left[K_h(U_{ijt} - u) K_h(U_{iks} - v) - E \{ K_h(U_{ijt} - u) K_h(U_{iks} - v) \} \right] \geq n\bar{T}_{jk}^2 \delta \right) \\ & \leq \exp \left(\frac{-c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right) \end{aligned}$$

which implies that

$$1 - \text{pr} \{ \Lambda_{jk,1}(\delta) \} \leq \exp \left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right). \quad (\text{A.26})$$

Combining (A.25) and (A.26), we obtain that there exists some constant $c > 0$ such that for any $\delta > 0$,

$$\text{pr} \left\{ \hat{Z}_{jk5}(u, v) \geq \delta \right\} \leq 2 \exp \left\{ -\frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right\},$$

which leads to

$$\text{pr} \left\{ |\hat{Z}_{jk5}(u, v)| \geq \delta \right\} \leq 4 \exp \left\{ -\frac{c\nu_{n,T,h,jk}\delta^2}{1 + \delta} \right\}.$$

It follows from the above, (A.22) and (A.23) that for each $\delta > 0$ and $(u, v) \in \mathcal{U}^2$,

$$\text{pr} \left\{ \left| \widehat{Z}_{jk1}(u, v) - E\{\widehat{Z}_{jk1}(u, v)\} \right| \geq \delta \right\} \leq 6 \exp \left\{ -\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta} \right\}.$$

Define the event $\Lambda_{jk,2}(\delta) = \{\|\widehat{\Xi}_{jk}(u, v) - E\{\widehat{\Xi}_{jk}(u, v)\}\|_{\text{F}} \leq \delta/2\}$. Note that $E\{\widehat{\Xi}_{jk}(u, v)\}$ is positive definite. On the event $\Lambda_{jk,2}(\delta)$ with $\delta \in (0, 1]$, we obtain that

$$\|\widehat{\Xi}_{jk}(u, v)\|_{\min} \geq c(1 - \delta/2). \quad (\text{A.27})$$

By (A.19), we have

$$1 - \text{pr}\{\Lambda_{jk,2}(\delta)\} \leq 18 \exp \left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta} \right). \quad (\text{A.28})$$

Define the event $\Lambda_{jk,3}(\delta) = \{\|\widehat{\mathbf{Z}}_{jk}(u, v) - E\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\| \leq \delta\}$. Note that, under Assumption 3 with $w_{ijk} = (n\bar{T}_{jk}^2)^{-1}$, $\sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} E\{K_h(U_{ijt} - u)K_h(U_{iks} - v)\} \leq c$, hence $\|E\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\|$ is uniformly bounded over \mathcal{U}^2 . On the event $\Omega_{jk,3}(\delta)$, we have

$$\|\widehat{\mathbf{Z}}_{jk}(u, v)\| \leq c(1 + \delta). \quad (\text{A.29})$$

On the event $\Lambda_{jk,2}(\delta) \cap \Lambda_{jk,3}(\delta)$ with $\delta \in (0, 1]$, it follows from (A.18), (A.27) and (A.29) that

$$|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \leq c\delta + c(1 - \delta/2)^{-1}(1 + \delta)\delta \leq c_3\delta.$$

This together with concentration inequalities in (A.21) and (A.28) imply that there exist some positive universal constants c_1 and c_2 such that for any $\delta \in (0, 1]$ and $(u, v) \in \mathcal{U}^2$,

$$\text{pr} \left\{ |\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \geq \delta \right\} \leq c_2 \exp(-c_1\nu_{n,T,h,jk}\delta^2),$$

which completes the proof of local concentration inequality for the covariance estimator.

A.2.3 Concentration inequality in Hilbert–Schmidt norm

We will derive the L_2 concentration inequality of $\|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_{\mathcal{S}}$. Let

$$\widetilde{\Xi}_{jk}(u, v) = (n\bar{T}_{jk}^2)^{-1} \sum_{i=1}^n \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \widetilde{U}_{ij kts} \widetilde{U}_{ij kts}^{\text{T}} K_h(U_{ijt} - u) K_h(U_{iks} - v).$$

Then we have that $\|\widehat{\Xi}_{jk}(u, v)\|_{\min} \geq c\|\widetilde{\Xi}_{jk}(u, v)\|_{\min}$. Similar to Appendix A.1.3, we will give a lower bound on $\|\widetilde{\Xi}_{jk}(u, v)\|_{\min}$. Denote $\widetilde{W} = \sup_{(u,v) \in \mathcal{U}^2} \|\widetilde{\Xi}_{jk}(u, v) - E\{\widetilde{\Xi}_{jk}(u, v)\}\|_{\text{F}}$.

In an analogy to Lemma 13.5 and Theorem 11.10 of [Boucheron et al. \(2014\)](#), we can show that $E(\widetilde{W}) \leq c(\nu_{n,T,h,jk})^{-1/2}$. Moreover, it follows from the facts $|(U_{ijt} - u)^a h^{-a} (U_{iks} - v)^b h^{-b} K_h(U_{ijt} - u) K_h(U_{iks} - v)| - E\{|(U_{ijt} - u)^a h^{-a} (U_{iks} - v)^b h^{-b} K_h(U_{ijt} - u) K_h(U_{iks} - v)|\} = 0$, $E[(n\bar{T}_{jk}^2)^{-2} \{(U_{ijt} - u)^a h^{-a} (U_{iks} - v)^b h^{-b} K_h(U_{ijt} - u) K_h(U_{iks} - v)\}^2] \leq c(n\bar{T}_{jk}^2)^{-2} h^{-2} \leq 1$ for $a, b = 0, 1, 2$, and Theorem 11.10 of [Boucheron et al. \(2014\)](#) that the variance of \widetilde{W} is bounded by $c(\nu_{n,T,h,jk})^{-1}$. Applying Theorem 12.5 of [Boucheron et al. \(2014\)](#), we obtain that there exist some positive constant c such that, for any $\delta > 0$,

$$\text{pr}\left\{\widetilde{W} - E(\widetilde{W}) > \delta\right\} \leq 9 \exp\left(-\frac{c\nu_{n,T,h,jk}\delta^2}{1+\delta}\right). \quad (\text{A.30})$$

Define the event $\Lambda_{jk,4}(\delta) = \{\sup_{(u,v) \in \mathcal{U}^2} \|\widetilde{\mathbf{\Xi}}_{jk}(u, v) - E\{\widetilde{\mathbf{\Xi}}_{jk}(u, v)\}\|_{\text{F}} \leq \delta/2\}$ with $\delta \in (0, 1]$. By [\(A.30\)](#), we obtain that there exists some constant $c > 0$ such that, for any $\delta \in (0, 1]$,

$$1 - \text{pr}\{\Lambda_{jk,4}(\delta)\} \leq 9 \exp(-c\nu_{n,T,h,jk}\delta^2).$$

On the event $\Lambda_{jk,4} = \Lambda_{jk,4}(\tilde{\delta}_1)$ with $c(\nu_{n,T,h,jk})^{-1/2} < \tilde{\delta}_1 \leq 1$, we have $\|\widehat{\mathbf{\Xi}}_{jk}(u, v)\|_{\min} \geq c\|\widetilde{\mathbf{\Xi}}_{jk}(u, v)\|_{\min} \geq c(1 - \tilde{\delta}_1/2) \geq c/2$. Note that $E\{\widehat{\mathbf{\Xi}}_{jk}(u, v)\}$ is positive definite and $\|E\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\|$ is uniformly bounded over \mathcal{U}^2 . On the event $\Lambda_{jk,4}$, it thus follows from [\(A.18\)](#) and $\|\widehat{\mathbf{Z}}_{jk}(u, v)\| \leq \|\widehat{\mathbf{Z}}_{jk}(u, v) - E\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\| + \|E\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\|$ that

$$|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)| \leq c\|\widehat{\mathbf{Z}}_{jk}(u, v) - E\{\widehat{\mathbf{Z}}_{jk}(u, v)\}\| + c\|\widehat{\mathbf{\Xi}}_{jk}(u, v) - E\{\widehat{\mathbf{\Xi}}_{jk}(u, v)\}\|_{\text{F}}. \quad (\text{A.31})$$

Combining [\(A.31\)](#) with [\(A.19\)](#), [\(A.21\)](#) and applying the first part of Lemma 6 in [Guo and Qiao \(2023\)](#) yields that, for any $(u, v) \in \mathcal{U}^2$ and integer $q \geq 1$,

$$E\left\{|\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)|^{2q} \mid \Lambda_{jk,4}\right\} \leq q!c\left(\frac{4}{c\nu_{n,T,h,jk}}\right)^q + (2q)!c\left(\frac{4}{c\nu_{n,T,h,jk}}\right)^{2q}.$$

Applying the second part of Lemma 6 in [Guo and Qiao \(2023\)](#), we can show that, for each $\delta \in (0, 1]$, [\(6\)](#) in Theorem 2 holds, which completes the proof of concentration inequality for the covariance estimator in Hilbert–Schmidt norm.

A.2.4 Concentration inequality in supremum norm

We will derive the uniform concentration bound of $\sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u, v) - \widetilde{\Sigma}_{jk}(u, v)|$. We partition the interval $\mathcal{U} = [0, 1]$ into N subintervals I_s for $s \in [N]$ of equal length. Let u_s

and $v_{s'}$ be the centers of I_s and $I_{s'}$, respectively, then we have

$$\begin{aligned} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)| &\leq \max_{s,s' \in [N]} \left[|\widehat{\Sigma}_{jk}(u_s, v_{s'}) - \widetilde{\Sigma}_{jk}(u_s, v_{s'})| \right. \\ &\quad \left. + |\{\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u_s, v_{s'})\} - \{\widetilde{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u_s, v_{s'})\}| \right]. \end{aligned}$$

We need to bound the second term. By some calculations, it suffices to bound $\left| \{\widehat{Z}_{jkm}(u,v) - \widehat{Z}_{jkm}(u_s, v_{s'})\} - [E\{\widehat{Z}_{jkm}(u,v)\} - E\{\widehat{Z}_{jkm}(u_s, v_{s'})\}] \right|$ and $\left| \{\widehat{\Xi}_{jklm}(u,v) - \widehat{\Xi}_{jklm}(u_s, v_{s'})\} - [E\{\widehat{\Xi}_{jklm}(u,v)\} - E\{\widehat{\Xi}_{jklm}(u_s, v_{s'})\}] \right|$ for $m, l = 1, 2, 3$, which means that we need to bound $|\widehat{Z}_{jkm}(u,v) - \widehat{Z}_{jkm}(u_s, v_{s'})|$ and $|\widehat{\Xi}_{jklm}(u,v) - \widehat{\Xi}_{jklm}(u_s, v_{s'})|$. Let $(u,v) \in I_s \times I_{s'}$ and consider $|\widehat{Z}_{jk1}(u,v) - \widehat{Z}_{jk1}(u_s, v_{s'})|$ for the case of $j \neq k$ first. The results for the case of $j = k$ can be proved in a similar fashion. Define the event $\Lambda_{Z,jk1} = \{|\sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \Theta_{ijktt'}| \leq E(|\sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \Theta_{ijktt'}|) + 1\}$. On this event, it follows from Assumption 6(ii) that

$$\begin{aligned} &\left| \{\widehat{Z}_{jk1}(u,v) - \widehat{Z}_{jk1}(u_s, v_{s'})\} \right| \\ &\leq \left| \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \Theta_{ijktt'} \left[\{K_h(U_{ijt} - u) - K_h(U_{ijt} - u_s)\} K_h(U_{ikt'} - v) \right. \right. \\ &\quad \left. \left. + \{K_h(U_{ikt'} - v) - K_h(U_{ikt'} - v_{s'})\} K_h(U_{ijt} - u_s) \right] \right| \\ &\leq \frac{c(|u - u_s| \vee |v - v_{s'}|)}{h} \left| \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \Theta_{ijktt'} \{K_h(U_{ikt'} - v) + K_h(U_{ijt} - u_s)\} \right| \\ &\leq \frac{c}{Nh} \left\{ E \left(\left| \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{t'=1}^{T_{ik}} \Theta_{ijktt'} \right| \right) + 1 \right\} \leq \frac{c}{Nh}. \end{aligned}$$

Applying similar techniques as above, we can define events $\Lambda_{Z,jkm}$ and $\Lambda_{\Xi,jklm}$ for $m, l = 1, 2, 3$. On the intersection of these events, we can obtain that $|\widehat{Z}_{jkm}(u,v) - \widehat{Z}_{jkm}(u_s, v_{s'})| \leq c(Nh)^{-1}$ and $|\widehat{\Xi}_{jklm}(u,v) - \widehat{\Xi}_{jklm}(u_s, v_{s'})| \leq c(Nh)^{-1}$. Combing the above results, we have

$$\sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)| \leq \max_{s,s' \in [N]} |\widehat{\Sigma}_{jk}(u_s, v_{s'}) - \widetilde{\Sigma}_{jk}(u_s, v_{s'})| + \frac{c}{Nh}.$$

By the Bernstein inequality, we have $\text{pr}(\Lambda_{Z,jkm}^c) \leq c \exp(-cn) \leq c \exp(-c\nu_{n,T,h,jk})$ and $\text{pr}(\Lambda_{\Xi,jklm}^c) \leq c \exp(-cn) \leq c \exp(-c\nu_{n,T,h,jk})$ for $m, l = 1, 2, 3$. It follows from the above results and the union bound of probability with the choice of $N = c(h\delta)^{-1}$ that there exists some positive constants c_3 and c_4 such that, for any $\delta \in (0, 1]$,

$$\text{pr} \left\{ \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)| \geq \delta \right\} \leq \frac{c_4}{h^2 \delta^2} \exp(-c_3 \nu_{n,T,h,jk} \delta^2). \quad (\text{A.32})$$

Take arbitrarily small $\epsilon_2 > 0$. If $n^{\epsilon_2} \nu_{n,T,h,jk} \delta^2 \geq 1$, the right side of (A.32) reduces to $c_4 n^{\epsilon_2} \nu_{n,T,h,jk} h^{-2} \exp(-c_3 \nu_{n,T,h,jk} \delta^2)$. If $n^{\epsilon_2} \nu_{n,T,h,jk} \delta^2 \leq 1$, we can choose c_4 and $n^{\epsilon_2} > c$ such that $c_4 \exp(-c_3 c^{-1}) \geq 1$ and hence the same bound $c_4 n^{\epsilon_2} \nu_{n,T,h,jk} h^{-2} \exp(-c_3 \nu_{n,T,h,jk} \delta^2)$ can still be used. We complete the proof of (7) in Theorem 2, the concentration inequality for the covariance estimator in supremum norm.

A.3 Proof of Theorem 3

Note that $\|\hat{\mu}_j - \mu_j\| \leq \|\hat{\mu}_j - \tilde{\mu}_j\| + \|\tilde{\mu}_j - \mu_j\|$, it suffices to bound $\|\tilde{\mu}_j - \mu_j\|$. By (A.1), for any $u \in \mathcal{U}$,

$$\tilde{\mu}_j(u) - \mu_j(u) = \mathbf{e}_0^\top \left[E\{\hat{\mathbf{S}}_j(u)\} \right]^{-1} E \left[\hat{\mathbf{R}}_j(u) - \hat{\mathbf{S}}_j(u) \{\mu_j(u), 0\}^\top \right].$$

By the Taylor expansion, we have

$$E_\epsilon \left[\hat{\mathbf{R}}_j(u) - \hat{\mathbf{S}}_j(u) \{\mu_j(u), 0\}^\top \middle| V_j \right] = \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} K_{h_{\mu,j}}(U_{ijt} - u) \{\mu_j(U_{ijt}) - \mu_j(u)\} := \mathbf{J}_1 + \mathbf{J}_2,$$

with

$$\begin{aligned} \mathbf{J}_1 &= \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} K_{h_{\mu,j}}(U_{ijt} - u) \frac{U_{ijt} - u}{h_{\mu,j}} h_{\mu,j} \frac{\partial \mu_j(u)}{\partial u}, \\ \mathbf{J}_2 &= \frac{1}{2} \sum_{i=1}^n v_{ij} \sum_{t=1}^{T_{ij}} \tilde{\mathbf{U}}_{ijt} K_{h_{\mu,j}}(U_{ijt} - u) \left(\frac{U_{ijt} - u}{h_{\mu,j}} \right)^2 h_{\mu,j}^2 \frac{\partial \mu_j^2(\delta_{ijt})}{\partial u^2}, \end{aligned}$$

where $\delta_{ijt} \in [u - h_{\mu,j}, u + h_{\mu,j}]$ and E_ϵ denotes the expectation over $\{Y_{ijt}\}$ in (1) conditional on the event $V_j = \{U_{ijt}, t \in [T_{ij}], i \in [n]\}$. First consider \mathbf{J}_1 , which equals to the second column of $\hat{\mathbf{S}}_j(u)$ multiplied by $h_{\mu,j} \partial \mu_j(u) / \partial u$, hence

$$E_U \left[\mathbf{e}_0^\top \left[E\{\hat{\mathbf{S}}_j(u)\} \right]^{-1} \mathbf{J}_1 \right] = h_{\mu,j} \frac{\partial \mu_j(u)}{\partial u} \mathbf{e}_0^\top \left[E\{\hat{\mathbf{S}}_j(u)\} \right]^{-1} E\{\hat{\mathbf{S}}_j(u)\} (0, 1)^\top = 0,$$

where E_U denotes the expectation over V_j . Consider \mathbf{J}_2 next. Under Assumption 7, we have $K_1 = \sup_{j \in [p], \xi \in \mathcal{U}} |\partial \mu_j^2(\xi) / \partial u^2| < \infty$. Each entry of $|\mathbf{J}_2|$ is bounded by the (1, 1)th entry of $\hat{\mathbf{S}}_j(u)$ multiplied by $K_1 h_{\mu,j}^2 / 2$, and by $E\{K_h(U_{ijt} - u)\} \leq 1$ we have that the (1, 1)th entry of $E\{\hat{\mathbf{S}}_j(u)\}$ is bounded by 1. Note that $E\{\hat{\mathbf{S}}_j(u)\}$ is positive definite. These results together yield that

$$|E_U(\mathbf{e}_0^\top \left[E\{\hat{\mathbf{S}}_j(u)\} \right]^{-1} \mathbf{J}_2)| \leq \|E\{\hat{\mathbf{S}}_j(u)\}^{-1}\|_{\min} E_U(|\mathbf{J}_2|) \leq c K_1 h_{\mu,j}^2,$$

which implies that $|\tilde{\mu}_j(u) - \mu_j(u)| \leq cK_1 h_{\mu,j}^2$ for any $u \in \mathcal{U}$. Hence

$$\sup_{u \in \mathcal{U}} |\tilde{\mu}_j(u) - \mu_j(u)| = O(h_{\mu,j}^2) \quad \text{and} \quad \|\tilde{\mu}_j - \mu_j\| = O(h_{\mu,j}^2). \quad (\text{A.33})$$

For $M > 0$ with the choice of $\delta = (\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} M \leq 1$, it follows from the union bound of probability and (4) in Theorem 1 that

$$\begin{aligned} & \text{pr} \left\{ \frac{\max_{j \in [p]} \|\hat{\mu}_j - \tilde{\mu}_j\|}{(\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2}} \geq M \right\} \\ & \leq \sum_{j=1}^p \text{pr} \left\{ \|\hat{\mu}_j - \tilde{\mu}_j\| \geq M (\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} \right\} \\ & \leq \sum_{j=1}^p c \exp \left(-c \gamma_{n,T,h,j} M^2 \frac{\log p}{\min_j \gamma_{n,T,h,j}} \right) \leq c \exp \{ (1 - cM^2) \log p \}. \end{aligned} \quad (\text{A.34})$$

We can choose a large M such that $1 - cM^2 < 0$, the right-side of (A.34) tends to 0. Hence $\max_{j \in [p]} \|\hat{\mu}_j - \tilde{\mu}_j\| = O_P \{ (\log p)^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} \}$. Combing this with (A.33) yields that

$$\max_{j \in [p]} \|\hat{\mu}_j - \mu_j\| = O_P \left\{ \left(\frac{\log p}{\min_j \gamma_{n,T,h,j}} \right)^{1/2} + \max_j h_{\mu,j}^2 \right\},$$

which completes the proof of (8).

The rate of convergence in (9) can be proved following the similar procedure. Let $h_{\mu,\min} = \min_j h_{\mu,j}$, we assume that $h_{\mu,\min} \asymp \{\log(p \vee n)/n\}^{\kappa_1}$ where $\kappa_1 \in (0, 1/2]$. Consider $h_{\mu,\min}$ with some $\kappa_1 > 1/2$, the corresponding rate is slower than that with $\kappa_1 \in (0, 1/2]$. To be specific, under the sparse design, the rate of $\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)|$ is at least $\{\log(p \vee n)/n\}^{1/4}$, which is slower than $\{\log(p \vee n)/n\}^{2/5}$ with $\kappa_1 = 1/5$. Under the dense design with $\bar{T}_\mu \{\log(p \vee n)/n\}^{1/4} \rightarrow 0$, the rate is at least $\{\log(p \vee n)/n\}^{1/4} \bar{T}_\mu^{-1/2}$, which is slower than $\{\log(p \vee n)/n\}^{2/5} \bar{T}_\mu^{-2/5}$ with $\kappa_1 \in (1/5, 1/4)$. Under the dense design with $\bar{T}_\mu \{\log(p \vee n)/n\}^{1/4} \rightarrow \tilde{c}$ or ∞ , the rate is at least $\{\log(p \vee n)/n\}^{1/4} \bar{T}_\mu^{-1/2}$, which is slower than $\{\log(p \vee n)/n\}^{1/2}$ with $\kappa_1 \in (1/4, 1/2)$. Based on the above three cases, if $\kappa_1 > 1/2$, the corresponding rate is slower than that with some $\kappa_1 \in (0, 1/2]$ and hence the κ_1 that corresponds with the optimal bandwidth under sparse or dense design is in $(0, 1/2]$. For $M > 0$ with the choice of $\delta = \{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} M \leq 1$, by the union bound

of probability and (5) in Theorem 1, we have

$$\begin{aligned}
& \text{pr} \left\{ \frac{\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)|}{\{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2}} \geq M \right\} \\
& \leq \sum_{j=1}^p \text{pr} \left\{ \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \tilde{\mu}_j(u)| \geq M \{\log(p \vee n)\}^{1/2} (\min_j \gamma_{n,T,h,j})^{-1/2} \right\} \\
& \leq \sum_{j=1}^p \frac{c(n^{\epsilon_1} \gamma_{n,T,h,j})^{1/2}}{h_{\mu,\min}} \exp \left\{ -c \gamma_{n,T,h,j} M^2 \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\} \\
& \leq c \exp \left\{ \left(\frac{3 + \epsilon_1}{2} + \kappa_1 - cM^2 \right) \log(p \vee n) \right\}. \tag{A.35}
\end{aligned}$$

We can choose a large M such that $3 + \epsilon_1 + 2\kappa_1 - 2cM^2 < 0$, the right-side of (A.35) tends to 0. Combing this with (A.33) yields that

$$\max_{j \in [p]} \sup_{u \in \mathcal{U}} |\hat{\mu}_j(u) - \mu_j(u)| = O_P \left[\left\{ \frac{\log(p \vee n)}{\min_j \gamma_{n,T,h,j}} \right\}^{1/2} + \max_j h_{\mu,j}^2 \right],$$

which completes the proof of (9).

A.4 Proof of Theorem 4

Note $\|\hat{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}} \leq \|\hat{\Sigma}_{jk} - \tilde{\Sigma}_{jk}\|_{\mathcal{S}} + \|\tilde{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}}$, it suffices to bound $\|\tilde{\Sigma}_{jk} - \Sigma_{jk}\|_{\mathcal{S}}$. By (A.17), for any $(u, v) \in \mathcal{U}^2$,

$$\tilde{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v) = \tilde{\mathbf{e}}_0^{\text{T}} \left[E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \right]^{-1} E \left[\hat{\mathbf{Z}}_{jk}(u, v) - \hat{\mathbf{\Xi}}_{jk}(u, v) \{ \Sigma_{jk}(u, v), 0, 0 \}^{\text{T}} \right].$$

By Taylor expansion, we have

$$\begin{aligned}
& E_{\tilde{\epsilon}} \left[\hat{\mathbf{Z}}_{jk}(u, v) - \hat{\mathbf{\Xi}}_{jk}(u, v) \{ \Sigma_{jk}(u, v), 0, 0 \}^{\text{T}} | \tilde{V}_{jk} \right] \\
& = \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijt} K_{h_{\Sigma,jk}}(U_{ijt} - u) \{ \Sigma_{jk}(U_{ijt}, U_{iks}) - \Sigma_{jk}(u, v) \} := \mathbf{L}_1 + \mathbf{L}_2 + \mathbf{L}_3,
\end{aligned}$$

where

$$\begin{aligned}
\mathbf{L}_1 &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijt} K_{h_{\Sigma,jk}}(U_{ijt} - u) \frac{U_{ijt} - u}{h_{\Sigma,jk}} h_{\Sigma,jk} \frac{\partial \Sigma_{jk}}{\partial u}(u, v), \\
\mathbf{L}_2 &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijt} K_{h_{\Sigma,jk}}(U_{ijt} - u) \frac{U_{iks} - v}{h_{\Sigma,jk}} h_{\Sigma,jk} \frac{\partial \Sigma_{jk}}{\partial v}(u, v), \\
\mathbf{L}_3 &= \sum_{i=1}^n w_{ijk} \sum_{t=1}^{T_{ij}} \sum_{s=1}^{T_{ik}} \tilde{\mathbf{U}}_{ijt} K_{h_{\Sigma,jk}}(U_{ijt} - u) \tilde{L}_{ijk}, \\
\tilde{L}_{ijk} &= \frac{1}{2} h_{\Sigma,jk}^2 \left(\frac{U_{ijt} - u}{h_{\Sigma,jk}} \right)^2 \frac{\partial^2 \Sigma_{jk}}{\partial u^2} (\delta_{ij kts1}, \delta_{ij kts2}) + \frac{1}{2} h_{\Sigma,jk}^2 \left(\frac{U_{iks} - v}{h_{\Sigma,jk}} \right)^2 \frac{\partial^2 \Sigma_{jk}}{\partial v^2} (\delta_{ij kts1}, \delta_{ij kts2}) \\
&\quad + h_{\Sigma,jk}^2 \frac{U_{ijt} - u}{h_{\Sigma,jk}} \frac{U_{iks} - v}{h_{\Sigma,jk}} \frac{\partial^2 \Sigma_{jk}}{\partial u \partial v} (\delta_{ij kts1}, \delta_{ij kts2}),
\end{aligned}$$

$(\delta_{ij kts1}, \delta_{ij kts2}) \in [u - h_{\Sigma,jk}, u + h_{\Sigma,jk}] \times [v - h_{\Sigma,jk}, v + h_{\Sigma,jk}]$ and the event $\tilde{V}_{jk} = \{(U_{ijt}, U_{iks}), t \in [T_{ij}], s \in [T_{ik}], i \in [n]\}$. First consider L_1 , which equals to the second column of $\hat{\mathbf{\Xi}}_{jk}(u, v)$ multiplied by $h_{\Sigma,jk} \partial \Sigma_{jk}(u, v) / \partial u$, hence $E_U \left(\tilde{\mathbf{e}}_0^T \left[E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \right]^{-1} \mathbf{L}_1 \right)$ equals to

$$h_{\Sigma,jk} \frac{\partial \Sigma_{jk}}{\partial u}(u, v) \tilde{\mathbf{e}}_0^T \left[E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \right]^{-1} E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} (0, 1, 0)^T = 0,$$

where E_U denotes the expectation over \tilde{V}_{jk} . Following the similar procedure, we can show that

$$E_U \left(\tilde{\mathbf{e}}_0^T \left[E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \right]^{-1} \mathbf{L}_2 \right) = 0.$$

Then consider \mathbf{L}_3 . By Assumption 7, we have

$$K_2 = \sup_{(j,k) \in [p]^2, (u,v) \in \mathcal{U}^2} \{ \partial^2 \Sigma_{jk}(u, v) / \partial v^2, \partial^2 \Sigma_{jk}(u, v) / \partial u^2, \partial^2 \Sigma_{jk}(u, v) / \partial u \partial v \} < \infty.$$

Each entry of $|\mathbf{L}_3|$ is bounded by the (1, 1)th entry of $\hat{\mathbf{\Xi}}_{jk}(u, v)$ multiplied by $2K_2 h_{\Sigma,jk}^2$, and by $E \{ K_h(U_{ijt} - u) K_h(U_{iks} - v) \} \leq 1$, the (1, 1)th entry of $E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \}$ is bounded by 1. Note that $E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \}$ is positive definite. Combining these results, we have

$$\left| E_U \left(\tilde{\mathbf{e}}_0^T \left[E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \right]^{-1} \mathbf{L}_3 \right) \right| \leq \| E \{ \hat{\mathbf{\Xi}}_{jk}(u, v) \} \|_{\min}^{-1} \| E_U(|\mathbf{L}_3|) \| \leq cK_2 h_{\Sigma,jk}^2,$$

which implies that $|\tilde{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)| \leq cK_2 h_{\Sigma,jk}^2$ for any $(u, v) \in \mathcal{U}^2$. Hence

$$\sup_{(u,v) \in \mathcal{U}^2} |\tilde{\Sigma}_{jk}(u, v) - \Sigma_{jk}(u, v)| = O(h_{\Sigma,jk}^2) \quad \text{and} \quad \|\tilde{\Sigma}_{jk} - \Sigma_{jk}\| = O(h_{\Sigma,jk}^2). \quad (\text{A.36})$$

For $M > 0$ with the choice of $\delta = (\log p)^{1/2}(\min_{j,k} \nu_{n,T,h,jk})^{-1/2}M \leq 1$, it follows from the union bound of probability and (6) in Theorem 2 that

$$\begin{aligned}
& \Pr \left\{ \frac{\max_{j,k \in [p]} \|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_S}{(\log p)^{1/2}(\min_{j,k} \nu_{n,T,h,jk})^{-1/2}} \geq M \right\} \\
& \leq \sum_{j=1}^p \sum_{k=1}^p \Pr \left\{ \|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_S \geq M(\log p)^{1/2}(\min_{j,k} \nu_{n,T,h,jk})^{-1/2} \right\} \\
& \leq \sum_{j=1}^p \sum_{k=1}^p c \exp \left(-c\nu_{n,T,h,jk}M^2 \frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right) \leq c \exp \{(2 - cM^2) \log p\}.
\end{aligned} \tag{A.37}$$

We can choose a large M such that $2 - cM^2 < 0$, the right-side of (A.37) tends to 0. Hence $\max_{j,k \in [p]} \|\widehat{\Sigma}_{jk} - \widetilde{\Sigma}_{jk}\|_S = O_P\{(\log p)^{1/2}(\min_{j,k} \nu_{n,T,h,jk})^{-1/2}\}$. Combing this with (A.36) yields that

$$\max_{j,k \in [p]} \|\widehat{\Sigma}_{jk} - \Sigma_{jk}\|_S = O_P \left\{ \left(\frac{\log p}{\min_{j,k} \nu_{n,T,h,jk}} \right)^{1/2} + \max_{j,k} h_{\Sigma,jk}^2 \right\}.$$

which completes the proof of (10).

The rate of convergence in (11) can be proved following the similar procedure. Let $h_{\Sigma,\min} = \min_{j,k} h_{\Sigma,jk}$, we can assume that $h_{\Sigma,\min} \asymp \{\log(p \vee n)/n\}^{\kappa_2}$ where $\kappa_2 \in (0, 1/2]$. Consider $h_{\Sigma,\min}$ with $\kappa_2 > 1/2$, the corresponding rate is slower than that with some $\kappa_2 \in (0, 1/2]$. Specifically, under the sparse design, the rate of $\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)|$ is at least 1, which is slower than $\{\log(p \vee n)/n\}^{1/3}$ with $\kappa_2 = 1/6$. Under the dense design with $\bar{T}_\Sigma \{\log(p \vee n)/n\}^{1/4} \rightarrow 0$, the rate is at least \bar{T}_Σ^{-1} , which is slower than $\{\log(p \vee n)/n\}^{1/3} \bar{T}_\Sigma^{-1/3}$ with $\kappa_2 \in (1/6, 1/4)$. Under the dense design with $\bar{T}_\Sigma \{\log(p \vee n)/n\}^{1/4} \rightarrow \tilde{c}$ or ∞ , the rate is at least \bar{T}_Σ^{-1} , which is slower than $\{\log(p \vee n)/n\}^{1/2}$ with $\kappa_2 \in (1/4, 1/2)$. Based on the above three cases, if $\kappa_2 > 1/2$, the corresponding rate is slower than that with some $\kappa_2 \in (0, 1/2]$ and hence κ_2 that corresponds with the optimal bandwidth under sparse or dense design is in $(0, 1/2]$. For $M > 0$ with the choice of $\delta = \{\log(p \vee n)\}^{1/2}(\min_j \gamma_{n,T,h,j})^{-1/2}M \leq 1$, by

the union bound of probability and (7) in Theorem 2, we have

$$\begin{aligned}
& \text{pr} \left\{ \frac{\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)|}{\log(p \vee n)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2}} \geq M \right\} \\
& \leq \sum_{j=1}^p \sum_{k=1}^p \text{pr} \left\{ \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)| \geq M \{\log(p \vee n)\}^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2} \right\} \\
& \leq \sum_{j=1}^p \sum_{k=1}^p \frac{c n^{\epsilon_2} \nu_{n,T,h,jk}}{h_{\Sigma, \min}^2} \exp \left\{ -c \nu_{n,T,h,jk} M^2 \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\} \\
& \leq c \exp \{ (3 + \epsilon_2 + 2\kappa_2 - cM^2) \log(p \vee n) \}.
\end{aligned}$$

We can choose a large M such that $3 + \epsilon_2 + 2\kappa_2 - cM^2 < 0$, the right-side of the above inequality tends to 0. Hence

$$\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \widetilde{\Sigma}_{jk}(u,v)| = O_P \{ \log(p \vee n)^{1/2} (\min_{j,k} \nu_{n,T,h,jk})^{-1/2} \}.$$

Combing this with (A.36) yields that

$$\max_{j,k \in [p]} \sup_{(u,v) \in \mathcal{U}^2} |\widehat{\Sigma}_{jk}(u,v) - \Sigma_{jk}(u,v)| = O_P \left[\left\{ \frac{\log(p \vee n)}{\min_{j,k} \nu_{n,T,h,jk}} \right\}^{1/2} + \max_{j,k} h_{\Sigma, jk}^2 \right].$$

which completes the proof of (11).

B Verification of the claim in Section 4

For the mean estimator, define the set of r candidate bandwidths $\mathcal{H}_\mu = \{h_\mu^{(1)}, \dots, h_\mu^{(r)}\}$. In our simulations, the bandwidth for each dimension can be chosen from \mathcal{H}_μ freely and hence there are r^p possible outcomes. The targeted evaluation metric is $\text{global}_{\text{opt}} = \min_{(m_1, \dots, m_p) \in [r]^p} \max_{j \in [p]} \text{MISE}(\widehat{\mu}_j, h_\mu^{(m_j)})$. We will show that $\text{global}_{\text{opt}} = \text{MaxMISE}(\mu)$, the right side of which is much easier to calculate as it only takes into account pr cases. On one hand, it is obvious that $\text{global}_{\text{opt}} \leq \text{MaxMISE}(\mu)$. On the other hand, for fixed $(j, m_j) \in [p] \times [r]$, $\text{MISE}(\widehat{\mu}_j, h_\mu^{(m_j)}) \geq \min_{m \in [r]} \text{MISE}(\widehat{\mu}_j, h_\mu^{(m)})$, and hence $\text{global}_{\text{opt}} \geq$

$$\min_{(m_1, \dots, m_p) \in [r]^p} \max_{j \in [p]} \min_{m \in [r]} \text{MISE}(\widehat{\mu}_j, h_\mu^{(m)}) = \max_{j \in [p]} \min_{m \in [r]} \text{MISE}(\widehat{\mu}_j, h_\mu^{(m)}) = \text{MaxMISE}(\mu).$$

Combining the above results yields $\text{global}_{\text{opt}} = \text{MaxMISE}(\mu)$. The corresponding claim for the covariance estimator can be verified in the same way.

References

- Boucheron, S., Lugosi, G. and Massart, P. (2014). *Concentration inequalities: A Nonasymptotic Theory of Independence*, Oxford University Press.
- Fan, J. and Marron, J. S. (1994). Fast implementations of nonparametric curve estimators, *Journal of Computational and Graphical Statistics* **3**: 35–56.
- Fan, Y., Foutz, N., James, G. and Jank, W. (2014). Functional response additive model estimation with online virtual stock markets, *The Annals of Applied Statistics* **8**: 2435–2460.
- Fan, Y., James, G. and Radchenko, P. (2015). Functional additive regression, *The Annals of Statistics* **43**: 2296–2325.
- Fang, Q., Guo, S. and Qiao, X. (2023). Adaptive functional thresholding for sparse covariance function estimation in high dimensions, *Journal of the American Statistical Association*, *in press* .
- Guo, S. and Qiao, X. (2023). On consistency and sparsity for high-dimensional functional time series with application to autoregressions, *Bernoulli* **29**: 451–472.
- Kong, D., Xue, K., Yao, F. and Zhang, H. (2016). Partially functional linear regression in high dimensions, *Biometrika* **103**: 147159.
- Lee, K.-Y., Ji, D., Li, L., Constable, T. and Zhao, H. (2022). Conditional functional graphical models, *Journal of the American Statistical Association*, *in press* .
- Li, B. and Solea, E. (2018). A nonparametric graphical model for functional data with application to brain networks based on fMRI, *Journal of the American Statistical Association* **113**: 1637–1655.
- Li, Y. and Hsing, T. (2010). Uniform convergence rates for nonparametric regression and principal component analysis in functional/longitudinal data., *The Annals of Statistics* **38**: 33213351.
- Luo, R. and Qi, X. (2017). Function-on-function linear regression by signal compression, *Journal of the American Statistical Association* **112**: 690705.
- Qiao, X., Guo, S. and James, G. (2019). Functional graphical models, *Journal of the American Statistical Association* **114**: 211–222.
- Qiao, X., Qian, C., James, G. and Guo, S. (2020). Doubly functional graphical models in high dimensions, *Biometrika* **107**: 415–431.
- Solea, E. and Li, B. (2022). Copula Gaussian graphical models for functional data, *Journal of the American Statistical Association* **117**: 781–793.
- Yao, F., Müller, H. and Wang, J.-L. (2005a). Functional data analysis for sparse longitudinal data, *Journal of the American Statistical Association* **100**: 577590.

- Yao, F., Müller, H. and Wang, J.-L. (2005b). Functional linear regression analysis for longitudinal data, *The Annals of Statistics* **33**: 2873–2903.
- Zapata, J., Oh, S. Y. and Petersen, A. (2022). Partial separability and functional graphical models for multivariate gaussian processes, *Biometrika* **109**: 665681.
- Zhang, J. and Chen, J. (2007). Statistical inferences for functional data, *The Annals of Statistics* **35**: 10521079.
- Zhang, X. and Wang, J.-L. (2016). From sparse to dense functional data and beyond, *The Annals of Statistics* **44**: 2281–2321.
- Zhu, H., Strawn, N. and Dunson, D. B. (2016). Bayesian graphical models for multivariate functional data, *Journal of Machine Learning Research* **17**: 1–27.