

# On stability of solitons for the Maxwell–Lorentz equations with spinning particle <sup>1</sup>

A.I. Komech

*Faculty of Mathematics, Vienna University*

alexander.komech@univie.ac.at

E.A. Kopylova<sup>2</sup>

*Faculty of Mathematics, Vienna University*

elena.kopylova@univie.ac.at

## Abstract

We consider stability of solitons of the Maxwell–Lorentz equations with extended charged rotating particle. The solitons are solutions which correspond to a particle moving with a constant velocity  $v \in \mathbb{R}^3$  and rotating with a constant angular velocity  $\omega \in \mathbb{R}^3$ . We consider spherically symmetric charge density of the particle. In this case the nonzero solitons with finite energy exist only for  $|v| < 1$  and  $\omega$  satisfying either  $\omega \parallel v$  or  $\omega \perp v$ .

Our main results are i) the *orbital stability* of moving solitons with  $|v| < 1$  and  $\omega = 0$  and ii) the *linear stability* of standing rotating solitons with  $v = 0$  and  $\omega \in \mathbb{R}^3$ .

We use the Hamilton–Poisson structure of the Maxwell–Lorentz system obtained in previous authors' paper. The structure is degenerate and admits the Casimir invariants. We construct the Lyapunov function as a linear combination of the Hamiltonian with a suitable Casimir invariant. The key points are a lower bound for this function. The proof of the bound in the case  $\omega \neq 0$  relies on angular momentum conservation and suitable spectral arguments including the Heinz inequality.

---

<sup>1</sup>Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

<sup>2</sup>Supported partly by Austrian Science Fund (FWF) P34177

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>The Maxwell potentials and the well-posedness</b>	<b>6</b>
<b>3</b>	<b>The Hamilton–Poisson structure</b>	<b>7</b>
<b>4</b>	<b>The canonical transformation to the comoving frame</b>	<b>9</b>
<b>5</b>	<b>The reduced system</b>	<b>10</b>
<b>6</b>	<b>The solitons</b>	<b>11</b>
<b>7</b>	<b>The Lyapunov function and bound below for solitons with <math>\omega = 0</math></b>	<b>13</b>
<b>8</b>	<b>Orbital stability of solitons with <math>\omega = 0</math></b>	<b>15</b>
<b>9</b>	<b>Linear stability of solitons with <math>v = 0</math> and <math>\omega \neq 0</math></b>	<b>16</b>
<b>10</b>	<b>Proof of Proposition 9.4</b>	<b>20</b>
<b>11</b>	<b>Proof of Proposition 9.5</b>	<b>22</b>
<b>A</b>	<b>Comments</b>	<b>22</b>
<b>B</b>	<b>Proof of Lemma 6.2</b>	<b>23</b>

## 1 Introduction

The paper concerns the orbital stability and instability of solitons of the Maxwell–Lorentz equations with an extended charged spinning nonrelativistic particle. We choose the units where the speed of light is  $c = 1$ . The equations read as follows (see [43, 46]):

$$\left\{ \begin{array}{ll} \dot{E}(x, t) = \operatorname{curl} B(x, t) - w(x, t)\rho(x - q(t)), & \operatorname{div} E(x, t) = \rho(x - q(t)) \\ \dot{B}(x, t) = -\operatorname{curl} E(x, t), & \operatorname{div} B(x, t) = 0 \\ m\dot{q}(t) = p(t), \quad \dot{p}(t) = \langle E(x, t + w(x, t)\wedge B(x, t)), \rho(x - q(t)) \rangle \\ I\dot{\omega}(t) = \langle (x - q(t))\wedge [E(x, t) + w(x, t)\wedge B(x, t)], \rho(x - q(t)) \rangle \end{array} \right., \quad (1.1)$$

where  $\rho(x - q)$  is the charge distribution of the extended particle centered at a point  $q \in \mathbb{R}^3$ ,  $m > 0$  is the mass of the particle and  $I > 0$  is its moment of inertia,  $\omega(t)$  is the angular velocity of the particle rotation and  $w(x, t) := \dot{q}(t) + \omega(t)\wedge(x - q(t))$  is the velocity field. The brackets  $\langle \cdot, \cdot \rangle$  here and below denote the inner product in the Hilbert spaces  $L^2 := L^2(\mathbb{R}^3) \otimes \mathbb{C}^n$  with any  $n \geq 1$ .

This model was introduced by Abraham in 1903–1905 (see [1, 2]) for the description of the classical extended electron coupled to its own Maxwell field. The model allows one to avoid the “ultraviolet divergence”, that is, the infiniteness of the own energy and mass of electrons in the case  $\rho(x) = \delta(x)$ . Using this model, Abraham was the first to discover the mass-energy equivalence up to the doubtful factor  $4/3$  (see [1, 2] and also [26, Section 12.9]), which was a *precursor* of the Einstein theory. This system also served as the classical Landé model of spin in Old Quantum Mechanics (1900–1924): see [7] (and also [26, Chapter 14] and [27, Appendix A]). Various approximations of such systems were used to explain the famous *radiation damping* in classical electrodynamics: the *Lorentz–Dirac equation* (with *runaway solutions*) as introduced by Dirac in [10], and many other approximations, see [23, Chapter 16]. The detailed account on the genesis and early investigations of the system by Dirac, Poincaré, Sommerfeld, and others can be found in [46, Chapter 3].

The system (1.1) was introduced about 1903, however, the first results of the corresponding long-time behavior were obtained only in 1990’, see below the comments on related works. The system plays a crucial role in a rigorous analysis of radiation by moving particles, see [6, 16, 46] and [28]–[39]. The mathematical analysis of the system is useful in connection to the related problems of nonrelativistic QED, see the survey [46]. In particular, the similarity in the renormalization of mass was pointed out by Hiroshima and Spohn [13].

We suppose that the charge density  $\rho(x)$  is smooth, spherically-symmetric and not identically zero:

$$\rho \in C_0^\infty(\mathbb{R}^2), \quad \rho(x) = \rho_1(|x|); \quad \rho(x) = 0 \text{ for } |x| \geq R_\rho; \quad \rho(x) \not\equiv 0. \quad (1.2)$$

**Remark 1.1.** The spherical symmetry of  $\rho(x)$  is used always in the proofs of conservation laws and of the Hamilton least action principle for the system (1.1), [19, 20, 25, 43, 46]. We suppose that this requirement corresponds to the spherical symmetry of the moment of inertia in the last equation of (1.1).

The system (1.1) is invariant under the groups of translations and rotations of the space  $\mathbb{R}^3$  and admits solitons  $\mathcal{S}_{v,\omega}$  discovered by Spohn [46]. The solitons are solutions moving with constant speed  $v \in \mathbb{R}^3$  and rotating with constant angular velocity  $\omega \in \mathbb{R}^3$ . As is pointed out in [46], for  $\rho(x) \not\equiv 0$ , the solitons with finite energy exist only for  $(v, \omega) \in \Sigma$ , where

$$\Sigma = \{(v, \omega) \in \mathbb{R}^3 \times \mathbb{R}^3 : |v| < 1 \text{ and either } \omega \parallel v, \text{ or } \omega \perp v\}. \quad (1.3)$$

**Remark 1.2.** In the case  $\rho(x) \equiv 0$  we have  $\ddot{q}(t) = \dot{\omega}(t) = 0$  and the Maxwell equations do not depend on the motion of the particle, so the energy of the Maxwell field is conserved. The soliton solutions read as  $(E(x,t), B(x,t), q(t), p(t), \omega(t)) = (0, 0, vt + q_0, mv, \omega)$  with arbitrary  $q_0, v, \omega \in \mathbb{R}^3$ , and they are obviously orbitally stable.

Our main results are i) the orbital stability of solitons  $\mathcal{S}_{v,0}$  with  $|v| < 1$  under conditions (1.2) and

ii) the *linear stability* of standing solitons with  $v = 0$  and  $\omega \in \mathbb{R}^3$  which are solutions to the system

$$\left\{ \begin{array}{ll} \dot{E}(x, t) = \operatorname{curl} B(x, t) - w(x, t)\rho(x), & \operatorname{div} E(x, t) = \rho(x) \\ \dot{B}(x, t) = -\operatorname{curl} E(x, t), & \operatorname{div} B(x, t) = 0 \\ I\dot{\omega}(t) = \langle x \wedge [E(x, t) + w(x, t) \wedge B(x, t)], \rho(x) \rangle \end{array} \right. . \quad (1.4)$$

The system (1.4) is equivalent to (1.1) with  $q(t) \equiv p(t) \equiv 0$  in the case of antisymmetric  $E(x, t)$  and symmetric  $B(x, t)$ , i.e.,  $E(-x, t) = -E(x, t)$ ,  $B(-x, t) = B(x, t)$ . However, we will consider the system (1.1) for general fields without the symmetry/antisymmetry restrictions.

We also have justified the restrictions (1.3) on the parameters of solitons. and calculated effective moment of inertia of solitons in Lemmas 6.1 and 6.2.

The *linear stability* means the stability for the linearized dynamics on the tangent space to the manifold of states with constant angular momentum. The manifold is not well defined, but the tangent space does, see Section 9. So, the nonlinear stability problem on the manifold is not correct contrary to the linearized problem on the tangent space. The tangent space *formally* is invariant with respect to the linearized dynamics, see Appendix A.

Let us comment on our approach. We develop methods of Geometrical Mechanics that arose in the seminal works of Arnold, Marsden, Weinstein, Holm, and others, on the stability in Hydrodynamics, Maxwell–Vlasov system, equations of magnetohydrodynamics, and others, see [3, 4, 15, 41] for surveys and references. The theory relies on the reduction of the systems by the action of the corresponding symmetry groups developing the ideas of Lie and Poincaré [40, 44].

In [31], we have applied the reduction by the symmetry groups of translations and rotations to construct the Hamilton–Poisson structure for the system (1.1). The construction relies on the Hamilton least action principle and the Lie–Poincaré calculus [3, 15]. The structure is degenerate and admits *Casimir invariants* (3.6). Hence, the theory of orbital stability [12] is not applicable in our case. We reduce the system via the canonical transformation to a comoving frame. The Hamilton functional of the reduced system depends on the parameter  $P$  which is the conserved total momentum. Thus, we restrict the Hamiltonian to the states with the fixed  $P$ .

To prove the orbital stability in the case  $\omega = 0$ , we use the Hamiltonian as the Lyapunov function and prove an appropriate lower bound. To prove the *linear stability* in the case  $\omega \neq 0$ , we construct the Lyapunov function as the sum of the Hamiltonian with a Casimir functional. Such strategy is known as the energy–Casimir method [14, 42]. We establish a lower bound for the Lyapunov function restricted to the *tangent space* to the manifold of states with constant angular momentum. The bound is proved by suitable spectral arguments including the Heinz inequality from the theory of interpolation [24] and the closed graph theorem.

Let us comment on related works. The Maxwell–Lorentz system (1.1) was in the focus of theoretical physics since 1903 till the discovery of Quantum Electrodynamics in 1927. However, the corresponding mathematical theory started only after Nodvick’s paper [43], where the Hamilton least action principle and conservation laws for the system (1.1) have been established in the Euler angles representation. The coordinate-free proof of the conservation laws was given by Kiessling

[25]. The coordinate-free proof of the Hamilton principle was given in [19] using the technique of [5, 44] which relies on the Lie algebras and the Poincaré equation. This technique was developed in [20], where the general theory of invariants was constructed for the Poincaré equations on manifolds and applied to the system (1.1).

In [6], Bambusi and Galgani proved the existence and orbital stability of solitons for the Maxwell–Lorentz system (1.1) without spinning ( $\omega(t) \equiv 0$ ). The proof relies on the transition to a comoving frame and a lower bound for the reduced Hamiltonian. The global convergence to solitons for the same system with relativistic kinetic energy of the particle was proved in [16]. In [36], the global attraction to stationary states was established for such system in presence of an external confining potential. Similar results were established in [35, 37] for a particle coupled to the scalar field instead of the Maxwell field. All these results were obtained under the *Wiener condition* on the Fourier transform of the charge density:

$$\hat{\rho}(k) := (2\pi)^{-3/2} \int e^{ikx} \rho(x) dx \neq 0, \quad k \in \mathbb{R}^3 \quad (1.5)$$

which allows us to apply the Wiener Tauberian theorem. The condition means a strong coupling of the charged particle to the eigenfunctions  $e^{ikx}$  of the continuous spectrum of the free field. The results [16, 35, 36, 37] provide the first rigorous proof of the *radiation damping* in classical electrodynamics. For the surveys, see [46] and [28, 29].

The global convergence to rotating solitons was established in [17] for the spinning charge remaining at rest ( $q(t) \equiv 0$ ) in the case of sufficiently small charge density  $\rho$ . In [38], Kunze extended the result to the same system without the smallness and the Wiener conditions. The conditions are replaced by the assumption that the mass of the particle does not belong to a sequence of resonances.

In [18], the asymptotic stability of the solitary manifold was proved for the Maxwell–Lorentz equations without spinning: any solution with initial state, close to the solitary manifold, for large times is close to the sum of a soliton and a dispersive wave. The proof relies on the Buslaev–Perelman–Sulem method of symplectic projection onto solitary manifold [8, 9] and use the Wiener condition (1.5). Similar results were obtained for the particle without spinning coupled to the scalar, Klein–Gordon, and Dirac fields [21, 22, 33].

The adiabatic effective dynamics of solitons was proved in [34] for a relativistic particle coupled to a scalar field in presence of a small external field: for initial states, close to the solitary manifold, the solution remains forever in a small vicinity of the manifold, and its projection onto the manifold moves for large time according to an effective Newton equation. The result was extended in [39] to the Maxwell–Lorentz equations (1.1) without spinning.

Let us comment on related open questions. The orbital stability of solitons  $S_{v,\omega}$  with general  $(v,\omega) \in \Sigma$  when  $\omega \neq 0$  remains an open question. We suppose that the *solitary manifold*  $\mathcal{S} = \{S_{v,\omega} : (v,\omega) \in \Sigma\}$  is a global attractor of the system (1.1), and at least the component of  $\mathcal{S}^{\parallel}$  with  $v \parallel \omega$  is asymptotically stable in a local convergence. We expect that an adiabatic effective dynamics holds for solutions close to  $\mathcal{S}^{\parallel}$  in presence of a weak external magnetic field. We suppose that this

dynamics agrees, on the classical level, with the precession of quantum spin in a magnetic field *orthogonal to the velocity*. In this case, the precession was observed experimentally and shows that the directions of the spin and velocity coincide [45, p. 115], [11].

## 2 The Maxwell potentials and the well-posedness

As is well known [23], the second line of the system (1.1) implies the expressions of the Maxwell fields in the potentials  $A(x, t) = (A_1(x, t), A_2(x, t), A_3(x, t))$  and  $\Phi(x, t)$ :

$$B(x, t) = \text{curl } A(x, t), \quad E(x, t) = -\dot{A}(x, t) - \nabla\Phi(x, t). \quad (2.1)$$

We choose the Coulomb gauge

$$\text{div } A(x, t) = 0. \quad (2.2)$$

Now the first line of the system (1.1) becomes

$$-\ddot{A}(x, t) - \nabla\dot{\Phi}(x, t) = -\Delta A(x, t) - w(x, t)\rho(x - q(t)), \quad \Delta\Phi(x, t) = \rho(x - q(t)). \quad (2.3)$$

Hence,  $\Phi(x, t)$  is the Coulombic potential  $\Phi(x, t) = -\frac{1}{4\pi} \int \frac{\rho(y - q(t))}{|x - y|} dy$ . The gradient  $\nabla\dot{\Phi}(x, t)$  in the first equation of (2.3) can be eliminated by the application of the orthogonal projection  $\mathcal{P}$  in  $L^2$  onto the divergence-free vector fields. The system (1.1) after the substitution of the expressions (2.1) becomes

$$\left\{ \begin{array}{l} \ddot{A}(x, t) = \Delta A(x, t) + \mathcal{P}([\dot{q}(t) + \omega(t) \wedge (x - q(t))] \rho(x - q(t))) \\ m\ddot{q}(t) = \langle -\dot{A}(x, t) + [\dot{q} + \omega \wedge (x - q)] \wedge \text{curl } A(x, t), \rho(x - q(t)) \rangle \\ I\dot{\omega}(t) = \langle (x - q(t)) \wedge [-\dot{A}(x, t) + [\dot{q}(t) + \omega \wedge (x - q(t))] \wedge \text{curl } A(x, t)], \rho(x - q(t)) \rangle \end{array} \right. \quad (2.4)$$

In the last two equations, the terms with  $\Phi(x, t)$  cancel because

$$\langle \nabla\Phi(x, t), \rho(x - q) \rangle = \langle -ik \frac{\hat{\rho}(k)}{k^2}, \hat{\rho}(k) \rangle = 0, \quad \langle (x - q) \nabla\Phi(x, t), \rho(x - q) \rangle = -\langle \nabla \wedge k \frac{\hat{\rho}(k)}{k^2}, \hat{\rho}(k) \rangle = 0.$$

Indeed, the first identity holds since the function  $\hat{\rho}(k)$  is even. The second identity follows from the rotation-invariance (1.2) since  $\nabla \wedge k = (\partial_{\varphi_1}, \partial_{\varphi_2}, \partial_{\varphi_3})$ , where  $\varphi_j$  is the angle of rotation about the axis  $k_j$ .

Denote the Sobolev spaces of real functions  $H^s = H^s(\mathbb{R}^3) \otimes \mathbb{R}^3$  with  $s \in \mathbb{R}$ , and  $\dot{H}^1 = \dot{H}^1(\mathbb{R}^3) \otimes \mathbb{R}^3$  being the completion of  $C_0^\infty := C_0^\infty(\mathbb{R}^3) \otimes \mathbb{R}^3$  with the norm  $\|A(x)\|_{\dot{H}^1} = \|\nabla A(x)\|_{L^2}$ .

**Definition 2.1.** *The Hilbert spaces  $\mathcal{F}^0 := \{A \in L^2 : \text{div } A(x) \equiv 0\}$ , and  $\dot{\mathcal{F}}^1 := \{A \in \dot{H}^1 : \text{div } A(x) \equiv 0\}$ .*

Denote the Hilbert spaces

$$\mathbb{Y} := \dot{\mathcal{F}}^1 \oplus \mathcal{F}^0 \oplus \mathbb{R}^3 \oplus \mathbb{R}^3 \oplus \mathbb{R}^3, \quad \mathbb{V} = L^2 \oplus H^{-1} \oplus \mathbb{R}^3 \oplus \mathbb{R}^3 \oplus \mathbb{R}^3. \quad (2.5)$$

The following proposition was proved in [31].

**Proposition 2.2.** i) For any initial state  $\mathcal{Y}(0) = (A(x, 0), \dot{A}(x, 0), q(0), p(0), \omega(0)) \in \mathbb{Y}$ , the system (2.4) admits a unique solution

$$\mathcal{Y}(t) = (A(x, t), \dot{A}(x, t), q(t), p(t), \omega(t)) \in C(\mathbb{R}, \mathbb{Y}) \cap C^1(\mathbb{R}, \mathbb{V}).$$

ii) The map  $W(t) : \mathcal{Y}(0) \mapsto \mathcal{Y}(t)$  is continuous in  $\mathbb{Y}$  for every  $t \in \mathbb{R}$ , and the energy and momentum are conserved:

$$\mathcal{E}(t) = \frac{1}{2} \int [\dot{A}^2(x, t) + |\operatorname{curl} A(x, t)|^2] dx + \frac{1}{2} m \dot{q}^2(t) + \frac{1}{2} I \omega^2(t) = \text{const}, \quad t \in \mathbb{R}, \quad (2.6)$$

$$P(t) = -\langle \dot{A}(x, t), \nabla_* A(x, t) \rangle + m \dot{q}(t) + \langle \rho(x - q(t)), A(x, t) \rangle = \text{const}, \quad t \in \mathbb{R}. \quad (2.7)$$

Here the differentiation  $\nabla_*$  is defined by

$$\langle \dot{A}, \nabla_* A \rangle_n := \langle \dot{A}, \nabla_n A \rangle, \quad n = 1, 2, 3. \quad (2.8)$$

The differentiation (2.8) is a suitable version of the basic operation [3, (10)] from the theory of coadjoint representation.

### 3 The Hamilton–Poisson structure

In this section we recall the Hamilton–Poisson structure of the system (2.4) which has been constructed in [31]. Denote by  $p$  and  $\pi$  the momentum and angular momentum of the particle in the Maxwell field,

$$p = m \dot{q} + \langle A(q + y), \rho(y) \rangle, \quad \pi = I \omega + \langle y \wedge A(q + y), \rho(y) \rangle. \quad (3.1)$$

In [19, 43], the Hamilton least action principle is established for the system (2.4). Using this principle and the Lie–Poincaré technique [3, 15], we have shown in [31] that the system (2.4) is equivalent to

$$\left\{ \begin{array}{l} \dot{A} = \Pi, \quad \dot{\Pi} = \Delta A + \mathcal{P}[\dot{q} + \omega \wedge (x - q)] \rho(x - q) \\ m \dot{q} = p - \langle A(q + y), \rho(y) \rangle, \quad \dot{p} = \langle [\dot{q} + \omega \wedge y] \rho(y), \nabla_* A(q + y) \rangle \\ \dot{\pi} = \omega \wedge \pi \end{array} \right. \quad (3.2)$$

Let us denote the Hamiltonian

$$\begin{aligned} H(A, \Pi, q, p, \pi) &= \frac{1}{2} \langle \Pi, \Pi \rangle + \frac{1}{2} \langle \operatorname{curl} A, \operatorname{curl} A \rangle + \frac{1}{2} m \dot{q}^2 + \frac{1}{2} I \omega^2 \\ &= \frac{1}{2} \int [\Pi^2 + (\operatorname{curl} A)^2] dx + \frac{1}{2m} [p - \langle A(x), \rho(x - q) \rangle]^2 + \frac{1}{2I} [\pi - \langle (x - q) \wedge A(x), \rho(x - q) \rangle]^2, \end{aligned} \quad (3.3)$$

which coincides with the energy (2.6). The Hamiltonian is well defined and Fréchet differentiable on the Hilbert phase space  $\mathbb{Y}$  defined in (2.5). It is easy to check that the system (3.2) admits the

representation in the Hamiltonian form

$$\dot{Y} = \mathcal{J}(Y)DH(Y), \quad Y = (A, \Pi, q, p, \pi), \quad (3.4)$$

where  $\mathcal{J}(Y)$  is a skew-symmetric operator

$$\mathcal{J}(Y) = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\pi \wedge \end{pmatrix}. \quad (3.5)$$

Note that the operator  $\mathcal{J}(Y)$  is not invertible for all  $Y = (A, \Pi, q, p, \pi)$ , and

$$\dim \text{Ker } \mathcal{J}(Y) = \begin{cases} 1, & \pi \neq 0 \\ 3, & \pi = 0 \end{cases}$$

In detail, the system (3.4) reads as

$$\begin{cases} \dot{A} = D_{\Pi}H, & \dot{\Pi} = -D_A H \\ \dot{q} = D_p H, & \dot{p} = -D_q H \\ \dot{\pi} = -\pi \wedge D_{\pi} H \end{cases}$$

The system admits the *Casimir invariants*

$$C(Y) = |\pi|, \quad Y = (A, \Pi, q, p, \pi). \quad (3.6)$$

Their presence is due to the fact that the matrix  $J(Y)$  is not invertible. It suffices to prove the conservation of  $C(Y)$  for  $\pi \neq 0$ . In this case  $D_{\pi}C = \frac{\pi}{|\pi|}$ , and so

$$\begin{aligned} \partial_t C(Y(t)) &= (DC(Y(t)), \dot{Y}(t)) = (DC(Y(t)), \mathcal{J}(Y(t))DH(Y(t))) \\ &= -(\mathcal{J}^*(Y(t))DC(Y(t)), DH(Y(t))) = 0 \end{aligned} \quad (3.7)$$

since the structural operator  $\mathcal{J}(Y)$  is skew-symmetric by (3.5), and

$$\mathcal{J}^*(Y)DC(Y) = \mathcal{J}^*(Y) \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \frac{\pi}{|\pi|} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ -\pi \wedge \frac{\pi}{|\pi|} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$



## 4 The canonical transformation to the comoving frame

Denote the fields in the comoving frame

$$\mathbf{A}(y, t) := A(q + y, t), \quad \mathbf{\Pi}(y, t) := \Pi(q + y, t). \quad (4.1)$$

We are going to change the variables in the system (3.4) via the map

$$T : (A, \Pi, q, p, \pi) \mapsto (\mathbf{A}, \mathbf{\Pi}, q, P, \pi). \quad (4.2)$$

Here  $P$  is the conserved total momentum (2.7): using (2.8),  $P$  can be written as

$$P = p - \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle. \quad (4.3)$$

Hence, we can define

$$\mathbf{H}(\mathbf{A}, \mathbf{\Pi}, q, P, \pi) := H(A, \Pi, q, p, \pi), \quad p = P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle. \quad (4.4)$$

**Lemma 4.1.** *The system (3.4) in the new variables is equivalent to similar system with the Hamiltonian  $\mathbf{H}$ :*

$$\dot{\mathbf{Y}}(t) = \mathcal{J}(\mathbf{Y}(t)) D\mathbf{H}(\mathbf{Y}(t)), \quad \mathbf{Y}(t) = (\mathbf{A}(t), \mathbf{\Pi}(t), q(t), P(t), \pi(t)), \quad (4.5)$$

where  $\mathcal{J}(\mathbf{Y})$  is given by (3.5).

*Proof.* First, the map (4.2) is canonical, i.e., it leaves the canonical form unchanged:

$$\langle \mathbf{\Pi}, \dot{\mathbf{A}} \rangle + P \cdot \dot{q} + \pi \cdot \omega = \langle \Pi, \dot{A} \rangle + p \cdot \dot{q} + \pi \cdot \omega. \quad (4.6)$$

Indeed, differentiating  $\mathbf{A}(y, t) := A(q(t) + y, t)$  in time, we obtain

$$\dot{\mathbf{A}}(y, t) = \dot{A}(q + y, t) + \dot{q}(t) \cdot \nabla A(q + y, t).$$

Hence, the identity (4.6) reduces to

$$\langle \Pi(q + y), \dot{q} \cdot \nabla A(q + y) \rangle + P \cdot \dot{q} = p \cdot \dot{q},$$

which holds by (4.3). Second, the identities (4.6) and (4.4) imply that the Lagrangians corresponding to the Hamiltonians  $\mathbf{H}$  and  $H$  are related via the map  $T$ :

$$\mathbf{l}(\mathbf{A}, q, \dot{\mathbf{A}}, \dot{q}, \omega) \equiv l(A, q, \dot{A}, \dot{q}, \omega).$$

Therefore, the Lagrangian actions also coincide:

$$\int_a^b \mathbf{l}(\mathbf{A}, q, \dot{\mathbf{A}}, \dot{q}, \omega) dt = \int_a^b l(A, q, \dot{A}, \dot{q}, \omega) dt.$$

Hence, the solutions of the systems (3.4) and (4.5) also are related by this transformation since they are critical trajectories of the action.  $\square$

**Remark 4.2.** Similar canonical transformations have been applied in [6, 16] to the systems of type (1.1) without spinning and in [35] to the system of the particle coupled to the scalar field.

## 5 The reduced system

The Hamiltonian (4.4) does not depend on the variable  $q$ . Hence, the equations for  $q$  and  $P$  in the system (4.5) reduce to

$$\dot{q} = D_P \mathbf{H}, \quad \dot{P} = 0,$$

which correspond to the conservation of the conjugate momenta  $P$ . As the result, the system (4.5) reduces to the following family of the Hamiltonian systems with the parameter  $P \in \mathbb{R}^3$ :

$$\dot{\mathbf{Z}}(t) = \mathbb{J}(\mathbf{Z}(t)) D_{\mathbf{Z}} \mathbf{H}_P(\mathbf{Z}(t)), \quad \mathbf{Z}(t) = (\mathbf{A}(t), \mathbf{\Pi}(t), \pi(t)), \quad (5.1)$$

where  $\mathbf{H}_P$  denotes the reduced Hamiltonian  $\mathbf{H}_P(\mathbf{A}, \mathbf{\Pi}, \pi) := \mathbf{H}(\mathbf{A}, \mathbf{\Pi}, q, P, \pi)$

$$\mathbb{J}(\mathbf{Z}) = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -\pi \wedge \end{pmatrix}, \quad \mathbf{Z} = (\mathbf{A}, \mathbf{\Pi}, \pi).$$

According to (4.4) and (3.3),

$$\mathbf{H}_P(\mathbf{A}, \mathbf{\Pi}, \pi) = \frac{1}{2} \int [\mathbf{\Pi}^2 + (\text{curl } \mathbf{A})^2] dy + \frac{1}{2m} [P + \langle \mathbf{\Pi}, \nabla_* \mathbf{A} \rangle - \langle \mathbf{A}, \rho \rangle]^2 + \frac{1}{2I} [\pi - \langle y \wedge \mathbf{A}(y), \rho(y) \rangle]^2. \quad (5.2)$$

In detail, the system (5.1) reads as

$$\left\{ \begin{array}{l} \dot{\mathbf{A}}(y, t) = D_{\mathbf{\Pi}} \mathbf{H}_P(\mathbf{Z}(t)) = \mathbf{\Pi}(y, t) + (v(t) \cdot \nabla) \mathbf{A}(y, t) \\ \dot{\mathbf{\Pi}}(y, t) = -D_{\mathbf{A}} \mathbf{H}_P(\mathbf{Z}(t)) = \Delta \mathbf{A}(y, t) + (v(t) \cdot \nabla) \mathbf{\Pi}(y, t) + \mathcal{P}[v(t) \rho(y) + y \wedge \omega(t) \rho(y)] \\ \dot{\pi}(t) = -\pi(t) \wedge D_{\pi} \mathbf{H}_P(\mathbf{Z}(t)) = -\pi(t) \wedge \omega(t) \end{array} \right\}, \quad (5.3)$$

where  $v(t)$  and  $\omega(t)$  are given by

$$v(t) = \frac{1}{m} [P - \langle \mathbf{\Pi}(t), \nabla_* \mathbf{A}(t) \rangle - \langle \mathbf{A}(y, t), \rho(y) \rangle], \quad \omega(t) = \frac{1}{I} [\pi(t) - \langle y \wedge \mathbf{A}(y, t), \rho(y) \rangle]. \quad (5.4)$$

The Hamiltonian (5.2) is well defined and Fréchet differentiable on the reduced phase space

$$\mathbb{Z} = \mathcal{F}^1 \oplus \mathcal{F}^0 \oplus \mathbb{R}^3,$$

and it is conserved along trajectories of the system (5.3).

## 6 The solitons

Here we calculate the solitons of the system (3.2) which are solutions of the form

$$S_{v,\omega}(t) = (\mathbf{A}_{v,\omega}(x - vt), \mathbf{\Pi}_{v,\omega}(x - vt), vt, p_v, \pi_{v,\omega}). \quad (6.1)$$

The second line of (3.2) and the definition (3.1) imply that

$$p_v = mv + \langle \mathbf{A}_{v,\omega}(y), \rho(y) \rangle, \quad \pi_{v,\omega} = I\omega + \langle y \wedge A_{v,\omega}(q + y), \rho(y) \rangle. \quad (6.2)$$

The corresponding conserved (total) momentum is given by (2.7):

$$P_{v,\omega} = -\langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega}(t) \rangle + mv + \langle \rho(y), \mathbf{A}_{v,\omega}(y) \rangle. \quad (6.3)$$

In the comoving frame, the solitons become stationary solutions

$$\mathbf{S}_{v,\omega} = (\mathbf{A}_{v,\omega}, \mathbf{\Pi}_{v,\omega}, \pi_{v,\omega}) \in \mathbb{Z}$$

of the reduced system (5.3):

$$\left\{ \begin{array}{l} 0 = D_{\mathbf{\Pi}} \mathbf{H}_{P_{v,\omega}}(\mathbf{S}_{v,\omega}) = \mathbf{\Pi}_{v,\omega}(y) + (v \cdot \nabla) \mathbf{A}_{v,\omega}(y) \\ 0 = -D_{\mathbf{A}} \mathbf{H}_{P_{v,\omega}}(\mathbf{S}_{v,\omega}) = \Delta \mathbf{A}_{v,\omega}(y) + (v \cdot \nabla) \mathbf{\Pi}_{v,\omega}(y) + \mathcal{P}[v\rho(y) + y \wedge \omega \rho(y)] \\ 0 = -\pi_{v,\omega} \wedge D_{\pi} \mathbf{H}_{P_{v,\omega}}(\mathbf{S}_{v,\omega}) = -\pi_{v,\omega} \wedge \omega \end{array} \right. \quad (6.4)$$

The first two equations imply that

$$\Delta \mathbf{A}_{v,\omega}(y) - (v \cdot \nabla)^2 \mathbf{A}_{v,\omega}(y) = -\mathcal{P}[v\rho(y) + \omega \wedge y\rho(y)]. \quad (6.5)$$

Note that  $\mathcal{P}[y \wedge \omega \rho(y)] = y \wedge \omega \rho(y)$  due to the spherical symmetry (1.2), and in the Fourier transform  $\widehat{\mathcal{P}[v\rho]}(k) = (v - \frac{k(v \cdot k)}{k^2}) \hat{\rho}(k)$ . Hence, solving (6.5) in the Fourier transform, we obtain

$$\hat{\mathbf{A}}_{v,\omega}(k) = \frac{(v - \frac{k(v \cdot k)}{k^2}) - i\omega \wedge \nabla \hat{\rho}(k)}{k^2 - (v \cdot k)^2}. \quad (6.6)$$

The conditions (1.2) imply that

$$\mathbf{A}_{v,\omega} \in \mathcal{F}^1, \quad \text{for } |v| < 1, \omega \in \mathbb{R}^3. \quad (6.7)$$

**Lemma 6.1.** *Let conditions (1.2) hold,  $v, \omega \in \mathbb{R}^3$  and  $|v| \geq 1$ . Then  $\mathbf{A}_{v,\omega} \notin \mathcal{F}^1$ .*

*Proof.* The real part of the function (6.6) equals  $R(k) = \frac{f(k)}{g(k)} \hat{\rho}(k)$ , where  $f(k) = v - \frac{k(v \cdot k)}{k^2}$  and  $g(k) = k^2 - (v \cdot k)^2 = k^2(1 - v^2 \cos^2(\widehat{k, v}))$ . It suffices to show that  $R \notin L_{\text{loc}}^2(\mathbb{R}^3 \setminus 0)$ .

In the case  $|v| > 1$ , the denominator  $g(k)$  vanishes on the cone  $|\cos(\widehat{k, v})| = 1/|v|$  while the nominator  $f(k)$  vanish only on the line  $k \parallel v$ . Hence,  $R \in L_{\text{loc}}^2(\mathbb{R}^3 \setminus 0)$  only if  $\rho(k) = 0$  on the cone

which implies that  $\hat{\rho}(k) \equiv 0$  by the spherical symmetry (1.2). This contradicts the last condition of (1.2).

In the case  $|v| = 1$ , both functions  $f(k)$  and  $g(k)$  vanish on the same line  $k \parallel v$ . However,

$$|f(k)| = \left| v - \frac{k(v \cdot k)}{k^2} \right| = \frac{|v|k| - k \cos(\widehat{k, v})|}{|k|} = |\sin(\widehat{k, v})|,$$

while  $|g(k)| = k^2 - (v \cdot k)^2 = k^2(1 - \cos^2(\widehat{k, v})) = k^2 \sin^2(\widehat{k, v})$ . Hence,  $R \in L_{\text{loc}}^2(\mathbb{R}^3 \setminus 0)$  only if  $\rho(k) = 0$  for  $k \parallel v$ , so  $\hat{\rho}(k) \equiv 0$  by the spherical symmetry (1.2).  $\square$

Further, the first equation of (6.4) implies that

$$\mathbf{\Pi}_{v, \omega}(y) = -v \cdot \nabla \mathbf{A}_{v, \omega}(y). \quad (6.8)$$

Hence, (6.7) gives

$$(\mathbf{A}_{v, \omega}, \mathbf{\Pi}_{v, \omega}) \in \mathcal{F}, \quad v, \omega \in \mathbb{R}^3, \quad |v| < 1.$$

It remains to check the last equation of (3.2). It is equivalent to the relation

$$\pi_{v, \omega} \parallel \omega, \quad (6.9)$$

where  $\pi_{v, \omega}$  is given by (6.2). We will deduce these relations from the identity

$$\pi_{v, \omega} = I_{\text{eff}}(v)\omega, \quad (6.10)$$

which holds for  $(v, \omega) \in \Sigma$  and fails for  $(v, \omega) \notin \Sigma$ . Here  $I_{\text{eff}}(v)$  is the effective moment of inertia of the soliton expressed by

$$I_{\text{eff}} = I_{\text{eff}}(v) := I + \delta I(v), \quad \delta I(v) = \begin{cases} \delta I^{\parallel}(v) = \int \frac{k_3^2 + k_2^2}{[k^2 - |v|^2 k_1^2] |k|^2} |\check{\rho}'_1(|k|)|^2 dk, & \omega \parallel v \\ \delta I^{\perp}(v) = \int \frac{k_3^2 + k_2^2}{[k^2 - |v|^2 k_2^2] |k|^2} |\check{\rho}'_1(|k|)|^2 dk, & \omega \perp v \end{cases}, \quad (6.11)$$

where  $\check{\rho}_1(|k|) := \hat{\rho}(k)$ . By (1.2),

$$I_{\text{eff}} > I. \quad (6.12)$$

The next lemma is proved in Appendix B.

**Lemma 6.2.** *Let all conditions (1.2) hold. Then*

- i) *For  $(v, \omega) \in \Sigma$ , the relation (6.10) holds, and hence, the soliton  $\mathbf{S}_{v, \omega} \in \mathbb{Z}$  exists and is unique.*
- ii) *The solitons do not exist if  $(v, \omega) \notin \Sigma$ .*

## 7 The Lyapunov function and bound below for solitons with $\omega = 0$

To prove the stability of a soliton  $\mathbf{S}_{v,\omega}$ , we must construct a Lyapunov function  $\Lambda(\mathbf{Z})$  which is an invariant and admits the lower bound

$$\delta\Lambda := \Lambda(\mathbf{S}_{v,\omega} + \delta\mathbf{Z}) - \Lambda(\mathbf{S}_{v,\omega}) \geq \varkappa \|\delta\mathbf{Z}\|_{\mathbb{Z}}^2, \quad \delta\mathbf{Z} \in \mathbb{Z}, \quad \|\delta\mathbf{Z}\|_{\mathbb{Z}} \ll 1 \quad (7.1)$$

with some  $\varkappa = \varkappa(v) > 0$ . In particular, the soliton must be critical point. In the case  $\omega = 0$ , the solitons  $\mathbf{S}_{v,0}$  are critical points of the Hamiltonian  $\mathbf{H}_{v,0} := \mathbf{H}_{P_{v,0}}$ :

$$D_{\mathbf{A}}\mathbf{H}_{v,0}(\mathbf{S}_{v,0}) = 0, \quad D_{\mathbf{\Pi}}\mathbf{H}_{v,0}(\mathbf{S}_{v,0}) = 0, \quad D_{\pi}\mathbf{H}_{v,0}(\mathbf{S}_{v,0}) = 0. \quad (7.2)$$

Indeed, the first and second identities hold due to the first two equations of (6.4). The last identity follows from (5.2), (6.2) and (6.10):

$$D_{\pi}\mathbf{H}_{P_{v,\omega}}(\mathbf{S}_{v,\omega}) = \frac{1}{I}[\pi_{v,\omega} - \langle y \wedge A_{v,\omega}(y), \rho(y) \rangle] = \omega. \quad (7.3)$$

So, in the case  $\omega = 0$  we take the Hamiltonian as the Lyapunov function.

**Lemma 7.1.** *Let conditions (1.2) hold,  $\omega = 0$  and  $|v| < 1$ . Then (7.1) holds for  $\Lambda = \mathbf{H}_{v,0}$ .*

*Proof.* First, let us calculate  $\delta\mathbf{H}_{v,\omega} = \mathbf{H}_{v,\omega}(\mathbf{S}_{v,\omega} + \delta\mathbf{Z}) - \mathbf{H}_{v,\omega}(\mathbf{S}_{v,\omega})$ . Denote  $\delta\mathbf{Z} = (\alpha, \beta, \gamma)$ , so  $\mathbf{A} = \mathbf{A}_{v,\omega} + \alpha$ ,  $\mathbf{\Pi} = \mathbf{\Pi}_{v,\omega} + \beta$ ,  $\pi = \pi_{v,\omega} + \gamma$ , where

$$\operatorname{div} \alpha(y) = \operatorname{div} \beta(y) = 0 \quad (7.4)$$

by (2.2). We have

$$\begin{aligned} \delta\mathbf{H}_{v,\omega} &= \frac{1}{2} \int [|\mathbf{\Pi}_{v,\omega} + \beta|^2 + |\operatorname{curl}(\mathbf{A}_{v,\omega} + \alpha)|^2] dy - \frac{1}{2} \int [|\mathbf{\Pi}_{v,\omega}|^2 + |\operatorname{curl} \mathbf{A}_{v,\omega}|^2] dy \\ &\quad + \frac{1}{2m} [P_{v,\omega} + \langle \mathbf{\Pi}_{v,\omega} + \beta, \nabla_*(\mathbf{A}_{v,\omega} + \alpha) \rangle - \langle \mathbf{A}_{v,\omega}(y) + \alpha, \rho(y) \rangle]^2 \\ &\quad - \frac{1}{2m} [P_{v,\omega} + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \rangle - \langle \mathbf{A}_{v,\omega}(y), \rho(y) \rangle]^2 \\ &\quad + \frac{1}{2I} [\pi_{v,\omega} + \gamma - \langle y \wedge (\mathbf{A}_{v,\omega}(y) + \alpha), \rho(y) \rangle]^2 - \frac{1}{2I} [\pi_{v,\omega} - \langle y \wedge \mathbf{A}_{v,\omega}(y), \rho(y) \rangle]^2. \end{aligned}$$

After rearrangements, we obtain

$$\begin{aligned} \delta\mathbf{H}_{v,\omega} &= \frac{1}{2} \int (|\beta|^2 + |\operatorname{curl} \alpha|^2) dy + \int (\mathbf{\Pi}_{v,\omega} \cdot \beta + \operatorname{curl} \mathbf{A}_{v,\omega} \cdot \operatorname{curl} \alpha) dy \\ &\quad + \frac{1}{2m} [(p_{v,\omega} + \delta p)^2 - p_{v,\omega}^2] + \frac{1}{2I} [(M_{v,\omega} + \delta M)^2 - M_{v,\omega}^2], \end{aligned} \quad (7.5)$$

where

$$\left\{ \begin{array}{l} p_{v,\omega} := P_{v,\omega} + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \mathbf{A}_{v,\omega} \rangle - \langle \mathbf{A}_{v,\omega}(y), \rho(y) \rangle = mv \\ \delta p := \langle \beta, \nabla_* \mathbf{A}_{v,\omega} \rangle + \langle \mathbf{\Pi}_{v,\omega}, \nabla_* \alpha \rangle + \langle \beta, \nabla_* \alpha \rangle - \langle \alpha(y), \rho(y) \rangle \\ M_{v,\omega} := \pi_{v,\omega} - \langle y \wedge \mathbf{A}_{v,\omega}(y), \rho(y) \rangle = I\omega \\ \delta M := \gamma - \langle y \wedge \alpha, \rho(y) \rangle \end{array} \right. \quad (7.6)$$

due to (6.3) and (6.2). Using the first two equations of (6.4) and taking into account (7.4), we obtain

$$\begin{aligned} & \int (\mathbf{\Pi}_{v,\omega} \cdot \beta + \operatorname{curl} \mathbf{A}_{v,\omega} \cdot \operatorname{curl} \alpha) dy = \langle \mathbf{\Pi}_{v,\omega}, \beta \rangle - \langle \Delta \mathbf{A}_{v,\omega}, \alpha \rangle \\ & = -\langle (v \cdot \nabla) \mathbf{A}_{v,\omega}, \beta \rangle - \langle \mathbf{\Pi}_{v,\omega}, (v \cdot \nabla) \alpha \rangle + \langle v \rho(y) + y \wedge \omega \rho(y), \alpha \rangle \\ & = -v \cdot \delta p - \omega \cdot \delta M + \omega \cdot \gamma + \langle \beta, (v \cdot \nabla) \alpha \rangle. \end{aligned}$$

Substituting into (7.5), after rearrangements, we get

$$\delta \mathbf{H}_{v,\omega} = J_1 + J_2, \quad (7.7)$$

where

$$J_1 = \frac{1}{2} \int (|\beta|^2 + |\nabla \alpha|^2) dy + \langle \beta, v \cdot \nabla \alpha \rangle, \quad (7.8)$$

$$J_2 = \frac{m}{2} [(v + \delta p/m)^2 - v^2 - 2v \cdot \delta p/m] + \frac{I}{2} [(\omega + \delta M/I)^2 - \omega^2 - 2\omega \delta M/I] + \omega \cdot \gamma. \quad (7.9)$$

The first term  $J_1$  admits the lower bound

$$J_1 \geq \frac{1-|v|}{2} \int (|\beta|^2 + |\nabla \alpha|^2) dy \quad (7.10)$$

since  $\operatorname{div} \alpha = 0$ . Using (7.6), the second term reads as

$$J_2 = \frac{(\delta p)^2}{2m} + \frac{(\delta M)^2}{2I} = \frac{(\delta p)^2}{2m} + \frac{1}{2I} [\gamma - \langle y \wedge \alpha, \rho(y) \rangle]^2 + \omega \cdot \gamma. \quad (7.11)$$

Now we can finish the proof of Lemma 7.1. Due to (7.7)–(7.11) with  $\omega = 0$ , it suffices to check that

$$\frac{1-|v|}{2} \int |\nabla \alpha|^2 dy + \frac{1}{2I} [\gamma - \langle y \wedge \alpha(y), \rho(y) \rangle]^2 \geq \varkappa (\|\nabla \alpha\|_{L^2}^2 + \gamma^2) \quad (7.12)$$

with some  $\varkappa > 0$ . Denote  $\mu := \gamma - \langle x \wedge \alpha(x), \rho(x) \rangle$ . Then (7.12) can be rewritten equivalently as

$$\|\nabla \alpha\|_{L^2}^2 + [\mu + \langle y \wedge \alpha(y), \rho(y) \rangle]^2 \leq C(\rho, v, I) \left( \int |\nabla \alpha(y)|^2 dy + \mu^2 \right).$$

It remains to note that

$$|\langle y \wedge \alpha(y), \rho(y) \rangle| = |\langle \nabla \hat{\rho}(k) \wedge \hat{\alpha}(k) \rangle| = \left| \left\langle \frac{\nabla \hat{\rho}(k)}{|k|} \wedge |k| \hat{\alpha}(k) \right\rangle \right| \leq \|\nabla \alpha\|_{L^2} \int \frac{|\nabla \hat{\rho}(k)|^2}{k^2} dk. \quad \square$$

## 8 Orbital stability of solitons with $\omega = 0$

Let us formulate the definition of the orbital stability of solitons (6.1) in terms of reduced trajectories

$$\mathbf{Z}(t) := (\mathbf{A}(y, t), \mathbf{\Pi}(y, t), \pi(t)) \in C(\mathbb{R}, \mathbb{Z}),$$

corresponding to solutions  $Y(t) = (A(x, t), \Pi(x, t), q(t), p(t), \pi(t)) \in C(\mathbb{R}, \mathbb{Y})$  of the system (3.2). Introduce the distance

$$d(Y, S_{v,\omega}(0)) := \|\mathbf{Z} - \mathbf{S}_{v,\omega}\|_{\mathbb{Z}} + |\dot{q} - v|, \quad Y = (A, \Pi, q, p, \pi), \quad \mathbf{Z} = (\mathbf{A}, \mathbf{\Pi}, \pi), \quad (8.1)$$

where  $\mathbf{A}$  and  $\mathbf{\Pi}$  are defined according to (4.1):

$$\mathbf{A}(y) := A(q + y), \quad \mathbf{\Pi}(y) := \Pi(q + y).$$

Suppose that the initial data  $Y(0)$  is close to  $S_{v,\omega}(0)$  in the following sense:

$$d(Y(0), S_{v,\omega}(0)) < \varepsilon. \quad (8.2)$$

**Definition 8.1.** *The soliton  $S_{v,\omega}(\cdot)$  is orbitally stable if for any  $r > 0$  there is an  $\varepsilon > 0$  such that, for any solution  $Y(t)$  of the system (3.2), the inequality (8.2) implies*

$$d(Y(t), S_{v,\omega}(0)) < r, \quad t \in \mathbb{R}. \quad (8.3)$$

Obviously, for  $\rho(x) \equiv 0$  all solitons are orbitally stable, see Remark 1.2. The main result of the present paper is the following theorem.

**Theorem 8.2.** *Let all conditions (1.2) hold, and  $|v| < 1$ . Then the soliton  $S_{v,0}$  is orbitally stable.*

*Proof.* The proof requires a modification of the general scheme of [6], [35] and [16]. Namely, the inequality (8.2) with sufficiently small  $\varepsilon > 0$  implies that the momentum  $P$  of the solution  $Y(t)$  is close to  $P_{v,0}$ :

$$|P - P_{v,0}| \rightarrow 0, \quad \varepsilon \rightarrow 0. \quad (8.4)$$

Lemma 7.1 implies that the soliton  $\mathbf{S}_{v,0}$  is a nondegenerate local minimizer of  $\Lambda_{v,0} = \mathbf{H}_P$ . This point is nondegenerate since the Hessian is positive definite by the lower bound (7.1). Now consider the Hamiltonian  $\Lambda_P(\mathbf{Z})$  corresponding to the momentum  $P$ . It is important that

- i)  $\mathbf{H}_P(\mathbf{Z})$  is the invariant of the system (5.3) with the parameter  $P$ .
- ii) for small  $|P - P_{v,0}|$ , there exists a critical point  $\mathbf{S}_P = (\mathbf{A}_P, \mathbf{\Pi}_P, \pi_P)$  of  $\mathbf{H}_P$ , which is close to the soliton  $\mathbf{S}_{v,0}$ :

$$D\Lambda_P(\mathbf{S}_P) = 0, \quad \|\mathbf{S}_P - \mathbf{S}_{v,0}\|_{\mathbb{Z}} \rightarrow 0, \quad \delta \rightarrow 0. \quad (8.5)$$

**Lemma 8.3.** *For sufficiently small  $\varepsilon > 0$ , the point  $\mathbf{S}_P$  is a strict local minimizer of  $\mathbf{H}_P$ , and the*

lower bound of type (7.1) holds:

$$\mathbf{H}_P(\mathbf{S}_P + \delta\mathbf{Z}) - \mathbf{H}_P(\mathbf{S}_P) \geq \varkappa \|\delta\mathbf{Z}\|_{\mathbb{Z}}^2, \quad \delta\mathbf{Z} \in \mathbb{Z}, \quad \|\delta\mathbf{Z}\|_{\mathbb{Z}} \ll 1, \quad (8.6)$$

where  $\varkappa > 0$ .

*Proof.* All equations of solitons (6.4) hold with  $\mathbf{S}_{v,\omega}$  replaced by  $\mathbf{S}_P$ . Moreover,  $\omega = 0$  by (7.3). Hence,  $\mathbf{S}_P$  is a soliton  $\mathbf{S}_{v,0}$  with  $v$  defined by (6.3) with  $P_{v,\omega}, \mathbf{A}_{v,\omega}, \mathbf{\Pi}_{v,\omega}$  replaced by  $P, \mathbf{A}_P, \mathbf{\Pi}_P$  respectively. Hence, for small  $\varepsilon > 0$ , the bound (8.6) follows by the arguments from the proof of Lemma 7.1.  $\square$

Combining (8.5) with (8.2), we obtain

$$\|\mathbf{Z}(0) - \mathbf{S}_P\|_{\mathbb{Z}} \rightarrow 0, \quad \varepsilon \rightarrow 0.$$

Therefore,

$$0 \leq \Lambda_P(\mathbf{Z}(0)) - \Lambda_P(\mathbf{S}_P) \leq \varepsilon_1, \quad \varepsilon_1 \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Now the conservation of  $\Lambda_P(\mathbf{Z}(t))$  for the system (5.3) with the parameter  $P$  implies that

$$0 \leq \Lambda_P(\mathbf{Z}(t)) - \Lambda_P(\mathbf{S}_P) \leq \varepsilon_1, \quad t \in \mathbb{R}.$$

Hence, by the lower bound (8.6),

$$\sup_{t \in \mathbb{R}} \|\mathbf{Z}(t) - \mathbf{S}_P\|_{\mathbb{Z}} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Finally, combined with (8.5), this inequality gives

$$\sup_{t \in \mathbb{R}} \|\mathbf{Z}(t) - \mathbf{S}_{v,0}\|_{\mathbb{Z}} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (8.7)$$

It remains to show that the last term in (8.1) is small uniformly in time:

$$\sup_{t \in \mathbb{R}} |\dot{q}(t) - v| \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (8.8)$$

This follows from (5.4) together with (8.4) and (8.7).  $\square$

## 9 Linear stability of solitons with $v = 0$ and $\omega \neq 0$

Now we consider the system (1.4), whose Hamiltonian form is

$$\left\{ \begin{array}{l} \dot{\mathbf{A}}(y, t) = \mathbf{\Pi}(y, t), \quad \dot{\mathbf{\Pi}}(y, t) = \Delta \mathbf{A}(y, t) + \omega(t) \wedge y \rho(y) \\ \dot{\pi}(t) = -\pi(t) \wedge \omega(t) \end{array} \right\}. \quad (9.1)$$



This system is identical to the one (5.3) with  $v(t) \equiv 0$  and  $\omega(t)$  given by (5.4). The Hamiltonian reads as

$$\mathbf{H}(\mathbf{A}, \mathbf{\Pi}, \pi) = \frac{1}{2} \int [\mathbf{\Pi}^2 + (\text{curl } \mathbf{A})^2] dx + \frac{1}{2I} [\pi - \langle x \wedge \mathbf{A}(x), \rho(x) \rangle]^2, \quad (9.2)$$

which, by (5.4), is identical to (5.2) with  $v(t) \equiv 0$ . The system (9.1) also can be written in the form (5.1). The Casimir  $|\pi|$  is conserved that follows just as in (3.7).

**Remark 9.1.** The Hamiltonian (9.2) is a quadratic form unlike (5.2), however the last equation in (9.1) is nonlinear.

Solitons of the system are their stationary states  $\mathbf{S}_\omega := \mathbf{S}_{0,\omega}$ , so

$$\mathbf{S}_\omega = (\mathbf{A}_\omega, \mathbf{\Pi}_\omega, \pi_\omega) = (\mathbf{A}_{0,\omega}, 0, \pi_{0,\omega}) \quad (9.3)$$

since  $\mathbf{\Pi}_{0,\omega} = 0$  by (6.8). However, in the case  $\omega \neq 0$ , the solitons are not critical points of the Hamiltonian  $\mathbf{H}$  since  $D_\pi \mathbf{H}(\mathbf{S}_\omega) = \omega$  by (7.3). Now we construct the Lyapunov function as a perturbation of the Hamiltonian by a Casimir invariant:

$$\Lambda(\mathbf{Z}) = \mathbf{H}(\mathbf{Z}) - \frac{\nu_{\text{eff}}}{2} \pi^2, \quad \mathbf{Z} = (\mathbf{A}, \mathbf{\Pi}, \pi), \quad \nu_{\text{eff}} := 1/I_{\text{eff}}(0). \quad (9.4)$$

Obviously, this function is an invariant for the system (5.3) and the solitons  $\mathbf{S}_\omega$  are its critical points since  $D_\pi \mathbf{H}(\mathbf{S}_\omega) = \omega$  and  $[D_\pi \frac{\nu_{\text{eff}}}{2} \pi^2]|_{\pi=\pi_{0,\omega}} = \frac{\pi_{0,\omega}}{I_{\text{eff}}(0)} = \omega$  by (6.10). Repeating the calculations (7.5)–(7.11) with the Lyapunov function (9.4) and  $v = 0$ , we obtain

$$\begin{aligned} \delta\Lambda &:= \Lambda(\mathbf{S}_{v,\omega} + \delta\mathbf{Z}) - \Lambda(\mathbf{S}_{v,\omega}) = \delta\mathbf{H} - \frac{\nu_{\text{eff}}}{2} ((\pi + \gamma)^2 - \pi^2) \\ &= \frac{1}{2} \int (|\beta|^2 + |\nabla\alpha|^2) dy + \frac{\nu}{2} [\gamma - \langle y \wedge \alpha, \rho(y) \rangle]^2 - \frac{\nu_{\text{eff}}}{2} \gamma^2, \quad \nu := 1/I. \end{aligned} \quad (9.5)$$

The presence of the last negative term breaks down the positivity of the form, so the estimate of type (7.1) now does not hold. Indeed, this follows immediately from the fact that the solitary manifold  $\mathcal{S}_* := \{\mathbf{S}_\omega : \omega \in \mathbb{R}^3\}$  is the set of critical points of  $\Lambda$  and  $\delta\Lambda$ , and hence,  $\delta\Lambda(\mathbf{S}_\omega) = 0$  for all  $\omega \in \mathbb{R}^3$ . Note that any modification of the Lyapunov function as  $\Lambda(\mathbf{Z}) = \mathbf{H}(\mathbf{Z}) - f(|\pi|)$  does not help, see Appendix A.

We will prove the *linear stability* of all solitons  $\mathbf{S}_\omega$  applying angular momentum conservation for solutions to (9.1): formally,

$$J(\mathbf{Z}(t)) := \langle \mathbf{A}(x, t) \wedge \mathbf{\Pi}(x, t) \rangle - \langle (x \wedge \nabla)_* \mathbf{A}(x, t), \mathbf{\Pi}(x, t) \rangle + \pi(t) = \text{const}, \quad t \in \mathbb{R}. \quad (9.6)$$

The conservation holds for a class of solutions which must be specified. The problem is the convergence of the middle term with  $x \wedge \nabla$ . In particular, the convergence and the conservation hold for solutions with compact supports. Indeed, denote

$$\mathbb{Z}_c = \{\mathbf{Z} = (\mathbf{A}, \mathbf{\Pi}, \pi) \in \mathbb{Z} : \mathbf{A}(x) = \mathbf{\Pi}(x) = 0, \quad |x| > \text{const}\}.$$

Then for solutions with  $\mathbf{Z} \in \mathbb{Z}_c$  we have

$$\mathbf{A}(x, t) = \mathbf{\Pi}(x, t) = 0, \quad |x| > \text{const} + |t|. \quad (9.7)$$

For such solutions, the conservation (9.6) follows by the method of [25] or by the application of momentum map [32].

To prove the linear stability, it suffices to verify the bound (7.1) with  $v = 0$  on the manifold of states with constant angular momentum:

$$\mathbb{M}_\omega := \{\mathbf{Z} \in \mathbb{Z} : J(\mathbf{Z}) = J(\mathbf{S}_\omega)\}. \quad (9.8)$$

However, the manifold is not well defined as we mentioned above. On the other hand, the tangent space  $\mathbb{T}_\omega := T_{\mathbf{S}_\omega}\mathbb{M}_\omega$  does. Indeed, the tangent space can be *defined* by the equation

$$\mathbb{T}_\omega := \{\delta\mathbf{Z} \in \mathbb{Z} : (dJ(\mathbf{S}_\omega), \delta\mathbf{Z}) = 0\}. \quad (9.9)$$

By (9.6), the differential  $dJ(\mathbf{S}_\omega)$  acts on the vector  $\delta\mathbf{Z} = (\alpha, \beta, \gamma) \in \mathbb{Z}$  as follows:

$$\begin{aligned} \langle dJ(\mathbf{S}_\omega), \delta\mathbf{Z} \rangle &= \langle \mathbf{A}_\omega(x) \wedge \beta(x) \rangle - \langle (x \wedge \nabla)_* \mathbf{A}_\omega(x), \beta(x) \rangle + \gamma \\ &= \langle \hat{\mathbf{A}}_\omega(k) \wedge \hat{\beta}(k) \rangle + \langle (\nabla \wedge k)_* \hat{\mathbf{A}}_\omega(k), \hat{\beta}(k) \rangle + \gamma \end{aligned} \quad (9.10)$$

because  $\mathbf{\Pi}_\omega = 0$  by (9.3). It remains to note that the functional  $dJ(\mathbf{S}_\omega)$  is defined on the entire space  $\mathbb{Z}$  and even on the Hilbert space  $\mathbb{Z}^0 := \mathcal{F}^0 \oplus \mathcal{F}^0 \oplus \mathbb{R}^3$  since  $\mathbf{A}_\omega(x) \in \dot{\mathcal{F}}^1 \cap L^2$  and  $(x \wedge \nabla)_n \mathbf{A}_\omega(x) \in \dot{\mathcal{F}}^1 \cap L^2$ . This follows from the asymptotics

$$\hat{\mathbf{A}}_\omega = \mathcal{O}(|k|^{-1}), \quad (\nabla \wedge k)_* \hat{\mathbf{A}}_\omega = \mathcal{O}(|k|^{-1}), \quad k \rightarrow 0 \quad (9.11)$$

which hold by (6.6) with  $v = 0$ . Expanding  $dJ(\mathbf{S}_\omega) = \sum_1^3 e_n dJ_n(\mathbf{S}_\omega)$  (with  $e_1 = (1, 0, 0)$ , etc), we rewrite (9.9) as

$$\mathbb{T}_\omega := \{\delta\mathbf{Z} \in \mathbb{Z} : (E_n, \delta\mathbf{Z}) = 0, \quad n = 1, 2, 3\}, \quad E_n := e_n dJ_n(\mathbf{S}_\omega) \in \mathbb{Z} \cap \mathbb{Z}^0. \quad (9.12)$$

Hence, the vectors  $E_n$  generate the three-dimensional orthogonal space  $\mathbb{N}_\omega$  to  $\mathbb{T}_\omega$  in  $\mathbb{Z}^0$ . Thus,  $\mathbb{T}_\omega$  is the closed linear subspace of codimension three in  $\mathbb{Z}$ . Denote by  $\mathbb{T}_\omega^0$  the closure of  $\mathbb{T}_\omega$  in  $\mathbb{Z}^0$ . Then  $\mathbb{T}_\omega^0$  is the closed linear subspace of codimension three in  $\mathbb{Z}^0$ . On the other hand, formula (6.6) with  $v = 0$  shows that  $\mathcal{S}_*$  is a linear three-dimensional subspace in  $\mathbb{Z}$  and  $\mathbb{Z}^0$ .

The following theorem is our main result for solitons with  $\omega \neq 0$ . It implies, in particular, that the space  $\mathcal{S}_*$  is transversal to  $\mathbb{T}_\omega$ . Denote by  $P_\omega$  the orthogonal projection of  $\mathbb{Z}^0$  onto  $\mathbb{T}_\omega^0$ :

$$P_\omega = 1 - \sum_1^3 p_n |E_n\rangle \langle E_n|, \quad (9.13)$$

where  $p_n$  are suitable normalization factors.

**Theorem 9.2.** *Let conditions (1.2) hold. Then*

$$\begin{aligned} \delta\Lambda &= \frac{1}{2} \int [\beta^2(y) + |\nabla\alpha(y)|^2] dy + \frac{1}{2} \nu [\gamma - \langle y \wedge \alpha(y), \rho(y) \rangle]^2 - \frac{1}{2} \nu_{\text{eff}} \gamma^2 \\ &\geq \varkappa \left[ \frac{1}{2} \int [\beta^2(y) + |\nabla\alpha(y)|^2] dy + \gamma^2 \right], \quad (\alpha, \beta, \gamma) \in \mathbb{T}_\omega. \end{aligned} \quad (9.14)$$

**Remark 9.3.** The bound implies that the solitons are stable for the equations (9.1) linearized at the soliton  $\mathbf{S}_\omega$  and restricted to the *invariant subspace*  $\mathbb{T}_\omega$ , see Appendix A.

Note that the quadratic form  $\delta\Lambda$  is continuous on the space  $\mathbb{Z}$ . Hence, it suffices to prove the bound (9.14) on the space of smooth functions  $\mathbb{T}_\omega^\infty := \mathbb{T}_\omega \cap [C_0^\infty \oplus C_0^\infty \oplus \mathbb{R}^3]$  which is dense in  $\mathbb{T}_\omega$  in the norm of  $\mathbb{Z}$ . For  $X = (\alpha, \beta, \gamma) \in \mathbb{T}_\omega^\infty$ , the quadratic form  $\delta\Lambda$  can be written as  $\delta\Lambda = (X, QX)$ , where  $Q$  is a linear selfadjoint operator in the Hilbert space  $\mathbb{Z}^0$  and the brackets denote the inner product in  $\mathbb{Z}^0$ . For the smooth functions, (9.14) becomes

$$(X, QX) \geq \varkappa (X, Q_0 X), \quad X \in \mathbb{T}_\omega^\infty, \quad (9.15)$$

where we denote (using the ‘‘Dirac notation’’),

$$Q = \begin{pmatrix} -\Delta + \nu |m^*(x)\rho(x)\rangle\langle\rho(y)m(y)| & 0 & -\nu |m^*(x)\rho(x)\rangle \\ 0 & 1 & 0 \\ -\nu \langle\rho(y)m(y)| & 0 & \delta \end{pmatrix}, \quad Q_0 = \begin{pmatrix} -\Delta & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \delta \end{pmatrix}, \quad (9.16)$$

where  $\delta := \nu - \nu_{\text{eff}} > 0$  by (6.12), and  $m(x)$  denotes the operator  $\alpha \mapsto x \wedge \alpha$ ,  $\alpha \in \mathbb{R}^3$ . The corresponding skew-symmetric matrix reads as

$$m(x) = \begin{pmatrix} 0 & -x_3 & x_2 \\ x_3 & 0 & -x_1 \\ -x_2 & x_1 & 0 \end{pmatrix}, \quad x \in \mathbb{R}^3.$$

The operator  $Q_0$  is positive definite and selfadjoint in  $\mathbb{Z}^0$  with a domain  $\mathbb{D}^0$ . Hence,  $Q$  also is the selfadjoint operator in  $\mathbb{Z}^0$  with the same domain since the difference  $Q - Q_0$  is a bounded finite-rank operator in  $\mathbb{Z}^0$  by (9.16) and (1.2). The common domain  $\mathbb{Z}^0$  of both operators  $Q$  and  $Q_0$  contains the space

$$\mathbb{D}^\infty = \mathbb{Z} \cap [C_0^\infty \oplus C_0^\infty \oplus \mathbb{R}^3] \subset \mathbb{Z}^0.$$

The domain is dense in  $\mathbb{Z}^0$ . Further we denote by  $\mathbb{T}_\omega^0$  the Hilbert space which is the closure of  $\mathbb{T}_\omega$  in  $\mathbb{Z}^0$  endowed with the inner product of  $\mathbb{Z}^0$ . In the next section, we will prove two following propositions.

**Proposition 9.4.** *Let conditions (1.2) hold. Then*

i) the operator  $Q$  is nonnegative:

$$(X, QX) \geq 0, \quad X \in \mathbb{D}^\infty. \quad (9.17)$$

ii)  $\ker Q \cap \mathbb{T}_\omega^0 = 0$ .

Consider the restrictions of the quadratic forms  $(X, QX)$  and  $(X, Q_0X)$  onto  $\mathbb{T}_\omega^0$ , and denote by  $\tilde{Q}$  and  $\tilde{Q}_0$  the corresponding nonnegative selfadjoint operators in  $\mathbb{T}_\omega^0$ :

$$(X, QX) = (X, \tilde{Q}X), \quad (X, Q_0X) = (X, \tilde{Q}_0X), \quad X \in \mathbb{D}^\infty \cap \mathbb{T}_\omega = \mathbb{T}_\omega^\infty. \quad (9.18)$$

The space  $\mathbb{T}_\omega^\infty$  is dense in  $\mathbb{T}_\omega^0$  and the operators can be expressed as

$$\tilde{Q} = P_\omega Q P_\omega, \quad \tilde{Q}_0 = P_\omega Q_0 P_\omega. \quad (9.19)$$

where  $P_\omega$  is the projection (9.13).

**Proposition 9.5.** *Let conditions (1.2) hold. Then  $\tilde{Q} \gg \mu \tilde{Q}_0$  with sufficiently small  $\mu > 0$ , i.e.*

$$\|\tilde{Q}X\|_{\mathbb{X}^0} \geq \mu \|\tilde{Q}_0X\|_{\mathbb{X}^0}, \quad X \in \mathbb{T}_\omega^\infty. \quad (9.20)$$

**Proof of Theorem 9.2.** Proposition 9.4 implies that  $\tilde{Q}$  is a selfadjoint nonnegative operator in  $\mathbb{T}_\omega^0$ , similarly to  $\tilde{Q}_0$ . Therefore, Proposition 9.5 and the Heinz inequality [24, Theorem 2] imply that  $\tilde{Q}^s \gg \mu^s \tilde{Q}_0^s$  for all  $s \in [0, 1]$ , i.e.,

$$\|\tilde{Q}^s X\|_{\mathbb{X}^0} \geq \mu^s \|\tilde{Q}_0^s X\|_{\mathbb{X}^0}, \quad X \in \mathbb{T}_\omega^\infty. \quad (9.21)$$

Finally, this bound with  $s = 1/2$  obviously implies (9.15) with  $\varkappa = \mu$ .  $\square$

## 10 Proof of Proposition 9.4

It suffices to prove the nonnegativity (9.17) for the case  $\beta = 0$ :

$$\mathcal{B} := \int |\nabla \alpha|^2 dy + \nu [\gamma - \langle m(y)\alpha(y), \rho(y) \rangle]^2 - \nu_{\text{eff}} \gamma^2 \geq 0, \quad (\alpha, \gamma) \in \mathbb{Y}^\infty := [\dot{\mathcal{F}}^1 \cap C_0^\infty] \oplus \mathbb{R}^3. \quad (10.1)$$

This form can be written as

$$\mathcal{B}(Y) = (Y, BY), \quad Y = (\alpha, \gamma) \in \mathbb{Y}^\infty,$$

where  $B$  is a selfadjoint operator in the Hilbert space  $\mathbb{Y}^0 := \mathcal{F}^0 \oplus \mathbb{R}^3$ , and the brackets  $(\cdot, \cdot)$  now denote the inner product in  $\mathbb{Y}^0$ . From (9.16), we obtain

$$B = \begin{pmatrix} -\sigma \Delta + \nu |m^* \rho(x)\langle \rho(y)m | & -\nu |m^* \rho(x)\rangle \\ -\nu \langle \rho m(y) | & \delta \end{pmatrix}. \quad (10.2)$$

Proposition 9.4 i) follows from the next lemma.

**Lemma 10.1.** *Let conditions (1.2) hold. Then the operator  $B$  is nonnegative.*

*Proof.* It suffices to show that the resolvent  $R(\lambda) = (B - \lambda)^{-1} : \mathbb{Y}^0 \rightarrow \mathbb{Y}^0$  is a bounded operator for  $\lambda < 0$ . The operator  $B - \lambda$  is a finite-dimensional perturbation of an invertible for  $\lambda < 0$  operator. Hence, it remains to check that

$$\ker(B - \lambda) = 0, \quad \lambda < 0. \quad (10.3)$$

The calculation of the kernel reduces to a finite-dimensional problem. Indeed, the equation  $(B - \lambda)Y = 0$  with  $Y = (\alpha, \gamma) \in \mathbb{Y}^0$  reads as

$$\begin{cases} (-\Delta - \lambda)\alpha + \nu m^* \rho \langle \rho m | \alpha \rangle - \nu m^* \rho \gamma & = 0 \\ -\nu \langle \rho m | \alpha \rangle + (\delta - \lambda)\gamma & = 0 \end{cases}. \quad (10.4)$$

Using the last equation, we rewrite the first one as

$$(-\Delta - \lambda)\alpha(x) = -m^*(x)\rho(x)(\delta - \lambda)\gamma + \nu m^*(x)\rho(x)\gamma = m^*(x)\rho(x)(\nu_{\text{eff}} + \lambda)\gamma, \quad \lambda \in \mathbb{R}. \quad (10.5)$$

Now

$$\alpha = (-\Delta - \lambda)^{-1}[m^*(x)\rho(x)(\nu_{\text{eff}} + \lambda)\gamma], \quad \lambda < 0. \quad (10.6)$$

**Remark 10.2.** For any vector  $w \in \mathbb{R}^3$ , the vector field  $m^*(x)\rho(x)w$  is divergence free by the spherical symmetry (1.2).

Substituting (10.6) into the second equation of (10.4), we obtain that

$$A(\lambda)\gamma = 0, \quad (10.7)$$

where we denote the matrix

$$A(\lambda) := -\nu \langle \rho(y)m(y), (-\sigma\Delta - \lambda)^{-1}[m^*(y)\rho(y)] \rangle (\nu_{\text{eff}} + \lambda) + (\delta - \lambda). \quad (10.8)$$

The matrix is Hermitian for  $\lambda \in \mathbb{R}$ , so it is defined uniquely by its quadratic form:

$$\begin{aligned} \gamma \cdot (A(\lambda)\gamma) &= -\nu \langle m^*(\nabla \hat{\rho}(k))\gamma, (\sigma k^2 - \lambda)^{-1} m^*(\nabla \hat{\rho}(k))\gamma \rangle (\nu_{\text{eff}} + \lambda) + (\delta - \lambda)\gamma^2 \\ &= -\nu \langle \nabla \hat{\rho}(k) \wedge \gamma, (\sigma k^2 - \lambda)^{-1} \nabla \hat{\rho}(k) \wedge \gamma \rangle (\nu_{\text{eff}} + \lambda) + (\delta - \lambda)\gamma^2 \\ &= -\nu \gamma^2 \int \frac{|\nabla \hat{\rho}(k)|^2 \sin^2(\widehat{k}, \gamma)}{\sigma k^2 - \lambda} dk (\nu_{\text{eff}} + \lambda) + (\delta - \lambda)\gamma^2 \\ &= -a_-(\lambda)\gamma^2 + a_+(\lambda)\gamma^2. \end{aligned} \quad (10.9)$$

Thus, the matrix  $A(\lambda)$  is the scalar  $-a_-(\lambda) + a_+(\lambda)$ , and (10.3) is equivalent to

$$a_-(\lambda) \neq a_+(\lambda), \quad \lambda < 0. \quad (10.10)$$

Note that the function  $a_-(\lambda)$  is strictly increasing, while  $a_+(\lambda)$  is strictly decreasing for  $\lambda \leq 0$ , and both functions are continuous. Hence, (10.10) is equivalent to the inequality

$$a_-(0) \leq a_+(0) \quad (10.11)$$

or

$$\nu\nu_{\text{eff}} \int \frac{|\nabla\hat{\rho}(k)|^2 \widehat{\sin^2(k, \gamma)}}{\sigma k^2} dk \leq \delta. \quad (10.12)$$

For  $\gamma = (0, 0, |\gamma|)$ , this inequality can be written as

$$\frac{1}{I(I + \delta I)} \int \frac{|\nabla\hat{\rho}(k)|^2 (k_1^2 + k_2^2)}{\sigma |k|^4} dk \leq \frac{1}{I} - \frac{1}{I + \delta I} = \frac{\delta I}{I(I + \delta I)}. \quad (10.13)$$

Now (6.11) with  $v = 0$  shows that both sides of the inequality (10.13) coincide.  $\square$

**Proof of Proposition 9.4.** The nonnegativity (9.17) follows from Lemma 10.1. For  $(\alpha, \beta, \gamma) \in \ker Q$  we have  $\beta = 0$ . Now (9.9) and (9.10) imply that  $\gamma = 0$  for  $(\alpha, \beta, \gamma) \in \ker Q \cap \mathbb{T}_\omega$ . Then also  $\alpha = 0$  by (10.5) with  $\lambda = 0$ . As the result,  $\ker Q \cap \mathbb{T}_\omega = 0$ .

## 11 Proof of Proposition 9.5

It suffices to prove that the operator  $\tilde{Q}_0 \tilde{Q}^{-1} : \mathbb{T}_\omega^0 \rightarrow \mathbb{T}_\omega^0$  is bounded. Indeed,  $\tilde{Q}_0 = \tilde{Q}_0 \tilde{Q}^{-1} \tilde{Q}$ , so we obtain  $\|\tilde{Q}_0 X\|_{\mathbb{X}^0} \leq C \|\tilde{Q} X\|_{\mathbb{X}^0}$  for  $X \in \mathbb{T}_\omega^\infty$  which is equivalent to (9.20).

The difference  $Q - Q_0$  is a finite-rank bounded operator in  $\mathbb{Z}^0$ . Hence,  $\tilde{Q} - \tilde{Q}_0$  is a finite-rank bounded operator in the Hilbert space  $\mathbb{T}_\omega^0$  by (9.19) and (9.13) since  $E_n \in \mathbb{Z}^0$  by (9.12). Therefore, the selfadjoint operators  $\tilde{Q}$  and  $\tilde{Q}_0$  have an identical domain  $\tilde{\mathbb{D}}^0$ . Moreover, Proposition 9.4 implies that  $\tilde{Q}$  is positive definite, and hence, it is a bijection of  $\tilde{\mathbb{D}}^0$  onto  $\mathbb{T}_\omega^0$ . Therefore, the operator  $\tilde{Q}_0 \tilde{Q}^{-1}$  is well defined on the entire Hilbert space  $\mathbb{T}_\omega^0$ , so the operator is bounded by the closed graph theorem.

## A Comments

I. In the case  $v = 0$  and  $\omega \neq 0$  the lower bound (7.1) fails because of the three-dimensional kernel of the quadratic form (9.5). One can try to improve the Lyapunov function (9.4) as follows:

$$\Lambda(\mathbf{Z}) = \mathbf{H}(\mathbf{Z}) - f(|\pi|), \quad \mathbf{Z} = (\mathbf{A}, \mathbf{\Pi}, \pi). \quad (\text{A.1})$$

The identity (7.3) implies that the function  $f$  must satisfy the identity  $D_\pi f(|\pi_\omega|) = \omega$ : using (6.10) with  $v = 0$ , we obtain

$$f'(|\pi_\omega|) \frac{\pi_\omega}{|\pi_\omega|} = f'(|\pi_\omega|) \frac{\omega}{|\omega|} = \omega, \quad (\text{A.2})$$

that is,  $f'(|\pi_\omega|) = |\omega|$ . In (9.4), we have chosen a particular function satisfying this condition, however, any modification of  $f$  does not help. Indeed, the difference  $f(|\pi_\omega + \gamma|) - f(|\pi_\omega|) = 0$  on the sphere  $|\pi_\omega + \gamma| - |\pi_\omega| = 0$  which contains the point  $\gamma = 0$ . The tangent plane to this sphere at the point  $\gamma = 0$  is orthogonal to  $\omega$  by (6.10). Accordingly, for any  $f$ , the function (A.1), restricted to the plane  $\{\mathbf{S}_\omega + (0, 0, \gamma) : \gamma \perp \omega\}$  has a degenerate Hessian: for  $\gamma \perp \omega$ , we have

$$\begin{aligned} [\partial_\varepsilon^2 \Lambda(\mathbf{A}_\omega, \mathbf{\Pi}_\omega, \pi_\omega + \varepsilon\gamma)]|_{\varepsilon=0} &= \partial_\varepsilon^2 [\mathbf{H}(\mathbf{A}_\omega, \mathbf{\Pi}_\omega, \pi_\omega + \varepsilon\gamma) - f(|\pi_\omega + \varepsilon\gamma|)]|_{\varepsilon=0} \\ &= \partial_\varepsilon [\omega \cdot \varepsilon\gamma]|_{\varepsilon=0} - \partial_\varepsilon^2 f(|\pi_\omega + \varepsilon\gamma|)|_{\varepsilon=0} = 0. \end{aligned} \quad (\text{A.3})$$

II. The tangent space  $\mathbb{T}_\omega$  *formally* is invariant with respect to the dynamics (9.1) linearized at the stationary point  $\mathbf{S}_\omega$ . Indeed, the linearized equation reads  $\dot{\xi} = dF(\mathbf{S}_\omega)\xi$ . The invariance means that the identity  $(dJ(\mathbf{S}_\omega), \xi) = 0$  implies

$$(dJ(\mathbf{S}_\omega), dF(\mathbf{S}_\omega)\xi) = 0. \quad (\text{A.4})$$

Let us write the system (9.1) as  $\dot{\mathbf{Z}}(t) = F(\mathbf{Z}(t))$ . Then the conservation of angular momentum (9.6) implies that

$$(dJ(\mathbf{Z}(t)), \dot{\mathbf{Z}}(t)) = (dJ(\mathbf{Z}(t)), F(\mathbf{Z}(t))) = 0.$$

Hence,  $(dJ(\mathbf{Z}), F(\mathbf{Z})) = 0$  for  $\mathbf{Z} \in \mathbb{M}_\omega$ . Applying the differentiation  $L_\xi$  in the direction of a vector  $\xi \in \mathbb{T}_\omega$ , we obtain  $(L_\xi dJ(\mathbf{S}_\omega), F(\mathbf{S}_\omega)) + (dJ(\mathbf{S}_\omega), L_\xi F(\mathbf{S}_\omega)) = 0$ . However, the soliton  $\mathbf{S}_\omega$  is the stationary solution, so  $F(\mathbf{S}_\omega) = 0$  and  $(dJ(\mathbf{S}_\omega), L_\xi F(\mathbf{S}_\omega)) = 0$  which implies (A.4) since  $dF(\mathbf{S}_\omega)\xi := L_\xi F(\mathbf{S}_\omega)$ .

III. The bound (9.15) follows from the estimate (9.21) with  $s = 1/2$  for the square roots  $\tilde{Q}^{1/2}$  and  $\tilde{Q}_0^{1/2}$  which are pseudodifferential operators. Our method of proof of Theorem 9.2 reduces the problem to calculations with differential operators and allows us to avoid calculations of pseudodifferential ones.

## B Proof of Lemma 6.2

**ad i).** Substituting (6.6) into (6.2), we obtain

$$\begin{aligned} \pi_{v,\omega} &= \omega - \langle \mathbf{A}_{v,\omega}(y) \wedge y \rho(y) \rangle = I\omega - \left\langle \frac{\omega \wedge \nabla \hat{\rho}(k)}{k^2 - (v \cdot k)^2} \wedge \nabla \hat{\rho}(k) \right\rangle = I\omega - \int \frac{(\omega \wedge k) \wedge k}{k^2 - (v \cdot k)^2} \frac{|\check{\rho}'_1(|k|)|^2 dk}{k^2} \\ &= \omega \left( I + \int \frac{|\check{\rho}'_1(|k|)|^2 dk}{k^2 - (v \cdot k)^2} \right) - \int \frac{k(\omega \cdot k) |\check{\rho}'_1(|k|)|^2}{k^2(k^2 - (v \cdot k)^2)} dk = \omega \left( I + \int \frac{|\check{\rho}'_1(|k|)|^2 dk}{k^2 - (v \cdot k)^2} \right) - q_{v,\omega}, \end{aligned} \quad (\text{B.1})$$

where we denote  $\check{\rho}_1(|k|) := \hat{\rho}(k)$ . We consider two cases  $\omega \parallel v$  and  $\omega \perp v$  separately.

**The case  $\omega \parallel v$ .** We may assume that  $\omega = (|\omega|, 0, 0)$  and  $v = (\pm|v|, 0, 0)$ . Then

$$q_{v,\omega} := \int \frac{k(\omega \cdot k) |\check{\rho}'_1(|k|)|^2}{k^2(k^2 - (v \cdot k)^2)} dk = |\omega| \int \frac{k_1 k |\check{\rho}'_1(|k|)|^2}{k^2(k^2 - v^2 k_1^2)} dk = |\omega| e_1 \int \frac{k_1^2 |\check{\rho}'_1(|k|)|^2}{k^2(k^2 - v^2 k_1^2)} dk. \quad (\text{B.2})$$

Substituting into (B.1), we obtain

$$\pi_{v,\omega} = I_{\text{eff}}^{\parallel}(v)\omega, \quad I_{\text{eff}}^{\parallel}(v) = I + \left\langle \frac{k_2^2 + k_3^2}{k^2(k^2 - v^2k_1^2)}, |\check{\rho}'_1(|k|)|^2 \right\rangle. \quad (\text{B.3})$$

**The case  $\omega \perp v$ .** We may assume that  $\omega = (|\omega|, 0, 0)$  and  $v = (0, |v|, 0)$ . In this case

$$q_{v,\omega} = |\omega|e_1 \int \frac{k_1^2 |\check{\rho}'_1(|k|)|^2}{k^2(k^2 - v^2k_2^2)} dk$$

and

$$\pi_{v,\omega} = I_{\text{eff}}^{\perp}(v)\omega, \quad I_{\text{eff}}^{\perp}(v) = I + \left\langle \frac{k_2^2 + k_3^2}{k^2(k^2 - v^2k_2^2)}, |\check{\rho}'_1(|k|)|^2 \right\rangle \quad (\text{B.4})$$

by (B.1). Now Lemma 6.2 i) is proved.

**ad ii).** Now we consider the case  $(v, \omega) \notin \Sigma$ . We may assume that  $v = (|v|, 0, 0)$  with  $|v| \neq 0$ , and  $\omega = (\omega_1, \omega_2, 0)$ , where  $\omega_1 \neq 0$  and  $\omega_2 \neq 0$ . In this case

$$q_{v,\omega} = \int \begin{pmatrix} k_1^2 \omega_1 \\ k_2^2 \omega_2 \\ 0 \end{pmatrix} \frac{|\check{\rho}'_1(|k|)|^2 dk}{k^2(k^2 - v^2k_1^2)} = \begin{pmatrix} \omega_1 \alpha_1 \\ \omega_2 \alpha_2 \\ 0 \end{pmatrix}, \quad \alpha_j = \int \frac{k_j^2 |\check{\rho}'_1(|k|)|^2 dk}{k^2(k^2 - v^2k_1^2)}, \quad j = 1, 2.$$

It remains to show that  $\alpha_1 > \alpha_2$  since then  $q_{v,\omega} \not\parallel \omega$ , so (6.9) breaks down by (B.1). Calculating in spherical coordinates, we get

$$\alpha_1 = \frac{2\pi C_\rho}{|v|^3} \left( \ln \frac{1+|v|}{1-v} - 2|v| \right), \quad \alpha_2 = \frac{\pi C_\rho}{|v|^3} \left( (v^2 - 1) \ln \frac{1+|v|}{1-|v|} + 2|v| \right),$$

where  $C_\rho > 0$  due to the last condition of (1.2). Hence,

$$\alpha_1 - \alpha_2 = \frac{\pi C_\rho}{|v|^3} \left( (3 - |v|^2) \ln \frac{1+|v|}{1-|v|} - 6|v| \right).$$

Expanding into the Taylor series for  $|v| < 1$ , we obtain

$$\begin{aligned} (3 - |v|^2) \ln \frac{1+|v|}{1-|v|} - 6|v| &= (3 - |v|^2) \left( 2|v| + 2\frac{|v|^3}{3} + 2\frac{|v|^5}{5} + 2\frac{|v|^7}{7} + \dots \right) - 6|v| \\ &= \left( \frac{6}{5} - \frac{2}{3} \right) |v|^5 + \left( \frac{6}{7} - \frac{2}{5} \right) |v|^7 + \dots + 2 \left( \frac{3}{2k+1} - \frac{1}{2k-1} \right) |v|^{2k+1} + \dots > 0 \end{aligned}$$

since  $\frac{3}{2k+1} - \frac{1}{2k-1} = \frac{4k-4}{(2k+1)(2k-1)} > 0$  for  $k \geq 2$ . Hence,  $\alpha_1 - \alpha_2 > 0$ .

**Remark B.1.** The inequality  $\alpha_1 > \alpha_2$  and the formulas (B.3), (B.4) easily imply that  $I_{\text{eff}}^{\perp}(v) > I_{\text{eff}}^{\parallel}(v)$  for  $\rho(x) \neq 0$ .



## References

- [1] M. Abraham, Prinzipien der Dynamik des Elektrons, *Ann. Phys. (Leipz.)* **10** (1903) 105–179.
- [2] M. Abraham, Theorie der Elektrizität, Bd.2: Elektromagnetische Theorie der Strahlung, Teubner, Leipzig, 1905.
- [3] V. I. Arnold, Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits, *Ann. Institut Fourier* **16** (1966), no. 1, 319–361.
- [4] V.I. Arnold, B.S. Khesin, Topological Methods in Hydrodynamics, Springer, New York, 1998.
- [5] V.I. Arnold, V.V. Kozlov, A.I. Neishtadt, Mathematical Aspects of Classical and Celestial Mechanics, Springer, Berlin, 1997.
- [6] D. Bambusi, L. Galgani, Some rigorous results on the Pauli–Fierz model of classical electrodynamics, *Ann. de l'I.H.P., section A* **58**(1993), no. 2, 155–171.
- [7] M. Born, Atomic Physics, Blackie & Son, London–Glasgow, 1951.
- [8] V.S. Buslaev, G. Perelman, On the stability of solitary waves for nonlinear Schrödinger equations, *Amer. Math. Soc. Trans.* **164** (1995), no. 2, 75–98.
- [9] V.S. Buslaev, C. Sulem, On asymptotic stability of solitary waves for nonlinear Schrödinger equations, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **20** (2003), no. 3, 419–475.
- [10] P. A. M. Dirac, Classical theory of radiating electrons, *Proc. R. Soc. Lond. A* **167** (1938), 148–169.
- [11] E. Gibney, Muons' big moment could fuel new physics, *Nature* **544** (2017), 145–146. <https://doi.org/10.1038/544145a>
- [12] M. Grillakis, J. Shatah, W.A. Strauss, Stability theory of solitary waves in the presence of symmetry, I; II. *J. Func. Anal.* **74** (1987), 160-197; **94** (1990), 308-348.
- [13] F. Hiroshima, H. Spohn, Mass renormalization in nonrelativistic quantum electrodynamics, *Journal of Mathematical Physics* **46** (2005), 042302.
- [14] D. Holm, J. E. Marsden, T. S. Ratiu, A. Weinstein, Nonlinear stability of fluid and plasma equilibria *Physics Reports* **123** (1985), 1–116.
- [15] D. Holm, T. Schmah, C. Stoica, Geometric Mechanics and Symmetry, Oxford University Press, Oxford, 2009.
- [16] V. Imaykin, A. Komech, N. Mauser, Soliton-type asymptotics for the coupled Maxwell–Lorentz equations, *Ann. Inst. Poincaré, Phys. Theor.* **5** (2004) 1117–1135.

- [17] V. Imaykin, A. Komech, H. Spohn, Rotating charge coupled to the Maxwell field: scattering theory and adiabatic limit, *Monatsh. Math.* **142** (2004), no. 1–2, 143–156.
- [18] V. Imaykin, A. Komech, H. Spohn, Scattering asymptotics for a charged particle coupled to the Maxwell field, *J. Math. Phys.* **52** (2011), no. 4, 042701.
- [19] V. Imaykin, A. Komech, H. Spohn, On the Lagrangian theory for rotating charge in the Maxwell field, *Physics Letters A* **379** (2015), no. 1–2, 5–10.
- [20] V. Imaykin, A. Komech, H. Spohn, On invariants for the Poincaré equations and applications, *J. Math. Phys.* **58** (2017), no. 1, 012901-1 – 012901-13.
- [21] V. Imaikin, A. Komech, B. Vainberg, On scattering of solitons for the Klein-Gordon equation coupled to a particle, *Comm. Math. Phys.* **268** (2006), no. 2, 321–367.
- [22] V. Imaikin, A. Komech, B. Vainberg, Scattering of solitons for coupled wave-particle equations, *J. Math. Analysis and Appl.* **389** (2012), no. 2, 713–740.
- [23] J.D. Jackson, *Classical Electrodynamics*, 3rd edition, Wiley, New York, 1999.
- [24] T. Kato, Notes on some inequalities for linear operators, *Math. Ann.* **125** (1952), no. 1, 208–212.
- [25] M. Kiessling, Classical electron theory and conservation laws, *Phys. Lett. A* **258** (1999), 197–204.
- [26] A. Komech, *Quantum Mechanics: Genesis and Achievements*, Springer, Dordrecht, 2013.
- [27] A. Komech, *Lectures on Quantum Mechanics and Attractors*, World Scientific, Singapore, 2022.
- [28] A. Komech, E. Kopylova, Attractors of nonlinear Hamiltonian partial differential equations, *Russ. Math. Surv.* **75** (2020), no. 1, 1–87.
- [29] A. Komech, E. Kopylova, *Attractors of Hamiltonian Nonlinear Partial Differential Equations*, Cambridge University Press, Cambridge, 2022.
- [30] A. Komech, E. Kopylova, On orbital stability of solitons for 2D Maxwell–Lorentz equations with spinning particle, submitted to *Comm. Pure Appl. Anal.*, 2023.
- [31] A. Komech, E. Kopylova, On the Hamilton–Poisson structure and solitons for the Maxwell–Lorentz equations with spinning particle, *J. Math. Anal. Appl.* **522**, no. 2, 15 June 2023, 126976.
- [32] A. Komech, E. Kopylova, Momentum map for the Maxwell–Lorentz equations with spinning particle, submitted to *Dynamics PDE*, 2023.
- [33] A. Komech, E. Kopylova, H. Spohn, Scattering of solitons for Dirac equation coupled to a particle, *J. Math. Analysis and Appl.* **383** (2011), no. 2, 265–290.

- [34] A. Komech, M. Kunze, H. Spohn, Effective dynamics for a mechanical particle coupled to a wave field, *Comm. Math. Phys.* **203** (1999), 1–19.
- [35] A. Komech, H. Spohn, Soliton-like asymptotics for a scalar particle interacting with a scalar wave field, *Nonlinear Analysis* **33** (1998), no. 1, 13–24.
- [36] A.I. Komech, H. Spohn, Long-time asymptotics for the coupled Maxwell–Lorentz equations, *Comm. Partial Differential Equations* **25** (2000), 559–584.
- [37] A.I. Komech, H. Spohn, M. Kunze, Long-time asymptotics for a classical particle interacting with a scalar wave field, *Comm. Partial Differential Equations* **22** (1997), 307–335.
- [38] M. Kunze, On the absence of radiationless motion for a rotating classical charge *Advances in Mathematics* **223**, no. 5, 1632–1665.
- [39] M. Kunze, H. Spohn, Adiabatic limit for the Maxwell-Lorentz equations, *Ann. Henri Poincaré* **1** (2000), 625–653.
- [40] S. Lie, *Theorie der Transformationsgruppen*, Teubner, Leipzig, 1890.
- [41] J.E. Marsden, G. Misiolek, J.-P. Ortega, M. Perlmutter, T.S. Ratiu, *Hamiltonian Reduction by Stages*, Springer, 2007.
- [42] E.T. Matsui, An equivariant Liapunov stability test and the energy-momentum-Casimir method, *Symplectic Geometry* **1** (2002), no. 4, 683–693.
- [43] J.S. Nodvik, A covariant formulation of classical electrodynamics for charges of finite extensions, *Ann. Phys.* **28** (1964), 225–319.
- [44] H. Poincaré, Sur une forme nouvelle des équations de la mécanique, *C. R. Acad. Sci.* **132** (1901), 369–371.
- [45] J.J. Sakurai, *Advanced Quantum Mechanics*, Pearson Education, Incorporated, 2006.
- [46] H. Spohn, *Dynamics of Charged Particles and Their Radiation Field*, Cambridge University Press, Cambridge, 2004.