

**HÖLDER REGULARITY OF WEAK SOLUTIONS  
TO NONLOCAL  $p$ -LAPLACIAN TYPE SCHRÖDINGER EQUATIONS  
WITH  $A_1^p$ -MUCKENHOUPT POTENTIALS**

YONG-CHEOL KIM

ABSTRACT. In this paper, using the De Giorgi-Nash-Moser method, we obtain an interior Hölder continuity of weak solutions to nonlocal  $p$ -Laplacian type Schrödinger equations given by an integro-differential operator  $L_K^p$  ( $p > 1$ ) as follows;

$$\begin{cases} L_K^p u + V|u|^{p-2}u = 0 & \text{in } \Omega, \\ u = g & \text{in } \mathbb{R}^n \setminus \Omega \end{cases}$$

where  $V = V_+ - V_-$  with  $(V_-, V_+) \in L_{loc}^1(\mathbb{R}^n) \times L_{loc}^q(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  and  $0 < s < 1$  is a potential such that  $(V_-, V_+^{b,i})$  belongs to the  $(A_1, A_1)$ -Muckenhoupt class and  $V_+^{b,i}$  is in the  $A_1$ -Muckenhoupt class for all  $i \in \mathbb{N}$  (here,  $V_+^{b,i} := V_+ \max\{b, 1/i\}/b$  for an almost everywhere positive bounded function  $b$  on  $\mathbb{R}^n$  with  $V_+/b \in L_{loc}^q(\mathbb{R}^n)$ ,  $g \in W^{s,p}(\mathbb{R}^n)$  and  $\Omega \subset \mathbb{R}^n$  is a bounded domain with Lipschitz boundary.) In addition, we get the local boundedness of weak subsolutions of the nonlocal  $p$ -Laplacian type Schrödinger equations. In a different way from [DKP1], we obtain the logarithmic estimate of the weak supersolutions which play a crucial role in proving the Hölder regularity of the weak solutions.

In particular, we note that all the above results are still working for any non-negative potential in  $L_{loc}^q(\mathbb{R}^n)$  ( $q > \frac{n}{ps} > 1, 0 < s < 1$ ).

CONTENTS

1. Introduction	1
2. Preliminaries	7
3. Weighted norm inequalities on the $A_p$ -Muckenhoupt class	11
4. Local properties of weak subsolutions	15
5. The logarithm of a weak solution is a locally bounded mean oscillation function	22
6. Interior Hölder regularity	27
References	33

1. INTRODUCTION

The recent research on fractional and nonlocal partial differential equations has extensively and actively been performed not only in pure mathematics but also in scientific areas that require its concrete applications. This type of problems appear in various applications such as continuum mechanics, phase transition phenomena that are related to a nonlocal version of the classical Allen-Cahn equation, population dynamics, nonlocal minimal surfaces, a nonlocal version of

Schrödinger equations for standing waves ( see [BV, CK1, CK2] ), game theory and also constrained variational problems with fractional diffusion arising in the quasi-geostrophic flow model, anomalous diffusions and American options with jump processes ( see [CV, C, S] ).

For  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ), let  $\mathcal{P}_q^{s,p}(\mathbb{R}^n)$  be the class of all potentials  $V = V_+ - V_-$  such that (i)  $V_- \in L_{loc}^1(\mathbb{R}^n)$ , (ii)  $V_+ \in L_{loc}^q(\mathbb{R}^n)$ , (iii) there is an almost everywhere positive bounded function  $b$  on  $\mathbb{R}^n$  so that

$V_+/b \in L_{loc}^q(\mathbb{R}^n)$ ,  $(V_-, V_+^{b,i})$  belongs to the  $(A_1, A_1)$ -Muckenhoupt class and  $V_+^{b,i}$  is in the  $A_1$ -Muckenhoupt class for all  $i \in \mathbb{N}$ , where

$$V_+^{b,i} := \frac{\max\{b, 1/i\}}{b} V_+.$$

If  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ), then we say that  $V$  is a  $A_1^p$ -Muckenhoupt potential. When  $p = 2$ , we call it  $A_1$ -Muckenhoupt potential.

The aim of this paper is to establish an interior Hölder regularity of weak solutions of nonlocal  $p$ -Laplacian type Schrödinger equations with  $A_1^p$ -Muckenhoupt potentials and to additionally obtain the local boundedness of weak subsolutions of the nonlocal equation.

For  $p > 1$ , let  $\mathcal{K}_p$  be the collection of all positive symmetric kernels satisfying the uniformly ellipticity assumption

$$(1.1) \quad \frac{c_{n,p,s} \lambda}{|y|^{n+ps}} \leq K(y) = K(-y) \leq \frac{c_{n,p,s} \Lambda}{|y|^{n+ps}}, \quad 0 < s < 1, \quad y \in \mathbb{R}^n \setminus \{0\},$$

where  $c_{n,p,s} > 0$  is the normalization constant given by

$$c_{n,p,s} = \frac{\Gamma(\frac{n+p}{2}) p (1-s)}{\pi^{\frac{n-1}{2}} \Gamma(\frac{p+1}{2})}.$$

For  $K \in \mathcal{K}_p$  ( $p > 1$ ), we consider integro-differential operators  $L_K^p$  given by

$$(1.2) \quad L_K^p u(x, t) = \text{p.v.} \int_{\mathbb{R}^n} H_p(u(x) - u(y)) K(x - y) dy$$

where  $H_p(t) = |t|^{p-2}t$  for  $t \in \mathbb{R}$ . If  $p = 2$ , then we write  $L_K^p = L_K$ . In particular, if  $K(y) = c_{n,p,s}|y|^{-n-ps}$ , then  $L_K^p = (-\Delta)_p^s$  is the fractional  $p$ -Laplacian and it is well-known [IN] that

$$\lim_{s \rightarrow 1^-} (-\Delta)_p^s u = -\Delta_p u$$

for any function  $u$  in the Schwartz space  $\mathcal{S}(\mathbb{R}^n)$ , where

$$\Delta_p u = \text{div}(|\nabla u|^{p-2} \nabla u)$$

is the classical  $p$ -Laplacian.

We are interested in the Dirichlet problem for the nonlocal  $p$ -Laplacian type Schrödinger equation

$$(1.3) \quad \begin{cases} L_K^p u + V|u|^{p-2}u = 0 & \text{in } \Omega \\ u = g & \text{in } \mathbb{R}^n \setminus \Omega \end{cases}$$

where  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$  and  $0 < s < 1$ ),  $g \in W^{s,p}(\mathbb{R}^n)$  and  $\Omega \subset \mathbb{R}^n$  is a bounded domain with Lipschitz boundary. The existence and uniqueness of weak solution to the above nonlocal equation was obtained in [K3] by applying

standard technique of calculus of variations. More precisely speaking about the problem, by employing the De Giorgi-Nash-Moser theory, we obtain an interior Hölder regularity of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equations, and also we get the local boundedness of weak subsolutions of the nonlocal equation. Here, we note that the boundary condition is imposed on  $\mathbb{R}^n \setminus \Omega$  with nonlocality. In fact, from the probabilistic point of view, it conforms to the natural phenomenon that a discontinuous Lévy process on the domain  $\Omega$  can exit  $\Omega$  for the first time jumping to any point in  $\mathbb{R}^n \setminus \Omega$ .

When  $p = 2$ , the research on the above nonlocal equations was strongly motivated by the study of standing wave solutions of the form

$$\Psi(x, t) = e^{-i\omega t} u(x)$$

of the time-dependent nonlocal Schrödinger equations

$$i \frac{\partial \Psi}{\partial t} = L_K \Psi + V(x) \Psi$$

which is a fundamental equation of fractional quantum mechanics and fractional quantum physics. This equation was used for the first time in the literature by Laskin ( see [L] ).

It turns out in Section 3 that any potential  $V$  in  $\mathcal{P}_q^{s,p}(\mathbb{R}^n)$  satisfies

$$(1.4) \quad \int_{\mathbb{R}^n} |\varphi(y)| V_-(y) dy \leq \int_{\mathbb{R}^n} |\varphi(y)| V_+(y) dy$$

and

$$(1.5) \quad \int_{\mathbb{R}^n} |\varphi(y)|^p V_-(y) dy \leq \int_{\mathbb{R}^n} |\varphi(y)|^p V_+(y) dy$$

for every  $\varphi \in Y_0^{s,p}(\Omega)$ , whenever  $q > \frac{n}{ps} > 1$  ( $p > 1$  and  $0 < s < 1$ ). As a matter of fact, the inequality (1.4) is a useful tool for the proof of nonlocal Caccioppoli type inequality to be given in Theorem 1.5, and also the inequality (1.5) makes it possible to prove in Lemma 3.3 below that  $Y_0^{s,p}(\Omega)$  is a quasi-Banach space.

**Notations.** We introduce the notations briefly for the readers as follows.

- For  $r > 0$ ,  $x_0 \in \mathbb{R}^n$  and  $s \in (0, 1)$ , let us denote by  $B_r^0 = B_r(x_0)$ ,  $B_r = B_r(0)$ . For  $n \in \mathbb{N}$ , denote by  $|S^{n-1}|$  the  $(n-1)$ -dimensional surface measure on the unit sphere  $S^{n-1}$  of  $\mathbb{R}^n$ .

- For two quantities  $a$  and  $b$ , we write  $a \lesssim b$  (resp.  $a \gtrsim b$ ) if there is a universal constant  $C > 0$  (depending only on  $\lambda, \Lambda, n, p, s$  and  $\Omega$ ) such that  $a \leq C b$  (resp.  $b \leq C a$ ).

- For  $a, b \in \mathbb{R}$ , we denote by

$$a \vee b = \max\{a, b\} \quad \text{and} \quad a \wedge b = \min\{a, b\}.$$

- Let  $\mathcal{F}^n$  be the family of all real-valued Lebesgue measurable functions on  $\mathbb{R}^n$ .

- For  $u \in C(B_r^0)$ , we consider the norm

$$\|u\|_{C(B_r^0)} = \sup_{x \in B_r^0} |u(x)|.$$

For  $\gamma \in (0, 1)$ , the  $\gamma^{\text{th}}$  Hölder seminorm of  $u$  on  $B_r^0$  is defined by

$$[u]_{C^\gamma(B_r^0)} = \sup_{x, y \in B_r^0, x \neq y} \frac{|u(x) - u(y)|}{|x - y|^\gamma}$$

and the  $\gamma^{\text{th}}$  Hölder norm of  $u$  on  $B_r^0$  is defined by

$$\|u\|_{C^\gamma(B_r^0)} = \|u\|_{C(B_r^0)} + [u]_{C^\gamma(B_r^0)}.$$

• For  $x_0 \in \Omega$ ,  $p > 1$  and  $r > 0$  with  $B_r^0 \subset \Omega$ , the *nonlocal tails* of the function  $u$  in  $B_r^0 \subset \Omega$  is defined by

$$(1.6) \quad \mathcal{T}_r(u; x_0) = \left( \frac{ps}{|S^{n-1}|} r^{ps} \int_{\mathbb{R}^n \setminus B_r(x_0)} \frac{|u(y)|^{p-1}}{|y-x_0|^{n+ps}} dy \right)^{\frac{1}{p-1}}.$$

We now state one of our main results which is called the local boundedness of weak subsolutions to the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), as follows.

**Theorem 1.1.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$ ,  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$ ,  $s \in (0, 1)$ ) and  $B_{2r}^0 \subset \Omega$ . If  $u \in Y_g^{s,p}(\Omega)^-$  is a weak subsolution of nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), then there is a constant  $C_0 > 0$  depending only on  $n, s, p, \lambda, \Lambda$  and  $\Omega$  such that*

$$\sup_{B_r^0} u \leq \delta \mathcal{T}_r(u_+; x_0) + C_0 \delta^{-\frac{(p-1)n}{sp^2}} \left( \int_{B_{2r}^0} u_+^p dx \right)^{\frac{1}{p}}$$

for any  $\delta \in (0, 1]$ .

**Remark.** (a) If  $u \in Y_g^{s,p}(\Omega)^-$  is a weak subsolution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.4) and  $g \in W^{s,p}(\mathbb{R}^n)$  for  $s \in (0, 1)$ , then we see that  $u \in L^p(\Omega)$  and  $u \leq g$  on  $\mathbb{R}^n \setminus \Omega$ , and thus  $u_+ \leq g_+$  there. Then it follows from Hölder's inequality and fractional Sobolev inequalities (2.4) and (2.5) that

$$\begin{aligned} [\mathcal{T}_r(u_+; x_0)]^{p-1} &\leq \frac{ps r^{ps}}{|S^{n-1}|} \left( \int_{\mathbb{R}^n \setminus \Omega} \frac{g_+^{p-1}(y)}{|y-x_0|^{n+ps}} dy + \int_{\Omega \setminus B_r(x_0)} \frac{|u(y)|^{p-1}}{|y-x_0|^{n+ps}} dy \right) \\ &\lesssim \frac{ps r^{-n(1-\frac{1}{p})}}{(((p-1)n + p^2s)|S^{n-1}|)^{1/p}} \left( \|g\|_{W^{s,p}(\mathbb{R}^n)}^{p-1} + \|u\|_{L^p(\Omega)}^{p-1} \right) < \infty. \end{aligned}$$

(b) If  $u \in Y_g^{s,p}(\Omega)^+$  is a weak supersolution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.4) and  $g \in W^{s,p}(\mathbb{R}^n)$  for  $s \in (0, 1)$ , then  $-u$  is its weak subsolution and  $u \geq g$  on  $\mathbb{R}^n \setminus \Omega$ , and so  $u_- \leq g_-$  there. Then, as in the above (a), we obtain that

$$[\mathcal{T}_r(u_-; x_0)]^{p-1} \lesssim \frac{ps r^{-n(1-\frac{1}{p})}}{(((p-1)n + p^2s)|S^{n-1}|)^{1/p}} \left( \|g\|_{W^{s,p}(\mathbb{R}^n)}^{p-1} + \|u\|_{L^p(\Omega)}^{p-1} \right) < \infty.$$

Then, from Theorem 1.1, we easily have that

$$-\inf_{B_r^0} u = \sup_{B_r^0} (-u) \leq \delta \mathcal{T}_r(u_-; x_0) + C_0 \delta^{-\frac{(p-1)n}{sp^2}} \left( \int_{B_{2r}^0} u_-^p dx \right)^{\frac{1}{p}}$$

for any  $\delta \in (0, 1]$ .

(c) If  $u \in Y_g^{s,p}(\Omega)^+$  is a weak solution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.4) and  $g \in W^{s,p}(\mathbb{R}^n)$  for  $s \in (0, 1)$ , then it follows from (a), (b) and Theorem 1.1 that

$$\text{osc}_{B_r^0} u \leq 2\delta \mathcal{T}_r(u; x_0) + 2C_0 \delta^{-\frac{(p-1)n}{sp^2}} \left( \int_{B_{2r}^0} |u|^p dx \right)^{\frac{1}{p}}$$

for any  $\delta \in (0, 1]$ .

The following *logarithmic estimate* plays a crucial role in proving the Hölder regularity of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation and in showing that the logarithm of such weak solution is a function with locally bounded mean oscillation. In a different way from [DKP1], we obtain the logarithmic estimate. We now state it as follows.

**Theorem 1.2.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  and  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$  and  $0 < s < 1$ ). If  $u \in Y_g^{s,p}(\Omega)^+$  is a weak supersolution of nonlocal  $p$ -Laplacian type Schrödinger equation (1.3) with  $u \geq 0$  in  $B_R^0 \subset \Omega$ , then there is a constant  $c_0 > 0$  depending only on  $n, s, p, \lambda, \Lambda$  and  $\Omega$  such that*

$$\begin{aligned} \iint_{B_r^0 \times B_r^0} \left| \ln \left( \frac{u(x) + b}{u(y) + b} \right) \right|^p d_K(x, y) \\ \leq c_0 r^{n-ps} \left[ \frac{1}{b^{p-1}} \left( \frac{r}{R} \right)^{ps} [\mathcal{T}_R(u_-; x_0)]^{p-1} + (1 + \|V_+\|_{L^q(\Omega)}) \right] \end{aligned}$$

for any  $b \in (0, 1)$  and  $r \in (0, R/2)$ , where  $d_K(x, y) = K(x - y) dx dy$ .

Employing the De Giorgi-Nash-Moser theory and using Theorem 1.1 and 1.2, we obtain the following Hölder continuity of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation, and also we can easily derive Corollary 1.4 as a natural by-product of Theorem 1.3.

**Theorem 1.3.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$ ,  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ), and let  $B_{2R}^0 \subset \Omega$ . If  $u \in Y_g^{s,p}(\Omega)$  is a weak solution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), then there exist constants  $\eta_0^- \in (0, \frac{ps}{2(p-1)})$  and  $\eta_0^+ \in (\frac{ps}{2(p-1)}, \frac{ps}{p-1})$  such that  $u$  is locally  $\eta$ -Hölder continuous in  $\Omega$  for any  $\eta \in (0, \eta_0^-] \cup [\eta_0^+, \frac{ps}{p-1})$ . Furthermore, for each  $x_0 \in \Omega$  and for each  $\eta \in (0, \eta_0^-] \cup [\eta_0^+, \frac{ps}{p-1})$ , we have that*

$$(1.7) \quad \text{osc}_{B_r^0} u \lesssim \left( \frac{r}{R} \right)^\eta \left[ \mathcal{T}_R(u; x_0) + \left( \int_{B_{2R}^0} |u(x)|^p dx \right)^{\frac{1}{p}} \right]$$

for any  $r \in (0, R/2)$ . Here it turns out that there exist universal constants  $c_0, c_* > 0$  such that

$$\eta_0^\pm = \frac{\ln \left( \frac{1 \pm \sqrt{1 - 4\delta^{\frac{ps}{p-1}}}}{2} \right)}{\ln \delta} \quad \text{for } \delta = e^{-(c_0/c_*)(1+\|V_+\|_{L^q(\Omega)})^{1/p}} \wedge \left( \frac{1}{4} \right)^{\frac{p-1}{ps}}.$$

The next corollary can easily be obtained by applying Theorem 1.3 and employing the interpolation on Hölder spaces between  $C^{\eta_0^-}(B_r^0)$  and  $C^{\eta_0^+}(B_r^0)$ , which eventually fill up an interior  $\eta$ -Hölder continuity of  $u$  in  $\Omega$  for all  $\eta \in (\eta_0^-, \eta_0^+)$ .

**Corollary 1.4.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$ ,  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ), and let  $B_{2R}^0 \subset \Omega$ . If  $u \in Y_g^{s,p}(\Omega)$  is a weak solution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), then we have the following estimate*

$$(1.8) \quad \sup_{r \in (0, R/2)} \|u\|_{C^\eta(B_r^0)} \lesssim \frac{1}{R^\eta} \left[ \mathcal{T}_R(u; x_0) + \left( \int_{B_{2R}^0} |u(x)|^p dx \right)^{\frac{1}{p}} \right]$$

for any  $\eta \in (0, \frac{ps}{p-1})$ .

**Remark.** If  $\frac{p-1}{p} < s < 1$ , then we can expect the better regularity, i.e.  $C^{1,\alpha}$ -estimate for some  $\alpha \in (0, 1)$ .

As a basic tool for our main results, we show that any weak subsolution of the nonlocal  $p$ -Laplacian type Schrödinger equations enjoys the following *nonlocal Caccioppoli type inequality*.

**Theorem 1.5.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$ ,  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$ ,  $s \in (0, 1)$ ), and let  $B_{2r}^0 \subset \Omega$ . If  $u \in Y_g^{s,p}(\Omega)^-$  is a weak subsolution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), then for any nonnegative  $\zeta \in C_c^\infty(B_r^0)$  we have the following estimate*

$$\begin{aligned} & \int_{B_r^0} [w(y)\zeta(y)]^p V(y) dy + \iint_{B_r^0 \times B_r^0} |\zeta(x)w(x) - \zeta(y)w(y)|^p d_K(x, y) \\ & \leq 2^{2p+1} \left( \frac{1}{4} + c_p \right) \iint_{B_r^0 \times B_r^0} [w(x) \vee w(y)]^p |\zeta(x) - \zeta(y)|^p d_K(x, y) \\ & \quad + 2^{p+2} \left( \sup_{x \in \text{supp}(\zeta)} \int_{\mathbb{R}^n \setminus B_r^0} w^{p-1}(y) K(x-y) dy \right) \|w\zeta^p\|_{L^1(B_r^0)} \end{aligned}$$

where  $w = (u - M)_+$  for  $M \in (0, \infty)$  and  $c_p = \frac{1}{2}[2(p-1)]^{p-1}$ .

**Remark.** (a) When  $p = 2$  and  $s = 1$ , the study of the classical Schrödinger operator, i.e. local Schrödinger operator  $-\Delta + V$  has been ongoing actively and widely in analysis area in Mathematics and Mathematical Physics (refer to [AS, CFG, F, S1, S2, Si]).

(b) In case that  $p = 2$  and  $V \in L_{\text{loc}}^q(\mathbb{R}^n)$  with  $q > \frac{n}{2s}$  ( $0 < s < 1$ ) is nonnegative, it is known in [CK1] that a fundamental solution for nonlocal Schrödinger operator  $L_K + V$  exists and its decay can be obtained. Also, under an additional restriction that the potential  $V$  is in a reverse Hölder class  $\text{RH}^\tau$  for  $\tau > \frac{n}{2s} > 1$  ( $0 < s < 1$ ), the  $L^a - L^b$  estimate for the Schrödinger operator  $L_K + V$  was obtained inside certain trapezoidal region  $\mathcal{Z}$  which is consist of  $(\frac{1}{a}, \frac{1}{b})$  and also the weak type  $L^a - L^b$  estimate was partially obtained on the boundary of the region  $\mathcal{Z}$  (see [CK2]).

(c) When  $p = 2$ ,  $0 < s < 1$  and  $V$  is an  $A_1$ -Muckenhoupt potential, it was shown in [K3] that a fundamental solution for nonlocal Schrödinger operator  $L_K + V$  exists and its decay can be obtained. Also, Hölder continuity and nonlocal Harnack inequalities for  $L_K + V$  were obtained in [K1] and [K2].

(d) When  $V = 0$  and  $0 < s < 1$ , the result of this problem was obtained by Di Castro, Kuusi and Palatucci [DKP]; as a matter of fact, when  $p \in (1, \infty)$ , they proved nonlocal Harnack inequalities for elliptic nonlocal  $p$ -Laplacian equations there, and also they obtained Hölder regularity in [DKP1].

(e) When  $p = 2$  and  $0 < s < 1$ , nonlocal Harnack inequalities for locally nonnegative weak solutions of nonlocal heat equations was obtained in [K] by applying the De Giorgi-Nash-Moser theory.

The paper is organized as follows. In Section 2, we furnish the function spaces and the definition of weak solutions of the nonlocal Schrödinger equation given in (1.3), and also give a well-known lemma which is useful in applying the De Giorgi-Nash-Moser theory. In Section 3, we give a brief introduction about weighted norm inequalities and the  $A_p$ -Muckenhoupt class. Additionally, we furnish several examples about sign-changing potentials in the class  $\mathcal{P}_q^{s,p}(\mathbb{R}^n)$  ( $q > \frac{n}{ps} > 1$ ,  $p > 1$ ,  $0 < s < 1$ ). In Section 4, we obtain a sort of *nonlocal Caccioppoli type inequality* and

several useful local properties of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation by using it. In Section 5, we show that the logarithm of a weak solution to the nonlocal  $p$ -Laplacian type Schrödinger equation becomes a function with locally bounded mean oscillation. In Section 6, we get an interior Hölder continuity of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation by applying the results obtained in Section 4 and Section 5.

## 2. PRELIMINARIES

Let  $\Omega \subset \mathbb{R}^n$  be a bounded domain with Lipschitz boundary and let  $K \in \mathcal{K}_p$  for  $p > 1$ . For  $p > 1$  and  $0 < s < 1$ , let  $X^{s,p}(\Omega)$  be the linear function space of all Lebesgue measurable functions  $v \in \mathcal{F}^n$  such that  $v|_\Omega \in L^p(\Omega)$  and

$$\iint_{\mathbb{R}_\Omega^{2n}} \frac{|v(x) - v(y)|^p}{|x - y|^{n+ps}} dx dy < \infty$$

where  $\mathbb{R}_S^{2n} := \mathbb{R}^{2n} \setminus (S^c \times S^c)$  for a set  $S \subset \mathbb{R}^n$ . We also set

$$(2.1) \quad X_0^{s,p}(\Omega) = \{v \in X^{s,p}(\Omega) : v = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega\}$$

Since  $C_0^2(\Omega) \subset X_0^{s,p}(\Omega)$ , we see that  $X^{s,p}(\Omega)$  and  $X_0^{s,p}(\Omega)$  are nonempty. Then we see that  $(X^{s,p}(\Omega), \|\cdot\|_{X^{s,p}(\Omega)})$  is a normed space with the norm  $\|\cdot\|_{X^{s,p}(\Omega)}$  given by

$$(2.2) \quad \|v\|_{X^{s,p}(\Omega)} = \|v\|_{L^p(\Omega)} + \left( \iint_{\mathbb{R}_\Omega^{2n}} \frac{|v(x) - v(y)|^p}{|x - y|^{n+ps}} dx dy \right)^{\frac{1}{p}} < \infty$$

for  $v \in X^{s,p}(\Omega)$ . For  $p \geq 1$ , we denote by  $W^{s,p}(\Omega)$  the usual fractional Sobolev space with the norm

$$(2.3) \quad \|v\|_{W^{s,p}(\Omega)} := \|v\|_{L^p(\Omega)} + [v]_{W^{s,p}(\Omega)} < \infty$$

where the seminorm  $[\cdot]_{W^{s,p}(\Omega)}$  is defined by

$$[v]_{W^{s,p}(\Omega)} = \left( \iint_{\Omega \times \Omega} \frac{|v(x) - v(y)|^p}{|x - y|^{n+ps}} dx dy \right)^{\frac{1}{p}}.$$

When  $\Omega = \mathbb{R}^n$  in (2.3), similarly we define the spaces  $W^{s,p}(\mathbb{R}^n)$  for  $p \geq 1$  and  $s \in (0, 1)$ .

If  $p \geq 1$  and  $s \in (0, 1)$  satisfy  $ps < n$ , then it is well-known [DPV] that there exists a universal constant  $c = c(n, p, s, \Omega) > 0$  such that

$$(2.4) \quad \|f\|_{L^\tau(\Omega)} \leq c \|f\|_{W^{s,p}(\Omega)}$$

for any  $f \in W^{s,p}(\Omega)$  and  $\tau \in [p, p_*]$ , where  $p_* = \frac{pn}{n-ps}$ . Furthermore, there is a universal constant  $c = c(n, p, s) > 0$  such that

$$(2.5) \quad \|f\|_{L^\tau(\mathbb{R}^n)} \leq c \|f\|_{W^{s,p}(\mathbb{R}^n)}, \forall \tau \in [p, p_*] \text{ and } \|f\|_{L^{p_*}(\mathbb{R}^n)} \leq c [f]_{W^{s,p}(\mathbb{R}^n)}$$

for any  $f \in W^{s,p}(\mathbb{R}^n)$ . Using (2.5), we easily see that there exists a constant  $c > 1$  depending only on  $n, p, s$  and  $\Omega$  such that

$$(2.6) \quad \|u\|_{X_0^{s,p}(\Omega)} \leq \|u\|_{X^{s,p}(\Omega)} \leq c \|u\|_{X_0^{s,p}(\Omega)}$$

for any  $u \in X_0^{s,p}(\Omega)$ , where

$$(2.7) \quad \|u\|_{X_0^{s,p}(\Omega)} := \left( \iint_{\mathbb{R}_\Omega^{2n}} \frac{|u(x) - u(y)|^p}{|x - y|^{n+ps}} dx dy \right)^{\frac{1}{p}}.$$

Thus  $\|\cdot\|_{X_0^{s,p}(\Omega)}$  is a norm on  $X_0^{s,p}(\Omega)$  which is equivalent to (2.2). By using the change of variables, we can easily derive the following version of the *fractional Sobolev inequality* (2.4).

**Proposition 2.1.** *Let  $B_R$  be a ball with radius  $R > 0$ . If  $s \in (0, 1)$  and  $p \in [1, \infty)$  with  $sp < n$ , then there is a constant  $c = c(n, p, s) > 0$  such that*

$$\|f\|_{L^\tau(B_R)} \leq c R^{-n(\frac{1}{p}-\frac{1}{\tau})} \|f\|_{L^p(B_R)} + c R^{-n(\frac{1}{p}-\frac{1}{\tau})+s} [f]_{W^{s,p}(B_R)}$$

for any  $\tau \in [p, p_*]$ . In particular, if  $\tau = p_* := \frac{pn}{n-ps}$ , then we have that

$$(2.8) \quad \|f\|_{L^{p_*}(B_R)} \leq c R^{-s} \|f\|_{L^p(B_R)} + c [f]_{W^{s,p}(B_R)}.$$

For  $g \in W^{s,p}(\mathbb{R}^n)$ , we consider the convex subsets of  $X^{s,p}(\Omega)$  by

$$\begin{aligned} X_g^{s,p}(\Omega)^\pm &= \{v \in X^{s,p}(\Omega) : (g-v)_\pm \in X_0^{s,p}(\Omega)\}, \\ X_g^{s,p}(\Omega) &:= X_g^{s,p}(\Omega)^+ \cap X_g^{s,p}(\Omega)^- = \{v \in X^{s,p}(\Omega) : g-v \in X_0^{s,p}(\Omega)\}. \end{aligned}$$

For  $g \in W^{s,p}(\mathbb{R}^n)$  and a potential  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  with  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ), let

$$Y^{s,p}(\Omega) = X^{s,p}(\Omega) \cap L_V^p(\Omega) \quad \text{and} \quad Y_g^{s,p}(\Omega) = X_g^{s,p}(\Omega) \cap L_V^p(\Omega)$$

where  $L_V^p(\Omega)$  is the weighted  $L^p$  class of all real-valued measurable functions  $u$  on  $\mathbb{R}^n$  satisfying

$$\begin{aligned} -\infty < \|u\|_{L_V^p(\Omega)}^p &:= \int_\Omega |u(y)|^p V_+(y) dy - \int_\Omega |u(y)|^p V_-(y) dy \\ &:= \|u\|_{L_{V_+}^p(\Omega)}^p - \|u\|_{L_{V_-}^p(\Omega)}^p < \infty. \end{aligned}$$

That is, we see that  $u \in L_V^p(\Omega)$  if and only if  $u \in L_{V_+}^p(\Omega) \cap L_{V_-}^p(\Omega)$ . Here, we note that  $\|u\|_{L_V^p(\Omega)}^p$  need not be nonnegative, and so the class  $L_V^p(\Omega)$  is not always a normed space.

Also we consider function spaces  $Y_g^{s,p}(\Omega)^+$  and  $Y_g^{s,p}(\Omega)^-$  defined by

$$Y_g^{s,p}(\Omega)^\pm = \{u \in Y^{s,p}(\Omega) : (g-u)_\pm \in Y_0^{s,p}(\Omega)\}.$$

Then we see that

$$Y_g^{s,p}(\Omega) = Y_g^{s,p}(\Omega)^+ \cap Y_g^{s,p}(\Omega)^-.$$

If  $u = g = 0$  in  $\mathbb{R}^n \setminus \Omega$ , then we easily know that  $Y_0^{s,p}(\Omega) = X_0^{s,p}(\Omega) \cap L_V^p(\Omega)$  need not be a Banach space. However, if  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ), then it turns out in Lemma 3.2 below that the class  $Y_0^{s,p}(\Omega)$  is a quasi-Banach space with the quasinorm  $\|\cdot\|_{Y_0^{s,p}(\Omega)}$  given by

$$\|u\|_{Y_0^{s,p}(\Omega)}^p := \|u\|_{X_0^{s,p}(\Omega)}^p + \int_\Omega |u(y)|^p V(y) dy, \quad u \in Y_0^{s,p}(\Omega),$$

$Y_0^{s,p}(\Omega) = X_0^{s,p}(\Omega)$  and they are quasinorm-equivalent.

In order to define weak solutions of the nonlocal equation (1.3), we consider a bilinear form  $\langle \cdot, \cdot \rangle_{H_p, K} : X^{s,p}(\Omega) \times X^{s,p}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\langle u, v \rangle_{H_p, K} = \iint_{\mathbb{R}^n \times \mathbb{R}^n} H_p(u(x) - u(y))(v(x) - v(y)) d_K(x, y)$$

where  $d_K(x, y) := K(x - y) dx dy$ .



**Definition 2.2.** Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  and  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1, 0 < s < 1$ ). Then we say that a function  $u \in Y_g^{s,p}(\Omega)^-$  ( $u \in Y_g^{s,p}(\Omega)^+$ ) is a weak subsolution (weak supersolution) of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), if it satisfies

$$(2.9) \quad \langle u, \varphi \rangle_{H_p, K} + \int_{\mathbb{R}^n} V(x)|u(x)|^{p-2}u(x)\varphi(x) dx \leq 0 \quad (\geq 0)$$

for any nonnegative  $\varphi \in Y_0^{s,p}(\Omega)$ . Also, we say that a function  $u$  is a weak solution of the nonlocal equation (1.3), if it is both a weak subsolution and a weak supersolution, i.e.

$$(2.10) \quad \langle u, \varphi \rangle_{H_p, K} + \int_{\mathbb{R}^n} V(x)|u(x)|^{p-2}u(x)\varphi(x) dx = 0 \quad \text{for any } \varphi \in Y_0^{s,p}(\Omega).$$

To prove our results, we need a well-known lemma [GT] that is useful in applying the De Giorgi-Nash-Moser method.

**Lemma 2.3.** Let  $\{N_k\}_{k=0}^\infty \subset \mathbb{R}$  be a sequence of positive numbers such that

$$N_{k+1} \leq d_0 e_0^k N_k^{1+\eta} \quad \text{for all } k \in \mathbb{N} \cup \{0\}$$

where  $d_0, \eta > 0$  and  $e_0 > 1$ . If  $N_0 \leq d_0^{-1/\eta} e_0^{-1/\eta^2}$ , then we have that  $N_k \leq e_0^{-k/\eta} N_0$  for any  $k = 0, 1, \dots$  and moreover  $\lim_{k \rightarrow \infty} N_k = 0$ .

We need several elementary inequalities which are useful in proving Theorem 1.2 and Theorem 1.5.

**Lemma 2.4.** (a) If  $\alpha, \beta \in \mathbb{R}$  and  $A, B \geq 0$ , then we have the inequality

$$|\beta - \alpha|^{p-2}(\beta - \alpha)(\beta B^p - \alpha A^p) \geq -c_p (|\alpha| + |\beta|)^p |B - A|^p$$

for any  $p > 1$ , where  $c_p = \frac{(p-1)^{p-1}}{2}$ .

(b) If  $\alpha, \beta \in \mathbb{R}$  with  $\beta \geq \alpha$  and  $A, B \geq 0$ , then we have the inequality

$$(\beta - \alpha)^{p-1}(\beta B^p - \alpha A^p) \geq \frac{1}{4}(\beta - \alpha)^p(A^p + B^p) - d_p (|\alpha| + |\beta|)^p |B - A|^p$$

for any  $p > 1$ , where  $d_p = \frac{1}{2}[2(p-1)]^{p-1}$ .

(c) If  $A \geq B \geq 0$  and  $p > 1$ , then  $(A - B)^{p-1} \geq b_p A^{p-1} - B^{p-1}$  for  $p > 1$ , where  $b_p = \mathbb{1}_{(1,2]}(p) + 2^{-(p-1)} \mathbb{1}_{(2,\infty)}(p)$ .

*Proof.* (a) Without loss of generality, we may assume that  $\alpha \neq \beta$ ; otherwise, the inequality is trivial. By simple calculation, we have that

$$\begin{aligned} I &:= |\beta - \alpha|^{p-2}(\beta - \alpha)(\beta B^p - \alpha A^p) \\ &= |\beta - \alpha|^p B^p + |\beta - \alpha|^{p-2}(\beta - \alpha)\alpha(B^p - A^p) \\ &\geq |\beta - \alpha|^p B^p - |\beta - \alpha|^{p-1} |\alpha| |B^p - A^p|. \end{aligned}$$

Similarly, we have the following inequality

$$I \geq |\beta - \alpha|^p A^p - |\beta - \alpha|^{p-1} |\beta| |B^p - A^p|.$$

This yields that

$$(2.11) \quad 2I \geq |\beta - \alpha|^p(A^p + B^p) - |\beta - \alpha|^{p-1} (|\alpha| + |\beta|) |B^p - A^p|.$$

Without loss of generality assuming that  $B \geq A$ , then it follows that

$$(2.12) \quad \begin{aligned} B^p - A^p &= \int_A^B \frac{d}{dt}(t^p) dt = p \int_A^B t^{p-1} dt \\ &\leq p B^{p-1} (B - A) \leq p (A^p + B^p)^{\frac{p-1}{p}} (B - A). \end{aligned}$$

Thus, by (2.12) and Young's inequality with any  $\varepsilon > 0$ , we have that

$$(2.13) \quad \begin{aligned} |B^p - A^p| &\leq p (A^p + B^p)^{\frac{p-1}{p}} |B - A| \\ &\leq p\varepsilon (A^p + B^p) + \frac{1}{\left(\varepsilon \frac{p}{p-1}\right)^{p-1}} |B - A|^p. \end{aligned}$$

From (2.11) and (2.13), we have that

$$(2.14) \quad \begin{aligned} 2I &\geq |\beta - \alpha|^{p-1} (A^p + B^p) [|\beta - \alpha| - p\varepsilon (|\alpha| + |\beta|)] \\ &\quad - \frac{1}{\left(\varepsilon \frac{p}{p-1}\right)^{p-1}} |\beta - \alpha|^{p-1} (|\alpha| + |\beta|) |B - A|^p. \end{aligned}$$

Taking  $\varepsilon = \frac{|\beta - \alpha|}{p(|\alpha| + |\beta|)} > 0$  in (2.14), the required inequality can be obtained.

(b) Without loss of generality, we may assume that  $\alpha \neq \beta$ ; for, if not, the inequality is obvious. By simple calculation, we have that

$$\begin{aligned} J &:= (\beta - \alpha)^{p-1} (\beta B^p - \alpha A^p) \\ &= (\beta - \alpha)^p B^p + (\beta - \alpha)^{p-1} \alpha (B^p - A^p) \\ &\geq (\beta - \alpha)^p B^p - (\beta - \alpha)^{p-1} |\alpha| |B^p - A^p|. \end{aligned}$$

Similarly, we have the following inequality

$$J \geq (\beta - \alpha)^p A^p - (\beta - \alpha)^{p-1} |\beta| |B^p - A^p|.$$

This yields that

$$(2.15) \quad 2J \geq (\beta - \alpha)^p (A^p + B^p) - (\beta - \alpha)^{p-1} (|\alpha| + |\beta|) |B^p - A^p|.$$

Assuming  $B \geq A$  without loss of generality, then it follows that

$$(2.16) \quad \begin{aligned} B^p - A^p &= \int_A^B \frac{d}{dt}(t^p) dt = p \int_A^B t^{p-1} dt \\ &\leq p B^{p-1} (B - A) \leq p (A^p + B^p)^{\frac{p-1}{p}} (B - A). \end{aligned}$$

Thus, by (2.16) and Young's inequality with any  $\varepsilon > 0$ , we have that

$$(2.17) \quad \begin{aligned} |B^p - A^p| &\leq p (A^p + B^p)^{\frac{p-1}{p}} |B - A| \\ &\leq p\varepsilon (A^p + B^p) + \frac{1}{\left(\varepsilon \frac{p}{p-1}\right)^{p-1}} |B - A|^p. \end{aligned}$$

From (2.15) and (2.17), we have that

$$(2.18) \quad \begin{aligned} 2J &\geq (\beta - \alpha)^{p-1}(A^p + B^p) [(\beta - \alpha) - p\varepsilon(|\alpha| + |\beta|)] \\ &\quad - \frac{1}{\left(\varepsilon \frac{p}{p-1}\right)^{p-1}} (\beta - \alpha)^{p-1} (|\alpha| + |\beta|) |B - A|^p. \end{aligned}$$

Taking  $\varepsilon = \frac{\beta - \alpha}{2p(|\alpha| + |\beta|)} > 0$  in (2.18), the required inequality can be obtained.

(c) If  $a, b \geq 0$ , then it easily follows from the fact that  $(a + b)^{p-1} \leq a^{p-1} + b^{p-1}$  for  $1 < p \leq 2$  and  $(a + b)^{p-1} \leq 2^{p-1}(a^{p-1} + b^{p-1})$  for  $p > 2$ . Set  $a = A - B$  and  $b = B$ .  $\square$

**Lemma 2.5.** ([DKP1]) *If  $p \geq 1$ ,  $\varepsilon \in (0, 1]$  and  $a, b \geq 0$ , then the inequality holds*

$$a^p \leq b^p + c_p \varepsilon b^p + (1 + c_p \varepsilon) \varepsilon^{1-p} |a - b|^p,$$

where  $c_p = (p - 1)\Gamma(1 \vee (p - 2))$  for the standard Gamma function  $\Gamma$ .

### 3. WEIGHTED NORM INEQUALITIES ON THE $A_p$ -MUCKENHOUPT CLASS

In this section, we briefly introduce the  $A_p$ -Muckenhoupt class for  $p \geq 1$  and we prove that any potential  $V$  in  $\mathcal{P}_q^{s,p}(\mathbb{R}^n)$  ( $q > \frac{n}{ps} > 1$ ,  $p > 1$ ,  $0 < s < 1$ ) satisfies certain weighted norm inequalities related with  $(V_-, V_+)$ .

By a *weight* on  $\mathbb{R}^n$  equipped with Lebesgue measure, we mean a locally integrable function on  $\mathbb{R}^n$  taking value in  $[0, \infty)$  almost everywhere. For  $f \in L_{loc}^1(\mathbb{R}^n)$  and  $x \in \mathbb{R}^n$ , the Hardy-Littlewood maximal function  $\mathcal{M}f$  is defined by

$$\mathcal{M}f(x) = \sup_{Q^x} \int_{Q^x} |f(y)| dy,$$

where the supremum is taken over all cubes  $Q^x$  with center  $x$ . For a pair of weights  $(v, \omega)$ , the quantity  $[v, \omega]_{(A_p, A_p)}$  is defined by

$$[v, \omega]_{(A_p, A_p)} = \begin{cases} \sup_Q \left( \int_Q v(y) dy \right) \left( \int_Q \omega(y)^{-\frac{1}{p-1}} dy \right)^{p-1}, & 1 < p < \infty, \\ \sup_Q \left( \int_Q v(y) dy \right) \|\omega^{-1}\|_{L^\infty(Q)}, & p = 1, \end{cases}$$

where the supremum is taken over all cubes  $Q \subset \mathbb{R}^n$ . For  $p \in [1, \infty)$ , we say that  $(v, \omega) \in (A_p, A_p)$  if  $[v, \omega]_{(A_p, A_p)} < \infty$ , and  $\omega \in A_p$  if  $[\omega]_{A_p} := [\omega, \omega]_{(A_p, A_p)} < \infty$ . For  $1 \leq p < \infty$ , it is well-known in [Gr] that  $(A_1, A_1) \subset (A_p, A_p)$ ,  $A_1 \subset A_p$  and  $(v, \omega) \in (A_p, A_p)$  is equivalent to the mapping property that  $\mathcal{M} : L_\omega^p(\mathbb{R}^n) \rightarrow L_v^{p,\infty}(\mathbb{R}^n)$  is bounded; that is, there is a universal constant  $C_{n,p} > 0$  such that

$$(3.1) \quad \sup_{t>0} [t v(\{x \in \mathbb{R}^n : \mathcal{M}f(x) > t\})^{\frac{1}{p}}] \leq C_{n,p} \left( \int_{\mathbb{R}^n} |f(y)|^p \omega(y) dy \right)^{\frac{1}{p}}$$

for any  $f \in L_\omega^p(\mathbb{R}^n)$ . Here, we denote by  $v(E) = \int_E v(y) dy$  for a set  $E \subset \mathbb{R}^n$ . The reader can also refer to [Gr] for these concepts in Fourier analysis.

We shall now furnish several examples in the class  $\mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$ ,  $0 < s < 1$ ) mentioned in the above introduction:

(a) If  $h_\eta(x) = |x|^\eta$  for  $\alpha \in \mathbb{R}$ , then it is easy to check that

$$[h_\eta]_{A_1} < \infty \text{ if and only if } -n < \eta \leq 0.$$

If we consider a sign-changing potential

$$V(x) = |x|^{\eta/q} \cos(|x|)$$

with  $\eta \in (-n, 0]$  and  $q > \frac{n}{ps} > 1$  ( $p > 1$ ,  $0 < s < 1$ ), then we can easily check that  $V_+ \in L_{\text{loc}}^q(\mathbb{R}^n)$ ,  $V_- \in L_{\text{loc}}^1(\mathbb{R}^n)$ ,  $V_+/b \in L_{\text{loc}}^q(\mathbb{R}^n)$ ,  $(V_-, V_+^{b,i}) \in (A_1, A_1)$  and  $V_+^{b,i} \in A_1$  for all  $i \in \mathbb{N}$ , where  $b(x) = \cos^+(|x|)$ .

(b) Let  $\nu$  be a Borel measure on  $\mathbb{R}^n$  satisfying that

$$\mathfrak{M}\nu(x) := \sup_{Q^x} \frac{\nu(Q^x)}{|Q^x|} \leq C \text{ for a.e. } x \in \mathbb{R}^n,$$

where the supremum is taken over all cubes  $Q^x$  with center  $x \in \mathbb{R}^n$ . If we set  $g_\gamma(x) = [\mathfrak{M}\nu(x)]^\gamma$  for  $\gamma \in (0, 1)$ , then it is known in [GR] that  $[g_\gamma]_{A_1} < \infty$ . We consider the following sign-changing potential

$$V(x) = g_\gamma(x) \sin(1/|x|)$$

for  $\gamma \in (0, 1)$ . If  $q > \frac{n}{ps} > 1$  for  $p > 1$  and  $0 < s < 1$ , then it is easy to check that  $V_+ \in L_{\text{loc}}^q(\mathbb{R}^n)$ ,  $V_- \in L_{\text{loc}}^1(\mathbb{R}^n)$ ,  $V_+/b \in L_{\text{loc}}^q(\mathbb{R}^n)$ ,  $(V_-, V_+^{b,i}) \in (A_1, A_1)$  and  $V_+^{b,i} \in A_1$  for all  $i \in \mathbb{N}$ , where  $b(x) = \sin^+(1/|x|)$ .

(c) Let  $\omega(x) = \ln(1/|x|)\chi_{B(0;1/e)} + \chi_{\mathbb{R}^n \setminus B(0;1/e)}$ . Then it is easy to check that  $\omega \in A_1$ . We consider the following sign-changing potential

$$V(x) = \left[ \ln(1/|x|)\chi_{B(0;1/e)} + \chi_{\mathbb{R}^n \setminus B(0;1/e)} \right] \cos(1/|x|).$$

From properties of the Gamma function, if  $q > \frac{n}{ps} > 1$  for  $p > 1$  and  $0 < s < 1$  then we can easily check that  $V_+ \in L_{\text{loc}}^q(\mathbb{R}^n)$ ,  $V_- \in L_{\text{loc}}^1(\mathbb{R}^n)$ ,  $V_+/b \in L_{\text{loc}}^q(\mathbb{R}^n)$ ,  $(V_-, V_+^{b,i}) \in (A_1, A_1)$  and  $V_+^{b,i} \in A_1$  for all  $i \in \mathbb{N}$ , where  $b(x) = \cos^+(1/|x|)$ .

Next, we need several fundamental lemmas which is useful in proving Theorem 3.3.

**Lemma 3.1.** *If  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$ ,  $p > 1$  and  $0 < s < 1$ , then*

$$\int_{\mathbb{R}^n} |\varphi(y)|^p V_-(y) dy \leq \int_{\mathbb{R}^n} |\varphi(y)|^p V_+(y) dy$$

for any  $\varphi \in Y_0^{s,p}(\Omega)$ .

*Proof.* From the exactly same way as the proof of Theorem 3.6 [K2], we see that

$$(3.2) \quad \int_{\mathbb{R}^n} |\varphi(y)|^p V(y) dy \geq 0$$

for any  $\varphi \in C_c^\infty(\mathbb{R}^n)$ . Take any  $\varphi \in Y_0^{s,p}(\Omega)$ . Since  $C_c^\infty(\Omega)$  is dense in  $X_0^{s,p}(\Omega)$  (see [FSV] and Theorem 1.4.2.2 in [G]), we can take a sequence  $\{\varphi_k\} \subset C_c^\infty(\Omega)$  such that

$$\varphi_k \rightarrow \varphi \text{ in } X_0^{s,p}(\Omega).$$

Since  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$ , there is a nonnegative bounded function  $b$  on  $\mathbb{R}^n$  such that

$$(3.3) \quad V_+^b := \frac{V_+}{b} \in L_{\text{loc}}^q(\mathbb{R}^n), (V_-, V_+^{b,i}) \in (A_1, A_1) \text{ and } V_+^{b,i} \in A_1$$

for all  $i \in \mathbb{N}$ . Then we claim that

$$(3.4) \quad \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_k(y)|^p V_+(y) dy = \int_{\mathbb{R}^n} |\varphi(y)|^p V_+(y) dy;$$

indeed, we note that  $\|\varphi_k\|_{X_0^{s,p}(\Omega)} \leq 2\|\varphi\|_{X_0^{s,p}(\Omega)}$  for all sufficiently large  $k \in \mathbb{N}$ , and also we see that, for any  $y \in \mathbb{R}^n$ ,

$$\begin{aligned} |\varphi_k(y)|^p - |\varphi(y)|^p &= \int_{|\varphi(y)|}^{|\varphi_k(y)|} \frac{d}{d\tau} \tau^p d\tau \\ &\leq p \left( |\varphi_k(y)| - |\varphi(y)| \right) \left( |\varphi_k(y)| \vee |\varphi(y)| \right)^{p-1}. \end{aligned}$$

Thus it follows from the fractional Sobolev inequality and Hölder's inequality that

$$\begin{aligned} (3.5) \quad & \left| \int_{\mathbb{R}^n} |\varphi_k(y)|^p V_+(y) dy - \int_{\mathbb{R}^n} |\varphi(y)|^p V_+(y) dy \right| \\ & \leq \int_{\mathbb{R}^n} \left| |\varphi_k(y)|^p - |\varphi(y)|^p \right| V_+(y) dy \\ & \leq p \left( \int_{\Omega} \left| |\varphi_k(y)| - |\varphi(y)| \right|^p V_+(y) dy \right)^{\frac{1}{p}} \\ & \quad \times \left( \int_{\Omega} \left( |\varphi_k(y)| \vee |\varphi(y)| \right)^p V_+(y) dy \right)^{\frac{p-1}{p}} \\ & \lesssim \|\varphi\|_{X_0^{s,p}(\Omega)} \|\varphi_k - \varphi\|_{X_0^{s,p}(\Omega)} \|V_+\|_{L^q(\Omega)} \rightarrow 0 \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Also, we claim that

$$(3.6) \quad \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_k(y)|^p V_-(y) dy = \int_{\mathbb{R}^n} |\varphi(y)|^p V_-(y) dy;$$

indeed, we have that

$$V_-(y) \leq [V_-, V_+^{b,1}]_{(A_1, A_1)} V_+^{b,1}(y) \quad \text{a.e. } y \in \mathbb{R}^n,$$

because  $(V_-, V_+^{b,1}) \in (A_1, A_1)$  by (3.3). Since  $V_+^b \in L_{\text{loc}}^q(\mathbb{R}^n)$  by (3.3), as in (3.5) we obtain that

$$\begin{aligned} & \left| \int_{\mathbb{R}^n} |\varphi_k(y)|^p V_-(y) dy - \int_{\mathbb{R}^n} |\varphi(y)|^p V_-(y) dy \right| \\ & \leq \int_{\mathbb{R}^n} \left| |\varphi_k(y)|^p - |\varphi(y)|^p \right| V_-(y) dy \\ & \leq \left( \|b\|_{L^\infty(\mathbb{R}^n)} \vee 1 \right) [V_-, V_+^{b,1}]_{(A_1, A_1)} \\ & \quad \times \int_{\Omega} \left| |\varphi_k(y)|^p - |\varphi(y)|^p \right| V_+^b(y) dy \rightarrow 0 \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Thus, by (3.2), (3.4) and (3.6), we obtain that

$$\begin{aligned} \int_{\mathbb{R}^n} |\varphi(y)|^p V_-(y) dy &= \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_k(y)|^p V_-(y) dy \\ &\leq \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_k(y)|^p V_+(y) dy = \int_{\mathbb{R}^n} |\varphi(y)|^p V_+(y) dy. \end{aligned}$$

Hence we are done.  $\square$

**Lemma 3.2.** *If  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$ ,  $0 < s < 1$ ), then  $Y_0^{s,p}(\Omega)$  is a quasi-Banach space with the quasinorm  $\|\cdot\|_{Y_0^{s,p}(\Omega)}$  given by*

$$\|u\|_{Y_0^{s,p}(\Omega)}^p := \|u\|_{X_0^{s,p}(\Omega)}^p + \int_{\Omega} |u(y)|^p V(y) dy, \quad u \in Y_0^{s,p}(\Omega).$$

Moreover,  $Y_0^{s,p}(\Omega) = X_0^{s,p}(\Omega)$  and they are quasinorm-equivalent.

*Proof.* It follows from Lemma 3.1 and (2.5) that

$$\begin{aligned} \|u\|_{X_0^{s,p}(\Omega)}^p &\leq \|u\|_{Y_0^{s,p}(\Omega)}^p \leq \|u\|_{X_0^{s,p}(\Omega)}^p + \|u\|_{L_{V_+}^p(\Omega)}^p \\ &\leq (1 + |\Omega|^{\frac{ps}{n} - \frac{1}{q}}) \|u\|_{X_0^{s,p}(\Omega)}^p. \end{aligned}$$

Since  $X_0^{s,p}(\Omega)$  is a Banach space, this implies the required results.  $\square$

**Theorem 3.3.** *If  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  and  $0 < s < 1$ , then*

$$(3.7) \quad \int_{\mathbb{R}^n} |\varphi(y)| V_-(y) dy \leq \int_{\mathbb{R}^n} |\varphi(y)| V_+(y) dy$$

for any  $\varphi \in Y_0^{s,p}(\Omega)$ .

*Proof.* Take any  $\varphi \in Y_0^{s,p}(\Omega)$ . Since  $C_c^\infty(\Omega)$  is dense in  $X_0^{s,p}(\Omega)$  (see [G] and [FSV]),  $Y_0^{s,p}(\Omega) = X_0^{s,p}(\Omega)$  and they are quasinorm-equivalent by Lemma 3.2, we can take a sequence  $\{\varphi_i\} \subset C_c^\infty(\Omega)$  such that  $\varphi_i \rightarrow \varphi$  in  $Y_0^{s,p}(\Omega)$ . So by (2.5) we also have that  $\varphi_i \rightarrow \varphi$  in  $L^{p_*}(\Omega)$ , where  $p_* = \frac{pn}{n-ps}$ . So we can choose a subsequence  $\{\varphi_{i_k}\}$  such that

$$(3.8) \quad \varphi_{i_k} \rightarrow \varphi \text{ a.e. in } \Omega.$$

Also we have that

$$(3.9) \quad \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_{i_k}(y)| V_+(y) dy = \int_{\mathbb{R}^n} |\varphi(y)| V_+(y) dy;$$

indeed, it follows from the fractional Sobolev inequality and Hölder's inequality that

$$\begin{aligned} &\left| \int_{\mathbb{R}^n} |\varphi_{i_k}(y)| V_+(y) dy - \int_{\mathbb{R}^n} |\varphi(y)| V_+(y) dy \right| \\ &\leq \int_{\Omega} \left| |\varphi_{i_k}(y)| - |\varphi(y)| \right| V_+(y) dy \\ &\leq \left( \int_{\Omega} \left| |\varphi_{i_k}(y)| - |\varphi(y)| \right|^p V_+(y) dy \right)^{\frac{1}{p}} \left( \int_{\Omega} V_+(y) dy \right)^{\frac{p-1}{p}} \\ &\leq |\Omega|^{\frac{s}{n} + \frac{1}{q'} - \frac{1}{p}} \|\varphi_{i_k} - \varphi\|_{Y_0^{s,p}(\Omega)} \|V_+\|_{L^q(\Omega)} \rightarrow 0 \text{ as } k \rightarrow \infty \end{aligned}$$

where  $q'$  is the dual exponent of  $q$ . Hence, by Fatou's lemma, Lemma 3.1, Lemma 3.5 in [K2], (3.2), (3.4), (3.8) and (3.9), we conclude that

$$\begin{aligned} \int_{\mathbb{R}^n} |\varphi(y)| V_-(y) dy &= \lim_{p \rightarrow 1^+} \int_{\mathbb{R}^n} |\varphi(y)|^p V_-(y) dy \\ &\leq \lim_{p \rightarrow 1^+} \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_{i_k}(y)|^p V_-(y) dy \\ &\leq \lim_{p \rightarrow 1^+} \liminf_{k \rightarrow \infty} \int_{\mathbb{R}^n} |\varphi_{i_k}(y)|^p V_+(y) dy \\ &\leq \lim_{p \rightarrow 1^+} \int_{\mathbb{R}^n} |\varphi(y)|^p V_+(y) dy \\ &= \int_{\mathbb{R}^n} |\varphi(y)| V_+(y) dy. \end{aligned}$$

Therefore we complete the proof.  $\square$

#### 4. LOCAL PROPERTIES OF WEAK SUBSOLUTIONS

In this section, we shall obtain certain local properties for weak subsolutions of the nonlocal  $p$ -Laplacian type Schrödinger equation. These results play a crucial role in establishing a nonlocal Hölder continuity for weak solutions of the nonlocal equation (1.3). In order to establish the result, we need several steps.

**Lemma 4.1.** *For  $p > 1$  and  $M > 0$ , let  $h(t) = t^{p-1}(t - M)_+ - (t - M)_+^p \geq 0$ . Then  $h$  is Lipschitz continuous on  $\mathbb{R}$ .*

*Proof.* We note that  $h(t) = 0$  for  $t < M$ ,  $h$  is in  $C^1(M, \infty)$ ,

$$\lim_{t \rightarrow M^+} \frac{h(t) - h(M)}{t - M} = 0 \quad \text{and} \quad \lim_{t \rightarrow M^-} \frac{h(t) - h(M)}{t - M} = M^{p-1}.$$

Also it is easy to check that  $\lim_{t \rightarrow \infty} h(t) = 0$ , because

$$\lim_{t \rightarrow \infty} \frac{t^{p-1}(t - M)_+}{(t - M)_+^p} = 1.$$

Moreover, in order to check the differentiability of  $h$  at the infinity, set  $g(t) = h(1/t)$ . Then we have that

$$\lim_{t \rightarrow 0^+} \frac{g(t) - g(0)}{t} = \lim_{t \rightarrow 0^+} \left[ \frac{1}{t^p} \left( \frac{1}{t} - M \right) - \frac{1}{t} \left( \frac{1}{t} - M \right)^p \right] = 0,$$

because

$$\lim_{t \rightarrow 0^+} \left[ \frac{1}{t^p} \left( \frac{1}{t} - M \right) / \frac{1}{t} \left( \frac{1}{t} - M \right)^p \right] = \lim_{t \rightarrow 0^+} \frac{1 - tM}{(1 - tM)^p} = 1.$$

Thus this implies the required result.  $\square$

**Corollary 4.2.** *If  $M > 0$  and  $u \in X^{s,p}(\Omega)$  for  $p > 1$  and  $0 < s < 1$ , then*

- (a)  $|u|^{p-2}u(u - M)_+ - (u - M)_+^p \in X^{s,p}(\Omega)$  and it is nonnegative in  $\mathbb{R}^n$ , and
- (b)  $\left[ |u|^{p-2}u(u - M)_+ - (u - M)_+^p \right] \zeta^p \in X_0^{s,p}(\Omega)$  for any nonnegative  $\zeta \in C_c^\infty(\Omega)$ .

*Proof.* (a) Note that  $|u|^{p-2}u(u - M)_+ - (u - M)_+^p = h \circ u$  for the function  $h$  given in Lemma 4.1. Since  $h$  is Lipschitz continuous on  $\mathbb{R}$  by Lemma 4.1, it is obvious that  $0 \leq h \circ u \in X^{s,p}(\Omega)$ .

(b) By the mean value theorem, we have that

$$\begin{aligned} |\zeta^p(x) - \zeta^p(y)| &= \left| \int_{\zeta(y)}^{\zeta(x)} \frac{d}{d\tau} \tau^p d\tau \right| \leq p(\zeta^{p-1}(x) \vee \zeta^{p-1}(y)) |\zeta(x) - \zeta(y)| \\ &\leq 2p \|\zeta\|_{L^\infty(\mathbb{R}^n)}^{p-1} |\zeta(x) - \zeta(y)| \end{aligned}$$

for all  $x, y \in \mathbb{R}^n$ . This means that  $\zeta^p$  is Lipschitz continuous on  $\mathbb{R}^n$ . So we can easily derive the result. Hence we complete the proof.  $\square$

Next we need the following *nonlocal Caccioppoli type inequality* which is Theorem 1.5. This is a very useful tool in proving an interior Hölder regularity of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation.

**[Proof of Theorem 1.5.]** For simplicity of the proof, we assume that  $x_0 = 0$ . Let  $w = (u - M)_+$  for  $M \in [0, \infty)$  and take any nonnegative  $\zeta \in C_c^\infty(B_r)$ . We use  $\varphi = w\zeta^p$  as a testing function in the weak formulation of the equation. Then we have that

$$(4.1) \quad \langle u, \varphi \rangle_{H_p, K} + \int_{\mathbb{R}^n} |u(y)|^{p-2} u(y) \varphi(y) V(y) dy \leq 0,$$

where  $d_K(x, y) = K(x - y) dx dy$  and

$$\langle u, \varphi \rangle_{H_p, K} = \iint_{\mathbb{R}^n \times \mathbb{R}^n} H_p(u(x) - u(y)) (\varphi(x) - \varphi(y)) d_K(x, y) \text{ for } p \geq 1.$$

The first term in the left-hand side of the above inequality can be decomposed into two parts as follows;

$$(4.2) \quad \begin{aligned} \langle u, \varphi \rangle_{H_p, K} &= \iint_{B_r \times B_r} H_p(u(x) - u(y)) (\varphi(x) - \varphi(y)) d_K(x, y) \\ &\quad + 2 \iint_{(\mathbb{R}^n \setminus B_r) \times B_r} H_p(u(x) - u(y)) \varphi(x) d_K(x, y) \\ &:= I_1 + 2I_2. \end{aligned}$$

For the estimate of  $I_1$ , without loss of generality we may assume that  $u(x) \geq u(y)$ . Then we first observe that  $w(x) \geq w(y)$  and

$$(4.3) \quad H_p(u(x) - u(y)) (\varphi(x) - \varphi(y)) \geq (w(x) - w(y))^{p-1} (\varphi(x) - \varphi(y))$$

whenever  $x, y \in B_r$ ; indeed, it can easily be checked by considering three possible occasions (i)  $u(x), u(y) > M$ , (ii)  $u(x) > M, u(y) \leq M$ , and (iii)  $u(y) \leq u(x) \leq M$ . Also we observe that

$$(4.4) \quad \begin{aligned} |\zeta(x)w(x) - \zeta(y)w(y)|^p &\leq 2^{p-1} |w(x) - w(y)|^p (\zeta^p(x) + \zeta^p(y)) \\ &\quad + 2^{p-1} (w^p(x) + w^p(y)) |\zeta(y) - \zeta(x)|^p. \end{aligned}$$

By (b) of Lemma 2.4, (4.3) and (4.4), we have that

$$(4.5) \quad \begin{aligned} &H_p(u(x) - u(y)) (\varphi(x) - \varphi(y)) \\ &\geq \frac{1}{4} |w(x) - w(y)|^p (\zeta^p(x) + \zeta^p(y)) \\ &\quad - d_p (w^p(x) + w^p(y)) |\zeta(y) - \zeta(x)|^p \\ &\geq \frac{1}{2^{p+1}} |\zeta(x)w(x) - \zeta(y)w(y)|^p \\ &\quad - \left(\frac{1}{4} + d_p\right) (w^p(x) + w^p(y)) |\zeta(y) - \zeta(x)|^p. \end{aligned}$$

Thus it follows from (4.5) that

$$(4.6) \quad \begin{aligned} I_1 &\geq \frac{1}{2^{p+1}} \iint_{B_r \times B_r} |\zeta(x)w(x) - \zeta(y)w(y)|^p d_K(x, y) \\ &\quad - 2^p \left(\frac{1}{4} + d_p\right) \iint_{B_r \times B_r} (w^p(x) + w^p(y)) |\zeta(y) - \zeta(x)|^p d_K(x, y). \end{aligned}$$



For the estimate of  $I_2$ , we note that

$$\begin{aligned} H_p(u(x) - u(y))\varphi(x) &\geq -(u(y) - u(x))_+^{p-1}(u(x) - M)_+\zeta^p(x) \\ &\geq -(u(y) - M)_+^{p-1}(u(x) - M)_+\zeta^p(x) \\ &= -w^{p-1}(y)w(x)\zeta^p(x) \end{aligned}$$

and thus we have that

$$\begin{aligned} (4.7) \quad I_2 &\geq - \iint_{(\mathbb{R}^n \setminus B_r) \times B_r} w^{p-1}(y)w(x)\zeta^p(x) d_K(x, y) \\ &\geq - \left( \sup_{x \in \text{supp}(\zeta)} \int_{\mathbb{R}^n \setminus B_r} w^{p-1}(y)K(x-y) dy \right) \int_{B_r} w(x)\zeta^p(x) dx. \end{aligned}$$

Finally, we claim that

$$(4.8) \quad \int_{\mathbb{R}^n} |u(y)|^{p-2}u(y)\varphi(y)V(y) dy \geq \int_{\mathbb{R}^n} w^p(y)\zeta^p(y)V(y) dy.$$

This is equivalent to the following inequality

$$\begin{aligned} &\int_{\mathbb{R}^n} \left[ |u(y)|^{p-2}u(y)w(y) - w^p(y) \right] \zeta^p(y)V_+(y) dy \\ &\geq \int_{\mathbb{R}^n} \left[ |u(y)|^{p-2}u(y)w(y) - w^p(y) \right] \zeta^p(y)V_-(y) dy, \end{aligned}$$

whose proof is just a direct application of Lemma 3.2, Theorem 3.3 and Corollary 4.2. Hence the required inequality can be obtained from (4.1), (4.2), (4.6), (4.7) and (4.8).  $\square$

Next, we shall obtain the local boundedness of such weak subsolutions which is Theorem 1.1 and a relation between the nonlocal tail terms of the positive part and the negative part of weak solutions of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3) in the following theorems.

**[Proof of Theorem 1.1.]** Take any  $\zeta \in C_c^\infty(B_r^0)$  such that  $|\nabla\zeta| \leq c/r$  on  $\mathbb{R}^n$ . Let  $w = (u - M)_+$  for  $M \in (0, \infty)$ . By Lemma 3.2, Theorem 1.6 and the mean value theorem, we have that

$$(4.9) \quad \begin{aligned} &\iint_{B_r^0 \times B_r^0} |\zeta(x)w(x) - \zeta(y)w(y)|^p d_K(x, y) \\ &\lesssim r^{p-ps} \|\nabla\zeta\|_{L^\infty(B_r^0)}^p \|w\|_{L^p(B_r^0)}^p + \mathcal{A}(w, \zeta, r, s) \|w\|_{L^1(B_r^0)} \end{aligned}$$

where  $\mathcal{A}(w, \zeta, r, s)$  is the value given by

$$\mathcal{A}(w, \zeta, r, s) = \sup_{x \in \text{supp}(\zeta)} \int_{\mathbb{R}^n \setminus B_r^0} w^{p-1}(y)K(x-y) dy.$$

Applying Proposition 2.1 to (4.9), we obtain that

$$(4.10) \quad \begin{aligned} \left( \int_{B_r^0} |w\zeta|^{p\gamma} dx \right)^{\frac{1}{\gamma}} &\lesssim \left( r^{p-ps} \|\nabla\zeta\|_{L^\infty(B_r^0)}^p + r^{-ps} \right) r^{ps} \int_{B_r^0} |w|^p dx \\ &\quad + \mathcal{A}(w, \zeta, r, s) r^{ps} \int_{B_r^0} w dx \end{aligned}$$

where  $\gamma = \frac{n}{n-ps} > 1$ . For  $k = 0, 1, 2, \dots$ , we set

$$r_k = (1 + 2^{-k})r, \quad r_k^* = \frac{r_k + r_{k+1}}{2},$$

$$M_k = M + (1 - 2^{-k})M_*, \quad M_k^* = \frac{M_k + M_{k+1}}{2},$$

$w_k = (u - M_k)_+$  and  $w_k^* = (u - M_k^*)_+$  for a constant  $M_* > 0$  to be determined later. In (4.9), for  $k = 0, 1, \dots$ , we choose a function  $\zeta_k \in C_c^\infty(B_{r_k}^0)$  with  $\zeta_k|_{B_{r_{k+1}}^0} \equiv 1$  such that  $0 \leq \zeta_k \leq 1$  and

$$|\nabla \zeta_k| \leq c 2^{k+2}/r \text{ in } \mathbb{R}^n.$$

For  $k = 0, 1, 2, \dots$ , we set

$$N_k = \left( \int_{B_{r_k}^0} |w_k|^p dx \right)^{\frac{1}{p}}.$$

Since  $w_k^* \geq w_{k+1}$  and

$$w_k^*(x) \geq M_{k+1} - M_k^* = 2^{-k-2}M_*$$

whenever  $u(x) \geq M_{k+1}$ , we then have that

$$(4.11) \quad \begin{aligned} N_{k+1} &\leq \left( \frac{1}{|B_{r_{k+1}}^0|} \int_{B_{r_{k+1}}^0} \frac{w_{k+1}^p (w_k^*)^{p(\gamma-1)}}{(M_{k+1} - M_k^*)^{p(\gamma-1)}} dx \right)^{\frac{1}{p}} \\ &\lesssim \left( \frac{2^k}{M_*} \right)^{\gamma-1} \left( \int_{B_{r_k}^0} |w_k^* \zeta_k|^{p\gamma} dx \right)^{\frac{1}{p}}. \end{aligned}$$

Since  $\zeta_k \in C_c^\infty(B_{r_k}^0)$ ,  $w_k^* \leq w_0$  for all  $k$  and

$$|y - x| \geq |y - x_0| - |x - x_0| \geq \left(1 - \frac{r_k^*}{r}\right) |y - x_0| \geq 2^{-k-2} |y - x_0|$$

for any  $x \in B_{r_k^*}^0$  and  $y \in \mathbb{R}^n \setminus B_{r_k^*}^0$ , we easily obtain that

$$(4.12) \quad \mathcal{A}(w_k^*, \zeta_k, r_k, s) \leq c 2^{k(n+ps)} r^{-ps} [\mathcal{T}_r(w_0; x_0)]^{p-1}.$$

Since  $0 \leq w_k^* \leq w_k$  and  $w_k(x) \geq M_k^* - M_k = 2^{-k-2}M_*$  if  $u(x) \geq M_k^*$ , it follows from (4.10), (4.11) and (4.12) that

$$\begin{aligned} \left( \frac{2^k}{M_*} \right)^{-\frac{p(\gamma-1)}{\gamma}} N_{k+1}^{\frac{p}{\gamma}} &\leq c 2^{pk} \int_{B_{r_k}^0} |w_k^*|^p dx \\ &\quad + c 2^{k(n+ps)} \left( \frac{r_k}{r} \right)^{ps} [\mathcal{T}_r(w_0; x_0)]^{p-1} \int_{B_{r_k}^0} w_k^* dx \\ &\leq c 2^{pk} N_k^p + c [\mathcal{T}_r(w_0; x_0)]^{p-1} 2^{k(n+ps)} \int_{B_{r_k}^0} \frac{w_k^* w_k^{p-1}}{(M_k^* - M_k)^{p-1}} dx \\ &\leq c \left( 2^{pk} + 2^{k(n+ps)} \left( \frac{2^k}{M_*} \right)^{p-1} [\mathcal{T}_r(w_0; x_0)]^{p-1} \right) N_k^p. \end{aligned}$$

Taking  $M^*$  in the above so that

$$M_* \geq \delta \mathcal{T}_r(w_0; x_0) \text{ for } \delta \in (0, 1],$$

we obtain that

$$\frac{N_{k+1}}{M_*} \leq d_0 a^k \left( \frac{N_k}{M_*} \right)^{1+\eta}$$

where  $d_0 = c^{\frac{\gamma}{p}} \delta^{-\frac{p-1}{p}\gamma} > 0$ ,  $a = 2^{\frac{\gamma}{p}(n+2s+p-1) + \frac{ps}{n-ps}} > 1$  and  $\eta = \gamma - 1 > 0$ .

If  $N_0 \leq d_0^{-\frac{1}{\eta}} a^{-\frac{1}{\eta^2}} M_*$ , then we set

$$M_* = \delta \mathcal{T}_r(w_0; x_0) + c_0 \delta^{-\frac{(p-1)n}{sp^2}} a^{\frac{(n-ps)^2}{p^2s^2}} N_0$$

where  $c_0 = c^{\frac{n}{sp^2}}$ . By Lemma 2.3, we conclude that

$$\begin{aligned} \sup_{B_r^0} u &\leq M + M_* \\ &\leq M + \delta \mathcal{T}_r(w_0; x_0) + c_0 \delta^{-\frac{(p-1)n}{sp^2}} a^{\frac{(n-ps)^2}{p^2s^2}} \left( \int_{B_{2r}^0} (u - M)_+^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

Hence, taking  $M \downarrow 0$  in the above estimate, we obtain the required result.  $\square$

The third one is a lemma which furnishes a relation between the nonlocal tails of the positive and negative part of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation.

**Lemma 4.3.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$ ,  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1$ ,  $s \in (0, 1)$ ). If  $u \in Y_g^{s,p}(\Omega)$  is a weak solution of the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3) such that  $u \geq 0$  in  $B_R^0 \subset \Omega$ , then we have the estimate*

$$\mathcal{T}_r(u_+; x_0) \lesssim \left( 1 + \|V_+\|_{L^q(\Omega)} \right) \sup_{B_r^0} u + \left( \frac{r}{R} \right)^{\frac{ps}{p-1}} \mathcal{T}_R(u_-; x_0)$$

for any  $r \in (0, R)$ .

*Proof.* Without loss of generality, we may assume that  $x_0 = 0$ . Let  $M = \sup_{B_r} u$  and  $\varphi(x) = w(x)\zeta^p(x)$  where  $w(x) = u(x) - 2M$  and  $\zeta \in C_c^\infty(B_{3r/4})$  is a function satisfying that  $\zeta|_{B_{r/2}} \equiv 1$ ,  $0 \leq \zeta \leq 1$  and  $|\nabla \zeta| \leq c/r$  in  $\mathbb{R}^n$ . Then we have that

$$\begin{aligned} 0 &= \iint_{B_r \times B_r} H_p(u(x) - u(y))(\varphi(x) - \varphi(y)) d_K(x, y) \\ &\quad + 2 \int_{\mathbb{R}^n \setminus B_r} \int_{B_r} H_p(u(x) - u(y))(u(x) - 2M)\zeta^p(x) d_K(x, y) \\ &\quad + \int_{\mathbb{R}^n} V(x)|u(x)|^{p-2}u(x)(u(x) - 2M)\zeta^p(x) dx \\ &:= J_1 + J_2 + J_3. \end{aligned} \tag{4.13}$$

Since the fact that

$$-2M \leq w(x) := u(x) - 2M \leq -M \tag{4.14}$$

for any  $x \in B_r$ , by (a) of Lemma 2.4 we have that

$$H_p(w(x) - w(y))(w(x)\zeta^p(x) - w(y)\zeta^p(y)) \geq -c_p 4^p M^p (\zeta(x) - \zeta(y))^p$$

for any  $x, y \in B_r$ , it follows from simple calculation that

$$(4.15) \quad \begin{aligned} J_1 &\geq -c_p 4^p M^p \iint_{B_r \times B_r} (\zeta(x) - \zeta(y))^p d_K(x, y) \\ &\geq -M^p r^{-ps} |B_r|. \end{aligned}$$

The lower estimate on  $J_2$  can be splitted as follows;

$$\begin{aligned} J_2 &\geq 4 \int_{\mathbb{R}^n \setminus B_r} \int_{B_r} M(u(y) - M)_+^{p-1} \zeta^p(x) d_K(x, y) \\ &\quad - 4M \int_{E_M} \int_{B_r} (u(x) - u(y))_+^{p-1} \zeta^p(x) d_K(x, y) \\ &:= J_{2,1} - J_{2,2}, \end{aligned}$$

where  $E_M = \{y \in \mathbb{R}^n \setminus B_r : u(y) < M\}$ . Since  $(u(y) - M)_+ \geq u_+(y) - M$ , it follows from (c) of Lemma 2.4 that

$$(u(y) - M)_+^{p-1} \geq b_p u_+^{p-1}(y) - M^{p-1}$$

where  $b_p = \mathbb{1}_{(1,2]}(p) + 2^{-(p-1)} \mathbb{1}_{(2,\infty)}(p)$ . Thus the lower estimate on  $J_{2,1}$  can be obtained as

$$(4.16) \quad J_{2,1} \geq d_2 M r^{-ps} |B_r| \left[ \mathcal{T}_r(u_+; 0) \right]^{p-1} - d_3 M^p r^{-ps} |B_r|$$

with universal constants  $d_2, d_3 > 0$ . If  $x \in B_r$  and  $y \in E_M$ , then we observe that

$$\begin{aligned} (u(x) - u(y))_+^{p-1} &\leq a_p (|u(x) - M|^{p-1} + |M - u(y)|^{p-1}) \\ &\leq a_p M^{p-1} + a_p (M + u_-(y) - u_+(y))^{p-1} \\ &\leq a_p M^{p-1} + a_p (M + u_-(y))^{p-1} \\ &\leq a_p (1 + a_p) M^{p-1} + a_p^2 [u_-(y)]^{p-1} \end{aligned}$$

where  $a_p = \mathbb{1}_{(1,2]}(p) + 2^{p-1} \mathbb{1}_{(2,\infty)}(p)$ , because  $u_+(y) < M + u_-(y)$  for any  $y \in E_M$ . Since  $u_-(y) = 0$  for all  $y \in B_R$ , the upper estimate on  $J_{2,2}$  can thus be achieved by

$$(4.17) \quad \begin{aligned} J_{2,2} &\leq 4a_p (1 + a_p) M^p \int_{\mathbb{R}^n \setminus B_r} \int_{B_r} \zeta^p(x) d_K(x, y) \\ &\quad + 4a_p M \int_{\mathbb{R}^n \setminus B_R} \int_{B_r} [u_-(y)]^{p-1} \zeta^2(x) d_K(x, y) \\ &\leq d_4 M^p r^{-ps} |B_r| + d_5 M R^{-ps} |B_r| \left[ \mathcal{T}_R(u_-; 0) \right]^{p-1} \end{aligned}$$

with universal constants  $d_4, d_5 > 0$ . Thus, by (4.16) and (4.17), we have that

$$(4.18) \quad \begin{aligned} J_2 &\geq -d M^p r^{-ps} |B_r| - d M R^{-ps} |B_r| \left[ \mathcal{T}_R(u_-; 0) \right]^{p-1} \\ &\quad + e M r^{-ps} |B_r| \left[ \mathcal{T}_r(u_+; 0) \right]^{p-1} \end{aligned}$$

where  $d, e > 0$  are some universal constants depending only on  $n, s, \lambda$  and  $\Lambda$ .

Finally, it follows from (4.14), Hölder's inequality and the fractional Sobolev inequality that

$$\begin{aligned}
(4.19) \quad J_3 &\geq -2M^p \int_{\mathbb{R}^n} V_+(x) \zeta^p(x) dx \\
&\geq -2M^p \|V_+\|_{L^q(\Omega)} \left( \int_{\Omega} \zeta^{pq'}(x) dx \right)^{\frac{1}{q'}} \\
&\geq -2M^p \|V_+\|_{L^q(\Omega)} \left( \int_{\Omega} \zeta^{\frac{pq'}{n-ps}}(x) dx \right)^{\frac{n-ps}{n}} |\Omega|^{\frac{1}{q'} - \frac{n-ps}{n}} \\
&\geq -2M^p \|V_+\|_{L^q(\Omega)} |\Omega|^{\frac{1}{q'} - \frac{n-ps}{n}} \left( r^{-ps} \|\zeta\|_{L^p(B_r)}^p + [\zeta]_{W^{s,p}(B_r)}^p \right) \\
&\geq -\|V_+\|_{L^q(\Omega)} M^p r^{-ps} |B_r|
\end{aligned}$$

where  $q > \frac{n}{ps} > 1$  and  $1 < q' < \frac{n}{n-ps}$  with

$$\frac{1}{q} + \frac{1}{q'} = 1.$$

Hence the estimates (4.13), (4.15), (4.18) and (4.19) give the required estimate.  $\square$

Next we shall obtain the local boundedness for nonnegative weak subsolutions of the nonlocal equation (1.3) by employing Theorem 1.1 and Lemma 4.3. It is interesting that this estimate no longer depends on the nonlocal tail term, whose proof is pretty simple.

**Theorem 4.4.** *Let  $V \in \mathcal{P}_q^{s,p}(\mathbb{R}^n)$  and  $g \in W^{s,p}(\mathbb{R}^n)$  for  $q > \frac{n}{ps} > 1$  ( $p > 1, s \in (0, 1)$ ). If  $u \in Y_g^{s,p}(\Omega)$  is a nonnegative weak solution of the nonlocal  $p$ -Laplace type Schrödinger equation (1.3), then we have the estimate*

$$\sup_{B_r^0} u \leq C \left( \int_{B_{2r}^0} u^p(x) dx \right)^{\frac{1}{p}}$$

for any  $r > 0$  with  $B_{2r}^0 \subset \Omega$ .

*Proof.* We choose some  $\delta \in (0, 1]$  so that  $1 - \delta d_0 > 0$  and take any  $r > 0$  with  $B_{2r}^0 \subset \Omega$  where

$$d_0 = c_0(1 + \|V_+\|_{L^q(\Omega)}) > 0$$

for the universal constant  $c_0 > 0$  given in Lemma 4.3. Then it follows from Theorem 1.1 and Lemma 4.3 that

$$\sup_{B_r^0} u \leq \delta d_0 \left[ \sup_{B_r^0} u + \mathcal{T}_{2r}(u^-; x_0) \right] + C_0 \delta^{-\frac{(p-1)n}{sp^2}} \left( \int_{B_{2r}^0} u^p(x) dx \right)^{\frac{1}{p}}$$

Since  $\mathcal{T}_{2r}(u^-; x_0) = 0$ , we can easily derive the required result by taking

$$C = \frac{C_0 \delta^{-\frac{(p-1)n}{sp^2}}}{1 - \delta d_0}.$$

Hence we complete the proof.  $\square$

5. THE LOGARITHM OF A WEAK SOLUTION IS A LOCALLY BOUNDED MEAN OSCILLATION FUNCTION

In this section, we prove that the logarithm of a weak supersolution to the nonlocal equation (1.3) is a function with locally bounded mean oscillation. To do this, the following tool which is called the *fractional Poincaré inequality* is very useful.

Let  $n \geq 1, p \geq 1, s \in (0, 1)$  and  $sp < n$ . For a ball  $B \subset \mathbb{R}^n$ , let  $u_B$  denote the average of  $u \in W^{s,p}(B)$  over  $B$ , i.e.

$$u_B = \fint_B u(y) dy.$$

Then it was shown in [BBM, MS] that

$$(5.1) \quad \|u - u_B\|_{L^p(B)}^p \leq \frac{c_{n,p}(1-s)|B|^{\frac{sp}{n}}}{(n-sp)^{p-1}} [u]_{W^{s,p}(B)}^p$$

with a universal constant  $c_{n,p} > 0$  depending only on  $n$  and  $p$ , which is usually very useful in getting the logarithmic estimate of weak supersolutions. Of course, the logarithmic estimate could be obtained as in [DKP1], but we will not apply their approach to achieve it. Our method to realize the logarithmic estimate is easier and more simple than their method.

**[Proof of Theorem 1.2.]** For simplicity, we set  $x_0 = 0$ . So, in what follows, we write  $B_r := B_r^0$  for  $r > 0$ . Take any  $r > 0$  so that  $B_{2r} \subset B_R$  where  $B_R \subset \Omega$ . Consider a radial function  $\zeta \in C_c^\infty(B_{3r/2})$  with values in  $[0, 1]$  such that  $\zeta|_{B_r} \equiv 1, \zeta|_{\mathbb{R}^n \setminus B_{2r}} \equiv 0$  and

$$|\nabla \zeta| \lesssim \frac{1}{r} \text{ in } \mathbb{R}^n.$$

We use the function

$$\varphi(x) = \frac{\zeta^p(x)}{u_b^{p-1}(x)}$$

as a testing function to the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3), where  $u_b(x) = u(x) + b$ . Then we have that

$$(5.2) \quad \begin{aligned} 0 &\leq \iint_{\mathbb{R}^n \times \mathbb{R}^n} H_p(u_b(x) - u_b(y))(\varphi(x) - \varphi(y)) d_K(x, y) + \int_{\mathbb{R}^n} V(x)H_p(u(x))\varphi(x) dx \\ &= \iint_{B_{2r} \times B_{2r}} H_p(u_b(x) - u_b(y))(\varphi(x) - \varphi(y)) d_K(x, y) \\ &\quad + 2 \int_{\mathbb{R}^n \setminus B_{2r}} \int_{B_{2r}} H_p(u_b(x) - u_b(y)) \varphi(x) d_K(x, y) \\ &\quad + \int_{\mathbb{R}^n} V(x)|u(x)|^{p-2}u(x)\varphi(x) dx \\ &:= H(u, \varphi) + I(u, \varphi) + J(u, \varphi). \end{aligned}$$

Without loss of generality, we may assume that

$$u_b(x) \geq u_b(y)$$

for the estimate  $H(u, \varphi)$ ; for, by symmetry, the other case  $u_b(x) < u_b(y)$  can be treated in the exactly same way. Then we have two possible occasions; (a)  $u_b(x) \leq 2u_b(y)$ , (b)  $u_b(x) > 2u_b(y)$ .

[Case (a) :  $u_b(y) \leq u_b(x) \leq 2u_b(y)$ ] By the mean value theorem, we note that

$$(5.3) \quad \begin{aligned} \zeta(x) \geq \zeta(y) &\Rightarrow \zeta^p(x) - \zeta^p(y) = p \int_{\zeta(y)}^{\zeta(x)} \tau^{p-1} d\tau \leq p\zeta^{p-1}(x)(\zeta(x) - \zeta(y)), \\ \zeta(x) < \zeta(y) &\Rightarrow \zeta^p(x) - \zeta^p(y) = p \int_{\zeta(x)}^{\zeta(y)} (-\tau^{p-1}) d\tau \leq p\zeta^{p-1}(x)(\zeta(x) - \zeta(y)). \end{aligned}$$

Then it follows from (5.3) that

$$(5.4) \quad \begin{aligned} \varphi(x) - \varphi(y) &= \frac{\zeta^p(x) - \zeta^p(y)}{u_b^{p-1}(y)} + \zeta^p(x) \left( \frac{1}{u_b^{p-1}(x)} - \frac{1}{u_b^{p-1}(y)} \right) \\ &\leq \frac{p\zeta^{p-1}(x)(\zeta(x) - \zeta(y))}{u_b^{p-1}(y)} \\ &\quad + \zeta^p(x) \int_0^1 \frac{d}{d\tau} \left( \frac{1}{[\tau(u_b(x) - u_b(y)) + u_b(y)]^{p-1}} \right) d\tau \\ &\leq \frac{p\zeta^{p-1}(x)(\zeta(x) - \zeta(y))}{u_b^{p-1}(y)} \\ &\quad - (p-1) \frac{\zeta^p(x)(u_b(x) - u_b(y))}{u_b^p(x)} \\ &\leq \frac{p\zeta^{p-1}(x) |\zeta(x) - \zeta(y)| u_b(y)}{u_b^p(y)} \\ &\quad - \frac{(p-1) \zeta^p(x)(u_b(x) - u_b(y))}{2^p u_b^p(y)}. \end{aligned}$$

Applying Young's inequality with indices  $p' = \frac{p}{p-1}, p, \varepsilon$ , it follows from (5.2) that

$$(5.5) \quad \begin{aligned} H(u, \varphi) &\leq c_{n,p,s} \Lambda p \\ &\iint_{B_{2r} \times B_{2r}} \frac{\varepsilon(u_b(x) - u_b(y))^p \zeta^p(x) + c_\varepsilon |\zeta(x) - \zeta(y)|^p u_b^p(y)}{u_b^p(y)} \frac{dx dy}{|x - y|^{n+ps}} \\ &\quad - \frac{c_{n,p,s} \lambda (p-1)}{2^p} \iint_{B_{2r} \times B_{2r}} \frac{(u_b(x) - u_b(y))^p \zeta^p(x)}{u_b^p(y)} \frac{dx dy}{|x - y|^{n+ps}}. \end{aligned}$$

If we choose  $\varepsilon = \frac{\lambda(p-1)}{2^{p+1}p\Lambda}$  in (5.5), then we have that

$$(5.6) \quad \begin{aligned} H(u, \varphi) &\lesssim - \iint_{B_{2r} \times B_{2r}} \zeta^p(x) \frac{(u_b(x) - u_b(y))^p}{u_b^p(y)} \frac{dx dy}{|x - y|^{n+ps}} \\ &\quad + \iint_{B_{2r} \times B_{2r}} \frac{|\zeta(x) - \zeta(y)|^p}{|x - y|^{n+ps}} dx dy \\ &\lesssim - \iint_{B_{2r} \times B_{2r}} \zeta^p(x) \frac{(u_b(x) - u_b(y))^p}{u_b^p(y)} \frac{dx dy}{|x - y|^{n+ps}} + r^{n-ps}. \end{aligned}$$

because  $x, y \in B_{2r}$ . Since  $0 \leq u_b(x) - u_b(y) \leq u_b(y)$ , we have that

$$(5.7) \quad \begin{aligned} |\ln u_b(x) - \ln u_b(y)|^p &= \left( \int_0^1 \frac{u_b(x) - u_b(y)}{\tau(u_b(x) - u_b(y)) + u_b(y)} d\tau \right)^p \\ &\leq \frac{(u_b(x) - u_b(y))^p}{u_b^p(y)}. \end{aligned}$$

Thus by (5.6) and (5.7) we have that

$$(5.8) \quad \begin{aligned} H(u, \varphi) &\lesssim - \iint_{B_{2r} \times B_{2r}} \zeta^p(x) \left| \ln \left( \frac{u_b(x)}{u_b(y)} \right) \right|^p \frac{dx dy}{|x - y|^{n+ps}} + r^{n-ps} \\ &\lesssim - \iint_{B_r \times B_r} \left| \ln \left( \frac{u_b(x)}{u_b(y)} \right) \right|^p \frac{dx dy}{|x - y|^{n+ps}} + r^{n-ps}. \end{aligned}$$

[Case (b) :  $u_b(x) > 2u_b(y)$ ] It follows from the inequality in Lemma 2.5 with  $\varepsilon = (2^{p-1} - 1)/2$  that

$$(5.9) \quad \begin{aligned} \varphi(x) - \varphi(y) &= \frac{\zeta^p(x) - \zeta^p(y)}{u_b^{p-1}(x)} + \zeta^p(y) \left( \frac{1}{u_b^{p-1}(x)} - \frac{1}{u_b^{p-1}(y)} \right) \\ &\leq \frac{\zeta^p(x) - \zeta^p(y)}{u_b^{p-1}(x)} + \zeta^p(y) \left( \frac{1}{2^{p-1}u_b^{p-1}(y)} - \frac{1}{u_b^{p-1}(y)} \right) \\ &\leq \frac{\varepsilon \zeta^p(y) + c_\varepsilon |\zeta(x) - \zeta(y)|^p}{u_b^{p-1}(x)} - (1 - 2^{-p+1}) \frac{\zeta^p(y)}{u_b^{p-1}(y)} \\ &\leq \frac{c |\zeta(x) - \zeta(y)|^p}{u_b^{p-1}(x)} - \left( \frac{1}{2} - \frac{1}{2^p} \right) \frac{\zeta^p(y)}{u_b^{p-1}(y)}. \end{aligned}$$

Since  $u_b(x) \geq u_b(x) - u_b(y) \geq u_b(y)$ , by (5.2) and (5.9) we have that

$$(5.10) \quad \begin{aligned} \frac{H(u, \varphi)}{c_{n,p,s}} &\leq c\Lambda \iint_{B_{2r} \times B_{2r}} \frac{|\zeta(x) - \zeta(y)|^p}{|x - y|^{n+ps}} dx dy \\ &\quad - c\lambda \left( \frac{1}{2} - \frac{1}{2^p} \right) \iint_{B_{2r} \times B_{2r}} \zeta^p(y) \frac{(u_b(x) - u_b(y))^{p-1}}{u_b^{p-1}(y)} \frac{dx dy}{|x - y|^{n+ps}}. \end{aligned}$$

Since we know that

$$(\ln t)^p \leq c(t-1)^{p-1} \quad \text{for } t > 2,$$

we have that

$$(5.11) \quad \begin{aligned} |\ln u_b(x) - \ln u_b(y)|^p &\leq c \left( \frac{u_b(x) - u_b(y)}{u_b(y)} \right)^{p-1} \\ &= c \frac{(u_b(x) - u_b(y))^{p-1}}{u_b^{p-1}(y)}. \end{aligned}$$



Combining (5.10) with (5.11), we have that

$$\begin{aligned}
(5.12) \quad H(u, \varphi) &\leq - \iint_{B_{2r} \times B_{2r}} \zeta^p(y) \left| \ln \left( \frac{u_b(x)}{u_b(y)} \right) \right|^p \frac{dx dy}{|x-y|^{n+ps}} \\
&\quad + \iint_{B_{2r} \times B_{2r}} \frac{|\zeta(x) - \zeta(y)|^p}{|x-y|^{n+ps}} dx dy \\
&\lesssim - \iint_{B_r \times B_r} \left| \ln \left( \frac{u_b(x)}{u_b(y)} \right) \right|^p \frac{dx dy}{|x-y|^{n+ps}} + r^{n-ps}.
\end{aligned}$$

Hence, by (5.8) and (5.12), we conclude that

$$(5.13) \quad H(u, \varphi) \lesssim - \iint_{B_r \times B_r} \left| \ln \left( \frac{u_b(x)}{u_b(y)} \right) \right|^p \frac{dx dy}{|x-y|^{n+ps}} + r^{n-ps}.$$

For the estimate of  $I(u, \varphi)$ , we note that (i)  $u(y) \geq 0$  and  $u(x) - u(y) \leq u(x)$  for  $(x, y) \in B_{2r} \times (B_R \setminus B_{2r})$  and (ii)  $(u(x) - u(y))_+ \leq u(x) + u_-(y)$  for  $(x, y) \in B_{2r} \times (\mathbb{R}^n \setminus B_R)$ . Since  $\zeta$  is supported in  $B_{3r/2}$ , the above observations (i) and (ii) yield that

$$\begin{aligned}
(5.14) \quad I(u, \varphi) &\leq 2 c_{n,p,s} \Lambda \int_{B_{3r/2}} \int_{\mathbb{R}^n \setminus B_{2r}} \frac{1}{|y-x_0|^{n+ps}} dy dx \\
&\quad + 2 c_{n,p,s} \Lambda \int_{B_{3r/2}} \int_{\mathbb{R}^n \setminus B_R} \frac{u_-^{p-1}(y)}{u_b^{p-1}(x)} \frac{dy dx}{|y-x_0|^{n+ps}} \\
&\lesssim r^{n-ps} + \frac{r^{n-ps}}{b^{p-1}} \left( \frac{r}{R} \right)^{ps} [\mathcal{T}_R(u_-; x_0)]^{p-1}
\end{aligned}$$

because we see  $|y-x| \geq |y-x_0| - |x-x_0| \geq |y-x_0|/4$  for any  $(x, y) \in B_{3r/2} \times (\mathbb{R}^n \setminus B_{2r})$ . Also, it follows from Hölder's inequality and Proposition 2.1 that

$$\begin{aligned}
(5.15) \quad J(u, \varphi) &\leq \int_{\mathbb{R}^n} V_+(x) \zeta^p(x) dx \\
&\leq \|V_+\|_{L^\tau(\Omega)} \left( \int_{\Omega} \zeta^{p\tau'}(x) dx \right)^{\frac{1}{\tau}} \\
&\leq \|V_+\|_{L^q(\Omega)} \left( \int_{\Omega} \zeta^{\frac{pn}{n-ps}}(x) dx \right)^{\frac{n-ps}{n}} |\Omega|^{\frac{1}{q'} - \frac{n-ps}{n}} \\
&\leq \|V_+\|_{L^q(\Omega)} |\Omega|^{\frac{1}{q'} - \frac{n-ps}{n}} \left( r^{-ps} \|\zeta\|_{L^p(B_r)}^p + [\zeta]_{W^{s,p}(B_r)}^p \right) \\
&\leq \|V_+\|_{L^q(\Omega)} |\Omega|^{\frac{1}{q'} - \frac{n-ps}{n}} r^{n-ps} \lesssim \|V_+\|_{L^q(\Omega)} r^{n-ps}
\end{aligned}$$

where  $q > \frac{n}{ps} > 1$ ,  $p > 1$  and  $1 < q' < \frac{n}{n-ps}$  with

$$\frac{1}{q} + \frac{1}{q'} = 1.$$

By (5.13), (5.14) and (5.15), we obtain that

$$\begin{aligned}
&\iint_{B_r \times B_r} \left| \ln \left( \frac{u(x)+b}{u(y)+b} \right) \right|^p \frac{dx dy}{|x-y|^{n+ps}} \\
&\lesssim \frac{r^{n-ps}}{b^{p-1}} \left( \frac{r}{R} \right)^{ps} [\mathcal{T}_R(u_-; x_0)]^{p-1} + r^{n-ps} (1 + \|V_+\|_{L^q(\Omega)})
\end{aligned}$$

for any  $b \in (0, 1)$  and  $r \in (0, R/2)$ , since  $x, y \in B_{2r}$ . Hence we complete the proof by applying (1.1).  $\square$

We now introduce a kind of local BMO spaces on  $B_R^0 \subset \Omega$ , i.e.  $\text{BMO}^p(B_R^0)$  for  $p > 0$ . We define the norm  $\|\cdot\|_{\text{BMO}^p(B_R^0)}$  by

$$\|f\|_{\text{BMO}^p(B_R^0)} = \sup_{r \in (0, R/2)} \left( \int_{B_r^0} |f(y) - f_{B_r^0}|^p dy \right)^{\frac{1}{p}}$$

and the space

$$\text{BMO}^p(B_R^0) = \{f \in L^1_{\text{loc}}(\mathbb{R}^n) : \|f\|_{\text{BMO}^p(B_R^0)} < \infty\}.$$

When  $p = 1$ , we write  $\text{BMO}^p(B_R^0) = \text{BMO}(B_R^0)$ . Then we easily see that

$$(5.16) \quad \| |f| \|_{\text{BMO}(B_R^0)} \leq \|f\|_{\text{BMO}(B_R^0)}$$

because  $||f| - |f_{B_r^0}|| \leq |f - f_{B_r^0}|$ , and also

$$(5.17) \quad \|f \pm g\|_{\text{BMO}(B_R^0)} \leq \|f\|_{\text{BMO}(B_R^0)} + \|g\|_{\text{BMO}(B_R^0)}.$$

We now observe that

$$a \wedge b = \frac{a + b - |a - b|}{2} \quad \text{and} \quad a \vee b = \frac{a + b + |a - b|}{2}$$

for any  $a, b \in \mathbb{R}$ . This implies that

$$(5.18) \quad \begin{aligned} \|f \vee g\|_{\text{BMO}(B_R^0)} &\leq \|f\|_{\text{BMO}(B_R^0)} + \|g\|_{\text{BMO}(B_R^0)}, \\ \|f \wedge g\|_{\text{BMO}(B_R^0)} &\leq \|f\|_{\text{BMO}(B_R^0)} + \|g\|_{\text{BMO}(B_R^0)}. \end{aligned}$$

In addition, we can obtain the following *John-Nirenberg inequality* (as in [Gr]) by using the Calderón-Zygmund decomposition in harmonic analysis as follows; there exists some constants  $b_1, b_2 > 0$  depending only on the dimension  $n$  such that

$$|\{x \in B_r^0 : |f(x) - f_{B_r^0}| > \lambda\}| \leq b_1 e^{-(b_2/\|f\|_{\text{BMO}(B_R^0)})\lambda} |B_r^0|$$

for any  $f \in \text{BMO}(B_R^0)$ , every  $r > 0$  with  $B_{2r}^0 \subset B_R^0$  and  $B_r^0 \subset \Omega$ , and every  $\lambda > 0$ . By standard analysis, this inequality makes it possible to easily show the following fact;

$$(5.19) \quad \begin{aligned} &\text{If } f \in \text{BMO}(B_R^0) \text{ for } B_R^0 \subset \Omega \text{ and } 1 < p < \infty, \text{ then } \|\cdot\|_{\text{BMO}(B_R^0)} \text{ is} \\ &\text{norm-equivalent to } \|\cdot\|_{\text{BMO}^p(B_R^0)}. \end{aligned}$$

**Lemma 5.1.** *If we set*

$$v(x) = \ln\left(\frac{a+b}{u(x)+b}\right) \text{ for } a, b \in (0, 1)$$

where the function  $u$  satisfies the same assumption as Theorem 1.2, then we have that

$$\int_{B_r^0} |v(x) - v_{B_r^0}|^p dx \lesssim \mathfrak{R}_{b,r,R}(u_-; x_0)$$

for any  $r \in (0, R/2)$ , where  $\mathfrak{R}_{b,r,R}(u_-; x_0)$  is given by

$$\mathfrak{R}_{b,r,R}(u_-; x_0) = \frac{1}{b^{p-1}} \left(\frac{r}{R}\right)^{ps} [\mathcal{T}_R(u_-; x_0)]^{p-1} + (1 + \|V_+\|_{L^q(\Omega)}).$$

It follows from this that  $v \in \text{BMO}(B_R^0)$  and moreover

$$\|v\|_{\text{BMO}(B_R^0)} \leq [\mathfrak{R}_{b,r,R}(u_-; x_0)]^{\frac{1}{p}} < \infty.$$

*Proof.* The first part easily follows from the fractional Poincaré inequality (5.1) and Theorem 1.2. Also the second part can be shown by applying the Remark of Theorem 1.1 and Hölder's inequality because  $u \in W^{s,p}(\mathbb{R}^n)$ .  $\square$

**Corollary 5.2.** *If we set*

$$\bar{v} = (v \vee 0) \wedge d \text{ for } d > 0$$

*with the same  $v$  as Lemma 5.1, then we have that*

$$\int_{B_r^0} |\bar{v}(x) - \bar{v}_{B_r^0}|^p dx \lesssim \mathfrak{R}_{b,r,R}(u_-; x_0)$$

*for any  $r \in (0, R/2)$ , where  $\mathfrak{R}_{b,r,R}(u_-; x_0)$  is given by*

$$\mathfrak{R}_{b,r,R}(u_-; x_0) = \frac{1}{b^{p-1}} \left( \frac{r}{R} \right)^{ps} [\mathcal{T}_R(u_-; x_0)]^{p-1} + (1 + \|V_+\|_{L^q(\Omega)}).$$

*It follows from this that  $\bar{v} \in \text{BMO}(B_R^0)$  and moreover*

$$\|\bar{v}\|_{\text{BMO}(B_R^0)} \leq [\mathfrak{R}_{b,r,R}(u_-; x_0)]^{\frac{1}{p}} < \infty.$$

*Proof.* Without loss of generality, assume that  $x_0 = 0$ . By Lemma 5.1, we have that

$$\int_{B_r} ||v(x)| - |v|_{B_r}|^p dx \lesssim \mathfrak{R}_{b,r,R}(u_-; 0),$$

because we see that

$$||v(x)| - |v|_{B_r}| \leq |v(x) - v_{B_r}|.$$

Then we can easily derive from (5.17) and (5.19) that

$$\int_{B_r} |\bar{v}(x) - \bar{v}_{B_r}| dx \lesssim [\mathfrak{R}_{b,r,R}(u_-; 0)]^{\frac{1}{p}}.$$

Finally, the second part can be done as in Lemma 5.1. Hence we complete the proof.  $\square$

## 6. INTERIOR HÖLDER REGULARITY

In this section, we establish an interior Hölder regularity of weak solutions to the nonlocal  $p$ -Laplacian type Schrödinger equation (1.3) by applying the previous results obtained in Section 4 and Section 5.

**[Proof of Theorem 1.3]** Fix any  $p > 1$  and  $0 < s < 1$  and take any  $R > 0$  with  $B_R(x_0) \subset \Omega$ . For simplicity, without loss of generality, we may assume that  $x_0 = 0$ . For any  $k \in \mathbb{N} \cup \{0\}$  and  $r \in (0, R/2)$ , we set

$$r_k = \frac{\delta^k r}{2} \text{ for } \delta \in \left(0, \left(\frac{1}{4}\right)^{\frac{p-1}{ps}}\right), B_k = B_{r_k} \text{ and } B_k^* = B_{2r_k}.$$

Let us set

$$\Xi(r_0) = 2\mathcal{T}_{r/2}(u; 0) + 2C_0 \left( \int_{B_r} |u|^p dx \right)^{\frac{1}{p}}$$

where  $C_0 > 1$  is the constant given in Theorem 1.1. As in the proof of Theorem 1.2, without loss of generality we may assume that

$$\|u\|_{L^\infty(\mathbb{R}^n)} = \frac{1}{4C_0} < 1.$$

So we see that

$$0 \leq \Xi(r_0) < 1.$$

For  $k \in \mathbb{N} \cup \{0\}$ , we set

$$\Xi(r_k) = \left(\frac{r_k}{r_0}\right)^\eta \Xi(r_0)$$

where  $\eta \in (0, \frac{ps}{p-1})$  is some constant to be determined later. For our proof, we have only to prove that

$$(6.1) \quad \operatorname{osc}_{B_k} u \leq \Xi(r_k)$$

for any  $k \in \mathbb{N} \cup \{0\}$ .

We proceed the proof by using the mathematical induction. By the remark of Theorem 1.1, we see that

$$\operatorname{osc}_{B_0} u \leq \Xi(r_0).$$

Assume that (6.1) holds for all  $k \in \{0, 1, \dots, m\}$ . Then we will show that (6.1) is still true for  $m + 1$ . For this proof, we consider two possible cases; either

$$(6.2) \quad \frac{|B_{k+1}^* \cap \{u \geq \inf_{B_k} u + \Xi(r_k)/2\}|}{|B_{k+1}^*|} \geq \frac{1}{2}$$

or

$$(6.3) \quad \frac{|B_{k+1}^* \cap \{u \leq \inf_{B_k} u + \Xi(r_k)/2\}|}{|B_{k+1}^*|} \geq \frac{1}{2}.$$

If (6.2) is true, then we set  $u_k = u - \inf_{B_k} u$ , and if (6.3) is true, then we set

$$u_k = \Xi(r_k) - \left(u - \inf_{B_k} u\right).$$

In these two cases, we see that  $u_k \geq 0$  in  $B_k$  and

$$(6.4) \quad \frac{|B_{k+1}^* \cap \{u_k \geq \Xi(r_k)/2\}|}{|B_{k+1}^*|} \geq \frac{1}{2}.$$

Furthermore,  $u_k$  is a weak solution satisfying that

$$(6.5) \quad \sup_{B_k} |u_m| \leq 2 \Xi(r_k)$$

for all  $k \in \{0, 1, \dots, m\}$ . Under the induction hypothesis, if  $m \geq 1$ , then we now claim that

$$(6.6) \quad [\mathcal{T}_{r_k}(u_k; 0)]^{p-1} \leq c \delta^{-(p-1)\eta} [\Xi(r_k)]^{p-1}$$

for all  $k \in \{0, 1, \dots, m\}$ . Indeed, by (6.5) and the fact that

$$\begin{aligned} \int_{\mathbb{R}^n \setminus B_0} \frac{|u_m(x)|^{p-1}}{|x|^{n+ps}} dx &\lesssim r_0^{-ps} \sup_{B_0} |u|^{p-1} \\ &\quad + r_0^{-ps} [\Xi(r_0)]^{p-1} + \int_{\mathbb{R}^n \setminus B_0} \frac{|u(x)|^{p-1}}{|x|^{n+ps}} dx \\ &\lesssim r_1^{-ps} \Xi(r_0), \end{aligned}$$

we have the following estimate

$$\begin{aligned}
[\mathcal{T}_{r_m}(u_m; 0)]^{p-1} &= c r_m^{ps} \sum_{k=1}^m \int_{B_{k-1} \setminus B_k} \frac{|u_m(x)|^{p-1}}{|x|^{n+ps}} dx + c r_m^{ps} \int_{\mathbb{R}^n \setminus B_0} \frac{|u_m(x)|^{p-1}}{|x|^{n+ps}} dx \\
&\lesssim r_m^{ps} \sum_{k=1}^m \left[ \sup_{B_{k-1}} |u_m| \right]^{p-1} \int_{\mathbb{R}^n \setminus B_k} \frac{1}{|x|^{n+ps}} dx + r_m^{ps} \int_{\mathbb{R}^n \setminus B_0} \frac{|u_m(x, t)|^{p-1}}{|x|^{n+ps}} dx \\
&\leq \sum_{k=1}^m \left( \frac{r_m}{r_k} \right)^{ps} [\Xi(r_{k-1})]^{p-1} = [\Xi(r_0)]^{p-1} \sum_{k=1}^m \left( \frac{r_m}{r_k} \right)^{ps} \left( \frac{r_{k-1}}{r_0} \right)^{(p-1)\eta} \\
&= [\Xi(r_0)]^{p-1} \left( \frac{r_m}{r_0} \right)^{(p-1)\eta} \sum_{k=1}^m \left( \frac{r_m}{r_k} \right)^{ps-(p-1)\eta} \left( \frac{r_{k-1}}{r_k} \right)^{(p-1)\eta} \\
&= [\Xi(r_m)]^{p-1} \delta^{-(p-1)\eta} \sum_{k=1}^m \delta^{(m-k)[ps-(p-1)\eta]} \\
&\leq \delta^{-(p-1)\eta} \frac{\delta^{(m-1)[ps-(p-1)\eta]}}{1 - \delta^{-(ps-(p-1)\eta)}} [\Xi(r_m)]^{p-1} \lesssim \delta^{-(p-1)\eta} [\Xi(r_m)]^{p-1}.
\end{aligned}$$

For  $k$  and  $d > 0$ , we set

$$v_k = \left[ \ln \left( \frac{\Xi(r_k)/2 + b}{u_k + b} \right) \vee 0 \right] \wedge d.$$

Applying Corollary 5.2 with  $a = \Xi(r_k)/2$ ,  $b \in (0, 1)$  and  $d > 0$ , we have that

$$(6.7) \quad \int_{B_{k+1}^*} |v_k(x) - (v_k)_{B_{k+1}^*}|^p dx \lesssim \mathfrak{R}_{b, r_{k+1}, r_k}((u_k)_-; 0)$$

where  $\mathfrak{R}_{b, r, R}(u_-; x_0)$  is given by

$$\mathfrak{R}_{b, r, R}(u_-; x_0) = \frac{1}{b^{p-1}} \left( \frac{r}{R} \right)^{ps} [\mathcal{T}_R(u_-; x_0)]^{p-1} + (1 + \|V_+\|_{L^q(\Omega)}).$$

If we set  $b = \delta^{\frac{ps}{p-1}-\eta} \Xi(r_k)$  in (6.7), then by (5.19) and (6.6) we obtain that

$$\begin{aligned}
(6.8) \quad & \int_{B_{k+1}^*} |v_k(x) - (v_k)_{B_{k+1}^*}| dx \\
& \leq \left( \int_{B_{k+1}^*} |v_k(x) - (v_k)_{B_{k+1}^*}|^p dx \right)^{\frac{1}{p}} \\
& \leq c (1 + \|V_+\|_{L^q(\Omega)})^{\frac{1}{p}}
\end{aligned}$$

where  $c > 0$  is the constant depending only on  $n, s, p, \eta, \lambda$  and  $\Lambda$ . From (6.4), we can derive the following estimate

$$\begin{aligned}
(6.9) \quad d &= \frac{1}{|B_{k+1}^* \cap \{u_k \geq \Xi(r_k)/2\}|} \int_{B_{k+1}^* \cap \{u_k \geq \Xi(r_k)/2\}} d dx \\
&= \frac{1}{|B_{k+1}^* \cap \{u_k \geq \Xi(r_k)/2\}|} \int_{B_{k+1}^* \cap \{v_k=0\}} d dx \\
&\leq \frac{2}{|B_{k+1}^*|} \int_{B_{k+1}^*} (d - v_k) dx = 2(d - (v_k)_{B_{k+1}^*}).
\end{aligned}$$

The estimates (6.8) and (6.9) make it possible to obtain the estimate

$$\begin{aligned}
(6.10) \quad \frac{|B_{k+1}^* \cap \{v_k = d\}|}{|B_{k+1}^*|} d &\leq \frac{2}{|B_{k+1}^*|} \int_{B_{k+1}^* \cap \{v_k = d\}} (d - (v_k)_{B_{k+1}^*}) dx \\
&\leq \frac{2}{|B_{k+1}^*|} \int_{B_{k+1}^* \cap \{v_k = d\}} (v_k - (v_k)_{B_{k+1}^*}) dx \\
&\lesssim (1 + \|V_+\|_{L^q(\Omega)})^{\frac{1}{p}}.
\end{aligned}$$

We now set

$$d = d_* := \ln\left(\frac{\Xi(r_k)/2 + \delta^{2s-\gamma} \Xi(r_k)}{3 \delta^{2s-\gamma} \Xi(r_k)}\right).$$

Then we see that  $d_* \sim \ln(1/\delta)$ . By (6.10), we have that

$$\begin{aligned}
(6.11) \quad \frac{|B_{k+1}^* \cap \{u_k \leq 2 \delta^{\frac{ps}{p-1}-\eta} \Xi(r_k)\}|}{|B_{k+1}^*|} &\leq \frac{c(1 + \|V_+\|_{L^q(\Omega)})^{\frac{1}{p}}}{d_*} \\
&\leq \frac{c_0(1 + \|V_+\|_{L^q(\Omega)})^{\frac{1}{p}}}{\ln(1/\delta)}.
\end{aligned}$$

Now we proceed the next step with a well-known iteration process as follows. For  $i \in \mathbb{N} \cup \{0\}$ , we set

$$\rho_i = (1 + 2^{-i})r_{k+1}, \bar{\rho}_i = \frac{\rho_i + \rho_{i+1}}{2}, B_i = B_{\rho_i} \text{ and } \bar{B}_i = B_{\bar{\rho}_i}.$$

For  $i \in \mathbb{N} \cup \{0\}$ , we consider a function  $\zeta_i \in C_c^\infty(B_{\bar{\rho}_i})$  with  $\zeta_i|_{B_{\rho_{i+1}}} \equiv 1$  such that  $0 \leq \zeta_i \leq 1$  and  $|\nabla \zeta_i| \leq c\rho_i^{-1}$  in  $\mathbb{R}^n$ . Moreover we set  $d_i = (1 + 2^{-i})\delta^{\frac{ps}{p-1}-\eta} \Xi(r_k)$  and  $w_i = (d_i - u_k)_+$  and

$$N_i = \frac{|B_i \cap \{u_k \leq d_i\}|}{|B_i|} = \frac{|B_i \cap \{w_i \geq 0\}|}{|B_i|}.$$

From (6.11), we see that

$$(6.12) \quad N_0 \leq \frac{c_0(1 + \|V_+\|_{L^q(\Omega)})^{\frac{1}{p}}}{\ln(1/\delta)}.$$

By Theorem 1.5, we have that

$$\begin{aligned}
[w_i \zeta_i]_{W^{s,p}(B_{\rho_i})}^p &\lesssim \iint_{B_{\rho_i} \times B_{\rho_i}} [w_i(x) \vee w_i(y)]^p |\zeta_i(x) - \zeta_i(y)|^p d_K(x, y) \\
&\quad + \left( \sup_{x \in \text{supp}(\zeta_i)} \int_{\mathbb{R}^n \setminus B_{\rho_i}} [w_i(y)]^{p-1} K(x-y) dy \right) \|w_i \zeta_i\|_{L^1(B_i)}^p \\
&:= A(\rho_i, w_i, \zeta_i) + B(\rho_i, w_i, \zeta_i).
\end{aligned}$$

Then we have the following estimate

$$\begin{aligned}
(6.13) \quad A(\rho_i, w_i, \zeta_i) &\lesssim d_i^p \int_{B_{\rho_i}} \int_{B_{\rho_i} \cap \{u_k \leq d_i\}} \frac{\sup_{\mathbb{R}^n} |\nabla \zeta_i|^p}{|x-y|^{n+ps-p}} dx dy \\
&\lesssim d_i^p \left(\frac{1}{\rho_i}\right)^p \int_{B_{\rho_i} \cap \{u_k \leq d_i\}} \int_{B_{2\rho_i}} \frac{1}{|y|^{n+2s-2}} dy dx \\
&\lesssim d_i^p \rho_i^{-ps} |B_i \cap \{u_k \leq d_i\}|.
\end{aligned}$$

From the fact that

$$|y-x| \geq |y|-|x| \geq \left(1 - \frac{\bar{\rho}_i}{\rho_i}\right)|y| \geq 2^{-i-2}|y|$$

for any  $y \in \mathbb{R}^n \setminus B_{\rho_i}$  and  $x \in B_{\bar{\rho}_i}$ , we obtain that

$$\begin{aligned}
(6.14) \quad B(\rho_i, w_i, \zeta_i) &\lesssim d_i 2^{i(n+ps)} |B_i \cap \{u_k \leq d_i\}| \int_{\mathbb{R}^n \setminus B_{\rho_i}} \frac{|w_i(y)|^{p-1}}{|y|^{n+ps}} dy \\
&\lesssim 2^{i(n+ps)} d_i \rho_i^{-ps} |B_i \cap \{u_k \leq d_i\}| [\mathcal{T}_{r_{k+1}}(w_i; 0)]^{p-1}.
\end{aligned}$$

Thus it follows from (6.13) and (6.14) that

$$\begin{aligned}
(6.15) \quad [w_i \zeta_i]_{W^{s,p}(B_{\rho_i})}^p &\lesssim \left(d_i^p + 2^{i(n+ps)} d_i [\mathcal{T}_{r_{k+1}}(w_i; 0)]^{p-1}\right) \rho_i^{-ps} |B_i \cap \{u_k \leq d_i\}|.
\end{aligned}$$

From (6.6) and the facts that  $w_i \leq 2\delta^{\frac{ps}{p-1}-\eta} \Xi(r_k)$  in  $B_k$  and  $w_i \leq |u_k| + 2\delta^{\frac{ps}{p-1}-\eta} \Xi(r_k)$  in  $\mathbb{R}^n$ , we can derive that

$$\begin{aligned}
(6.16) \quad [\mathcal{T}_{r_{k+1}}(w_i; 0)]^{p-1} &\lesssim r_{k+1}^{ps} \int_{B_k \setminus B_{k+1}} \frac{|w_i(y, t)|^{p-1}}{|y|^{n+ps}} dy + \left(\frac{r_{k+1}}{r_k}\right)^{ps} [\mathcal{T}_{r_k}(w_i; 0)]^{p-1} \\
&\lesssim \delta^{ps-(p-1)\eta} [\Xi(r_k)]^{p-1} + \delta^{ps} [\mathcal{T}_{r_k}(u_k; (0, 0))]^{p-1} \lesssim d_i^{p-1}.
\end{aligned}$$

Thus by (6.15) and (6.16), we have that

$$(6.17) \quad [w_i \zeta_i]_{W^{s,p}(B_{\rho_i})}^p \lesssim 2^{i(n+ps)} d_i^p \rho_i^{-ps} |B_i \cap \{u_k \leq d_i\}|.$$

By applying (6.17) and the fractional Sobolev's inequality with the exponent

$$\gamma = \frac{n}{n-ps},$$

we can deduce the following inequalities

$$\begin{aligned}
(6.18) \quad \left(\int_{B_{i+1}} |w_i|^{p\gamma} dx\right)^{\frac{1}{\gamma}} &\leq \left(\int_{B_i} |w_i \zeta_i|^{\frac{pn}{n-ps}} dx\right)^{\frac{n-ps}{n}} \\
&\lesssim [w_i \zeta_i]_{W^{s,p}(B_{\rho_i})}^p + \rho_i^{-ps} \|w_i \zeta_i\|_{L^p(B_{\rho_i})}^p \\
&\lesssim 2^{i(n+ps)} d_i^p \rho_i^{-ps} |B_i \cap \{u_k \leq d_i\}|.
\end{aligned}$$

Since  $|B_{i+1}| \sim \rho_i^n \sim |B_i|$  and

$$w_i = (d_i - u_k)_+ \geq (d_i - d_{i+1}) \mathbb{1}_{\{u_k \leq d_{i+1}\}} \geq 2^{-i-2} d_i \mathbb{1}_{\{u_k \leq d_{i+1}\}},$$

this estimate (6.18) yields that

$$\begin{aligned} (d_i - d_{i+1})^p \left( \frac{|B_{i+1} \cap \{u_k \leq d_{i+1}\}|}{|B_{i+1}|} \right)^{\frac{1}{\gamma}} &\leq \frac{\rho_i^{ps}}{|B_i|} \left( \int_{B_{i+1}} |w_i|^{p\gamma} dx dt \right)^{\frac{1}{\gamma}} \\ &\lesssim 2^{i(n+ps)} d_i^p \frac{|B_i \cap \{u_k \leq d_i\}|}{|B_i|}, \end{aligned}$$

which gives that

$$N_{i+1}^{\frac{1}{\gamma}} \leq c \frac{2^{i(n+ps)} d_i^p}{(d_i - d_{i+1})^p} N_i \leq c 2^{i(n+ps+p)} N_i.$$

This leads us to obtain that

$$N_{i+1} \leq c_1 2^{i\gamma(n+ps+p)} N_i^{1+\frac{ps}{n-ps}}.$$

If we could show that

$$(6.19) \quad N_0 = \frac{|B_0 \cap \{u_k \leq 2\delta^{\frac{ps}{p-1}-\eta} \Xi(r_k)\}|}{|B_0|} \leq c_1^{-\frac{n-ps}{ps}} 2^{-\frac{\gamma(n-ps)^2(n+ps+p)}{p^2s^2}} := c_*,$$

then by Lemma 2.2 we conclude that  $\lim_{i \rightarrow \infty} N_i = 0$ , i.e.

$$\inf_{B_{k+1}} u_k > \delta^{\frac{ps}{p-1}-\eta} \Xi(r_k).$$

In order to guarantee (6.19), by (6.12) we observe that

$$c_* \geq \frac{c_0 \left(1 + \|V_+\|_{L^q(\Omega)}\right)^{\frac{1}{p}}}{\ln(1/\delta)} \Leftrightarrow \delta \leq e^{-(c_0/c_*)(1+\|V_+\|_{L^q(\Omega)})^{1/p}},$$

and so we choose  $\delta > 0$  as

$$\delta = e^{-(c_0/c_*)(1+\|V_+\|_{L^q(\Omega)})^{1/p}} \wedge \left(\frac{1}{4}\right)^{\frac{p-1}{ps}}.$$

If  $u_k = u - \inf_{B_k} u$ , then by (6.1) we have that

$$(6.20) \quad \operatorname{osc}_{B_{k+1}} u = \operatorname{osc}_{B_{k+1}} u_k \leq \operatorname{osc}_{B_k} u - \inf_{B_{k+1}} u_k \leq (1 - \delta^{\frac{ps}{p-1}-\eta}) \Xi(r_k).$$

If  $u_k = \Xi(r_k) - (u - \inf_{B_k} u)$ , then we have that

$$(6.21) \quad \operatorname{osc}_{B_{k+1}} u = \operatorname{osc}_{B_{k+1}} u_k = \Xi(r_k) - \inf_{B_{k+1}} u + \inf_{B_k} u - \inf_{B_{k+1}} u_k \leq (1 - \delta^{\frac{ps}{p-1}-\eta}) \Xi(r_k).$$

From (6.20) and (6.21), we obtain that

$$\begin{aligned} (6.22) \quad \operatorname{osc}_{B_{k+1}} u &\leq (1 - \delta^{\frac{ps}{p-1}-\eta}) \Xi(r_k) = (1 - \delta^{\frac{ps}{p-1}-\eta}) \left(\frac{r_k}{r_{k+1}}\right)^\eta \Xi(r_{k+1}) \\ &= (1 - \delta^{\frac{ps}{p-1}-\eta}) \delta^{-\eta} \Xi(r_{k+1}). \end{aligned}$$

To find  $\eta$  so that  $(1 - \delta^{\frac{ps}{p-1}-\eta}) \delta^{-\eta} \leq 1$ , we consider the function

$$\xi(\eta) = \delta^\eta + \delta^{\frac{ps}{p-1}-\eta}.$$

We note that

$$\xi'(\eta) = \delta^\eta \ln \delta \left(1 - \delta^{\frac{ps}{p-1}-2\eta}\right) = 0 \Leftrightarrow \eta = \frac{ps}{2(p-1)} := \eta_{p,s},$$



$\xi'(\eta) < 0$  for  $\eta < \eta_{p,s}$  and  $\xi'(\eta) > 0$  for  $\eta > \eta_{p,s}$ . These facts imply that the graph of  $\xi$  is going down from the point  $(0, 1 + \delta^{2\eta_{p,s}})$  to the point  $(\eta_{p,s}, 1 + 2\delta^{\eta_{p,s}})$  and is going up the point  $(2\eta_{p,s}, 1 + \delta^{2\eta_{p,s}})$  right after that. Since we see that

$$2\delta^{\eta_{p,s}} < 1 + \delta^{2\eta_{p,s}} \quad \text{and} \quad 2\delta^{\eta_{p,s}} < 1,$$

we can find exactly two  $\eta$ 's inside  $(0, \frac{ps}{p-1})$  so that

$$(6.23) \quad \delta^\eta + \delta^{\frac{ps}{p-1} - \eta} = 1.$$

If we set  $Y = \delta^\eta$ , then the above equation (6.23) will be transformed into

$$\frac{Y^2 - Y + \delta^{\frac{ps}{p-1}}}{Y} = 0$$

and its solutions are

$$\delta^\eta = Y = \frac{1 \pm \sqrt{1 - 4\delta^{\frac{ps}{p-1}}}}{2} \in (0, 1),$$

because  $\delta < (1/4)^{\frac{p-1}{ps}}$ . Hence it turns out that the solutions of (6.23) are

$$\eta_0^\pm = \frac{\ln\left(\frac{1 \pm \sqrt{1 - 4\delta^{\frac{ps}{p-1}}}}{2}\right)}{\ln \delta}.$$

Then we see that  $\xi(\eta) \geq 1$ , i.e.  $(1 - \delta^{\frac{ps}{p-1} - \eta})\delta^{-\eta} \leq 1$  for all  $\eta \in (0, \eta_0^-] \cup [\eta_0^+, 2\eta_{p,s})$ . Thus, if  $\eta \in (0, \eta_0^-] \cup [\eta_0^+, 2\eta_{p,s})$ , then by (6.22) we conclude that

$$\text{osc}_{B_{k+1}} u \leq \Xi(r_{k+1}).$$

Therefore we complete the proof.  $\square$

**Acknowledgement.** Yong-Cheol Kim was supported by School of Education, Korea University Grant in 2023.

#### REFERENCES

- [AS] M. Aizenman and B. Simon, *Brownian Motions and Harnack inequality for Schrödinger Operators*, Comm. pure and appl. Math. **35**, 1982, 209–273.
- [BBM] J. Bourgain, H. Brezis and P. Mironescu, *Limiting embedding theorems for  $W^{s,p}$  when  $s \uparrow 1$  and applications*, J. Anal. Math. **87**, 2002, 77–101.
- [BV] C. Bucur and E. Valdinoci, *Nonlocal Diffusion and Applications*, Lecture Notes of the Unione Matematica Italiana, vol. 20, Springer, 2016.
- [CFG] F. Chiarenza, E. Fabes and N. Garofalo, *Harnack's inequality for Schrödinger operators and the continuity of solutions*, Proc. Amer. Math. Soc. **98(3)**, 1986, 415–425.
- [CK1] W. Choi and Y.-C. Kim, *The Malgrange-Ehrenspreis theorem for nonlocal Schrödinger operators with certain potentials*, Comm. Pure. & Appl. Anal. **17(5)**, 2018, 1993–2010.
- [CK2] W. Choi and Y.-C. Kim,  *$L^p$ -mapping properties for nonlocal Schrödinger operators with certain potentials*, Discrete Contin. Dyn. Syst. - A **38**, 2018, 5811–5834.
- [CV] L. A. Caffarelli and A. Vasseur, *Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation*, Ann. Math. **171(3)**, 2010, 1903–1930.
- [C] P. Constantin, *Euler equations, Navier-Stokes equations and turbulence*, in: *Mathematical Foundations of Turbulent Viscous Flows*, in: *Lecture Notes in Math. vol. 1871*, Springer, Berlin, 2006, pp. 1–43.
- [DKP] A. Di Castro, T. Kuusi and G. Palatucci, *Nonlocal Harnack inequalities*, Jour. of Funct. Anal. **267**, 2014, 1807–1836.
- [DKP1] A. Di Castro, T. Kuusi and G. Palatucci, *Local behavior of fractional  $p$ -minimizers*, Ann. Inst. H. Poincaré Anal. Non Linéaire **33(5)**, 2016, 1279–1299.

- [DPV] E. Di Nezza, G. Palatucci and E. Valdinoci *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136**, 2012, 521–573.
- [F] C. Fefferman, *The uncertainty principle*, Bull. Amer. Math. Soc.(N.S.) **9**, 1983, 129–206.
- [FSV] A. Fiscella, R. Servadei and E. Valdinoci, *Density properties for fractional Sobolev spaces*, Ann. Acad. Sci. Fenn. Math. **40**, 2015, 235–253.
- [GR] J. Garcia-Cuerva and J. L. Rubio De Francia, *Weighted norm inequalities and related topics*, North-Holland, 1985.
- [GT] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 224. Springer-Verlag, Berlin, 1983. xiii+513 pp.
- [Gr] Loukas Grafakos, *Classical and Modern Fourier Analysis* Prentice Hall, 2004, Pearson Education Inc.
- [G] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Monographs and Studies in Mathematics, vol 24 Pitman (Advanced Publishing Program), 1985, Boston.
- [IN] H. Ishii and G. Nakamura, *A class of integral equations and approximation of  $p$ -Laplace equations*, Calc. Var. **37**, 2010, 485–522.
- [K] Y.-C. Kim, *Nonlocal Harnack inequalities for nonlocal heat equations*, J. Differential Equations **267**, 2019, 6691–6757.
- [K1] Y.-C. Kim, *Nonlocal Harnack inequalities for nonlocal Schrödinger operators with  $A_1$ -Muckenhoupt potentials*, J. Math. Anal. Appl. **507**, 2022, 125746.
- [K2] Y.-C. Kim, *Hölder continuity of weak solutions to nonlocal Schrödinger equations with  $A_1$ -Muckenhoupt potentials*, to appear in Potential Analysis.
- [K3] Y.-C. Kim, *Nonlocal Harnack inequalities for nonlocal  $p$ -Laplacian type Schrödinger operators with  $A_1^p$ -Muckenhoupt potentials*, submitted.
- [L] N. Laskin, *Fractional Schrödinger equation*, Phys. Rev. E **66**, 2002, 056108.
- [Mi] N. Miller, *Weighted Sobolev spaces and Pseudodifferential operators with smooth symbols*, Trans. Amer. Math. Soc. **269**(1), 1982, 91–109.
- [SV] R. Servadei and E. Valdinoci, *Variational methods for nonlocal operators of elliptic type*, Discrete Contin. Dyn. Syst. **33**, 2013, 2105–2137.
- [S1] Z. Shen,  *$L^p$  estimates for Schrödinger operators with certain potentials*, Annales de l'institut Fourier **45**(2), 1995, 513–546.
- [S2] Z. Shen, *On the Neumann Problem for Schrödinger operators in Lipschitz Domains*, Indiana Univ. Math. Jour. **43**(1), 1994, 143–176.
- [S] L. Silvestre, *The regularity of the obstacle problem for a fractional power of the Laplace operator*, Commun. Pure Appl. Math. **60**(1), 2007, 67–112.
- [Si] B. Simon, *Schrödinger semigroups*, Bull. Amer. Math. Soc. **7**(3), 1982, 447–526.

• YONG-CHEOL KIM : DEPARTMENT OF MATHEMATICS EDUCATION, KOREA UNIVERSITY, SEOUL 02841, REPUBLIC OF KOREA & SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 02455, REPUBLIC OF KOREA  
 Email address: ychkim@korea.ac.kr