

INVERSE PROBLEMS OF IDENTIFYING THE TIME-DEPENDENT SOURCE COEFFICIENT FOR SUBELLIPTIC HEAT EQUATIONS

MANSUR I. ISMAILOV, TOHRU OZAWA, AND DURVUDKHAN SURAGAN

ABSTRACT. We discuss inverse problems of determining the time-dependent source coefficient for a general class of subelliptic heat equations. We show that a single data at an observation point guarantees the existence of a (smooth) solution pair for the inverse problem. Moreover, additional data at the observation point implies an explicit formula for the time-dependent source coefficient. We also explore an inverse problem with nonlocal additional data, which seems a new approach even in the Laplacian case.

CONTENTS

1. Introduction	1
2. Single datum	2
3. Double data	6
4. Nonlocal data	8
5. Particular cases	9
5.1. Laplacian	9
5.2. Sub-Laplacians	10
5.3. Baouendi-Grushin operator	10
References	10

1. INTRODUCTION

Let $X = (X_1, \dots, X_m)$ be a system of real smooth vector fields defined over a subset W of \mathbb{R}^d , $d \geq 2$, satisfying the Hörmander condition (H): There exists a natural number r such that the vector fields X_1, \dots, X_m together with their r commutators span the tangent space at each point of W .

Let $\Omega \subset W$ be a bounded connected open subset with smooth boundary $\partial\Omega$ non-characteristic of $X = (X_1, \dots, X_m)$. In Ω , we consider the following inverse problem

2010 *Mathematics Subject Classification.* 35R30, 35K20, 65M22.

Key words and phrases. Inverse problem, control parameter, subelliptic heat equation, Hörmander vector field.

This work was supported by the NU grant 20122022FD4105. The second author is supported in part by JSPS Kakenhi 18KK0073, 19H00644. No new data was collected or generated during the course of the research.

of finding a pair (u, p) :

$$\begin{cases} \partial_t u(x, t) - \Delta_X u(x, t) = p(t)u(x, t) + f(x, t), & \text{in } \Omega \times (0, T), \\ u(x, 0) = \varphi(x), & x \in \Omega, \\ u(x, t) = 0, & \text{on } \partial\Omega \times (0, T), \end{cases} \quad (1.1)$$

with an additional condition. Here

$$\Delta_X := - \sum_{i=1}^m X_i^* X_i,$$

with $X_i^* = -X_i - \operatorname{div} X_i$, is a self-adjoint subelliptic operator (see [2]).

In the classical case when $X_i = \partial_{x_i}$, $i = 1, \dots, d$, one has the usual (elliptic) Laplacian Δ instead of the subelliptic operator Δ_X in (1.1) and there is a vast of literature on such parabolic inverse problems, especially, in 1D-cases. See, e.g. [6] and references therein. Note that, in the classical case, the existence and uniqueness result of this inverse problem was established by Cannon, Lin and Wang [4] and [5], by using the method of retardation of the time variable and a priori estimates. Without any claim of completeness, we refer an interested reader to [3] and [7] for analysis of related inverse problems in the classical setting.

The present paper aims to analyse inverse problems of recovering the time-dependent source parameter $p(t)$ in the Cauchy-Dirichlet problem for the subelliptic heat equation (1.1). First, in order to find a pair (u, p) , we fix a point $q \in \Omega$ as an observation point for some time-dependent quantity. So, by using this additional data we recover the time-dependent source parameter $p(t)$. Interestingly, we discover that this method works well to study an inverse problem for the same model but with nonlocal additional data. The latter approach seems new even in the Laplacian case. Our proofs rely on subelliptic spectral theory arguments.

Moreover, we also state that the solution to the inverse problem can be found explicitly in the case of an increase in the number of overdetermined data by potential theory techniques.

We organize our paper as follows. In Section 2, we prove the existence and uniqueness result with a single datum at an observation point by using a spectral theory approach. In Section 3, the time-dependent source parameter is found in an explicit form by applying the potential theory arguments. In Section 4, our technique from Section 2 is applied to treat a nonlocal case. Some interesting particular models are discussed in Section 5.

2. SINGLE DATUM

Let Ω be a bounded connected open subset with smooth boundary $\partial\Omega$ non-characteristic of $X = (X_1, \dots, X_m)$ and let $|H| > 0$, where the set H is defined in (2.4). Consider the following inverse problem of finding a pair (u, p) :

$$\begin{cases} \partial_t u(x, t) - \Delta_X u(x, t) = p(t)u(x, t) + f(x, t), & \text{in } \Omega \times (0, T), \\ u(x, 0) = \varphi(x), & x \in \Omega, \\ u(x, t) = 0, & \text{on } \partial\Omega \times (0, T), \end{cases} \quad (2.1)$$

where $\varphi \in C^{2k}(\bar{\Omega})$ and $f \in C^{2k,0}(\bar{\Omega} \times [0, T])$ for integer $k > \frac{\tilde{\nu}}{4} + 1$. Here and after we understand $g \in C^1(\Omega)$ if $Xg \in C(\Omega)$.

The operator Δ_X is well-defined on $\{u \in H_{X,0}^1(\Omega) : \Delta_X u \in L^2(\Omega)\}$. Recall that $H_X^1(\Omega) = \{u \in L^2(\Omega) : X_i u \in L^2(\Omega), 1 \leq i \leq m\}$ and $H_{X,0}^1(\Omega)$ is the closure of $C_0^\infty(\Omega)$ in $H_X^1(\Omega)$.

For any $x \in \bar{\Omega}$ and given $1 \leq k \leq r$, let $V_k(x)$ be the subspaces of the tangent space at x spanned by all commutators of X_1, \dots, X_m with length at most k . Note that the number r in the statement (H) in the introduction is called the Hörmander index. The Hausdorff dimension ν of Ω (or it can be also called the Métivier index of Ω) is defined as

$$\nu := \sum_{k=1}^r k(\nu_k - \nu_{k-1})$$

with $\nu_0 = 0$. Here it is assumed that for each $x \in \bar{\Omega}$, $\dim V_k(x)$ is a constant denoted by ν_k in a neighborhood of x .

Let us consider the following Dirichlet spectral problem of finding a nontrivial function ϕ and eigenvalue λ :

$$\begin{cases} -\Delta_X \phi(x) = \lambda \phi(x), & \text{in } \Omega, \\ \phi(x) = 0, & \text{on } \partial\Omega. \end{cases}$$

By using the spectral theorem for compact self-adjoint operators, it can be shown that the eigenspaces are finite-dimensional and that the Dirichlet eigenvalues λ are real, positive, and have no limit point. The eigenspaces are orthogonal in the space of square-integrable functions and consist of smooth functions. In fact, the system of eigenfunctions $\phi_n(x)$, $n \in \mathbb{N}$, are complete orthonormal system in $L_2(\Omega)$.

The eigenvalues can be arranged in increasing order:

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \rightarrow \infty,$$

where each eigenvalue is counted according to its geometric multiplicity. It is well-known that Weyl's asymptotic formula for the Dirichlet Laplacian ($\nu = d$) holds

$$\lambda_n \sim n^{\frac{2}{\nu}}. \quad (2.2)$$

The asymptotic formula (2.2) fails to hold for general Hörmander vector fields not satisfying the so-called Métivier condition. However, recently, in [2] for the operator Δ_X it was proved that Weyl's asymptotic formula

$$\lambda_n \sim n^{\frac{2}{\tilde{\nu}}} \quad (2.3)$$

holds if and only if $|H| > 0$. Here

$$H := \{x \in \Omega \mid \nu(x) = \tilde{\nu}\} \quad (2.4)$$

and $\nu(x) := \sum_{j=1}^r j(\nu_j(x) - \nu_{j-1}(x))$ (with $\nu_0(x) := 0$) is a pointwise homogeneous dimension and

$$\tilde{\nu} := \max_{x \in \Omega} \nu(x). \quad (2.5)$$

Clearly, $\tilde{\nu} = \nu$ if $\dim V_k(x)$ is a constant in a neighborhood of x .

Let $q \in \Omega$ be a point such that $\phi_n(q)$, $n = 1, 2, \dots$, is bounded with $\phi_n(q) \neq 0$ for some $n_0 \in \mathbb{N}$. Let $\mathbb{N}_q \subset \mathbb{N}$ be maximal set such that $\phi_n(q) \neq 0$ for all $n \in \mathbb{N}_q$. We use this point $q \in \Omega$ as an observation point for the quantity:

$$w(t) = u(q, t), \quad t \in [0, T]. \quad (2.6)$$

Set $\varphi_n = \int_{\Omega} \varphi(x) \phi_n(x) dx$ and $f_n(t) = \int_{\Omega} f(t, x) \phi_n(x) dx$.

Lemma 1. *If $\varphi \in C^{2k}(\bar{\Omega})$ and $\Delta_X^m \varphi = 0$, $m = 0, \dots, k-1$, on $x \in \partial\Omega$ with $k > \frac{\bar{\nu}}{4} + 1$, then we have*

$$\sum_{n=1}^{\infty} \lambda_n |\varphi_n| \leq c^{\frac{1}{2}} \|\Delta_X^k \varphi\|_{L_2},$$

where $c = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^{2(k-1)}}$.

Proof. From Green's identities for the subelliptic operator Δ_X we get

$$\varphi_n = (-1)^k \frac{1}{\lambda_n^k} (\Delta_X^k \varphi)_n,$$

where $(\Delta_X^k \varphi)_n = \int_{\Omega} \Delta_X^k \varphi(x) \phi_n(x) dx$. By the Cauchy-Schwartz and Bessel inequalities we have

$$\sum_{n=1}^{\infty} \lambda_n |\varphi_n| = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^{k-1}} |(\Delta_X^k \varphi)_n| \leq \left(\sum_{n=1}^{\infty} \frac{1}{\lambda_n^{2(k-1)}} \right)^{\frac{1}{2}} \left(\sum_{n=1}^{\infty} |(\Delta_X^k \varphi)_n|^2 \right)^{\frac{1}{2}} \leq c^{\frac{1}{2}} \|\Delta_X^k \varphi\|_{L_2}.$$

The series $c = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^{2(k-1)}}$ is convergent by the Weyl-type asymptotic formula (2.3), since $k > \frac{\bar{\nu}}{4} + 1$. \square

Let us introduce:

$$D_k(\Omega) = \{ \varphi \in C^{2k}(\bar{\Omega}) : \Delta_X^m \varphi = 0, \quad m = 0, \dots, k-1 \text{ on } x \in \partial\Omega \}$$

for some $k > \frac{\bar{\nu}}{4} + 1$.

We have the following assumptions about the given functions.

- (1) $\varphi \in D_k(\Omega)$ with $\varphi_n \phi_n(q) \geq 0$ for $\forall n \in \mathbb{N}_q$ and $\varphi_{n_0} \phi_{n_0}(q) > 0$ for some $n_0 \in \mathbb{N}_q$;
- (2) $f \in C(\Omega \times [0, T])$ and $f(x, t) \in D_k(\Omega)$ with $f_n(t) \phi_n(q) \geq 0$ for $\forall t \in [0, T]$ and for $\forall n \in \mathbb{N}_q$;
- (3) $w \in C[0, T]$ with $w(t) \neq 0$ for $\forall t \in [0, T]$ and $w(0) = \varphi(q)$.

The following theorem is valid for the existence and uniqueness of the inverse problem.

Theorem 1. *Assuming conditions (1)-(3) hold, a unique smooth pair (u, p) exists that solves the inverse problem (2.1) with (2.6).*

Proof of Theorem 1. The solution of (2.1) has the form:

$$u(x, t) = \sum_{n=1}^{\infty} \left(\varphi_n e^{-\lambda_n t + \int_0^t p(\tau) d\tau} + \int_0^t f_n(\tau) e^{-\lambda_n(t-\tau) + \int_{\tau}^t p(s) ds} d\tau \right) \phi_n(x)$$

with $\varphi_n = \int_{\Omega} \varphi(x) \phi_n(x) dx$ and $f_n(t) = \int_{\Omega} f(t, x) \phi_n(x) dx$.

Let $r(t) = e^{-\int_0^t p(\tau)d\tau}$. So, we have

$$r(t)u(x, t) = \sum_{n=1}^{\infty} \left(\varphi_n e^{-\lambda_n t} + \int_0^t f_n(\tau) e^{-\lambda_n(t-\tau)} r(\tau) d\tau \right) \phi_n(x).$$

From the additional condition (2.6) it follows that

$$r(t)u(q, t) = \sum_{n=1}^{\infty} \left(\varphi_n e^{-\lambda_n t} + \int_0^t f_n(\tau) e^{-\lambda_n(t-\tau)} r(\tau) d\tau \right) \phi_n(q) = r(t)w(t)$$

or

$$r(t) = \frac{\sum_{n=1}^{\infty} \varphi_n e^{-\lambda_n t} \phi_n(q)}{w(t)} + \frac{1}{w(t)} \int_0^t \left(\sum_{n=1}^{\infty} \phi_n(q) f_n(\tau) e^{-\lambda_n(t-\tau)} \right) r(\tau) d\tau \quad (2.7)$$

in the case $w(t) \neq 0$.

From Green's identities for the subelliptic operator Δ_X we get

$$\varphi_n = \frac{(-1)^k}{\lambda_n^k} \int_{\Omega} \Delta_X^k \varphi(x) \phi_n(x) dx$$

for $\varphi \in C^{2k}(\bar{\Omega})$ and $\Delta_X^m \varphi = 0$, $m = 0, \dots, k-1$, on $x \in \partial\Omega$. We have

$$|\varphi_n| = \frac{1}{\lambda_n^k} \left| \int_{\Omega} \Delta_X^k \varphi(x) \phi_n(x) dx \right|.$$

Let $|\phi_n(q)| \leq M$ for some $M = \text{const} > 0$. The majorant of the series

$$\sum_{n=1}^{\infty} \varphi_n e^{-\lambda_n t} \phi_n(q)$$

is $\sum_{n=1}^{\infty} |\varphi_n|$, that is, by the Cauchy-Schwartz inequality we have

$$\sum_{n=1}^{\infty} |\varphi_n| = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^k} \left| \int_{\Omega} \Delta_X^k \varphi(x) \phi_n(x) dx \right| \leq \left(\sum_{n=1}^{\infty} \frac{1}{\lambda_n^{2k}} \right)^{\frac{1}{2}} \left(\sum_{n=1}^{\infty} \left| \int_{\Omega} \Delta_X^k \varphi(x) \phi_n(x) dx \right|^2 \right)^{\frac{1}{2}}.$$

Now we apply the Bessel inequality

$$\sum_{n=1}^{\infty} \frac{1}{\lambda_n^{2k}} \sum_{n=1}^{\infty} \left| \int_{\Omega} \Delta_X^k \varphi(x) \phi_n(x) dx \right|^2 \leq \sum_{n=1}^{\infty} \frac{1}{\lambda_n^{2k}} \|\Delta_X^k \varphi\|_{L_2}^2.$$

This series is convergent by the Weyl-type asymptotic formula (2.3).

The same inequality is true for the second series in (2.7). Then the second kind Volterra equation has a unique continuous solution in $[0, T]$ and it must be positive. Then

$$r(t) = e^{-\int_0^t p(\tau)d\tau} \implies p(t) = -\frac{r'(t)}{r(t)}.$$

The function $r(t)$ is continuously differentiable if the series $\sum_{n=1}^{\infty} \varphi_n \lambda_n e^{-\lambda_n t} \phi_n(q)$ is uniformly convergent. It is the case since the majorant series $M \sum_{n=1}^{\infty} \lambda_n |\varphi_n|$ is convergent for $k > \frac{\tilde{\nu}}{4} + 1$ by Lemma 1. \square

3. DOUBLE DATA

Let $\Omega \subset \mathbb{R}^d$ be bounded open set with piecewise smooth boundary $\partial\Omega$. Consider the inverse problem of identifying a pair (u, p) :

$$\begin{cases} \partial_t u(x, t) - \Delta_X u(x, t) = p(t)u(x, t) + f(x, t), & \text{in } \Omega \times (0, T), \\ u(x, 0) = \varphi(x), & x \in \Omega, \\ u(x, t) = 0, & \text{on } \partial\Omega \times (0, T), \end{cases} \quad (3.1)$$

where $u_0 \in C_0^2(\Omega)$ and $f \in C_0^{2,0}(\Omega \times [0, T])$ are given functions. Assume that there exists $q \in \Omega$ such that $f(q, t) \in C^1[0, T]$ is continuously differentiable function with $f(q, t) \neq 0$. We now fix $q \in \Omega$ as an observation point for two time-dependent quantities:

$$w_1(t) := v_1(t, q), \quad w_2(t) := v_2(t, q), \quad t \in [0, T]. \quad (3.2)$$

Here

$$\begin{cases} \partial_t v_1(x, t) - \Delta_X v_1(x, t) = p(t)v_1(x, t) + f(x, t), & \text{in } \Omega \times (0, T), \\ v_1(x, 0) = \varphi(x), & x \in \Omega, \\ v_1(x, t) = 0, & \text{on } \partial\Omega \times (0, T), \end{cases} \quad (3.3)$$

and

$$\begin{cases} \partial_t v_2(x, t) - \Delta_X v_2(x, t) = p(t)v_2(x, t) + \Delta_X f(x, t), & \text{in } \Omega \times (0, T), \\ v_2(x, 0) = \Delta_X \varphi(x), & x \in \Omega, \\ v_2(x, t) = 0, & \text{on } \partial\Omega \times (0, T). \end{cases} \quad (3.4)$$

Theorem 2. *Let $\varphi \in C_0^2(\Omega)$ and $f \in C_0^{2,0}(\Omega \times [0, T])$. Let $q \in \Omega$ be such that $f(q, t) \in C^1[0, T]$ is continuously differentiable function with $f(q, t) \neq 0$. Let w_1 and w_2 defined in (3.2) be the observation data. Then there exists a unique smooth solution pair (u, p) for the inverse problem (3.1) with*

$$p(t)w_1(t) = w_1'(t) - w_2(t) - f(q, t).$$

Note that if p is found, then the solution u of the direct problem has the following representation

$$\begin{aligned} u(x, t) = \exp\left(\int_0^t p(\tau)d\tau\right) \int_0^t \int_{\Omega} h_D(x, y, t - \tau) \exp\left(-\int_0^\tau p(s)ds\right) f(y, \tau) dy d\tau \\ + \exp\left(\int_0^t p(\tau)d\tau\right) \int_{\Omega} h_D(x, y, t) \varphi(y) dy, \end{aligned}$$

where h_D is the subelliptic heat kernel [2, Theorem 1.1] for the Cauchy-Dirichlet problem for the subelliptic heat equation in the cylindrical domain $\Omega \times [0, T]$.

Proof of Theorem 2. Let us recall the following transformation from [4]:

$$v(x, t) = r(t)u(x, t), \quad r(t) = \exp\left(-\int_0^t p(\tau)d\tau\right),$$

that is,

$$p(t) = -\frac{r'(t)}{r(t)}, \quad u(x, t) = \frac{v(x, t)}{r(t)}.$$

Thus, we have

$$\begin{cases} \partial_t v(x, t) - \Delta_X v(x, t) = r(t)f(x, t), & \text{in } \Omega \times (0, T), \\ u(x, 0) = \varphi(x), & x \in \Omega, \\ u(x, t) = 0, & \text{on } \partial\Omega \times (0, T). \end{cases} \quad (3.5)$$

We now fix a point $q \in \Omega$ as an observation point for two time dependent quantities:

$$\tilde{w}_1(t) := w_1(t, q), \quad \tilde{w}_2(t) := w_2(t, q), \quad t \in [0, T].$$

Here

$$\begin{cases} \partial_t w_1(x, t) - \Delta_X w_1(x, t) = r(t)f(x, t), & \text{in } \Omega \times (0, T), \\ w_1(x, 0) = \varphi(x), & x \in \Omega, \\ w_1(x, t) = 0, & \text{on } \partial\Omega \times (0, T). \end{cases} \quad (3.6)$$

and

$$\begin{cases} \partial_t w_2(x, t) - \Delta_X w_2(x, t) = r(t)\Delta_X f(x, t), & \text{in } \Omega \times (0, T), \\ w_2(x, 0) = \Delta_X \varphi(x), & x \in \Omega, \\ w_2(x, t) = 0, & \text{on } \partial\Omega \times (0, T). \end{cases} \quad (3.7)$$

It is known that the solutions of Cauchy-Dirichlet problems (3.6) and (3.7), correspondingly, can be presented by

$$w_1(x, t) = \int_0^t \int_{\Omega} h_D(x, y, t - \tau) r(\tau) f(y, \tau) dy + \int_{\Omega} h_D(x, y, t) \varphi(y) dy \quad (3.8)$$

and

$$w_2(x, t) = \int_0^t \int_{\Omega} h_D(x, y, t - \tau) r(\tau) \Delta_X f(y, \tau) dy + \int_{\Omega} h_D(x, y, t) \Delta_X \varphi(y) dy. \quad (3.9)$$

That is, for $q \in \Omega$ and for all $t \in [0, T]$ we have

$$\begin{aligned} w_1(q, t) &= \int_0^t \int_{\Omega} h_D(q, y, t - \tau) r(\tau) f(y, \tau) dy d\tau + \int_{\Omega} h_D(q, y, t) \varphi(y) dy = \tilde{w}_1(t); \\ w_2(q, t) &= \int_0^t \int_{\Omega} h_D(q, y, t - \tau) r(\tau) \Delta_X f(y, \tau) dy d\tau + \int_{\Omega} h_D(q, y, t) \Delta_X \varphi(y) dy = \tilde{w}_2(t). \end{aligned}$$

By differentiating \tilde{w}_1 and applying the Leibniz integral rule, then using $\partial_t h_D(x, y, t - \tau) = \Delta_X h_D(x, y, t - \tau)$, $t > \tau$, $\partial_t h_D(x, y, t) = \Delta_X h_D(x, y, t)$, $t > 0$, and Green's

second identity for the operator Δ_X , we get

$$\begin{aligned}
\tilde{w}'_1(t) &= \int_0^t \int_{\Omega} \partial_t h_D(q, y, t - \tau) r(\tau) f(y, \tau) dy d\tau + r(t) f(q, t) \\
&+ \int_{\Omega} \partial_t h_D(q, y, t) \varphi(y) dy = \int_0^t \int_{\Omega} \Delta_X h_D(q, y, t - \tau) r(\tau) f(y, \tau) dy d\tau \\
&+ r(t) f(q, t) + \int_{\Omega} \Delta_X h_D(q, y, t) \varphi(y) dy \\
&= \int_0^t \int_{\Omega} h_D(q, y, t - \tau) r(\tau) \Delta_X f(y, \tau) dy d\tau + r(t) f(q, t) \\
&+ \int_{\Omega} h_D(q, y, t) \Delta_X \varphi(y) dy = \tilde{w}_2(t) + r(t) f(q, t).
\end{aligned} \tag{3.10}$$

Thus, we arrive at

$$\tilde{w}_2(t) = \tilde{w}'_1(t) - r(t) f(q, t). \tag{3.11}$$

Moreover, we have the following relations

$$\tilde{w}_1(t) := r(t) w_1(t), \quad \tilde{w}_2(t) := r(t) w_2(t), \quad t \in [0, T].$$

Plugging in (3.11) we obtain

$$p(t) w_1(t) = w'_1(t) - w_2(t) - f(q, t).$$

Also, the direct Cauchy-Dirichlet problems (3.5), (3.6), and (3.7) have a unique solution. It implies that there exists a unique classical pair (v, r) , that is, (u, p) for the inverse problem (3.1). The proof is complete. \square

4. NONLOCAL DATA

Consider the following problem:

$$\begin{cases} \partial_t u(x, t) = \Delta_X u(x, t) + r(t) f(x, t), & \text{in } \Omega \times (0, T), \\ u(x, 0) = \varphi(x), & x \in \Omega, \\ u(x, t) = 0, & \text{on } \partial\Omega \times (0, T), \end{cases} \tag{4.1}$$

where $\varphi \in C^{2k}(\Omega)$ and $f \in C^{2k, 0}(\Omega \times [0, T])$ for some integer $k > \frac{\bar{\nu}}{4} + 1$.

Note that the equation (4.1) is equivalent to (2.1) (see the proof of Theorem 2). Now we consider the inverse problem of finding (u, r) from (4.1) with nonlocal datum

$$\int_{\Omega} \omega(x) u(x, t) dx = w(t), \quad t \in [0, T], \tag{4.2}$$

where $\omega(x) \in L_2(\Omega)$.

One could also discuss the case when (4.2) is replaced by the measurement in Section 2:

$$u(q, t) = w(t), \quad t \in [0, T],$$

where q is a given space observation point in Ω such that the sequence $\varphi_n(q)$, $n = 1, 2, \dots$, is bounded. This is retrieved when one considers the Dirac delta distribution

$\delta(x - q)$ centred at q in equation (4.2).

The following theorem is valid for the existence and uniqueness of the inverse problem.

Theorem 3. *Let the following conditions hold:*

- (1) $\varphi \in D_k(\Omega)$;
- (2) $f \in C(\Omega \times [0, T])$, $f(x, t) \in D_k(\Omega)$ and $\int_{\Omega} \omega(x) f(x, t) dx \neq 0$ for $\forall t \in [0, T]$;
- (3) $w \in C^1[0, T]$ with $w(t) \neq 0$ for $\forall t \in [0, T]$ and $w(0) = \int_{\Omega} \omega(x) \varphi(x) dx$.

Then there exists a unique smooth pair (u, r) for the inverse problem (4.1)-(4.2).

Proof. The solution of (4.1) has the form:

$$u(x, t) = \sum_{n=1}^{\infty} \left(\varphi_n e^{-\lambda_n t} + \int_0^t f_n(\tau) e^{-\lambda_n(t-\tau)} r(\tau) d\tau \right) \phi_n(x),$$

where $\varphi_n = \int_{\Omega} \varphi(x) \phi_n(x) dx$ and $f_n(t) = \int_{\Omega} f(x, t) \phi_n(x) dx$. So, we have

$$u_t(x, t) = \sum_{n=1}^{\infty} \left(-\lambda_n \varphi_n e^{-\lambda_n t} - \lambda_n \int_0^t f_n(\tau) e^{-\lambda_n(t-\tau)} r(\tau) d\tau + f_n(t) r(t) \right) \phi_n(x).$$

From the over-determination condition (4.2) it follows that

$$\begin{aligned} \int_{\Omega} \omega(x) u_t(x, t) dx &= \sum_{n=1}^{\infty} \left(-\lambda_n \varphi_n e^{-\lambda_n t} - \lambda_n \int_0^t f_n(\tau) e^{-\lambda_n(t-\tau)} r(\tau) d\tau \right) \int_{\Omega} \omega(x) \phi_n(x) dx \\ &\quad + r(t) \int_{\Omega} \omega(x) \sum_{n=1}^{\infty} f_n(t) \phi_n(x) dx = \omega'(t) \end{aligned}$$

or

$$\begin{aligned} r(t) &= \frac{\omega'(t) + \sum_{n=1}^{\infty} \lambda_n \varphi_n e^{-\lambda_n t} \int_{\Omega} \omega(x) \phi_n(x) dx}{\int_{\Omega} \omega(x) f(x, t) dx} \\ &\quad + \frac{1}{\int_{\Omega} \omega(x) f(x, t) dx} \int_0^t \left(\sum_{n=1}^{\infty} \lambda_n f_n(\tau) e^{-\lambda_n(t-\tau)} \int_{\Omega} \omega(x) \phi_n(x) dx \right) r(\tau) d\tau \end{aligned} \tag{4.3}$$

since $\int_{\Omega} \omega(x) \sum_{n=1}^{\infty} f_n(t) \phi_n(x) dx = \int_{\Omega} \omega(x) f(x, t) dx$.

Let $\int_{\Omega} \omega^2(x) dx \leq M$ in (4.2) for some $M = \text{const} > 0$. The series

$$\sum_{n=1}^{\infty} \lambda_n \varphi_n e^{-\lambda_n t} \int_{\Omega} \omega(x) \phi_n(x) dx$$

is uniformly convergent since the majorant series $M \sum_{n=1}^{\infty} \lambda_n |\varphi_n|$ is convergent by Lemma 1. Thus, the Volterra integral equation (4.3) has a unique continuous solution. \square

5. PARTICULAR CASES

5.1. Laplacian. In the classical case, when $X_i = \partial_{x_i}$, $i = 1, \dots, d$, one has the usual (elliptic) Laplacian Δ instead of the subelliptic operator Δ_X in (1.1). Our approach seems new even in this classical case.

5.2. Sub-Laplacians. Let $\mathbb{G} = (\mathbb{R}^d, \circ)$ be a stratified Lie group and $\{X_1, \dots, X_m\}$, $m \leq d$, be a system of generators for \mathbb{G} , i.e. a basis for the first strata of \mathbb{G} . It satisfies the Hörmander condition (H). The sum of squares operator

$$\Delta_{\mathbb{G}} = - \sum_{i=1}^m X_i^2$$

is called the sub-Laplacian on \mathbb{G} . $\Delta_{\mathbb{G}}$ is elliptic if and only if \mathbb{G} is same as $(\mathbb{R}^d, +)$. It is clear that the above results are valid on stratified Lie groups. Note that the class of stratified Lie groups includes the groups of Iwasawa type and the H-type groups. A simple example in \mathbb{R}^3 is the operator

$$(\partial_y + 2x\partial_z)^2 + (\partial_x - 2y\partial_z)^2.$$

5.3. Baouendi-Grushin operator. Let $z := (x, y) := (x_1, \dots, x_m, y_1, \dots, y_k) \in \mathbb{R}^m \times \mathbb{R}^k$ with $m, k \geq 1$ and $m + k = d$. Let us consider the vector fields

$$X_i = \frac{\partial}{\partial x_i}, \quad i = 1, \dots, m, \quad Y_j = |x|^\gamma \frac{\partial}{\partial y_j}, \quad \gamma \geq 0, \quad j = 1, \dots, k.$$

The Baouendi-Grushin operator on \mathbb{R}^{m+k} is defined by

$$\Delta_\gamma = \sum_{i=1}^m X_i^2 + \sum_{j=1}^k Y_j^2 = \Delta_x + |x|^{2\gamma} \Delta_y, \quad (5.1)$$

where Δ_x and Δ_y stand for the standard Laplacians in the variables $x \in \mathbb{R}^m$ and $y \in \mathbb{R}^k$, respectively. Recall that the Baouendi-Grushin operator for a positive even integer γ is a sum of squares of vector fields satisfying the Hörmander condition (H). In that case, the results of the present paper hold for the Baouendi-Grushin operator (cf. [1]). A simple example in \mathbb{R}^2 is the following operator

$$\partial_{xx}^2 + x^2 \partial_{yy}^2.$$

REFERENCES

- [1] K. Beauchard, P. Cannarsa, M. Yamamoto, Inverse source problem and null controllability for multidimensional parabolic operators of Grushin type, *Inverse Problems* 30, (2014), 025006.
- [2] H. Chen and H.-G. Chen, Estimates of Dirichlet eigenvalues for a class of sub-elliptic operators, *Proc. London Math. Soc.* (3) 122 (2021) 808–847.
- [3] P. Cannarsa, J. Tort, M. Yamamoto. Determination of source terms in a degenerate parabolic equation, *Inverse Problems* 26, (2010) 105003 (20pp).
- [4] J. R. Cannon, Y. Lin, S. Wang. Determination of a control parameter in a parabolic partial differential equation, *J. Aust. Math. Soc. Ser. B* 33, 149–163, 1991.
- [5] J. R. Cannon, Y. Lin, S. Wang. Determination of source parameter in parabolic equations, *Meccanica* 27, 85–94, 1992.
- [6] A. Hazanee, D. Lesnic, M. I. Ismailov, N. B. Kerimov. Inverse time-dependent source problems for the heat equation with nonlocal boundary conditions, *Applied Mathematics and Computation* 346, 800–815, 2019.
- [7] M. Karazym, T. Ozawa, D. Suragan, Multidimensional inverse Cauchy problems for evolution equations, *Inverse Probl. Sci. En.*, 28:11 (2020), 1582–1590.

MANSUR I. ISMAILOV:
DEPARTMENT OF MATHEMATICS
GEBZE TECHNICAL UNIVERSITY

TURKEY
AND DEPARTMENT OF MATHEMATICS
KHAZAR UNIVERSITY
BAKU AZ1096
AZERBAIJAN
E-mail address MISMAILOV@GTU.EDU.TR

TOHRU OZAWA:
DEPARTMENT OF APPLIED PHYSICS
WASEDA UNIVERSITY
TOKYO 169-8555
JAPAN
E-mail address TXOZAWA@WASEDA.JP

DURVUDKHAN SURAGAN:
DEPARTMENT OF MATHEMATICS
NAZARBAYEV UNIVERSITY
ASTANA 010000
KAZAKHSTAN
E-mail address DURVUDKHAN.SURAGAN@NU.EDU.KZ