

**THE UNIFORM DIVERSIFICATION STRATEGY IS OPTIMAL  
FOR EXPECTED UTILITY MAXIMIZATION UNDER HIGH  
MODEL AMBIGUITY**

LAURENCE CARASSUS AND JOHANNES WIESEL

ABSTRACT. We investigate an expected utility maximization problem under model uncertainty in a one-period financial market. We formalize the model uncertainty by taking a Wasserstein ball of radius  $k$  around  $\mathbb{P}$  and consider the corresponding Wasserstein distributionally robust optimization problem. We show that optimal solutions converge to the uniform diversification strategy when uncertainty is increasingly large, i.e. when the radius  $k$  tends to infinity.

1. INTRODUCTION

In this article we consider a one period financial market with  $d$  risky assets and a riskless asset. We assume that all assets are traded, and without loss of generality we express their prices in discounted units. We refer to Section 4 for a more detailed description of the financial market under consideration. We investigate the *Wasserstein distributionally robust optimization problem*

$$(1) \quad u(x_0) := \sup_{w \in \mathbb{R}^d, \langle w, \mathbf{1} \rangle \geq a} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)]$$

for a concave increasing utility function  $U : \mathbb{R} \rightarrow \mathbb{R}$  and an initial capital  $x_0 \neq 0$ . In the above,  $\langle \cdot, \cdot \rangle$  denotes the scalar product on  $\mathbb{R}^d$  and the vector  $w \in \mathbb{R}^d$  denotes the market weights of the  $d$  risky assets. We work under the constraint that we have positive exposure to the risky assets, i.e.  $a \in (0, 1]$ . The probability measures  $\mathbb{P}$  and  $\mathbb{Q}$  describe the laws of the vector of returns of the  $d$  risky assets, and  $X : \mathbb{R}^d \rightarrow \mathbb{R}^d$  denotes the identity map. We assume that  $\mathbb{E}_{\mathbb{P}}[|X|^p] < \infty$  for some  $p \geq 1$  and denote by  $B_k(\mathbb{P})$  the  $p$ -Wasserstein ball of radius  $k > 0$  around  $\mathbb{P}$ .

Defining

$$(2) \quad u(k, w) := \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)]$$

for  $w \in \mathbb{R}^d$  and

$$w_k := \arg \max_{w \in \mathbb{R}^d, \langle w, \mathbf{1} \rangle \geq a} u(k, w),$$

the main result of this paper is the following:

---

*Date:* June 5, 2023.

JW acknowledges support by NSF Grant DMS-2205534.

**Theorem 1.1.** *Let  $U : \mathbb{R} \rightarrow \mathbb{R}$  be a non-decreasing, non-constant, differentiable and concave function and assume that there exist some constants  $C, \underline{x} > 0$  such that  $U(-\underline{x}) < 0$  and for all  $x \leq -\underline{x}$*

$$(3) \quad U(x) \geq -C(1 + |x|^p).$$

Furthermore assume that one of the three conditions below holds true:

- (i)  $U$  is bounded from above.
- (ii)  $AE_{-\infty}(U) := \liminf_{x \rightarrow -\infty} \frac{xU'(x)}{U(x)} > 1$ .
- (iii)  $AE_{+\infty}(U) := \limsup_{x \rightarrow +\infty} \frac{xU'(x)}{U(x)} < 1$ .

Then  $w_k$  is well-defined and

$$(4) \quad \lim_{k \rightarrow \infty} w_k = w^u,$$

where

$$w^u := \frac{a}{d} \mathbf{1} := \frac{a}{d} (1, \dots, 1)$$

is the rescaled unit vector satisfying  $\langle w^u, \mathbf{1} \rangle = a$ .

**Remark 1.2.** *If condition (3) is not postulated, the problem is ill-posed (see Lemma 5.3). The conditions (ii) and (iii) of Reasonable Asymptotic Elasticity in Theorem 1.1 are classical, see the discussion in Section 5. One may instead assume the milder conditions (22) and (23), see Lemma 5.7 and Proposition 6.2 below.*

**Remark 1.3.** *We prove in Theorem 1.1 that  $\lim_{k \rightarrow \infty} \arg \max_{w \in \mathbb{R}^d: \langle w, \mathbf{1} \rangle \geq a} u(k, w) = w^u$ . One may wonder if  $w^u = \arg \max_{w \in \mathbb{R}^d: \langle w, \mathbf{1} \rangle \geq a} \lim_{k \rightarrow \infty} u(k, w)$  too. We show in Lemma 5.4 that in fact  $\lim_{k \rightarrow \infty} u(k, w) = -\infty$ , so that any  $w$  is an optimizer.*

**Example 1.4.** *We now provide examples where the different cases of Theorem 1.1 apply. Define*

$$U(x) := -\frac{1}{\delta} [(x+1)^{-\delta} - 1] 1_{\{x>0\}} - \frac{1}{p} [(1-x)^p - 1] 1_{\{x \leq 0\}}$$

for some  $\delta > 0$ . Then  $U$  is concave, strictly increasing, continuously differentiable, bounded from above and it satisfies Assumption (3). Similarly we can define

$$U(x) := \log(x) 1_{\{x>1\}} + [x-1] 1_{\{x \leq 1\}}.$$

In this case  $U$  is concave, strictly increasing, continuously differentiable and it satisfies Assumption (3) and (iii) with  $AE_{+\infty}(U) = 0$ . Lastly, (ii) can only occur if  $p > 1$ , see Remark 28. So let us assume that  $p > 1$  and take  $q \in (1, p]$  and define

$$U(x) := x 1_{\{x>0\}} - \frac{1}{q} [(1-x)^q - 1] 1_{\{x \leq 0\}}.$$

Then  $U$  is concave, strictly increasing, continuously differentiable and it satisfies Assumption (3) and (ii) with  $AE_{-\infty}(U) = q$ .

## 2. LITERATURE REVIEW AND DISCUSSION OF THEOREM 1.1

The study of optimal investment problems under a fixed probabilistic model  $\mathbb{P}$  goes back at least to Merton [1969]. More recently, it has been widely acknowledged that agents might face some degree of uncertainty about the choice of  $\mathbb{P}$ . This is often referred to as *Knightian uncertainty* after Knight [1921] and has led to the max-min approach to optimal investment, as studied in detail in Gilboa and Schmeidler [1989], Maccheroni et al. [2006], Schied [2007], Schied et al. [2009]. These works consider an agent who is averse to model uncertainty and thus solves the worst utility maximisation problem

$$\max_w \min_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)];$$

we refer to Oblój and Wiesel [2021] for a more complete discussion of this approach with historical references.

There are many ways to model Knightian uncertainty through the ambiguity set  $\mathcal{P}$  above: we refer to Maenhout [2004], Calafiore [2007], Kerkhof et al. [2010], Hansen and Sargent [2011], and the references therein. In this article we define  $\mathcal{P}$  to be a ball  $B_k(\mathbb{P})$  in the *Wasserstein distance*  $\mathcal{W}_p$  around  $\mathbb{P}$ . The Wasserstein distance has seen a recent spike of interest in the mathematical finance and statistics literature. One reason for this is the fact that Wasserstein balls can be considered non-parametric neighbourhoods of  $\mathbb{P}$ ; indeed  $B_k(\mathbb{P})$  contains infinitely many measures, which are not absolutely continuous with respect to each other. Let us also mention the recent statistical advances of e.g. Dereich et al. [2013], Fournier and Guillin [2015] on the convergence speed of the Wasserstein distance between the true and empirical measure, as well as (dual) reformulations of  $u(x_0)$ , or more generally Wasserstein distributionally robust optimization problems, see e.g. Mohajerin Esfahani and Kuhn [2018], Blanchet and Murthy [2019], Gao and Kleywegt [2022] and the survey articles Pflug and Pohl [2018], Rahimian and Mehrotra [2019].

In this work we are interested in an agent's optimal investment decision when facing an *increasingly large* amount of Knightian uncertainty. Our main contribution is that, under minimal assumptions on  $U$ , the optimal portfolio choice then converges to the uniform diversification strategy. This can be contrasted with the recent works Bartl et al. [2020] and Oblój and Wiesel [2021], where the authors compute first-order sensitivities of the value function  $u(x_0)$  for  $k \rightarrow 0$ , i.e. for infinitesimally small amounts of model uncertainty. It is important to point out, that in order to analyze the other end of the asymptotic regime—namely infinitesimally large amounts of model uncertainty—as we do in this paper, we actually cannot rely on the analytic theory derived in these works. Instead we follow a different strategy, which can be summarized through the following three-step procedure: first, we prove that in  $u(x_0)$  we can restrict to bounded weights (see Proposition 6.2 and Lemma 6.4). Then we give a reformulation of  $u(x_0)$ , which essentially reduces  $u(k, w)$  to the following optimization problem

$$(5) \quad u(k, w) = \inf_{X^\mu, Z^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0| |Z^\mu| |w|)],$$

where  $X^\mu$  has law  $\mathbb{P}$  under  $\mu$  and  $Z^\mu$  is such that  $\mathbb{E}_\mu[|Z^\mu|^p]^{1/p} \leq k$  for some atomless measure  $\mu$ , see Lemma 7.2. Finally, we analyze this problem when  $k \rightarrow \infty$ . It turns out that while the distribution of  $X^\mu$  does not change, the radially symmetric part  $|Z^\mu|$  will dominate in (5) and if  $|w| > |w^u|$ ,  $u(k, w) - u(k, w^u)$  goes to  $-\infty$ , see Proposition 7.3. It is thus sufficient to make the simple observation that  $w^u$  is the unique vector in the set  $\{w : \langle w, \mathbf{1} \rangle \geq a\}$  minimizing the Euclidean distance to the origin. Indeed, by Hölder's inequality, we have for all  $w \in \mathbb{R}^d$  such that  $\langle w, \mathbf{1} \rangle \geq a$  that

$$(6) \quad |w| = \left( \sum_{i=1}^d w_i^2 \right)^{1/2} \geq \frac{1}{\sqrt{d}} \sum_{i=1}^d w_i \geq \frac{a}{\sqrt{d}} = |w^u|,$$

where the inequality is strict if and only if  $w \neq w^u$ . Following this idea of proof enables us to keep the assumptions on  $U$  and  $\mathbb{P}$  minimal.

Note that (5) also implies that the domain of  $U$  necessarily is the whole real line and not only  $(0, \infty)$ . When  $U$  is defined on  $\mathbb{R}$ , the existence of an optimal strategy has been proved in a one period model in the PhD thesis of R. Blanchard, see Blanchard [2017]. The generalization to the multiperiod case is provided in Carassus and Ferhoun [2023]. In the specific case of this article, the proof simplifies significantly. Indeed, existence holds true under minimal conditions on  $U$  and the quasi-sure no arbitrage condition of Bouchard and Nutz [2015] for  $B_k(\mathbb{P})$ , which is in fact always satisfied (see Proposition 4.3).

Probably the paper closest to the results presented here is Pflug et al. [2012], where the authors consider a robust maximisation problem for risk measures. Their main examples concern the Markowitz functional and the conditional value at risk. However, their setting does not easily translate to ours: indeed, [Pflug et al., 2012, Proposition 1] essentially assumes that the map  $\mathbb{Q} \mapsto \|U'(X)\|_{L^p(\mathbb{Q})}$  is constant on  $B_k(\mathbb{P})$ , which is clearly not satisfied for general utility functions of interest. Furthermore, results similar to Pflug et al. [2012] are proved in Sass and Westphal [2021] for the power and logarithmic utility functions when there is drift uncertainty in a multivariate continuous-time Black-Scholes type financial market. However, to the best of our knowledge, the case of general utility maximization has not been addressed, even in a one-period setup.

Lastly, let us mention that, while an extension of our main result to multiple periods seems natural to expect, the methods developed in this article do not seem adequate to provide a direct route for this extension. Indeed we have failed to derive a dynamic programming principle for  $u(k, w)$  as the one step associated problems do not carry our minimal assumptions, in particular monotonicity of  $x_0 \mapsto u(x_0)$ . In consequence we leave the extension of Theorem 1.1 to multiple periods for future research.

The rest of the paper is organized as follows. Section 3 summarizes the notation used in the paper. After describing how we model the financial market in Section 4 and detail the no arbitrage condition, we give a preliminary discussion of the

assumptions of Theorem 1.1 in Section 5. We then show existence of optimizers in different settings in Section 6. These results enable us to state the proof of Theorem 1.1, which can be found in Section 1.1. We conduct a more in depth analysis of the function  $u(k, w)$  in Appendix A, where we detail sufficient criteria for the existence and uniqueness of worst-case measures in the ball  $B_k(\mathbb{P})$ .

### 3. NOTATION

Let us quickly summarize the notation used throughout the article:  $C$  denotes a positive constant (which might change from occurrence to occurrence),  $\langle \cdot, \cdot \rangle$  denotes the scalar product,  $|\cdot|$  the Euclidean distance and  $X : \mathbb{R}^l \rightarrow \mathbb{R}^l$  is the identity mapping for all  $l \geq 1$ . Let  $p \geq 1$ . We denote by  $\mathcal{P}_p(\mathbb{R}^d)$  the set of probability measures  $\mathbb{Q}$  on  $\mathbb{R}^d$  satisfying  $\mathbb{E}_{\mathbb{Q}}[|X|^p] < \infty$ . Let  $\mathbb{P} \in \mathcal{P}_p(\mathbb{R}^d)$  and define  $C_{\mathbb{P}} := (\mathbb{E}_{\mathbb{P}}[|X|^p])^{\frac{1}{p}}$ . We define the ball  $B_k(\mathbb{P})$  around  $\mathbb{P}$  of radius  $k > 0$  with respect to the Wasserstein metric

$$\mathcal{W}_p(\mathbb{P}, \mathbb{Q}) = \left( \inf_{\pi \in \Pi(\mathbb{P}, \mathbb{Q})} \mathbb{E}_{\pi}[|X - Y|^p] \right)^{1/p},$$

where  $\Pi(\mathbb{P}, \mathbb{Q})$  is the set of probability measures on  $\mathbb{R}^d \times \mathbb{R}^d$  such that  $X_*\pi = \mathbb{P}$  and  $Y_*\pi = \mathbb{Q}$ . Here we denote by  $X : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  and  $Y : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^d$  the canonical projections to the first and second coordinate respectively, by a slight abuse of notation (which is standard in the optimal transport literature). The expression  $f(X, Y)_*\pi$  stands for the push-forward measure of  $\pi$  under the function  $(x, y) \mapsto f(x, y)$ , i.e.  $f_*\pi(A) = \pi(f(X, Y) \in A)$  for all Borel sets  $A$ . An application of the triangle inequality directly yields

$$(7) \quad \mathbb{E}_{\mathbb{Q}}[|X|^p]^{\frac{1}{p}} = \mathcal{W}_p(\mathbb{Q}, \delta_0) \leq \mathcal{W}_p(\mathbb{P}, \delta_0) + \mathcal{W}_p(\mathbb{Q}, \mathbb{P}) \leq C_{\mathbb{P}} + k < \infty$$

for all  $\mathbb{Q} \in B_k(\mathbb{P})$ .

### 4. FINANCIAL MARKET

We consider a one period financial market with  $d + 1$  traded assets. The prices of these assets are given by  $(S_t^i)_{t=0,1}$  for  $i \in \{0, \dots, d\}$ . Assume that  $S^0$  is a numeraire asset, i.e.  $S_t^0 > 0$ . The returns  $R^i$ ,  $i \in \{0, \dots, d\}$  represent the profit (or loss) created tomorrow from investing one dollar's worth of asset  $i$  today and are defined as

$$R^i = \frac{S_1^i}{S_0^i} - \frac{S_1^0}{S_0^0}.$$

An agent can trade in the market using strategies  $\phi = (\phi^0, \dots, \phi^d) \in \mathbb{R}^{d+1}$ , where

$$\sum_{i=0}^d \phi^i S_0^i = x_0$$

and  $x_0 \neq 0$  is a given initial wealth. We obtain

$$S_0^0 \phi^0 = x_0 - \sum_{i=1}^d \phi^i S_0^i.$$

The portfolio value at time one is given by

$$\begin{aligned}
\sum_{i=0}^d \phi^i S_1^i &= \left( x_0 - \sum_{i=1}^d \phi^i S_0^i \right) \frac{S_1^0}{S_0^0} + \sum_{i=1}^d \phi^i S_1^i \\
&= x_0 \frac{S_1^0}{S_0^0} + \sum_{i=1}^d \phi^i \left( S_1^i - S_0^i \frac{S_1^0}{S_0^0} \right) \\
&= x_0 \frac{S_1^0}{S_0^0} + \sum_{i=1}^d \phi^i S_0^i R^i = x_0 \frac{S_1^0}{S_0^0} + x_0 \sum_{i=1}^d w^i R^i =: V^{x_0, w},
\end{aligned}$$

where we define the portfolio weights  $w^i := \frac{\phi^i S_0^i}{x_0}$  as the fraction of initial wealth  $x_0$  held in asset  $i$ , recalling that  $x_0 \neq 0$ . By definition we obtain

$$(8) \quad \sum_{i=0}^d w^i = 1.$$

For notational simplicity we assume that  $S_1^0 = S_0^0$  for the remainder of the paper, i.e.  $R^0 = 0$  and  $R^i = S_1^i/S_0^i - 1$ . This yields

$$(9) \quad V^{x_0, w} = x_0 + x_0 \sum_{i=1}^d w^i R^i = x_0 \left( 1 + \sum_{i=1}^d w^i R^i \right).$$

As stated in the Introduction, we model the distribution of  $R = (R^1, \dots, R^d)$  through a probability measure  $\mathbb{P}$ , i.e. the distribution of  $R$  is equal to the distribution of the identity map  $X$  under  $\mathbb{P}$ . By a slight abuse of notation we then write  $V^{x_0, w} := x_0 + x_0 \langle w, X \rangle$  for  $w \in \mathbb{R}^d$ . We can now state our definition of efficient markets relative to the prior  $\mathbb{P}$ :

**Definition 4.1.** *We say that the market has no arbitrage  $\text{NA}(\mathbb{P})$ , if*

$$\forall x_0 \neq 0: \quad V^{x_0, w} \geq x_0 \quad \mathbb{P}\text{-a.s.} \quad \Rightarrow \quad V^{x_0, w} = x_0 \quad \mathbb{P}\text{-a.s.}$$

A similar definition is used in Rásonyi and Stettner [2006] [(3) and Proposition 1]. Definition 4.1 is sometimes also called relative arbitrage with respect to the numeraire portfolio. Using (9) and  $x_0 \neq 0$  we obtain that  $\text{NA}(\mathbb{P})$  is equivalent to

$$\langle w, X \rangle \geq 0 \quad \mathbb{P}\text{-a.s.} \quad \Rightarrow \quad \langle w, X \rangle = 0 \quad \mathbb{P}\text{-a.s.}$$

We also define the following natural generalization to a set of priors  $\mathcal{Q}$ , going back to the seminal work of Bouchard and Nutz [2015]:

**Definition 4.2.** *For a set  $\mathcal{Q}$  of probability measures on  $\mathbb{R}^d$  we say that the quasi-sure no arbitrage  $\text{NA}(\mathcal{Q})$  holds, if for all  $w \in \mathbb{R}^d$*

$$\langle w, X \rangle \geq 0 \quad \mathcal{Q}\text{-q.s.} \quad \Rightarrow \quad \langle w, X \rangle = 0 \quad \mathcal{Q}\text{-q.s.}$$

The next proposition shows that the quasi-sure notion of no arbitrage is very weak when applied to Wasserstein balls  $B_k(\mathbb{P})$ . Indeed, it holds independently if  $\text{NA}(\mathbb{P})$  is true or not:

**Proposition 4.3.**  *$\text{NA}(B_k(\mathbb{P}))$  holds for any  $\mathbb{P} \in \mathcal{P}_p(\mathbb{R}^d)$  and any  $k > 0$ .*

*Proof.* Take  $w \in \mathbb{R}^d$  such that

$$(10) \quad \langle w, X \rangle \geq 0 \quad B_k(\mathbb{P})\text{-q.s.}$$

We claim that this implies

$$(11) \quad \langle w, x \rangle \geq 0 \quad \text{for all } x \in \mathbb{R}^d.$$

Let  $x \in \mathbb{R}^d$ . Lemma 4.4 states that there exists  $\alpha > 0$  such that  $\mathbb{P}_x = (1 - \alpha)\mathbb{P} + \alpha\delta_x \in B_k(\mathbb{P})$ . By (10)

$$\langle w, X \rangle \geq 0 \quad \mathbb{P}_x\text{-a.s.}$$

holds, so that we must thus have  $\langle w, x \rangle \geq 0$ . As  $x \in \mathbb{R}^d$  was arbitrary, (11) follows. But (11) clearly implies that  $w = 0$ , showing the claim.  $\square$

We have used the following lemma:

**Lemma 4.4.** *For any  $\tilde{x} \in \mathbb{R}^d$  and  $\alpha \in \left(0, \left(\frac{k}{C_{\mathbb{P}} + |\tilde{x}|}\right)^p\right)$*   

$$\mathbb{P}_{\tilde{x}} := (1 - \alpha)\mathbb{P} + \alpha\delta_{\tilde{x}} \in B_k(\mathbb{P}).$$

*Proof.* Let  $\alpha > 0$ . Choosing the coupling  $\pi := (1 - \alpha)\left(x \mapsto (x, x)\right)_* \mathbb{P} + \alpha(\mathbb{P} \otimes \delta_{\tilde{x}}) \in \Pi(\mathbb{P}, \mathbb{P}_{\tilde{x}})$  we see that

$$\mathcal{W}_p(\mathbb{P}_{\tilde{x}}, \mathbb{P}) \leq (\alpha \mathbb{E}_{\mathbb{P}}[|X - \tilde{x}|^p])^{1/p} \leq \alpha^{1/p}(\mathbb{E}_{\mathbb{P}}[|X|^p]^{1/p} + |\tilde{x}|) = \alpha^{1/p}(C_{\mathbb{P}} + |\tilde{x}|).$$

Setting the last expression less or equal to  $k$  and solving for  $\alpha$  yields

$$\alpha \leq \left(\frac{k}{C_{\mathbb{P}} + |\tilde{x}|}\right)^p.$$

$\square$

**Definition 4.5.** *For  $\mathbb{Q} \in \mathcal{P}_p(\mathbb{R}^d)$  and  $\mathcal{Q} \subseteq \mathcal{P}_p(\mathbb{R}^d)$  we define*

$$\begin{aligned} \text{supp}(\mathbb{Q}) &:= \bigcap \{A \subset \mathbb{R}^d, \text{ closed}, \mathbb{Q}(A) = 1\} \\ \text{supp}(\mathcal{Q}) &:= \bigcap \{A \subset \mathbb{R}^d, \text{ closed}, \mathbb{Q}(A) = 1 \forall \mathbb{Q} \in \mathcal{Q}\}. \end{aligned}$$

*We denote the affine hull of a set  $A \subseteq \mathbb{R}^d$  by  $\text{Aff}(A)$ .*

**Lemma 4.6.** *We have*

$$\text{supp}(B_k(\mathbb{P})) = \mathbb{R}^d.$$

*Proof.* For any  $x \in \mathbb{R}$ , taking the measures  $\mathbb{P}_x \in B_k(\mathbb{P})$  from Lemma 4.4 we realize that

$$\text{supp}(B_k(\mathbb{P})) \supseteq \bigcup_{x \in \mathbb{R}^d} \text{supp}(\mathbb{P}_x) \supseteq \bigcup_{x \in \mathbb{R}^d} \{x\} = \mathbb{R}^d.$$

This yields the claim.  $\square$

The next proposition asserts that there exists a measure  $\mathbb{P}^*$  in the Wasserstein ball satisfying the no arbitrage condition and having full affine hull support. The measure  $\mathbb{P}^*$  will be a key ingredient in the proof of Theorem 1.1.

**Proposition 4.7.** *There exists  $\mathbb{P}^* \in B_k(\mathbb{P})$  such that  $\text{Aff}(\text{supp}(\mathbb{P}^*)) = \mathbb{R}^d$  and  $NA(\mathbb{P}^*)$  holds.*

Proposition 4.7 is proved for general robust utility maximization problems in a one-period setup by Bayraktar and Zhou [2017][Lemma 2.2] and in the multi-period one by Blanchard and Carassus [2020][Theorem 3.29]. The proof in our context is very simple and is given for sake of completeness.

*Proof.* Denote the  $i$ th unit vector by  $e_i = (0, \dots, 0, 1, 0, \dots, 0)$  and consider the probability measure

$$\mathbb{P}^* := \frac{1}{2d}(\mathbb{P}_{-e_1} + \mathbb{P}_{e_1} + \mathbb{P}_{-e_2} + \dots + \mathbb{P}_{e_d}),$$

where  $\mathbb{P}_{\pm e_i} \in B_k(\mathbb{P})$  is defined as in Lemma 4.4 for  $\tilde{x} = \pm e_i$  and  $\alpha \in \left(0, \left(\frac{k}{C_{\mathbb{P}}+1}\right)^p\right)$ . By convexity of the Wasserstein distance (see e.g. [Villani, 2008, Theorem 4.8]) and  $\mathbb{P}_{\pm e_i} \in B_k(\mathbb{P})$  we conclude that  $\mathbb{P}^* \in B_k(\mathbb{P})$ . Furthermore

$$(12) \quad \text{supp}(\mathbb{P}^*) \supseteq \{\pm e_i : i = 1, \dots, d\},$$

so that  $\text{Aff}(\text{supp}(\mathbb{P}^*)) = \mathbb{R}^d$ . As (12) also implies that  $0 \in \text{int}(\text{conv}(\text{supp}(\mathbb{P}^*)))$ , we conclude that  $\text{NA}(\mathbb{P}^*)$  holds true, see Jacod and Shiryaev [1998][Theorem 3g].  $\square$

**Lemma 4.8.** *Let  $\mathbb{P}^*$  be as in Proposition 4.7. There exists  $\beta^* \in (0, 1]$  such that for any  $w \in \mathbb{R}^d \setminus \{0\}$ ,*

$$(13) \quad \mathbb{P}^* (\langle w, X \rangle < -\beta^*|w|) \geq \beta^*.$$

Condition 13 is called the quantitative no arbitrage condition and was proved in Blanchard and Carassus [2020][Proposition 3.35]. Again, the proof is easy in our context and is reported for sake of completeness.

*Proof.* Recall from Proposition 4.7 that (12) and  $\text{NA}(\mathbb{P}^*)$  holds. Then for every  $w \in \mathbb{R}^d$ ,

$$(14) \quad \begin{aligned} \langle w, X \rangle \geq 0 \quad \mathbb{P}^*\text{-a.s.} &\Rightarrow \langle w, X \rangle = 0 \quad \mathbb{P}^*\text{-a.s.} \\ &\Rightarrow w \in \{w \in \mathbb{R}^d : \langle w, x \rangle = 0, \forall x \in \text{supp}(\mathbb{P}^*)\} \Rightarrow w = 0. \end{aligned}$$

We introduce for  $n \geq 1$

$$(15) \quad A_n := \left\{ w \in \mathbb{R}^d : |w| = 1, \mathbb{P}^* \left( \langle w, X \rangle < -\frac{1}{n} \right) < \frac{1}{n} \right\}, \quad n_0 := \inf\{n \geq 1 : A_n = \emptyset\}$$

with the convention that  $\inf \emptyset = +\infty$ . We prove by contradiction that  $n_0 < \infty$ . Assume that  $n_0 = \infty$  and choose  $w_n \in A_n$  for all  $n \geq 1$ . By passing to a subsequence we can assume that  $w_n$  tends to some  $w^* \in \mathbb{R}^d$  with  $|w^*| = 1$ . Let  $B_n := \{\langle w_n, X \rangle < -1/n\}$ . Then  $\{\langle w^*, X \rangle < 0\} \subseteq \liminf_n B_n$  and Fatou's Lemma implies that

$$\mathbb{P}^* (\langle w^*, X \rangle < 0) \leq \mathbb{E}_{\mathbb{P}^*} [\mathbb{1}_{\liminf_{n \rightarrow \infty} B_n}] \leq \liminf_{n \rightarrow \infty} \mathbb{E}_{\mathbb{P}^*} [\mathbb{1}_{B_n}] = 0.$$

So  $\langle w^*, X \rangle \geq 0$   $\mathbb{P}^*$ -a.s. and (14) implies that  $w^* = 0$ , which contradicts  $|w^*| = 1$ . Thus  $n_0 < \infty$  and we can set  $\beta^* = 1/n_0 \in (0, 1]$ . By definition of  $A_{n_0}$ , (13) holds true.  $\square$

## 5. DISCUSSION OF THEOREM 1.1 AND ITS ASSUMPTIONS

Let us first single out (3) in the statement of Theorem 1.1.

**Assumption 5.1.** *There exists  $\underline{x} > 0$  and  $C > 0$  such that  $U(-\underline{x}) < 0$  and*

$$U(x) \geq -C(1 + |x|^p) \quad \text{for all } x \leq -\underline{x},$$

We now show that Assumption 5.1 is necessary to guarantee that  $u(k, w) > -\infty$ . For this, we first provide an equivalent formulation.

**Lemma 5.2.** *Assume that  $U$  is non-decreasing and  $\lim_{x \rightarrow -\infty} U(x) < 0$ . Assumption 5.1 is equivalent to the following condition: there exists  $C > 0$  such that*

$$(16) \quad U^-(x) \leq C(1 + |x|^p) \quad \text{for all } x \in \mathbb{R},$$

where  $U^-$  denotes the negative part of  $U$ .

If furthermore  $U$  is concave, Assumption 5.1 implies that there exists  $C > 0$  such that

$$(17) \quad |U(x)| \leq C(1 + |x|^p) \quad \text{for all } x \in \mathbb{R}.$$

*Proof.* Assume that Assumption 5.1 holds true. Let  $x \in \mathbb{R}$ . If  $U(x) \geq 0$ , then  $U^-(x) = 0$  and (16) holds true. Assume that  $U(x) \leq 0$ . Then  $U^-(x) = -U(x)$ . If  $x \leq -\underline{x}$ , then (16) holds true. Else, as  $U$  is non-decreasing  $U^-(x) = -U(x) \leq -U(-\underline{x})$ . So, (16) holds true for all  $x$  with the constant  $C - U(-\underline{x}) > 0$ .

Now, assume that (16) holds true. As  $\lim_{x \rightarrow -\infty} U(x) < 0$ , there exists  $\underline{x} > 0$  such that  $U(-\underline{x}) < 0$ . If  $x \leq -\underline{x}$ , then  $U(x) \leq U(-\underline{x}) < 0$  and Assumption 5.1 holds true, recalling that  $U^-(x) = -U(x)$ .

Now we prove (17). If  $U(z) \leq 0$ , then  $|U(z)| = -U(z) = U^-(z)$  and (16) implies (17). Let now  $y \in \mathbb{R}$  be the smallest number satisfying  $U(y) > 0$ . By continuity of  $U$  we have  $U(y) = 0$ . If  $z \leq y$ , the monotonicity of  $U$  implies that  $U(z) \leq U(y) = 0$  and (17) holds true. If  $z > y$ , setting  $x = z - y > 0$ , concavity and monotonicity of  $U$  clearly imply that

$$0 = U(y) \geq \frac{1}{2}(U(y+x) + U(y-x)) = \frac{1}{2}(U^+(y+x) - U^-(y-x)),$$

so that  $U^+(y+x) \leq U^-(y-x)$ . In particular (16) thus implies

$$\begin{aligned} U^+(z) &= U^+(y+x) \leq U^-(y-x) \\ &\leq C(1 + |y-x|^p) \\ &\leq C(1 + 2^{p-1}(|2y|^p + |y+x|^p)) \\ &\leq C(1 + 2^{p-1}|2y|^p + 2^{p-1})(1 + |y+x|^p) \\ &= \tilde{C}(1 + |z|^p), \end{aligned}$$

writing  $\tilde{C} := C(1 + 2^{p-1}|2y|^p + 2^{p-1}) > C$ . Thus, for all  $z \in \mathbb{R}$

$$|U(z)| = U^+(z) + U^-(z) \leq 2\tilde{C}(1 + |z|^p).$$

□

The following lemma shows that Assumption 5.1 is necessary to guarantee that  $u(k, w)$  is well-defined.

**Lemma 5.3.** *Assume that  $U$  is non-decreasing and  $\lim_{x \rightarrow -\infty} U(x) = -\infty$ . If Assumption 5.1 is not satisfied, then*

$$u(k, w) = \inf_{\mathbb{Q} \in B_k(\mathbb{F})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] = -\infty$$

for all  $w \in \mathbb{R}^d$  satisfying  $\langle w, \mathbf{1} \rangle \geq a$  and  $\mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] < \infty$ .

*Proof.* Let  $w \in \mathbb{R}^d$  such that  $\langle w, \mathbf{1} \rangle \geq a$  and  $\mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] < \infty$ . As  $\lim_{x \rightarrow -\infty} U(x) = -\infty$ , there exists some  $\underline{x} > 0$  such that  $U(-\underline{x}) < 0$  and if Assumption 5.1 is not satisfied, there exists a sequence  $(y_n)_n$  such that  $y_n \leq -\underline{x}$  and

$$(18) \quad U(y_n) \leq -n(1 + |y_n|^p) \quad \forall n \in \mathbb{N}.$$

We argue that  $(y_n)_n$  are necessarily such that  $\inf_n y_n = -\infty$ . Indeed, otherwise  $\inf_n y_n \geq c$  for some  $c \in \mathbb{R}$ . Then by monotonicity of  $U$ , we get that

$$U(c) \leq U(y_n) \leq -n(1 + |y_n|^p) \leq -n(1 + \underline{x}^p),$$

a contradiction for  $n \in \mathbb{N}$  large enough, so  $\inf_n y_n = -\infty$  follows. Taking a subsequence if necessary we can thus assume that  $(y_n)_n$  are negative, decreasing and  $\lim_{n \rightarrow \infty} y_n = -\infty$ . Next we define the sequence  $(x_n)_n$  of vectors in  $\mathbb{R}^d$  via

$$x_n = \frac{y_n}{|w|^2 x_0} w - \frac{w}{|w|^2},$$

recalling that  $w \neq 0, x_0 \neq 0$ . By definition we have  $x_0 + x_0 \langle w, x_n \rangle = y_n$  for all  $n \in \mathbb{N}$ . We now define

$$\alpha_{x_n} := \frac{k^p}{2^{p-1}(|x_n|^p + C_{\mathbb{P}}^p)} \leq \left( \frac{k}{C_{\mathbb{P}} + |x_n|} \right)^p, \quad \mathbb{P}_{x_n} := (1 - \alpha_{x_n})\mathbb{P} + \alpha_{x_n} \delta_{x_n}.$$

Then  $\mathbb{P}_{x_n} \in B_k(\mathbb{F})$  by Lemma 4.4 and we obtain

$$\begin{aligned} \inf_{\mathbb{Q} \in B_k(\mathbb{F})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] &\leq \mathbb{E}_{\mathbb{P}_{x_n}}[U(x_0 + x_0 \langle w, X \rangle)] \\ &\leq (1 - \alpha_{x_n})\mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] + \alpha_{x_n} U(y_n). \end{aligned}$$

Recalling (18) and noting that  $|x_n|^p \leq C(|y_n|^p + 1)$  for some  $C > 0$  we thus obtain

$$\begin{aligned} &\inf_{\mathbb{Q} \in B_k(\mathbb{F})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] \\ &\leq (1 - \alpha_{x_n})\mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] + \frac{k^p}{2^{p-1}(|x_n|^p + C_{\mathbb{P}}^p)} U(y_n) \\ &\leq \max(0, \mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)]) - \frac{k^p}{2^{p-1}(|x_n|^p + C_{\mathbb{P}}^p)} n(1 + |y_n|^p) \\ &\leq \max(0, \mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)]) - \frac{k^p}{2^{p-1}(C(|y_n|^p + 1) + C_{\mathbb{P}}^p)} n(1 + |y_n|^p) \\ &\rightarrow -\infty \end{aligned}$$

for  $n \rightarrow \infty$ , showing the claim.  $\square$

**Lemma 5.4.** *Assume that  $\lim_{x \rightarrow -\infty} U(x) = -\infty$ . Then*

$$\lim_{k \rightarrow \infty} u(k, w) = -\infty$$

for all  $w \in \mathbb{R}^d$  satisfying  $\langle w, \mathbf{1} \rangle \geq a$  and  $\mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] < \infty$ .

*Proof.* The proof is very similar to the proof of Lemma 5.3. Let  $w \in \mathbb{R}^d$  such that  $\langle w, \mathbf{1} \rangle \geq a$  and  $\mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] < \infty$ . Let for all  $k \in \mathbb{N}$

$$x_k := -k \text{sign}(x_0) \frac{w}{|w|^2},$$

recalling that  $w \neq 0$ . By definition we have  $x_0 + x_0 \langle w, x_k \rangle = x_0 - k|x_0|$ . As  $\langle w, \mathbf{1} \rangle \geq a$ ,  $|w| \geq \frac{a}{\sqrt{d}}$ , see (6) and  $|x_k| \leq \frac{k\sqrt{d}}{a}$ . We now define

$$\alpha_{x_k} := \frac{k^p}{2^{p-1}(k^p \frac{d^{\frac{p}{2}}}{a^p} + C_{\mathbb{P}}^p)} \leq \left( \frac{k}{C_{\mathbb{P}} + |x_k|} \right)^p, \quad \mathbb{P}_{x_k} := (1 - \alpha_{x_k})\mathbb{P} + \alpha_{x_k} \delta_{x_k}.$$

Then  $\mathbb{P}_{x_k} \in B_k(\mathbb{P})$  by Lemma 4.4 and we obtain

$$\begin{aligned} & \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] \\ & \leq \mathbb{E}_{\mathbb{P}_{x_k}}[U(x_0 + x_0 \langle w, X \rangle)] \\ & \leq (1 - \alpha_{x_k}) \mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)] + \alpha_{x_k} U(x_0 + x_0 \langle w, x_k \rangle) \\ & \leq \max(0, \mathbb{E}_{\mathbb{P}}[U(x_0 + x_0 \langle w, X \rangle)]) + \frac{k^p}{2^{p-1}(k^p \frac{d^{\frac{p}{2}}}{a^p} + C_{\mathbb{P}}^p)} U(x_0 - k|x_0|) \\ & \rightarrow -\infty \end{aligned}$$

for  $k \rightarrow \infty$ , showing the claim.  $\square$

We now continue our discussion with a few remarks on the assumptions on asymptotic elasticity. From an economics point of view, asymptotic elasticity can be interpreted as the ratio of the marginal utility  $U'(x)$  and average utility,  $\frac{U(x)}{x}$ . In the classical continuous time case of a fixed probability measure  $\mathbb{P}$ , a utility function with domain equal to  $\mathbb{R}$  must have a Reasonable Asymptotic Elasticity (RAE) at  $\pm\infty$ , i.e.

$$\liminf_{x \rightarrow -\infty} \frac{xU'(x)}{U(x)} > 1 \quad \text{and} \quad \limsup_{x \rightarrow \infty} \frac{xU'(x)}{U(x)} < 1,$$

in order to guarantee that the utility maximisation problem admits a solution, see Schachermayer [2001]. Having a RAE at  $+\infty$  means that the marginal utility should be smaller than the average utility for large positive financial positions, while a RAE at  $-\infty$  means that marginal utility should be larger than the average utility for large negative positions. In discrete time it is enough to postulate RAE at  $+\infty$  or  $-\infty$ , see Rásonyi and Stettner [2005]. RAE at  $+\infty$  or  $-\infty$  provides a control on  $U$  at  $+\infty$  or  $-\infty$  and concavity provides the other, see Lemma 5.7. In the case of multiple priors it is known that these conditions are not sufficient, see Blanchard and Carassus [2018] and Carassus and Ferhoun [2023].

We now state a number of implications of concavity and RAE. We start with the following result, which is similar to Carassus and Ferhoun [2023][Lemma 5.1].

**Lemma 5.5.** *Assume that  $U$  is concave, non-decreasing and differentiable and that there exists some positive constants  $\underline{x}$  and  $\bar{x}$  such that  $U(-\underline{x}) < 0$  and  $U(\bar{x}) > 0$ . Then for all  $x \in \mathbb{R}$  and  $\lambda \geq 1$*

$$(19) \quad U(\lambda x) \leq \lambda(U(x) + C),$$

where  $C := U^-(-\underline{x}) + U^+(\bar{x}) > 0$ .

*Proof.* Assume first that  $-\underline{x} \leq x \leq 0$ . Then  $\lambda x \leq 0 \leq \bar{x}$  and

$$U(\lambda x) \leq U(\bar{x}) = U^+(\bar{x}).$$

Then, as  $U$  is non-decreasing,  $C = -U(-\underline{x}) + U^+(\bar{x})$  and  $\lambda \geq 1$ , we obtain that

$$\lambda(U(x) + C) \geq \lambda(U(-\underline{x}) + C) = \lambda U^+(\bar{x}) \geq U^+(\bar{x})$$

and (19) holds true for  $-\underline{x} \leq x \leq 0$ .

Assume now that  $x \geq 0$ . Then, for all  $\lambda \geq 1$ ,  $-\underline{x} < 0 \leq x \leq \lambda x$ . As  $U$  is concave

$$\frac{U(\lambda x) - U(-\underline{x})}{\lambda x + \underline{x}} \leq \frac{U(x) - U(-\underline{x})}{x + \underline{x}}.$$

As  $\frac{\lambda x + \underline{x}}{x + \underline{x}} \leq \lambda$ ,  $U(x) \geq U(-\underline{x})$  and  $U(-\underline{x}) \leq 0$ ,

$$\begin{aligned} U(\lambda x) &\leq (U(x) - U(-\underline{x})) \frac{\lambda x + \underline{x}}{x + \underline{x}} + U(-\underline{x}) \\ &\leq \lambda(U(x) + U^-(-\underline{x})) \leq \lambda(U(x) + C) \end{aligned}$$

and (19) holds true for  $x \geq 0$ .

Now assume that  $x \leq -\underline{x}$ . Then, for all  $\lambda \geq 1$ ,  $\lambda x \leq x \leq -\underline{x}$ . As  $U$  is concave

$$\frac{U(-\underline{x}) - U(x)}{-\underline{x} - x} \leq \frac{U(-\underline{x}) - U(\lambda x)}{-\underline{x} - \lambda x}.$$

Thus,

$$U(\lambda x) - U(-\underline{x}) \leq (U(x) - U(-\underline{x})) \frac{-\underline{x} - \lambda x}{-\underline{x} - x}.$$

As  $\frac{-\underline{x} - \lambda x}{-\underline{x} - x} \geq \lambda$ ,  $U(x) \leq U(-\underline{x})$  and  $U(-\underline{x}) \leq 0$ ,

$$U(\lambda x) \leq \lambda(U(x) + U^-(-\underline{x})) \leq \lambda(U(x) + C)$$

and (19) holds true for  $x \leq -\underline{x}$ . □

Results similar to Lemmas 5.6 and 5.7 are proved in Carassus and Ferhoun [2023][Proposition 5.3] under some slightly different assumptions.

**Lemma 5.6.** *Assume that  $U$  is concave, non-decreasing and differentiable. Assume that  $AE_{+\infty}(U) < 1$  and  $\lim_{x \rightarrow \infty} U(x) > 0$ . Then there exists some constants  $\bar{x} > 0$  and  $\gamma < 1$  such that  $U(\bar{x}) > 0$  and for all  $x \geq \bar{x}$  and  $\lambda \geq 1$*

$$(20) \quad U(\lambda x) \leq \lambda^\gamma U(x).$$

Moreover,  $AE_{+\infty}(U)$  is the infimum of  $\gamma$  such that (20) holds true.

Assume that  $AE_{-\infty}(U) > 1$ . Then there exist some constants  $\underline{x} > 0$  and  $\gamma' > 1$  such that  $U(-\underline{x}) < 0$  and for all  $x \leq -\underline{x}$  and  $\lambda \geq 1$

$$(21) \quad U(\lambda x) \leq \lambda^{\gamma'} U(x).$$

Moreover,  $AE_{-\infty}(U)$  is the supremum of  $\gamma'$  such that (21) holds true.

*Proof.* The first assertion is proved in Kramkov and Schachermayer [1999][Lemma 6.3]. The second one can be proved similarly, see also Carassus and Ferhoun [2023][Proposition 5.3].  $\square$

**Lemma 5.7.** *Assume that  $U$  is concave, non-decreasing and differentiable. Assume that either  $AE_{+\infty}(U) < 1$  and  $\lim_{x \rightarrow \infty} U(x) > 0$  or  $AE_{-\infty}(U) > 1$ .*

*Then there exist some strictly positive constants  $\underline{x}, \bar{x}, \underline{\gamma}, \bar{\gamma}$  satisfying  $\underline{\gamma} \leq 1 \leq \bar{\gamma}$  and  $\underline{\gamma} < \bar{\gamma}$  and such that  $U(-\underline{x}) < 0$  and  $U(\bar{x}) > 0$  and for all  $x \in \mathbb{R}$  and  $\lambda \geq 1$*

$$(22) \quad U(\lambda x) \leq \lambda^{\underline{\gamma}}(U(x) + C)$$

$$(23) \quad U(\lambda x) \leq \lambda^{\bar{\gamma}}(U(x) + C),$$

where  $C := U^(-\underline{x}) + U^+(\bar{x}) > 0$ .

*Proof.* Assume that  $AE_{+\infty}(U) < 1$  and  $\lim_{x \rightarrow \infty} U(x) > 0$ . Then (20) holds true. Note that if  $x \leq \bar{x}$  and  $\lambda \geq 1$ , as  $U$  is non-decreasing, (20) implies that

$$(24) \quad U(\lambda x) \leq U(\lambda \bar{x}) \leq \lambda^{\bar{\gamma}} U(\bar{x}).$$

We claim that (20) and (24) above imply the following: for all  $x \in \mathbb{R}$  and  $\lambda \geq 1$

$$(25) \quad U(\lambda x) \leq \lambda^{\gamma}(U(x) + C).$$

If  $x \geq \bar{x}$ , (25) is a direct consequence of (20). If  $-\underline{x} \leq x \leq \bar{x}$ ,

$$U(x) + C \geq U(-\underline{x}) + C = U(\bar{x})$$

and (24) implies that

$$U(\lambda x) \leq \lambda^{\bar{\gamma}} U(\bar{x}) \leq \lambda^{\bar{\gamma}}(U(x) + C).$$

If  $x < -\underline{x} \leq 0$ , then as  $U$  is non-decreasing, we also have  $U(\lambda x) < 0$ . We distinguish two cases: (i)  $U(x) + C \geq 0$ : in this case (25) is trivial. (ii)  $U(x) + C < 0$ : recalling that  $\lambda \geq \lambda^{\gamma}$  we conclude from (19) that

$$U(\lambda x) \leq \lambda(U(x) + C) \leq \lambda^{\gamma}(U(x) + C).$$

This shows (25). Recalling (19), we can choose  $\underline{\gamma} = \gamma < 1 = \bar{\gamma}$  and (22) and (23) are proved.

Assume that  $AE_{-\infty}(U) > 1$ . Then (21) holds true and implies that for all  $x \leq -\underline{x}$  and  $\lambda \geq 1$

$$U(\lambda x) \leq \lambda^{\gamma'} U(x) \leq \lambda^{\gamma'}(U(x) + C).$$

Assume now that  $x \geq 0$ . Then, for all  $\lambda \geq 1$ ,  $-\underline{x} < 0 \leq x \leq \lambda x$ . As  $U$  is concave

$$\frac{U(\lambda x) - U(-\underline{x})}{\lambda x + \underline{x}} \leq \frac{U(x) - U(-\underline{x})}{x + \underline{x}}.$$

As  $\frac{\lambda x + \underline{x}}{x + \underline{x}} \leq \lambda \leq \lambda^{\gamma'}$  and  $U(-\underline{x}) \leq 0$ ,

$$U(\lambda x) \leq \lambda^{\gamma'}(U(x) + U^(-\underline{x})) \leq \lambda^{\gamma'}(U(x) + C).$$

Assume that  $-\underline{x} \leq x \leq 0$ . Then  $\lambda x \leq 0 \leq \bar{x}$  and

$$U(\lambda x) \leq U(\bar{x}) = U^+(\bar{x}).$$

As  $U$  is non-decreasing and  $\lambda^{\gamma'} \geq 1$ , we obtain that

$$\lambda^{\gamma'} (U(x) + C) \geq \lambda^{\gamma'} (U(-\underline{x}) + C) = \lambda^{\gamma'} U^+(\bar{x}) \geq U^+(\bar{x}).$$

So we have proved that for all  $x \in \mathbb{R}$  and  $\lambda \geq 1$

$$(26) \quad U(\lambda x) \leq \lambda^{\gamma'} (U(x) + C).$$

Recalling (19), we can choose  $\underline{\gamma} = 1 < \gamma' = \bar{\gamma}$  and (22) and (23) are proved.  $\square$

**Corollary 5.8.** *Assume that  $U$  is non decreasing and satisfies (22) or (23). Set  $\gamma = \underline{\gamma}$  if (22) holds true and  $\gamma = \bar{\gamma}$  if (23) holds true. Then there exists some  $D > 0$  such that*

$$(27) \quad U(x) \leq D(|x|^\gamma + 1) \quad \text{for all } x \in \mathbb{R}.$$

Assume that there exist constants  $\underline{x} > 0$  and  $\gamma' > 1$  such that  $U(-\underline{x}) < 0$  and for all  $x \leq -\underline{x}$  and  $\lambda \geq 1$

$$U(\lambda x) \leq \lambda^{\gamma'} U(x).$$

Then there exists some  $C' > 0$  such that

$$(28) \quad U(x) \leq -C'|x|^{\gamma'} \text{ for all } x \leq -\underline{x}.$$

However, the converses do not hold.

**Remark 5.9.** *If  $AE_{-\infty}(U) > 1$  and Assumption 5.1 holds true, i.e. that for all  $x \leq -\underline{x}$ ,  $U(x) \geq -C(1 + |x|^p)$ , Lemma 5.6 and (28) imply that  $\gamma' \leq p$  and taking supremum over  $\gamma'$  we conclude that*

$$1 < AE_{-\infty}(U) \leq p.$$

Thus in case (ii) of Theorem 1.1, we must have that  $AE_{-\infty}(U) \in (1, p]$ .

*Proof of Corollary 5.8.* We prove that (27) holds true. By assumption, for all  $x \in \mathbb{R}$  and  $\lambda \geq 1$

$$(29) \quad U(\lambda x) \leq \lambda^\gamma (U(x) + C)$$

where  $C := U^(-\underline{x}) + U^+(\bar{x}) > 0$ . Let  $x \geq \bar{x}$ . Choosing  $\lambda = x/\bar{x} \geq 1$

$$U(x) \leq \left(\frac{x}{\bar{x}}\right)^\gamma (U(\bar{x}) + C) = |x|^\gamma \frac{2U(\bar{x}) + U^-(\underline{x})}{\bar{x}^\gamma}.$$

If  $x \leq \bar{x}$  then  $U(x) \leq U(\bar{x})$ . Thus trivially, for all  $x \in \mathbb{R}$

$$U(x) \leq |x|^\gamma \frac{2U(\bar{x}) + U^-(\underline{x})}{\bar{x}^\gamma} + U(\bar{x}),$$

so that the first assertion is true for  $D := (2U(\bar{x}) + U^-(\underline{x}))/\bar{x}^\gamma + U(\bar{x})$ .

For the second assertion let  $x \leq -\underline{x}$ . Choosing  $\lambda = -x/\underline{x} \geq 1$

$$U(x) \leq \left(\frac{-x}{\underline{x}}\right)^{\gamma'} U(-\underline{x}) = -|x|^{\gamma'} \frac{U^-(\underline{x})}{\underline{x}^{\gamma'}}$$

and we set  $C' := U^-(\underline{x})/\underline{x}^{\gamma'}$ .  $\square$

6. EXISTENCE OF BOUNDED OPTIMAL SOLUTIONS OF  $u(x_0)$ 

We are now in a position to show that under assumptions made in Theorem 1.1, we can restrict to weights  $w$  in a compact set  $K \subseteq \mathbb{R}^d$  and that  $w_k = \arg \max_{w: \langle w, \mathbf{1} \rangle \geq a} u(k, w)$  exists. We start with an easy lemma.

**Lemma 6.1.** *Assume that there exists some  $K = K(x_0) \geq \frac{a}{\sqrt{d}} > 0$ , such that for all  $w \in \mathbb{R}^d$  with  $\langle w, \mathbf{1} \rangle \geq a$  and  $|w| \geq K$ ,*

$$(30) \quad \begin{aligned} u(k, w) &= \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w, X \rangle)] \\ &\leq \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w^u, X \rangle)] = u(k, w^u). \end{aligned}$$

Then

$$(31) \quad u(x_0) = \sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \leq K} u(k, w).$$

*Proof.* By (30)

$$\sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \geq K} u(k, w) \leq u(k, w^u) \leq \sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \leq K} u(k, w),$$

as  $|w^u| = \frac{a}{\sqrt{d}} \leq K$ . In conclusion (31) holds true.  $\square$

**6.1. Existence of bounded optimal solutions under RAE.** The following lemma holds true without assuming that  $U$  is concave.

**Proposition 6.2.** *Assume that  $U$  is non-decreasing and that there exists some strictly positive constants  $x^*$ ,  $C$ ,  $\underline{\gamma}$ ,  $\bar{\gamma}$  satisfying  $\underline{\gamma} \leq 1 \leq \bar{\gamma}$  and  $\underline{\gamma} < \bar{\gamma}$  and such that  $U(-x^*) \leq -(1 + C)$  and for all  $x \in \mathbb{R}$  and  $\lambda \geq 1$*

$$(32) \quad U(\lambda x) \leq \lambda^{\underline{\gamma}} (U(x) + C),$$

$$(33) \quad U(\lambda x) \leq \lambda^{\bar{\gamma}} (U(x) + C).$$

Finally assume that

$$\sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w^u, X \rangle)] < \infty.$$

Then there exists  $K = K(x_0) \geq \frac{a}{\sqrt{d}} > 0$ , such that

$$(34) \quad u(x_0) = \sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \leq K} u(k, w).$$

*Proof.* We show the following: there exists  $K = K(x_0) \geq \frac{a}{\sqrt{d}} > 0$ , such that for  $w \in \mathbb{R}^d$  with  $\langle w, \mathbf{1} \rangle \geq a$  and  $|w| \geq K$ ,

$$(35) \quad \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w, X \rangle)] \leq \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w^u, X \rangle)].$$

The statement of the proposition then follows from Lemma 6.1. The proof of (35) follows ideas of Rásonyi and Stettner [2005] in the monoprior case and [Blanchard and Carassus, 2018, Proposition 3.24] in the robust case.

Let

$$\mathbb{P}^* = \frac{1}{2d} (\mathbb{P}_{-e_1} + \mathbb{P}_{e_1} + \mathbb{P}_{-e_2} + \cdots + \mathbb{P}_{e_d}) \in B_k(\mathbb{P})$$

be as in Proposition 4.7, where we choose

$$\alpha := \alpha_{\pm e_i} = \left( \frac{k}{2(C_{\mathbb{P}} + 1)} \right)^p$$

for each  $\mathbb{P}_{\pm e_i}$  (recall the construction in Lemma 4.4). We calculate

$$\begin{aligned} L^* &:= \mathbb{E}_{\mathbb{P}^*} [U^+(1 + |X|)] = \frac{1}{2d} \sum_{i=1}^d 2((1 - \alpha) \mathbb{E}_{\mathbb{P}} [U^+(1 + |X|)] + \alpha U^+(2)) \\ (36) \quad &= (1 - \alpha) \mathbb{E}_{\mathbb{P}} [U^+(1 + |X|)] + \alpha U^+(2) < \infty. \end{aligned}$$

Indeed, using (27)

$$(37) \quad U^+(1 + |X|) \leq D(1 + |X|^{\underline{\gamma}}) \mathbf{1}_{\{U(1+|X|) \geq 0\}} \leq D(1 + |X|^{\underline{\gamma}})$$

and as  $\underline{\gamma} < 1 \leq p$  and  $\mathbb{E}_{\mathbb{P}}[|X|^{\underline{\gamma}}] \leq 1 + \mathbb{E}_{\mathbb{P}}[|X|^p] = 1 + C_{\mathbb{P}}^p < \infty$  and also  $U^+(2) < \infty$ .

As  $\underline{\gamma} < \bar{\gamma}$  there exists some  $0 < \eta < 1$  such that  $\underline{\gamma} < \eta \bar{\gamma}$ . Fix some  $x_0 \in \mathbb{R}$  and take  $w \in \mathbb{R}^d$  such that

$$|w| \geq K_0(x_0) := \max \left( 1, \frac{1}{|x_0|^{\frac{1}{\eta}}}, \frac{1}{\beta^*} \left( 1 + \frac{x^*}{|x_0|} \right), \left( \frac{1 + x^*}{\beta^*} \right)^{\frac{1}{1-\eta}} \right).$$

Let  $\underline{\lambda} := |x_0||w|$  and  $\bar{\lambda} := |x_0||w|^{\eta}$ . As  $w \geq 1$  and  $0 < \eta < 1$ ,

$$\underline{\lambda} = |x_0||w| \geq \bar{\lambda} = |x_0||w|^{\eta} \geq 1$$

as we have chosen  $|w| \geq 1/|x_0|^{1/\eta}$ .

We start with the estimation of  $U^+$ . Using that  $U$  is non-decreasing, applying the Cauchy-Schwarz inequality and then (32), we get that

$$\begin{aligned} U(x_0 + x_0 \langle w, X \rangle) &\leq U(|x_0| + |x_0||w||X|) \\ &= U \left( \underline{\lambda} \left( \frac{|x_0|}{\underline{\lambda}} + \frac{|x_0|}{\underline{\lambda}} |w||X| \right) \right) \\ &\leq \underline{\lambda}^{\underline{\gamma}} \left( U \left( \frac{|x_0|}{\underline{\lambda}} + \frac{|x_0|}{\underline{\lambda}} |w||X| \right) + C \right) \\ &= |x_0|^{\underline{\gamma}} |w|^{\underline{\gamma}} \left( U \left( \frac{1}{|w|} + |X| \right) + C \right) \\ &\leq |x_0|^{\underline{\gamma}} |w|^{\underline{\gamma}} (U^+(1 + |X|) + C), \end{aligned}$$

as  $|w| \geq 1$ . Thus

$$(38) \quad \mathbb{E}_{\mathbb{P}^*} \left[ \mathbf{1}_{\{U(x_0 + x_0 \langle w, X \rangle) \geq 0\}} U(x_0 + x_0 \langle w, X \rangle) \right] \leq |x_0|^{\underline{\gamma}} |w|^{\underline{\gamma}} (L^* + C),$$

For the estimation of  $U^-$  we introduce the event

$$(39) \quad B := \left\{ \frac{\text{sign}(x_0)}{|w|} \langle w, X \rangle < -\beta^* \right\},$$

where  $\text{sign}(x) = +1$  if  $x \geq 0$  and  $\text{sign}(x) = -1$  if  $x < 0$ .

Recall that  $\bar{\lambda} \geq 1$ . We start with the elementary inequality on  $B$

$$(40) \quad \begin{aligned} x_0 + x_0 \langle w, X \rangle &\leq \bar{\lambda} \left( \frac{1}{|w|^\eta} + |w|^{1-\eta} \frac{\text{sign}(x_0)}{|w|} \langle w, X \rangle \right) \\ &\leq \bar{\lambda} \left( \frac{1}{|w|^\eta} - \beta^* |w|^{1-\eta} \right). \end{aligned}$$

As  $|w| \geq 1$  and  $\eta > 0$ ,  $|w|^\eta \geq 1$  and  $\frac{1}{|w|^\eta} - \beta^* |w|^{1-\eta} \leq 1 - \beta^* |w|^{1-\eta}$ . As we have assumed that  $|w| \geq \left(\frac{1+x^*}{\beta^*}\right)^{\frac{1}{1-\eta}}$ , we get that

$$\frac{1}{|w|^\eta} - \beta^* |w|^{1-\eta} \leq -x^*.$$

So using that  $U$  is increasing, (40) and (33), we get that

$$\begin{aligned} U(x_0 + x_0 \langle w, X \rangle) \mathbf{1}_B &\leq U \left( \bar{\lambda} \left( \frac{1}{|w|^\eta} - \beta^* |w|^{1-\eta} \right) \right) \mathbf{1}_B \\ &\leq \bar{\lambda}^{\bar{\gamma}} \left( U \left( \frac{1}{|w|^\eta} - \beta^* |w|^{1-\eta} \right) + C \right) \mathbf{1}_B \\ &= |x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} \left( U \left( \frac{1}{|w|^\eta} - \beta^* |w|^{1-\eta} \right) + C \right) \mathbf{1}_B \\ &\leq |x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} (U(-x^*) + C) \mathbf{1}_B. \end{aligned}$$

In conclusion, recalling that  $U(-x^*) \leq -(1+C)$ , we obtain that

$$(41) \quad U(x_0 + x_0 \langle w, X \rangle) \mathbf{1}_B \leq -|x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} \mathbf{1}_B.$$

Now, we show that  $B \subseteq \{x_0 + x_0 \langle w, X \rangle \leq -x^*\}$ . Indeed, on  $B$  we have that

$$\begin{aligned} x_0 + x_0 \langle w, X \rangle &\leq |x_0| \left( 1 + |w| \frac{\text{sign}(x_0)}{|w|} \langle w, X \rangle \right) \\ &\leq |x_0| (1 - \beta^* |w|) \leq -x^*, \end{aligned}$$

as we have assumed that  $|w| \geq \frac{1}{\beta^*} \left( 1 + \frac{x^*}{|x_0|} \right)$ . This shows the claim. Clearly then  $B \subseteq \{U(x_0 + x_0 \langle w, X \rangle) < 0\}$  follows, as  $U$  is non decreasing and  $U(-x^*) < 0$ . So recalling (41)

$$U(x_0 + x_0 \langle w, X \rangle) \mathbf{1}_{\{U(x_0 + x_0 \langle w, X \rangle) < 0\}} \leq -|x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} \mathbf{1}_B.$$

Recalling (13) we have  $\mathbb{P}^*(B) > \beta^*$ . Combining these with the estimate (38), we have for  $|w| \geq K_0$

$$\mathbb{E}_{\mathbb{P}^*} [U(x_0 + x_0 \langle w, X \rangle)] \leq |x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} (L^* + C) - \beta^* |x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}}.$$

We note that  $\inf_{\mathbb{Q} \in \mathcal{B}_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w, X \rangle)] \leq \mathbb{E}_{\mathbb{P}^*} [U(x_0 + x_0 \langle w, X \rangle)]$  and

$$\inf_{\mathbb{Q} \in \mathcal{B}_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w^u, X \rangle)] \geq - \sup_{\mathbb{Q} \in \mathcal{B}_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w^u, X \rangle)].$$

Thus, (35) holds, if

$$(42) \quad \begin{aligned} & |x_0|^\gamma |w|^\gamma (L^* + C) - \frac{\beta^*}{2} |x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} \leq 0 \\ & -\frac{\beta^*}{2} |x_0|^{\bar{\gamma}} |w|^{\eta \bar{\gamma}} + \sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w^u, X \rangle)] \leq 0. \end{aligned}$$

As  $\eta \bar{\gamma} - \gamma > 0$ , this holds true if

$$|w| \geq K_1(x_0) := \max \left( \left( \frac{2(L^* + C)}{\beta^* |x_0|^{\bar{\gamma} - \gamma}} \right)^{\frac{1}{\eta \bar{\gamma} - \gamma}}, \left( \frac{2 \sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w^u, X \rangle)]}{\beta^* |x_0|^{\bar{\gamma}}} \right)^{\frac{1}{\eta \bar{\gamma}}} \right).$$

We define  $K(x_0) := \max \left( K_0(x_0), K_1(x_0), \frac{a}{\sqrt{d}} \right)$ . This shows (35) and concludes the proof.  $\square$

**Corollary 6.3.** *In the setting of Proposition 6.2 we have the following: if  $U$  is continuous,  $w_k = \arg \max_{w: \langle w, \mathbf{1} \rangle \geq a} u(k, w)$  exists and  $|w_k| \leq K$ .*

*Proof.* Take a sequence of maximizers  $(w_n)_n$  of

$$u(x_0) = \sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \leq K} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w, X \rangle)],$$

see (34). Then, as  $|w_n| \leq K$ , after taking a subsequence  $\lim_{n \rightarrow \infty} w_n = w^*$ ,  $|w^*| \leq K$  and  $\langle w^*, \mathbf{1} \rangle \geq a$ . Then, for any  $\mathbb{Q} \in B_k(\mathbb{P})$ ,

$$\begin{aligned} u(x_0) &= \limsup_{n \rightarrow \infty} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U(x_0 + x_0 \langle w_n, X \rangle)] \\ &\leq \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}} [U^+(x_0 + x_0 \langle w_n, X \rangle)] - \liminf_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w_n, X \rangle)] \\ &\leq \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}} [U^+(x_0 + x_0 \langle w_n, X \rangle)] - \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w^*, X \rangle)], \end{aligned}$$

applying the Fatou lemma for  $U^-$  which is continuous. Now using the Cauchy-Schwarz inequality and (27) we get that

$$\begin{aligned} U^+(x_0 + x_0 \langle w_n, X \rangle) &= (U(x_0 + x_0 \langle w_n, X \rangle)) \mathbf{1}_{U(x_0 + x_0 \langle w_n, X \rangle) \geq 0} \\ &\leq D(|x_0 + x_0 \langle w_n, X \rangle|^\gamma + 1) \mathbf{1}_{U(x_0 + x_0 \langle w_n, X \rangle) \geq 0} \\ &\leq D(2^{\gamma-1} |x_0|^\gamma (1 + |w_n|^\gamma |X|^\gamma) + 1) \\ &\leq D(2^{\gamma-1} |x_0|^\gamma (1 + K^\gamma |X|^\gamma) + 1). \end{aligned}$$

Recall that  $\gamma < 1 \leq p$  and that  $\mathbb{E}_{\mathbb{Q}}[|X|^p] < \infty$  for all  $\mathbb{Q} \in B_k(\mathbb{P})$ , see (7). So,

$$\mathbb{E}_{\mathbb{Q}}[|X|^\gamma] \leq 1 + \mathbb{E}_{\mathbb{Q}}[|X|^p] < \infty.$$

The dominated convergence theorem together with continuity of  $U^+$  shows that

$$\limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}} [U^+(x_0 + x_0 \langle w_n, X \rangle)] = \mathbb{E}_{\mathbb{Q}} [U^+(x_0 + x_0 \langle w^*, X \rangle)].$$

Thus,  $u(x_0) \leq \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w^*, X \rangle)]$ . Taking the infimum over all  $\mathbb{Q} \in B_k(\mathbb{P})$ , we find that

$$u(x_0) \leq \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w^*, X \rangle)] \leq u(x_0)$$

and  $w^*$  is an optimizer for  $u(x_0)$ .  $\square$

## 6.2. Existence of optimal solutions when $U$ is bounded from above.

**Lemma 6.4.** *Assume that  $U$  is non-decreasing, continuous and bounded from above, as well as  $\lim_{x \rightarrow -\infty} U(x) = -\infty$  and  $u(k, w^u) > -\infty$ . Then there exists  $K = K(x_0) \geq \frac{a}{\sqrt{d}} > 0$ , such that*

$$(43) \quad u(x_0) = \sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \leq K} u(k, w).$$

Furthermore,  $w_k = \arg \max_{w: \langle w, \mathbf{1} \rangle \geq a} u(k, w)$  exists and  $|w_k| \leq K$ .

*Proof.* We follow the same ideas as in the proof of Proposition 6.2. In particular we show that there exists  $K = K(x_0) \geq \frac{a}{\sqrt{d}} > 0$ , such that for  $w \in \mathbb{R}^d$  with  $\langle w, \mathbf{1} \rangle \geq a$  and  $|w| \geq K$ ,

$$(44) \quad \begin{aligned} u(k, w) &= \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] \\ &\leq \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w^u, X \rangle)] = u(k, w^u) \end{aligned}$$

and (43) follows again from Lemma 6.1. Take  $\mathbb{P}^* \in B_k(\mathbb{P})$  as in Proposition 4.7. Then (13) implies that

$$\mathbb{P}^*(\langle w, X \rangle < 0) > 0 \quad \text{for all } w \in \mathbb{R}^d \setminus \{0\}.$$

To prove (44), we argue by contradiction: assume that for all  $n \in \mathbb{N}$  there exists  $w_n \in \mathbb{R}^d$  with  $\langle w_n, \mathbf{1} \rangle \geq a$  and  $|w_n| \geq n$  and  $u(k, w_n) > u(k, w^u)$ . Then  $\lim_{n \rightarrow \infty} |w_n| = \infty$ . After taking a subsequence  $\lim_{n \rightarrow \infty} w_n/|w_n| = w^*$ , and  $|w^*| = 1$ . On the other hand,  $U$  is bounded from above by a constant  $C > 0$ , and so

$$\begin{aligned} \limsup_{n \rightarrow \infty} u(k, w_n) &= \limsup_{n \rightarrow \infty} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w_n, X \rangle)] \\ &\leq \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{P}^*}[U(x_0 + x_0 \langle w_n, X \rangle)] \\ &= \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{P}^*}[U(x_0 + x_0 |w_n| \langle \frac{w_n}{|w_n|}, X \rangle)] \\ &\leq \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{P}^*}[U(x_0 + x_0 |w_n| \langle \frac{w_n}{|w_n|}, X \rangle) \mathbf{1}_{\{x_0 \langle w^*, X \rangle < 0\}}] + C. \end{aligned}$$

Applying the reverse Fatou lemma to the last expression above and recalling that  $U$  is continuous and  $\lim_{x \rightarrow -\infty} U(x) = -\infty$  shows that

$$\limsup_{n \rightarrow \infty} u(k, w_n) = -\infty,$$

a contradiction to  $\limsup_{n \rightarrow \infty} u(k, w_n) \geq u(k, w^u) > -\infty$ .

We now prove that  $w_k$  exists. Take a sequence of maximizers  $(w_n)_n$  of

$$u(x_0) = \sup_{\langle w, \mathbf{1} \rangle \geq a, |w| \leq K} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)].$$

Then, as  $|w_n| \leq K$ , after taking a subsequence  $\lim_{n \rightarrow \infty} w_n = w^*$ , where  $|w^*| \leq K$  and  $\langle w^*, 1 \rangle \geq a$ . Then, for any  $\mathbb{Q} \in B_k(\mathbb{P})$ ,

$$\begin{aligned} u(x_0) &= \limsup_{n \rightarrow \infty} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w_n, X \rangle)] \\ &\leq \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w_n, X \rangle)] \leq \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w^*, X \rangle)], \end{aligned}$$

applying the reverse Fatou lemma and using that  $U$  is continuous as before. So taking the infimum over all  $\mathbb{Q} \in B_k(\mathbb{P})$ , we find that

$$u(x_0) \leq \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w^*, X \rangle)] \leq u(x_0)$$

and  $w^*$  is an optimizer for  $u(x_0)$ .  $\square$

## 7. PROOF OF THEOREM 1.1

To provide better intuition, it is helpful to rewrite the optimization problem  $u(k, w)$  in a different way.

**Lemma 7.1.** *Fix a Polish probability space  $(\Omega, \mathcal{F}, \mu)$  without atom. Then*

$$(45) \quad u(k, w) = \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)],$$

where  $X^\mu : \Omega \rightarrow \mathbb{R}^d$  is a measurable map, the notation  $X^\mu \sim \mathbb{P}$  means that  $X^\mu$  has law  $\mathbb{P}$  under  $\mu$  and  $B_k^\mu$  is the set of measurable maps  $Z^\mu : \Omega \rightarrow \mathbb{R}^d$  such that

$$\|Z^\mu\|_{L_\mu^p(\mathbb{R}^d)} := (\mathbb{E}_\mu[|Z^\mu|^p])^{1/p} \leq k.$$

*Proof.* We only show the “ $\geq$ ”-inequality; the “ $\leq$ ”-inequality follows very similarly. We fix  $\varepsilon > 0$ . Then there exists  $\mathbb{Q}_\varepsilon \in B_k(\mathbb{P})$  such that

$$u(k, w) = \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] \geq \mathbb{E}_{\mathbb{Q}_\varepsilon}[U(x_0 + x_0 \langle w, X \rangle)] - \varepsilon.$$

There exists some coupling  $\pi_\varepsilon \in \Pi(\mathbb{P}, \mathbb{Q}_\varepsilon)$  satisfying  $(\mathbb{E}_{\pi_\varepsilon}[|Y - X|^p])^{1/p} \leq k$  and thus

$$\begin{aligned} u(k, w) &\geq \mathbb{E}_{\pi_\varepsilon}[U(x_0 + x_0 \langle w, Y \rangle)] - \varepsilon \\ &= \mathbb{E}_{\pi_\varepsilon}[U(x_0 + x_0 \langle w, X + (Y - X) \rangle)] - \varepsilon. \end{aligned}$$

Applying the [Villani, Ch.1, the measurable isomorphism p.7]<sup>1</sup>, we can find a measurable map  $T_\varepsilon : (\Omega, \mu) \rightarrow (\mathbb{R}^d \times \mathbb{R}^d, (X, Y - X)_* \pi_\varepsilon)$  such that  $T_{\varepsilon*} \mu = (X, Y - X)_* \pi_\varepsilon$ . By a slight abuse of notation we call its first component  $X_\varepsilon^\mu$  and its second component  $Z_\varepsilon^\mu$ . By definition,  $X_\varepsilon^\mu$  has law  $\mathbb{P}$  under  $\mu$  and  $(\mathbb{E}_\mu[|Z_\varepsilon^\mu|^p])^{1/p} = (\mathbb{E}_{\pi_\varepsilon}[|Y - X|^p])^{1/p} \leq k$ . We then write

$$\begin{aligned} u(k, w) &\geq \mathbb{E}_{\pi_\varepsilon}[U(x_0 + x_0 \langle w, X + (Y - X) \rangle)] - \varepsilon \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X_\varepsilon^\mu + Z_\varepsilon^\mu \rangle)] - \varepsilon \\ &\geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] - \varepsilon. \end{aligned}$$

This concludes the proof.  $\square$

<sup>1</sup>As we have not assumed that  $(\mathbb{R}^d \times \mathbb{R}^d, (X, Y - X)_* \pi_\varepsilon)$  is atomless, the map  $T_\varepsilon$  is not necessary an isomorphism.

The following observation is crucial:

**Lemma 7.2.** Fix  $w \neq 0$  and  $\varepsilon > 0$ . Assume that  $U$  is non-decreasing. Then there exists an  $\varepsilon$ -optimizer  $(X^\varepsilon, Z^\varepsilon)$  of (45) such that  $X^\varepsilon \sim \mathbb{P}$  under  $\mu$ ,  $Z^\varepsilon \in B_k^\mu$  and

$$(46) \quad Z^\varepsilon = -\text{sign}(x_0)|Z^\varepsilon|\frac{w}{|w|}.$$

In particular

$$(47) \quad \begin{aligned} u(k, w) &= \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] \\ &= \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0||Z^\mu||w|)]. \end{aligned}$$

*Proof.* Take any  $\varepsilon$ -optimizer  $(X^\varepsilon, \tilde{Z}^\varepsilon)$  of (45), i.e.  $X^\varepsilon \sim \mathbb{P}$  under  $\mu$ ,  $\tilde{Z}^\varepsilon \in B_k^\mu$  and

$$\mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon + \tilde{Z}^\varepsilon \rangle)] \leq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] + \varepsilon.$$

Define

$$(48) \quad Z^\varepsilon := -\text{sign}(x_0)|\tilde{Z}^\varepsilon|\frac{w}{|w|}.$$

Then trivially  $|Z^\varepsilon| = |\tilde{Z}^\varepsilon|$  and (46) holds true. Moreover,

$$(49) \quad \begin{aligned} &\inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] + \varepsilon \\ &\geq \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon + \tilde{Z}^\varepsilon \rangle)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon \rangle + x_0 \langle w, \tilde{Z}^\varepsilon \rangle)] \\ &\stackrel{\text{(CS)}}{\geq} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon \rangle - |x_0||w||\tilde{Z}^\varepsilon|)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon \rangle - |x_0|\langle w, \frac{w}{|w|}|\tilde{Z}^\varepsilon|)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon \rangle + x_0 \langle w, Z^\varepsilon \rangle)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon + Z^\varepsilon \rangle)]. \end{aligned}$$

This shows that  $(X^\varepsilon, Z^\varepsilon)$  is also an  $\varepsilon$ -minimizer. Next we show (47). First, as  $\tilde{Z}^\varepsilon \in B_k^\mu$  (49) shows that

$$\begin{aligned} &\inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] + \varepsilon \\ &\geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0||w||Z^\mu|)]. \end{aligned}$$

So letting  $\varepsilon \rightarrow 0$  yields

$$\begin{aligned} &\inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] \\ &\geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0||w||Z^\mu|)]. \end{aligned}$$

Next we show

$$\begin{aligned} &\inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0||w||Z^\mu|)] \\ &\geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)]. \end{aligned}$$

This follows by the same argument as above: we choose an  $\varepsilon$ -optimizer  $(X^\varepsilon, Z^\varepsilon)$  for the left hand side and define  $Z^*$  as in (48). This yields

$$\begin{aligned} & \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0| |w| |Z^\mu|)] + \varepsilon \\ & \geq \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon \rangle - |x_0| |w| |Z^\varepsilon|)] \\ & = \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\varepsilon + Z^* \rangle)] \\ & \geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)]. \end{aligned}$$

Taking  $\varepsilon \rightarrow 0$  concludes the proof.  $\square$

Recall

$$w_k := \arg \max_{w: \langle w, \mathbf{1} \rangle \geq a} u(k, w).$$

The following proposition is central for the proof of Theorem 1.1.

**Proposition 7.3.** *Let  $U$  be concave, non-decreasing, differentiable, non-constant and let Assumption 5.1 holds. Fix a compact subset  $K \subseteq \{w \in \mathbb{R}^d : \langle w, \mathbf{1} \rangle \geq a\}$ . Then the following holds: for any  $\delta > 0$  there exists  $k_0 \in \mathbb{N}$  such that for all  $w \in K$  satisfying  $|w| - |w^u| \geq \delta$  and for all  $k \geq k_0$  we have*

$$(50) \quad u(k, w) = \inf_{\mathbb{Q} \in B_k(\mathbb{P})} E_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)]$$

$$(51) \quad < \inf_{\mathbb{Q} \in B_k(\mathbb{P})} E_{\mathbb{Q}}[U(x_0 + x_0 \langle w^u, X \rangle)] = u(k, w^u).$$

*Proof.* Recall that by Lemma 7.2 we can rewrite the claim as

$$\begin{aligned} & \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0| |Z^\mu| |w|)] \\ & = \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] \\ & < \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w^u, X \rangle)] \\ & = \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w^u, X^\mu \rangle - |x_0| |Z^\mu| |w^u|)]. \end{aligned}$$

Let  $(X^u(k), Z^u(k))$  be an  $\varepsilon$ -optimizer for  $u(k, w^u)$ , so that by definition

$$\begin{aligned} & \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle - |x_0| |Z^\mu| |w|)] \\ & \quad - \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w^u, X^\mu \rangle - |x_0| |Z^\mu| |w^u|)] \\ & \leq \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^u(k) \rangle - |x_0| |Z^u(k)| |w|) - U(x_0 + x_0 \langle w^u, X^u(k) \rangle - |x_0| |Z^u(k)| |w^u|)] + \varepsilon. \end{aligned}$$

To simplify notation we set

$$A_k(w) := x_0 + x_0 \langle w, X^u(k) \rangle - |x_0| |Z^u(k)| |w|.$$

Fix  $\delta > 0$ . Using this notation, the claim follows if we show that there exists  $k_0 \in \mathbb{N}$  such that for all  $w \in K$  satisfying  $|w| - |w^u| \geq \delta$  and for all  $k \geq k_0$  we have

$$(52) \quad \mathbb{E}_\mu[U(A_k(w)) - U(A_k(w^u))] < -\varepsilon.$$

Let

$$C_k := x_0 + x_0 \langle w^u, X^u(k) \rangle + x_0 \langle |w^u| \frac{w^u - w}{\delta}, X^u(k) \rangle.$$

By concavity of  $U$

$$(53) \quad U(A_k(w)) - U(A_k(w^u)) \leq (A_k(w) - A_k(w^u))U'(A_k(w^u))$$

$$(54) \quad U(A_k(w^u)) - U(C_k) \geq (A_k(w^u) - C_k)U'(A_k(w^u)).$$

Recall that  $|w| - |w^u| \geq \delta$  by assumption, and thus

$$(55) \quad \begin{aligned} A_k(w^u) - C_k &= -|x_0||Z^u(k)||w^u| + x_0 \langle |w^u| \frac{w - w^u}{\delta}, X^u(k) \rangle \\ &= \frac{|w^u|}{\delta} (-|x_0||Z^u(k)|\delta + x_0 \langle w - w^u, X^u(k) \rangle) \\ A_k(w) - A_k(w^u) &= x_0 \langle w - w^u, X^u(k) \rangle - |x_0||Z^u(k)|(|w| - |w^u|) \\ &\leq x_0 \langle w - w^u, X^u(k) \rangle - |x_0||Z^u(k)|\delta = \frac{\delta}{|w^u|} (A_k(w^u) - C_k). \end{aligned}$$

As  $U' \geq 0$  it follows from (53), (55) and (54) that

$$\begin{aligned} U(A_k(w)) - U(A_k(w^u)) &\leq \frac{\delta}{|w^u|} (A_k(w^u) - C_k)U'(A_k(w^u)) \\ &\leq \frac{\delta}{|w^u|} (U(A_k(w^u)) - U(C_k)). \end{aligned}$$

To show (52) it thus remains to argue that the expectation of the last expression becomes small for large  $k \in \mathbb{N}$ ; in fact we will show that it diverges to  $-\infty$ . For this we first note that by definition of  $Z^u(k)$  and Lemma 7.2 we have

$$\begin{aligned} \mathbb{E}_\mu[U(A_k(w^u))] &\leq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w^u, X^\mu \rangle - |x_0||Z^\mu||w^u|)] + \varepsilon \\ &= \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_\mathbb{Q}[U(x_0 + x_0 \langle w^u, X \rangle)] + \varepsilon. \end{aligned}$$

Define now

$$\mathbb{Q}^* := \left( x \mapsto x - k \operatorname{sign}(x_0) \frac{w^u}{|w^u|} \right)_* \mathbb{P}.$$

Using the coupling

$$\pi = \left( x \mapsto \left( x, x - k \operatorname{sign}(x_0) \frac{w^u}{|w^u|} \right) \right)_* \mathbb{P} \in \Pi(\mathbb{P}, \mathbb{Q}^*)$$

yields the upper bound

$$\mathcal{W}_p(\mathbb{P}, \mathbb{Q}^*) \leq \mathbb{E}_\mathbb{P} \left[ \left| k \frac{w^u}{|w^u|} \right|^p \right]^{1/p} = k$$

and thus  $\mathbb{Q}^* \in B_k(\mathbb{P})$ . In particular

$$\begin{aligned} \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_\mathbb{Q}[U(x_0 + x_0 \langle w^u, X \rangle)] &\leq \mathbb{E}_{\mathbb{Q}^*}[U(x_0 + x_0 \langle w^u, X \rangle)] \\ &= \mathbb{E}_\mathbb{P}[U(x_0 + x_0 \langle w^u, X \rangle - k|x_0||w^u|)] \\ &\leq U(x_0 + x_0 \mathbb{E}_\mathbb{P}[\langle w^u, X \rangle] - k|x_0||w^u|) \\ &\leq U(x_0 + |x_0||w^u| \mathbb{E}_\mathbb{P}[|X|] - k|x_0||w^u|) \rightarrow -\infty \end{aligned}$$

for  $k \rightarrow \infty$ , using Jensen's and Cauchy-Schwarz inequalities,  $\mathbb{E}_{\mathbb{P}}[|X|] \leq 1 + \mathbb{E}_{\mathbb{P}}[|X|^p] < \infty$ , see (7) and  $\lim_{x \rightarrow -\infty} U(x) = -\infty$ . This last limit follows from the fact that  $U$  is concave, non-decreasing and non-constant. In conclusion

$$(56) \quad \lim_{k \rightarrow \infty} \mathbb{E}_{\mu}[U(A_k(w^u))] = -\infty.$$

On the other hand, as we assume that  $w \in K$  and that  $U$  is non-decreasing, the Cauchy-Schwarz, triangle inequalities and (17) yield

$$(57) \quad \begin{aligned} \mathbb{E}_{\mu}[U(C_k)] &\geq \mathbb{E}_{\mu}[U(-|x_0| - |x_0| |w^u + |w^u| \frac{w^u - w}{\delta}|X^u(k)|)] \\ &\geq \mathbb{E}_{\mu}[U(-|x_0| - |x_0| |w^u| (1 + \frac{|w^u| + |w|}{\delta}) |X^u(k)|)] \\ &\geq \mathbb{E}_{\mu}[U(-|x_0| - \tilde{C} |X^u(k)|)] \\ &\geq -C \left(1 + 2^{p-1} (|x_0|^p + \tilde{C}^p \mathbb{E}_{\mu}[|X^u(k)|^p])\right) > -\infty \end{aligned}$$

for a constant  $\tilde{C} = \tilde{C}(x_0, \delta, K) > 0$ , where the last inequality follows from  $X^u(k) \sim \mathbb{P}$  under  $\mu$ , i.e.

$$\mathbb{E}_{\mu}[|X^u(k)|^p] = \mathbb{E}_{\mathbb{P}}[|X|^p] < \infty.$$

Combining (56)-(57) shows that

$$\lim_{k \rightarrow \infty} \mathbb{E}_{\mu}[U(A_k(w^u)) - U(C_k)] = -\infty,$$

as claimed.  $\square$

**Corollary 7.4.** *In the setting of Proposition 7.3 assume furthermore that  $w^u \in K$ . Fix  $\delta > 0$  and let  $k_0 \in \mathbb{N}$  as in Proposition 7.3. If  $w_k$  exists and belongs to  $K$ , we have that*

$$(58) \quad 0 \leq |w_k| - |w^u| < \delta \quad \text{for all } k \geq k_0.$$

and

$$\lim_{k \rightarrow \infty} w_k = w^u.$$

*Proof.* Fix  $\delta > 0$  and take  $k \geq k_0$ . Assume that  $|w_k| - |w^u| \geq \delta$ . Then as  $w_k \in K$  (50) implies  $u(k, w_k) < u(k, w^u)$ , which contradicts the definition of  $w_k$ . Thus,  $|w_k| - |w^u| < \delta$ . This concludes the proof of (58) recalling (6). Thus,  $\lim_{k \rightarrow \infty} |w_k| = |w^u|$ . As  $(w_k)_k \subseteq K$ , there exists a subsequence of  $(w_k)_k$  (which we still denote by  $(w_k)_k$ ), converging to some  $w^*$ . By continuity of  $w \mapsto |w|$  we have

$$|w^*| = \lim_{k \rightarrow \infty} |w_k| = \lim_{k \rightarrow \infty} |w_k| = |w^u|.$$

As  $w^u$  is the unique element of  $K$  having norm  $|w^u|$  (recall (6)), we obtain  $w^* = w^u$  and the claim follows.  $\square$

*Proof. of Theorem 1.1* We assume that  $U : \mathbb{R} \rightarrow \mathbb{R}$  is non-decreasing, non-constant, differentiable, concave and satisfies Assumption 5.1. Let  $w \in \mathbb{R}^d$  be such that  $\langle w, 1 \rangle \geq a$ . Then (16) implies that

$$\begin{aligned} u(k, w) &= \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] \geq - \sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U^-(x_0 + x_0 \langle w, X \rangle)] \\ &\geq -C \sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[1 + |x_0 + x_0 \langle w, X \rangle|^p]. \end{aligned}$$

Using the Cauchy-Schwarz inequality, we get that

$$1 + |x_0 + x_0 \langle w, X \rangle|^p \leq 1 + |x_0|^p (1 + |w| |X|)^p \leq 1 + |x_0|^p 2^{p-1} (1 + |w|^p |X|^p).$$

So, using (7) we obtain

$$(59) \quad \begin{aligned} u(k, w) &\geq - \sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U^-(x_0 + x_0 \langle w, X \rangle)] \\ &\geq -C \left( 1 + |x_0|^p 2^{p-1} (1 + |w|^p \sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[|X|^p]) \right) > -\infty. \end{aligned}$$

Assume first that  $U$  is bounded from above. We can apply Lemma 6.4 as  $U$  is continuous,  $\lim_{x \rightarrow -\infty} U(x) = -\infty$  and  $u(k, w^u) > -\infty$ , see (59). Thus  $w_k$  exists,  $|w_k| \leq K$  and Corollary 7.4 applies with the compact  $\{w \in \mathbb{R}^d : \langle w, 1 \rangle \geq a, |w| \leq K\}$  (recall that  $K \geq \frac{a}{\sqrt{d}}$ .)

Assume now that either  $\text{AE}_{+\infty}(U) < 1$  and  $\lim_{x \rightarrow \infty} U(x) > 0$  or  $\text{AE}_{-\infty}(U) > 1$ . We can apply Lemma 5.7 and there exists some strictly positive constants  $\underline{x}, \bar{x}, \underline{\gamma}, \bar{\gamma}$  satisfying  $\underline{\gamma} \leq 1 \leq \bar{\gamma}$  and  $\underline{\gamma} < \bar{\gamma}$  and such that  $U(-\underline{x}) < 0$  and  $U(\bar{x}) > 0$ , (22) and (23) hold true with  $C := U^-(\underline{x}) + U^+(\bar{x}) > 0$ . As  $\lim_{x \rightarrow -\infty} U(x) = -\infty$ , there exists some  $\lambda^* \geq 1$  such that  $U(-\lambda^* \underline{x}) \leq -(1 + C)$ . As

$$\sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}} [U^-(x_0 + x_0 \langle w^u, X \rangle)] < \infty,$$

see (59), we can apply Corollary 6.3 with  $x^* = \lambda^* \underline{x}$ . We obtain that  $w_k$  exists and  $|w_k| \leq K$  and again Corollary 7.4 applies with the same compact as above.  $\square$

#### APPENDIX A. FURTHER RESULTS ON $u(k, w)$

In this appendix we dive deeper into the analysis of the max-min problem  $u(k, w)$ ; in particular we are interested in understanding when the infimum are attained. It turns out that this is the case under some slightly stricter growth assumptions than those stated in Assumption 5.1 together with a constraint on the measure  $\mathbb{P}$ .

**Assumption A.1.** *Assume that  $p > 1$  and that there exist  $C > 0$  and  $1 \leq r < p$  such that*

$$U^-(x) \leq C(1 + |x|^r) \quad \text{for all } x \in \mathbb{R}.$$

As before we conclude by concavity of  $U$  that Assumption A.1 implies

$$|U(x)| \leq C(1 + |x|^r) \quad \text{for all } x \in \mathbb{R}.$$

**Lemma A.2.** *Fix  $w \neq 0$ . Under Assumption A.1 there exists an optimizer  $\mathbb{Q}^*$  of (2), i.e.*

$$u(k, w) = \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] = \mathbb{E}_{\mathbb{Q}^*}[U(x_0 + x_0 \langle w, X \rangle)].$$

Moreover, there exists an optimizer  $(X^*, Z^*)$  of (45), i.e.

$$u(k, w) = \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] = \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + Z^* \rangle)],$$

such that  $X^* \sim \mathbb{P}$  under  $\mu$ ,  $Z^* \in B_k^\mu$  and

$$(60) \quad Z^* = -\text{sign}(x_0) |Z^*| \frac{w}{|w|}.$$

*Proof.* We first prove existence of optimizers for (2). Take a sequence  $(\mathbb{Q}_n)_n$  in  $B_k(\mathbb{P})$  of  $1/n$ -optimizers of  $u(k, w)$ , i.e.

$$u(k, w) \leq \mathbb{E}_{\mathbb{Q}_n}[U(x_0 + x_0 \langle w, X \rangle)] \leq u(k, w) + \frac{1}{n}.$$

As  $B_k(\mathbb{P})$  is compact wrt.  $W_r$  by Lemma A.3 below, there exists a convergent subsequence, which we still call  $(\mathbb{Q}_n)_n$ , converging to some  $\mathbb{Q}^* \in B_k(\mathbb{P})$  in  $W_r$ . By Assumption A.1 and Cauchy-Schwarz inequality

$$\begin{aligned} |U(x_0 + x_0 \langle w, X \rangle)| &\leq C(1 + |x_0 + x_0 \langle w, X \rangle|^r) \\ &\leq C(1 + |x_0|^r 2^{r-1}(1 + |w|^r |X|^r)) \\ &\leq \tilde{C}(1 + |X|^r) \end{aligned}$$

and [Villani, 2008, Def. 6.8 p108] shows that

$$u(k, w) = \lim_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_n}[U(x_0 + x_0 \langle w, X \rangle)] = \mathbb{E}_{\mathbb{Q}^*}[U(x_0 + x_0 \langle w, X \rangle)].$$

This concludes the proof of existence of optimizers for (2).

Using the same arguments as in the proof of Lemma 7.1 there exists  $(X^*, \tilde{Z})$  such that  $X^* \sim \mathbb{P}$  under  $\mu$ ,  $\tilde{Z} \in B_k^\mu$  and

$$(61) \quad u(k, w) = \mathbb{E}_{\mathbb{Q}^*}[U(x_0 + x_0 \langle w, X \rangle)] = \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + \tilde{Z} \rangle)].$$

Define

$$Z^* := -\text{sign}(x_0) |\tilde{Z}| \frac{w}{|w|}.$$

Then trivially  $|Z^*| = |\tilde{Z}|$  and (60) holds true. Using Lemma 7.1 and (61), we get that

$$\begin{aligned} u(k, w) &= \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + \tilde{Z} \rangle)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle + x_0 \langle w, \tilde{Z} \rangle)] \\ &\stackrel{\text{(CS)}}{\geq} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle - |x_0| |w| |\tilde{Z}|)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle - |x_0| \langle w, \frac{w}{|w|} |\tilde{Z}| \rangle)] \\ &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle + x_0 \langle w, Z^* \rangle)] \\ &\geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)]. \end{aligned}$$

This shows that  $(X^*, Z^*)$  is also a minimizer and concludes the proof.  $\square$

**Lemma A.3.** *Let  $1 \leq r < p$ . Then the  $p$ -Wasserstein ball  $B_k(\mathbb{P})$  is compact wrt.  $W_r$ .*

*Proof.* Recall that (7) shows that  $\sup_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[|X|^p]^{1/p} < \infty$ . Then an application of Prokhorov's theorem shows that  $B_k(\mathbb{P})$  is weakly precompact. Hence, for every sequence of measures  $(\mathbb{Q}_n)_n$  in  $B_k(\mathbb{P})$  there exists a subsequence, which we also call  $(\mathbb{Q}_n)_n$  and a measure  $\mathbb{Q}$  such that  $\mathbb{Q}_n$  converges weakly to  $\mathbb{Q}$ . As  $W_p$  is weakly lower semicontinuous (see [Villani, 2008, Rem. 6.12, p.109]), this implies

$\mathbb{Q} \in B_k(\mathbb{P})$ . It remains to show that  $\mathbb{Q}_n$  converge to  $\mathbb{Q}$  in  $W_r$ . By [Villani, 2008, Definition 6.8 and Theorem 6.9] it is sufficient to show that  $\lim_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_n}[|X|^r] = \mathbb{E}_{\mathbb{Q}}[|X|^r]$ . For this we note that

$$\begin{aligned}
\lim_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_n}[|X|^r] &= \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} (\mathbb{E}_{\mathbb{Q}_n}[|X|^r \mathbf{1}_{\{|X| \leq l\}}] + \mathbb{E}_{\mathbb{Q}_n}[|X|^r \mathbf{1}_{\{|X| > l\}}]) \\
&\leq \limsup_{l \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_n}[(|X| \wedge l)^r] + \limsup_{l \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{E}_{\mathbb{Q}_n}[|X|^r \mathbf{1}_{\{|X| > l\}}] \\
&\leq \limsup_{l \rightarrow \infty} \mathbb{E}_{\mathbb{Q}}[(|X| \wedge l)^r] + \limsup_{l \rightarrow \infty} \sup_{n \in \mathbb{N}} \mathbb{E}_{\mathbb{Q}_n}[|X|^r \mathbf{1}_{\{|X| > l\}}] \\
&\leq \mathbb{E}_{\mathbb{Q}}[|X|^r] + \limsup_{l \rightarrow \infty} \sup_{n \in \mathbb{N}} \left( \mathbb{E}_{\mathbb{Q}_n}[|X|^p] \right)^{r/p} \mathbb{Q}_n(|X| > l)^{(p-r)/p} \\
&\leq \mathbb{E}_{\mathbb{Q}}[|X|^r] + \limsup_{l \rightarrow \infty} \sup_{n \in \mathbb{N}} \left( \mathbb{E}_{\mathbb{Q}_n}[|X|^p] \right)^{r/p} \left( \frac{\mathbb{E}_{\mathbb{Q}_n}[|X|^p]}{l^p} \right)^{(p-r)/p} \\
&\leq \mathbb{E}_{\mathbb{Q}}[|X|^r] + \limsup_{l \rightarrow \infty} \left( \frac{\sup_{n \in \mathbb{N}} \mathbb{E}_{\mathbb{Q}_n}[|X|^p]}{l^{p-r}} \right) \\
&\leq \mathbb{E}_{\mathbb{Q}}[|X|^r] + \limsup_{l \rightarrow \infty} \left( \frac{(C_{\mathbb{P}} + k)^p}{l^{p-r}} \right) \\
&= \mathbb{E}_{\mathbb{Q}}[|X|^r],
\end{aligned}$$

where we have used weak convergence for the second inequality, the dominated convergence theorem and Hölder's inequality for the third inequality, Markov's inequality for the fourth one and (7) for the last one. Furthermore,

$$\begin{aligned}
\lim_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_n}[|X|^r] &= \lim_{l \rightarrow \infty} \lim_{n \rightarrow \infty} (\mathbb{E}_{\mathbb{Q}_n}[|X|^r \mathbf{1}_{\{|X| \leq l\}}] + |X|^r \mathbf{1}_{\{|X| > l\}}]) \\
&\geq \liminf_{l \rightarrow \infty} \liminf_{n \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_n}[(|X| \wedge l)^r] \\
&= \mathbb{E}_{\mathbb{Q}}[|X|^r].
\end{aligned}$$

This concludes the proof.  $\square$

Under slightly stricter assumptions we can strengthen Lemma 7.2 in the following way:

**Lemma A.4.** *Fix  $w \neq 0$  and assume that  $U$  is strictly increasing and satisfies Assumption A.1. Then any optimizer  $(X^*, Z^*)$  of (45) satisfies*

$$Z^* = -\text{sign}(x_0) |Z^*| \frac{w}{|w|}.$$

Furthermore  $\|Z^*\|_{L^p_{\mu}(\mathbb{R}^d)} = k$  holds for any optimizer of (45).

*Proof.* Take an optimizer  $(X^*, Z^*)$  of (45), which exists by Lemma A.2, define

$$A := \left\{ Z^* \neq -\text{sign}(x_0) |Z^*| \frac{w}{|w|} \right\}$$

and assume that  $\mu(A) > 0$ . Recall from the Cauchy-Schwarz inequality that we have

$$(62) \quad x_0 \langle w, z \rangle > -|x_0| |w| |z| = x_0 \langle w, -\text{sign}(x_0) |z| \frac{w}{|w|} \rangle$$

for all  $z \in \mathbb{R}^d$  such that  $z \neq -\text{sign}(x_0)|z|w/|w|$ . Similarly to the proof of Lemma A.2 we thus have

$$\begin{aligned}
& \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + Z^* \rangle)] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + Z^* \rangle) \mathbf{1}_A] + \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + Z^* \rangle) \mathbf{1}_{A^c}] \\
&> \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* - \text{sign}(x_0) |Z^*| \frac{w}{|w|} \rangle) \mathbf{1}_A] + \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + Z^* \rangle) \mathbf{1}_{A^c}] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* - \text{sign}(x_0) |Z^*| \frac{w}{|w|} \rangle)] \\
&\geq \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)],
\end{aligned}$$

where we used (62) and that  $U$  is strictly increasing for the strict inequality and also that

$$\| -\text{sign}(x_0) |Z^*| \frac{w}{|w|} \|_{L_\mu^p(\mathbb{R}^d)} = \|Z^*\|_{L_\mu^p(\mathbb{R}^d)} \leq k.$$

This leads to a contradiction to  $\mu(A) > 0$ . Furthermore, from the fact that  $U$  is strictly increasing it is clear that  $0 \neq Z^*$  for any optimizer  $(X^*, Z^*)$ . Indeed, otherwise for any  $\delta > 0$

$$\begin{aligned}
u(k, w) &= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle)] \\
&> \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle - |x_0| \delta)] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle - x_0 \langle w, \delta \frac{w}{|w|^2} \text{sign}(x_0) \rangle)].
\end{aligned}$$

As  $(X^* - \delta \frac{w}{|w|^2} \text{sign}(x_0))_* \mu \in B_k(\mathbb{P})$  for  $\delta > 0$  small enough, we get a contradiction. Lastly, if  $\|Z^*\|_{L_\mu^p(\mathbb{R}^d)} < k$  then we have

$$\begin{aligned}
u(k, w) &= \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu + Z^\mu \rangle)] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* + Z^* \rangle)] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle + x_0 \langle w, -\text{sign}(x_0) |Z^*| \frac{w}{|w|} \rangle)] \\
&= \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle - |x_0| |Z^*| |w|)] \\
&> \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^* \rangle - |x_0| \frac{k}{\|Z^*\|_{L_\mu^p(\mathbb{R}^d)}} |Z^*| |w|)].
\end{aligned}$$

and

$$\left\| \frac{k}{\|Z^*\|_{L_\mu^p(\mathbb{R}^d)}} |Z^*| \right\|_{L_\mu^p(\mathbb{R}^d)} = k,$$

leading to another contradiction. This concludes the proof.  $\square$

**Lemma A.5.** *Fix  $w \neq 0$  and assume that  $U$  is strictly increasing and satisfies Assumption A.1. Assume also that  $\langle \frac{w}{|w|}, X \rangle_* \mathbb{P}$  is absolutely continuous wrt. the Lebesgue measure on  $\mathbb{R}$ . Then the problem*

$$u(z, w) = \inf_{\mathbb{Q} \in B_k(\mathbb{P})} E_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)]$$

has a unique minimizer.

We conjecture that the same claim is true without the assumption that  $\langle \frac{w}{|w|}, X \rangle_* \mathbb{P}$  is absolutely continuous wrt. the Lebesgue measure on  $\mathbb{R}$ . We leave this for future research.

*Proof.* We first show that (2) can be rewritten as a one-dimensional optimization problem. Let

$$\mathbb{P}^w := \langle \frac{w}{|w|}, X \rangle_* \mathbb{P}.$$

We claim that

$$(63) \quad u(k, w) = \inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] = \inf_{\mathbb{Q} \in B_k(\mathbb{P}^w)} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 |w| X)].$$

Indeed, Lemma 7.1 shows that

$$\inf_{\mathbb{Q} \in B_k(\mathbb{P})} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 \langle w, X \rangle)] = \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle + x_0 \langle w, Z^\mu \rangle)].$$

For any  $X^\mu$  such that  $X^\mu \sim \mathbb{P}$  under  $\mu$ , define

$$X^w := \langle \frac{w}{|w|}, X^\mu \rangle$$

Then  $X^w \sim \mathbb{P}^w$  under  $\mu$ . Clearly, we have that

$$\langle w, X^\mu \rangle + \langle w, Z^\mu \rangle = |w| \left( X^w + \langle \frac{w}{|w|}, Z^\mu \rangle \right),$$

and

$$|\langle \frac{w}{|w|}, Z^\mu \rangle| \leq |Z^\mu|$$

for any  $Z^\mu \in L_\mu^p(\mathbb{R}^d)$ . On the other hand for any  $Y^\mu \in L_\mu^p(\mathbb{R})$  we have

$$|w|(X^w + Y^\mu) = \langle w, X^\mu \rangle + \langle w, \frac{w}{|w|} Y^\mu \rangle \quad \text{and} \quad |\frac{w}{|w|} Y^\mu| = |Y^\mu|.$$

This shows

$$\begin{aligned} & \inf_{X^\mu \sim \mathbb{P}, Z^\mu \in B_k^\mu} \mathbb{E}_\mu[U(x_0 + x_0 \langle w, X^\mu \rangle + x_0 \langle w, Z^\mu \rangle)] \\ &= \inf_{X^w \sim \mathbb{P}^w, \|Y^\mu\|_{L_\mu^p(\mathbb{R})} \leq k} \mathbb{E}_\mu[U(x_0 + x_0 |w|(X^w + Y^\mu))] \\ &= \inf_{\mathbb{Q} \in B_k(\mathbb{P}^w)} \mathbb{E}_{\mathbb{Q}}[U(x_0 + x_0 |w| X)], \end{aligned}$$

where the last equality follows from similar arguments as in Lemma 7.1. So, (63) is proved. From Lemma A.2 an optimizer  $\mathbb{Q}^* \in B_k(\mathbb{P})$  of (2) exists, so an optimizer  $\mathbb{Q}^* \in B_k(\mathbb{P}^w)$  for (63) also exists. Assume now that there exist  $\mathbb{Q}^1 \neq \mathbb{Q}^2 \in B_k(\mathbb{P}^w)$  attaining the infimum. Then clearly also the probability measure  $(\mathbb{Q}^1 + \mathbb{Q}^2)/2$  attains the infimum, but

$$\bar{\mathbb{Q}} := \frac{1}{2}(\mathbb{Q}^1 + \mathbb{Q}^2) \in \text{int}(B_k(\mathbb{P}^w)).$$

Indeed, take optimal transport plans  $\pi^1$  and  $\pi^2$  for  $(\mathbb{P}^w, \mathbb{Q}^1)$  and  $(\mathbb{P}^w, \mathbb{Q}^2)$ . As  $\mathbb{P}^w$  is absolutely continuous with respect to the Lebesgue measure, we know by [Santambrogio, 2015, Theorem 2.9], that the unique optimal coupling of  $\mathbb{P}^w$  and  $\bar{\mathbb{Q}}$  for  $\mathcal{W}_p$  is supported on a graph of a function. As  $(\pi^1 + \pi^2)/2 \in \Pi(\mathbb{P}^w, \bar{\mathbb{Q}})$  is not

supported on a graph of a function (recalling that  $\mathbb{Q}^1 \neq \mathbb{Q}^2$ ), we conclude that it is not optimal. Thus

$$\mathcal{W}_p(\mathbb{P}^w, \bar{\mathbb{Q}})^p < \frac{1}{2}(\mathcal{W}_p(\mathbb{P}^w, \mathbb{Q}^1)^p + \mathcal{W}_p(\mathbb{P}^w, \mathbb{Q}^2)^p) \leq k^p,$$

so that indeed  $\bar{\mathbb{Q}} \in \text{int}(B_k(\mathbb{P}^w))$  as claimed. As  $U$  is strictly increasing, it then follows that  $\mathbb{Q}^1$  and  $\mathbb{Q}^2$  do not attain the infimum (using an argument similar to  $Z^* \neq 0$  in the proof of Lemma A.4), a contradiction.  $\square$

#### REFERENCES

- Daniel Bartl, Samuel Drapeau, Jan Obloj, and Johannes Wiesel. Robust uncertainty sensitivity analysis. *arXiv preprint arXiv:2006.12022*, 2020.
- E. Bayraktar and Z. Zhou. On arbitrage and duality under model uncertainty and portfolio constraints. *Mathematical Finance*, 27(4):998–1012, 2017.
- R. Blanchard. *Stochastic control applied in the theory of decision in a discrete time non-dominated multiple-priors framework*. PhD thesis, Universite de Reims Champagne-Ardenne, 2017.
- R. Blanchard and L. Carassus. Multiple-priors investment in discrete time for unbounded utility function. *Annals of Applied Probability*, 88(2):241–281, 2018.
- R. Blanchard and L. Carassus. No arbitrage with multiple-priors in discrete time. *Stochastic Process and Applications*, 130(11):6657–6688, 2020.
- Jose Blanchet and Karthyek Murthy. Quantifying distributional model risk via optimal transport. *Mathematics of Operations Research*, 44(2):565–600, 2019.
- B. Bouchard and M. Nutz. Arbitrage and duality in nondominated discrete-time models. *The Annals of Applied Probability*, 25:823–859, 2015.
- Giuseppe C Calafiore. Ambiguous risk measures and optimal robust portfolios. *SIAM Journal on Optimization*, 18(3):853–877, 2007.
- L. Carassus and M. Ferhoun. Discrete time optimal investment under model uncertainty. Preprint, 2023.
- Steffen Dereich, Michael Scheutzow, and Reik Schottstedt. Constructive quantization: Approximation by empirical measures. In *Annales de l’IHP Probabilités et statistiques*, volume 49, pages 1183–1203, 2013.
- Nicolas Fournier and Arnaud Guillin. On the rate of convergence in wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3-4):707–738, 2015.
- Rui Gao and Anton Kleywegt. Distributionally robust stochastic optimization with wasserstein distance. *Mathematics of Operations Research*, 2022.
- Itzhak Gilboa and David Schmeidler. Maxmin expected utility with non-unique prior. *Journal of mathematical economics*, 18(2):141–153, 1989.
- Lars Peter Hansen and Thomas J Sargent. Robustness. In *Robustness*. Princeton university press, 2011.
- J. Jacod and A. N. Shiryaev. Local martingales and the fundamental asset pricing theorems in the discrete-time case. *Finance Stoch.*, 2:259–273, 1998.
- Jeroen Kerkhof, Bertrand Melenberg, and Hans Schumacher. Model risk and capital reserves. *Journal of Banking & Finance*, 34(1):267–279, 2010.
- F. Knight. *Risk, uncertainty and profit*. Boston: Houghton Mifflin, 1921.

- D. O. Kramkov and W. Schachermayer. The asymptotic elasticity of utility functions and optimal investment in incomplete markets. *Ann. Appl. Probab.*, 9: 904–950, 1999.
- Fabio Maccheroni, Massimo Marinacci, and Aldo Rustichini. Ambiguity aversion, robustness, and the variational representation of preferences. *Econometrica*, 74 (6):1447–1498, 2006.
- Pascal J Maenhout. Robust portfolio rules and asset pricing. *Review of financial studies*, 17(4):951–983, 2004.
- Robert C Merton. Lifetime portfolio selection under uncertainty: The continuous-time case. *The review of Economics and Statistics*, pages 247–257, 1969.
- Peyman Mohajerin Esfahani and Daniel Kuhn. Data-driven distributionally robust optimization using the wasserstein metric: Performance guarantees and tractable reformulations. *Mathematical Programming*, 171(1-2):115–166, 2018.
- Jan Oblój and Johannes Wiesel. Distributionally robust portfolio maximization and marginal utility pricing in one period financial markets. *Mathematical Finance*, 31(4):1454–1493, 2021.
- Georg Ch Pflug and Mathias Pohl. A review on ambiguity in stochastic portfolio optimization. *Set-Valued and Variational Analysis*, 26:733–757, 2018.
- Georg Ch Pflug, Alois Pichler, and David Wozabal. The 1/n investment strategy is optimal under high model ambiguity. *Journal of Banking & Finance*, 36(2): 410–417, 2012.
- Hamed Rahimian and Sanjay Mehrotra. Distributionally robust optimization: A review. *arXiv preprint arXiv:1908.05659*, 2019.
- M. Rásonyi and L. Stettner. On the utility maximization problem in discrete-time financial market models. *Ann. Appl. Probab.*, 15:1367–1395, 2005.
- M. Rásonyi and L. Stettner. On the existence of optimal portfolios for the utility maximization problem in discrete time financial models. In: *Kabanov, Y.; Lipster, R.; Stoyanov, J. (Eds), From Stochastic Calculus to Mathematical Finance*, Springer., pages 589–608, 2006.
- Filippo Santambrogio. Optimal transport for applied mathematicians. *Birkhäuser*, NY, 55(58-63):94, 2015.
- Jörn Sass and Dorothee Westphal. Robust utility maximization in a multivariate financial market with stochastic drift. *International Journal of Theoretical and Applied Finance*, 24(04):2150020, 2021.
- W. Schachermayer. Optimal investment in incomplete markets when wealth may become negative. *Ann. Appl. Probab.*, 11:694–734, 2001.
- Alexander Schied. Optimal investments for risk-and ambiguity-averse preferences: a duality approach. *Finance and Stochastics*, 11(1):107–129, 2007.
- Alexander Schied, Hans Föllmer, and Stefan Weber. Robust preferences and robust portfolio choice. *Handbook of Numerical Analysis*, 15:29–87, 2009.
- C. Villani. *Optimal transport: old and new*, volume 338. Springer Berlin, 2008.

LAURENCE CARASSUS

LÉONARD DE VINCI PÔLE UNIVERSITAIRE, RESEARCH CENTER, 92 916 PARIS LA DÉFENSE, FRANCE  
AND LMR, UMR 9008, UNIVERSITÉ REIMS CHAMPAGNE-ARDENNE, FRANCE

*Email address:* `laurence.carassus@devinci.fr`

JOHANNES WIESEL

COLUMBIA UNIVERSITY, DEPARTMENT OF STATISTICS  
1255 AMSTERDAM AVENUE  
NEW YORK, NY 10027, USA

*Email address:* `johannes.wiesel@columbia.edu`