STABILITY OF ROSE WINDOW GRAPHS

MILAD AHANJIDEH¹, ISTVÁN KOVÁCS^{2*}, KLAVDIJA KUTNAR³

ABSTRACT. A graph Γ is said to be stable if for the direct product $\Gamma \times \mathbf{K}_2$, $\operatorname{Aut}(\Gamma \times \mathbf{K}_2)$ is isomorphic to $\operatorname{Aut}(\Gamma) \times \mathbb{Z}_2$; otherwise, it is called unstable. An unstable graph is called non-trivially unstable when it is not bipartite and no two vertices have the same neighborhood. Wilson described nine families of unstable Rose Window graphs and conjectured that these contain all non-trivially unstable Rose Window graphs (2008). In this paper we show that the conjecture is true.

1. Introduction

Let Γ be a finite and simple graph. The direct product $\Gamma \times \mathbf{K}_2$ of Γ with the complete graph \mathbf{K}_2 , also known as the canonical double cover of Γ , has vertex set $V(\Gamma) \times \{0,1\}$ and edges $\{(u,0),(v,1)\}$, where $\{u,v\} \in E(\Gamma)$. Following [22], here we shall rather use the notation $B(\Gamma)$ for $\Gamma \times \mathbf{K}_2$. The graph $B(\Gamma)$ admits some natural automorphisms, namely the permutation

$$(v,i) \mapsto (v,1-i), \ v \in V(\Gamma), \ i = 0,1;$$
 (1)

and for every $\alpha \in Aut(\Gamma)$, the permutation

$$(v,i) \mapsto (\alpha(v),i), \ v \in V(\Gamma), \ i = 0,1.$$
 (2)

The above permutations can be easily seen to form a group, which is isomorphic to $\operatorname{Aut}(\Gamma) \times \mathbb{Z}_2$. The graph Γ is said to be stable if $\operatorname{Aut}(\operatorname{B}(\Gamma)) \cong \operatorname{Aut}(\Gamma) \times \mathbb{Z}_2$, and $\operatorname{unstable}$ otherwise. This concept of stability was introduced by Marušič et al. [16]. It plays a key role in the investigation of the automorphisms of the direct products of arbitrary graphs (see [23]) and also in the study of regular embeddings of canonical double covers on orientable surfaces (see [17]). Arc-transitive graphs in terms of their stability were investigated by Surowski [20]. Recently, this topic has been studied by several authors [5, 8, 9, 10, 18, 19].

It is well known that Γ is unstable whenever it is disconnected, or bipartite having a non-trivial automorphism group, or it has two vertices with the same neighborhood (see, e.g., [22]), and in each of these cases Γ is also called *trivially unstable*. Necessary and sufficient conditions for a graph to be non-trivially unstable were given by Wilson [22], and in the same paper these conditions were also applied to four infinite classes of graphs:

- the circulant graphs,
- the generalized Petersen graphs,

²⁰²⁰ Mathematics Subject Classification. 05C25, 20B25.

Key words and phrases. Rose Window graph, stable graph, edge-transitive.

¹ Supported by the Slovenian Research Agency (reserach project N1-0208).

² Supported by the Slovenian Research Agency (research program P1-0285 and research projects J1-1695, N1-0140, J1-2451, N1-0208, J1-3001).

³ Supported by the Slovenian Research Agency (research program P1-0285 and research projects J1-1695, J1-1715, N1-0140, J1-2451, J1-2481, N1-0209, J1-3001).

^{*} Corresponding author.

family	(n, a, r)	conditions
W1	(4m,2m,r)	_
W2	(2m, m+2, m+1)	_
W3	(m,2d,1)	$m \text{ is odd, } m \ge 5, d^2 \equiv \pm 1 \pmod{m}, d \not\equiv \pm 1 \pmod{m}.$
W4	(2m, a, m-1)	$a \neq m$
W5	(8m, a, 2m)	a is even
W6	(4m, m, r),	$m, r \text{ are odd}, m \geq 3$
W7	(2m,m,m-1)	m is odd
W8	(2m,m,r)	m is odd, r is even, $r^2 \equiv \pm 1 \pmod{m}$, $r \not\equiv \pm 1 \pmod{m}$
W9	(2m, a, m - s)	$s^2 \equiv 1 \pmod{n}, \ sa \equiv -a \pmod{n}, \ a < m.$

Table 1. The nine families of unstable Rose Window graphs $R_n(a,r)$.

- the Rose Window graphs,
- the toroidal graphs.

Various families of unstable graphs were constructed for each class and Wilson conjectured that these families cover all non-trivially unstable graphs in the given class. The conjecture has been confirmed for the generalized Petersen graphs [19] and for the toroidal graphs [24], whereas it has been disproved for the class of circulant graphs [18]. To the best of our knowledge, a characterization of the unstable circulant graphs of order n and valency k is known only in special cases: n is odd [5, 18], n = 2p for a prime p [9] and $k \leq 7$ [10]. Up to now, no progress has been made on the Rose Window graphs. The aim of this paper is to fill in this gap and show that the conjecture for the Rose Window graphs is also true.

Definition 1.1 ([21]). Let n be an integer such that $n \geq 3$ and let $a, r \in \mathbb{Z}_n$, $a, r \neq 0$. The Rose Window graph $R_n(a, r)$ is defined to have the vertex set $\{u_i, v_i : i \in \mathbb{Z}_n\}$, and edges

$$\{u_i, u_{i+1}\}, \{v_i, v_{i+r}\}, \{u_i, v_i\}, \{u_{i+a}, v_i\}, i \in \mathbb{Z}_n,$$

where the additions in the subscripts are done in \mathbb{Z}_n .

Here and in what follows, \mathbb{Z}_n denotes the ring of residue classes of integers modulo n. Throughout the paper we regard elements of \mathbb{Z}_n simultaneously as integers. The Rose Window graphs can be regarded as tetravalent analogues of the well-studied generalized Petersen graphs. Definition 1.1 is due to Wilson [21], who was interested primarily in constructing edge-transitive Rose Window graphs and rotary maps with underlying Rose Window graphs. It was shown later that all edge-transitive Rose Window graphs are covered by Wilson's constructions [12]. For various interesting properties of Rose Window graphs, we refer to [4, 7, 11, 12].

Regarding stability, Wilson found nine families of unstable Rose Window graphs, which are listed in Table 1 (see [22, Theorems R.1–R.9]). He checked by computer that these families contain all non-trivially unstable Rose Window graphs on up to 200 vertices and posed the following conjecture.

Conjecture 1.2 ([22]). Every non-trivially unstable Rose Window graph is isomorphic to a graph in one of the families W1-W9 given in Table 1.

In this paper we show that the conjecture is true by proving the following theorem.

Theorem 1.3. Let $\Gamma = R_n(a, r)$ be a non-trivially unstable Rose Window graph.

- (i) If n is odd, then Γ is isomorphic to a graph in the family W3.
- (ii) If n is even and $B(\Gamma)$ is not edge-transitive, then Γ is isomorphic to a graph in one of the families W1 and W4-W9.
- (iii) If n is even and $B(\Gamma)$ is edge-transitive, then Γ is isomorphic to a graph in one of the families W2 and W4.

2. Preliminaries

All groups in this paper will be finite and all graphs will be finite and simple. If Γ is a graph, then $V(\Gamma)$, $E(\Gamma)$ and $Aut(\Gamma)$ denote its vertex set, edge set and automorphism group, respectively. Given two vertices $u, v \in V(\Gamma)$, we write $u \sim v$ when u and v are adjacent. For $v \in V(\Gamma)$, the set of vertices adjacent with v is denoted by $\Gamma(v)$, and for a non-empty subset $S \subseteq V(\Gamma)$, the subgraph of Γ induced by S is denoted by $\Gamma[S]$. Suppose that Γ is bipartite, i.e., $V(\Gamma)$ can be partitioned into two sets V_1 and V_2 such that no edge of Γ lies entirely in either V_1 or V_2 . Throughout the paper we will refer to the sets V_1 and V_2 as biparts of Γ . By \mathbf{K}_n and \mathbf{C}_n we denote the complete graph and the cycle on n vertices, respectively, and the symbol $n\Gamma$ will be used for n disjoint copies of Γ .

If $G \leq \operatorname{Aut}(\Gamma)$, then the stabilizer of v in G is denoted by G_v , the orbit of v under G by $\operatorname{Orb}_G(v)$, and the set of all orbits under G by $\operatorname{Orb}(G, V(\Gamma))$. If $S \subseteq V(\Gamma)$, then the setwise and the pointwise stabilizers of S in G are denoted by $G_{\{S\}}$ and $G_{(S)}$, respectively.

Let $s \geq 1$ be an integer. An s-arc of Γ is an (s+1)-tuple (v_1, \ldots, v_{s+1}) of its vertices such that v_i is adjacent with v_{i+1} for each $1 \leq i \leq s$, and $v_i \neq v_{i+2}$ for each $1 \leq i \leq s-1$. The graph Γ is said to be (G, s)-arc-transitive if $G \leq \operatorname{Aut}(\Gamma)$ and G is transitive on the set of all s-arcs of Γ . A (G, s)-arc-transitive graph is called (G, s)-transitive if it is not (G, s+1)-arc-transitive. An $(\operatorname{Aut}(\Gamma), s)$ -transitive graph is also called s-transitive, and an $(\operatorname{Aut}(\Gamma), 1)$ -arc-transitive graph is simply called arc-transitive.

We shall need the following result about (G, s)-transitive tetravalent graphs.

Proposition 2.1 ([1, Chapter 17]). Let Γ be a connected (G, s)-transitive tetravalent graph with $s \leq 2$ and let $v \in V(\Gamma)$. Then exactly one of following statements is true.

- (i) s = 1 and G_v is a 2-group.
- (ii) s = 2 and $G_v \cong A_4$ or S_4 .

Let $G \leq \operatorname{Aut}(B(\Gamma))$ be the group generated by the automorphisms defined in (1) and (2). The elements in G are referred to as the *expected automorphisms* of $B(\Gamma)$.

Proposition 2.2 ([8]). Let Γ be a connected and non-bipartite graph. An automorphism σ of $B(\Gamma)$ is expected if and only if for every $v \in V(\Gamma)$, there exists some $w \in V(\Gamma)$ such that

$$\sigma(\{(v,0),(v,1)\}) = \{(w,0),(w,1)\}.$$

We review next four results about Rose Window graphs obtained in [4, 12, 21]. The first one establishes some obvious isomorphisms.

Proposition 2.3 ([21]). A Rose Window graph $R_n(a,r)$ is isomorphic to each of the graphs: $R_n(-a,r)$, $R_n(a,-r)$ and $R_n(-a,-r)$.

As a corollary of Proposition 2.3, up to graph isomorphism, one may always assume that $1 \le a, r \le n/2$ for every graph $R_n(a, r)$.

It was pointed out in [21] that any Rose Window graph Γ admits at most 3 edge-orbits under $\operatorname{Aut}(\Gamma)$. The next two results provide information about the order of $\operatorname{Aut}(\Gamma)$ when the number of edge-orbits is 3 or 2.

Proposition 2.4 ([4, Corollary 3.5(a)]). Suppose that $\Gamma = R_n(a, r)$ admits three edgeorbits under $\operatorname{Aut}(\Gamma)$. If $2a \not\equiv 0 \pmod{n}$, then $|\operatorname{Aut}(\Gamma)| = 2n$.

Proposition 2.5 ([4, Corollary 3.9]). Suppose that $\Gamma = R_n(a, r)$ admits two edge-orbits under $\operatorname{Aut}(\Gamma)$. If $2a \not\equiv 0 \pmod{n}$, then $|\operatorname{Aut}(\Gamma)| = 4n$.

Proposition 2.6 ([4, Lemma 2.1 & Theorem 2.3]). Suppose that $\Gamma = R_n(a, r)$ is not edge-transitive. Then there are two edge-orbits under $\operatorname{Aut}(\Gamma)$ if and only if

$$r^2 \equiv 1 \pmod{n}$$
 and $ra \equiv \pm a \pmod{n}$.

The following four families of edge-transitive graphs were introduced by Wilson [21].

Proposition 2.7 ([12, Theorem 1.2]). Let $\Gamma = R_n(a, r)$ be an edge-transitive graph and suppose that $1 \le a, r \le n/2$. Then one of the following holds.

- (a) $\Gamma = R_n(2,1)$,
- (b) $\Gamma = R_{2m}(m-2, m-1),$
- (c) $\Gamma = R_{12m}(3m+2,3m-1)$ or $R_{12m}(3m-2,3m+1)$,
- (d) $\Gamma = R_{2m}(2b, r), b^2 \equiv \pm 1 \pmod{m}, r \in \{1, m-1\} \text{ and } r \text{ is odd.}$

Moreover, in family (d), $H \triangleleft Aut(\Gamma)$, where H is generated by the permutation

$$(u_0, u_2, \ldots, u_{2m-2})(u_1, u_3, \ldots, u_{2m-1})(v_0, v_2, \ldots, v_{2m-2})(v_1, v_3, \ldots, v_{2m-1}).$$

A subgroup H of a group G is *core-free* if it contains no non-trivial normal subgroup of G. The following result of Lucchini [15], independently discovered by Herzog and Kaplan [6], about cyclic core-free subgroups in finite groups will be needed in the proof of the main result of this paper.

Proposition 2.8 ([6, 15]). If C is a core-free cyclic proper subgroup of a group G, then $|C|^2 < |G|$.

3. On the canonical double covers of Rose Window graphs

In this section we collect a couple of elementary properties of the graphs in the title. Throughout the section we let $\Gamma = R_n(a, r)$.

The canonical double cover $B(\Gamma)$ has vertex set

$$\{u_{i,j}, v_{i,j} : i \in \mathbb{Z}_n, j \in \mathbb{Z}_2\},\$$

and edges

$$\{u_{i,j}, u_{i+1,j+1}\}, \{v_{i,j}, v_{i+r,j+1}\}, \{u_{i,j}, v_{i,j+1}\}, \{u_{i+a,j}, v_{i,j+1}\},$$

where $i \in \mathbb{Z}_n, j \in \mathbb{Z}_2$ and the additions in the subscripts are done in \mathbb{Z}_n and \mathbb{Z}_2 , respectively.

3.1. The rim, the hub and the spoke cycles. The subgraph of $B(\Gamma)$ induced by the edges $\{u_{i,j}, u_{i+1,j+1}\}$ $(i \in \mathbb{Z}_n, j \in \mathbb{Z}_2)$ is a cycle of length 2n if n is odd, and two cycles of the same length n if n is even. These cycles will be referred to as the rim cycles of $B(\Gamma)$.

If $2r \not\equiv 0 \pmod{n}$, then the subgraph induced by the edges $\{v_{i,j}, v_{i+r,j+1}\}$ $(i \in \mathbb{Z}_n, j \in \mathbb{Z}_2)$ is a union of disjoint cycles of the same length and these will be referred to as the hub cycles.

Finally, the subgraph induced by the remaining edges $\{u_{i,j}, v_{i,j+1}\}, \{u_{i+a,j}, v_{i,j+1}\}\ (i \in \mathbb{Z}_n, j \in \mathbb{Z}_2)$ is a union of cycles of the same length and these will be referred to as the spoke cycles. The next lemma is straightforward.

Lemma 3.1. Let $\Gamma = R_n(a,r)$ such that $2r \not\equiv 0 \pmod{n}$, and let ℓ_h and ℓ_s be the length of the hub and the spoke cycles of $B(\Gamma)$, respectively. Then

$$\ell_h = \frac{2^{\varepsilon} n}{\gcd(r, n)} \quad and \quad \ell_s = \frac{2n}{\gcd(a, n)},$$
 (3)

where $\varepsilon = 0$ if $n/\gcd(r, n)$ is even and $\varepsilon = 1$ otherwise.

3.2. The partition $\{S_i : 1 \leq i \leq 4\}$. Suppose that n is even. The following sets of vertices are well-defined:

$$S_1 = \{u_{i,j} : i \equiv j \pmod{2}\}, \ S_2 = \{u_{i,j} : i \not\equiv j \pmod{2}\},$$
(4)

$$S_3 = \{v_{i,j} : i \equiv j \pmod{2}\}, \ S_4 = \{v_{i,j} : i \not\equiv j \pmod{2}\}.$$
 (5)

The above sets form a partition of $V(B(\Gamma))$. Note that the vertices visited by the two rim cycles of $B(\Gamma)$ form exactly the sets S_1 and S_2 , respectively

3.3. Connectivity.

Lemma 3.2. The following conditions are equivalent for any Rose Window graph $\Gamma = R_n(a, r)$.

- (i) Γ is bipartite.
- (ii) $B(\Gamma)$ is disconnected.
- (iii) n, a are even and r is odd.

Proof. (i) \Rightarrow (ii): This follows from the known fact that the canonical double cover of any bipartite graph is disconnected (see, e.g. [22]).

(ii) \Rightarrow (iii): The number n must be even, for otherwise there is only one rim cycle and $B(\Gamma)$ is clearly connected. If a were odd, then one can easily check that every spoke cycle would contain vertices from both rim cycles, so $B(\Gamma)$ would be connected. This shows that a is even. Note that, if X is a spoke cycle, then $V(X) \subseteq S_1 \cup S_4$ or $V(X) \subseteq S_2 \cup S_3$.

Finally, if r were even, then every hub cycle would contain vertices from both sets S_3 and S_4 , and so B(Γ) would be connected. This shows that r is odd.

(iii) \Rightarrow (i): It is straightforward to check that Γ is bipartite with biparts V_1 and V_2 defined as $V_1 = \{u_i, v_{i+1} : i \equiv 0 \pmod{2}\}$ and $V_2 = \{u_i, v_{i+1} : i \equiv 1 \pmod{2}\}$.

3.4. Some expected automorphisms. Define the permutations ρ , μ and β of $V(B(\Gamma))$ as follows:

$$\rho = \prod_{j \in \mathbb{Z}_2} (u_{0,j}, u_{1,j}, \dots, u_{n-1,j})(v_{0,j}, v_{1,j}, \dots, v_{n-1,j}).$$
 (6)

$$\rho = \prod_{j \in \mathbb{Z}_2} (u_{0,j}, u_{1,j}, \dots, u_{n-1,j})(v_{0,j}, v_{1,j}, \dots, v_{n-1,j}).$$

$$\mu = \prod_{i \in \mathbb{Z}_n, j \in \mathbb{Z}_2} (u_{i,j}, u_{-i,j})(v_{i,j}, v_{-i-a,j})$$

$$\beta = \prod_{i \in \mathbb{Z}_n} (u_{i,0}, u_{i,1})(v_{i,0}, v_{i,1}).$$
(8)

$$\beta = \prod_{i \in \mathbb{Z}_n} (u_{i,0}, u_{i,1})(v_{i,0}, v_{i,1}). \tag{8}$$

It is easy to verify that all these permutations are expected automorphism of $B(\Gamma)$. Also, $\langle \rho, \mu \rangle \cong D_{2n}$ and β commutes with both ρ and μ . Here D_{2n} denotes the dihedral group of order 2n.

The next result can be obtained directly from Proposition 2.2.

Lemma 3.3. Let $\Gamma = R_n(a,r)$ be a connected and non-bipartite graph and let β be the automorphism of $B(\Gamma)$ defined in (8). An automorphism σ of $B(\Gamma)$ is expected if and only if $\sigma\beta = \beta\sigma$.

Lemma 3.4. Let $\Gamma = R_n(a,r)$ be a graph such that n is even and one of the following conditions hold.

- (i) a, r are even, $2a \not\equiv 0 \pmod{n}$ and $4r \not\equiv 0 \pmod{n}$.
- (ii) a is odd and $4a \not\equiv 0 \pmod{n}$.
- (iii) a is odd, $4a \equiv 0 \pmod{n}$, $2a \not\equiv 0 \pmod{n}$ and r is even.

Then the only automorphism of $B(\Gamma)$ fixing S_1 pointwise and S_2 setwise is the identity permutation, where S_1 and S_2 are the sets defined in (4).

Proof. Let σ be an automorphism of B(Γ) fixing S_1 pointwise and S_2 setwise. Note that σ maps any rim/hub/spoke cycle to a rim/hub/spoke cycle.

- (i): Let X be any spoke cycle having a vertex from S_1 . Since a is even, it follows that $|V(X) \cap S_1| = |V(X) \cap S_4| = \ell_s/2$, where S_4 is the set defined in (5). As $2a \not\equiv 0$ $(\text{mod } n), \ell_s > 4 \text{ due to } (3), \text{ we see that } \sigma \text{ fixes every vertex in } S_4 \text{ too. Now, as } r \text{ is even,}$ for every hub cycle X, it holds that $|V(X) \cap S_3| = |V(X) \cap S_4| = \ell_h/2$, where S_3 is the set defined in (5). As $4r \not\equiv 0 \pmod{n}$, $\ell_h > 4$ due to (3), and this yields that σ fixes every vertex in S_3 too. It is easy to see that σ is the identity permutation.
- (ii): In this case, $|X \cap S_1| = \ell_s/4$ for every spoke cycle X. Since a is odd and $4a \not\equiv 0$ (mod n), it follows from (3) that $\ell_s > 8$, which shows that σ is the identity permutation.
 - (iii): Fix an arbitrary vertex $u_{i,j} \in S_2$ and let X be the spoke cycle through $u_{i,j}$. Then

$$X = (u_{i,j}, v_{i,j+1}, u_{i+a,j}, v_{i+a,j+1}, u_{i+2a,j}, v_{i+2a,j+1}, u_{i+3a,j}, v_{i+3a,j+1}).$$

Then $u_{i+a,j}, u_{i+3a,j} \in S_1$, hence σ maps X to itself, and either it fixes both $u_{i,j}$ and $u_{i+2a,j}$ or it swaps them. Using also that $u_{i,j}$ and $u_{i+2a,j}$ form an antipodal pair in the rim cycle consisting of the vertices in S_2 , we conclude that σ is either the identity permutation, or it acts on X as the unique reflection fixing $u_{i+a,j}$ and $u_{i+3a,j}$ and this does not depend on the choice of the vertex $u_{i,j}$ from S_2 . We finish the proof by showing that the latter possibility leads to a contradiction.

Indeed, then for every $i \in \mathbb{Z}_n$ and $j \in \mathbb{Z}_2$,

$$\sigma(v_{i,j}) = \begin{cases} v_{i+a,j} & \text{if } i \equiv j \pmod{2} \\ v_{i-a,j} & \text{if } i \not\equiv j \pmod{2}. \end{cases}$$

As r is even and $v_{0,0} \sim v_{r,1}$, we get $v_{a,0} = \sigma(v_{0,0}) \sim \sigma(v_{r,1}) = v_{r-a,1}$. This shows that $r - 2a \equiv \pm r \pmod{n}$. Since $2a \not\equiv 0 \pmod{n}$, it follows that $2r \equiv 2a \pmod{n}$. Using also that a is odd and $4a \equiv 0 \pmod{n}$, we find that 4 divides n, and hence $2r \equiv 2a \pmod{4}$. This contradicts the assumption that a is odd and r is even.

3.5. **Edge-transitive graphs.** The lemma below can be retrieved from the proof of [12, Lemma 3.3].

Lemma 3.5. Let $\Gamma = R_n(a,r)$ be a non-bipartite graph such that n > 4 and $B(\Gamma)$ is edge-transitive. Then $B(\Gamma)$ is s-transitive for $s \leq 2$.

3.6. Quotient graphs. Let Σ be an arbitrary graph and $\pi = \{C_1, \ldots, C_k\}$ be a partition of $V(\Sigma)$. The quotient graph Σ/π is defined to have the vertex set π and edges $\{C_i, C_j\}$, where $C_i \neq C_j$ and there is an edge $\{u, v\} \in E(\Sigma)$ with $u \in C_i$ and $v \in C_j$. In the case when π is formed by the orbits under a subgroup $H \leq \operatorname{Aut}(\Sigma)$, we also write Σ/H for Σ/π .

For a divisor k of n, let $\phi_{n,k}: \mathbb{Z}_n \to \mathbb{Z}_k$ be the group homorphism for which $\phi_{n,k}(1) = 1$ (here the integer 1 represents an element in \mathbb{Z}_n in the left side, whereas it represents an element in \mathbb{Z}_k in the right side). Furthermore, let $\phi_{n,k}^*$ be the mapping from $\{u_{i,j}, v_{i,j} : i \in \mathbb{Z}_n, j \in \mathbb{Z}_2\}$ to $\{u_{i,j}, v_{i,j} : i \in \mathbb{Z}_k, j \in \mathbb{Z}_2\}$ defined as follows:

$$\phi_{n,k}^*(u_{i,j}) = u_{\phi_{n,k}(i),j} \text{ and } \phi_{n,k}^*(v_{i,j}) = v_{\phi_{n,k}(i),j}, \ i \in \mathbb{Z}_n, \ j \in \mathbb{Z}_2.$$
 (9)

The following lemma can be easily derived after one applies well-known classical results about quotients of arc-transitive graphs (see, e.g., [15]).

Lemma 3.6. Let $\Gamma = R_n(a,r)$, $G = \operatorname{Aut}(B(\Gamma))$, and let $\rho \in G$ be the automorphism defined in (6). Suppose that $B(\Gamma)$ is connected, edge-transitive, and that $\operatorname{Orb}(H, V(B(\Gamma)))$ is a block system for G for some subgroup $H < \langle \rho \rangle$, $H \neq 1$.

- (i) If |H| = n/2, then $G_{u_{0,0}}$ is a 2-group and $B(\Gamma)/H \cong C_8$.
- (ii) If |H| < n/2, then $H \triangleleft G$ and

$$B(\Gamma)/H \cong B(R_k(a',r')),$$

where k = n/|H|, $a' = \phi_{n,k}(a)$ and $r' = \phi_{n,k}(r)$.

- (iii) With the notation in (ii), the mapping $\phi_{n,k}^*$ defined in (9) is a homomorphism from $B(\Gamma)$ onto $B(R_k(a',r'))$.
- (iv) Letting $\Sigma = B(R_k(a', r'))$ in (ii), there is an epimorphism

$$\varphi: G/H \to \operatorname{Aut}(\Sigma)$$

such that $\varphi(\rho)$ is the automorphism of Σ defined in (6). Furthermore, Σ is edgetransitive.

4. Proof of case (I) of Theorem 1.3

Let $\Gamma = R_n(a, r)$ be a non-trivially unstable Rose Window graph and assume that n is odd. We show below that Γ is isomorphic to a graph in W3.

The key idea is to represent B(Γ) as a Rose Window graph. Let $\theta: \mathbb{Z}_n \times \mathbb{Z}_2 \to \mathbb{Z}_{2n}$ be the unique isomorphism mapping (1,1) to 1. Define the mapping $f: V(B(\Gamma)) \to \{u_i, v_i : i \in \mathbb{Z}_{2n}\}$ as follows:

$$f(u_{i,j}) = u_{\theta((i,j))}$$
 and $f(v_{i,j}) = v_{\theta((i,j))+n}, i \in \mathbb{Z}_n, j \in \mathbb{Z}_2.$

A straightforward check shows that f is an isomorphism from $B(\Gamma)$ to $R_{2n}(a', r')$, where $a' = \theta((a, 0))$ and $r' = \theta((r, 1))$. Equivalently,

$$a' = \begin{cases} a & \text{if } a \text{ is even,} \\ a+n & \text{if } a \text{ is odd,} \end{cases} \quad r' = \begin{cases} r+n & \text{if } r \text{ is even,} \\ r & \text{if } r \text{ is odd.} \end{cases}$$
 (10)

Note that a' is always even and r' is always odd.

Let $G = \operatorname{Aut}((R_{2n}(a', r')))$ and denote by ℓ the number of edge-orbits under G. Then $\ell \in \{1, 2, 3\}$.

Assume first that $\ell = 3$. Since a' is even, $a' \neq n$. Thus |G| = 4n due to Proposition 2.4. This contradicts the assumption that $R_n(a, r)$ is unstable.

Now assume that $\ell = 2$. As $a' \neq n$, |G| = 8n due to Proposition 2.5. Furthermore, by Proposition 2.6, $(r')^2 \equiv 1 \pmod{2n}$ and $r'a' \equiv \pm a' \pmod{2n}$. Since $r' \equiv r \pmod{n}$ and $a' \equiv a \pmod{n}$, it follows that $r^2 \equiv 1 \pmod{n}$ and $ra \equiv \pm a \pmod{n}$. If $R_n(a,r)$ were edge-transitive, then $R_{2n}(a',r')$ would be edge-transitive as well, which is not the case. After applying Propositions 2.5 and 2.6 to $R_n(a,r)$, we obtain that $|\operatorname{Aut}(R_n(a,r))| = 4n$. This contradicts the assumption that $R_n(a,r)$ is unstable.

Thus $\ell = 1$, i.e., $R_{2n}(a', r')$ is edge-transitive. Let

$$a'' = \min(a', 2n - a')$$
 and $r'' = \min(r', 2n - r')$,

and put $\Sigma = R_{2n}(a'', r'')$. By Proposition 2.3, $\Sigma \cong R_{2n}(a', r')$, in particular, Σ is edge-transitive. We obtain that Σ belongs to one of the families (a)–(d) in Proposition 2.7.

The graph Σ is not in the family (a). For otherwise, (a'', r'') = (2, 1). If a is even, then a' = a < n, hence a'' = a' = a, so a = 2; if a is odd, then a' = a + n, hence a'' = 2n - a' = n - a, and so a = n - 2. A similar argument yields that r = 1 or n - 1. If a = 2, then the vertices u_0 and v_{-1} have the same neighborhood; and if a = n - 2, then u_0 and v_1 share the same neighborhood. In either case, $\Gamma = R_n(a, r)$ has two vertices having the same neighborhood, which is impossible.

The graph Σ is not in the family (b). For otherwise, r'' = n - 1, which is impossible as r'' is odd. Also, Σ is neither in the family (c) because n is odd.

Thus $\Sigma = R_{2n}(a'', r'')$ is in the family (d), i.e., a'' = 2d such that $d^2 \equiv \pm 1 \pmod{n}$, $2 \leq a'' \leq n$ and $r'' \in \{1, n-1\}$ is odd. It follows that r'' = 1. We obtain a = 2d or n-2d and r = 1 or n-1. Finally, $d \not\equiv \pm 1 \pmod{n}$, for otherwise, $R_n(a, r)$ has two vertices with the same neighborhood, and $R_n(a, r)$ is indeed isomorphic to a graph in W3.

5. Proof of case (II) of Theorem 1.3

For this section we set the following assumptions.

Hypothesis 5.1. $\Gamma = R_n(a,r)$ is a non-trivially unstable graph such that n = 2m, $1 \le a, r \le m$ and $B(\Gamma)$ is not edge-transitive. Furthermore,

$$G = \operatorname{Aut}(B(\Gamma)),$$

 $C = \langle \rho \rangle$, where ρ is defined in (6),

 σ is an unexpected automorphism of B(Γ).

To settle case (ii) of Theorem 1.3, we have to show that under the above assumptions Γ is isomorphic to a graph in one of W1 and W4–W9. This will be done in three lemmas.

Lemma 5.2. Assuming Hypothesis 5.1, suppose that σ maps $S_1 \cup S_2$ to itself. Then Γ is isomorphic to a graph in one of W1, W5 and W6.

Proof. Since $\langle \rho, \beta \rangle$ is transitive on $S_1 \cup S_2$, $\sigma(u_{0,0}) = \alpha(u_{0,0})$ for some $\alpha \in \langle \rho, \beta \rangle$, where β is the automorphism of B(Γ) defined in (8). As $\alpha^{-1}\sigma$ is unexpected, we may assume that $\sigma(u_{0,0}) = u_{0,0}$.

For $\alpha \in G$ such that $\alpha(S_1) = S_1$, let α^* denote the permutation of S_1 induced by the restriction of α to S_1 . Clearly, α^* is an automorphism of the rim cycle containing $u_{0,0}$. Thus $\sigma^* \in \langle \mu^* \rangle$, hence we may assume that $\sigma \in G_{(S_1)}$. Here μ is the automorphism of $B(\Gamma)$ defined in (7), in particular, $\mu(S_1) = S_1$ and μ^* is well-defined.

Assume first that a is even. Then r is even due to Lemma 3.2. If a = m, then Γ is in W1. Let $a \neq m$. Suppose that $4r \not\equiv 0 \pmod{n}$. Then Lemma 3.4(i) yields that σ is the identity, which is impossible, hence $4r \equiv 0 \pmod{n}$.

Assume for the moment that r=m. Denote by ℓ_s the length of the spoke cycles of B(Γ). Since $a \neq m$, it follows that $\ell_s \geq 6$ according to (3). Let X be a spoke cycle. Then $|V(X) \cap (S_3 \cup S_4)| = \frac{1}{2}\ell_s$ and

$$|V(X) \cap S_1| = \frac{1}{2}\ell_s \text{ or } |V(X) \cap S_2| = \frac{1}{2}\ell_s.$$

Let us call X of type 1 in the former case and of type 2 in the latter case. It is clear that σ fixes all vertices of a spoke cycle of type 1. Furthermore, the hub edges $\{v_{i,j}, v_{i+m,j+1}\}$, where $i \in \mathbb{Z}_n, j \in \mathbb{Z}_2$, form a matching, and the vertices $v_{i,j}$ and $v_{i+m,j+1}$ belong to spoke cycles of distinct types. This shows that σ must fix every vertex in $S_3 \cup S_4$, and consequently, also in S_2 . This contradicts the assumption σ is not the identity permutation, and thus $r \neq m$. Using also that r is even and $4r \equiv 0 \pmod{n}$, we conclude that $r = \frac{n}{4}$ and $r = \frac{n}{4}$ and $r = \frac{n}{4}$ is divisible by 8, hence that $r = \frac{n}{4}$ is in W5.

Assume second that a is odd. Then $4a \equiv 0 \pmod{n}$ follows from Lemma 3.4(ii) and the condition that $\sigma \in G_{(S_1)}$. We show below that $a \neq m$. On the contrary, assume that a = m. Fix an arbitrary vertex $v_{i,j}$ and let X be the spoke cycle containing $v_{i,j}$. Then

$$X = (v_{i,j}, u_{i,j+1}, v_{i+m,j}, u_{i+m,j+1}).$$

This shows that σ fixes both $u_{i,j+1}$ and $u_{i+m,j+1}$, and $\sigma(v_{i,j}) = v_{i,j}$ or $v_{i+m,j}$. In view of Proposition 2.2, we may assume that

$$\sigma(v_{i,j}) = v_{i,j}, \ \sigma(v_{i+m,j}) = v_{i+m,j}, \ \sigma(v_{i,j+1}) = v_{i+m,j+1} \text{ and } \sigma(v_{i+m,j+1}) = v_{i,j+1}.$$
 (11)

Let Y be the hub cycle containing $v_{i,j}$. Then $\sigma(Y) = Y$, and thus $\sigma(v_{i',j'}) = v_{i',j'}$, where $v_{i',j'}$ is the vertex antipodal to $v_{i,j}$ in Y. Then $v_{i',j'} = v_{i,j+1}$ if r has odd order in \mathbb{Z}_n , and $v_{i',j'} = v_{i+m,j+1}$ otherwise. In either case we obtain a contradiction with the assumptions in (11), and by this we have shown that $a \neq m$. Finally, Lemma 3.4(iii) shows that r is odd, and so Γ is in W6.

For the rest of the section it will be assumed that $\sigma(S_1 \cup S_2) \neq S_1 \cup S_2$. Consequently, G is transitive on $V(B(\Gamma))$ and divides $E(B(\Gamma))$ into two orbits (recall that $B(\Gamma)$ is assumed to be non-edge-transitive). There are two hub cycles of the same length n, so by (3),

$$gcd(r, n) \in \{1, 2\}, \text{ and if } gcd(n, r) = 2, \text{ then } m \text{ is odd.}$$
 (12)

Denote by X_1 and X_2 the two rim cycles and by X_3 and X_4 the two hub cycles so that $V(X_i) = S_i$ for i = 1, 2 and $v_{0,0} \in V(X_3)$, where S_1 and S_2 are defined in (4). If r is even, then

$$V(X_3) = \{v_{i,j} : i \equiv 0 \pmod{2}\} \text{ and } V(X_4) = \{v_{i,j} : i \equiv 1 \pmod{2}\};$$
 (13)

and if r is odd, then

$$V(X_3) = S_3 \text{ and } V(X_4) = S_4.$$
 (14)

Clearly, the subsets

$$S_1 = V(X_1), S_2 = V(X_2), V(X_3), V(X_4)$$
 (15)

form a block system for G.

Lemma 5.3. Assuming Hypothesis 5.1, suppose that $\sigma(S_1 \cup S_2) \neq S_1 \cup S_2$ and let K be the kernel of the action of G on the block system defined in (15). If $K_{(S_1)} \neq 1$, then Γ is isomorphic to a graph in one of W1, W5 and W6.

Proof. Fix $\tau \in K_{(S_1)}$ such that τ is not the identity permutation. Clearly, τ fixes S_1 pointwise and S_2 setwise.

Thus if a is even, then the proof of Lemma 5.2 can be repeated with τ playing the role of σ .

Let a be odd. If a=m, then every spoke cycle contains exactly one vertex from S_1 , S_2 , $V(X_3)$ and $V(X_4)$, respectively, implying that τ is the identity permutation, which is impossible. Hence $a \neq m$, and the argument used in the proof of Lemma 5.2 can be applied again.

Lemma 5.4. Assuming Hypothesis 5.1, suppose that $\sigma(S_1 \cup S_2) \neq S_1 \cup S_2$ and let K be the kernel of the action of G on the block system defined in (15). If $K_{(S_1)} = 1$, then Γ is isomorphic to a graph in one of W7, W8 and W9.

Proof. Since $\langle \rho, \beta \rangle$ is transitive on $S_3 \cup S_4$, $\sigma(u_{0,0}) = \alpha(v_{0,1})$ for some $\alpha \in \langle \rho, \beta \rangle$, where β is the automorphism of B(Γ) defined in (8). As $\alpha^{-1}\sigma$ is unexpected, we may assume that $\sigma(u_{0,0}) = v_{0,1}$. Then the hub cycle containing $v_{0,1}$ is not preserved by σ , hence we have that $\sigma(v_{0,1}) = u_{0,0}$ or $u_{a,0}$. Replacing σ with $\rho^a \mu \sigma$ in the latter case, we may also assume that σ interchanges $u_{0,0}$ and $v_{0,1}$.

Furthermore, replacing Γ with $R_n(a, -r)$ if necessary (see Proposition 2.3), we may also assume that $\sigma(u_{1,1}) = v_{r,0}$ holds. Using that σ is an isomorphism from X_1 to the hub cycle containing $v_{0,1}$, which maps the arc $(u_{0,0}, u_{1,1})$ to the arc $(v_{0,1}, v_{r,0})$, we deduce that

$$\sigma(u_{i,j}) = v_{ri,j+1} \text{ for every } u_{i,j} \in S_1.$$
 (16)

Since $K_{(S_1)} = 1$, it follows that K is isomorphic to a subgroup of D_{2n} . It follows from (13) and (14) that $\rho^2 \in K$.

Suppose that n = 4, so $1 \le a, r \le 2$. Then r = 1 by (12). Since Γ has no two vertices with the same neighborhood, it follows that a = 1, and so Γ is isomorphic to the graph in W9 with m = 2, a = 1 and s = 3 due to Proposition 2.3.

For the rest of the proof we assume that n > 4. Then $\langle \rho^2 \rangle$ is characteristic in K. Using also that $K \triangleleft G$, we obtain that $\langle \rho^2 \rangle \triangleleft G$. This implies that there is an integer k such that $\gcd(k,m) = 1$ and $\sigma \rho^2 = \rho^{2k} \sigma$. Thus

$$\sigma(u_{2,0}) = \sigma \rho^2(u_{0,0}) = \rho^{2k} \sigma(u_{0,0}) = v_{2k,1}.$$

On the other hand, $\sigma(u_{2,0}) = v_{2r,1}$ due to (16), hence $2k \equiv 2r \pmod{n}$, $k \equiv r \pmod{m}$, and therefore, we may assume that k = r. Note that, if i is even, then $\sigma \rho^i \sigma^{-1} = (\sigma \rho^2 \sigma^{-1})^{\frac{i}{2}} = \rho^{ri}$. The following identity holds.

$$\sigma \rho^i = \rho^{ri} \sigma \text{ for every } i \in \mathbb{Z}_n, \ i \equiv 0 \pmod{2}.$$
 (17)

Case 1. r is even.

By (12), m is odd, so $u_{m,1}$ is the vertex antipodal to $u_{0,0}$ in X_1 . Then $\sigma^2(u_{m,1}) = u_{m,1}$. Using also (16), we obtain that $\sigma(v_{0,0}) = \sigma^2(u_{m,1}) = u_{m,1}$. Thus by (17), $\sigma(v_{r,0}) = u_{m,1}$.




FIGURE 1. The path **P** and its image under σ .

 $\sigma \rho^r(v_{0,0}) = \rho^{r^2} \sigma(v_{0,0}) = u_{r^2+m,1}$. On the other hand, $\sigma(v_{r,0}) \sim \sigma(v_{0,1}) = u_{0,0}$, implying that $r^2 \equiv \pm 1 \pmod{m}$.

Assume first that a is odd. If a = m, then Γ belongs to W8 if $r \not\equiv \pm 1 \pmod{m}$, and it belongs to W7 if $r \equiv \pm 1 \pmod{m}$. Assume that a < m. Observe that $\{u_{a,1}\} = B(\Gamma)(v_{0,0}) \cap S_1$. Since r is even, $v_{0,1} \in V(X_3)$, see (13), hence $\sigma(S_1) = V(X_3)$. Using also (16) and (17), we have that

$$\{v_{ra,0}\} = \{\sigma(u_{a,1})\} = B(\Gamma)(\sigma(v_{0,0})) \cap V(X_3) = B(\Gamma)(u_{m,1}) \cap V(X_3) = \{v_{m-a,0}\}.$$

This shows that $ra \equiv m - a \pmod{n}$. Then $r^2a \equiv -ra \equiv m + a \pmod{n}$, and using also that $r^2 \equiv \pm 1 \pmod{m}$ and a < m, we deduce that $r^2 \equiv 1 \pmod{m}$. Set s = m + r. Then $\Gamma = R_{2m}(a, m + s) = R_{2m}(a, m - s)$, $s^2 \equiv 1 \pmod{n}$ and $sa \equiv -a \pmod{n}$, hence Γ belongs to W9.

Now let a be even. Then $u_{0,0} = \sigma(v_{0,1}) \sim \sigma(u_{a,0}) = v_{ra,1}$, implying that $ra \equiv -a \pmod{n}$. It follows that $r^2 \equiv 1 \pmod{m}$, and taking s = m + r, we find that Γ belongs to W9.

Case 2. r is odd.

Then $V(X_3) = S_3$ and $V(X_4) = S_4$, where S_3 and S_4 are the sets defined in (5), see (14). By Lemma 3.2, a is odd. Consider the path

$$\mathbf{P} = (v_{-a,0}, u_{0,1}, v_{0,0}, u_{a,1}, v_{a,0})$$

and its image $\sigma(\mathbf{P})$, see Figure 1. The vertices of \mathbf{P} (from the left to the right) belong to the sets S_4, S_2, S_3, S_1 and S_4 , respectively. Therefore, the their images are from S_1, S_3, S_2, S_4 and S_1 , respectively. Furthermore, by (16), $\sigma(u_{a,1}) = v_{ra,0}$, so $x_4 = ra$. All these imply in turn that, $x_5 = ra$, $x_3 = ra + a$, $x_2 = ra + a$ and $x_1 = ra + 2a$. Using (17), we find that

$$u_{ra+2a,1} = \sigma(v_{-a,0}) = \sigma \rho^{-a-1}(v_{1,0}) = \rho^{-ra-r} \sigma(v_{1,0})$$

$$u_{ra,1} = \sigma(v_{a,0}) = \sigma \rho^{a-1}(v_{1,0}) = \rho^{ra-r} \sigma(v_{1,0}).$$

Thus $2(ra + a) \equiv 0 \pmod{n}$, whence ra + a = 0 or m in \mathbb{Z}_n .

Assume first that $ra + a \equiv 0 \pmod{n}$. We show below that this possibility cannot occur. Notice that, then

$$\sigma(u_{0,1}) = v_{0,0}, \ \sigma(v_{0,0}) = u_{0,1} \text{ and } \sigma(v_{1,0}) = u_{r,1}.$$
 (18)

By the initial assumptions on σ we also have

$$\sigma(u_{0,0}) = v_{0,1}, \ \sigma(u_{1,1}) = v_{r,0} \text{ and } \sigma(v_{0,1}) = u_{0,0}.$$
 (19)

Now $B(\Gamma)(v_{0,1}) \cap S_2 = \{u_{a,0}\}$ yields

$$\{\sigma(u_{a,0})\} = B(\Gamma)(\sigma(v_{0,1})) \cap S_3 = B(\Gamma)(u_{0,0}) \cap S_3 = v_{-a,1},$$

from which it follows that $v_{-a,1} = \sigma(u_{a,0}) = \sigma \rho^{a-1}(u_{1,0}) = \rho^{ra-r}\sigma(u_{1,0})$, so $\sigma(u_{1,0}) = v_{r,1}$. It can be shown in the same way that $\sigma(v_{-a,1}) = u_{a,0}$, and deduce from this that $\sigma(v_{1,1}) = u_{r,0}$. To sum up,

$$\sigma(u_{1,0}) = v_{r,1} \text{ and } \sigma(v_{1,1}) = u_{r,0}.$$
 (20)

The conditions in (18)–(20) together with (17) yield that σ commutes with β . This, however, contradicts the assumption that σ is unexpected due to Lemma 3.3. We verify that $\sigma\beta(u_{i,j}) = \beta\sigma(u_{i,j})$ and $\sigma\beta(v_{i,j}) = \beta\sigma(v_{i,j})$ for the case when i is even and j = 0 only, the remaining cases can be handled in the same way.

$$\sigma\beta(u_{i,0}) = \sigma(u_{i,1}) = \sigma\rho^{i}(u_{0,1}) = \rho^{ri}\sigma(u_{0,1}) = \rho^{ri}(v_{0,0}) = \beta\rho^{ri}(v_{0,1})
= \beta\rho^{ri}\sigma(u_{0,0}) = \beta\sigma\rho^{i}(u_{0,0}) = \beta\sigma(u_{i,0}).$$

and

$$\sigma\beta(v_{i,0}) = \sigma(v_{i,1}) = \sigma\rho^{i}(v_{0,1}) = \rho^{ri}\sigma(v_{0,1}) = \rho^{ri}(u_{0,0}) = \beta\rho^{ri}(u_{0,1})
= \beta\rho^{ri}\sigma(v_{0,0}) = \beta\sigma\rho^{i}(v_{0,0}) = \beta\sigma(v_{i,0}).$$

We have shown that $ar + a \not\equiv 0 \pmod{n}$, and therefore, $ar \equiv m - a \pmod{n}$. It follows that m is even, in particular, $a \neq m$. Then $u_{0,0} = \sigma(v_{0,1}) \sim \sigma(v_{r,1}) = \rho^{r^2 - r} \sigma(v_{1,1}) = u_{r^2 + m, 1}$, implying that $r^2 \equiv \pm 1 \pmod{m}$. Using also that $ra \equiv -a \pmod{m}$ and $a \neq m$, we have that $r^2 \equiv 1 \pmod{m}$. Finally, taking s = m + r, we find that Γ belongs to W9.

6. Proof of case (III) of Theorem 1.3

For this section we set the following assumptions.

Hypothesis 6.1. $\Gamma = R_n(a, r)$ is a non-trivially unstable graph, n = 2m, $1 \le a, r \le m$ and $B(\Gamma)$ is edge-transitive. Furthermore,

$$G = Aut(B(\Gamma)),$$

 G^+ is the subgroup of G preserving the biparts of $B(\Gamma)$,

 $C = \langle \rho \rangle$, where ρ is defined in (6),

$$L = \langle \rho^m \rangle$$
.

Note that, if n > 4, then by Lemma 3.5, B(Γ) is s-transitive for $s \leq 2$, in particular, it is vertex-transitive. These properties will be used a couple of times in the section.

In order to settle case (iii) of Theorem 1.3, we have to show that under the above assumptions Γ is isomorphic to a graph in one of W2 and W4. We deal first with the case when $C \triangleleft G$.

Lemma 6.2. Assuming Hypothesis 6.1, suppose that $C \triangleleft G$. Then m is odd and Γ is isomorphic to a graph in the family W4.

Proof. Fix $\sigma \in G_{u_{0,0}}$ such that $\sigma(u_{1,1}) = v_{0,1}$. There is an integer k such that $\gcd(k,n) = 1$ and $\sigma \rho^i = \rho^{ki} \sigma$ for every $i \in \mathbb{Z}_n$. Then

$$\sigma(u_{-1,1}) = \sigma \rho^{-2}(u_{1,1}) = \rho^{-2k} \sigma(u_{1,1}) = v_{-2k,1}.$$

This implies that $v_{-2k,1} = v_{-a,1}$, hence $a \equiv 2k \pmod{n}$. Note that a is even. Then $\sigma(v_{0,1}) = u_{1,1}$ or $u_{-1,1}$. Here we consider only the first case as the other case can be

handled in the same way. For every $i \in \mathbb{Z}_n$,

$$\sigma(u_{i,0}) = \sigma \rho^{i}(u_{0,0}) = \rho^{ki} \sigma(u_{0,0}) = u_{ki,0},
\sigma(u_{i,1}) = \sigma \rho^{i-1}(u_{1,1}) = \rho^{ki-k} \sigma(u_{1,1}) = v_{ki-k,1},
\sigma(v_{i,1}) = \sigma \rho^{i}(v_{0,1}) = \rho^{ki} \sigma(v_{0,1}) = u_{ki+1,1}.$$

Note that $\operatorname{Orb}_C(v_{0,0})$ is mapped to itself by σ . Thus $u_{-kr+1,1} = \sigma(v_{-r,1})$ and $u_{kr+1,1} = \sigma(v_{r,1})$ share a common neighbor in $\operatorname{Orb}_C(v_{0,0})$. This implies that $2kr \equiv \pm a \equiv \pm 2k \pmod{n}$, whence $r \equiv \pm 1 \pmod{m}$. Since a is even, r must be even as well because Γ is not bipartite. Thus m is odd and r = m - 1 or m + 1. We conclude that Γ is isomorphic to a graph in the family W4.

The case when C is not normal in G requires three preparatory lemmas. We denote the center of G by Z(G), and for a subgroup $H \leq G$ and an element $g \in G$, by $C_G(H)$ and $C_G(g)$ the centralizers of H and g, respectively, in G.

Lemma 6.3. Assuming Hypothesis 6.1, $L \triangleleft G$, consequently, $\rho^m \in Z(G)$.

Proof. For the sake of simplicity, let $V = V(B(\Gamma))$. We proceed by induction on n. With the help of the computer algebra package MAGMA [2] we checked that the lemma holds if $n \leq 48$. From now on it is assumed that n > 48.

Case 1. $G_{u_{0,0}}$ is not a 2-group.

As the graph B(Γ) is s-arc-transitive for $s \leq 2$, see the remark after Hypothesis 6.1, it follows from Proposition 2.1 that $G_{u_{0,0}} \cong A_4$ or S_4 . Thus $|G^+| \leq 48n < |C|^2$ because of n > 48. By Proposition 2.8, G^+ contains a normal subgroup N such that $1 < N \leq C$. If |N| is even, then L is a characteristic subgroup of N, and we find that $L \triangleleft G$.

Assume now that |N| is odd. Let \mathcal{B} be a minimal non-trivial block system for G, which is a refinement of the block system $\mathrm{Orb}(N,V)$. It is easy to show that $\mathcal{B}=\mathrm{Orb}(H,V)$ for some subgroup $1 < H \le N$.

If |H| = n/2, then $G_{u_{0,0}}$ is a 2-group due to Lemma 3.6(i), which is impossible. Thus |H| < n/2, and by part (ii) of the same lemma,

$$B(\Gamma)/H \cong B(R_k(\phi_{n,k}(a), \phi_{n,k}(r))),$$

where k = n/|H|. As |H| is odd, k is even and the induction hypothesis can be applied to B $(R_k(\phi_{n,k}(a), \phi_{n,k}(r)))$. Lemma 3.6(iii) shows that $HL/H \triangleleft G/H$, whence $HL \triangleleft G$. This implies that $L \triangleleft G$, as claimed.

Case 2. $G_{u_{0,0}}$ is a 2-group.

Let T be a Sylow 2-subgroup of G^+ such that $G_{u_{0,0}} \leq T$. Then $\mathrm{Orb}_T(u_{0,0})$ is a block for G (see [3, Theorem 1.5A]). Let Δ be a minimal block contained in the latter orbit such that $u_{0,0} \in \Delta$ and $\Delta \neq \{u_{0,0}\}$, and let \mathcal{B} be the block system generated by Δ . As $|\Delta|$ divides $|\mathrm{Orb}_T(u_{0,0})|$, $|\Delta|$ is a power of 2. Since $G_{u_{0,0}}$ is a 2-group, we obtain that Δ contains a vertex v such that $v \neq u_{0,0}$ and $G_{u_{0,0}} = G_v$. It is known that the subset Δ' of V, defined as

$$\Delta' = \{w : w \in V \text{ and } G_w = G_{u_{0,0}}\}$$

is a block for G (see, e.g., [3, Excercise 1.6.5]). Since $\Delta \cap \Delta'$ is also a block and Δ is minimal, we obtain $\Delta \subseteq \Delta'$. Using also that $\mu \in G_{u_{0,0}}$, see (7), we obtain that Δ is one of the following sets:

$$\{u_{0,0}, u_{m,0}\}, \{u_{0,0}, v_{b,0}\}, \text{ and } \{u_{0,0}, u_{m,0}, v_{b,0}, v_{b+m,0}\},$$
 (21)

where $a \equiv -2b \pmod{n}$. If $\Delta = \{u_{0,0}, u_{m,0}\}$, then $\mathcal{B} = \operatorname{Orb}(L, V)$. By Lemma 3.6(ii), $L \triangleleft G$. Therefore, to finish the proof it sufficient to exclude the last two sets in (21). Assume first that

$$\Delta = \{u_{0,0}, u_{m,0}, v_{b,0}, v_{b+m,0}\}.$$

Since Δ is a minimal block for G, it follows that the setwise stabilizer $G_{\{\Delta\}}$ acts primitively on Δ . Denote by K and $(G_{\{\Delta\}})^*$ the kernel and the image, respectively, of the action of $G_{\{\Delta\}}$ on Δ . We have that $(G_{\{\Delta\}})^* \cong G_{\{\Delta\}}/K$. As $G_{u_{0,0}}$ is a 2-group, $G_{\{\Delta\}}$ is also a 2-group. Therefore, there exists a subgroup $M \leq G_{\{\Delta\}}$ such that K < M, |M/K| = 2 and $M/K < Z(G_{\{\Delta\}}/K)$. We obtain that $M \lhd G_{\{\Delta\}}$ and there are two M-orbits on Δ , each of size 2. This implies that $G_{\{\Delta\}}$ is imprimitive on Δ , a contradiction.

Assume second that $\Delta = \{u_{0,0}, v_{b,0}\}$. Consider the induced subgraph $B(\Gamma)[\Delta_1 \cup \Delta_2]$, where $\Delta_1, \Delta_2 \in \mathcal{B}$ and Δ_1 and Δ_2 are adjacent in Γ/\mathcal{B} . Denote the latter subgraph by Σ . As $B(\Gamma)$ is arc-transitive, see the remark after Hypothesis 6.1, Σ does not depend on the choice of Δ_1 and Δ_2 . There are three options for Σ , namely, $\Sigma \cong \mathbf{K}_2$ or $2\mathbf{K}_2$ or \mathbf{C}_4 . Choosing $\Delta_1 = \Delta = \{u_{0,0}, v_{b,0}\}$ and $\Delta_2 = (\beta \rho^{-b})(\Delta) = \{u_{-b,1}, v_{0,1}\}$, one can observe that $\Sigma \cong 2\mathbf{K}_2$ or \mathbf{C}_4 . Now choosing Δ_2 to be $(\beta \rho)(\Delta)$, we conclude that $\{v_{b,0}, v_{b+1,1}\}$ is an edge, whence $r \equiv \pm 1 \pmod{n}$, in particular, r is odd. On the other hand, $a \equiv -2b \pmod{n}$, so a is even. By Lemma 3.2, Γ is bipartite, which is a contradiction. \square

We describe next a situation, which we will encounter a couple of times in the rest of the paper. Suppose that d is an odd divisor of m such that

$$d > 3$$
, $\langle \rho^d \rangle \triangleleft G$, $\phi_{n,d}(a) = \pm 2$ and $\phi_{n,d}(r) = \pm 1$.

Let $\Sigma = B\left(R_d(\phi_{n,d}(a), \phi_{n,d}(r))\right)$ and let

$$\varepsilon = \begin{cases} 1 & \text{if } \phi_{n,d}(a) = -2, \\ -1 & \text{if } \phi_{n,d}(a) = 2. \end{cases}$$

By Lemma 3.6(iii), $\phi_{n,d}^*$ is a homomorphism from B(Γ) onto Σ , where $\phi_{n,d}^*$ is defined in (9). Observe that, for every $i \in \mathbb{Z}_d$ and $j \in \mathbb{Z}_2$, $v_{i+\varepsilon,j}$ is the unique vertex of Σ for which

$$\Sigma(u_{i,j}) = \Sigma(v_{i+\varepsilon,j}).$$

This shows that sets $\{u_{i,j}, v_{i+\varepsilon,j}\}$, $i \in \mathbb{Z}_d$, $j \in \mathbb{Z}_2$, form a block system for $\operatorname{Aut}(\Sigma)$. Consequently, the sets

$$\Delta_{i,j} := \left\{ (\phi_{n,d}^*)^{-1}(u_{i,j}), (\phi_{n,d}^*)^{-1}(v_{i+\varepsilon,j}) \right\}
= \left\{ u_{i+xd,j}, v_{i+xd+\varepsilon,j} : 0 \le x \le n/d - 1 \right\}, \ 0 \le i \le d - 1 \text{ and } j = 0, 1,$$
(22)

form a block system \mathcal{B} for G. Note that the quotient graph $B(\Gamma)/\mathcal{B}$ is a cycle of length 2d

In the next lemma we exclude certain possibilities for (a, r).

Lemma 6.4. Assuming Hypothesis 6.1, suppose that m = 2l for some odd number l. Then

$$(a,r) \notin \{(l+2,l+1),(l+2,l-1),(l-2,l+1),(l-2,l-1)\}.$$

Proof. On the contrary, assume that (a,r) is one of the pairs given in the lemma. By Lemma 6.3, $L \triangleleft G$. As $0 < a,r \le m$, $\phi_{n,m}(a) = a$ and $\phi_{n,m}(r) = r$, and hence $B(\Gamma)/L \cong B(R_m(a,r))$ due to Lemma 3.6(ii). Applying Lemma 6.3 to $B(R_m(a,r))$, we obtain $\langle \rho^l \rangle / L \triangleleft G/L$, and hence $\langle \rho^l \rangle \triangleleft G$.




FIGURE 2. The 4-cycles X_1, X_2, X_3 and X_4 in the graph $B(R_{4l}(l+2, l+1))$.

If l=3, then using the computer algebra package MAGMA [2], we find that if (a,r)=(l+2,l+1)=(5,4) or (a,r)=(l-2,l-1)=(1,2), then B(Γ) is not edge-transitive; and if (a,r)=(l+2,l-1)=(5,2) or (a,r)=(l-2,l+1)=(1,4), then B(Γ) is edge-transitive, but Γ is stable. Both cases are impossible, hence in the rest of the proof we assume that l>3. Then G admits the block system \mathcal{B} with blocks $\Delta_{i,j}$ defined in (22) with d=l. Let K be the kernel of the action of G on \mathcal{B} .

Case 1.
$$(a,r) = (l+2, l+1)$$
 or $(l-2, l-1)$.

We discuss only the case when (a,r)=(l+2,l+1) because the other case can be handled by copying the same argument. For $i \in \mathbb{Z}_l$ and $j \in \mathbb{Z}_2$,

$$B(\Gamma)[\Delta_{i,j} \cup \Delta_{i+1,j+1}] \cong 4\mathbf{C}_4.$$

Clearly, the automorphisms in K permute these 4-cycles. Consider the 4-cycles X_1, X_2, X_3 and X_4 shown in Figure 2. Choose $\sigma \in G_{u_{0,0}}$ such that σ swaps $u_{1,1}$ and $v_{0,1}$. Then $\sigma \in K$, it maps each of X_1 and X_2 to itself, and it swaps X_3 and X_4 . Figure 2 shows us that $\sigma(u_{l+1,1}) = v_{3l,1}$. On the other hand, as X_1 is preserved by σ , $\sigma(v_{-1,0}) = v_{-1,0}$. These give rise to a contradiction because $u_{l+1,1} \sim v_{-1,0}$ but $v_{3l,1} \not\sim v_{-1,0}$.

Case 2.
$$(a,r) = (l+2, l-1)$$
 or $(l-2, l+1)$.

We discuss only the case when (a,r)=(l+2,l-1) because the other case can be handled in the same way. For $i \in \mathbb{Z}_l$ and $j \in \mathbb{Z}_2$,

$$B(\Gamma)[\Delta_{i,j} \cup \Delta_{i+1,j+1}] \cong 2\mathbf{C}_8.$$

The subgraph of $B(\Gamma)$ induced by the set $\Delta_{0,0} \cup \cdots \cup \Delta_{12,0}$ is shown in Figure 3. Label the vertices in the blocks $\Delta_{i,j}$ with numbers from 1 to 8 as shown in the figure. For $\sigma \in K$ and for an integer $k \geq 0$, let σ_k be the permutation of $\{1,\ldots,8\}$ induced by the restriction of σ to $\Delta_{i,j}$, where $k \equiv i \pmod{l}$ and $k \equiv j \pmod{2}$. Note that $\sigma_i = \sigma_j$ holds if $i \equiv j \pmod{2l}$. The 8-cycles intersect at either no vertex or a pair of antipodal vertices. Moreover, each 8-cycle between $\Delta_{i,j}$ and $\Delta_{i+1,j+1}$ has every other vertex in $\Delta_{i,j}$. These imply that σ_{i+1} is determined by σ_i .

Choose $\sigma \in G_{u_{0,0}} \cap G_{u_{1,1}}$. Clearly, $\sigma \in K$ and

$$\sigma_0(i) = \sigma_1(i) = i \text{ if } 1 \le i \le 4.$$




FIGURE 3. The subgraph of the canonical double cover $B(R_{4l}(l+2, l-1))$ induced by the set $\Delta_{0,0} \cup \cdots \cup \Delta_{12,0}$.

Denote by X_1 and X_2 the two 8-cycles between $\Delta_{1,1}$ and $\Delta_{2,0}$ such that $u_{1,1} \in V(X_1)$. Then σ maps each of X_1 and X_2 to itself, and it is easy to see that it acts on these cycles either as the identity or a specific reflection. Denote by X_3 the 8-cycle between $\Delta_{0,0}$ and $\Delta_{1,1}$ not through $u_{0,0}$. It follows that a vertex of X_3 is either fixed by σ or it is switched with the vertex antipodal to it in X_3 . This implies that σ acts on X_3 either as the identity or the reflection with respect to the center of X_3 . Therefore, σ_0 is the identity permutation or $\sigma_0 = (5,7)(6,8)$.

Assume for the moment that $\sigma_0 = (5,7)(6,8)$. Then

$$\sigma_0 = \sigma_1$$
 and $\sigma_2 = (1,6)(2,5)(3,8)(4,7)$.

A tedious, although straightforward computation, yields that

$$\sigma_3 = (1,5)(2,8)(3,7)(4,6), \ \sigma_4 = (1,5)(2,8)(3,7)(4,6), \ \sigma_5 = (1,8)(2,7)(3,6)(4,5),$$

$$\sigma_6 = \sigma_7 = (1,3)(2,4), \ \sigma_8 = (1,8)(2,7)(3,6)(4,5), \ \sigma_9 = (1,7)(2,6)(3,5)(4,8),$$

$$\sigma_{10} = (1,7)(2,6)(3,5)(4,8), \ \sigma_{11} = \sigma_2, \ \sigma_{12} = \sigma_0.$$

This shows that $\sigma_i = \sigma_j$ if and only if $i \equiv j \pmod{12}$. On the other hand, $\sigma_0 = \sigma_{2l}$, contradicting our assumption that l is odd. We conclude that σ_0 must be the identity, implying that $G_{u_{0,0}} \cap G_{u_{1,1}} = 1$, whence $|G_{u_{0,0}}| = 4$.

It can be shown in the same way as above that the existence of a non-identity automorphism in $K_{u_{0,0}}$ forces that 3 divides l. But then $\Gamma = R_{12s}(3s + 2, 3s - 1)$, where l = 3s, and therefore, it is edge-transitive due to Proposition 2.7. This and the fact that $|G_{u_{0,0}}| = 4$ show that Γ is stable, a contradiction.

Lemma 6.5. Assuming Hypothesis 6.1, suppose that m is even. Then Γ/L is unstable.

Proof. Let $M = \langle \rho^m, \beta \rangle$, where β is the automorphism of B(Γ) defined in (8). On the contrary, assume that Γ/L is stable. It follows from this and Lemma 3.3 that $M \triangleleft G$. By Lemma 6.3, $\rho^m \in Z(G)$, implying that $C_G(M) = C_G(\beta)$. Consider the action of G on itself by conjugaction. By the orbit-stabilizer lemma, $|G| = |C_G(\beta)| \cdot |\beta^G|$, where β^G

denotes the conjugacy class of G containing β . Using also that $\beta \notin Z(G)$ and $M \triangleleft G$, we find that $\beta^G = \{\beta, \beta \rho^m\}$, and therefore, $C_G(M) = C_G(\beta)$ has index 2 in G. Note that the group $C_G(\beta)$ is the group of expected automorphisms of $B(\Gamma)$ due to Lemma 3.3, in particular,

$$C_G(M) \cong \operatorname{Aut}(\Gamma) \times \mathbb{Z}_2.$$
 (23)

Since $B(R_m(a,r))$ is edge-transitive and $R_m(a,r)$ is stable, it follows that $R_m(a,r)$ is edge-transitive. Let $\Sigma = R_m(a',r')$, where $a' = \min(a,m-a)$ and $r' = \min(r,m-r)$. Then Σ is one of the graphs in the families (a)–(d) in Proposition 2.7. If $R_m(a,r)$ is bipartite, then so is $\Gamma = R_n(a,r)$, see Lemma 3.2, which is not the case. Thus $R_m(a,r)$ is not bipartite, which yields that it is neither in the family (a) nor in (d). Lemma 6.4 implies that it cannot be in the family (b). Thus Σ is in the family (c), i.e., m = 12l and (a',r') = (3l+2,3l-1) or (3l-2,3l+1). Then

$$(a,r) \in \{ (3l+2, 3l-1), (3l-2, 3l+1), (9l-2, 9l+1), (9l+2, 9l-1) \}.$$
 (24)

Note that l must be odd because $\Gamma = R_n(a, r)$ is not bipartite. One can check that $r^2 \not\equiv 1 \pmod{n}$. Combining this with Propositions 2.4–2.7 yields $|\operatorname{Aut}(\Gamma)| = 2n$. Using also (23), we find $C_G(M) = \langle \rho, \mu, \beta \rangle$, where μ is the automorphism of B(Γ) defined in (7).

Choose $\sigma \in G_{u_{0,0}}$ such that $\sigma(u_{1,1}) = v_{0,1}$. Then $\sigma \notin C_G(M)$. Recall that G^+ is the subgroup of G that preserves the biparts of $B(\Gamma)$. Then $G^+ \cap C_G(M) = \langle \rho, \mu \rangle$. Since $\sigma \in G^+$, it follows that σ normalizes $\langle \rho, \mu \rangle$, and therefore, it normalizes also $C = \langle \rho \rangle$. The proof of Lemma 6.2 shows that a is even. This is in contradiction with (24) and the fact that l is odd.

We are ready to handle the case when C is not normal in G.

Lemma 6.6. Assuming Hypothesis 6.1, suppose that $C \not\subset G$. Then m is odd and (a,r)=(m-2,m-1) or (2,m-1).

Note that if (a, r) = (m-2, m-1), then Γ is in the family W2, and if (a, r) = (2, m-1), then it is in the family W4.

Proof. We proceed by induction on n. With the help of the computer algebra package MAGMA [2] we checked that the lemma holds if $n \leq 12$. From now on it is assumed that n > 12.

Claim. m is odd

Proof of the claim. On the contrary, assume that m is even, say m = 2l. By Lemma 6.5, $R_m(a,r)$ is unstable. The graph $R_m(a,r)$ is not bipartite, for otherwise, so is $\Gamma = R_n(a,r)$ due to Lemma 3.2, which is contrary to Hypothesis 6.1.

Assume for the moment that $R_m(a,r)$ has two vertices with the same neighborhood. Consequently, $B(R_m(a,r))$ has two vertices having identical neighborhood too. As $B(R_m(a,r))$ is vertex-transitive, see the remark after Hypothesis 6.1, there exists a vertex of $B(\Gamma)$, which is different from $u_{0,0}$ and has the same neighborhood as $u_{0,0}$. On the other hand, a direct check shows that $u_{i,j}$ and $u_{0,0}$ have different neighborhood if $(i,j) \neq (0,0)$; whereas $v_{i,j}$ and $u_{0,0}$ share the same neighborhood if and only if $i \equiv \pm 1 \pmod{m}$, j = 0, $a \equiv \pm 2 \pmod{m}$ and $r \equiv \pm 1 \pmod{m}$. But then a is even and r is odd, contradicting the fact that $R_m(a,r)$ is not bipartite. To sum up, $R_m(a,r)$ is non-trivially unstable.




FIGURE 4. Part of the 16-cycles X_1, X_2 and X_3 in the graph $B(R_{4l}(2, l-1))$.

Note that C/L is not normal in G/L, and therefore, the induction hypothesis can be applied to $R_m(a, r)$. As a result, l is odd and

$$(\min(a, m-a), \min(r, m-r)) = (l-2, l-1) \text{ or } (2, l-1).$$

Combining this with Lemma 6.3 yields

$$(a,r) \in \{(2,l-1), (2,l+1), (m-2,l-1), (m-2,l+1)\}.$$

We consider only the case when (a, r) = (2, l - 1), the other pairs can be handled in the same way. The first lines in the proof of Lemma 6.4 can be applied again and this results in the block system \mathcal{B} for G with blocks $\Delta_{i,j}$ defined in (22) with d = l (now l > 3 because n > 12). The quotient graph $B(\Gamma)/\mathcal{B}$ is a cycle of length 2l. Let K be the kernel of the action of G on \mathcal{B} . For $i \in \mathbb{Z}_l$, $j \in \mathbb{Z}_2$,

$$B(\Gamma)[\Delta_{i,j} \cup \Delta_{i+1,j+1}] \cong \mathbf{C}_{16}.$$

Consider part of the 16-cycles X_1, X_2 and X_3 shown in Figure 4. Choose $\sigma \in G_{u_{0,0}}$ such that σ swaps $u_{1,1}$ and $v_{0,1}$. Then σ maps each of X_1, X_2 and X_3 to itself. Considering the action of σ on X_1 , we find $\sigma(u_{2,0}) = u_{2,0}$. Then considering its action on X_2 yields $\sigma(v_{1,0}) = v_{3l+1,0}$. Finally, considering its action on X_3 yields $\sigma(u_{3,1}) = v_{2,1}$. This gives rise to a contradiction because $v_{1,0} \sim u_{3,1}$ and $v_{3l+1,0} \not\sim v_{2,1}$.

Since m is odd, it follows that

$$B(\Gamma/L) \cong B(R_m(a,r)) \cong R_{2m}(a',r'),$$

where a' and r' are defined in (10). Let $\Sigma = R_{2m}(a'', r'')$ where $a'' = \min(a', 2m - a')$ and $r'' = \min(r', 2m - r')$. By Proposition 2.3, Σ is edge-transitive, hence it is one of the graphs in the families (a)–(d) of Theorem 2.7.

The fact that Σ is bipartite shows that Σ cannot be in the family (b). As m is odd, it is clear that it cannot be in the family (c) neither. If Σ is in the family (d), then it follows that $C \triangleleft G$; this is impossible. We conclude that Σ is in the family (a), i.e., $a' \equiv \pm 2 \pmod{n}$ and $r' \equiv \pm 1 \pmod{n}$. If a is even, then $a' = a \leq m$, hence a = 2; and if a is odd, then a' = a + m > m, hence a = m - 2. Similarly, we find that r = 1 or m - 1. As $(a, r) \neq (2, 1)$, because $R_n(a, r)$ is not bipartite, we have that

$$(a,r) \in \{(2,m-1),(m-2,1),(m-2,m-1)\}.$$




FIGURE 5. The subgraph of $B(R_{2m}(m-2,1))$ induced by the set $\Delta_{0,0} \cup \cdots \cup \Delta_{4,0}$.

We finish the proof by excluding the possibility (a,r)=(m-2,1). On the contrary, assume that (a,r)=(m-2,1). Then G admits the block system \mathcal{B} with blocks $\Delta_{i,j}$ defined in (22) with l=m. Let K be the kernel of the action of G on \mathcal{B} . Label the vertices in $\Delta_{i,j}$ with numbers from 1 to 4 as shown in Figure 5. For $\sigma \in K$ and for an integer $k \geq 0$, let σ_k be the permutation of $\{1,2,3,4\}$ induced by the restriction of σ to $\Delta_{i,j}$, where $k \equiv i \pmod{m}$ and $k \equiv j \pmod{2}$. Note that $\sigma_i = \sigma_j$ holds if $i \equiv j \pmod{2m}$. Choose $\sigma \in G_{u_{0,0}}$ such that σ swaps $u_{1,1}$ and $v_{m+2,1}$. Then we compute

$$\sigma_0 = (2,4), \ \sigma_1 = (1,2)(3,4), \ \sigma_2 = (1,3), \ \sigma_3 = (1,4)(2,3), \ \sigma_4 = \sigma_0.$$

This shows that $\sigma_i = \sigma_j$ if and only if $i \equiv j \pmod{4}$. On the other hand, $\sigma_0 = \sigma_{2m}$, contradicting the fact that m is odd.

Acknowledgement

The authors are grateful to the anonymous referees for their valuable comments which helped to improve the text considerably.

REFERENCES

- [1] N. Biggs, Algebraic graph theory, Cambridge Univ. Press, New York 1992.
- [2] W. Bosma, C. Cannon, C. Playoust, The Magma algebra system I: The user language, J. Symbolic Comput., 24 (1997), 235–265.
- [3] J. D. Dixon, B. Mortimer, *Permutation groups*, Graduate Text in Mathematics 163, Springer-Verlag, New York 1996.
- [4] E. Dobson, I. Kovács, Š. Miklavič, The automorphism groups of non-edge-transitive rose window graphs, Ars. Math. Contemp., 9 (2015), 63–75.
- [5] B. Fernandez, A. Hujdurović, Canonical double covers of circulants, J. Comb. Theory Ser. B, 154 (2022), 49–59.
- [6] M. Herzog, G. Kaplan, Large cyclic subgroups contain non-trivial normal subgroups, J. Group Theory, 4 (2001), 247–253.
- [7] I. Hubard, A. Ramos-Riviera, P. Šparl, Arc-transitive maps with underlying Rose Window graphs, J. Graph Theory, 96 (2021), 203–230.
- [8] A. Hujdurović, D. Mitrović, Some conditions implying stability of graphs, J. Graph Theory, 105 (2024), 98–109.
- [9] A. Hujdurović, D. Mitrović, D. Witte Morris, On the automorphisms of the double cover of a circulant graph, Electron. J. Combin., 28 (2021), #P4.43.

- [10] A. Hujdurović, D. Mitrović, D. Witte Morris, Automorphisms of the double cover of a circulant graph of valency at most 7, Algbr. Comb., 6 (2023), 1235–1271.
- [11] D. Jokanović, Š. Miklavič, M. Milićević, P. Šparl, Domination in Rose window graphs, Bull. Malays. Math. Sci. Soc., 44 (2021), 509–526.
- [12] I. Kovács, K. Kutnar, D. Marušič, Classification of edge-transitive rose window graphs, J. Graph Theory, 65 (2010), 216–231.
- [13] I. Kovács, K. Kutnar, J. Ruff, Rose window graphs underlying rotary maps, Discrete Math., 310 (2010), 1802–1811.
- [14] P. Lorimer, Vertex-transitive graphs: symmetric graphs of prime valency, J. Graph Theory, 8 (1984), 55–68.
- [15] A. Lucchini, On the order of transitive permutation groups with cyclic point-stabiliser, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei, (9) Mat. Appl., 9 (1998), 241–243.
- [16] D. Marušič, R. Scapellato, N. Zagaglia Salvi, A characterization of particular symmetric (0,1) matrices, Linear Algebra Appl., 119 (1989), 153–162.
- [17] R. Nedela, M. Škoviera, Regular embeddings of canonical double coverings of graphs, J. Combin. Theory Ser. B, 67 (1996), 249–277.
- [18] Y. L. Qin, B. Xia, S. Zhou, Stability of circulant graphs, J. Combin. Theory Ser. B, 136 (2019), 154–169.
- [19] Y. L. Qin, B. Xia, S. Zhou, Canonical double covers of generalized Petersen graphs, and double generalized Petersen graphs, J. Graph Theory, 97 (2021), 70–81.
- [20] D. B. Surowski, Stability of arc-transitive graphs, J. Graph Theory, 38 (2001), 95–110.
- [21] S. Wilson, Rose Window graphs, Ars Math. Contemp., 1 (2008), 7–19.
- [22] S. Wilson, Unexpected symmetries in unstable graphs, J. Comb. Theory Ser. B, 98 (2008), 359–383.
- [23] D. Witte Morris, On automorphisms of direct products of Cayley graphs on abelian groups, Electron. J. Combin., 28 (2021), #P3.5.
- [24] D. Witte Morris, Automorphisms of the canonical double cover of a toroidal grid, preprint arXiv:2301.05396 [math.CO] (2023).

M. Ahanjideh, I. Kovács, K. Kutnar

UP IAM, University of Primorska, Muzejski trg 2, SI-6000 Koper, Slovenia

UP FAMNIT, University of Primorska, Glagoljaska ulica 8, SI-6000 Koper, Slovenia

Email address: milad.ahanjideh@upr.si
Email address: istvan.kovacs@upr.si
Email address: klavdija.kutnar@upr.si