

DOTTED LIMITS

JOANNA KO

ABSTRACT. Marked limits, or Cartesian quasi-limits introduced by Gray, give an alternative approach to **Cat**-weighted limits in 2-category theory. This was first established by Street, and we aim to give a new approach to this result using marked codescent objects of marked coherence data which we introduce in this article. We then propose the notion of dotted limits, which is a natural generalisation of marked limits to the enhanced 2-categorical setting. We establish that dotted limits and \mathcal{F} -weighted limits both have the same expressive power.

1. INTRODUCTION

Marked limits¹ of 2-functors were first introduced by Gray in [Gra74], under the name *Cartesian quasi-limits*. Later in [Szy19], Szyld investigated the notion under the name σ -*s-limits*. In Mesiti's recent work [Mes23], this notion was studied in a more restrictive sense under the name *lax normal conical 2-limits*.

Marked limits are gaining attention in 2-category theory, mainly due to their convenience and simplicity. In contrast, the notion of **Cat**-weighted limits, though captures all the 2-categorical limits that we are interested in, is sometimes convoluted, as it is designed for categories enriched in any arbitrary closed symmetric monoidal category \mathbf{V} , but not exclusively for **Cat**. On the other hand, marked limits often provide a clearer and a more spread-out presentation of 2-dimensional limits, which makes them more convenient to track and handle. Indeed, marked limits are very similar to lax limits, in the sense that both of them consider non-strict natural transformations; while the latter considers lax natural transformations alone, the former considers lax natural transformations that restricts to 2-natural transformations with respect to a specific class of morphisms in the domain 2-category. This makes the notion more flexible, by allowing different kind of shapes for the cones rather than only triangles, and is the reason why marked limits should be able to capture all 2-categorical limits of interest.

Seeing that marked limits possess some advantages over **Cat**-weighted limits, it is then crucial to show that marked limits are as expressive as **Cat**-weighted limits, so that any 2-categorical limits of interest can be described using marked limits.

It is well-known that a **Cat**-weighted limit can be expressed as a marked-lax limit with the same universal property. This has first been discussed in Street's paper [Str76], and also in the recent work [Mes23]. The converse, which says that an marked-lax limit can be turned into a **Cat**-weighted limit with the same universal property, has also been established in [Str76]. In this article, we provide an alternative proof via a new perspective.

Our proof is done by constructing the weak morphism classifier, which will then give us the left adjoint $(\)^\ddagger$ to the inclusion $[\mathcal{C}, \mathbf{Cat}] \hookrightarrow [\mathcal{C}, \mathbf{Cat}]_{l, \Sigma}$ of the category of 2-functors $\mathcal{C} \rightarrow \mathbf{Cat}$ and 2-natural transformations to that of marked-lax natural transformations. The construction of the weak morphism classifier relies on the notion of *marked codescent objects of marked coherence data*, which is a modified version of lax codescent objects of strict coherence data studied by Lack in [Lac02].

Key words and phrases. dotted limit; marked limit; codescent objects; 2-monads; enhanced 2-category theory; \mathcal{F} -categories.

¹We prefer the name marked limits, because it hints that certain collection of morphisms are special and recorded. We borrow this terminology from marked simplicial sets. Apart from that, we discovered that in [GHL22], the authors used the name *marked limits* for the case which replaces the identity 2-cells in our case with invertible 2-cells. This notion was first considered in [Gra74], and recently in [DDS18]. We believe that this terminology is convenient and accurate in depicting the situation.

Next, we introduce the notion of *dotted limits*, which is the enhanced 2-categorical counterpart of marked limits.

The study of enhanced 2-category theory began in [LS12] by Lack and Shulman. The main reason for developing the theory is that, indeed, many interesting categorical structures in Categorical Algebra are T -algebras, for a 2-monad T on a 2-category \mathcal{A} ; most of the time, however, the morphisms between them are the pseudo or lax T -morphisms, instead of the strict T -morphisms. For instance, lax monoidal functors and pseudo (strong) monoidal functors between monoidal categories are more common than strict monoidal functors. Therefore, studying $\mathbf{T}\text{-Alg}_w$ for $w = l, p, c$ becomes salient. In particular, whether or not $\mathbf{T}\text{-Alg}_w$ admits all the limits that \mathcal{A} possesses is a key question. In [Lac05], Lack has shown that the answer to the question in the case of $\mathbf{T}\text{-Alg}_l$ depends heavily on the relationship between $\mathbf{T}\text{-Alg}_s$ and $\mathbf{T}\text{-Alg}_l$. Thus, instead of viewing them as separate 2-categories, we should combine $\mathbf{T}\text{-Alg}_s$ and $\mathbf{T}\text{-Alg}_l$ into one single structure, which is the starting point of \mathcal{F} -categories and enhanced 2-category theory.

In our paper, we propose the notion of dotted limits, which, very much in the same spirit of marked limits, often provides a more convenient expression and simpler description than \mathcal{F} -weighted limits. To illustrate its convenience, we describe several examples of \mathcal{F} -categorical limits as dotted limits.

Our main results in the paper:

Main Theorem 5.3.6. The dotted-lax limit of an \mathcal{F} -functor $S: \mathbb{D} \rightarrow \mathbb{A}$ has the same universal property as the \mathcal{F} -weighted limit $\{\Delta(1)^\#, S\}$.

Main Theorem 5.3.9. Let $\Phi: \mathbb{D} \rightarrow \mathbb{F}$ be an \mathcal{F} -weight and $S: \mathbb{D} \rightarrow \mathbb{A}$ be an \mathcal{F} -functor. Let $\Sigma := \{(d \in \text{mor } \mathbb{D}, 1)\}$ be the class of morphisms, and $T := \{(D, \delta \in (\Phi D)_\tau)\}$ be a collection of objects in the \mathcal{F} -category of elements $\mathbb{E}\mathbb{1}\Phi$ of Φ . The \mathcal{F} -weighted limit $\{\Phi, S\}$ has the same universal property as the dotted-lax limit of the \mathcal{F} -functor $SP: (\mathbb{E}\mathbb{1}\Phi, \Sigma, T) \rightarrow \mathbb{A}$.

say that any dotted limits can be equivalently expressed as \mathcal{F} -weighted limits, and vice versa. The latter can be done in a similar fashion as in the 2-categorical case, whereas the former is done by constructing the tight part for the left adjoint, using mainly left Kan extensions and embedded image factorisations.

The outline of the paper goes as follows. In Section 2, we briefly recall the preliminary notions, particularly the basics of 2-monad theory. In Section 3, we begin by recalling marked limits and move on to introduce the notion of marked codescent objects of marked coherence data, and then give a complete and detailed proof of that any marked limit has the same universal property as a **Cat**-weighted limit. In Section 4, we recall the basics of enhanced 2-category theory, including notions such as \mathcal{F} -categories and \mathcal{F} -weights. In Section 5, we introduce the notion of dotted limits, and present several important examples; last but not least, we show the equivalence of dotted limits and \mathcal{F} -weighted limits.

ACKNOWLEDGEMENT

The author would like to thank John Bourke for his clear guidance, beneficent suggestions and inspiring opinions, which helps her get more familiar with enhanced 2-category theory and advances her knowledge of category theory to a huge extent. The author is also grateful to Giacomo Tendas for his helpful comments and revisions.

2. PRELIMINARIES ON BASIC CATEGORY THEORY, ENRICHED CATEGORY THEORY AND 2-CATEGORY THEORY

2.1. Weighted limits. A standard reference for enriched category theory is Kelly's book [Kel82].

Let \mathcal{V} be a closed symmetric monoidal category, \mathcal{C}, \mathcal{D} be two \mathcal{V} -categories, $W: \mathcal{C} \rightarrow \mathcal{V}$ be a \mathcal{V} -weight, and $F: \mathcal{C} \rightarrow \mathcal{D}$ be a \mathcal{V} -functor.

Definition 2.1.1. *The \mathcal{V} -weighted limit $\{W, F\}$ of F by W is characterised by an isomorphism*

$$(2.1) \quad \mathcal{D}(D, \{W, F\}) \cong [\mathcal{C}, \mathcal{V}](W, \mathcal{D}(D, F-))$$

Let \mathcal{C} be a 2-category. It is worth mentioning how the presheaf 2-category $[\mathcal{C}, \mathbf{Cat}]$ corresponds to the 2-category of (strict) T -algebras, strict T -morphisms, and T -transformations, whereas $[\mathcal{C}, \mathbf{Cat}]_l$ corresponds to the 2-category of (strict) T -algebras, lax T -morphisms, and T -transformations, for some 2-monad T , because this is the key motivation to our proposal of codescent objects of marked coherence data in Section 3.2.

Let $E: \text{ob } \mathcal{C} \rightarrow \mathcal{C}$ denote the canonical identity-on-object 2-functor.

There is a standard result:

Lemma 2.2.6. *The left Kan extension along E is left 2-adjoint to the pre-composition with E :*

$$\begin{array}{ccc} & \text{Lan}_E & \\ & \curvearrowright & \\ [\mathcal{C}, \mathbf{Cat}] & \perp & [\text{ob } \mathcal{C}, \mathbf{Cat}] \\ & \curvearrowleft & \\ & E^* & \end{array} .$$

Moreover, for any 2-functor $X: \mathcal{C} \rightarrow \mathbf{Cat}$, there is a formula:

$$(2.3) \quad \text{Lan}_E X = \sum_{d \in \mathcal{C}} \mathcal{C}(d, -) \times X(d).$$

Proof. The adjunction follows immediately from the universal property of the left Kan extension of any functor $\mathcal{C} \rightarrow \mathbf{Cat}$ along E . \square

Now, we define a 2-monad $T := E^* \text{Lan}_E: [\text{ob } \mathcal{C}, \mathbf{Cat}] \rightarrow [\text{ob } \mathcal{C}, \mathbf{Cat}]$, with multiplication and unit given by

$$\begin{aligned} (\mu_X)_c: \sum_{d_2 \in \mathcal{C}} \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_2, c) \times \mathcal{C}(d_1, d_2) \times \mathcal{C}(d, d_1) \times X(d) &\rightarrow \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times X(d), \\ (f: d_1 \rightarrow c, g: d \rightarrow d_1, x \in X(d)) &\mapsto (fg: d \rightarrow c, x \in X(d)), \end{aligned}$$

and

$$\begin{aligned} (\eta_X)_c: X(c) &\rightarrow \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times X(d), \\ x &\mapsto (1_c, x), \end{aligned}$$

respectively.

Our later construction of marked codescent objects of marked coherence data depends on the correspondence between the data of lax T -morphisms and that of lax natural transformations.

Theorem 2.2.7 ([BKP89, Example 6.6]). *The 2-category $\mathbf{T}\text{-Alg}_l$ of (strict) T -algebras, lax T -morphisms, and T -transformations is isomorphic to $[\mathcal{C}, \mathbf{Cat}]_l$.*

Similarly, the 2-category $\mathbf{T}\text{-Alg}_s$ of (strict) T -algebras, strict T -morphisms, and T -transformations is isomorphic to $[\mathcal{C}, \mathbf{Cat}]$.

Proof. A strict T -algebra is a pair $(A \in [\text{ob } \mathcal{C}, \mathbf{Cat}], a: TA \rightarrow A)$ satisfying

$$\begin{array}{ccc} T^2 A & \xrightarrow{Ta} & TA & & A & \xrightarrow{\eta_A} & TA \\ \mu_A \downarrow & & \downarrow a & , & \searrow 1_A & & \downarrow a \\ TA & \xrightarrow{a} & A & & & & A \end{array} .$$

Note that $a: TA \rightarrow A = \sum_{d \in \mathcal{C}} \mathcal{C}(d, -) \times A(d) \rightarrow A$ is a 2-natural transformation with component $a_c: \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) \rightarrow A(c)$, which is simply a functor $a_{c,d}: \mathcal{C}(d, c) \times A(d) \rightarrow A(c)$ for a choice of $d \in \text{ob } \mathcal{C}$. Now by the tensor-hom adjunction, this is equivalent to a functor

$$\mathcal{C}(d, c) \rightarrow \mathbf{Cat}(A(d), A(c)).$$

So the data of a structure morphism $a: TA \rightarrow A$ is equivalent to a functor $A(f): A(d) \rightarrow A(c)$ induced by a morphism $f: d \rightarrow c$ in \mathcal{C} , and a natural transformation

$$\begin{aligned} A(\gamma): A(f_1) &\rightarrow A(f_2), \\ (x_1 \mapsto y_1) &\mapsto (x_2 \mapsto y_2), \end{aligned}$$

induced by a 2-cell $\gamma: f_1 \Rightarrow f_2$ in \mathcal{C} , where $f_i: d \rightarrow c$ and $x_i \in A(d)$, for $i = 1, 2$. These are precisely the data of a 2-functor $A: \mathcal{C} \rightarrow \mathbf{Cat}$.

Next, for an object $(f: d_1 \rightarrow c, g: d \rightarrow d_1, x \in A(d))$ of $\sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \mathcal{C}(d, d_1) \times A(d)$, the commutative diagrams above amount to

$$(a \cdot Ta)_c(f, g, x) = a(f, A(g)(x)) = A(fg)(x),$$

and similarly, $A(f)(A(g))(\chi) = A(fg)(\chi)$ for a morphism $\chi: x_1 \rightarrow x_2$ in $A(d)$; and amount to

$$(a \cdot \eta_A)_c(x) = a(1_c, x) = A(1_c)(x) = 1_{A(c)}(x),$$

for an object x of $A(c)$, and similarly, $A(1_c)(\chi) = 1_{A(c)}(\chi)$ for a morphism $\chi: x_1 \rightarrow x_2$ in $A(c)$. These are precisely the functoriality and unity of the 2-functor $A: \mathcal{C} \rightarrow \mathbf{Cat}$.

Therefore, a strict T -algebra (A, a) corresponds to a 2-functor $A: \mathcal{C} \rightarrow \mathbf{Cat}$.

Let $(F: A \rightarrow B, \overline{F}: b \cdot TF \Rightarrow F \cdot a)$ be a lax T -morphism. Then, we have the coherence

$$(2.4) \quad \begin{array}{ccc} T^2 B & \xrightarrow{ Tb } & TB \\ T^2 F \nearrow & \mu_B & \searrow b \\ T^2 A & \xrightarrow{ TF } & TB \\ \mu_A \searrow & & \nearrow F \\ TA & \xrightarrow{ a } & A \end{array} \quad = \quad \begin{array}{ccc} T^2 B & \xrightarrow{ Tb } & TB \\ T^2 F \nearrow & \downarrow T\overline{F} & \searrow b \\ T^2 A & \xrightarrow{ TF } & TA \\ \mu_{TA} \searrow & & \nearrow F \\ TA & \xrightarrow{ a } & A \end{array} \quad = \quad \begin{array}{ccc} TB & \xrightarrow{ b } & B \\ \mu_B \nearrow & \downarrow \overline{F} & \searrow F \\ TA & \xrightarrow{ a } & A \\ \mu_A \searrow & & \nearrow F \\ A & \xrightarrow{ a } & A \end{array} \quad ,$$

$$(2.5) \quad \begin{array}{ccc} B & \xrightarrow{ 1 } & B \\ F \nearrow & \eta_B & \searrow b \\ A & \xrightarrow{ TF } & TB \\ \eta_A \searrow & & \nearrow F \\ TA & \xrightarrow{ a } & A \end{array} \quad = \quad \begin{array}{ccc} B & \xrightarrow{ 1 } & B \\ F \nearrow & \downarrow \overline{F} & \searrow F \\ A & \xrightarrow{ TF } & TA \\ \eta_A \searrow & & \nearrow F \\ TA & \xrightarrow{ a } & A \end{array} \quad .$$

Note that the modification $\overline{F}: b \cdot TF \Rightarrow F \cdot a$ has its component at $c \in \text{ob } \mathcal{C}$ being a natural transformation $\overline{F}_c: (b \cdot TF)_c \rightarrow (F \cdot a)_c$, which means we have

$$\begin{array}{ccc} \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) & \xrightarrow{ a_c } & A(c) \\ (TF)_c \downarrow & \cong_{\overline{F}_c} & \downarrow F(c) \\ \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times B(d) & \xrightarrow{ b_c } & B(c) \end{array}$$

which is equivalent to the following natural transformation $\overline{F}_{c,d}$ for a chosen $d \in \text{ob } \mathcal{C}$

$$\begin{array}{ccc} \mathcal{C}(d, c) \times A(d) & \xrightarrow{ a_{c,d} } & A(c) \\ \mathcal{C}(d, c) \times F(d) \downarrow & \cong_{\overline{F}_{c,d}} & \downarrow F(c) \\ \mathcal{C}(d, c) \times B(d) & \xrightarrow{ b_{c,d} } & B(c) \end{array}$$

Let (f, x) be an object of $\mathcal{C}(d, c) \times A(d)$. The above commutative diagram then gives

$$B(f)(F(d)(x)) = F(c)(A(f)(x)).$$

So the component $\overline{F}_{c,d(f,x \in A(d))}: b_{c,d} \cdot (\mathcal{C}(d, c) \times F(d))(f, x) \rightarrow F(c) \cdot a_{c,d}(f, x)$ amounts to the component $B(f)(F(d)(x)) \rightarrow F(c)(A(f)(x))$ of a natural transformation

$$\begin{array}{ccc} A(d) & \xrightarrow{ F(d) } & B(d) \\ A(f) \downarrow & \cong_{\overline{F}_{c,d,f}} & \downarrow B(f) \\ A(c) & \xrightarrow{ F(c) } & B(c) \end{array}$$

and this is precisely the data of a lax natural transformation $F: A \rightarrow B$.

From Equation (2.5), we obtain $\overline{F}_{c,c,1_c} = 1$ as $(a \cdot \eta_A)_c = A(1_c)$. Now, for an object $(f: d \rightarrow c, g: e \rightarrow d, y \in A(e))$ of $\mathcal{C}(d, c) \times \mathcal{C}(e, d) \times A(e)$, we have

$$(\overline{F} \cdot \mu_A)_{c,d,e(f,g,y)} = \overline{F}_{c,e,fg(y)}, \quad ((\overline{F} \cdot T a) \circ (b \cdot T \overline{F}))_{c,d,e(f,g,y)} = (\overline{F}_{c,d,f} \circ A(g))_y \circ (B(f) \circ \overline{F}_{d,e,g})_y.$$

By Equation (2.4), we obtain $\overline{F}_{c,e,fg} = (\overline{F}_{c,d,f} \circ A(g)) \circ (B(f) \circ \overline{F}_{d,e,g})$. Altogether, we recover the lax unity and the lax naturality of F .

Hence, a lax T -morphism (F, \overline{F}) corresponds to a lax natural transformation $F: A \rightarrow B$. If in addition (F, \overline{F}) is strict, then $\overline{F} = 1$ and hence it corresponds to a 2-natural transformation $F: A \rightarrow B$. \square

3. MARKED LIMITS AND THEIR EQUIVALENCE TO **Cat**-WEIGHTED LIMITS

3.1. Marked limits.

Definition 3.1.1. Let \mathcal{A} be a 2-category. Let Σ be a class of morphisms in \mathcal{A} , which contains all the identities and is closed under composition. The pair (\mathcal{A}, Σ) is called a marked 2-category.

Definition 3.1.2 ([Gra74, I,2, p.14, (i)]). Let (\mathcal{A}, Σ) be a marked 2-category. Let $F, G: \mathcal{A} \rightrightarrows \mathcal{B}$ be 2-functors. A marked-lax natural transformation $\alpha: F \rightarrow G$ between F and G is a lax natural transformation $\alpha: F \rightarrow G$ such that for any $f \in \Sigma$, the 2-component $\alpha_f = 1$.

Remark 3.1.3. It is important to point out that this notion is more general than the lax normal natural transformation considered in [Mes23], even though any 2-category \mathcal{A} can be recovered by considering the Grothendieck construction of $\Delta(1): \mathcal{A} \rightarrow \mathbf{Cat}$. We allow *different* classes Σ of morphisms in \mathcal{A} , as opposed to fixing the specific class $\Sigma = \{(f, 1)\}$ of morphisms in $\int^{\text{op}} \Delta(1)$.

Notation 3.1.4. We denote by $A \rightrightarrows B$ a morphism from A to B in Σ .

Remark 3.1.5. Similarly, we can talk about *marked-pseudo* or *marked-colax* natural transformations.

Notation 3.1.6. We denote the 2-category of 2-functors $\mathcal{A} \rightarrow \mathcal{B}$, marked-lax natural transformations, and modifications by $[\mathcal{A}, \mathcal{B}]_{l,\Sigma}$.

Notation 3.1.7. We denote by $\Sigma(a, b)$ the full subcategory of $\mathcal{A}(a, b)$ consisting of morphisms in Σ that have fixed source a and fixed target b .

Definition 3.1.8 ([Gra74, I,7, p.23, (iii)], [Szy19, Definition 2.4]). Let (\mathcal{A}, Σ) be a small marked 2-category, i.e., $\text{ob } \mathcal{A}$ is small, and let \mathcal{B} be a 2-category. The marked-lax limit $m_l \lim F$ of a 2-functor $F: \mathcal{A} \rightarrow \mathcal{B}$ is characterised by an isomorphism

$$(3.1) \quad \mathcal{B}(B, m_l \lim F) \cong [\mathcal{A}, \mathcal{B}]_{l,\Sigma}(\Delta(B), F)$$

in \mathbf{Cat} , which is natural in $B \in \text{ob } \mathcal{B}$.

Remark 3.1.9. We can talk about marked-colax or marked-pseudo limits, by replacing the marked-lax natural transformations with marked-colax or marked-pseudo natural transformations, respectively.

3.2. Equivalence to **Cat-weighted limits.** We now begin to show that a marked limit can be equivalently expressed as a **Cat**-weighted limit, which has first been shown in [Str76, Theorem 14], and we are going to provide a more detailed proof of the result.

To achieve our goal, we introduce the concept of marked codescent objects of marked coherence data, which is a modified version of lax codescent objects of strict coherence data studied in [Lac02].

Let (A, a) and (B, b) be strict T -algebras for a 2-monad T . In [Lac02], Lack has shown that a lax T -morphism is equivalent to a lax codescent cocone of the strict coherence data

$$T^3 A \begin{array}{c} \xrightarrow{\mu_{TA}} \\ \xrightarrow{T\mu_A} \\ \xrightarrow{T^2 a} \end{array} T^2 A \begin{array}{c} \xrightarrow{\mu_A} \\ \xrightarrow{T\eta_A} \\ \xrightarrow{Ta} \end{array} T A .$$

Now, let \mathcal{C} be an arbitrary 2-category, and $E: \text{ob } \mathcal{C} \rightarrow \mathcal{C}$ be the canonical identity-on-object 2-functor. Define $T := E^* \text{Lan}_E: [\text{ob } \mathcal{C}, \mathbf{Cat}] \rightarrow [\text{ob } \mathcal{C}, \mathbf{Cat}]$. Then T is a 2-monad on $[\text{ob } \mathcal{C}, \mathbf{Cat}]$, and from Theorem 2.2.7, a lax T -morphism in this case is a lax natural transformation in $[\mathcal{C}, \mathbf{Cat}]$. We now investigate how the corresponding lax codescent cocone of the strict coherence data given above looks like concretely when $T = E^* \text{Lan}_E$.

The lax codescent cocone $(B, G := b \cdot TF: TA \rightarrow B, \overline{G} := b \cdot T\overline{F}: G \cdot \mu_A \Rightarrow G \cdot Ta)$

$$\begin{array}{ccc} & TA & \\ \mu_A \nearrow & & \searrow G \\ T^2A & \Downarrow \overline{G} & B \\ Ta \nearrow & & \searrow G \\ & TA & \end{array}$$

has its component at $c \in \text{ob } \mathcal{C}$ given by

$$\begin{array}{ccc} & \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) & \\ \mu_{Ac} \nearrow & & \searrow G(c) \\ \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \mathcal{C}(d, d_1) \times A(d) & \Downarrow \overline{G}_c & B(c) \\ Ta_c \nearrow & & \searrow G(c) \\ & \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) & \end{array}$$

in our case $T = E^* \text{Lan}_E$. The component of \overline{G}_c at $(h: d_1 \rightarrow c, k: d \rightarrow d_1, x \in A(d))$ is given by

$$\overline{G}_{c, d_1, d(h, k, x)} = (B(h) \cdot F_k)_{c, d_1, d(h, k, x)}: B(hk)(F_d(x)) \rightarrow B(h)(F_{d_1}(A(k)(x))).$$

So the component of \overline{G}_c at (h, k, x) corresponds to $B(h) \cdot F_k$ in the following diagram of 2-cells:

$$(3.2) \quad \begin{array}{ccc} A(d) & \xrightarrow{F_d} & B(d) \\ A(k) \downarrow & \swarrow_{F_k} & \downarrow B(k) \\ A(d_1) & \xrightarrow{F_{d_1}} & B(d_1) \\ A(h) \downarrow & \swarrow_{F_h} & \downarrow B(h) \\ A(c) & \xrightarrow{F_c} & B(c) \end{array} \cdot$$

Now let $F: A \rightarrow B$ be a marked-lax natural transformation, then for any morphism k in $\Sigma \subseteq \mathcal{C}$, $F_k = 1$. So from Diagram 3.2, we then have $\overline{G}_{c(h, k, x)} = B(h) \cdot F_k = 1$. We may express this situation as follows:

$$(3.3) \quad \begin{array}{ccc} & \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) & \\ \mu_{Ac} \nearrow & & \searrow G(c) \\ \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \Sigma(d, d_1) \times A(d) \xrightarrow{\text{inc}} \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \mathcal{C}(d, d_1) \times A(d) & \Downarrow \overline{G}_c & B(c) \\ Ta_c \nearrow & & \searrow G(c) \\ & \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) & \end{array} = 1.$$

Since all identities are included in Σ , there is a factorisation

$$(3.4) \quad \begin{array}{ccc} \sum_{d \in \mathcal{C}} \mathcal{C}(d, c) \times A(d) & \xleftarrow{\quad} & \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \mathcal{C}(d, d_1) \times A(d) \\ & \searrow & \nearrow \\ & \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \Sigma(d, d_1) \times A(d) & \end{array}$$

Diagram 3.4 motivates an abstract diagram

$$\begin{array}{ccc} X_0 & \xleftarrow{\quad} & X_1 \\ & \searrow & \nearrow \\ & X_\sigma & \end{array}$$

in \mathbf{Cat} , whereas Equation (3.3) motivates a coherence condition

$$X_\sigma \xrightarrow{\exists} X_1 \begin{array}{ccc} \xrightarrow{s} & X_0 & \xrightarrow{y} \\ \Downarrow v & & \nearrow y \\ \xrightarrow{t} & X_0 & \xrightarrow{y} \end{array} = X_\sigma \xrightarrow{y_\sigma} Y$$

where $X_\sigma, \mathfrak{z}: X_\sigma \rightarrow X_1, y: X_0 \rightarrow Y$, and $y_\sigma: X_\sigma \rightarrow X$ are still left to be defined.

Definition 3.2.1. The category Δ_σ is generated by a set $\{[0], [1], [2], [\sigma]\}$ of objects and a set $\{p, m, q, s, t, i, j, k\}$ of morphisms, depicted as follows:

$$\begin{array}{ccccc} & \xleftarrow{p} & & \xleftarrow{s} & \\ [2] & \xleftarrow{m} & [1] & \xleftarrow{t} & [0] \\ & \xleftarrow{q} & & \xrightarrow{i} & \\ & & & \swarrow k & \searrow j \\ & & & & [\sigma] \end{array},$$

with relations given by

$$is = it = 1, ps = ms, qt = mt, pt = qs, \text{ and } i = jk,$$

which are exactly the simplicial relations with an extra equation $i = jk$. We call these the marked relations.

Definition 3.2.2. A collection of marked coherence data \mathfrak{X}_σ is a 2-functor $\mathfrak{X}_\sigma: \Delta_\sigma^{\text{op}} \rightarrow \mathcal{C}$. This means \mathfrak{X}_σ is a diagram

$$\begin{array}{ccccc} & \xrightarrow{p} & & \xrightarrow{s} & \\ X_2 & \xrightarrow{m} & X_1 & \xrightarrow{t} & X_0 \\ & \xrightarrow{q} & & \xleftarrow{i} & \\ & & & \swarrow \mathfrak{z} & \searrow i \\ & & & & X_\sigma \end{array}$$

in \mathcal{C} , that satisfies the marked identities

$$si = ti = 1, sp = sm, tq = tm, tp = sq, \text{ and } i = j\mathfrak{z}.$$

Definition 3.2.3. A marked codescent cocone of the marked coherence data \mathfrak{X}_σ defined in Definition 3.2.2 consists of a triple $(Y \in \text{ob } \mathcal{C}, y_\sigma: X_\sigma \rightarrow Y, v: ys \Rightarrow yt)$:

$$\begin{array}{ccc} & X_0 & \\ s \nearrow & & \searrow y := y_\sigma i \\ X_1 & \Downarrow v & Y \\ t \searrow & & \nearrow y := y_\sigma i \\ & X_0 & \end{array}$$

where $y := y_\sigma i: X_0 \rightarrow Y$, that satisfies the multiplicative equation

$$\begin{array}{ccccc} & X_1 & \xrightarrow{s} & X_0 & \xrightarrow{y} \\ p \nearrow & & \searrow s & & \searrow y \\ X_2 & \xrightarrow{m} & X_1 & \Downarrow v & Y \\ q \searrow & & \nearrow t & & \nearrow y \\ & X_1 & \xrightarrow{t} & X_0 & \end{array} = \begin{array}{ccccc} & X_1 & \xrightarrow{s} & X_0 & \xrightarrow{y} \\ p \nearrow & & \searrow t & & \searrow y \\ X_2 & \xrightarrow{m} & X_1 & \Downarrow v & Y \\ q \searrow & & \nearrow s & & \nearrow y \\ & X_1 & \xrightarrow{t} & X_0 & \end{array}$$

and the marked equation

$$X_\sigma \xrightarrow{\mathfrak{z}} X_1 \begin{array}{ccc} \nearrow s & X_0 & \searrow y \\ \Downarrow v & & \nearrow y \\ \searrow t & X_0 & \end{array} = X_\sigma \xrightarrow{y_\sigma} Y.$$

Definition 3.2.4. Let $(Y_1, y_{\sigma_1}: X_\sigma \rightarrow Y_1, v_1: y_{\sigma_1} s \Rightarrow y_{\sigma_1} t)$ and $(Y_2, y_{\sigma_2}: X_\sigma \rightarrow Y_2, v_2: y_{\sigma_2} s \Rightarrow y_{\sigma_2} t)$ be two marked codescent cocones. A morphism of marked codescent cocones from (Y_1, y_{σ_1}, v_1) to (Y_2, y_{σ_2}, v_2) consists of a morphism $f: Y_1 \rightarrow Y_2$ and a 2-cell $\theta: fy_{\sigma_1} \Rightarrow y_{\sigma_2}$ in \mathcal{C} , such that

$$\theta t \circ fv_1 = v_2 \circ \theta s.$$

Remark 3.2.5. In particular, if $Y_1 = Y_2$, then a morphism of lax codescent cocones reduces to a 2-cell $\theta: y_{\sigma_1} \Rightarrow y_{\sigma_2}$ satisfying

$$\theta t \circ v_1 = v_2 \circ \theta s.$$

Notation 3.2.6. We denote by $\mathbf{MarkedCocone}(\mathfrak{X}_\sigma, Y)$ the category of marked codescent cocones of \mathfrak{X}_σ with a fixed nadir $Y \in \text{ob } \mathcal{C}$ and the morphisms of marked codescent cocones as in Remark 3.2.5.

Definition 3.2.7. A marked codescent object Z of the marked coherence data in Definition 3.2.2 is the universal marked codescent cocone, i.e., it is characterised by

$$\mathbf{MarkedCocone}(\mathfrak{X}_\sigma, Y) \cong \mathcal{C}(Z, Y).$$

The following proposition is crucial in establishing our main result in this subsection. It tells us that a marked codescent cocone (Y, y_σ, ν) of a marked coherence data \mathfrak{X}_σ corresponds bijectively to a 2-natural transformation from the marked weight W to the 2-presheaf $\mathcal{C}(\mathfrak{X}_\sigma-, Y)$; in other words, a marked codescent object is described equivalently as a weighted colimit. This is the main reason that we can 'strictify' a marked-lax natural transformation into a 2-natural transformation.

Notation 3.2.8. We view a poset $\{0 < 1 < 2 < \dots < n\}$ as a category, which is denoted as $\mathbf{n} := \{0 \rightarrow 1 \rightarrow 2 \rightarrow \dots \rightarrow n\}$.

Proposition 3.2.9. *Marked codescent objects can be described by weighted colimits.*

More precisely, there exists a weight $W: \Delta_\sigma \rightarrow \mathbf{Cat}$, which we call the marked weight, such that

$$[\Delta_\sigma, \mathbf{Cat}](W, \mathcal{C}(\mathfrak{X}_\sigma-, Y)) \cong \mathbf{MarkedCocone}(\mathfrak{X}_\sigma, Y).$$

Proof. We define the marked weight W as follows. Let $W: \Delta_\sigma \rightarrow \mathbf{Cat}$ be a 2-functor, which acts as the usual embedding of the 2-truncated simplex category Δ_2 into \mathbf{Cat} on the objects $\{[0], [1], [2]\}$ and the morphisms $\{p, m, q, s, t, i\}$, and in addition, $W([\sigma]) = \mathbf{0}$, $W(k): \mathbf{1} \rightarrow \mathbf{0}$ is the constant map at 0, and $W(j): \mathbf{0} \rightarrow \mathbf{0}$ is the identity on $\mathbf{0}$. This means W is a diagram

$$\begin{array}{ccccc} & \xleftarrow{W_p} & & \xleftarrow{W_s} & \\ \mathbf{2} & \xleftarrow{W_m} & \mathbf{1} & \xleftarrow{W_t} & \mathbf{0} \\ & \xleftarrow{W_q} & & \xrightarrow{W_i} & \\ & & & \searrow^{W_k} & \nearrow_{W_j} \\ & & & \mathbf{0} & \end{array}$$

in \mathbf{Cat} , satisfying the marked relations.

A standard checking verifies that any 2-natural transformation $\gamma: W \rightarrow \mathcal{C}(\mathfrak{X}_\sigma-, Y)$ corresponds to a marked codescent cocone, and a modification $\gamma_1 \rightarrow \gamma_2$ is equivalently a morphism from (Y, y_1, ν_1) to (Y, y_2, ν_2) . \square

Now, let us consider the weighted colimit $W * \mathfrak{X}_\sigma =: X^\ddagger$.

Lemma 3.2.10. *There exists a morphism $x^\ddagger: X_0 \rightarrow X^\ddagger$ and a 2-cell $\chi^\ddagger: x^\ddagger \mathfrak{s} \Rightarrow x^\ddagger \mathfrak{t}$, such that $(X^\ddagger, x^\ddagger, \chi^\ddagger)$ is a marked codescent cocone of the marked coherence data \mathfrak{X}_σ .*

Proof. There is an isomorphism of categories

$$\mathcal{C}(X^\ddagger, Y) \cong [\Delta_\sigma, \mathbf{Cat}](W, \mathcal{C}(\mathfrak{X}_\sigma-, Y))$$

natural in $Y \in \text{ob } \mathcal{C}$. The result follows directly from Proposition 3.2.9. \square

Proposition 3.2.11. *The marked codescent cocone $(X^\ddagger, x^\ddagger, \chi^\ddagger)$ constructed in Lemma 3.2.10 is the marked codescent object of the marked coherence data \mathfrak{X}_σ , i.e., it is the universal marked codescent cocone.*

Proof. Suppose that $(B, G: X_0 \rightarrow B, \overline{G}: G\mathfrak{s} \rightarrow G\mathfrak{t})$ is a marked codescent cocone of \mathfrak{X}_σ .

In the proof of Lemma 3.2.10, we see that a marked codescent cocone of \mathfrak{X}_σ is equivalent to a 2-natural transformation $\beta: W \rightarrow \mathcal{C}(\mathfrak{X}_\sigma-, B)$ with $\beta_{[0]}(0) = G$, $\beta_{[1]}(\iota) = \overline{G}$, and the naturality amounts to the multiplicative and marked equations.

A standard argument shows that a morphism between marked codescent cocones (B, G_1, \overline{G}_1) and (B, G_2, \overline{G}_2) is equivalent to a modification $\beta_1 \rightarrow \beta_2$. And the universal property of the marked codescent cocone $(X^\ddagger, x^\ddagger, \chi^\ddagger)$ follows from that of the weighted colimit. \square

Recall that we have the 2-monad $T = E^* \text{Lan}_E: [\text{ob } \mathcal{C}, \mathbf{Cat}] \rightarrow [\text{ob } \mathcal{C}, \mathbf{Cat}]$, and we know that a strict T -algebra $(A, a: TA \rightarrow A)$ is equivalent to a 2-functor $A: \mathcal{C} \rightarrow \mathbf{Cat}$.

Let $\mathfrak{A}_\sigma: \Delta_\sigma^{\text{op}} \rightarrow [\mathcal{C}, \mathbf{Cat}]$ be a collection of marked coherence data in $[\mathcal{C}, \mathbf{Cat}]$ as follows:

$$T^3 A \begin{array}{c} \xrightarrow{\mu_{TA}} \\ \xrightarrow{T\mu_A} \\ \xrightarrow{T^2 a} \end{array} T^2 A \begin{array}{c} \xrightarrow{\mu_A} \\ \xrightarrow{Ta} \\ \xrightarrow{T\eta_A} \end{array} TA \begin{array}{c} \xrightarrow{\mu_A} \\ \xrightarrow{Ta} \\ \xrightarrow{T\eta_A} \end{array} A_\sigma \quad ,$$

where $A_\sigma := \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, -) \times \Sigma(d, d_1) \times A(d)$.

Then, a marked codescent cocone of \mathfrak{A}_σ consists of a morphism $G_\sigma: A_\sigma \rightarrow B$, where $B: \mathcal{C} \rightarrow \mathbf{Cat}$ is a 2-functor, and a 2-cell

$$\begin{array}{ccc} & TA & \\ \mu_A \nearrow & & \searrow G \\ T^2 A & \Downarrow \overline{G} & B \\ Ta \searrow & & \nearrow G \\ & TA & \end{array}$$

in $[\mathcal{C}, \mathbf{Cat}]$, where $G := G_\sigma T\eta_A$, satisfying the multiplicative equation

$$\begin{array}{ccc} \mu_{TA} \nearrow T^2 A \xrightarrow{\mu_A} TA \searrow G & & \mu_{TA} \nearrow T^2 A \xrightarrow{\mu_A} TA \searrow G \\ T^3 A \xrightarrow{T\mu_A} T^2 A \Downarrow \overline{G} B & = & T^3 A \xrightarrow{T\mu_A} T^2 A \xrightarrow{Ta} T^2 A \xrightarrow{\mu_A} TA \searrow G \\ T^2 a \searrow T^2 A \xrightarrow{Ta} TA \nearrow G & & T^2 a \searrow T^2 A \xrightarrow{Ta} TA \nearrow G \end{array}$$

and the marked equation

$$A_\sigma \xrightarrow{\iota} \begin{array}{ccc} & TA & \\ \mu_A \nearrow & & \searrow G \\ T^2 A & \Downarrow \overline{G} & B \\ Ta \searrow & & \nearrow G \\ & TA & \end{array} = A_\sigma \xrightarrow{G_\sigma} B .$$

The following proposition justifies our notion of marked codescent cocones and shows that it is really the bridge to marked-lax natural transformations.

Proposition 3.2.12. *There is an isomorphism of categories*

$$\mathbf{MarkedCocone}(\mathfrak{A}_\sigma, B) \cong [\mathcal{C}, \mathbf{Cat}]_{\iota, \Sigma}(A, B),$$

natural in B .

Proof. In [Lac02], it is shown that a lax codescent cocone

$$\begin{array}{ccc} & TA & \\ \mu_A \nearrow & & \searrow G \\ T^2 A & \Downarrow \overline{G} & B \\ Ta \searrow & & \nearrow G \\ & TA & \end{array}$$

is equivalent to a lax T -morphism $(F: A \rightarrow B, \overline{F}: b \cdot Tf \Rightarrow F \cdot a)$, in which we have $G = b \cdot TF$ and $\overline{G} = b \cdot T\overline{F}$.

Similarly, we can express a marked codescent cocone $(B, G_\sigma: A_\sigma \rightarrow B, \overline{G})$ in terms of (F, \overline{F}) , except that our marked equation replaces the original unital equation. It remains to transform the marked equation into an equation in terms of (F, \overline{F}) .

To ask

$$A_\sigma \xrightarrow{\iota} \begin{array}{ccc} & TA & \\ \mu_A \nearrow & & \searrow G \\ T^2 A & \Downarrow \overline{G} & B \\ Ta \searrow & & \nearrow G \\ & TA & \end{array} = A_\sigma \xrightarrow{G_\sigma} B .$$

is equivalently to ask

$$\begin{array}{c}
 \begin{array}{ccccc}
 & & T^2 B & & \\
 & T^2 F \nearrow & & \searrow T b & \\
 A_\sigma & \xrightarrow{\iota} T^2 A & \Downarrow T \overline{F} & \rightarrow T B & \xrightarrow{b} B \\
 & T a \searrow & & \nearrow T F & \\
 & & T A & &
 \end{array} \\
 = A_\sigma \xrightarrow{G_\sigma} B .
 \end{array}$$

The component of the left hand side at $c \in \mathcal{C}$ is given by

$$\begin{array}{c}
 \mathcal{C}(d_1, c) \times \Sigma(d, d_1) \times A(d) \xrightarrow{\text{in}_c} \mathcal{C}(d_1, c) \times \mathcal{C}(d, d_1) \times A(d) \\
 \begin{array}{ccc}
 \xrightarrow{T^2 F_c} & \mathcal{C}(d_1, c) \times \mathcal{C}(d, d_1) \times B(d) & \xrightarrow{T b_c} \\
 \Downarrow T \overline{F}_c & & \Downarrow T \overline{F}_c \\
 \xrightarrow{T a_c} & \mathcal{C}(d_1, c) \times A(d_1) & \xrightarrow{T F_c}
 \end{array} \\
 \xrightarrow{b_c} \mathcal{C}(d_1, c) \times B(d_1) \xrightarrow{b_c} B(c) ,
 \end{array}$$

hence the equation is precisely to ask for any $(f, e, x) \in \sum_{d_1 \in \mathcal{C}} \sum_{d \in \mathcal{C}} \mathcal{C}(d_1, c) \times \Sigma(d, d_1) \times A(d)$, the 2-cell $b_c \cdot T \overline{F}_{c(f, e, x)} = B(f) \cdot \overline{F}_c$ is the identity, depicted below:

$$\begin{array}{ccc}
 \begin{array}{ccc}
 A(d) & \xrightarrow{F_d} & B(d) \\
 A(e) \downarrow & \swarrow \overline{F}_e & \downarrow B(e) \\
 A(d_1) & \xrightarrow{F_{d_1}} & B(d_1) \xrightarrow{B(f)} B(c)
 \end{array} \\
 = A(d) \xrightarrow{F_d} B(d) \xrightarrow{B(f)} B(c) ,
 \end{array}$$

which is equivalent to ask that $\overline{F}_e = 1$. All in all, this asks for any $e \in \Sigma$, $\overline{F}_e = 1$, which is exactly the condition for a marked-lax natural transformation.

It is left to show that a morphism of marked codescent cocones corresponds to a modification between two marked-lax natural transformations.

Following [Lac02], it is clear that a morphism $(B, G_1: TA \rightarrow B, \overline{G}_1: G_1 \cdot \mu_A \Rightarrow G_1 \cdot Ta) \rightarrow (B, G_2: TA \rightarrow B, \overline{G}_2: G_2 \cdot \mu_A \Rightarrow G_2 \cdot Ta)$ of marked codescent cocones given by a 2-cell $\theta: G_1 \Rightarrow G_2$ satisfying $\theta \cdot Ta \cdot \overline{G}_1 = \overline{G}_2 \cdot \theta \cdot \mu_A$, is equivalent to a T -transformation $(F_1: A \rightarrow B, \overline{F}_1: b \cdot T F_1 \Rightarrow F_1 \cdot a) \rightarrow (F_2: A \rightarrow B, \overline{F}_2: b \cdot T F_2 \Rightarrow F_2 \cdot a)$ given by a 2-cell $\rho: F_1 \Rightarrow F_2$ satisfying

$$\begin{array}{ccc}
 \begin{array}{ccc}
 TA & \xrightarrow{TF_1} & TB \\
 \Downarrow T \rho & & \Downarrow T \rho \\
 a \downarrow & TF_2 & \downarrow b \\
 A & \xrightarrow{F_2} & B
 \end{array} \\
 = \begin{array}{ccc}
 TA & \xrightarrow{TF_1} & TB \\
 \overline{F}_1 \swarrow & & \swarrow \overline{F}_1 \\
 a \downarrow & F_1 & \downarrow b \\
 A & \xrightarrow{F_1} & B \\
 \Downarrow \rho & & \Downarrow \rho \\
 A & \xrightarrow{F_2} & B
 \end{array} ,
 \end{array}$$

where $G = b \cdot T F$, $\overline{G} = b \cdot T \overline{F}$ and $\theta = b \cdot T \rho$. Consider the components at $c \in \mathcal{C}$, the above equation then says

$$\begin{array}{ccc}
 \begin{array}{ccc}
 \mathcal{C}(d, c) \times A(d) & \xrightarrow{TF_{1c}} & \mathcal{C}(d, c) \times B(d) \\
 \Downarrow T \rho_c & & \Downarrow T \rho_c \\
 a_c \downarrow & TF_{2c} & \downarrow b_c \\
 A(c) & \xrightarrow{F_{2c}} & B(c)
 \end{array} \\
 = \begin{array}{ccc}
 \mathcal{C}(d, c) \times A(d) & \xrightarrow{TF_{1c}} & \mathcal{C}(d, c) \times B(d) \\
 \overline{F}_{1c} \swarrow & & \swarrow \overline{F}_{1c} \\
 a_c \downarrow & F_{1c} & \downarrow b_c \\
 A(c) & \xrightarrow{F_{1c}} & B(c) \\
 \Downarrow \rho_c & & \Downarrow \rho_c \\
 A(c) & \xrightarrow{F_{2c}} & B(c)
 \end{array} .
 \end{array}$$

This means for $(f: d \rightarrow c, x \in A(d))$,

$$\rho_c(A(f)(x)) \circ \overline{F}_{1f} = \overline{F}_{2f} \circ B(f)(\rho_d(x)),$$

which is exactly the modification axiom. \square

We achieve the following desired theorem:

Proposition 3.2.13. *There is an isomorphism of categories*

$$[\mathcal{C}, \mathbf{Cat}](A^\dagger, B) \cong [\mathcal{C}, \mathbf{Cat}]_{l, \Sigma}(A, B),$$

natural in B .

In other words, there is a left adjoint $(\)^\ddagger: [\mathcal{A}, \mathbf{Cat}]_{l,\Sigma} \rightarrow [\mathcal{A}, \mathbf{Cat}]$ to the inclusion.

Proof. By Proposition 3.2.9, Proposition 3.2.12, and the universal property of the weighted colimit $A^\ddagger = W * \mathfrak{A}_\sigma$, we have the following chain of isomorphisms:

$$\begin{aligned} [\mathcal{C}, \mathbf{Cat}](W * \mathfrak{A}_\sigma, B) &\cong [\Delta_\sigma, \mathbf{Cat}](W, [\mathcal{C}, \mathbf{Cat}](\mathfrak{A}_\sigma -, B)) \\ &\cong \mathbf{MarkedCocone}(\mathfrak{A}_\sigma, B) \\ &\cong [\mathcal{C}, \mathbf{Cat}]_{l,\Sigma}(A, B). \end{aligned} \quad \square$$

Theorem 3.2.14. *Let (\mathcal{A}, Σ) be a marked 2-category. The marked-lax limit of a 2-functor $F: \mathcal{A} \rightarrow \mathcal{B}$ has the same universal property as the \mathbf{Cat} -weighted limit $\{\Delta(1)^\ddagger, F\}$.*

Proof. We have

$$\begin{aligned} \mathcal{B}(B, \mathfrak{m}_l \lim F) &\cong [\mathcal{A}, \mathcal{B}]_{l,\Sigma}(\Delta(B), F) \\ &\cong [\mathcal{A}, \mathbf{Cat}]_{l,\Sigma}(\Delta(1), \mathcal{B}(B, F-)) \\ &\cong [\mathcal{A}, \mathbf{Cat}](\Delta(1)^\ddagger, \mathcal{B}(B, F-)) \\ &\cong \mathcal{B}(B, \{\Delta(1)^\ddagger, F\}). \end{aligned} \quad \square$$

3.3. \mathbf{Cat} -weighted limits as marked limits. The other direction that any \mathbf{Cat} -weighted limit can be equivalently expressed as a marked-lax limit with the same universal property is well-known. This has first been discussed in Street's paper [Str76], and also in the recent work [Mes23]. Nonetheless, we will have to recall the construction for this result in later sections, hence we decide to present the proof.

Let \mathcal{A}, \mathcal{D} be 2-categories, let $W: \mathcal{D} \rightarrow \mathbf{Cat}$ be a weight and $R: \mathcal{D} \rightarrow \mathcal{A}$ be a 2-functor.

The 2-category of elements of W consists of

- objects those pairs (D, δ) ;
- morphisms $(d, \omega): (D_1, \delta_1) \rightarrow (D_2, \delta_2)$ given by pairs $(d: D_1 \rightarrow D_2, \omega: Wd\delta_1 \rightarrow \delta_2)$;
- 2-cells $\alpha: (d, \omega) \Rightarrow (d', \omega')$ given by the 2-cells $\alpha: d \Rightarrow d'$ in \mathcal{D} such that $\omega'W\alpha\delta_1 = \omega$,

and the composition of two morphisms $(h: D_1 \rightarrow D_2, \eta: Wh\delta_1 \rightarrow \delta_2)$ and $(k: D_2 \rightarrow D_3, \kappa: Wk\delta_2 \rightarrow \delta_3)$ is given by

$$(kh: D_1 \rightarrow D_3, \kappa \circ (Wk \cdot \eta): WkWh\delta_1 \rightarrow \delta_3).$$

We also have the projection of $\mathcal{E}lW$ onto \mathcal{D} :

$$\begin{aligned} P: \mathcal{E}lW &\rightarrow \mathcal{D}, \\ (D, \delta) &\mapsto D. \end{aligned}$$

Remark 3.3.1. Following [Mes23] $\mathcal{E}lW$ is the lax comma object of $\Delta(1): 1 \rightarrow \mathbf{Cat}$ and $W: \mathcal{D} \rightarrow \mathbf{Cat}$

$$\begin{array}{ccc} \mathcal{E}lW & \xrightarrow{P} & \mathcal{D} \\ \downarrow \text{!} & \nearrow \mu & \downarrow W \\ 1 & \xrightarrow{\Delta(1)} & \mathbf{Cat} \end{array}$$

We are going to show that there is a certain marked-lax limit possessing the same universal property of the weighted limit $\{W, R\}$.

Proposition 3.3.2. *Let $\Sigma := \{(d \in \text{mor } \mathcal{D}, 1)\}$ be the class of morphisms in $\mathcal{E}lW$. Then*

$$[\mathcal{D}, \mathbf{Cat}](W, \mathcal{A}(A, R-)) \cong [\mathcal{E}lW, \mathcal{A}]_{l,\Sigma}(\Delta(A), RP).$$

Proof. It suffices to show

$$[\mathcal{D}, \mathbf{Cat}](W, \mathcal{A}(A, R-)) \cong [\mathcal{E}lW, \mathbf{Cat}]_{l,\Sigma}(\Delta(1), \mathcal{A}(A, RP-)).$$

Let $\beta: W \rightarrow \mathcal{A}(A, R-)$ be a 2-natural transformation, we can then form the composition of μ and βP

$$\left(\begin{array}{ccc} \mathcal{E}lW & \xrightarrow{P} & \mathcal{D} \\ \downarrow ! & \nearrow \mu & W \downarrow \xrightarrow{\beta} \\ 1 & \xrightarrow{\Delta(1)} & \mathbf{Cat}_{\leftarrow} \end{array} \right) \mathcal{A}(A, R-).$$

Since β is 2-natural, the 2-component of the composition $\beta P \circ \mu$ with respect to a morphism in Σ is the identity, hence $\beta P \circ \mu$ is indeed a marked-lax natural transformation. Thus, whiskering with P and pre-composing with μ defines a functor

$$(3.5) \quad \begin{aligned} G: [\mathcal{D}, \mathbf{Cat}](W, \mathcal{A}(A, R-)) &\rightarrow [\mathcal{E}lW, \mathbf{Cat}]_{l, \Sigma}(\Delta(1), \mathcal{A}(A, RP-)), \\ \beta &\mapsto \beta P \circ \mu, \\ \Gamma: \beta_1 \rightarrow \beta_2 &\mapsto \Gamma P \circ \mu. \end{aligned}$$

Next, we would like to construct the inverse of G .

Let $\alpha: \Delta(1)! \rightarrow \mathcal{A}(A, RP-)$ be a marked-lax natural transformation. For any object D of \mathcal{D} , define a functor

$$\begin{aligned} \alpha_{(\Delta(D), -)}: WD &\rightarrow \mathcal{A}(A, RD), \\ \delta &\mapsto \alpha_{(D, \delta)}, \\ \omega: \delta_1 \rightarrow \delta_2 &\mapsto \alpha_{(1_D, \omega)}. \end{aligned}$$

Since α is marked natural, for a morphism $(d: D_1 \rightarrow D_2, 1: Wd\delta_1 \rightarrow Wd\delta_1) \in \Sigma$, $\alpha_{(d,1)} = 1$. This implies that for any $\delta_1 \in WD_1$, $Rd \cdot \alpha_{(D_1, \delta_1)} = \alpha_{(D_2, Wd\delta_1)}$. So we have a commutative diagram

$$\begin{array}{ccc} WD_1 & \xrightarrow{Wd} & WD_2 \\ \alpha_{(\Delta(D_1), -)} \downarrow & & \downarrow \alpha_{(\Delta(D_2), -)} \\ \mathcal{A}(A, RD_1) & \xrightarrow{(Rd)_*} & \mathcal{A}(A, RD_2) \end{array}$$

Now, let $\theta: d \Rightarrow d'$ be a 2-cell in \mathcal{D} , and let $(d: D_1 \rightarrow D_2, W\theta\delta_1: Wd\delta_1 \rightarrow Wd'\delta_1)$ and $(d': D_1 \rightarrow D_2, 1: Wd'\delta_1 \rightarrow Wd'\delta_1)$ be morphisms from (D_1, δ_1) to $(D_2, Wd'\delta_1)$ in $\mathcal{E}lW$. Then it is clear that θ is also a 2-cell $(d, W\theta\delta_1) \Rightarrow (d', 1)$ in $\mathcal{E}lW$. Since α is natural, we have

$$\alpha_{(d, W\theta\delta_1)} = (R\theta)_* \cdot \alpha_{(D_1, \delta_1)},$$

which gives

$$\alpha_{(\Delta(D_2), -)} \cdot W\theta = (R\theta)_* \cdot \alpha_{(\Delta(D_1), -)}.$$

Altogether, we are able to define a 2-natural transformation $\beta: W \rightarrow \mathcal{A}(A, R-)$, whose 1-component is given by $\beta_D := \alpha_{(\Delta(D), -)}$, and 2-component is given by a natural transformation β_d , where its component at $\delta \in WD$ is given by $\beta_{d\delta} = \alpha_{(d,1)}$, which is the identity because α is marked-lax natural.

Next, let $\Lambda: \alpha_1 \rightarrow \alpha_2$ be a modification between the marked-lax natural transformations α_1 and α_2 . Consider the 2-components of the transformations with respect to the morphism $(d: D_1 \rightarrow D_2, 1: Wd\delta_1 \rightarrow Wd\delta_1)$, we get $\Lambda_{(D_2, Wd\delta_1)} = Rd \cdot \Lambda_{(D_1, \delta_1)}$ from the modification axiom. Therefore, if we construct a map $\Theta: \beta_1 \rightarrow \beta_2$, where β_1 and β_2 are the 2-natural transformations constructed from α_1 and α_2 , by setting its component at D as a natural transformation $\Theta_D := \xi^D: \alpha_{1(\Delta(D), -)} \rightarrow \alpha_{2(\Delta(D), -)}$ with component at $\delta \in WD$ given by $\xi_\delta^D := \Lambda_{(D, \delta)}: \alpha_{1_{D, \delta}} \rightarrow \alpha_{2_{D, \delta}}$, we then obtain a modification Θ which satisfies the modification axiom

$$\Theta_{D_2} \cdot Wd = (Rd)_* \cdot \Theta_{D_1}.$$

The above constructions of β and Θ define a functor

$$\begin{aligned} G': [\mathcal{E}lW, \mathbf{Cat}]_{l, \Sigma}(\Delta(1), \mathcal{A}(A, RP-)) &\rightarrow [\mathcal{D}, \mathbf{Cat}](W, \mathcal{A}(A, R-)), \\ \alpha &\mapsto \beta = \{\alpha_{(\Delta(D), -)}\}_{D \in \mathcal{D}}, \\ \Lambda: \alpha_1 \rightarrow \alpha_2 &\mapsto \Theta = \{\{\Lambda_{(D, \delta)}\}_{\delta \in WD}\}_{D \in \mathcal{D}}. \end{aligned}$$

A straightforward checking shows that G and G' are inverses of each other. \square

Theorem 3.3.3. *Let \mathcal{A}, \mathcal{D} be 2-categories. Let $W: \mathcal{D} \rightarrow \mathbf{Cat}$ be a weight and $R: \mathcal{D} \rightarrow \mathcal{A}$ be a 2-functor. Let $\Sigma := \{(d \in \text{mor } \mathcal{D}, 1)\}$ be the class of morphisms in the 2-category of elements $\mathcal{E}IW$ of W , so that $(\mathcal{E}IW, \Sigma)$ is a marked 2-category. The \mathbf{Cat} -weighted limit $\{W, R\}$ has the same universal property as the marked-lax limit of the 2-functor $RP: \mathcal{E}IW \rightarrow \mathcal{A}$.*

Proof. By Proposition 3.3.2, we have a chain of isomorphisms

$$\begin{aligned} \mathcal{A}(A, \{W, R\}) &\cong [\mathcal{D}, \mathbf{Cat}](W, \mathcal{D}(A, R-)) \\ &\cong [\mathcal{E}IW, \mathcal{A}]_{l, \Sigma}(\Delta(A), RP) \\ &\cong \mathcal{A}(A, \text{mlim} RP). \end{aligned} \quad \square$$

4. PRELIMINARIES ON ENHANCED 2-CATEGORY THEORY

4.1. \mathcal{F} -categories. We recall the basics of enhanced 2-category theory, which was first proposed by Lack and Shulman in [LS12]. A gentle introduction can also be found in [Bou14] by Bourke.

Definition 4.1.1. *Let \mathcal{F} be the full subcategory of the arrow category \mathbf{Cat}^2 of the category \mathbf{Cat} , determined by the fully faithful and injective-on-object functors, i.e., the full embeddings.*

In other words, an object of \mathcal{F} is a full embedding

$$A_\tau \xrightarrow{j_A} A_\lambda,$$

a morphism f from j_A to j_B in \mathcal{F} is given by two functors $f_\tau: A_\tau \rightarrow B_\tau$ and $f_\lambda: A_\lambda \rightarrow B_\lambda$ making the following square commutes

$$\begin{array}{ccc} A_\tau & \xrightarrow{j_A} & A_\lambda \\ f_\tau \downarrow & & \downarrow f_\lambda \\ B_\tau & \xrightarrow{j_B} & B_\lambda \end{array}.$$

We call A_τ the *tight* part of A , and A_λ the *loose* part of A ; similarly, we apply this terminology to f and α .

Remark 4.1.2. \mathcal{F} is (co)complete and Cartesian closed.

An \mathcal{F} -category \mathbb{A} is a category \mathbb{A} enriched in \mathcal{F} . This means \mathbb{A} has

- objects x, y, \dots ;
- hom-objects $\mathbb{A}(x, y)$ in \mathcal{F} , each consists of a full embedding $\mathbb{A}(x, y)_\tau \hookrightarrow \mathbb{A}(x, y)_\lambda$ of the tight part into the loose part.

We can form a 2-category \mathcal{A}_τ as follows:

- \mathcal{A}_τ has all the objects of \mathbb{A} ;
- the hom-categories $\mathcal{A}_\tau(x, y)$ for any objects x, y are $\mathbb{A}(x, y)_\tau$.

Similarly, we can form a 2-category \mathcal{A}_λ by setting the hom-categories $\mathcal{A}_\lambda(x, y)$ as $\mathbb{A}(x, y)_\lambda$ for any objects x, y .

Since for each pair of objects x, y , $\mathcal{A}_\tau(x, y) \hookrightarrow \mathcal{A}_\lambda(x, y)$ is a full-embedding, we obtain a 2-functor

$$J_{\mathbb{A}}: \mathcal{A}_\tau \rightarrow \mathcal{A}_\lambda.$$

By construction, $J_{\mathbb{A}}$ is identity-on-object, faithful, and locally fully faithful.

Remark 4.1.3. We may identify an \mathcal{F} -category \mathbb{A} with $J_{\mathbb{A}}$. Indeed, any 2-functor which is identity-on-object, faithful, and locally fully faithful uniquely determined an \mathcal{F} -category.

The morphisms in \mathcal{A}_τ are called the *tight* morphisms, whereas those in \mathcal{A}_λ are called *loose*.

Notation 4.1.4. We write $A \rightarrow B$ for a tight morphism from A to B , and $A \rightsquigarrow B$ for a loose morphism from A to B .

Example 4.1.5 (2-categories). A 2-category can be viewed as a *chordate* \mathcal{F} -category, namely, all the morphisms are assumed to be tight. On the other hand, a 2-category can also be viewed as an *inchordate* \mathcal{F} -category, namely, only the identities are tight.

Example 4.1.6 (\mathbb{F}). Since any monoidal closed category is self-enriched, \mathcal{F} is also self-enriched, and we denote this \mathcal{F} -category by \mathbb{F} .

More precisely, \mathbb{F} consists of

- objects the full embeddings $A_\tau \hookrightarrow A_\lambda$;
- tight morphisms the tightness-preserving functors, depicted as

$$\begin{array}{ccc} A_\tau & \xrightarrow{j_A} & A_\lambda \\ f_\tau \downarrow & & \downarrow f_\lambda \\ B_\tau & \xrightarrow{j_B} & B_\lambda \end{array};$$

- loose morphisms the functors $A_\lambda \rightarrow B_\lambda$;
- 2-cells the natural transformations between the loose morphisms with tight components, or equivalently, the pair of two natural transformations α_τ and α_λ making the following diagram commute

$$\begin{array}{ccc} A_\tau & \xrightarrow{j_A} & A_\lambda \\ g_\tau \left(\begin{array}{c} \Leftarrow \\ \alpha_\tau \end{array} \right) f_\tau & & g_\lambda \left(\begin{array}{c} \Leftarrow \\ \alpha_\lambda \end{array} \right) f_\lambda \\ B_\tau & \xrightarrow{j_B} & B_\lambda \end{array},$$

as j_B is fully faithful.

Example 4.1.7 (T -algebras). We can combine $T\text{-Alg}_s$ and $T\text{-Alg}_l$ into an \mathcal{F} -category $T\text{-Alg}_{s,l}$. The colax and the pseudo cases are also similar.

4.2. \mathcal{F} -functors and \mathcal{F} -natural transformations. Let \mathbb{A} and \mathbb{B} be two \mathcal{F} -categories. An \mathcal{F} -functor $F: \mathbb{A} \rightarrow \mathbb{B}$ is a functor enriched in \mathcal{F} , which precisely means that F consists of 2-functors $F_\tau: \mathcal{A}_\tau \rightarrow \mathcal{B}_\tau$ and $F_\lambda: \mathcal{A}_\lambda \rightarrow \mathcal{B}_\lambda$ making the following diagram commute

$$\begin{array}{ccc} \mathcal{A}_\tau & \xrightarrow{J_A} & \mathcal{A}_\lambda \\ F_\tau \downarrow & & \downarrow F_\lambda \\ \mathcal{B}_\tau & \xrightarrow{J_B} & \mathcal{B}_\lambda \end{array}.$$

Remark 4.2.1. F_τ is uniquely determined by F_λ : an \mathcal{F} -functor $F: \mathbb{A} \rightarrow \mathbb{B}$ is a 2-functor $F_\lambda: \mathcal{A}_\lambda \rightarrow \mathcal{B}_\lambda$ which preserves tightness, i.e., F_λ sends a tight morphism in \mathbb{A} to a tight morphism in \mathbb{B} .

When $\mathbb{B} = \mathbb{F}$, an \mathcal{F} -functor $F: \mathbb{D} \rightarrow \mathbb{F}$ is called an \mathcal{F} -weight. Such an \mathcal{F} -weight $\Phi: \mathbb{D} \rightarrow \mathbb{F}$ amounts to a triple $(\Phi_\tau, \Phi_\lambda, \varphi)$, where $\Phi_\tau: \mathcal{D}_\tau \rightarrow \mathbf{Cat}$ and $\Phi_\lambda: \mathcal{D}_\lambda \rightarrow \mathbf{Cat}$ are 2-functors, and $\varphi: \Phi_\tau \rightarrow \Phi_\lambda J_{\mathbb{D}}$ is a 2-natural transformation

$$\begin{array}{ccc} \mathcal{D}_\tau & \xrightarrow{\Phi_\tau} & \mathbf{Cat} \\ & \searrow J_{\mathbb{D}} & \downarrow \varphi \\ & & \mathcal{D}_\lambda \\ & & \nearrow \Phi_\lambda \end{array}$$

such that all the components are full-embeddings.

Let $F, G: \mathbb{A} \rightrightarrows \mathbb{B}$ be two \mathcal{F} -functors. An \mathcal{F} -natural transformation $\alpha: F \rightarrow G$ consists of 2-natural transformations $\alpha_\tau: F_\tau \rightarrow G_\tau$ and $\alpha_\lambda: F_\lambda \rightarrow G_\lambda$ making the following diagram of 2-cells commutes

Notation 5.1.7. We denote the above \mathcal{F} -category of \mathcal{F} -functors $\mathbb{D} \rightarrow \mathbb{A}$, lax natural transformations between loose parts and dotted-lax natural transformations, and modifications by $[\mathbb{D}, \mathbb{A}]_{l, \Sigma, T}$.

Notation 5.1.8. We denote by $\Sigma_\lambda(a, b)$ the full subcategory of $\mathcal{D}_\lambda(a, b)$ consisting of loose morphisms in Σ that have fixed source a and fixed target b , and by $\Sigma_\tau(a, b)$ the full subcategory of $\mathcal{D}_\tau(a, b)$ consisting of tight morphisms in Σ that have fixed source a and fixed target b .

It is clear that we have 2-categories Σ_λ and Σ_τ , which are wide sub-2-categories of \mathcal{D}_λ and \mathcal{D}_τ , respectively, and hence there are inclusions $J_{\Sigma_\lambda} : \Sigma_\lambda \hookrightarrow \mathcal{D}_\lambda$ and $J_{\Sigma_\tau} : \Sigma_\tau \hookrightarrow \mathcal{D}_\tau$.

Definition 5.1.9. *Let (\mathbb{D}, Σ, T) be a small dotted \mathcal{F} -category, and \mathbb{A} be an \mathcal{F} -category. The dotted(-marked)-lax limit $d_l \lim S$ of an \mathcal{F} -functor $S : \mathbb{D} \rightarrow \mathbb{A}$ is characterised by an isomorphism*

$$(5.1) \quad \mathbb{A}(A, d_l \lim S) \cong [\mathbb{D}, \mathbb{A}]_{l, \Sigma, T}(\Delta(A), S)$$

in \mathbb{F} , which is natural in $A \in \text{ob } \mathbb{A}$.

Remark 5.1.10. By replacing dotted-lax natural transformations with dotted-colax or dotted-pseudo natural transformations, we obtain the notions of dotted-colax or dotted-pseudo limits, respectively.

5.2. Examples of dotted limits. To illustrate the convenience and practicality of dotted limits, we describe several examples of important \mathcal{F} -weighted limits in the form of dotted limits. Many of these \mathcal{F} -weighted limits have several variations: the l -rigged, c -rigged, and p -rigged versions, which were first discussed in [LS12].

In the forthcoming examples, we see that the presentation of the \mathcal{F} -categorical limits in dotted limits are simpler than in \mathcal{F} -weighted limits. In fact, the indexing dotted \mathcal{F} -categories that we give are almost the same as the indexing \mathcal{F} -categories given in [LS12], except that we are marking some morphisms and dotting some objects. In other words, the \mathcal{F} -weights are eliminated completely. The data of the dotted \mathcal{F} -categories alone already suffice to capture the \mathcal{F} -categorical limits of interests.

In each of the following examples, we denote by L the corresponding dotted limit of an \mathcal{F} -functor $S : \mathbb{D} \rightarrow \mathbb{A}$. Besides, we provide the description of p -rigged and c -rigged limits in terms of dotted-lax limits, whereas that of l -rigged limits in terms of dotted-colax limits, as this formulation facilitates our later discussion on the lifting theorem.

We first investigate the examples such that in their indexing dotted \mathcal{F} -categories, Σ coincides with \mathcal{D}_τ .

Example 5.2.1 (w -rigged inserters). Recall in [LS12] that an l -rigged inserter can be formed with the indexing \mathcal{F} -category

$$\mathbb{B} = \left\{ x \begin{array}{c} \xrightarrow{g} \\ \rightsquigarrow_f \end{array} y \right\},$$

where the image under the loose part Φ_λ of the weight Φ is

$$\mathbf{1} \begin{array}{c} \xrightarrow{0} \\ \xrightarrow{1} \end{array} \mathbf{2},$$

and that under the tight part Φ_τ is the identity

$$\mathbf{1} \longrightarrow \mathbf{1},$$

and where φ has components at x and y being the identity and 0, respectively. Therefore, the object 0 in $\mathbf{2}$ is tight.

Let

$$\mathbb{D} = \left\{ \dot{x} \begin{array}{c} \xrightarrow{\dot{g}} \\ \rightsquigarrow_{\dot{f}} \end{array} \dot{y} \right\},$$

which is indeed the same as \mathbb{B} , by noticing that Σ coincides with \mathcal{D}_τ , except that we now set the objects x and y to be dotted.

A dotted-colax natural transformation $\alpha : \Delta(L) \rightarrow S_\lambda$ clearly corresponds to

$$\begin{array}{c}
\alpha_x \nearrow Sx \xrightarrow{Sg} \\
L \searrow \Downarrow \alpha_f \nearrow Sy \\
\alpha_x \searrow Sx \xrightarrow{Sf}
\end{array}$$

an l -rigged inserter, as desired.

Similarly, a c -rigged inserter can be described using dotted limits with the same \mathbb{D} for l -rigged inserters as above, except that we replace dotted-colax natural transformations with dotted-lax natural transformations.

Recall in [LS12] that a p -rigged inserter can be formed with the indexing \mathcal{F} -category

$$\mathbb{B} = \{ x \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{f} \end{array} y \},$$

where the image under the loose part Φ_λ of the weight Φ is

$$\mathbf{1} \begin{array}{c} \xrightarrow{0} \\ \xrightarrow{1} \end{array} \mathbf{2},$$

and that $\Phi_\tau(x) = \mathbf{1}$, and $\Phi_\tau(y) = \emptyset$, and where φ has components at x and y being the identity and the unique morphism $\emptyset \rightarrow \mathbf{2}$, respectively. This implies that no objects in $\mathbf{2}$ are tight.

Now, consider the dotted \mathcal{F} -category

$$\mathbb{D} = \{ \dot{x} \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{f} \end{array} \dot{y} \},$$

which is very much the same as \mathbb{B} , except that $g \in \Sigma$ and $x \in T$.

A dotted-lax natural transformation $\alpha: \Delta(L) \rightarrow S_\lambda$ clearly corresponds to

$$\begin{array}{c}
\alpha_x \nearrow Sx \xrightarrow{Sf} \\
L \searrow \Downarrow \alpha_f \nearrow Sy \\
\alpha_x \searrow Sx \xrightarrow{Sg}
\end{array}$$

giving a p -rigged inserter.

Example 5.2.2 (l -rigged l -descent and c -rigged c -descent objects). We start with the l -rigged l -descent objects.

Let \mathbb{D} be the locally discrete sub- \mathcal{F} -category such that \mathcal{D}_λ is generated by

$$\mathbf{i} \begin{array}{c} \xrightarrow{\delta_\rho} \\ \xleftarrow{\sigma} \end{array} \mathbf{2} \begin{array}{c} \xrightarrow{\delta_\rho} \\ \xrightarrow{\delta_1} \end{array} \mathbf{3},$$

whereas \mathcal{D}_τ is generated by

$$\mathbf{i} \begin{array}{c} \xrightarrow{\delta_\rho} \\ \xleftarrow{\sigma} \end{array} \mathbf{2} \begin{array}{c} \xrightarrow{\delta_\rho} \\ \xrightarrow{\delta_1} \end{array} \mathbf{3}.$$

This dotted \mathcal{F} -category is indeed the same as the indexing \mathcal{F} -category provided in [LS12], except that we have dotted objects. Note that the corresponding \mathcal{F} -weight given there is non-trivial.

Let the image of an \mathcal{F} -functor $S: \mathbb{D} \rightarrow \mathbb{A}$ takes the form

$$A_0 \begin{array}{c} \xrightarrow{\delta_0^A} \\ \xleftarrow{\sigma^A} \end{array} A_1 \begin{array}{c} \xrightarrow{\delta_1^A} \\ \xrightarrow{\delta_2^A} \end{array} A_2,$$

where $A_i := \mathbf{S}i + \mathbf{1}$ and $\delta_i^A := S\delta_i$ for $i = 0, 1, 2$, and $\sigma^A := S\sigma$. Then a dotted-colax natural transformation $\alpha: \Delta(L) \rightarrow S_\lambda$ actually gives

$$\begin{array}{c}
a_0 \nearrow A_0 \xrightarrow{\delta_0^A} \\
L \searrow \Downarrow \alpha_{\delta_1} \nearrow A_1 \\
a_0 \searrow A_0 \xrightarrow{\delta_1^A}
\end{array}$$

that satisfies the equations for l -descent objects, where all the projections $A \xrightarrow{a_i} A_i$ are tight because **1**, **2** and **3** are all in T .

Since δ_0^A 's are already tight, this means that the projection a_0 is tight and detects tightness.

Now for c -rigged c -descent objects, we again adapt the same dotted \mathcal{F} -category \mathbb{D} as above, and consider dotted-lax natural transformations instead.

Example 5.2.3 (w -rigged equifiers). We start with the l -rigged equifiers.

Let

$$\mathbb{D} = \left\{ \begin{array}{c} f \\ \begin{array}{ccc} \dot{x} & \alpha \Downarrow \Psi \beta & \dot{y} \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & g \end{array} \end{array} \right\},$$

which again is the same as the \mathcal{F} -category used to describe l -rigged equifiers by \mathcal{F} -weighted limits, except that we have dotted objects.

By the naturality of γ , we have

$$\begin{array}{c} L \\ \begin{array}{ccc} \gamma_x \swarrow & Sf & \searrow \gamma_y \\ Sx & \Psi S\alpha & Sy \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & Sg \end{array} \end{array} = \begin{array}{c} L \\ \begin{array}{ccc} \gamma_x \swarrow & \Leftarrow \gamma_g & \searrow \gamma_y \\ Sx & \text{~~~~~} & Sy \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & Sg \end{array} \end{array} = \begin{array}{c} L \\ \begin{array}{ccc} \gamma_x \swarrow & Sf & \searrow \gamma_y \\ Sx & \Psi S\beta & Sy \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & Sg \end{array} \end{array} .$$

So the above data assemble to the equifier in \mathcal{A}_λ :

$$L \xrightarrow{\gamma_x} \begin{array}{c} Sf \\ \begin{array}{ccc} Sx & \alpha \Downarrow \Psi \beta & Sy \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & Sg \end{array} \end{array} .$$

In addition, the projections γ_x and γ_y are tight, and γ_x detects tightness.

For c -rigged equifiers, consider this time when

$$\mathbb{D} = \left\{ \begin{array}{c} f \\ \begin{array}{ccc} \dot{x} & \alpha \Downarrow \Psi \beta & \dot{y} \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & g \end{array} \end{array} \right\}.$$

This is also the same as the \mathcal{F} -category used in describing the limit as an \mathcal{F} -weighted limit.

Now, a dotted-lax natural transformation $\gamma: \Delta(L) \rightarrow S_\lambda$ gives our desired c -rigged equifiers.

For p -rigged equifiers, let

$$\mathbb{D} = \left\{ \begin{array}{c} f \\ \begin{array}{ccc} \dot{x} & \alpha \Downarrow \Psi \beta & \dot{y} \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & g \end{array} \end{array} \right\}.$$

Again, the only difference between the \mathcal{F} -category given in [LS12] and this dotted \mathcal{F} -category is that $g \in \Sigma$.

A dotted lax-natural transformation $\gamma: \Delta(L) \rightarrow S_\lambda$ gives the equifier of S_λ in \mathcal{A}_λ :

$$L \xrightarrow{\gamma_x} \begin{array}{c} Sf \\ \begin{array}{ccc} Sx & \alpha \Downarrow \Psi \beta & Sy \\ \text{~~~~~} & \text{~~~~~} & \text{~~~~~} \\ & \text{~~~~~} & Sg \end{array} \end{array} .$$

The projection γ_x is tight and detects tightness.

Example 5.2.4 (Alternating projective limits). This is an example of a limit which is not PIE but lifts to the \mathcal{F} -category of algebras, as shown in [LS12, Example 6.8].

Consider the opposite poset of natural numbers

$$\dot{1} \leftarrow 2 \leftarrow 3 \leftarrow 4 \cdots ,$$

and denote this \mathcal{F} -category by \mathbb{D} , where all the odd numbers are in T , all the morphisms are in Σ , and the unique morphism $n \rightarrow m$ is tight precisely when it is the identity or when n is even but m is odd.

A dotted-lax natural transformation $\alpha: \Delta(L) \rightarrow S_\lambda$ gives

$$\begin{array}{ccccccc}
 & & & L & & & \\
 & & & \downarrow & & & \\
 \dots & \rightsquigarrow & S(4) & \longrightarrow & S(3) & \rightsquigarrow & S(2) \longrightarrow S(1)
 \end{array}$$

which is the projective limit of S_λ in \mathcal{A}_λ , in addition, the projections at odd numbers are tight.

The projections at odd numbers are tight and detects tightness.

We will discuss more on the above examples in Section 6.

5.3. The equivalence between \mathcal{F} -limits and dotted limits. Motivated by the 2-categorical case, we expect that dotted limits are as expressive as \mathcal{F} -limits. Our main goal in this section is to show that the two notions are equivalent.

Notation 5.3.1. In this section, we denote a surjective-on-objects functor by \twoheadrightarrow , and a full embedding by \hookrightarrow .

Remark 5.3.2. Indeed, surjective-on-objects functors and full embeddings form an orthogonal factorisation system on **Cat**:

For any arbitrary functor $F: A \rightarrow B$, the *embedded image* $\widetilde{\text{im}} F$ of F is the full subcategory of B whose objects are in the image of F .

Now, it is clear that we have a factorisation

$$\begin{array}{ccc}
 A & \xrightarrow{F} & B \\
 \searrow E & & \swarrow M \\
 & \widetilde{\text{im}} F &
 \end{array}$$

of F through $\widetilde{\text{im}} F$, where by default E is a surjective-on-objects functor, and M is a full embedding.

The uniqueness of lifts of surjective-on-objects functors against full embeddings is straightforward.

We first show that any dotted-lax limit has the same universal property as an \mathcal{F} -weighted limit.

Similar to the 2-categorical case in section 3.2, we aim to construct a left adjoint $(\)^\# : [\mathbb{D}, \mathbb{F}]_{l,\Sigma,T} \rightarrow [\mathbb{D}, \mathbb{F}]$ to the inclusion \mathcal{F} -functor, whose universal property is given by

$$(5.2) \quad [\mathbb{D}, \mathbb{F}]_{l,\Sigma,T}(F, G) \cong [\mathbb{D}, \mathbb{F}](F^\#, G).$$

With this left adjoint $(\)^\#$, we can then deduce that for any \mathcal{F} -functor $S: \mathbb{D} \rightarrow \mathbb{A}$ and an object $A \in \mathbb{A}$,

$$\begin{aligned}
 [\mathbb{D}, \mathbb{A}]_{l,\Sigma,T}(\Delta(A), S-) &\cong [\mathbb{D}, \mathbb{F}]_{l,\Sigma,T}(\Delta(1), \mathbb{A}(A, S-)) \\
 &\cong [\mathbb{D}, \mathbb{F}](\Delta(1)^\#, \mathbb{A}(A, S-)),
 \end{aligned}$$

as desired. Indeed, from Proposition 3.2.13, we obtain a 2-categorical left adjoint $(\)_\lambda^\# := (\)^\ddagger : [\mathcal{D}_\lambda, \mathbf{Cat}]_{l,\Sigma} \rightarrow [\mathcal{D}_\lambda, \mathbf{Cat}]$, and so we already have

$$[\mathcal{D}_\lambda, \mathbf{Cat}]_{l,\Sigma}(F_\lambda, G_\lambda) \cong [\mathcal{D}_\lambda, \mathbf{Cat}](F_\lambda^\#, G_\lambda),$$

which gives exactly the loose part of our desired isomorphism (5.2). That means our goal is to construct, for any \mathcal{F} -weight $(F_\tau, F_\lambda, \theta)$, a 2-functor $F_\tau^\# : \mathcal{D}_\tau \rightarrow \mathbf{Cat}$ such that $(F_\tau^\#, F_\lambda^\#, \varphi)$ is an \mathcal{F} -weight, and that (5.2) is fulfilled.

Construction 5.3.3. Let $\mathbb{D} = (\mathbb{D}, \Sigma, T)$ be a small dotted \mathcal{F} -category. Let $F = (F_\tau, F_\lambda, \theta: F_\tau \rightarrow F_\lambda J_{\mathbb{D}})$ be an \mathcal{F} -weight.

We view T as a full sub-2-category of Σ_τ , i.e., T is a 2-category where every morphism is tight and is in Σ . Denote by $J_T^{\mathcal{D}_\tau} : T \hookrightarrow \Sigma_\tau \hookrightarrow \mathcal{D}_\tau$ the inclusion of T into \mathcal{D}_τ .

Since **Cat** is cocomplete and T is small, the left Kan extension $L := \text{Lan}_{J_T^{\mathcal{D}_\tau}} F_\tau J_T^{\mathcal{D}_\tau}$ of $F_\tau J_T^{\mathcal{D}_\tau}$ along $J_T^{\mathcal{D}_\tau}$

$$\begin{array}{ccc}
 T & \xrightarrow{J_T^{\mathcal{D}_\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\
 \searrow J_T^{\mathcal{D}_\tau} & & \downarrow \pi \\
 & & \mathcal{D}_\tau \xrightarrow{L} \mathbf{Cat}
 \end{array}$$

exists. On the other hand, we have 2-natural transformations $\theta \cdot J_T^{\mathcal{D}\tau} : F_\tau J_T^{\mathcal{D}\tau} \rightarrow F_\lambda J_{\mathbb{D}} J_T^{\mathcal{D}\tau}$ and $\eta_F \cdot J_{\mathbb{D}} \cdot J_T^{\mathcal{D}\tau}$, where $\eta_F : F_\lambda \rightarrow F_\lambda^\#$ denotes the component of the unit of the 2-categorical adjunction in Proposition 3.2.13. Therefore, we have a composite

$$\begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\ & \searrow^{J_T^{\mathcal{D}\tau}} & \downarrow^{F_\lambda J_{\mathbb{D}}} \\ & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat} \end{array}$$

$\theta \cdot J_T^{\mathcal{D}\tau} \Downarrow$ $\eta_F \cdot J_{\mathbb{D}} \Downarrow$

of 2-natural transformations, and by the universal property of the left Kan extension, there exists a unique 2-natural transformation $l : L \rightarrow F_\lambda^\# J_{\mathbb{D}}$ such that

$$\begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\ & \searrow^{J_T^{\mathcal{D}\tau}} & \downarrow^{F_\lambda J_{\mathbb{D}}} \\ & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat} \end{array} = \begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\ & \searrow^{J_T^{\mathcal{D}\tau}} & \downarrow^{\pi} \\ & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat} \end{array},$$

$\theta \cdot J_T^{\mathcal{D}\tau} \Downarrow$ $\eta_F \cdot J_{\mathbb{D}} \Downarrow$ $l \Downarrow$

that is to say, $(l \cdot J_T^{\mathcal{D}\tau}) \circ \pi = (\eta_F \cdot J_{\mathbb{D}} \cdot J_T^{\mathcal{D}\tau}) \circ (\theta \cdot J_T^{\mathcal{D}\tau})$. As a result, for each object d in \mathbb{D} , we have a functor $l_d : Ld \rightarrow F_\lambda^\# d$.

Nevertheless, this is not yet the end of our story, as l_d is not necessarily a full embedding. To resolve the issue, let us consider the embedded image $\widetilde{\text{im}} l_d$. Following Remark 5.3.2, we have the factorisation

$$\begin{array}{ccc} Ld & \xrightarrow{l_d} & F_\lambda^\# d \\ & \searrow^{p_d} & \uparrow^{\widetilde{l}_d} \\ & & \widetilde{L}d \end{array}$$

of l_d through $\widetilde{L}d$, where by default p_d is a surjective-on-objects functor, and \widetilde{l}_d is a full embedding. Using the orthogonality of surjective-on-objects functors and full embeddings, \widetilde{L} uniquely extends to a 2-functor such that p_d and \widetilde{l}_d are the 1-components of the 2-natural transformations $p : L \rightarrow \widetilde{L}$ and $\widetilde{l} : \widetilde{L} \rightarrow F_\lambda^\# J_{\mathbb{D}}$, respectively.

By setting $F_\tau^\# := L$ and $\varphi := \widetilde{l}$, we finish our construction of the \mathcal{F} -weight $F^\# = (F_\tau^\#, F_\lambda^\#, \varphi)$.

Now, let $G = (G_\tau, G_\lambda, \gamma : G_\tau \rightarrow G_\lambda J_{\mathbb{D}})$ be another \mathcal{F} -weight. Since $J_{\mathbb{A}}$ is locally fully faithful, if we view T as a full sub-2-category of Σ_τ , then the condition that the 1-component α_t is tight for $t \in T$ can be seen as the existence of a 2-natural transformation $\alpha_\delta : F_\tau J_T^{\mathcal{D}\tau} \rightarrow G_\tau J_T^{\mathcal{D}\tau}$, satisfying (4.1):

$$(5.3) \quad (\alpha_\lambda \cdot J_{\mathbb{D}} J_T^{\mathcal{D}\tau}) \circ (\theta \cdot J_T^{\mathcal{D}\tau}) = (\gamma \cdot J_T^{\mathcal{D}\tau}) \circ \alpha_\delta.$$

More precisely, a dotted-lax natural transformation $\alpha : F \rightarrow G$ in $[\mathbb{D}, \mathbb{F}]_{l, \Sigma}(F, G)$ consists of

- a lax natural transformation $\alpha_\lambda : F_\lambda \rightarrow G_\lambda$ between the loose parts;
- a 2-natural transformation $\alpha_\lambda \cdot J_{\Sigma_\lambda} : F_\lambda J_{\Sigma_\lambda} \rightarrow G_\lambda J_{\Sigma_\lambda}$;
- a 2-natural transformation $\alpha_\delta : F_\tau J_T^{\mathcal{D}\tau} \rightarrow G_\tau J_T^{\mathcal{D}\tau}$,

that fulfill (5.3), i.e.,

$$\begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\ & \searrow^{J_T^{\mathcal{D}\tau}} & \downarrow^{F_\lambda J_{\mathbb{D}}} \\ & & \mathcal{D}_\tau \xrightarrow{G_\lambda J_{\mathbb{D}}} \mathbf{Cat} \end{array} = \begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\ & \searrow^{J_T^{\mathcal{D}\tau}} & \downarrow^{\alpha_\delta} \\ & & \mathcal{D}_\tau \xrightarrow{G_\lambda J_{\mathbb{D}}} \mathbf{Cat} \end{array},$$

$\theta \cdot J_T^{\mathcal{D}\tau} \Downarrow$ $\alpha_\lambda \cdot J_{\mathbb{D}} \Downarrow$ $\alpha_\delta \Downarrow$ $\gamma \Downarrow$

here $\alpha_\lambda \cdot J_{\mathbb{D}} J_T^{\mathcal{D}\tau} : F_\lambda J_{\mathbb{D}} J_T^{\mathcal{D}\tau} \rightarrow G_\lambda J_{\mathbb{D}} J_T^{\mathcal{D}\tau}$ is a 2-natural transformation.

In other words, our goal is to show that $\alpha : F \rightarrow G$ corresponds bijectively to an \mathcal{F} -natural transformation $\beta : F^\# \rightarrow G$, or equivalently, a pair of 2-natural transformations $\beta_\lambda : F_\lambda^\# \rightarrow G_\lambda$ and $\beta_\tau : F_\tau^\# \rightarrow G_\tau$, satisfying (4.1). Since $\beta_\lambda : F_\lambda^\# \rightarrow G_\lambda$ is obtained immediately from the 2-categorical adjunction in Proposition 3.2.13, our task is now reduced to finding an appropriate 2-natural transformation $\beta_\tau : F_\tau^\# \rightarrow G_\tau$.

To begin with, we check that in the 2-categorical adjunction, any unit component and the transposes extend from marked-lax natural transformations to dotted-lax natural transformations.

Proposition 5.3.4. *The component $\eta_F: F_\lambda \rightarrow F_\lambda^\#$ of the unit of the adjunction in Proposition 3.2.13 is a dotted-lax natural transformation.*

In addition, given an \mathcal{F} -natural transformation $\beta: F^\# \rightarrow G$, its transpose obtained by pre-composing with the unit η_F is also a dotted-lax natural transformation.

Proof. It suffices to show that for any $t \in T$, η_{Ft} is a tight morphism in \mathbb{F} , i.e., a functor preserving tight parts.

From the construction of $l = \varphi \circ p$ in Construction 5.3.3, we have $((\varphi \circ p) \cdot J_T^{\mathcal{D}\tau}) \circ \pi = (l \cdot J_T^{\mathcal{D}\tau}) \circ \pi = (\eta_F \cdot J_{\mathbb{D}} \cdot J_T^{\mathcal{D}\tau}) \circ (\theta \cdot J_T^{\mathcal{D}\tau})$. To wit, we have a square

$$\begin{array}{ccc} F_\tau t & \xrightarrow{\theta_t} & F_\lambda t \\ \pi_t \downarrow & & \downarrow \eta_{Ft} \\ Lt & & \\ p_t \downarrow & & \\ F_\tau^\# t & \xrightarrow{\varphi_t} & F_\lambda^\# t \end{array},$$

which means η_F preserves tight parts.

Now, since $\beta_{\tau t}$ is clearly tight for all $t \in T$, β is always a dotted-lax natural transformation. Therefore, the composition $\beta \circ \eta_F$ is also a dotted-lax natural transformation. \square

We attain our main theorem as follows.

Proposition 5.3.5. *There is an isomorphism of categories*

$$[\mathbb{D}, \mathbb{F}]_{l, \Sigma, T}(F, G) \stackrel{\eta_F^*}{\cong} [\mathbb{D}, \mathbb{F}](F^\#, G)$$

given by the pre-composition by η_F , natural in F and G .

In other words, there is a left adjoint $(\)^\#: [\mathbb{D}, \mathbb{F}]_{l, \Sigma, T} \rightarrow [\mathbb{D}, \mathbb{F}]$ to the inclusion.

Proof. Our goal is to prove that given a dotted-lax natural transformation $\alpha: F \rightarrow G$, there exists a 2-natural transformation $\varsigma: F_\tau^\# \rightarrow G_\tau$ such that the transpose $\beta_\lambda: F_\lambda^\# \rightarrow G_\lambda$ of α_λ together with ς become an \mathcal{F} -natural transformation $\beta: F^\# \rightarrow G$, and that $\beta \circ \eta_F = \alpha$; moreover, for any two \mathcal{F} -natural transformation $\beta, \beta': F^\# \rightarrow G$, if $\beta \circ \eta_F = \beta' \circ \eta_F$, then $\beta = \beta'$.

By the universal property of the left Kan extension L in Construction 5.3.3, there exists a unique 2-natural transformation $g: L \rightarrow G_\tau$ such that

$$\begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau & \xrightarrow{F_\tau} & \mathbf{Cat} \\ & \searrow J_T^{\mathcal{D}\tau} & \alpha_\delta \downarrow & \nearrow G_\tau & \\ & & \mathcal{D}_\tau & & \end{array} = \begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau & \xrightarrow{F_\tau} & \mathbf{Cat} \\ & \searrow J_T^{\mathcal{D}\tau} & \pi \downarrow & \nearrow L & \nearrow \exists! g \\ & & \mathcal{D}_\tau & & \nearrow G_\tau \end{array},$$

i.e., $(g \cdot J_T^{\mathcal{D}\tau}) \circ \pi = \alpha_\delta$. From the 2-categorical adjunction, we have $\beta_\lambda \circ \eta_F = \alpha_\lambda$, so by (5.3), the equation then becomes

$$\begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau & \xrightarrow{F_\tau} & \mathbf{Cat} \\ & \searrow J_T^{\mathcal{D}\tau} & \theta \cdot J_T^{\mathcal{D}\tau} \downarrow & \nearrow F_\lambda J_{\mathbb{D}} & \nearrow \eta_F \cdot J_{\mathbb{D}} \\ & & \mathcal{D}_\tau & \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} & \mathbf{Cat} \\ & & & \nearrow \beta_\lambda \cdot J_{\mathbb{D}} & \nearrow G_\lambda J_{\mathbb{D}} \end{array} = \begin{array}{ccc} T & \xrightarrow{J_T^{\mathcal{D}\tau}} & \mathcal{D}_\tau & \xrightarrow{F_\tau} & \mathbf{Cat} \\ & \searrow J_T^{\mathcal{D}\tau} & \pi \downarrow & \nearrow L & \nearrow g \\ & & \mathcal{D}_\tau & \xrightarrow{F_\tau} & \mathbf{Cat} \\ & & & \nearrow G_\tau & \nearrow \gamma \cdot J_{\mathbb{D}} \\ & & & & \nearrow G_\lambda J_{\mathbb{D}} \end{array}.$$

Besides, from the construction of l in Construction 5.3.3, we have

$$\begin{array}{ccc}
 T & \xrightarrow{J_T^{\mathcal{D}_\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\
 & \searrow^{J_T^{\mathcal{D}_\tau}} & \downarrow^{\theta \cdot J_T^{\mathcal{D}_\tau}} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda J_{\mathbb{D}}} \mathbf{Cat} \\
 & & \downarrow^{\eta_{F \cdot J_{\mathbb{D}}}} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat}
 \end{array}
 =
 \begin{array}{ccc}
 T & \xrightarrow{J_T^{\mathcal{D}_\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\
 & \searrow^{J_T^{\mathcal{D}_\tau}} & \downarrow^{\pi} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat} \\
 & & \downarrow^{\beta_\lambda \cdot J_{\mathbb{D}}} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat}
 \end{array}
 .$$

Combining the two equations, we obtain

$$\begin{array}{ccc}
 T & \xrightarrow{J_T^{\mathcal{D}_\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\
 & \searrow^{J_T^{\mathcal{D}_\tau}} & \downarrow^{\pi} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat} \\
 & & \downarrow^{\beta_\lambda \cdot J_{\mathbb{D}}} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat}
 \end{array}
 =
 \begin{array}{ccc}
 T & \xrightarrow{J_T^{\mathcal{D}_\tau}} & \mathcal{D}_\tau \xrightarrow{F_\tau} \mathbf{Cat} \\
 & \searrow^{J_T^{\mathcal{D}_\tau}} & \downarrow^{\pi} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat} \\
 & & \downarrow^{\beta_\lambda \cdot J_{\mathbb{D}}} \\
 & & \mathcal{D}_\tau \xrightarrow{F_\lambda^\# J_{\mathbb{D}}} \mathbf{Cat}
 \end{array}
 .$$

Now the universal property of L forces $\gamma \circ g = (\beta_\lambda \cdot J_{\mathbb{D}}) \circ l$. This means that for any object d in \mathbb{D} , we have a commutative square

$$\begin{array}{ccc}
 Ld & \xrightarrow{g_d} & G_\tau d \\
 p_d \downarrow & \nearrow^{\exists! \varsigma_d} & \downarrow \gamma_d \\
 F_\tau^\# d & \xrightarrow{\varphi_d} F_\lambda^\# d & \xrightarrow{\beta_\lambda d} G_\lambda d
 \end{array}
 ,$$

and since p_d is surjective-on-objects and γ_d is a full embedding, there exists a unique lift $\varsigma_d: F_\tau^\# d \rightarrow G_\tau d$.

Let $f, f_i: c \rightarrow d$ be tight morphisms in \mathcal{D}_τ for $i = 1, 2$, and $m: f_1 \Rightarrow f_2$ be a 2-cell in \mathbb{D} . In the following diagram:

$$\begin{array}{ccccc}
 F_\tau^\# c & \xrightarrow{\varphi_c} & F_\lambda^\# c & \xrightarrow{\beta_\lambda c} & G_\tau c \\
 \downarrow^{F_\tau^\# f_1} & \searrow^{F_\tau^\# m} & \downarrow^{F_\lambda^\# f_1} & \searrow^{F_\lambda^\# m} & \downarrow^{G_\tau f_1} \\
 F_\tau^\# f_1 & \xrightarrow{F_\tau^\# m} & F_\tau^\# f_2 & \xrightarrow{F_\lambda^\# m} & G_\tau f_2 \\
 \downarrow^{F_\tau^\# f_1} & \searrow^{F_\tau^\# m} & \downarrow^{F_\lambda^\# f_1} & \searrow^{F_\lambda^\# m} & \downarrow^{G_\tau f_1} \\
 F_\tau^\# d & \xrightarrow{\varphi_d} & F_\lambda^\# d & \xrightarrow{\beta_\lambda d} & G_\tau d \\
 \downarrow^{F_\tau^\# f_1} & \searrow^{F_\tau^\# m} & \downarrow^{F_\lambda^\# f_1} & \searrow^{F_\lambda^\# m} & \downarrow^{G_\tau f_1} \\
 F_\tau^\# f_1 & \xrightarrow{F_\tau^\# m} & F_\tau^\# f_2 & \xrightarrow{F_\lambda^\# m} & G_\tau f_2 \\
 \downarrow^{F_\tau^\# f_1} & \searrow^{F_\tau^\# m} & \downarrow^{F_\lambda^\# f_1} & \searrow^{F_\lambda^\# m} & \downarrow^{G_\tau f_1} \\
 F_\tau^\# d & \xrightarrow{\varphi_d} & F_\lambda^\# d & \xrightarrow{\beta_\lambda d} & G_\tau d \\
 \downarrow^{F_\tau^\# f_1} & \searrow^{F_\tau^\# m} & \downarrow^{F_\lambda^\# f_1} & \searrow^{F_\lambda^\# m} & \downarrow^{G_\tau f_1} \\
 F_\tau^\# f_1 & \xrightarrow{F_\tau^\# m} & F_\tau^\# f_2 & \xrightarrow{F_\lambda^\# m} & G_\tau f_2 \\
 \downarrow^{F_\tau^\# f_1} & \searrow^{F_\tau^\# m} & \downarrow^{F_\lambda^\# f_1} & \searrow^{F_\lambda^\# m} & \downarrow^{G_\tau f_1} \\
 F_\tau^\# d & \xrightarrow{\varphi_d} & F_\lambda^\# d & \xrightarrow{\beta_\lambda d} & G_\tau d
 \end{array}
 ,$$

all the parallelograms in the front, i.e., those constituted by solid arrows, commute, because φ , $\beta_\lambda \cdot J_{\mathbb{D}}$, and γ are all 2-natural transformations, and the 1-components of φ and γ are monomorphisms in \mathbf{Cat} . Thus, the rectangle at the back, i.e., that constituted by two dashed arrows, also commutes. This amounts to $G_\tau f \varsigma_c = \varsigma_d F_\tau^\# f$ and $G_\tau m \varsigma_c = \varsigma_d F_\tau^\# m$. Consequently, $\{\varsigma_d\}_{d \in \mathbb{D}}$ assemble to a 2-natural transformation $\varsigma: F_\tau^\# \rightarrow G_\tau$.

From the above, we have $(\beta_\lambda \cdot J_{\mathbb{D}}) \circ \varphi = \gamma \circ \varsigma$, which is exactly (4.1). Therefore, β_λ and ς together form an \mathcal{F} -natural transformation $\beta: F^\# \rightarrow G$.

Next, we check that for $t \in T$, $\varsigma_c \eta_{F_t} = \alpha_{\lambda t}$. From the 2-categorical adjunction, we already have $\beta_\lambda \circ \eta_F = \alpha_\lambda$.

Indeed, for any $t \in T$, we have the following commutative diagram:

$$\begin{array}{ccc}
F_\tau t & \xrightarrow{\theta_t} & F_\lambda t \\
p_t \pi_t \downarrow & & \downarrow \eta_{F_t} \\
F_\tau^\# t & \xrightarrow{\varphi_t} & F_\lambda^\# t \\
\varsigma_t \downarrow & & \downarrow \beta_{\lambda t} \\
G_\tau t & \xrightarrow{\gamma_t} & G_\lambda t
\end{array},$$

where the top diagram is constructed in the proof of Proposition 5.3.4, and the bottom diagram follows immediately from our above construction of ς_t . Together with (5.3), we obtain

$$\gamma_t \alpha_{\delta_t} = \alpha_{\lambda_t} \theta_t = \beta_{\lambda_t} \eta_{F_t} \theta_t = \gamma_t \varsigma_t p_t \pi_t,$$

now since γ_t is a monomorphism, we deduce that

$$\alpha_{\delta_t} = \varsigma_t p_t \pi_t.$$

Altogether, we obtain $\beta \circ \eta_F = \alpha$.

Finally, let $\beta, \beta': F^\# \rightarrow G$ be two \mathcal{F} -natural transformations such that $\beta \circ \eta_F = \beta' \circ \eta_F$.

In particular, $\beta_\lambda \circ \eta_F = \beta'_\lambda \circ \eta_F$, which implies $\beta_\lambda = \beta'_\lambda$. Hence, we have for any $d \in \text{ob } \mathbb{D}$,

$$\gamma_d \beta_{\tau_d} = \beta_{\lambda_d} \varphi_d = \beta'_{\lambda_d} \varphi_d = \gamma_d \beta'_{\tau_d}.$$

Since γ_d is monic, this implies $\beta_{\tau_d} = \beta'_{\tau_d}$; as d is arbitrary, we obtain $\beta_\tau = \beta'_\tau$. As a consequence, $\beta = \beta'$. \square

Theorem 5.3.6. *Let (\mathbb{D}, Σ, T) be a dotted \mathcal{F} -category. The dotted-lax limit of an \mathcal{F} -functor $S: \mathbb{D} \rightarrow \mathbb{A}$ has the same universal property as the \mathcal{F} -weighted limit $\{\Delta(1)^\#, S\}$.*

Proof. We have

$$\begin{aligned}
\mathbb{A}(A, \text{d}_l \lim S) &\cong [\mathbb{D}, \mathbb{A}]_{l, \Sigma, T}(\Delta(A), S) \\
&\cong [\mathbb{D}, \mathbb{F}]_{l, \Sigma, T}(\Delta(1), \mathbb{A}(A, S-)) \\
&\cong [\mathbb{D}, \mathbb{F}](\Delta(1)^\#, \mathbb{A}(A, S-)) \\
&\cong \mathbb{A}(A, \{\Delta(1)^\#, S\}).
\end{aligned}$$

\square

We proceed to show the converse: any \mathcal{F} -weighted limit can be equivalently viewed as a dotted limit.

Let \mathbb{A}, \mathbb{D} be \mathcal{F} -categories, $\Phi: \mathbb{D} \rightarrow \mathbb{F}$ be an \mathcal{F} -weight and $S: \mathbb{D} \rightarrow \mathbb{A}$ be an \mathcal{F} -functor. Consider the \mathcal{F} -category of elements of Φ with

- objects the pairs $(D \in \mathbb{D}, \delta \in (\Phi D)_\lambda)$;
- loose morphisms $(D_1, \delta_1) \rightsquigarrow (D_2, \delta_2)$ given by the pairs $(d: D_1 \rightsquigarrow D_2 \in \mathcal{D}_\lambda, \omega: Wd\delta_1 \rightarrow \delta_2)$;
- tight morphisms $(D_1, \delta_1) \rightarrow (D_2, \delta_2)$ given by the pairs $(d: D_1 \rightarrow D_2 \in \mathcal{D}_\tau, \omega: Wd\delta_1 \rightarrow \delta_2)$;
- 2-cells $(d, \omega) \Rightarrow (d', \omega')$ given by the 2-cells $\alpha: d \Rightarrow d'$ such that $\omega' W\alpha\delta_1 = \omega$,

where the composition of two morphisms is given by the same rule as in the 2-categorical case. Similar to the 2-category of elements, we also have a projection \mathcal{F} -functor

$$P: \mathbb{E}l\Phi \rightarrow \mathbb{D}.$$

Remark 5.3.7. In [Lam20], Lambert presented a proof of 2-categories, 2-functors, lax natural transformations, and modifications forming a lax 3-category $\mathbf{2-Cat}_l$. Using a very similar argument, we see that \mathcal{F} -categories, \mathcal{F} -functors, lax natural transformations between loose parts, and modifications form a lax 3-category $\mathcal{F}\text{-Cat}_{l, \lambda}$.

Following the definition of lax comma objects in [Mes23], one can easily check that $\mathbb{E}l\Phi$ is the lax comma object of $\Delta(1): 1 \rightarrow \mathbb{F}$ and $\Phi: \mathbb{D} \rightarrow \mathbb{F}$

$$\begin{array}{ccc}
\mathbb{E}l\Phi & \xrightarrow{P} & \mathbb{D} \\
! \downarrow & \nearrow \mu & \downarrow \Phi \\
1 & \xrightarrow{\Delta(1)} & \mathbb{F}
\end{array}$$

in $\mathcal{F}\text{-Cat}_{l,\lambda}$.

Proposition 5.3.8. *Let $\Sigma := \{(d \in \text{mor } \mathbb{D}, 1)\}$ be the class of morphisms, and $T := \{(D, \delta \in (\Phi D)_\tau)\}$ be the class of objects in $\mathbb{E}\mathbb{I}\Phi$. There is an isomorphism*

$$[\mathbb{D}, \mathbb{F}](\Phi, \mathbb{A}(A, S-)) \cong [\mathbb{E}\mathbb{I}\Phi, \mathbb{A}]_{l,\Sigma,T}(\Delta(A), SP)$$

in \mathbb{F} , or equivalently, there is an isomorphism of categories

$$[\mathbb{D}, \mathbb{F}]_\lambda(\Phi, \mathbb{A}(A, S-)) \cong [\mathbb{E}\mathbb{I}\Phi, \mathbb{A}]_{l,\Sigma,T,\lambda}(\Delta(A), SP),$$

which restricts to the tight parts

$$[\mathbb{D}, \mathbb{F}]_\tau(\Phi, \mathbb{A}(A, S-)) \cong [\mathbb{E}\mathbb{I}\Phi, \mathbb{A}]_{l,\Sigma,T,\tau}(\Delta(A), SP).$$

Proof. The isomorphism for the loose parts are given by Proposition 3.3.2. It remains to check that the tight morphisms on both sides correspond to each other.

Recall that in the proof of Proposition 3.3.2, we defined a pair of inverse functors

$$\begin{aligned} G: [\mathcal{D}_\lambda, \mathbf{Cat}](\Phi_\lambda, \mathcal{A}_\lambda(A, S_\lambda-)) &\rightarrow [\mathcal{E}\ell\Phi, \mathbf{Cat}]_{l,\Sigma}(\Delta(1), \mathcal{A}_\lambda(A, S_\lambda P-)), \\ \beta &\mapsto \beta P \circ \mu, \\ \Gamma: \beta_1 &\rightarrow \beta_2 \mapsto \Gamma P \circ \mu, \end{aligned}$$

and

$$\begin{aligned} G': [\mathcal{E}\ell\Phi, \mathbf{Cat}]_{l,\Sigma}(\Delta(1), \mathcal{A}_\lambda(A, S_\lambda P-)) &\rightarrow [\mathcal{D}_\lambda, \mathbf{Cat}](\Phi_\lambda, \mathcal{A}_\lambda(A, S_\lambda-)), \\ \alpha &\mapsto \beta = \{\alpha_{(\Delta(D), -)}\}_{D \in \mathcal{D}}, \\ \Lambda: \alpha_1 &\rightarrow \alpha_2 \mapsto \Theta = \{\{\Lambda_{(D,\delta)}\}_{\delta \in WD}\}_{D \in \mathcal{D}}. \end{aligned}$$

Now suppose $\beta: \Phi_\lambda \rightarrow \mathcal{A}_\lambda(A, S_\lambda-)$ is tight, i.e., β is a 2-natural transformation such that for any $\delta \in (\Phi D)_\tau$, $\beta_D(\delta): A \rightarrow S_\lambda D$ is tight in \mathbb{A} . Clearly, $G\beta = (\beta \circ P) \circ \mu =: \alpha$ is a marked-lax nat transformation $\Delta(1) \xrightarrow{\alpha} \mathcal{A}_\lambda(A, S_\lambda P-)$. Note that $\alpha_{(D,\delta)}$ is given by the composite $\beta_D \circ \mu_{(D,\delta)}$. If $(D, \delta) \in T$, then that means $\mu_{(D,\delta)}$ always picks out an object in $(\Phi D)_\tau$, therefore $\beta_D(\delta)$ is tight by the assumption, hence is $\alpha_{(D,\delta)}$.

Conversely, let $\alpha: \Delta(1) \rightarrow \mathcal{A}_\lambda(A, S_\lambda P-)$ be a dotted natural transformation, and so $G'(\alpha) =: \beta_\alpha$ is a 2-natural transformation. Now for any $\delta \in (\Phi D)_\tau$, we have $(D, \delta) \in T$, thus $\beta_{\alpha_D}(d) = \alpha_{(D,\delta)}$ is tight. \square

Theorem 5.3.9. *Let \mathbb{A}, \mathbb{D} be \mathcal{F} -categories. Let $\Phi: \mathbb{D} \rightarrow \mathbb{F}$ be an \mathcal{F} -weight and $S: \mathbb{D} \rightarrow \mathbb{A}$ be an \mathcal{F} -functor. Let $\Sigma := \{(d \in \text{mor } \mathbb{D}, 1)\}$ be the class of morphisms, and $T := \{(D, \delta \in (\Phi D)_\tau)\}$ be the collection of objects in the \mathcal{F} -category of elements $\mathbb{E}\mathbb{I}\Phi$ of Φ , so that $(\mathbb{E}\mathbb{I}\Phi, \Sigma, T)$ is a dotted \mathcal{F} -category. The \mathcal{F} -weighted limit $\{\Phi, S\}$ has the same universal property as the dotted-lax limit of the \mathcal{F} -functor $SP: \mathbb{E}\mathbb{I}\Phi \rightarrow \mathbb{A}$.*

Proof. By Proposition 5.3.8, we have a chain of isomorphisms

$$\begin{aligned} \mathbb{A}(A, \{\Phi, S\}) &\cong [\mathbb{D}, \mathbb{F}](\Phi, \mathbb{D}(A, S-)) \\ &\cong [\mathbb{E}\mathbb{I}\Phi, \mathbb{A}]_{l,\Sigma}(\Delta(A), SP) \\ &\cong \mathbb{A}(A, \text{d}_l \lim SP). \end{aligned} \quad \square$$

6. DISCUSSIONS

In [Szy19], Szyld showed the lifting theorem for 2-categorical limits using marked limits, which provides extra information on the projections than the classical proof. Indeed, Szyld's theorem can be rephrased in a cleaner manner in terms of enhanced 2-category theory and dotted limits, for the case of strict and lax (or colax) T -morphisms.

Definition 6.0.1. A dotted \mathcal{F} -category $(\mathbb{D}, \Sigma, \Gamma)$ is said to be weakly PIE-indexing precisely when each connected component of the full sub- \mathcal{F} -category $\Sigma \subseteq \mathbb{D}$ has an initial object, which is contained in Γ .

A weakly PIE-indexing dotted \mathcal{F} -category is said to be strongly PIE-indexing precisely if furthermore the unique morphism from the initial object to any object is always tight.

Notation 6.0.2. We denote by N the collection of these initial objects.

Now, [Szy19, Theorem 3.3] can be reformulated as

Theorem 6.0.3. Let $(\mathbb{D}, \Sigma, \Gamma)$ be a weakly PIE-indexing dotted \mathcal{F} -category, and let T be a 2-monad on a 2-category \mathcal{A} , viewed as a chordate \mathcal{F} -category. Let $S: \mathbb{D} \rightarrow \mathbf{T}\text{-Alg}_{s,p}$ be an \mathcal{F} -functor. Denote by $U_{s,p}: \mathbf{T}\text{-Alg}_{s,p} \rightarrow \mathcal{A}$ the underlying \mathcal{F} -functor. If $\mathrm{d}_l \lim U_{s,p} S$ exists in \mathcal{A} , then $\mathrm{d}_l \lim S$ exists in $\mathbf{T}\text{-Alg}_{s,p}$, and is preserved by $U_{s,p}$.

Moreover, the projections $\{\pi_n\}_{n \in N}$ of $\mathrm{d}_l \lim S$ are tight and jointly detect tightness.

Theorem 6.0.4. Let $(\mathbb{D}, \Sigma, \Gamma)$ be a strongly PIE-indexing dotted \mathcal{F} -category, and let T be a 2-monad on a 2-category \mathcal{A} , viewed as a chordate \mathcal{F} -category. Let $S: \mathbb{D} \rightarrow \mathbf{T}\text{-Alg}_{s,c}$ be an \mathcal{F} -functor. Denote by $U_{s,c}: \mathbf{T}\text{-Alg}_{s,c} \rightarrow \mathcal{A}$ the underlying \mathcal{F} -functor. If $\mathrm{d}_l \lim U_{s,c} S$ exists in \mathcal{A} , then $\mathrm{d}_l \lim S$ exists in $\mathbf{T}\text{-Alg}_{s,c}$, and is preserved by $U_{s,c}$.

Moreover, the projections $\{\pi_n\}_{n \in N}$ of $\mathrm{d}_l \lim S$ are tight and jointly detect tightness.

Dually, If $\mathrm{d}_c \lim U_{s,l} S$ exists in \mathcal{A} , then $\mathrm{d}_c \lim S$ exists in $\mathbf{T}\text{-Alg}_{s,l}$, and is preserved by $U_{s,l}$, and the projections of $\mathrm{d}_c \lim S$ are tight and jointly detect tightness.

It is shown in [LS12] that w -rigged \mathcal{F} -weighted limits such as w -rigged inserters, w -rigged equifiers, w -rigged w -descent objects and w -rigged comma objects all lift to $\mathbf{T}\text{-Alg}_{s,w'}$, where $w' = c$ if $w = l$ and vice versa. Indeed, they are precisely the limits that lift. Looking at the examples of p -rigged limits illustrated in Section 5.2, it is clear that the assumptions in Theorem 6.0.3 are met; looking at the examples of l -rigged and c -rigged limits illustrated in Section 5.2, it is immediate that the assumptions in Theorem 6.0.4 are fulfilled. In other words, through the lens of dotted limits, we see that Szyld's lifting theorem also applies to some of the \mathcal{F} -categorical limits.

Nevertheless, the alternating projective limits in Example 5.2.4, which is shown to lift in [LS12], do not satisfy the assumptions: there is no initial object in the indexing dotted \mathcal{F} -category, as m tends to infinity.

A possible direction and further exploration on dotted limits is to characterise w -rigged \mathcal{F} -weighted limits in the language of dotted limits, so that a more elementary and explicit description of rigged limits can be found.

REFERENCES

- [BKP89] Robert Blackwell, Gregory Maxwell Kelly, and John Power. Codescent objects and coherence. *Journal of Pure and Applied Algebra*, 59:1–41, 1989.
- [Bou10] John Bourke. *Codescent objects in 2-dimensional universal algebra*. PhD thesis, University of Sydney, Carlsaw Building F07, Eastern Ave, Camperdown NSW 2006, Australia, 8 2010.
- [Bou14] John Bourke. Two-dimensional monadicity. *Advances in Mathematics*, 252:708–747, 2014.
- [DDS18] María Emilia Descotte, Eduardo Dubuc, and Martin Szyld. Sigma limits in 2-categories and flat pseudofunctors. *Advances in Mathematics*, 333:266–313, 2018.
- [GHL22] Andrea Gagna, Yonatan Harpaz, and Edoardo Lanari. Bilimits are bifinal objects. *Journal of Pure and Applied Algebra*, 226(12):107137, 2022.
- [Gra74] John Walker Gray. *Formal category theory: adjointness for 2-categories*, volume 391. Springer Lecture Notes in Mathematics, 1974.
- [Kel82] Gregory Maxwell Kelly. *Basic concepts of enriched category theory*. Cambridge University Press, 1982.
- [Lac02] Stephen Lack. Codescent objects and coherence. *Journal of Pure and Applied Algebra*, 175(1–3):223–241, 2002.
- [Lac05] Stephen Lack. Limits for lax morphisms. *Applied Categorical Structures*, 13(3):189–203, 2005.
- [LS12] Stephen Lack and Michael Shulman. Enhanced 2-categories and limits for lax morphismss. *Advances in Mathematics*, 229(1):294–356, 2012.

- [Lam20] Michael Lambert. Discrete 2-fibrations, 2020. <https://arxiv.org/abs/2001.11477>.
- [Mes23] Luca Mesiti. The 2-set-enriched grothendieck construction and the lax normal conical 2-limits, 2023. <https://arxiv.org/abs/2302.04566>.
- [Str76] Ross Street. Limits indexed by category-valued 2-functors. *Journal of Pure and Applied Algebra*, 8:149–181, 1976.
- [Szy19] Martin Szyld. Lifting pie limits with strict projections. *Theory and Applications of Categories*, 34(1):1–12, 2019.
- Email address:* joanna.ko.maths@gmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, MASARYK UNIVERSITY, KOTLÁŘSKÁ 2, BRNO 61137, CZECH REPUBLIC