

ALMOST SURE AVERAGING FOR EVOLUTION EQUATIONS DRIVEN BY FRACTIONAL BROWNIAN MOTIONS*

B. PEI[†], B. SCHMALFUSS[‡], AND Y. XU[§]

Abstract. We apply the averaging method to a coupled system consisting of two evolution equations which has a slow component driven by fractional Brownian motion (FBM) with the Hurst parameter $H_1 > \frac{1}{2}$ and a fast component driven by additive FBM with the Hurst parameter $H_2 \in (1 - H_1, 1)$. The main purpose is to show that the slow component of such a couple system can be described by a stochastic evolution equation with averaged coefficients. Our first result provides a pathwise mild solution for the system of mixed stochastic evolution equations. Our main result deals with an averaging procedure which proves that the slow component converges almost surely to the solution of the corresponding averaged equation using the approach of time discretization. To do this we generate a stationary solution by a exponentially attracting random fixed point of the random dynamical system generated by the fast component.

Key words. Almost sure averaging, Random fixed points, Fractional Brownian motion, Slow-fast systems

MSC codes. 60G22, 60H05, 60H15, 34C29.

1. Introduction. The aim of this article is to address the almost sure averaging for the problem of slow-fast stochastic evolution equation in a separable Hilbert space V

$$(1.1) \quad dX^\epsilon(t) = AX^\epsilon(t) dt + f(X^\epsilon(t), Y^\epsilon(t)) dt + h(X^\epsilon(t)) dB^{H_1}(t), X^\epsilon(0) = X_0 \in V,$$

$$(1.2) \quad dY^\epsilon(t) = \frac{1}{\epsilon}BY^\epsilon(t) dt + \frac{1}{\epsilon}g(X^\epsilon(t), Y^\epsilon(t)) dt + dB^{H_2}(t/\epsilon), Y^\epsilon(0) = Y_0 \in V$$

where $0 < \epsilon \ll 1$ is a small parameter, $X^\epsilon(t) \in V$ and $Y^\epsilon(t) \in V$ are the state variables, B^{H_1}, B^{H_2} are the trace class V -valued fractional Brownian motions (FBMs) with $H_1 \in (1/2, 1)$, $H_2 \in (1 - H_1, 1)$, f, h, g are sufficiently regular.

Very often a complex system can be viewed as a combination of slow and fast motions [12, 22], see (1.1)-(1.2) for instance, which leads to two widely separated time scales equations and is extremely difficult to analyze directly. It is highly desirable to obtain a simplified equation capturing the dynamics of the system at the slow time scale. Averaging plays an important role to extract effective macroscopic dynamics which can describe approximately the slow motion (see [3, 37] for the deterministic case and [9, 11, 12, 20] for the stochastic case). It is worth mentioning that this idea was exploited in atmospheric science where, for instance, in the study of the climate variability by Hasselmann [19] who got the Nobel Prize in Physics 2021 applying

*Submitted to the editors 02.20.2023

Funding: B. Pei was partially supported by National Natural Science Foundation of China (NSFC) under Grant No.12172285, Guangdong Basic and Applied Basic Research Foundation under Grant No.2214050001158, and Fundamental Research Funds for the Central Universities. Y. Xu was partially supported by Key International (Regional) Joint Research Program of NSFC under Grant No.12120101002. B. Pei thanks the Alexander von Humboldt Foundation (Germany) for support.

[†]School of Mathematics and Statistics, Northwestern Polytechnical University, 127 West Youyi Road, 710072, Xi'an, China and Institut für Stochastik, Friedrich Schiller Universität, Jena, Ernst Abbe Platz 2, D-77043, Jena, Germany (binpei@nwpu.edu.cn).

[‡]Institut für Stochastik, Friedrich Schiller Universität, Jena, Ernst Abbe Platz 2, D-77043, Jena, Germany (bjoern.schmalfuss@uni-jena.de).

[§]School of Mathematics and Statistics, Northwestern Polytechnical University, 127 West Youyi Road, 710072, Xi'an, China (hsux3@nwpu.edu.cn).

the stochastic averaging framework considering climate and weather as slow and fast motions, respectively. Later, Arnold [2] has recast Hasselmann's program of reducing complex deterministic multiscale climate models to simpler stochastic models for the slow variables. Kifer [21] contained a short survey of stochastic averaging methods for random dynamical system (RDS) with application to climate models. Eichinger et al. [10] studied the sample paths estimates for stochastic fast-slow systems driven by FBM and also illustrated their results in an example arising in climate modeling.

The theory of stochastic averaging has a popular history which can be traced back to the work of Khasminskii [20]. We here mention only a few relevant references. Freidlin and Wentzell [12] provided a mathematically rigorous overview of fundamental stochastic averaging procedures. Pavilotis and Stuart [30] covered stochastic averaging and homogenization results obtained by perturbation analysis, see, e.g. [16, 26, 38, 39, 40, 35] (and the references therein) for further generalizations. Most of the known results in the literature mainly considered the case of perturbation by Brownian motion (BM). While slow-fast systems with FBM have seen a tremendous spike of interest in the last two years [18, 25, 24, 32, 33, 29]. We mention only the most relevant result in our case here. Hairer and Li [18] considered slow-fast systems where the slow system is driven by FBM and proved the convergence to the averaged solution took place in probability. Pei, Inahama and Xu [32, 34] answered affirmatively that an averaging principle still holds for fast-slow mixed stochastic differential equations (SDEs) if disturbances involve both BM and FBM $H \in (1/3, 1)$ in the mean square sense. The result [32] was extended in [33] to establish an averaging principle in the mean square sense for stochastic partial differential equation (SPDEs) driven by FBM with an additional fast-varying diffusion process.

One naturally wonders what happens to the averaging principle of this type when the driving noise of fast motion does not have (semi) martingale property. A first attempt to study the long time behaviour of SDEs driven by FBM was made by Hairer [17]. The approach is closer to the usual Markovian semigroup approach to invariant measures than the theory of RDS. Note that FBM does not define the Markov process as in the case of usual BM. Therefore, it is not possible to apply standard methods to show existence of stationary solutions. This research direction is fairly new and there are not many papers at the moment. Li and Sieber [25] established a quantitative quenched ergodic theorem on the conditional evolution of the process of the fast dynamics with frozen slow input and proved a fractional averaging principle for interacting slow-fast systems including a non-Markovian fast dynamics in probability. However, we will show that the fast motion is shown to define RDS which has the unique, exponentially attracting random fixed point and our intention in this article is to study the almost sure averaging replacing the invariant measures by the random fixed points of fast motion which are pathwise exponentially attracting. Note that the almost sure averaging for evolution equations driven by two FBMs is still an open problem in both finite dimensional and infinite dimensional state spaces. The key idea to solve the problem is to replace the stationary solution of a Markov semigroup by the attracting random fixed point of a RDS, which can exist in the case of non-white noise.

To investigate the almost sure averaging of system (1.1)-(1.2), it is essential to obtain the unique solution. Another simple understanding on (1.1)-(1.2) is to rewrite

this system in the following form

$$\begin{aligned} \begin{pmatrix} dX^\epsilon(t) \\ dY^\epsilon(t) \end{pmatrix} &= \begin{pmatrix} A & O \\ O & \frac{1}{\epsilon}B \end{pmatrix} \begin{pmatrix} X^\epsilon(t) \\ Y^\epsilon(t) \end{pmatrix} dt + \begin{pmatrix} f(X^\epsilon(t), Y^\epsilon(t)) \\ \frac{1}{\epsilon}g(X^\epsilon(t), Y^\epsilon(t)) \end{pmatrix} dt \\ &+ \begin{pmatrix} h(X^\epsilon(t)) & O \\ O & \text{id} \end{pmatrix} \begin{pmatrix} dB^{H_1}(t) \\ dB^{H_2}(t/\epsilon) \end{pmatrix}. \end{aligned}$$

Existence and uniqueness of solution to this kind of equation have been established, for instance, Maslowski and Nualart [28], Garrido-Atienza, et.al. [13], Chen, et.al. [4] and Pei, et.al. [33]. In this last paper, the authors were able to overcome the lack of the regularity relying on a pathwise approach, a stopping time technique and an approximation for the fractional noise. However, our technique differs qualitatively from the method mentioned above.

The paper is organized as follows. In Section 2 we formulate the basic properties of RDE and of stochastic integrals with respect to the FBM that are used in the paper. Section 3 contains the existence and uniqueness of a pathwise mild solution to the nonlinear infinite-dimensional evolution equations. In Section 4, we prove that almost sure averaging for evolution equations (1.1)-(1.2). The random fixed points for the RDS generated by (1.2) are concluded in Section 4.2.

2. Preliminaries on the random perturbations and pathwise stochastic integrals. In this section we review some basic concepts of pathwise stochastic integrals that will be used later.

2.1. Random perturbations. Let $(V, (\cdot, \cdot))$ be a separable Hilbert space and its norm is denoted by $\|\cdot\|$. Let $C([T_1, T_2]; V)$ be the space of continuous functions on $[T_1, T_2]$ with values in V equipped with the usual norm

$$\|u\|_\infty = \|u\|_{\infty, T_1, T_2} = \sup_{s \in [T_1, T_2]} \|u(s)\|.$$

We consider a V -valued continuous trace class FBM on some interval $[0, T]$ denoted B^{H_1} with Hurst parameter $H_1 \in (1/2, 1)$. The distribution \mathbb{P}_{H_1} of this process is determined by the covariance

$$(2.1) \quad \mathbb{E}[B^{H_1}(t) \otimes B^{H_1}(s)] = Q_1(|t|^{2H_1} + |s|^{2H_1} - |t-s|^{2H_1}), \quad \mathbb{E}[B^{H_1}(t)] = 0$$

where \mathbb{P}_{H_1} is defined on $\mathcal{B}(C_0([0, T]; V))$ the Borel- σ -algebra of the space of continuous functions $C_0([0, T]; V)$ which are zero at zero. Q_1 be a symmetric positive operator of finite trace. In addition, for H_2 in $(1-H_1, 1)$ let B^{H_2} be a continuous trace class FBM on \mathbb{R} with distribution \mathbb{P}_{H_2} defined on $\mathcal{B}(C_0(\mathbb{R}; V))$ where $C_0(\mathbb{R}; V)$ is the set of continuous paths on \mathbb{R} with values in V which are zero at zero equipped with the compact open topology. The covariance of this stochastic process can be defined by (2.1) with a trace class operator Q_2 and replace H_1 by H_2 . We assume that B^{H_1} and B^{H_2} are independent.

We define $C^\beta([T_1, T_2]; V)$ to be the Banach space of Hölder continuous functions on $[T_1, T_2]$ with exponent $0 < \beta < 1$ having values in V . A norm of this space is given by

$$\|u\|_\beta = \|u\|_{\beta, T_1, T_2} = \|u\|_{\infty, T_1, T_2} + \|u\|_{\beta, T_1, T_2}$$

with

$$\|u\|_\beta = \|u\|_{\beta, T_1, T_2} = \sup_{T_1 \leq s < t \leq T_2} \frac{\|u(t) - u(s)\|}{|t-s|^\beta}.$$

By Kolmogorov's theorem we know that B^{H_1} has a version $C^\beta([0, T]; V)$ where we assume that $\beta \in (1/2, H_1)$. B^{H_2} has a version so that on any interval $[T_1, T_2]$, $T_1 < T_2$ we have $B^{H_2}|_{[T_1, T_2]} \in C^{\gamma'}([T_1, T_2]; V)$ where $\gamma' < H_2$ and

$$(2.2) \quad H_1 > 1/2, \quad H_2 \in (1 - H_1, 1).$$

Let us consider canonical versions of B^{H_1} and B^{H_2} :

$$B^{H_1}(\omega_1) = \omega_1, \quad \omega_1 \in C_0([0, T]; V), \quad B^{H_2}(\omega_2) = \omega_2, \quad \omega_2 \in C_0(\mathbb{R}; V).$$

Taking into account that we have Hölder continuous versions for B^{H_1} and B^{H_2} we describe the canonical versions as follows: Let

$$\Omega_1 = C_0([0, T]; V) \cap C^\beta([0, T]; V), \quad \Omega_2 = C_0(\mathbb{R}; V) \cap C^{\gamma'}(\mathbb{R}; V)$$

Then the canonical processes are given by

$$(\Omega_1, \mathcal{B}(C_0([0, T]; V)) \cap \Omega_1, \mathbb{P}'_{H_1}), \quad (\Omega_2, \mathcal{B}(C_0(\mathbb{R}; V)) \cap \Omega_2, \mathbb{P}'_{H_2})$$

where \mathbb{P}'_{H_1} now stands for $\mathbb{P}_{H_1}(\cdot \cap \Omega_1)$ and similar for \mathbb{P}'_{H_2} .

We consider now the metric dynamical system (MDS)

$$(C(\mathbb{R}, V), \mathcal{B}(C(\mathbb{R}, V)), \mathbb{P}_{H_2}, \theta)$$

with the measurable flow $\theta = (\theta_t)_{t \in \mathbb{R}}$ given by the shift operators $\theta_t \omega_2(\cdot) = \omega_2(\cdot + t) - \omega_2(t)$, see Arnold [1, p. 546]. The measure \mathbb{P}_{H_2} is ergodic with respect to $(\theta_t)_{t \in \mathbb{R}}$. For details we refer for instance to [15]. Since Ω_2 is $(\theta_t)_{t \in \mathbb{R}}$ -invariant and has full measure that we can conclude $(\Omega_2, \mathcal{B}(C_0(\mathbb{R}, V)) \cap \Omega_2, \mathbb{P}'_{H_2}, \theta)$ is also ergodic.

Introduce the product measure $\mathbb{P} := \mathbb{P}_{H_1} \times \mathbb{P}_{H_2}$ on

$$\begin{aligned} & (C_0([0, T]; V) \times C_0(\mathbb{R}; V), \mathcal{B}(C_0([0, T]; V)) \otimes \mathcal{B}(C_0(\mathbb{R}; V)), \mathbb{P}) \\ & = (C_0([0, T]; V) \times C_0(\mathbb{R}; V), \mathcal{B}(C_0([0, T]; V) \times C_0(\mathbb{R}; V)), \mathbb{P}). \end{aligned}$$

We set $\Omega := \Omega_1 \times \Omega_2$ and

$$\mathcal{F} = \mathcal{B}(C_0([0, T]; V) \times C_0(\mathbb{R}; V)) \cap \Omega$$

being the trace σ -algebra w.r.t. Ω equipped with the probability measure $\mathbb{P}'(\cdot) = \mathbb{P}(\cdot \cap \Omega) = \mathbb{P}'_{H_1} \times \mathbb{P}'_{H_2}$. Let us denote for the following the measures \mathbb{P}' , \mathbb{P}'_{H_1} , \mathbb{P}'_{H_2} by \mathbb{P} , \mathbb{P}_{H_1} , \mathbb{P}_{H_2} . Then we can describe the Hölder continuous and canonical version of (B^{H_1}, B^{H_2}) by the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with paths $\omega = (\omega_1, \omega_2) \in \Omega$.

2.2. Pathwise stochastic integrals. Although this construction has already been done in the recent paper [4] (see also Maslowski et al. [28]), we present it here for the sake of completeness. We begin this subsection by introducing some function spaces. Let $C^{\beta, \sim}([T_1, T_2]; V) \subset C([T_1, T_2]; V)$ be the set of functions with the finite norm

$$\|u\|_{\beta, \sim} = \|u\|_{\beta, \sim, T_1, T_2} = \|u\|_{\infty, T_1, T_2} + \sup_{T_1 < s < t \leq T_2} (s - T_1)^\beta \frac{\|u(t) - u(s)\|}{|t - s|^\beta}.$$

For $\rho > 0$ we can consider the equivalent norm

$$\|u\|_{\beta, \rho, \sim} = \|u\|_{\beta, \rho, \sim, T_1, T_2} = \sup_{s \in [T_1, T_2]} e^{-\rho(s - T_1)} \|u(s)\|$$

$$+ \sup_{T_1 < s < t \leq T_2} (s - T_1)^\beta e^{-\rho(t-T_1)} \frac{\|u(t) - u(s)\|}{|t - s|^\beta}.$$

It is known that $C^{\beta, \sim}([T_1, T_2]; V)$ is a Banach space, see [4], Lunardi [27, p.123].

Now we want to define the stochastic integral with $\omega_1 \in \Omega_1$ as integrator. The definition that we use throughout this article is given by Zähle [41] generalized to infinite dimensional case in Chen et al. [4]. Consider the separable Hilbert space $L_2(V)$ of Hilbert Schmidt operators from V into V with the usual norm $\|\cdot\|_{L_2(V)}$ and inner product $(\cdot, \cdot)_{L_2(V)}$. A base $(E_{ji})_{j, i \in \mathbb{N}}$ in this space is given by

$$E_{ji}e_k = \begin{cases} 0 & i \neq k, \\ e_j, & i = k \end{cases}$$

where $(e_k)_{k \in \mathbb{N}}$ is a complete orthonormal system in V . Consider the mapping $\Psi : [0, T] \rightarrow L_2(V)$ and suppose that $\psi_{ji} := (\Psi(\cdot), E_{ji})_{L_2(V)} \in I_{T_1+}^\alpha(L^p((T_1, T_2); \mathbb{R}))$ and $\psi_{ji}(T_1+)$ (the right-side limit of ψ_{ji} at T_1) exists and $\alpha p < 1$. Moreover, assume that $\zeta_{iT_2-} := (\omega_{1, T_2-}(t), e_i)_V \in I_{T_2-}^{1-\alpha}(L^{p'}((T_1, T_2); \mathbb{R}))$ such that $\frac{1}{p} + \frac{1}{p'} \leq 1$, and the mapping

$$[T_1, T_2] \ni r \mapsto \|D_{T_1+}^\alpha \Psi[r]\|_{L_2(V)} \|D_{T_2-}^{1-\alpha} \omega_{1, T_2-}[r]\| \in L^1((T_1, T_2); \mathbb{R})$$

where

$$(2.3) \quad \begin{aligned} D_{T_1+}^\alpha \Psi[r] &= \frac{1}{\Gamma(1-\alpha)} \left(\frac{\Psi(r)}{(r-T_1)^\alpha} + \alpha \int_{T_1}^r \frac{\Psi(r) - \Psi(q)}{(r-q)^{1+\alpha}} dq \right), \\ D_{T_2-}^{1-\alpha} \omega_{1, T_2-}[r] &= \frac{(-1)^{1-\alpha}}{\Gamma(\alpha)} \left(\frac{\omega_1(r) - \omega_1(T_2-)}{(T_2-r)^{1-\alpha}} + (1-\alpha) \int_r^{T_2} \frac{\omega_1(r) - \omega_1(q)}{(q-r)^{2-\alpha}} dq \right) \end{aligned}$$

are Weyl fractional derivatives, being $\omega_{1, T_2-}(r) = \omega_1(r) - \omega_1(T_2-)$, with $\omega_1(T_2-)$ the left side limit of ω_1 at T_2 . For the definition of the space $I_{T_1+}^\alpha(L^p((T_1, T_2); \mathbb{R}))$ and $I_{T_2-}^\alpha(L^{p'}((T_1, T_2); \mathbb{R}))$ we refer to Samko et al. [36] and Zähle [41, p. 337].

We then introduce

$$(2.4) \quad \int_{T_1}^{T_2} \Psi d\omega_1 := (-1)^\alpha \int_{T_1}^{T_2} D_{T_1+}^\alpha \Psi[r] D_{T_2-}^{1-\alpha} \omega_{1, T_2-}[r] dr.$$

Due to Pettis' theorem and the separability of V , the integrand is weakly measurable and, hence, measurable, and

$$\begin{aligned} \left\| \int_{T_1}^{T_2} \Psi(r) d\omega_1(r) \right\| &= \left(\sum_{j=1}^{\infty} \left\| \sum_{i=1}^{\infty} \int_{T_1}^{T_2} D_{T_1+}^\alpha \psi_{ji}[r] D_{T_2-}^{1-\alpha} \zeta_{iT_2-}[r] dr \right\|^2 \right)^{\frac{1}{2}} \\ &\leq \int_{T_1}^{T_2} \|D_{T_1+}^\alpha \Psi[r]\|_{L_2(V)} \|D_{T_2-}^{1-\alpha} \omega_{1, T_2-}[r]\| dr. \end{aligned}$$

LEMMA 2.1. *Suppose that $\Psi \in C^\gamma([T_1, T_2]; L_2(V))$ and $\omega_1 \in \Omega_1$ such that $1 - \beta < \alpha < \gamma, 1/2 < \beta$. Then (2.4) is well defined and there exists a positive constant c such that*

$$\left\| \int_{T_1}^{T_2} \Psi(r) d\omega_1(r) \right\| \leq c \|\Psi\|_\gamma \|\omega_1\|_\beta (T_2 - T_1)^\beta.$$

3. Pathwise mild solution of stochastic evolution equations driven by two FBMs.

3.1. Ornstein-Uhlenbeck processes driven by the FBM. Let V be a separable Hilbert space and let $-B$ be a symmetric positive operator with compact inverse. Then B generates the strongly continuous analytic semigroup S_B and $-B$ is closed. The eigenelements of $-B$ generate a complete orthonormal system $(e_{B,i})_{i \in \mathbb{N}}$ with spectrum $0 < \lambda_{B,1} \leq \lambda_{B,2} \leq \dots$ where these eigenvalues have finite multiplicity and tend to infinity.

Consider $\omega_2 \in \Omega_2$. For simplicity we assume that this random process can be presented by the orthonormal system $(e_{B,i})_{i \in \mathbb{N}}$ generated by the linear operator B :

$$\omega_2(t) = \sum_{i=1}^{\infty} (q_{ii}^2)^{\frac{1}{2}} e_{B,i} \omega_2^i(t), \quad \sum_{i=1}^{\infty} q_{ii}^2 < \infty, \quad q_{ij}^2 = 0 \quad \text{for } i \neq j.$$

Then ω_2^i are twosided one dimensional standard FBM which are iid where q_{ij}^2 are the representations of Q_2 w.r.t. the base $(e_{B,i})_{i \in \mathbb{N}}$.

Let Ω_2^* be the set of $\omega_2 \in \Omega_2$ which are subexponentially growing:

$$\Omega_2^* = \bigcap_{m \in \mathbb{N}} \Omega_{2,m}^*, \quad \omega_2 \in \Omega_{2,m}^* \text{ iff } \lim_{t \rightarrow \pm\infty} \|\omega_2(t)\| e^{-\frac{1}{m}|t|} = 0.$$

This set is straightforwardly $(\theta_t)_{t \in \mathbb{R}}$ -invariant.

LEMMA 3.1. $\Omega_2^* \in \mathcal{F}_2$ has measure one.

Proof. Note that $\|\omega_2(t)\| e^{-\frac{1}{m}|t|} = 0$ if and only if

$$(3.1) \quad \lim_{n \rightarrow \infty} \sup_{s \in [n, n+1]} \|\omega_2(s)\| e^{-\frac{1}{m}|s|} = 0$$

for $t \rightarrow +\infty$ and similar for $t \rightarrow -\infty$. Indeed, if (3.1) does not hold, then there exists a subsequence $(n_i)_{i \in \mathbb{N}}$ and an $\epsilon > 0$ so that

$$\sup_{s \in [n_i, n_i+1]} \|\omega_2(s)\| e^{-\frac{1}{m}|s|} > \epsilon \quad \text{for all } i \in \mathbb{N}.$$

Hence there exists a sequence $(n_i)_{i \in \mathbb{N}} : [n_i, n_i + 1] \ni t_i \rightarrow \infty$ so that

$$\limsup_{i \rightarrow \infty} \|\omega_2(t_i)\| e^{-\frac{1}{m}|t_i|} \geq \epsilon.$$

The mappings

$$\Omega_2 \ni \omega_2 \mapsto \sup_{s \in [n, n+1]} \|\omega_2(s)\|, \quad \sup_{s \in [-n-1, -n]} \|\omega_2(s)\|$$

are $(\mathcal{F}, \mathcal{B}(\mathbb{R}^+))$ -measurable. Hence $\Omega_{2,m}^*$ and thus Ω_2^* is measurable.

For $t \rightarrow +\infty$ we have an asymptotically linearly bounded growths:

$$\|\omega_2(t)\| \leq \sum_{i=0}^{\lfloor t \rfloor} \sup_{s \in [0,1]} \|\theta_i \omega_2(s)\| \sim \mathbb{E} \sup_{s \in [0,1]} \|\theta_i \omega_2(s)\| \lfloor t \rfloor$$

with probability one by the ergodic theorem where the right hand side is finite, see Kunita [23, Theorem 1.4.1]. Similarly we can argue for $t \rightarrow -\infty$. For ω_2 from this set we have

$$\limsup_{t \rightarrow \pm\infty} \frac{\|\omega_2(t)\|}{|t|} < \infty.$$

Hence Ω_2^* contains a subset of measure one so that $\mathbb{P}_{H_2}(\Omega_2^*) = 1$. \square

We consider the equation

$$(3.2) \quad dZ(t) = BZ(t) dt + d\omega_2(t), \quad Z(0) = Z_0 \in V, \quad t \geq 0$$

interpreted in mild form.

Let π_p be the orthonormal projection with respect to $(e_{B,i})_{i=1,\dots,p}$. Then by Cheridito et al. [5, Proposition A.1] for some $Z_0 \in V$ we have a p -dimensional Ornstein Uhlenbeck process (O-U process) generated by the finite dimensional FBM $\pi_p\omega_2$:

$$Z_p(t, \omega) = S_B(t)\pi_p Z_0 + \pi_p\omega_2(t) + B \int_0^t S_B(t-r)\pi_p\omega_2(r)dr$$

Note that π_p commutes with B and $S_B(t)$. $\pi_p\omega(r)$ converges pointwise to $\omega_2(r)$ on any interval $[0, t]$ and since $\pi_p\omega_2$ has a uniform bounded γ' Hölder-norm on $[0, t]$ this convergence is uniform. On the other hand for $\gamma < \gamma'$ applying Maslowski and Nualart [28, (4.27)] we have the convergence of $(\pi_p\omega_2)_{p \in \mathbb{N}}$ to ω_2 with respect to the γ -Hölder norm. Then by the proof of Pazy [31, Lemma 4.3.4, Theorem 4.3.5 (iii)] or Lunardi [27, Theorem 4.3.1 (III)] we obtain that

$$\begin{aligned} & S_B(t)\pi_p Z_0 + \pi_p\omega_2(t) + B \int_0^t S_B(t-r)\omega_2(r)dr \\ & \xrightarrow{p \rightarrow \infty} S_B(t)Z_0 + \omega_2(t) + B \int_0^t S_B(t-r)\omega_2(r)dr \end{aligned}$$

which we consider to be the solution of (3.2) formally written as

$$S(t)Z_0 + \int_0^t S(t-r)d\omega_2(r).$$

This holds for every $t > 0$.

We replace now ω_2 by $\theta_{-t}\omega_2$:

$$(3.3) \quad \begin{aligned} & \theta_{-t}\omega_2(t) + B \int_0^t S_B(t-r)\theta_{-t}\omega_2(r)dr \\ & = B \int_0^t S_B(t-r)\omega_2(r-t)dr - S_B(t)\omega_2(-t) \\ & = B \int_{-t}^0 S_B(-r)\omega_2(r)dr - S_B(t)\omega_2(-t) \end{aligned}$$

by

$$B \int_0^t S_B(t-r)\omega_2(-t)dr = S_B(t)\omega_2(-t) - \omega_2(-t).$$

We show that the right hand side of (3.3) is uniformly bounded for $t > 1$. We have

$$(3.4) \quad \left\| B \int_{-t}^0 S_B(-r)\omega_2(r)dr \right\| \leq \left\| B \int_{-1}^0 S_B(-r)\omega_2(r)dr \right\| + \left\| B \int_{-t}^{-1} S_B(-r)\omega_2(r)dr \right\|.$$

In particular we can estimate the first term

$$\left\| B \int_{-1}^0 S_B(-r)\omega_2(r)dr \right\| \leq \left\| B \int_0^1 S_B(1-r)\theta_{-1}\omega_2(r)dr \right\| + \|S_B(1)\omega_2(-1)\| + \|\omega_2(-1)\| < \infty.$$

The right hand side is finite by Pazy [31, Theorem 4.3.5]. For the second norm we have that for $\omega_2 \in \Omega_2^*$ for any $\lambda_{B,1} > \lambda_B > 2\zeta > 0$ there exists a $C(\zeta, \omega_2)$ so that

$$\|\omega_2(t)\| \leq C(\zeta, \omega_2)e^{\zeta|t|} \quad \text{for } t \in \mathbb{R}.$$

Note that there is a constant C_ζ so that

$$\|BS_B(t)\| \leq C_\zeta \frac{1}{t} e^{-(\lambda_B - \zeta)t}, \quad \text{for } t > 0.$$

Thus the last norm in (3.4) is bounded for any $t > 1$ by

$$\frac{C_\zeta C(\zeta, \omega_2)}{\lambda_B - 2\zeta}.$$

Hence the random variable

$$\begin{aligned} Z(\omega_2) &= B \int_{-\infty}^0 S_B(-q)\omega_2(q) dq \\ &= \lim_{t \rightarrow \infty} \left(B \int_{-t}^0 S_B(-q)\omega_2(q) dq - S_B(t)\omega_2(-t) + S_B(t)Z_0 \right) \end{aligned}$$

is well defined.

Consider the mild form of (3.2) with initial time $r \in \mathbb{R}$ and $t > r$:

$$(3.5) \quad \begin{aligned} Z(t, \omega_2) &= S_B(t-r)Z_0 + \int_r^t S_B(t-q) d\omega_2(q) \\ &= S_B(t-r)Z_0 + \int_0^{t-r} S_B(t-r-q) d\theta_r \omega_2(q) \\ &= S_B(t-r)Z_0 + B \int_0^{t-r} S_B(t-r-q)\theta_r \omega_2(q) dq + \theta_r \omega_2(t-r). \end{aligned}$$

In particular for $Z_0 = Z(\theta_r \omega_2)$ we have

$$Z(\theta_t \omega_2) = Z(t, \omega_2)$$

is a stationary solution to (3.2). In addition by Pazy [31, Theorem 4.3.5 (iii)] the second and the third term on the right hand side of (3.5) have a finite γ -Hölder norm

with respect to $t \in [r, T]$. Then the right hand side has a finite γ -Hölder norm with respect to $t \in [r + \delta, T]$, $0 < \delta < T - r$ so that $Z(\theta_t \omega_2)$ has a finite γ -Hölder norm on any compact interval.

Let $\omega_{2,\epsilon}(\cdot)$ be the scaled function $\omega_2(\frac{\cdot}{\epsilon})$. Over the probability space $(\Omega_2, \mathcal{F}_2, \mathbb{P}_{H_2})$ this is an FBM which has the same distribution of $\frac{1}{\epsilon^{H_2}} \omega_2(\cdot)$. We consider the stationary mild solution of

$$(3.6) \quad dZ^\epsilon(t) = \frac{1}{\epsilon} B Z^\epsilon(t) dt + d\omega_{2,\epsilon}(t).$$

Similar to above this solution process is given by

$$Z^\epsilon(\theta_r \omega_2) = \frac{1}{\epsilon} B \int_{-\infty}^0 S_{\frac{B}{\epsilon}}(-q) \theta_r \omega_{2,\epsilon}(q) dq, \quad r \in \mathbb{R}$$

is a continuous random process which solves (3.6). We note that for any $\epsilon \in (0, 1]$ $\omega_{2,\epsilon} \in \Omega_2^*$ if and only if $\omega_2 \in \Omega_2^*$.

The following lemma describes the relation between Z and Z^ϵ .

LEMMA 3.2. *Let $\omega_{2,\epsilon}(\cdot) = \omega_2(\frac{\cdot}{\epsilon})$. Then we have*

$$Z(\theta_r \omega_2) = Z^\epsilon(\theta_r \omega_2) \quad r \in \mathbb{R}.$$

Proof. We have

$$\begin{aligned} Z(\theta_r \omega_2) &= B \int_{-\infty}^0 S_B(-q) \theta_r \omega_2(q) dq \\ &= B \int_{-\infty}^0 S_B(-q) (\omega_{2,\epsilon}(r + \epsilon q) - \omega_{2,\epsilon}(r)) dq \\ &= \frac{1}{\epsilon} B \int_{-\infty}^0 S_{\frac{B}{\epsilon}}(-q) (\omega_{2,\epsilon}(r + q) - \omega_{2,\epsilon}(r)) dq \\ &= \frac{1}{\epsilon} B \int_{-\infty}^0 S_{\frac{B}{\epsilon}}(-q) \theta_r \omega_{2,\epsilon}(q) dq = Z^\epsilon(\theta_r \omega_2). \end{aligned}$$

We note that $S_{\frac{B}{\epsilon}}(t) = S_B(\frac{t}{\epsilon})$ for $t \geq 0$. □

LEMMA 3.3. *We have:*

- (1) $\mathbb{E}[\sup_{s \in [0, T]} \|Z(\theta_s \omega_2)\|] < \infty$.
- (2) Let $T > 0$. We have for $\epsilon \rightarrow 0$ on a $(\theta_t)_{t \in \mathbb{R}}$ -invariant set of full measure.

$$\sup_{s \in [0, T]} \|Z^\epsilon(\theta_s \omega_2)\| = o(|\epsilon|^{-1}).$$

Proof. (1) To see that $\mathbb{E}[\sup_{[0, T]} \|Z(\theta_t \omega_2)\|] < \infty$ we have

$$\begin{aligned} \sup_{[0, T]} \|Z(\theta_t \omega_2)\| &\leq \sup_{t \in [0, T]} \|Z(\theta_t \omega_2) - Z(\omega_2)\| + \|Z(\omega_2)\| \\ &\leq \sup_{s < t \in [0, T]} \|Z(\theta_t \omega_2) - Z(\theta_s \omega_2)\| + \|Z(\omega_2)\| \end{aligned}$$

Then we have by Lunardi [27, Theorem 4.3.1 (III)]

$$\mathbb{E}[\sup_{s < t \in [0, T]} \|Z(\theta_t \omega_2) - Z(\theta_s \omega_2)\|] \leq C \mathbb{E}[\|\omega_2\|_\gamma] T^\gamma.$$

By Kunita [23, Theorem 1.4.1] we have a random variable $K(\omega_2)$ such that

$$\mathbb{E}[\|\omega_2\|_\gamma] \leq \mathbb{E}[K(\omega_2)] \leq (\mathbb{E}[K(\omega_2)^{2n}])^{1/(2n)} < \infty$$

when $2nH_2 > 1$. The finiteness of $\mathbb{E}[\|Z(\omega)\|]$ follows by taking the expectation of the right hand side of (3.3) having in mind $\mathbb{E}[\|\omega_2(t)\|^2] \leq (\sum_i q_{ii}^2 |t|^{2H_2})$.

(2) By (1) we can apply Arnold [1, Proposition 4.1.3] and know that $\|Z(\theta_q \omega_2)\|$ is sublinear growing i.e.

$$\lim_{q \rightarrow \pm\infty} \|Z(\theta_q \omega_2)\| \cdot |q|^{-1} = 0$$

on a $(\theta_t)_{t \in \mathbb{R}}$ invariant set of full measure. Suppose the assertion does not hold for ω_2 from the invariant set mentioned above. Then there exists a $\delta > 0$ and a sequence $(\epsilon_j)_{j \in \mathbb{N}}$, $\epsilon_j > 0$ tending to zero so that by Lemma 3.2

$$(3.7) \quad \sup_{s \in [0, T]} \|Z^{\epsilon_j}(\theta_s \omega_2)\| \epsilon_j = \sup_{s \in [0, T]} \|Z(\theta_{\frac{s}{\epsilon_j}} \omega_2)\| \epsilon_j > \delta$$

for every j . Let s_{ϵ_j} be the largest element in $[0, T/\epsilon_j]$ so that

$$\sup_{q \in [0, T/\epsilon_j]} \|Z(\theta_q \omega_2)\| = \|Z(\theta_{s_{\epsilon_j}} \omega_2)\|.$$

For $j \rightarrow \infty$ the sequence (s_{ϵ_j}) tends to ∞ . Hence

$$(3.8) \quad 0 = \lim_{j \rightarrow \infty} \|Z(\theta_{s_{\epsilon_j}} \omega_2)\| \cdot \frac{1}{s_{\epsilon_j}} \geq \lim_{j \rightarrow \infty} \|Z(\theta_{s_{\epsilon_j}} \omega_2)\| \cdot \frac{\epsilon_j}{T} = 0$$

which is a contradiction to (3.7). \square

3.2. Description of the problem of stochastic evolution equations. Let $u(t) = (u_1(t), u_2(t)) \in V \times V$ be the solution of

$$(3.9) \quad du(t) = Ju(t) dt + F(u(t)) dt + G(u(t)) (d\omega_1(t), d\omega_2(t))$$

where $u(0) = u_0 = (u_{01}, u_{02}) \in V \times V$, ω_1 is a path of the canonical FBM with Hurst exponent H_1 , and ω_2 is a path of the canonical FBM with Hurst exponent H_2 so that (2.2) holds. In contrast to the equation considered in Maslowski et al. [28] and Chen et al. [4], (3.9) contains the term ω_2 which does not have the Hölder regularity of ω_1 .

We describe the assumptions regarding the operator J and nonlinear terms F and G as follows

- (H1) Let $-J$ be a closed positive symmetric operator with compact inverse. Then, J generates an exponential analytic semigroup S_J on $V \times V$, such that $\|S_J(t)\| \leq e^{-\lambda_J t}$, $\lambda_J > 0$, for $t \geq 0$.
- (H2) The operator $F : V \times V \rightarrow V \times V$ is Lipschitz continuous with Lipschitz constant c_{DF} .
- (H3) The operator is given by

$$G = \begin{pmatrix} h(u_1) & 0 \\ 0 & \text{id} \end{pmatrix}$$

where $h : V \rightarrow L_2(V)$. The latter space is the space of Hilbert Schmidt operators on V . h has bounded first and second derivatives with bounds c_{Dh} and c_{D^2h} .

Let $-J$ from (H1). We can assume that $-J$ has a positive spectrum of finite multiplicity $(\lambda_{J,i})_{i \in \mathbb{N}}$ so that $\lim_{i \rightarrow \infty} \lambda_{J,i} = \infty$. The associated eigenelements $(e_{J,i})_{i \in \mathbb{N}}$ are chosen so that they form a complete orthonormal system.

For $\alpha \geq 0$ define the Banach spaces $\tilde{V}_\alpha = D((-J)^\alpha)$ where the norm of this space is given in the following definition

$$\tilde{V}_\alpha = \left\{ u \in \tilde{V} : \|u\|_\alpha^2 = \sum_{i=1}^{\infty} \tilde{\lambda}_i^{2\alpha} |\hat{u}_i|^2 < \infty, \quad u = \sum_{i=1}^{\infty} \hat{u}_i e_{J,i} \right\} \quad \text{with } \tilde{V} = \tilde{V}_0.$$

Here \tilde{V} stands for $V \times V$ which has the orthonormal basis $(e_{J,i})_{i \in \mathbb{N}} \in \tilde{V}$ coming from the eigenvalues of $-J$ related to the eigenvectors $\tilde{\lambda}_i = \lambda_{J,i}$. Then $\tilde{V}_1 = D(-J)$ is the domain of J . However, later considering the operators $-A, -B$ we set $\tilde{V} = V$ and the eigenvalues and eigenvectors $\lambda_{A,i}, e_{A,i}$ and $\lambda_{B,i}, e_{B,i}$ are given by the eigenvalues, eigenvectors of $-A, -B$. The properties of B has been used at the beginning of Section 3.1. The operators A, B generate an exponential semigroup S_A, S_B like with similar properties as J but defined on $\tilde{V} = V$. Let $L(\tilde{V}_v, \tilde{V}_\zeta)$ denote the space of continuous linear operators from \tilde{V}_v into \tilde{V}_ζ . There exists a constant $c > 0$, such that for $0 \leq s < t \leq T$, we have

$$(3.10) \quad \|S_J(t)\|_{L(\tilde{V}, \tilde{V}_\sigma)} \leq ct^{-\sigma} e^{-\lambda_J t}, \quad \text{for } \sigma > 0$$

$$(3.11) \quad \|S_J(t-s) - \text{id}\|_{L(\tilde{V}_\sigma, \tilde{V})} \leq c(t-s)^\sigma, \quad \text{for } \sigma \in [0, 1].$$

In (3.10) notice that λ_J is a positive constant. We also note that, for $0 < \sigma \leq 1$, there exists $c > 0$ such that for $0 \leq q \leq r \leq s \leq t$, we derive

$$(3.12) \quad \|S_J(t-r) - S_J(t-q)\|_{L(\tilde{V})} \leq c(r-q)^\sigma (t-r)^{-\sigma}$$

and for $\varrho, \nu \in (0, 1]$

$$(3.13) \quad \begin{aligned} & \|S_J(t-r) - S_J(s-r) - S_J(t-q) + S_J(s-q)\|_{L(\tilde{V})} \\ & \leq c(t-s)^\varrho (r-q)^\nu (s-r)^{-(\varrho+\nu)}. \end{aligned}$$

Remark 3.4. By (H3), it is easy to obtain the estimate (see for instance Maslowski et al. [28])

$$\begin{aligned} \|h(u)\|_{L_2(V)} & \leq c_h + c_{Dh} \|u\|, \\ \|h(u) - h(v)\|_{L_2(V)} & \leq c_{Dh} \|u - v\|, \\ \|h(u_1) - h(u_2) - h(v_1) + h(v_2)\|_{L_2(V)} & \leq c_{Dh} \|u_1 - v_1 - (u_2 - v_2)\| \\ & \quad + c_{D^2h} \|u_1 - u_2\| + (\|u_1 - v_1\| + \|u_2 - v_2\|) \end{aligned}$$

for all $u, v, u_i, v_i \in V, i = 1, 2$.

3.3. Pathwise mild solution. Let $t \rightarrow Z(\theta_t \omega_2)$ be the stationary O-U process defined in Section 3.1.

In the different formulas, c will denote a generic constant that may differ from line to line. Sometimes we will write c_T when we want to stress the dependence on T .

Let now

$$J = \begin{pmatrix} A & 0 \\ 0 & B \end{pmatrix}.$$

We interpret the equation (3.9) in mild form:

$$\begin{aligned}
(3.14) \quad u(t) &= S_J(t)u_0 + \int_0^t S_J(t-r)F(u(r))dr + \int_0^t S_J(t-r)G(u(r))d\omega(r) \\
&= \begin{pmatrix} S_A(t)u_{01} \\ S_B(t)(u_{02} - Z(\omega_2)) \end{pmatrix} + \begin{pmatrix} 0 \\ Z(\theta_t\omega_2) \end{pmatrix} + \int_0^t S_J(t-r)F(u(r))dr + \\
(3.15) \quad &+ \begin{pmatrix} \int_0^t S_A(t-r)h(u_1(r))d\omega_1(r) \\ 0 \end{pmatrix}.
\end{aligned}$$

To estimate the stochastic integral of the last expression in (3.14), we refer to [4, 28, 41] where a similar estimate is derived. $u \in C^{\gamma, \sim}([0, T], V \times V)$. In addition, from (2.2) it follows that we find α, β, γ so that we assume that $\beta > 1/2$, $0 < \alpha < \gamma < \beta$, $\beta + \alpha > 1$. The condition $\alpha < \gamma$ allows to define $D_{0+}^\alpha h(u_1(\cdot))[r]$ and $\alpha + \beta > 1$ allows to define $D_{T-}^{1-\alpha} \omega_{1,T}(\cdot)[r]$, see (2.3). Then we have

$$\int_0^t S_A(t-r)h(u_1(r))d\omega_1(r) = (-1)^\alpha \int_0^t D_{0+}^\alpha S_A(t-\cdot)h(u_1(\cdot))[r]D_{T-}^{1-\alpha} \omega_{1,T}(\cdot)[r]dr.$$

We firstly introduce the operator $u \mapsto \mathcal{T}(u, \omega_1, \omega_2, u_0)$ for fixed u_0 with domain $C^{\gamma, \sim}([0, T]; V \times V)$ and $\omega_1 \in C^{\beta, \sim}([0, T]; V)$, $\omega_2 \in C^{\gamma', \sim}([0, T]; V)$. This operator is defined by

$$\begin{aligned}
\mathcal{T}(u, \omega_1, \omega_2, u_0)[t] &:= \begin{pmatrix} S_A(t)u_{01} \\ S_B(t)(u_{02} - Z(\omega_2)) \end{pmatrix} + \begin{pmatrix} 0 \\ Z(\theta_t\omega_2) \end{pmatrix} \\
&+ \int_0^t S_J(t-r)F(u(r))dr + \begin{pmatrix} \int_0^t S_A(t-r)h(u_1(r))d\omega_1(r) \\ 0 \end{pmatrix}.
\end{aligned}$$

LEMMA 3.5. *Let (H1)-(H3) hold and for any $T > 0$ there exists a $c_T > 0$ such that under (2.2) we can ensure the above condition on $\alpha, \beta, \gamma < \gamma' < H_2$. For $\rho > 0, \omega_1 \in C^\beta([0, T], V)$, $\omega_2 \in C^{\gamma'}([0, T], V)$ and $u \in C^{\gamma, \sim}([0, T], V \times V)$, we have*

$$\begin{aligned}
(3.16) \quad &\|\mathcal{T}(u, \omega_1, \omega_2, u_0)\|_{\gamma, \rho, \sim} \\
&\leq c_T(\|u_{01}\| + \|u_{02}\| + \|Z(\theta_t\omega_2)\|_\gamma) + C(\rho, \omega_1, T)(1 + \|u\|_{\gamma, \rho, \sim})
\end{aligned}$$

where $C(\rho, \omega_1, T) > 0$ such that $\lim_{\rho \rightarrow \infty} C(\rho, \omega_1, T) = 0$.

Proof. Despite the fact that a quite similar result was proved in [4, Lemma 10] (but in that paper there is not any drift and ω_2 -term), for the sake of completeness we give the proof in Appendix 4.4. \square

LEMMA 3.6. *Let (H1)-(H3) and (2.2) hold and for any $T > 0$ there exists a $c_T > 0$ such that for $\rho > 0, \omega_1 \in C^\beta([0, T], V)$, $\omega_2 \in C^{\gamma'}([0, T], V)$ and $u^1, u^2 \in C^{\gamma, \sim}([0, T]; V \times V)$,*

$$\begin{aligned}
&\|\mathcal{T}(u^1, \omega_1, \omega_2, u_0) - \mathcal{T}(u^2, \omega_1, \omega_2, u_0)\|_{\gamma, \rho, \sim} \\
&\leq c_T(1 + \|u^1\|_{\gamma, \rho, \sim} + \|u^2\|_{\gamma, \rho, \sim})\tilde{K}(\rho)\|u^1 - u^2\|_{\gamma, \rho, \sim}
\end{aligned}$$

where $\lim_{\rho \rightarrow \infty} \tilde{K}(\rho) = 0$.

Proof. The proof follows by using the same techniques as in [4, Lemma 11], because the Z -term is cancelled. \square

LEMMA 3.7. *Let (H1)-(H3) and (2.2) hold and $u_0 \in V \times V$. Then, for every $T > 0$, (3.9) has a unique solution u in $C^{\gamma, \sim}([0, T]; V \times V)$.*

Proof. According to (3.16), for sufficiently large ρ the centered and closed ball in $C^{\gamma, \sim}([0, T]; V \times V)$, $\|\cdot\|_{\gamma, \rho, \sim}$ which is mapped by $\mathcal{T}(\cdot, \omega_1, \omega_2, u_0)$ into itself. Then, by Lemma 3.6 by Theorem 12 in [4], (3.9) has a unique solution u in $C^{\gamma, \sim}([0, T]; V \times V)$. \square

4. Almost Sure Averaging for Fast-Slow Evolution Equations.

4.1. Description of the averaging problem. In this subsection, our intention is to convert the original system (1.1)-(1.2) into reduced systems without fast component. Thus, we are interested in solving the following stochastic system

$$(4.1) \quad dX^\epsilon(t) = AX^\epsilon(t) dt + f(X^\epsilon(t), Y^\epsilon(t)) dt + h(X^\epsilon(t)) d\omega_1(t), X^\epsilon(0) = X_0 \in V,$$

$$(4.2) \quad dY^\epsilon(t) = \frac{1}{\epsilon}BY^\epsilon(t) dt + \frac{1}{\epsilon}g(X^\epsilon(t), Y^\epsilon(t)) dt + d\omega_{2, \epsilon}(t), Y^\epsilon(0) = Y_0 \in V$$

where ω_1 is a path of the canonical FBM with Hurst exponent H_1 , ω_2 is a path of the canonical FBM with Hurst exponent H_2 , and $H_1 \in (\frac{1}{2}, 1), H_2 \in (1 - H_1, 1)$, which have been given in Section 2. Then $\omega_{2, \epsilon}(\cdot) = \omega_2(\frac{\cdot}{\epsilon})$. By the solution of (4.1)-(4.2) on $[0, T]$, we mean a process (X^ϵ, Y^ϵ) which satisfies

$$(4.3) \quad \begin{aligned} X^\epsilon(t) &= S_A(t)X_0 + \int_0^t S_A(t-r)f(X^\epsilon(r), Y^\epsilon(r)) dr \\ &\quad + \int_0^t S_A(t-r)h(X^\epsilon(r)) d\omega_1(r) \end{aligned}$$

and

$$(4.4) \quad \begin{aligned} Y^\epsilon(t) &= S_{\frac{B}{\epsilon}}(t)Y_0 + \frac{1}{\epsilon} \int_0^t S_{\frac{B}{\epsilon}}(t-r)g(X^\epsilon(r), Y^\epsilon(r)) dr \\ &\quad + \int_0^t S_{\frac{B}{\epsilon}}(t-r) d\omega_{2, \epsilon}(r). \end{aligned}$$

We assume that the following conditions for the coefficients of the system are fulfilled.

(A1) We assume for the operators A, B the conditions of (H1).

(A2) The coefficient $F(x, y)$ from (H2) is now specialized by $(f(x, y), 1/\epsilon g(x, y))$. The coefficients $f(x, y) : V \times V \rightarrow V$ of (1.1) and $g(x, y) : V \times V \rightarrow V$ of (1.2) are globally Lipschitz continuous in x, y i.e., there exists a positive constant C_1 and let $C_2 = \|g(0, 0)\|$ such that

$$\begin{aligned} \|f(x_1, y_1) - f(x_2, y_2)\| + \|g(x_1, y_1) - g(x_2, y_2)\| &\leq C_1(\|x_1 - x_2\| + \|y_1 - y_2\|) \\ \|g(x, y)\| &\leq C_1(\|x\| + \|y\|) + C_2 \end{aligned}$$

for all $x_1, y_1, x_2, y_2, x, y \in V$.

(A3) h satisfies the conditions of (H3).

(A4) f is bounded.

LEMMA 4.1. *Let (A1)-(A3) and (2.2) hold. For any $X_0 \in V, Y_0 \in V$ and $T > 0$, there is a unique solution (X^ϵ, Y^ϵ) to (4.3)-(4.4) in $C^{\gamma, \sim}([0, T]; V \times V)$.*

Proof. This is just the special case of Lemma 3.7, we omit the proof here. \square

We present now the main result of this article.

THEOREM 4.2. *Let (A1)-(A4) and (2.2) hold and assume further that $\lambda_B > C_1$. For any $X_0 \in V$, as $\epsilon \rightarrow 0$ the solution of (4.1) converges to \bar{X} which solves following (4.5). That is, we have almost surely*

$$\lim_{\epsilon \rightarrow 0} \|X^\epsilon - \bar{X}\|_{\gamma, \sim} = 0$$

where \bar{X} is the mild solution to the averaged equation

$$(4.5) \quad \bar{X}(t) = S_A(t)X_0 + \int_0^t S_A(t-r)\bar{f}(\bar{X}(r))dr + \int_0^t S_A(t-r)h(\bar{X}(r))d\omega_1(r)$$

where the Lipschitz continuous function \bar{f} will be given in (4.12) later.

To prove Theorem 4.2, we first obtain the random fixed point for the RDS generated by (4.2) in Section 4.2. Then, we give the proof of Theorem 4.2 in Section 4.3.

4.2. Random fixed points. To describe the behavior of the fast variable we have to introduce a RDS. Let B be a separable Banach space, and $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}}, \theta)$ be an ergodic MDS. A measurable mapping

$$\varphi : \mathbb{R}^+ \times \hat{\Omega} \times B \rightarrow B$$

is called RDS if the cocycle property holds

$$\varphi(t + \tau, \omega, b) = \varphi(t, \theta_\tau \omega, \cdot) \circ \varphi(\tau, \omega, b) \quad \text{for } t, \tau \in \mathbb{R}^+, \omega \in \hat{\Omega}, b \in B.$$

and $\varphi(0, \omega, \cdot) = \text{id}_B$. For details we refer to Arnold [1].

Consider the parameterized equation

$$(4.6) \quad dY^{\epsilon, x}(t) = \frac{1}{\epsilon} B Y^{\epsilon, x}(t) dt + \frac{1}{\epsilon} g(x, Y^{\epsilon, x}(t)) dt + d\omega_{2, \epsilon}(t)$$

where $x \in V$ is a fixed but arbitrary element. Straightforwardly this equation interpreted in mild sense generates an RDS for every $x \in V$. We would like to show that under particular conditions on g the RDS generated by this equation has a random fixed point.

DEFINITION 4.3. *Let φ be an RDS over an ergodic metric dynamical system $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}}, \theta)$ with values in the separable Banach space B . A random variable $Y : \omega \rightarrow B$ is called random fixed point for φ if*

$$\varphi(t, \omega, Y(\omega)) = Y(\theta_t \omega)$$

for all $t \geq 0$ on a $(\theta_t)_{t \in \mathbb{R}}$ -invariant set of full measure.

In particular when Y is a random fixed point for an RDS generated by a differential equation then the function $t \rightarrow Y(\theta_t \omega)$ is a stationary solution of the equation (4.6).

We formulate conditions for the existence of a random fixed point. Recall that a random variable $X(\omega) \geq 0$ is called tempered on a $(\theta_t)_{t \in \mathbb{R}}$ -invariant set of full measure if

$$\lim_{t \rightarrow \pm\infty} \frac{\log^+ X(\theta_t \omega)}{|t|} = 0.$$

A family of sets $(C(\omega))_{\omega \in \hat{\Omega}}$, $C(\omega) \neq \emptyset$ and closed is called tempered random set if $\text{distance}_B(y, C(\omega))$ for all $y \in B$ is measurable, it is called tempered if

$$X(\omega) = \sup_{x \in C(\omega)} \|x\|_B$$

is tempered. We note that for every random set there exists a sequence of random variables $(x_n)_{n \in \mathbb{N}}$ so that

$$C(\omega) = \overline{\bigcup_{n \in \mathbb{N}} \{x_n(\omega)\}}.$$

Hence the above supremum defines a random variable. Random variables $y(\omega) \in C(\omega)$ for $\omega \in \hat{\Omega}$ are called selectors of C .

Let us present here an existence theorem random fixed points.

LEMMA 4.4. *Suppose that the RDS φ has a random forward invariant set closed C which is tempered:*

$$\varphi(t, \omega, C(\omega)) \subset C(\theta_t \omega) \quad \text{for } t \geq 0, \omega \in \hat{\Omega}.$$

Let

$$k(\omega) = \sup_{x \neq y \in C(\omega)} \log \left(\frac{\|\varphi(1, \omega, x) - \varphi(1, \omega, y)\|}{\|x - y\|} \right)$$

so that $\mathbb{E}k < 0$. The random variable

$$(4.7) \quad \omega \mapsto \sup_{t \in [0, 1]} \|\varphi(t, \theta_{-t} \omega, y(\theta_{-t} \omega))\|$$

is assumed to be tempered for any selector y from C . Then the RDS φ has a random fixed point $Y(\omega) \in C(\omega)$ which is unique. In addition $\|Y(\omega)\|$ is tempered. This random fixed point is pullback and forward attracting:

$$(4.8) \quad \lim_{t \rightarrow \infty} \|\varphi(t, \theta_{-t} \omega, y(\theta_{-t} \omega)) - Y(\omega)\| = 0, \quad \lim_{t \rightarrow \infty} \|\varphi(t, \omega, y(\omega)) - Y(\theta_t \omega)\| = 0$$

with exponential speed for every measurable selector $y(\omega) \in C(\omega)$.

For the definition and proof we refer to Chueshov and Schmalfuss [7, Definition 3.1.21, Theorem 4.2.1]. Let us check if the assumptions for our problem are satisfied.

THEOREM 4.5. *Consider (4.6) interpreted in the mild sense. Then for every $x \in V$ and $\epsilon > 0$ the RDS generated by (4.6) has a tempered pullback and forward exponentially attracting random fixed point $Y_F^\epsilon(\omega_2, x)$. The exponential rate of forward and pullback convergence to the random fixed point is given by $\frac{\lambda_B - C_1}{\epsilon}$.*

Proof. We set

$$(4.9) \quad \tilde{g}_\epsilon(x, y, \omega_2) = g(x, y + Z^\epsilon(\omega_2)) = g\left(x, y + \frac{1}{\epsilon} \int_{-\infty}^0 BS_{\frac{B}{\epsilon}}(-q)\omega_{2, \epsilon}(q) dq\right).$$

Then the equation

$$d\tilde{Y}^{\epsilon, x}(t) = \frac{1}{\epsilon} B\tilde{Y}^{\epsilon, x}(t) dt + \frac{1}{\epsilon} \tilde{g}_\epsilon(x, \tilde{Y}^{\epsilon, x}(t), \theta_t \omega_2) dt$$

has a unique mild solution forming the RDS $\tilde{\varphi}^{\epsilon, x}$. This RDS generates by conjugation an RDS for the mild version of (4.6). In particular we have

$$Y^{\epsilon, x}(t, \omega_2) - Z^\epsilon(\theta_t \omega_2) = \tilde{Y}^{\epsilon, x}(t, \omega_2).$$

We obtain by (A2)

$$\|\tilde{g}_\epsilon(x, y, \omega_2)\| \leq C_1\|y\| + \|\tilde{g}_\epsilon(x, 0, \omega_2)\|$$

where

$$\|\tilde{g}_\epsilon(x, 0, \omega_2)\| \leq C_1(\|Z^\epsilon(\omega_2)\| + \|x\|) + C_2.$$

Let $\tilde{y}^{\epsilon, x}$ be the solution of the one dimensional equation

$$\tilde{y}^{\epsilon, x}(t) = e^{-\frac{\lambda_B}{\epsilon}t}y_0 + \int_0^t e^{-\frac{\lambda_B}{\epsilon}(t-r)}\frac{1}{\epsilon}\left(C_1\tilde{y}^{\epsilon, x}(r) + C_1(\|Z^\epsilon(\omega_2)\| + \|x\|) + C_2\right)dr$$

which generates an one dimensional RDS with tempered forward and random fixed point $\tilde{y}_F^{\epsilon, x}(\theta_t\omega_2)$, see Chueshov and Schmalfuss [7, Theorem 3.1.23]. In addition, by a comparison argument

$$\|\tilde{Y}^{\epsilon, x}(t)\| \leq \tilde{y}^{\epsilon, x}(t) \quad \text{if } t \geq 0, y_0 \geq \|\tilde{Y}(0)\|.$$

Hence the ball with radius

$$(4.10) \quad \tilde{R}^{\epsilon, x}(\omega_2) = 2 \int_{-\infty}^0 e^{\frac{(\lambda_B - C_1)r}{\epsilon}} \frac{1}{\epsilon} (C_1(\|Z^\epsilon(\theta_r\omega_2)\| + \|x\|) + C_2) dr$$

and center 0 defines a tempered forward and pullback absorbing set $C^{\epsilon, x}(\omega_2)$. The temperedness follows by Lemma 3.3.

Consider

$$\begin{aligned} & \|\tilde{\varphi}^{\epsilon, x}(t, \omega_2, \tilde{Y}_{01}) - \tilde{\varphi}^{\epsilon, x}(t, \omega_2, \tilde{Y}_{02})\| \\ & \leq e^{-\frac{\lambda_B}{\epsilon}t} \|\tilde{Y}_{0,1} - \tilde{Y}_{0,2}\| + \int_0^t e^{-\frac{\lambda_B}{\epsilon}(t-r)} \frac{C_1}{\epsilon} \|\tilde{\varphi}^{\epsilon, x}(r, \omega_2, \tilde{Y}_{01}) - \tilde{\varphi}^{\epsilon, x}(r, \omega_2, \tilde{Y}_{02})\| dr. \end{aligned}$$

Then a Grönwall Lemma argument gives

$$\|\tilde{\varphi}^{\epsilon, x}(1, \omega_2, \tilde{Y}_{01}) - \tilde{\varphi}^{\epsilon, x}(1, \omega_2, \tilde{Y}_{02})\| \leq e^{-\frac{\lambda_B - C_1}{\epsilon}} \|\tilde{Y}_{01} - \tilde{Y}_{02}\|$$

so that the logarithm of the contraction constant is given by $k = \frac{1}{\epsilon}(-\lambda_B + C_1)$. Note that this constant k is nonrandom, less than 0 and the estimate is true for any pair $\tilde{Y}_{0i} \in V$, $i = 1, 2$. We have for some measurable selector y with $y(\omega_2) \in B(0, \tilde{R}^{\epsilon, x}(\omega_2))$

$$\begin{aligned} \|\tilde{\varphi}^{\epsilon, x}(q, \theta_{-t}\omega_2, y(\theta_{-t}\omega_2))\| & \leq e^{-\frac{\lambda_B}{\epsilon}q} \|y(\theta_{-t}\omega_2)\| \\ & \quad + \int_0^q e^{-\frac{\lambda_B}{\epsilon}(q-r)} \left(\frac{C_1}{\epsilon} \|\tilde{\varphi}^{\epsilon, x}(r, \theta_{-t}\omega_2, y(\theta_{-t}\omega_2))\| \right. \\ & \quad \left. + \frac{1}{\epsilon} (C_1(\|Z^\epsilon(\theta_{r-t}\omega_2)\| + \|x\|) + C_2) \right) dr. \end{aligned}$$

Then by a Grönwall Lemma argument for $t \in [0, 1]$

$$\begin{aligned} & \|\tilde{\varphi}^{\epsilon, x}(t, \theta_{-t}\omega_2, y(\theta_{-t}\omega_2))\| \\ & \leq \|y(\theta_{-t}\omega_2)\| + \int_0^t e^{-\frac{(\lambda_B - C_1)(t-r)}{\epsilon}} \frac{1}{\epsilon} (C_1(\|Z^\epsilon(\theta_{r-t}\omega_2)\| + \|x\|) + C_2) dr \\ & \leq \|y(\theta_{-t}\omega_2)\| + \int_{-t}^0 e^{\frac{(\lambda_B - C_1)r}{\epsilon}} \frac{1}{\epsilon} (C_1(\|Z^\epsilon(\theta_r\omega_2)\| + \|x\|) + C_2) dr \\ & \leq \|y(\theta_{-t}\omega_2)\| + \int_{-1}^0 e^{\frac{(\lambda_B - C_1)r}{\epsilon}} \frac{1}{\epsilon} (C_1(\|Z^\epsilon(\theta_r\omega_2)\| + \|x\|) + C_2) dr \\ & \leq \tilde{R}^{\epsilon, x}(\theta_{-t}\omega_2) + \int_{-1}^0 \frac{1}{\epsilon} (C_1(\|Z^\epsilon(\theta_r\omega_2)\| + \|x\|) + C_2) dr. \end{aligned}$$

However integrals and suprema w.r.t compact time intervals of tempered random variables are tempered, see Chueshov and Schmalfuss [7, Remark 3.1.8, p. 186] \square

THEOREM 4.6. *For fixed ω_2 , $\epsilon > 0$ this fixed point depends Lipschitz continuously on x with Lipschitz constant $\frac{C_1}{\lambda_B - C_1}$.*

Proof. We deal with the Lipschitz continuity of the fixed points $\tilde{Y}_F^\epsilon(\omega_2, x)$. We have for any $t \geq 0$

$$\begin{aligned} & \tilde{Y}_F^\epsilon(\omega_2, x_1) - \tilde{Y}_F^\epsilon(\omega_2, x_2) \\ &= \tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_1)) - \tilde{\varphi}^{\epsilon, x_2}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) \\ &= \tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_1)) - \tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) \\ &\quad + \tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) - \tilde{\varphi}^{\epsilon, x_2}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) \\ &=: K_1 + K_2. \end{aligned}$$

We set $t = n + t'$, $t' = t'(t) \in [0, 1)$. When $\|x_2\| \leq \|x_1\|$ then $\tilde{Y}_F^\epsilon(\omega_2, x_2) \in \tilde{C}^{\epsilon, x_1}(\omega_2)$. Then

$$\begin{aligned} \|K_1\| &= \|\tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_1)) - \tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2))\| \\ &\leq \sup_{y_1 \neq y_2 \in C^{\epsilon, x_1}(\theta_{-1}\omega_2)} \frac{\|\tilde{\varphi}^{\epsilon, x_1}(1, \theta_{-1}\omega_2, y_1) - \tilde{\varphi}^{\epsilon, x_1}(1, \theta_{-1}\omega_2, y_2)\|}{\|y_1 - y_2\|} \times \dots \times \\ &\quad \times \sup_{y_1 \neq y_2 \in C^{\epsilon, x_1}(\theta_{-n}\omega_2)} \frac{\|\tilde{\varphi}^{\epsilon, x_1}(1, \theta_{-n}\omega_2, y_1) - \tilde{\varphi}^{\epsilon, x_1}(1, \theta_{-n}\omega_2, y_2)\|}{\|y_1 - y_2\|} \\ &\quad \times \|\tilde{Y}_F^\epsilon(\theta_{-n-t'}\omega_2, x_1) - \tilde{Y}_F^\epsilon(\theta_{-n-t'}\omega_2, x_2)\| \\ &\leq e^{-n(\lambda_B - C_1)/\epsilon} \sup_{t' \in [0, 1)} \|\tilde{Y}_F^\epsilon(\theta_{-n-t'}\omega_2, x_1) - \tilde{Y}_F^\epsilon(\theta_{-n-t'}\omega_2, x_2)\|. \end{aligned}$$

Here, $\tilde{Y}_F^\epsilon(\omega_2, x_1), \tilde{Y}_F^\epsilon(\omega_2, x_2) \in C^{\epsilon, x_1}(\omega_2)$ are tempered, see (4.7), hence

$$\sup_{s \in [0, 1]} \|Y_F^\epsilon(\theta_{-s}\omega_2, x_1)\|, \sup_{s \in [0, 1]} \|Y_F^\epsilon(\theta_{-s}\omega_2, x_2)\|$$

is tempered, see Chueshov and Schmalfuss [7, Remark 3.1.8, p. 186]. Thus the right hand side can be made arbitrarily small when n is sufficiently large. We then have that K_1 is zero for $t \rightarrow \infty$. We deal with K_2 .

$$\begin{aligned} \|K_2\| &= \|\tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) - \tilde{\varphi}^{\epsilon, x_2}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2))\| \\ &\leq \int_0^t e^{-\frac{\lambda_B(t-r)}{\epsilon}} \frac{C_1}{\epsilon} (\|\tilde{\varphi}^{\epsilon, x_1}(r, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) \\ &\quad - \tilde{\varphi}^{\epsilon, x_2}(r, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2))\| + \|x_1 - x_2\|) dr. \end{aligned}$$

Then by the Grönwall lemma

$$\begin{aligned} & \|\tilde{\varphi}^{\epsilon, x_1}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2)) - \tilde{\varphi}^{\epsilon, x_2}(t, \theta_{-t}\omega_2, \tilde{Y}_F^\epsilon(\theta_{-t}\omega_2, x_2))\| \\ &\leq \int_0^t e^{-\frac{(\lambda_B - C_1)(t-r)}{\epsilon}} \frac{C_1}{\epsilon} \|x_1 - x_2\| dr \\ &\leq \frac{C_1}{\lambda_B - C_1} \|x_1 - x_2\| \end{aligned}$$

so that the Lipschitz constant is independent of ϵ and ω_2 . Finally we have

$$\|\tilde{Y}_F^\epsilon(\omega_2, x_1) - \tilde{Y}_F^\epsilon(\omega_2, x_2)\| \leq \frac{C_1}{\lambda_B - C_1} \|x_1 - x_2\|. \quad \square$$

LEMMA 4.7. *We have*

$$Y_F^\epsilon(\theta_r \omega_2, x) = Y_F^1(\theta_{\frac{r}{\epsilon}} \omega_2, x)$$

Proof. We prove the equality of $\tilde{Y}_F^\epsilon(\theta_r \omega_2, x)$, $\tilde{Y}_F^1(\theta_{\frac{r}{\epsilon}} \omega_2, x)$ for any $\epsilon > 0$. Then by Lemma 3.2, $Z(\theta_{1/\epsilon} \omega_2)$ and $Z^\epsilon(\theta_r \omega_2)$ are equal. This causes for $r' = \frac{r}{\epsilon}$

$$\begin{aligned} \tilde{Y}_F^\epsilon(\omega_2, x) &= \int_{-\infty}^0 S_B(-r) \frac{1}{\epsilon} \tilde{g}_\epsilon(x, \tilde{Y}_F^\epsilon(\theta_r \omega_2, x), \theta_r \omega) dr \\ &= \int_{-\infty}^0 S_B(-\frac{r}{\epsilon}) \frac{1}{\epsilon} g(x, \tilde{Y}_F^\epsilon(\theta_r \omega_2, x) + Z^\epsilon(\theta_r \omega_2)) dr \\ &= \int_{-\infty}^0 S_B(-\frac{r}{\epsilon}) \frac{1}{\epsilon} g(x, \tilde{Y}_F^\epsilon(\theta_r \omega_2, x) + Z(\theta_{\frac{r}{\epsilon}} \omega_2)) dr \\ &= \int_{-\infty}^0 S_B(-r') g(x, \tilde{Y}_F^\epsilon(\theta_{r'} \omega_2, x) + Z(\theta_{r'} \omega_2)) dr' \\ &= \int_{-\infty}^0 S_B(-r') \tilde{g}_1(x, \tilde{Y}_F^\epsilon(\theta_{r'} \omega_2, x), \theta_{r'} \omega_2) dr'. \end{aligned}$$

On the other hand we have by the uniqueness of the fixed point

$$\tilde{Y}_F^1(\omega_2, x) = \int_{-\infty}^0 S_B(-r') \tilde{g}_1(x, \tilde{Y}_F^1(\theta_{r'} \omega_2, x), \theta_{r'} \omega_2) dr'.$$

Note that a random fixed point can be presented by such an integral over an infinite domain. This follows by

$$\tilde{Y}_F^1(\omega_2, x) = S_B(t) \tilde{Y}_F^1(\theta_{-t} \omega_2, x) + \int_{-t}^0 S_B(-r') \tilde{g}_1(x, \tilde{Y}_F^1(\theta_{r'} \omega_2, x), \theta_{r'} \omega_2) dr'.$$

The temperedness of $\|\tilde{Y}_F^1(\omega_2, x)\|$ and the exponential decay of S_B ensure that the right hand side converges to the integral over the infinite domain.

We have

$$\tilde{g}_\epsilon(x, y, \theta_r \omega_2) = g(x, y + Z^\epsilon(\theta_r \omega_2))$$

and hence

$$\varphi^{\epsilon, x}(t, \omega_2, y) = Z^\epsilon(\theta_t \omega_2) + \tilde{\varphi}^{\epsilon, x}(t, \omega_2, y - Z^\epsilon(\omega_2)).$$

$\varphi^{\epsilon, x}(t, \omega_2, \cdot)$ is the RDS version of the solution to (4.6). There exists a random fixed point

$$(4.11) \quad Y_F^\epsilon(\omega_2, x) = \tilde{Y}_F^\epsilon(\omega_2, x) + Z^\epsilon(\omega_2).$$

The tempered set $\tilde{C}^{\epsilon, x}(\omega_2)$ is changed to

$$C^{\epsilon, x}(\omega_2) = \tilde{C}^{\epsilon, x}(\omega_2) + Z^\epsilon(\omega_2)$$

which is a ball with center $Z^\epsilon(\omega_2)$ and radius $\tilde{R}^{\epsilon, x}(\omega_2)$. \square

LEMMA 4.8. *The mapping $r \rightarrow Y_F^\epsilon(\theta_r \omega_2, x)$ is γ -Hölder continuous for $\gamma < H_2$ and $x \in V$.*

Proof. Let $r > r' \in \mathbb{R}$. We consider

$$\begin{aligned} \|\tilde{Y}_F^\epsilon(\theta_r \omega_2, x) - \tilde{Y}_F^\epsilon(\theta_{r'} \omega_2, x)\| &\leq \|S_{\frac{\epsilon}{2}}(r - r') - \text{id}\| \tilde{Y}_F^\epsilon(\theta_{r'} \omega_2, x) \| \\ &\quad + \left\| \int_{r'}^r S_{\frac{\epsilon}{2}}(r - s) \frac{1}{\epsilon} \tilde{g}_\epsilon(x, \tilde{Y}_F^\epsilon(\theta_s \omega_2, x), \theta_s \omega) ds \right\|. \end{aligned}$$

Since $s \mapsto \tilde{Y}_F^\epsilon(\theta_s \omega_2, x)$ and $s \mapsto Z^\epsilon(\theta_s \omega_2)$ are continuous the Lebesgue integral can be estimated by $C_{\epsilon, T} |r - r'|$ and by (3.10), (3.11) the first term on the right hand side of the last inequality causes Hölder continuity on any interval $r \in [r' + \delta, r' + T]$, $0 < \delta < T$ and $x \in V$.

Then by the transformation (4.11) and by Lemma 3.3 again, $r \mapsto Y_F^\epsilon(\theta_r \omega_2, x)$ is γ -Hölder continuous. \square

Remark 4.9. For the random variable $\tilde{R}^{\epsilon, x}$ introduced in (4.10) we can prove that $\sup_{r \in [0, T]} \tilde{R}^{\epsilon, x}(\theta_r \omega_2)$ is defined on $(\theta_t)_{t \in \mathbb{R}}$ invariant set of full measure independent of $x \in V$ and $\epsilon > 0$. This random variable is tempered.

4.3. An ergodic theorem for separable Hilbert-spaces. In this subsection we formulate an ergodic theorem in separable Hilbert spaces. We assume that f is bounded. Define

$$(4.12) \quad \bar{f}(x) = \mathbb{E}[f(x, Y_F^1(\omega_2, x))]$$

where this Hilbert-space valued expectation is determined by $\mathbb{E}[(f(x, Y_F^1(\omega_2, x)), y)]$ for all y in a dense countable set of V .

LEMMA 4.10. \bar{f} is Lipschitz continuous.

Proof. Because Y_F^ϵ depends Lipschitz continuously on x with Lipschitz constant $\frac{C_1}{\lambda_B - C_1}$, for all $x_1, x_2 \in V$, we have

$$\begin{aligned} \|\bar{f}(x_1) - \bar{f}(x_2)\| &\leq \mathbb{E}[\|f(x_1, Y_F^1(\omega_2, x_1)) - f(x_2, Y_F^1(\omega_2, x_2))\|] \\ &\leq C_1 \mathbb{E}[\|x_1 - x_2\| + \|Y_F^1(\omega_2, x_1) - Y_F^1(\omega_2, x_2)\|] \\ &\leq (C_1 + \frac{C_1^2}{\lambda_B - C_1}) \|x_1 - x_2\| =: C' \|x_1 - x_2\|. \end{aligned}$$

Thus, the desired result is obtained. \square

LEMMA 4.11. Let $\nu > 0$. There exists a $(\theta_t)_{t \in \mathbb{R}}$ invariant set in Ω_2 of full measure we have for ω_2 from this set and for $x \in V$

$$\lim_{T \rightarrow \pm\infty} \frac{1}{|T|} \left\| \int_0^T (-A)^{-\nu} (f(x, Y_F^1(\theta_r \omega_2, x)) - \bar{f}(x)) dr \right\| = 0.$$

Proof. Consider the operator $(-A)^{-\nu}$ which is a compact operator on V . Let $\Omega_{2, x}$ be the $(\theta_t)_{t \in \mathbb{R}}$ -invariant set so that for $\omega_2 \in \Omega_{2, x}$

$$\lim_{T \rightarrow \pm\infty} \frac{1}{|T|} \left\| \int_0^T (-A)^{-\nu} (f(x, Y_F^1(\theta_r \omega_2, x)) - \bar{f}(x)) dr \right\| = 0$$

which follows for every $x \in V$ by Chueshov at al. [6, Section 2] and for the invariance assertion see Arnold [1, Appendix 1]. For a dense and countable set $D \subset V$ the set

$$\bigcap_{x \in D} \Omega_{2, x} =: \tilde{\Omega}_2$$

is $(\theta_t)_{t \in \mathbb{R}}$ invariant and has full measure. Let $x \notin D$ and $(x_n)_{n \in \mathbb{N}}$ be a sequence in D so that

$$\lim_{n \rightarrow \infty} \|x - x_n\| = 0.$$

By the Lipschitz continuity and the uniform Lipschitz constant of $x \mapsto Y_F^1(\omega_2, x)$ with respect to ω_2 and $\epsilon > 0$ we have by Lemma 4.10 and Theorem 4.5 for any $\zeta > 0$ and $\tilde{n} \in \mathbb{N}$ so that

$$\begin{aligned} & \|(-A)^{-\nu}((f(x_{\tilde{n}}, Y_F^1(\theta_r \omega_2, x_{\tilde{n}})) - \bar{f}(x_{\tilde{n}})) - (f(x_{\tilde{n}}, Y_F^1(\theta_r \omega_2, x)) - \bar{f}(x)))\| \\ & \leq 2C' \|(-A)^{-\nu}\| \|x - x_{\tilde{n}}\| \leq \frac{\zeta}{2}. \end{aligned}$$

On the other hand we choose an $T_0 = T_0(\omega_2, \zeta) > 0$ so that for all $|T| > T_0$ we have that

$$\lim_{T \rightarrow \pm\infty} \frac{1}{|T|} \left\| \int_0^T (-A)^{-\nu} (f(x_{\tilde{n}}, Y_F^1(\theta_r \omega_2, x_{\tilde{n}})) - \bar{f}(x_{\tilde{n}})) dr \right\| \leq \frac{\zeta}{2}$$

on $\tilde{\Omega}_2$ [6]. Hence

$$\begin{aligned} & \frac{1}{|T|} \left\| \int_0^T (-A)^{-\nu} (f(x, Y_F^1(\theta_r \omega_2, x)) - \bar{f}(x)) dr \right\| \\ & \leq \frac{1}{|T|} \left\| \int_0^T (-A)^{-\nu} (f(x, Y_F^1(\theta_r \omega_2, x_{\tilde{n}})) - \bar{f}(x_{\tilde{n}})) dr \right\| \\ & \quad + 2 \|(-A)^{-\nu}\| C'_1 \|x - x_{\tilde{n}}\| \leq \zeta \end{aligned}$$

for $|T| > T_0$ and for $\omega_2 \in \tilde{\Omega}_2$. \square

4.4. Proof of Theorem 4.2. Following the discretization techniques inspired by Khasminskii in [20], we divide $[0, T]$ into intervals of size δ , where $\delta \in (0, 1)$ is a fixed number. Then, we construct an auxiliary process \hat{Y}^ϵ with initial value $\hat{Y}^\epsilon(0) = Y^\epsilon(0) = Y_0$, and for $t \in [k\delta, \min\{(k+1)\delta, T\})$,

$$\begin{aligned} \hat{Y}^\epsilon(t) &= S_{\frac{B}{\epsilon}}(t - k\delta) \hat{Y}^\epsilon(k\delta) + \frac{1}{\epsilon} \int_{k\delta}^t S_{\frac{B}{\epsilon}}(t - s) g(X^\epsilon(k\delta), \hat{Y}^\epsilon(s)) ds \\ & \quad + \int_{k\delta}^t S_{\frac{B}{\epsilon}}(t - s) d\omega_{2,\epsilon}(s) \end{aligned} \tag{4.13}$$

i.e.

$$\begin{aligned} \hat{Y}^\epsilon(t) &= S_{\frac{B}{\epsilon}}(t) Y_0 + \frac{1}{\epsilon} \int_0^t S_{\frac{B}{\epsilon}}(t - s) g(X^\epsilon(s_\delta), \hat{Y}^\epsilon(s)) ds \\ & \quad + \int_0^t S_{\frac{B}{\epsilon}}(t - s) d\omega_{2,\epsilon}(s) \end{aligned} \tag{4.14}$$

where $s_\delta = \lfloor s/\delta \rfloor \delta$ is the nearest breakpoint preceding s . Also, we define the process \hat{X}^ϵ , by

$$\begin{aligned} \hat{X}^\epsilon(t) &= S_A(t) X_0 + \int_0^t S_A(t - r) f(X^\epsilon(r_\delta), \hat{Y}^\epsilon(r)) ds \\ & \quad + \int_0^t S_A(t - r) h(X^\epsilon(r)) d\omega_1(r). \end{aligned} \tag{4.15}$$

LEMMA 4.12. *Assume (A1)-(A4). Then, for all $T > 0$, we have for large ρ have*

$$\|X^\epsilon\|_{\gamma,\rho,\sim} \leq C(\|X_0\| + 1)$$

where $C > 0$ is a constant which is independent of ϵ .

Proof. The proof of this result can be done by a slight generalization of [4, Lemma 9]. It is easy to see

$$\|X^\epsilon\|_{\gamma,\rho,\sim} \leq c_T \|X_0\| + c_T K(\rho) \|\omega_1\|_\beta (1 + \|X^\epsilon\|_{\gamma,\rho,\sim})$$

then, taking ρ big enough such that $c_T K(\rho) \|\omega_1\|_\beta < \frac{1}{2}$, we have

$$\|X^\epsilon\|_{\gamma,\rho,\sim} \leq 2c_T \|X_0\| + 1.$$

Here $K(\rho)$ is a positive function tending to zero for $\rho \rightarrow \infty$. □

Remark 4.13. Due to the boundedness of f , Y^ϵ does not have any effect on the estimate for $\|X^\epsilon\|_{\gamma,\rho,\sim}$.

Remark 4.14. Assume (A1)-(A4). Then, for all $T > 0$, we have

$$\|\hat{X}^\epsilon\|_{\gamma,\rho,\sim} + \|\bar{X}\|_{\gamma,\rho,\sim} \leq C(\|X_0\| + 1)$$

where $C > 0$ is a constant which is independent of ϵ . We obtain by the same method an similar estimate for $\|\hat{X}^\epsilon\|_{\gamma,\rho,\sim}$.

LEMMA 4.15. *For any solution Y^ϵ of (4.2) and any solution \hat{Y}^ϵ of (4.14), $t \in [0, T]$, we have*

$$\|Y^\epsilon\|_\infty + \|\hat{Y}^\epsilon\|_\infty \leq C(1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1}))$$

where C is a constant which is independent of ϵ .

Proof. For $t \in [0, T]$, from (4.2), one has

$$\begin{aligned} Y^\epsilon(t) &= S_{\frac{B}{\epsilon}}(t)Y_0 + \frac{1}{\epsilon} \int_0^t S_{\frac{B}{\epsilon}}(t-r)g(X^\epsilon(r), Y^\epsilon(r)) dr + \int_0^t S_{\frac{B}{\epsilon}}(t-r) d\omega_{2,\epsilon}(r) \\ &= S_{\frac{B}{\epsilon}}(t)(Y_0 - Z^\epsilon(\omega_2)) + Z^\epsilon(\theta_t \omega_2) + \frac{1}{\epsilon} \int_0^t S_{\frac{B}{\epsilon}}(t-r)g(X^\epsilon(r), Y^\epsilon(r)) dr. \end{aligned}$$

Then, we have

$$\begin{aligned} \|Y^\epsilon(t)\| &\leq \|S_{\frac{B}{\epsilon}}(t)\| \|Y_0 - Z^\epsilon(\omega_2)\| + \|Z^\epsilon(\theta_t \omega_2)\| \\ &\quad + \left\| \frac{1}{\epsilon} \int_0^t S_{\frac{B}{\epsilon}}(t-r)g(X^\epsilon(r), Y^\epsilon(r)) dr \right\| \\ &\leq e^{-\frac{\lambda B t}{\epsilon}} \|Y_0 - Z^\epsilon(\omega_2)\| + \|Z^\epsilon(\theta_t \omega_2)\| \\ &\quad + \frac{1}{\epsilon} \int_0^t e^{-\frac{\lambda B}{\epsilon}(t-r)} (C_2 + C_1(\|X^\epsilon(r)\| + \|Y^\epsilon(r)\|)) dr. \end{aligned}$$

By Lemma 4.12 and (A2), it is easy to know

$$\begin{aligned} \sup_{t \in [0, T]} \|Y^\epsilon(t)\| &\leq \|Y_0\| + 2 \sup_{t \in [0, T]} \|Z^\epsilon(\theta_t \omega_2)\| \\ &\quad + \sup_{t \in [0, T]} \frac{1}{\epsilon} \int_0^t e^{-\frac{\lambda B}{\epsilon}(t-r)} (C_2 + C_1 \|X^\epsilon(r)\|) ds \end{aligned}$$

$$\begin{aligned}
& + \sup_{t \in [0, T]} \frac{C_1}{\epsilon} \int_0^t e^{-\frac{\lambda_B}{\epsilon}(t-r)} \|Y^\epsilon(r)\| ds \\
& \leq C(1 + \|X_0\| + \|Y_0\| + \sup_{t \in [0, T]} \|Z^\epsilon(\theta_t \omega_2)\|) + \frac{C_1}{\lambda_B} \sup_{t \in [0, T]} \|Y^\epsilon(r)\|.
\end{aligned}$$

Then, by $\lambda_B > C_1$ and Lemma 3.3, we have

$$\begin{aligned}
\sup_{t \in [0, T]} \|Y^\epsilon(t)\| & \leq C(1 + \|X_0\| + \|Y_0\| + \sup_{t \in [0, T]} \|Z^\epsilon(\theta_t \omega_2)\|) \\
& \leq C(1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})).
\end{aligned}$$

Indeed we have for $\epsilon \rightarrow 0$

$$\sup_{t \in [0, T]} \|Z^\epsilon(\theta_t \omega_2)\| = \sup_{t \in [0, T]} \|Z(\theta_{\frac{t}{\epsilon}} \omega_2)\| = o(\epsilon^{-1})$$

by Lemma 3.3. The estimate for $\|\hat{Y}^\epsilon\|_\infty$ can be obtained in a similar way. \square

LEMMA 4.16. *For the stationary solution and any solution of (4.13), for $t \in [k\delta, \min\{(k+1)\delta, T\}]$, we have*

$$\int_{k\delta}^{(k+1)\delta} \|\hat{Y}^\epsilon(t) - Y_F^\epsilon(\theta_t \omega_2, X^\epsilon(k\delta))\| dt \leq C(1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1}))\epsilon$$

where C is a constant independent of ϵ and δ .

Proof. By the Grönwall lemma argument, we have

$$\|\hat{Y}^\epsilon(t) - Y_F^\epsilon(\theta_t \omega_2, X^\epsilon(k\delta))\| \leq e^{-\frac{\lambda_B - C_1}{\epsilon}(t-k\delta)} \|\hat{Y}^\epsilon(k\delta) - Y_F^\epsilon(\theta_{k\delta} \omega_2, X^\epsilon(k\delta))\|$$

Integrate above inequality from $k\delta$ to $(k+1)\delta$, due to $\lambda_B > C_1$, we have

$$\begin{aligned}
& \int_{k\delta}^{(k+1)\delta} \|\hat{Y}^\epsilon(t) - Y_F^\epsilon(\theta_t \omega_2, X^\epsilon(k\delta))\| dt \\
& \leq C \|\hat{Y}^\epsilon(k\delta) - Y_F^\epsilon(\theta_{k\delta} \omega_2, X^\epsilon(k\delta))\| \frac{\epsilon}{\lambda_B - C_1} (1 - e^{-\frac{(\lambda_B - C_1)\delta}{\epsilon}}) \\
& \leq C(\|\hat{Y}^\epsilon(k\delta)\| + \frac{C_1}{\lambda_B - C_1} \|X^\epsilon(k\delta)\| + \|Y_F^\epsilon(\theta_{k\delta} \omega_2, 0)\|) \frac{\epsilon}{\lambda_B - C_1} \\
& \leq C \sup_{r \in [0, T]} (\|X^\epsilon(r)\| + \|Y^\epsilon(r)\| + R^{\epsilon, 0}(\theta_r \omega_2) + \|Z^\epsilon(\theta_r \omega_2)\|) \frac{\epsilon}{\lambda_B - C_1}.
\end{aligned}$$

Thus, by Lemmas 3.3, 4.12 and 4.15, the desired result is obtained. \square

LEMMA 4.17. *For the solution \hat{Y}^ϵ of (4.13) and the solution Y^ϵ of (4.2), $s \in [k\delta, \min\{(k+1)\delta, T\}]$, $s < t \leq T$, $\rho > 1$, $k \geq 1$, ϵ small enough, we have*

$$e^{-\rho t} \int_{k\delta}^{(k+1)\delta} \|Y^\epsilon(s) - \hat{Y}^\epsilon(s)\| ds \leq C\delta^{1+\gamma}(1 + (k\delta)^{-\gamma})$$

where C is a constant which is independent of ϵ and δ .

Proof. For $s \in [k\delta, \min\{(k+1)\delta, T\}]$, by Lemma 4.15, one has

$$\|Y^\epsilon(s) - \hat{Y}^\epsilon(s)\| \leq e^{-\frac{\lambda_B}{\epsilon}(s-k\delta)} \|Y^\epsilon(k\delta) - \hat{Y}^\epsilon(k\delta)\|$$

$$\begin{aligned}
& \left\| \frac{1}{\epsilon} \int_{k\delta}^s S_{\frac{B}{\epsilon}}(s-r) (g(X^\epsilon(r), Y^\epsilon(r)) - g(X^\epsilon(r_\delta), \hat{Y}^\epsilon(r))) dr \right\| \\
& \leq C e^{-\frac{\lambda_B}{\epsilon}(s-k\delta)} (\|Y^\epsilon\|_\infty + \|\hat{Y}^\epsilon\|_\infty) \\
& \quad + \frac{C_1}{\epsilon} \int_{k\delta}^s e^{-\frac{\lambda_B}{\epsilon}(s-r)} \|X^\epsilon(r) - X^\epsilon(r_\delta)\| dr \\
& \quad + \frac{C_1}{\epsilon} \int_{k\delta}^s e^{-\frac{\lambda_B}{\epsilon}(s-r)} \|Y^\epsilon(r) - \hat{Y}^\epsilon(r)\| dr.
\end{aligned}$$

Then, multiplying both sides of the above equation by $e^{\frac{\lambda_B}{\epsilon}s}$, we have

$$\begin{aligned}
e^{\frac{\lambda_B}{\epsilon}s} \|Y^\epsilon(s) - \hat{Y}^\epsilon(s)\| & \leq C e^{\frac{\lambda_B}{\epsilon}k\delta} (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) \\
& \quad + \frac{C_1}{\epsilon} \int_{k\delta}^s e^{\frac{\lambda_B}{\epsilon}r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\| dr \\
& \quad + \frac{C_1}{\epsilon} \int_{k\delta}^s e^{\frac{\lambda_B}{\epsilon}r} \|Y^\epsilon(r) - \hat{Y}^\epsilon(r)\| dr.
\end{aligned}$$

By the Grönwall inequality [8, p.37] and [9, p.13], we have

$$\begin{aligned}
\|Y^\epsilon(s) - \hat{Y}^\epsilon(s)\| & \leq C e^{-\frac{\lambda_B}{\epsilon}(s-k\delta)} (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) e^{\frac{C_1}{\epsilon}(s-k\delta)} \\
& \quad + \frac{C_1}{\epsilon} \int_{k\delta}^s e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-r)} \|X^\epsilon(r) - X^\epsilon(r_\delta)\| dr.
\end{aligned}$$

Next, multiplying both sides of the above equation by $e^{-\rho t}$ with $t > s, \rho > 1$, we have

$$\begin{aligned}
& e^{-\rho t} \|Y^\epsilon(s) - \hat{Y}^\epsilon(s)\| \\
& \leq C e^{-\rho t} e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-k\delta)} (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) \\
& \quad + \frac{C_1}{\epsilon} \int_{k\delta}^s e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-r)} (r - r_\delta)^\gamma r_\delta^{-\gamma} e^{-\rho(t-r)} \frac{r_\delta^\gamma e^{-\rho r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\|}{(r - r_\delta)^\gamma} dr \\
& \leq C e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-k\delta)} (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) \\
& \quad + \delta^\gamma \|X^\epsilon\|_{\gamma, \rho, \sim} \frac{C_1}{\epsilon} \int_{k\delta}^s e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-r)} r_\delta^{-\gamma} dr.
\end{aligned}$$

Integrate the above inequality from $k\delta$ to $(k+1)\delta$, by Lemma 4.12, we have

$$\begin{aligned}
& e^{-\rho t} \int_{k\delta}^{(k+1)\delta} \|Y^\epsilon(s) - \hat{Y}^\epsilon(s)\| ds \\
& \leq C \int_{k\delta}^{(k+1)\delta} e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-k\delta)} (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) ds \\
& \quad + \int_{k\delta}^{(k+1)\delta} \delta^\gamma \|X^\epsilon\|_{\gamma, \rho, \sim} \frac{C_1}{\epsilon} \int_{k\delta}^s e^{-\frac{(\lambda_B - C_1)}{\epsilon}(s-r)} r_\delta^{-\gamma} dr ds \\
& \leq C\epsilon (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) + C\delta^\gamma \|X^\epsilon\|_{\gamma, \rho, \sim} \int_{k\delta}^{(k+1)\delta} s_\delta^{-\gamma} ds \\
& \leq C\epsilon (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) + C(1 + \|X_0\|) \delta^{1+\gamma} (k\delta)^{-\gamma} \\
& \leq C\delta^{1+\gamma} (1 + (k\delta)^{-\gamma})
\end{aligned}$$

where we take $\epsilon(1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) \leq \delta^{1+\gamma}$ for ϵ small enough. \square

LEMMA 4.18. *Let (A1)-(A4) and (2.2) hold. For any $X_0 \in V$, as $\epsilon \rightarrow 0$ the solution of (4.15) converges to \bar{X} which solves (4.5)*

$$\lim_{\epsilon \rightarrow 0} \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \sim} = 0$$

where this norm is considered with respect to a fixed interval $[0, T]$.

Proof. For the following we fix $\gamma < \sigma < 1 - \sigma'', \sigma' < 1 - \gamma$ and define $\tilde{\sigma} = \min\{\sigma', \sigma'', \gamma\}$. We will show that for almost every (ω_1, ω_2) and every $\mu > 0$ there exists an $\epsilon_0 > 0$ so that for $\epsilon < \epsilon_0, \rho > \rho_0$ we have

$$(4.16) \quad \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim} \leq \mu.$$

Note that the norm here is equivalent to the norm in the conclusion. In the following proof a constant C appears. This constant can change from inequality to inequality. C may depend on $T, \omega_1, \omega_2, \sigma', \sigma'', \gamma$ and other parameters like the Lipschitz constant of f and of $x \mapsto Y_F^\epsilon(\omega_2, x)$. But C does not depend on $\mu, \epsilon, \rho, \delta$. Here $\delta \in (0, 1)$ is a parameter depending on μ . To estimate all the terms in the following inequality we have to consider 3 cases. For the first case the right hand side will be absorbed by the left hand side of the inequality when ρ is sufficiently large. The second case includes terms providing estimates like $C\delta^{\tilde{\sigma}}, \tilde{\sigma} > 0$ where C is a priori determined by $T, \omega_1, \omega_2, \sigma', \sigma'', \gamma$ but independent of $\mu, \epsilon, \rho, \delta$, then we choose *fixed* δ so that $C\delta^{\tilde{\sigma}} < \lambda\mu, \lambda > 0$ sufficiently small. The third case contains terms providing an estimate $C\delta^{-\tilde{\sigma}}, \tilde{\sigma} > 0$ which can be made arbitrarily small when ϵ is sufficiently small, taking in account that δ is fixed.

By applying triangle inequality to $\|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim}$, we obtain

$$\begin{aligned} & \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim} \\ & \leq \left\| \int_0^\cdot S_A(\cdot - r)(f(X^\epsilon(r_\delta), \hat{Y}^\epsilon(r)) - f(X^\epsilon(r_\delta), Y_F^\epsilon(\theta_r \omega_2, X^\epsilon(r_\delta)))) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)\Delta_f(X^\epsilon(r_\delta); X^\epsilon(r)) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)\Delta_f(X^\epsilon(r); \hat{X}^\epsilon(r)) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)\Delta_f(\hat{X}^\epsilon(r); \bar{X}(r)) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)\Delta_f(\bar{X}(r); \bar{X}(r_\delta)) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)(f(\bar{X}(r_\delta), Y_F^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta))) - \bar{f}(\bar{X}(r_\delta))) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)(\bar{f}(\bar{X}(r_\delta)) - \bar{f}(\bar{X}(r))) dr \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)(h(X^\epsilon(r)) - h(\hat{X}^\epsilon(r))) d\omega_1(r) \right\|_{\gamma, \rho, \sim} \\ & \quad + \left\| \int_0^\cdot S_A(\cdot - r)(h(\hat{X}^\epsilon(r)) - h(\bar{X}(r))) d\omega_1(r) \right\|_{\gamma, \rho, \sim} =: \sum_{i=1}^9 I_i \end{aligned}$$

where $r_\delta = \lfloor r/\delta \rfloor \delta$ is the nearest breakpoint preceding r and for $U(r), \hat{U}(r) \in V$

$$\Delta_f(U(r); \hat{U}(r)) := f(U(r), Y_F^\epsilon(\theta_r \omega_2, U(r))) - f(\hat{U}(r), Y_F^\epsilon(\theta_r \omega_2, \hat{U}(r))).$$

To proceed, we adapt the approach used in the proof of Lemma 3.5 to estimate I_2 .

$$\begin{aligned}
I_2 &\leq \sup_{t \in [0, T]} e^{-\rho t} \left\| \int_0^t S_A(t-r) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\| \\
&\quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_s^t S_A(t-r) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\| \\
&\quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_0^s (S_A(t-r) - S_A(s-r)) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\| \\
&=: I_{21} + I_{22} + I_{23}.
\end{aligned}$$

For I_{21} , if $0 \leq t < \delta$, it is easy to see $I_{21} \leq C\delta$, then we consider that $\delta \leq t$ and by the Lipschitz continuity of f , Y_F^ϵ and Lemma 4.12, also the boundedness of f ,

$$\begin{aligned}
I_{21} &\leq C \left(\sup_{t \in [0, \delta]} e^{-\rho t} \int_0^t \|\Delta_f(X^\epsilon(r); X^\epsilon(r_\delta))\| dr + \sup_{t \in [\delta, T]} e^{-\rho t} \int_0^t \|\Delta_f(X^\epsilon(r); X^\epsilon(r_\delta))\| dr \right) \\
&\leq C\delta + C \sup_{t \in [\delta, T]} e^{-\rho t} \left(\int_0^\delta \|\Delta_f(X^\epsilon(r); X^\epsilon(r_\delta))\| dr + \int_\delta^t \|\Delta_f(X^\epsilon(r); X^\epsilon(r_\delta))\| dr \right) \\
&\leq C\delta + C\delta + C \sup_{t \in [\delta, T]} \int_\delta^t e^{-\rho(t-r)} (r-r_\delta)^\gamma r_\delta^{-\gamma} \frac{r_\delta^\gamma e^{-\rho r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\|}{(r-r_\delta)^\gamma} dr \\
&\leq C\delta + C\delta^\gamma \sup_{t \in [\delta, T]} \left(\int_\delta^t r_\delta^{-\gamma} dr \right) \|X^\epsilon\|_{\gamma, \rho, \sim} \\
&\leq C\delta + C\delta^\gamma \sup_{t \in [\delta, T]} \left(\int_{t_\delta}^t r_\delta^{-\gamma} dr + \sum_{k=1}^{\lfloor t/\delta \rfloor - 1} \int_{k\delta}^{(k+1)\delta} (k\delta)^{-\gamma} dr \right) \|X^\epsilon\|_{\gamma, \rho, \sim} \\
&\leq C\delta + C\delta^\gamma \sup_{t \in [\delta, T]} \left(\delta^{1-\gamma} + \delta^{1-\gamma} \sum_{k=1}^{\lfloor t/\delta \rfloor - 1} \int_{k-1}^k k^{-\gamma} dv \right) \|X^\epsilon\|_{\gamma, \rho, \sim} \\
&\leq C\delta + C\delta^\gamma \sup_{t \in [\delta, T]} \left(\delta^{1-\gamma} + \delta^{1-\gamma} \int_0^{\lfloor t/\delta \rfloor - 1} v^{-\gamma} dv \right) \|X^\epsilon\|_{\gamma, \rho, \sim} \leq C\delta^\gamma.
\end{aligned}$$

Here, by Lemma 4.12, $\|X^\epsilon\|_{\gamma, \rho, \sim}$ can be estimated independently of $\epsilon > 0$.

For I_{22} and I_{23} we divide the estimate into two cases. Consider at first the integral under the supremum of I_{22} for $0 < s < t \leq T$ and $s < 2\delta$: If $s < t \leq 2\delta$, it is easy to see that this integral is less than $C\delta^{1-\gamma}$, then, we consider $s < 2\delta < t$

$$\begin{aligned}
&\sup_{0 < s < 2\delta < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_s^{2\delta} S_A(t-r) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\| \\
&\quad + \sup_{0 < s < 2\delta < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_{2\delta}^t S_A(t-r) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\| \\
&\leq \sup_{0 < s < 2\delta < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} (2\delta-s)^\gamma (2\delta-s)^{1-\gamma} \\
&\quad + C \sup_{0 < s < 2\delta < t \leq T} (t-s)^{-\gamma} \int_{2\delta}^t e^{-\rho(t-r)} s^\gamma r_\delta^{-\gamma} (r-r_\delta)^\gamma \frac{r_\delta^\gamma e^{-\rho r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\|}{(r-r_\delta)^\gamma} dr \\
&\leq C\delta^\gamma (1 + \|X^\epsilon\|_{\gamma, \rho, \sim}) \leq C\delta^\gamma
\end{aligned}$$

where $s^\gamma r_\delta^{-\gamma} \leq 1$ for every $\delta > 0$ and the Lipschitz continuity of f , Y_F^ϵ , Lemma 4.12, and the boundedness of f . For the integral under supremum of I_{23} we have

$$\sup_{\substack{0 < s < t \leq T \\ s < 2\delta}} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_0^s (S_A(t-r) - S_A(s-r)) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\|$$

$$\leq C \sup_{\substack{0 < s < t \leq T \\ s < 2\delta}} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \int_0^s (t-s)^\gamma (s-r)^{-\gamma} dr \leq C\delta$$

where we use the fact that f is bounded.

We continue with the area $0 < s < t \leq T$ and $2\delta \leq s$ for I_{22} :

$$\begin{aligned} & \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_s^t S_A(t-r) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\| \\ & \leq \sup_{2\delta \leq s < t \leq T} \int_s^t \frac{\|S_A(t-r)\| s^\gamma r_\delta^{-\gamma} e^{-\rho(t-r)} r_\delta^\gamma e^{-\rho r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\|}{(t-s)^\gamma (r-r_\delta)^{-\gamma} (r-r_\delta)^\gamma} dr \\ & \leq C \sup_{2\delta \leq s < t \leq T} \int_s^t \frac{s^\gamma r_\delta^{-\gamma} e^{-\rho(t-r)}}{(t-s)^\gamma (r-r_\delta)^{-\gamma}} dr \|X^\epsilon\|_{\gamma, \rho, \sim} \leq C\delta^\gamma. \end{aligned}$$

where $s^\gamma r_\delta^{-\gamma} \leq c$ independently of $\delta > 0$ and by the Lipschitz continuity of f , Y_F^ϵ and Lemma 4.12. Next, for I_{23} we have

$$\begin{aligned} & \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \frac{\left\| \int_0^{2\delta} (S_A(t-r) - S_A(s-r)) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ & + \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \frac{\left\| \int_{2\delta}^s (S_A(t-r) - S_A(s-r)) \Delta_f(X^\epsilon(r); X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ & \leq \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \frac{\int_0^{2\delta} \|(S(t-s) - \text{id})S_A(s-r)\| \|\Delta_f(X^\epsilon(r); X^\epsilon(r_\delta))\| dr}{(t-s)^\gamma} \\ & + \sup_{2\delta \leq s < t \leq T} \frac{\int_{2\delta}^s s^\gamma \|(S(t-s) - \text{id})S_A(s-r)\| e^{-\rho(t-r)} e^{-\rho r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\| dr}{(t-s)^\gamma} \\ & \leq C \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \int_0^{2\delta} (t-s)^\gamma (s-r)^{-\gamma} dr \\ & + C \sup_{2\delta \leq s < t \leq T} \int_{2\delta}^s \frac{s^\gamma r_\delta^{-\gamma} (t-s)^\gamma (s-r)^{-\gamma} e^{-\rho(t-r)} r_\delta^\gamma e^{-\rho r} \|X^\epsilon(r) - X^\epsilon(r_\delta)\|}{(t-s)^\gamma (r-r_\delta)^{-\gamma} (r-r_\delta)^\gamma} dr \\ & \leq C \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \int_0^{2\delta} (2\delta-r)^{-\gamma} dr + C \sup_{2\delta \leq s < t \leq T} s^\gamma \delta^\gamma \int_{2\delta}^s \left(\frac{r}{r_\delta}\right)^\gamma r^{-\gamma} (s-r)^{-\gamma} dr \\ & \leq C\delta^{1-\gamma} + C \sup_{2\delta \leq s < t \leq T} s^\gamma \delta^\gamma \int_0^s r^{-\gamma} (s-r)^{-\gamma} dr \leq C\delta^{\bar{\sigma}} \end{aligned}$$

where we use the Lipschitz continuity of f , Y_F^ϵ and Lemma 4.12, also the boundedness of f . Thus, putting above estimates together, we have

$$I_2 \leq C\delta^{\bar{\sigma}}.$$

Based on the Lipschitz continuity of f, \bar{f}, Y_F^ϵ and boundedness of f, \bar{f} , Remark 4.14 we can apply the estimates for I_2 to estimate I_5 and I_7 . We have

$$I_5 + I_7 \leq C\delta^\gamma.$$

Then, by the Lipschitz continuity of f, Y_F^ϵ and Remark 4.14 again, we have

$$\begin{aligned} I_4 & \leq \sup_{t \in [0, T]} e^{-\rho t} \left\| \int_0^t S_A(t-r) \Delta_f(\hat{X}^\epsilon(r); \bar{X}(r)) dr \right\| \\ & + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_s^t S_A(t-r) \Delta_f(\hat{X}^\epsilon(r); \bar{X}(r)) dr \right\| \end{aligned}$$

$$\begin{aligned}
& + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} (t-s)^{-\gamma} \left\| \int_0^s (S_A(t-r) - S_A(s-r)) \Delta_f(\hat{X}^\epsilon(r); \bar{X}(r)) dr \right\| \\
& =: I_{41} + I_{42} + I_{43}.
\end{aligned}$$

For the first term above, we have

$$I_{41} \leq C \sup_{t \in [0, T]} \int_0^t e^{-\rho(t-r)} e^{-\rho r} \|\hat{X}^\epsilon(r) - \bar{X}(r)\| dr \leq C \rho^{-1} \sup_{r \in [0, T]} e^{-\rho r} \|\hat{X}^\epsilon(r) - \bar{X}(r)\|.$$

Next, for I_{42} , by Lemma 4.21, one has

$$\begin{aligned}
I_{42} & \leq C \sup_{0 < s < t \leq T} s^\gamma \frac{\int_s^t \|S_A(t-r)\| e^{-\rho(t-r)} e^{-\rho r} \|\hat{X}^\epsilon(r) - \bar{X}(r)\| dr}{(t-s)^\gamma} \\
& \leq C \sup_{0 < s < t \leq T} \int_s^t e^{-\rho(t-r)} (t-r)^{-\gamma} dr \sup_{r \in [0, T]} e^{-\rho r} \|\hat{X}^\epsilon(r) - \bar{X}(r)\| \\
& \leq C \sup_{0 < s < t \leq T} \int_0^t e^{-\rho(t-r)} (t-r)^{-\gamma} dr \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim} \\
& \leq C \rho^{-1+\gamma} \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim}.
\end{aligned}$$

The third integral on the right hand of I_4 can be estimated by Lemma 4.21, and taking $\sigma > \gamma$, that is, we have by Lemmas 3.10 and 3.11,

$$\begin{aligned}
I_{43} & \leq \sup_{0 < s < t \leq T} s^\gamma \frac{\int_0^s e^{-\rho(t-r)} \|(S(t-s) - \text{id}) S_A(s-r)\| e^{-\rho r} \|\hat{X}^\epsilon(r) - \bar{X}(r)\| dr}{(t-s)^\gamma} \\
& \leq C \sup_{0 < s < t \leq T} s^\gamma \int_0^s e^{-\rho(t-r)} (s-r)^{-\sigma} (t-s)^{-\gamma+\sigma} dr \sup_{r \in [0, T]} e^{-\rho r} \|\hat{X}^\epsilon(r) - \bar{X}(r)\| \\
& \leq C \rho^{-1+\sigma} \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim}.
\end{aligned}$$

Thus, taking ρ large enough, we have

$$(4.17) \quad I_4 \leq \frac{1}{3} \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim}.$$

Dealing with I_6 we need ergodic theory. Denote

$$\Delta_{f, \bar{f}}^\epsilon(\omega_2, x) := f(x, Y_F^\epsilon(\omega_2, x)) - \bar{f}(x), \quad \Delta_{f, \bar{f}}^1(\omega_2, x) := f(x, Y_F^1(\omega_2, x)) - \bar{f}(x)$$

where x is $\bar{X}(r_\delta)$ or $\bar{X}(k\delta)$, then, we have

$$\begin{aligned}
I_6 & \leq \sup_{t \in [0, T]} e^{-\rho t} \left\| \int_0^t S_A(t-r) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\| \\
& \quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} \left\| \frac{\int_s^t S_A(t-r) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr}{(t-s)^\gamma} \right\| \\
& \quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} \left\| \frac{\int_0^s (S_A(t-r) - S_A(s-r)) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr}{(t-s)^\gamma} \right\| \\
& =: I_{61} + I_{62} + I_{63}.
\end{aligned}$$

The first term above can be written

$$I_{61} \leq \sup_{t \in [0, T]} \left\| \int_0^t (S_A(t-r) - S_A(t-r_\delta)) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|$$

$$\begin{aligned}
& + \sup_{t \in [0, T]} \left\| \int_0^t S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\| \\
& =: I_{611} + I_{612}.
\end{aligned}$$

By the boundedness property of f, \bar{f} and Lemmas 3.10 and 3.11, we have

$$\begin{aligned}
I_{611} & \leq \sup_{t \in [0, T]} \int_0^t \|(S_A(t-r) - S_A(t-r_\delta))\| \|\Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta))\| dr \\
& \leq C\delta^\sigma \sup_{t \in [0, T]} \int_0^t \|(-A)^\sigma S_A(t-r)\| dr \leq C\delta^\sigma.
\end{aligned}$$

Then, one has

$$\begin{aligned}
I_{612} & \leq \sup_{t \in [0, T]} \left\| \int_{t_\delta}^t S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\| \\
& + \sup_{t \in [0, T]} \left\| \sum_{k=0}^{\lfloor t/\delta \rfloor - 1} \int_{k\delta}^{(k+1)\delta} S_A(t-k\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\| \\
& \leq \sup_{t \in [0, T]} \int_{t_\delta}^t \|S_A(t-r_\delta)\| \|\Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta))\| dr \\
& + \sup_{t \in [0, T]} \sum_{k=0}^{\lfloor t/\delta \rfloor - 1} \|(-A)^\sigma S_A(t-k\delta)\| \left\| \int_{k\delta}^{(k+1)\delta} (-A)^{-\sigma} \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \\
& \leq C\delta + C\delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \left\| \int_{k\delta}^{(k+1)\delta} (-A)^{-\sigma} \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \\
& \leq C\delta + C\delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \left\| \int_{k\delta}^{(k+1)\delta} (-A)^{-\sigma} \Delta_{f, \bar{f}}^1(\theta_{\frac{x}{\epsilon} \omega_2}, \bar{X}(k\delta)) dr \right\| \\
& \leq C\delta + C\delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \left\| \epsilon \int_{\frac{k\delta}{\epsilon}}^{\frac{(k+1)\delta}{\epsilon}} (-A)^{-\sigma} \Delta_{f, \bar{f}}^1(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \\
& \leq C\delta + C\delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \frac{T\epsilon}{\delta(k+1)} \left\| \int_0^{\frac{(k+1)\delta}{\epsilon}} (-A)^{-\sigma} \Delta_{f, \bar{f}}^1(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \\
& + C\delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \frac{T\epsilon}{\delta k} \left\| \int_0^{\frac{k\delta}{\epsilon}} (-A)^{-\sigma} \Delta_{f, \bar{f}}^1(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\|
\end{aligned}$$

where we use the fact that

$$(4.18) \quad \sup_{t \in [0, T]} \sum_{k=0}^{\lfloor t/\delta \rfloor - 1} \|(-A)^\sigma S_A(t-k\delta)\| \leq C\delta^{-1}, \quad \sigma \in (0, 1),$$

see Page 28 in Pei et al. [33]. We have for $\epsilon \rightarrow 0$, $\frac{(k+1)\delta}{\epsilon} \rightarrow +\infty$ for any $k, 1 \leq k \leq \lfloor T/\delta \rfloor - 1$. In addition we take the maximum over finitely many elements determined by the fixed number δ given and T . Following Lemma 4.11, we have for every element under the maximum

$$(4.19) \quad \frac{\epsilon}{\delta(k+1)} \left\| \int_0^{\frac{(k+1)\delta}{\epsilon}} (-A)^{-\sigma} \Delta_{f, \bar{f}}^1(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0$$

almost surely. We note that by Lemma 4.11 we can consider as an argument the random variable $\bar{X}(k\delta)$ inside the integrand of the last integral because the exceptional

set for the convergence is independent of x . Thus, we have for ϵ sufficiently small depending on (ω_1, ω_2) almost surely and the δ given

$$(4.20) \quad I_{61} \leq C\delta^\sigma.$$

Next, we turn to estimate I_{62} :

$$\begin{aligned} I_{62} &\leq C \sup_{0 < s < t \leq T} \frac{\left\| \int_s^t (S_A(t-r) - S_A(t-r_\delta)) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ &\quad + C \sup_{0 < s < t \leq T} \frac{\left\| \int_s^t S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} =: I_{621} + I_{622}. \end{aligned}$$

For above estimate, let us begin with I_{621} . Taking $\sigma' < 1 - \gamma$ into account, by the boundedness property of f, \bar{f} , we have

$$\begin{aligned} I_{621} &\leq C \sup_{0 < s < t \leq T} \left\{ (t-s)^{-\gamma} \int_s^t \left\| (S_A(t-r) - S_A(t-r_\delta)) \right\| \left\| \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) \right\| dr \right\} \\ &\leq C \sup_{0 < s < t \leq T} \left\{ (t-s)^{-\gamma} \delta^{\sigma'} \int_s^t (t-r)^{-\sigma'} dr \right\} \\ &\leq C \sup_{0 < s < t \leq T} \left\{ (t-s)^{-\gamma+1-\sigma'} \delta^{\sigma'} \right\} \leq C\delta^{\sigma'}. \end{aligned}$$

Now, we deal with I_{622} . Consider $\ell_t := \{s < t : t < (\lfloor \frac{s}{\delta} \rfloor + 2)\delta\}$, $\ell_t^c = \{s < t : t \geq (\lfloor \frac{s}{\delta} \rfloor + 2)\delta\}$. Note that we have for $s \in \ell_t$ that $t-s < 2\delta$ and for $s \in \ell_t^c$ that $t-s \geq \delta$.

$$\begin{aligned} I_{622} &\leq C \sup_{0 < s < t \leq T} \left\{ \frac{\left\| \int_s^t S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} \mathbf{1}_{\ell_t}(s) \right\} \\ &\quad + C \sup_{0 < s < t \leq T} \left\{ \frac{\left\| \int_s^{(\lfloor s\delta^{-1} \rfloor + 1)\delta} S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \\ &\quad + C \sup_{0 < s < t \leq T} \left\{ \frac{\left\| \int_{\lfloor t\delta^{-1} \rfloor \delta}^t S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \\ &\quad + C \sup_{0 < s < t \leq T} \left\{ \frac{\left\| \int_{(\lfloor s\delta^{-1} \rfloor + 1)\delta}^{\lfloor t\delta^{-1} \rfloor \delta} S_A(t-r_\delta) \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\}. \end{aligned}$$

The first three expressions on the right hand side of the last inequality can be estimated by $C\delta^{1-\gamma}$. Thus, we have

$$\begin{aligned} I_{622} &\leq C\delta^{1-\gamma} \\ &\quad + C \sup_{0 < s < t \leq T} \left\{ \frac{\left\| \sum_{k=\lfloor s\delta^{-1} \rfloor + 1}^{\lfloor t\delta^{-1} \rfloor} S_A(t-r_\delta) \int_{k\delta}^{(k+1)\delta} \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \\ &\leq C\delta^{1-\gamma} + C\delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \left\| \int_{k\delta}^{(k+1)\delta} (-A)^{-\sigma} \Delta_{f, \bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \end{aligned}$$

where we apply (4.18). Using the ergodic theorem again, the remaining term on the right hand side can be estimated similar to I_{612} , see (4.19). We have

$$I_{62} \leq C\delta^{\bar{\sigma}}$$

for sufficiently small $\epsilon > 0$.

The next term is

$$\begin{aligned} I_{63} &\leq C \sup_{0 < s < t \leq T} \frac{\int_0^s \|(S_A(t-s) - \text{id})(S_A(s-r) - S_A(s-r_\delta))\| \|\Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta))\| dr}{(t-s)^\gamma} \\ &\quad + C \sup_{0 < s < t \leq T} \frac{\left\| \int_0^s (S_A(t-s) - \text{id}) S_A(s-r_\delta) \Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^\gamma} =: I_{631} + I_{632}. \end{aligned}$$

For I_{631} , taking $\gamma < \sigma < 1 - \sigma''$, by the boundedness property of f, \bar{f} and $r - r_\delta \leq \delta$, we have

$$\begin{aligned} I_{631} &\leq C \sup_{0 < s < t \leq T} \int_0^s (t-s)^{\sigma-\gamma} \|(-A)^\sigma (S_A(s-r) - S_A(s-r_\delta))\| dr \\ &\leq C \delta^{\sigma''} \sup_{0 < s < t \leq T} \int_0^s (t-s)^{\sigma-\gamma} \|(-A)^{\sigma+\sigma''} S_A(s-r)\| dr \\ &\leq C \delta^{\sigma''} \sup_{0 < s < t \leq T} \int_0^s (t-s)^{\sigma-\gamma} (s-r)^{-\sigma-\sigma''} dr \leq C \delta^{\sigma''} \end{aligned}$$

and for I_{632} and $\gamma < \sigma < 1 - \sigma''$,

$$\begin{aligned} I_{632} &\leq C \sup_{0 < s < t \leq T} \frac{\left\| \int_{\lfloor \frac{s}{\delta} \rfloor \delta}^s (-A)^\sigma S_A(s - \lfloor \frac{r}{\delta} \rfloor \delta) \Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta)) dr \right\|}{(t-s)^{\gamma-\sigma}} \\ &\quad + C \sup_{0 < s < t \leq T} \frac{\left\| \sum_{k=0}^{\lfloor \frac{s}{\delta} \rfloor - 1} \int_{k\delta}^{(k+1)\delta} (-A)^\sigma S_A(s - k\delta) \Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\|}{(t-s)^{\gamma-\sigma}} \\ &\leq C \sup_{0 < s < t \leq T} \left\{ (t-s)^{-\gamma+\sigma} \int_{\lfloor \frac{s}{\delta} \rfloor \delta}^s (s - \lfloor \frac{r}{\delta} \rfloor \delta)^{-\sigma} dr \right\} \\ &\quad + C \sup_{0 < s < t \leq T} \frac{\left\| \sum_{k=0}^{\lfloor \frac{s}{\delta} \rfloor - 1} (-A)^{\sigma+\sigma''} S_A(s - k\delta) \int_{k\delta}^{(k+1)\delta} (-A)^{-\sigma''} \Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\|}{(t-s)^{\gamma-\sigma}} \\ &\leq C \delta^{1-\sigma} + C \delta^{-1} \max_{0 \leq k \leq \lfloor T/\delta \rfloor - 1} \left\| \int_{k\delta}^{(k+1)\delta} (-A)^{-\sigma''} \Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(k\delta)) dr \right\| \end{aligned}$$

where we apply (4.18) again. Using ergodic theorem and the estimate similar to (4.19) again and taking ϵ small enough, we have

$$I_{63} \leq C \delta^{\bar{\sigma}}.$$

To deal with I_1 , by replacing $\Delta_{f,\bar{f}}^\epsilon(\theta_r \omega_2, \bar{X}(r_\delta))$ in I_6 by

$$(f(X^\epsilon(r_\delta), \hat{Y}^\epsilon(r)) - f(X^\epsilon(r_\delta), Y_F^\epsilon(\theta_r \omega_2, X^\epsilon(r_\delta))))$$

we can apply the techniques to estimate I_6 . But instead the ergodic theory argument we apply Lemma 4.16 so that

$$I_1 \leq C \delta^{\bar{\sigma}} + C \delta^{-1} (1 + \|X_0\| + \|Y_0\| + o(\epsilon^{-1})) \epsilon \leq C \delta^{\bar{\sigma}}$$

for $\epsilon < \epsilon_0$ and δ given we have that $\delta^{-1} o(\epsilon^{-1}) \epsilon < C \delta^\gamma$.

The estimates for I_8 and I_9 follow by using the same techniques in Appendix and [4, Lemma 9]. Thus, we have

$$I_8 + I_9 \leq C \|\omega_1\|_\beta (1 + \|X^\epsilon\|_{\gamma,\rho,\sim} + \|\hat{X}^\epsilon\|_{\gamma,\rho,\sim}) K(\rho) \|X^\epsilon - \hat{X}^\epsilon\|_{\gamma,\rho,\sim}$$

$$(4.21) \quad +C \|\omega_1\|_\beta (1 + \|\bar{X}\|_{\gamma,\rho,\sim} + \|\hat{X}^\epsilon\|_{\gamma,\rho,\sim})K(\rho)\|\hat{X}^\epsilon - \bar{X}\|_{\gamma,\rho,\sim}$$

where $\lim_{\rho \rightarrow \infty} K(\rho) = 0$.

For $\|X^\epsilon - \hat{X}^\epsilon\|_{\gamma,\rho,\sim}$, by (4.3) and (4.15), it is easy to see

$$(4.22) \quad \begin{aligned} \|X^\epsilon - \hat{X}^\epsilon\|_{\gamma,\rho,\sim} &\leq \left\| \int_0^\cdot S_A(\cdot - r)(f(X^\epsilon(r), Y^\epsilon(r)) - f(X^\epsilon(r_\delta), Y^\epsilon(r))) dr \right\|_{\gamma,\rho,\sim} \\ &\quad + \left\| \int_0^\cdot S_A(\cdot - r)(f(X^\epsilon(r_\delta), Y^\epsilon(r)) - f(X^\epsilon(r_\delta), \hat{Y}^\epsilon(r))) dr \right\|_{\gamma,\rho,\sim} \\ &=: J_1 + J_2. \end{aligned}$$

By the same techniques for I_2 , it is easy to see the J_1 is less than $C\delta^\gamma$. For the second term of the right side of (4.22), we can apply the similar techniques used in the estimate of I_6 , replacing $\Delta_{f,\bar{f}}^\epsilon(X^\epsilon(r_\delta))$ by

$$\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) := f(X^\epsilon(r_\delta), Y^\epsilon(r)) - f(X^\epsilon(r_\delta), \hat{Y}^\epsilon(r)).$$

Thus, we have

$$\begin{aligned} J_2 &\leq \sup_{t \in [0, T]} e^{-\rho t} \int_0^t \|(S_A(t-r) - S_A(t-r_\delta))\| \|\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr \\ &\quad + \sup_{t \in [0, T]} e^{-\rho t} \left\| \int_0^t S_A(t-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\| \\ &\quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} \frac{\int_s^t \|(S_A(t-r) - S_A(t-r_\delta))\| \|\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr}{(t-s)^\gamma} \\ &\quad + \sup_{\substack{0 < s < t \leq T \\ s < 2\delta}} s^\gamma e^{-\rho t} \frac{\left\| \int_s^t S_A(t-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ &\quad + \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \frac{\left\| \int_s^t S_A(t-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ &\quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} \frac{\int_0^s \|(S_A(t-s) - \text{id})(S_A(s-r) - S_A(s-r_\delta))\| \|\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr}{(t-s)^\gamma} \\ &\quad + \sup_{\substack{0 < s < t \leq T \\ s < 2\delta}} s^\gamma e^{-\rho t} \frac{\left\| \int_0^s (S_A(t-s) - \text{id}) S_A(s-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ &\quad + \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \frac{\left\| \int_0^{2\delta} (S_A(t-s) - \text{id}) S_A(s-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} \\ &\quad + \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \frac{\left\| \int_{2\delta}^s (S_A(t-s) - \text{id}) S_A(s-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\|}{(t-s)^\gamma} = \sum_{i=1}^9 J_{2i}. \end{aligned}$$

For the terms $J_{21}, J_{23}, J_{24}, J_{26}, J_{27}, J_{28}$, it is easy to know

$$J_{21} + J_{23} + J_{24} + J_{26} + J_{27} + J_{28} \leq C\delta^{\bar{\sigma}}.$$

Then, using same method with the estimates of I_{612}, I_{622} and I_{632} , we apply Lemma 4.17 instead the ergodic theory argument. By the Lipschitz continuity and

boundedness property of f and Lemma 4.17, using same method with the estimate of I_{21} , we have

$$\begin{aligned}
J_{22} &\leq \sup_{t \in [0, T]} e^{-\rho t} \int_0^\delta \|S_A(t-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr \\
&\quad + \sup_{t \in [0, T]} e^{-\rho t} \int_{t_\delta}^t \|S_A(t-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr \\
&\quad + \sup_{t \in [0, T]} e^{-\rho t} \left\| \sum_{k=1}^{\lfloor t/\delta \rfloor - 1} \int_{k\delta}^{(k+1)\delta} S_A(t-k\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr \right\| \\
&\leq C\delta + C \sup_{t \in [0, T]} \sum_{k=1}^{\lfloor t/\delta \rfloor - 1} e^{-\rho t} \int_{k\delta}^{(k+1)\delta} \|Y^\epsilon(r) - \hat{Y}^\epsilon(r)\| dr \\
&\leq C\delta + C \sup_{t \in [0, T]} \sum_{k=1}^{\lfloor t/\delta \rfloor - 1} \delta^{1+\gamma} (1 + (k\delta)^{-\gamma}) \\
&\leq C\delta^\gamma + C \sup_{t \in [0, T]} \sum_{k=1}^{\lfloor t/\delta \rfloor - 1} \delta \int_{k-1}^k k^{-\gamma} dv \leq C\delta^\gamma.
\end{aligned}$$

To proceed, for J_{25} and J_{29} , by the Lipschitz continuity and boundedness property of f and Lemma 4.17, one has

$$\begin{aligned}
J_{25} &\leq \sup_{2\delta \leq s < t \leq T} \left\{ s^\gamma e^{-\rho t} \frac{\int_s^t \|S_A(t-r_\delta)\| \|\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr}{(t-s)^\gamma} \mathbf{1}_{\ell_t}(s) \right\} \\
&\quad + \sup_{2\delta \leq s < t \leq T} \left\{ s^\gamma e^{-\rho t} \frac{\int_s^{s_\delta + \delta} \|S_A(t-r_\delta)\| \|\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \\
&\quad + \sup_{2\delta \leq s < t \leq T} \left\{ s^\gamma e^{-\rho t} \frac{\int_{t_\delta}^t \|S_A(t-r_\delta)\| \|\Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta))\| dr}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \\
&\quad + C \sup_{2\delta \leq s < t \leq T} \left\{ s^\gamma e^{-\rho t} \frac{\sum_{k=\lfloor s\delta^{-1} \rfloor + 1}^{\lfloor t\delta^{-1} \rfloor - 1} \int_{k\delta}^{(k+1)\delta} \|Y^\epsilon(r) - \hat{Y}^\epsilon(r)\| dr}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \\
&\leq C\delta^{1-\gamma} + C \sup_{2\delta \leq s < t \leq T} \left\{ \frac{\sum_{k=\lfloor s\delta^{-1} \rfloor + 1}^{\lfloor t\delta^{-1} \rfloor - 1} s^\gamma \delta^{1+\gamma} (1 + (k\delta)^{-\gamma})}{(t-s)^\gamma} \mathbf{1}_{\ell_t^c}(s) \right\} \leq C\delta^{\bar{\sigma}}
\end{aligned}$$

where the first three terms are less than $C\delta^{1-\gamma}$ by the boundedness property of f and

$$\begin{aligned}
\sum_{k=\lfloor s\delta^{-1} \rfloor + 1}^{\lfloor t\delta^{-1} \rfloor - 1} s^\gamma (1 + (k\delta)^{-\gamma}) &\leq \sum_{k=\lfloor s\delta^{-1} \rfloor + 1}^{\lfloor t\delta^{-1} \rfloor - 1} \left(s^\gamma + \left(\frac{s}{k\delta}\right)^\gamma \right) \leq \sum_{k=\lfloor s\delta^{-1} \rfloor + 1}^{\lfloor t\delta^{-1} \rfloor - 1} (T^\gamma + 1) \\
&\leq C(\lfloor t\delta^{-1} \rfloor - 1 - (\lfloor s\delta^{-1} \rfloor + 1) + 1) \leq C\delta^{-1}(t-s)
\end{aligned}$$

has been used in the last term. Then, we have

$$\begin{aligned}
J_{29} &\leq \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \left\| \frac{\int_{s_\delta}^s (S_A(t-s) - \text{id}) S_A(s-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr}{(t-s)^\gamma} \right\| \\
&\quad + \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \left\| \frac{\int_{2\delta}^{s_\delta} (S_A(t-s) - \text{id}) S_A(s-r_\delta) \Delta_f^{Y^\epsilon, \hat{Y}^\epsilon}(X^\epsilon(r_\delta)) dr}{(t-s)^\gamma} \right\|
\end{aligned}$$

$$\begin{aligned}
&\leq C\delta^{1-\gamma} + C \sup_{2\delta \leq s < t \leq T} s^\gamma e^{-\rho t} \int_{2\delta}^{s\delta} (s-r_\delta)^{-\gamma} \|Y^\epsilon(r) - \hat{Y}^\epsilon(r)\| dr \\
&\leq C\delta^{1-\gamma} + C \sup_{2\delta \leq s < t \leq T} \sum_{k=2}^{\lfloor s\delta^{-1} \rfloor - 1} s^\gamma (s-k\delta)^{-\gamma} e^{-\rho t} \int_{k\delta}^{(k+1)\delta} \|Y^\epsilon(r) - \hat{Y}^\epsilon(r)\| dr \\
&\leq C\delta^{1-\gamma} + C \sup_{2\delta \leq s < t \leq T} \sum_{k=2}^{\lfloor s\delta^{-1} \rfloor - 1} s^\gamma (s-k\delta)^{-\gamma} \delta^{1+\gamma} (1+(k\delta)^{-\gamma}) \\
&\leq C\delta^{1-\gamma} + C\delta^\gamma + C \sup_{2\delta \leq s < t \leq T} \sum_{k=2}^{\lfloor s\delta^{-1} \rfloor - 1} \int_{k\delta}^{(k+1)\delta} s^\gamma \delta^\gamma (s-k\delta)^{-\gamma} (k\delta)^{-\gamma} dr \\
&\leq C\delta^{1-\gamma} + C\delta^\gamma + C\delta^\gamma \sup_{2\delta \leq s < t \leq T} \int_{2\delta}^{s\delta} s^\gamma (s-r_\delta)^{-\gamma} r_\delta^{-\gamma} dr \\
&\leq C\delta^{1-\gamma} + C\delta^\gamma + C\delta^\gamma \sup_{2\delta \leq s < t \leq T} \int_{2\delta}^s s^\gamma (s-r)^{-\gamma} r^{-\gamma} \frac{r_\delta^{-\gamma}}{r^{-\gamma}} dr \\
&\leq C\delta^{1-\gamma} + C\delta^\gamma + C\delta^\gamma \sup_{2\delta \leq s < t \leq T} \int_0^s s^\gamma (s-r)^{-\gamma} r^{-\gamma} dr \leq C\delta^{\tilde{\sigma}}.
\end{aligned}$$

Thus, we have

$$(4.23) \quad \|X^\epsilon - \hat{X}^\epsilon\|_{\gamma, \rho, \sim} \leq C\delta^{\tilde{\sigma}}.$$

Then, by (4.23) and taking ρ large enough and δ small enough, we have

$$(4.24) \quad I_8 + I_9 \leq C\delta^{\tilde{\sigma}} + \frac{1}{3} \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim}.$$

To deal with I_3 , we can apply the similar techniques used in the estimate of I_4 . By the Lipschitz continuity of f , (A2), Lemma 4.12, Lemma 4.17 and (4.23), it is easy to see

$$I_3 \leq C \|X^\epsilon - \hat{X}^\epsilon\|_{\gamma, \rho, \sim} \leq C\delta^{\tilde{\sigma}}.$$

Thus, putting above estimates together, we have for sufficiently small $\epsilon > 0$

$$(4.25) \quad \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim} \leq C\delta^{\tilde{\sigma}} + \frac{2}{3} \|\hat{X}^\epsilon - \bar{X}\|_{\gamma, \rho, \sim}$$

so that (4.16) holds. \square

LEMMA 4.19. *Let (A1)-(A4) and (2.2) hold. For any $X_0 \in V$, as $\epsilon \rightarrow 0$ the solution of (4.15) converges to X^ϵ which solves (4.3). That is, we have almost surely*

$$\lim_{\epsilon \rightarrow 0} \|X^\epsilon - \hat{X}^\epsilon\|_{\gamma, \sim} = 0$$

where this norm is considered with respect to a fixed interval $[0, T]$.

Proof. Note that the norm in (4.23) is equivalent to the norm in the conclusion. By (4.23), similar to the argument of (4.16), the desired result will be obtained. \square

To close this section, we note that Lemma 4.18 and Lemma 4.19 yield Theorem 4.2. This completes the proof of Theorem 4.2.

Appendix: Several auxiliary technical lemmas. We recall the following technical lemma from [4, 13].

LEMMA 4.20. [4, Lemma 8] *Let $a > -1, b > -1$ and $a + b \geq -1, d > 0$ and $t \in [0, T]$. For $\rho > 0$ we define*

$$K(\rho) := \sup_{t \in [0, T]} t^d \int_0^1 e^{-\rho t(1-v)} v^a (1-v)^b dv,$$

then we have that $\lim_{\rho \rightarrow \infty} K(\rho) = 0$.

LEMMA 4.21. [14, Lemma 14] *For any non-negative a and d such that $a + d < 1$, and for any $\rho \geq 1$, there exists a positive constant c such that*

$$\int_0^t e^{-\rho(t-r)} (t-r)^{-a} r^{-d} dr \leq c \rho^{a+d-1}.$$

Proof of Lemma 3.5 By the definition of the norm and of \mathcal{T} ,

$$\begin{aligned} \|\mathcal{T}(u, \omega_1, \omega_2, u_0)\|_{\gamma, \rho, \sim} &\leq \|S_A(\cdot)u_{01}\|_{\gamma, \rho, \sim} + \|S_B(\cdot)(u_{02} - Z(\omega_2))\|_{\gamma, \rho, \sim} \\ &\quad + \|Z(\theta, \omega_2)\|_{\gamma, \rho, \sim} + \left\| \int_0^\cdot S_J(\cdot - r)F(u(r)) dr \right\|_{\gamma, \rho, \sim} \\ &\quad + \left\| \int_0^\cdot S_A(\cdot - r)h(u_1(r)) d\omega_1(r) \right\|_{\gamma, \rho, \sim} \\ &=: \mathbf{I}_1 + \mathbf{I}_2 + \mathbf{I}_3 + \mathbf{I}_4 + \mathbf{I}_5. \end{aligned}$$

By Lemma 4.8, we begin with estimate for $\mathbf{I}_1 + \mathbf{I}_2 + \mathbf{I}_3$

$$\begin{aligned} \mathbf{I}_1 + \mathbf{I}_2 + \mathbf{I}_3 &= \sup_{t \in [0, T]} e^{-\rho t} \|S_A(t)u_{01}\| + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} \frac{\|S_A(t)u_{01} - S_A(s)u_{01}\|}{(t-s)^\gamma} \\ &\quad + \sup_{t \in [0, T]} e^{-\rho t} \|S_B(t)(u_{02} - Z(\omega_2))\| + \|Z(\theta, \omega_2)\|_{\gamma, \rho, \sim} \\ &\quad + \sup_{0 < s < t \leq T} s^\gamma e^{-\rho t} \frac{\|S_B(t)(u_{02} - Z(\omega_2)) - S_B(s)(u_{02} - Z(\omega_2))\|}{(t-s)^\gamma} \\ &\leq c_T (\|u_{01}\| + \|u_{02}\| + \|Z(\theta, \omega_2)\|_\gamma) \end{aligned}$$

where we use $\|Z(\theta, \omega_2)\|_{\gamma, \rho, \sim} \leq c_T \|Z(\theta, \omega_2)\|_\gamma$ and Lemma 4.8.

Then, by [15, Lemma 4, (9)], for \mathbf{I}_4 , we have

$$\mathbf{I}_4 \leq c_T \bar{K}(\rho) (1 + \|u\|_{\gamma, \rho, \sim})$$

where \bar{K} has similar properties like K .

Now, we show the $\|\cdot\|_{\gamma, \rho, \sim}$ -norm of the stochastic integral.

$$\begin{aligned} \mathbf{I}_5 &= \sup_{0 < s < t \leq T} \frac{s^\gamma e^{-\rho t}}{(t-s)^\gamma} \left\| \int_s^t S_A(t-r)h(u_1(r)) d\omega_1(r) \right\| \\ &\quad + \sup_{0 < s < t \leq T} \frac{s^\gamma e^{-\rho t}}{(t-s)^\gamma} \left\| \int_0^s (S_A(t-r) - S_A(s-r))h(u_1(r)) d\omega_1(r) \right\| \\ (4.26) \quad &\quad + \sup_{t \in [0, T]} e^{-\rho t} \left\| \int_0^t S_A(t-r)h(u_1(r)) d\omega_1(r) \right\| =: \mathbf{I}_{51} + \mathbf{I}_{52} + \mathbf{I}_{53}. \end{aligned}$$

Since $\|D_t^{1-\alpha}\omega_{1,t-}[r]\| \leq c\|\omega_1\|_\beta (t-r)^{\alpha+\beta-1}$, by using the inequality of (3.12) and Remark 3.4 we get

$$\begin{aligned}
& s^\gamma e^{-\rho t} \left\| \int_s^t S_A(t-r)h(u_1(r)) d\omega_1(r) \right\| \\
& \leq c s^\gamma e^{-\rho t} \int_s^t \left(\frac{\|S_A(t-r)\|_{L(V)} \|h(u_1(r))\|_{L_2(V)}}{(r-s)^\alpha} \right. \\
& \quad + \int_s^r \frac{\|S_A(t-r) - S_A(t-q)\|_{L(V)} \|h(u_1(r))\|_{L_2(V)}}{(r-q)^{1+\alpha}} dq \\
& \quad \left. + \int_s^r \frac{\|S_A(t-q)\|_{L(V)} \|h(u_1(r)) - h(u_1(q))\|_{L_2(V)}}{(r-q)^{1+\alpha}} dq \right) \frac{\|\omega_1\|_\beta}{(t-r)^{-\alpha-\beta+1}} dr \\
& \leq c T^\gamma \|\omega_1\|_\beta \left(\int_s^t e^{-\rho(t-r)} \frac{(c_h + c_{Dh}|u_1(r)|)e^{-\rho r}}{(r-s)^\alpha} (t-r)^{\alpha+\beta-1} dr \right. \\
& \quad + \int_s^t \int_s^r e^{-\rho(t-r)} \frac{e^{-\rho r} (c_h + c_{Dh}|u_1(r)|)(r-q)^\gamma}{(t-r)^\gamma (r-q)^{1+\alpha}} dq (t-r)^{\alpha+\beta-1} dr \\
& \quad \left. + \int_s^t \int_s^r e^{-\rho(t-r)} \frac{e^{-\rho r} c_{Dh}|u_1(r) - u_1(q)|q^\gamma (r-q)^\gamma}{(r-q)^{1+\alpha} q^\gamma (r-q)^\gamma} dq (t-r)^{\alpha+\beta-1} dr \right) \\
& \leq c T^\gamma \|\omega_1\|_\beta (t-s)^\beta (1 + \|u_1\|_{\gamma,\rho,\sim}) \int_s^t e^{-\rho(t-r)} (r-s)^{-\alpha} (t-r)^{\alpha-1} dr \\
& \quad + c T^\gamma \|\omega_1\|_\beta (1 + \|u_1\|_{\gamma,\rho,\sim}) \int_s^t e^{-\rho(t-r)} (r-s)^{\gamma-\alpha} (t-r)^{\alpha+\beta-1-\gamma} dr \\
(4.27) \quad & + c T^\gamma \|\omega_1\|_\beta (t-s)^\beta \|u_1\|_{\gamma,\rho,\sim} \int_s^t e^{-\rho(t-r)} (r-s)^{-\alpha} (t-r)^{\alpha-1} dr.
\end{aligned}$$

By a change of variable, $\gamma < \beta$, it is easy to see that

$$\begin{aligned}
& (t-s)^\beta \int_s^t e^{-\rho(t-r)} (r-s)^{-\alpha} (t-r)^{\alpha-1} dr \\
& = (t-s)^\gamma (t-s)^{\beta-\gamma} \int_0^1 e^{-\rho(t-s)(1-v)} v^{-\alpha} (1-v)^{\alpha-1} dv \leq (t-s)^\gamma K(\rho),
\end{aligned}$$

taking in Lemma 4.20 $a = -\alpha, b = \alpha - 1, d = \beta - \gamma$ and $t - s$ as the corresponding t there. The second integral on the right side may be rewritten in the same way, since

$$\begin{aligned}
& \int_s^t e^{-\rho(t-r)} (r-s)^{\gamma-\alpha} (t-r)^{\alpha+\beta-1-\gamma} dr \\
& \leq (t-s)^\gamma (t-s)^{\beta-\gamma} \int_s^t e^{-\rho(t-r)} (r-s)^{-\alpha} (t-r)^{\alpha-1} dr.
\end{aligned}$$

Thus, we have

$$(4.28) \quad \mathbf{I}_{51} \leq c_T K(\rho) \|\omega_1\|_\beta (1 + \|u\|_{\gamma,\rho,\sim})$$

For \mathbf{I}_{52} , we should follow similar steps than before when obtaining (4.27). Now we need to replace the estimates for $\|S_A(t-r)\|_{L(V)}$ and $\|S_A(t-r) - S_A(t-q)\|_{L(V)}$ by estimates for $\|S_A(t-r) - S_A(s-r)\|_{L(V)}$ and $\|S_A(t-r) - S_A(t-q) - (S_A(s-r) - S_A(s-q))\|_{L(V)}$ respectively, for which we use (3.12) and (3.13) for appropriate parameters. Then it is not hard to see that for $\alpha' + \gamma < \alpha + \beta, 0 < \alpha < \alpha' < 1$:

$$s^\gamma e^{-\rho t} \left\| \int_0^s (S_A(t-r) - S_A(s-r))h(u_1(r)) d\omega_1(r) \right\|$$

$$\begin{aligned}
&\leq c(t-s)^\gamma \|\omega_1\|_\beta T^\gamma \left(\int_0^s e^{-\rho(t-r)} \frac{(c_h + c_{Dh}|u_1(r)|)e^{-\rho r}}{r^\alpha (s-r)^\gamma} (s-r)^{\alpha+\beta-1} dr \right. \\
&\quad + \int_0^s \int_0^r e^{-\rho(t-r)} \frac{e^{-\rho r} (c_h + c_{Dh}|u_1(r)|)(r-q)^{\alpha'}}{(s-r)^{\alpha'+\gamma} (r-q)^{1+\alpha}} dq (s-r)^{\alpha+\beta-1} dr \\
&\quad \left. + \int_0^s \int_0^r e^{-\rho(t-r)} \frac{e^{-\rho r} c_{Dh}|u_1(r) - u_1(q)|q^\gamma}{(s-r)^\gamma (r-q)^{1+\alpha} q^\gamma} dq (s-r)^{\alpha+\beta-1} dr \right) \\
&\leq c(t-s)^\gamma T^\gamma \|\omega_1\|_\beta (1 + \|u_1\|_{\gamma,\rho,\sim}) \int_0^s e^{-\rho(t-r)} r^{-\alpha} (s-r)^{\alpha+\beta-1-\gamma} dr \\
&\quad + c(t-s)^\gamma T^\gamma \|\omega_1\|_\beta (1 + \|u_1\|_{\gamma,\rho,\sim}) \int_0^s e^{-\rho(t-r)} r^{\alpha'-\alpha} (s-r)^{\alpha+\beta-1-\alpha'-\gamma} dr \\
&\quad + c(t-s)^\gamma T^\gamma \|\omega_1\|_\beta \|u_1\|_{\gamma,\rho,\sim} \int_0^s e^{-\rho(t-r)} r^{-\alpha} (s-r)^{\alpha-\gamma+\beta-1} dr.
\end{aligned}$$

The third integral on the right hand side of the last inequality can be estimated by

$$s^{\beta-\gamma} \int_0^1 e^{-\rho s(1-v)} v^{-\alpha} (1-v)^{\alpha-1} dv$$

and in a similar manner the other integrals. We have

$$\mathbf{I}_5 \leq c_T K(\rho) \|\omega_1\|_\beta (1 + \|u_1\|_{\gamma,\rho,\sim}) \leq c_T K(\rho) \|\omega_1\|_\beta (1 + \|u\|_{\gamma,\rho,\sim}).$$

In a similar manner than before for the first expression on \mathbf{I}_{51} we obtain

$$\mathbf{I}_{53} \leq c_T \|\omega_1\|_\beta K(\rho) (1 + \|u\|_{\gamma,\rho,\sim}).$$

All the previous estimates imply that

$$\mathbf{I}_5 \leq c_T \|\omega_1\|_\beta K(\rho) (1 + \|u\|_{\gamma,\rho,\sim}).$$

Collecting all the above estimates we have a constant $C(\rho, \omega_1, T) > 0$ such that $\lim_{\rho \rightarrow \infty} C(\rho, \omega_1, T) = 0$ and

$$\|\mathcal{T}(u, \omega_1, \omega_2, u_0)\|_{\gamma,\rho,\sim} \leq c_T (\|u_{01}\| + \|u_{02}\| + \|Z(\theta, \omega_2)\|_\gamma) + C(\rho, \omega_1, T) (1 + \|u\|_{\gamma,\rho,\sim}).$$

REFERENCES

- [1] L. ARNOLD, *Random Dynamical Systems*, Springer, 1998.
- [2] ———, *Hasselmann's Program Revisited: The Analysis of Stochasticity in Deterministic Climate Models*, Springer, 2001.
- [3] N. BOGOLYUBOV AND Y. MITROPOLSKII, *Asymptotic Methods in the Theory of Nonlinear Oscillations*, Moscow: Gostekhteorizdat, 1955.
- [4] Y. CHEN, H. GAO, M. GARRIDO-ATIENZA, AND B. SCHMALFUSS, *Pathwise solutions of SPDEs driven by Hölder-continuous integrators with exponent larger than 1/2 and random dynamical systems*, *Discrete and Continuous Dynamical Systems*, 34 (2014), pp. 79–98.
- [5] P. CHERIDITO, H. KAWAGUCHI, AND M. MAEJIMA, *Fractional Ornstein-Uhlenbeck processes*, *Electronic Journal of Probability*, 8 (2003), pp. 1–14.
- [6] I. CHUESHOV AND B. SCHMALFUSS, *Averaging of attractors and inertial manifolds for parabolic PDE with random coefficients*, *Advanced Nonlinear Studies*, 5 (2005), pp. 461–492.
- [7] I. CHUESHOV AND B. SCHMALFUSS, *Synchronization in Infinite-Dimensional Deterministic and Stochastic Systems*, Springer, 2020.
- [8] E. CODDINGTON AND N. LEVINSON, *Theory of Ordinary Differential Equations*, McGraw Hill, 1955.
- [9] J. DUAN AND W. WANG, *Effective Dynamics of Stochastic Partial Differential equations*, Elsevier, 2014.

- [10] K. EICHINGER, C. KUEHN, AND A. NEAMȚU, *Sample paths estimates for stochastic fast-slow systems driven by fractional Brownian motion*, Journal of Statistical Physics, 179 (2020), pp. 1222–1266.
- [11] M. ENGEL, M. A. GKOGKAS, AND C. KUEHN, *Homogenization of coupled fast-slow systems via intermediate stochastic regularization*, Journal of Statistical Physics, 183 (2021), p. 25.
- [12] M. FREIDLIN AND A. WENTZELL, *Random Perturbations of Dynamical Systems*, Springer, 2012.
- [13] M. GARRIDO-ATIENZA, K. LU, AND B. SCHMALFUSS, *Random dynamical systems for stochastic partial differentialequations driven by a fractional Brownian motion*, Discrete and Continuous Dynamical Systems-B, 14 (2010), pp. 473–493.
- [14] M. GARRIDO-ATIENZA, K. LU, AND B. SCHMALFUSS, *Unstable invariant manifolds for stochastic PDEs driven by a fractional Brownian motion*, Journal of Differential Equations, 248 (2010), pp. 1637–1667.
- [15] M. GARRIDO-ATIENZA, B. SCHMALFUSS, AND J. VALERO, *Setvalued dynamical systems for stochastic evolution equations driven by fractional noise*, Journal of Dynamics and Differential Equations, 34 (2022), pp. 79–105.
- [16] D. GIVON, *Strong convergence rate for two-time-scale jump-diffusion stochastic differential systems*, Multiscale Modeling & Simulation, 6 (2007), pp. 577–594.
- [17] M. HAIRER, *Ergodicity of stochastic differential equations driven by fractional Brownian motion*, The Annals of Probability, 33 (2005), pp. 703–758.
- [18] M. HAIRER AND X.-M. LI, *Averaging dynamics driven by fractional Brownian motion*, The Annals of Probability, 48 (2020), pp. 1826–1860.
- [19] K. HASSELMANN, *Stochastic climate models part I. theory*, Tellus, 28 (1976), pp. 473–485.
- [20] R. KHAMINSKII, *On an averaging principle for Itô stochastic differential equations*, Kibernetika, 4 (1968), pp. 260–279.
- [21] Y. KIFER, *Averaging and climate models*, Progress in Probability, 49 (2001), pp. 171–188.
- [22] C. KUEHN, *Multiple Time Scale Dynamics*, Springer Cham, 2015.
- [23] H. KUNITA, *Stochastic Flows and Stochastic Differential Equations*, Cambridge University Press, 1990.
- [24] X.-M. LI AND J. SIEBER, *Mild stochastic sewing lemma, SPDE in random environment, and fractional averaging*, Stochastics and Dynamics, 22 (2022), p. 2240025.
- [25] ———, *Slow-fast systems with fractional environment and dynamics*, The Annals of Applied Probability, 32 (2022), pp. 3964–4003.
- [26] W. LIU, M. RÖCKNER, X. SUN, AND Y. XIE, *Averaging principle for slow-fast stochastic differential equations with time dependent locally lipschitz coefficients*, Journal of Differential Equations, 268 (2020), pp. 2910–2948.
- [27] A. LUNARDI, *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, Springer Science & Business Media, 2012.
- [28] B. MASLOWSKI AND D. NUALART, *Evolution equations driven by a fractional Brownian motion*, Journal of Functional Analysis, 202 (2003), pp. 277–305.
- [29] A. NEAMȚU AND C. KUEHN, *Rough center manifolds*, SIAM Journal on Mathematical Analysis, 53 (2021), pp. 3912–3957.
- [30] G. PAVLIOTIS AND A. STUART, *Multiscale Methods: Averaging and Homogenization*, Springer Science & Business Media, 2008.
- [31] A. PAZY, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer Science & Business Media, 2012.
- [32] B. PEI, Y. INAHAMA, AND Y. XU, *Averaging principles for mixed fast-slow systems driven by fractional Brownian motion*, arXiv preprint arXiv:2001.06945, (2020).
- [33] ———, *Pathwise unique solutions and stochastic averaging for mixed stochastic partial differential equations driven by fractional Brownian motion and Brownian motion*, arXiv preprint arXiv:2004.05305, (2020).
- [34] ———, *Averaging principle for fast-slow system driven by mixed fractional Brownian rough path*, Journal of Differential Equations, 301 (2021), pp. 202–235.
- [35] B. PEI, Y. XU, AND G. YIN, *Stochastic averaging for a class of two-time-scale systems of stochastic partial differential equations*, Nonlinear Analysis, 160 (2017), pp. 159–176.
- [36] S. SAMKO, A. KILBAS, AND O. MARICHEV, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach, Switzerland, 1993.
- [37] V. VOLOSOV, *Averaging in systems of ordinary differential equations*, Russian Mathematical Surveys, 17 (1962), pp. 1–126.
- [38] Y. XU, J. DUAN, AND W. XU, *An averaging principle for stochastic dynamical systems with Lévy noise*, Physica D: Nonlinear Phenomena, 240 (2011), pp. 1395–1401.
- [39] Y. XU, B. PEI, AND R. GUO, *Stochastic averaging for slow-fast dynamical systems with fractional Brownian motion*, Discrete and Continuous Dynamical Systems-B, 20 (2015),

- pp. 2257–2267.
- [40] Y. XU, B. PEI, AND J.-L. WU, *Stochastic averaging principle for differential equations with non-lipschitz coefficients driven by fractional Brownian motion*, *Stochastics and Dynamics*, 17 (2017), p. 1750013.
- [41] M. ZÄHLE, *Integration with respect to fractal functions and stochastic calculus. I*, *Probability Theory and Related Fields*, 111 (1998), pp. 333–374.