

# V-MINIMAL SUBMANIFOLDS

MONICA ALICE APRODU

**ABSTRACT.** We introduce the notion of  $V$ -minimality, for  $V$  a smooth vector field on a Riemannian manifold, a natural extension of the classical notion of minimality, and we prove several basic properties. One featured example is given for locally conformal Kaehler (l.c.K) manifolds. It is well-known that in general, complex submanifolds in non-Kaehler l.c.K manifolds are not minimal. We prove that, however, they are  $V$ -minimal for  $V$  a suitable multiple of the Lee vector field. Extending some results from [1], to emphasis the utility of this notion, we prove that a PHH submersion is  $V$ -harmonic if and only if it has minimal fibres and a PHH  $V$ -harmonic submersion pulls back complex submanifolds to  $V$  minimal submanifolds.

## 1. INTRODUCTION

Let  $(M, g)$  and  $(N, h)$  be compact Riemannian manifolds,  $V$  a smooth vector field on  $M$  and  $\varphi : M \rightarrow N$  a smooth map. In [20], [5] the authors introduced the notion of  $V$ -harmonic map, see Definition 2.1, that naturally generalizes the classical notion of harmonic map. Unlike the latter,  $V$ -harmonicity is not defined via a variational problem, but rather by imposing the vanishing of a modified tension field, called the  $V$ -tension field of  $\varphi$ . If  $V = 0$ , the two notions coincide, more generally, the same is true if  $V$  is vertical.

In this generalized context,  $V$ -harmonic morphisms appear naturally, [20], and are directly connected to the minimality of the fibres, exactly as in the classical case, see Theorem 2.6 in [20].

In the case of Kähler target manifolds, harmonic morphisms generalize to pseudo-harmonic morphisms [13]. Furthermore, if a natural extra-condition that the naturally almost complex structure on the horizontal distribution satisfies a Kähler-type condition, then the harmonicity implies the minimality of the fibres. These maps, called pseudo-horizontally homothetic (PHH) enjoy other geometric properties, [1].

In the presence of a vector field  $V$  on a manifold  $M$ , one can naturally ask whether the definition of minimality can be perturbed to underline the influence of  $V$ . The main goal of this paper is to propose a natural definition which we call  $V$ -minimality, Definition 2.7. As for  $V$ -harmonicity, minimality corresponds to the case  $V = 0$ . One specific example is of particular interest. It is known that a complex submanifold  $K$  in a locally conformal Kähler manifold  $M$  is not minimal, unless the Lee vector field is tangent to

---

2020 *Mathematics Subject Classification.* 58E20,53C43,53C42, 32Q15.

*Key words and phrases.*  $V$ -harmonic maps,  $V$ -harmonic morphisms, PHWC maps, Kähler manifolds, locally conformal Kähler manifolds, minimal submanifolds.

This work was supported by Romanian Ministry of Education, Program PN-III, Project number PN-III-P4-ID-PCE-2020-0025, Contract 30/04.02.2021 and "Dunarea de Jos" University of Galati Grant, Project number RF 2488/31.05.2024.

the subvariety, Theorem 12.1, [9]. However, we notice that  $K$  is  $V$ -minimal, for  $V$  a suitable integer multiple of the Lee vector field of  $M$ , see Theorem 2.13. Furthermore, we connect  $V$ -minimality to PHH maps and their  $V$ -harmonicity.

The outline of the paper is as follows. Firstly, we recall the notions needed for our aim,  $V$ -harmonic maps, PHH and PHWC maps, l.c.K manifolds and we briefly review some of their properties. We then define  $V$ -minimal submanifolds, Definition 2.7 and  $V$ -pseudo harmonic morphisms, Definition 3.3. The main results of the paper are Theorem 2.13, Theorem 3.4, Theorem 3.6 and Theorem 3.7. Theorem 3.6 shows that for a PHH submersion,  $V$ -harmonicity is equivalent to having  $V$ -minimal fibres. In Theorem 3.7, we prove that a PHH  $V$ -harmonic submersion pulls back complex submanifolds to  $V$ -minimal submanifolds.

## 2. $V$ - HARMONIC MORPHISMS AND $V$ - MINIMAL SUBMANIFOLDS

**2.1.  $V$  - Harmonic Maps and Morphisms.** We remind the reader of some basic facts on  $V$ -Harmonic Maps and Morphisms that we shall use in the sequel [20],[5], [6], [7], [16].

**Definition 2.1** (see [5], [20]). Let  $(M^m, g)$  and  $(N^n, h)$  be two Riemannian manifolds,  $V$  a smooth vector field on  $M$ , and  $\varphi : M \rightarrow N$  a smooth map. The map  $\varphi$  is called  $V$ -harmonic if satisfies:

$$(1) \quad \tau_V(\varphi) := \tau(\varphi) + d\varphi(V) = 0,$$

where  $\tau(\varphi)$  is the tension field of the map  $\varphi$ . Since the differential  $d\varphi$  of  $\varphi$  can be viewed as a section of the bundle  $T^*M \otimes \varphi^{-1}TN$ ,  $d\varphi(V)$  is a section of the bundle  $\varphi^{-1}TN$ . The tension  $\tau_V(\varphi)$  is called the  $V$ -tension field of  $\varphi$ .

**Remark 2.2** (cf. [5], [20]). In particular,

1. For  $V = 0$ , the map  $\varphi$  is harmonic.
2. A smooth function  $f : M \rightarrow \mathbf{R}$  is said to be  $V$ -harmonic if:

$$\Delta_V(f) := \Delta(f) + \langle V, \nabla f \rangle = 0.$$

By taking the trace of the second fundamental form, we obtain the tension field of the map  $\varphi$ ,  $\tau(\varphi) = \text{trace} \nabla d\varphi$ , which is a section of  $\varphi^{-1}TN$ . In local coordinates  $(x_i)_{i=\overline{1,m}}$  on  $M$  and  $(y_\alpha)_{\alpha=\overline{1,n}}$  on  $N$ , respectively, it has the following expression (see for example [4], [10]):

$$(2) \quad \tau(\varphi) = \sum_{\alpha=1}^n \tau(\varphi)^\alpha \frac{\partial}{\partial y_\alpha},$$

where, denoting by  ${}^M\Gamma_{ij}^k$  and  ${}^N L_{\beta\gamma}^\alpha$  the Christoffel symbols of  $M$  and  $N$ , and  $\varphi^\alpha = \varphi \circ y_\alpha$ ,

$$(3) \quad \tau(\varphi)^\alpha = \sum_{i,j=1}^m g^{ij} \left( \frac{\partial^2 \varphi^\alpha}{\partial x_i \partial x_j} - \sum_{k=1}^m {}^M\Gamma_{ij}^k \frac{\partial \varphi^\alpha}{\partial x_k} + \sum_{\beta,\gamma=1}^n {}^N L_{\beta\gamma}^\alpha \frac{\partial \varphi^\beta}{\partial x_i} \frac{\partial \varphi^\gamma}{\partial x_j} \right).$$

For  $V$  a smooth vector field on  $M$ , given in local coordinates on  $M$  by  $V = \sum_{i=1}^m V_i \frac{\partial}{\partial x_i}$ , the  $V$ -tension field of the map  $\varphi$  has the following expression in local coordinates:

$$(4) \quad \tau_V(\varphi) = \sum_{\alpha=1}^n \tau(\varphi)^\alpha \frac{\partial}{\partial y_\alpha} + \sum_{\alpha=1}^n \sum_{i=1}^m V_i \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial}{\partial y_\alpha}.$$

Using the properties of the second fundamental form and of the tension field of the composition of two maps, (see [4]), we have the following lemma:

**Lemma 2.3** (see [20]). *The  $V$ -tension field of the composition of two maps  $\varphi : M \rightarrow N$  and  $\psi : N \rightarrow P$  is given by:*

$$(5) \quad \tau_V(\psi \circ \varphi) = d\psi(\tau_V(\varphi)) + \text{trace} \nabla d\psi(d\varphi, d\varphi).$$

As in the case of harmonic maps, Zhao (Definition 1.2 in [20]) defined  $V$ -harmonic morphisms as maps  $\varphi : M \rightarrow N$ , between Riemannian manifolds, which pulls back local harmonic functions on  $N$  to local  $V$ -harmonic functions on  $M$ .

The following results characterize the  $V$ -harmonic morphisms.

**Theorem 2.4** (see [20]). *Let  $\varphi : (M^m, g) \rightarrow (N^n, h)$  be a smooth map between Riemannian manifolds. Then the following conditions are equivalent:*

- 1)  $\varphi$  is a  $V$ -harmonic morphism;
- 2)  $\varphi$  is a horizontally weakly conformal  $V$ -harmonic map;
- 3)  $\forall \psi : W \rightarrow P$  a smooth map from an open subset  $W \subset N$  with  $\varphi^{-1}(W) \neq \emptyset$ , to a Riemann manifold  $P$ , we have:  $\tau_V(\psi \circ \varphi) = \lambda^2 \tau(\psi)$ , for some smooth function  $\lambda^2 : M \rightarrow [0, \infty)$ ;
- 4)  $\forall \psi : W \rightarrow P$  a harmonic map from an open subset  $W \subset N$  with  $\varphi^{-1}(W) \neq \emptyset$ , to a Riemann manifold  $P$ , the map  $\psi \circ \varphi$  is a  $V$ -harmonic map;
- 5)  $\exists \lambda^2 : M \rightarrow [0, \infty)$  a smooth function such that:  $\Delta_V(f \circ \varphi) = \lambda^2 \Delta f$ , for any function  $f$  defined on an open subset  $W$  of  $N$  with  $\varphi^{-1}(W) \neq \emptyset$ .

**Corollary 2.5** (see [20]). *For  $\varphi : M \rightarrow N$  a  $V$ -harmonic morphism with dilation  $\lambda$  and  $\psi : N \rightarrow P$  a harmonic morphism with dilation  $\theta$ , the composition  $\psi \circ \varphi$  is a  $V$ -harmonic morphism with dilation  $\lambda(\theta \circ \varphi)$ .*

**Theorem 2.6** (see [20]). *For a horizontally weakly conformal map  $\varphi : M \rightarrow N$  with dilation  $\lambda$ , any two of the following conditions imply the third:*

- 1)  $\varphi$  is a  $V$ -harmonic map (and so a  $V$ -harmonic morphism);
- 2)  $V + \nabla \log(\lambda^{2-n})$  is vertical at regular points;
- 3) the fibres of  $\varphi$  are minimal at regular points.

**2.2.  $V$  - minimal submanifolds.** Let  $(M, g)$  be a Riemannian manifold,  $V$  a smooth vector field on  $M$ , and  $K$  a submanifold of  $M$ . For any  $x \in K$ , we have the orthogonal decomposition of the tangent bundle  $T_x M = T_x K \oplus T_x K^\perp$ , with respect to  $g_x$ . According to this decomposition, we have

$$\nabla_X^M Y = \nabla_X^K Y + A(X, Y), \text{ for all } X, Y \in \Gamma(TK).$$

The symmetric bilinear map  $A : TK \times TK \rightarrow TK^\perp$  is the second fundamental form of the submanifold  $K$ .

**Definition 2.7.** The submanifold  $K$  of  $M$  is called  $V$ -minimal if

$$\text{trace}(A) - V \in \Gamma(TK).$$

**Remark 2.8.** Note that, as the trace of the second fundamental form has the image in the normal bundle, the  $V$ -minimality can be rephrased as

$$\text{trace}(A) - V^\perp = 0.$$

**Remark 2.9.** Assume  $\varphi : M \rightarrow N$  is a differentiable map and  $K$  is a fibre of  $\varphi$ . Then the  $V$ -minimality condition translates to:  $\text{trace}(A) - V$  is a vertical vector. We refer to the subsequent Proposition 2.11 for further aspects of  $V$ -minimality of the fibers.

**Proposition 2.10.** *Let  $K$  be a closed submanifold of a Riemann manifold  $(M, g)$ . Then there exists  $V$  a smooth vector field on  $M$ , such that  $K$  is  $V$ -minimal.*

*Proof.* Let  $A : TK \times TK \rightarrow TK^\perp$  be the second fundamental form of the submanifold  $K$ .

Consider on  $K$  the vector field  $\text{trace}(A)$ . It is known that it is a normal vector field on  $K$ . Using a known result (see [12] Lemma 8.6 and Exercise 8-15) there exists a smooth vector field  $V$  on  $M$ , such that  $V|_K = \text{trace}(A)$ . This implies that  $V|_K = V|_K^\perp$  and, moreover,  $\text{trace}(A) - V^\perp = 0$ . In conclusion,  $K$  is  $V$ -minimal.  $\square$

This proposition can be adapted to Riemann submersions such that the result holds true for a vector field  $V$  that is universal across all fibers, as follows.

Recall that (see[15]) for a Riemannian submersion  $\varphi : M \rightarrow N$ , can be defined two tensors, one of which is the second fundamental form of all the fibers.

If  $\mathcal{H}$  and  $\mathcal{V}$  denote the horizontal and the vertical distribution on  $M$ , then the second fundamental form of all fibers  $\varphi^{-1}(y)$ ,  $y \in N$ , gives rise to a (1,2)-tensor field on  $M$ , defined by:

$$T_X Y = \mathcal{H}\nabla_{\mathcal{V}X}^M(\mathcal{V}Y) + \mathcal{V}\nabla_{\mathcal{V}X}^M(\mathcal{H}Y),$$

for all arbitrary vector field  $X, Y$  in  $M$  and  $\nabla^M$  the covariant derivative on  $M$ .

It is known that the tensor  $T$  has the following properties:

- 1)  $T_X$  is a skew-symmetric linear operator on  $TM$  and it reverses the horizontal and vertical subspaces.
- 2)  $T_X = T_{\mathcal{V}X}$ , for all  $X$  in  $M$ .
- 3)  $T_X Y = T_Y X$ , for all  $X, Y$  vertical vector fields on  $M$ .
- 4) If  $X, Y$  are horizontal vector fields, and  $V, W$  are vertical vector fields on  $M$ , then:  $\nabla_V^M W = T_V W + \mathcal{V}\nabla_V^M W$  and  $\nabla_V^M X = \mathcal{H}\nabla_V^M X + T_V X$ .

In this context, we prove the following.

**Proposition 2.11.** *Let  $\varphi : M \rightarrow N$  be a Riemannian submersion. Then there exists a smooth vector field  $V$  on  $M$  such that any fiber of  $\varphi$  is  $V$ -minimal.*

*Proof.* Let us consider  $T_X Y = \mathcal{H}\nabla_{\mathcal{V}X}^M(\mathcal{V}Y) + \mathcal{V}\nabla_{\mathcal{V}X}^M(\mathcal{H}Y)$ , the (1,2)-tensor field on  $M$ , described above.

Choose the vector field  $V$  on  $M$ , to be the trace( $\mathcal{H}\nabla_{\mathcal{V}X}^M(\mathcal{V}Y)$ ).

For any  $y \in N$ , denote by  $K$  the fiber  $\varphi^{-1}(y)$ . Then,

$$V|_K = \text{trace}(\mathcal{H}\nabla_X^M Y), \forall X, Y \in \Gamma(TK).$$

But, the horizontal component of  $\nabla_X^M Y, \forall X, Y \in \Gamma(TK)$  is exactly the second fundamental form of  $K$  in  $M$ , so is equal to  $A(X, Y)$ .

Thus,  $\text{trace}(A) - V_{|K}^\perp = \text{trace}(A) - V_{|K} = 0$ , so the fiber  $K$  is  $V$ -minimal.  $\square$

This propositions provide a plethora of examples. However, the most interesting ones are obtained when  $V$  is a naturally defined vector field on  $M$ , as exemplified in the next subsection.

**2.3.  $V$ -minimality and locally conformal Kähler manifolds.** We recall here briefly the definitions of a locally conformal Kähler manifold and a locally conformal Kähler submersion.

**Definition 2.12.** ([9], [18]) A complex  $n$ -dimensional Hermitian manifold  $(M^{2n}, J, g)$ , where  $J$  denotes its complex structure and  $g$  its Hermitian metric is called a *locally conformal Kähler (l.c.K.) manifold* if there is an open cover  $\{U_i\}_{i \in I}$  of  $M$  and a family of  $C^\infty$  functions  $f_i : U_i \rightarrow \mathbf{R}, i \in I$ , such that each local metric:

$$(6) \quad g_i = \exp(-f_i)g|_{U_i}$$

is Kählerian, where  $g|_{U_i} = \iota_i^*$ , and  $\iota_i : U_i \rightarrow M$  is the inclusion.

The manifold is called *globally conformal Kähler (g.c.K) manifold* if there exist a  $C^\infty$  function  $f : M \rightarrow \mathbf{R}$ , such that the metric  $\exp(-f)g$  is Kähler.

Let  $\Omega, \Omega_i$  be the 2-forms associated with  $(J, g)$  and  $(J, g_i)$ , respectively ( $\Omega(\cdot, \cdot) = g(\cdot, J\cdot)$ ). Then (6) implies:

$$(7) \quad \Omega_i = \exp(-f_i)\Omega|_{U_i}.$$

A Hermitian manifold  $(M^{2n}, J, g)$  is l.c.K manifold if and only if there exists a globally defined closed 1-form  $\omega$  on  $M$ , such that  $d\Omega = \omega \wedge \Omega$  (see [9], Theorem 1.1). The closed 1-form  $\omega$  is called the *Lee form of the manifold*. When the manifold is g.c.K, the Lee form is exact and when the manifold is Kähler, the Lee form is zero.

The vector field  $B = \omega^\sharp$  on  $M$ , described by the condition by  $\omega(X) = g(X, B)$ , for all  $X$  in  $M$ , is called the *Lee vector field* of  $M$ .

A complex submanifolds of a l.c.K. manifold is seldom minimal, see Theorem 12.1, [9]. However, it is  $V$ -minimal for a suitable vector field  $V$ , as shown below.

**Theorem 2.13.** *With the notation as above, let  $K$  be a complex submanifold of a l.c.K manifold  $M$  of complex dimension  $m$ , and  $V$  a smooth vector field on  $M$ . Then  $K$  is a  $V$ -minimal submanifold on  $M$ , for  $V = -mB$ .*

*Proof.* Denote as usual by  $A$  the second fundamental form of the complex submanifold  $K \subset M$ . Using eq.(12.26), Theorem 12.1, [9], and choosing an orthonormal frame on  $K$ ,  $\{E_i, JE_i\}_{i=1, \dots, m}, m = \dim_{\mathbf{C}} K$ ,

$$\begin{aligned} \text{trace}(A) &= \sum_{i=1}^m (A(E_i, E_i) + A(JE_i, JE_i)) \\ (8) \quad &= -\sum_{i=1}^m g(E_i, E_i)B^\perp \\ &= -nB^\perp. \end{aligned}$$

If we choose the vector field on  $M$  to be  $V = -mB$ , then  $\text{trace}(A) - V = -m(B^\perp - B) \in \Gamma(TK)$ , which means that  $K \subset M$  is a  $V = -mB$  minimal submanifold.  $\square$

**Definition 2.14.** ([9], [14]) Let  $(M, J, g)$  and  $(M', J', g')$  be two almost Hermitian manifolds and  $\varphi : M \rightarrow M'$  a Riemannian submersion, which is holomorphic (i.e.  $d\varphi \circ J = J' \circ d\varphi$ ). Moreover, if  $(M, J, g)$  is a l.c.K manifold, then  $\varphi$  is called a *l.c.K submersion*.

Another example is obtained when considering l.c.K submersions, as proven in the following result.

**Proposition 2.15.** *Let  $\varphi : M \rightarrow M'$  a l.c.K submersion, and  $V$  a vector field on  $M$ , where  $M$  is compact. Then, the fibers of  $\varphi$  are  $V$ -minimal if and only if the Lee vector field  $B$  of  $M$  satisfies  $V + mB$  is a vertical vector field, where  $m$  is the dimension of the fiber.*

*Proof.* Let  $y \in M'$ , and denote by  $K$ , the fiber  $\varphi^{-1}(y)$  which is an immersed complex submanifold of  $M$  ( $K \xrightarrow{i} M$ ).

The l.c.K metric  $g$  on  $M$  induces a l.c.K metric  $g_K$  on  $K$  with the Lee form  $\omega_K = i^*\omega$ , and the Lee vector field  $B_K$  given by the decomposition  $B = B_K + B^\perp$ ,  $B_K \in \Gamma(TK)$ ,  $B^\perp \in \Gamma(TK^\perp)$ .

Using Gauss-Weingarten equation of the submanifold  $K$  we have:

$$\begin{aligned}\nabla_X^M B_K &= \nabla_X^K B_K + A(X, B_K), \\ \nabla_X^M B^\perp &= -\mathcal{A}_{B^\perp}(X) + D_X B^\perp,\end{aligned}$$

for all  $X \in \Gamma(TK)$ , where  $A$  is the second fundamental form of  $K$  in  $M$ ,  $D$  is the induced connection of the normal bundle of  $K$ , and  $-\mathcal{A}_{B^\perp}(X)$  is the tangential component of  $\nabla_X^M B^\perp$ . Moreover, for any  $X, Y \in \Gamma(TK)$ ,

$$g(\mathcal{A}_{B^\perp}(X), Y) = g(A(X, Y), B^\perp),$$

and the mean curvature vector of  $K$  is

$$(9) \quad \eta = \frac{1}{2m} \text{trace}(A).$$

If  $K$  is  $V$ -minimal, then  $\text{trace}(A) - V$  is a vertical vector field.

Keeping in mind that  $\text{trace}(A) \in \Gamma(TK^\perp)$ , and decompose  $V = V^v + V^\perp$  into tangent and normal component with respect to  $K$ , we have

$$(10) \quad \text{trace}(A) - V^\perp = 0.$$

As  $M$  is a l.c.K manifold, let us denote by  $\tilde{A}$  the (local) second fundamental form of  $K$  with respect to the local Kähler metric of the l.c.K structure on  $M$ , and  $\tilde{\eta}$  the corresponding mean curvature vector. Then, (see [9], Theorem 1.2 or [19] or [18]),  $\tilde{A}(X, Y) = A(X, Y) + \frac{1}{2}g(X, Y)B^\perp$ , for all  $X, Y \in \Gamma(TK)$ , and  $\tilde{\eta} = \eta + \frac{1}{2}B^\perp$ .

Since complex submanifolds of Kähler manifolds are minimal,  $\tilde{\eta} = 0$ , and consequently

$$(11) \quad \eta = -\frac{1}{2}B^\perp.$$

Using (9), (10), (11), we obtain:  $B^\perp = -\frac{1}{m}V^\perp$ , which implies that  $V + mB$  is a vertical vector field.

Conversely, if  $V + mB$  is a vertical vector field, then  $B^\perp = -\frac{1}{m}V^\perp$ . Since  $\tilde{\eta} = 0$ , and using (11), we get  $\eta = \frac{1}{2m}V^\perp$  and, from (9),  $\text{trace}(A) - V^\perp = 0$ . So  $K$  is  $V$ -minimal.  $\square$

**Remark 2.16.** Let  $\varphi : M \rightarrow M'$ ,  $M$  compact, a l.c.K submersion, and  $V$  a vector field on  $M$ . Denote by  $\omega, \omega'$  the Lee forms of  $M$  and  $M'$ , and by  $B$  and  $B'$  the corresponding Lee vector fields. From [9], Proposition 10.1,  $\omega(X) = \omega'(X') \circ \varphi$ , for any horizontal vector field  $X$  in  $M$ , which is  $\varphi$  related to  $X' \in \Gamma(TM')$ , i.e  $d\varphi(X) = X'$ , and  $\mathcal{H}(B)$  is a horizontal vector field  $\varphi$  related to  $B'$ , given by the decomposition  $B = \mathcal{V}(B) + \mathcal{H}(B)$  of  $B$  into vertical and horizontal component with respect to  $M$ .

If the fibers of  $\varphi$  are  $V$ -minimal, then  $V + mB$  is a vertical vector and  $d\varphi(B) = -\frac{1}{m}d\varphi(V)$ .

Then,  $-\frac{1}{m}d\varphi(V) = d\varphi(\mathcal{V}(B) + \mathcal{H}(B)) = d\varphi(\mathcal{H}(B)) = B'$ . So the Lie vector field on  $M'$  is  $-\frac{1}{m}d\varphi(V)$ .

### 3. V-PSEUDO HARMONIC MORPHISMS

#### 3.1. Pseudo Harmonic Morphisms and Pseudo Horizontally Homothetic Maps.

The notion of harmonic morphisms can be generalized when the target manifold is endowed with a Kähler structure (see [13], [8]).

Let us consider a smooth map  $\varphi : (M^m, g) \rightarrow (N^{2n}, J, h)$  from a Riemannian manifold to a Kähler manifold. The map  $\varphi$  is said to be a *pseudo-harmonic morphism* (shortening PHM) if and only if it pulls back local holomorphic functions on  $N$  to local harmonic maps from  $M$  to  $\mathbf{C}$ .

For any  $x \in M$ , denote by  $d\varphi_x^* : T_{\varphi(x)}N \rightarrow T_xM$  the adjoint map of the tangent linear map  $d\varphi_x : T_xM \rightarrow T_{\varphi(x)}N$ .

If  $X$  is a local section on the pull-back bundle  $\varphi^{-1}TN$ , then  $d\varphi^*(X)$  is a local horizontal vector field on  $M$ .

**Definition 3.1** (see [13]). The map  $\varphi$  is called *pseudo-horizontally (weakly) conformal (shortening PHWC)* at  $x \in M$  if  $[d\varphi_x \circ d\varphi_x^*, J] = 0$ .

The map  $\varphi$  is called *pseudo-horizontally (weakly) conformal* if it is pseudo-horizontally (weakly) conformal at every point of  $M$ .

Then, *pseudo-harmonic morphism* can also be characterised as harmonic, pseudo-horizontally (weakly) conformal maps (see [13], [8]).

The local description of PHWC condition is given by the following (see [13]): let  $(x_i)_{i=1, \dots, m}$  be real local coordinates on  $M$ ,  $(z_\alpha)_{\alpha=1, \dots, n}$  be complex local coordinates on  $N$ , and  $\varphi^\alpha = z_\alpha \circ \varphi$ ,  $\forall \alpha = 1, \dots, n$ . Then the PHWC condition for  $\varphi$  reads:

$$(12) \quad \sum_{i,j=1}^m g_M^{ij} \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} = 0$$

for all  $\alpha, \beta = 1, \dots, n$ .

A special class of pseudo-horizontally weakly conformal maps, are pseudo-horizontally homothetic maps.

**Definition 3.2** (see [1]). A map  $\varphi : (M^m, g) \rightarrow (N^{2n}, J, h)$  is called *pseudo-horizontally homothetic at  $x$*  (shortening PHH) if is PHWC at a point  $x \in M$  and satisfy:

$$(13) \quad d\varphi_x \left( (\nabla_v^M d\varphi^*(JY))_x \right) = J_{\varphi(x)} d\varphi_x \left( (\nabla_v^M d\varphi^*(Y))_x \right),$$

for any horizontal tangent vector  $v \in T_x M$  and any vector field  $Y$  locally defined on a neighbourhood of  $\varphi(x)$ .

A PHWC map  $\varphi$  is called *pseudo-horizontally homothetic*, if and only if

$$(14) \quad d\varphi(\nabla_X^M d\varphi^*(JY)) = Jd\varphi(\nabla_X^M d\varphi^*(Y)),$$

for any horizontal vector field  $X$  on  $M$  and any vector field  $Y$  on  $N$ .

The condition (14) is true for every horizontal vector field  $X$  on  $M$  and any section  $Y$  of  $\varphi^{-1}TN$ . So, instead of working with vector fields on  $N$ , we work with the larger space of sections in the pull back bundle  $\varphi^{-1}TN$ .

One of the basic properties of pseudo-horizontally homothetic maps (Proposition 3.3, [1]) shows that a PHH submersion is a harmonic map if and only if it has minimal fibres. Also, pseudo-horizontally homothetic maps are good tools to construct minimal submanifolds (Theorem 4.1, [1]).

**3.2.  $V$ -Pseudo Harmonic Morphisms.** Generalizing the class of harmonic maps and morphisms, respectively to  $V$ -harmonic maps and pseudo harmonic morphisms we obtain  $V$ -pseudo harmonic morphisms with a description similar to pseudo harmonic morphisms.

**Definition 3.3.** Let  $(M^m, g)$  be a Riemannian manifold of real dimension  $m$ ,  $(N^{2n}, J, h)$  a Hermitian manifold of complex dimension  $n$ ,  $\varphi : M \rightarrow N$  a smooth map and  $V$  a smooth vector field on  $M$ . The map  $\varphi$  is called  *$V$ -pseudo harmonic morphism* (shortening  $V$ -PHM) if  $\varphi$  is  *$V$ -harmonic* and *pseudo horizontally weakly conformal*.

The characterization of *pseudo harmonic morphism* given in [13] remain true in the general case of  *$V$ -harmonic* maps.

**Theorem 3.4.** Let  $\varphi : M \rightarrow N$  be a smooth map from a Riemannian manifold  $(M^m, g)$  to a Kähler one  $(N^{2n}, J, h)$  and  $V$  a smooth vector field on  $M$ . Then  $\varphi$  is  *$V$ -pseudo harmonic morphism ( $V$ -PHM)* if and only if it pulls back local complex valued holomorphic functions on  $N$  to local  $V$ -harmonic functions on  $M$ .

*Proof.* We adapt the proof of Proposition 2 of [13] to our context. Let us consider  $p \in M$  a given point,  $\{x_i\}_{i=1, \overline{m}}$  local coordinates at  $p$ ,  $\{z_\alpha\}_{\alpha=1, \overline{n}}$  local complex coordinates at  $\varphi(p) \in N$ , and  $\varphi^\alpha = z_\alpha \circ \varphi$ . By  ${}^M \Gamma_{ij}^k$  and  ${}^N L_{\beta\gamma}^\alpha$  are denoted the Christoffel symbols on  $M$  and  $N$ , respectively. The vector field  $V$ , in local coordinates in  $M$ , reads:  $V = \sum_{i=1}^m V_i \frac{\partial}{\partial x_i}$ .

Suppose that  $\varphi$  is  *$V$ -PHM* and  $f : N \rightarrow \mathbf{C}$  is a local holomorphic function on  $N$ .

Using the  $V$ -tension field of composition of two maps (Lemma 2.3) and the  $V$ -harmonicity of the map  $\varphi$  (Definition 2.1):  $\tau_V(f \circ \varphi) = \text{trace} \nabla df(d\varphi, d\varphi)$ .

In order to prove that  $\tau_V(f \circ \varphi) = 0$ , at point  $p$ , since  $d\varphi(\frac{\partial}{\partial x_i}) = \sum_{\alpha=1}^n \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial}{\partial z_\alpha}$ , we have:



$$(15) \quad \nabla df \left( d\varphi \left( \frac{\partial}{\partial x_i} \right), d\varphi \left( \frac{\partial}{\partial x_j} \right) \right) = \sum_{\alpha, \beta=1}^n \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} \nabla df \left( \frac{\partial}{\partial z_\alpha}, \frac{\partial}{\partial z_\beta} \right).$$

To compute  $\nabla df \left( \frac{\partial}{\partial z_\alpha}, \frac{\partial}{\partial z_\beta} \right)$ , first let us remark that  $f^{-1}T\mathbf{C} = N \times \mathbf{C}$  is the trivial vector bundle and the fibre  $(f^{-1}T\mathbf{C})_{\varphi(p)} = \mathbf{C}$ . The induced connection on the pull-back bundle  $f^{-1}T\mathbf{C}$  is defined by:

$$\nabla_X^f \sigma := X(\sigma), \forall \sigma \in \Gamma(f^{-1}T\mathbf{C}), (\sigma : N \rightarrow \mathbf{C} \text{ is a map})$$

So,

$$(16) \quad \nabla df \left( \frac{\partial}{\partial z_\alpha}, \frac{\partial}{\partial z_\beta} \right) = \nabla_{\frac{\partial}{\partial z_\alpha}}^f df \left( \frac{\partial}{\partial z_\beta} \right) - df \left( \nabla_{\frac{\partial}{\partial z_\alpha}}^N \frac{\partial}{\partial z_\beta} \right) = \frac{\partial^2 f}{\partial z_\alpha \partial z_\beta} - \sum_{\gamma=1}^n N L_{\alpha\beta}^\gamma \frac{\partial f}{\partial z_\gamma}$$

From the equations (15) and (16) we obtain:

$$\nabla df \left( d\varphi \left( \frac{\partial}{\partial x_i} \right), d\varphi \left( \frac{\partial}{\partial x_j} \right) \right) = \sum_{\alpha, \beta=1}^n \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} \frac{\partial^2 f}{\partial z_\alpha \partial z_\beta} - \sum_{\alpha, \beta, \gamma=1}^n \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} N L_{\alpha\beta}^\gamma \frac{\partial f}{\partial z_\gamma}$$

and

$$(17) \quad \begin{aligned} & \text{trace} \nabla df \left( d\varphi \left( \frac{\partial}{\partial x_i} \right), d\varphi \left( \frac{\partial}{\partial x_j} \right) \right) = \\ & = \sum_{\alpha, \beta=1}^n \left( \sum_{i, j=1}^m g^{ij} \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} \right) \frac{\partial^2 f}{\partial z_\alpha \partial z_\beta} - \sum_{\alpha, \beta, \gamma=1}^n \left( \sum_{i, j=1}^m g^{ij} \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} \right) N L_{\alpha\beta}^\gamma \frac{\partial f}{\partial z_\gamma} \end{aligned}$$

As  $\varphi$  is PHWC (see (12)), the term  $\sum_{i, j=1}^m g^{ij} \frac{\partial \varphi^s}{\partial x_i} \frac{\partial \varphi^k}{\partial x_j}$ , in relation (17), vanishes, and hence

$$\text{trace} \nabla df \left( d\varphi \left( \frac{\partial}{\partial x_i} \right), d\varphi \left( \frac{\partial}{\partial x_j} \right) \right) = 0.$$

Conversely, consider  $f : N \rightarrow \mathbf{C}$  a local complex holomorphic function,  $V$  a smooth vector field on  $M$ , and  $\varphi : M \rightarrow N$  a smooth map, such that  $\tau_V(f \circ \varphi) = 0$ .

Applying the chain rule for  $V$ -harmonic maps (Lemma 2.3), we get:

$$(18) \quad 0 = df(\tau_V(\varphi)) + \text{trace} \nabla df(d\varphi, d\varphi)$$

We compute the above equality in local coordinates, using the local description of the  $V$ -tension field (4) and the equation (17):

$$(19) \quad \begin{aligned} 0 &= \sum_{\alpha=1}^n \tau(\varphi)^\alpha \frac{\partial f}{\partial z_\alpha} + \sum_{\alpha=1}^n \sum_{i=1}^m V_i \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial f}{\partial z_\alpha} \\ &+ \sum_{\alpha, \beta=1}^n \left( \sum_{i, j=1}^m g^{ij} \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} \right) \frac{\partial^2 f}{\partial z_\alpha \partial z_\beta} - \sum_{\alpha, \beta, \gamma=1}^n \left( \sum_{i, j=1}^m g^{ij} \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} \right) N L_{\alpha\beta}^\gamma \frac{\partial f}{\partial z_\gamma} \end{aligned}$$

Choosing in (19) particular holomorphic functions  $f$ , for example  $f(z) = z_k, \forall k = \overline{1, n}$ , and normal coordinates centred at  $p$  (respectively, at  $\varphi(p)$ ), then  $\frac{\partial^2 f}{\partial z_\alpha \partial z_\beta} = 0$ ,  $\frac{\partial f}{\partial z_\gamma} = \delta_{k\gamma}$ , and the Christoffel symbols of  $N$  vanish. Then,

$\tau_V(f \circ \varphi) = 0$  is equivalent to:

$$\tau(\varphi)^k + \sum_{i=1}^m V_i \frac{\partial \varphi^k}{\partial x_i} = 0, \forall k = \overline{1, n},$$

where the term  $\tau(\varphi)^k + \sum_{i=1}^m V_i \frac{\partial \varphi^k}{\partial x_i} = 0$  is the  $k$  component of the  $\tau_V(\varphi)$ .

This proves that  $\varphi$  is a  $V$ -harmonic map. It follows that the equation (18) reduces to:  $\text{trace} \nabla df(d\varphi, d\varphi) = 0$ .

In this last equality, choosing  $f(z) = z_\alpha z_\beta, \forall \alpha, \beta = \overline{1, n}$  and normal coordinates in  $M$  and  $N$  respectively,  $\frac{\partial^2(z_\alpha z_\beta)}{\partial z_\alpha \partial z_\beta} = 1$ , and  ${}^N L_{\alpha\beta}^\gamma = 0$ , implies

$$\sum_{i,j=1}^m g^{ij} \frac{\partial \varphi^\alpha}{\partial x_i} \frac{\partial \varphi^\beta}{\partial x_j} = 0$$

which means that  $\varphi$  is PHWC.  $\square$

The following result gives a relation between  $V$ -harmonic morphisms, pseudo harmonic morphisms and  $V$ -pseudo harmonic morphisms, compare to Proposition 3 of [13].

**Proposition 3.5.** *Let  $(M, g)$  and  $(N, h)$  be two Riemannian manifolds,  $V$  a smooth vector field on  $M$ ,  $(P, J, p)$  a Kähler manifold and  $\psi : M \rightarrow N$  and  $\varphi : N \rightarrow P$  two smooth maps. If  $\psi$  is  $V$ -harmonic morphism and  $\varphi$  is pseudo harmonic morphism (PHM), then  $\varphi \circ \psi$  is  $V$ -pseudo harmonic morphism ( $V$ -PHM).*

*Proof.* As  $\varphi$  is a pseudo harmonic morphism (see Section 1.2), for any  $f : W \rightarrow \mathbf{C}$  a local complex valued holomorphic function defined in an open subset  $W \subset P$ , with  $\varphi^{-1}(W) \subset N$  non-empty,  $f \circ \varphi : \varphi^{-1}(W) \rightarrow \mathbf{C}$  is a local harmonic function on  $N$ .

Using Corollary 2.5 for the  $V$ -harmonic morphism  $\psi$  and the local harmonic function  $f \circ \varphi : \varphi^{-1}(W) \rightarrow \mathbf{C}$ , on  $N$ , we get  $f \circ \varphi \circ \psi : \psi^{-1}(\varphi^{-1}(W)) \rightarrow \mathbf{C}$ , a local  $V$ -harmonic function on  $M$ .

So,  $\varphi \circ \psi : M \rightarrow P$  pulls-back local holomorphic functions on  $P$  to local  $V$ -harmonic functions on  $M$ . By Theorem 3.4,  $\varphi \circ \psi$  is  $V$ -PHM.  $\square$

**Theorem 3.6.** *Let  $\varphi : (M^m, g) \rightarrow (N^{2n}, J, h), n \geq 2$ , be a pseudo horizontally homothetic submersion (PHH). Then  $\varphi$  is  $V$ -harmonic if and only if  $\varphi$  has  $V$ -minimal fibres.*

*Proof.* Similar to Proposition 3.3, [1], we can choose in the pull-back bundle  $\varphi^{-1}TN$  a local frame  $\{e_1, e_2, \dots, e_n, Je_1, Je_2, \dots, Je_n\}$  such that

$$\{d\varphi^*(e_1), \dots, d\varphi^*(e_n), d\varphi^*(Je_1), \dots, d\varphi^*(Je_n)\}$$

is an orthogonal frame in the horizontal distribution: choose  $e_1 \in \Gamma(\varphi^{-1}TN)$  a non-vanishing section and, using the PHWC property of  $\varphi$  and the fact that  $d\varphi^*(e_1)$  is an horizontal vector, we get the orthogonality of  $d\varphi^*(e_1)$  and  $d\varphi^*(Je_1)$ . At step  $k$ , take  $e_k$  orthogonal on both vectors  $d\varphi \circ d\varphi^*(e_i)$  and  $d\varphi \circ d\varphi^*(Je_i), \forall i \leq k-1$ .

If we denote by  $E_k = d\varphi^*(e_k), E'_k = d\varphi^*(Je_k)$ , then we have an orthogonal frame in the horizontal distribution  $\mathcal{H}(TM), \{E_1, E_2, \dots, E_n, E'_1, \dots, E'_n\}$ . Note that, by the PHWC condition of  $\varphi$ ,

$$(20) \quad g(E_i, E_i) = g(E'_i, E'_i).$$

It was proved in [1] that, the property of the induced connection  $\nabla^\varphi$  on  $\varphi^{-1}TN$  (Lemma 1.16, [17]) and the PHH condition, imply:

$$(21) \quad \nabla_{E'_i}^\varphi d\varphi(E'_i) = -\nabla_{E_i}^\varphi d\varphi(E_i) + Jd\varphi([E'_i, E_i])$$

and

$$(22) \quad \frac{1}{g(E_i, E_i)} Jd\varphi([E'_i, E_i]) = \frac{1}{g(E'_i, E'_i)} \nabla_{E'_i}^\varphi d\varphi(E'_i) + \frac{1}{g(E_i, E_i)} \nabla_{E_i}^\varphi d\varphi(E_i)$$

$$(23) \quad d\varphi(\nabla_{E'_i}^M E'_i) = -d\varphi(\nabla_{E_i}^M E_i) + Jd\varphi([E'_i, E_i]).$$

and

$$(24) \quad \frac{1}{g_M(E_i, E_i)} Jd\varphi([E'_i, E_i]) = \frac{1}{g_M(E_i, E_i)} d\varphi(\nabla_{E_i}^M E_i) + \frac{1}{g_M(E'_i, E'_i)} d\varphi(\nabla_{E'_i}^M E'_i).$$

We choose  $\{u_1, u_2, \dots, u_s\}$  an orthonormal basis for the vertical distribution  $\mathcal{V}(TM)$ . With this notation and using (21), (22), (23), (24), it was shown in Proposition 3.3. in [1], that

$$(25) \quad \tau(\varphi) = -d\varphi\left(\sum_{j=1}^s \nabla_{u_j}^M u_j\right).$$

If  $\varphi$  is a  $V$ -harmonic map, then  $\tau_V(\varphi) = 0$  and, from (25) we obtain:

$$0 = -d\varphi\left(\sum_{j=1}^s \nabla_{u_j}^M u_j\right) + d\varphi(V),$$

which is equivalent with the  $V$ -minimality of the fibres.

Conversely, suppose  $\varphi$  has  $V$ -minimal fibres. As  $\varphi$  is a PHH submersions, in the above constructed frame  $\{u_1, \dots, u_s, E_1, \dots, E_n, E'_1, \dots, E'_n\}$ , the  $V$ -tension field of  $\varphi$  reads:

$$\tau_V(\varphi) = -d\varphi\left(\sum_{j=1}^s \nabla_{u_j}^M u_j\right) + d\varphi(V)$$

The  $V$ -minimality of the fibres imply  $d\varphi(\text{trace}(A) - V) = 0$ .

Let us choose  $y \in N$ , and denote the fibre by  $K = \varphi^{-1}(y)$ .

Using (20), we have

$$\text{trace}(A) - V = \sum_{i=1}^s A(u_i, u_i) - V = \frac{1}{g_M(u_i, u_i)} \sum_{i=1}^s (\nabla_{u_i}^M u_i - \nabla_{u_i}^K u_i) - V$$

Since  $\nabla_{u_i}^K u_i$  is vertical,  $d\varphi(\nabla_{u_i}^K u_i) = 0$ , and

$$d\varphi(\text{trace}(A) - V) = \sum_{i=1}^s d\varphi(\nabla_{u_i}^M u_i) - d\varphi(V),$$

and hence  $\varphi$  is  $V$ -harmonic. □

The construction of minimal submanifolds was done for horizontally homothetic harmonic morphisms (see [3]) and generalised for the case of pseudo-horizontally homothetic harmonic submersions (see [1]). Replacing harmonicity by  $V$ -harmonicity, a similar result can be proved.

**Theorem 3.7.** *Let  $(M^m, g)$  be a Riemannian manifold,  $(N^{2n}, J, h)$  be a Kähler manifold,  $V$  a smooth vector field on  $M$  and  $\varphi : M \rightarrow N$  be a pseudo-horizontally homothetic (PHH),  $V$ -harmonic submersion.*

*If  $P^{2p} \subset N^{2n}$  is a complex submanifold of  $N$ , then  $\varphi^{-1}(P) \subset M$  is a  $V$ -minimal submanifold of  $M$ .*

*Proof.* We shall use an orthogonal local basis in the tangent bundle of  $\varphi^{-1}(P)$ , as in ([1]). Specifically, let us consider the decomposition of  $TM$  into the vertical distribution  $V$  and the horizontal one  $H$ ,  $T_x M = V_x \oplus H_x$ , for any point  $x \in M$ . Denote  $\varphi^{-1}(P)$ , by  $K$ ,  $H_1 = TK \cap H$  and, for any  $x \in K$ ,  $H_x = H_{1x} \oplus H_{2x}$ . Also,  $T_x K = V_x \oplus H_{1x}$ ,  $H_{1x} = \{u \in H_x, d\varphi_x(u) \in T_{\varphi(x)} P\}$ ,  $\forall x \in K$ .

We choose a linearly independent system of local sections  $\{e_1, \dots, e_p, Je_1, \dots, Je_p\}$  in  $\varphi^{-1}TN$ , such that, if we denote by  $E_i = d\varphi^*(e_i)$  and  $E'_i = d\varphi^*(Je_i)$ , the restriction of the system to  $K$  is a local orthonormal basis in  $H_1$ .

The system was chosen in the following way. Since  $d\varphi_{x|_{H_x}} \circ d\varphi_x^*$  is a linear isomorphism, take  $v_1$  a local section of  $\varphi^{-1}TP$ , non-vanishing along  $K$  and choose the section  $e_1$  of  $\varphi^{-1}TN$ , such that  $(d\varphi|_H \circ d\varphi^*)(e_1) = v_1$ . Using the PHWC condition of  $\varphi$ ,  $d\varphi(d\varphi^*(Je_1))$  is also a section of  $\varphi^{-1}TP$  and  $d\varphi^*(e_1)$  and  $d\varphi^*(Je_1)$  are orthogonal local vector fields in  $H$  and also in  $H_1$ . By induction, at step  $k$ , we take  $e_k$  perpendicular on both  $(d\varphi \circ d\varphi^*)(e_i)$  and  $(d\varphi \circ d\varphi^*)(Je_i)$ ,  $\forall 1 \leq i \leq k-1$ , such that  $(d\varphi \circ d\varphi^*)(e_i)$  is a section in  $\varphi^{-1}TP$ .

In the sequel, the computations are performed on points of  $K$  only.

Consider  $\{u_1, u_2, \dots, u_r\}$  an orthonormal local basis in the vertical distribution  $V$  such that  $\{u_1, u_2, \dots, u_r, E_1, \dots, E_p, E'_1, \dots, E'_p\}$  is an orthogonal local basis in  $TK$ . Denote by  $A$  the second fundamental form of the submanifold  $K$ .

The  $V$ -minimality of  $K$  is equivalent to  $\text{trace}(A) - V \in \Gamma(TK)$ , which reads:

$$\begin{aligned} & \sum_{i=1}^r \left[ \frac{1}{g_M(u_i, u_i)} d\varphi(\nabla_{u_i}^M u_i - \nabla_{u_i}^K u_i) \right] + \sum_{i=1}^p \left[ \frac{1}{g_M(E_i, E_i)} d\varphi(\nabla_{E_i}^M E_i - \nabla_{E_i}^K E_i) \right] \\ & + \sum_{i=1}^p \left[ \frac{1}{g_M(E'_i, E'_i)} d\varphi(\nabla_{E'_i}^M E'_i - \nabla_{E'_i}^K E'_i) \right] - d\varphi(V) \text{ is a section of } \varphi^{-1}TP. \end{aligned}$$

Using relations (20) and (23), we compute:

$$\begin{aligned} & \sum_{i=1}^p \left[ \frac{1}{g_M(u_i, u_i)} d\varphi(\nabla_{u_i}^M u_i - \nabla_{u_i}^K u_i) \right] + \sum_{i=1}^p \left[ \frac{1}{g_M(E_i, E_i)} d\varphi(\nabla_{E_i}^M E_i) + \frac{1}{g_M(E'_i, E'_i)} d\varphi(\nabla_{E'_i}^M E'_i) \right] \\ & - \sum_{i=1}^p \left[ \frac{1}{g_M(E_i, E_i)} (\nabla_{E_i}^K E_i) + \frac{1}{g_M(E'_i, E'_i)} (\nabla_{E'_i}^K E'_i) \right] - d\varphi(V) \\ & = \sum_{i=1}^p d\varphi(\nabla_{u_i}^M u_i) - \sum_{i=1}^p d\varphi(\nabla_{u_i}^K u_i) - d\varphi(V) \end{aligned}$$

As  $\varphi$  is a pseudo-horizontally homothetic  $V$ -harmonic submersion, from Theorem (3.6),  $\varphi$  has  $V$ -minimal fibres, which is equivalent to:  $d\varphi(\sum_{i=1}^p \nabla_{u_i}^M u_i) = d\varphi(V)$ .

Hence,  $\text{trace}(A) - V \in \Gamma(TK)$  is equivalent to  $\sum_{i=1}^p d\varphi(\nabla_{u_i}^K u_i)$  is a section of  $\varphi^{-1}TP$ .  $\square$

**Acknowledgements.** This work was partially supported by Romanian Ministry of Education, Program PN-III, Project number PN-III-P4-ID-PCE-2020-0025, Contract 30/04.02.2021 and by "Dunarea de Jos" University of Galati Grant, Project number RF 2488/31.05.2024.

#### REFERENCES

- [1] Aprodu Monica Alice, Aprodu Marian, Brânzănescu Vasile, *A Class of Harmonic Submersions and Minimal Submanifolds*, International Journal of Mathematics, Vol. 11, No. 9, (2000), 1177-1191.
- [2] Baird Paul, Eells James, *A conservation law for harmonic maps*, Geometry Symposium Utrecht 1980, Lecture Notes in Mathematics, 894 Springer-Verlag (1981), 1-25.
- [3] Baird Paul, Gudmundsson Sigmundur, *p-harmonic maps and minimal submanifolds*, Math. Ann, No. 294 (1992), 611 - 624.
- [4] Baird Paul, Wood John C., *Harmonic Morphisms Between Riemannian Manifolds*, Clarendon Press-Oxford (2003), ISBN 0-19-850362-8.
- [5] Chen Qun, Jost Jürgen, Wang Guofang, *A Maximum Principle for Generalizations of Harmonic Maps in Hermitian, Affine, Weyl, and Finsler Geometry*, J. Geom. Anal., No. 25 (2015), 2407-2426.
- [6] Chen Qun, Jost Jürgen, Qiu Hongbing, *Existence and Liouville theorems for V-harmonic maps from complete manifolds*, Ann. Global Anal. Geom. Vol. 42 (2012), No. 4, 565-584.
- [7] Chen Qun, Qiu Hongbing, *Rigidity of self-shrinkers and translating solitons of mean curvature flows*, Adv. Math. Vol. 294, (2016), 517-531.
- [8] Chen Jingyi, *Structures of Certain Harmonic Maps into Kähler Manifolds*, International Journal of Mathematics, Vol. 8, No. 5, (1997), 573-581.
- [9] Dragomir Sorin, Ornea Liviu, *Locally Conformal Kähler Geometry*, Progress in Mathematics 155, Birkhäuser Boston (1998), ISBN 978-1-4612-7387-5.
- [10] Eells James, Lemaire Luc, *A report on harmonic maps*, Bulletin of the London Mathematical Society 10(1) (1978), 1 - 68.
- [11] Hsiang Wu-Yi, Lawson H.Blaine, *Minimal submanifolds of low cohomogeneity*, J. Differential Geom. No. 5(1-2) (1971), 1-38.
- [12] Lee John M., *Introduction to Smooth Manifolds*, Springer, Graduate Texts in Mathematics 218 (2013), ISBN 978-1-4419-9981-8.
- [13] Loubeau Eric, *Pseudo Harmonic Morphisms*, International Journal of Mathematics, Vol. 8, No. 7, (1997), 943-957.
- [14] Marrero Juan Carlos, Rocha Juan, *Locally Conformal Kähler Submersions*, Geometriae Dedicata, Vol. 52, (1994), 271-289.
- [15] O'Neill Barret, *The fundamental equation of a submersion*, Michigan Math. J. Vol. 13, No. 4, (1966), 459-469.
- [16] Qiu Hongbing, *The heat flow of V-harmonic maps from complete manifolds into regular balls*, Proc. Amer. Math. Soc. Vol. 145, (2017), No. 5, 2271-2280.
- [17] Urakawa Hajime, *Calculus of Variations and Harmonic Maps*, Translations of Mathematical Monographs 132 AMS (1993), ISBN 0-8218-4581-0.
- [18] Vaisman Izu, *Generalized Hopf Manifolds*, Geometriae Dedicata, Vol. 13, (1982), 231-255.
- [19] Vaisman Izu, *On locally conformal almost Kähler manifolds*, Israel J.Math, Vol. 24, (1976), 338-351.
- [20] Zhao Guangwen, *V-Harmonic Morphisms Between Riemannian Manifolds*, Proc. Amer. Math. Soc., Vol.148, No. 3, (2020), 1351-1361.

- [21] White Brian, *The Space of Minimal Submanifolds for Varying Riemannian Metrics*, Indiana University Mathematics Journal, Vol. 40, No. 1 (1991), 161 - 200.
- [22] Wood John C., *Harmonic morphisms, foliations and Gauss map*, Contemp. Math.49 (1986), 145-184.

Department of Mathematics and Computer Science, Faculty of Sciences and Environmental,  
University "Dunărea de Jos" of Galați, 111 Științei Str., RO-800189, Galați, Romania.  
*Email address:* Monica.Aprodu@ugal.ro