COMPLEX COBORDISM MODULO c₁-SPHERICAL COBORDISM AND RELATED GENERA

M. BAKURADZE

Abstract. We prove that the ideal in complex cobordism ring \mathbf{MU}^* generated by the polynomial generators $S = (x_1, x_k, k \ge 3)$ of c_1 -spherical cobordism ring W^* , viewed as elements in \mathbf{MU}^* by forgetful map is prime. Using the Baas-Sullivan theory of cobordism with singularities we define a commutative complex oriented cohomology theory $\mathbf{MU}_S^*(-)$, complex cobordism modulo c_1 -spherical cobordism, with the coefficient ring \mathbf{MU}^*/S . Then any $\Sigma \subseteq S$ is also regular in \mathbf{MU}^* and therefore gives a multiplicative complex oriented cohomology theory $\mathbf{MU}_{\Sigma}^*(-)$. The generators of $W^*[1/2]$ can be specified in such a way that for $\Sigma = (x_k, k \ge 3)$ the corresponding cohomology is identical to the Abel cohomology, previously constructed in [14]. Another example corresponding to $\Sigma = (x_k, k \ge 5)$ gives the coefficient ring of the universal Buchstaber formal group law after tensored by $\mathbb{Z}[1/2]$, i.e., is identical to the scalar ring of the Krichever-Hoehn complex elliptic genus [18], [17].

1. INTRODUCTION AND STATEMENTS

The theory of complex cobordism $\mathbf{MU}^*(-)$ and special unitary cobordism $\mathbf{MSU}^*(-)$ play an important role in cobordism theory [25]. The ring \mathbf{MSU}^* is torsion free after localized away from 2

(1.1)
$$\mathbf{MSU}^*[1/2] = \mathbb{Z}[1/2][y_2, y_3, \cdots], \quad |y_i| = 2i.$$

With SU-structure forgetful inclusion in mind

$$\mathbf{MSU}^*[1/2] \subset \mathbf{MU}^*[1/2] = \mathbb{Z}[1/2][a_1, a_2, a_3, \cdots], \ |a_i| = 2i,$$

the generators y_i can be treated as elements in $\mathbf{MU}^*[1/2]$.

In particular y_i is a *SU*-manifold if and only if all Chern numbers of y_i having factor c_1 are zero. Then we have to check the main Chern number $s_i(y_i)$ for Novikov's criteria [23] (see (3.1) below) for the membership of the set of polynomial generators in **MSU**^{*}[1/2].

In [16] Conner and floyd introduced c_1 -spherical cobordism groups $W^{2n} \subset \mathbf{MU}^{2n}$. A complex cobordism class belongs to W^{2n} if and only if every Chern number involving c_1^2 vanishes.

 W^* is not a subring of \mathbf{MU}^* . However, with respect to some complex orientation and some * multiplication (see Section 2), the ring W^* is polynomial on generators in every positive even degree except 4, [15], [20]:

(1.2)
$$W^* = Z[x_1, x_k : k \ge 3], \ x_1 = \mathbf{CP}_1, \ x_k \in W_{2k}.$$

The polynomial generators x_k can be specified by the condition for the main Chern numbers $s_k(x_k) = d(k)d(k-1)$ for $k \ge 3$, where

$$d(k) = \begin{cases} p, & \text{if } k+1 = p^s \text{ for some prime } p, \\ 1, & \text{otherwise.} \end{cases}$$

²⁰¹⁰ Mathematics Subject Classification. 55N22; 55N35.

Key words and phrases. Complex bordism, SU-bordism, Formal group law, Complex elliptic genus.

The author was supported by EU fellowships for Georgian researchers, 2023 (57655523).

By [15] the cohomology theory $W^*(-)$ is complex oriented and (by Proposition 3.12) the ring $W^*[1/2]$ is generated by the coefficients of the corresponding formal group law F_W . In particular

(1.3) $W^*[1/2]$ is a free $MSU^*[1/2]$ module generated by 1 and $[CP_1]$.

Our main observation is that the sequence of polynomial generators of W^* , $(x_k, k \ge 3)$ in (1.2) viewed as elements in \mathbf{MU}^* by forgetful map, generate the same ideal in $\mathbf{MU}^*[1/2]$ as the coefficients $\alpha_{ij}, i, j \ge 2$ of the universal formal group law.

We prove in Section 3 one of our main results

Theorem 1.1. Let $S = (x_1, x_k, k \ge 3)$ be a sequence of polynomial generators of c_1 -spherical cobordism W^* . Then

i) S is regular in \mathbf{MU}^* .

ii) any subsequence Σ of S is regular in \mathbf{MU}^* .

Then the use of Baas-Sullivan theory of cobordism with singularities defines a multiplicative cohomology theory $\mathbf{MU}_{\Sigma}^{*}(-)$, with the scalar ring $\mathbf{MU}^{*}/(\Sigma)$.

The obstruction to commutativity in $\mathbf{MU}_{\Sigma}^{*}(-)$ studied in [21]. In some cases all obstructions vanish only for dimensional reasons. So is for $\Sigma = S$ in Theorem 1.1.

In Section 3 we define $\mathbf{MU}_{S}^{*}(-)$, the complex cobordism modulo c_{1} -spherical cobordism.

Theorem 1.2. Let $\Sigma = S$ in Theorem 1.1. There is a commutative complex oriented cohomology theory $\mathbf{MU}_{S}^{*}(-)$, with the coefficient ring \mathbf{MU}^{*}/S .

Let \mathbf{CP}_i be the cobordism class of complex projective space of dimension 2i and

$$F_U = \sum \alpha_{ij} x^i y^j$$

be the universal formal group law of complex cobordism. One can choose the following polynomial generators y_2, y_3 and y_4 in $\mathbf{MSU}^*[1/2]$ viewed as elements in $\mathbf{MU}^*[1/2]$ (see Section 3.5)

(1.4)
$$y_2 = \mathbf{CP}_2 - \frac{9}{8}\mathbf{CP}_1^2, \quad y_3 = -\alpha_{22}, \quad y_4 = -\alpha_{23} + \frac{3}{2}\alpha_{22}\mathbf{CP}_1.$$

We need the following result on quasi-toric representatives in the SU-bordism ring.

Theorem ([20], Thm 10.8.) There are quasi-toric SU-manifolds M^{2i} , $i \ge 5$, with $s_i(M^{2i}) = d(i)d(i-1)$ for odd i and $s_i(M^{2i}) = 2d(i)d(i-1)$ for even i, and they have the minimum possible characteristic numbers s_i and represent polynomial generators of $\mathbf{MSU}^*[1/2]$.

Let us specify the generators in (1.1) as

$$y_i = [M^{2i}], i \ge 5.$$

The motivation to this is that the Krichever complex elliptic genus (see [10], § E.5) vanishes on these generators.

We could also choose $y_i, i \ge 5$ based on the following result

Theorem ([26], Thm 6.1.) The kernel of the complex elliptic genus on $MSU^*[1/2]$ is equal to the ideal of *SU*-flops.

One interesting example of Theorem 1.1 is the Abel cohomology h_{Ab}^* constructed in [14].

Theorem 1.3. Let $\Sigma = (x_i, i \ge 3)$, where $x_i = y_i$ is specified above. Then $\mathbf{MU}_{\Sigma}^*(-)[1/2]$ is identical to the Abel cohomology $h_{Ab}^*[1/2]$.

Another example of Theorem 1.1 is related to the Buchstaber formal group law F_B [6], [7]. After tensored with rationals the corresponding classifying map f_B is identical [3] to Krichver-Hoehn complex elliptic genus [18], [17]. The scalar ring Λ_B of F_B is calculated in [11]. In particular, it has only 2-torsion and the quotient

$$\Lambda_{\mathcal{B}} = \Lambda_B / torsion$$

is an integral domain. We will consider this example in Section 3 and prove the following

Theorem 1.4. Let $\Sigma = (x_i, i \ge 5)$, where $x_i = y_i$ is specified above. Let Λ_B be the scalar ring of the universal Buchstaber formal group law. Then cohomology theory $\mathbf{MU}_{\Sigma}^*(-)[1/2]$ is multiplicative, commutative, complex oriented and has the coefficient ring identical to $\mathbf{MU}^*[1/2]/\Sigma := \Lambda_B[1/2]$.

In Section 4 we discuss a genus on c_1 -spherical cobordism W^*

(1.5)
$$\phi_W: W^*[1/2] \to [Z/2][x_1, x_3, x_4].$$

The following questions remains open.

What about integral versions of Theorem 1.3 and Theorem 1.4?

Is the cohomology theory $\mathbf{MU}^*/(x_i, i \ge 5)$ commutative? The same question arises for other examples of Theorem 1.1 ii).

2. Preliminaries

2.1. Krichever complex elliptic genus. The Krichever complex elliptic genus (also called Krichever-Höhn genus) has been defined in [18] and [17]. Krichever wrote down its characteristic power series Q(x) using Baker-Akhiezer function. In [17], Höhn defined four variable elliptic genus

(2.1)
$$\phi_{KH} : \mathbf{MU}^* \to \mathbb{Q}[p_1, p_2, p_3, p_4], \ |p_i = 2i|.$$

determined by the following property: if one denotes by f the exponent of the corresponding formal group F_{KH} , then the series

$$h(x) := \frac{f'(x)}{f(x)}$$

satisfies the differential equation

(2.2)
$$(h'(x))^2 = S(h(x)),$$

where S is the generic monic polynomial of degree 4:

$$S(t) = p_4 + p_3 t + p_2 t^2 + p_1 t^3 + t^4.$$

By definition [8] the universal Krichever formal group law F_K is of the form

(2.3)
$$F_K(x,y) = xw(y) + yw(x) - w'(0)xy + \frac{w(x)\beta(x) - w(y)\beta(y)}{xw(y) - yw(x)}x^2y^2,$$

where

$$w = \frac{\partial F_K(x,y)}{\partial x}(x,0)$$

is the invariant differential form of F_K and $\beta(x) = \frac{w'(x) - w'(0)}{2x}$.

The universal Krichever genus on ϕ_K on \mathbf{MU}^* is the homomorphism classifying the formal group law F_K .

Lemma 1. ([3], Thm 2.3) The exponent of F_K satisfies (2.3), i.e., ϕ_K is a specialization of ϕ_{KH} .

The universal Buchstaber formal group law F_B is defined [5] as a universal example among formal group laws of the form

$$\frac{x^2A(y) - y^2A(x)}{xB(y) - yB(x)},$$

where A(0) = B(0) = 1.

If A'(0) = B'(0) then B(x) is identical to the image of $\omega(x)$ under the classifying map f_B .

One can suggest kernel of ϕ_K directly in terms of the universal formal group law F_U as follows [4],[5]

Lemma 2. Let $A(x,y) = \sum A_{ij}x^iy^j$ be defined by

$$A(x,y) = F_U(x,y)(xw(y) - yw(x)), \quad where \quad w(x) = \frac{\partial F_U(x,y)}{\partial x}(x,0)$$

is the invariant differential form of F_U . Then F_k is identical to F_B and the classifying map $f_B = f_K$ is the obvious quotient map

$$\mathbf{MU}^* \to \mathbf{MU}^*/(A_{ij}, i, j \ge 3).$$

Remark 1. As shown in [11], all torsion elements of Λ_B , the scalar ring of F_B have order two. Furthermore, there is a set of multiplicative generators for Λ_B with one generator in every degree of the form p^k , where p is a prime and $k \ge 0$, and one generator in every degree $2^n - 2$, where $n \ge 3$; the latter generators generate the torsion subring.

2.2. c_1 -spherical cobordism. in [16] Conner and floyd introduced the groups $W^{2n} \subset \mathbf{MU}^{2n}$. A complex cobordism class belongs to W^{2n} if and only if every Chern number involving c_1^2 vanishes.

Recall the idempotent in [9], [15]

$$\pi_0: \mathbf{MU}^* \to \mathbf{MU}^*: \ \pi_0 = 1 + \sum_{k \ge 2} \alpha_{1k} \partial_k$$

and the projection $\pi_0 : \mathbf{MU}^* \to W^* = Im\pi_0$.

By [15] Proposition 2.15 the multiplication * in W^* is given by

$$a * b = \pi_0(a \cdot b) = ab + 2[V]\partial a\partial b,$$

where $[V] = \alpha_{12} \in \mathbf{MU}^4$ is the cobordism class $\mathbf{CP}_1^2 - \mathbf{CP}_2$.

Then W^* is a subgroup, not a subring in \mathbf{MU}^* . With respect to the multiplication * the ring W^* is polynomial on generators in every positive even degree except 4 as in (1.2).

2.3. The Abel formal group law. By definition [12] the universal Abel formal group law formal group is the universal example of the formal group laws of the form

$$F_{Ab} = x + y + \alpha_{11}xy + \sum_{i \ge 2} \alpha_{1i}(xy^i + x^iy)$$

By Quillen [24] and Novikov [23] the scalar ring of the universal Lazard formal group law is identical to \mathbf{MU}^* , the ring of complex cobordism. So to put it another way F_{Ab} is classified from the universal formal group law F_U over \mathbf{MU}^*

$$F_U = x + y + \sum \alpha_{ij} x^i y^j$$

by the quotient of \mathbf{MU}^* modulo the ideal I_{Ab} generated by all coefficients $\alpha_{ij} \in \mathbf{MU}^*$, but only $\alpha_{1i} = \alpha_{i1}$

(2.4)
$$I_{Ab} = (\alpha_{ij} : i, j \ge 2).$$

Buchstaber named the group law F_{Ab} after Abel, motivated by Abel's article: "Abel N. H. Methode generale pour trouver des fonctions d'une seule quantite variable lorsqu'une propriete de ces fonctions est exprimee par une equation entre deux variables oeuvres completes, I. Christiania, 1881. p. 1-10."

In that article, Abel gave a general solution of a functional equation for the exponent series of F_{Ab} .

Remark 2. For our purposes, let us recall from [12] that the scalar ring F_{Ab} , i.e.,

$$\Lambda_{Ab} = \mathbf{MU}^* / I_{Ab}$$

is non polynomial infinitely generated ring. There is a set of multiplicative generators for $\Lambda_{Ab} \otimes \mathbb{F}_2$ with one generator in every complex degree 2^k , $k \ge 0$.

 Λ_{Ab} is an integral domain. After tensored with rational numbers it is a subring of the polynomial ring in two parameters of degree 2 and 4, i.e.,

(2.5)
$$\Lambda_{Ab} \otimes \mathbb{Q} \subset \mathbb{Q}[\alpha, \beta], \ |\alpha| = 2, |\beta| = 4.$$

in [14] proved that F_{Ab} can be realized by commutative complex oriented cohomology theory h_{Ab}^* also called after Abel.

We will also need

Lemma 3. The classifying map $f_{Ab} : \mathbf{MU}^* \to \Lambda_{Ab}$ followed by $\Lambda_{Ab} \otimes \mathbb{Q}$ in is a specialization of the Krichever-Hoehn complex elliptic genus ϕ_{KH} .

Proof. A direct proof of this fact could include a verification of the defining property (2.3) of the Krichever-Höhn genus in case of F_{Ab} . Leaving this to the reader, we derive the lemma from the fact that F_{Ab} is obviously a specialization of F_K as $w(x) = 1 + \sum_{\geq 1} \alpha_{1i} x^i$ and ϕ_K is a specialization of ϕ_{KH} by Lemma 1.

2.4. Some auxiliary combinatorial definitions. Before we continue about the construction of the Abel cohomology, we need to recall some combinatorial facts.

By Euclid's algorithm for the natural numbers m_1, m_2, \cdots, m_k one can find integers $\lambda_1, \lambda_2, \cdots, \lambda_k$ such that

(2.6)
$$\lambda_1 m_1 + \lambda_2 m_2 + \dots + \lambda_k m_k = gcd(m_1, m_2, \dots, m_k).$$

Let

(2.7)
$$d(m) = gcd\left\{ \binom{m+1}{1}, \binom{m+1}{2}, \cdots, \binom{m+1}{m-1} | m \ge 1 \right\}.$$

By [19] one has

$$d(m) = \begin{cases} p, & \text{if } m+1 = p^s \text{ for some prime } p, \\ 1, & \text{otherwise} \end{cases}$$

therefore the elements

(2.8)
$$e_m = \lambda_1 \alpha_{1\,m} + \lambda_2 \alpha_{2\,m-1} + \dots + \lambda_m \alpha_{m\,1}$$

may be chosen as the polynomial generators in \mathbf{MU}^* .

Let $m \ge 4$ and let $\lambda_2, \cdots, \lambda_{m-2}$ are such integers that

(2.9)
$$d_2(m) := \sum_{i=2}^{m-2} \lambda_i \binom{m+1}{i} = gcd\left\{\binom{m+1}{2}, \cdots, \binom{m+1}{m-2}\right\}$$

Then by [11] Lemma 9.7 one has for $m \ge 3$

(2.10)
$$d_2(m) = d(m)d(m-1).$$

Let us apply Euclid's algorithm for the Chern numbers

$$s_{m-1}(\alpha_{i,m-i}) = \binom{m+1}{i}$$

in (2.9) and let

$$z_k = \sum_{i=2}^{k-1} \lambda_i \alpha_{i\,k+1-i}, \ k \ge 3.$$

Let us recall from [14] the constriction of abelian cohomology $h_{Ab}^*(-)$.

Lemma 4. Let I_{Ab} be as in (2.4) and $I_{Ab}(l) \subset I_{Ab}$ be generated by those elements of \mathbf{MU}^* whose degree is less or equal 2l. Let $I = (z_3, \cdots)$ and $I(l) = (z_3, \cdots, z_l)$. Then

i)
$$I_{Ab} = I;$$

ii) $I_{Ab}(l) = I(l).$

Let's put $\mathbf{MU}^* = \mathbb{Z}[e_1, e_2, \cdots]$ for e_m in (2.8) and let

$$A_l = \mathbb{Z}[e_1, e_2, \cdots, e_l]$$
 and $A^{l+1} = \mathbb{Z}[e_{l+1}, e_{l+2}, \cdots],$

i.e.,

$$\mathbf{MU}^* = A_l \otimes A^{l+1}.$$

The preimage of I(l) by obvious inclusion defines the ideal of A_l denoted by same symbol so that

(2.11)
$$\mathbf{MU}^*/I(l) = A_l/I(l) \otimes A^{l+1}.$$

Remark 3. We note here that by definition I(l) is generated by those polynomials in $\mathbb{Z}[e_1, e_2, \cdots, e_l]$ that are in the kernel of the quotient map $\mathbf{MU}^* \to \mathbf{MU}_*/I_{Ab}$.

The construction of abelian cohomology is based on the following key result [12], [14].

Lemma 5. The ideals I and I(l) are prime, or equivalently \mathbf{MU}^*/I and $\mathbf{MU}^*/I(l)$ are integral domains.

Using Baas-Sullivan theory the commutative complex oriented cohomology $h_{Ab}^*(-)$ with coefficient ring \mathbf{MU}^*/I was defined in [14].

Remark 4. We recall that for a commutative ring R, a regular sequence is a sequence (r_1, \dots, r_n) in R such that r_i is not a zero-divisor on $R/(r_1, \dots, r_{i-1})$, for $i = 1, \dots, n$.

If R is a graded ring and the members of a regular sequence are homogeneous of positive degree, then any permutation of a regular sequence is regular [13]. Therefore we can regard our regular sequences as regular sets.

3. Proofs

Lemma 6. Let J_W be the ideal in \mathbf{MU}^* generated by $y_2 = w_1 * w_1 = 8\mathbf{CP}_2 - 9\mathbf{CP}_1^2$ and polynomial generators $(x_i, i \ge 3)$ of W^* as elements in \mathbf{MU}_* by group inclusion and $J_W(l)$ be (y_2, x_3, \dots, x_l) . Let I and I(l) be the ideals in Lemma 4, $\tilde{I} = (y_2, I)$ and $\tilde{I}(l) = (y_2, I(l))$. Then

i) $J_W(l) = \tilde{I}(l), \forall l \ge 3;$

ii) The sequence y_2, x_3, \cdots is regular in \mathbf{MU}^* .

Proof. i) First let us prove $\tilde{J}_W \subset \tilde{I}$. We have $I = I_{AB}$ by Lemma 4. Also $\Lambda_{Ab} = \mathbf{MU}^*/I$ is an integral domain by Lemma 5. Moreover $\mathbf{MU}^*/I \otimes \mathbb{Q}$ is polynomial ring in two parameters $\mathbf{CP}_1, \mathbf{CP}_2$ by (3). Therefore it suffices to check $x_k \in \tilde{I}$ viewed as elements in $\mathbf{MU}^*[1/2]$.

For the latter we need two facts.

By SU representatives in Introduction $J_{SU} \subset I$, where J_{SU} is the ideal in $\mathbf{MU}^*[1/2]$ generated by y_5, y_6, \cdots , special unitary flops in $\mathbf{MSU}^*[1/2]$ of dimensions ≥ 10 . These generators vanish under the Krichever genus and therefore vanish under its specialization, the classifying map of the Abel formal group law. Then $y_3, y_4 \in I$ by (1.4) therefore $(y_2, y_3, y_4, J_{SU}) \subset \tilde{I}$.

Then by [15] $W^*[1/2]$ is a free $\mathbf{MSU}^*[1/2]$ module generated by 1 and \mathbf{CP}_1 . So after inverting 2 each generator $x_k, k \leq l$, of $J_W(l)$ as element in $\mathbf{MU}^*[1/2]$ is in kernel of the quotient map to the integral domain $\mathbf{MU}^*/(I(l), x_2)$. Thus $x_k \in \tilde{I}(l)$ and $J_W \subset \tilde{I}$ as the ideals in \mathbf{MU}^* . Here we use Remark 4 for $\tilde{I}(l)$.

Now let's prove $I(l) \subset J_W(l), \forall l \geq 2$ by induction on l. For l = 2 and l = 3 this is true because $y_2 = 8\mathbf{CP}_2 - 9\mathbf{CP}_1^2$ and $x_3 = y_3 = z_3 = -\alpha_{22}$. Let $I(l-1) \subset J_W(l-1)$ by induction hypothesis. To prove $z_l \in (J_W(l-1), x_l)$ recall that the main Chern number $s_l(z_l - x_l) = 0$ by the definitions of z_l, x_l , i.e., (1.2) and (2.10). This means that $z_l - x_l = P(e_1, e_2, \dots, e_{l-1})$ as decomposable element in \mathbf{MU}^* . Here e_i are as in (2.8). On the other hand $x_l \in J_W(l) \subset \tilde{I}(l)$ hence $P = x_l - z_l$ is in the kernel of the quotient map $\mathbf{MU}^* \to \mathbf{MU}^*/\tilde{I}(l-1)$, i.e. $P \in \tilde{I}(l-1) = J_W(l-1)$. The induction hypothesis and Remark 3 completes the proof.

ii) By [14] again I is prime ideal, therefore so is $\tilde{I} = J_W$ by Remark 3.

3.1. **Proof of Theorem 1.1.** i) As $x_3, \dots, x_l \subset J_W(l)$ by Lemma 6 ii) and Remark 4 the ring $R = \mathbf{MU}^*/(x_3, \dots, x_l)$ is an integral domain. In particular, the multiplication by $x_1 = \mathbf{CP}_1$ is mono in R. By Remark 4 again (x_1, x_3, \dots, x_l) is regular in $\mathbf{MU}^* \forall l$ and one has ii).

3.2. **Proof of Theorem 1.3.** We use Lemma 6 and apply the specified generators x_3, \cdots in $W^*[1/2]$ such that $(x_3, \cdots) = (z_3, \cdots)$ in $\mathbf{MU}^*[1/2]$.

 $\overline{7}$

3.3. **Proof of Theorem 1.2.** We have to prove that all obstructions to commutativity vanish. This follows by dimension argument and definition of obstruction elements in [21]. In particular, the obstruction elements to commutativity $\tilde{\mathcal{P}}_2(x_i)$ defined by certain cohomology operation [21] Theorem 3.4

$$ilde{\mathcal{P}}_2: \mathbf{MU}^{-2n} o \mathbf{MU}^{-4n-2} \otimes \mathbb{F}_2$$

So that all obstruction elements are of odd complex dimension -2i - 1. Therefore they vanish in the scalar ring $\mathbf{MU}^*/(x_1, x_3, \cdots) \otimes \mathbb{F}_2$ which is evenly generated as a quotient of Λ_{Ab} by Remark 2.

Alternatively, we can use Busato's long proof [14] of commutativity of $h_{Ab}^*(-)$. The proof is that $\tilde{\mathcal{P}}_2(\alpha_{ij})$ projects to 0 in $\Lambda_{Ab} \otimes \mathbb{F}_2$, for α_{ij} , $i, j \geq 2$. In our case, we have to consider one more element $\alpha_{11} = x_1$. By [21] Theorem 3.4 we have $\tilde{\mathcal{P}}_2(\alpha_{11}) = \alpha_{22} = x_3$ modulo α_{11} . So that the obstruction element $\tilde{\mathcal{P}}_2(x_1)$ vanishes.

3.4. **Proof of Theorem 1.4.** One can use the fact that $\mathbf{MSU}^*[1/2]$ is a subring in $W^*[1/2]$ and follow [2] where we proved the statement for generators $(y_i, i \ge 5)$ in $\mathbf{MSU}^*[1/2]$. The scalar ring Λ_B of F_B and its classifying map f_B is calculated in [11] (see also Remark 1). In particular, Λ_B has the ideal of elements of order two generated by $f_B(e_{2^n-2}), n \ge 3$. Here e_i are as in (2.8). The quotient by torsion

$$\Lambda_{\mathcal{B}} = \Lambda_B / (f_B(e_{2^n - 2}), \ n \ge 3).$$

is an integral domain.

Remark 5. Using the arguments in the proof of Theorem 1.2 one can prove that for Σ containing all generators x_n of W^* with odd indices $n = 2i - 1 \ge 1$, the cohomology theory h_{Σ}^* is commutative.

3.5. **Proof of** (1.4). We have to check that y_2, y_3, y_4 are represented by SUmanifolds i.e., all Chern numbers of y_i having factor c_1 are zero. Then we have to check the main Chern number $s_i(y_i)$ for Novikov's criteria [23] for the membership of the set of polynomial generators in $\mathbf{MSU}^*[1/2]$.

In particular, SU manifold x_n is a polynomial generator if and only if

(3.1)
$$s_n(y_n) = \begin{cases} \pm 2^k p & \text{if } n = p^l, \ p \text{ is odd prime,} \\ \pm 2^k p & \text{if } n+1 = p^l, \ p \text{ is odd prime,} \\ \pm 2^k & \text{otherwise.} \end{cases}$$

Rewrite the elements y_i in \mathbf{MU}^* as follows

$$y_{2} = \mathbf{CP}_{2} - \frac{9}{8}\mathbf{CP}_{1}^{2};$$

$$2y_{3} = -3\mathbf{CP}_{3} + 8\mathbf{CP}_{1}\mathbf{CP}_{2} - 5\mathbf{CP}_{1}^{3};$$

$$y_{4} = -2\mathbf{CP}_{1}^{4} + 7\mathbf{CP}_{1}^{2}\mathbf{CP}_{2} - 3\mathbf{CP}_{2}^{2} - 4\mathbf{CP}_{1}\mathbf{CP}_{3} + 2\mathbf{CP}_{4}.$$

The proof follows from the following calculations for the main Chern numbers.

$$c_1c_1[\mathbf{CP}_1^2] = 8, \quad c_1c_1[\mathbf{CP}_2] = 9, \quad c_2[\mathbf{CP}_1^2] = 4, \quad c_2[\mathbf{CP}_2] = 3.$$

This implies that that $c_1c_1(y_2) = 0$. There are no more Chern numbers involving c_1 and $s_2(y_2) = 3$. Therefore y_2 forms a generator of $\mathbf{MSU}^4[1/2]$.

For y_3 :

X	$c_3(X)$	$c_1c_2(X)$	$c_1c_1c_1(X)$
\mathbf{CP}_3	4	24	64
$\mathbf{CP}_1\mathbf{CP}_2$	6	24	54
\mathbf{CP}_1^3	8	24	48.

It follows all Chern numbers of y_3 involving c_1 are zero. Then y_3 forms a generator in $\mathbf{MSU}^6[1/2]$ as $s_3(y_3) = -2 \cdot 3$ satisfies the Novikov criteria. Similarly for y_4 : we have $s_4(y_4) = 2 \cdot 5$ and

X	$c_1c_1c_1c_1(X)$	$c_1c_1c_2(X)$	$c_1c_3(X)$
\mathbf{CP}_1^4	384	192	64
$\mathbf{CP}_1^2\mathbf{CP}_2$	432	204	60
\mathbf{CP}_2^2	486	216	54
$\mathbf{CP}_1\mathbf{CP}_3$	512	224	56
\mathbf{CP}_4	625	250	50.

4. c_1 -spherical cobordism modulo quasi-toric manifolds

Recall again from [26] that in special unitary cobordism $MSU^*[1/2]$ one can construct polynomial generators y_n , $n \ge 5$ using Euclid's algorithm and SU-flops or equivalently generators specified in Introduction.

The restriction of the Krichever complex elliptic genus

$$\mathbf{MU}^* \to \mathbb{Q}[p_1, p_2, p_3, p_4]$$

on $\mathbf{MSU}^*[1/2]$ gives the genus

$$\mathbf{MSU}^*[1/2] \to \mathbb{Z}[1/2][y_2, y_3, y_4],$$

as it is zero on all $y_i, i \ge 5$.

As $W^*[1/2]$ is a free $\mathbf{MSU}^*[1/2]$ module generated by 1 and $x_1 = \mathbf{CP}_1$ one has the following

Theorem 4.1. SU-flops of dimension ≥ 10 as elements in the ring $W^*[1/2]$ with * product generate the kernel ideal of the genus

$$\phi_W: W^*[1/2] \to \mathbb{Z}[1/2][x_1, x_3, x_4].$$

References

- 1. N. Baas, On bordism theory of manifolds with singularities, Math. Scand., 33(1973), 279–302.
- 2. M. Bakuradze, Polynomial generators of MSU^{*}[1/2] related to classifying maps of certain formal group laws, Homology, Homotopy and Applications, to appear.
- M. Bakuradze, V.V. Vershinin On addition theorems related to elliptic integrals, Proc. Steklov Math. Inst., 305(2019), 22-32.
- M. Bakuradze, On the Buchstaber formal group law and some related genera, Proc. Steklov Math. Inst., 286(2014), 7-21.
- M. Bakuradze, Formal group laws by Buchstaber, Krichever and Nadiradze coincide, Russian Math. Surveys, 68:3 (2013), 571–573.
- 6. M. Bakuradze, Computing the Krichever genus, J. Homotopy Relat. Struct, 9, 1(2014), 85-93.
- M. Bakuradze, Cohomological realization of the Buchstaber formal group law, Russian Math. Surv., 77:5 (2022), 943-945.

- V. M. Buchstaber, E. Yu. Netay, CP(2)-multiplicative Hirzebruch genera and elliptic cohomology, Russian Math. Surveys, 69:4 (2014), 757-759.
- V. M. Buchstaber, Projectors in unitary cobordisms that are related to SU-theory, Uspekhi Mat. Nauk, 27:6(168)(1972), 231-232.
- V. M. Buchstaber and T. E. Panov, *Toric topology*, Math. Surveys Monogr., vol. 204, Amer. Math. Soc., Providence, RI 2015, xiv+518 pp.
- V. Buchstaber, K. Ustinov, Coefficient rings of Buchstaber formal group laws, Math. Notices, 206:11(2015), 19-60.
- Bukhshtaber V.M., Kholodov A.N., Formal groups, functional equations, and generalized cohomology theories, (English. Russian original), Math. USSR Sbornik 69:1 (1991), 77–97.
- 13. W. Bruns, J. Herzog, Cohen-Macaulay rings, Cambridge University Press, (1993).
- 14. Ph. Busato, *Realization of Abel's universal formal group law*, Math. Z., **239**(2002), 527–561.
- G. Chernykh, T. Panov, SU-linear operations in complex cobordism and the c₁-spherical bordism theory, Izvestiya Ross. Akad. Nauk Ser. Mat. 87 (2023), no. 4.
- 16. P.E Conner, E.E. Floyd Torsion in SU-bordism, Mem. Amer. Math. Soc. 60 (1966).
- G. Höhn, Komplexe elliptische Geschlechter und S¹-äquivariante Kobordismustheorie, 2004, https://arxiv.org/abs/math/0405232
- I. Krichever, Generalized elliptic genera and Baker-Akhiezer functions, Math. Notes, 47(1990), 132–142.
- E. E. Kummer, Uber die Erganzungssatze zu den allgemeinen Reciprocitatsgesetzen, J. Reine Angew. Math., 44 (1852), 93–146.
- I. Yu. Limonchenko, T. E. Panov, and G. S. Chernykh SU-bordism: structure results and geometric representatives, Russian Math. Surv., 74:3(2019), 461–524.
- O. K. Mironov, Multiplications in cobordism theories with singularities, and Steenrod tom Dieck operations, Math. USSR Izvestija 13(1979), 89–106;
- S. P. Novikov, Homotopy properties of Thom complexes (Russian), Mat. Sb. 57(1962), 407–442.
- S. P. Novikov, The methods of algebraic topology from the viewpoint of cobordism theory (in Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* **31** (1967), no. 4, 855–951; translation in *Math. USSR-Izv.* **1** (1967), no. 4, 827–913.
- D. Quillen, On the formal group laws of unoriented and complex cobordism theory, Bull. Amer. Math. Soc. 75 (1969), 1293–1298.
- R. E. Stong, Notes On Cobordism Theorey, Princeton University Press and University of Tokyo Press, (1968)
- B. Totaro, Chern numbers for singular varieties and elliptic homology, Annals of Mathematics, 151(2000), 757–792.

FACULTY OF EXACT AND NATURAL SCIENCES, A. RAZMADZE MATH. INSTITUTE, IV. JAVAKHISHVILI TBILISI STATE UNIVERSITY, GEORGIA

Email address: malkhaz.bakuradze@tsu.ge

10