# Calculus of Fractions for Quasicategories

Daniel Carranza

Krzysztof Kapulkin

Zachery Lindsey

June 6, 2023

#### Abstract

We describe a generalization of Gabriel and Zisman's Calculus of Fractions to quasicategories, showing that the two essentially coincide for the nerve of a category. We then prove that the marked Ex-functor can be used to compute the localization of a marked quasicategory satisfying our condition and that the appropriate (co)completeness properties of the quasicategory carry over to its localization. Finally, we present an application of these results to discrete homotopy theory.

# Contents

Prelim	linaries	
1	Classical calculus of fractions	4
2	Quasicategories	7
3	Marked simplicial sets and localization	10
$(\infty, 1)$ -0	calculus of fractions	
4	$(\infty, 1)$ -calculus of fractions — definition $\ldots \ldots \ldots$	14
5	$(\infty, 1)$ -calculus of fractions vs. classical calculus of fractions $\ldots \ldots \ldots \ldots \ldots \ldots \ldots$	17
6	Examples of calculus of fractions	20
Model	for the localization	
7	The marked Ex functor	29
8	Localization at equivalences via marked Ex	33
9	Localization via marked Ex	35
Proper	rties of the localization	
10	Mapping spaces in the localization	41
11	(Co)limits in the localization	49
Applic	ation to discrete homotopy theory	
12	Background on discrete homotopy theory	54
13	Finite limits in $N_{\Box}$ Graph	57
14	$(\infty, 1)$ -categorical setting for discrete homotopy theory	60
Appen	dices	
Α	Cubical sets	62
В	Cubical categories	65

2020 Mathematics Subject Classification: 18N55, 18N60, 18N40, 55U35, 18N50.

## Introduction

The localization of a category at a class of maps (usually thought of as some sort of 'weak equivalences') refers to a universal or free way of inverting these maps in the category. Computing it is in general hard, as the morphisms in the localization are classes of finite zigzags, e.g.,  $\cdot \leftarrow \cdot \rightarrow \cdot \leftarrow \cdots \rightarrow \cdot$  where the 'backwards' maps are weak equivalences. It is perhaps not surprising that even if the category in question is locally small, its localization need not be. Likewise, it is unreasonable to expect the localization to possess any interesting categorical structure, like limits or colimits, and indeed very few examples of localizations do, even when the category being localized is itself complete and cocomplete.

Introduced in [GZ67], calculus of fractions is a set of conditions on a marked category, i.e., a pair  $(\mathcal{C}, W)$  consisting of a category and a class of maps, which simplifies this task. Shall  $(\mathcal{C}, W)$  satisfy (or admit) calculus of fractions, the set of maps between any two objects can be described using zigzags of length 2 instead of general zigzags. In addition, several (co)completeness properties can be carried over from  $\mathcal{C}$  to its localization. As such, calculus of fractions is a powerful tool for computing localization, see, e.g., [Mar83]. We should note here that there are two sets of conditions one could impose: calculus of *left* fractions and calculus of *right* fractions. Throughout this introduction, we speak of the former, but all statements have dual versions, given in the paper, involving the latter.

In this paper, we present a generalization of the calculus of fractions to the theory of  $(\infty, 1)$ -categories, taken here to be quasicategories. Given a marked quasicategory  $(\mathcal{C}, W)$ , we can express our conditions as a certain lifting property against a family of poset inclusions of a half-cube into a whole cube. This resembles the familiar condition from the usual calculus of fractions [GZ67, Def. 2.2] which asks that every (co)span be completed to a square.

As a test case, we consider quasicategories arising as nerves of categories and show that the nerve of a marked category  $(\mathcal{C}, W)$  satisfies (our quasicategorical) calculus of fractions exactly when  $(\mathcal{C}, W)$  satisfies proper calculus of fractions. The properness condition is a mild strengthening of the definition found in [GZ67] and is satisfied by most interesting examples. In particular, it is implied by the 2-out-of-3 property.

As indicated above, the key result proven in [GZ67, Ch. 1] is that if  $(\mathcal{C}, W)$  satisfies calculus of left fractions, then the localization of  $\mathcal{C}$  at W is given by the category of left fractions  $W^{-1}\mathcal{C}$ , i.e., its objects are zigzags  $\cdot \to \cdot \stackrel{\sim}{\leftarrow} \cdot$  of length 2. Our replacement of it is the marked Ex-functor. Recall that the Ex-functor [Kan57] is a functor on simplicial sets given by  $(\text{Ex } X)_n = \mathsf{sSet}(\mathsf{sd}[n], X)$ , where  $\mathsf{sd}[n]$  is the poset of non-empty subsets of the linear order  $[n] = \{0 \leq 1 \leq \ldots \leq n\}$  ordered by inclusion. We 'upgrade' it to a functor  $\text{Ex}_+$  from marked simplicial sets to simplicial sets by equipping  $\mathsf{sd}[n]$  with a marking: a 1-simplex  $A_0 \subseteq A_1$  is marked if and only if  $\max A_0 = \max A_1$ . (We note that the marked sd functor was previously used by Szumiło [Szu17] to prove that the homotopy theories of cofibration categories and cocomplete quasicategories are equivalent.)

With this definition, we can state our first main theorem:

**Theorem** (cf. Theorems 7.5 and 9.6). If a marked quasicategory  $(\mathcal{C}, W)$  satisfies calculus of left fractions, then  $\operatorname{Ex}_+(\mathcal{C}, W)$  is a quasicategory and the canonical map  $\max^* \colon \mathcal{C} \to \operatorname{Ex}_+(\mathcal{C}, W)$ , induced by  $\max \colon \operatorname{sd}[n] \to [n]$ , is the localization of  $\mathcal{C}$  at W.

We note here that our calculus of fractions condition is satisfied by a wide range of marked quasicategories. For example, all reflective localizations (referred to simply as localizations in [Lur09, Def. 5.2.7.2]) satisfy calculus of fractions.

We then generalize another theorem of Gabriel and Zisman's, namely that if a marked category  $(\mathcal{C}, W)$  satisfies calculus of left fractions, then the canonical functor from  $\mathcal{C}$  to its localization preserves finite colimits, and moreover if  $\mathcal{C}$  is cocomplete, then so is its localization. In [GZ67], this is proven by establishing that the hom-sets in the category of fractions are filtered colimits of those in  $\mathcal{C}$ . The same line of argument applies in our setting, although the proofs are less straightforward:

**Theorem** (cf. Corollaries 10.9 and 11.5 and Theorem 11.7). Suppose that a marked quasicategory  $(\mathcal{C}, W)$  satisfies calculus of left fractions and W is closed under 2-out-of-3.

- 1. The mapping space in  $\text{Ex}_+(\mathcal{C}, W)$  from x to y is a filtered colimit of mapping spaces from x to y', indexed by marked 1-simplices  $y \xrightarrow{\sim} y'$ .
- 2. If C is finitely cocomplete, then so is  $Ex_+(C, W)$ . Moreover,  $max^* \colon C \to Ex_+(C, W)$  preserves finite colimits.

We conclude the paper with an application of our results to discrete homotopy theory, a young area of mathematics which uses ideas from homotopy theory to study combinatorics (cf. [BL05]). Our main application is in establishing the relevant  $(\infty, 1)$ -category of graphs, which is done using calculus of right fractions. We can then prove that the graph nerve functor constructed in [CK22a] ascends to a representable functor from the quasicategory of graphs to that of spaces. This allows us to state a conjecture:

**Conjecture** (cf. Conjecture 14.3). The simplicial map induced by the graph nerve functor is an equivalence of quasicategories, thus establishing an equivalence between the discrete and classical homotopy theory.

Our quasicategorical setting seems to provide the appropriate tools to prove this conjecture.

In order to establish our desired application, we make extensive use of cubical sets, cubical categories, and the homotopy coherent nerve (both cubical [KV20] and simplicial [Lur09, Ch. 1]). Many results pertaining to these objects might be well-known in folklore or unsurprising to say the least, however precise references are hard to find. We therefore collect the relevant facts about them in an appendix. Two results may be of independent interest:

Theorem (cf. Theorem B.16, Proposition B.17, and Theorem B.20).

- 1. The category  $Cat_{\Box}$  of cubical categories carries a Bergner-style model structure which is Quillen equivalent to the Joyal model structure on simplicial sets.
- The quasicategory given by the cubical homotopy coherent nerve of the cubical category of cubical Kan complexes N<sub>□</sub>Kan<sub>□</sub> is equivalent in the Joyal model structure to the homotopy coherent nerve of the simplicial category of (simplicial) Kan complexes N<sub>Δ</sub>Kan<sub>Δ</sub>, thus making the former a model for spaces.

**Relation to** [Cis19]. Ideas similar to those of Section 10 are considered in [Cis19, §7.2]. One of the main results of Section 10 is that if  $(\mathcal{C}, W)$  satisfies calculus of (right) fractions then the mapping spaces in the localization may be computed as a filtered colimit of mapping spaces in  $\mathcal{C}$ , indexed by the "marked slice" over the domain (as indicated above). For us, this is a not-so-easy consequence of the definition of the marked Ex-functor. In [Cis19, Thm. 7.2.8], an analogous colimit formula (which is no longer filtered in general) is proven for any sufficiently well-behaved model of the marked slice. Moreover in [Cis19, Rem. 7.2.10], a model for this colimit is given whose 0-simplices are spans/right fractions, just as in the mapping spaces of our marked Ex (though the two do not coincide in general). In the language of [Cis19, Def. 7.2.6], a well-behaved model for the marked slice at an object x is called a *calculus of right fractions* at x.

**Comments on style.** As mentioned before, calculus of fractions appears in two forms: calculus of left fractions and calculus of right fractions. For the readers' convenience, we give most statements in both forms; the only exceptions being technical lemmas that are only needed for a proof of another theorem and that we do not expect to be of independent interest.

In addition, we assume that the reader is familiar with the basics of simplicial homotopy theory [GJ99] and the main ideas of quasicategory theory [Cis19, Joy09, Lur09]. While we have included some background material on these in Section 2, it is rather brief. We try to give precise references, notably

to [Cis19, Joy02, Joy09, Lur09] for all facts used in the paper, but we realize that this does not replace a comprehensive introduction.

**Organization of the paper.** In the first three sections, we collect the necessary background on calculus of fractions (Section 1), following [GZ67], quasicategories (Section 2), and marked simplicial sets (Section 3). As indicated above, throughout the paper, we assume familiarity with foundational notions and results of simplicial homotopy theory, so this review is rather brief and intended mostly to establish the necessary notation. We then define calculus of fractions for quasicategories (Section 4), compare it to the classical version (Section 5), and provide a wide range of examples (Section 6). We then move towards describing the localizations of marked quasicategories admitting a calculus of fractions. In Section 7, we define the marked Ex-functor and show that it takes marked quasicategories satisfying calculus of fractions to quasicategories. Before showing that it computes the localization of an arbitrary marked quasicategory in Section 9, we first prove (Section 8) this result when the quasicategory is marked at equivalences. We then turn our attention to properties of the resulting localization. In Section 10, we give a formula for the mapping spaces in the localization as a filtered colimit of those in the quasicategory itself; and in Section 11, we prove results about existence of finite (co)limits in the localization as well as preservation thereof by the localization functor. We then apply our results to construct the quasicategory of graphs and its 'global sections' functor to spaces, which we conjecture to be an equivalence of quasicategories. We are not assuming prior familiarity with discrete homotopy theory, and hence we begin this part with an introduction to the field in Section 12. In Section 13, we show that the naive localization of the category of graphs (at homotopy equivalences) has all finite limits. This result along with those proven in previous sections allows us then to construct the quasicategory associated to discrete homotopy theory in Section 14. In the last three sections, we make heavy use of some constructions on and facts about cubical sets and cubical categories, not all of which we were able to find in the literature. For this reason, we have collected them in Appendices A and B, respectively.

Acknowledgements. This paper grew out of the third author's Ph.D. thesis written in 2018 at Indiana University under the supervision of Mike Mandell. The three of us are deeply grateful to Mike for suggesting this research direction to us. We would also like to thank Andrew Blumberg and Mike Mandell for writing [BM19], which provided the initial motivation and an application of some of the results found in this paper.

This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930 while the first two authors participated in a program hosted by the Mathematical Sciences Research Institute in Berkeley, California, during the 2022–23 academic year.

#### Preliminaries

#### 1 Classical calculus of fractions

In this section, we recall the classical results on calculus of fractions following [GZ67, Ch. 1]. The main results of interest are Theorems 1.7 and 1.8. These results both have analogues in the quasicategorical case and our exposition in this section is intended to reflect the structure of the paper going forward. In particular, the analogue of Theorem 1.7 is proven in Section 9, and the analogue of Theorem 1.8 is proven in Sections 10 and 11.

We first define marked categories, which package the data of a category with the collection of morphisms one wishes to localize at.

**Definition 1.1.** A marked category is a pair  $(\mathcal{C}, W)$  where  $\mathcal{C}$  is a category and W is a collection of morphisms in  $\mathcal{C}$ .

In a marked category  $(\mathcal{C}, W)$ , we write either  $x \xrightarrow{\sim} y$  or  $x \xrightarrow{\sim} y$  to indicate that the morphism  $x \to y$ is in W. For a category  $\mathcal{C}$ , we write  $\mathcal{C}^{\natural}$  for the pair  $\mathcal{C}$  marked at all isomorphisms of  $\mathcal{C}$ . Given marked categories  $(\mathcal{C}, W), (\mathcal{C}', W')$ , we write  $\mathsf{Cat}_+((\mathcal{C}, W), (\mathcal{C}', W'))$  for the full subcategory of the functor category from  $\mathcal{C}$  to  $\mathcal{D}$  consisting of functors which sends morphisms in W to morphisms in W'.

We recall the definition of the localization of C at a class W.

**Definition 1.2.** Let  $(\mathcal{C}, W)$  be a marked category. The *localization* of  $\mathcal{C}$  at W is a functor

$$\gamma\colon \mathcal{C}\to \mathcal{C}[W^{-1}]$$

into a category  $\mathcal{C}[W^{-1}]$  such that

- the functor  $\gamma$  sends morphisms in W to isomorphisms in  $\mathcal{C}[W^{-1}]$ ; and
- for any category  $\mathcal{D}$ , the pre-composition functor

$$\gamma^* \colon \mathsf{Cat}(\mathcal{C}[W^{-1}], \mathcal{D}) \to \mathsf{Cat}_+((\mathcal{C}, W), \mathcal{D}^{\natural})$$

is an equivalence of categories.

In [GZ67], a construction is given for the localization of a pair  $(\mathcal{C}, W)$  which satisfies *calculus of fractions*. This condition allows one to construct a workable model of the localization in which morphisms between objects become (co)spans of morphisms rather than arbitrary zigzags.

**Definition 1.3.** A marked category  $(\mathcal{C}, W)$  satisfies *calculus of left fractions* (or *CLF*) if the following conditions hold:

- 1. W is closed under composition and contains identities;
- 2. any span  $(f: X \to Y, w: X \to X')$  with  $w \in W$  can be completed to a commutative square

$$\begin{array}{ccc} X & \xrightarrow{w} & X \\ f \downarrow & & \downarrow \\ Y & \xrightarrow{w'} & Y \end{array}$$

with  $w' \in W$ ;

3. given parallel morphisms  $f, g: X \to Y$ , if there exists  $w: X' \to X$  in W such that fw = gw then there exists  $v: Y \to Y'$  in W such that vf = vg (as in the diagram)

$$X' \xrightarrow{w} X \xrightarrow{f} Y \xrightarrow{v} Y'$$

We also introduce a slight strengthening of calculus of fractions.

**Definition 1.4.** A marked category  $(\mathcal{C}, W)$  satisfies *proper CLF* if it satisfies conditions (1) and (3) of Definition 1.3, as well as the following strengthening of condition (2):

2'. Any span  $(f: X \to Y, w: X \to X')$  with  $w \in W$  can be completed to a commutative square

$$\begin{array}{ccc} X & \xrightarrow{w} & X' \\ f \downarrow & & \downarrow f' \\ Y & \xrightarrow{w'} & Y' \end{array}$$

with  $w' \in W$ . Moreover, if f is in W then so is f'.

**Definition 1.5.** A marked category  $(\mathcal{C}, W)$  satisfies calculus of right fractions (or CRF) if  $(\mathcal{C}^{op}, W)$  satisfies CLF. It satisfies proper CRF if  $(\mathcal{C}^{op}, W)$  satisfies proper CLF.

In Section 4, we will introduce the notion of calculus of fractions for quasicategories. When viewing an ordinary category as a quasicategory, this notion will agree with proper CLF (respectively, proper CRF) but not CLF (respectively, CRF).

**Remark 1.6.** Most examples which satisfy CLF (or CRF) also satisfy proper CLF (respectively, proper CRF). For instance, if W is closed under 2-out-of-3 then the two are equivalent. The diagram



depicts an example of a marked category which satisfies CLF but not proper CLF.

We describe the model of the localization constructed in [GZ67] for a pair  $(\mathcal{C}, W)$  satisfying CLF.

Define a category  $CW^{-1}$  whose objects are those of C. The set of morphisms from x to y in  $CW^{-1}$  is the set of cospans

$$x \xrightarrow{f} y' \xleftarrow{w} y$$

where w is a weak equivalence, subject to the equivalence relation defined by  $(f, w) \sim (g, v)$  if there exists a commutative diagram



In particular, conditions (2) and (3) of CLF (Definition 1.3) are required to define composition of morphisms. With this, there is an evident functor  $\mathcal{C} \to \mathcal{C}W^{-1}$  which is identity on objects and which sends a morphism f to the equivalence class of the co-span  $(f, \mathrm{id})$ .

Dually, if  $(\mathcal{C}, W)$  satisfies CRF then there is an analogous construction  $W^{-1}\mathcal{C}$  whose morphisms are given by spans.

**Theorem 1.7** ([GZ67, Prop. 2.4]). Let  $(\mathcal{C}, W)$  be a marked category.

- 1. If  $(\mathcal{C}, W)$  satisfies calculus of left fractions then the functor  $\mathcal{C} \to \mathcal{C}W^{-1}$  is the localization of  $\mathcal{C}$  at W.
- 2. If  $(\mathcal{C}, W)$  satisfies calculus of right fractions then the functor  $\mathcal{C} \to W^{-1}\mathcal{C}$  is the localization of  $\mathcal{C}$  at W.

Moreover, under certain conditions on  $(\mathcal{C}, W)$ , the category  $\mathcal{C}W^{-1}$  is locally small and admits finite colimits (dually, the category  $W^{-1}\mathcal{C}$  is locally small and admits finite limits). In the following statement, we write  $x \downarrow W$  (dually  $W \downarrow x$ ) for the full subcategory of the slice category  $x \downarrow \mathcal{C}$  (dually  $\mathcal{C} \downarrow x$ ) comprised of weak equivalences  $x \xrightarrow{\sim} x'$  (dually  $x' \xrightarrow{\sim} x$ ).

**Theorem 1.8** ([GZ67, Prop. 3.1 & Cor. 3.2]). Let x, y be objects in a locally small marked category  $(\mathcal{C}, W)$ .

- 1. Suppose  $(\mathcal{C}, W)$  satisfies CLF and that  $y \downarrow W$  is finally small. Then,
  - the set of maps in  $CW^{-1}$  from x to y is computed by the filtered colimit

$$\mathcal{C}W^{-1}(x,y) \cong \operatorname{colim}_{y' \in y \downarrow W} \mathcal{C}(x,y');$$

- the functor  $\mathcal{C} \to \mathcal{C}W^{-1}$  preserves finite colimits; and
- if C admits all finite colimits then so does  $CW^{-1}$ .

2. Suppose  $(\mathcal{C}, W)$  satisfies CRF and that  $W \downarrow x$  is finally small. Then,

• the set of maps in  $W^{-1}\mathcal{C}$  from x to y is computed by the filtered colimit

$$\mathcal{C}W^{-1}(x,y) \cong \operatorname{colim}_{x' \in (W \downarrow x)^{\operatorname{op}}} \mathcal{C}(x',y);$$

- the functor  $\mathcal{C} \to W^{-1}\mathcal{C}$  preserves finite limits; and
- if C admits all finite limits then so does  $W^{-1}C$ .

#### 2 Quasicategories

In this section, we review aspects of quasicategory theory such as: detecting equivalences, the Joyal model structure, joins, slices, mapping spaces, and (co)limits. Throughout, we will assume familiarity with the basics of simplicial homotopy theory, including the Kan–Quillen model structure.

We write sSet for the category of simplicial sets. Arbitrary simplicial sets will typically be denoted with capital letters e.g.  $X, Y, \ldots$  We follow standard notation, including writing [n] for the linear order  $\{0 \le 1 \le \cdots \le n\}$  and  $\Delta$  for the category of finite non-empty linear orders. We write

$$\partial_i \colon [n-1] \to [n], \quad \sigma_i \colon [n] \to [n-1]$$

for the face and degeneracy poset maps, respectively, omitting the data of the domain and codomain from our notation. We write  $\Delta^n$  for the representable *n*-simplex,  $\partial\Delta^n$  for its boundary, and  $\Lambda^n_k$  for the (n, k)horn. The action of simplicial operators will be written on the right, e.g.  $x\sigma_0$  denotes the 0-degeneracy of an *n*-simplex  $x \in X_n$ .

**Definition 2.1.** A quasicategory is a simplicial set  $\mathcal{C}$  with the right lifting property against the set

$$\{\Lambda_k^n \hookrightarrow \Delta^n \mid n \ge 2, \ 0 < k < n\}.$$

**Example 2.2.** If C is a category then its nerve  $NC \in \mathsf{sSet}$  is a quasicategory.

We adopt the convention of referring to 0-simplices in a quasicategory as *objects* and 1-simplices in a quasicategory as *morphisms*.

Let  $E^1$  denote the nerve of the contractible category with two objects 0, 1.

**Definition 2.3.** A 1-simplex f in a simplicial set X is an  $E^1$ -equivalence if, as a map  $f: \Delta^1 \to X$ , it admits a lift to a map  $E^1 \to X$ .

$$\begin{array}{ccc} \Delta^1 & \stackrel{f}{\longrightarrow} X \\ \downarrow & & \\ E^1 \end{array}$$

**Example 2.4.** If C is a category then an  $E^1$ -equivalence in its nerve is exactly an isomorphism in C.

Let J denote the simplicial set



Unless otherwise specified, any map  $\Delta^1 \hookrightarrow J$  will denote the inclusion of the 1-simplex from 0 to 1.

**Definition 2.5.** A 1-simplex f in a simplicial set X is a J-equivalence if, as a map  $f: \Delta^1 \to X$ , it admits a lifts to a map  $J \to X$ .



We recall that  $E^1$ -equivalences and J-equivalence coincide in a quasicategory.

**Proposition 2.6.** If X is a quasicategory then a 1-simplex in X is a J-equivalence if and only if it is an  $E^1$ -equivalence.

In light of Proposition 2.6, we will refer to both  $E^1$ -equivalences and J-equivalences in a quasicategory as simply equivalences (or invertible).

The simplicial sets  $E^1$  and J give well-behaved interval objects for defining a notion of homotopy.

#### Definition 2.7.

- For simplicial maps  $f, g: X \to Y$ , an  $E^1$ -homotopy (or natural equivalence) from f to g is a map  $\alpha: X \times E^1 \to Y$  such that  $\alpha|_{X \times \{0\}} = g$  and  $\alpha|_{X \times \{1\}} = g$ .
- A map  $f: X \to Y$  is an  $E^1$ -homotopy equivalence if there exists  $g: Y \to X$  along with  $E^1$ -homotopies  $gf \sim \operatorname{id}_X$  and  $fg \sim \operatorname{id}_Y$ .
- For simplicial maps  $f, g: X \to Y$ , a *J*-homotopy from f to g is a map  $\alpha: X \times J \to Y$  such that  $\alpha|_{X \times \{0\}} = f$  and  $\alpha|_{X \times \{1\}} = g$ .
- A map  $f: X \to Y$  is a *J*-homotopy equivalence if there exists  $g: Y \to X$  along with *J*-homotopies  $gf \sim id_X$  and  $fg \sim id_Y$ .

If Y is a quasicategory then any map  $\alpha \colon X \times \Delta^1 \to Y$  whose restriction  $\alpha|_{\{x\} \times \Delta^1} \colon \{x\} \times \Delta^1 \to Y$ along every 0-simplex  $x \in X_0$  is invertible lifts to a natural equivalence.

Recall that quasicategories are the fibrant objects in the Joyal model structure on simplicial sets.

**Theorem 2.8** ([Joy09, Thm. 6.12]). The category of simplicial sets sSet admits a model structure where

- cofibrations are monomorphisms;
- fibrant objects are quasi-categories; fibrations between fibrant objects are inner isofibrations;
- weak equivalences are weak categorical equivalences.

In particular, both  $E^1$ -homotopy equivalences and J-homotopy equivalences are weak categorical equivalences.

We also recall the definition of left fibrations, which we use in Section 11 to compute colimits of  $\infty$ -groupoids.

**Definition 2.9.** A *left fibration* is a simplicial map  $f: X \to Y$  with the right lifting property against the set

$$\{\Lambda_k^n \hookrightarrow \Delta^n \mid n \ge 1, \ 0 \le k < n\}.$$

In particular, a left fibration between quasicategories is a fibration ([Joy09, Prop. 4.10]).

In Section 11, we also make use of the join and slice constructions, which we now define. Let  $\Delta_{aug}$  denote the category obtained by adjoining an initial object [-1] to  $\Delta$ . Define the *ordinal sum* functor  $\Delta_{aug} \times \Delta_{aug} \rightarrow \Delta_{aug}$  by

$$([m], [n]) \mapsto [m+n+1]$$

By Day convolution, this gives a monoidal product on the presheaf category, which we denote by

 $-*-:\mathsf{sSet}_{\mathsf{aug}}\times\mathsf{sSet}_{\mathsf{aug}}\to\mathsf{sSet}_{\mathsf{aug}}.$ 

The inclusion  $\Delta \subseteq \Delta_{aug}$  induces a forgetful functor on presheaf categories  $sSet_{aug} \rightarrow sSet$  (which we do not introduce notation for). This functor admits a right adjoint  $X \mapsto \hat{X}$  defined by

$$\widehat{X}_n := \begin{cases} \{*\} & n = -1 \\ X_n & \text{otherwise.} \end{cases}$$

**Definition 2.10.** The *join* of simplicial sets is the functor  $-*-sSet \times sSet \rightarrow sSet$  defined by

$$X * Y := \widehat{X} * \widehat{Y}.$$

**Proposition 2.11** ([Joy09, Prop. 3.2]). For simplicial sets X, Y, the n-simplices of X \* Y are given by the formula

$$(X * Y)_n := X_n \sqcup Y_n \sqcup \coprod_{i+j+1=n} X_i \times Y_j.$$

If X and Y are quasicategories then X \* Y is again a quasicategory ([Joy09, Cor. 3.23]). The simplicial set X \* Y admits two natural inclusions

$$i_0: X \hookrightarrow (X * Y) \quad i_1: Y \hookrightarrow (X * Y).$$

Thus, we may view the functors - X and X - a taking values in the slice under X, i.e.

$$- * X, X * -: \mathsf{sSet} \to X \downarrow \mathsf{sSet}.$$

In this sense, both functors admit right adjoints ([Joy09, Prop. 3.12]). These are the *slice over* and *slice under* functors, respectively. Explicitly, these functors send an object  $(Y, f: X \to Y)$  to the simplicial sets  $f \downarrow Y$  and  $Y \downarrow f$  defined by

$$\begin{array}{ll} (f \downarrow Y)_n &:= X \downarrow \mathsf{sSet} \left( (X \ast \Delta^n, i_0), (Y, f) \right) \\ (Y \downarrow f)_n &:= X \downarrow \mathsf{sSet} \left( (\Delta^n \ast X, i_1), (Y, f) \right). \end{array}$$

There are natural projection maps

$$f \downarrow Y \to Y \quad Y \downarrow f \to Y$$

defined by restriction along the inclusion  $\Delta^n \to \Delta^n * X$ .

If Y is a quasicategory then both the slice over and the slice under Y are quasicategories ([Joy09, Cor. 3.20]). As such, we occasionally refer to them as *slice quasicategories*.

Recall that *mapping spaces* in a quasicategory may be constructed in multiple equivalent ways, one of which is as a certain fiber of the slice under the domain (or over the codomain).

**Definition 2.12.** Let x, y be objects in a quasicategory C.

1. The left mapping space  $\operatorname{map}_{L}(\mathcal{C}, x, y)$  from x to y in  $\mathcal{C}$  is the pullback

$$\begin{array}{ccc} \operatorname{map}_{L}(\mathcal{C}, x, y) & \longrightarrow x \downarrow \mathcal{C} \\ & \downarrow & & \downarrow \\ & \Delta^{0} & \overset{\mathcal{Y}}{\longrightarrow} & \mathcal{C} \end{array}$$

2. The right mapping space map<sub>R</sub>( $\mathcal{C}, x, y$ ) from x to y in  $\mathcal{C}$  is the pullback

$$\operatorname{map}_{R}(\mathcal{C}, x, y) \longrightarrow \mathcal{C} \downarrow y \\ \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \\ \Delta^{0} \xrightarrow{x} \longrightarrow \mathcal{C}$$

**Proposition 2.13** ([Lur09, Prop. 1.2.2.3 & Cor. 4.2.1.8]). The left and right mapping spaces are homotopyequivalent Kan complexes.

Using slice quasicategories, one may define the notion of cones over (and under) a diagram, as well as limits and colimits in a quasicategory.

**Definition 2.14.** Let  $\mathcal{C}$  be a quasicategory and  $f: K \to \mathcal{C}$  be a simplicial map.

- 1. An *initial object* in C is an object  $x \in C$  such that any map  $f : \partial \Delta^n \to C$  satisfying  $f|_{\{0\}} = x$  can be lifted to a map  $\Delta^n \to C$ .
- 2. A cone under f is a lift of f to a map  $K * \Delta^0 \to \mathcal{C}$ , i.e. an object of  $f \downarrow \mathcal{C}$ .
- 3. The *colimit cone* of f is the initial object of  $f \downarrow C$ .

These definitions dualize in a straightforward way.

**Definition 2.15.** Let  $\mathcal{C}$  be a quasicategory and  $f: K \to \mathcal{C}$  be a simplicial map.

- 1. A terminal object in C is an object  $x \in C$  such that any map  $f: \partial \Delta^n \to C$  satisfying  $f|_{\{n+1\}} = x$  can be lifted to a map  $\Delta^n \to C$ .
- 2. A cone over f is a lift of f to a map  $\Delta^0 * K \to C$ , i.e. an object of  $C \downarrow f$ .
- 3. The *limit cone* of f is the terminal object of  $\mathcal{C} \downarrow f$ .

Note that the mapping space between any two initial (or terminal) objects is contractible [Joy02, Prop. 4.4], hence any two colimits (or limits) of f are equivalent.

## 3 Marked simplicial sets and localization

Our final section of preliminaries describes the analogues of marked categories and localization for quasicategories. Mainly, we are interested in the category of marked simplicial sets and the model structure thereon. We begin with the definition of a marked simplicial set.

**Definition 3.1.** A marked simplicial set is a pair (X, W) where X is a simplicial set and  $W \subseteq X_1$  is a subset of 1-simplices which contains all degenerate 1-simplices.

We write  $sSet_+$  for the category of marked simplicial sets and simplicial maps which preserve marked 1-simplices. This category is Cartesian closed, and we write  $(-)^{(X,W)}: sSet_+ \rightarrow sSet_+$  for the left adjoint to the functor  $(X, W) \times -: sSet_+ \rightarrow sSet_+$ . Moreover, if (X', W') is a marked quasicategory then  $(X', W')^{(X,W)}$  is a quasicategory for any  $(X, W) \in sSet_+$  [Lur09, Rem. 3.1.3.1].

We make use of the following functors between simplicial sets and marked simplicial sets.

**Proposition 3.2.** There is an adjoint quadruple

$$((-)^{\flat} \dashv U \dashv (-)^{\#} \dashv \operatorname{Core}_{+})$$

between marked simplicial sets and simplicial sets where

- $(-)^{\flat}$ : sSet  $\rightarrow$  sSet<sub>+</sub> is the minimal marking functor which sends X to the pair  $(X, \sigma_0 X_0)$ , where  $\sigma_0 X_0$  denotes the collection of degenerate 1-simplices in X;
- U: sSet<sub>+</sub> → sSet is the forgetful functor which sends a marked simplicial set (X, W) to its underlying simplicial set X.
- $(-)^{\#}$ : sSet  $\rightarrow$  sSet<sub>+</sub> is the maximal marking functor which sends X to the pair  $(X, X_1)$ ; and
- Core<sub>+</sub>: sSet<sub>+</sub> → sSet is the marked core which sends (X, W) to the maximal simplicial subset of X whose 1-simplices are all marked.

Additionally, we have the *natural marking* functor

$$(-)^{\natural} : \mathsf{sSet} \to \mathsf{sSet}_+$$

which sends X to the pair (X, W) where W is the set of J-equivalences in X. The natural marking is functorial as simplicial maps preserve J-equivalences.

**Remark 3.3.** The minimal marking functor  $(-)^{\flat}$ :  $sSet \to sSet_+$  admits a further left adjoint which sends a marked simplicial set (X, W) to the quotient X/W. That is, the pushout

However, we will not make use of this functor.

The category  $sSet_+$  is a reflective subcategory of a presheaf category. The representable presheaves are the minimally marked simplices  $(\Delta^0)^{\flat}, (\Delta^1)^{\flat}, (\Delta^2)^{\flat}, \ldots$  along with the maximally marked 1-simplex  $(\Delta^1)^{\#}$ . In particular, every marked simplicial set is a canonical colimit of representable presheaves.

We now discuss generalizations of closure under composition and closure under 2-out-of-3 to the quasicategorical setting.

**Definition 3.4.** Let (X, W) be a marked simplicial set.

- 1. The set W is weakly closed under composition if every map  $(\Lambda_1^2)^{\#} \to (X, W)$  lifts to a map  $(\Delta^2)^{\#} \to (X, W)$ .
- 2. The set W is strongly closed under composition if, given any 2-simplex  $u \in X_2$  whose  $\partial_0$  and  $\partial_2$ -faces are marked, we have that its  $\partial_1$ -face is also marked.

3. The set W is closed under 2-out-of-3 if, given a 2-simplex  $u \in X_2$  such that any two faces of u are marked, the third face is also marked.

**Remark 3.5.** We can rephrase the above properties in terms of properties of the marked core of (X, W). Namely, the set W is

- 1. weakly closed under composition if and only if the marked core  $\operatorname{Core}_+(X, W)$  has the right lifting property with respect to the horn inclusion  $\Lambda_1^2 \hookrightarrow \Delta^2$ ;
- 2. strongly closed under composition if and only if the marked core inclusion  $\operatorname{Core}_+(X, W) \hookrightarrow X$  is an inner fibration;
- 3. closed under 2-out-of-3 if and only if the marked core inclusion  $\operatorname{Core}_+(X, W) \hookrightarrow X$  is a Kan fibration.

In particular, closure under 2-out-of-3 implies strong closure under composition.

If X is a quasicategory then condition (1) is equivalent to  $\operatorname{Core}_+(X, W)$  being a quasicategory. Thus, if W is strongly closed under composition then it is weakly closed under composition. This is not true in general, with the maximally marked horn  $(\Lambda_1^2)^{\#}$  providing a counter-example.

Note the join of simplicial sets naturally ascends to a monoidal product on marked simplicial sets as follows: given marked simplicial sets (X, W) and (X', W'), we write (X, W) \* (X', W') for the marking on X \* X' given by the images of W and W' under the join inclusions, respectively. The induced endofunctors

$$-*(X,W), (X,W)*-: \mathsf{sSet}_+ \to \mathsf{sSet}_+$$

have right adjoint defined on an object  $(X', W', f: (X, W) \to (X', W'))$  by

$$\begin{array}{l} (f \downarrow X', W')_n := (X, W) \downarrow \mathsf{sSet}_+ \left( (\Delta^n * X, i_2), (Y, f) \right) \\ (X' \downarrow f \ , W')_n := (X, W) \downarrow \mathsf{sSet}_+ \left( (X * \Delta^n, i_1), (Y, f) \right) \end{array}$$

where a 1-simplex  $(X, W) * \Delta^1 \to (X', W')$  (or  $\Delta^1 * (X, W) \to (X', W')$ ) is marked if it ascends to a map  $(X, W) * (\Delta^1)^{\#} \to (X', W')$  (or  $(\Delta^1)^{\#} * (X, W) \to (X', W')$ ). That is, the marked simplices are created by the pre-image of W under the projection map  $(f \downarrow X', W') \to (X', W')$ .

We now define localization of quasicategories.

**Definition 3.6.** Let  $(\mathcal{C}, W)$  be a marked quasicategory. The *localization* of  $\mathcal{C}$  at W is a marked map

$$\gamma \colon (\mathcal{C}, W) \to \mathcal{C}[W^{-1}]^{\natural}$$

from  $\mathcal{C}$  to some quasicategory  $\mathcal{C}[W^{-1}]$  with the property that, for any quasicategory  $\mathcal{D}$ , the induced map

$$\gamma^* : \mathsf{sSet}(\mathcal{C}[W^{-1}], \mathcal{D}) \to \mathsf{sSet}_+((\mathcal{C}, W), \mathcal{D}^{\natural})$$

is an equivalence of quasicategories.

**Remark 3.7.** We distinguish two "edge-cases" of localization. First, when  $(\mathcal{C}, W) = \mathcal{C}^{\natural}$ , the localization is an equvalence in the Joyal model structure. Second, when  $(\mathcal{C}, W) = \mathcal{C}^{\#}$ , the localization is a fibrant replacement of  $\mathcal{C}$  in the Kan–Quillen model structure. This follows from the proof of [Cis19, Prop. 7.1.3], which shows that the localization fits into (and can be computed as) a homotopy pushout

$$\begin{array}{ccc} \operatorname{Core}_+(\mathcal{C}, W) & & \longrightarrow \mathcal{C} \\ & & & & \downarrow^{\gamma} \\ \operatorname{Ex}^{\infty} \left( \operatorname{Core}_+(\mathcal{C}, W) \right) & \longrightarrow \mathcal{C}[W^{-1}] \end{array}$$

of quasicategories. This also shows the localization always exists and is unique up to categorical equivalence. For an ordinary marked category  $(\mathcal{C}, W)$ , it is generally not the case that the " $\infty$ -localization" of  $(N\mathcal{C}, W)$  (in the sense of Definition 3.6) is categorically-equivalent to the nerve of its "1-localization" (in the sense of Definition 1.2). For instance, every  $\infty$ -groupoid arises as the localization of a maximally-marked poset, but not every  $\infty$ -groupoid is weakly equivalent (in the Joyal model structure) to the nerve of a category. However, when passing to homotopy categories, the  $\infty$ -localization of  $(N\mathcal{C}, W)$  becomes the 1-localization of (Ho  $N\mathcal{C}, W) = (\mathcal{C}, W)$  (cf. [Cis19, Rem. 7.1.6]).

The corresponding notion of homotopy for marked simplicial sets is given by marked homotopies.

#### Definition 3.8.

- 1. For marked maps  $f, g: (X, W) \to (X', W')$ , a marked homotopy from f to g is a marked map  $\alpha: (X, W) \times (\Delta^1)^{\#} \to (X', W')$  such that  $\alpha|_{X \times \{0\}} = f$  and  $\alpha|_{X \times \{1\}} = g$ .
- 2. A map  $f: (X, W) \to (X', W')$  is a marked homotopy equivalence if there exists  $g: (X', W') \to (X, W)$ along with marked homotopies  $gf \sim id_{(X,W)}$  and  $fg \sim id_{(X',W')}$ .

The following result shows that marked homotopies between preorders are preserved by the join.

**Proposition 3.9.** For any marked simplicial set (X, W), the functors

$$-*(X,W), (X,W)*-: \mathsf{sSet}_+ o \mathsf{sSet}_+$$

preserve marked homotopies between preorders (and hence marked homotopy equivalences between preorders).

*Proof.* We prove the result for - \*X, as the proof for X \* - is analogous.

For  $m \ge 0$  and a preorder P, consider the map

$$\alpha \colon (P \ast (\Delta^m)^{\flat}) \times (\Delta^1)^{\#} \to (P \times (\Delta^1)^{\#}) \ast (\Delta^m)^{\flat}$$

defined, as a preorder map, by

$$\alpha(x,t) := \begin{cases} (x,t) & x \in P \\ x & x \in \Delta^m \end{cases}$$

It is straightforward to verify this map preserves markings and induces a natural transformation

via extension by colimits. With this, we have that for any marked homotopy  $H: P \times (\Delta^1)^{\#} \to Q$  from f to g and  $(X, W) \in \mathsf{sSet}_+$ , the composite

$$(P * (X, W)) \times (\Delta^1)^{\#} \xrightarrow{\alpha_{(X, W)}} (P \times (\Delta^1)^{\#}) * (X, W) \xrightarrow{H * (X, W)} Q * (X, W)$$

is a marked homotopy from f \* (X, w) to g \* (X, W).

Recall from [Lur09] that  $sSet_+$  admits a model structure (the model structure for Cartesian fibrations over  $\Delta^0$ ).

**Theorem 3.10** ([Lur09, Props. 3.1.3.7 & 3.1.4.1]). The category sSet<sub>+</sub> admits a model structure where

- cofibrations are monomorphisms;
- fibrant objects are quasicategories marked at equivalences.

We also note that marked homotopy equivalences are weak equivalences in this model structure. The key property of this model structure which we wish to use is the following.

**Proposition 3.11** ([Hin16, Prop. 1.1.3]). A marked map of the form  $(\mathcal{C}, W) \to \mathcal{D}^{\natural}$  between quasicategories is a weak equivalence if and only if it is the localization of  $\mathcal{C}$  at W.

This model structure is Quillen equivalent to the Joyal model structure via the minimal marking adjunction.

**Theorem 3.12** ([Lur09, Prop. 3.1.5.3]). The adjunction

$$(-)^{\flat}$$
: sSet  $\rightleftharpoons$  sSet<sub>+</sub> : U

is a Quillen equivalence between the Joyal model structure on sSet and the model structure on  $sSet_+$ .

### $(\infty, 1)$ -calculus of fractions

# 4 $(\infty, 1)$ -calculus of fractions — definition

In this section, we introduce the definition of calculus of fractions for quasicategories. Namely, the marked quasicategories satisfying calculus of (left or right) fractions will be those that satisfy a certain right lifting property.

To describe this right lifing property, we first give an auxiliary construction.

#### Definition 4.1.

1. For  $n \ge 0$  and  $0 \le k \le n$ , define a subset  ${}^*L_k^n$  of the powerset  $\mathcal{P}[n]$ , viewed as a simplicial set, by

$${}^{*}L_{k}^{n} := \{A \subseteq [n] \mid k \in A\}.$$

A 1-simplex  $A_0 \subseteq A_1$  in  ${}^*L_k^n$  is marked if  $\max(A_0) = \max(A_1)$ .

- 2. For  $n \ge 0$  and  $0 \le k \le n$ , let  ${}^*R_k^n$  be the opposite simplicial set of  ${}^*L_k^n$ , where a 1-simplex  $A_0 \supseteq A_1$  is marked if min  $A_0 = \min A_1$ .
- 3. For  $n \ge 0$  and  $0 \le k \le n$ , let  $L_k^n$  be the maximal simplicial subset of  $L_k^n$  which omits the 0-simplex [n].
- 4. For  $n \ge 0$  and  $0 \le k \le n$ , let  $\mathbb{R}^n_k$  be the maximal simplicial subset of  $\mathbb{R}^n_k$  which omits the 0-simplex [n].

**Example 4.2.** We depict the pairs  $L_1^2 \subseteq {}^*L_1^2$ ,  $L_2^2 \subseteq {}^*L_2^2$ , and  $L_1^3 \subseteq {}^*L_1^3$ , respectively.

The inclusion  $L_1^2 \subseteq {}^*\!L_1^2$ .

The inclusion  $L_2^2 \subseteq {}^*\!L_2^2$ .



The inclusion  $L_1^3 \subseteq {}^*\!L_1^3$ .

We also depict the pairs  $R_0^2 \subseteq {}^*\!R_0^2$ ,  $R_1^2 \subseteq {}^*\!R_1^2$ , and  $R_2^3 \subseteq {}^*\!R_2^3$ , respectively.

$$\begin{cases} 0,1\} & \{0,1,2\} \xrightarrow{\sim} \{0,1\} & \{0,1,2\} \xrightarrow{\sim} \{0,1\} \\ \downarrow^{\sim} & \subseteq & \sim \downarrow & \downarrow^{\sim} & \downarrow & \downarrow \\ \{0,2\} \xrightarrow{\sim} \{0\} & \{0,2\} \xrightarrow{\sim} \{0\} & \{1,2\} \xrightarrow{\sim} \{1\} & \{1,2\} \xrightarrow{\sim} \{1\} \end{cases}$$

The inclusion 
$$R_0^2 \subseteq {}^*\!R_0^2$$

The inclusion  $R_1^2 \subseteq {}^*\!R_1^2$ .



The inclusion  $R_2^3 \subseteq {}^*\!R_2^3$ .

The isomorphism  $F: [n]^{\text{op}} \to [n]$  defined by F(i) = n - i induces a map

$$F_* \colon ({}^*\!L^n_k)^{\mathrm{op}} \to {}^*\!R^n_{n-k}$$

which sends a subset to its image under F.

**Proposition 4.3.** The map  $F_*: ({}^*L_k^n)^{\mathrm{op}} \to {}^*R_{n-k}^n$  is an isomorphism. Moreover,  $F_*$  restricts to an isomorphism

$$F_*|_{L_k^n} \colon (L_k^n)^{\mathrm{op}} \xrightarrow{=} R_{n-k}^n.$$

**Definition 4.4.** Let  $(\mathcal{C}, W)$  be a marked quasicategory.

1. We say  $(\mathcal{C}, W)$  satisfies *calculus of left fractions* (or CLF) if W is weakly closed under composition and  $(\mathcal{C}, W)$  has the right lifting property against the set

$$\{L_k^n \hookrightarrow {}^*\!L_k^n \mid n \ge 2, \ 0 < k \le n\}$$

That is, for  $n \ge 2$  and  $0 < k \le n$ , any map  $L_k^n \to (\mathcal{C}, W)$  lifts to a map  ${}^*\!L_k^n \to (\mathcal{C}, W)$ .



2. We say  $(\mathcal{C}, W)$  satisfies *calculus of right fractions* (or CRF) if W is weakly closed under composition and  $(\mathcal{C}, W)$  has the right lifting property against the set

$$\{R_k^n \hookrightarrow R_k^n \mid n \ge 2, \ 0 \le k < n\}.$$

**Remark 4.5.** It follows from Proposition 4.3 that  $(\mathcal{C}, W)$  satisfies CLF if and only if  $(\mathcal{C}^{op}, W)$  satisfies CRF.

A priori, the quasicategorical definition of calculus of fractions may not agree with the classical definition when viewing an ordinary category as a quasicategory. We will show in Theorem 5.2 that the quasicategorical notion agrees with that of proper calculus of fractions (Definition 1.4). As our focus going forward will be on quasicategories, we use the terms calculus of left fractions and CLF (or calculus of right fractions and CRF, respectively) to mean the quasi-categorical notion unless otherwise specified.

The following proposition establishes that the k = n lifting condition in CLF (and the k = 0 condition in CRF) is redundant.

**Proposition 4.6.** Let (X, W) be a marked simplicial set.

- 1. If (X, W) has the right lifting property against  $\{L_{n-1}^n \to {}^*\!L_{n-1}^n \mid n \ge 2\}$  then it has the right lifting property against  $\{L_n^n \to {}^*\!L_n^n \mid n \ge 2\}$ .
- 2. If (X, W) has the right lifting property against  $\{R_1^n \to {}^*\!R_1^n \mid n \ge 2\}$  then it has the right lifting property against  $\{R_0^n \to {}^*\!R_0^n \mid n \ge 2\}$ .

*Proof.* We prove (1), as (2) is dual. Given  $n \ge 2$ , the inclusion  ${}^*L_n^n \to {}^*L_n^{n+1}$  defined, as a poset map, by  $A \mapsto A \cup \{n+1\}$  has a retraction  $r : {}^*L_n^{n+1} \to {}^*L_n^n$  defined by

$$r(A) := \begin{cases} \sigma_n(S) & \text{if } n+1 \in S\\ \{n\} & \text{otherwise.} \end{cases}$$

This map restricts so that the diagram

$$\begin{array}{cccc} L_n^n & & & L_n^{n+1} & \stackrel{r|_{L_n^{n+1}}}{\longrightarrow} & L_n^n \\ & & & \downarrow & & \downarrow \\ *L_n^n & & & *L_n^{n+1} & \stackrel{r}{\longrightarrow} & *L_n^n \end{array}$$

commutes. Thus,  $L_n^n \to {}^*\!L_n^n$  is a retract of  $L_n^{n+1} \to {}^*\!L_n^{n+1}$ .

### 5 $(\infty, 1)$ -calculus of fractions vs. classical calculus of fractions

We now show that the quasicategorical definition of calculus of fractions agrees with that of *proper* calculus of fractions, as stated in Definition 1.4 (cf. Theorem 5.2).

To show this, we use the following technical lemma which encapsulates our usage of the "co-equalising" condition from classical CLF (i.e. condition (3) of Definition 1.3).

**Lemma 5.1.** Let  $(\mathcal{C}, W)$  be a marked category satisfying CLF. For  $m \geq 1$ , let

$$\{f_i, g_i \colon x_i \rightrightarrows y \mid i = 1, \dots, m\}$$

be a set of m parallel pairs of morphisms with identical codomains. If, for each i, there exists a weak equivalence  $w_i$  such that  $f_iw_i = g_iw_i$  then there exists a weak equivalence u such that  $uf_i = ug_i$  for all i = 1, ..., m.

*Proof.* We fix m and construct u by induction over i. For i = 1, this holds by definition of CLF.

Fix *i* and suppose there exists a weak equivalence  $u^i$  such that  $u^i f_j = u^i g_j$  for all j = 1, ..., i. As  $u^i f_{i+1} w_{i+1} = u^i g_{i+1} w_{i+1}$  and  $(\mathcal{C}, W)$  satisfies CLF, there exists a weak equivalence u' such that  $u'u^i f_{i+1} = u'u^i g_{i+1}$ . Setting  $u = u'u^i$  gives that  $uf_j = ug_j$  for j = 1, ..., i+1.

**Theorem 5.2.** Let  $(\mathcal{C}, W)$  be a category with weak equivalenes.

- 1. The pair  $(\mathcal{C}, W)$  satisfies proper CLF if and only if  $(N\mathcal{C}, W)$  satisfies CLF.
- 2. The pair  $(\mathcal{C}, W)$  satisfies proper CRF if and only if  $(N\mathcal{C}, W)$  satisfies CRF.

*Proof.* We show (1), as (2) is dual.

Suppose  $(N\mathcal{C}, W)$  satisfies CLF. Given a diagram,



a filler exists as  $(N\mathcal{C}, W)$  has the right lifting property against  $L_1^2 \hookrightarrow {}^*L_1^2$  and  $L_0^2 \hookrightarrow {}^*L_0^2$ . For  $f, g \in \mathcal{C}$  and  $w \in W$  such that fw = gw, define a map  $L_1^3 \to (N\mathcal{C}, W)$  by the diagram:



This map admits a lift  ${}^*L_1^3 \to (N\mathcal{C}, W)$  as in the diagram:



The right face witnesses that  $u_1 f = u_2 g$ . The maps  $u_2$  and  $u_3$  are weak equivalences by assumption. The front and bottom faces witness that  $u_1 = u_2 = u_3$ , which suffices.

Now, suppose  $(\mathcal{C}, W)$  satisfies proper CLF. Weak equivalences are (strongly) closed under composition by assumption, so it remains to show  $(N\mathcal{C}, W)$  has the right lifting property with respect to the set of maps

$$\{L_k^n \to {}^*\!L_k^n \mid n \ge 2, \ 0 < k \le n\}.$$

We proceed by induction on n. The base case n = 2 follows by definition of proper CLF.

Fix  $f: L_k^{n+1} \to (\mathcal{C}, W)$  and suppose  $(N\mathcal{C}, W)$  has the right lifting property against

$$\{L_k^n \to {}^*\!L_k^n \mid 0 < k \le n\}.$$

It suffices to construct an object  $x \in C$  and, for  $i \neq k$ , a morphism

$$\varphi_i \colon f([n+1] - \{i\}) \to x$$

which is a weak equivalence for  $i \neq n+1$  and such that the square

commutes for all i, j = 0, ..., n + 1 where  $i, j \neq k$ .

Let d denote the value  $\partial_k^n(n)$ . That is,

$$d := \begin{cases} n & \text{if } k = n+1\\ n+1 & \text{otherwise.} \end{cases}$$

The inclusion  $L^n_n \to L^{n+1}_k$  defined by  $S \mapsto \partial^n_k(S) \cup \{k\}$  gives a lifting problem

which admits a lift  $g_0: {}^*L_n^n \to (\mathcal{C}, W)$  by the inductive hypothesis. Let  $x_0$  denote the object  $g_0|_{\{[n]\}}$  and, for  $i = 0, \ldots, n+1$  such that  $i \neq k, d$ , let

$$\varphi_{0,i} \colon f([n+1] - \{i\}) \to x_0$$

denote the image under  $g_0$  of the 1-simplex  $[n] - \{(\partial_k^n)^{-1}(i)\} \subseteq [n]$ . Note this map is a weak equivalence except when d = n and i = n + 1.

The diagram

$$f([n+1] - \{0, d\}) \longrightarrow f([n+1] - \{0\}) \xrightarrow{\varphi_{0,0}} x_0$$

$$\sim \downarrow$$

$$f([n+1] - \{d\})$$

admits a lift

thus defining a map  $\varphi_{1,d}$ . If k = n + 1 then d = n, which gives that the top map (and hence the bottom map) is a weak equivalence. For  $i \neq k, d$ , let

$$\varphi_{1,i} \colon f([n+1] - \{i\}) \to x_1$$

denote the composite  $\varphi_{0,i} \circ w$ . Note that  $\varphi_{1,i}$  is a weak equivalence for  $i \neq n+1$ .

Fix i, j = 0, ..., n + 1 such that  $i, j \neq k$ . If  $i, j \neq d$  then the square

commutes by construction. Otherwise, we have that every face except the front face in the cube

$$\begin{array}{c} f([n+1] - \{0, i, d\}) & \longrightarrow f([n+1] - \{0, i\}) \\ & & \downarrow \\ & & f([n+1] - \{i, d\}) & \longrightarrow f([n+1] - \{i\}) \\ & & & \downarrow \\ & & & f([n+1] - \{0, d\}) & \longrightarrow f([n+1] - \{0\}) \\ & & & & \downarrow \\ & & & & f([n+1] - \{d\}) & \longrightarrow x_1 \end{array}$$

commutes. Thus, the front face commutes after pre-composition with the top-left weak equivalence. By Lemma 5.1, there exists a weak equivalence

$$u \colon x_1 \to x$$

such that x and  $\varphi_i = u \circ \varphi_{1,i}$  assemble to give a lift of  $f: L_k^{n+1} \to (\mathcal{C}, W)$ .

**Remark 5.3.** The proof of the reverse direction in Theorem 5.2 proves a slightly stronger result: given a marked quasicategory  $(\mathcal{C}, W)$ , if

- W is weakly closed under composition; and
- $(\mathcal{C}, W)$  has the right lifting property against the maps  $L_1^2 \hookrightarrow {}^*\!L_1^2$  and  $L_1^3 \hookrightarrow {}^*\!L_1^3$ ,

then its homotopy category Ho  $\mathcal{C}$ , when marked at all morphisms which are identified with a morphism in W, satisfies classical CLF. If  $(\mathcal{C}, W)$  additionally has the right lifting property against  $L_2^2 \hookrightarrow {}^*L_2^2$  then Ho  $\mathcal{C}$  at this marking satisfies proper CLF. The dual statement holds for (proper) CRF.

#### 6 Examples of calculus of fractions

This section is devoted to giving examples of marked quasicategories (beyond nerves of 1-categories) which satisfy calculus of fractions. In particular, analogues of the examples given in [GZ67, Ch. 1] are covered in this section. These include: marking at equivalences (Corollary 6.2), marking at the inverse image of equivalences under a limit-preserving functor (Theorem 6.3), and pulling back CLF/CRF from an adjunction (Theorem 6.5). Note that this last example applies to all reflective localizations (Corollary 6.6).

We also introduce the notion of a simple inner horn decomposition in Definition 6.7. This is a technical condition for showing when a marked map has the left lifting property against all marked quasicategories  $(\mathcal{C}, W)$  where W is weakly closed under composition (see Lemma 6.8). In particular, we use this both in the proof of Theorem 6.5 and in Section 7 to show that our analogue of the category of fractions produces a quasicategory (Theorem 7.5). We conclude by showing that if a marked quasicategory satisfies calculus of fractions then the full subcategory of the arrow category spanned by marked arrows also satisfies calculus of fractions (Corollary 6.12). This requires a lemma relating right lifting properties against the maps  $L_k^n \hookrightarrow *L_k^n$  between a marked quasicategory and its marked arrow category (see Proposition 10.2).

To show that a quasicategory marked at equivalences satisfies CLF (and CRF), we make use of the following result.

#### Proposition 6.1.

- 1. a) For  $n \ge 1$  and  $0 \le k \le n$ , the map max:  ${}^{*}L_{k}^{n} \to (\Delta^{\{k,\dots,n\}})^{\flat}$  is a marked homotopy equivalence. b) For  $n \ge 2$  and  $0 < k \le n$ , the inclusion  $L_{k}^{n} \to {}^{*}L_{k}^{n}$  is an acyclic cofibration.
- 2. a) For  $n \ge 1$  and  $0 \le k \le n$ , the map min:  ${}^{*}R_{k}^{n} \to (\Delta^{k})^{\flat}$  is a marked homotopy equivalence. b) For  $n \ge 2$  and  $0 \le k < n$ , the inclusion  $R_{k}^{n} \to {}^{*}R_{k}^{n}$  is an acyclic cofibration.

*Proof.* We prove (1), as (2) is formally dual.

- a) The map max:  ${}^{*}L_{k}^{n} \to (\Delta^{\{k,\dots,n\}})^{\flat}$  has a section defined by  $i \mapsto \{0,\dots,i\}$ . For  $A \in {}^{*}L_{k}^{n}$ , the 1-simplex  $A \subseteq \{0,\dots,\max A\}$  is marked, giving the desired homotopy.
- b) The inclusion  $\partial_0: {}^*L_{k-1}^{n-1} \to L_k^n$  defined by sending a subset to its image under  $\partial_0$  admits a retraction defined by  $A \mapsto \sigma_0(A \{0\})$ . For  $A \in L_k^n$ , the 1-simplex  $\partial_0(\sigma_0(A \{0\})) \subseteq A$  is marked as  $\max A \ge k > 0$  (hence  $A \{0\}$  contains  $\max A$ ). Thus,  $\partial_0$  is a marked homotopy equivalence. The diagram

commutes, hence the top right map is a weak equivalence by 2-out-of-3 (applied twice).

**Corollary 6.2.** Any quasicategory marked at equivalences satisfies both CLF and CRF.

*Proof.* Follows from Proposition 6.1, as quasicategories marked at equivalences are fibrant objects in the model structure on  $sSet_+$ .

Our second example of when CLF is satisfied is in the case where weak equivalences are created by a colimit-preserving functor.

**Theorem 6.3.** Let  $F: \mathcal{C} \to \mathcal{D}$  be a functor between quasicategories and  $W \subseteq \mathcal{C}_1$  be the 1-simplices which are sent to equivalences under F.

- 1. If C admits all finite colimits and F preserves finite colimits then  $(\mathcal{C}, \mathcal{W})$  satisfies CLF.
- 2. If C admits all finite limits and F preserves finite limits then (C, W) satisfies CRF.

Before proving Theorem 6.3, we introduce an additional construction. This construction is only used to prove Theorem 6.3 and will not be needed going forward (though a related construction is introduced in Section 11).

For  $n \geq 2$  and  $0 \leq k \leq n$ , define  $\widehat{L_k^n}$  to be the nerve of the preorder whose underlying set is

$$\{A \subseteq [n] \mid k \in A \text{ and } A \neq [n]\}$$

and where  $A_0 \leq A_1$  if max  $A_0 \leq \max A_1$ . There is an evident inclusion  $L_k^n \subseteq \widehat{L_k^n}$ . The  $E^1$ -equivalences in  $\widehat{L_k^n}$  are exactly the 1-simplices  $A_0 \leq A_1$  where max  $A_0 = \max A_1$ . In particular, the map max:  $\widehat{L_k^n} \to \Delta^{\{k,\dots,n\}}$  is an  $E^1$ -equivalence with inverse given by  $i \mapsto \{k, i\}$ .

**Lemma 6.4.** Let (X, W) be a marked simplicial set. For  $n \ge 2$  and  $0 < k \le n$ , the map

$$L_k^n * (X, W) \hookrightarrow \widehat{L_k^n}^{\natural} * (X, W)$$

is an acyclic cofibration of marked simplicial sets.

Proof. Applying Proposition 3.9 gives that the top and left maps in

are marked homotopy equivalences. By 2-out-of-3, the right map is a weak equivalence. Similarly, max:  $\widehat{L_k^n} \to \Delta^{\{k,\dots,n\}}$  is an  $E^1$ -homotopy equivalence, hence a weak equivalence of marked simplicial sets. The result follows by 2-out-of-3 on the triangle



We now proceed with the proof of Theorem 6.3.

Proof of Theorem 6.3. We show (1), as (2) is analogous. Fix a map  $D: L_k^n \to \mathcal{C}$ . The underlying simplicial set of  ${}^*L_k^n$  is the join of the underlying simplicial set of  $L_k^n$  with  $\Delta^0$ . Thus, the colimit cone  $\lambda_D$  gives a lift



of underlying simplicial sets. It remains to show that if  $A \in L_k^n$  is such that  $n \in A$  then the component map  $D(A) \to \operatorname{colim} D$  is a weak equivalence. As F preserves finite colimits and creates weak equivalences, it suffices to show  $FD(A) \to \operatorname{colim}(FD)$  is an equivalence in  $\mathcal{D}$ .

By assumption,  $FD: L_k^n \to \mathcal{D}$  sends marked simplicies to equivalences. In the commutative triangle,



the right map is an  $E^1$ -homotopy equivalence and the bottom map is a weak equivalence by Proposition 6.1. Thus, the left inclusion is an acyclic cofibration of marked simplicial sets. Hence, there exists  $\overline{FD}$ :  $\widehat{L_k^n} \to \mathcal{D}$  such that the triangle



commutes. The inclusion  $L^n_k \hookrightarrow \widehat{L^n_k}$  induces a map

 $f \colon \overline{FD} \downarrow \mathcal{D} \to FD \downarrow \mathcal{D}$ 

by pre-composition. It suffices to show f is a trivial fibration: this would imply that the colimit of  $\overline{FD}$  (which is an initial object of  $\overline{FD} \downarrow D$ ) is equivalent to the colimit of FD. A subset  $A \in L_k^n$  such that  $n \in A$  is a terminal object in  $\widehat{L_k^n}$ , hence the component map

$$FD(A) = \overline{FD}(A) \to \operatorname{colim}(\overline{FD}) = \operatorname{colim}(FD)$$

is an equivalence in  $\mathcal{D}$ .

Given a commutative square

$$\begin{array}{ccc} \partial \Delta^n & \stackrel{u}{\longrightarrow} & \overline{FD} \downarrow \mathcal{D} \\ & & & \downarrow^f \\ \Delta^n & \stackrel{v}{\longrightarrow} & FD \downarrow \mathcal{D} \end{array}$$

the maps u and v may be identified as maps

$$u \colon \widehat{L}_k^n * \partial \Delta^n \to \mathcal{D}, \quad v \colon L_k^n * \Delta^n \to \mathcal{D}$$

making the triangles



commute. These maps assemble into a square



of marked maps, which induces a map  $P \to \mathcal{D}^{\natural}$  from the pushout P. In the square of marked maps

the left and right maps are acyclic cofibrations by Lemma 6.4. Thus, the induced map from the pushout into  $\widehat{L_k^n}^{\natural} * (\Delta^n)^{\flat}$  is as well, since it is both a weak equivalence (by 2-out-of-3) and a monomorphism (by direct verification). Thus, the map  $P \to \mathcal{D}^{\natural}$  lifts to a map  $\widehat{L_k^n}^{\natural} * (\Delta^n)^{\flat} \to \mathcal{D}$ . This gives a lift of the starting square when viewed as a map  $\Delta^n \to \overline{FD} \downarrow \mathcal{D}$ .

The following result gives a condition for when CLF (or CRF) may be "pulled back" from an adjunction.

**Theorem 6.5.** Let  $L: \mathcal{C} \to \mathcal{D}$  be a map between quasicategories with right adjoint  $R: \mathcal{D} \to \mathcal{C}$ .

- 1. Suppose  $W \subseteq \mathcal{D}_1$  is a marking on  $\mathcal{D}$  such that  $(\mathcal{D}, W)$  satisfies CLF. If every component of the unit is sent by L to a marked 1-simplex  $L\eta_x \colon Lx \to LRLx$  then  $(\mathcal{C}, L^{-1}W)$  satisfies CLF.
- 2. Suppose  $W \subseteq C_1$  is a marking on C such that (C, W) satisfies CRF. If every component of the counit is sent by R to a marked 1-simplex  $R\varepsilon_y \colon RLRy \to Ry$  then  $(\mathcal{D}, R^{-1}W)$  satisfies CRF.

This result will require a more technical proof, in particular, we will construct explicit cellular decompositions in the marked model structure. Before doing so, we recall a consequence of Theorem 6.5 which follows from Corollary 6.2.

**Corollary 6.6.** Let  $i: \mathcal{D} \to \mathcal{C}$  be a fully faithful functor between quasicategories.

- 1. If i admits a left adjoint  $L: \mathcal{C} \to \mathcal{D}$  then  $(\mathcal{C}, L^{-1}(\operatorname{Equiv} \mathcal{D}))$  satisfies CLF, where  $L^{-1}(\operatorname{Equiv} \mathcal{D})$  denotes the set of morphisms in  $\mathcal{C}$  which are sent to equivalences in  $\mathcal{D}$ .
- 2. If i admits a right adjoint  $R: \mathcal{C} \to \mathcal{D}$  then  $(\mathcal{C}, R^{-1}(\text{Equiv }\mathcal{D}))$  satisfies CRF, where  $R^{-1}(\text{Equiv }\mathcal{D})$ denotes the set of morphisms in  $\mathcal{C}$  which are sent to equivalences in  $\mathcal{D}$ .

As for the proof of Theorem 6.5, we begin by introducing the definition of a *simple inner horn decomposition*, which will also be used in Section 7.

**Definition 6.7.** For an injective map  $f: (X, W) \hookrightarrow (Y, V)$  between marked simplicial sets, a *simple inner* horn decomposition for f consists of

- for  $n \ge 0$ , a partition of the set of non-degenerate *n*-simplices in Y which are not in the image of f into a pair of sets  $A^n \sqcup B^n$  such that  $A^1$ ,  $A^0$ , and  $B^0$  are empty;
- for  $n \ge 1$ , a pair of finite partitions

$$A^{n} = A_{1}^{n} \sqcup \cdots \sqcup A_{a(n)}^{n}, \quad B^{n} = B_{1}^{n} \sqcup \cdots \sqcup B_{b(n)}^{n}$$

such that b(1) = 1 and a(n+1) = b(n);

• for  $n \ge 2$ , a function  $d: \{1, ..., a(n)\} \to \{1, ..., n-1\};$ 

such that for  $n \ge 2$  and  $k \in \{1, \ldots, a(n)\},\$ 

1. the d(k)-th face map restricts to a bijection

$$\partial_{d(k)}|_{A_k^n} \colon A_k^n \xrightarrow{\cong} B_k^{n-1};$$

- 2. for  $u \in A_1^2$ , if  $u\partial_1$  is marked then every face of u is marked;
- 3. for  $u \in A_k^n$  and  $i \neq d(k)$ , writing  $u\partial_i$  as a degeneracy  $\vec{\sigma} \colon \Delta^{n-1} \to \Delta^p$  of some non-degenerate *p*-simplex  $v \colon \Delta^p \to Y$ , one of the following holds:
  - v is contained in the image of f;
  - v is contained in  $A_i^p$  for some  $i \in \{1, \ldots, a(p)\};$
  - p < n-1 and v is contained in  $B_i^p$  for some  $i \in \{1, \dots, b(p)\};$
  - p = n 1 and v is contained in  $B_i^{n-1}$  for some  $i \in \{1, \dots, k-1\}$ .

We say f admits a simple inner horn decomposition if there exists a simple inner horn decomposition for f.

**Lemma 6.8.** Let  $f: (X, W) \to (Y, V)$  be an injective map of marked simplicial sets. If f admits a simple inner horn decomposition then f is in the saturation of the set

$$\{\left(\Lambda_k^n\right)^{\flat} \hookrightarrow \left(\Delta^n\right)^{\flat} \mid n \ge 2, \ 0 < k < n\} \cup \{\left(\Lambda_1^2\right)^{\#} \hookrightarrow \left(\Delta^2\right)^{\#}\}.$$

*Proof.* Let  $Y_1^1 \subseteq (Y, V)$  denote the image of f, viewed as a marked simplicial subset of (Y, V). For  $n \ge 2$  and  $k \in \{1, \ldots, a(n)\}$ , define a sequence of marked simplicial subsets of Y inductively by

$$Y_k^n := \begin{cases} Y_{a(n)}^{n-1} \cup B_1^{n-1} \cup A_1^n & \text{if } k = 1\\ Y_{k-1}^n \cup B_k^{n-1} \cup A_k^n & \text{if } k > 1. \end{cases}$$

The conditions for a simple inner horn decomposition assert that each  $Y_k^n$  is a well-defined simplicial subset of Y, i.e. that if u is a non-degenerate simplex of  $Y_k^n$  then every face of u is also contained in  $Y_k^n$ . By construction, there is a sequence of inclusions

$$Y_1^1 \subseteq Y_1^2$$
$$\subseteq Y_1^3 \subseteq \ldots \subseteq Y_{a(3)}^3$$
$$\vdots$$
$$\subseteq Y_1^n \subseteq \ldots \subseteq Y_{a(n)}^n$$
$$\vdots$$

and the colimit of this sequence is Y. We show each inclusion in this sequence is a pushout of a coproduct of maps in the set

$$\{\left(\Lambda_k^n\right)^{\flat} \hookrightarrow \left(\Delta^n\right)^{\flat} \mid n \ge 2, \ 0 < k < n\} \cup \{\left(\Lambda_1^2\right)^{\#} \hookrightarrow \left(\Delta^2\right)^{\#}\}$$

For a 2-simplex  $u \in A_1^2$ , the restriction  $u|_{\Lambda_1^2} \colon \Lambda_1^2 \to Y$  to the 1-horn factors through  $Y_1^1$  by definition of a simple inner horn decomposition. Moreover, if  $u\partial_1 \in B_1^1$  is marked then every face of u is marked. As  $\partial_1|_{B_1^2} \colon A_1^2 \to B_1^1$  is a bijection, it follows that the square



is a pushout in  $sSet_+$ . Similarly, for  $n \ge 3$  and k > 1, the squares



are pushouts.

We highlight an important benefit of Lemma 6.8, which is that marked quasicategories which are weakly closed under composition are exactly the objects in  $\mathbf{sSet}_+$  with the right lifting property against the set  $\{(\Lambda_k^n)^{\flat} \hookrightarrow (\Delta^n)^{\flat} \mid n \geq 2, \ 0 < k < n\} \cup \{(\Lambda_1^2)^{\#} \hookrightarrow (\Delta^2)^{\#}\}.$ 

The following lemma, which makes use of simple horn decompositions, will then allow us to prove (a generalization of) Theorem 6.5.

**Lemma 6.9.** For a marked poset (P, W) and any subset  $Q \subseteq P$ , viewed as a discrete set, the induced map f from the pushout in the diagram



is in the saturation of the set

$$\{(\Lambda_k^n)^{\flat} \hookrightarrow (\Delta^n)^{\flat} \mid n \ge 2, 0 < k < n\} \cup \{(\Lambda_1^2)^{\#} \hookrightarrow (\Delta^2)^{\#}\},\$$

*Proof.* By Lemma 6.8, it suffices to construct a simple inner horn decomposition for f.

Note the codomain of f is the poset  $(P \times [1]) * [0]$ . We write the cone point as  $\top$  and all other elements as  $(x, \varepsilon)$  for some  $x \in P$  and  $\varepsilon = 0, 1$ . A non-degenerate *n*-simplex in  $(P \times [1]) * [0]$  is an alternating chain. An *n*-chain is not contained in the image of f if and only if the first element in the chain is of the form (x, 0) for some  $x \in P$  and the last element in the chain is the cone point. Let  $S^n$  denote the set of such *n*-chains.

For  $n \ge 2$  and  $k = 1, \ldots, n-1$ , define subsets  $A_k^n \subseteq S^n$  and  $B_k^{n-1} \subseteq S^{n-1}$  by

$$A_k^n := \left\{ ((x_0, 0) \le (x_1, \varepsilon_1) \le \dots \le (x_{n-1}, \varepsilon_{n-1}) \le \top) \in S^n \mid \begin{array}{c} \varepsilon_{k-1} = 0\\ \varepsilon_k = 1\\ x_k = x_{k-1} \end{array} \right\}$$
$$B_k^{n-1} := \left\{ ((x_0, 0) \le (x_1, \varepsilon_1) \le \dots \le (x_{n-2}, \varepsilon_{n-2}) \le \top) \in S^n \mid \begin{array}{c} \varepsilon_{k-1} = 0\\ \varepsilon_k = 1\\ x_k \ne x_{k-1} \end{array} \right\}.$$

(For  $B_{n-1}^{n-1}$ , our convention is that the second and third conditions are vacuous.) The sets  $A_k^n$  and  $B_k^n$  are all mutually disjoint, and the pair

$$A^n := A_1^n \sqcup \cdots \sqcup A_{n-1}^n, \quad B^n := B_1^n \sqcup \cdots \sqcup B_{n-1}^n$$

partitions  $S^n$ . We show the tuple

$$(\{A_k^n\}_{k=1}^{n-1}, \{B_k^n\}_{k=1}^n, \mathrm{id})$$

is a simple inner horn decomposition for f.

Fix  $n \ge 2$  and  $k \in \{1, ..., n-1\}$ .

1. For a chain  $((x_0, 0) \leq (x_1, \varepsilon_1) \leq \cdots \leq (x_{n-1}, \varepsilon_{n-1}) \leq \top)$  in  $A_k^n$ , we have that  $\varepsilon_{k+1} = 1$ . As this chain is non-degenerate and  $\varepsilon_k = \varepsilon_{k+1}$ , we have that  $x_{k+1} \neq x_k = x_{k-1}$ . Thus, the k-th face of  $((x_0, 0) \leq (x_1, \varepsilon_1) \leq \cdots \leq (x_{n-1}, \varepsilon_{n-1}) \leq \top)$  is an element of  $B_k^{n-1}$ . The inverse function is given by

$$((x_0, 0) \le \dots \le (x_{k-1}, 0) \le (x_k, 1) \le \dots \le (x_{n-2}, 1) \le \top) \mapsto ((x_0, 0) \le \dots \le (x_{k-1}, 0) \le (x_{k-1}, 1) \le (x_k, 1) \le \dots \le (x_{n-2}, 1) \le \top).$$

- 2. For  $((x_0,0) \leq (x_0,1) \leq \top) \in A_1^2$ , the  $\partial_1$ -face  $(x_0,0) \leq \top$  is marked if  $x_0 \in Q$ . This gives that  $(x_0,1) \leq \top$  is marked. The 1-simplex  $(x_0,0) \leq (x_0,1)$  is marked as a product of marked 1-simplices.
- 3. For an *n*-simplex  $((x_0, 0) \le (x_1, \varepsilon_1) \le \cdots \le (x_{n-1}, \varepsilon_{n-1}) \le \top) \in A_k^n$  and  $i \ne k$ , the *i*-th face of this *n*-simplex is non-degenerate. We proceed by case analysis on *i*:
  - if i < k 1 then the *i*-th face is an element of  $A_{k-1}^{n-1}$ ;
  - if i = k 1 then we proceed by case analysis on k:
    - for k = 1, the 0-th face does not contain any elements of the form (x, 0). Thus, it is in the image of f;
    - for  $k \ge 2$ , we have that  $x_{k-2} \ne x_{k-1}$  as  $((x_0, 0) \le (x_1, \varepsilon_1) \le \cdots \le (x_{n-1}, \varepsilon_{n-1}) \le \top)$  is non-degenerate and  $\varepsilon_{k-2} = \varepsilon_{k-1} = 0$ . As  $x_k = x_{k-1}$ , the (k-1)-th face is an element of  $B_{k-1}^{n-1}$ ;
  - if i > k then we proceed by case analysis on k:
    - if k < n-1 then  $\varepsilon_{n-1} = 1$ , thus the *i*-th face is an element of  $A_k^{n-1}$ .
    - if k = n 1 then i = n. The *n*-th face does not contain  $\top$ , hence is contained in the image of f. □

The following theorem is a generalization of Theorem 6.5, giving a condition for when a calculus of fractions within a quasicategory C may be pulled back from a smaller class to a larger class.

**Theorem 6.10.** Let  $W_0 \subseteq W$  be two markings on a quasicategory C where W is weakly closed under composition and let  $F: (C, W) \to (C, W_0)$  be an endofunctor which takes W to  $W_0$ .

- 1. If  $(\mathcal{C}, W_0)$  has the right lifting property against the set  $\{L_k^n \hookrightarrow {}^*L_k^n \mid n \ge 2, \ 0 < k \le n\}$  and there exists a marked homotopy  $\alpha : (\mathcal{C}, W) \times (\Delta^1)^{\#} \to (\mathcal{C}, W)$  from id to F then  $(\mathcal{C}, W)$  satisfies CLF.
- 2. If  $(\mathcal{C}, W_0)$  has the right lifting property against the set  $\{R_k^n \hookrightarrow *R_k^n \mid n \geq 2, 0 \leq k < n\}$  and there exists a marked homotopy  $\alpha : (\mathcal{C}, W) \times (\Delta^1)^{\#} \to (\mathcal{C}, W)$  from F to id then  $(\mathcal{C}, W)$  satisfies CRF.

*Proof.* We prove (1), as (2) is formally dual.

It suffices to show  $(\mathcal{C}, W)$  has the right lifting property against  $\{L_k^n \hookrightarrow {}^*L_k^n \mid n \ge 2, \ 0 < k \le n\}$ . Thus, we fix a map  $f: L_k^n \to (\mathcal{C}, W)$ . The composite  $Ff: L_k^n \to (\mathcal{C}, W_0)$  admits a lift  $\overline{Ff}: {}^*L_k^n \to (\mathcal{C}, W_0)$  by assumption. This gives a commutative square

$$\begin{array}{ccc} L_k^n & \stackrel{i_1}{\longrightarrow} & L_k^n \times (\Delta^1)^{\#} \\ & & & & \downarrow \\ & & & \downarrow \\ {}^*\!L_k^n & \stackrel{\overline{Ff}}{\longrightarrow} & (\mathcal{C}, W_0) \end{array}$$

We apply Lemma 6.9 with  $(P, W) = L_k^n$  and  $Q \subseteq P$  being the collection of subsets which contain n. Thus, this square induces a map from the pushout which admits a lift to a map

$$\overline{[\alpha,\overline{Ff}]}: \left( (P \times \Delta^1) * \Delta^0, (W \times \Delta^1_1) \cup (Q \times \{0,1\}) * \Delta^0 \right) \to (\mathcal{C},W).$$

We have an inclusion from  $L_k^n$  to the domain of  $\overline{[\alpha, \overline{Ff}]}$  defined, as a poset map, by sending [n] to the cone point and all other subsets A to (A, 0). The composite

$${}^{*}L_{k}^{n} \hookrightarrow \left( (P \times \Delta^{1}) * \Delta^{0}, (W \times \Delta_{1}^{1}) \cup (Q \times \{0,1\}) * \Delta^{0} \right) \xrightarrow{[\alpha, \overline{Ff}]} (\mathcal{C}, W)$$

is the desired lift.

From Theorem 6.10, we deduce Theorem 6.5.

Proof of Theorem 6.5. For (1), we apply Theorem 6.10 to the inclusion of markings  $F(W) \subseteq L^{-1}(W)$  and the endofunctor  $FL: \mathcal{C} \to \mathcal{C}$ . The required homotopy is given by the unit  $\eta: id \to RL$ . 

Statement (2) is dual.

We show that if a quasicategory satisfies CLF then its arrow category does as well. To do this, we make use of the following lemma.

**Lemma 6.11.** Let (X, W) be a marked simplicial set.

1. For  $n \ge 2$  and  $0 \le k \le n$ , the marked simplicial set (X, W) has the right lifting property against  $L_{k+1}^{n+1} \xrightarrow{-} {}^{*}L_{k+1}^{n+1}$  if and only if the map

$$i_0^* \colon (X, W)^{(\Delta^1)^{\#}} \to (X, W)$$

has the right lifting property against  $L_k^n \hookrightarrow {}^*L_k^n$ .

2. For  $n \ge 2$  and  $0 \le k \le n$ , the following are equivalent, the marked simplicial set (X, W) has the right lifting property against  $R_k^{n+1} \hookrightarrow *R_k^{n+1}$  if and only if the map

$$i_1^* \colon (X, W)^{(\Delta^1)^{\#}} \to (X, W)$$

has the right lifting property against  $R_{k+1}^n \hookrightarrow {}^*\!R_{k+1}^n$ .

*Proof.* We prove (1), as (2) is dual.

The data of a commutative square

$$\begin{array}{ccc} L^n_k & \stackrel{u}{\longrightarrow} & (X,W)^{(\Delta^1)^{\#}} \\ & & & & \downarrow^{i^*_0} \\ & & & \downarrow^{i^*_0} \\ & ^*\!L^n_k & \stackrel{v}{\longrightarrow} & (X,W) \end{array}$$

is exactly the data of a map  $\overline{u}: L_k^n \times (\Delta^1)^{\#} \to (X, W)$  and a map  $v: {}^*L_k^n \to (X, W)$  such that v is a lift of the map  $\overline{u} \circ i_0: L_k^n \to (X, W)$ . Such a square admits a lift if and only if the map from the pushout

$$[\overline{u},v]: {}^{*}L_{k}^{n} \cup_{L_{k}^{n}} L_{k}^{n} \times (\Delta^{1})^{\#} \to (X,W)$$

admits a lift to a map  ${}^{*}L_{k}^{n} \times (\Delta^{1})^{\#} \to (X, W)$ . We show the inclusion

$${}^{*}L_{k}^{n} \cup_{L_{k}^{n}} L_{k}^{n} \times (\Delta^{1})^{\#} \hookrightarrow {}^{*}L_{k}^{n} \times (\Delta^{1})^{\#}$$

is naturally isomorphic to the inclusion  $L_{k+1}^{n+1} \hookrightarrow {}^*\!L_{k+1}^{n+1}$ , from which the result follows. Define a map  $f : {}^*\!L_k^n \times (\Delta^1)^{\#} \to {}^*\!L_{k+1}^{n+1}$  by

$$f(A, i) := \begin{cases} \partial_0(A) & \text{if } i = 0\\ \partial_0(A) \cup \{0\} & \text{if } i = 1. \end{cases}$$

This map has an inverse defined by

$$A \mapsto \begin{cases} (\sigma_0(A), 1) & \text{if } 0 \in A \\ (\sigma_0(A), 0) & \text{if } 0 \notin A. \end{cases}$$

This gives an isomorphism  ${}^*L_k^n \times (\Delta^1)^{\#} \cong {}^*L_{k+1}^{n+1}$ . Moreover, the map f restricts to a map

$$f|_{L_k^n \times (\Delta^1)^\#} \colon L_k^n \times (\Delta^1)^\# \to L_{k+1}^{n+1}$$

It remains to show the square

$$\begin{array}{ccc} L_k^n & \stackrel{i_0}{\longrightarrow} & L_k^n \times (\Delta^1)^{\#} \\ & & & & \downarrow^{f|_{L_k^n \times (\Delta^1)^{\#}}} \\ & *L_k^n & \stackrel{\partial_0}{\longrightarrow} & L_{k+1}^{n+1} \end{array}$$

is a pushout. A non-degenerate *n*-simplex  $(A_0 \subsetneq \cdots \subsetneq A_n)$  in  $L_{k+1}^{n+1}$  is in the image of  $\partial_0 \colon {}^*L_k^n \to L_{k+1}^{n+1}$  if and only if  $A_n$  does not contain 0. Moreover, if  $A_n \neq [n+1] - \{0\}$  then this *n*-simplex is also in the image of  $f_i_0$ . This shows the intersection of the images of the bottom and right maps is the image of the composite map. If  $A_n$  contains 0 then  $(A_0 \subsetneq \cdots \subsetneq A_n)$  in  $L_{k+1}^{n+1}$  is in the image under f of  $L_k^n \times (\Delta^1)^{\#}$ , which shows the union of the images of the bottom and right maps is all of  $L_{k+1}^{n+1}$ , which suffices.

**Corollary 6.12.** Let  $(\mathcal{C}, W)$  be a marked quasicategory such that W is strongly closed under composition.

- 1. If  $(\mathcal{C}, W)$  satisfies CLF then the marked arrow category  $(\mathcal{C}, W)^{(\Delta^1)^{\#}}$  satisfies CLF.
- 2. If  $(\mathcal{C}, W)$  satisfies CRF then the marked arrow category  $(\mathcal{C}, W)^{(\Delta^1)^{\#}}$  satisfies CRF.

*Proof.* Marked 1-simplices are weakly closed under composition by assumption. For (1), Lemma 6.11 shows the marked arrow category has the right lifting property with respect to the set  $\{L_k^n \hookrightarrow {}^*L_k^n \mid n \geq 2, 0 < k \leq n\}$ . Statement (2) is dual.

# Model for the localization

#### 7 The marked Ex functor

We now describe the analogue of the category of fractions  $CW^{-1}$  (or  $W^{-1}C$ ) for quasicategories. Given a marked simplicial set (X, W), we form a simplicial set whose 0-simplices are those of X and whose 1simplices are (co)spans of the form  $x \to y' \leftarrow y$  (or  $x \leftarrow x' \to y$ ). This is done using a marked variant of the Ex functor. The main theorem of this section is Theorem 7.5, which generalizes the analogous result for classical calculus of fractions: if  $(\mathcal{C}, W)$  is a marked quasicategory where W is weakly closed under composition then calculus of fractions is a sufficient *and necessary* condition for the "category of fractions" to be a quasicategory.

For  $n \ge 0$ , let  $\mathrm{sd}_+[n]$  be the poset  $\mathcal{P}_{\neq \varnothing}[n]$  of non-empty subsets of [n] (regarded as a simplicial set). We view a 1-simplex  $A_0 \subseteq A_1$  as marked if  $\max A_0 = \max A_1$ . This gives the data of a cosimplicial object  $\Delta \to \mathsf{sSet}_+$ , with simplicial operators defined using the image.

**Example 7.1.** We give example depictions of  $sd_{+}[0]$ ,  $sd_{+}[1]$ , and  $sd_{+}[2]$ , respectively.



The marked simplicial sets  $sd_+[0]$ ,  $sd_+[1]$ , and  $sd_+[2]$ .

**Definition 7.2.** The marked subdivision functor  $Sd_+: sSet \to sSet_+$  is the extension by colimits of the cosimplicial object  $sd_+: \Delta \to sSet_+$ .

$$\begin{array}{c} \Delta \xrightarrow{\mathrm{sd}_+} \mathrm{sSet}_+ \\ \downarrow & \stackrel{\gamma}{\underset{\mathrm{Sd}_+}{}} \mathrm{sSet} \end{array}$$

The marked Ex-functor is its right adjoint  $Ex_+: sSet_+ \rightarrow sSet$ , defined by

$$\operatorname{Ex}_{+}(X, W)_{n} := \mathsf{sSet}_{+}(\operatorname{sd}_{+}[n], (X, W)).$$

Remark 7.3. The composite

$$sSet \xrightarrow{(-)^{\#}} sSet_{+} \xrightarrow{Ex_{+}} sSet$$

recovers Kan's original Ex-functor [Kan57].

We write  $\operatorname{sd}_{+}^{\operatorname{op}}[n]$  for the opposite poset of  $\operatorname{sd}_{+}[n]$ . We view a 1-simplex  $A_0 \supseteq A_1$  of  $\operatorname{sd}_{+}^{\operatorname{op}}[n]$  as marked if  $\min A_0 = \min A_1$ .

**Example 7.4.** Below are example depictions of  $sd_{+}^{op}[0]$ ,  $sd_{+}^{op}[1]$ , and  $sd_{+}^{op}[2]$ , respectively.



The marked simplicial sets  $sd_{\pm}^{op}[0]$ ,  $sd_{\pm}^{op}[1]$ , and  $sd_{\pm}^{op}[2]$ .

The respective extension by colimits is denoted  $\mathrm{Sd}_{+}^{\mathrm{op}}$ :  $\mathsf{sSet} \to \mathsf{sSet}_{+}$ , and its right adjoint  $\mathrm{Ex}_{+}^{\mathrm{op}}$ :  $\mathsf{sSet}_{+} \to \mathsf{sSet}$  is defined by

$$\operatorname{Ex}_{+}^{\operatorname{op}}(X, W)_{n} := \mathsf{sSet}_{+}(\operatorname{sd}_{+}^{\operatorname{op}}[n], (X, W)).$$

By construction, we have a natural isomorphism  $\operatorname{Sd}^{\operatorname{op}}_+\Delta^n \cong (\operatorname{Sd}_+(\Delta^n)^{\operatorname{op}})^{\operatorname{op}}$  of cosimplicial objects. This induces isomorphisms

 $\operatorname{Sd}_{+}^{\operatorname{op}} X \cong (\operatorname{Sd}_{+} X^{\operatorname{op}})^{\operatorname{op}}, \quad \operatorname{Ex}_{+}^{\operatorname{op}}(X, W) \cong (\operatorname{Ex}_{+}(X^{\operatorname{op}}, W))^{\operatorname{op}}$ 

natural in  $X \in \mathsf{sSet}$  and  $(X, W) \in \mathsf{sSet}_+$ , respectively.

The goal of this section is to prove the following result.

**Theorem 7.5.** Let (X, W) be a marked simplicial set.

- 1. a) If  $\text{Ex}_+(X, W)$  is a quasicategory then (X, W) has the right lifting property with respect to the set  $\{L_k^n \hookrightarrow {}^*L_k^n \mid n \ge 2, \ 0 < k \le n\}$ .
  - b) If X is a quasicategory and (X, W) satisfies CLF then  $Ex_+(X, W)$  is a quasicategory.
- 2. a) If  $\operatorname{Ex}_{+}^{\operatorname{op}}(X,W)$  is a quasicategory then (X,W) has the right lifting property with respect to the set  $\{R_{k}^{n} \hookrightarrow {}^{*}R_{k}^{n} \mid n \geq 2, \ 0 \leq k < n\}$ .
  - b) If X is a quasicategory and (X, W) satisfies CRF then  $\operatorname{Ex}^{\operatorname{op}}_+(X, W)$  is a quasicategory.

To prove this, we establish some auxiliary definitions and results. As Theorem 7.5 is our primary focus in this section, we work with calculus of left fractions and  $Sd_+: sSet \rightarrow sSet_+$ , omitting the dual statements.

We identify  $\operatorname{Sd}_+ \Lambda_k^n$  with the maximal simplicial subset of  $\operatorname{Sd}_+ \Delta^n$  which omits the 0-simplices [n] and  $[n] - \{k\}$ . Note that there are inclusions  ${}^*L_k^n \subseteq \operatorname{Sd}_+ \Delta^n$  and  $L_k^n \subseteq \operatorname{Sd}_+ \Lambda_i^n$ .

**Proposition 7.6.** For  $n \ge 1$  and  $0 \le k \le n$ , the inclusion  $L_k^n \hookrightarrow {}^*L_k^n$  is a retract of the inclusion  $\operatorname{Sd}_+ \Lambda_k^n \hookrightarrow \operatorname{Sd}_+ \Delta^n$ .

*Proof.* The inclusion  ${}^{*}L_{k}^{n} \subseteq \operatorname{Sd}_{+}\Delta^{n}$  admits a retraction defined, as a poset map, by  $A \mapsto A \cup \{k\}$ . The restriction of this retraction to  $\operatorname{Sd}_{+}\Lambda_{k}^{n}$  factors through  $L_{k}^{n}$ , as desired.

**Definition 7.7.** For  $n \ge 0$  and 0 < k < n,

- 1. let  $J_k^n \subseteq \operatorname{Sd}_+ \Delta^n$  be the marked simplicial subset which contains  $\operatorname{Sd}_+ \Lambda_i^n$  and, for  $k \ge 0$ , all *m*-simplices  $(A_0 \subseteq \cdots \subseteq A_m)$  such that  $k \in A_0$ ; and
- 2. let  $K_k^n \subseteq \text{Sd}_+ \Delta^n$  be the maximal marked simplicial subset which omits the 0-simplex  $[n] \{k\}$ .

By definition, there is a sequence of inclusions

$$\operatorname{Sd}_{+}\Lambda_{k}^{n} \subseteq J_{k}^{n} \subseteq K_{k}^{n} \subseteq \operatorname{Sd}_{+}\Delta^{n}$$

Note there is an inclusion  ${}^*L_k^n \subseteq J_k^n$  which restricts to an inclusion  $L_k^n \subseteq \operatorname{Sd}_+ \Lambda_i^n$ .

**Proposition 7.8.** For  $n \ge 2$  and  $0 \le k \le n$ , the square

$$\begin{array}{ccc} L_k^n & \longrightarrow & \mathrm{Sd}_+ \Lambda_k^n \\ & & & & \downarrow \\ *L_k^n & \longrightarrow & J_k^n \end{array}$$

is a pushout.

*Proof.* Follows since the simplices in  $J_k^n$  which are not contained in  $\mathrm{Sd}_+ \Lambda_i^n$  are exactly the chains  $A_0 \subseteq \cdots \subseteq A_m$  where  $A_m = [n]$  and all subsets contain k (i.e.  $k \in A_0$ ).

**Corollary 7.9.** Let (X, W) be a marked simplicial set. If (X, W) has the right lifting property against  $\{L_k^n \hookrightarrow {}^*L_k^n \mid n \ge 2, \ 0 < k \le n\}$  then it has the right lifting property against

$$\{ \mathrm{Sd}_+ \Lambda_k^n \hookrightarrow J_k^n \mid n \ge 2, \ 0 < k \le n \}$$

**Proposition 7.10.** For  $n \ge 2$  and  $0 \le k < n$ , the inclusion  $K_k^n \subseteq \text{Sd}_+ \Delta^n$  admits a retraction.

*Proof.* The retraction is defined by

$$S \mapsto \begin{cases} [n] & \text{if } S = [n] - \{i\} \\ S & \text{otherwise.} \end{cases}$$

This map preserves marked 1-simplices as  $i \neq n$  by assumption.

**Lemma 7.11.** The inclusion  $J_k^n \subseteq K_k^n$  is in the saturation of the set

$$\{(\Lambda_k^n)^{\flat} \hookrightarrow (\Delta^n)^{\flat} \mid n \ge 2, \ 0 < k < n\} \cup \{(\Lambda_1^2)^{\#} \hookrightarrow (\Delta^2)^{\#}\}.$$

*Proof.* We construct a simple inner horn decomposition and apply Lemma 6.8.

The simplices of  $K_k^n$  which are not contained in  $J_k^n$  are exactly chains  $(A_0 \subseteq \cdots \subseteq A_m)$  such that  $A_m = [n]$  and  $k \notin A_0$ . For  $m \ge 2$  and  $j = 1, \ldots, m-1$ , define subsets  $S_j^m \subseteq (K_k^n)_m$  of non-degenerate *m*-simplices by

$$S_j^m := \left\{ (A_0 \subsetneq \cdots \subsetneq A_m) \in (K_k^n)_m \mid \begin{array}{c} A_m = [n] \\ k \not\in A_{j-1} \\ A_j = A_{j-1} \cup \{k\} \end{array} \right\}.$$

For  $m \ge 1$  and  $j = 1, \ldots, m$ , define subsets  $T_j^m \subseteq (K_k^n)_m$  of non-degenerate *m*-simplices by

$$T_j^m = \left\{ (A_0 \subsetneq \cdots \subsetneq A_m) \in (K_k^n)_m \mid \begin{array}{c} A_m = [n] \\ k \not\in A_{j-1} \\ A_j \supsetneq A_{j-1} \cup \{k\} \end{array} \right\}.$$

We show that

$$(\{S_j^m\}_{j=1}^{m-1}, \{T_j^m\}_{j=1}^m, \mathrm{id}_{\{1,\dots,m-1\}})$$

is a simple inner horn decomposition for the inclusion  $J_k^n \subseteq K_k^n$ . Fix  $m \ge 2$  and  $j \in \{1, \ldots, m-1\}$ .

1. Every simplex  $(A_0 \subsetneq \cdots \subsetneq A_m)$  in  $S_j^m$  is non-degenerate, thus  $A_{j+1}$  contains an element which is not contained in  $A_j = A_{j-1} \cup \{k\}$ . This shows the image of  $\partial_j|_{S_j^m}$  is contained in  $T_j^{m-1}$ . The inverse function  $T_j^{m-1} \to S_j^m$  is defined by

$$(A_0 \subsetneq \cdots \subsetneq A_{m-1}) \mapsto (A_0 \subsetneq \cdots \subsetneq A_{j-1} \subsetneq A_{j-1} \cup \{i\} \subsetneq A_j \subsetneq \cdots \subsetneq A_{m-1}).$$

- 2. If  $(A_0 \subsetneq A_1 \subsetneq A_2)$  is such that  $\max A_0 = \max A_2$ , then  $\max A_0 \le \max A_1$  and  $\max A_1 \le \max A_2$  implies  $(A_0 \subsetneq A_1 \subsetneq A_2)$  is maximally-marked.
- 3. For  $u \in S_j^m$  and  $l \neq j$ , we proceed by case analysis on  $u\partial_l$ :
  - if l < j then  $u\partial_l \in S_{j-1}^{m-1}$ ;
  - if l > j then  $f \partial_l \in S_j^{m-1}$ ;
  - if l = m then  $u\partial_l \in \mathrm{Sd}_+ \Lambda_i^n$  since, writing  $u = (A_0 \subseteq \cdots \subseteq A_m)$ , we have that  $A_{m-1} \neq [n]$  as u is non-degenerate.

With this, we may prove Theorem 7.5.

Proof of Theorem 7.5. Statement (1) follows from Proposition 7.6.

For (2), fix a map  $\Lambda_k^n \to \text{Ex}_+(X, W)$  for  $n \ge 2$  and 0 < i < n. By adjointness, it suffices to show the transpose  $\text{Sd}_+ \Lambda_k^n \to (X, W)$  admits a lift to  $\text{Sd}_+ \Delta^n \to (X, W)$ . A lift from  $\text{Sd}_+ \Lambda_i^n$  to  $J_k^n$  exists by Corollary 7.9. A lift from  $J_k^n$  to  $K_k^n$  exists by Lemma 7.11. As  $\text{Sd}_+ \Delta^n$  retracts onto  $K_k^n$  by Proposition 7.10, this gives the desired lift.



## 8 Localization at equivalences via marked Ex

In this section, we move towards showing that if  $(\mathcal{C}, W)$  satisfies CLF then  $\operatorname{Ex}_+(\mathcal{C}, W)$  is a model for the localization of  $\mathcal{C}$  at W. We construct the canonical map  $\mathcal{C} \to \operatorname{Ex}_+(\mathcal{C}, W)$  (or  $\mathcal{C} \to \operatorname{Ex}_+^{\operatorname{op}}(\mathcal{C}, W)$ ) and show that it sends weak equivalences to equivalences (Proposition 8.1). We then consider the case where W is exactly the set of equivalences in  $\mathcal{C}$ . In this case, one has that the localization is a categorical equivalence (Corollary 9.5), which we prove by adapting the argument given in [LTW79, Thm. 4.1] to show that  $X \to \operatorname{Ex} X$  is a weak homotopy equivalence.

The map of posets max:  $\mathrm{sd}_+[n] \to [n]$  is natural in n; that is, it induces a natural transformation between functors  $\Delta \to \mathrm{sSet}_+$ . For a simplicial set X, this induces a map

$$\max_{!} \colon \operatorname{Sd}_{+} X \to X^{\natural}$$

natural in X via extension by colimits. For a marked simplicial set (X, W), pre-composition by max induces a map

$$\max^* \colon X \to \operatorname{Ex}_+(X, W)$$

which sends an *n*-simplex  $f: \Delta^n \to X$  of X to the *n*-simplex  $f \circ \max: \operatorname{Sd}_+ \Delta^n \to (X, W)$  of  $\operatorname{Ex}_+(X, W)$ . Dually, the map min:  $\operatorname{sd}_+^{\operatorname{op}}[n] \to [n]$  induces a map

$$\min_{!} \colon \operatorname{Sd}_{\perp}^{\operatorname{op}} X \to X^{\flat}$$

and a map

$$\min^* \colon X \to \operatorname{Ex}^{\operatorname{op}}_+(X, W).$$

The map  $\max^* \colon \mathcal{C} \to \operatorname{Ex}_+(\mathcal{C}, W)$  (and its dual) will be our candidate for the localization map. Note that for a weak equivalence  $w \in W$ , the diagram



defines a map  $J \to \text{Ex}_+(X, W)$  which witnesses that  $(\max^*)_1(w) \in \text{Ex}_+(X, W)_1$  is invertible. Dualizing this diagram gives that

**Proposition 8.1.** For a marked simplicial set (X, W), the maps

$$\max^* \colon X \to \operatorname{Ex}_+(X, W), \quad \min^* \colon X \to \operatorname{Ex}_+^{\operatorname{op}}(X, W)$$

send weak equivalences to J-equivalences.

Via Proposition 8.1, the maps

$$\max^* \colon X \to \operatorname{Ex}_+(X, W), \quad \min^* \colon X \to \operatorname{Ex}_+^{\operatorname{op}}(X, W)$$

ascend to marked maps

$$\overline{\max^*} \colon (X, W) \to \operatorname{Ex}_+(X, W)^{\natural}, \quad \overline{\min^*} \colon (X, W) \to \operatorname{Ex}_+^{\operatorname{op}}(X, W)^{\natural}$$

A further corollary of Proposition 8.1 is that the functors  $Ex_+, Ex_+^{op}: \mathsf{sSet}_+ \to \mathsf{sSet}$  send marked homotopy equivalances to *J*-homotopy equivalences.

**Proposition 8.2.** The functors  $Ex_+, Ex_+^{op}$ :  $sSet_+ \rightarrow sSet$  send marked homotopy equivalences to *J*-homotopy equivalences.

*Proof.* For  $Ex_+$ , the map

$$J \to \operatorname{Ex}_+ (\Delta^1)^{\#}$$

which witnesses the invertibility of the 1-simplex  $(\max^*)_1(\mathrm{id}_{(\Delta^1)^{\#}}) \in \mathrm{Ex}_+((\Delta^1)^{\#})_1$  preserves endpoints. Thus, a marked homotopy

$$\alpha \colon (X, W) \times (\Delta^1)^{\#} \to (X', W')$$

induces a *J*-homotopy

$$\operatorname{Ex}_{+}(X,W) \times J \to \operatorname{Ex}_{+}(X,W) \times \operatorname{Ex}_{+}(\Delta^{1})^{\#} \to \operatorname{Ex}_{+}(X',W').$$

The argument for  $Ex^{op}_{+}$  is analogous.

In order to show  $\text{Ex}_+(\mathcal{C}, W)$  computes the localization, we first consider the case where W is exactly the equivalences in  $\mathcal{C}$ . In this case, we wish to show  $\text{Ex}_+ \mathcal{C}^{\natural}$  is weakly equivalent to  $\mathcal{C}$ . To show this, we prove the following lemma.

**Lemma 8.3.** For  $n \ge 0$ , the map

max: 
$$\operatorname{Sd}_+ \Delta^n \to (\Delta^n)^{\flat}$$

is a marked homotopy equivalence.

*Proof.* The map of posets max:  $sd_+[n] \to [n]$  has a section  $f: [n] \to sd_+[n]$  defined by

$$f(i) := \{0, \ldots, i\}.$$

For  $A \in \mathrm{sd}_+[n]$ , the inclusion  $A \subseteq \{0, \ldots, \max A\}$  induces a marked homotopy  $\mathrm{Sd}_+ \Delta^n \times (\Delta^1)^{\#} \to \mathrm{Sd}_+ \Delta^n$  from the composite  $f \circ \max$  to the identity map on  $\mathrm{Sd}_+ \Delta^n$ .

Our proof that  $\max^* \colon \mathcal{C} \to \operatorname{Ex}_+ \mathcal{C}^{\natural}$  is a categorical equivalence uses an analogous argument to [LTW79, Thm. 4.1], which shows  $X \to \operatorname{Ex} X$  is a Kan–Quillen weak equivalence. In particular, we make use of the generalized diagonal lemma of [CKW23, Thm. 2.5] and specifically its instantiation to the Joyal model structure [CKW23, Ex. 3.6].

**Theorem 8.4.** For a quasicategory C, the maps

$$\max^* \colon \mathcal{C} \to \operatorname{Ex}_+ \mathcal{C}^{\natural}, \quad \min^* \colon \mathcal{C} \to \operatorname{Ex}_+^{\operatorname{op}} \mathcal{C}^{\natural}$$

are equivalences of quasicategories.

*Proof.* We show the map max<sup>\*</sup> is an equivalence. The argument for min<sup>\*</sup> is dual. We regard the square

as a diagram of bisimplicial sets by varying m and n. Fixing n, the left map in the square

$$\begin{array}{c} \mathcal{C} & \longrightarrow \operatorname{Ex}_{+} C^{\natural} \\ \downarrow & \downarrow \\ U((\mathcal{C}^{\natural})^{(\Delta^{n})^{\#}}) & \longrightarrow \operatorname{Ex}_{+}((\mathcal{C}^{\natural})^{(\Delta^{n})^{\#}}) \end{array}$$

is obtained by applying the forgetful functor  $U: \mathsf{sSet}_+ \to \mathsf{sSet}$  to the map

$$!^* \colon (\mathcal{C}^{\natural})^{\Delta^0} \to (\mathcal{C}^{\natural})^{(\Delta^n)^{\#}}$$

This map is a marked homotopy equivalence as  $(\mathcal{C}^{\natural})^{-}: (\mathsf{sSet}_{+})^{\mathrm{op}} \to \mathsf{sSet}_{+}$  preserves marked homotopies. The codomain is a quasicategory since  $\mathcal{C}$  is. A marked 1-simplex of the codomain  $H: (\Delta^{n})^{\#} \times (\Delta^{1})^{\#} \to \mathcal{C}^{\natural}$  is a marked homotopy. As  $\mathcal{C}$  is a quasicategory marked at equivalences, every such H ascends to an  $E^{1}$ -homotopy, thus the codomain is marked at exactly equivalences. This gives that !\* is a weak equivalance between fibrant-cofibrant objects in  $\mathsf{sSet}_{+}$ , hence the left map is an equivalence of quasicategories. The right map is obtained by applying  $\mathrm{Ex}_{+}: \mathsf{sSet}_{+} \to \mathsf{sSet}$  to !\*. By Proposition 8.2, the right map is a weak equivalence.

Fixing m, the bottom map

$$\begin{array}{c} & & & & \\ \downarrow & & & \downarrow \\ \operatorname{Core}_{+}\left( (\mathcal{C}^{\natural})^{(\Delta^{m})^{\flat}} \right) \longrightarrow \operatorname{Core}_{+}\left( (\mathcal{C}^{\natural})^{\operatorname{Sd}_{+}\Delta^{m}} \right)
\end{array}$$

is obtained by applying  $\operatorname{Core}_+: \mathsf{sSet}_+ \to \mathsf{sSet}$  to the map

$$(\mathcal{C}^{\natural})^{\max} \colon (\mathcal{C}^{\natural})^{(\Delta^m)^{\flat}} \to (\mathcal{C}^{\natural})^{\mathrm{Sd}_{+}\Delta^m}.$$

This map is a marked homotopy equivalence by Lemma 8.3. It is, moreover, a map between quasicategories marked at equivalences, hence this map is an equivalence of quasicategories. Thus, it is an equivalence of categories after applying Core<sub>+</sub>.

Applying the diagonal functor to this square,

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \operatorname{Ex}_{+} C^{\natural} \\ \sim & & \downarrow \sim \\ \cdot & & & \ddots \\ \cdot & & & \cdot \end{array}$$

the left, right, and bottom maps are weak equivalences by [CKW23, Ex. 3.6]. Thus, the top map is a weak equivalence by 2-out-of-3.  $\hfill \Box$ 

#### 9 Localization via marked Ex

In this section, we show that  $\text{Ex}_{+}(\mathcal{C}, W)$  and  $\text{Ex}_{+}^{\text{op}}(\mathcal{C}, W)$  are models for the localization for an arbitrary W, using the "base case" where W is the set of equivalences. Our main theorem is Theorem 9.6, which is the quasicategorical analogue of Theorem 1.7. In particular, this result requires multiple auxiliary lemmas, most important of which is Lemma 9.3. As these statements are more technical in nature and are only used to prove the main theorem of this section (Theorem 9.6), we state Lemmas 9.1 and 9.2 and Corollaries 9.4 and 9.5 only for  $\text{Ex}_{+}$ , omitting the dual versions.

**Lemma 9.1.** For a marked simplicial set (X, W), the composite

$$\operatorname{Sd}_+ X \xrightarrow{\max_!} X^{\flat} \hookrightarrow (X, W)$$

is adjoint transpose to the map

$$X \xrightarrow{\max} \operatorname{Ex}_+(X, W)$$

under the  $(Sd_+ \dashv Ex_+)$ -adjunction.

*Proof.* We wish to show the diagram

$$\begin{array}{c} \operatorname{Sd}_{+} X \xrightarrow{\max_{!}} X^{\flat} \\ \operatorname{Sd}_{+} \max^{*} \downarrow & \downarrow \\ \operatorname{Sd}_{+} \operatorname{Ex}_{+}(X, W) \xrightarrow{\varepsilon_{(X, W)}} (X, W) \end{array}$$

commutes. It suffices to show it commutes after applying U. That is, it suffices to show the triangle

$$U\operatorname{Sd}_{+} X \xrightarrow{\max_{!}} X$$

$$\operatorname{Sd}_{+} \operatorname{max}^{*} U\operatorname{Sd}_{+} \operatorname{Ex}_{+}(X, W)$$

$$\varepsilon_{(X, W)}$$

commutes.

Recall the *n*-simplices of  $Sd_+ X$  are pairs (u, v) where

- $u: \Delta^m \to X$  is an *m*-simplex of X for some  $m \ge 0$ ; and
- $v: \Delta^n \to \operatorname{Sd}_+ \Delta^m$  is an *n*-simplex of  $\operatorname{Sd}_+ \Delta^m$

subject to the identification  $(u\varphi, v) = (u, \mathrm{Sd}_+(\varphi) \circ v)$  for any map  $\varphi \colon [m] \to [m']$  in the simplex category. The top map is given by the formula

$$(\max_{!})_{n}(u,v) = u \circ \max \circ v.$$

We calculate the bottom composite as

$$(\varepsilon_{(X,W)})_m((\mathrm{Sd}_+\max^*)_m(u,v)) = (\varepsilon_{(X,W)})_m(u \circ \max, v)$$
$$= (u \circ \max) \circ v.$$

**Lemma 9.2.** For a marked simplicial set (X, W), the squares

and
commute for all  $n \geq 0$ .

Proof. For the first square, the top square in

commutes by definition and the bottom square commutes by naturality of adjunction bijections. The desired square is identical to the outer square in this diagram.

For the second square, we show that each sub-diagram of

$$sSet(\Delta^{n}, Ex_{+}(X, W)) \xrightarrow{Ex_{+}(\eta_{(X,W)})_{*}} sSet(\Delta^{n}, Ex_{+} X^{\#}) \xrightarrow{Ex_{+}((\max^{*})^{\#})_{*}} sSet(\Delta^{n}, Ex_{+}(Ex_{+}(X, W)^{\#})) \xrightarrow{\cong \downarrow} sSet_{+}(Sd_{+} \Delta^{n}, (X, W)) \xrightarrow{(\eta_{(X,W)})_{*}} sSet_{+}(Sd_{+} \Delta^{n}, X^{\#}) \xrightarrow{((\max^{*})^{\#})_{*}} sSet_{+}(Sd_{+} \Delta^{n}, Ex_{+}(X, W)^{\#}) \xrightarrow{\cong \downarrow \varphi} sSet_{+}(U(Sd_{+} \Delta^{n}), X) \xrightarrow{(\max^{*})_{*}} sSet_{+}(U(Sd_{+} \Delta^{n}), Ex_{+}(X, W)) \xrightarrow{\cong \downarrow \psi} \theta_{\downarrow} \cong sSet_{+}((USd_{+} \Delta^{n})^{\flat}, (X, W)) \xrightarrow{(\max_{*})^{*}} sSet_{+}(Sd_{+}(USd_{+} \Delta^{n}), (X, W))$$

commutes. The desired equality is exactly the outer composite.

The top left and right squares commute by naturality of adjunction bijections. The middle right square commutes by definition. We show the bottom left triangle and bottom right square commute by explicit computation.

For the bottom left triangle, fix  $f: \operatorname{Sd}_+ \Delta^n \to (X, W)$ . We apply the forgetful functor  $U: \operatorname{sSet}_+ \to \operatorname{sSet}$ and calculate

$$U(\psi\varphi(\eta_{(X,W)} \circ f)) = U(\psi(\varphi(\eta_{(X,W)}) \circ Uf))$$
  
=  $U(\psi(Uf))$   
=  $Uf$  (\*)  
=  $Uf \circ \mathrm{id}$   
=  $Uf \circ U\varepsilon_{\mathrm{Sd}_{+}\Delta^{n}}$   
=  $U(f \circ \varepsilon_{\mathrm{Sd}_{+}\Delta^{n}}),$ 

where (\*) holds since applying U to any map g obtained by the adjunction bijection  $\psi$  is exactly g. For the bottom right square, fix  $f: U \operatorname{Sd}_+ \Delta^n \to X$ . We calculate

$$\begin{split} \theta(\max^* \circ f) &= \theta(\max^*) \circ \operatorname{Sd}_+ f \\ &= \varepsilon_X \circ \max_! \circ \operatorname{Sd}_+ f \quad \text{by Lemma 9.1} \\ &= \varepsilon_X \circ f \circ \max_! \qquad \text{by naturality} \\ &= \theta(f) \circ \max_!, \end{split}$$

where  $\varepsilon_X$  denotes the counit  $X^{\flat} \to (X, W)$  of the minimal marking adjunction.

**Lemma 9.3.** For a marked simplicial set (X, W),

1. the maps

$$\operatorname{Ex}_{+}(X,W) \xrightarrow{\operatorname{Ex}_{+}(\operatorname{\overline{\max}}^{*})} \operatorname{Ex}_{+} \left( \operatorname{Ex}_{+}(X,W)^{\natural} \right)$$

are J-homotopic; and

2. the maps

$$\operatorname{Ex}^{\operatorname{op}}_{+}(X,W) \xrightarrow[]{\operatorname{Ex}_{+}(\operatorname{\overline{min}^{*}})}_{\operatorname{\overline{min}^{*}}} \operatorname{Ex}^{\operatorname{op}}_{+}\left(\operatorname{Ex}^{\operatorname{op}}_{+}(X,W)^{\natural}\right)$$

are J-homotopic.

*Proof.* We show (1), as (2) is dual.

For  $n \ge 0$ , consider the (non-commuting) triangle

$$\operatorname{Sd}_{+}(U\operatorname{Sd}_{+}\Delta^{n}) \xrightarrow{\operatorname{Sd}_{+}(U\operatorname{max})} \operatorname{Sd}_{+}\Delta^{n}$$

$$(*)$$

$$(U\operatorname{Sd}_{+}\Delta^{n})^{\flat}$$

The underlying simplicial set of the right and middle terms is the poset of subsets of [n] ordered by containment. The underlying simplicial set of the left term is the poset of chains  $A_0 \subseteq \cdots \subseteq A_k$  of subsets of [n] ordered by containment. We identify the (underlying) maps above as monotone functions



defined by

$$\operatorname{Sd}_{+}(U\max)(A_{0} \subseteq \cdots \subseteq A_{k}) = \{\max A_{0}, \dots, \max A_{k}\}$$
$$\max_{l}(A_{0} \subseteq \cdots \subseteq A_{k}) = A_{k}.$$

The natural containment  $\{\max A_0, \ldots, \max A_k\} \subseteq A_k$  induces a marked homotopy

 $\operatorname{Sd}_+(U\operatorname{Sd}_+\Delta^n) \times (\Delta^1)^{\#} \to \operatorname{Sd}_+\Delta^n$ 

between the composites in Diagram (\*). The map

$$\operatorname{Sd}_+ J \to (\Delta^1)^{\#}$$

which witnesses invertibility of  $(\max^*)_1(id_{(\Delta^1)^{\#}}) \in (Ex_+ (\Delta^1)^{\#})_1$  induces a map

$$\operatorname{Sd}_+(U\operatorname{Sd}_+J) \xrightarrow{\max_1} \operatorname{Sd}_+J \longrightarrow (\Delta^1)^{\#}.$$

Let  $\alpha$  denote the composite

$$\mathrm{Sd}_{+}(U\,\mathrm{Sd}_{+}(\Delta^{n}\times J)) \longrightarrow \mathrm{Sd}_{+}(U\,\mathrm{Sd}_{+}\,\Delta^{n})\times\mathrm{Sd}_{+}(U\,\mathrm{Sd}_{+}\,J) \longrightarrow \mathrm{Sd}_{+}(U\,\mathrm{Sd}_{+}\,\Delta^{n})\times(\Delta^{1})^{\#} \longrightarrow \mathrm{Sd}_{+}\Delta^{n}$$

(where the first map is induced by the universal property of the product). Pre-composition by  $\alpha$  induces a map

$$\alpha^* \colon \operatorname{Ex}_+(X, W) \to \operatorname{Ex}_+\left(\operatorname{Ex}_+(X, W)^{\#}\right)^J.$$

By Lemma 9.2, this map witnesses that the diagram

$$\operatorname{Ex}_{+}(X,W) \xrightarrow{\max^{*}} \operatorname{Ex}_{+}\left(\operatorname{Ex}_{+}(K,W)^{\#}\right) \\
 \overbrace{\operatorname{Ex}_{+}\eta_{(X,W)}} \xrightarrow{\operatorname{Ex}_{+}(X^{\#})} \\
 \operatorname{Ex}_{+}X^{\#} \xrightarrow{\operatorname{Ex}_{+}((\max^{*})^{\#})}$$

commutes up to *J*-homotopy.

Note that, for  $f: \operatorname{Sd}_+ \Delta^n \to (X, W)$ , the map

$$f\alpha\colon \mathrm{Sd}_+(U\,\mathrm{Sd}_+\,\Delta^n)\to(X,W)$$

sends weak equivalences to equivalences as f does by Proposition 8.1. Thus,  $\alpha^*$  factors as

$$\operatorname{Ex}_{+}(X,W) \times J \xrightarrow{\alpha^{*}} \operatorname{Ex}_{+}(\operatorname{Ex}_{+}(X,W)^{\#})$$

$$\beta \xrightarrow{\beta} \operatorname{Ex}_{+}(\operatorname{Ex}_{+}(X,W)^{\natural})$$

We show that  $\beta$  is the desired homotopy by verifying its endpoints evaluate to max<sup>\*</sup> and Ex<sub>+</sub>(max<sup>\*</sup>), respectively. The diagram

$$\operatorname{Ex}_{+}(X,W) \xrightarrow[\max_{\operatorname{Ex}_{+}(X,W)^{\sharp}}]{\operatorname{Ex}_{+}(X,W)^{\sharp}} \operatorname{Ex}_{+}(\operatorname{Ex}_{+}(X,W)^{\sharp})$$
$$\xrightarrow{\operatorname{Ex}_{+}(X,W)^{\#}} \operatorname{Ex}_{+}(\operatorname{Ex}_{+}(X,W)^{\#})$$

commutes by naturality of max<sup>\*</sup>:  $U \Rightarrow Ex_+$  applied to the inclusion  $Ex_+(X, W)^{\natural} \hookrightarrow Ex_+(X, W)^{\#}$ . From this, we have that

$$\operatorname{Ex}_{+} \eta \circ \max_{\operatorname{Ex}_{+}(X,W)^{\natural}}^{*} = \max_{\operatorname{Ex}_{+}(X,W)^{\#}}^{*} = \alpha^{*}|_{\operatorname{Ex}_{+}(X,W) \times \{0\}} = \operatorname{Ex}_{+} \eta \circ \beta|_{\operatorname{Ex}_{+}(X,W) \times \{0\}},$$

which implies  $\max_{\text{Ex}_{+}(X,W)^{\natural}}^{*} = \beta|_{\text{Ex}_{+}(X,W) \times \{0\}}$  as  $\text{Ex}_{+} \eta$  is monic. For the other endpoint, the diagram

$$\begin{array}{ccc} (X,W) & \xrightarrow{\max^*} & \operatorname{Ex}_+(X,W)^{\natural} \\ & & & & & & \\ \eta' & & & & & & \\ X^{\#} & \xrightarrow{(\max^*)^{\#}} & \operatorname{Ex}_+(X,W)^{\#} \end{array}$$

commutes as it commutes after applying  $U: \mathsf{sSet}_+ \to \mathsf{sSet}$ . Thus, this diagram commutes after applying  $\mathrm{Ex}_+:\mathsf{sSet}_+ \to \mathsf{sSet}$ . Hence,

$$\operatorname{Ex}_{+} \eta \circ \operatorname{Ex}_{+} \overline{\operatorname{max}^{*}} = \operatorname{Ex}_{+} (\operatorname{max}^{*})^{\#} \circ \operatorname{Ex}_{+} \eta' = \alpha^{*}|_{\operatorname{Ex}_{+}(X,W) \times \{1\}} = \operatorname{Ex}_{+} \eta \circ \beta|_{\operatorname{Ex}_{+}(X,W) \times \{1\}},$$

which implies  $\operatorname{Ex}_{+} \overline{\operatorname{max}^{*}} = \beta|_{\operatorname{Ex}_{+}(X,W) \times \{1\}}$  as  $\operatorname{Ex}_{+} \eta$  is monic.

**Corollary 9.4.** For a marked quasicategory  $(\mathcal{C}, W)$  satisfying CLF, the (underlying) maps in the diagram

$$\operatorname{Ex}_{+}(\mathcal{C},W)^{\natural} \xrightarrow{\operatorname{Ex}_{+}(\overline{\max^{*}})^{\natural}} \operatorname{Ex}_{+} \left( \operatorname{Ex}_{+}(\mathcal{C},W)^{\natural} \right)^{\natural}$$

are  $E^1$ -homotopic.

*Proof.* The underlying maps are J-homotopic by Lemma 9.3. The result follows since the codomain is a quasicategory.  $\Box$ 

**Corollary 9.5.** For a marked quasicategory  $(\mathcal{C}, W)$  satisfying CLF, the map

$$\operatorname{Ex}_{+}(\operatorname{\overline{max}}^{*})\colon \operatorname{Ex}_{+}(\mathcal{C},W) \to \operatorname{Ex}_{+}\left(\operatorname{Ex}_{+}(\mathcal{C},W)^{\natural}\right)$$

is an equivalence of categories.

*Proof.* This map is  $E^1$ -homotopic to max<sup>\*</sup> (Corollary 9.4), which is an equivalence (Theorem 8.4).

With this, we may prove the main theorem of this section.

**Theorem 9.6.** Let  $(\mathcal{C}, W)$  be a marked quasicategory

- 1. If  $(\mathcal{C}, W)$  satisfies CLF then  $\max^* \colon \mathcal{C} \to \operatorname{Ex}_+(\mathcal{C}, W)$  is the localization of  $\mathcal{C}$  at W.
- 2. If  $(\mathcal{C}, W)$  satisfies CRF then  $\min^* \colon \mathcal{C} \to \operatorname{Ex}^{\operatorname{op}}_+(\mathcal{C}, W)$  is the localization of  $\mathcal{C}$  at W.

*Proof.* We prove (1), as (2) is formally dual.

Let  $\gamma \colon \mathcal{C} \to \mathcal{C}[W^{-1}]$  be the localization of  $\mathcal{C}$  at W. By Proposition 8.1, we have the following diagram of maps



which commutes up to natural equivalence. This implies the map

$$\max_{\mathcal{C}[W^{-1}]}^* \colon \mathcal{C}[W^{-1}] \to \operatorname{Ex}_+(\mathcal{C}[W^{-1}]^{\natural})$$

is  $E^1$ -homotopic to the composite  $Ex_+(\gamma) \circ F$  since

$$(\mathrm{Ex}_{+}(\gamma) \circ F \circ \gamma) \sim (\mathrm{Ex}_{+}(\gamma) \circ \mathrm{max}^{*}) = (\mathrm{max}^{*} \circ \gamma),$$

where the equality follows from naturality. Moreover, this diagram ascends to a diagram of marked simplicial sets

$$\begin{array}{c} (\mathcal{C}, W) \xrightarrow{\overline{\max^*}} \operatorname{Ex}_+(\mathcal{C}, W)^{\natural} \\ \hline \overline{\gamma} \\ \hline \varphi \\ \mathcal{C} [W^{-1}]^{\natural} \end{array}$$

whose underlying maps commute up to  $E^1$ -homotopy. By Proposition 8.2, the bottom right triangle in

$$\begin{array}{ccc} \mathcal{C}[W^{-1}] & & \xrightarrow{F} & \operatorname{Ex}_{+}(\mathcal{C}, W) \\ & & & & & \downarrow^{\operatorname{Ex}_{+}(\overline{\gamma})} & & \downarrow^{\operatorname{Ex}_{+}(\overline{\max^{*}})} \\ & & & & \operatorname{Ex}_{+}(\mathcal{C}[W^{-1}]^{\natural}) & \xrightarrow{F} & \operatorname{Ex}_{+}(\operatorname{Ex}_{+}(\mathcal{C}, W)^{\natural}) \end{array}$$

commutes up to  $E^1$ -homotopy. As the vertical maps are equivalences (by Theorem 8.4 and Corollary 9.5, respectively), the map F is an equivalence by 2-out-of-6.

**Corollary 9.7.** Let C be a quasicategory. Then ExC is a Kan complex if and only if  $C^{\#}$  admits the calculus of left fractions.

*Proof.* If  $\text{Ex} \mathcal{C}$  is a Kan complex, then by the first part of Theorem 7.5, we get that  $\mathcal{C}^{\#}$  admits the calculus of left fractions. The converse follows from Theorem 9.6 and Remark 7.3.

**Remark 9.8.** A special case of Corollary 9.7 for the nerve of a category was observed in [LTW79, Rmk. 5.8] (where it is attributed to Fritsch and Latch) and proven in [MO15, Thm. 2.1]. Note that a fully marked category  $C^{\#}$  satisfies CLF if and only if it satisfies proper CLF, and hence the result above indeed generalizes those of [LTW79, MO15].

**Example 9.9.** Recall that a localization  $\gamma: \mathcal{C} \to \mathcal{C}[W^{-1}]$  is *reflective* is  $\gamma$  admits a full and faithful right adjoint. These are occasionally referred to simply as *localizations*, e.g., [Lur09, Def. 5.2.7.2]. In view of Corollary 6.6, these are therefore necessarily of the form max<sup>\*</sup>:  $\mathcal{C} \to \text{Ex}_+(\mathcal{C}, W)$  (as described in Theorem 9.6) where W is the collection of maps in  $\mathcal{C}$  inverted by  $\gamma$ . Dually, a coreflective localization is of the form min<sup>\*</sup>:  $\mathcal{C} \to \text{Ex}_+^{\circ p}(\mathcal{C}, W)$ .

# Properties of the localization

### 10 Mapping spaces in the localization

In this section, we give explicit descriptions for the mapping spaces in  $\operatorname{Ex}_+(\mathcal{C}, W)$  and  $\operatorname{Ex}_+^{\operatorname{op}}(\mathcal{C}, W)$ . To do so, we give two constructions of the "marked slice" category. Mirroring the usual slice construction, both are defined by mapping out of "marked cones" over an *n*-simplex. We form one version of the marked cone using the product; the other is formed using the join. These allow us to define the *simplicial set* of fractions between two objects in a quasicategory (Definition 10.4), whic yields our first description of the mapping space (Theorem 10.5). Our second description of the mapping spaces in  $\operatorname{Ex}_+(\mathcal{C}, W)$  is as a colimit of mapping spaces in  $\mathcal{C}$  (Corollary 10.9), indexed by the marked slice category.

Define a cosimplicial object  $C^L: \Delta \to (\mathsf{sSet}_+)_*$  by the pushout



We regard  $C^L \Delta^n$  as pointed with distinguished basepoint  $[i_0]$ .

Geometrically, one thinks of  $C^L \Delta^n$  as a prism  $\Delta^n \times \Delta^1$  which has been quotiented to a cone at the 0-endpoint. The 1-simplices which extend from the cone point to the original *n*-simplex are exactly the marked 1-simplices of  $C^L \Delta^n$ .

This induces a functor  $C^L : \mathsf{sSet} \to (\mathsf{sSet}_+)_*$  via extension by colimits which sends a simplicial set X to the pushout

$$\begin{array}{c} X \longrightarrow \Delta^{0} \\ \downarrow_{i_{0}} \downarrow & \downarrow_{[i_{0}]} \\ X \times (\Delta^{1})^{\#} \longrightarrow C^{L} X \end{array}$$

with distinguished basepoint  $[i_0]$ . The additional assignment

$$C^{L}([1]^{\#}) := \operatorname{colim} \begin{pmatrix} (\Delta^{1})^{\#} & \longrightarrow \Delta^{0} \\ i_{0} \downarrow & & \\ (\Delta^{1})^{\#} \times (\Delta^{1})^{\#} & \end{pmatrix}$$

extends  $C^L$  to a functor  $\mathsf{sSet}_+ \to (\mathsf{sSet}_+)_*$ . Pre-composition gives a functor  $-\downarrow -: (\mathsf{sSet}_+)_* \to \mathsf{sSet}_+$  defined, on an object (X, W, x), by

$$(x \downarrow W)_n := (\mathsf{sSet}_+)_* (C^L \Delta^n, (X, W, x)).$$

We refer to  $x \downarrow W$  as the marked slice under x, as it is the simplicial subset of the "fat" slice under x (cf. [Lur09, §4.2.1]) spanned by 0-simplices  $(y, x \to y)$  such that  $x \to y$  is in W. Explicitly, an n-simplex of  $x \downarrow W$  is a map  $u: \Delta^n \times \Delta^1 \to X$  such that

- the restriction  $u|_{\Delta^n \times \{0\}}$  is constant at the 0-simplex x; and
- for  $i = 0, \ldots, n$ , the 1-simplex  $u|_{(i,0) \leq (i,1)} \colon \Delta^1 \to X$  is marked.

A 1-simplex  $u: (\Delta^1)^2 \to X$  is marked if the restriction  $u|_{(0,1)\leq (1,1)}: \Delta^1 \to X$  is a marked 1-simplex. Note the map

$$(\Delta^n)^{\flat} \times (\Delta^1)^{\#} \to C^L \Delta^n$$

induces a natural inclusion

$$x \downarrow W \to (X, W)^{(\Delta^1)^{\#}}$$

by pre-composition. We write  $\Pi_x$  for the composite

$$x \downarrow W \hookrightarrow (X, W)^{(\Delta^1)^{\#}} \xrightarrow{i_1^*} (X, W)$$

which we refer to as the *projection map*.

We similarly define a cosimplicial object  $C^R \Delta^n$  by the pushout,

$$\begin{array}{ccc} \left(\Delta^{n}\right)^{\flat} & \longrightarrow & \Delta^{0} \\ & & & & \\ i_{1} \downarrow & & & & \\ \left(\Delta^{n}\right)^{\flat} \times \left(\Delta^{1}\right)^{\#} & \longrightarrow & C^{R} \Delta^{n} \end{array}$$

which one thinks of as a prism  $(\Delta^n)^{\flat} \times (\Delta^1)^{\#}$  which has been quotiented to a cone at the 1-endpoint. This defines a functor  $C^R: \mathsf{sSet}_+ \to (\mathsf{sSet}_+)_*$  via extension by colimits (with an analogous definition on  $\Delta^{1^{\#}}$ ). Its right adjoint  $-\downarrow -: (\mathsf{sSet}_+)_* \to \mathsf{sSet}_+$  is defined by

$$(W \downarrow x)_n := (\mathsf{sSet}_+)_* (C^R \Delta^n, (X, W, x)).$$

We refer to  $W \downarrow x$  as the marked slice over x. This functor comes with a natural projection map

$$W \downarrow x \hookrightarrow (X, W)^{(\Delta^1)^{\#}} \xrightarrow{i_0^*} (X, W)$$

Note that for  $(X, W, x) \in (\mathsf{sSet}_+)_*$ , there is an isomorphism

$$(x \downarrow W)^{\mathrm{op}} \cong W^{\mathrm{op}} \downarrow x$$

natural in (X, W, x).

The following proposition follows by construction.

**Proposition 10.1.** For a marked simplicial set (X, W) and  $x \in X_0$ , the squares

$$\begin{array}{cccc} x \downarrow W & \longleftrightarrow & (X,W)^{(\Delta^{1})^{\#}} & & W \downarrow x & \longleftrightarrow & (X,W)^{(\Delta^{1})^{\#}} \\ \downarrow & & \downarrow^{i_{0}} & & \downarrow & \downarrow^{i_{1}} \\ \Delta^{0} & \underbrace{ & }_{x} & (X,W) & & \Delta^{0} & \underbrace{ & }_{x} & (X,W) \end{array}$$

are pullbacks.

**Proposition 10.2.** Let  $x \in X$  be a 0-simplex in a marked simplicial set (X, W) such that W is strongly closed under composition.

- 1. If X is a quasicategory then  $x \downarrow W$  and  $W \downarrow x$  are quasicategories.
- 2. If (X, W) satisfies CLF then  $x \downarrow W$  satisfies CLF.
- 3. If (X, W) satisfies CRF then  $W \downarrow x$  satisfies CRF.

*Proof.* For (1), applying [Lur09, Cor. 2.4.7.12] to the map  $\Delta^0 \xrightarrow{x} X$  gives that  $x \downarrow W \to \Delta^0$  is a fibration. Dually,  $W \downarrow x \to \Delta^0$  is a fibration.

For (2), the result for  $x \downarrow W$  follows from Lemma 6.11, as  $x \downarrow W \rightarrow \Delta^0$  is a pullback of a map which has the right lifting property against  $\{L_k^n \hookrightarrow {}^*L_k^n \mid n \ge 2, 0 < k \le n\}$  by Proposition 10.1.

The proof of (3) is dual.

We now describe an alternate construction of the marked cone and slice using the join rather than the product. Let  $C_*^L: \mathsf{sSet} \to (\mathsf{sSet}_+)_*$  denote the functor which sends a simplicial set X to the simplicial set  $\Delta^0 * X$  (whose distinguished point is the cone point), where every 1-simplex connected to the cone point is marked. This functor has a right adjoint  $(\mathsf{sSet}_+)_* \to \mathsf{sSet}$  which sends (X, W, x) to the simplicial set

$$(x\downarrow^* W)_n := (\mathsf{sSet}_+)_* (C^L_* \Delta^n, (X, W, x))_*$$

Explicitly,  $x \downarrow^* W$  is the simplicial subset of the ordinary slice category under x spanned by 0-simplices  $(y, x \to y)$  such that  $x \to y$  is in W. The map  $C^L \Delta^n \to C^L_* \Delta^n$  induced by the square

$$\begin{array}{ccc} (\Delta^n)^{\flat} & \longrightarrow & \Delta^0 \\ & & & \downarrow^{\perp} \\ (\Delta^n)^{\flat} \times (\Delta^1)^{\#} & \xrightarrow{(i,0)\mapsto i} & C^L_* \Delta^n \end{array}$$

commutes with faces and degeneracies. Thus, pre-composition induces a map

 $x\downarrow^* W \to x\downarrow W$ 

which is natural in (X, W, x). Analogously, one defines a functor  $C^R_*: \mathsf{sSet} \to (\mathsf{sSet}_+)_*$  and a map  $C^R \Delta^n \to C^R_* \Delta^n$ , which induces a map

$$W \downarrow^* x \to W \downarrow x.$$

**Proposition 10.3.** Let x be an object in a marked quasicategory  $(\mathcal{C}, W)$ . The maps

$$x \downarrow^* W \to x \downarrow W, \quad W \downarrow^* x \to W \downarrow x$$

are categorical equivalences.

*Proof.* These maps are restrictions onto full subcategories of the canonical equivalences from the ordinary slice to the "fat" slice ([Lur09, Prop. 4.2.1.5]).

For the remainder of this section, we work with the marked "fat" slice  $x \downarrow W$  rather than the marked ordinary slice  $x \downarrow^* W$ .

With this, we introduce the following auxiliary construction.

**Definition 10.4.** Let (X, W) be a marked simplicial set and  $x, y \in X$  be 0-simplices.

1. The simplicial set  $LF_{(X,W)}(x,y)$  of left fractions from x to y is the pullback

in  $sSet_+$ .

2. The simplicial set  $LF_{(X,W)}(x,y)$  of right fractions from x to y is the pullback

$$RF_{(X,W)}(x,y) \longrightarrow (X \downarrow y,W)$$

$$\downarrow \qquad \qquad \downarrow \Pi_{y}$$

$$W \downarrow x \xrightarrow{\Pi_{x}} (X,W)$$

in  $sSet_+$ .

Our interest in the simplicial set of left (and right) fractions is in modelling the mapping space of the localization.

**Theorem 10.5.** Let (X, W) be a marked simplicial set and x, y be 0-simplices. There are isomorphisms

$$\operatorname{map}_{L}(\operatorname{Ex}_{+}(X,W),x,y) \cong \operatorname{Ex}_{+}(\operatorname{LF}_{(X,W)}(x,y)),$$
$$\operatorname{map}_{R}(\operatorname{Ex}_{+}^{\operatorname{op}}(X,W),x,y) \cong \operatorname{Ex}_{+}^{\operatorname{op}}(\operatorname{RF}_{(X,W)}(x,y))$$

natural in (X, W) and x, y.

The proof of Theorem 10.5 will require some additional definitions and lemmas. We present these only for  $Sd_+$  and  $Ex_+$ , omitting their dual statements, as their primary purpose is in proving Theorem 10.5.

Observe that the left mapping space of a quasicategory arises as pre-composition by a cosimplicial object. For  $n \ge 0$ , let  $\Delta^{n+1}/\partial_0$  denote the simplicial set obtained by the pushout

$$\begin{array}{ccc} \Delta^n & \longrightarrow & \Delta^0 \\ \partial_0 & & & & \downarrow [\partial_0] \\ \Delta^{n+1} & \longrightarrow & \Delta^{n+1} / \partial_0 \end{array}$$

Let q denote the quotient map  $\Delta^{n+1} \to \Delta^{n+1}/\partial_0$ . We regard  $\Delta^{n+1}/\partial_0$  as bi-pointed at 0 and  $[\partial_0]$ . The left mapping space map<sub>L</sub>: sSet<sub>\*\*</sub>  $\to$  sSet is exactly the induced functor given by pre-composition, i.e.

$$\operatorname{map}_{L}(X, x, y)_{n} := \mathsf{sSet}_{**}((\Delta^{n+1}/\partial_{0}, 0, [\partial_{0}]), (X, x, y)).$$

Define a map  $f: \Delta^0 * \operatorname{Sd}_+ \Delta^n \to \operatorname{Sd}_+ \Delta^{n+1}$  by

$$f(A) := \begin{cases} \{0\} & A = \bot \\ \partial_0(A) \cup \{0\} & \text{otherwise.} \end{cases}$$

This formula defines a map on the underlying simplicial sets as the domain and codomain are nerves of posets. One verifies this map sends marked 1-simplices to marked 1-simplices. We also define a map  $g: \operatorname{Sd}_+ \Delta^n \times (\Delta^1)^{\#} \to \operatorname{Sd}_+ \Delta^{n+1}$  by

$$g(A,t) := \begin{cases} \partial_0(A) & \text{if } t = 0\\ \partial_0(A) \cup \{0\} & \text{if } t = 1. \end{cases}$$

Lemma 10.6. The square

$$\begin{array}{ccc} \operatorname{Sd}_{+}\Delta^{n} \xrightarrow{\operatorname{id} \times \{1\}} & \operatorname{Sd}_{+}\Delta^{n} \times (\Delta^{1})^{\#} \\ & & i_{1} \\ & & \downarrow^{g} \\ \Delta^{0} * \operatorname{Sd}_{+}\Delta^{n} \xrightarrow{f} & \operatorname{Sd}_{+}\Delta^{n+1} \end{array}$$

is a pushout.

*Proof.* The square commutes by definition. Given a non-degenerate simplex  $(A_0 \subsetneq \cdots \subsetneq A_n)$  of  $\mathrm{Sd}_+ \Delta^{n+1}$ , we show this simplex is in the image of f or g by case analysis on  $A_0$ . Moreover, we show the intersection of the images is the image of the composite map in the square. This suffices as each map in the square is injective.

If  $A_0 = \{0\}$  then this simplex is in the image of f as

$$(A_0 \subsetneq A_1 \subsetneq \cdots \subsetneq A_n) = f(\bot \subsetneq \sigma_0(A_1 - \{0\}) \subsetneq \cdots \subsetneq \sigma_0(A_n - \{0\})).$$

Note this simplex is not contained in the image of g as  $\{0\}$  is not contained in the image of g.

If  $A_0$  does not contain 0 then let i = 0, ..., n be maximal such that  $0 \notin A_i$ . Applying g to the n-simplex

$$(\sigma_0(A_0), 0) \subsetneq \cdots \subsetneq (\sigma_0(A_i), 0) \subsetneq (\sigma_0(A_{i+1}), 1) \subsetneq \cdots \subsetneq (\sigma_0(A_n), 1)$$

gives exactly  $(A_0 \subsetneq \cdots \subsetneq A_n)$ . Note this simplex is not contained in the image of f as the 0-simplices in the image of f are subsets that contain 0.

The simplices that remain in our case analysis are exactly those in the set of non-degenerate simplices defined by

$$\{(A_0 \subsetneq \cdots \subsetneq A_n) \mid \{0\} \subsetneq A_0\}.$$

Note the previous arguments show that the intersection of the images of f and g must be contained in this set. However, this set is contained in the image of the composite map since, for an *n*-simplex  $(A_0 \subsetneq \cdots \subsetneq A_n)$  in this set, we have equalities

$$(f \circ i_1)(\sigma_0(A_0 - \{0\}) \subsetneq \cdots \subsetneq \sigma_0(A_n - \{0\})) = \partial_0 \sigma_0(A_0 - \{0\}) \cup \{0\} \subsetneq \cdots \subsetneq \partial_0 \sigma_0(A_n - \{0\}) \cup \{0\})$$
$$= (A_0 \subsetneq \cdots \subsetneq A_n),$$

and

$$(g \circ (\mathrm{id} \times \{1\}))(\sigma_0(A_0 - \{0\}) \subsetneq \cdots \subsetneq \sigma_0(A_n - \{0\}))$$
  
=  $\partial_0 \sigma_0(A_0 - \{0\}) \cup \{0\} \subsetneq \cdots \subsetneq \partial_0 \sigma_0(A_n - \{0\}) \cup \{0\})$   
=  $(A_0 \subsetneq \cdots \subsetneq A_n).$ 

This gives both that the intersection of the images is the image of the composite and that every nondegenerate *n*-simplex of  $\mathrm{Sd}_+ \Delta^{n+1}$  is contained in either the image of f or the image of g.

The square

$$\begin{array}{c} \operatorname{Sd}_{+}\Delta^{n} & \longrightarrow & \Delta^{0} \\ & & & \downarrow \\ & & & \downarrow [\partial_{0}] \\ \operatorname{Sd}_{+}\Delta^{n} \times (\Delta^{1})^{\#} \xrightarrow{g} \operatorname{Sd}_{+}\Delta^{n+1} \xrightarrow{q} \operatorname{Sd}_{+}(\Delta^{n+1}/\partial_{0}) \end{array}$$

induces a map  $[g]: C^L(\mathrm{Sd}_+ \Delta^n) \to \mathrm{Sd}_+(\Delta^{n+1}/\partial_0)$  by universal property of the pushout. This map gives a commutative square

in the arrow category  $\mathbf{sSet}_{+}^{[1]}$ . One verifies these maps are natural in n, so that for a marked simplicial set (X, W) and  $x, y \in X$ , this induces a commutative square

by pre-composition, where one verifies that  $\operatorname{Ex}_{+} \Pi_{x}$  is identical to the pre-composition map with  $(\emptyset \hookrightarrow \operatorname{Sd}_{+} \Delta^{n}) \hookrightarrow (\{\bot\} \to \Delta^{0} * \operatorname{Sd}_{+} \Delta^{n})$  and  $\operatorname{Ex}_{+} \Pi_{y}$  is identical to the pre-composition map with  $(\emptyset \hookrightarrow \operatorname{Sd}_{+} \Delta^{n}) \hookrightarrow (\{[i_{0}]\} \to C^{L}(\operatorname{Sd}_{+} \Delta^{n})).$ 

Lemma 10.7. 1. The square

is a pushout in the arrow category  $sSet_{+}^{[1]}$ .

2. For a marked simplicial set (X, W) and  $x, y \in X$ , the square

is a pullback in sSet.

1

*Proof.* For (1), it is clear that the domains form a pushout square. For the codomains, we consider the diagram

$$\begin{array}{cccc} \operatorname{Sd}_{+}\Delta^{n} & \longrightarrow & \Delta^{0} \\ & & & & & \downarrow^{[i_{0}]} \\ \operatorname{Sd}_{+}\Delta^{n} & \stackrel{i_{1}}{\longrightarrow} & \operatorname{Sd}_{+}\Delta^{n} \times (\Delta^{1})^{\#} & \longrightarrow & C^{L}(\operatorname{Sd}_{+}\Delta^{n}) \\ & & & & \downarrow^{[g]} \\ \Delta^{0} * \operatorname{Sd}_{+}\Delta^{n} & \stackrel{f}{\longrightarrow} & \operatorname{Sd}_{+}\Delta^{n+1} & \longrightarrow & \operatorname{Sd}_{+}(\Delta^{n+1}/\partial_{0}) \end{array}$$

The bottom left square is a pushout by Lemma 10.6. The top right square is a pushout by definition of  $C^L: \mathbf{sSet} \to (\mathbf{sSet}_+)_*$ . The right composite square is a pushout as  $\mathrm{Sd}_+$  preserves colimits, hence the bottom right square is a pushout. This gives that the bottom composite square is a pushout.

For statement (2), fix  $n \ge 0$  and consider the product of co-spans

where the right horizontal arrows are given by applying the domain projection  $sSet_{+}^{[1]} \rightarrow sSet_{+}$ . The pullback of each horizontal co-span gives the co-span

As limits commute with limits, this pullback is computed by the pullback of the pullbacks of the vertical co-spans. The co-span

$$s\mathsf{Set}_{+}^{[1]}((\{0, [\partial_{0}]\} \to \mathrm{Sd}_{+}(\Delta^{n+1}/\partial_{0})), (\{x, y\} \to (X, W)))$$

$$\downarrow$$

$$\overset{\perp \mapsto x}{\underset{[i_{0}] \mapsto y}{\longrightarrow}} s\mathsf{Set}_{+}(\{\bot, [i_{0}]\} \to \{x, y\})$$

is the pullback of the vertical co-spans as the contravariant Yoneda embedding takes colimits to limits. In particular, our computation of the top right corner follows from statement (1). This pullback is exactly the *n*-simplices of the left mapping space of  $\text{Ex}_+(X, W)$  from x to y.

With this, we may prove Theorem 10.5, giving our first description of the mapping spaces in the localization.

Proof of Theorem 10.5. The first isomorphism follows from Proposition 10.1 and item (2) of Lemma 10.7, as  $\text{Ex}_+: \text{sSet}_+ \to \text{sSet}$  preserves pullbacks. The second is dual.

For our second description of the mapping spaces in the localization, which will mirror the classical description of mapping spaces (cf. Theorem 1.8), we make use of the following colimit computation.

**Theorem 10.8.** For a Grothendieck universe  $\mathcal{U}$ , let x, y be objects in a locally  $\mathcal{U}$ -small quasicategory  $\mathcal{C}$ .

1. Suppose  $U(y \downarrow W)$  is finally  $\mathcal{U}$ -small. Then, the Kan fibrant replacement of the (underlying) simplicial set of left fractions  $LF_{(\mathcal{C},W)}(x,y)$  from x to y models the  $\infty$ -colimit of the functor

$$U(y \downarrow W) \xrightarrow{U\Pi} \mathcal{C} \xrightarrow{\mathcal{C}(x,-)} \mathcal{S}.$$

That is, there is an equivalence of Kan complexes

$$\operatorname{colim} \left( \mathcal{C}(x, -) \circ U \Pi \circ i \right) \simeq U(\operatorname{LF}_{(\mathcal{C}, W)}(x, y))[\operatorname{LF}_{(\mathcal{C}, W)}(x, y)_1^{-1}].$$

2. Suppose  $U(W \downarrow x)$  is finally  $\mathcal{U}$ -small. Then, the Kan fibrant replacement of the (underlying) simplicial set of right fractions  $\operatorname{RF}_{(\mathcal{C},W)}(x,y)$  from x to y models the  $\infty$ -colimit of the functor

$$U(W \downarrow x)^{\mathrm{op}} \xrightarrow{(U\Pi)^{\mathrm{op}}} \mathcal{C}^{\mathrm{op}} \xrightarrow{\mathcal{C}(-,y)} \mathcal{S}.$$

That is, there is an equivalence of Kan complexes

$$\operatorname{colim} \left( \mathcal{C}(-, y) \circ (U\Pi)^{\operatorname{op}} \circ i \right) \simeq U(\operatorname{RF}_{(\mathcal{C}, W)}(x, y))[\operatorname{RF}_{(\mathcal{C}, W)}(x, y)_1^{-1}].$$

*Proof.* We prove (1), as (2) is formally dual.

By [Lur09, Cor. 3.3.4.6], the colimit of any functor  $F: \mathcal{C} \to \mathcal{S}$  is computed by the Kan fibrant replacement of the domain of the left fibration classified by F. Thus, it suffices to show the functor  $\mathcal{C}(x, -) \circ \Pi$ classifies the left fibration  $LF_{(\mathcal{C},W)}(x, y) \to y \downarrow W$ .

Applying [Cis19, Cor. 5.3.21], the right square in

is a homotopy pullback (in the Joyal model structure) as C(x, -) classifies the left fibration  $x \downarrow C \rightarrow C$ . The left square is a homotopy pullback as a pullback along a left fibration. Thus, the composite square is a homotopy pullback by [Cis19, Cor. 5.3.21], which suffices.

This gives a description of mapping spaces in the localization as a filtered colimit of mapping spaces in C indexed by the marked slice category.

**Corollary 10.9.** For a Grothendieck universe  $\mathcal{U}$ , let x, y be objects in a marked quasicategory  $(\mathcal{C}, W)$  which is locally  $\mathcal{U}$ -small.

1. If  $U(y \downarrow W)$  is finally  $\mathcal{U}$ -small and  $(\mathcal{C}, W)$  satisfies CLF then there is an equivalence

$$\operatorname{map}_{L}(\operatorname{Ex}_{+}(\mathcal{C}, W), x, y) \simeq \operatorname{colim}(\mathcal{C}(x, -) \circ U\Pi)$$

which is natural in x, y and  $(\mathcal{C}, W)$ .

2. If  $U(W \downarrow x)$  is finally U-small and  $(\mathcal{C}, W)$  satisfies CRF then there is an equivalence

 $\operatorname{map}_{R}(\operatorname{Ex}^{\operatorname{op}}_{+}(\mathcal{C}, W), x, y) \simeq \operatorname{colim}(\mathcal{C}(-, y) \circ U\Pi^{\operatorname{op}})$ 

which is natural in x, y and  $(\mathcal{C}, W)$ .

*Proof.* We prove (1), as (2) is analogous.

By Theorem 10.5, we have an isomorphism

$$\operatorname{map}_{L}(\operatorname{Ex}_{+}(\mathcal{C}, W), x, y) \cong \operatorname{Ex}_{+}(\operatorname{LF}_{(\mathcal{C}, W)}(x, y)),$$

natural in  $(\mathcal{C}, W)$ , x, and y. As the left simplicial set is Kan, the right one is. The marked 1-simplices in  $LF_{(\mathcal{C},W)}(x,y)$  are closed under composition as W is strongly closed under composition. Thus, Theorem 7.5 gives that  $LF_{(\mathcal{C},W)}(x,y)$  satisfies CLF. By Theorem 9.6, the map

$$\max^* : ULF_{(\mathcal{C},W)}(x,y) \to Ex_+(LF_{(\mathcal{C},W)}(x,y))$$

is the localization of  $LF_{(\mathcal{C},W)}(x,y)$ . As the codomain is Kan, this map is also the localization of  $ULF_{(\mathcal{C},W)}(x,y)$  at all maps, i.e. the Kan fibrant replacement of  $ULF_{(\mathcal{C},W)}(x,y)$ . The result then follows by Theorem 10.8.

## 11 (Co)limits in the localization

This section fully establishes the quasicategorical analogue of Theorem 1.8, giving conditions for the existence of (co)limits in the localization (Theorem 11.7). Following the argument given in [GZ67, Ch. 1], we prove this by showing the categories  $x \downarrow W$  and  $W \downarrow x$  are filtered if W is closed under 2-out-of-3 (Corollary 11.5). In fact, we show more generally that, in the presence of either CLF (or CRF) on a maximally-marked quasicategory  $C^{\#}$ , being (co)filtered is *equivalent* to being weakly contractible (Corollary 11.4). From this, one concludes that the mapping spaces in  $\text{Ex}_+(\mathcal{C}, W)$  are *filtered* colimits of the mapping spaces in  $\mathcal{C}$ .

Define a cosimplicial object  $\widehat{sd}: \Delta \to \mathsf{sSet}_+$  which sends [n] to the preorder on the set

$$\{A \subseteq [n] \mid A \neq \emptyset\}$$

where

$$A_0 \le A_1 := \max A_0 \le \max A_1.$$

We view this preorder as a simplicial set marked at  $E^1$ -equivalences (i.e. a 1-simplex  $A_0 \leq A_1$  is marked if  $\max A_0 = \max A_1$ ). Let  $\widehat{Sd}: \mathsf{sSet} \to \mathsf{sSet}_+$  denote its extension by colimits. Note the commutative triangle of cosimplicial objects



induces a commutative triangle



where each map is natural in X.

**Proposition 11.1.** For a simplicial set X,

- 1. the map max<sub>1</sub>:  $\widehat{Sd} X \to X^{\flat}$  is an  $E^1$ -homotopy equivalence;
- 2. each map in the triangle



is a weak equivalence.

- Proof. 1. The map  $f: [n] \to \widehat{sd}[n]$  defined by  $i \mapsto \{i\}$  commutes with faces and degeneracies. Moreover, for  $A \in \widehat{sd}[n]$ , we have that  $A \leq \{\max A\}$  is an equivalence. This induces a homotopy  $\widehat{sd}[n] \times E[1] \to \widehat{sd}[n]$  from the identity to  $f \circ \max$  which also commutes with faces and degeneracies. Thus, this map induces a section  $X^{\flat} \to \widehat{Sd} X$  to the map  $\max_{1}: \widehat{Sd} X \to X^{\flat}$ , along with a homotopy  $\widehat{Sd} X \times E^{1} \to \widehat{Sd} X$  from the identity to  $f_{1} \circ \max_{1}$ .
  - 2. Lemma 8.3 shows the map max:  $sd_+[n] \rightarrow [n]$  is a homotopy equivalence. By [Cis19, Prop. 3.1.14], the left map in the triangle is a weak equivalence. The right map is a weak equivalence by (1). Thus, the top map is by 2-out-of-3.

Corollary 11.2. For a simplicial set X, each map in the triangle



is a weak equivalence.

*Proof.* Follows from Proposition 3.9 using an argument analogous to the proof of Proposition 11.1.

Recall (cf. [Lur09, Def. 5.3.1.7]) that a simplicial set X is *filtered* if every map  $K \to X$  from a finite simplicial set K (i.e. K has finitely many non-degenerate simplices) admits a lift to a map  $K * \Delta^0 \to X$ . Dually, X is *cofiltered* if every such map  $K \to X$  admits a lift to a map  $\Delta^0 * K \to X$ .

Lemma 11.3. If a quasicategory has the right lifting property against the set of maps

$$\{\operatorname{Sd} \partial \Delta^n \hookrightarrow \operatorname{Sd} \Delta^n \mid n \ge 0\}$$

then it is filtered.

*Proof.* Let C be a quasi-category with the right lifting property against  $\{\operatorname{Sd} \partial \Delta^n \to \operatorname{Sd} \Delta^n \mid n \geq 0\}$ . By [Lur09, Lem. 5.3.1.12], it suffices to check that every map  $\partial \Delta^n \to C$  admits a lift to a map  $\partial \Delta^n * \Delta^0 \to C$ . That is, it suffices to show C has the right lifting property against the set of maps

$$\{\partial \Delta^n \hookrightarrow (\partial \Delta^n) * \Delta^0 \mid n \ge 0\}.$$

The square

$$\begin{array}{ccc} \operatorname{Sd}_{+} \partial \Delta^{n} & \longrightarrow & \widehat{\operatorname{Sd}} \partial \Delta^{n} \\ & & & & \downarrow \\ & & & & \downarrow \\ (\operatorname{Sd}_{+} \partial \Delta^{n}) * \Delta^{0} & \longmapsto & (\widehat{\operatorname{Sd}} \partial \Delta^{n}) * \Delta^{0} \end{array}$$

induces a map from the pushout, which we denote by  $f: P \to (\widehat{\mathrm{Sd}} \partial \Delta^n) * \Delta^0$ . The horizontal maps are weak equivalences by Proposition 11.1 and Corollary 11.2, respectively. Thus, f is a weak equivalence by 2-out-of-3. One verifies f is a monomorphism, thus it is an acyclic cofibration.

As C has the right lifting property against the set  $\{ \operatorname{Sd} \partial \Delta^n \hookrightarrow \operatorname{Sd} \Delta^n \mid n \ge 0 \}$ , its maximal marking  $C^{\#}$  has the right lifting property against the set of maps

$$\{\mathrm{Sd}_+ \partial \Delta^n \hookrightarrow (\mathrm{Sd}_+ \partial \Delta^n) * \Delta^0 \mid n \ge 0\}$$

by adjointness (as  $\operatorname{Sd} \Delta^n \cong U(\operatorname{Sd}_+ \partial \Delta^n * \Delta^0)$ ). Thus, it has the right lifting property against the map  $\widehat{\operatorname{Sd}} \partial \Delta^n \hookrightarrow P$ .

Given a map  $u: \partial \Delta^n \to \mathcal{C}$ , this gives that the composite

$$\widehat{\mathrm{Sd}}\,\partial\Delta^n\xrightarrow{\mathrm{max}_!}(\partial\Delta^n)^\flat\xrightarrow{u^\flat}\mathcal{C}^\flat\hookrightarrow\mathcal{C}^\#$$

lifts to a map

$$v\colon P\to \mathcal{C}^{\#}$$

As this composite sends marked 1-simplices to degeneracies and  $\widehat{\mathrm{Sd}} \partial \Delta^n \hookrightarrow P$  is surjective on 1-simplices, we may view v as a map  $P \to \mathcal{C}^{\natural}$ .

The map  $P \to (\widehat{\mathrm{Sd}} \partial \Delta^n) * \Delta^0$  is an acyclic cofibration and  $\mathcal{C}^{\sharp}$  is fibrant, hence v lifts to a map

$$\overline{v} \colon (\widehat{\mathrm{Sd}}\,\partial\Delta^n) * \Delta^0 \to \mathcal{C}^{\natural}.$$

Applying naturality to the section  $i: (\partial \Delta^n)^{\flat} \to \widehat{\mathrm{Sd}} \partial \Delta^n$  defined in Proposition 11.1 gives a commutative square:

$$\begin{array}{ccc} \left(\partial\Delta^{n}\right)^{\flat} & & \stackrel{i}{\longrightarrow} \widehat{\mathrm{Sd}} \,\partial\Delta^{n} \\ & & & \downarrow \\ \left(\partial\Delta^{n}\right)^{\flat} * \Delta^{0} & \stackrel{i*\Delta^{0}}{\longrightarrow} \left(\widehat{\mathrm{Sd}} \,\partial\Delta^{n}\right) * \Delta^{0} \end{array}$$

The (underlying) map  $\overline{v} \circ (i * \Delta^0)$  is the desired lift.



Corollary 11.4. Let C be a quasicategory.

1. If  $C^{\#}$  satisfies CLF then C is filtered if and only if the Kan fibrant replacement of C is contractible.

2. If  $C^{\#}$  satisfies CRF then C is cofiltered if and only if the Kan fibrant replacement of C is contractible.

*Proof.* We prove (1), as (2) is formally dual.

For the forward direction, apply [Lur09, Lem. 5.3.1.18]. For the converse, Theorem 9.6 shows that  $\operatorname{Ex}_{+} \mathcal{C}^{\#} \cong \operatorname{Ex} \mathcal{C}$  is a Kan fibrant replacement of  $\mathcal{C}$ . By adjointness,  $\operatorname{Ex} \mathcal{C}$  is contractible if and only if  $\mathcal{C}$  has the right lifting property with respect to the set of maps  $\{\operatorname{Sd} \partial \Delta^n \hookrightarrow \operatorname{Sd} \Delta^n \mid n \geq 0\}$ . This then follows from Lemma 11.3.

We can now show that if W is closed under 2-out-of-3 then the marked slice category is filtered.

**Corollary 11.5.** Let x be an object in a marked quasicategory  $(\mathcal{C}, W)$  such that W is closed under 2-outof-3.

- 1. The underlying simplicial set  $U(x \downarrow W)$  of the marked slice under x is filtered.
- 2. The underlying simplicial set  $U(W \downarrow x)$  of the marked slice over x is cofiltered.

*Proof.* We prove (1), as (2) is formally dual.

As W is closed under 2-out-of-3, every 1-simplex of  $x \downarrow W$  is marked. That is,  $x \downarrow W = U(x \downarrow W)^{\#}$ . As  $x \downarrow W$  satisfies CLF by Proposition 10.2 and Corollary 11.4, it suffices to show the Kan fibrant replacement of  $U(x \downarrow W)$  is contractible. This follows as  $x \downarrow W$  is a quasicategory with an initial object  $(x, x \xrightarrow{x\sigma_1} x)$ .

From this, it follows that the localization functor  $\mathcal{C} \to \text{Ex}_+(\mathcal{C}, W)$  preserves colimits. The proofs presented here follow the arguments given in [GZ67, Ch. 1].

**Lemma 11.6.** Let  $(\mathcal{C}, W)$  be a marked quasicategory.

- 1. If  $(\mathcal{C}, W)$  satisfies CLF then every co-equalizer diagram  $\Delta^1 \cup_{\partial \Delta^1} \Delta^1 \to \operatorname{Ex}_+(\mathcal{C}, W)$  is  $E^1$ -homotopic to one which factors through  $\max^* \colon \mathcal{C} \to \operatorname{Ex}_+(\mathcal{C}, W)$ .
- 2. If  $(\mathcal{C}, W)$  satisfies CRF then every equalizer diagram  $\Delta^1 \cup_{\partial \Delta^1} \Delta^1 \to \operatorname{Ex}^{\operatorname{op}}_+(\mathcal{C}, W)$  is  $E^1$ -homotopic to one which factors through  $\min^* \colon \mathcal{C} \to \operatorname{Ex}^{\operatorname{op}}_+(\mathcal{C}, W)$ .

*Proof.* We prove (1), as (2) is formally dual.

Fix a co-equaliser diagram F, which we view as a map  $\operatorname{Sd}_+\Delta^1 \cup_{\{0,1\}} \operatorname{Sd}_+\Delta^1 \to (\mathcal{C}, W)$  as in the diagram:



As  $(\mathcal{C}, W)$  satisfies CLF, the span  $(w_1, w_2)$  may be completed to a commutative square as in the diagram:

$$\begin{array}{c|c} y & \xrightarrow{w_1} & y_1 \\ \hline & & & \\ w_2 \\ \downarrow & & & \\ y_2 & \xrightarrow{v_0} & \downarrow \\ & & & \\ y_2 & \xrightarrow{v_2} & y_3 \end{array}$$

Let  $v_1f_1: x \to y_3$  be a composite of  $f_1$  and  $v_1$  and let  $v_2f_2: x \to y_3$  be a composite of  $f_2$  and  $v_2$ . The diagram



depicts a marked homotopy  $(\mathrm{Sd}_+ \Delta^1 \cup_{\{0,1\}} \mathrm{Sd}_+ \Delta^1) \times (\Delta^1)^{\#} \to (\mathcal{C}, W)$  from F to  $[v_1 f_1, v_2 f_2] \circ \max_{\{0,1\}}$ (note the outer vertical 1-simplices in each component of the diagram are identified). This induces an  $E^1$ -homotopy in  $\mathrm{Ex}_+(\mathcal{C}, W)$  from F to  $\max^* \circ [v_1 f_1, v_2 f_2]$ .

With this, we may state a complete quasicategorical analogue of Theorem 1.8, giving a criterion for the existence of colimits in the localization.

**Theorem 11.7.** Let  $(\mathcal{C}, W)$  be a marked quasicategory which is locally  $\mathcal{U}$ -small where W is closed under 2-out-of-3.

- 1. Suppose  $(\mathcal{C}, W)$  satisfies CLF and that, for any object  $x \in \mathcal{C}$ , the marked slice  $U(x \downarrow W)$  over x is finally  $\mathcal{U}$ -small. Then,
  - a) for any finite colimit cone  $\lambda \colon K * \Delta^0 \to \mathcal{C}$ , the composite

$$\max^* \circ \lambda \colon K * \Delta^0 \to \operatorname{Ex}_+(\mathcal{C}, W)$$

is a colimit cone; and

- b) if C admits all finite colimits then  $Ex_+(C, W)$  admits all finite colimits.
- 2. Suppose  $(\mathcal{C}, W)$  satisfies CRF and that, for any object  $x \in \mathcal{C}$ , the marked slice  $U(W \downarrow x)$  under x is finally  $\mathcal{U}$ -small. Then,
  - a) for any finite limit cone  $\lambda \colon K * \Delta^0 \to \mathcal{C}$ , the composite

$$\min^* \circ \lambda \colon K \ast \Delta^0 \to \operatorname{Ex}^{\operatorname{op}}_+(\mathcal{C}, W)$$

is a limit cone; and

b) if C admits all finite limits then  $\operatorname{Ex}_{+}^{\operatorname{op}}(\mathcal{C}, W)$  admits all finite limits.

*Proof.* We prove (1), as (2) is formally dual.

For (a), let F denote the restriction  $\lambda|_K \colon K \to \mathcal{C}$  of  $\lambda$  to K and let  $\operatorname{colim} F$  denote the value of  $\lambda$  at the cone point. It suffices to show  $\operatorname{colim} F$ , as an object in  $\operatorname{Ex}_+(\mathcal{C}, W)$ , represents the functor of cones under  $\max^* \circ F$ . That is, for every object  $x \in \operatorname{Ex}_+(\mathcal{C}, W)$ , we show that the canonical map

$$\varphi \colon \operatorname{Ex}_{+}(\mathcal{C}, W)(\operatorname{colim} F, x) \to \lim \left( K^{\operatorname{op}} \xrightarrow{\max^{*} \circ F} \operatorname{Ex}_{+}(\mathcal{C}, W)^{\operatorname{op}} \xrightarrow{\operatorname{Ex}_{+}(\mathcal{C}, W)(-, x)} S \right)$$

is an equivalence of Kan complexes.

Let  $G: K^{\mathrm{op}} \times U(x \downarrow W) \to \mathcal{S}$  denote the composite map

$$K^{\mathrm{op}} \times U(x \downarrow W) \xrightarrow{F^{\mathrm{op}} \times \Pi_x} \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \xrightarrow{\mathcal{C}(-,-)} \mathcal{S}.$$

We show there is a natural  $E^1$ -homotopy from  $\varphi$  to the canonical map

$$\operatorname{colim}_{U(x \downarrow W)}(\lim_{K^{\operatorname{op}}} G) \to \lim_{K^{\operatorname{op}}}(\operatorname{colim}_{U(x \downarrow W)} G).$$

As  $U(x \downarrow W)$  is filtered (Corollary 11.5), the result would then follow since filtered colimits commute with finite limits (cf. ([Lur09, Prop. 5.3.3.3])).

Comparing the codomains of  $\varphi$  and the canonical map, the diagram  $\operatorname{colim}_{U(x \downarrow W)} G \colon K^{\operatorname{op}} \to S$  is naturally  $E^1$ -homotopic to the diagram

$$K^{\mathrm{op}} \xrightarrow{\max^* \circ F} \mathrm{Ex}_+(\mathcal{C}, W) \xrightarrow{\mathrm{Ex}_+(\mathcal{C}, W)(-, x)} \mathcal{S}$$

by Corollary 10.9. Comparing the domains, the diagram  $\lim_{K^{op}} G: U(x \downarrow W) \to S$  is naturally  $E^1$ -homotopic to the diagram

$$U(x \downarrow W) \xrightarrow{\Pi_x} \mathcal{C} \xrightarrow{\mathcal{C}(\operatorname{colim} F, -)} \mathcal{S}$$

as the Yoneda embedding  $Y: \mathcal{C}^{\mathrm{op}} \to \mathcal{S}^{\mathcal{C}}$  takes colim  $F = \lim(F^{\mathrm{op}})$  to  $\lim(Y \circ F^{\mathrm{op}})$  (cf. [Lur09, Prop. 5.1.3.2]). Thus, the colimit of  $\lim_{K^{\mathrm{op}}} G$  is naturally equivalent to the mapping space  $\operatorname{Ex}_{+}(\mathcal{C}, W)(\operatorname{colim} F, x)$  by Corollary 10.9.

For (b), this follows from (a) by Lemma 11.6.

# Application to discrete homotopy theory

In the remainder of the paper, we describe an application of our theory to the area of discrete homotopy theory, which is a combinatorially meaningful homotopy theory of graphs. We begin with an overview of the field of discrete homotopy theory, which culminates with the construction of an  $(\infty, 1)$ -category of graphs  $N_{\Box}$ Graph. This  $(\infty, 1)$ -category is not quite the correct object, being the localization of the category of graphs at the class of homotopy equivalences, not weak equivalences. We show however that the pair ( $N_{\Box}$ Graph, weak equiv's) satisfies calculus of right fractions and hence Theorem 9.6 can be used to compute the localization. This then allows us to prove that the functor N: Graph  $\rightarrow$  cSet constructed in [CK22a] ascends to a representable functor between the corresponding  $(\infty, 1)$ -categories.

Throughout, we freely use the language of cubical homotopy theory, including the model structures on cubical sets and cubical categories. The reader unfamiliar with these notions may wish to consult Appendices A and B, where the requisite background is collected.

### 12 Background on discrete homotopy theory

Discrete homotopy theory, introduced by Barcelo and collaborators in [BKLW01, BBdLL06], is a novel approach to the homotopy theory of graphs. The usual viewpoint on graphs in algebraic topology is that they are 1-dimensional CW-complexes and hence their homotopy type is completely determined by the Euler characteristic  $\chi(X)$  of each connected component, defined as the number of vertices minus the number of edges. The corresponding homotopy type is then the coproduct of Eilenberg–Mac Lane spaces  $K(F_{1-\chi(X)}, 1)$ , where  $F_{1-\chi(X)}$  is the free group on  $1 - \chi(X)$  generators. As indicated, discrete homotopy theory departs from this view, instead defining homotopy groups of a graph to carry combinatorial meaning. Consequently, it is possible for graphs to have non-trivial homotopy groups in any number of degrees.

The origins of discrete homotopy theory go back to the work of Atkin [Atk74, Atk76], a physicist, who began developing homotopy theory of simplicial complexes to model certain social and technological networks. Kramer and Laubenbacher made these ideas mathematically precise [KL98], while Barcelo [BKLW01, BBdLL06] realized that the natural framework for them lies in the category of graphs, which initiated a systematic study of the field.

Since then, the area of discrete homotopy theory has found a number of applications, including in combinatorics (to hyperplane arrangements and matroid theory) and, more recently, in topological data analysis [BCW14, MZ19].

We begin our introduction to the subject with the definition of a graph:

#### Definition 12.1.

- 1. A graph is a set G with a symmetric and reflexive relation  $\sim$ . The elements of G are called vertices and two vertices v and w are connected by an edge if  $v \sim w$ .
- 2. A graph map  $f: (G, \sim) \to (H, \sim)$  is a function  $f: G \to H$  between sets preserving the relation.

3. The category of graphs and graph maps is denoted Graph.

In other words, our graphs are reflexive graphs without multiple edges. When depicting graphs we will omit the loops on each vertex. We will generally omit the relation from the notation used for graphs, denoting graphs simply with capital letters  $G, H, K, \ldots$ 

We mention the following examples of graphs:

### Example 12.2.

1. The graph  $I_m$  has, as vertices, the integers  $0, 1, \ldots, m$  and  $i \sim j$  whenever |i - j| = 1. For instance, we can depict  $I_3$  as follows:

$$\underbrace{ \begin{array}{ccc} 0 & 1 & 2 & 3 \\ \bullet & \bullet & \bullet & \bullet \\ \end{array} }$$

Note in particular that  $I_0$  is the graph with a single vertex and hence the terminal object in the category of graphs.

- 2. The graph  $I_{\infty}$  has, as vertices, all integers with  $i \sim j$  whenever |i j| = 1.
- 3. The graph  $C_m$  is obtained from  $I_m$  by identifying the vertices 0 and m. For instance, we can depict  $C_3$  as



**Definition 12.3.** The box product of graphs G and H is a graph  $G \otimes H$  whose set of vertices is  $G \times H$  and where  $(v, w) \sim (v', w')$  if either

- 1. v = v' and  $w \sim w'$ ; or
- 2.  $v \sim v'$  and w = w'.

One easily verifies that the box product equips the category of graphs with a symmetric monoidal structure. This structure is moreover closed: the functor  $G \otimes -:$  Graph  $\rightarrow$  Graph admits a right adjoint given by the internal hom graph hom<sup> $\otimes$ </sup>(G, H) whose vertices are graph maps  $G \rightarrow H$  with  $f \sim g$  whenever  $f(v) \sim g(v)$  for every vertex  $v \in G$ .

#### Definition 12.4.

1. Let  $f, g: G \to H$  be graph maps. A homotopy from f to g is a graph map

$$\alpha\colon G\otimes I_m\to H$$

for some integer  $m \ge 0$  such that  $\alpha|_{G \otimes \{0\}} = f$  and  $\alpha|_{G \otimes \{m\}} = g$ .

2. A graph map  $f: G \to H$  is a homotopy equivalence if there is a map  $g: H \to G$  along with homotopies

$$\alpha: gf \sim \mathrm{id}_X$$
 and  $\beta: fg \sim \mathrm{id}_Y$ .

**Example 12.5.** With the definitions above, it can be shown that  $I_m \to I_0$  is a homotopy equivalence, but the map  $I_{\infty} \to I_0$  is not.

Similarly, we can speak of pointed graphs (i.e. graphs with a distinguished vertex), pointed maps (i.e. maps preserving the distinguished vertex), and based homotopies (i.e. homotopies constant on the distinguished point).

**Definition 12.6.** Let  $(G, v_0)$  be a pointed graph. The *n*-th homotopy group of  $(G, v_0)$  is

$$A_n(G, v_0) = \{f \colon I_{\infty}^{\otimes n} \to G \mid f(\vec{m}) = v_0 \text{ for all but finitely many } \vec{m} \in I_{\infty}^{\otimes n}\} / \sim_*,$$

where the equivalence relation  $\sim_*$  is based homotopy.

One can verify that  $A_n(G, v_0)$  are groups for  $n \ge 1$  and are abelian for  $n \ge 2$ . These homotopy groups carry important information about the graph and are at the forefront of the aforementioned applications of discrete homotopy theory to matroid theory, hyperplane arrangements, and topological data analysis. Unfortunately, computing them in general is difficult and little is known about them.

**Definition 12.7.** A graph map  $f: G \to H$  is a *weak equivalence* if for all vertices  $v \in G$  and all non-negative integers  $n \ge 0$ , the induced map

$$f_*: A_n(G, v) \to A_n(H, f(v))$$

is an isomorphism.

**Example 12.8.** It can be shown that all cycles  $C_m$  are weakly equivalent for  $m \ge 5$ . Furthermore,  $I_{\infty} \to I_0$  is a weak equivalence.

One can ask whether there is a construction associating a space to a graph in such a way that the usual homotopy groups of the space agree with the homotopy groups of a graph as defined above. A tentative construction of that kind was given in [BBdLL06] and proven to work in [CK22a]. Here, we will present a different construction, N: Graph  $\rightarrow$  cSet called the *nerve* of a graph, also introduced in [CK22a] for the purpose of proving the conjecture of [BBdLL06]. Our model for spaces is therefore the category cSet of cubical sets. Readers unfamiliar with cubical sets are welcome to consult Appendix A for the necessary background or one of the references [Cis06, DKLS20, CK22b].

Before introducing the functor, we need a preliminary notion of stability of a graph map. To state it, we fix a graph G and  $n \ge 1$ . For i = 0, ..., n and  $\varepsilon = 0, 1$ , a map  $f: I_{\infty}^{\otimes n} \to G$  is stable in direction  $(i, \varepsilon)$ if there exists  $M \ge 0$  so that for  $m_i > M$ , we have

$$f(m_1,\ldots,(2\varepsilon-1)m_i,\ldots,m_n)=f(m_1,\ldots,(2\varepsilon-1)M,\ldots,m_n).$$

We are now ready to define the nerve functor N:  $\mathsf{Graph} \to \mathsf{cSet}$ .

**Definition 12.9.** The *nerve* of a graph G is a cubical set NG whose *n*-cubes are given by

 $(NG)_n = \{f \colon I_{\infty}^{\otimes n} \to G \mid f \text{ is stable in all directions}\}$ 

The key property of cubical sets arising as nerves of graphs is that they are always fibrant, i.e., Kan complexes.

**Theorem 12.10** ([CK22a, Thm. 4.5]). For any graph G, its nerve NG is a Kan complex.

Moreover, the homotopy groups of the nerve (and hence the geometric realization thereof) agree with the homotopy groups of the graph. Here, the homotopy groups of a cubical sets are as defined, e.g., in [Kan55] and investigated further in [CK22b].

**Theorem 12.11** ([CK22a, Thm. 4.6]). For any pointed graph  $(G, v_0)$ , we have canonical isomorphisms:

$$A_n(G, v_0) \cong \pi_n(\mathrm{N}G, v_0), \cong \pi_n(|\mathrm{N}G|, v_0).$$

As a closed monoidal category, the category of graphs is enriched over itself. Since the nerve functor N: Graph  $\rightarrow$  cSet is lax monoidal [CK22a, Lem. 5.16.(2)], we may apply Example B.8 to obtain that Graph (or N<sub>•</sub>Graph, in the notation of Appendix B) is a cubical category with the mapping cubical set from G to H given by N hom<sup> $\otimes$ </sup>(G, H).

It follows from Theorem 12.10 that the cubical category **Graph** is locally Kan, i.e., that all mapping cubical sets are Kan complexes. Therefore, its (cubical) homotopy coherent nerve  $N_{\Box}$  **Graph** is a quasicategory (Proposition B.12). The reader unfamiliar with the (cubical) homotopy coherent nerve  $N_{\Box}$ : Cat<sub> $\Box$ </sub>  $\rightarrow$  sSet, as defined in [KV20], is welcome to consult Appendix B, where we recall its definition and main properties.

#### Corollary 12.12.

- 1. The cubical category Graph is locally Kan.
- 2. The simplicial set  $N_{\Box}$  Graph is a quasicategory.

It is easy to see that the quasicategory  $N_{\Box}$ Graph is the localization of the category of graphs at the class of homotopy equivalences. However, for applications in discrete homotopy theory, one is interested in the localization of Graph at the class of weak equivalences. To identify it, we will show that the marked quasicategory ( $N_{\Box}$ Graph, weak equiv's) satisfies CRF, which then establishes  $Ex^{op}_{+}(N_{\Box}$ Graph, weak equiv's) as the required localization.

# **13** Finite limits in $N_{\Box}$ Graph

To verify that  $(N_{\Box}Graph$ , weak equiv's) satisfies CRF, we wish to appeal to Theorem 6.3 and hence we must in particular establish that  $N_{\Box}Graph$  has all finite limits. To this end, our next goal is to show that  $N_{\Box}Graph$  has pullbacks.

We begin by recalling the path graph functor, which we denote by  $(-)^{I}$ : Graph  $\rightarrow$  Graph. For a graph G, the path graph  $G^{I}$  has

- as vertices, graph maps  $I_{\infty} \to G$  which are stable in both directions;
- an edge  $f \sim g$  if  $f(m) \sim g(m)$  for all  $m \in I_{\infty}$ .

As well, for a graph map  $f: G \to K$ , we write  $P_f$  for the double mapping path space of  $id_K$  and f, i.e. the pullback

Fix a cospan  $(f: G \to K, g: H \to K)$  of graphs and let P denote the limit of the diagram



We denote the limit projections into the path space of K as  $\pi_1, \pi_2 \colon P \to K^I$ , respectively. We also write  $\pi_G, \pi_H, \pi_K$  for the respective limit projections, where  $\pi_K$  denotes the middle projection in the diagram; that is,  $\pi_K = \partial_{1,0}^* \pi_1 = \partial_{1,0}^* \pi_2$ .

The map  $\pi_1: P \to K^I$  is a 0-cube in N hom<sup> $\otimes$ </sup>(P,  $K^I$ ). Via the isomorphism N hom<sup> $\otimes$ </sup>(P,  $K^I$ )  $\cong$  hom<sub>L/R</sub>( $\Box^1$ , N hom<sup> $\otimes$ </sup>(P, K)) of [CK22a, Prop. 3.10], we may identify  $\pi_1$  as a 1-cube in N hom<sup> $\otimes$ </sup>(P, K)

from  $\pi_K$  to  $f\pi_G$ . Likewise, we regard the map  $\pi_2 \colon P \to K^I$  as a 1-cube in N hom<sup> $\otimes$ </sup>(P,K) from  $\pi_K$  to  $g\pi_H$ , giving the data of a cone  $\Pi \colon \Delta^0 * \Lambda_2^2 \to \mathbb{N}_{\Box}$ Graph over (f,g).



**Theorem 13.1.** For any co-span  $(f,g): \Lambda_2^2 \to N_{\Box}\mathsf{Graph}$ , the square  $\Pi: \Delta^0 * \Lambda_2^2 \to N_{\Box}\mathsf{Graph}$ 



is a pullback in  $N_{\Box}$ Graph.

Before proving this theorem, we note the following computation.

**Proposition 13.2.** For  $n \ge 1$ , identifying  $\partial \Delta^n * \Delta^1$  as a simplicial subset of  $\Delta^{n+2}$ , we have that

$$\mathfrak{C}(\partial \Delta^n * \Delta^1)(i,j) \cong \begin{cases} \Box^{j-i-1} & \text{for } i > 0 \text{ or } i = 0, j < n \\ \partial \Box^{n-1} & \text{for } i = 0, j = n \\ \Box^n_{n,1} & \text{for } i = 0, j = n+1 \\ \Box^n_{n,1} \otimes \Box^1 & \text{for } i = 0, j = n+2. \end{cases}$$

Proof of Theorem 13.1. Fix a diagram  $\lambda: \partial \Delta^n * \Lambda_2^2 \to \mathcal{N}_{\Box}$ Graph such that  $\lambda|_{\{n\}*\Lambda_2^2} = \Pi$ . Identifying  $\Delta^n * \Lambda_2^2$  as the pushout

and  $\partial \Delta^n * \Lambda_2^2$  as the pushout

$$\begin{array}{c|c} \partial \Delta^n * \Delta^0 \xrightarrow{\operatorname{id}_{\partial \Delta^n} * \{1\}} \partial \Delta^n * \Delta^1 \\ \downarrow^{\operatorname{id}_{\partial \Delta^n} * \{1\}} & & \downarrow^{i_2} \\ \partial \Delta^n * \Delta^1 \xrightarrow{\quad i_1 \quad \to \quad \partial \Delta^n * \Lambda_2^2} \end{array}$$

it suffices to construct fillers

$$F, G: \Delta^n * \Delta^1 \to \mathcal{N}_\square$$
 Graph

for the maps

$$\lambda i_1, \lambda i_2 \colon \partial \Delta^n * \Delta^1 \to \mathcal{N}_\square \mathsf{Graph}$$

such that  $F|_{\Delta^n * \partial_0} = G|_{\Delta^n * \partial_0}$ . We identify  $\Delta^n * \Delta^1$  with  $\Delta^{n+2}$  and  $\partial \Delta^n * \Delta^1$  with a simplicial subset of  $\Delta^{n+2}$  so that we may restate this equality as  $F\partial_{n+1} = G\partial_{n+1}$  where F, G are maps  $\Delta^{n+2} \to \mathbb{N}_{\Box}$ Graph.

Let

# $D,D'\colon \mathfrak{C}(\partial\Delta^n\ast\Delta^1)\to\mathsf{Graph}$

denote the adjoint transposes of  $\lambda i_1$  and  $\lambda i_2$  under the  $(\mathfrak{C} \dashv N_{\Box})$ -adjunction, respectively. Functoriality gives cubical maps

 $D_{0,n+1} \colon \sqcap_{n,1}^{n} \to \operatorname{N} \hom^{\otimes}(D(0), G)$  $D'_{0,n+1} \colon \sqcap_{n,1}^{n} \to \operatorname{N} \hom^{\otimes}(D'(0), H)$  $D_{0,n+2} \colon \sqcap_{n,1}^{n} \otimes \square^{1} \to \operatorname{N} \hom^{\otimes}(D(0), K)$  $D'_{0,n+2} \colon \sqcap_{n,1}^{n} \otimes \square^{1} \to \operatorname{N} \hom^{\otimes}(D'(0), K).$ 

where computation of the mapping spaces of  $\mathfrak{C}(\partial \Delta^n * \Delta^1)$  follows from Proposition 13.2. As  $\lambda i_1|_{\partial \Delta^n * \{1\}} = \lambda i_2|_{\partial \Delta^n * \{1\}}$  by assumption, we have that D(0) = D'(0) and  $D_{0,n+2}|_{\bigcap_{n,1}^n \otimes \{1\}} = D'_{0,n+2}|_{\bigcap_{n,1}^n \otimes \{1\}}$ . As the codomains are all Kan, we define maps

- $u_G: \square^n \to \operatorname{N} \hom^{\otimes}(D(0), G)$ , which is a lift of  $D_{0,n+1}$ ;
- $u_H \colon \Box^n \to \operatorname{N} \hom^{\otimes}(D(0), H)$ , which is a lift of  $D'_{0,n+1}$ ; and
- $u_K : \square^n \to \operatorname{N} \operatorname{hom}^{\otimes}(D(0), K)$ , which is a lift of  $D_{0,n+2}|_{\bigcap_{n,1}^n \otimes \{1\}} = D'_{0,n+2}|_{\bigcap_{n,1}^n \otimes \{1\}}$ .

These lifts assemble into maps  $t_1, t_2 \colon \sqcap_{n,1}^{n+1} \to \operatorname{N} \hom^{\otimes}(D(0), K)$  defined by the face assignments

$$t_1 \partial_{i,\varepsilon} := \begin{cases} D_{0,n+2} \partial_{i,\varepsilon} & \text{for } i < n \text{ or } (i,\varepsilon) = (n,0) \\ u_K & \text{for } (i,\varepsilon) = (n+1,0) \\ f_* u_G & \text{for } (i,\varepsilon) = (n+1,1) \end{cases}$$

and

$$t_2 \partial_{i,\varepsilon} := \begin{cases} D'_{0,n+2} \partial_{i,\varepsilon} & \text{for } i < n \text{ or } (i,\varepsilon) = (n,0) \\ u_K & \text{for } (i,\varepsilon) = (n+1,0) \\ g_* u_H & \text{for } (i,\varepsilon) = (n+1,1), \end{cases}$$

where

$$f_* \colon \operatorname{N} \hom^{\otimes}(D(0), G) \to \operatorname{N} \hom^{\otimes}(D(0), K) \quad g_* \colon \operatorname{N} \hom^{\otimes}(D(0), H) \to \operatorname{N} \hom^{\otimes}(D(0), K)$$

are the maps induced by post-composition. The maps  $t_1, t_2$  admits lifts which we denote by

$$u_1, u_2 \colon \Box^{n+1} \to \operatorname{N} \hom^{\otimes}(D(0), K).$$

By adjointness, we identify  $u_1$  and  $u_2$  as maps

$$u_1, u_2 \colon \Box^n \to \hom_L(\Box^1, \operatorname{N} \hom^{\otimes}(D(0), K)) \cong \operatorname{N}\left(\hom^{\otimes}(D(0), K)^I\right),$$

where the isomorphism follows from [CK22a, Prop. 3.10]. There is a natural monomorphism

$$\hom^{\otimes}(D(0), K)^{I} \to \hom^{\otimes}(D(0), K^{I})$$

which we post-compose with to obtain

$$\overline{u_1}, \overline{u_2} \colon \Box^n \to \operatorname{N} \hom^{\otimes}(D(0), K^I).$$

As N hom<sup> $\otimes$ </sup>(D(0), -): Graph  $\rightarrow$  cSet preserves finite limits, the cone



induces a map  $u: \square^n \to \operatorname{N} \hom^{\otimes}(X_0, P)$ . By construction, u is a filler for the maps

 $D(0,n), D'(0,n): \partial \Box^n \to \operatorname{N} \hom^{\otimes}(D(0), P).$ 

Thus, the functors  $D, D' : \mathfrak{C}(\partial \Delta^n * \Delta^1) \to \mathsf{Graph}$  lift to functors  $\mathfrak{C}[n+2] \to \mathsf{Graph}$  satisfying the required equalities, which suffices by adjointness.

**Corollary 13.3.** The quasicategory  $N_{\Box}$  Graph has finite limits.

*Proof.* It has pullbacks by Theorem 13.1. As  $I_0$  is a terminal object, this suffices by [Lur09, Cor. 4.4.2.4].

## 14 $(\infty, 1)$ -categorical setting for discrete homotopy theory

In this section, we apply earlier results to develop the  $(\infty, 1)$ -categorical foundations for discrete homotopy theory. More precisely, we construct the quasicategory of graphs, which is the localization of the category of graphs at weak equivalences (Theorem 14.2), show that the graph nerve functor is representable, represented by  $I_0$  (Proposition 14.1), and conjecture that it is an equivalence of quasicategories (Conjecture 14.3).

The nerve functor N: Graph  $\rightarrow$  Kan $_{\Box}$  extends to a cubical functor  $\overline{N}$ : Graph  $\rightarrow$  Kan $_{\Box}$  between cubical categories by Example B.5. The action on mapping spaces N hom<sup> $\otimes$ </sup>(G, H)  $\rightarrow$  hom<sub>L</sub>(NG, NH) is the adjoint transpose of the composite

$$\operatorname{N} \operatorname{hom}^{\otimes}(G, H) \otimes \operatorname{N} G \hookrightarrow \operatorname{N}(\operatorname{hom}^{\otimes}(G, H) \otimes G) \xrightarrow{\operatorname{N} \varepsilon} \operatorname{N} H.$$

In Theorem B.20, we construct a canonical equivalence  $N_{\Box}\mathsf{Kan}_{\Box} \to N_{\Delta}\mathsf{Kan}_{\Delta}$  between cubical Kan complexes and simplicial Kan complexes. Thus,  $N_{\Box}\mathsf{Kan}_{\Box}$  serves as a model for  $\infty$ -groupoids, allowing us to phrase the question of whether the functor  $N_{\Box}\overline{N}: N_{\Box}\mathsf{Graph} \to N_{\Box}\mathsf{Kan}_{\Box}$  is representable (meaning that post-composing  $\overline{N}$  with the canonical equivalence produces a representable functor).

Proposition 14.1. 1. The composite map

$$\mathrm{N}_{\Box}\mathsf{Graph} \xrightarrow{\mathrm{N}_{\Box}\mathrm{N}} \mathrm{N}_{\Box}\mathsf{Kan}_{\Box} \to \mathrm{N}_{\Delta}\mathsf{Kan}_{\Delta}$$

is representable, represented by  $I_0$ .

2. The map

 $\mathrm{N}_{\Box}\overline{\mathrm{N}} \colon \mathrm{N}_{\Box}\mathsf{Graph} \to \mathrm{N}_{\Box}\mathsf{Kan}_{\Box}$ 

preserves finite limits.

*Proof.* For (1), we note that the cubical functors

$$\mathsf{Graph} \xrightarrow[\mathsf{Graph}(I_0,-)]{\overline{\mathsf{Kan}}} \mathsf{Kan}_{\square}$$

are E[1]-homotopic. This suffices by statement (2) of Theorem B.20.

Item (2) follows from item (1) by [Lur09, Prop. 5.1.3.2].

#### Theorem 14.2.

- 1. The quasicategory  $N_{\Box}$  Graph marked at the collection W of weak homotopy equivalences satisfies CRF.
- 2. The localization  $N_{\Box}$ Graph $[W^{-1}]$  of  $N_{\Box}$ Graph at weak homotopy equivalences has all finite limits.
- 3. The induced map

$$N_{\Box} \operatorname{Graph}[W^{-1}] \xrightarrow{N} N_{\Box} \operatorname{Kan}$$

preserves finite limits.

4. The composite

$$\mathrm{N}_{\Box}\mathsf{Graph}[W^{-1}] \xrightarrow{\overline{\mathrm{N}}} \mathrm{N}_{\Box}\mathsf{Kan}_{\Box} o \mathrm{N}_{\Delta}\mathsf{Kan}_{\Delta}$$

is representable.

*Proof.* For (1), we have that  $N_{\Box}$ Graph has all finite limits (Corollary 13.3) and  $N_{\Box}\overline{N}$  preserves them (Proposition 14.1). Thus, the pair ( $N_{\Box}$ Graph, W) satisfies CRF by Theorem 6.3. Item (2) then follows from Theorem 11.7.

We prove (4), from which (3) follows. By the universal property of the localization, we wish to show the composite map

$$\mathrm{N}_{\Box}\mathsf{Graph} \xrightarrow{\mathrm{max}^*} \mathrm{Ex}^{\mathrm{op}}_+(\mathrm{N}_{\Box}\mathsf{Graph}, W) \xrightarrow{\overline{\mathrm{N}}} \mathrm{N}_{\Box}\mathsf{Kan}_{\Box} \to \mathrm{N}_{\Delta}\mathsf{Kan}_{\Delta}$$

is representable. By Proposition 14.1, it suffices to show the triangle

$$\begin{array}{c|c} \mathbf{N}_{\Box}\mathsf{Graph} & \xrightarrow{\mathbf{N}_{\Box}\mathsf{Graph}(I_{0},-)} & \mathbf{N}_{\Box}\mathsf{Kan}_{\Box} \\ & \xrightarrow{\mathbf{max}^{*}} & \xrightarrow{\mathbf{Fx}^{\mathrm{op}}_{+}(\mathbf{N}_{\Box}\mathsf{Graph},W)(I_{0},-)} \\ & \mathrm{Ex}^{\mathrm{op}}_{+}(\mathbf{N}_{\Box}\mathsf{Graph},W) \end{array}$$

commutes up to  $E^1$ -homotopy. For a graph G, the mapping space  $\operatorname{Ex}_+^{\operatorname{op}}(\mathbb{N}_{\Box}\operatorname{\mathsf{Graph}}, W)(I_0, G)$  is a filtered colimit of mapping spaces in  $\mathbb{N}_{\Box}\operatorname{\mathsf{Graph}}$  indexed over  $(W \downarrow I_0)^{\operatorname{op}}$  (Corollary 10.9), hence it suffices to show  $I_0$  is an initial object in the marked "fat" slice  $W \downarrow I_0$ . By Proposition 10.3, it suffices to show  $I_0$  is an initial object in the marked ordinary slice  $W \downarrow^* I_0$ .

Fix a map  $\partial \Delta^n \to W \downarrow^* I_0$  whose value at 0 is  $I_0$ . We identify this map as a cubical functor  $u: \mathfrak{C}(\Lambda_{n+1}^{n+1}) \to \text{Graph}$  such that

- $u(0) = u(n+2) = I_0$ ; and
- for i = 0, ..., n + 1, the map

$$u(i \le n+2) \colon u(i) \to I_0$$

is a weak homotopy equivalence.

We wish to lift u to a functor  $\mathfrak{C}[n+1] \to \mathsf{Graph}$ , which corresponds to the data of lifts  $v_1, v_2$ 

such that  $v_2\partial_{n+1,1} = u(n+1 \le n+2)_*(v_1)$ . As  $\hom^{\otimes}(I_0, u(n+1)) \cong u(n+1)$  is weakly contractible by assumption, the lift  $v_1$  exists. As  $\hom^{\otimes}(I_0, I_0) \cong I_0$ , the lift  $v_2$  exists and the required equality is immediate.

We conclude with a conjecture that the functor  $\overline{N}: N_{\Box}\mathsf{Graph}[W^{-1}] \to N_{\Box}\mathsf{Kan}_{\Box}$  is an equivalence of quasicategories.

Conjecture 14.3. The functor

$$\overline{\mathbf{N}} \colon \mathbf{N}_{\Box}\mathsf{Graph}[W^{-1}] \to \mathbf{N}_{\Box}\mathsf{Kan}_{\Box}$$

is an equivalence of quasicategories.

# Appendices

### A Cubical sets

In this section, we recall the definitions and results regarding the homotopy theory of cubical sets which are used in Sections 12, 13 and 14. The most significant of these is the *Grothendieck* model structure on cubical sets, its fibrant objects (i.e. the *cubical Kan complexes*), and the *triangulation* Quillen equivalence (Theorem A.5). Our primary reference is [DKLS20] — other references on the topic include [Cis06],[Cis14], [Jar06], and [KV20].

We begin by defining the cube category  $\Box$ . The objects of  $\Box$  are posets of the form  $[1]^n = \{0 \le 1\}^n$  and the maps are generated (inside the category of posets) under composition by the following four special classes:

• faces  $\partial_{i,\varepsilon}^n \colon [1]^{n-1} \to [1]^n$  for  $i = 1, \ldots, n$  and  $\varepsilon = 0, 1$  given by:

 $\partial_{i,\varepsilon}^{n}(x_{1}, x_{2}, \dots, x_{n-1}) = (x_{1}, x_{2}, \dots, x_{i-1}, \varepsilon, x_{i}, \dots, x_{n-1});$ 

• degeneracies  $\sigma_i^n \colon [1]^n \to [1]^{n-1}$  for  $i = 1, 2, \dots, n$  given by:

$$\sigma_i^n(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n);$$

• negative connections  $\gamma_{i,0}^n \colon [1]^n \to [1]^{n-1}$  for  $i = 1, 2, \ldots, n-1$  given by:

$$\gamma_{i,0}^n(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, \max\{x_i, x_{i+1}\}, x_{i+2}, \dots, x_n).$$

• positive connections  $\gamma_{i,1}^n: [1]^n \to [1]^{n-1}$  for  $i = 1, 2, \dots, n-1$  given by:

$$\gamma_{i,1}^n(x_1, x_2, \dots, x_n) = (x_1, x_2, \dots, x_{i-1}, \min\{x_i, x_{i+1}\}, x_{i+2}, \dots, x_n).$$

These maps obey the following *cubical identities*:

$$\begin{split} \partial_{j,\varepsilon'}\partial_{i,\varepsilon} &= \partial_{i+1,\varepsilon}\partial_{j,\varepsilon'} \quad \text{for } j \leq i; \\ \sigma_{i}\sigma_{j} &= \sigma_{j}\sigma_{i+1} \quad \text{for } j \leq i; \\ \gamma_{j,\varepsilon'}\partial_{i,\varepsilon} &= \begin{cases} \partial_{i-1,\varepsilon}\sigma_{j} & \text{for } j < i; \\ \text{id} & \text{for } j = i; \\ \partial_{i,\varepsilon}\sigma_{j-1} & \text{for } j > i; \\ \gamma_{j,\varepsilon'}\gamma_{i,\varepsilon} &= \begin{cases} \gamma_{i,\varepsilon}\gamma_{j+1,\varepsilon'} & \text{for } j > i; \\ \gamma_{i,\varepsilon}\gamma_{i+1,\varepsilon} & \text{for } j = i, \varepsilon' = \varepsilon; \\ \partial_{j,\varepsilon}\sigma_{j} & \text{for } j = i-1, i, \varepsilon = \varepsilon'; \\ \partial_{j,\varepsilon}\sigma_{j} & \text{for } j = i-1, i, \varepsilon = 1-\varepsilon'; \\ \partial_{i,\varepsilon}\gamma_{j-1,\varepsilon'} & \text{for } j > i; \end{cases} \\ \sigma_{j}\gamma_{i,\varepsilon} &= \begin{cases} \gamma_{i-1,\varepsilon}\sigma_{j} & \text{for } j < i; \\ \sigma_{i}\sigma_{i} & \text{for } j = i; \\ \gamma_{i,\varepsilon}\sigma_{j+1} & \text{for } j > i. \end{cases} \end{split}$$

Our convention is to write cubical operators on the right e.g. given an *n*-cube  $x \in X_n$  of a cubical set X, we write  $x\partial_{1,0}$  for the  $\partial_{1,0}$ -face of x.

#### Definition A.1.

- 1. For  $n \ge 0$ , the combinatorial n-cube  $\square^n$  is the representable functor  $\square(-, [1]^n) : \square^{\text{op}} \to \mathsf{Set}$ .
- 2. The boundary of the n-cube  $\partial \Box^n$  is the subobject of  $\Box^n$  defined by

$$\partial \Box^n := \bigcup_{\substack{j=1,\dots,n\\\eta=0,1}} \operatorname{Im} \partial_{j,\eta}$$

3. given i = 0, ..., n and  $\varepsilon = 0, 1$ , the  $(i, \varepsilon)$ -open box  $\sqcap_{i,\varepsilon}^{n}$  is the subobject of  $\partial \square^{n}$  defined by

$$\sqcap_{i,\varepsilon}^n:=\bigcup_{(j,\eta)\neq (i,\varepsilon)}\operatorname{Im}\partial_{j,\eta}$$

Cubical sets are related to simplicial sets via the following adjunction.

**Definition A.2.** Define a functor  $\Box \to \mathsf{sSet}$  which sends  $[1]^n$  to  $(\Delta^1)^n$ . Left Kan extension along the Yoneda embedding gives the *triangulation* functor

whose right adjoint  $U: \mathsf{sSet} \to \mathsf{cSet}$  is defined by

$$(UX)_n := \mathsf{sSet}\left((\Delta^1)^n, X\right).$$

By construction, there is an isomorphism  $|X|_{\Box} \cong |T(X)|_{\Delta}$  between the cubical geometric realization of X and the simplicial geometric realization of T(X) (which is natural in X).

Define a monoidal product  $- \otimes -: \Box \times \Box \to \Box$  on the cube category by  $[1]^m \otimes [1]^n = [1]^{m+n}$ . Postcomposing with the Yoneda embedding and left Kan extending gives the *geometric product* of cubical sets.



This product is biclosed: for a cubical set X, we write  $\hom_L(X, -)$ :  $\mathsf{cSet} \to \mathsf{cSet}$  and  $\hom_R(X, -)$ :  $\mathsf{cSet} \to \mathsf{cSet}$  for the right adjoints to the functors  $- \otimes X$  and  $X \otimes -$ , respectively.

Moreover, the triangulation functor is strong monoidal, i.e. it sends geometric products to products.

### Proposition A.3.

- 1. The triangulation functor  $T: cSet \rightarrow sSet$  is strong monoidal.
- 2. Its right adjoint  $U: \mathsf{sSet} \to \mathsf{cSet}$  is lax monoidal.

*Proof.* For (1), it suffices to check on representables, on which it follows by definition as

$$T(\Box^m \otimes \Box^n) = T(\Box^{m+n})$$
  
=  $(\Delta^1)^{m+n}$   
=  $(\Delta^1)^m \times (\Delta^1)^n$   
=  $T(\Box^m) \times T(\Box^n).$ 

Item (2) follows formally from item (1).

The following result gives an explicit description of cubes in the geometric product, which we use implicitly throughout Appendix B

**Proposition A.4** ([DKLS20, Prop. 1.24]). Let X, Y be cubical sets.

- 1. For  $k \ge 0$ , the k-cubes of  $X \otimes Y$  consists of all pairs  $(x \in X_m, y \in Y_n)$  such that m + n = k, subject to the identification  $(x\sigma_{m+1}, y) = (x, y\sigma_1)$ .
- 2. For  $x \in X_m$  and  $y \in Y_n$ , the faces, degeneracies, and connections of the (m + n)-cube (x, y) are computed by

$$\begin{split} (x,y)\partial_{i,\varepsilon} &= \begin{cases} (x\partial_{i,\varepsilon},y) & 1 \leq i \leq m\\ (x,y\partial_{i-m,\varepsilon}) & m+1 \leq i \leq m+n; \end{cases} \\ (x,y)\sigma_i &= \begin{cases} (x\sigma_i,y) & 1 \leq i \leq m+1\\ (x,y\sigma_{i-m}) & m+1 \leq i \leq m+n; \end{cases} \\ (x,y)\gamma_{i,\varepsilon} &= \begin{cases} (x\gamma_{i,\varepsilon},y) & 1 \leq i \leq m\\ (x,y\gamma_{i-m,\varepsilon}) & m+1 \leq i \leq n. \end{cases} \end{split}$$

The category of cubical sets carries the *Groethendieck* model structure (also known as the *Cisinski* model structure), which models the homotopy theory of spaces.

**Theorem A.5** (Cisinski, cf. [DKLS20, Thms. 1.34 & 6.26]). The category of cubical sets carries a model structure where

- cofibrations are monomorphisms;
- fibrations are maps with the right lifting property against all open box inclusions, i.e. the set

$$\left\{ \begin{array}{c} n \geq 1 \\ \sqcap_{i,\varepsilon}^{n} \hookrightarrow \square^{n} \mid i = 1, \dots, n \\ \varepsilon = 0, 1 \end{array} \right\};$$

and

• weak equivalences are created by triangulation, i.e. a map  $f: X \to Y$  is a weak equivalence if  $Tf: TX \to TY$  is a weak equivalence in the Kan-Quillen model structure.

Moreover, the triangulation adjunction

$$T: \mathsf{cSet} \rightleftharpoons \mathsf{sSet} : U$$

gives a Quillen equivalence with the Kan-Quillen model structure on simplicial sets.

We refer to fibrant cubical sets as cubical Kan complexes.

**Definition A.6.** A *cubical Kan complexes* (or simply *Kan complex*) is a cubical set with the right lifting property with respect to all open box inclusions.

## **B** Cubical categories

The purpose of this section is to establish the background on cubical categories used in the application of our results to discrete homotopy theory (see Sections 12, 13 and 14). The main result of this section (Theorem B.20) establishes an equivalence of categories between the cubical homotopy-coherent nerve of the category of cubical Kan complexes and the simplicial homotopy-coherent nerve of the category of simplicial Kan complexes. To do so, we make use of the model structure on cubical categories (Theorem B.16).

We recall some basic results about enriched categories, of which cubical and simplicial categories are examples.

**Definition B.1.** Let  $(\mathcal{V}, \otimes, 1)$  be a monoidal category. A  $\mathcal{V}$ -enriched category  $\mathsf{C}$  (or a category enriched in  $(\mathcal{V}, \otimes, 1)$ ) consists of

- a class of objects ob C;
- for  $x, y \in ob \mathsf{C}$ , a hom-object  $\mathsf{C}(x, y) \in \mathcal{V}$ ;
- for  $x, y, z \in ob C$ , a composition morphism

$$\circ \colon \mathsf{C}(y,z) \otimes \mathsf{C}(x,y) \to \mathsf{C}(x,z);$$

and

• for  $x \in ob C$ , an *identity element* morphism  $\underline{id}_x : 1 \to C(x, x)$ ,

such that, for all  $x, y, z, w \in \text{ob} \mathsf{C}$ , the diagrams

$$\begin{array}{c} \mathcal{C}(z,w)\otimes\mathcal{C}(y,z)\otimes\mathcal{C}(x,y) \xrightarrow{\mathcal{C}(z,w)\otimes\circ} \mathcal{C}(z,w)\otimes\mathcal{C}(x,z) \\ & & & & \downarrow \circ \\ & & & \downarrow \circ \\ & & \mathcal{C}(y,w)\otimes\mathcal{C}(x,y) \xrightarrow{\circ} \mathcal{C}(x,w) \end{array} \\ \mathcal{C}(x,y) \xrightarrow{\mathcal{C}(x,y)\otimes\underline{\mathrm{id}}_x} \mathcal{C}(x,y)\otimes\mathcal{C}(x,x) & & \mathcal{C}(y,y)\otimes\mathcal{C}(x,y) \xleftarrow{\underline{\mathrm{id}}_y\otimes\mathcal{C}(x,y)} \mathcal{C}(x,y) \\ & & & \downarrow \circ \\ & & & \downarrow \circ \\ & & & \downarrow \circ \\ & & & & \downarrow \circ \\ & & & & \mathcal{C}(x,y) \end{array} \end{array}$$

commute.

**Example B.2** ([EK66, Thms. I.5.2 & II.6.4]). If  $(\mathcal{V}, \otimes, 1)$  is left (or right) closed monoidal then, writing hom $(x, -): \mathcal{V} \to \mathcal{V}$  for the right adjoint to the functor  $- \otimes x$  (or  $x \otimes -$ ), the category  $\mathcal{V}$  ascends to a  $\mathcal{V}$ -enriched category where the hom-objects are given by hom(x, y).

**Example B.3** ([EK66, Prop. II.6.3]). If  $\Phi: \mathcal{W} \to \mathcal{V}$  is a lax monoidal functor then any  $\mathcal{W}$ -enriched category D gives rise to a  $\mathcal{V}$ -category  $\Phi_{\bullet}D$  with the same objects. The hom-object from x to y is given by  $\Phi(\mathsf{D}(x, y))$ .

**Definition B.4.** Let  $(\mathcal{V}, \otimes, 1)$  be a monoidal category. A  $\mathcal{V}$ -enriched functor  $F: \mathsf{C} \to \mathsf{D}$  between  $\mathcal{V}$ enriched categories consists of

- an assignment on objects  $ob C \rightarrow ob D$ ;
- for  $x, y \in ob \mathsf{C}$ , a morphism

$$F_{x,y}\colon \mathsf{C}(x,y)\to\mathsf{D}(Fx,Fy)$$

in  $\mathcal{V}$ 

such that, for  $x, y, z \in ob C$ , the diagrams

$$\begin{array}{ccc} \mathsf{C}(y,z)\otimes\mathsf{C}(x,y) & \stackrel{\circ}{\longrightarrow} \mathsf{C}(x,z) & \square^{0} & \stackrel{\underline{\operatorname{id}}_{x}}{\longrightarrow} \mathsf{C}(x,x) \\ F_{y,z}\otimes F_{x,y} \downarrow & \downarrow F_{x,z} & & \downarrow F_{x,x} \\ \mathsf{D}(Fy,Fz)\otimes\mathsf{D}(Fx,Fy) & \stackrel{\circ}{\longrightarrow} \mathsf{D}(Fx,Fz) & & \mathsf{D}(Fx,Fx) \end{array}$$

commute.

We write  $Cat_{\mathcal{V}}$  for the category of  $\mathcal{V}$ -enriched categories and  $\mathcal{V}$ -enriched functors.

**Example B.5** ([EK66, Thms. I.6.6 & II.6.4]). Any lax monoidal functor  $\Phi: \mathcal{W} \to \mathcal{V}$  between closed categories ascends to a  $\mathcal{V}$ -enriched functor

$$\overline{\Phi} \colon \Phi_{\bullet} \mathcal{W} \to \mathcal{V}$$

which sends an object  $x \in \Phi_{\bullet} \mathcal{W}$  to  $\Phi x \in \mathcal{V}$ . The action  $\overline{\Phi}_{x,y} \colon \Phi \mathcal{W}(x,y) \to \mathcal{V}(\Phi x, \Phi y)$  on hom-objects is adjoint transpose to the map

$$\Phi(\mathcal{W}(x,y)) \otimes \Phi x \to \Phi(\mathcal{W}(x,y) \otimes x) \xrightarrow{\Phi(\operatorname{eval}_y)} \Phi y.$$

**Example B.6** ([EK66, Prop. II.6.3]). The construction given in Example B.3 is functorial, i.e. any lax monoidal functor  $\Phi: \mathcal{W} \to \mathcal{V}$  induces a functor

$$\Phi_{\bullet} \colon \mathsf{Cat}_{\mathcal{W}} \to \mathsf{Cat}_{\mathcal{V}}.$$

In particular, any W-enriched functor  $\mathsf{C} \to \mathsf{D}$  gives rise to a V-enriched functor  $\Phi_{\bullet}\mathsf{C} \to \Phi_{\bullet}\mathsf{D}$ .

Given another lax monoidal functor  $\Psi: \mathcal{U} \to \mathcal{W}$ , the functor  $(\Phi\Psi)_{\bullet}$  is naturally isomorphic to  $\Phi_{\bullet}\Psi_{\bullet}$ .

**Definition B.7.** Given  $\mathcal{V}$ -enriched functors  $F, G: \mathsf{C} \to \mathsf{D}$ , a  $\mathcal{V}$ -enriched natural transformation  $\alpha$  from F to G consists of, for each object  $x \in \mathsf{C}$ , a morphism  $\alpha_x: 1 \to \mathsf{D}(Fx, Gx)$  in  $\mathcal{V}$  such that, for  $x, y \in \mathsf{C}$ , the

square



commutes.

**Example B.8** ([EK66, Prop. II.6.3]). Any monoidal natural transformation  $\eta$  between lax monoidal functor  $\Phi, \Psi: \mathcal{W} \to \mathcal{V}$  induces a  $\mathcal{V}$ -enriched natural transformation  $\eta_{\bullet}$  from  $\Phi_{\bullet}$  to  $\Psi_{\bullet}$ .

#### Cubical and simplicial Kan complexes

Our interest in enriched categories lies in studying cubical and simplicial categories, which we now define.

#### Definition B.9.

- A *cubical category* is a category enriched in (cSet, ⊗, □<sup>0</sup>). A *cubical functor* is a cSet-enriched functor between cubical categories.
- A simplicial category is a category enriched in (sSet, ×, Δ<sup>0</sup>). A simplicial functor is an sSet-enriched functor between simplicial categories.

Let  $Cat_{\Box}$  denote the category of cubical categories, and  $Cat_{\Delta}$  denote the category of simplicial categories.

In a cubical or simplicial category, we refer to the hom-object from x to y as the mapping space from x to y. Following Example B.2, we write cSet for the cubical category of cubical sets with mapping spaces given by hom<sub>L</sub>(x, y). Likewise, sSet denotes the simplicial category of simplicial sets. We write Kan<sub>□</sub> for the full cubical subcategory of cSet spanned by cubical Kan complexes, and Kan<sub>△</sub> for the full simplicial subcategory of simplicial Kan complexes.

Following Example B.6, the triangulation adjunction  $T: \mathsf{cSet} \rightleftharpoons \mathsf{sSet}: U$  induces an adjunction

$$T_{\bullet} \colon \mathsf{Cat}_{\Box} \rightleftarrows \mathsf{Cat}_{\Delta} : U_{\bullet}.$$

Simplicial and cubical categories which are locally Kan give examples of quasi-categories via their *homotopy-coherent nerve*. This construction is defined via pre-composition with a cosimplicial object (in  $Cat_{\Box}$  or  $Cat_{\Delta}$ , respectively), which we now define.

**Remark B.10.** Our primary reference for results regarding the cubical homotopy-coherent nerve is [KV20]. Although this reference is written using the categorical product on cubical sets as the base for enrichment (as opposed to the geometric product), any results used apply to both products.

For  $n \ge 0$ , define a cubical category  $\mathfrak{C}[n]$  as follows:

• the set of objects of  $\mathfrak{C}[n]$  is  $\{0, \ldots, n\}$ ;

• for  $i, j \in \{0, ..., n\}$ , the mapping space from i to j is defined by

$$\mathfrak{C}[n](i,j) := \begin{cases} \Box^{j-i-1} & i < j \\ \{ \mathrm{id}_i \} & i = j \\ \varnothing & i > j \end{cases}$$

• for  $i, j, k \in \{0, ..., n\}$ , the composition map when i < j < k is given by

$$\Box^{k-j-1} \otimes \Box^{j-i-1} \cong \Box^{k-i-2} \xrightarrow{\partial_{j,1}} \Box^{k-i-1}.$$

If i = j or j = k then composition is defined to be the natural isomorphisms

$$\mathfrak{C}[n](j,k) \otimes \{\mathrm{id}_j\} \xrightarrow{\cong} \mathfrak{C}[n](i,k) \xleftarrow{\cong} \{\mathrm{id}_j\} \otimes \mathfrak{C}[n](i,j),$$

respectively. Otherwise, the composition map is trivial as the domain becomes the empty cubical set.

We omit definitions of the simplicial operators of  $\mathfrak{C}[n]$ , which may be found in [KV20, Lem. 2.2]. Extension by colimits yields a functor  $\mathfrak{C}: \mathsf{sSet} \to \mathsf{Cat}_{\Box}$ . The *cubical homotopy-coherent nerve*  $N_{\Box}: \mathsf{Cat}_{\Box} \to \mathsf{sSet}$  is its right adjoint, which is defined by

$$(\mathbb{N}_{\Box}\mathsf{C})_n := \mathsf{Cat}_{\Box}(\mathfrak{C}[n],\mathsf{C}).$$

The simplicial homotopy-coherent nerve  $N_\Delta \colon \mathsf{Cat}_\Delta \to \mathsf{sSet}$  is defined by

$$(N_{\Delta}D)_n := Cat_{\Box}(T_{\bullet}\mathfrak{C}[n], D).$$

In particular, there is a natural isomorphism  $N_{\Delta} \cong N_{\Box} \circ U_{\bullet}$  ([KV20, Prop. 2.8]).

## Definition B.11.

- 1. A cubical category C is *locally Kan* if for all  $x, y \in ob C$ , the mapping space from x to y is a cubical Kan complex.
- 2. A simplicial category D is *locally Kan* if for all  $x, y \in ob C$ , the mapping space from x to y is a simplicial Kan complex.

### Proposition B.12.

- 1. If C is a locally Kan cubical category then  $N_{\Box}C$  is a quasicategory.
- 2. If D is a locally Kan simplicial category then  $N_{\Delta}D$  is a quasicategory.

Proof. Item (1) is [KV20, Thm. 2.6]. Item (2) is [Lur09, Prop. 1.1.5.10].

Locally Kan simplicial categories are the fibrant objects in the *Bergner model structure* on simplicial categories. The simplicial homotopy-coherent nerve is the right adjoint in a Quillen equivalence between the Bergner model structure and the Joyal model structure ([Lur09, Thm. 2.2.5.1]). We move towards defining an analogous model structure on cubical categories which is Quillen equivalent to these. This will require some auxiliary definitions and results.

The adjunction  $\pi_0: \mathsf{cSet} \rightleftharpoons \mathsf{Set} : \mathsf{Sk}_0$  induces an adjunction

$$(\pi_0)_{ullet} \colon \mathsf{Cat}_{\Box} \rightleftarrows \mathsf{Cat} : (\mathrm{Sk}_0)_{ullet}$$

We write Ho:  $\operatorname{Cat}_{\Box} \to \operatorname{Cat}$  for the functor  $(\pi_0)_{\bullet}$  and refer to Ho C as the homotopy category of C. Its right adjoint  $(\operatorname{Sk}_0)_{\bullet}$  is a full subcategory inclusion, hence any ordinary category may be viewed as a cubical category with discrete mapping spaces. For an ordinary category  $\mathcal{C}$ , we often write the cubical category  $(\operatorname{Sk}_0)_{\bullet}(\mathcal{C})$  as simply  $\mathcal{C}$ .

Proposition B.13. For a cubical category C, there is an equivalence of categories

$$\operatorname{Ho}(\mathsf{C}) \simeq \operatorname{Ho}(\mathsf{N}_{\Box}\mathsf{C})$$

natural in C.

Before proving Proposition B.13, we give a brief explicit description of the functor  $\mathfrak{C}$ : sSet  $\rightarrow$  Cat<sub> $\Box$ </sub>. Similar descriptions may be found in [RZ18, Rem. 5.2], and in the simplicial case in [Rie14, Thm. 16.4.1] and [DS11, Cor. 4.4] for the functor  $\mathfrak{C}_{\Delta}$ : sSet  $\rightarrow$  Cat<sub> $\Delta$ </sub>.

For a simplicial set X with 0-simplices x, y, the k-cubes of the mapping space  $\mathfrak{C}X(x, y)$  are finite tuples of pairs

$$((s_n, f_n), \ldots, (s_1, f_1))$$

where

• each  $s_i$  is an  $(m_i)$ -simplex of X for some  $m_i \ge 1$  such that

$$- s_1|_{\{0\}} = x \text{ and } s_n|_{\{m_n\}} = y; \text{ and} - \text{ for } i = 1, \dots, n-1, s_i|_{\{m_i\}} = s_{i+1}|_{\{0\}};$$

• each  $f_i$  is a  $(d_i)$ -cube of the mapping space  $\mathfrak{C}[m_i](0, m_i) \cong \Box^{m_i - 1}$  for some  $d_i$ , where  $d_1 + \cdots + d_n = k$ ,

subject to the equivalence relation generated by

• for  $i = 1, \ldots, n$  and  $j = 1, \ldots, m_i$ , the relation

$$\left((s_n, f_n), \ldots, (s_i\partial_j, f_i), \ldots, (s_1, f_1)\right) \sim \left((s_n, f_n), \ldots, (s_i, \mathfrak{C}(\partial_j)_{0, m_i - 1} \circ f_i), \ldots, (s_1, f_1)\right);$$

• for  $i = 1, \ldots, n$  and  $j = 1, \ldots, m_i - 1$ , the relation

$$((s_n, f_n), \dots, (s_i \sigma_j, f_i), \dots, (s_1, f_1)) \sim ((s_n, f_n), \dots, (s_i, \mathfrak{C}(\sigma_j)_{0,m_i+1} \circ f_i), \dots, (s_1, f_1));$$

and

• for  $i = 1, \ldots, n - 1$ , the relation

$$((s_n, f_n), \dots, (s_{i+1}, f_{i+1}), (s_i, f_i \sigma_{m_i+1}), \dots, (s_1, f_1)) \sim ((s_n, f_n), \dots, (s_{i+1}, f_{i+1} \sigma_1), (s_i, f_i), \dots, (s_1, f_1)).$$

If x = y then we identify the empty tuple with the 0-cube  $id_x \in \mathfrak{C}X(x, x)$ . Otherwise, we consider all tuples to be non-empty. Composition is given by concatenation of tuples, hence if each  $d_i = 0$  for all i (that is, each  $f_i$  is a 0-cube) then

$$[((s_n, f_n), \dots, (s_1, f_1))] = [(s_n, f_n)] \circ \dots \circ [(s_1, f_1)]$$

in  $\pi_0(\mathfrak{C}(X)(x,y))$ .

If X is the homotopy-coherent nerve of a cubical category C, an  $m_i$ -simplex  $s_i$  of  $X = \mathbb{N}_{\Box}\mathsf{C}$  is a cubical functor  $s_i \colon \mathfrak{C}[m_i] \to \mathsf{C}$ . With this, the counit map  $(\varepsilon_{\mathsf{C}})_{x,y} \colon \mathfrak{C}\mathbb{N}_{\Box}\mathsf{C}(x,y) \to \mathsf{C}(x,y)$  has an explicit formula

$$((s_n, f_n), \ldots, (s_1, f_1)) \mapsto (s_n f_n) \circ \cdots \circ (s_1 f_1),$$

where  $s_i f_i$  denotes the composite

$$\Box^{d_i} \xrightarrow{f_i} \mathfrak{C}[m_i](0,m_i) \xrightarrow{(s_i)_{0,m_i}} \mathsf{C}(s_i(0),s_i(m_i)).$$

Proof of Proposition B.13. The triangle of left adjoints



commutes (up to natural isomorphism) when restricted to representable presheaves, hence it commutes. Instantiating at  $N_{\Box}C$  gives an isomorphism

$$\operatorname{Ho}(\mathbb{N}_{\square}\mathsf{C})\cong\operatorname{Ho}(\mathfrak{C}\mathbb{N}_{\square}\mathsf{C})$$

natural in C. It suffices to show the counit  $\mathfrak{CN}_{\Box}C \to C$  induces a bijection on connected components in each mapping space, hence an equivalence of homotopy categories  $\operatorname{Ho}(\mathfrak{CN}_{\Box}C) \simeq \operatorname{Ho} C$ .

Fix objects  $x, y \in C$  and consider the cubical map

$$(\varepsilon_{\mathsf{C}})_{x,y} \colon \mathfrak{CN}_{\Box}\mathsf{C}(x,y) \to \mathsf{C}(x,y).$$

It is clear this map is surjective on 0-cubes, hence it is surjective on connected components. For injectivity, fix  $u, v \in (\mathfrak{CN}_{\Box}\mathsf{C}(x, y))_0$  such that  $[(\varepsilon_{\mathsf{C}})_{x,y}u] = [(\varepsilon_{\mathsf{C}})_{x,y}v]$  in  $\pi_0\mathsf{C}(x, y)$ . We write

$$u = ((s_m, f_m), \dots, (s_1, f_1))$$
  
$$v = ((s'_n, f'_n), \dots, (s'_1, f'_1)),$$

where each  $f_i$  is a 0-cube in  $\mathfrak{C}[m_i](0, m_i)$  and each  $f'_i$  is a 0-cube in  $\mathfrak{C}[n_i](0, n_i)$ . As the equivalence relation defining  $\pi_0$  is generated by 1-cubes, we may assume there exists a 1-cube  $t \in \mathsf{C}(x, y)_1$  from  $(s_m f_m) \circ \cdots \circ (s_1 f_1)$  to  $(s'_n f'_n) \circ \cdots \circ (s'_1 f'_1)$ . Note that t induces a cubical functor  $\overline{t}: \mathfrak{C}[2] \to \mathsf{C}$  whose assignment on objects is

$$\begin{aligned} t(0) &= x\\ \overline{t}(1) &= \overline{t}(2) = y \end{aligned}$$

and whose action on the mapping space

$$\mathfrak{C}[2](0,2) \to \mathsf{C}(x,y)$$

is given by  $t: \Box^1 \to \mathsf{C}(x, y)$ . Given a 2-simplex

Given a 2-simplex

$$a_0 \xrightarrow{s_{01}} s_{02} \xrightarrow{s_{12}} a_2$$

in a simplicial set X, the isomorphism  $\tau_1 X \cong (\pi_0)_{\bullet} \mathfrak{C}(X)$  induces an equality

$$[(s_{12},*)] \circ [(s_{01},*)] = [(s_{02},*)]$$

in  $\pi_0 \mathfrak{C}(X)(a_0, a_2)$ , where \* denotes the unique 0-cube of  $\mathfrak{C}[1](0, 1)$ . An induction argument shows that, given an *n*-simplex, writing  $s_{ij}$  for the restriction of this *n*-simplex to the 1-simplex  $i \leq j$  gives an equality

$$[(s_{(n-1)n}, *)] \circ \cdots \circ [(s_{01}, *)] = [(s_{0n}, *)]$$

in  $\pi_0 \mathfrak{C}(X)(a_0, a_n)$ .

The composable sequence of arrows  $(s_m f_m), \ldots, (s_1 f_1)$  gives a cubical functor  $[m] \to \mathsf{C}$ . The composite

 $\mathfrak{C}[m] \to [m] \to \mathsf{C}$ 

defines an *m*-simplex  $r \in (\mathbb{N}_{\Box} \mathbb{C})_m$  whose restriction to the 1-simplex  $0 \leq m$  is the composite  $(s_m f_m) \circ \cdots \circ (s_1 f_1)$ . Analogously, the composable sequence  $(s'_n f'_n), \ldots, (s'_1, f'_1)$  defines an *n*-simplex  $r' \in (\mathbb{N}_{\Box} \mathbb{C})_n$  whose restriction to the 1-simplex  $0 \leq n$  is the composite  $(s'_n f'_1) \circ \cdots \circ (s'_1 f'_1)$ . The tuple  $(t, \varphi)$  (where  $\varphi$  denotes the unique non-degenerate 1-cube in  $\mathfrak{C}[2](0,2)$ ) defines a 1-cube from  $(r|_{0\leq m},*)$  to  $(r'|_{0\leq n},*)$ . Therefore,

$$[((s_m, f_m), \dots, (s_1, f_1))] = [(s_m, f_m)] \circ \dots \circ [(s_1, f_1)]$$
  
=  $[(r|_{0 \le m}, *)]$   
=  $[(r'|_{0 \le n}, *)]$   
=  $[(s'_n, f'_n)] \circ \dots \circ [(s'_1, f'_1)]$   
=  $[((s'_n, f'_n), \dots, (s'_1, f'_1))].$ 

**Remark B.14.** This is a cubical analogue of the result stated in [Lur09, Warn. 1.2.3.3]. Here, we are able to prove it without the additional assumption that C is locally Kan.

**Corollary B.15.** Given an arrow  $f: [1] \to C$  in a locally Kan cubical category C, the following are equivalent:

- 1. f becomes an isomorphism in HoC;
- 2. f lifts to a map  $\mathfrak{C}(E^1) \to \mathsf{C}$ ;
- 3. the adjoint transpose  $\widetilde{f} \colon \Delta^1 \to N_{\Box} \mathsf{C}$  of f lifts to a map  $E^1 \to N_{\Box} \mathsf{C}$ .

*Proof.* The homotopy-coherent nerve of C is a quasicategory by Proposition B.12. Thus, we have implications  $(1) \implies (3) \implies (2) \implies (1)$ , where the first implication follows from Proposition B.13.

Define a functor  $\Sigma_{\Box} : \mathsf{cSet} \to \mathsf{Cat}_{\Box}$  which sends a cubical set X to the cubical category  $\Sigma_{\Box} X$  with two objects 0, 1 and whose mapping spaces are

$$\begin{split} \Sigma_{\Box} X(0,1) &:= X & \Sigma_{\Box} X(1,0) := \varnothing \\ \Sigma_{\Box} X(0,0) &:= \{ \mathrm{id}_0 \} & \Sigma_{\Box} X(1,1) := \{ \mathrm{id}_1 \}. \end{split}$$

**Theorem B.16.** The category of cubical categories carries a model structure where:

• cofibrations are the saturation of the set

$$\{ \varnothing \hookrightarrow [0] \} \cup \{ \Sigma_{\Box} \partial \Box^n \hookrightarrow \Sigma_{\Box} \Box^n \mid n \ge 0 \};$$

• fibrant objects are locally Kan cubical categories; fibrations between fibrant objects are maps with the right lifting property against the set

$$\{[0] \hookrightarrow \mathfrak{C}(E^1)\} \cup \{\Sigma_{\Box} \sqcap_{i,\varepsilon}^n \hookrightarrow \Sigma_{\Box} \square^n \mid n \ge 1, \ i = 1, \dots, n \ \varepsilon = 0, 1\};$$

- weak equivalences are DK-equivalences, i.e. cubical functors  $F: \mathsf{C} \to \mathsf{D}$  such that
  - 1. Ho  $F: \operatorname{Ho} \mathsf{C} \to \operatorname{Ho} \mathsf{D}$  is an equivalence of categories; and
  - 2. for all  $x, y \in C$ , the cubical map

$$\mathsf{F}_{x,y} \colon \mathsf{C}(x,y) \to \mathsf{D}(x,y)$$

is a weak equivalence in the Grothendieck model structure.

Moreover, the triangulation adjunction induces a Quillen equivalence

$$T_{ullet}: \mathsf{Cat}_{\Box} \rightleftarrows \mathsf{Cat}_{\Delta}: U_{ullet}$$

with the Bergner model structure on simplicial categories.

*Proof.* For existence of the model structure as well as the description of the cofibrations and weak equivalences, we apply [Lur09, Prop. A.3.2.4]. The Quillen equivalence follows from [Lur09, Rem. A.3.2.6].

For the description of fibrant objects, we apply [Sta14, Prop. 2.3], where I is the set of generating cofibrations in the theorem statement and J is the set

$$J = \{[0] \hookrightarrow \mathfrak{C}(E^1)\} \cup \{\Sigma_{\Box} \sqcap_{i,\varepsilon}^n \hookrightarrow \Sigma_{\Box} \square^n \mid n \ge 1, \ i = 1, \dots, n, \ \varepsilon = 0, 1\}.$$

One verifies that a locally Kan cubical category is exactly a cubical category with the right lifting property with respect to J. We show that a weak equivalence between locally Kan cubical categories that has the right lifting property with respect to J is an acyclic fibration; the remaining hypotheses of [Sta14, Prop. 2.3] are immediate.

To this end, fix such a map  $F: C \xrightarrow{\sim} D$ . It suffices to show this map has the right lifting property with respect to the set of generating cofibrations

$$\{\emptyset \hookrightarrow [0]\} \cup \{\Sigma_{\Box} \partial \Box^n \hookrightarrow \Sigma_{\Box} \Box^n \mid n \ge 0\}$$

The maps  $\Sigma_{\Box} \partial \Box^n \hookrightarrow \Sigma_{\Box} \Box^n$  are immediate, so it remains to show F has the right lifting property against the map  $\emptyset \hookrightarrow [0]$ . Given a square



there exists  $x \in C$  and an isomorphism  $f: E[1] \to \text{Ho} D$  from Ho F(x) to y. As D is locally Kan, applying Corollary B.15 gives a cubical functor  $\overline{f}: \mathfrak{C}(E^1) \to D$  such that  $\text{Ho} \overline{f} = f$ . The square

$$\begin{bmatrix} 0 \end{bmatrix} \xrightarrow{x} \mathsf{C} \\ \downarrow & \downarrow^{F} \\ \mathfrak{C}(E^{1}) \xrightarrow{\overline{f}} \mathsf{D} \end{bmatrix}$$

admits a lift  $g: \mathfrak{C}(E^1) \to \mathsf{C}$  by assumption. The object g(1) is a lift of the starting square.

Proposition B.17. The adjunction

$$\mathfrak{C} \colon \mathsf{sSet} \rightleftarrows \mathsf{Cat}_{\Box} : \mathrm{N}_{\Box}$$

is a Quillen equivalence.

*Proof.* We first show that  $\mathfrak{C} \dashv N_{\Box}$  is a Quillen adjunction. By [JT07, Prop. 7.15], it suffices to show  $\mathfrak{C}: \mathsf{sSet} \to \mathsf{Cat}_{\Box}$  sends monomorphisms to cofibrations and  $N_{\Box}: \mathsf{Cat}_{\Box} \to \mathsf{sSet}$  sends fibrations between fibrant objects to fibrations.

As  $\mathfrak{C}$  is a left adjoint, it suffices to show it takes boundary inclusions to cofibrations. This follows as the square
is a pushout for all  $n \ge 1$ . The inclusion  $\mathfrak{C}(\partial \Delta^0) \hookrightarrow \mathfrak{C}[0]$  is a cofibration by definition.

The functor  $N_{\Box}$  sends locally Kan quasicategories to quasicategories by Proposition B.12. One verifies that it sends fibrations between fibrant objects to inner isofibrations between quasicategories, thus  $N_{\Box}$  preserves fibrations between fibrant objects.

In the commutative triangle of adjunctions



the top adjunction is a Quillen equivalence by [Lur09, Thm. 2.2.5.1]. The bottom right adjunction is a Quillen equivalence by Theorem B.16. Thus,  $\mathfrak{C} \dashv N_{\Box}$  is a Quillen equivalence by 2-out-of-3.

Our main goal is to contruct an equivalence of quasicategories

$$N_{\Box} Kan_{\Box} \rightarrow N_{\Delta} Kan_{\Delta}$$
.

To do so, we use the following fact about cubical natural transformations.

**Proposition B.18.** Let  $F, G: C \to D$  be cubical functors. Any natural transformation  $\alpha$  from F to G induces a natural transformation

$$\alpha \colon \mathbb{N}_{\Box}\mathsf{C} \times \Delta^1 \to \mathsf{D}$$

from  $N_{\Box}F$  to  $N_{\Box}G$ .

*Proof.* We construct a natural transformation

 $\alpha\colon \mathrm{N}_{\Box}\mathsf{C}\times\Delta^1\to\mathrm{N}_{\Box}\mathsf{D}$ 

from  $N_{\Box}F$  to  $N_{\Box}G$  by skeletal induction on  $N_{\Box}C$ , with the property that the adjoint transpose factors as

$$\mathfrak{C}(\mathbf{N}_{\Box}\mathsf{C}\times\Delta^{1})\xrightarrow{\mathfrak{C}(\alpha)}\mathfrak{C}\times[1] \xrightarrow{\mathfrak{C}(\alpha)}\mathfrak{C}\times[1]$$

We define  $\alpha_0$ : Sk<sub>0</sub> N<sub> $\square$ </sub>C ×  $\Delta^1 \rightarrow$  N<sub> $\square$ </sub>D on a 1-simplex ( $x, 0 \leq 1$ ) to be the 1-simplex  $\alpha_x$ :  $Fx \rightarrow Gx$  of N<sub> $\square$ </sub>D. The adjoint transpose of this map factors as in the above triangle as the left map is identity-on-objects.

Fix  $\alpha_n$ :  $\operatorname{Sk}_n \operatorname{N}_{\Box} \mathsf{C} \times \Delta^1 \to \operatorname{N}_{\Box} \mathsf{D}$ . To construct  $\alpha_{n+1}$ , we fix a non-degenerate (n+1)-simplex u of  $\operatorname{N}_{\Box} \mathsf{C}$  and give a lift as in the diagram

By adjointness, this is equivalent to a lift as in the diagram

By the inductive hypothesis, the top map factors as

Thus, it suffices to give a lift as in

$$\mathfrak{C}(\partial \Delta^{n+1}) \times [1] \xrightarrow{\beta_n \mathfrak{C}(u)} \mathsf{D}$$

$$\int$$

$$\mathfrak{C}[n+1] \times [1]$$

since we may concatenate with a naturality square on the left to obtain a lift

Let  $\varphi$  denote the map  $\beta_n \mathfrak{C}(u)$ . The inclusion  $\mathfrak{C}(\partial \Delta^{n+1}) \hookrightarrow \mathfrak{C}[n+1]$  is a bijection on objects. As mapping spaces commute with products, one verifies that it suffices to construct lifts as in

such that

$$\varphi|_{(n+1,0)\leq (n+1,1)} \circ v_1 = v_3 = v_2 \circ \varphi|_{(0,0)\leq (0,1)}.$$

For i = 0, ..., n + 1, recall that  $\varphi|_{(i,0) \leq (i,1)} = \alpha_{\mathfrak{C}(u)(i)}$  by construction. Setting  $v_1 = F_{\mathfrak{C}(u)(0),\mathfrak{C}(u)(n+1)}u$ and  $v_2 = G_{\mathfrak{C}(u)(0),\mathfrak{C}(u)(n+1)}u$ , the required equality follows from naturality.

Given a product-preserving fibrant replacement functor on simplicial sets (e.g.  $Ex^{\infty}$ ), we write

$$T^{\mathrm{f}} \colon \mathsf{cSet} o \mathsf{Kan}_\Delta$$

for the composition of the triangulation functor with such a fibrant replacement.

For any cubical category  $\mathsf{C},$  there is a cubical functor

$$(\widetilde{\eta}_{\bullet})_{\mathsf{C}} \colon \mathsf{C} \to (UT^{\dagger})_{\bullet}\mathsf{C}$$

which is identity-on-objects and whose action on mapping spaces

$$\widetilde{\eta}_{\mathsf{C}(x,y)} \colon \mathsf{C}(x,y) \to UT^{\mathrm{f}}(\mathsf{C}(x,y))$$

is defined to be the derived unit of the triangulation adjunction. Moreover, this functor is natural in C (cf. Example B.8). Analogously, for any simplicial category D, there is a simplicial functor

$$(\widetilde{\varepsilon}_{\bullet})_{\mathsf{D}} \colon (TU)_{\bullet}\mathsf{D} \to \mathsf{D}$$

natural in D.

## Proposition B.19.

1. For any cubical category C, the cubical functor

$$(\widetilde{\eta}_{\bullet})_{\mathsf{C}} \colon \mathsf{C} \to (UT^{\mathrm{f}})_{\bullet}\mathsf{C}$$

is a DK-equivalence.

2. For a locally Kan simplicial category D, the simplicial functor

$$(\widetilde{\varepsilon}_{\bullet})_{\mathsf{D}} \colon (TU)_{\bullet}\mathsf{D} \to \mathsf{D}$$

is a DK-equivalence.

*Proof.* Follows since  $(T \dashv U)$  is a Quillen equivalence.

Following Example B.5, the functor  $T^{f}: \mathsf{cSet} \to \mathsf{Kan}_{\Delta}$  is lax monoidal, hence induces a simplicial functor

$$\overline{T^{\mathrm{f}}} \colon (T^{\mathrm{f}})_{\bullet} \mathsf{cSet} \to \mathsf{Kan}_{\Delta}.$$

Applying  $U_{\bullet}$  gives a cubical functor  $U_{\bullet}\overline{T^{f}}$ :  $(UT^{f})_{\bullet}cSet \rightarrow U_{\bullet}Kan_{\Delta}$ . We further apply  $N_{\Box}$ :  $Cat_{\Box} \rightarrow sSet$  to the composite

$$\mathsf{Kan}_{\Box} \hookrightarrow \mathsf{cSet} \xrightarrow{(\widetilde{\eta}_{\bullet})_{\mathsf{cSet}}} (UT^{\mathrm{f}})_{\bullet} \mathsf{cSet} \xrightarrow{U_{\bullet}(T^{\mathrm{f}})} U_{\bullet} \mathsf{Kan}_{\Delta}$$

which gives a map  $N_{\Box}Kan_{\Box} \rightarrow N_{\Delta}Kan_{\Delta}$  (as  $N_{\Delta} \cong N_{\Box} \circ U_{\bullet}$ ). We refer to this map as the *canonical* equivalence of cubical Kan complexes and simplicial Kan complexes.

## Theorem B.20.

1. The canonical equivalence

$$\mathrm{N}_{\Box}\mathsf{Kan}_{\Box}\to\mathrm{N}_{\Delta}\mathsf{Kan}_{\Delta}$$

is a categorical equivalence.

2. For a locally Kan cubical category C and  $X \in C$ , the composite

$$\mathrm{N}_{\Box}\mathsf{C} \xrightarrow{\mathrm{N}_{\Box}(\mathsf{C}(X,-))} \mathrm{N}_{\Box}\mathsf{Kan}_{\Box} \to \mathrm{N}_{\Delta}\mathsf{Kan}_{\Delta}$$

is a model for the representable functor represented by X.

*Proof.* Let F denote the composite

$$\mathsf{Kan}_{\Box} \hookrightarrow \mathsf{cSet} \xrightarrow{\widetilde{\eta}_{\bullet}} (UT^{\mathrm{f}})_{\bullet} \mathsf{cSet} \xrightarrow{U_{\bullet}(\overline{T^{\mathrm{f}}})} U_{\bullet}\mathsf{Kan}_{\Delta},$$

so that  $N_{\Box}F: N_{\Box}Kan_{\Box} \rightarrow N_{\Delta}Kan_{\Delta}$  is the canonical equivalence. For (1), the lax monoidal functor  $U: sSet \rightarrow cSet$  induces a cubical functor

$$\overline{U}: U_{\bullet}\mathsf{Kan}_{\Delta} \to \mathsf{Kan}_{\Box}.$$

We show  $N_{\Box}\overline{U}$  is an inverse of  $N_{\Box}F$  up to  $E^1$ -homotopy.

The derived unit of the triangulation adjunction defines a natural transformation from  $\mathrm{id}_{\mathsf{Kan}_{\square}}$  to  $\overline{U} \circ F$ . Applying  $N_{\square}$  gives a natural transformation

$$N_{\Box}Kan_{\Box} \times \Delta^1 \to N_{\Box}Kan_{\Box}$$

from  $\operatorname{id}_{N_{\Box}\mathsf{Kan}_{\Box}}$  to  $N_{\Box}\overline{U} \circ N_{\Box}F$ . Each component of this natural transformation is a weak equivalence between Kan complexes, hence invertible. Thus, this natural transformation is an  $E^1$ -homotopy. An analogous argument shows the image under  $\operatorname{Ex}^{\infty}$  of the derived counit defines an  $E^1$ -homotopy

$$N_{\Box}U_{\bullet}Kan_{\Delta} \times E^{1} \rightarrow N_{\Box}U_{\bullet}Kan_{\Delta}$$

from  $\operatorname{id}_{\mathbb{N}_{\square}U_{\bullet}\mathsf{Kan}_{\Delta}}$  to  $\mathbb{N}_{\square}F \circ \mathbb{N}_{\square}\overline{U}$ .

For  $(\overline{2})$ , the triangle of simplicial functors



commutes by definition of  $\overline{T^{f}}$ . Applying N<sub> $\Box$ </sub> yields the  $E^{1}$ -homotopy commutative diagram



where the left square is given by naturality of  $\tilde{\eta}_{\bullet}$  and Proposition B.18. The left horizontal maps are categorical equivalences by Proposition B.19. As the top right map is a model for the representable functor at  $X \in N_{\Delta}T^{f}_{\bullet}C$ , the top composite is a model for the representable functor at  $X \in N_{\Box}C$ . The composite  $N_{\Box}Kan_{\Box} \rightarrow N_{\Delta}Kan_{\Delta}$  along the left and bottom is the canonical equivalence.

**Remark B.21.** We do not state a dual version of item (2) in Theorem B.20 for the contravariant representable

$$(\mathrm{N}_{\Box}\mathsf{C})^{\mathrm{op}} \xrightarrow{\mathrm{N}_{\Box}(\mathsf{C}(-,X))} \mathrm{N}_{\Box}\mathsf{Kan}_{\Box} \to \mathrm{N}_{\Delta}\mathsf{Kan}_{\Delta}.$$

This is because the geometric product of cubical sets is not symmetric, hence the composition map of C does not dualize to a composition in (what would be)  $C^{\text{op}}$ . As there is no opposite category, there cannot be a contravariant (representable) functor.

## References

[Atk74] R. H. Atkin, An algebra for patterns on a complex. I, Internat. J. Man-Mach. Stud. 6 (1974), 285–307.

[Atk76] R. H. Atkin, An algebra for patterns on a complex. II, Internat. J. Man-Mach. Stud. 8 (1976), no. 5, 483-498.

- [BBdLL06] E. Babson, H. Barcelo, M. de Longueville, and R. Laubenbacher, Homotopy theory of graphs, J. Algebraic Combin. 24 (2006), no. 1, 31–44.
- [BCW14] H. Barcelo, V. Capraro, and J. A. White, Discrete homology theory for metric spaces, Bull. Lond. Math. Soc. 46 (2014), no. 5, 889–905.
- [BKLW01] H. Barcelo, X. Kramer, R. Laubenbacher, and C. Weaver, Foundations of a connectivity theory for simplicial complexes, Adv. Appl. Math. 26 (2001), no. 1, 97–128.
  - [BL05] H. Barcelo and R. Laubenbacher, Perspectives on A-homotopy theory and its applications, Discrete Math. 298 (2005), no. 1-3, 39–61.

- [BM19] A. J. Blumberg and M. A. Mandell, E<sub>2</sub> structures and derived Koszul duality in string topology, Algebr. Geom. Topol. 19 (2019), no. 1, 239–279.
- [Cis06] D.-C. Cisinski, Les préfaisceaux comme modèles des types d'homotopie, Astérisque 308 (2006), xxiv+390.
- [Cis14] D.-C. Cisinski, Univalent universes for elegant models of homotopy types, 2014.
- [Cis19] D.-C. Cisinski, Higher categories and homotopical algebra, Cambridge Studies in Advanced Mathematics, vol. 180, Cambridge University Press, Cambridge, 2019.
- [CK22a] D. Carranza and K. Kapulkin, Cubical setting for discrete homotopy theory, revisited, 2022. preprint.
- [CK22b] D. Carranza and K. Kapulkin, Homotopy groups of cubical sets, 2022. preprint.
- [CKW23] D. Carranza, K. Kapulkin, and L. Z. Wong, *Diagonal lemma for presheaves on Eilenberg–Zilber categories*, 2023. preprint.
- [DKLS20] B. Doherty, K. Kapulkin, Z. Lindsey, and C. Sattler, *Cubical models of*  $(\infty, 1)$ -categories, 2020. to appear in Mem. Amer. Math. Soc.
  - [DS11] D. Dugger and D. I. Spivak, Rigidification of quasi-categories, Algebr. Geom. Topol. 11 (2011), no. 1, 225–261.
  - [EK66] S. Eilenberg and G. M. Kelly, Closed categories, Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965), 1966, pp. 421–562.
  - [GJ99] P. G. Goerss and J. F. Jardine, Simplicial homotopy theory, Progress in Mathematics, vol. 174, Birkhäuser Verlag, Basel, 1999.
  - [GZ67] P. Gabriel and M. Zisman, Calculus of fractions and homotopy theory, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 35, Springer-Verlag New York, Inc., New York, 1967.
  - [Hin16] V. Hinich, Dwyer-Kan localization revisited, Homology Homotopy Appl. 18 (2016), no. 1, 27–48.
  - [Jar06] J. F. Jardine, Categorical homotopy theory, Homology Homotopy Appl. 8 (2006), no. 1, 71–144.
  - [Joy02] A. Joyal, Quasi-categories and Kan complexes, J. Pure Appl. Algebra 175 (2002), no. 1-3, 207–222. Special volume celebrating the 70th birthday of Professor Max Kelly.
  - [Joy09] A. Joyal, *The theory of quasi-categories and its applications*, 2009. Vol. II of course notes from Simplicial Methods in Higher Categories, Centra de Recerca Matemàtica, Barcelona, 2008.
  - [JT07] A. Joyal and M. Tierney, Quasi-categories vs Segal spaces, Categories in algebra, geometry and mathematical physics, 2007, pp. 277–326.
  - [Kan55] D. M. Kan, Abstract homotopy. I, Proc. Nat. Acad. Sci. U.S.A. 41 (1955), 1092–1096.
  - [Kan57] D. M. Kan, On c. s. s. complexes, Amer. J. Math. 79 (1957), 449-476.
  - [KL98] X. H. Kramer and R. C. Laubenbacher, Combinatorial homotopy of simplicial complexes and complex information systems, Applications of computational algebraic geometry (San Diego, CA, 1997), 1998, pp. 91–118.
  - [KV20] K. Kapulkin and V. Voevodsky, A cubical approach to straightening, J. Topol. 13 (2020), no. 4, 1682–1700.
- [LTW79] D. M. Latch, R. W. Thomason, and W. S. Wilson, Simplicial sets from categories, Math. Z. 164 (1979), no. 3, 195–214.
- [Lur09] J. Lurie, Higher topos theory, Annals of Mathematics Studies, vol. 170, Princeton University Press, Princeton, NJ, 2009.
- [Mar83] H. R. Margolis, Spectra and the Steenrod algebra, North-Holland Mathematical Library, vol. 29, North-Holland Publishing Co., Amsterdam, 1983. Modules over the Steenrod algebra and the stable homotopy category.
- [MO15] L. Meier and V. Ozornova, Fibrancy of partial model categories, Homology Homotopy Appl. **17** (2015), no. 2, 53–80.
- [MZ19] F. Mémoli and L. Zhou, Persistent homotopy groups of metric spaces, 2019.
- [Rie14] E. Riehl, *Categorical homotopy theory*, New Mathematical Monographs, vol. 24, Cambridge University Press, Cambridge, 2014.
- [RZ18] M. Rivera and M. Zeinalian, Cubical rigidification, the cobar construction and the based loop space, Algebr. Geom. Topol. 18 (2018), no. 7, 3789–3820.
- [Sta14] A. E. Stanculescu, Constructing model categories with prescribed fibrant objects, Theory Appl. Categ. 29 (2014), No. 23, 635–653.
- [Szu17] K. Szumiło, Frames in cofibration categories, J. Homotopy Relat. Struct. 12 (2017), no. 3, 577-616.