## Multiplicative topological indices: Analytical properties and application to random networks

R. Aguilar-Sánchez<sup>1</sup>, J. A. Méndez-Bermúdez<sup>2†</sup>, José M. Rodríguez<sup>3†</sup> and José M. Sigarreta<sup>4\*†</sup>

<sup>1</sup>Facultad de Ciencias Químicas, Benemérita Universidad Autónoma de Puebla, Puebla, 72570, Puebla, Mexico.

<sup>2</sup>Instituto de Física, Benemérita Universidad Autónoma de

Puebla, Apartado Postal J-48, Puebla, 72570, Puebla, Mexico.

<sup>3</sup>Departamento de Matemáticas, Universidad Carlos III de Madrid, Avenida de la Universidad 30, Leganés, 28911, Madrid, Spain.

<sup>4</sup>Facultad de Matemáticas, Universidad Autónoma de Guerrero, Carlos E. Adame No.54 Col. Garita, Acapulco, 610101, Guerrero, Mexico.

> \*Corresponding author(s). E-mail(s): josemariasigarretaalmira@hotmail.com; Contributing authors: ras747698@gmail.com; jmendezb@ifuap.buap.mx; jomaro@math.uc3m.es; †These authors contributed equally to this work.

#### Abstract

We make use of multiplicative degree-based topological indices  $X_{\Pi}(G)$  to perform a detailed analytical and statistical study of random networks G = (V(G), E(G)). We consider two classes of indices:  $X_{\Pi}(G) = \prod_{u \in V(G)} F_V(d_u)$  and  $X_{\Pi}(G) = \prod_{uv \in E(G)} F_E(d_u, d_v)$ , where uv denotes the edge of G connecting the vertices u and v,  $d_u$  is the degree of the vertex u, and  $F_V(x)$  and  $F_E(x, y)$  are functions of the vertex degrees. Specifically, we find analytical inequalities involving these multiplicative indices. Also, we apply  $X_{\Pi}(G)$  on three models of random networks: Erdös-Rényi networks, random geometric graphs, and

bipartite random networks. We show that  $\langle \ln X_{\Pi}(G) \rangle$ , normalized to the order of the network, scale with the corresponding average degree; here  $\langle \cdot \rangle$  denotes the average over an ensemble of random networks.

 ${\bf Keywords:}$  random networks, degree–based topological indices, scaling approach

### 1 Introduction

We can identify two types of graph invariants which are currently been studied in chemical graph theory, namely

$$X_{\Sigma}(G) = X_{\Sigma, F_{V}}(G) = \sum_{u \in V(G)} F_{V}(d_{u}) \quad \text{or}$$
  
$$X_{\Sigma}(G) = X_{\Sigma, F_{E}}(G) = \sum_{uv \in E(G)} F_{E}(d_{u}, d_{v})$$
(1)

and

$$X_{\Pi}(G) = X_{\Pi, F_V}(G) = \prod_{u \in V(G)} F_V(d_u) \quad \text{or}$$
  
$$X_{\Pi}(G) = X_{\Pi, F_E}(G) = \prod_{uv \in E(G)} F_E(d_u, d_v).$$
(2)

Here uv denotes the edge of the graph G = (V(G), E(G)) connecting the vertices u and v,  $d_u$  is the degree of the vertex u, and  $F_V(x)$  and  $F_V(x, y)$  are appropriate chosen functions, see e.g. [1]. While both  $X_{\Sigma}(G)$  and  $X_{\Pi}(G)$  are referred as topological indices in the literature, to make a distinction between them, here we name  $X_{\Pi}(G)$  as multiplicative topological indices (MTIs).

Among the vast amount of topological indices of the form given by Eq. (1) we can refer to some prominent examples: the first and second Zagreb indices [2]

$$M_1(G) = \sum_{u \in V(G)} d_u^2 = \sum_{uv \in E(G)} d_u + d_v$$
(3)

and

$$M_2(G) = \sum_{uv \in E(G)} d_u d_v, \tag{4}$$

respectively, the Randić connectivity index [3]

$$R(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}},\tag{5}$$

the harmonic index [4]

$$H(G) = \sum_{uv \in E(G)} \frac{2}{d_u + d_v},\tag{6}$$

the sum-connectivity index [5]

$$\chi(G) = \sum_{uv \in E(G)} \frac{1}{\sqrt{d_u + d_v}},\tag{7}$$

and the inverse degree index [4]

$$ID(G) = \sum_{u \in V(G)} \frac{1}{d_u} = \sum_{uv \in E(G)} \left( \frac{1}{d_u^2} + \frac{1}{d_v^2} \right).$$
(8)

In fact, within a statistical approach on random networks, it has been recently shown that the average values of indices of the type  $X_{\Sigma}(G)$ , normalized to the order of the network n, scale with the average degree  $\langle k \rangle$ ; see e.g. [6–8]. That is,  $\langle X_{\Sigma}(G) \rangle / n$  is a function of  $\langle k \rangle$  only. Moreover, it was also found that  $\langle X_{\Sigma}(G) \rangle$ , for indices like R(G) and H(G), is highly correlated with the Shannon entropy of the eigenvectors of the adjacency matrix of random networks [9]. This is a notable result because it puts forward the application of topological indices beyond mathematical chemistry. Specifically, given que equivalence of the Hamiltonian of a tight-binding network (in the proper setup) and the corresponding network adjacency matrix, either  $\langle R(G) \rangle$ or  $\langle H(G) \rangle$  could be used to determine the eigenvector localization and delocalization regimes which in turn determine the insulator and metallic regimes of quantum transport.

Therefore, motivated by their potential applications, in this work we study MTIs, see Eq. (2), on random networks.

Indeed, several MTIs have been already deeply analyzed in the literature (see e.g. the work of V. R. Kulli), among the most relevant we have to mention the Narumi-Katayama index [10]

$$NK(G) = \prod_{u \in V(G)} d_u, \tag{9}$$

and multiplicative versions of the Zagreb indices [11]:

$$\Pi_1(G) = \prod_{u \in V(G)} d_u^2,\tag{10}$$

$$\Pi_2(G) = \prod_{uv \in E(G)} d_u d_v, \tag{11}$$

and

$$\Pi_1^*(G) = \prod_{uv \in E(G)} d_u + d_v.$$
(12)

In addition to the MTIs of Eqs. (9-12), we also consider multiplicative versions of the indices in Eqs. (5-8): the multiplicative Randić connectivity index

$$R_{\Pi}(G) = \prod_{uv \in E(G)} \frac{1}{\sqrt{d_u d_v}},\tag{13}$$

the multiplicative harmonic index

$$H_{\Pi}(G) = \prod_{uv \in E(G)} \frac{2}{d_u + d_v},$$
(14)

the multiplicative sum-connectivity index

$$\chi_{\Pi}(G) = \prod_{uv \in E(G)} \frac{1}{\sqrt{d_u + d_v}},\tag{15}$$

and the multiplicative inverse degree index

$$ID_{\Pi}(G) = \prod_{uv \in E(G)} \left( \frac{1}{d_u^2} + \frac{1}{d_v^2} \right).$$
(16)

Below we apply the MTIs of Eqs. (9-16) on three models of random networks: Erdös-Rényi (ER) networks, random geometric (RG) graphs, and bipartite random (BR) networks. ER networks [12–14]  $G_{\text{ER}}(n, p)$  are formed by n vertices connected independently with probability  $p \in [0, 1]$ . While RG graphs [15, 16]  $G_{\text{RG}}(n, r)$  consist of n vertices uniformly and independently distributed on the unit square, where two vertices are connected by an edge if their Euclidean distance is less or equal than the connection radius  $r \in [0, \sqrt{2}]$ . In addition we examine BR networks  $G_{\text{BR}}(n_1, n_2, p)$  composed by two disjoint sets, set 1 and set 2, with  $n_1$  and  $n_2$  vertices each such that there are no adjacent vertices within the same set, being  $n = n_1 + n_2$  the total number of vertices in the bipartite network. The vertices of the two sets are connected randomly with probability  $p \in [0, 1]$ .

The purpose of this work is threefold. First, from an analytical viewpoint, we find several inequalities that relate multiplicative indices to their additive versions; second, we want to establish the statistical analysis of MTIs as a generic tool for studying average properties of random networks; and third, we perform for the first time (to our knowledge), a scaling study of MTIs on random networks.

## 2 Some inequalities involving multiplicative topological indices

We obtain in this section several analytical inequalities involving general multiplicative topological indices  $X_{\Pi}$ . There are many papers studying particular multiplicative topological indices (see, e.g., [19], [22] and the references therein).

The following equalities are direct:

$$X_{\Pi,F_{V}}(G) = \prod_{u \in V(G)} F_{V}(d_{u}) = e^{\sum_{u \in V(G)} \log F_{V}(d_{u})} = e^{X_{\Sigma,\log F_{V}}(G)},$$
  
$$X_{\Pi,F_{E}}(G) = \prod_{uv \in E(G)} F_{E}(d_{u},d_{v}) = e^{\sum_{uv \in E(G)} \log F_{E}(d_{u},d_{v})} = e^{X_{\Sigma,\log F_{E}}(G)}.$$

Since the geometric mean is at most the arithmetic mean (by Jensen's inequality), we have the following inequalities.

**Proposition 1** Let G be a graph with n vertices and m edges. Then

$$X_{\Pi, F_V}(G)^{1/n} \le \frac{1}{n} X_{\Sigma, F_V}(G), \qquad X_{\Pi, F_E}(G)^{1/m} \le \frac{1}{m} X_{\Sigma, F_E}(G)$$

In [18] appears the following Jensen-type inequality.

**Theorem 2** Let  $\mu$  be a probability measure on the space X and  $a \leq b$  real constants. If  $f: X \to [a,b]$  is a measurable function and  $\varphi$  is a convex function on [a,b], then f and  $\varphi \circ f$  are  $\mu$ -integrable functions and

$$\varphi\left(a+b-\int_X f\,d\mu\right) \le \varphi(a)+\varphi(b)-\int_X \varphi\circ f\,d\mu.$$

Theorem 2 allows to find the following converse version of Proposition 1.

**Theorem 3** Let G be a graph with n vertices and m edges,  $a_V \leq b_V$ ,  $a_E \leq b_E$  be real constants and  $F_V$ ,  $F_E$  be functions satisfying

 $a_V \leq F_V(d_u) \leq b_V \quad \forall u \in V(G), \qquad a_E \leq F_E(d_u, d_v) \leq b_E \quad \forall uv \in E(G).$ 

Then

$$\frac{1}{n} X_{\Sigma, F_V}(G) \le e^{a_V} + e^{b_V} - \frac{e^{a_V + b_V}}{X_{\Pi, F_V}(G)^{1/n}},$$
  
$$\frac{1}{m} X_{\Sigma, F_E}(G) \le e^{a_E} + e^{b_E} - \frac{e^{a_E + b_E}}{X_{\Pi, F_E}(G)^{1/m}}.$$

5

*Proof* Since  $\varphi(x) = e^x$  is a convex function on  $\mathbb{R}$ , Theorem 2 gives

$$e^{a_V + b_V - \frac{1}{n} \sum_{u \in V(G)} \log F_V(d_u)} \le e^{a_V} + e^{b_V} - \frac{1}{n} \sum_{u \in V(G)} e^{\log F_V(d_u)},$$

$$e^{a_V + b_V} e^{-\log(\prod_{u \in V(G)} F_V(d_u))^{1/n}} \le e^{a_V} + e^{b_V} - \frac{1}{n} \sum_{u \in V(G)} F_V(d_u),$$

$$\frac{e^{a_V + b_V}}{X_{\Pi, F_V}(G)^{1/n}} \le e^{a_V} + e^{b_V} - \frac{1}{n} X_{\Sigma, F_V}(G).$$

In a similar way,

$$e^{a_E + b_E - \frac{1}{n} \sum_{uv \in E(G)} \log F_E(d_u, d_v)} \le e^{a_E} + e^{b_E} - \frac{1}{n} \sum_{uv \in E(G)} e^{\log F_E(d_u, d_v)},$$

$$e^{a_E + b_E} e^{-\log(\prod_{uv \in E(G)} F_E(d_u, d_v))^{1/n}} \le e^{a_E} + e^{b_E} - \frac{1}{n} \sum_{uv \in E(G)} F_E(d_u, d_v),$$

$$\frac{e^{a_E + b_E}}{X_{\Pi, F_E}(G)^{1/m}} \le e^{a_E} + e^{b_E} - \frac{1}{m} X_{\Sigma, F_E}(G).$$

The following Kober's inequalities in  $\left[20\right]$  (see also  $\left[23,\ \mathrm{Lemma}\ 1\right]$ ) are useful.

Lemma 1 If  $a_j > 0$  for  $1 \leq j \leq k$ , then

$$\sum_{j=1}^{k} a_j + k(k-1) \left(\prod_{j=1}^{k} a_j\right)^{1/k} \le \left(\sum_{j=1}^{k} \sqrt{a_j}\right)^2 \le (k-1) \sum_{j=1}^{k} a_j + k \left(\prod_{j=1}^{k} a_j\right)^{1/k}$$

Lemma 1 has the following consequence.

**Theorem 4** Let G be a graph with n vertices and m edges. Then

$$X_{\Sigma,F_V^2}(G) + n(n-1)X_{\Pi,F_V}(G)^{2/n} \le X_{\Sigma,F_V}(G)^2 \le (n-1)X_{\Sigma,F_V^2}(G) + nX_{\Pi,F_V}(G)^{2/n},$$
  
$$X_{\Sigma,F_E^2}(G) + m(m-1)X_{\Pi,F_E}(G)^{2/m} \le X_{\Sigma,F_E}(G)^2 \le (m-1)X_{\Sigma,F_E^2}(G) + mX_{\Pi,F_E}(G)^{2/m}.$$

Proof Lemma 1 gives

$$\sum_{u \in V(G)} F_V(d_u)^2 + n(n-1) \Big(\prod_{u \in V(G)} F_V(d_u)^2\Big)^{1/n} \le \Big(\sum_{u \in V(G)} F_V(d_u)\Big)^2,$$
$$X_{\Sigma, F_V^2}(G) + n(n-1)X_{\Pi, F_V}(G)^{2/n} \le X_{\Sigma, F_V}(G)^2,$$

and

$$\left(\sum_{u \in V(G)} F_V(d_u)\right)^2 \le (n-1) \sum_{u \in V(G)} F_V(d_u)^2 + n \left(\prod_{u \in V(G)} F_V(d_u)^2\right)^{1/n},$$
$$X_{\Sigma, F_V}(G)^2 \le (n-1) X_{\Sigma, F_V^2}(G) + n X_{\Pi, F_V}(G)^{2/n}.$$

7

In a similar way,

$$\sum_{uv \in E(G)} F_E(d_u, d_v)^2 + m(m-1) \Big(\prod_{uv \in E(G)} F_E(d_u, d_v)^2\Big)^{1/m} \le \Big(\sum_{uv \in E(G)} F_E(d_u, d_v)\Big)^2,$$
$$X_{\Sigma, F_E^2}(G) + m(m-1) X_{\Pi, F_E}(G)^{2/m} \le X_{\Sigma, F_E}(G)^2,$$

and

$$\left(\sum_{uv\in E(G)} F_E(d_u, d_v)\right)^2 \le (m-1) \sum_{uv\in E(G)} F_E(d_u, d_v)^2 + m \left(\prod_{uv\in E(G)} F_E(d_u, d_v)^2\right)^{1/m} X_{\Sigma, F_E}(G)^2 \le (m-1) X_{\Sigma, F_E}(G) + m X_{\Pi, F_E}(G)^{2/m}.$$

Let us recall the following known Petrović inequality [21].

**Theorem 5** Let  $\varphi$  be a convex function on [0, a], and  $w_1, \ldots, w_n \ge 0$ . If  $t_1, \ldots, t_n \in [0, a]$  satisfy  $\sum_{k=1}^{n} t_k w_k \in (0, a]$ , and

$$\sum_{k=1}^{n} t_k w_k \ge t_j, \qquad j = 1, \dots, n,$$

then

$$\sum_{k=1}^{n} \varphi(t_k) w_k \le \varphi\Big(\sum_{k=1}^{n} t_k w_k\Big) + \Big(\sum_{k=1}^{n} w_k - 1\Big)\varphi(0).$$

Theorem 5 has the following consequence.

**Proposition 6** Let  $\varphi$  be a convex function on [0, a]. If  $t_1, \ldots, t_n \in [0, a]$  satisfy  $\sum_{k=1}^{n} t_k \in (0, a]$ , then

$$\sum_{k=1}^{n} \varphi(t_k) \le \varphi\Big(\sum_{k=1}^{n} t_k\Big) + (n-1)\varphi(0).$$

Proposition 6 allows to prove the following inequalities.

**Theorem 7** Let G be a graph with n vertices and m edges. Then

$$X_{\Sigma, F_V}(G) \le X_{\Pi, F_V}(G) + n - 1, \quad X_{\Sigma, F_E}(G) \le X_{\Pi, F_E}(G) + m - 1.$$

*Proof* Let us consider the convex function  $\varphi(x) = e^x$ . Proposition 6 gives

$$\begin{split} X_{\Sigma,F_{V}}(G) &= \sum_{u \in V(G)} F_{V}(d_{u}) = \sum_{u \in V(G)} e^{\log F_{V}(d_{u})} \\ &\leq e^{\sum_{u \in V(G)} \log F_{V}(d_{u})} + n - 1 = e^{\log \Pi_{u \in V(G)} F_{V}(d_{u})} + n - 1 \\ &= \Pi_{u \in V(G)} F_{V}(d_{u}) + n - 1 = X_{\Pi,F_{V}}(G) + n - 1. \end{split}$$

In a similar way,

$$\begin{split} X_{\Sigma,F_E}(G) &= \sum_{uv \in E(G)} F_E(d_u, d_v) = \sum_{uv \in E(G)} e^{\log F_E(d_u, d_v)} \\ &\leq e^{\sum_{uv \in E(G)} \log F_E(d_u, d_v)} + m - 1 = e^{\log \Pi_{uv \in E(G)} F_E(d_u, d_v)} + m - 1 \\ &= \Pi_{uv \in E(G)} F_E(d_u, d_v) + m - 1 = X_{\Pi,F_E}(G) + m - 1. \end{split}$$

**Theorem 8** Let G be a graph. Then

$$X_{\Pi, F_V}(G) \ge X_{\Sigma, \log F_V}(G) + 1, \qquad X_{\Pi, F_E}(G) \ge X_{\Sigma, \log F_E}(G) + 1$$

*Proof* The equality  $e^x \ge x + 1$  holds for every  $x \in \mathbb{R}$ . This inequality gives

$$\begin{aligned} X_{\Pi, F_V}(G) &= e^{\log X_{\Pi, F_V}(G)} \ge \log X_{\Pi, F_V}(G) + 1 \\ &= \log \left( \Pi_{u \in E(G)} F_V(d_u) \right) + 1 = \sum_{u \in E(G)} \log F_V(d_u) + 1 \\ &= X_{\Sigma, \log F_V}(G) + 1. \end{aligned}$$

The same argument gives the second inequality.

## 3 Statistical properties of multiplicative topological indices on random networks

## 3.1 Multiplicative topological indices on Erdös-Rényi random networks

Before computing MTIs on ER random networks we note that in the dense limit, i.e. when  $\langle d \rangle \gg 1$ , we can approximate  $d_u \approx d_v \approx \langle d \rangle$  in Eqs. (9-16), with

$$\langle d \rangle = (n-1)p. \tag{17}$$

Thus, for example, when  $np \gg 1$ , we can approximate  $NK_{\Pi}(G_{\text{ER}})$  as

$$NK(G_{\rm ER}) = \prod_{u \in V(G)} d_u \approx \prod_{u \in V(G)} \langle d \rangle \approx \langle d \rangle^n \,,$$

which leads us to

$$\ln NK(G_{\rm ER}) \approx n \ln \langle d \rangle$$

or

$$\frac{\ln NK(G_{\rm ER})}{n} \approx \ln \langle d \rangle \,. \tag{18}$$

A similar approximation gives

$$\frac{\ln \Pi_1(G_{\rm ER})}{n} \approx 2 \ln \langle d \rangle \,. \tag{19}$$

Also, for  $\Pi_2(G_{\text{ER}})$  we have that

$$\Pi_{2}(G_{\text{ER}}) = \prod_{uv \in E(G)} d_{u}d_{v} \approx \prod_{uv \in E(G)} \langle d \rangle \langle d \rangle = \prod_{uv \in E(G)} \langle d \rangle^{2} \approx \langle d \rangle^{n \langle d \rangle}$$

where we have used  $| E(G_{\text{ER}}) | = n \langle d \rangle / 2$ . Therefore

$$\ln \Pi_2(G_{\rm ER}) \approx n \langle d \rangle \ln \langle d \rangle$$

or

$$\frac{\ln \Pi_2(G_{\rm ER})}{n} \approx \langle d \rangle \ln \langle d \rangle \,. \tag{20}$$

Following this procedure, for the rest of the MTIs of Eqs. (12-16) we get:

$$\frac{\ln \Pi_1^*(G_{\rm ER})}{n} \approx \frac{1}{2} \langle d \rangle \ln(2 \langle d \rangle), \tag{21}$$

$$\frac{\ln R_{\Pi}(G_{\rm ER})}{n} \approx -\frac{1}{2} \left\langle d \right\rangle \ln \left\langle d \right\rangle, \tag{22}$$

$$\frac{\ln H_{\Pi}(G_{\rm ER})}{n} \approx -\frac{1}{2} \langle d \rangle \ln \langle d \rangle , \qquad (23)$$

$$\frac{\ln \chi_{\Pi}(G_{\rm ER})}{n} \approx -\frac{\ln 2}{4} \langle d \rangle - \frac{1}{4} \langle d \rangle \ln \langle d \rangle, \qquad (24)$$

and

$$\frac{\ln ID_{\Pi}(G_{\rm ER})}{n} \approx \frac{\ln 2}{2} \langle d \rangle - \langle d \rangle \ln \langle d \rangle \,. \tag{25}$$

From the approximate expressions above we note that the logarithm of the MTIs on ER random networks, normalized to the network size n, does not depend on n. That is, in the dense limit, we expect  $\ln X_{\Pi}(G_{\text{ER}})/n$  to depend on  $\langle d \rangle$  only; here X represents all the MTIs of Eqs. (9-16).

Having Eqs. (18-25) as a guide, in what follows we compute the average values of the logarithm of the MTIs listed in Eqs. (9-16). All averages are computed over ensembles of  $10^7/n$  ER networks characterized by the parameter pair (n, p).

In Fig. 1 we present the average logarithm of the eight MTIs of Eqs. (9-16) as a function of the probability p of ER networks of sizes  $n = \{125, 250, 500, 1000\}$ . Since  $\langle \ln X_{\Pi}(G_{\text{ER}}) \rangle < 0$  for  $R_{\Pi}$ ,  $H_{\Pi}$  and  $\chi_{\Pi}$  we conveniently plotted  $-\langle \ln X_{\Pi}(G_{\text{ER}}) \rangle$  in log scale to have a detailed view of the data for small p, see Figs. 1(e-g). Thus we observe that  $\langle \ln X_{\Pi}(G_{\text{ER}}) \rangle$  for NK,  $\Pi_{1,2}$  and  $\Pi_1^*$  [for  $R_{\Pi}$ ,  $H_{\Pi}$  and  $\chi_{\Pi}$ ] is a monotonically increasing [monotonically decreasing] function of p. In contrast,  $\langle \ln ID_{\Pi}(G_{\text{ER}}) \rangle$  is a nonmonotonic function of p, see Fig. 1(h).

Then, following Eqs. (18-25), in Fig. 2 we show again the average logarithm of the MTIs but now normalized to the network size,  $\langle \ln X_{\Pi}(G_{\text{ER}}) \rangle /n$ , as a function of  $\langle k \rangle$ . As can be clearly observed in this figure, the curves  $\langle \ln X_{\Pi}(G_{\text{ER}}) \rangle /n$  versus  $\langle k \rangle$  fall one on top of the other for different network

9





**Fig. 1** Average logarithm of the multiplicative topological indices of Eqs. (9-16) as a function of the probability p of Erdös-Rényi networks of size n. The inset in (h) is an enlargement for small p.

sizes; so these indices are properly scaled where  $\langle k \rangle$  works as the scaling parameter. That is,  $\langle k \rangle$  is the parameter that fixes the average values of the normalized MTIs on ER networks. It is remarkable that the scaling of  $\langle \ln X_{\Pi}(G_{\text{ER}}) \rangle / n$  with  $\langle k \rangle$ , expected for  $\langle k \rangle \gg 1$  according to Eqs. (18-25), works perfectly well for all values of  $\langle k \rangle$ ; even for  $\langle k \rangle \ll 1$ . Also, in Figs. 2, we show that Eqs. (18-25) (orange-dashed lines) indeed describe well the data (thick full curves) for  $\langle k \rangle \geq 10$ , which can be regarded as de dense limit of the ER model.

# 3.2 Multiplicative topological indices on random geometric graphs

As in the previous Subsection, here we start by exploring the dense limit. Indeed, for RG graphs in the dense limit, i.e. when  $\langle d \rangle \gg 1$ , we can



Fig. 2 Average logarithm of the multiplicative topological indices of Eqs. (9-16), normalized to the network size n, as a function of the average degree  $\langle k \rangle$  of Erdös-Rényi networks. Orange dashed lines in (a-h) are Eqs. (18-25), respectively. The vertical magenta dashed lines indicate  $\langle k \rangle = 10$ . The inset in (h) is an enlargement for small  $\langle k \rangle$ .

approximate  $d_u \approx d_v \approx \langle d \rangle$ , where

$$\langle d \rangle = (n-1)g(r) \tag{26}$$

and [17]

$$g(r) = \begin{cases} r^2 \left[ \pi - \frac{8}{3}r + \frac{1}{2}r^2 \right], & 0 \le r \le 1, \\ \frac{1}{3} - 2r^2 \left[ 1 - \arcsin(1/r) + \arccos(1/r) \right] & (27) \\ + \frac{4}{3}(2r^2 + 1)\sqrt{r^2 - 1} - \frac{1}{2}r^4, & 1 \le r \le \sqrt{2}. \end{cases}$$



Fig. 3 Average logarithm of the multiplicative topological indices of Eqs. (9-16) as a function of the connection radius r of random geometric graphs of size n. The inset in (h) is an enlargement for small r.

Thus, we obtain:

$$\frac{\ln NK(G_{\rm RG})}{n} \approx \ln \left\langle d \right\rangle,\tag{28}$$

$$\frac{\ln \Pi_1(G_{\rm RG})}{n} \approx 2 \ln \langle d \rangle \,, \tag{29}$$

$$\frac{\ln \Pi_2(G_{\rm RG})}{n} \approx \langle d \rangle \ln \langle d \rangle , \qquad (30)$$

$$\frac{\ln \Pi_1^*(G_{\rm RG})}{n} \approx \frac{1}{2} \langle d \rangle \ln(2 \langle d \rangle), \tag{31}$$

$$\frac{\ln R_{\Pi}(G_{\rm RG})}{n} \approx -\frac{1}{2} \langle d \rangle \ln \langle d \rangle, \qquad (32)$$

$$\frac{\ln H_{\Pi}(G_{\rm RG})}{n} \approx -\frac{1}{2} \left\langle d \right\rangle \ln \left\langle d \right\rangle, \tag{33}$$



Fig. 4 Average logarithm of the multiplicative topological indices of Eqs. (9-16), normalized to the graph size n, as a function of the average degree  $\langle k \rangle$  of random geometric graphs. Orange dashed lines in (a-h) are Eqs. (28-35), respectively. The vertical magenta dashed lines indicate  $\langle k \rangle = 10$ . The inset in (h) is an enlargement for small  $\langle k \rangle$ .

$$\frac{\ln \chi_{\Pi}(G_{\rm RG})}{n} \approx -\frac{\ln 2}{4} \left\langle d \right\rangle - \frac{1}{4} \left\langle d \right\rangle \ln \left\langle d \right\rangle, \tag{34}$$

and

$$\frac{\ln ID_{\Pi}(G_{\rm RG})}{n} \approx \frac{\ln 2}{2} \langle d \rangle - \langle d \rangle \ln \langle d \rangle.$$
(35)

Remarkably, the approximate Eqs. (28-35) for RG graphs are exactly the same as the corresponding equations for ER graphs, see Eqs. (18-25); however note that the definition of  $\langle d \rangle$  is different for both models, i.e. compare Eqs. (17) and (26-27).

Then, in Fig. 3 we present the average logarithm of the eight MTIs of Eqs. (9-16) as a function of the connection radius r of RG graphs of sizes  $n = \{125, 250, 500, 1000\}$ . All averages are computed over ensembles of  $10^7/n$  random graphs, each ensemble is characterized by a fixed parameter pair (n, r).





**Fig. 5** Average logarithm of the multiplicative topological indices of Eqs. (9-16) as a function of the probability p of bipartite random networks of sizes  $n_1$  and  $n_2$ . In all panels  $n_1 = n_2 = \{125, 250, 500, 1000\}$ . The inset in (h) is an enlargement for small p.

For comparison purposes, Fig. 1 for ER networks and Fig. 3 for RG graphs have the same format and contents. In fact, all the observations made in the previous Subsection for ER networks are also valid for RG graphs by replacing  $G_{\rm ER} \rightarrow G_{\rm RG}$  and  $p \rightarrow g(r)$ . Therefore, in Fig. 4 we plot  $\langle \ln X_{\Pi}(G_{\rm RG}) \rangle /n$  as a function of  $\langle k \rangle$  showing that all curves are properly scaled; that is, curves for different graph sizes fall on top of each other. Also, in Fig. 4, we show that Eqs. (28-35) (orange-dashed lines) indeed describe well the data (thick full curves) for  $\langle k \rangle \geq 10$ , which can be regarded as de dense limit of RG graphs.



**Fig. 6** Average logarithm of the multiplicative topological indices of Eqs. (9-16), normalized to the network size n, as a function of the average degree  $\langle k \rangle$  of bipartite random networks of sizes  $n_1$  and  $n_2$ . In all panels  $n_1 = n_2 = \{125, 250, 500, 1000\}$ . Orange dashed lines in (a,b) are Eqs. (18,19), respectively. Orange dashed lines in (c-h) are Eqs. (37-42), respectively. The vertical magenta dashed lines indicate  $\langle k \rangle = 10$ . The inset in (h) is an enlargement for small  $\langle k \rangle$ .

#### 3.3 Multiplicative topological indices on bipartite random networks

We start by writing approximate expressions for the MTIs on BR networks in the dense limit. Moreover, since edges in a bipartite network join vertices of different sets, and we are labeling here the sets as set 1 and set 2, we replace  $d_u$  by  $d_1$  and  $d_v$  by  $d_2$  in the expression for the MTIs. Thus, when  $n_1p \gg 1$ and  $n_2p \gg 1$ , we can approximate  $d_u = d_1 \approx \langle d_1 \rangle$  and  $d_v = d_2 \approx \langle d_2 \rangle$  in Eqs. (9-16), with

$$\langle d_{1,2} \rangle = n_{2,1}p. \tag{36}$$

Therefore, in the dense limit, the MTIs of Eqs. (11-16) on BR networks are well approximated by:

$$\frac{\ln \Pi_2(G_{\rm BR})}{n_{1,2}} \approx \langle d_{1,2} \rangle \ln(\langle d_1 \rangle \langle d_2 \rangle), \tag{37}$$

$$\frac{\ln \Pi_1^*(G_{\rm BR})}{n_{1,2}} \approx \langle d_{1,2} \rangle \ln(\langle d_1 \rangle + \langle d_2 \rangle), \tag{38}$$

$$\frac{\ln R_{\Pi}(G_{\rm BR})}{n_{1,2}} \approx -\frac{1}{2} \langle d_{1,2} \rangle \ln \left( \langle d_1 \rangle \langle d_2 \rangle \right), \tag{39}$$

$$\frac{\ln H_{\Pi}(G_{\rm BR})}{n_{1,2}} \approx \langle d_{1,2} \rangle \left[ \ln 2 - \ln \left( \langle d_1 \rangle + \langle d_2 \rangle \right) \right],\tag{40}$$

$$\frac{\ln \chi_{\Pi}(G_{\rm BR})}{n_{1,2}} \approx -\frac{1}{2} \left\langle d_{1,2} \right\rangle \ln \left( \left\langle d_1 \right\rangle + \left\langle d_2 \right\rangle \right),\tag{41}$$

and

$$\frac{\ln ID_{\Pi}(G_{\rm BR})}{n_{1,2}} \approx \langle d_{1,2} \rangle \ln \left( \frac{1}{\langle d_1 \rangle^2} + \frac{1}{\langle d_2 \rangle^2} \right). \tag{42}$$

Above we used  $|E(G_{BR})| = n_1 n_2 p = n_{1,2} \langle d_{1,2} \rangle$ . We note that we are not providing approximate expressions for  $NK(G_{BR})$  nor  $\Pi_1(G_{BR})$ .

Now we compute MTIs on ensembles of  $10^7/n$  BR networks. In contrast to ER and RG networks now the BR network ensembles are characterized by three parameters:  $n_1$ ,  $n_2$ , and p. For simplicity, but without lost of generality, in our numerical calculations we consider  $n_1 = n_2$ . It is remarkable to notice that in the case of  $n_1 = n_2 = n/2$ , where  $\langle k_1 \rangle = \langle k_2 \rangle = \langle k \rangle = np/2$ , Eqs. (37-42) reproduce Eqs. (20-25).

Then, in Fig. 5 we present the average logarithm of the eight MTIs of Eqs. (9-16) as a function of the probability p of BR networks of sizes  $n_1 = n_2 = \{125, 250, 500, 1000\}$ . Therefore, by plotting  $\langle \ln X_{\Pi}(G_{\rm BR}) \rangle /n$  as a function of  $\langle k \rangle$ , see Fig. 6, we confirm that the curves of the MTIs on BR networks are properly scaled, as predicted by Eqs. (37-42); see the orange dashed lines in Fig. 6(c-h). Here, we can also say that  $\langle k \rangle \geq 10$  can be regarded as de dense limit of BR networks. We have verified (not shown here) that we arrive to equivalent conclusions when  $n_1 \neq n_2$ .

Note that in panels (a,b) of Fig. 6 we included Eqs. (18,19) as orange dashed lines. Those equations were obtained for ER networks, however they reproduce perfectly well both  $NK(G_{\rm BR})$  and  $\Pi_1(G_{\rm BR})$  when  $\langle k \rangle \geq 10$  since we are considering  $n_1 = n_2 = n/2$ .

## 4 Conclusions

In this work we have performed a thorough analytical and statistical study of multiplicative topological indices (MTIs) on random networks. As models of random networks we have used Erdös-Rényi networks, random geometric graphs, and bipartite random networks.



Fig. 7 Average logarithm of (a) NK(G), (b)  $\chi_{\Pi}(G)$ , and (c)  $ID_{\Pi}(G)$ , normalized to the network size n, as a function of the average degree  $\langle k \rangle$  of random geometric graphs of size n = 500, Erdös-Rényi networks of size n = 250 and bipartite random networks of size  $n = 2n_1 = 2n_2 = 2000$ . Orange dashed lines are (a) Eq. (18), (b) Eq. (24), and (c) Eq. (25). The inset in (c) is an enlargement for small  $\langle k \rangle$ .

From an analytical viewpoint, we find several inequalities that relate multiplicative indices to their additive versions.

Within a statistical approach, we show that the average logarithm of the MTIs,  $\langle \ln X_{\Pi}(G) \rangle$ , normalized to the order of the network, scale with the network average degree  $\langle k \rangle$ . Thus, we conclude that  $\langle k \rangle$  is the parameter that fixes the average values of the logarithm of the MTIs on random networks. Moreover, the equivalence among Eqs. (18-25) for Erdös-Rényi networks, Eqs. (28-35) for random geometric graphs, and Eqs. (37-42) for bipartite random networks (when the subsets are equal in size) allows us to state a scaling law that connects the three graph models. That is, given the multiplicative topological index  $X_{\Pi}$ , the average of its logarithm divided by the network size is the same function of the average degree regardless the network model:

$$\frac{\langle \ln X_{\Pi}(G_{\rm ER}) \rangle}{n} \approx \frac{\langle \ln X_{\Pi}(G_{\rm RG}) \rangle}{n} \approx \frac{\langle \ln X_{\Pi}(G_{\rm BR}) \rangle}{n} \approx f(\langle k \rangle).$$
(43)

To validate Eq. (43) we choose the Narumi-Katayama index, see Eq. (9), apply it to Erdös-Rényi networks, random geometric graphs and bipartite random networks and plot  $\langle \ln NK_{\Pi}(G) \rangle / n$  in Fig. 7(a). There, we can clearly see that the curves corresponding to the three network models fall one on top of the other, thus validating Eq. (43). Also notice that we are using networks of different sizes to stress the scaling law in Eq. (43). We observe the same scaling behavior in all the other MTIs of Eqs. (10-16), as can be seen in Figs. 7(b,c) where we present  $\langle \ln \chi_{\Pi}(G) \rangle / n$  and  $\langle \ln ID_{\Pi}(G) \rangle / n$ , respectively.

Finally, it is fair to mention that we found that the multiplicative version of the geometric-arithmetic index  $GA_{\Pi}(G)$  on random networks does not scale with the average degree. This situation is quite unexpected for us since all other MTIs we study here scale with  $\langle k \rangle$ . In addition, our previous statistical and theoretical studies of topological indices of the form  $X_{\Sigma}(G)$ , see Eq. (1), showed that the normalized geometric-arithmetic index  $GA_{\Sigma}(G)$  scales with

 $\langle k \rangle$  [7]. Thus, we believe that the multiplicative index  $GA_{\Pi}(G)$  on random graphs requires further analysis.

We hope that our work may motivate further analytical as well as computational studies of multiplicative topological indices on random networks.

Acknowledgments. The research of J.M.R. and J.M.S. was supported by a grant from Agencia Estatal de Investigación (PID2019-106433GBI00/AEI/10.13039/501100011033), Spain. J.M.R. was supported by the Madrid Government (Comunidad de Madrid-Spain) under the Multiannual Agreement with UC3M in the line of Excellence of University Professors (EPUC3M23), and in the context of the V PRICIT (Regional Programme of Research and Technological Innovation).

## Declarations

The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

### References

- I. Gutman, Degree-based topological indices, Croat. Chem. Acta 86, 351–361 (2013).
- [2] I. Gutman and N. Trinajstić, Chem. Phys. Lett. 17 (1972) 535.
- [3] Randić, M. On characterization of molecular branching. J. Am. Chem. Soc. 1975, 97 6609–6615.
- [4] S. Fajtlowicz, On conjectures of Graffiti–II, Congr. Numer. 60 (1987) 187–197.
- [5] B. Zhou and N. Trinajstić, J. Math. Chem. 46 (2009) 1252.
- [6] C. T. Martínez-Martínez, J. A. Mendez-Bermudez, J. M. Rodríguez, and J. M. Sigarreta, Computational and analytical studies of the Randić index in Erdös–Rényi models, Appl. Math. Comput. 377, 125137 (2020).
- [7] R. Aguilar-Sanchez, I. F. Herrera-Gonzalez, J. A. Mendez-Bermudez, and J. M. Sigarreta, Computational properties of general indices on random networks. Symmetry 12, 1341 (2020).
- [8] C. T. Martínez-Martínez, J. A. Mendez-Bermudez, J. M. Rodríguez, and J. M. Sigarreta, Computational and analytical studies of the harmonic index in Erdös–Rényi models, MATCH Commun. Math. Comput. Chem. 85, 395–426 (2021).

- [9] R. Aguilar-Sanchez, J. A. Mendez-Bermudez, F. A. Rodrigues, and J. M. Sigarreta-Almira, Topological versus spectral properties of random geometric graphs, Phys. Rev. E 102, 042306 (2020).
- [10] H. Narumi and M. Katayama, Mem. Fac. Engin. Hokkaido Univ. 16 (1984) 209.
- [11] R. Todeschini and V. Consonni, MATCH Commun. Math. Comput. Chem. 64 (2010) 359.
- [12] R. Solomonoff and A. Rapoport, Connectivity of random nets. Bull. Math. Biophys. 13, 107–117 (1951).
- [13] P. Erdös and A. Rényi, On random graphs. Publ. Math. (Debrecen) 6, 290–297 (1959).
- [14] P. Erdös and A. Rényi, On the evolution of random graphs, Inst. of the Hung. Acad. of Sci. 5, 17–61 (1960); P. Erdös and A. Rényi, On the strength of connectedness of a random graph, Acta Mathematica Hungarica 12, 261–267 (1961).
- [15] J. Dall and M. Christensen, Random geometric graphs, Phys. Rev. E 66, 016121 (2002).
- [16] M. Penrose, Random Geometric Graphs; (Oxford University Press, Oxford, 2003).
- [17] E. Estrada and M. Sheerin, Random rectangular graphs, Phys Rev. E 91, 042805 (2015).
- [18] P. Bosch, Y. Quintana, J. M. Rodríguez, J. M. Sigarreta, Jensen-type inequalities for m-convex functions, to appear in *Open Math.*
- [19] I. Gutman, I. Milovanović, E. Milovanović, Relations between Ordinary and Multiplicative Degree-Based Topological Indices, *Filomat* 32:8 (2018) 3031–3042.
- [20] H. Kober, On the arithmetic and geometric means and on Hölder's inequality, Proc. Amer. Math. Soc. 9 (1958) 452–459.
- [21] M. Petrović's, Sur Une Fonctionnelle, Publ. Math. Univ. Belgrade 1 (1932) 146–149.
- [22] T. Réti, I. Gutman, Relations between ordinary and multiplicative Zagreb indices, Bull. Inter. Math. Virtual Inst. 2 (2012) 133–140.
- [23] B. Zhou, I. Gutman, T. Aleksić, A note on Laplacian energy of graphs, MATCH Commun. Math. Comput. Chem. 60 (2008) 441–446.