FLAT COMODULES AND CONTRAMODULES AS DIRECTED COLIMITS, AND COTORSION PERIODICITY

LEONID POSITSELSKI

ABSTRACT. This paper is a follow-up to [47]. We consider two algebraic settings of comodules over a coring and contramodules over a topological ring with a countable base of two-sided ideals. These correspond to two (noncommutative) algebraic geometry settings of certain kind of stacks and ind-affine ind-schemes. In the context of a coring C over a noncommutative ring A, we show that all A-flat C-comodules are \aleph_1 -directed colimits of A-countably presentable A-flat C-comodules. In the context of a complete, separated topological ring \mathfrak{R} with a countable base of neighborhoods of zero consisting of two-sided ideals, we prove that all flat \Re -contramodules are \aleph_1 -directed colimits of countably presentable flat \Re -contramodules. We also describe arbitrary complexes, short exact sequences, and pure acyclic complexes of A-flat C-comodules and flat \mathfrak{R} -contramodules as \aleph_1 -directed colimits of similar complexes of countably presentable objects. The arguments are based on a very general category-theoretic technique going back to an unpublished 1977 preprint of Ulmer and rediscovered in [38]. Applications to cotorsion periodicity and coderived categories of flat objects in the respective settings are discussed. In particular, in any acyclic complex of cotorsion R-contramodules, all the contramodules of cocycles are cotorsion.

Contents

Introduction	2
1. Preliminaries on Accessible Categories	5
2. Preliminaries on Corings and Comodules	9
3. A-Flat C-Comodules as Directed Colimits	12
4. Exact Sequences of A-Flat C-Comodules as Directed Colimits	14
5. Cotorsion Periodicity for Comodules	16
6. Exact Sequences of A-Flat C-Comodules II	20
7. Preliminaries on Topological Rings and Contramodules	23
8. Separated and Flat Contramodules	25
9. Countably Presentable Flat Contramodules	29
10. Flat Contramodules as Directed Colimits	32
11. Exact Sequences of Flat Contramodules as Directed Colimits	34
12. Cotorsion Periodicity for Contramodules	36
13. Exact Sequences of Flat Contramodules II	40
References	42

INTRODUCTION

The classical Govorov-Lazard theorem [18, 23] says that all flat modules (over an arbitrary associative ring R) are directed colimits of projective modules, and in fact, even of finitely generated free R-modules. In the context of algebraic geometry, over a nonaffine scheme X, there are usually not enough projective objects in the abelian category of quasi-coherent sheaves X-qcoh, which makes the role of flat quasi-coherent sheaves even more important than the role of flat modules in module theory. What should a suitable version of the Govorov-Lazard theorem say about flat quasi-coherent sheaves?

One could try to use locally free or locally projective quasi-coherent sheaves in the role of projective modules. But it is still an open problem whether there are enough locally projective quasi-coherent sheaves on X under any reasonable assumptions on a scheme X [52]. On the other hand, it is known that there are enough flat quasi-coherent sheaves on any quasi-compact semi-separated scheme [25, Section 2.4], [12, Lemma A.1]. So describing flat quasi-coherent sheaves is a worthwhile undertaking.

The approach in this paper follows the idea that, in many homological algebra contexts, one can use objects of finite projective dimension in lieu of projective ones. It is known that any countably presentable flat *R*-module has projective dimension ≤ 1 [17, Corollary 2.23]. Thus our suggested answer to the question in the first paragraph is this: the Govorov–Lazard theorem in algebraic geometry should tell us that any flat sheaf is a directed colimit of locally countably presentable flat ones. Notice, for comparison, that any finitely presentable flat module is projective.

A partial result was obtained in the paper [16], where it was shown that, under certain additional assumptions on a quasi-compact semi-separated scheme X, any flat quasi-coherent sheaf on X is a directed colimit of locally countably generated flat quasi-coherent sheaves locally of projective dimension ≤ 1 [16, Theorem B or Theorem 4.9]. In full generality, the desired assertion for quasi-compact quasiseparated schemes (or even more generally, for countably quasi-compact, countably quasi-separated schemes) X was proved in the preprint [47]: any flat quasi-coherent sheaf on X is an \aleph_1 -directed colimit of locally countably presentable flat quasicoherent sheaves [47, Theorem 2.4 or Theorem 3.5]. Here an \aleph_1 -directed colimit means the directed colimit of a diagram indexed by an \aleph_1 -directed poset, i. e., a poset in which every countable subset has an upper bound.

The present paper aims to extend the results of [47] to two algebraic geometry settings more general than schemes: the stacks and the ind-schemes. In fact, we don't assume our rings to be commutative in this paper, so it also extends the results of [47] into certain noncommutative algebraic geometry realms.

The observation that certain noncommutative stacks X can be described by corings \mathcal{C} over noncommutative rings A is due to Kontsevich and Rosenberg [21, Section 2], [22]. The quasi-coherent sheaves on X are then interpreted as left \mathcal{C} -comodules (which form an abelian category whenever \mathcal{C} is a flat right A-module). The quasi-coherent sheaf corresponding to a \mathcal{C} -comodule \mathcal{M} is flat if and only if \mathcal{M} is a flat A-module.

In commutative algebraic geometry, this description applies to the stacks X admitting a flat affine surjective morphism $U \longrightarrow X$ from an affine scheme U. Specifically, one has $A = \mathcal{O}(U)$ and $\mathcal{C} = \mathcal{O}(U \times_X U)$. For example, if X is a quasi-compact semiseparated scheme covered by a finite collection of affine open subschemes $U_{\alpha} \subset X$, then one can take U to be the disjoint union $U = \coprod_{\alpha} U_{\alpha}$. Then $A = \bigoplus_{\alpha} \mathcal{O}(U_{\alpha})$ and $\mathcal{C} = \bigoplus_{\alpha,\beta} \mathcal{O}(U_{\alpha} \cap U_{\beta})$. The category of quasi-coherent sheaves on X is equivalent to the category of C-comodules, and the inverse image functor X-qcoh $\longrightarrow U$ -qcoh corresponds to the forgetful functor C-Comod $\longrightarrow A$ -Mod. The C-comodule structure on a given A-module describes the descent/gluing datum needed to glue a quasicoherent sheaf on X from a given collection of quasi-coherent sheaves on U_{α} .

Let C be a coring over a noncommutative ring A (in the sense of the paper [51] and the book [6]). Then we prove that any C-comodule that is flat as an A-module can be obtained as an \aleph_1 -directed colimit of C-comodules that are flat and countably presentable as A-modules (see Theorem 3.1). Similarly, any C-comodule is an \aleph_1 -directed colimit of C-comodules that are countably presentable as A-modules (Remark 3.2). Furthermore, any complex of A-flat C-comodules is an \aleph_1 -directed colimit of complexes of A-countably presentable A-flat C-comodules (by Proposition 3.3). We also show that any short exact sequence of A-flat C-comodules is an \aleph_1 -directed colimit of short exact sequences of A-countably presentable A-flat C-comodules (this is the result of our Proposition 4.4). More generally, any A-pure acyclic complex of A-flat C-comodules is an \aleph_1 -directed colimit of A-flat C-comodules (this is the result of A-flat C-comodules (see Corollary 4.5).

A left *R*-module *P* is said to be *cotorsion* if $\operatorname{Ext}_{R}^{1}(F, P) = 0$ for all flat left *R*-modules *F*. The *cotorsion periodicity theorem* of Bazzoni, Cortés-Izurdiaga, and Estrada [2] claims that, in any acyclic complex of cotorsion *R*-modules, all the modules of cocycles are also cotorsion [2, Theorem 1.2(2), Proposition 4.8(2), or Theorem 5.1(2)].

The following comodule version of cotorsion periodicity theorem is obtained in this paper. Assume that \mathcal{C} is a flat right A-module, all left C-comodules are quotients of A-flat left C-comodules, and all left C-comodules having finite projective dimension as left A-modules also have finite projective dimension as C-comodules. Let us say that a left C-comodule \mathcal{B} is cotorsion if $\operatorname{Ext}^1_{\mathcal{C}}(\mathcal{F}, \mathcal{B}) = 0$ for all A-flat left C-comodules \mathcal{F} . Here $\operatorname{Ext}^*_{\mathcal{C}}$ denotes the Ext groups in the abelian category of left C-comodules of cocycles are also cotorsion (see Theorem 5.4 and Corollary 5.5). As a corollary of this periodicity theorem, we conclude that the derived category of the abelian category of left C-comodules is equivalent to the derived category of the exact category of cotorsion left C-comodules (Corollary 5.6).

Under the same assumptions as in the previous paragraph, we also obtain the following description of A-pure acyclic complexes of A-flat left C-comodules. Such complexes are \aleph_1 -directed colimits of totalizations of finite acyclic complexes (of a certain fixed length) of complexes of A-countably presentable A-flat C-comodules. This result, based on [28, Lemma 2.1] and [36, proof of Proposition 8.8], provides

a comodule version of the well-known description of pure acyclic complexes of flat R-modules as directed colimits of finite contractible complexes of finitely generated projective R-modules [14, Theorem 2.4 (1) \Leftrightarrow (3)], [26, Theorem 8.6 (ii) \Leftrightarrow (iii)]. This is our Theorem 6.5.

The discussion of ind-schemes in this paper is mostly restricted to strict ind-affine \aleph_0 -ind-schemes, i. e., the ind-schemes represented by countable directed diagrams of closed immersions of affine schemes. The category of such ind-schemes \mathfrak{X} is anti-equivalent to the category of complete, separated topological commutative rings \mathfrak{R} with a countable base of neighborhoods of zero consisting of open ideals [4, Example 7.11.2(i)], [35, Example 1.6(2)]. Flat pro-quasi-coherent pro-sheaves on \mathfrak{X} (in the sense of [4, Section 7.11.3], [35, Section 3.4]) are described by flat \mathfrak{R} -contramodules, while the category of all pro-quasi-coherent pro-sheaves on \mathfrak{X} is equivalent to the category of separated \mathfrak{R} -contramodules [35, Examples 3.8(1-2)].

As a noncommutative generalization of this class of ind-schemes, we consider complete, separated topological associative rings \Re with a countable base of neighborhoods of zero consisting of open *two-sided* ideals. Contramodules over such topological rings \Re were discussed in the long preprint [29, Appendix E], while the more general case of a countable base of *right* ideals was studied in the paper [43].

In this context (assuming a countable topology base of two-sided ideals in \mathfrak{R}), we prove that any flat \mathfrak{R} -contramodule is an \aleph_1 -directed colimit of countably presentable flat \mathfrak{R} -contramodules (see Theorem 10.1). Furthermore, any complex of flat \mathfrak{R} -contramodules is an \aleph_1 -directed colimit of complexes of countably presentable flat \mathfrak{R} -contramodules (by Proposition 10.2). We also show that any short exact sequence of flat \mathfrak{R} -contramodules is an \aleph_1 -directed colimit of short exact sequences of countably presentable flat \mathfrak{R} -contramodules (Proposition 11.3), and any acyclic complex of flat \mathfrak{R} -contramodules with flat \mathfrak{R} -contramodules of cocycles is an \aleph_1 -directed colimit of such complexes of countably presentable flat \mathfrak{R} -contramodules (Corollary 11.4). It follows that any acyclic complexes of flat \mathfrak{R} -contramodules with flat \mathfrak{R} -contramodules of cocycles is an \aleph_1 -directed colimit of totalizations of short exact sequences of complexes of flat \mathfrak{R} -contramodules (Theorem 13.2).

Then we deduce the following contramodule version of cotorsion periodicity theorem. A left \mathfrak{R} -contramodule \mathfrak{B} is said to be *cotorsion* if $\operatorname{Ext}^{\mathfrak{R},1}(\mathfrak{F},\mathfrak{B}) = 0$ for all flat left \mathfrak{R} -contramodules \mathfrak{F} . Here $\operatorname{Ext}^{\mathfrak{R},*}$ denotes the Ext groups in the abelian category of left \mathfrak{R} -contramodules \mathfrak{R} -Contra. The theorem claims that, in any acyclic complex of cotorsion \mathfrak{R} -contramodules, the contramodules of cocycles are also cotorsion (see Theorem 12.3 and Corollary 12.4). As a corollary of this periodicity theorem, we deduce an equivalence between the derived category of the abelian category of left \mathfrak{R} -contramodules and the derived category of the exact category of cotorsion left \mathfrak{R} -contramodules (Corollary 12.8).

Periodicity theorems can be thought of as expressing special properties of directed colimit closures of exact categories (see the paper [39] for a discussion of the general concept of such directed colimit closure). We refer to the introduction to [3] for a quick survey on periodicity theorems and to the preprint [40, Sections 7.8 and 7.10]

for a discussion of the mentioned point of view on periodicity theorems. In particular, the category of flat modules is the directed colimit closure of the category of finitely generated projective modules, which is split exact. In this context, the flat/projective periodicity theorem [5, 26] can be interpreted as saying that for the exact category of flat modules, the contraderived category coincides with the derived category [40, Theorem 7.14], while the cotorsion periodicity theorem [2] is closely related to the assertion that for the exact category of flat modules, the coderived category coincides with the derived category [40, Theorem 7.18].

We refer the reader to the survey paper [37, Section 7] for a discussion of the history and philosophy of the coderived and contraderived categories (see also [45, Remark 9.2]). More advanced discussions of the coderived and contraderived categories in the context relevant to the present paper can be found in the papers [45, 46] (see [36] for a different point of view). In this paper we observe that, under suitable assumptions, the results describing the pure acyclic complexes of flat objects as the directed colimits of pure acyclic complexes of countably presentable flat objects can be interpreted as implying that for the exact category of such flat objects, the coderived category coincides with the derived category. The description of the coderived category obtained in [46, Corollary 0.5 or Proposition 8.13] is used as the reference point for the analogy or comparison here, in connection with our Theorems 6.5 and 13.2.

Let us say a few words about the proofs. The proofs of the main results of this paper are based on very general category-theoretic observations concerning preservation of κ -accessible categories and κ -presentable objects (for a regular cardinal κ) by category-theoretic constructions such as the pseudopullback [8, Proposition 3.1], [48, Pseudopullback Theorem 2.2], the equifier [38, Section 3], and the inserter [38, Section 4]. These results, going back to an unpublished 1977 preprint of Ulmer [53] and rediscovered in [38], depend on the assumption of existence of a smaller infinite cardinal $\lambda < \kappa$ such that the κ -accessible categories involved have colimits of λ -indexed chains and the functors involved preserve such colimits. In the situation at hand, we take $\kappa = \aleph_1$ and $\lambda = \aleph_0$. The key observation is that the classes of A-flat C-comodules and flat \Re -contramodules, are preserved by all directed colimits. This allows to prove that the respective categories are \aleph_1 -accessible, and describe their full subcategories of \aleph_1 -presentable objects.

Acknowledgement. I am grateful to Jan Sťovíček for helpful discussions. The author is supported by the GAČR project 23-05148S and the Czech Academy of Sciences (RVO 67985840).

1. Preliminaries on Accessible Categories

Let κ be a regular cardinal. We refer to [1, Definition 1.4, Theorem 1.5, Corollary 1.5, Definition 1.13(1), and Remark 1.21] for the discussion of κ -directed posets vs. κ -filtered small categories, and accordingly, κ -directed vs. κ -filtered colimits. Let C be a category with κ -directed (equivalently, κ -filtered) colimits. An object $S \in C$ is said to be κ -presentable if the functor $\operatorname{Hom}_{\mathsf{C}}(S, -) \colon \mathsf{C} \longrightarrow \mathsf{Sets}$ preserves κ -directed colimits (see [1, Definition 1.13(2)]). We will denote the class of all κ -presentable objects of C by $\mathsf{C}_{<\kappa} \subset \mathsf{C}$. The \aleph_0 -presentable objects are called finitely presentable (see [1, Definition 1.1]); and the \aleph_1 -presentable objects can be similarly called *countably presentable*.

This category-theoretic terminology is consistent with the module-theoretic one. Given an associative ring A, an A-module M is κ -presentable as an object of the category of left A-modules A-Mod (in the sense of the definition above) if and only if it is the cokernel of a morphism of free A-modules with less than κ generators.

Given an additive category E, we denote by Com(E) the category of (unbounded) cochain complexes in E. Clearly, if κ -directed colimits exist in E, then they also exist in Com(E). The following lemma is standard.

Lemma 1.1. (a) Let E be an additive category with directed colimits. Then any bounded complex of finitely presentable objects of E is a finitely presentable object in Com(E).

(b) Let κ be an uncountable regular cardinal and E be an additive category with κ -directed colimits. Then any complex of κ -presentable objects of E is a κ -presentable object in Com(E).

Proof. Part (a) holds, because directed colimits commute with finite limits in the category of abelian groups. Part (b) is true due to the fact that \aleph_1 -directed colimits commute with countable limits of abelian groups.

A category C with κ -directed colimits is said to be κ -accessible if there is a set of κ -presentable objects $S \subset C$ such that every object of C is a κ -directed colimit of objects from S (see [1, Definition 2.1]). If this is the case, then the κ -presentable objects of C are precisely all the retracts of objects from S. A κ -accessible category C is said to be *locally* κ -presentable if all colimits exist in C (see [1, Definition 1.17 and Theorem 1.20]). The \aleph_0 -accessible categories are called *finitely accessible* [1, Remark 2.2(1)], and the locally \aleph_0 -presentable categories are called *locally finitely* presentable [1, Definition 1.9 and Theorem 1.11].

We denote by $A-Mod_{flat} \subset A-Mod$ the full subcategory of flat left A-modules.

Lemma 1.2. For any ring A and any regular cardinal κ , the category of flat left A-modules A-Mod_{flat} is κ -accessible. The κ -presentable objects of A-Mod_{flat} are precisely all the κ -presentable flat A-modules (i. e., the flat A-modules that are κ -presentable in A-Mod).

Proof. The Govorov–Lazard description of flat A-modules as the directed colimits of finitely generated projective A-modules [18], [23], [17, Corollary 2.22] means that the category A–Mod_{flat} is finitely accessible, and the finitely generated projective A-modules are its finitely presentable objects. By [1, Theorem 2.11 and Example 2.13(1)], it follows that the category A–Mod_{flat} is κ -presentable for all regular cardinals κ ; and the argument in [1, proof of Theorem 2.11 (iv) \Rightarrow (i)] implies the desired description of κ -presentable objects. The point is that all flat A-modules that are κ -presentable in A-Mod are also κ -presentable in A-Mod_{flat}, since the full subcategory $A-Mod_{flat}$ is closed under directed colimits in A-Mod. On the other hand, any directed colimit of κ -presentable flat A-modules indexed by a directed poset of cardinality smaller than κ is a κ -presentable flat A-module again.

The following lemma illustrates the significance of countably presentable (i. e., \aleph_1 -presentable) flat modules.

Lemma 1.3. Any countably presentable flat module over an associative ring A is a countable directed colimit of finitely generated free A-modules. Consequently, the projective dimension of any countably presentable flat module is less than or equal to 1.

Proof. See [17, Corollary 2.23]. The second assertion of the lemma follows from the first one because the telescope construction of countable directed colimits provides a two-term projective resolution of any countable directed colimit of projective A-modules.

In the rest of this section, we discuss several category-theoretic constructions which will be used in this paper: the product, the pseudopullback, the isomorpher, the inserter, the equifier, and the diagram category. We recall results from the papers [53, [8, 48, 38] concerning κ -accessibility of the categories produced by such constructions and the descriptions of κ -presentable objects in these categories.

In almost all the contexts, we will consider a regular cardinal κ and a smaller infinite cardinal $\lambda < \kappa$ (so κ has to be uncountable). In the applications in the main body of the paper, we will use $\kappa = \aleph_1$ and $\lambda = \aleph_0$.

A λ -indexed chain (of objects and morphisms) in a category C is a functor $\lambda \longrightarrow C$. where λ is considered as an ordinal, and this ordinal, viewed as an ordered set, is considered as a small category. In other words, a λ -indexed chain is a directed diagram $(C_i \to C_i)_{0 \le i \le j \le \lambda}$ in C.

Proposition 1.4. Let κ be a regular cardinal and Ξ be a set of cardinality smaller than κ . Let $(\mathsf{K}_{\xi})_{\xi\in\Xi}$ be a family of κ -accessible categories indexed by the set Ξ . Then the Cartesian product $\mathsf{K} = \prod_{\xi \in \Xi} \mathsf{K}_{\xi}$ is also a κ -accessible category. The κ -presentable objects of K are precisely all the collections of objects $(S_{\xi} \in \mathsf{K}_{\xi})_{\xi \in \Xi}$ with $S_{\xi} \in (\mathsf{K}_{\xi})_{<\kappa}$ for all $\xi \in \Xi$.

Proof. This is a corrected version of [1, proof of Proposition 2.67]. See [38, Proposition 2.1 for the details. \square

Let K_1, K_2 , and L be three categories, and let $F_1: K_1 \longrightarrow L$ and $F_2: K_2 \longrightarrow L$ be two functors. The *pseudopullback* C of the pair of functors F_1 , F_2 is defined as the category of triples (K_1, K_2, θ) , where $K_1 \in \mathsf{K}_1$ and $K_2 \in \mathsf{K}_2$ are two objects, and θ : $F_1(K_1) \simeq F_2(K_2)$ is an isomorphism in L.

Theorem 1.5. Let κ be a regular cardinal and $\lambda < \kappa$ be a smaller infinite cardinal. Let K_1 , K_2 , and L be three κ -accessible categories where all λ -indexed chains have colimits. Assume that two functors $F_1: \mathsf{K}_1 \xrightarrow{\gamma} \mathsf{L}$ and $F_2: \mathsf{K}_2 \longrightarrow \mathsf{L}$ preserve κ -directed colimits and colimits of λ -indexed chains. Assume further that the functors F_1 and F_2 take κ -presentable objects to κ -presentable objects. Then the pseudopullback category C is κ -accessible. The κ -presentable objects of C are precisely all the triples $(S_1, S_2, \theta) \in C$ with $S_1 \in (K_1)_{<\kappa}$ and $S_2 \in (K_2)_{<\kappa}$.

Proof. This is [48, Pseudopullback Theorem 2.2], based on the argument in [8, proof of Proposition 3.1]. The assertion essentially goes back to [53, Remark 3.2(I), Theorem 3.8, Corollary 3.9, and Remark 3.11(II)]. See also [38, Corollary 5.1].

Let K and L be two categories, and let $F_1, F_2: K \Longrightarrow L$ be two parallel functors. The *isomorpher* C of the pair of functors F_1, F_2 is defined as the category of pairs (K, θ) , where $K \in K$ is an object and $\theta: F_1(K) \simeq F_2(K)$ is an isomorphism in L.

Theorem 1.6. Let κ be a regular cardinal and $\lambda < \kappa$ be a smaller infinite cardinal. Let K and L be two κ -accessible categories where all λ -indexed chains have colimits. Assume that two parallel functors F_1 , F_2 : $\mathsf{K} \rightrightarrows \mathsf{L}$ preserve κ -directed colimits and colimits of λ -indexed chains. Assume further that the functors F_1 and F_2 take κ -presentable objects to κ -presentable objects. Then the isomorpher category C is κ -accessible. The κ -presentable objects of C are precisely all the pairs $(S, \theta) \in \mathsf{C}$ with $S \in \mathsf{K}_{<\kappa}$.

Proof. This is an equivalent reformulation of Theorem 1.5; see [38, Remark 5.2]. The assertion can be also viewed as a particular case of [53, Remark 3.2(I), Theorem 3.8, Corollary 3.9, and Remark 3.11(II)].

Let K and L be two categories, and $F, G: K \Rightarrow L$ be two parallel functors. The *inserter* D of the pair of functors F, G is defined [1, Section 2.71] as the category of pairs (K, ϕ) , where $K \in K$ is an object and $\phi: F(K) \longrightarrow G(K)$ is a morphism in L.

Theorem 1.7. Let κ be a regular cardinal and $\lambda < \kappa$ be a smaller infinite cardinal. Let K and L be two κ -accessible categories where all λ -indexed chains have colimits. Assume that two parallel functors F, G: K \Rightarrow L preserve κ -directed colimits and colimits of λ -indexed chains. Assume further that the functor F takes κ -presentable objects to κ -presentable objects. Then the inserter category D is κ -accessible. The κ -presentable objects of D are precisely all the pairs $(S, \phi) \in D$ with $S \in K_{<\kappa}$.

Proof. This is [53, Theorem 3.8, Corollary 3.9, and Remark 3.11(II)] or [38, Theorem 4.1].

Let D and M be two categories, $F, G: D \Rightarrow M$ be two parallel functors, and ϕ , $\psi: F \Rightarrow G$ be two parallel natural transformations. The *equifier* E of the pair of natural transformations ϕ, ψ is defined [1, Lemma 2.76] as the full subcategory in D consisting of all objects $E \in D$ for which the two morphisms $\phi_E: F(E) \longrightarrow G(E)$ and $\psi_E: F(E) \longrightarrow G(E)$ are equal to each other, i. e., $\phi_E = \psi_E$.

Theorem 1.8. Let κ be a regular cardinal and $\lambda < \kappa$ be a smaller infinite cardinal. Let D and M be two κ -accessible categories where all λ -indexed chains have colimits. Assume that two parallel functors F, G: D \Rightarrow M preserve κ -directed colimits and colimits of λ -indexed chains. Assume further that the functor F takes κ -presentable objects to κ -presentable objects. Let ϕ , ψ : $F \rightrightarrows G$ be two parallel natural transformations. Then the equifier category E is κ -accessible. The κ -presentable objects of E are precisely all the objects of E that are κ -presentable as objects of D .

Proof. This is [53, Theorem 3.8, Corollary 3.9, and Remark 3.11(II)] or [38, Theorem 3.1].

Let k be a commutative ring. The notion of a κ -presented k-linear category spelled out in [38, Section 6] represents an intuitively clear idea of a quiver with less than κ vertices, less than κ arrows, and less than κ relations between k-linear combinations of various iterated compositions of the arrows. We refer to [38] for the details. Given a small k-linear category D and a k-linear category K, we denote by $\operatorname{Fun}_k(D, \mathsf{K})$ the category of k-linear functors $D \longrightarrow \mathsf{K}$.

Theorem 1.9. Let κ be a regular cardinal and $\lambda < \kappa$ be a smaller infinite cardinal. Let C be a κ -accessible k-linear category where all λ -indexed chains have colimits, and let D be a κ -presented k-linear category. Then the diagram category $Fun_k(D, C)$ is κ -accessible. The κ -presentable objects of $Fun_k(D, C)$ are precisely all the k-linear functors $D \longrightarrow C_{<\kappa}$.

Proof. This is [38, Theorem 6.2].

2. Preliminaries on Corings and Comodules

The following definitions seem to go back to Sweedler's paper [51]. A detailed exposition can be found in the book [6]. The book [28] and the survey paper [30, Sections 2.5 and 3.4] can be used as further references.

A coring over an associative ring A is a comonoid object in the monoidal category of A-A-bimodules with respect to the tensor product over A. In other words, a coring \mathbb{C} over A is an A-A-bimodule endowed with A-A-bimodule maps of comultiplication $\mu: \mathbb{C} \longrightarrow \mathbb{C} \otimes_A \mathbb{C}$ and counit $\epsilon: \mathbb{C} \longrightarrow A$ satisfying the usual coassociativity and counitality axioms. Specifically, the two compositions $(\mu \otimes id_{\mathbb{C}}) \circ \mu$ and $(id_{\mathbb{C}} \otimes \mu) \circ \mu$ must be equal to each other,

$$\mathcal{C} \longrightarrow \mathcal{C} \otimes_A \mathcal{C} \rightrightarrows \mathcal{C} \otimes_A \mathcal{C} \otimes_A \mathcal{C},$$

and both the compositions $(\epsilon \otimes id_{\mathcal{C}}) \circ \mu$ and $(id_{\mathcal{C}} \otimes \epsilon) \circ \mu$ must be equal to the identity map $id_{\mathcal{C}}$,

$$\mathcal{C} \longrightarrow \mathcal{C} \otimes_A \mathcal{C} \rightrightarrows \mathcal{C}.$$

The category of left A-modules A-Mod is a left module category (with respect to the tensor product operation) over the monoidal category of A-A-bimodules A-Bimod-A. A left C-comodule is a comodule object in the module category A-Mod over the comonoid object C in the monoidal category A-Bimod-A. In other words, a left comodule \mathcal{M} over C is a left A-module endowed with a left A-module map of left coaction $\nu \colon \mathcal{M} \longrightarrow \mathbb{C} \otimes_A \mathcal{M}$ satisfying the usual coassociativity and counitality

axioms together with the comultiplication map μ and the counit map ϵ of the coring C. Specifically, the two compositions $(\mu \otimes id_{\mathcal{M}}) \circ \nu$ and $(id_{\mathcal{C}} \otimes \nu) \circ \nu$ must be equal to each other,

$$\mathcal{M} \longrightarrow \mathcal{C} \otimes_A \mathcal{M} \rightrightarrows \mathcal{C} \otimes_A \mathcal{C} \otimes_A \mathcal{M},$$

and the composition $(\epsilon \otimes id_{\mathcal{M}}) \circ \nu$ must be equal to the identity map $id_{\mathcal{M}}$,

$$\mathcal{M} \longrightarrow \mathcal{C} \otimes_A \mathcal{M} \longrightarrow \mathcal{M}.$$

We denote the additive category of left C-comodules by C-Comod, and use the notation $\operatorname{Hom}_{\mathbb{C}}(-,-)$ for the groups of morphisms in C-Comod. A C-comodule is said to be A-flat if it is flat as an A-module. The full subcategory of A-flat left C-comodules is denoted by C-Comod_{A-flat} \subset C-Comod.

Lemma 2.1. Let \mathcal{C} be a coring over a ring A.

(a) All colimits exist in the additive category of left C-comodules. The forgetful functor C-Comod \longrightarrow A-Mod preserves colimits (in particular, coproducts).

(b) If \mathcal{C} is a flat right A-module, then the category \mathcal{C} -Comod is a Grothendieck abelian category, and the forgetful functor \mathcal{C} -Comod \longrightarrow A-Mod is exact.

(c) Conversely, if the category C-Comod is abelian and the forgetful functor C-Comod \longrightarrow A-Mod is exact, then C is a flat right A-module.

(d) There exists a coring C such that the category C-Comod is not abelian.

Proof. Part (a) follows from the fact that the tensor product functor $\mathbb{C} \otimes_A -$ preserves colimits in A-Mod. Parts (b-c) are covered by [6, Sections 18.6, 18.14 and 18.16]; see also [30, Proposition 2.12(a)]. A counterexample for part (d) can be found in [29, Example C.1.1].

When C is not a flat right A-module, exact sequences of arbitrary left C-comodules are not well-behaved. But there is still a well-behaved notion of a *short exact sequence* of A-flat left C-comodules, defined as a short sequence of A-flat left C-comodules that is exact in A-Mod. We refer to the survey [7] for background material on exact categories in the sense of Quillen.

Lemma 2.2. The additive category of A-flat C-comodules C-Comod_{A-flat} endowed with the class of short exact sequences defined above is an exact category.

Proof. Essentially, it suffices to check that the kernel of any surjective morphism of A-flat C-comodules exists in C-Comod_{A-flat} and agrees with the kernel of the same morphism computed in the category of flat A-modules. For this purpose, one needs to use the fact that the functors $C \otimes_A -$ and $C \otimes_A C \otimes_A -$ take short exact sequences of flat A-modules to short exact sequences of A-modules. \Box

More generally, an exact sequence of left A-modules is called *pure* if it stays exact after the functor $N \otimes_A -$ is applied to it, for every right A-module N. An A-pure short exact sequence of C-comodules is a short sequence of C-comodules that is pure exact as a short sequence of A-modules. The category of left C-comodules C-Comod with A-pure short exact sequences is also an exact category. A left C-comodule is said to be A-countably presentable if it is countably presentable as an A-module. Similarly, a C-comodule is said to be A-finitely presentable if it is finitely presentable as an A-module.

Lemma 2.3. Let \mathcal{C} be a coring over a ring A. Then

(a) all A-finitely presentable C-comodules are finitely presentable as objects of the additive category C-Comod;

(b) all bounded complexes of A-finitely presentable C-comodules are finitely presentable as objects of the additive category of complexes of C-comodules;

(c) all A-countably presentable C-comodules are countably presentable (i. e., \aleph_1 -presentable) as objects of the additive category C-Comod;

(d) all complexes of A-countably presentable C-comodules are countably presentable as objects of the additive category of complexes of C-comodules.

Proof. Part (b) follows from part (a) according to Lemma 1.1(a); and part (d) follows from part (c) in view of Lemma 1.1(b). Parts (a) and (c) follow from the facts that, for any two left C-comodules \mathcal{L} and \mathcal{M} , the abelian group Hom_c(\mathcal{L}, \mathcal{M}) can be computed as the kernel of (the difference of) a natural pair of maps

$$\operatorname{Hom}_{A}(\mathcal{L}, \mathcal{M}) \ \rightrightarrows \ \operatorname{Hom}_{A}(\mathcal{L}, \ \mathfrak{C} \otimes_{A} \mathcal{M}),$$

and the functor $\mathcal{C} \otimes_A$ – preserves colimits.

For the converse assertions to Lemma 2.3(c–d), see Remarks 3.2 and 3.4 below. It will be also explained there that the additive categories of left C-comodules C–Comod and complexes of left C-comodules Com(C–Comod) are locally \aleph_1 -presentable.

Lemma 2.4. Let C be a coring over a ring A. Assume that C is a flat left and right A-module, and the ring A has finite weak global dimension (or in other words, finite Tor-dimension). Then every C-comodule is a quotient comodule of an A-flat C-comodule.

Proof. This is [28, Lemma 1.1.3]. For a discussion of this construction in a module-theoretic context, see [34, Section 2.2]. \Box

Example 2.5. This example is due to Kontsevich and Rosenberg [21, Section 2], [22]. Let X be a quasi-compact semi-separated scheme, and let $X = \bigcup_{\alpha} U_{\alpha}$ be a finite covering of X by affine open subschemes. Let U denote the disjoint union $U = \coprod_{\alpha} U_{\alpha}$; so U is an affine scheme and there is a natural surjective flat affine morphism of schemes $U \longrightarrow X$. Let $A = \mathcal{O}(U)$ and $\mathcal{C} = \mathcal{O}(U \times_X U)$ denote the rings of functions on the affine schemes U and $U \times_X U$. Then the two projections $p_1, p_2: U \times_X U \rightrightarrows U$ induce two ring homomorphisms $A \rightrightarrows \mathcal{C}$, endowing \mathcal{C} with two A-module structures (which we view as the "left" and the "right" one).

The diagonal morphism $\delta: U \longrightarrow U \times_X U$ induces a ring homomorphism $\delta^{\vee}: \mathbb{C} \longrightarrow A$. *A*. Furthermore, we have $\mathbb{C} \otimes_A \mathbb{C} = \mathbb{O}((U \times_X U) \times_U (U \times_X U)) = \mathbb{O}(U \times_X U \times_X U)$. The projection $q: U \times_X U \times_X U \longrightarrow U \times_X U$ dropping the second (inner) coordinate induces a ring homomorphism $q^{\vee}: \mathbb{C} \longrightarrow \mathbb{C} \otimes_A \mathbb{C}$. The maps $\epsilon = \delta^{\vee}: \mathbb{C} \longrightarrow A$ and

 $\mu = q^{\vee} \colon \mathfrak{C} \longrightarrow \mathfrak{C} \otimes_A \mathfrak{C}$ that we have constructed endow the A-A-bimodule \mathfrak{C} with a coring structure (over the commutative ring A).

For any quasi-coherent sheaf N on X, the inverse image of N with respect to the morphism $U \longrightarrow X$ produces a quasi-coherent sheaf M on U, that is an A-module \mathcal{M} . The inverse images of M with respect to the two projections $U \times_X U \rightrightarrows U$ are naturally isomorphic as quasi-coherent sheaves on $U \times_X U$, that is $p_1^*M \simeq p_2^*M$. The corresponding bimodules of sections are $\mathcal{M} \otimes_A \mathbb{C}$ and $\mathbb{C} \otimes_A \mathcal{M}$; so we have a natural isomorphism of A-A-bimodules $\mathcal{M} \otimes_A \mathbb{C} \simeq \mathbb{C} \otimes_A \mathcal{M}$. Precomposing this isomorphism with the map $\mathcal{M} \longrightarrow \mathcal{M} \otimes_A \mathbb{C}$ induced by the "left" map $A \longrightarrow \mathbb{C}$ (induced by p_1), we obtain a map $\nu \colon \mathcal{M} \longrightarrow \mathbb{C} \otimes_A \mathcal{M}$, endowing \mathcal{M} with a left \mathbb{C} -comodule structure. The assignment $N \longmapsto \mathcal{M}$ is an equivalence of abelian categories X-qcoh $\simeq \mathbb{C}$ -Comod. Here X-qcoh denotes the abelian category of quasi-coherent sheaves on X.

Remark 2.6. A quasi-coherent sheaf F on a scheme X is said to be *flat* if the tensor product functor $F \otimes_{\mathcal{O}_X} - : X - \mathsf{qcoh} \longrightarrow X - \mathsf{qcoh}$ is exact, or equivalently, the $\mathcal{O}(U)$ -module F(U) is flat for every affine open subscheme $U \subset X$. It suffices to check the latter condition for the affine open subschemes $U = U_\alpha$ from a given affine open covering $X = \bigcup_\alpha U_\alpha$ of the scheme X.

Now let X be a quasi-compact semi-separated scheme. Then it follows that, in the context of Example 2.5, a quasi-coherent sheaf on X is flat if and only if the corresponding \mathcal{C} -comodule is A-flat.

By [25, Section 2.4] or [12, Lemma A.1], any quasi-coherent sheaf on a quasicompact semi-separated scheme X is a quotient sheaf of a flat quasi-coherent sheaf. Therefore, the corings \mathcal{C} appearing in Example 2.5 also have the property that any \mathcal{C} -comodule is a quotient comodule of an A-flat \mathcal{C} -comodule.

3. A-FLAT C-COMODULES AS DIRECTED COLIMITS

The following theorem describing A-flat C-comodules is the first main result of this paper. In view of Example 2.5 and Remark 2.6, it is a generalization of the assertion of [47, Theorem 2.4] for quasi-compact semi-separated schemes.

Theorem 3.1. Let \mathcal{C} be a coring over an associative ring A. Then the category \mathcal{C} -Comod_{A-flat} of A-flat left \mathcal{C} -comodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of \mathcal{C} -Comod_{A-flat} are precisely all the A-countably presentable A-flat \mathcal{C} -comodules. Consequently, every A-flat \mathcal{C} -comodule is an \aleph_1 -directed colimit of A-countably presentable A-flat \mathcal{C} -comodules.

Proof. All A-countably presentable C-comodules are \aleph_1 -presentable as objects of C-Comod by Lemma 2.3(c). Since the full subcategory C-Comod_{A-flat} is closed under directed colimits in C-Comod, it follows that all A-countably presentable A-flat C-comodules are \aleph_1 -presentable in C-Comod_{A-flat}.

The full assertion of the theorem is obtained by applying Theorems 1.7 and 1.8 for $\kappa = \aleph_1$ and $\lambda = \aleph_0$. One starts with the category $\mathsf{K} = A - \mathsf{Mod}_{\mathsf{flat}}$, the category

L = A-Mod, and the pair of parallel functors $F, G: K \Rightarrow L$ given by the rules F(M) = M and $G(M) = \mathbb{C} \otimes_A M$ for all $M \in A-Mod_{flat}$. Then the inserter category D is the category of all "noncoassociative, noncounital" A-flat left C-comodules, i. e., flat left A-modules \mathcal{D} endowed with a left A-module morphism $\nu: \mathcal{D} \longrightarrow \mathbb{C} \otimes_A \mathcal{D}$.

The category $\mathsf{K} = A - \mathsf{Mod}_{\mathsf{flat}}$ is \aleph_1 -accessible by Lemma 1.2; and the abelian category $A - \mathsf{Mod}$ is locally finitely presentable, hence locally κ -presentable for every regular cardinal κ [1, Remark 1.20]. Theorem 1.7 is applicable (notice that the functor F takes \aleph_1 -presentable objects to \aleph_1 -presentable objects, as it must; while the functor G does not, and it does not have to). The theorem tells us that the category D is \aleph_1 -accessible, and provides a description of the \aleph_1 -presentable objects of D .

Now let M be the Cartesian product of categories $M = A-Mod \times A-Mod_{flat}$. The functor $F: D \longrightarrow M$ assigns to a noncoassociative, noncounital A-flat C-comodule \mathcal{D} the pair of A-modules $(\mathcal{D}, \mathcal{D})$. The functor $G: D \longrightarrow M$ assigns to \mathcal{D} the pair of A-modules $(\mathcal{C} \otimes_A \mathcal{C} \otimes_A \mathcal{D}, \mathcal{D})$. The natural transformation $\phi: F \longrightarrow G$ assigns to \mathcal{D} the pair of compositions $(\mu \otimes id_{\mathcal{D}}) \circ \nu: \mathcal{D} \longrightarrow \mathcal{C} \otimes_A \mathcal{D} \longrightarrow \mathcal{C} \otimes_A \mathcal{C} \otimes_A \mathcal{D}$ and $(\epsilon \otimes id_D) \circ \nu: \mathcal{D} \longrightarrow \mathcal{C} \otimes_A \mathcal{D} \longrightarrow \mathcal{D}$. The natural transformation $\psi: F \longrightarrow G$ assigns to \mathcal{D} the pair of maps consisting of the composition $(id_{\mathcal{C}} \otimes \nu) \circ \nu: \mathcal{D} \longrightarrow \mathcal{C} \otimes_A \mathcal{D} \longrightarrow$ $\mathcal{C} \otimes_A \mathcal{C} \otimes_A \mathcal{D}$ and the identity map $id_{\mathcal{D}}: \mathcal{D} \longrightarrow \mathcal{D}$. Then the equifier category E is the category of coassociative, counital A-flat left C-comodules C-Comod_A-flat.

Proposition 1.4 tells us that the category M is \aleph_1 -accessible, and describes its full subcategory of \aleph_1 -presentable objects. Theorem 1.8 is applicable (once again, the functor F takes \aleph_1 -presentable objects to \aleph_1 -presentable objects, while the functor Gdoes not). The theorem tells us that the category $\mathsf{E} = \mathsf{C}-\mathsf{Comod}_{A-\mathsf{flat}}$ is \aleph_1 -accessible, and provides the desired description of the \aleph_1 -presentable objects of E . \Box

Remark 3.2. Replacing the original category $K = A-Mod_{flat}$ by K = A-Mod (and the category $M = A-Mod \times A-Mod_{flat}$ by $M = A-Mod \times A-Mod$) in the proof of Theorem 3.1, one obtains a proof of the assertion that the category of all comodules C-Comod is \aleph_1 -accessible for any coring C. In fact, all colimits exist in C-Comodby Lemma 2.1(a); so the additive category C-Comod is even locally \aleph_1 -presentable. Furthermore, Theorems 1.7 and 1.8 tell us that the \aleph_1 -presentable objects of C-Comodare precisely all the A-countably presentable C-comodules. So any C-comodule is an \aleph_1 -directed colimit of A-countably presentable C-comodules. Thus we obtain a much more precise version of [27, Theorem 10].

Let us also provide a description of arbitrary complexes of A-flat \mathcal{C} -comodules as directed colimits.

Proposition 3.3. Let C be a coring over an associative ring A. Then the category $Com(C-Comod_{A-flat})$ of complexes of A-flat left C-comodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of $Com(C-Comod_{A-flat})$ are precisely all the complexes of A-countably presentable A-flat C-comodules. Consequently, every complex of A-flat C-comodules is an \aleph_1 -directed colimit of complexes of A-countably presentable A-flat C-comodules. *Proof.* All complexes of A-countably presentable C-comodules are \aleph_1 -presentable as objects of Com(C-Comod) by Lemma 2.3(d). Since the full subcategory Com(C-Comod_{A-flat}) is closed under directed colimits in Com(C-Comod), it follows that all complexes of A-countably presentable A-flat C-comodules are \aleph_1 -presentable in Com(C-Comod_{A-flat}).

The full assertion of the proposition is obtained by combining the results of Theorems 3.1 and 1.9. All one needs to do is to produce a suitable \aleph_1 -presented \mathbb{Z} -linear category D describing complexes in additive categories. This simple construction can be found in [38, proof of Corollary 10.4].

Remark 3.4. Similarly to the proof of Proposition 3.3, it follows from Remark 3.2 that the category Com(C-Comod) of arbitrary complexes of C-comodules is locally \aleph_1 -presentable. Furthermore, the \aleph_1 -presentable objects of Com(C-Comod) are precisely all the complexes of A-countably presentable C-comodules. So any complex of C-comodules in an \aleph_1 -directed colimit of complexes of A-countably presentable C-countably presentable C-comodules. Alternatively, as the categories involved are locally presentable rather than only accessible, it suffices to use a suitable additive version of [19, Theorem 1.2] instead of Theorem 1.9 in this proof.

4. EXACT SEQUENCES OF A-FLAT C-COMODULES AS DIRECTED COLIMITS

We start with three module-theoretic lemmas.

Lemma 4.1. Let A be an associative ring. Then the kernel of any surjective morphism from a countably presentable flat A-module to a countably presentable flat A-module is a countably presentable (flat) A-module.

Proof. Notice that the kernel of a surjective morphism from a countably presentable module to a countably presentable module need not be countably presentable in general (unless the ring is countably coherent). The lemma claims that for *flat* modules it is.

Let F be a countably presentable flat A-module. By Lemma 1.3, there exists a countable directed diagram of finitely generated projective A-modules $P_0 \longrightarrow P_1 \longrightarrow P_2 \longrightarrow \cdots$ such that $F = \varinjlim_{n \ge 0} P_n$. Hence we have a short exact sequence of A-modules $0 \longrightarrow \bigoplus_{n=0}^{\infty} P_n \longrightarrow \bigoplus_{n=0}^{\infty} P_n \longrightarrow F \longrightarrow 0$. Put $P' = P'' = \bigoplus_{n=0}^{\infty} P_n$. Now let $0 \longrightarrow H \longrightarrow G \longrightarrow F \longrightarrow 0$ be a short exact sequence of A-modules.

Now let $0 \longrightarrow H \longrightarrow G \longrightarrow F \longrightarrow 0$ be a short exact sequence of A-modules. Since the A-module P' is projective, the A-module morphism $P' \longrightarrow F$ lifts over the surjective A-module morphism $G \longrightarrow F$. We obtain a commutative diagram of a morphism of short exact sequences acting by the identity on the rightmost terms



Consequently, there is a short exact sequence of A-modules $0 \longrightarrow P'' \longrightarrow H \oplus P' \longrightarrow G \longrightarrow 0$. Now we notice that the A-module P'' is countably presentable. If the A-module G is countably presentable as well, then it follows that the A-module $H \oplus P'$ is countably presentable (as the class of countably presentable A-modules is closed under extensions). Consequently, the A-module H is countably presentable as a direct summand of a countably presentable A-module.

For a generalization of Lemma 4.1 to arbitrary regular cardinals κ , see [38, Corollary 10.12] or [39, Corollary 4.7].

Lemma 4.2. Let A be an associative ring. Then the abelian category of three-term complexes of left A-modules $C^1 \longrightarrow C^2 \longrightarrow C^3$ is locally finitely presentable, and consequently, locally \aleph_1 -presentable. The \aleph_1 -presentable objects of this category are precisely all the three-term complexes of countably presentable A-modules.

Proof. The assertion holds for any regular cardinal κ in lieu of \aleph_1 . See [39, Proposition 1.1(c)] for a much more general result.

Lemma 4.3. For any ring A, the category of short exact sequences of flat A-modules is \aleph_1 -accessible. The \aleph_1 -presentable objects of this category are precisely all the short exact sequences of countably presentable flat A-modules.

Proof. Once again, the assertion holds for any regular cardinal κ . See [38, Corollary 10.13] or [39, Propositions 1.1(c) and 4.6].

Recall from the discussion in Section 2 that the category C-Comod need not be abelian for a coring C over an associative ring A in general; but one can still speak about *short exact sequences of A-flat C-comodules*. The following proposition describes such short exact sequences.

Proposition 4.4. Let C be a coring over an associative ring A. Then the category of short exact sequences of A-flat C-comodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of this category are precisely all the short exact sequences of A-countably presentable A-flat C-comodules. Consequently, every short exact sequence of A-flat C-comodules is an \aleph_1 -directed colimit of short exact sequences of A-countably presentable A-flat C-comodules.

Proof. Similar to the proof of Theorem 3.1, with the following changes. Use the category of short exact sequences of flat A-modules (from Lemma 4.3) in the role of K, the category of three-term complexes of A-modules (from Lemma 4.2) in the role of L, and put $M = L \times K$. The functors and natural transformations are given by the same formulas as in the proof of Theorem 3.1, applied to short exact sequences or three-term complexes instead of (co)modules.

The assertion of Proposition 4.4 (together with Theorem 3.1) can be restated by saying that the exact category of A-flat C-comodules $C-Comod_{A-flat}$ from Lemma 2.2 is a *locally* \aleph_1 -*coherent exact category* in the sense of [39, Section 1].

We refer to [7, Section 10] for the definition of an acyclic complex in an exact category. Acyclic complexes in the exact category $C-Comod_{A-flat}$ are called A-pure

acyclic complexes of A-flat C-comodules. Equivalently, the A-pure acyclic complexes of A-flat C-comodules are the complexes of C-comodules whose underlying complexes of A-modules are acyclic with flat A-modules of cocycles.

Corollary 4.5. Let \mathcal{C} be a coring over an associative ring A. Then the category of A-pure acyclic complexes of A-flat \mathcal{C} -comodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of this category are precisely all the A-pure acyclic complexes of A-countably presentable A-flat \mathcal{C} -comodules. Consequently, every A-pure acyclic complex of A-flat \mathcal{C} -comodules is an \aleph_1 -directed colimit of A-pure acyclic complexes of A-countably presentable A-flat \mathcal{C} -comodules.

Proof. This is provable similarly to Proposition 4.4 (using [38, Corollary 10.14] instead of Lemma 4.3, and the obvious version of Lemma 4.2 for unbounded complexes). This is akin to the argument used in [47, proof of Theorem 4.2]. But we prefer to spell out an argument deducing the corollary from the proposition.

The argument is similar to [38, proof of Corollary 10.14]. An A-pure acyclic complex of A-flat C-comodules \mathcal{F}^{\bullet} is the same thing as a collection of short exact sequences of A-flat C-comodules $0 \longrightarrow \mathcal{F}^n \longrightarrow \mathcal{F}^n \longrightarrow \mathcal{H}^n \longrightarrow 0$ together with a collection of isomorphisms of C-comodules $\mathcal{H}^n \simeq \mathcal{G}^{n+1}$, $n \in \mathbb{Z}$. Therefore, the category of A-pure acyclic complexes of A-flat C-comodules can be constructed from the category of short exact sequences of A-flat C-comodules using Cartesian products (as in Proposition 1.4) and the isomorpher construction (as in Theorem 1.6).

Specifically, take $\kappa = \aleph_1$ and $\lambda = \aleph_0$. Denote by K the Cartesian product of copies of the category of short exact sequences of A-flat C-comodules (from Proposition 4.4), indexed by all the integers $n \in \mathbb{Z}$. Denote by L the Cartesian product of copies of the category of A-flat C-comodules C-Comod_{A-flat}, also indexed by $n \in \mathbb{Z}$. Let $F_1: \mathsf{K} \longrightarrow \mathsf{L}$ be the functor taking a collection of short exact sequences $(0 \rightarrow \mathfrak{G}^n \rightarrow \mathfrak{F}^n \rightarrow \mathfrak{H}^n \rightarrow 0)_{n \in \mathbb{Z}}$ to the collection of flat comodules $(\mathfrak{H}^n)_{n \in \mathbb{Z}}$, and let $F_2: \mathsf{K} \longrightarrow \mathsf{L}$ be the functor taking the same collection of short exact sequences to the collection of flat comodules $(\mathfrak{G}^{n+1})_{n \in \mathbb{Z}}$. Then the isomorpher category C is the category of A-pure acyclic complexes of A-flat C-comodules.

Theorem 3.1 and Proposition 4.4 together with Proposition 1.4 tell us that the categories K and L are \aleph_1 -accessible, and describe their full subcategories of \aleph_1 -presentable objects. In view of these descriptions, Theorem 1.6 is applicable, telling us that the isomorpher category C is \aleph_1 -accessible, and describing its full subcategory of \aleph_1 -presentable objects. \Box

5. Cotorsion Periodicity for Comodules

In this section we deduce certain conditional results, based on the techniques of [47, Section 8] and generalizing some results from [47, Section 9].

Let \mathcal{C} be a coring over a ring A such that \mathcal{C} is a flat right A-module (so the category of left \mathcal{C} -comodules is abelian by Lemma 2.1(b)). Consider the following condition on the coring \mathcal{C} :

(*) Every left C-comodule is a quotient comodule of an A-flat left C-comodule.

Two classes of corings \mathcal{C} known to satisfy (*) are described in Remark 2.6 and Lemma 2.4 (geometrically, these correspond to quasi-compact semi-separated schemes and a certain kind of smooth stacks). We are not aware of any example of a coring \mathcal{C} over A that is flat as a (left and) right A-module but does not satisfy (*). We will also need a second condition:

(**) Let \mathcal{M} be a left \mathcal{C} -comodule such that the underlying A-module of \mathcal{M} has finite projective dimension as an object of the abelian category A-Mod. Then \mathcal{M} has a finite projective dimension as an object of the abelian category \mathcal{C} -Comod.

Remark 5.1. The corings C corresponding to quasi-compact semi-separated schemes X as per Example 2.5 satisfy (**). This is the result of [47, Theorem 5.3].

On the other hand, when A = k is a field and \mathcal{C} is a coalgebra over k, the condition (**) asks the comodule category \mathcal{C} -Comod to have finite homological dimension. This, of course, does *not* hold in general. For example, if \mathcal{C} is a finite-dimensional coalgebra over k, then the category of \mathcal{C} -comodules is equivalent to the category of modules over the k-vector space dual finite-dimensional algebra, \mathcal{C} -Comod $\simeq \mathcal{C}^*$ -Mod. Taking \mathcal{C}^* to be a finite-dimensional k-algebra of infinite global dimension, one obtains an example when (**) does not hold.

Informally speaking, (**) means that "the coring \mathcal{C} has finite homological dimension relative to the ring A". In the context of algebraic geometry, one can expect this to be satisfied for many stacks in characteristic zero, such as the quotient stacks X/G of algebraic varieties X by actions of algebraic groups G (because the homological dimension of the category of algebraic group representations does not exceed the dimension of the group, in characteristic zero). On the other hand, in prime characteristic p, the quotient of a point by the action of a finite group of the order divisible by p does not satisfy (**), as the previous paragraph explains.

See Remarks 5.7 and 6.6 below for a further discussion.

We denote by $\operatorname{Ext}^*_{\mathbb{C}}(-,-)$ the Ext groups in the abelian category \mathbb{C} -Comod. A left \mathbb{C} -comodule \mathbb{B} is said to be *cotorsion* [29, Section C.2] if $\operatorname{Ext}^1_{\mathbb{C}}(\mathcal{F}, \mathbb{B}) = 0$ for all A-flat left \mathbb{C} -comodules \mathcal{F} . Let $\mathbb{B} = \mathbb{C}$ -Comod^{cot} denote the full subcategory of cotorsion \mathbb{C} -comodules in the abelian category $\mathbb{C} = \mathbb{C}$ -Comod. Obviously, the full subcategory \mathbb{C} -Comod^{cot} is closed under extensions in \mathbb{C} -Comod, so it inherits an exact category structure from the abelian exact structure of \mathbb{C} -Comod.

We will say that a class of objects F in an abelian category C is *resolving* if F is closed under extensions and kernels of epimorphisms in C, and every object of C is a quotient object of an object from F.

The following corollary is our version of [47, Corollary 5.6 and Lemma 9.1]. Part (a) was already mentioned in the proof of Theorem 3.1.

Corollary 5.2. Let \mathcal{C} be a coring over a ring A.

(a) The class of all A-flat left C-comodules is closed under directed colimits in the additive category C-Comod.

(b) Assume that the condition (*) above on the coring \mathcal{C} is satisfied. Then the class of all A-flat left \mathcal{C} -comodules is resolving in the abelian category \mathcal{C} -Comod. Consequently, one has $\operatorname{Ext}^n_{\mathcal{C}}(\mathcal{F}, \mathcal{B}) = 0$ for all A-flat left \mathcal{C} -comodules \mathcal{F} , all cotorsion left \mathcal{C} -comodules \mathcal{B} , and all integers $n \geq 1$.

(c) Assume that the condition (**) above on the coring \mathcal{C} is satisfied. Then any A-flat left \mathcal{C} -comodule is an \aleph_1 -directed colimit of A-flat \mathcal{C} -comodules having finite projective dimensions as objects of \mathcal{C} -Comod.

Proof. Part (a) follows from Lemma 2.1(a), since the class of all flat modules is closed under directed colimits in A-Mod. In the context of part (b), condition (*) presumes that C is a flat right A-module. Then the class of all A-flat left C-comodules is closed under extensions and kernels of epimorphisms in C-Comod by Lemma 2.1(b), since the class of all flat modules is closed under extensions and kernels of epimorphisms in A-Mod. Every left C-comodule is an epimorphic image of an A-flat left C-comodule by the condition (*). The final assertion of (b) is provided by [47, Lemma 6.1].

In part (c), all countably presentable flat A-modules have projective dimensions ≤ 1 in A-Mod by Lemma 1.3. Under condition (**), it follows that all A-countably presentable A-flat C-comodules have finite projective dimensions in C-Comod. So it remains to refer to Theorem 3.1.

Asume that \mathcal{C} is a flat right A-module. Let $\mathsf{F} = \mathcal{C}-\mathsf{Comod}_{A-\mathsf{flat}}$ denote the full subcategory of all A-flat \mathcal{C} -comodules in the abelian category $\mathsf{C} = \mathcal{C}-\mathsf{Comod}$. Since the full subcategory F is closed under extensions in $\mathcal{C}-\mathsf{Comod}$, it inherits an exact category structure from the abelian exact structure of $\mathcal{C}-\mathsf{Comod}$. Since F is also closed under directed colimits in C , and the directed colimit functors are exact in $\mathsf{C} = \mathcal{C}-\mathsf{Comod}$, it follows that F is an *exact category with exact directed colimits* in the sense of [47, Section 7], i. e., all directed colimits exist and directed colimits of admissible short exact sequences are admissible short exact sequences in F .

Let C be an abelian category and $B \subset C$ be a class of objects. An object $M \in C$ is called B-*periodic* if there exists a short exact sequence $0 \longrightarrow M \longrightarrow B \longrightarrow M \longrightarrow 0$ in C with M as both the rightmost and the leftmost term, and a middle object $B \in B$.

The following theorem summarizes the technology of proving cotorsion periodicity theorems going back to [2] and developed in its present form in [47].

Theorem 5.3. Let C be an abelian category with coproducts and $F \subset C$ be a resolving subcategory closed under directed colimits. Assume that the directed colimit functors in F take admissible short exact sequences to admissible short exact sequences (in the exact category structure on F inherited from the abelian exact structure of C, i. e., the short sequences in F that are exact in C). Assume further that the category F coincides with the directed colimit closure of its full subcategory consisting of all the objects of F that have finite projective dimensions in C. Let $B = F^{\perp_1} \subset C$ be the full subcategory of all objects $B \in C$ such that $\operatorname{Ext}^1_C(F, B) = 0$ for all $F \in F$. Then any B-periodic object of C belongs to B.

Proof. This is a particular case of [47, Corollary 8.2].

Now we can prove our cotorsion periodicity theorem for comodules.

Theorem 5.4. Let C be a coring over an associative ring A. Assume that both the conditions (*) and (**) on the coring C are satisfied. Then any cotorsion-periodic left C-comodule is cotorsion.

Proof. This is our generalization of [47, Theorem 9.2]. The assumptions of Theorem 5.3 are satisfied by Corollary 5.2 (cf. the discussion above). \Box

Corollary 5.5. Let \mathcal{C} be a coring over an associative ring A. Assume that both the conditions (*) and (**) on the coring \mathcal{C} are satisfied. Let \mathcal{B}^{\bullet} be an acyclic complex in \mathcal{C} -Comod whose terms \mathcal{B}^n are cotorsion left \mathcal{C} -comodules for all $n \in \mathbb{Z}$. Then the \mathcal{C} -comodules of cocycles of the complex \mathcal{B}^{\bullet} are also cotorsion.

Proof. This is our generalization of [47, Corollary 9.4]. To deduce it from Theorem 5.4, we apply [47, Proposition 8.4]. It remains to check the assumptions of that proposition from [47].

Indeed, the abelian category C-Comod is Grothendieck by Lemma 2.1(b), so it has infinite products. The full subcategory B of cotorsion left C-comodules is closed under infinite products in the abelian category C = C-Comod by [10, Corollary 8.3] or [11, Corollary A.2]. It is also obvious that B is closed under direct summands in C.

Finally, any left C-comodule is a quotient comodule of A-flat C-comodule by the condition (*), and any A-flat C-comodule is a quotient comodule of a coproduct of A-countably presentable A-flat left C-comodules by Theorem 3.1. The underlying A-modules of the C-comodules of the latter kind have projective dimension ≤ 1 by Lemma 1.3. Such C-comodules have finite projective dimensions in C-Comod by the condition (**). Therefore, every left C-comodule is a quotient comodule of an A-flat left C-comodule having finite projective dimension in C-Comod. It remains to recall that $\operatorname{Ext}^n_{\mathbb{C}}(\mathcal{F}, \mathcal{B}) = 0$ for any A-flat left C-comodule \mathcal{F} , any left C-comodule $\mathcal{B} \in \mathsf{B}$, and all integers $n \geq 1$ by Corollary 5.2(b).

Corollary 5.6. Let \mathcal{C} be a coring over an associative ring A. Assume that both the conditions (*) and (**) on the coring \mathcal{C} are satisfied. Then the inclusion of exact/abelian categories $\mathcal{C}\text{-}\mathsf{Comod}^{\mathsf{cot}} \longrightarrow \mathcal{C}\text{-}\mathsf{Comod}$ induces an equivalence of their unbounded derived categories,

$$D(C-Comod^{cot}) \simeq D(C-Comod).$$

Proof. This is our generalization of [47, Corollary 9.7]. To deduce it from Corollary 5.5, it suffices to refer to [47, Proposition 9.6]. \Box

Remark 5.7. The assumption of condition (**) cannot be dropped in any one of the assertions of Theorem 5.4, Corollary 5.5, or Corollary 5.6. It suffices to choose a field A = k and a finite-dimensional coalgebra \mathcal{C} over k whose dual algebra \mathcal{C}^* is a Frobenius algebra of infinite global dimension (e. g., the algebra of dual numbers $\mathcal{C}^* = k[\epsilon]/(\epsilon^2)$). Then C-Comod = C*-Mod (cf. Remark 5.1), all C-comodules are A-flat, and the cotorsion C-comodules are the injective C-comodules, or equivalently the injective C*-modules. Pick any C*-module of infinite projective/injective dimension; then its two-sided projective-injective resolution is an unbounded acyclic complex of injective C-comodules refuting the conclusions of both Corollaries 5.5 and 5.6. Then it is clear from [9, proof of Proposition 7.6] or [15, Proposition 2] that the conclusion of Theorem 5.4 does not hold for the coalgebra/coring C, either.

6. EXACT SEQUENCES OF A-FLAT C-COMODULES II

In this section we also deduce some results conditional upon the assumptions (*) and (**) from Section 5 on a coring \mathcal{C} .

We start with an abstract category-theoretic discussion of absolutely acyclic complexes in exact categories of finite homological dimension. Given an additive category E, we denote by K(E) the triangulated homotopy category of (unbounded cochain complexes in) E.

Let E be an exact category. Any (termwise admissible) short exact sequence of complexes in E can be viewed as a bicomplex with three rows, so one can construct its total complex. A complex in E is said to be *absolutely acyclic* [28, Sections 2 and 4], [29, Section A.1], [36, Section 5.1] if it belongs to the minimal thick subcategory of the homotopy category K(E) containing the totalizations of short exact sequences in E. In other words, a complex in E is absolutely acyclic if and only if it can be constructed as a homotopy direct summand of a complex obtained from the totalizations of short exact sequences of complexes in E by passing to cones of closed morphisms of complexes a finite number of times.

Equivalently, the full subcategory of absolutely acyclic complexes in Com(E) can be constructed as the closure of the full subcategory of contractible complexes under extensions (in the termwise exact structure on Com(E)) and direct summands. This is essentially shown in [46, Proposition 8.12].

Denote by $\operatorname{Ext}_{\mathsf{E}}^*(-,-)$ the Yoneda Ext functor in an exact category E . The exact category E is said to have homological dimension $\leq d$ (where $d \geq -1$ is an integer) if $\operatorname{Ext}_{\mathsf{E}}^n(X,Y) = 0$ for all $X, Y \in \mathsf{E}$ and n > d.

Lemma 6.1. In an exact category of finite homological dimension, all acyclic complexes are absolutely acyclic.

Proof. This is [28, Lemma 2.1].

Proposition 6.2. Let E be an exact category of homological dimension $\leq d$, where d is a finite integer. Then any absolutely acyclic complex in E is a direct summand of the totalization of a (d+2)-term exact complex $0 \longrightarrow E_d^{\bullet} \longrightarrow E_{d-1}^{\bullet} \longrightarrow \cdots \longrightarrow E_0^{\bullet} \longrightarrow E_{-1}^{\bullet} \longrightarrow 0$ of complexes in E .

Proof. This result is a particular case of a similar assertion for *exact DG-categories*, which is essentially proved in [36, proof of Proposition 8.8]. More precisely, as a particular case of [36, proof of Proposition 8.8] (corresponding to [36, Examples 4.39 and 6.1(4)]) one obtains the claim that any absolutely acyclic complex A^{\bullet} in E is a *homotopy direct summand* of the totalization T^{\bullet} of a (d+2)-term exact complex of complexes in E. Here a "homotopy direct summand" means a direct summand in the

homotopy category K(E). It remains to show that a homotopy direct summand can be turned into a strict direct summand.

Now we have a pair of closed morphisms of complexes $i: A^{\bullet} \longrightarrow T^{\bullet}$ and $p: T^{\bullet} \longrightarrow A^{\bullet}$ such that the composition $pi: A^{\bullet} \longrightarrow A^{\bullet}$ is homotopic to the identity morphism id: $A^{\bullet} \longrightarrow A^{\bullet}$. It follows that the difference $pi - \text{id}: A^{\bullet} \longrightarrow A^{\bullet}$ factorizes as $A^{\bullet} \longrightarrow C^{\bullet} \longrightarrow A^{\bullet}$, where C^{\bullet} is the cone of identity morphism id: $A^{\bullet} \longrightarrow A^{\bullet}$ and $A^{\bullet} \longrightarrow C^{\bullet}$ is the natural closed morphism, while $C^{\bullet} \longrightarrow A^{\bullet}$ is some arbitrary closed morphism. Hence the identity morphism id: $A^{\bullet} \longrightarrow A^{\bullet}$ factorizes through the direct sum $T^{\bullet} \oplus C^{\bullet}$ as a morphism in the category of complexes in E. Finally, we point out that C^{\bullet} is the totalization of a 2-term exact complex $0 \longrightarrow A^{\bullet} \longrightarrow A^{\bullet} \longrightarrow 0$ of complexes in E. This takes care of all cases with $d \ge 0$, while in the trivial case of d = -1 we have $\mathsf{E} = 0$ and there is nothing to prove.

Let F be an exact category. A class of objects $G \subset F$ is said to be *self-generating* [47, Section 6] if for any admissible epimorphism $F \longrightarrow H$ in F with $H \in G$ there exists a morphism $G \longrightarrow F$ in F with $G \in G$ such that the composition $G \longrightarrow F \longrightarrow H$ is an admissible epimorphism in F. A class of objects G in F is said to be *self-resolving* [36, Section 7.1], [47, Section 8] if it is self-generating, closed under extensions, and closed under kernels of admissible epimorphisms.

Clearly, any self-resolving full subcategory G in an exact category F interits an exact category structure from F (since G is closed under extensions in F). The following lemma is straightforward and well-known.

Lemma 6.3. Let F be an exact category and $G \subset F$ be a self-resolving full subcategory. Then the inclusion of exact categories $G \longrightarrow F$ induces isomorphisms on the Ext groups,

$$\operatorname{Ext}^n_{\mathsf{G}}(X,Y) \simeq \operatorname{Ext}^n_{\mathsf{F}}(X,Y) \quad \text{for all } X, Y \in \mathsf{G} \text{ and } n \ge 0.$$

Proof. See [20, Section 12] or [29, Proposition A.2.1].

Now we return to comodules over a coring \mathcal{C} over an associative ring A.

Corollary 6.4. Let \mathcal{C} be a coring over an associative ring A such that \mathcal{C} is a flat right A-module.

(a) The full subcategory G of all A-countably presentable A-flat left C-comodules is self-resolving in the exact category F of all A-flat left C-comodules.

(b) If the conditions (*) and (**) on the coring C are satisfied, then the exact category G of A-countably presentable A-flat left C-comodules has finite homological dimension.

Proof. In part (a), the assertion that G is self-generating in F is an easy corollary of Theorem 3.1. Let \mathcal{H} be an A-countably presentable A-flat left C-comodule, \mathcal{F} be an A-flat C-comodule, and $\mathcal{F} \longrightarrow \mathcal{H}$ be a surjective C-comodule morphism. By Theorem 3.1, there exists an \aleph_1 -directed poset Ξ and a Ξ -indexed diagram of A-countably presentable A-flat C-comodules $(\mathcal{G}_{\xi})_{\xi \in \Xi}$ for which $\mathcal{F} = \varinjlim_{\xi \in \Xi} \mathcal{G}_{\xi}$. Then it is clear that there exists an index $\xi_0 \in \Xi$ such that the composition $\mathcal{G}_{\xi_0} \longrightarrow \mathcal{F} \longrightarrow \mathcal{H}$ is surjective. The full subcategory G is closed under kernels of surjective homomorphisms in

F by Lemma 4.1, and the assertion that G is closed under extensions in F is provable similarly to Corollary 5.2(b).

In part (b), the full subcategory G is self-resolving in F by part (a), and the full subcategory F is resolving in the abelian category C = C-Comod by Corollary 5.2(b). Consequently, G is self-resolving in C. The underlying A-modules of all the C-comodules from G have projective dimensions at most 1 by Lemma 1.3, hence the objects of G has finite projective dimensions in C by condition (**). Moreover, the latter projective dimensions must be uniformly bounded by some fixed integer d, since the class of objects G is closed under countable coproducts in C. Finally, Lemma 6.3 tells us that the Ext groups computed in G and in C agree. Thus the homological dimension of the exact category G cannot exceed d.

The following theorem is a comodule version of [14, Theorem 2.4(1) \Leftrightarrow (3)] or [26, Theorem 8.6(ii) \Leftrightarrow (iii)]. It should be also compared to [46, Corollary 0.5 or Proposition 8.13]. In the latter context, one can say that Theorem 6.5 means that (under the conditions (*) and (**)) for the exact category of A-flat C-comodules, the derived category coincides with the (suitably defined) coderived category.

Theorem 6.5. Let \mathcal{C} be a coring over an associative ring A. Assume that both the conditions (*) and (**) from Section 5 on the coring \mathcal{C} are satisfied. Then there exists a finite integer $d \ge 0$ such that the following classes of complexes of left \mathcal{C} -comodules coincide:

- (1) all A-pure acyclic complexes of A-flat C-comodules;
- (2) the closure of the class of all contractible complexes of A-countably presentable A-flat C-comodules under extensions and directed colimits;
- (3) all \aleph_1 -directed colimits of totalizations of (d+2)-term exact complexes of complexes of A-countably presentable A-flat C-comodules.

Proof. One can easily see that the totalization of a (d + 2)-term exact complex of complexes is a finitely iterated extension of d + 1 cones of identity endomorphisms of complexes. Furthermore, \aleph_1 -directed colimits are a subclass of directed colimits, the class of all A-pure acyclic complexes of A-flat C-comodules is closed under extensions and directed colimits, and the contractible complexes of A-flat C-comodules are A-pure acyclic. This proves the inclusions $(3) \subset (2) \subset (1)$.

Conversely, to prove that $(1) \subset (3)$, recall that all A-pure acyclic complexes of A-flat C-comodules are \aleph_1 -directed colimits of A-pure acyclic complexes of A-countably presentable A-flat C-comodules by Corollary 4.5. The exact category G of A-countably presentable A-flat C-comodules has finite homological dimension by Corollary 6.4(b). Hence the A-pure acyclic complexes of A-countably presentable A-flat C-comodules are absolutely acyclic by Lemma 6.1, and it remains to refer to Proposition 6.2.

Remark 6.6. Without the assumption (**), the conclusion of Theorem 6.5 certainly does *not* hold (cf. Remarks 5.1 and 5.7). In fact, for a coalgebra \mathcal{C} over a field A = k, if the global dimension of \mathcal{C} is infinite, then the coderived category of \mathcal{C} -comodules is usually quite different from their derived category [37, Section 7].

7. Preliminaries on Topological Rings and Contramodules

The material of this section goes back to [28, Remark A.3], [29, Appendix E], [30, Section 2.1], and [43, Section 5]. For more recent references, see [33, Section 2] or [32, Section 2]. The exposition in [44, Sections 6–7] may be the most accessible one for a beginner. In this paper, we are mostly interested in contramodules over complete, separated topological associative rings with a countable base of neighborhoods of zero consisting of open two-sided ideals.

Let \mathfrak{A} be a topological abelian group where open subgroups form a base of neighborhoods of zero. Consider the natural map to the directed limit $\lambda_{\mathfrak{A}} \colon \mathfrak{A} \longrightarrow \varprojlim_{\mathfrak{U} \subset \mathfrak{A}} \mathfrak{A}/\mathfrak{U}$, where \mathfrak{U} ranges over the open subgroups of \mathfrak{A} . The topological group \mathfrak{A} is said to be *separated* if the map $\lambda_{\mathfrak{A}}$ is injective, and *complete* if $\lambda_{\mathfrak{A}}$ is surjective.

Let \mathfrak{R} be a complete, separated topological ring with a base of neighborhoods of zero consisting of open right ideals. A right \mathfrak{R} -module \mathbb{N} is said to be *discrete* if the action map $\mathbb{N} \times \mathfrak{R} \longrightarrow \mathbb{N}$ is continuous in the given topology of \mathfrak{R} and the discrete topology on \mathbb{N} . Equivalently, a right \mathfrak{R} -module \mathbb{N} is discrete if, for every element $x \in \mathbb{N}$, the annihilator of x in \mathfrak{R} is an open right ideal. The full subcategory of discrete right \mathfrak{R} -modules $\mathsf{Discr}-\mathfrak{R} \subset \mathsf{Mod}-\mathfrak{R}$ is a Grothendieck abelian category. The inclusion functor $\mathsf{Discr}-\mathfrak{R} \longrightarrow \mathsf{Mod}-\mathfrak{R}$ is exact and preserves all colimits.

Given an abelian group A and a set X, we use $A[X] = A^{(X)}$ as a notation for the direct sum of X copies of A. The elements of A[X] are interpreted as finitely supported formal linear combinations $\sum_{x \in X} a_x x$ of elements of X with the coefficients $a_x \in A$ (where $a_x = 0$ for all but a finite subset of indices $x \in X$).

Let \mathfrak{A} be a complete, separated topological abelian group with a base of neighborhoods of zero consisting of open subgroups. Given a set X, we denote by $\mathfrak{A}[[X]]$ the directed limit of abelian groups $\lim_{\mathfrak{U} \subset \mathfrak{A}} (\mathfrak{A}/\mathfrak{U})[X]$ (where \mathfrak{U} ranges over the open subgroups of \mathfrak{A}). The elements of $\mathfrak{A}[[X]]$ are interpreted as infinite formal linear combinations $\sum_{x \in X} a_x x$ with the families of coefficients $a_x \in \mathfrak{A}$ converging to zero in the topology of \mathfrak{A} . The latter condition means that, for every open subgroup $\mathfrak{U} \subset \mathfrak{A}$, one has $a_x \in \mathfrak{U}$ for all but a finite subset of indices $x \in X$.

Given a map of sets $X \longrightarrow Y$, the induced (pushforward) map $\mathfrak{A}[[f]]: \mathfrak{A}[[X]] \longrightarrow \mathfrak{A}[[Y]]$ is defined by the rule $\sum_{x \in X} a_x x \longmapsto \sum_{y \in Y} b_y y$, where $b_y = \sum_{x \in X}^{f(x)=y} a_x$ for every $y \in Y$. Here the latter infinite sum (defining the element $b_y \in \mathfrak{A}$) is not formal; rather, it is understood as the limit of finite partial sums in the topology of \mathfrak{A} . Due to this construction, the assignment $X \longmapsto \mathfrak{A}[[X]]$ is a covariant functor Sets \longrightarrow Ab from the category of sets to the category of abelian groups. We will mostly view it as an endofunctor on the category of sets, $\mathfrak{A}[[-]]:$ Sets \longrightarrow Sets, ignoring the abelian group structures on the sets $\mathfrak{A}[[X]]$.

Let \mathfrak{R} be a complete, separated topological ring where open right ideals form a base of neighborhoods of zero. Then the functor $\mathfrak{R}[[-]]: \mathsf{Sets} \longrightarrow \mathsf{Sets}$ acquires a natural structure of a *monad* on the category of sets. We refer to [24, Chapter VI] (see also [44, Section 6]) for the background discussion of monads and algebras over monads. Specifically, for any set X, the monad unit $\epsilon_X: X \longrightarrow \mathfrak{R}[[X]]$ is the obvious natural map taking every element $x \in X$ to the formal linear combination $\sum_{y \in X} r_y y$, where $r_x = 1$ and $r_y = 0$ for all $y \neq x$.

The monad multiplication $\phi_X \colon \mathfrak{R}[[\mathfrak{R}[[X]]]] \longrightarrow \mathfrak{R}[[X]]$ is the natural "opening of parentheses" map producing a formal linear combination from a formal linear combination of formal linear combinations. This construction involves taking products of pairs of elements in \mathfrak{R} and infinite sums of zero-convergent families of elements. The conditions that \mathfrak{R} is complete, separated, and open right ideals form a base of neighborhoods of zero guarantee the convergence.

Left \mathfrak{R} -contramodules are defined as algebras over the monad $(\mathfrak{R}[[-]], \epsilon, \phi)$ on the category of sets. We usually prefer to speak of modules (rather than algebras) over monads in this context, as contramodules are more similar to modules over rings than to algebras. So, specifically, a left \mathfrak{R} -contramodule \mathfrak{P} is a set endowed with a *left* contraaction map $\pi : \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}$ satisfying the following contraassociativity and contraunitality axioms. The two compositions $\pi \circ \phi_{\mathfrak{P}}$ and $\pi \circ \mathfrak{R}[[\pi]]$ must be equal to each other,

$$\mathfrak{R}[[\mathfrak{R}[[\mathfrak{P}]]]] \rightrightarrows \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}$$

and the composition $\pi \circ \epsilon_{\mathfrak{P}}$ must be ideal to the identity map $\mathrm{id}_{\mathfrak{P}}$,

 $\mathfrak{P} \longrightarrow \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}.$

We denote the category of left \mathfrak{R} -contramodules by \mathfrak{R} -Contra.

Restricting the contraaction map $\pi: \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}$ to the subset of finitely supported formal linear combinations $\mathfrak{R}[\mathfrak{P}] \subset \mathfrak{R}[[\mathfrak{P}]]$, one constructs the underlying left \mathfrak{R} -module structure on a left \mathfrak{R} -contramodule \mathfrak{P} [44, Sections 6.1–6.2]. Hence the forgetful functor \mathfrak{R} -Contra $\longrightarrow \mathfrak{R}$ -Mod. The category of left \mathfrak{R} -contramodules is abelian. All set-indexed products and coproducts (hence also limits and colimits) exist in \mathfrak{R} -Contra. The forgetful functor \mathfrak{R} -Contra $\longrightarrow \mathfrak{R}$ -Mod is exact, faithful, and preserves infinite products. We use the notation $\operatorname{Hom}^{\mathfrak{R}}(-,-)$ for the groups of morphisms in the abelian category \mathfrak{R} -Contra.

For any set X, the set/abelian group $\mathfrak{R}[[X]]$ has a natural left \mathfrak{R} -contramodule structure with the contraaction map $\pi = \phi_X \colon \mathfrak{R}[[\mathfrak{R}[[X]]]] \longrightarrow \mathfrak{R}[[X]]$. The \mathfrak{R} -contramodules $\mathfrak{R}[[X]]$ are called the *free* left \mathfrak{R} -contramodules. For any left \mathfrak{R} -contramodule \mathfrak{P} , the \mathfrak{R} -contramodule morphisms $\mathfrak{R}[[X]] \longrightarrow \mathfrak{P}$ correspond bijectively to maps of sets $X \longrightarrow \mathfrak{P}$; so there is a natural isomorphism of abelian groups

$$\operatorname{Hom}^{\mathfrak{R}}(\mathfrak{R}[[X]],\mathfrak{P})\simeq \operatorname{Hom}_{\mathsf{Sets}}(X,\mathfrak{P}).$$

There are enough projective objects in the abelian category \Re -Contra; the projective \Re -contramodules are precisely the direct summands of the free ones.

Given a left \mathfrak{R} -contramodule \mathfrak{P} and a closed subgroup $\mathfrak{A} \subset \mathfrak{R}$, we denote by $\mathfrak{A} \prec \mathfrak{P} \subset \mathfrak{P}$ the image of the composition $\mathfrak{A}[[\mathfrak{P}]] \longrightarrow \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}$ of the natural inclusion $\mathfrak{A}[[\mathfrak{P}]] \hookrightarrow \mathfrak{R}[[\mathfrak{P}]]$ with the contraaction map $\pi : \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}$. So $\mathfrak{A} \prec \mathfrak{P}$ is a subgroup in \mathfrak{P} (since π is a left \mathfrak{R} -contramodule morphism, hence in particular an abelian group homomorphism, and $\mathfrak{A}[[\mathfrak{P}]]$ is subgroup in $\mathfrak{R}[[\mathfrak{P}]]$). If $\mathfrak{J} \subset \mathfrak{R}$ is a closed left ideal, then $\mathfrak{J} \prec \mathfrak{P}$ is an \mathfrak{R} -subcontramodule in \mathfrak{P} (since $\mathfrak{J}[[\mathfrak{P}]]$ is an \mathfrak{R} -subcontramodule in the free contramodule $\mathfrak{R}[[\mathfrak{P}]]$).

Let \mathbb{N} be a discrete right \mathfrak{R} -module and \mathfrak{P} be a left \mathfrak{R} -contramodule. Then the *contratensor product* $\mathbb{N} \odot_{\mathfrak{R}} \mathfrak{P}$ is the abelian group constructed as the cokernel of (the difference of) the natural pair of maps

$$\mathcal{N}\otimes_{\mathbb{Z}}\mathfrak{R}[[\mathfrak{P}]]\,
ightarrow\,\mathcal{N}\otimes_{\mathbb{Z}}\mathfrak{P}.$$

Here the first map $\mathcal{N} \otimes_{\mathbb{Z}} \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathcal{N} \otimes_{\mathbb{Z}} \mathfrak{P}$ is induced by the contraaction map $\pi \colon \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathfrak{P}$, while the second map $\mathcal{N} \otimes_{\mathbb{Z}} \mathfrak{R}[[\mathfrak{P}]] \longrightarrow \mathcal{N} \otimes_{\mathbb{Z}} \mathfrak{P}$ is defined by the rule

$$y \otimes \sum_{p \in \mathfrak{P}} r_p p \longmapsto \sum_{p \in \mathfrak{P}} y r_p \otimes p$$

for all $y \in \mathbb{N}$ and $\sum_{p \in \mathfrak{P}} r_p p \in \mathfrak{R}[[\mathfrak{P}]]$. Here the sum in the right-hand side is welldefined, because one has $yr_p = 0$ in \mathbb{N} for all but a finite subset of indices $p \in \mathfrak{P}$.

The contratensor product functor $\odot_{\mathfrak{R}}$: Discr $-\mathfrak{R} \times \mathfrak{R}$ -Contra \longrightarrow Ab preserves all colimits in both the arguments. For any discrete right \mathfrak{R} -module \mathfrak{N} and any set X, one has a natural isomorphism of abelian groups

$$\mathcal{N} \odot_{\mathfrak{R}} \mathfrak{R}[[X]] \simeq \mathcal{N}[X] = \mathcal{N}^{(X)}.$$

For any closed right ideal $\mathfrak{J} \subset \mathfrak{R}$, one has

$$\mathfrak{J} \wedge (\mathfrak{R}[[X]]) = \mathfrak{J}[[X]] \subset \mathfrak{R}[[X]]$$

For any open right ideal $\mathfrak{I} \subset \mathfrak{R}$ and any left \mathfrak{R} -contamodule \mathfrak{P} , one has

$$(\mathfrak{R}/\mathfrak{I})\odot_{\mathfrak{R}}\mathfrak{P}\simeq\mathfrak{P}/(\mathfrak{I}\prec\mathfrak{P}).$$

In particular, $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{R}[[X]] \simeq \mathfrak{R}[[X]]/(\mathfrak{I} \land \mathfrak{R}[[X]]) = \mathfrak{R}[[X]]/\mathfrak{I}[[X]] \simeq (\mathfrak{R}/\mathfrak{I})[X].$

For any open two-sided ideal $\mathfrak{I} \subset \mathfrak{R}$ and any left \mathfrak{R} -contramodule \mathfrak{P} , the quotient group $\mathfrak{P}/(\mathfrak{I} \prec \mathfrak{P})$ has a natural module structure over the quotient ring $\mathfrak{R}/\mathfrak{I}$. The subgroup $\mathfrak{I} \prec \mathfrak{P}$ is an \mathfrak{R} -subcontramodule in \mathfrak{P} , as mentioned above; and the quotient \mathfrak{R} -contramodule structure on $\mathfrak{P}/(\mathfrak{I} \prec \mathfrak{P})$ comes from the $\mathfrak{R}/\mathfrak{I}$ -module structure.

8. Separated and Flat Contramodules

In this section we discuss contramodules over a topological ring with a *countable* base of neighborhoods of zero. There are no new results in this section, which is based on [29, Section E.1] and [43, Section 6]. A brief survey is available in [30, Section 3.3]. A more recent exposition can be found in [32, Sections 4–5]. A connection with algebraic geometry is discussed in [35, Chapter 3].

Given a left \mathfrak{R} -contramodule \mathfrak{P} , consider the natural map to the directed limit $\lambda_{\mathfrak{R},\mathfrak{P}} \colon \mathfrak{P} \longrightarrow \varprojlim_{\mathfrak{I} \subset \mathfrak{R}} \mathfrak{P}/(\mathfrak{I} \land \mathfrak{P})$, where \mathfrak{I} ranges over the open right ideals of \mathfrak{R} . A left \mathfrak{R} -contramodule \mathfrak{P} is said to be *separated* if the map $\lambda_{\mathfrak{R},\mathfrak{P}}$ is injective, and *complete* if $\lambda_{\mathfrak{R},\mathfrak{P}}$ is surjective. We denote full subcategory of separated \mathfrak{R} -contamodules by \mathfrak{R} -Contra_{sep} $\subset \mathfrak{R}$ -Contra.

Lemma 8.1. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Then all left \mathfrak{R} -contramodules are complete (but they need not be separated).

Proof. This goes back, at least, to [28, Lemma A.2.3 and Remark A.3]. Counterexamples of nonseparated contramodules appeared in [28, Section A.1] (see also [49, Example 2.5], [54, Example 3.20], [30, Section 1.5], [31, Example 2.7(1)], and a further discussion in Remark 8.5 below). The completeness assertion for contramodules over topological rings with a countable topology base consisting of two-sided ideals can be found in [29, Lemma E.1.1], and the general case of a countable topology base of right ideals is covered by [43, Lemma 6.3(b)]. See also [30, Lemmas 3.6 and 3.9]. \Box

Lemma 8.2. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Then, for any nonzero left \mathfrak{R} -contramodule \mathfrak{P} , there exists an open right ideal $\mathfrak{I} \subset \mathfrak{R}$ such that $\mathfrak{P}/(\mathfrak{I} \land \mathfrak{P}) \neq 0$.

Proof. This assertion belongs to a class of results known as "contramodule Nakayama lemmas" and going back, at least, to [28, Lemma A.2.1]. The assertion for contramodules over topological rings with a countable topology base of two-sided ideals can be found in [29, Lemma E.1.2], and the general case of a countable base of right ideals is covered by [43, Lemma 6.14]. See also [30, Lemmas 2.1(b) and 3.22]. \Box

A left \mathfrak{R} -contramodule \mathfrak{F} is called *flat* if the contratensor product functor $- \odot_{\mathfrak{R}} \mathfrak{F}$: Discr $-\mathfrak{R} \longrightarrow \mathsf{Ab}$ is exact on the abelian category of discrete right \mathfrak{R} -modules. All projective \mathfrak{R} -contramodules are separated and flat. All directed colimits and coproducts of flat \mathfrak{R} -contramodules (computed in the category \mathfrak{R} -Contra) are flat. We denote the full subcategory of flat \mathfrak{R} -contramodules by \mathfrak{R} -Contra_{flat} $\subset \mathfrak{R}$ -Contra.

When \mathfrak{R} has a base of neighborhoods of zero consisting of open two-sided ideals \mathfrak{I} , the discrete right \mathfrak{R} -modules are the directed unions of modules over the quotient rings $\mathfrak{R}/\mathfrak{I}$. For any right $\mathfrak{R}/\mathfrak{I}$ -module N and any left \mathfrak{R} -contramodule \mathfrak{P} , one has

$$N \odot_{\mathfrak{R}} \mathfrak{P} \simeq N \otimes_{\mathfrak{R}/\mathfrak{I}} \mathfrak{P}/(\mathfrak{I} \checkmark \mathfrak{P}).$$

Hence, in this case, a left \mathfrak{R} -contramodule \mathfrak{F} is flat if and only if the left $\mathfrak{R}/\mathfrak{I}$ -module $\mathfrak{F}/(\mathfrak{I} \prec \mathfrak{F})$ is flat for every open two-sided ideal $\mathfrak{I} \subset \mathfrak{R}$.

Lemma 8.3. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Then all flat left \mathfrak{R} -contramodules are separated.

Proof. The case of topological rings with a countable topology base of two-sided ideals can be found in [29, Lemma E.1.7] (cf. the definition of a "flat contramodule" in [29, the paragraph before Lemma E.1.4], which is stated differently than our definition above). The general case of a countable base of right ideals is covered by [43, Corollary 6.15]. See also [30, Propositions 3.7(a) and 3.10(a)].

Lemma 8.4. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Then

(a) the class of all flat left \Re -contramodules is closed under extensions and kernels of epimorphisms in \Re -Contra;

(b) for any discrete right \mathfrak{R} -module \mathbb{N} , the contratensor product functor $\mathbb{N} \odot_{\mathfrak{R}} - :$ \mathfrak{R} -Contra \longrightarrow Ab takes short exact sequences of flat left \mathfrak{R} -contramodules to short exact sequences of abelian groups. *Proof.* The case of topological rings with a countable topology base of two-sided ideals can be found in [29, Lemmas E.1.4 and E.1.5]. The general case of a countable base of right ideals is covered by [43, Lemma 6.7 or 6.10, Corollaries 6.8, 6.13 and 6.15], with a summary in [43, Corollary 7.1(a–b)]. See also [30, Propositions 3.7(b) and 3.10(b)]. \Box

In particular, according to Lemma 8.4(a), the full subcategory of flat \Re -contramodules \Re -Contra_{flat} is closed under extensions in \Re -Contra when the topological ring \Re has a countable base of neighborhoods of zero. So the full subcategory \Re -Contra_{flat} inherits an exact category structure from the abelian exact structure of \Re -Contra in this case.

Remark 8.5. Lemma 8.4 tells us that the class of flat \Re -contramodules is wellbehaved (under the assumption of a countable topology base in \Re). But the class of separated \Re -contramodules is *not* well-behaved.

In suffices to consider the case of the ring of *p*-adic integers $\Re = \mathbb{Z}_p$ (in the *p*-adic topology), or the ring of formal Taylor power series $\Re = k[[t]]$ in one variable *t* over a field *k* (in the *t*-adic topology). In both cases, the forgetful functor \Re -Contra \longrightarrow \Re -Mod is fully faithful; in fact, even the forgetful functors \mathbb{Z}_p -Contra \longrightarrow Ab and k[[t]]-Contra $\longrightarrow k[t]$ -Mod are fully faithful. The essential images of these forgetful functors are described in [30, Section 2.2] and the references therein.

With these descriptions in mind, there is a now-classical counterexample showing that the full subcategory \Re -Contra_{sep} is not closed under extensions in \Re -Contra [49, Example 2.5] and the category \Re -Contra_{sep} is not abelian [31, Example 2.7(1)]. Furthermore, the full subcategory \Re -Contra_{sep} is not closed under directed colimits, and in fact, not even closed under coproducts in \Re -Contra [30, Section 1.5]. (See also [35, Example 3.1] and Example 8.9 below.)

However, separated contramodules are sometimes easier to work with than arbitrary ones, as the following lemma illustrates.

Lemma 8.6. Let \mathfrak{R} be a complete, separated topological ring, and let $\mathfrak{R} \supset \mathfrak{I}_1 \supset \mathfrak{I}_2 \supset \mathfrak{I}_3 \supset \cdots$ be a countable base of neighborhoods of zero in \mathfrak{R} consisting of open two-sided ideals $\mathfrak{I}_n \subset \mathfrak{R}$, $n \geq 1$.

(a) The full subcategory \mathfrak{R} -Contra_{sep} of separated \mathfrak{R} -contramodules \mathfrak{M} in \mathfrak{R} -Contra is equivalent to the category formed by the following sets of data:

- (1) for every $n \ge 1$, a left \Re/\Im_n -module M_n is given; and
- (2) for every $n \ge 1$, an isomorphism of left \Re/\Im_n -modules

$$M_n \simeq \mathfrak{R}/\mathfrak{I}_n \otimes_{\mathfrak{R}/\mathfrak{I}_{n+1}} M_{n+1}$$

is given.

(b) The full subcategory \mathfrak{R} -Contra_{flat} of flat \mathfrak{R} -contramodules \mathfrak{F} in \mathfrak{R} -Contra is equivalent to the category formed by the following sets of data:

- (1) for every $n \geq 1$, a flat left \Re/\Im_n -module F_n is given; and
- (2) for every $n \ge 1$, an isomorphism of left \Re/\Im_n -modules

$$F_n \simeq \Re/\Im_n \bigotimes_{\mathfrak{R}/\mathfrak{I}_{n+1}} F_{n+1}$$

is given.

Here, in both parts (a) and (b), morphisms between the described sets of data are defined in the obvious way as collections module morphisms forming commutative diagrams with the reduction isomorphisms.

Proof. In part (a), the equivalence of categories assigns to a separated left \Re -contramodule \mathfrak{M} the collection of modules $M_n = (\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{M} = \mathfrak{M}/(\mathfrak{I}_n \wedge \mathfrak{M})$, endowed with the obvious reduction isomorphisms. The inverse functor assigns to a collection of modules M_n endowed with reduction isomorphisms the separated \mathfrak{R} -contramodule $\mathfrak{M} = \varprojlim_{n\geq 1} M_n$. Here the $\mathfrak{R}/\mathfrak{I}_n$ -modules M_n are viewed as \mathfrak{R} -contramodules via the natural surjective ring homomorphisms $\mathfrak{R} \longrightarrow \mathfrak{R}/\mathfrak{I}_n$, and the directed limit can be computed in the category \mathfrak{R} -Contra (as the forgetful functor \mathfrak{R} -Contra \longrightarrow \mathfrak{R} -Mod preserves all limits). The assertion that these rules indeed define mutually inverse equivalences of categories is [29, Lemma E.1.3] (see [43, Corollary 6.4] for a generalization to topological rings with a countable base of right ideals). Lemma 8.1 needs to be used here.

The equivalence of categories in part (b) is obtained by restricting the equivalence of part (a) to the respective full subcategories on both sides. Lemma 8.3 plays an important role here. $\hfill \Box$

Lemma 8.7. Let \mathfrak{R} be a complete, separated topological ring, and let $\mathfrak{R} \supset \mathfrak{I}_1 \supset \mathfrak{I}_2 \supset \mathfrak{I}_3 \supset \cdots$ be a countable base of neighborhoods of zero in \mathfrak{R} consisting of open two-sided ideals $\mathfrak{I}_n \subset \mathfrak{R}$, $n \geq 1$. Then a left \mathfrak{R} -contramodule \mathfrak{P} is projective if and only if (it is separated and flat and) the corresponding reduction datum in Lemma 8.6 consists of projective modules, that is, if and only if the $\mathfrak{R}/\mathfrak{I}_n$ -module $\mathfrak{P}/(\mathfrak{I}_n \prec \mathfrak{P})$ is projective for every $n \geq 1$.

Proof. This is [29, Corollary E.1.10(a)]; see also [30, Proposition 3.8].

The category \Re -Contra is better behaved than \Re -Contra_{sep} in that \Re -Contra is abelian (cf. Remark 8.5). Still, the directed colimit functors are *not* exact in \Re -Contra, as one can see, e. g., from [43, Examples 4.4]. The next lemma tells us that the directed colimit functors *are* exact in the exact category \Re -Contra_{flat} (assuming a countable base of neighborhoods of zero in \Re).

Lemma 8.8. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Then the directed colimit of any directed diagram of short exact sequences of flat left \mathfrak{R} -contramodules (computed in the category \mathfrak{R} -Contra) is a short exact sequence of (flat) left \mathfrak{R} -contramodules.

Proof. In the case of a topological ring with a countable topology base of two-sided ideals, the assertion is easily deduced from the fact that the class of all flat left \Re -contramodules is closed under directed colimits in \Re -Contra and the description of flat left \Re -contramodules in Lemma 8.6(b) together with Lemma 8.4(b). The fact that countable directed limits of diagrams of surjective maps of abelian groups are exact functors needs to be used here. The general case of a countable base of right

ideals is covered by [43, Lemma 6.16] (see also [43, Proposition 6.17, Remark 6.18, and Corollary 7.1(c)] and [30, Proposition 3.10(c)]).

Example 8.9. In the context of algebraic geometry, a *strict ind-affine* \aleph_0 *-ind-scheme* \mathfrak{X} is an ind-object in the category of affine schemes representable by a countable directed diagram of closed immersions of affine schemes $X_1 \longrightarrow X_2 \longrightarrow X_2 \longrightarrow \cdots$. So one has $X_n = \operatorname{Spec} R_n$ for some commutative rings R_n , and the scheme morphisms $X_n \longrightarrow X_{n+1}$ correspond to surjective ring homomorphisms $R_{n+1} \longrightarrow R_n$.

The category of strict ind-affine \aleph_0 -ind-schemes is equivalent to the category of complete, separated topological commutative rings having a countable base of neighborhoods of zero consisting of two-sided ideals. To an ind-scheme $\mathfrak{X} = \underset{n\geq 1}{\underset{n\geq 1}{\boxtimes}} X_n$ as above, the topological ring $\mathfrak{R} = \varprojlim_{n\geq 1} R_n$ (with the topology of projective limit of discrete rings R_n) is assigned [4, Example 7.11.2(i)], [35, Example 1.6(2)].

In this context, the sets of data from Lemma 8.6(a) are called " \mathcal{O}^p -modules on \mathfrak{X} " in [4, Section 7.11.3] or pro-quasi-coherent pro-sheaves on \mathfrak{X} in [35, Section 3.1]. The sets of data from Lemma 8.6(b) are called "flat \mathcal{O}^p -modules on \mathfrak{X} " in [4, Section 7.11.3] or flat pro-quasi-coherent pro-sheaves on \mathfrak{X} in [35, Section 3.4]. So the category of pro-quasi-coherent pro-sheaves on \mathfrak{X} is equivalent to the category of separated \mathfrak{R} -contramodules, while the category of flat pro-quasi-coherent pro-sheaves on \mathfrak{X} is equivalent to the category of flat \mathfrak{R} -contramodules. We refer to [4, Remark 7.11.3(ii)] and [35, Examples 3.1 and 3.8] for further discussion. For a discussion of non-indaffine \aleph_0 -ind-schemes with further details, see Remark 10.3 below.

9. Countably Presentable Flat Contramodules

We refer to Section 1 above for the definitions of κ -presentable objects, locally κ -presentable categories, and κ -accessible categories, with the references to [1].

Proposition 9.1. Let κ be an uncountable regular cardinal and \Re be a complete, separated topological ring with a base of neighborhoods of zero of cardinality less than κ , consisting of open right ideals.

(a) The category of left \mathfrak{R} -contramodules \mathfrak{R} -Contra is locally κ -presentable. A left \mathfrak{R} -contramodule \mathfrak{P} is κ -presentable as an object of the category \mathfrak{R} -Contra if and only if it is the cokernel of a morphism of free left \mathfrak{R} -contramodules $\mathfrak{R}[[Y]] \longrightarrow \mathfrak{R}[[X]]$ with the sets X and Y having cardinalities less than κ .

(b) The category of complexes of left \mathfrak{R} -contramodules $\mathsf{Com}(\mathfrak{R}-\mathsf{Contra})$ is locally κ -presentable. A complex of left \mathfrak{R} -contramodules \mathfrak{P}^{\bullet} is κ -presentable as an object of the category $\mathsf{Com}(\mathfrak{R}-\mathsf{Contra})$ if and only if all its terms \mathfrak{P}^n , $n \in \mathbb{Z}$, are κ -presentable \mathfrak{R} -contramodules.

Proof. Part (a): the "if" assertion is easily provable using the fact that the forgetful functor \Re -Contra $\longrightarrow \Re$ -Mod preserves κ -directed colimits. The latter follows from the observation that any zero-convergent family of elements in \Re has cardinality less than κ . To prove the "only if" assertion, one then needs to check that all

left \Re -contramodules are κ -directed colimits of the cokernels of morphisms of free contramodules with less than κ generators, and the class of all such cokernels is closed under direct summands. The first assertion of part (a) follows immediately (cf. the discussion in [43, Sections 1.1 and 5] and [44, Sections 6.2 and 6.4]).

Part (b): the "if" assertion follows from the "if" assertion of part (a) by Lemma 1.1(b). To deduce the "only if", one can check directly that all complexes of left \Re -contramodules are κ -directed colimits of complexes whose terms are the cokernels of morphisms of free contramodules with less than κ generators. The first assertion is then also clear (cf. [45, second paragraph of the proof of Lemma 6.3]). Alternatively, one can refer to Theorem 1.9 or a suitable additive version of [19, Theorem 1.2], similarly to the argument in Proposition 3.3 and Remark 3.4.

Now let \mathfrak{R} be a complete, separated topological ring with a *countable* base of neighborhoods of zero consisting of open right ideals. Then the fully faithful inclusion functor \mathfrak{R} -Contra_{sep} $\longrightarrow \mathfrak{R}$ -Contra has a left adjoint functor $\Lambda_{\mathfrak{R}} \colon \mathfrak{R}$ -Contra $\longrightarrow \mathfrak{R}$ -Contra_{sep} assigning to every left \mathfrak{R} -contramodule \mathfrak{P} its maximal separated quotient \mathfrak{R} -contramodule

$$\Lambda_{\mathfrak{R}}(\mathfrak{P}) = \lim_{\mathfrak{I} \subset \mathfrak{R}} \mathfrak{P}/(\mathfrak{I} \land \mathfrak{P}).$$

It is explained in [43, Lemma 6.2] or [32, Proposition 4.2(b)] how to endow the directed limit $\varprojlim_{\mathfrak{I}\subset\mathfrak{R}}\mathfrak{P}/(\mathfrak{I}\prec\mathfrak{P})$ computed in the category of abelian groups with a left \mathfrak{R} -contramodule structure. The natural morphism $\lambda_{\mathfrak{R},\mathfrak{P}} \colon \mathfrak{P} \longrightarrow \varprojlim_{\mathfrak{I}\subset\mathfrak{R}}\mathfrak{P}/(\mathfrak{I}\prec\mathfrak{P})$ is the adjunction unit.

Lemma 9.2. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Let \mathfrak{P} be a left \mathfrak{R} -contramodule such that the separated left \mathfrak{R} -contramodule $\Lambda_{\mathfrak{R}}(\mathfrak{P})$ is flat. Then the \mathfrak{R} -contramodule \mathfrak{P} is flat and separated, and the natural morphism $\lambda_{\mathfrak{R},\mathfrak{P}} \colon \mathfrak{P} \longrightarrow \Lambda_{\mathfrak{R}}(\mathfrak{P})$ is an isomorphism.

Proof. This is an intermediate step in the proofs of [29, Corollary E.1.7] and [43, Corollary 6.15] (which are restated above as Lemma 8.3). The argument is based on the contramodule Nakayama lemma (Lemma 8.2 above). Alternatively, the same proof can be phrased as follows. One observes that, for any left \mathfrak{R} -contramodule \mathfrak{P} , the adjunction morphism $\lambda_{\mathfrak{R},\mathfrak{P}}$ induces an isomorphism of the contratensor product functors $- \odot_{\mathfrak{R}} \mathfrak{P} \simeq - \odot_{\mathfrak{R}} \Lambda_{\mathfrak{R}}(\mathfrak{P})$ (essentially, the nonseparated part of \mathfrak{P} is killed by contratensor products). This shows that flatness of $\Lambda_{\mathfrak{R}}(\mathfrak{P})$ implies flatness of \mathfrak{P} . Then the separatedness and completeness of \mathfrak{P} follow from Lemmas 8.1 and 8.3.

Proposition 9.3. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals, and let κ be an uncountable regular cardinal. Then the category of separated left \mathfrak{R} -contramodules \mathfrak{R} -Contra_{sep} is locally κ -presentable. A separated left \mathfrak{R} -contramodule \mathfrak{M} is κ -presentable as an object of the category \mathfrak{R} -Contra_{sep} if and only if it has the form $\mathfrak{M} = \Lambda_{\mathfrak{R}}(\mathfrak{P})$, where \mathfrak{P} is a κ -presentable object of the category \mathfrak{R} -Contra. Proof. One needs to observe that the full subcategory \mathfrak{R} -Contra_{sep} $\subset \mathfrak{R}$ -Contra is closed under \aleph_1 -directed colimits. Indeed, for any closed subgroup $\mathfrak{A} \subset \mathfrak{R}$, the functor $\mathfrak{P} \longrightarrow \mathfrak{A} \land \mathfrak{P}$ commutes with \aleph_1 -directed colimits (since the functor $X \longmapsto \mathfrak{A}[[X]]$: Sets \longrightarrow Sets does), and countable directed limits also commute with \aleph_1 -directed colimits. Consequently, the inclusion functor \mathfrak{R} -Contra_{sep} \longrightarrow \mathfrak{R} -Contra preserves κ -directed colimits, and it follows that the left adjoint functor $\Lambda_{\mathfrak{R}}: \mathfrak{R}$ -Contra $\longrightarrow \mathfrak{R}$ -Contra_{sep} takes κ -presentable objects to κ -presentable objects. This proves the "if" assertion.

Furthermore, the functor Λ_{\Re} , being a left adjoint, preserves all colimits. Hence all colimits exist in \Re -Contra_{sep} and can be constructed by applying Λ_{\Re} to the colimits in \Re -Contra. Since all objects of \Re -Contra are κ -directed colimits of κ -presentable objects by Proposition 9.1(a), so are all the objects of \Re -Contra_{sep}. Thus the category \Re -Contra_{sep} is locally κ -presentable.

To deduce the assertion "only if", we notice that all the objects of \mathfrak{R} -Contra_{sep} are κ -directed colimits of separated contramodules of the desired form $\mathfrak{M} = \Lambda_{\mathfrak{R}}(\mathfrak{P})$ with $\mathfrak{P} \in \mathfrak{R}$ -Contra_{< κ}, as we have just shown. It remains to check that the class of all separated contramodules of the desired form $\mathfrak{M} = \Lambda_{\mathfrak{R}}(\mathfrak{P})$ is closed under direct summands. Let us prove this class is closed under cokernels in \mathfrak{R} -Contra_{sep}.

Suppose given a morphism $g: \mathfrak{M} \longrightarrow \mathfrak{N}$, where $\mathfrak{M} = \Lambda_{\mathfrak{R}}(\mathfrak{P}), \ \mathfrak{N} = \Lambda_{\mathfrak{R}}(\mathfrak{Q})$, and $\mathfrak{P}, \ \mathfrak{Q} \in \mathfrak{R}$ -Contra_{< κ}. Then the natural (adjunction) morphisms $\mathfrak{P} \longrightarrow \mathfrak{M}$ and $\mathfrak{Q} \longrightarrow \mathfrak{N}$ are surjective. Let $\mathfrak{F} \longrightarrow \mathfrak{P}$ be a surjective morphism onto \mathfrak{P} from a free left \mathfrak{R} -contramodule with less than κ generators. Then the composition $\mathfrak{F} \longrightarrow \mathfrak{P} \longrightarrow$ $\mathfrak{M} \longrightarrow \mathfrak{N}$ can be lifted to a morphism $f: \mathfrak{F} \longrightarrow \mathfrak{Q}$. The cokernel of the morphism gin \mathfrak{R} -Contra_{sep} can be obtained by applying $\Lambda_{\mathfrak{R}}$ to the cokernel of the morphism fin \mathfrak{R} -Contra. The latter cokernel is κ -presentable in \mathfrak{R} -Contra. \Box

Lemma 9.4. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals, and let κ be an uncountable regular cardinal. Then a flat left \mathfrak{R} -contramodule is κ -presentable as an object of the abelian category \mathfrak{R} -Contra if and only if it is κ -presentable as an object of the category \mathfrak{R} -Contra.

Proof. The "only if" assertion holds, since the functor $\Lambda_{\mathfrak{R}}: \mathfrak{R}$ -Contra $\longrightarrow \mathfrak{R}$ -Contra_{sep} takes κ -presentable objects to κ -presentable objects, as explained in the proof of Proposition 9.3, and flat contramodules are separated by Lemma 8.3.

The proof of the "if" is based on Lemma 9.2 and the "only if" assertion of Proposition 9.3. If a left \mathfrak{R} -contramodule \mathfrak{F} is κ -presentable in \mathfrak{R} -Contra_{sep}, then Proposition 9.3 tells us that $\mathfrak{F} = \Lambda_{\mathfrak{R}}(\mathfrak{P})$, where \mathfrak{P} is κ -presentable in \mathfrak{R} -Contra. Now if \mathfrak{F} is flat, then $\mathfrak{P} \simeq \mathfrak{F}$ by Lemma 9.2.

Lemma 9.5. Let \mathfrak{R} be a complete, separated topological ring, and let $\mathfrak{R} \supset \mathfrak{I}_1 \supset \mathfrak{I}_2 \supset \mathfrak{I}_3 \supset \cdots$ be a countable base of neighborhoods of zero in \mathfrak{R} consisting of open two-sided ideals $\mathfrak{I}_n \subset \mathfrak{R}$, $n \geq 1$. Let κ be an uncountable regular cardinal. Then a separated \mathfrak{R} -contramodule \mathfrak{M} is κ -presentable as an object of \mathfrak{R} -Contra_{sep} if and only if the $\mathfrak{R}/\mathfrak{I}_n$ -module $M_n = \mathfrak{M}/(\mathfrak{I}_n \prec \mathfrak{M})$ is κ -presentable for every $n \geq 1$.

Consequently, a flat \mathfrak{R} -contramodule \mathfrak{F} is κ -presentable as an object of \mathfrak{R} -Contra (or of \mathfrak{R} -Contra_{sep}) if and only if the flat $\mathfrak{R}/\mathfrak{I}_n$ -module $F_n = \mathfrak{F}/(\mathfrak{I}_n \prec \mathfrak{F})$ is κ -presentable for every $n \geq 1$.

Proof. The first "only if" assertion follows easily from the "only if" assertion of Proposition 9.3. To prove the first "if", use the description of the category \Re -Contra_{sep} provided by Lemma 8.6(a) together with the fact that \aleph_1 -directed colimits commute with countable limits in the category of abelian groups. Then the remaining assertions (concerning flat contramodules) follow from Lemma 9.4.

Lemma 9.6. Let \mathfrak{R} be a complete, separated topological ring, and let $\mathfrak{R} \supset \mathfrak{I}_1 \supset \mathfrak{I}_2 \supset \mathfrak{I}_3 \supset \cdots$ be a countable base of neighborhoods of zero in \mathfrak{R} consisting of open twosided ideals $\mathfrak{I}_n \subset \mathfrak{R}, n \geq 1$. Then the projective dimension of a flat \mathfrak{R} -contramodule \mathfrak{F} (as an object of \mathfrak{R} -Contra) is equal to the supremum of the projective dimensions of the flat $\mathfrak{R}/\mathfrak{I}_n$ -modules $F_n = \mathfrak{F}/(\mathfrak{I}_n \prec \mathfrak{F})$ (as objects of $\mathfrak{R}/\mathfrak{I}_n$ -Mod), $n \geq 1$.

Proof. Let $\mathfrak{P}_{\bullet} \longrightarrow \mathfrak{F}$ be a projective resolution of \mathfrak{F} in \mathfrak{R} -Contra and \mathfrak{F}_i be the image of the differential $\mathfrak{P}_i \longrightarrow \mathfrak{P}_{i-1}$ (so $\mathfrak{F} = \mathfrak{F}_0$). Applying Lemma 8.4(a) iteratively and keeping in mind that projective contramodules are flat, one shows that the \mathfrak{R} -contramodules \mathfrak{F}_i are flat for all $i \geq 0$. Now Lemma 8.4(b) tells us that the functors $\mathfrak{Q} \longmapsto (\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{Q} = \mathfrak{Q}/(\mathfrak{I}_n \land \mathfrak{Q})$ preserve exactness of the short exact sequences $0 \longrightarrow \mathfrak{F}_i \longrightarrow \mathfrak{P}_i \longrightarrow \mathfrak{F}_{i-1} \longrightarrow 0$. Therefore, $(\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{P}_{\bullet}$ is a projective resolution of the $\mathfrak{R}/\mathfrak{I}_n$ -module $F_n = (\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{F}_i$ and the $\mathfrak{R}/\mathfrak{I}_n$ -module $(\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{F}_i$ is the image of the differential $(\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{P}_i \longrightarrow (\mathfrak{R}/\mathfrak{I}_n) \odot_{\mathfrak{R}} \mathfrak{P}_{i-1}$. It remains to use the characterization of projective \mathfrak{R} -contramodules provided by Lemma 8.7 in order to deduce the desired assertion.

Corollary 9.7. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then any countably presentable flat \mathfrak{R} -contramodule has projective dimension at most 1 in the abelian category \mathfrak{R} -Contra.

Proof. Recall that all countably presentable flat modules (over a discrete ring R) have projective dimension at most 1 in R-Mod, by Lemma 1.3. Then it remains to compare Lemma 9.5 (for flat contramodules and $\kappa = \aleph_1$) with Lemma 9.6.

Question 9.8. One would expect the assertion of Corollary 9.7 to hold for left contramodules over any complete, separated topological ring \Re with a countable base of neighborhoods of zero consisting of open *right* ideals. But we do not know whether this is true. In fact, we don't even know the answer to Question 11.2 below, which would seem to be easier.

10. FLAT CONTRAMODULES AS DIRECTED COLIMITS

The following theorem describing flat \Re -contramodules is the second main result of this paper. It should be compared with the discussion of the conventional finite,

rather than countable Govorov–Lazard theorem for flat contramodules in the paper [42, Section 10 and Example 13.4]. The counterexamples in [42] show that the finite Govorov–Lazard theorem fails for flat contramodules over topological rings *without* a countable base of neighborhoods of zero (it is still an open question whether it holds under the assumption of a countable base). Theorem 10.1 is also a contramodule analogue of [47, Theorems 2.4 and 3.5] and Theorem 3.1 above.

We refer to Lemmas 9.4 and 9.5 for an explication of what is meant by a countably presentable flat \Re -contramodule in Theorem 10.1 (i. e., countably presentable in \Re -Contra or \Re -Contra_{sep}, as described in Propositions 9.1(a) and 9.3).

Theorem 10.1. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the category \mathfrak{R} -Contra_{flat} of flat left \mathfrak{R} -contramodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of \mathfrak{R} -Contra_{flat} are precisely all the countably presentable flat \mathfrak{R} -contramodules. Consequently, every flat \mathfrak{R} -contramodule is an \aleph_1 -directed colimit of countably presentable flat \mathfrak{R} -contramodules.

Proof. The full subcategory \Re -Contra_{flat} is closed under directed colimits in \Re -Contra. Therefore, all countably presentable flat \Re -contramodules are \aleph_1 -presentable in \Re -Contra_{flat}.

The full assertion of the theorem is obtained by applying Theorem 1.6 for $\kappa = \aleph_1$ and $\lambda = \aleph_0$. The argument is based on the description of the category of flat \Re -contramodules provided by Lemma 8.6(b) and the description of countably presentable separated/flat \Re -countramodules provided by Lemma 9.5.

Let $\mathfrak{R} \supset \mathfrak{I}_1 \supset \mathfrak{I}_2 \supset \mathfrak{I}_3 \supset \cdots$ be a countable base of neighborhoods of zero in \mathfrak{R} consisting of open two-sided ideals. Let $\mathsf{K} = \mathsf{L}$ be the Cartesian product of the categories of flat left modules over the rings $\mathfrak{R}/\mathfrak{I}_n$, taken over the integers $n \geq 1$; so $\mathsf{K} = \mathsf{L} = \prod_{n=1}^{\infty} (\mathfrak{R}/\mathfrak{I}_n - \mathsf{Mod}_{\mathsf{flat}})$. Consider the following pair of parallel functors F_1 , $F_2 \colon \mathsf{K} \rightrightarrows \mathsf{L}$. The functor F_1 takes a collection of flat modules $(K_n \in \mathfrak{R}/\mathfrak{I}_n - \mathsf{Mod}_{\mathsf{flat}})_{n \geq 1}$ to the collection of flat modules $L_n = \mathfrak{R}/\mathfrak{I}_n \otimes_{\mathfrak{R}/\mathfrak{I}_{n+1}} K_{n+1}$. The functor F_2 is the identity functor.

Then Lemma 8.6(b) tells us that the isomorpher category C is equivalent to the category of flat left \mathfrak{R} -contramodules \mathfrak{R} -Contra_{flat}. The categories $\mathfrak{R}/\mathfrak{I}_n$ -Mod_{flat} are \aleph_1 -accessible by Lemma 1.2, and their Cartesian product $\prod_{n=1}^{\infty}(\mathfrak{R}/\mathfrak{I}_n-\mathsf{Mod}_{\mathsf{flat}})$ is \aleph_1 -accessible by Proposition 1.4. Theorem 1.6 is applicable, and it tells us that the category C is \aleph_1 -accessible. Comparing the description of the \aleph_1 -presentable objects of C provided by Theorem 1.6 with the description of the \aleph_1 -presentable flat \mathfrak{R} -contramodules provided by Lemma 9.5 one can see that these two classes of objects coincide, as desired.

Let us also provide a description of arbitrary complexes of flat \Re -contramodules as directed colimits.

Proposition 10.2. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the category $\mathsf{Com}(\mathfrak{R}-\mathsf{Contra}_{\mathsf{flat}})$ of complexes of flat left \mathfrak{R} -contramodules is \aleph_1 -accessible.

The \aleph_1 -presentable objects of $\mathsf{Com}(\mathfrak{R}-\mathsf{Contra}_{\mathsf{flat}})$ are precisely all the complexes of countably presentable flat \mathfrak{R} -contramodules. Consequently, every complex of flat \mathfrak{R} -contramodules is an \aleph_1 -directed colimit of complexes of countably presentable flat \mathfrak{R} -contramodules.

Proof. All complexes of countably presentable \Re -contramodules are \aleph_1 -presentable as objects of $\mathsf{Com}(\Re-\mathsf{Contra})$ by Proposition 9.1(b). Since the full subcategory $\mathsf{Com}(\Re-\mathsf{Contra}_{\mathsf{flat}})$ is closed under directed colimits in $\mathsf{Com}(\Re-\mathsf{Contra})$, it follows that all complexes of countably presentable flat \Re -contramodules are \aleph_1 -presentable in $\mathsf{Com}(\Re-\mathsf{Contra}_{\mathsf{flat}})$.

The full assertion of the proposition is obtained by combining the results of Theorems 10.1 and 1.9. The argument is similar to the proof of Proposition 3.3. \Box

Remark 10.3. One can restrict the results of this section to commutative topological rings \mathfrak{R} , interpret commutative topological rings as ind-affine ind-schemes (as per Example 8.9), and then generalize to non-ind-affine ind-schemes. To avoid a detailed discussion of the basics of the ind-scheme theory (see [35, Section 1.2] and the references therein), let us say that a *strict ind-quasi-compact ind-quasi-separated* \aleph_0 -*ind-scheme* \mathfrak{X} can be defined as an ind-object in the category of quasi-compact quasi-separated schemes representable by a countable directed diagram of closed immersions of quasi-compact quasi-separated schemes $X_1 \longrightarrow X_2 \longrightarrow X_3 \longrightarrow \cdots$.

An " \mathcal{O}^p -module on \mathfrak{X} " (in the terminology of [4, Section 7.11.3]) or a pro-quasicoherent pro-sheaf on \mathfrak{X} in our preferred terminology of [35, Section 3.1] can be then defined as a sequence of quasi-coherent sheaves M_n of the schemes X_n together with isomorphisms of quasi-coherent sheaves $M_n \simeq i_n^* M_{n+1}$ on X_n , where $i_n \colon X_n \longrightarrow X_{n+1}$ denote the closed immersion morphisms ($n \ge 1$). A pro-quasi-coherent pro-sheaf $(F_n)_{n\ge 1}$ is said to be *flat* if the quasi-coherent sheaf F_n on X_n is flat for every $n \ge 1$. The flat pro-quasi-coherent pro-sheaves form a well-behaved (exact) full subcategory in a badly behaved additive category of arbitrary pro-quasi-coherent pro-sheaves on \mathfrak{X} (see Example 8.9 and Remark 8.5).

The argument similar to the proof of Theorem 10.1 and based on the result of [47, Theorem 2.4] shows that the category of flat pro-quasi-coherent pro-sheaves on \mathfrak{X} is \aleph_1 -accessible, and the flat pro-quasi-coherent pro-sheaves $(F_n)_{n\geq 1}$ with locally countably presentable flat quasi-coherent sheaves F_n on X_n are the \aleph_1 -presentable objects of this category. The flat pro-quasi-coherent pro-sheaf version of Proposition 10.2 also holds, with the same proof. The same applies to the results of Proposition 11.3 and Corollary 11.4 below.

11. EXACT SEQUENCES OF FLAT CONTRAMODULES AS DIRECTED COLIMITS

Let us start with a contramodule analogue/generalization of Lemma 4.1.

Lemma 11.1. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the kernel of any

surjective morphism from a countably presentable flat \Re -contramodule to a countably presentable flat \Re -contramodule is a countably presentable (flat) \Re -contramodule.

Proof. The class of flat \mathfrak{R} -contramodules is closed under kernels of surjective morphisms in \mathfrak{R} -Contra by Lemma 8.4(a). Now if $0 \longrightarrow \mathfrak{H} \longrightarrow \mathfrak{G} \longrightarrow \mathfrak{F} \longrightarrow 0$ is a short exact sequence of flat left \mathfrak{R} -contramodules, then Lemma 8.4(b) tells us that $0 \longrightarrow (\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{H} \longrightarrow (\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{G} \longrightarrow (\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{F} \longrightarrow 0$ is a short exact sequence of left $\mathfrak{R}/\mathfrak{I}$ -modules for any open two-sided ideal $\mathfrak{I} \subset \mathfrak{R}$. If the \mathfrak{R} -contramodules \mathfrak{F} and \mathfrak{G} are countably presentable, then so are the $\mathfrak{R}/\mathfrak{I}$ -modules $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{F}$ and $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{G}$, by Lemma 9.5. So Lemma 4.1 tells us that the $\mathfrak{R}/\mathfrak{I}$ -module $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{H}$ is countably presentable, and it remains to apply Lemma 9.5 again in order to conclude that the \mathfrak{R} -contramodule \mathfrak{H} is countably presentable.

Alternatively, one can deduce the lemma from Corollary 9.7 essentially in the same way as the proof of Lemma 4.1 deduces it from Lemma 1.3. Then one needs to use the facts that the kernel of any surjective morphism of countably presentable \Re -contramodules is a countably generated \Re -contramodule, and any countably generated projective \Re -contramodule is countably presentable.

Question 11.2. Similarly to Question 9.8, we do *not* know, and it would be interesting to learn, whether the assertion of Lemma 11.1 holds for left contramodules over any complete, separated topological ring \Re with a countable base of neighborhoods of zero consising of open *right* ideals.

The following proposition describes short exact sequences of flat \Re -contramodules.

Proposition 11.3. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the category of short exact sequences of flat \mathfrak{R} -contramodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of this category are precisely all the short exact sequences of countably presentable flat \mathfrak{R} -contramodules. Consequently, every short exact sequence of flat \mathfrak{R} -contramodules is an \aleph_1 -directed colimit of short exact sequences of countably presentable flat \mathfrak{R} -contramodules.

Proof. The argument is similar to the proof of Theorem 10.1 and uses Lemmas 4.3 and 8.4(b) together with Lemmas 8.6(b) and 9.5. The assertion is obtained by applying Proposition 1.4 and Theorem 1.6 for $\kappa = \aleph_1$ and $\lambda = \aleph_0$.

Let $\mathfrak{R} \supset \mathfrak{I}_1 \supset \mathfrak{I}_2 \supset \mathfrak{I}_3 \supset \cdots$ be a countable base of neighborhoods of zero in \mathfrak{R} consisting of open two-sided ideals. Let $\mathsf{K} = \mathsf{L}$ be the Cartesian product of the categories of short exact sequences of flat left modules over the rings $\mathfrak{R}/\mathfrak{I}_n$, taken over the integers $n \geq 1$. Consider the pair of functors $F_1, F_2: \mathsf{K} \rightrightarrows \mathsf{L}$ similar to the one in the proof of Theorem 10.1.

Then Lemmas 8.4(b) and 8.6(b) imply that the isomorpher category C is equivalent to the category of short exact sequences of flat left \Re -contramodules (one also needs to use the fact that directed limits of countable sequences of surjective maps of short exact sequences of abelian groups are short exact sequences of abelian groups again). The categories K and L are \aleph_1 -accessible by Lemma 4.3 and Proposition 1.4. Theorem 1.6 is applicable; it tells us that the category C is \aleph_1 -accessible and provides a description of its full subcategory of \aleph_1 -presentable objects. To compare it with the description asserted in the proposition, one needs to use Lemma 9.5.

The assertion of Proposition 11.3 (together with Theorem 10.1) can be restated by saying that the exact category of flat \mathfrak{R} -contramodules \mathfrak{R} -Contra_{flat} is a *locally* \aleph_1 -coherent exact category in the sense of [39, Section 1].

Let us say that an acyclic complex of flat \Re -contramodules is *pure acyclic* if its \Re -contramodules of cocycles are flat. For topological rings \Re with a countable base of neighborhoods of zero, this is consistent with the notion of *contratensor purity* [33, Section 3], [42, Section 13]. It is worth noticing that any bounded above acyclic complex of flat contramodules is pure acyclic by Lemma 8.4(a).

Corollary 11.4. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the category of pure acyclic complexes of flat \mathfrak{R} -contramodules is \aleph_1 -accessible. The \aleph_1 -presentable objects of this category are precisely all the pure acyclic complexes of countably presentable flat \mathfrak{R} -contramodules. Consequently, every pure acyclic complex of flat \mathfrak{R} -contramodules is an \aleph_1 -directed colimit of pure acyclic complexes of countably presentable flat \mathfrak{R} -contramodules.

Proof. Similar to the proof of Corollary 4.5. One can either prove the assertion of the corollary by a direct argument similar to the proof of Proposition 11.3 and using [38, Corollary 10.14] instead of Lemma 4.3, or deduce the corollary from Proposition 11.3 using the argument with building pure acyclic complexes by splicing short exact sequences as in the proof of Corollary 4.5. \Box

12. Cotorsion Periodicity for Contramodules

For the sake of completeness of the exposition, we start with presenting a weak version of flat/projective periodicity for contramodules. The definition of a *periodic object* (with respect to a class of objects in an abelian category) was given in Section 5. The following proposition is a contramodule generalization of [5, Theorem 2.5] and [26, Remark 2.15] (see also [9, Proposition 7.6]). For a strong version of flat/projective periodicity theorem for contramodules (with a proof based on the results of this paper) see the recent preprint [41].

Proposition 12.1. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then

(a) any projective-periodic flat \Re -contramodule is projective;

(b) in any acyclic complex of projective \Re -contramodules with flat \Re -contramodules of cocycles, the contramodules of cocycles are actually projective (so the complex is contractible).

Proof. This is an easy corollary of the flat/projective periodicity for modules over a ring ([5, Theorem 2.5], [26, Remark 2.15]) together with the basic properties of flat and projective contramodules (Lemmas 8.4(b) and 8.7).

Part (a): let $0 \longrightarrow \mathfrak{F} \longrightarrow \mathfrak{F} \longrightarrow \mathfrak{F} \longrightarrow \mathfrak{F} \longrightarrow 0$ be a short exact sequence of flat left \mathfrak{R} -contramodules with a projective \mathfrak{R} -contramodule \mathfrak{P} . By Lemma 8.4(b), for any open two-sided ideal $\mathfrak{I} \subset \mathfrak{R}$, the short sequence of left $\mathfrak{R}/\mathfrak{I}$ -modules $0 \longrightarrow$ $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{F} \longrightarrow (\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{P} \longrightarrow (\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{F} \longrightarrow 0$ is exact. Clearly, the $\mathfrak{R}/\mathfrak{I}$ -module $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{F}$ is flat and the $\mathfrak{R}/\mathfrak{I}$ -module $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{P}$ is projective (see the discussion in Sections 7–8, cf. Lemmas 8.6(b) and 8.7). By [5, Theorem 2.5], it follows that the $\mathfrak{R}/\mathfrak{I}$ -module $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{F}$ is projective for every open two-sided ideal $\mathfrak{I} \subset \mathfrak{R}$. Using Lemma 8.7, we conclude that the \mathfrak{R} -contramodule \mathfrak{F} is projective.

Part (b): let \mathfrak{P}^{\bullet} be a pure acyclic complex of projective left \mathfrak{R} -contramodules. By Lemma 8.4(b), for any open two-sided ideal $\mathfrak{I} \subset \mathfrak{R}$, the complex of left $\mathfrak{R}/\mathfrak{I}$ -modules $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{P}^{\bullet}$ is (pure) acyclic. Moreover, if \mathfrak{Z}^n , $n \in \mathbb{Z}$, are the \mathfrak{R} -contramodules of cocycles of the complex \mathfrak{P}^{\bullet} , then $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{Z}^n$ are the $\mathfrak{R}/\mathfrak{I}$ -modules of cocycles of the complex $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{P}^{\bullet}$. The complex $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{P}^{\bullet}$ is also a complex of projective left $\mathfrak{R}/\mathfrak{I}$ -modules. By [26, Theorem 8.6 and Remark 2.15] or [9, Proposition 7.6], it follows that the complex $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}} \mathfrak{P}^{\bullet}$ is contractible and the $\mathfrak{R}/\mathfrak{I}$ -modules $(\mathfrak{R}/\mathfrak{I}) \odot_{\mathfrak{R}}$ \mathfrak{I}^n are projective. Once again, we use Lemma 8.7 in order to conclude that the \mathfrak{R} -contramodules \mathfrak{I}^n are projective and the complex \mathfrak{P}^{\bullet} is contractible. \Box

Now we pass to the cotorsion periodicity for \mathfrak{R} -contramodules, which is the main topic of this section. Let us denote by $\operatorname{Ext}^{\mathfrak{R},*}(-,-)$ the Ext groups in the abelian category \mathfrak{R} -Contra. A left \mathfrak{R} -contramodule \mathfrak{B} is said to be *cotorsion* [29, Section E.5], [43, Definition 7.3] if $\operatorname{Ext}^{\mathfrak{R},1}(\mathfrak{F},\mathfrak{B}) = 0$ for all flat left \mathfrak{R} -contramodules \mathfrak{F} .

Introduce the notation $B = \Re$ -Contra^{cot} for the full subcategory of cotorsion \Re -contramodules in the abelian category $C = \Re$ -Contra. Obviously, the full subcategory \Re -Contra^{cot} is closed under extensions in \Re -Contra, so it inherits an exact category structure from the abelian exact structure of \Re -Contra.

The following corollary summarizes some of our previous results and observations. It is a contramodule version of [47, Corollary 5.6 and Lemma 9.1] and Corollary 5.2 above.

Corollary 12.2. Let \mathfrak{R} be a complete, separated topological ring.

(a) Assume that \mathfrak{R} has a countable base of neighborhoods of zero consisting of open right ideals. Then the class all flat left \mathfrak{R} -contramodules is resolving and closed under directed colimits in \mathfrak{R} -Contra. Consequently, one has $\operatorname{Ext}^{\mathfrak{R},n}(\mathfrak{F},\mathfrak{B}) = 0$ for all flat left \mathfrak{R} -contramodules \mathfrak{F} , all cotorsion left \mathfrak{R} -contramodules \mathfrak{B} , and all integers $n \geq 1$. The directed colimit functors in the exact category of flat contramodules \mathfrak{R} -Contra_{flat} are exact (i. e., the directed colimits of admissible short exact sequences are admissible short exact sequences).

(b) Assume that \mathfrak{R} has a countable base of neighborhoods of zero consisting of open two-sided ideals. Then any flat \mathfrak{R} -contramodule is a directed (in fact, \aleph_1 -directed) colimit of flat \mathfrak{R} -contramodules having projective dimensions not exceeding 1 in \mathfrak{R} -Contra. *Proof.* Part (a): the full subcategory \Re -Contra_{flat} $\subset \Re$ -Contra is resolving by Lemma 8.4(a) and closed under directed colimits according to the discussion in Sections 7–8. The Ext vanishing assertion follows by [47, Lemma 6.1]. The directed colimits in the exact category \Re -Contra_{flat} (though not in the abelian category \Re -Contra!) are exact by Lemma 8.8.

In part (b), all flat \Re -contramodules are \aleph_1 -directed colimits of countably presentable flat \Re -contramodules by Theorem 10.1, and countably presentable flat \Re -contramodules have projective dimension at most 1 by Corollary 9.7.

The following result is a contramodule generalization of the cotorsion periodicity theorem of Bazzoni, Cortés-Izurdiaga, and Estrada [2, Theorem 1.2(2) or Proposition 4.8(2)]. It is also a contramodule analogue of [47, Theorem 9.2] and Theorem 5.4 above.

Theorem 12.3. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then any cotorsion-periodic \mathfrak{R} -contramodule is cotorsion.

Proof. This is another application of Theorem 5.3, whose assumptions are satisfied in the situation at hand (for $F = \Re$ -Contra_{flat} $\subset \Re$ -Contra = C) by Corollary 12.2.

Corollary 12.4. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Let \mathfrak{B}^{\bullet} be an acyclic complex in \mathfrak{R} -Contra whose terms \mathfrak{B}^n are cotorsion \mathfrak{R} -contramodules for all $n \in \mathbb{Z}$. Then the \mathfrak{R} -contramodules of cocycles of the complex \mathfrak{B}^{\bullet} are also cotorsion.

Proof. This is a contramodule generalization of [2, Theorem 5.1(2)], and an analogue of [47, Corollary 9.4] and Corollary 5.5 above. The infinite product functors are exact in \Re -Contra, and the class of cotorsion \Re -contramodules is closed under (direct summands and) infinite products. So the argument of [15, Proposition 2] is applicable, deducing the assertion of the corollary from Theorem 12.3.

The definition of a *coresolving subcategory* is dual to that of a resolving one. A class of objects B in an abelian category C is said to be *cogenerating* if every object of C is a subobject of an object from B. We will say that a class of objects $B \subset C$ is *coresolving* if it is cogenerating, closed under extensions, and closed under cokernels of monomorphisms in C.

Proposition 12.5. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open right ideals. Then the class $B = \mathfrak{R}$ -Contra^{cot} of all cotorsion left \mathfrak{R} -contramodules is coresolving in $C = \mathfrak{R}$ -Contra.

Proof. It is obvious from the definition that the class of all cotorsion \Re -contramodules is closed under extensions, and it follows from the higher Ext vanishing assertion of Corollary 12.2(a) that \Re -Contra^{cot} is also closed under cokernels of monomorphisms in \Re -Contra. The most nontrivial assertion in the proposition is that the class of cotorsion \Re -contramodules is cogenerating. This is a part of [43, Corollary 7.8]. (Notice that there are usually *not* enough injective objects in \Re -Contra.) \Box Recall the notation K(E) for the homotopy category of an additive category E. For an abelian (or exact) category E, we denote by $Ac(E) \subset K(E)$ the full subcategory of acyclic complexes. So the derived category D(E) is the Verdier quotient category D(E) = K(E)/Ac(E).

Proposition 12.6. Let C be an abelian category with exact functors of countable product, and let $B \subset C$ be a cogenerating subcategory closed under countable products. Then for any complex C^{\bullet} in C there exists a complex B^{\bullet} in B together with a quasi-isomorphism $C^{\bullet} \longrightarrow B^{\bullet}$ of complexes in C. Consequently, the inclusion of additive categories $B \longrightarrow C$ induces a triangulated equivalence of Verdier quotient categories

$$\frac{\mathsf{K}(\mathsf{B})}{\mathsf{K}(\mathsf{B})\cap\mathsf{Ac}(\mathsf{C})} \stackrel{\simeq}{\longrightarrow} \frac{\mathsf{K}(\mathsf{C})}{\mathsf{Ac}(\mathsf{C})} \,=\, \mathsf{D}(\mathsf{C}).$$

Proof. The proof of the first assertion is similar to the construction of K-projective and K-injective resolutions of complexes of modules in [50, Section 3]. Let us spell out the construction as follows. For every $n \in \mathbb{Z}$, consider the canonical truncation $\tau_{\geq n}C^{\bullet}$ of the complex C^{\bullet} . Since the complex $\tau_{\geq n}C^{\bullet}$ is bounded below and the subcategory $\mathsf{B} \subset \mathsf{C}$ is cogenerating, one can easily find a (bounded below) complex ${}^{(n)}D^{\bullet}$ in B together with a quasi-isomorphism $\tau_{\geq n}C^{\bullet} \longrightarrow {}^{(n)}D^{\bullet}$ in C . Moreover, one can choose the complex ${}^{(n)}D^{\bullet}$ so that the morphism of complexes $\tau_{\geq n}C^{\bullet} \longrightarrow {}^{(n)}D^{\bullet}$ is a termwise monomorphism. The composition $C^{\bullet} \longrightarrow \tau_{\geq n}C^{\bullet} \longrightarrow {}^{(n)}D^{\bullet}$ provides a morphism of complexes $C^{\bullet} \longrightarrow {}^{(n)}D^{\bullet}$.

Now the induced morphism into the product $C^{\bullet} \longrightarrow B^{0,\bullet} = \prod_{n \in \mathbb{Z}} {}^{(n)}D^{\bullet}$ is a termwise monomorphism of complexes in C inducing monomorphisms on all the cohomology objects. Furthermore, $B^{0,\bullet}$ is a complex in B. Next we apply the same construction to the cokernel of the morphism of complexes $C^{\bullet} \longrightarrow B^{0,\bullet}$, embedding the complex $B^{0,\bullet}/C^{\bullet}$ into a complex $B^{1,\bullet}$ in B in such a way that the morphism of complexes $B^{0,\bullet}/C^{\bullet} \longrightarrow B^{1,\bullet}$ is (not only termwise monic, but also) induces monomorphisms on all the cohomology objects. Iterating the procedure, we obtain an exact complex of complexes

$$0 \longrightarrow C^{\bullet} \longrightarrow B^{0,\bullet} \longrightarrow B^{1,\bullet} \longrightarrow B^{2,\bullet} \longrightarrow \cdots$$

such that passing to the cohomology objects produces an exact complex

$$0 \longrightarrow H^*(C^{\bullet}) \longrightarrow H^*(B^{0,\bullet}) \longrightarrow H^*(B^{1,\bullet}) \longrightarrow H^*(B^{2,\bullet}) \longrightarrow \cdots$$

of graded objects in C.

Finally, we totalize the bicomplex $B^{\bullet,\bullet}$ by taking infinite products along the diagonals. Put $B^{\bullet} = \text{Tot}^{\sqcap}(B^{\bullet,\bullet})$ to be the product totalization. Then B^{\bullet} is a complex in B. Furthermore, the resulting morphism of complexes $C^{\bullet} \longrightarrow B^{\bullet}$ is a quasi-isomorphism by the next Lemma 12.7. This proves the first assertion of the proposition, and the second one then follows by [47, Lemma 9.5]. An opposite version of this argument can be found in [29, proof of Proposition A.4.3].

The next lemma is a classical result of Eilenberg and Moore [13].

Lemma 12.7. Let C be an abelian category with exact functors of countable product, and let

 $0 \longrightarrow C^{0,\bullet} \longrightarrow C^{1,\bullet} \longrightarrow C^{2,\bullet} \longrightarrow C^{3,\bullet} \longrightarrow \cdots$

be a bounded below complex of complexes in C. Assume that, passing to the cohomology objects, we obtain an acyclic complex

 $0 \longrightarrow H^*(C^{0,\bullet}) \longrightarrow H^*(C^{1,\bullet}) \longrightarrow H^*(C^{2,\bullet}) \longrightarrow H^*(C^{3,\bullet}) \longrightarrow \cdots$

of graded objects in C. Then the product totalization $\operatorname{Tot}^{\sqcap}(C^{\bullet,\bullet})$ of the bicomplex $C^{\bullet,\bullet}$ is an acyclic complex in C.

Proof. This is a particular case of [13, Theorem 7.4] or (essentially) the dual version of [29, Lemma A.4.4]. \Box

The following corollary is a contramodule/ind-affine ind-scheme analogue of [47, Corollary 9.7] and Corollary 5.6 above.

Corollary 12.8. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the inclusion of exact/abelian categories \mathfrak{R} -Contra^{cot} $\longrightarrow \mathfrak{R}$ -Contra induces an equivalence of their unbounded derived categories,

$$D(\mathfrak{R}-Contra^{cot}) \simeq D(\mathfrak{R}-Contra).$$

Proof. It suffices to compare Propositions 12.5 and 12.6 with Corollary 12.4. \Box

13. Exact Sequences of Flat Contramodules II

In this short section we present contramodule/ind-affine ind-scheme analogues of the results of Section 6. We refer to Section 6 for the discussion of *self-resolving subcategories* in abelian/exact categories.

The following lemma is a contramodule analogue of Corollary 6.4.

Lemma 13.1. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then

(a) the full subcategory G of all countably presentable flat \Re -contramodules is self-resolving in the abelian category \Re -Contra;

(b) the homological dimension of the exact category G (with the exact structure inherited from \Re -Contra) does not exceed 1.

Proof. Part (a): the class of all flat \mathfrak{R} -contramodules is closed under extensions in \mathfrak{R} -Contra by Lemma 8.4(a). The assertion that the class of all countably presentable \mathfrak{R} -contramodules is closed under extensions is provable in the same way as the similar assertion for modules over a ring. Hence the class G of all countably presentable flat \mathfrak{R} -contramodules is closed under extensions in \mathfrak{R} -Contra. The class G is also closed under kernels of epimorphisms in \mathfrak{R} -Contra by Lemma 11.1. Finally, the class G is self-generating in \mathfrak{R} -Contra, since every object of G is a quotient object in \mathfrak{R} -Contra

of an object from G (namely, of the free \Re -contramodule with a countable set of generators) that is projective in \Re -Contra.

Part (b): the assertion follows directly from Corollary 9.7. One can also invoke Lemma 6.3 here to the effect that it follows from part (a) that the Ext groups computed in G agree with the ones computed in \Re -Contra.

The following theorem is a contramodule/ind-affine ind-scheme version of [14, Theorem 2.4 (1) \Leftrightarrow (3)] or [26, Theorem 8.6 (ii) \Leftrightarrow (iii)], and an analogue of Theorem 6.5 above. It should be also compared to [46, Corollary 0.5 or Proposition 8.13]. In the latter context, Theorem 13.2 can be interpreted as saying that (over a topological ring \mathfrak{R} with a countable topology base of two-sided ideals) for the exact category of flat \mathfrak{R} -contramodules, the derived category coincides with the (suitably defined) coderived category.

Theorem 13.2. Let \mathfrak{R} be a complete, separated topological ring with a countable base of neighborhoods of zero consisting of open two-sided ideals. Then the following classes of complexes of left \mathfrak{R} -contramodules coincide:

- (1) all pure acyclic complexes of flat \Re -contramodules;
- (2) the closure of the class of all contractible complexes of countably presentable flat ℜ-contramodules under extensions and directed colimits;
- (3) all \aleph_1 -directed colimits of totalizations of short exact sequences of complexes of countably presentable flat \Re -contramodules.

Proof. The totalization of a short exact sequence of complexes is an extension of the two cones of identity endomorphisms of the rightmost and leftmost complexes in the short exact sequence. Furthermore, the class of all pure acyclic complexes of flat \mathfrak{R} -contramodules is closed under extensions and directed colimits (by Lemmas 8.4(a) and 8.8), and the contractible complexes of flat \mathfrak{R} -contramodules are pure acyclic. This proves the inclusions (3) \subset (2) \subset (1).

Conversely, to prove that $(1) \subset (3)$, recall that all pure acyclic complexes of flat \mathfrak{R} -contramodules are \aleph_1 -directed colimits of pure acyclic complexes of countably presentable flat \mathfrak{R} -contramodules by Corollary 11.4. The exact category G of countably presentable flat \mathfrak{R} -contramodules has homological dimension ≤ 1 by Lemma 13.1(b). Hence the pure acyclic complexes of countably presentable flat \mathfrak{R} -contramodules are absolutely acyclic by Lemma 6.1, and it remains to refer to Proposition 6.2 (for d = 1).

Questions 13.3. The main results of this paper concerning contramodules, such as Theorems 10.1, 12.3, and and 13.2, as well as Propositions 10.2 and 11.3, and Corollary 11.4 assume existence of a countable topology base of two-sided ideals in the topological ring \Re . Are any of these theorems valid for complete, separated topological rings with a countable base of neighborhoods of zero consisting of open *right* ideals? It would be interesting to know; but in fact, even Questions 9.8 and 11.2 seem to be open.

References

- J. Adámek, J. Rosický. Locally presentable and accessible categories. London Math. Society Lecture Note Series 189, Cambridge University Press, 1994.
- [2] S. Bazzoni, M. Cortés-Izurdiaga, S. Estrada. Periodic modules and acyclic complexes. Algebras and Represent. Theory 23, #5, p. 1861–1883, 2020. arXiv:1704.06672 [math.RA]
- [3] S. Bazzoni, M. Hrbek, L. Positselski. Fp-projective periodicity. Journ. of Pure and Appl. Algebra 228, #3, article ID 107497, 24 pp., 2024. arXiv:2212.02300 [math.CT]
- [4] A. Beilinson, V. Drinfeld. Quantization of Hitchin's integrable system and Hecke eigensheaves. February 2000. Available from http://www.math.utexas.edu/~benzvi/Langlands.html or http://math.uchicago.edu/~drinfeld/langlands.html
- [5] D. J. Benson, K. R. Goodearl. Periodic flat modules, and flat modules for finite groups. *Pacific Journ. of Math.* 196, #1, p. 45–67, 2000.
- [6] T. Brzezinski, R. Wisbauer. Corings and comodules. London Mathematical Society Lecture Note Series, 309. Cambridge University Press, Cambridge, 2003.
- [7] T. Bühler. Exact categories. Expositiones Math. 28, #1, p. 1-69, 2010. arXiv:0811.1480
 [math.H0]
- [8] B. Chorny, J. Rosický. Class-locally presentable and class-accessible categories. Journ. of Pure and Appl. Algebra 216, #10, p. 2113-2125, 2012. arXiv:1110.0605 [math.CT]
- [9] L. W. Christensen, H. Holm. The direct limit closure of perfect complexes. Journ. of Pure and Appl. Algebra 219, #3, p. 449-463, 2015. arXiv:1301.0731 [math.RA]
- [10] R. Colpi, K. R. Fuller. Tilting objects in abelian categories and quasitilted rings. Trans. Amer. Math. Soc. 359, #2, p. 741–765, 2007.
- P. Čoupek, J. Šťovíček. Cotilting sheaves on Noetherian schemes. Math. Zeitschrift 296, #1-2, p. 275-312, 2020. arXiv:1707.01677 [math.AG]
- [12] A. I. Efimov, L. Positselski. Coherent analogues of matrix factorizations and relative singularity categories. Algebra and Number Theory 9, #5, p. 1159–1292, 2015. arXiv:1102.0261 [math.CT]
- [13] S. Eilenberg, J. C. Moore. Limits and spectral sequences. Topology 1, p. 1–23, 1962.
- [14] E. E. Enochs, J. R. García Rozas. Flat covers of complexes. Journ. of Algebra 210, #1, p. 86– 102, 1998.
- [15] S. Estrada, X. Fu, A. Iacob. Totally acyclic complexes. Journ. of Algebra 470, p. 300-319, 2017. arXiv:1603.03850 [math.AC]
- [16] S. Estrada, P. A. Guil Asensio, S. Odabaşi. A Lazard-like theorem for quasi-coherent sheaves. Algebras and Represent. Theory 16, #4, p. 1193-1205, 2013. arXiv:1109.0439 [math.AG]
- [17] R. Göbel, J. Trlifaj. Approximations and endomorphism algebras of modules. Second Revised and Extended Edition. De Gruyter Expositions in Mathematics 41, De Gruyter, Berlin–Boston, 2012.
- [18] V. E. Govorov. On flat modules (Russian). Sibir. Mat. Zh. 6, p. 300–304, 1965.
- [19] S. Henry. When does $\operatorname{Ind}_{\kappa}(C^{I}) \simeq \operatorname{Ind}_{\kappa}(C)^{I}$? Electronic preprint arXiv:2307.06664 [math.CT].
- [20] B. Keller. Derived categories and their uses. M. Hazewinkel, Ed., Handbook of Algebra, vol. 1, Elsevier, 1996, p. 671–701.
- [21] M. Kontsevich, A. Rosenberg. Noncommutative smooth spaces. The Gelfand Mathematical Seminars 1996-1999, p. 85-108, Birkhäuser Boston, Boston, MA, 2000. arXiv:math.AG/9812158
- [22] M. Kontsevich, A. Rosenberg. Noncommutative spaces and flat descent. Max-Planck-Institut für Mathematik (Bonn) preprint MPIM 2004-36.
- [23] D. Lazard. Autour de la platitude. Bull. Soc. Math. France 97, p. 81–128, 1969.
- [24] S. Mac Lane. Categories for the working mathematician. Second edition. Graduate Texts in Mathematics, 5. Springer-Verlag, New York, 1998.

- [25] D. Murfet. Derived categories of quasi-coherent sheaves. Notes, October 2006. Available from http://www.therisingsea.org/notes
- [26] A. Neeman. The homotopy category of flat modules, and Grothendieck duality. Inventiones Math. 174, #2, p. 255–308, 2008.
- [27] H.-E. Porst. On corings and comodules. Archivum Mathematicum 42, #4, p. 419–425, 2006.
- [28] L. Positselski. Homological algebra of semimodules and semicontramodules: Semi-infinite homological algebra of associative algebraic structures. Appendix C in collaboration with D. Rumynin; Appendix D in collaboration with S. Arkhipov. Monografie Matematyczne vol. 70, Birkhäuser/Springer Basel, 2010. xxiv+349 pp. arXiv:0708.3398 [math.CT]
- [29] L. Positselski. Contraherent cosheaves on schemes. Electronic preprint arXiv:1209.2995v18 [math.CT].
- [30] L. Positselski. Contramodules. Confluences Math. 13, #2, p. 93-182, 2021. arXiv:1503.00991 [math.CT]
- [31] L. Positselski. Contraadjusted modules, contramodules, and reduced cotorsion modules. Moscow Math. Journ. 17, #3, p. 385-455, 2017. arXiv:1605.03934 [math.CT]
- [32] L. Positselski. Flat ring epimorphisms of countable type. Glasgow Math. Journ. 62, #2, p. 383–439, 2020. arXiv:1808.00937 [math.RA]
- [33] L. Positselski. Contramodules over pro-perfect topological rings. Forum Mathematicum 34, #1, p. 1-39, 2022. arXiv:1807.10671 [math.CT]
- [34] L. Positselski. An explicit self-dual construction of complete cotorsion pairs in the relative context. *Rendiconti Semin. Matem. Univ. Padova* 149, p. 191-253, 2023. arXiv:2006.01778 [math.RA]
- [35] L. Positselski. Semi-infinite algebraic geometry of quasi-coherent sheaves on ind-schemes: Quasi-coherent torsion sheaves, the semiderived category, and the semitensor product. Birkhäuser/Springer Nature, Cham, Switzerland, 2023. xix+216 pp. arXiv:2104.05517 [math.AG]
- [36] L. Positselski. Exact DG-categories and fully faithful triangulated inclusion functors. Electronic preprint arXiv:2110.08237 [math.CT].
- [37] L. Positselski. Differential graded Koszul duality: An introductory survey. Bulletin of the London Math. Society 55, #4, p. 1551-1640, 2023. arXiv:2207.07063 [math.CT]
- [38] L. Positselski. Notes on limits of accessible categories. Electronic preprint arXiv:2310.16773 [math.CT], to appear in *Cahiers de topologie et géométrie différentielle catégoriques*.
- [39] L. Positselski. Locally coherent exact categories. Appl. Categorical Struct. 32, #4, article no. 20, 30 pp., 2024. arXiv:2311.02418 [math.CT]
- [40] L. Positselski. Philosophy of contraherent cosheaves. Electronic preprint arXiv:2311.14179 [math.AG].
- [41] L. Positselski. A contramodule generalization of Neeman's flat and projective module theorem. Electronic preprint arXiv:2408.10928 [math.RA].
- [42] L. Positselski, P. Příhoda, J. Trlifaj. Closure properties of lim C. Journ. of Algebra 606, p. 30– 103, 2022. arXiv:2110.13105 [math.RA]
- [43] L. Positselski, J. Rosický. Covers, envelopes, and cotorsion theories in locally presentable abelian categories and contramodule categories. *Journ. of Algebra* 483, p. 83–128, 2017. arXiv:1512.08119 [math.CT]
- [44] L. Positselski, J. Šťovíček. The tilting-cotilting correspondence. Internat. Math. Research Notices 2021, #1, p. 189-274, 2021. arXiv:1710.02230 [math.CT]
- [45] L. Positselski, J. Šťovíček. Derived, coderived, and contraderived categories of locally presentable abelian categories. *Journ. of Pure and Appl. Algebra* 226, #4, article ID 106883, 2022, 39 pp. arXiv:2101.10797 [math.CT]

- [46] L. Positselski, J. Šťovíček. Coderived and contraderived categories of locally presentable abelian DG-categories. *Math. Zeitschrift* **308**, #1, article no. 14, 70 pp., 2024. arXiv:2210.08237 [math.CT]
- [47] L. Positselski, J. Šťovíček. Flat quasi-coherent sheaves as directed colimits, and quasi-coherent cotorsion periodicity. Electronic preprint arXiv:2212.09639 [math.AG].
- [48] G. Raptis, J. Rosický. The accessibility rank of weak equivalences. Theory and Appl. of Categories 30, no. 19, p. 687-703, 2015. arXiv:1403.3042 [math.AT]
- [49] A.-M. Simon. Approximations of complete modules by complete big Cohen-Macaulay modules over a Cohen-Macaulay local ring. Algebras and Representation Theory 12, #2–5, p. 385–400, 2009.
- [50] N. Spaltenstein. Resolutions of unbounded complexes. Compositio Math. 65, #2, p.121–154, 1988.
- [51] M. Sweedler. The predual theorem to the Jacobson-Bourbaki theorem. Trans. of the Amer. Math. Soc. 213, p. 391–406, 1975.
- [52] B. Totaro. The resolution property for schemes and stacks. Journ. für die reine und angew. Mathematik 577, p. 1-22, 2004. arXiv:math.AG/0207210
- [53] F. Ulmer. Bialgebras in locally presentable categories. Preprint, University of Wuppertal, Summer 1977. Available from https://math.cas.cz/~positselski or https://ncatlab. org/nlab/files/Bialgebras_in_locally_presentable_categories.pdf
- [54] A. Yekutieli. On flatness and completion for infinitely generated modules over noetherian rings. Communicat. in Algebra 39, #11, p. 4221-4245, 2011. arXiv:0902.4378 [math.AC]

INSTITUTE OF MATHEMATICS, CZECH ACADEMY OF SCIENCES, ŽITNÁ 25, 115 67 PRAHA 1, CZECH REPUBLIC

Email address: positselski@math.cas.cz