Low energy spectrum of the XXZ model coupled to a magnetic field

Simone Del Vecchio*, Jürg Fröhlich[†], Alessandro Pizzo[‡], and Alessio Ranallo [§]

November 15, 2023

Abstract

It is shown that, for a class of Hamiltonians of *XXZ* chains in an external magnetic field that are small perturbations of an Ising Hamiltonian, the spectral gap above the ground-state energy remains strictly positive when the perturbation is turned on, uniformly in the length of the chain. The result is proven for both the *ferromagnetic* and the *antiferromagnetic* Ising Hamiltonian; in the latter case the external magnetic field is required to be small, and for an even number of sites the two-fold degenerate ground-state energy of the unperturbed Hamiltonian may split into two energy levels whose difference is small. This result is proven by using a new, quite subtle refinement of a method developed in earlier work and used to iteratively block-diagonalize Hamiltonians of ever larger subsystems with the help of local unitary conjugations. One novel ingredient of the method presented in this paper consists of the use of Lieb-Robinson bounds.

1 Introduction

In this paper we study short-range perturbations of the Hamiltonian of an Ising chain. An example covered by our analysis is the celebrated XXZ chain, whose Hamiltonian includes nearest-neighbour interactions of quantum spins (with spin 1/2) with coupling constants of two different strengths, a large "parallel" one, J, in interaction terms among z-components of neighboring spins, and a small, "perpendicular" one in interaction terms among perpendicular (x- and y-) components. In addition, interaction terms of the spins in the chain with an external magnetic field of strength h in the z-direction may be included in the Hamiltonian. In this paper we focus our attention on this particular class of models, because they have attracted quite a lot of interest. But our methods can be applied to a considerably more general family of models, as specified later on; see Remark 1.5. The results established in this paper cover perturbations of ferromagnetic (J > 0) and antiferromagnetic (J < 0) Ising Hamiltonians, provided that the perpendicular coupling constant is a small parameter as compared to |J|. With regard to the strength, h, of the magnetic field, the regimes we study in this paper depend on the sign of J:

^{*}Dipartimento di Matematica, Università degli Studi di Bari, Italy / email: simone.delvecchio@uniba.it [†]Institut für Theoretiche Physik, ETH-Zürich , Switzerland / email: juerg@phys.ethz.ch

[‡]Dipartimento di Matematica, Università di Roma "Tor Vergata", Italy / email: pizzo@mat.uniroma2.it

[§]Dipartimento di Matematica, Università di Roma "Tor Vergata", Italy / email: ranallo@mat.uniroma2.it

for J < 0, *h* is assumed to be small enough such that an antiferromagnetic ordering of spins is preserved.

Our analysis relies on an iterative block-diagonalization of local Hamiltonians supported in ever longer intervals (i.e., subsets of successive sites) of the lattice, with the help of unitary (Lie Schwinger) conjugations. The sequence of such conjugations applied to Hamiltonians of subsystems of the chain yields a "flow" of transformed Hamiltonians that contain effective potentials supported in intervals of arbitrary length. In this respect our method of analysis is an elaboration on the one introduced in [FP], where perturbations of *ultralocal* Hamiltonians of quantum chains have been considered; generalizations of the technique to higher dimensional lattices, complex coupling constants, and to bosonic Hamiltonians have appeared in [DFPR1], [DFPR2], [DFPR3], and [DFP]. The novelties introduced in the present paper enable us to study perturbations of Hamiltonians that are *not* ultralocal but involve short-range interactions. We also remark that the Hamiltonian of the antiferromagnetic chain does not satisfy all assumptions of most of the methods earlier introduced in the literature to study quantum spin chains.

The unperturbed Hamiltonian of *antiferromagnetic* chains, i.e., the sum of the Ising Hamiltonian and the interaction term with the external magnetic field, has a ground-state subspace which is two-dimensional under natural assumptions on the size of h and on the parity of the number of sites in the chain. But, in contrast to models such as the celebrated "AKLT model" (see [AKLT]), no *local quantum topological order* condition (see [MN]) holds.

Our analysis enables us to prove that if J > 0 (ferromagnetic Hamiltonian), uniformly in the length of the chain, a spectral gap (of order J + h) above the ground-state energy of the unperturbed Hamiltonian persists when the perpendicular interaction terms are added, provided the coupling constant of the latter is sufficiently small. For the antiferromagnetic model, with J < 0 and sufficiently small h, we show that, for chains with an odd number of sites there is a gap of order h above the ground-state energy provided the coupling constant t of the perpendicular interaction is sufficiently small compared to h, similarly to the ferromagnetic case. For antiferromagnetic chains with an even number of sites, the two-fold degenerate ground-state energy of the unperturbed Hamiltonian may split into two energy levels whose difference is, however, bounded above by a fractional power of the coupling constant t, which is assumed much smaller than |J| - h. This splitting is a boundary effect.

In order to cope with the feature that the unperturbed Hamiltonian consists of terms which are *local* but *not ultralocal* (i.e., not on-site), we make use of an argument exploiting Lieb Robinson bounds to control the flow of effective potentials. A similar idea is exploited in our analysis of the AKLT model (see [DFPRa]).

The analysis presented in this paper only involves spin operators, i.e., no domain-wall representation of the models is used. Thus, in the interaction terms of the spins with the magnetic field, no *non-local* operators appear. The connection of the models studied in this paper to models of interacting fermions can be made by using a Klein-Jordan-Wigner transformation to fermionic operators in our analysis of quantum spin chains. This enables us to draw conclusions on the low-lying energy spectrum of one-dimensional systems of (spinless) fermions with Hubbard-type interactions that can be either repulsive (corresponding to an antiferromagnetic *XXZ* spin chain) or attractive (corresponding to a ferromagnetic chain).

This paper can be viewed to be a contribution to a research area pertaining to the characterization of *"topological phases";* see, e.g., [BN, BH, BHM, DS, K, KT, LMY, NSY, NSY2, NSY3, H, O, S], which has been pursued very actively in recent years. In these studies, known techniques and novel ones have been tailored to the study of the low-lying energy spectrum of quantum lattice systems. To our knowledge, the antiferromagnetic *XXZ* chains in an external magnetic field have not been studied previously with mathematically rigorous techniques, except for some results within the range of Bethe ansatz techniques; see [F]. Recent numerical results based on tensor network renormalization techniques agree with Bethe ansatz benchmarks for some values of the parameters in the Hamiltonian of such chains; see [RW] and references therein.

For the ferromagnetic XXZ Hamiltonian of an infinite chain in the absence of an external magnetic field, and for an arbitrary ratio greater than 1 between the "parallel" and the "perpendicular" coupling constants, proofs of a strictly positive spectral energy gap above the ground-state energies can be found in [KN], for spin $\frac{1}{2}$, and in [KNS] for arbitrary spin. Further results on the low lying spectrum of this Hamiltonian have been derived in [NSt], [NSS], and [FS].

1.1 Definition of the model

We consider a one-dimensional lattice, Λ , consisting of an arbitrary number, $N < \infty$, of sites. With every site $j \in \Lambda$ we associate a Hilbert space $\mathcal{H}_j \simeq \mathbb{C}^2$. By $\sigma_j = (\sigma_j^x, \sigma_j^y, \sigma_j^z)$ we denote the Pauli matrices acting on \mathcal{H}_j , for every $j \in \{1, \ldots, N\}$. The Hilbert space of the entire chain is given by

$$\mathcal{H}^{(N)} := \bigotimes_{j=1}^{N} \mathcal{H}_j.$$
(1.1)

We consider small, finite-range perturbations of both the ferromagnetic (J > 0) and the antiferromagnetic (J < 0) Ising Hamiltonian, H^0_{Λ} , with a magnetic field of strength h > 0 in the -z-direction, where

$$H^{0}_{\Lambda} := -J \sum_{i=1}^{N-1} \sigma^{z}_{i} \sigma^{z}_{i+1} - h \sum_{i=1}^{N} \sigma^{z}_{i}.$$
(1.2)

In particular, we study the Hamiltonian of the *XXZ* chain in a magnetic field, which is given by

$$K_{\Lambda} \equiv K_{\Lambda}(t) := -J \sum_{i=1}^{N-1} \sigma_i^z \sigma_{i+1}^z + \frac{t}{2} \sum_{i=1}^{N-1} (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) - h \sum_{i=1}^N \sigma_i^z$$
(1.3)

where $t \in \mathbb{R}$ is a coupling constant with |t| small as compared to |J| and, in some cases, to *h*. (We could also consider tilting the external magnetic field a little.) More precisely, we make the following assumptions:

i) For the ferromagnetic coupling, J > 0,

$$|t| \ll J + h, \quad \sqrt{|t|} < \frac{h}{J}. \tag{1.4}$$

ii) For the antiferromagnetic coupling, J < 0,

$$|t| \ll |J| - h, \quad h < \frac{|J|}{2};$$
 (1.5)

if Λ has an odd number of sites, we may also consider $|t| \ll h$; see Remark 1.4.

1.1.1 Local Hamiltonians

To implement the iterative *local* Lie-Schwinger block-diagonalization method, which will be our main tool to study the low-energy spectrum of the Hamiltonian in (1.3), it is useful to define unperturbed *local* Hamiltonians associated with intervals $\mathcal{I} \subseteq \Lambda = \{1, ..., N\}$ of arbitrary length (where an interval is a subset of Λ consisting of successive sites). We say that a selfadjoint operator A is a *local "observable" supported* in the interval \mathcal{I} if

$$A = A' \otimes \mathbb{1} \tag{1.6}$$

where A' acts on $\bigotimes_{j \in I} \mathcal{H}_j$, and $\mathbb{1}$ is the identity operator on $\bigotimes_{j \notin I} \mathcal{H}_j$. The unperturbed local Hamiltonian associated with I is

$$H_I^0 := -J \sum_{i:i,i+1 \in I} \sigma_i^z \sigma_{i+1}^z - h \sum_{i \in I} \sigma_i^z.$$

$$(1.7)$$

Since $H_{\mathcal{I}}^0$ is *not* additive under taking the union of adjacent intervals, i.e.,

$$H^{0}_{I\cup I'} \neq H^{0}_{I} + H^{0}_{I'}$$
(1.8)

where $\mathcal{I} \cap \mathcal{I}'$ consists of one single site, we will need auxiliary Hamiltonians related to $H_{\mathcal{I}}^0$ ($\mathcal{I} \subset \Lambda$) but enjoying the additivity property, in order to control the effective interaction terms created by the block-diagonalization algorithm introduced below. To this end, we define

$$H_{I}^{C} := -\sum_{i:i,i+1 \in I} \left\{ J \sigma_{i}^{z} \sigma_{i+1}^{z} + \frac{h}{2} [\sigma_{i}^{z} + \sigma_{i+1}^{z}] \right\},$$
(1.9)

where the superscript C stands for "combinatorial". It is easily verified that

$$H_{I}^{0} = H_{I}^{C} - \frac{h}{2}\sigma_{i_{-}}^{z} - \frac{h}{2}\sigma_{i_{+}}^{z}$$
(1.10)

where i_{\pm} are the endpoint sites of the interval I.

For an interval I, we denote with card(I) the number of sites contained in I. By $|\uparrow\rangle$ and $|\downarrow\rangle$ we denote the eigenvectors of σ^z corresponding to the eigenvalues 1 and -1, respectively. Similarly, the symbols

$$|\uparrow\cdots\uparrow\rangle \ , \ |\downarrow\uparrow\cdots\uparrow\rangle \ , \ |\downarrow\uparrow\downarrow\uparrow\cdots\uparrow\rangle \ (1.11)$$

stand for vectors in $\mathcal{H}_I := \bigotimes_{i \in I} \mathcal{H}_i$ consisting of tensor products of M vectors $|\uparrow\rangle$ and/or $|\downarrow\rangle$.

In Propositions 1.1 and 1.2 below, we identify the ground-states and the spectral gaps above the ground-state energies of the Hamiltonians H_{I}^{0} and H_{I}^{C} .

Proposition 1.1. Under the assumption that h and J are positive, the Hamiltonians H_I^0 and H_I^C have only one ground-state, denoted Ψ_I , corresponding to the vector

$$|\uparrow\uparrow\cdots\uparrow\rangle. \tag{1.12}$$

Moreover, under the condition that $\frac{J}{h} + \frac{3}{2} < card(I)$, the spectral gaps above the ground-state energies of the Hamiltonians H_I^0 and H_I^C are equal to 2J + 2h and 2J + h, respectively.

The corresponding proposition for the Hamiltonians with antiferromagnetic exchange coupling constant (J < 0) reads as follows.

Proposition 1.2. Let card(I) be even. Under the assumption -J > h > 0, H_I^0 and H_I^C have both two ground-states, Ψ_I^A and Ψ_I^B , corresponding to the vectors

$$|\uparrow\downarrow\uparrow\downarrow\uparrow\cdots\uparrow\downarrow\rangle \quad , \quad |\downarrow\uparrow\downarrow\uparrow\downarrow\cdots\downarrow\uparrow\rangle, \tag{1.13}$$

respectively. The spectral gap above the ground-state energy is equal to 2|J| - 2h for $H_{\overline{I}}^0$ and to 2|J| - h for $H_{\overline{I}}^C$.

Let card(I) be odd. Under the assumption -J > h > 0, H_I^C has two ground-states, Ψ_I^A and Ψ_I^B , corresponding to the vectors

$$|\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\cdots\downarrow\uparrow\rangle \quad , \quad |\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\cdots\uparrow\downarrow\rangle, \tag{1.14}$$

respectively, whereas Ψ_I^A is the only ground-state of H_I^0 . For H_I^C , the spectral gap above the ground-state energy is equal to to 2|J| - h. When considering the Hamiltonian H_I^0 we call "spectral gap" the energy difference between the ground-state energy and the spectrum of $H_I^0 \upharpoonright_{V^\perp\{\Psi_I^A, \Psi_I^B\}}$, where $\bigvee^{\perp}\{\Psi_I^A, \Psi_I^B\}$ is the orthogonal complement of the subspace generated by Ψ_I^A and Ψ_I^B . This energy difference is given by 2|J|. Moreover, the distance between the eigenvalue of H_I^0 associated with Ψ_I^B and the spectrum of $H_I^0 \upharpoonright_{\bigvee^{\perp}\{\Psi_I^A, \Psi_I^B\}}$ is given by 2|J| - 2h.

The statements described in Propositions 1.1 and 1.2 are summarized in the table below¹

| | J > 0 | J < 0, odd # of sites | J < 0, even # of sites |
|---------------------------|---------------------------------|---|---|
| Ground-states of H_I^C | $ \uparrow\cdots\uparrow angle$ | $ \uparrow\downarrow\cdots\downarrow\uparrow\rangle$ and $ \downarrow\uparrow\cdots\uparrow\downarrow\rangle$ | $ \uparrow\downarrow\cdots\downarrow\rangle$ and $ \downarrow\uparrow\cdots\uparrow\rangle$ |
| Ground-states of H_I^0 | $ \uparrow\cdots\uparrow angle$ | $ \uparrow\downarrow\cdots\downarrow\uparrow angle$ | $ \uparrow\downarrow\cdots\downarrow\rangle$ and $ \downarrow\uparrow\cdots\uparrow\rangle$ |
| Spectral gap of H_I^C | 2J + h | 2 J - h | 2 J - h |
| Spectral gap of H^0_{T} | 2J + 2h | 2 J | 2 J - 2h |

Remark 1.3. We note that, for antiferromagnetic exchange couplings (J < 0), the Hamiltonian H^0_{Λ} does not have the property that $Ker(H^0_{\Lambda}) \subseteq Ker(H^0_{I})$, for all $I \subset \Lambda$. Indeed, the vectors Ψ^A_{Λ} and Ψ^B_{Λ} are eigenvectors of H^0_{I} but the corresponding eigenvalues coincide only if the number of sites of the interval I is even, i.e., they are different whenever the number of sites in I is odd. In contrast, upon subtracting a Λ - dependent constant, the Hamiltonian H^C_{Λ} is frustration free according to the usual definition (see [MN]).

1.2 Statement of the main result and organization of the paper

The results proven in this paper are summarized in the theorem below.

Theorem

We consider the Hamiltonian $K_{\Lambda}(t)$ of an *XXZ* model defined in in (1.3) on a chain Λ , with $\operatorname{card}(\Lambda) \equiv N$.

¹As specified in Proposition 1.2, for an antiferromagnetic chain with an odd number of sites the expression "spectral gap" of $H_{\mathcal{I}}^0$ refers to the energy difference between the ground-state energy and the spectrum of $H_{\mathcal{I}}^0 \upharpoonright_{\mathcal{I}}^{-1} (\Psi_{\mathcal{I}}^A, \Psi_{\mathcal{I}}^B)$.

- (a) If J > 0 there exists a constant $\bar{t} > 0$ depending on J and h, but independent of $N > \frac{J}{h} + \frac{3}{2}$, such that, for all $|t| < \bar{t}$, the ground-state energy E_{Λ} of the Hamiltonian K_{Λ} in (1.3) is non-degenerate and the spectral gap above the ground-state energy is bounded below by $2J + 2h O(\sqrt{|t|})$.
- (b) If J < 0 then, for |J| > 2h, there exists a t

 > 0 depending on J and h, but independent of N, such that, for all |t| < t
 the following statements hold.
 - If Λ has an odd number of sites, the set $\mathfrak{S} := \sigma(K_{\Lambda}) \cap [E_{\Lambda}, E_{\Lambda} + 2|J| O(\sqrt{|t|})]$, where $\sigma(K_{\Lambda})$ is the spectrum of K_{Λ} and E_{Λ} its ground-state energy, consists of two points, E_{Λ} and E'_{Λ} , with $E'_{\Lambda} - E_{\Lambda} = 2h - O(\sqrt{|t|})$, and the spectral projection associated with \mathfrak{S} is of rank 2;
 - If Λ has an even number of sites, the set $\mathfrak{S} := \sigma(K_{\Lambda}) \cap [E_{\Lambda}, E_{\Lambda}+2|J|-2h-O(\sqrt{|t|})]$ consists of at most two points, E_{Λ} and E'_{Λ} , with $|E'_{\Lambda}-E_{\Lambda}| \leq O(\sqrt{|t|})$, and the spectral projection associated with \mathfrak{S} is of rank 2.

Remark 1.4. In point (b) of the Theorem above, the dependence on *h* of \bar{t} is only required to ensure the gap $E'_{\Lambda} - E_{\Lambda}$ is given by $2h - O(\sqrt{|t|})$. More precisely, the allowed range of the coupling constant is $|t| \ll |J| - h$, hence our theorem also applies to a regime where the gap $E'_{\Lambda} - E_{\Lambda}$ may close.

Remark 1.5. For J < 0 (antiferromagnetic chain), similar results hold if the perturbation term $\sum_{i=1}^{N-1} (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y)$ is replaced by an arbitrary *translation-invariant*, *short-range* perturbation. For J > 0, the translation invariance of the perturbation term is not required.

Remark 1.6. The ferromagnetic *XXZ* model with h = 0 is not covered by the theorem formulated above, because our strategy (for the ferromagnetic chain) uses the non-degeneracy of the ground-state subspace of the local Hamiltonians, which holds for h > 0. However, we can easily deal with the ferromagnetic model in a vanishing magnetic field (h = 0) and show that the ground-state energy is doubly degenerate, and that the spectral gap above the ground-state energy is bounded below by $2J - O(\sqrt{|t|})$; see Remark 4.12.

Remark 1.7. The theorem stated above also holds for the antiferromagnetic *XXZ* model with spins coupled to a staggered magnetic field in the *z*-direction (see [R]), whose Hamiltonian can be obtained from the Hamiltonian K_{Λ} with ferromagnetic couplings (J > 0) by a unitary conjugation that flips the σ^z operators either on all sites with *i* even, or on all sites with *i* odd.

Remark 1.8. We expect that the techniques developed in this paper enable us to extend the theorem stated above to chains of quantum spins of arbitrary spin $s \ge 1$. Indeed, the spins Ising Hamiltonians with spins coupled to a magnetic field in the z-direction have a lowlying energy spectrum very similar to the ones described in Propositions 1.1 and 1.2 for the ferromagnetic and antiferromagnetic models with $s = \frac{1}{2}$, respectively. Since the properties stated in these propositions, as well as the Lieb-Robinson bounds considered later, are the only relevant ingredients that will be required, the whole procedure used to prove our main results should apply, word-by-word, to the more general class of models alluded to above.

Summary of contents. In Sect. 2, we begin with the definition of the local interaction terms (Sect. 2.1), supported in *t*-dependent subsets of lattice sites, on which we will apply our block-diagonalization procedure. In Sect. 2.2, we present a brief review of the method originally introduced for perturbations of ultra-local Hamiltonians. Next, in Sects. 2.3 and 2.4, we present the key ideas enabling us to cope with complications – as compared to the Hamiltonians treated in [FP] – arising in the implementation of a local block-diagonalization procedure, which are related to: i) the nearest-neighbour interaction structure of the Ising Hamiltonian; and ii) the

degeneracy (in the antiferromagnetic model) of the ground-state energy of the unperturbed Hamiltonians H_{τ}^{C} .

In Sect. 3, we introduce the algorithm that determines the effective potentials of the transformed Hamiltonians at each step of the block-diagonalization flow.

In Sect. 4, we quantitatively control the effective potentials produced along the blockdiagonalization flow (Sect. 4.2) and analyze the spectrum of the ferromagnetic and antiferromagnetic Hamiltonians $K_{\Lambda}(t)$, for arbitrary large N (Sect. 4.1), as described in the theorem above.

Notation

1) We use the same symbol for an operator $O_{\mathcal{I}}$ acting on $\otimes_{i \in \mathcal{I}} \mathcal{H}_i$ and the corresponding operator acting on the entire Hilbert space $\mathcal{H}^{(N)}$ that is obtained from $O_{\mathcal{I}}$ by tensoring with the identity matrix on the Hilbert spaces of all remaining sites.

2) With the symbol " \subset " we denote strict inclusion, otherwise we use the symbol " \subseteq ".

3) With the symbol $O(\sqrt{|t|})$ we denote a quantity which, in absolute value, is bounded above by $\sqrt{|t|}$ multiplied by a constant possibly depending on further parameters entering the definition of the models, but independent of the number of sites, *N*, of the chain.

4) We denote by $\langle \cdot, \cdot \rangle$ the scalar product on $\mathcal{H}^{(N)}$.

5) We denote the identity matrix by $\mathbb{1}$ or 1, interchangeably.

2 **Proof strategy**

For convenience of notation, and without loss of generality, we assume that t > 0.

2.1 Local interaction terms and projections

Since we consider the term

$$\frac{t}{2} \sum_{i=1}^{N-1} (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y)$$
(2.15)

as a small perturbation of the remaining part of the Hamiltonian K_{Λ} , we split the operator in (2.15) into terms localized in *N*-independent, but *t*-dependent intervals. For, our methods involve introducing a *macroscopic* lattice with a lattice spacing of order $\sqrt{t^{-1}}$, as explained below.

Without loss of generality, we assume t positive and such that

$$(N-1)\cdot\sqrt{t}\in\mathbb{N},\ \frac{\sqrt{t^{-1}}}{3}\in\mathbb{N}.$$
 (2.16)

For the antiferromagnetic Hamiltonian, $\sqrt{t^{-1}}$ is assumed to be an odd integer w.l.o.g. . We introduce a *macroscopic* (finite) lattice with left endpoint X = 1, right endpoint X = N, and spacing $\sqrt{t^{-1}}$. The M^{th} site of this lattice is the point

$$1 + (M-1) \cdot \sqrt{t^{-1}}$$
, with $1 \le M \le (N-1) \cdot \sqrt{t} + 1$. (2.17)

The set of successive sites I at *position* (i.e., starting from) J =: Q(I) and of *length* $K =: \ell(I)$, in units of $\sqrt{t^{-1}}$, is the interval whose endpoints coincide with the sites M = J and M = J + K of the macroscopic lattice.

Definition 2.1. We define \mathfrak{I} to be the set of intervals $I \subseteq \Lambda$ whose left endpoint is the site $1 + (J - 1)\sqrt{t^{-1}}$, for some $J \in \mathbb{N}$, and whose length is given by $K \cdot \sqrt{t^{-1}}$ (hence card(I) = $K \cdot \sqrt{t^{-1}} + 1$), for some $K \in \mathbb{N}$. We set

$$Q(I) := J \quad , \quad \ell(I) := K.$$

Hence the interval I, with Q(I) = J and $\ell(I) = K$, is

$$\left[1 + (Q(I) - 1)\sqrt{t^{-1}}, 1 + (Q(I) + \ell(I) - 1)\sqrt{t^{-1}}\right].$$
(2.18)

Figure 1: The picture illustrates the interval I, with Q(I) = J and $\ell(I) = 1$, and the subsequent one, I_{+1} .

For $I \in \mathfrak{I}$, with $\ell(I) = 1$, we define

$$V_{I} := \frac{1}{2\sqrt{t^{-1}}} \sum_{i:i,i+1 \in I} (\sigma_{i}^{x} \sigma_{i+1}^{x} + \sigma_{i}^{y} \sigma_{i+1}^{y}).$$
(2.19)

Our definition of intervals I, with $\ell(I) = 1$, imply the bound

$$\|V_{I}\| \le 1. (2.20)$$

In the implementation of the block-diagonalization procedure it is convenient to re-write the Hamiltonian $K_{\Lambda}(t)$ by making use of the definitions introduced above, namely

$$K_{\Lambda}(t) = H_{\Lambda}^{0} + \sqrt{t} \sum_{I \in \mathfrak{I}; \, \ell(I)=1} V_{I} \,.$$

$$(2.21)$$

In addition to (2.16), for the ferromagnetic models we require that

$$(N - \frac{3}{2} >) \sqrt{t^{-1}} > \frac{J}{h}, \qquad (2.22)$$

so that Proposition 1.1 holds for all the unperturbed Hamiltonians H_I^0 . The block-diagonalization used below is w.r.t. spectral projections supported in intervals I, which we define next.

Definition 2.2. By $P_{\mathcal{I}}^{(-)}$ we denote the orthogonal projection onto the ground-state subspace of $H_{\mathcal{I}}^C$, and we set

$$P_{I}^{(+)} := \mathbb{1} - P_{I}^{(-)}. \tag{2.23}$$

Analogous definitions will be employed for projections associated with other subsets of the lattice Λ .

For the following a total ordering relation on the set \Im will turn out to be useful according to which shorter intervals precede longer ones. This ordering relation is defined below.

Definition 2.3. An ordering relation ">" on \Im is specified as follows.

$$I > I'$$
 if $\ell(I) > \ell(I')$ or if $\ell(I) = \ell(I')$ and $Q(I) > Q(I')$. (2.24)

The symbol I_{-1} (I_{+1} , resp.) stands for the element of \Im preceding (following, resp.) I in the given ordering. For convenience, we define the symbol I_0 to be the element preceding the smallest element of \Im in the given ordering. Note that the biggest element in this ordering is the entire lattice Λ .

2.2 Outline of the block-diagonalization flow

The study of the low-energy spectrum of the XXZ Hamiltonians introduced in (1.3) is based on an extension and refinement of the local Lie Schwinger block-diagonalization procedure introduced in [FP]. Starting from the decomposition of the interaction terms into potentials V_I , and taking the ordering relation introduced in Definition 2.3 into account, we will construct an iterative block-diagonalization algorithm based on unitary (Lie-Schwinger) conjugations supported in intervals of the set \Im . These conjugations are denoted by e^{Z_I} .

In the very first step, corresponding to the interval I with Q(I) = 1 and $\ell(I) = 1$, the conjugation is such that

$$e^{Z_{I}} \left(H_{I}^{0} + \sqrt{t} V_{I} \right) e^{-Z_{I}} = H_{I}^{0} + \sqrt{t} V_{I}^{I}, \qquad (2.25)$$

where the new potential V_I^I is block-diagonal w.r.t. $P_I^{(-)}$ and $P_I^{(+)} := \mathbb{1} - P_I^{(-)}$, i.e.,

$$V_{I}^{I} = P_{I}^{(-)} V_{I}^{I} P_{I}^{(-)} + P_{I}^{(+)} V_{I}^{I} P_{I}^{(+)} .$$
(2.26)

It is evident that the action of the conjugation on the remaining terms of the Hamiltonian K_{Λ} may create new terms. For example, for \mathcal{I}' such that $Q(\mathcal{I}') = 2$ and $\ell(\mathcal{I}') = 1$, we have that

$$e^{Z_{I}} \sqrt{t} V_{I'} e^{-Z_{I}} = \sqrt{t} V_{I'} + \sqrt{t} \Delta V_{I \cup I'}(t)$$
(2.27)

where, in general, $\Delta V_{I \cup I'}(t)$ is a non-zero operator supported in the longer interval $I \cup I'$; indeed, $Q(I \cup I') = 1$ and $\ell(I \cup I') = 2$.

For a Hamiltonian whose unperturbed part is *ultralocal*, i.e., consists of on-site terms only, it is shown in [FP] how effective potentials, supported in intervals of arbitrary length belonging to \Im , are created in subsequent steps of the block-diagonalization, starting from the first sequence of steps in which the potentials associated with intervals of length $\ell(I) = 1$ are block-diagonalized. The control of their norms relies on the fact that the number of growth processes yielding an effective potential supported in an interval I' can be bounded by $const^{\ell(I')}$. In estimating the norm of a potential supported in the interval I'. Indeed, each factor of Z_I appearing

in commutators is proportional to \sqrt{t} . Hence, for *t* sufficiently small, the norm of any effective potential has power law decay in *t* with an exponent proportional to the length of the interval in which the potential under consideration is supported.

The block-diagonalization procedure terminates with a final Hamiltonian, unitarily conjugated to the original Hamiltonian of the chain, with the property that each effective potential appearing in the final Hamiltonian is block-diagonal w.r.t. to the projections in Definition 2.2 associated with the support of the potential. Hence the final Hamiltonian is block-diagonal w.r.t. to the projections $P_{\Lambda}^{(-)}$, $P_{\Lambda}^{(+)}$. Consequently, the low-energy spectrum of the original Hamiltonian can be controlled. For a more detailed overview of the block-diagonalization procedure for *ultralocal* unperturbed Hamiltonians see Section 2.1 in [DFPRa].

For the models studied in the present paper, there are several complications arising when one attempts to construct the block-diagonalization flow following the strategy in [FP]. Some of these complications become already visible in the study of the AKLT model (see [DFPRa]).

2.3 "Hooked" unperturbed terms and Lieb Robinson bounds

One complication stems from the property of the unperturbed local Hamiltonians H_I^0 considered in this paper that they are *not ultralocal*, i.e., do not consist only of on-site terms. This implies, for example, that in the very first step of the block-diagonalization procedure, i.e., in the step corresponding to the interval I with $Q(I) = \ell(I) = 1$, the conjugation

$$e^{Z_{\bar{I}}}\left\{K_{\Lambda} - (H^0_{\bar{I}} + \sqrt{t}V_{\bar{I}})\right\}e^{-Z_{\bar{I}}}$$
(2.28)

includes terms, such as

$$e^{Z_{I}} \left[-J\sigma_{i_{+}}^{z}\sigma_{i_{+}+1}^{z} \right] e^{-Z_{I}} = -J\sigma_{i_{+}}^{z}\sigma_{i_{+}+1}^{z} + \sqrt{t} \sum_{n=1}^{\infty} \frac{1}{n!} ad^{n} Z_{I} \left(\frac{-J\sigma_{i_{+}}^{z}\sigma_{i_{+}+1}^{z}}{\sqrt{t}} \right),$$
(2.29)

where i_+ is the right endpoint of \mathcal{I} in the microscopic lattice, and

$$adA(B) := [A, B], \qquad ad^{n}A(B) := [A, ad^{n-1}A(B)], \text{ for } n \ge 2.$$
 (2.30)

Hence a new potential is created whose support extends over the enlarged set $I \cup \{i_+ + 1\}$. In order to gain control over the flow, we have to verify that the contribution to the new potential

$$\sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_I(\frac{-J\sigma_{i_+}^z \sigma_{i_++1}^z}{\sqrt{t}})$$
(2.31)

that needs to be block-diagonalized, i.e., the off-diagonal part w.r.t. the two projections

$$P_{I\cup\{i_{+}+1\}}^{(-)}$$
 and $P_{I\cup\{i_{+}+1\}}^{(+)}$, (2.32)

has an operator norm bounded by $t^{\rho \cdot \ell(\mathcal{I}')}$, where ρ is a universal constant, and $\ell(\mathcal{I}') = 2$ is the length of the shortest interval in \Im containing the support of the effective potential created by the conjugation. In fact, this decay holds trivially for all the terms in the series with the exception of the leading one, i.e., except for the off-diagonal part of

$$ad Z_{I}(\frac{-J\sigma_{i_{+}}^{z}\sigma_{i_{+}+1}^{z}}{\sqrt{t}}),$$
 (2.33)

which we will refer to as a "*hooked*" unperturbed term created in the conjugation generated by the operator Z_I . To control the size of the term in (2.33) we are forced to change the strategy in [FP] by introducing intervals I^* and $\overline{I^*}$, which are *enlargements* of I, and defining suitable corresponding operators Z_{I^*} .

Definition 2.4. On \Im we define the operation * assigning to each interval $I \in \Im, I \neq \Lambda$, a larger interval I^* contained in the lattice. I^* is defined in the following way

$$I^* = \left\{ i \in \Lambda \cap \left[1 + (Q(I) - \frac{4}{3})\sqrt{t^{-1}}, 1 + (Q(I) - \frac{2}{3} + \ell(I))\sqrt{t^{-1}} \right] \right\}.$$
 (2.34)

Moreover, we denote by \mathfrak{I}^* the image of \mathfrak{I} under the map *.

Figure 2: The picture displays the relation between I, with Q(I) = J and $\ell(I) = 1$, and I^* .

Definition 2.5. For each $I^* \in \mathfrak{I}^*$ we define $\overline{I^*}$ the interval obtained from $I^* \in \mathfrak{I}^*$ by joining the nearest two sites (if present) belonging to the lattice, both on the right and on the left. We call these sites $i^*_+ + 1$, $i^*_+ + 2$, $i^*_- - 1$, and $i^*_- - 2$, where i^*_{\pm} are the sites corresponding to the right and the left endpoint of I^* , respectively.



Figure 3: The picture displays the relation between I^* and $\overline{I^*}$. The red dots are the sites (of the microscopic lattice) inside I^* , the blue ones are those added to yield $\overline{I^*}$.

The rationale behind the use of enlarged intervals is as follows. Consider the very first step as described above. In order to block-diagonalize a potential supported in I, we consider the unperturbed operator $H_{I^*}^0$ and the unitary operator Z_{I^*} (defined in Section 3.1) such that

$$e^{Z_{I^*}} \left(H^0_{I^*} + \sqrt{t} V_I \right) e^{-Z_{I^*}} =: H^0_{I^*} + \sqrt{t} V'_{I^*}$$
(2.35)

where by construction V'_{I^*} is block-diagonal w.r.t. $P^{(-)}_{I^*}$, $P^{(+)}_{I^*}$. The counterpart of the term in (2.33) is

$$ad Z_{I^*}(\frac{-J\sigma_{i_*}^{\zeta}\sigma_{i_*+1}^{\zeta}}{\sqrt{t}}).$$
(2.36)

Since the leading order term in the operator Z_{I^*} corresponds (in the ferromagnetic case) to

$$\frac{1}{H_{I^*}^0 - E_{I^*}} P_{I^*}^{(+)} \sqrt{t} V_I P_{I^*}^{(-)} - h.c.$$
(2.37)

where E_{I^*} is defined in (3.10), an argument based on Lieb-Robinson bounds shows that the norm of

$$P_{\overline{I^*}}^{(+)} \Big[\frac{1}{H_{I^*}^0 - E_{I^*}} P_{I^*}^{(+)} \sqrt{t} V_{I} P_{\overline{I^*}}^{(-)}, \frac{-J\sigma_{i_*}^{c_*} \sigma_{i_*}^{c_*} - 1}{\sqrt{t}} \Big] P_{\overline{I^*}}^{(-)}$$
(2.38)

decays as $O(t^{\frac{1}{4}} \cdot ||V_I||)$, as $t \to 0$, where it is crucial that the set I is at a distance (in the microscopic unit) of order at least $\sqrt{t^{-1}}$ from the endpoints of I^* . An analogous procedure holds for the subsequent block-diagonalization steps.

Definition 2.6. For each $I^* \in \mathfrak{I}^*$ we define $\widetilde{I^*} \in \mathfrak{I}$ the smallest interval in \mathfrak{I} containing the interval I^* .

Remark 2.7. Concerning the enlarged intervals introduced in Definitions 2.4, 2.5, and 2.6, in the next sections it is helpful to distinguish *boundary*- from *bulk-intervals*. The first set consists of those intervals I for which one of the endpoints coincides with an endpoint of Λ . For this reason the corresponding enlargements I^* , $\overline{I^*}$ and $\overline{I^*}$ involve only one side of the interval.



Figure 4: The picture shows how I, with Q(I) = J and $\ell(I) = 1$, relates to $\widetilde{I^*}$.

2.4 Degeneracy of the "bulk" ground-state energy level in the antiferromagnetic case

The second complication appearing in the study of the antiferromagnetic case is related to the fact that the ground-state subspace of H_I^0 is degenerate if the number of sites is even, but the ground-state energy splits into two levels as soon as the interval I is enlarged by one more site. This yields some technical difficulties to show that, by adding the perturbation (which is by assumption translation invariant; see Remark 1.5), if the two-degenerate energy level splits the resulting ones remain in fact very close one to each other, where the small gap is due to boundary effects.

The control of this possible energy splitting is carried out in detail in Section 4.1.2, since some definitions are needed. Here we just explain the underlying property we shall use. We call it degeneracy of the *bulk* ground-state energy.

Consider three intervals I, \mathcal{J} and \mathcal{J}' , with \mathcal{J} , $\mathcal{J}' \subset I$, and where \mathcal{J} , \mathcal{J}' are two successive intervals of same length, i.e.,

$$Q(\mathcal{J}) = J, \ Q(\mathcal{J}') = J + 1 \quad , \quad \ell(\mathcal{J}) = \ell(\mathcal{J}') \,. \tag{2.39}$$

Next we consider two operators $W_{\mathcal{J}}$ and $W_{\mathcal{J}'}$ with the property $W_{\mathcal{J}'} = \tau_1(W_{\mathcal{J}})$ where τ_n is the natural shift by *n* edges in the macroscopic lattice (for details, see Definition 3.6). Then, for a macroscopic unit corresponding to $\sqrt{t^{-1}}$ which is by assumption an odd natural number, the following property holds

$$\langle \Psi_{I}^{A}, W_{\mathcal{J}}\Psi_{I}^{A} \rangle = \langle \Psi_{I}^{B}, W_{\mathcal{J}'}\Psi_{I}^{B} \rangle \quad , \quad \langle \Psi_{I}^{B}, W_{\mathcal{J}}\Psi_{I}^{B} \rangle = \langle \Psi_{I}^{A}, W_{\mathcal{J}'}\Psi_{I}^{A} \rangle . \tag{2.40}$$

We assume that each potential $W_{\mathcal{T}}$ is block-diagonal w.r.t.

$$P_{\mathcal{J}}^{(-)A}, P_{\mathcal{J}}^{(-)B}, P_{\mathcal{J}}^{(+)},$$

i.e.,

$$W_{\mathcal{J}} = P_{\mathcal{J}}^{(-)A} W_{\mathcal{J}} P_{\mathcal{J}}^{(-)A} + P_{\mathcal{J}}^{(-)B} W_{\mathcal{J}} P_{\mathcal{J}}^{(-)B} + P_{\mathcal{J}}^{(+)} W_{\mathcal{J}} P_{\mathcal{J}}^{(+)}, \qquad (2.41)$$

where $P_{\mathcal{J}}^{(-)A}$ and $P_{\mathcal{J}}^{(-)B}$ are the spectral projections onto the subspaces generated by $\Psi_{\mathcal{J}}^{A}$ and $\Psi_{\mathcal{J}}^{B}$, respectively. The importance of having operators $W_{\mathcal{J}}$ block-diagonalized as displayed in (2.41) is discussed in Remark 3.1. Next we consider the Hamiltonian

$$\mathfrak{G}_{\mathcal{I}} := H^0_{\mathcal{I}} + \sum_{\mathcal{J} \subset \mathcal{I}}' W_{\mathcal{J}}, \qquad (2.42)$$

where the symbol ' means that we sum over an even number of intervals which are paired, in the sense that they come into pairs consisting of an interval \mathcal{J} and its translated \mathcal{J}' . Then, by using the property in (2.40), we have

$$\langle \Psi_{I}^{A}, \ \mathfrak{G}_{I}\Psi_{I}^{A} \rangle = \langle \Psi_{I}^{B}, \ \mathfrak{G}_{I}\Psi_{I}^{B} \rangle.$$

$$(2.43)$$

3 The block-diagonalization algorithm

We implement the procedure outlined in Sections 2.2 and 2.3 by an algorithm described in Section 3.2. In order to do this we need some definitions collected in the next section.

3.1 Conjugation formulae

The expressions that we are going to define enter the definition of the operators appearing in the transformed Hamiltonian in step I, which will turn out to be of the form

$$K_{\Lambda}^{I}(t) := H_{\Lambda}^{0} + \sqrt{t} \sum_{\mathcal{J} \leq I} V_{\mathcal{J}^{*}}^{I} + \sqrt{t} \sum_{\mathcal{J} > I} V_{\mathcal{J}}^{I}$$
(3.1)

where the reader can notice that the intervals labelling the potentials V are of two types:

- i) if $\mathcal{J} \leq \mathcal{I}$ then *V* is labeled by intervals $\overline{\mathcal{J}^*}$;
- ii) if $\mathcal{J} > \mathcal{I}$ the corresponding *V* is labeled by \mathcal{J} .

This distinction is due to the fact that the first type of potentials, i.e., those corresponding to $\mathcal{J} \leq \mathcal{I}$, are block-diagonalized, and the block-diagonalization is w.r.t. the two projections $P_{\overline{\mathcal{J}^*}}^{(-)}$, $P_{\overline{\mathcal{I}^*}}^{(+)}$ (see Definition 2.2), consequently they can be written as

$$V_{\overline{\mathcal{J}^{*}}}^{I} = P_{\overline{\mathcal{J}^{*}}}^{(+)} V_{\overline{\mathcal{J}^{*}}}^{I} P_{\overline{\mathcal{J}^{*}}}^{(+)} + P_{\overline{\mathcal{J}^{*}}}^{(-)} V_{\overline{\mathcal{J}^{*}}}^{I} P_{\overline{\mathcal{J}^{*}}}^{(-)}.$$
(3.2)

In Section 3.2 we explain how these potentials emerge, step by step, as byproducts of the blockdiagonalization flow. In this respect, the algorithm described in Definition 3.3 prescribes that, in step I, the potential $V_{I}^{I_{-1}}$ gets transformed to a block-diagonalized potential $V_{\overline{I^*}}^{I}$ which does not coincide but includes the leading order term of the Lie-Schwinger series (for details see point b) in Definition 3.3),

$$\sum_{j=1}^{\infty} t^{\frac{j-1}{2}} (V_{I^*}^{I_{-1}})_j^{\text{diag}}, \qquad (3.3)$$

where the operators $(V_{I^*}^{I_{-1}})_j$ are defined below; here "diag" means diagonal part w.r.t. to the two projections $P_{I^*}^{(+)}$, $P_{I^*}^{(-)}$. The key identity which we exploit is

$$e^{Z_{I^*}} \left(G_{I^*} + \sqrt{t} V_{I}^{I_{-1}} \right) e^{-Z_{I^*}} = G_{I^*} + \sqrt{t} \sum_{j=1}^{\infty} t^{\frac{j-1}{2}} \left(V_{I^*}^{I_{-1}} \right)_j^{\text{diag}}$$
(3.4)

where G_{I^*} , $(V_{I^*}^{I_{-1}})_j^{\text{diag}}$, and Z_{I^*} are defined below:

1)

$$G_{I^*} := H^0_{I^*} + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} V^{\overline{I}_{-1}}_{\overline{\mathcal{J}^*}}, \qquad (3.5)$$

for which we remark that if I is a bulk-interval (see Remark 2.7) then the sum in (3.5) does *not* include those \mathcal{J}^* sharing one of the endpoints with I^* , consequently G_{I^*} is localized² in I^* ;

2)

$$(V_{I^*}^{I_{-1}})_1 := V_I^{I_{-1}}, (3.6)$$

²If *I* is a boundary-interval the intervals $\mathcal{J} \subset I$ sharing one of the endpoints of Λ with *I* are included in the sum in (3.5), but G_{I^*} is still localized in I^* .

and, for $j \ge 2$,

$$(V_{I^{*}}^{I_{-1}})_{j} :=$$

$$\sum_{p \ge 2, r_{1} \ge 1, \dots, r_{p} \ge 1; r_{1} + \dots + r_{p} = j} \frac{1}{p!} \operatorname{ad} (Z_{I^{*}})_{r_{1}} \left(\operatorname{ad} (Z_{I^{*}})_{r_{2}} \dots \left(\operatorname{ad} (Z_{I^{*}})_{r_{p}} (G_{I^{*}}) \right) \dots \right) +$$

$$\sum_{p \ge 1, r_{1} \ge 1, \dots, r_{p} \ge 1; r_{1} + \dots + r_{p} = j-1} \frac{1}{p!} \operatorname{ad} (Z_{I^{*}})_{r_{1}} \left(\operatorname{ad} (Z_{I^{*}})_{r_{2}} \dots \left(\operatorname{ad} (Z_{I^{*}})_{r_{p}} ((V_{I^{*}}^{I_{-1}})_{1}) \dots \right) \right).$$

$$(3.7)$$

where the ad (adjoint action) has been defined in (2.30);

3)

$$Z_{I^*} := \sum_{j=1}^{\infty} t^{\frac{j}{2}} (Z_{I^*})_j, \qquad (3.8)$$

where the terms $(Z_{I^*})_j$ are defined accordingly to the rank of $P_{I^*}^{(-)}$, i.e., we distinguish two cases, J > 0 (ferromagnetic behavior) and J < 0 (antiferromagnetic behavior).

<u>J>0</u> In this case, the ground-state subspace of $H_{I^*}^C$ is one-dimensional, thus $P_{I^*}^{(-)}$ is a rank one orthogonal projection onto the subspace generated by Ψ_{I^*} . The $(Z_{I^*})_j$ are defined recursively as follows

$$(Z_{I^*})_j := \frac{1}{G_{I^*} - E_{I^*}} P_{I^*}^{(+)} (V_{I^*}^{I_{-1}})_j P_{I^*}^{(-)} - h.c.$$
(3.9)

where

$$E_{I^*} := \langle \Psi_{I^*}, \left\{ H^0_{I^*} + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} V^{\overline{I_{-1}}}_{\overline{\mathcal{J}^*}} \right\} \Psi_{I^*} \rangle.$$
(3.10)

<u>J<0</u> In this case, the ground-state subspace of $H_{I^*}^C$ is two-dimensional, thus $P_{I^*}^{(-)}$ is a rank two orthogonal projection, i.e., $P_{I^*}^{(-)} = P_{I^*}^{(-),A} + P_{I^*}^{(-),B}$ (see their definition below (2.41)), thus the operators $(Z_{I^*})_j$ are defined recursively as follows

$$(Z_{I^*})_j := \left\{ \frac{1}{G_{I^*} - E_{I^*}^A} P_{I^*}^{(+)} (V_{I^*}^{I_{-1}})_j P_{I^*}^{(-),A} + \frac{1}{G_{I^*} - E_{I^*}^B} P_{I^*}^{(+)} (V_{I^*}^{I_{-1}})_j P_{I^*}^{(-),B} \right\} - h.c.$$

$$(3.11)$$

where

$$E_{I^*}^A := \langle \Psi_{I^*}^A, \left\{ H_{I^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} V_{\overline{\mathcal{J}^*}}^{I_{-1}} \right\} \Psi_{I^*}^A \rangle, \tag{3.12}$$

$$E_{I^*}^B := \langle \Psi_{I^*}^B, \left\{ H_{I^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} V_{\overline{\mathcal{J}^*}}^{\overline{I}_{-1}} \right\} \Psi_{I^*}^B \rangle.$$
(3.13)

We shall prove inductively that $\Psi_{I^*}^A$ and $\Psi_{I^*}^B$ are eigenvectors of G_{I^*} ; this implies the identity in (3.4) with the given definition of Z_{I^*} ; see [DFFR].

We point out that the construction of Z_{I^*} requires the control of some (depending on the sign of *J*) of the following gaps:

for
$$J > 0$$
, inf spec $[(G_{I^*} - E_{I^*})P_{I^*}^{(+)}];$ (3.14)

for
$$J < 0$$
, inf spec $[(G_{I^*} - E^A_{I^*})P^{(+)}_{I^*}]$, inf spec $[(G_{I^*} - E^B_{I^*})P^{(+)}_{I^*}]$. (3.15)

We discuss these quantities in Section 4.1.

Remark 3.1. In the antiferromagnetic case, the definition of Z_{I^*} requires the decomposition of $P^{(-)}\mathcal{H}^N$ into the (one-dimensional) subspaces corresponding to the ranges of $P_{I^*}^{(-),A}$ and $P_{I^*}^{(-),B}$. Differently from the case where a *local quantum topological order* condition holds (see [DFPRa]), here we cannot design an algorithm such that $G_{I^*} P_{I^*}^{(-)} = \hat{E}_{I^*} P_{I^*}^{(-)}$, for some \hat{E}_{I^*} , so as to essentially reduce the block-diagonalization to the usual one where $P_{I^*}^{(-)}$ is of rank 1.

This feature makes the control of the two gaps in (3.15) challenging (see Lemma 4.6) and not feasible without a structure where the potentials $V_{\overline{J^*}}^{I_{-1}}$ in (3.5) are block-diagonalized also w.r.t. $P_{\overline{J^*}}^{(-),A}$ and $P_{\overline{J^*}}^{(-),B}$. The control of the two gaps in (3.15) is also possible thanks to the estimate of the energy difference $E_{I^*}^A - E_{I^*}^B$ which turns out to be $O(\sqrt{t})$; see Lemma 4.4. In the proof of Lemma 4.4, the block-diagonalization of the potentials $V_{\overline{J^*}}^{I_{-1}}$ w.r.t. $P_{\overline{J^*}}^{(-),A}$ and $P_{\overline{J^*}}^{(-),B}$ is crucial to exploit the argument explained in Section 2.4. Thanks to the structure of the *enlargements* described above, the block-diagonalization property (also of the Hamiltonian G_{I^*}) with respect to $P_{I^*}^{(-),A}$ and $P_{I^*}^{(-),B}$ is easily granted by prescription b) of the algorithm in Definition 3.3.

3.2 The algorithm: definition and consistency

Next Definition 3.2 provides the basis of the iteration yielding the effective potentials $V_{\mathcal{J}}^{\mathcal{I}}$, step by step, by applying the algorithm in Definition 3.3, where the "steps" are associated with the intervals $\mathcal{I} \in \mathfrak{I}$, and the ordering of the steps follows the ordering relation of the intervals set in Definition 2.3.

Definition 3.2. We set:

• for $\mathcal{J} \in \mathfrak{I}$ such that $\ell(\mathcal{J}) = 1$,

$$V_{\mathcal{J}}^{I_0} := V_{\mathcal{J}} \Big(= \frac{1}{2\sqrt{t^{-1}}} \sum_{i:i,i+1 \in \mathcal{J}} (\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y) \Big);$$
(3.16)

• for $\mathcal{J} \in \mathfrak{I}$ with $\ell(\mathcal{J}) \geq 2$,

a-1) if $I \prec \mathcal{J}$, and $I^* \not\subseteq \mathcal{J}$,

$$V_{\mathcal{J}}^{I_0} := 0.$$
 (3.17)

In the sequel, i_{-}^{*} , i_{+}^{*} are the sites in the microscopic lattice corresponding to the two endpoints of the interval \mathcal{I}^{*} . In (3.21), (3.28), and (3.29) one of the two hooked terms is absent whenever i_{-} or i_{+} (endpoints of \mathcal{I}) coincides with the left or with the right endpoint of Λ , respectively.

Definition 3.3. For the subsequent steps we set:

$$V_{\mathcal{J}}^{\mathcal{I}} := V_{\mathcal{J}}^{\mathcal{I}_{-1}}; \qquad (3.18)$$

a-2) if $I > \mathcal{J}$,

$$V_{\overline{\mathcal{J}^*}}^{I} := V_{\overline{\mathcal{J}^*}}^{I_{-1}}; \qquad (3.19)$$

b) if $I = \mathcal{J}$,

$$V_{\overline{J^*}}^{\overline{I}} := P_{\overline{I^*}}^{(+)} V_{\overline{I}}^{\overline{I}_{-1}} P_{\overline{I^*}}^{(+)} + P_{\overline{I^*}}^{(-)} V_{\overline{I}}^{\overline{I}_{-1}} P_{\overline{I^*}}^{(-)}$$
(3.20)

$$+P_{\overline{I^*}}^{(+)} \Big(ad Z_{\overline{I^*}} \Big(\frac{\sigma_{i_-^*-1}^{\zeta} \sigma_{i_-^*}^{\zeta}}{\sqrt{t}} + \frac{\sigma_{i_+^*}^{\zeta} \sigma_{i_+^*+1}^{\zeta}}{\sqrt{t}} \Big) \Big) P_{\overline{I^*}}^{(+)}; \qquad (3.21)$$

c) if I^{*} ⊂ J, first we introduce some symbols referring to three sets of intervals entering the formula for V^I_J and used in (3.24), (3.25), and (3.26), respectively,

$$[\mathcal{G}_{\mathcal{J}}^{I}]_{1} := \left\{ \mathcal{K} \in \mathfrak{I} \mid \mathcal{K} \succ I, \, \mathcal{K} \cap I^{*} \neq \emptyset, \\ \mathcal{K} \neq \mathcal{J}, \, \text{and} \, \widetilde{I^{*}} \cup \mathcal{K} = \mathcal{J} \right\}$$
(3.22)

$$[\mathcal{G}_{\mathcal{J}}^{I}]_{2} := \left\{ \mathcal{K}^{*} \in \mathfrak{I}^{*} \mid I > \mathcal{K}, \, \mathcal{K}^{*} \cap I^{*} \neq \emptyset, \, \mathcal{K}^{*} \notin I^{*} \\ \text{and} \, \widetilde{I^{*}} \cup \widetilde{\mathcal{K}^{*}} = \mathcal{J} \right\}$$

$$[\mathcal{G}_{\mathcal{J}}^{I}]_{3} := \left\{ \mathcal{K}^{*} \in \mathfrak{I}^{*} \mid \mathcal{K}^{*} \subset I^{*}, \, \overline{\mathcal{K}^{*}} \notin I^{*} \right\}.$$

Next, we write the definition of the potential $V_{\mathcal{J}}^{\mathcal{I}}$, which, apart from the term in (3.23), results from growth processes where some potentials are hooked in the conjugation implemented by $e^{Z_{\mathcal{I}}*}$ (see (3.24), (3.25), and (3.26)), and from collecting higher order terms both of the Lie-Schwinger series (see (3.27)) and of the hooked Ising terms (see (3.29) and (3.28)) which were not included in b):

$$V_{\mathcal{J}}^{I} := e^{Z_{I^{*}}} V_{\mathcal{J}}^{I_{-1}} e^{-Z_{I^{*}}}$$
(3.23)

$$+\sum_{\mathcal{K}\in[\mathcal{G}_{\mathcal{J}}^{I}]_{1}}\sum_{n=1}^{\infty}\frac{1}{n!}ad^{n}Z_{I^{*}}(V_{\mathcal{K}}^{I_{-1}})$$
(3.24)

$$+\sum_{\mathcal{K}^* \in [\mathcal{G}_{\mathcal{J}}^I]_2} \sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_{\mathcal{I}^*}(V_{\overline{\mathcal{K}^*}}^{I_{-1}})$$
(3.25)

$$+\delta_{\widetilde{I^*}=\mathcal{J}} \sum_{\mathcal{K}^* \in [\mathcal{G}_{\mathcal{J}}^I]_3} \sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_{I^*}(V_{\overline{\mathcal{K}^*}}^{I_{-1}})$$
(3.26)

$$+\delta_{\widetilde{I^*}=\mathcal{J}}\left[\sum_{m=2}^{\infty} t^{\frac{(m-1)}{2}} (V_{I^*}^{\mathcal{I}_{-1}})_m^{\text{diag}_{I^*}}\right]$$
(3.27)

$$+\delta_{\widetilde{I}^{*}=\mathcal{J}}\Big(\sum_{n=2}^{\infty}\frac{1}{n!}ad^{n}Z_{\widetilde{I}^{*}}(\frac{\sigma_{i_{-}^{*}=1}^{z}\sigma_{i_{-}^{*}}^{z}}{\sqrt{t}}+\frac{\sigma_{i_{+}}^{z}\sigma_{i_{+}^{*}+1}^{z}}{\sqrt{t}})\Big)$$
(3.28)

$$+\delta_{\widetilde{I}^{*}=\mathcal{J}}[P_{\overline{I^{*}}}^{(-)}\left(ad Z_{I^{*}}\left(\frac{\sigma_{i_{-}}^{z}\sigma_{i_{-}}^{z}}{\sqrt{t}}+\frac{\sigma_{i_{+}}^{z}\sigma_{i_{+}}^{z}+1}{\sqrt{t}}\right)\right)P_{\overline{I^{*}}}^{(+)}+h.c.].$$
(3.29)

Remark 3.4. The fact that the operators defined in the algorithm, namely $V_{\mathcal{J}}^{I}$, $V_{\overline{\mathcal{J}}^{*}}^{I}$, and $Z_{I^{*}}$, are all well-defined bounded operators requires the main technical results contained in this paper. More precisely, we iteratively use, at each step, Theorem 4.9, Lemma 4.10, and, in the antiferromagnetic case, Lemma 4.3.

The next theorem states that the Hamiltonian in (3.1) is obtained by successive conjugations, of K_{Λ} , generated by the operators $Z_{\mathcal{J}^*}$, with $\mathcal{J} \leq \mathcal{I}$, up to $\mathcal{J} = \mathcal{I}$. For this purpose, it is enough that, for each step of the block-diagonalization, the algorithm is consistent with the conjugation of the Hamiltonian $K_{\Lambda}^{\mathcal{I}_{-1}}(t)$ implemented by the operator $Z_{\mathcal{I}^*}$. **Theorem 3.5.** Given the algorithm in Definition 3.3, the Hamiltonian $K_{\Lambda}^{I}(t)$ defined in (3.1) satisfies

$$K_{\Lambda}^{I}(t) = e^{Z_{I^{*}}} K_{\Lambda}^{I_{-1}}(t) e^{-Z_{I^{*}}}.$$
(3.30)

Proof

We consider the conjugation of each term in the expression below

$$e^{Z_{I^*}} K_{\Lambda}^{I_{-1}}(t) e^{-Z_{I^*}}$$
(3.31)

$$= e^{Z_{I^*}} \left[H^0_{\Lambda} + \sqrt{t} \sum_{\mathcal{J} \prec I} V^{I_{-1}}_{\overline{\mathcal{J}^*}} + \sqrt{t} \sum_{\mathcal{J} \ge I} V^{I_{-1}}_{\mathcal{J}} \right] e^{-Z_{I^*}}$$
(3.32)

and we re-assemble the obtained operators according to the rules of Definition 3.3, so as to get the Hamiltonian $K_{\Lambda}^{I}(t)$. The final result follows from combining the observations below.

i) For $\mathcal{J} \cap \mathcal{I}^* = \emptyset$ and for $\mathcal{J}^* \cap \mathcal{I}^* = \emptyset$,

$$e^{Z_{I^*}} V_{\mathcal{J}}^{I_{-1}} e^{-Z_{I^*}} = V_{\mathcal{J}}^{I_{-1}} =: V_{\mathcal{J}}^{I}$$
(3.33)

$$e^{Z_{I^*}} V_{\overline{\mathcal{J}^*}}^{\mathcal{I}_{-1}} e^{-Z_{I^*}} = V_{\overline{\mathcal{J}^*}}^{\mathcal{I}_{-1}} =: V_{\overline{\mathcal{J}^*}}^{\mathcal{I}}$$
 (3.34)

hold respectively; the identities above follow from a-1) and a-2) in Definition 3.3.

ii) From the definition of the unperturbed Hamiltonian G_{I^*} (see (3.5)), we observe that

$$e^{Z_{I^*}} (H^0_{I^*} + \sqrt{t} \sum_{\mathcal{J} \subset I^*} V^{I_{-1}}_{\mathcal{J}^*} + \sqrt{t} V^{I_{-1}}_{I}) e^{-Z_{I^*}}$$
(3.35)

$$= e^{Z_{I^{*}}} \Big\{ H_{I^{*}}^{0} + \sqrt{t} \sum_{\overline{\mathcal{J}^{*} \subset I^{*}}} V_{\overline{\mathcal{J}^{*}}}^{I_{-1}} + \sqrt{t} V_{I}^{I_{-1}} + \sum_{\mathcal{J^{*} \subset I^{*}}; \overline{\mathcal{J}^{*} \not \subset I^{*}}} V_{\overline{\mathcal{J}^{*}}}^{I_{-1}} \Big\} e^{-Z_{I^{*}}} \\ + \sum_{\mathcal{J^{*} \subset I^{*}}; \overline{\mathcal{J}^{*} \not \subset I^{*}}} V_{\overline{\mathcal{J}^{*}}}^{I_{-1}} \Big\} e^{-Z_{I^{*}}} \\ = e^{Z_{I^{*}}} \Big\{ G_{I^{*}} + \sqrt{t} V_{I}^{I_{-1}} + \sqrt{t} \sum_{\mathcal{J^{*} \subset I^{*}}; \overline{\mathcal{J}^{*} \not \subset I^{*}}} V_{\overline{\mathcal{J}^{*}}}^{I_{-1}} \Big\} e^{-Z_{I^{*}}};$$
(3.36)

next we use (3.4) and get (recall that $G_{I^*} = H^0_{I^*} + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} V^{I_{-1}}_{\overline{\mathcal{J}^*}}$)

$$(3.35) = H_{I^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^* \subset I^*}} V_{\overline{\mathcal{J}^*}}^{I_{-1}} + \sqrt{t} \sum_{m=1}^{\infty} t^{(m-1)/2} (V_{I^*}^{I_{-1}})_m^{\text{diag}_{I^*}}$$

$$+ e^{Z_{I^*}} \sqrt{t} \sum_{\mathcal{J}^* \subset I^*; \overline{\mathcal{J}^*} \notin I^*} V_{\overline{\mathcal{J}^*}}^{I_{-1}} e^{-Z_{I^*}}$$

$$= H_{I^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} V_{\overline{\mathcal{J}^*}}^{\overline{I}}$$

$$(3.37)$$

$$= H_{I^*}^0 + \sqrt{t} \sum_{\mathcal{J} \subset I^*} V_{\overline{\mathcal{J}^*}}^I$$
(3.38)

$$+\sqrt{t}((3.20)) + \sqrt{t}((3.27)) + \sqrt{t}((3.26)), \qquad (3.39)$$

where for the second identity we have used that

$$\sqrt{t} \sum_{\mathcal{J} \subset \mathcal{I}^*} V_{\overline{\mathcal{J}^*}}^{\mathcal{I}} = \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset \mathcal{I}^*} V_{\overline{\mathcal{J}^*}}^{\mathcal{I}_{-1}} + \sqrt{t} \sum_{\mathcal{J}^* \subset \mathcal{I}^*; \overline{\mathcal{J}^*} \notin \mathcal{I}^*} V_{\overline{\mathcal{J}^*}}^{\mathcal{I}_{-1}}$$
(3.40)

and

=

$$\sqrt{t} \sum_{m=1}^{\infty} t^{(m-1)/2} (V_{I^*}^{I_{-1}})_m^{\text{diag}_{I^*}} + \left\{ e^{Z_{I^*}} \sqrt{t} \sum_{\mathcal{J}^* \subset I^*; \overline{\mathcal{J}^*} \not \subset I^*} V_{\overline{\mathcal{J}^*}}^{I_{-1}} e^{-Z_{I^*}} - \sqrt{t} \sum_{\mathcal{J}^* \subset I^*; \overline{\mathcal{J}^*} \not \subset I^*} V_{\overline{\mathcal{J}^*}}^{I_{-1}} \right\}$$

= $\sqrt{t}(3.20) + \sqrt{t}(3.27) + \sqrt{t}(3.26)$

where (3.27) and (3.26) are referred to case c) of Definition 3.3 for a potential $V_{\mathcal{J}'}^{\mathcal{I}}$ with $\mathcal{J}' \equiv \widetilde{I^*}$.

iii) With regard to the terms $V_{\mathcal{J}}^{\mathcal{I}_{-1}}$, with $\mathcal{I}^* \subset \mathcal{J}$, we have

$$e^{Z_{I^*}} V_{\mathcal{J}}^{\mathcal{I}_{-1}} e^{-Z_{I^*}} = (3.23).$$
 (3.41)

iv) With regard to the terms $V_{\mathcal{J}}^{\mathcal{I}_{-1}}$, with $\mathcal{I}^* \cap \mathcal{J} \neq \emptyset$ and $\mathcal{I}^* \notin \mathcal{J}, \mathcal{J} \notin \mathcal{I}^*$, it follows that

$$e^{Z_{I^*}} V_{\mathcal{J}}^{I_{-1}} e^{-Z_{I^*}} = V_{\mathcal{J}}^{I_{-1}} + \sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_{I^*}(V_{\mathcal{J}}^{I_{-1}}), \qquad (3.42)$$

and we observe that the first term on the right hand side is $V_{\mathcal{J}}^{I}$ (see case a-1) Definition 3.3), while the second term is a contribution to $V_{\mathcal{J}'}^{I}$, according to the rule in (3.24), for $\mathcal{J}' \equiv \mathcal{J} \cup \widetilde{I^*}$.

- v) With regard to the terms $V_{\overline{\mathcal{J}}^*}^{I_{-1}}$, we notice that they are present in (3.32) only for $I_{-1} \geq \mathcal{J}$. Hence, we have to consider the occurrences $I > \mathcal{J}$ which we discuss below:
 - the case $\mathcal{J}^* \cap \mathcal{I}^* = \emptyset$ has been already discussed in i);
 - the case $\mathcal{J}^* \subset \mathcal{I}^*$ has been already discussed in ii);
 - if $\mathcal{J}^* \cap I^* \neq \emptyset$ and $\mathcal{J}^* \not\subset I^*$ we write

$$e^{Z_{I^*}} V_{\overline{\mathcal{I}^*}}^{\underline{I}_{-1}} e^{-Z_{I^*}} = V_{\overline{\mathcal{I}^*}}^{\underline{I}_{-1}} + \sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_{\overline{I^*}}(V_{\overline{\mathcal{I}^*}}^{\underline{I}_{-1}})$$
(3.43)

where the first term corresponds to $V_{\overline{\mathcal{J}^*}}^I$ by a-2) of Definition 3.3, and the other terms, i.e.,

$$\sum_{n=1}^{\infty}\frac{1}{n!}\,ad^n Z_{\mathcal{I}^*}(V_{\overline{\mathcal{J}^*}}^{\mathcal{I}_{-1}})\,,$$

contribute to $V_{\mathcal{J}'}^{\mathcal{I}}$, with $\widetilde{\mathcal{I}^*} \cup \widetilde{\mathcal{J}^*} \equiv \mathcal{J}'$, according to (3.25).

vi) With regard to the terms in the unperturbed Hamiltonian H^0_{Λ} which are supported in a microscopic interval (i, i + 1) overlapping with \mathcal{I}^* , but not contained in it, we have

$$e^{Z_{I^*}} \left(\frac{\sigma_{i^*_{+}}^{z} \sigma_{i^*_{+}+1}^{z}}{\sqrt{t}} + \frac{\sigma_{i^*_{-}-1}^{z} \sigma_{i^*_{-}}^{z}}{\sqrt{t}} \right) e^{-Z_{I^*}}$$

$$= \frac{\sigma_{i^*_{+}}^{z} \sigma_{i^*_{+}+1}^{z}}{\sqrt{t}} + \sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_{I^* \equiv \mathcal{J}^*} \left(\frac{\sigma_{i^*_{+}}^{z} \sigma_{i^*_{+}+1}^{z}}{\sqrt{t}} \right)$$

$$= \frac{\sigma_{i^*_{+}}^{z} \sigma_{i^*_{+}+1}^{z}}{\sqrt{t}} + (3.21) + (3.28) + (3.29),$$

where the first term in the last line contributes (once multiplied by $-\sqrt{t} \cdot J$) to H^0_{Λ} .

vii) All the terms in H^0_{Λ} which are supported in intervals with empty intersection with I^* are left invariant by the conjugation.

In the final part of this section, we show that the potentials produced by the algorithm of Definition 3.3 are covariant under translations in the bulk.

Definition 3.6. Let $\mathbb{Z} \ni r \mapsto U_{[r]}$, where $U_{[r]}$ is a unitary operator on \mathcal{H}_{Λ} and [r] is the equivalence class of $r \mod N$ (i.e., $r \mapsto [r] \in \mathbb{Z}/N\mathbb{Z}$), be the action of the additive group of integers factorizing through the natural unitary action of the finite, cyclic translation group $\mathbb{Z}/N\mathbb{Z}$ on the Hilbert space of the chain, Λ , with N sites. We define

$$\tau_k(\cdot) := U_{[k \sqrt{t^{-1}}]}(\cdot) U^*_{[k \sqrt{t^{-1}}]},$$

with $k \in \mathbb{Z}$ and likewise $k \sqrt{t^{-1}}$ (recall that $\sqrt{t^{-1}} \in \mathbb{N}$). We also use the same symbol τ_k for the translation by $[k \sqrt{t^{-1}}] \in \mathbb{Z}/N\mathbb{Z}$ of the sites of the chain.

In the following, given an interval \mathcal{J} , we denote with $j_{\ell}(\mathcal{J})$ the leftmost site of the interval, analogously $j_r(\mathcal{J})$ stands for the rightmost site of the interval. This notation will be needed again later in Section 4.1.2.

Proposition 3.7. For any intervals I, \mathcal{J} , and for any $k \in \mathbb{Z} \cap [0, N-1]$ such that

$$\mathfrak{j}_{\ell}(\mathcal{J}) > 1$$
, and $\mathfrak{j}_{r}(\mathcal{J}) + k\sqrt{t^{-1}} < N$,

we have

$$\begin{aligned} i) \ \ \tau_k(V_{\mathcal{J}^*}^{I}) &= V_{\tau_k(\mathcal{J}^*)}^{(I)_{+k}} \qquad if \ I \geq \mathcal{J} \ ; \\ ii) \ \ \tau_k(V_{\mathcal{J}}^{I}) &= V_{\tau_k(\mathcal{J})}^{(I)_{+k}} \qquad if \ I \prec \mathcal{J} \ . \end{aligned}$$

Proof

We make use of an induction on the index I labelling the steps of the iteration. For the basis of the induction, namely the case $I = (I_0)_{+1}$, since for every \mathcal{J} satisfying the hypotheses of the statement it holds $(I_0)_{+1} \prec \mathcal{J}$, only *ii*) above is to be proven. In order to prove it, we invoke a-1) in Definition 3.3 and we deduce that $V_{\mathcal{J}}^{(I_0)_{+1}} = V_{\mathcal{J}}^{I_0}$, $V_{\tau_k(\mathcal{J})}^{(I_0)_{+k+1}} = V_{\tau_k(\mathcal{J})}^{I_0}$ for $\ell(\mathcal{J}) = 1$, and $V_{\mathcal{J}}^{(I_0)_{+1}} = V_{\tau_k(\mathcal{J})}^{(I_0)_{+1}} = 0$ for $\ell(\mathcal{J}) > 1$; thus the statement follows by the translation covariance of the potentials in the Hamiltonian K_{Λ} which yields $\tau_k(V_{\mathcal{J}}^{I_0}) = V_{\tau_k(\mathcal{J})}^{I_0}$. Now suppose that the statement holds in step I_{-1} . We observe that the various cases a), b),

Now suppose that the statement holds in step I_{-1} . We observe that the various cases a), b), and c) of the algorithm depend on the relative position between I and \mathcal{J} .

If I and \mathcal{J} are in a relative position such that case a) of the algorithm has to be used to express the potential $V_{\overline{\mathcal{J}^*}}^I$ in terms of $V_{\mathcal{J}^*}^{I_{-1}}$ (resp. $V_{\mathcal{J}}^I$ in terms of $V_{\mathcal{J}}^{I_{-1}}$), we only have to show that case a) also applies to $V_{\tau_k(\overline{\mathcal{J}^*})}^{(I)_{+k}}$ (resp. $V_{\tau_k(\mathcal{J})}^{(I)_{+k}}$), since then the statement will follow by the inductive hypothesis. Indeed, if $I > \mathcal{J}$ (i.e., case a-2) applies to $V_{\overline{\mathcal{J}^*}}^I$), the relation $(I)_{+k} > \tau_k(\mathcal{J})$ follows, thus case a-2) applies to $V_{\tau_k(\overline{\mathcal{J}^*})}^{(I)_{+k}}$. If $I < \mathcal{J}$ and $I^* \notin \mathcal{J}$ (i.e., case a-1) applies to $V_{\mathcal{J}}^I$), we want to show that $(I)_{+k} < \tau_k(\mathcal{J})$ and $(I)_{+k}^* \notin \tau_k(\mathcal{J})$ (recall the constraint on k in the statement). We check the latter for k = 1 so that the statement for general k follows by iteration. The relation $I_{+1} < \tau_1(\mathcal{J})$ is obvious, while for $I_{+1}^* \notin \tau_1(\mathcal{J})$ the only slightly nontrivial case is when I contains the right endpoint of the lattice. In this case $I_{+1}^* \notin \tau_1(\mathcal{J})$ holds since I_{+1}^* contains the left endpoint of the boundary of the lattice while $\tau_1(\mathcal{J})$ does not. If I and \mathcal{J} are in a relative position such that case b) of the algorithm has to be applied to $V_{\overline{\mathcal{J}^*}}^I$ then case b) applies also to $V_{\tau_k(\overline{\mathcal{J}^*})}^{(I)_{+k}}$ since in this case $\tau_k(I) = I_{+k}$. Consequently, the statement follows by inspecting formulas (3.20)-(3.21) together with (3.8)-(3.7) and by the inductive hypothesis; in this respect we remark that the endpoints of the lattice Λ cannot belong to the sets \mathcal{J} and $\tau_k(\mathcal{J})$ (as assumed in the statement) in order to exploit the translation covariance of formula (3.20)-(3.21).

The proof is analogous if I and \mathcal{J} are in a relative position such that case c) of the algorithm applies to $V_{\mathcal{T}}^{I}$. \Box

Remark 3.8. A result analogous to Proposition 3.7 holds for negative k.

4 Operator norms and control of the flow

The control of the algorithm designed in the previous section requires an elaborate proof by induction which concerns the operator norm of the effective potentials, the convergence of the series yielding the operators Z_{I^*} , and a bound from below on the spectral gaps in (3.14) and (3.15). For this purpose, we split the proof into different parts which will be merged as ingredients in the proof of Theorem 4.9. In Section 4.1 we provide the argument to control the gaps in (3.14) and (3.15), for which we have to distinguish the cases J > 0 and J < 0.

4.1 Gap estimates

In estimating the gap above the ground-state energy for the local Hamiltonians $G_{I_{+1}^*}$, for both the ferromagnetic and the antiferromagnetic case our standing assumption is

$$\|V_{\overline{\mathcal{J}^*}}^I\| \le C_{J,h} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{(\ell(\mathcal{J}))^2} \tag{4.1}$$

for all $\mathcal{J} \leq I$. Here $C_{J,h}$ is a quantity dependent on J and h; see Lemma 4.10. (The proof of this operator norm estimate will be the content of Theorem 4.9 and Lemma 4.10 in Section 4.2). Our results and the related arguments heavily depend on the features of the model; as said above we present them treating the ferromagnetic and antiferromagnetic cases separately.

4.1.1 Spectral Gap of $G_{I_{+1}^*}$ in the ferromagnetic case

In this case, the strategy for estimating the spectral gap of $G_{I_{+1}^*}$ is similar to the one in [FP] due to the nondegeneracy of the ground-state energy of the local unperturbed Hamiltonians. Nonetheless we spell it out in detail for the convenience of the reader. Our proof relies on the following considerations.

(i) For $\overline{\mathcal{J}^*} \subset \mathcal{I}^*_{+1}$, $V^{\overline{I}}_{\overline{\mathcal{J}^*}}$ is a block-diagonalized potential with respect to the projections $P^{(-)}_{\mathcal{I}^*_{+1}}, P^{(+)}_{\mathcal{I}^*_{+1}}$, i.e.,

$$V_{\overline{\mathcal{J}^*}}^{\underline{I}} := P_{I_{+1}^*}^{(+)} V_{\overline{\mathcal{J}^*}}^{\underline{I}} P_{I_{+1}^*}^{(+)} + P_{I_{+1}^*}^{(-)} V_{\overline{\mathcal{J}^*}}^{\underline{I}} P_{I_{+1}^*}^{(-)}$$

Indeed, by inspecting the definition of $V_{\overline{\mathcal{J}^*}}^I$, see formulae (3.20)-(3.21) where \mathcal{I} coincides with \mathcal{J} , this easily follows from $P_{\overline{\mathcal{J}^*}}^{(+)}P_{\mathcal{I}^*_{\pm 1}}^{(-)} = 0$ and

$$P_{\mathcal{I}_{+1}^{*}}^{(+)} P_{\mathcal{J}^{*}}^{(-)} V_{\mathcal{J}}^{\mathcal{J}_{-1}} P_{\mathcal{J}^{*}}^{(-)} P_{\mathcal{I}_{+1}^{*}}^{(-)} = P_{\mathcal{I}_{+1}^{*}}^{(+)} \langle V_{\mathcal{J}}^{\mathcal{J}_{-1}} \rangle_{\Psi_{\mathcal{J}^{*}}} P_{\mathcal{J}^{*}}^{(-)} P_{\mathcal{I}_{+1}^{*}}^{(-)} = \langle V_{\mathcal{J}}^{\mathcal{J}_{-1}} \rangle_{\Psi_{\mathcal{J}^{*}}} P_{\mathcal{I}_{+1}^{*}}^{(+)} P_{\mathcal{I}_{+1}^{*}}^{(-)} = 0,$$

where we recall that for $\langle V \rangle_{\Psi}$ stands for $\langle \Psi, V \Psi \rangle$, for any vector Ψ and operator V. In particular this implies that $G_{I_{+1}^*}$ is block-diagonal with respect to the projections $P_{I_{+1}^*}^{(-)}, P_{I_{+1}^*}^{(+)}$.

(ii) Denoting $(\mathcal{I}_{+1}^*)_K := \{\mathcal{J} \in \mathfrak{I} : \ell(\mathcal{J}) = K, \ \overline{\mathcal{J}^*} \subset \mathcal{I}_{+1}^*\} \text{ and } \mathscr{I}_K := \{i \in \bigcup_{\mathcal{J} \in (\mathcal{I}_{+1}^*)_K} \overline{\mathcal{J}^*}\}, \text{ we have$

have

$$\sum_{\mathcal{J}\in(\mathcal{I}^*_{+1})_{K}} (H^{C}_{\overline{\mathcal{J}^*}} - \langle H^{C}_{\overline{\mathcal{J}^*}} \rangle_{\Psi_{\overline{\mathcal{J}^*}}}) = \sum_{\mathcal{J}\in(\mathcal{I}^*_{+1})_{K}} \sum_{i,i+1\in\overline{\mathcal{J}^*}} (H^{C}_{i,i+1} - \langle H^{C}_{i,i+1} \rangle_{\Psi_{i,i+1}}) \quad (4.2)$$

$$\leq (K+1) \sum_{i,i+1 \in \mathscr{I}_{K}} (H_{i,i+1}^{C} - \langle H_{i,i+1}^{C} \rangle_{\Psi_{i,i+1}}) \qquad (4.3)$$
$$= (K+1) (H_{\mathscr{I}_{K}}^{C} - \langle H_{\mathscr{I}_{K}}^{C} \rangle_{\Psi_{\mathscr{I}_{K}}})$$

where $H_{i,i+1}^C$ and $\Psi_{i,i+1}$ are H_I^C and Ψ_I for $I \equiv \{i, i+1\}$, consequently $H_{\mathscr{I}_K}^C = \sum_{i,i+1 \in \mathscr{I}_K} H_{i,i+1}^C$ by definition, with $\Psi_{\mathscr{I}_K}$ its ground-state.

Next, assuming the bound in (4.1) and making use of the inequality

$$P_{\overline{\mathcal{J}^*}}^{(+)} \le \frac{1}{2J+h} \left(H_{\overline{\mathcal{J}^*}}^C - \langle H_{\overline{\mathcal{J}^*}}^C \rangle_{\Psi_{\overline{\mathcal{J}^*}}} \right), \tag{4.4}$$

where we have used that 2J + h is the spectral gap above the ground-state energy of $H_{\overline{\mathcal{J}^*}}^C$, we can estimate

$$\pm P_{\overline{\mathcal{J}^*}}^{(+)} V_{\overline{\mathcal{J}^*}}^{I} P_{\overline{\mathcal{J}^*}}^{(+)} \le \frac{C_{J,h}}{2J+h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(H_{\overline{\mathcal{J}^*}}^C - \left\langle H_{\overline{\mathcal{J}^*}}^C \right\rangle_{\Psi_{\overline{\mathcal{J}^*}}} \right).$$

$$(4.5)$$

The inequality in (4.5) combined with point (ii) above yields

$$\pm \sum_{\mathcal{J} \in (\mathcal{I}_{+1}^*)_K} P_{\mathcal{J}^*}^{(+)} V_{\mathcal{J}^*}^{\mathcal{I}} P_{\mathcal{J}^*}^{(+)} \le \frac{C_{J,h}}{2J+h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(K+1\right) \left(H_{\mathcal{I}_{+1}^*}^C - \langle H_{\mathcal{I}_{+1}^*}^C \rangle_{\Psi_{\mathcal{I}_{+1}^*}}\right), \tag{4.6}$$

since $\mathscr{I}_K \subset \mathcal{I}_{+1}^*$ for all *K* and $H_{\overline{\mathcal{J}^*}}^C - \langle H_{\overline{\mathcal{J}^*}}^C \rangle_{\Psi_{\overline{\mathcal{J}^*}}} \leq H_{\mathcal{I}_{+1}^*}^C - \langle H_{\mathcal{I}_{+1}^*}^C \rangle_{\Psi_{\mathcal{I}_{+1}^*}}$. With these ingredients, the proof of the next lemma can then be easily derived.

Lemma 4.1. Assuming that the bound in (4.1) holds in step I of the block-diagonalization, and choosing t > 0 small enough such that

$$\left\{1 - \frac{4C_{J,h}}{2J+h} \cdot \sqrt{t} - \frac{2C_{J,h}}{2J+h} \cdot \sqrt{t} \sum_{l=3}^{\infty} l \cdot t^{\frac{l-2}{16}}\right\} > 0, \qquad (4.7)$$

the inequality

$$P_{\mathcal{I}_{+1}^{*}}^{(+)}\left(G_{\mathcal{I}_{+1}^{*}}-E_{\mathcal{I}_{+1}^{*}}\right)P_{\mathcal{I}_{+1}^{*}}^{(+)} \geq (2J+2h)\left\{1-\frac{4C_{J,h}}{2J+h}\cdot\sqrt{t}-\frac{2C_{J,h}}{2J+h}\cdot\sqrt{t}\sum_{l=3}^{\infty}l\cdot t^{\frac{l-2}{16}}\right\}P_{\mathcal{I}_{+1}^{*}}^{(+)} \quad (4.8)$$

holds, where

$$E_{I_{+1}^*} := \langle G_{I_{+1}^*} \rangle_{\Psi_{I_{+1}^*}}.$$
(4.9)

Proof

By definition of $G_{\mathcal{I}_{+1}^*}$ and using the identity

$$V_{\overline{\mathcal{J}^*}}^{\underline{I}} = P_{\overline{\mathcal{J}^*}}^{(+)} V_{\overline{\mathcal{J}^*}}^{\underline{I}} P_{\overline{\mathcal{J}^*}}^{(+)} + P_{\overline{\mathcal{J}^*}}^{(-)} V_{\overline{\mathcal{J}^*}}^{\underline{I}} P_{\overline{\mathcal{J}^*}}^{(-)}$$

for all $\overline{\mathcal{J}^*} \subset \mathcal{I}^*_{+1}$, we can write

$$P_{I_{+1}^{*}}^{(+)} G_{I_{+1}^{*}} P_{I_{+1}^{*}}^{(+)} = P_{I_{+1}^{*}}^{(+)} (H_{I_{+1}^{*}}^{0} + \sqrt{t} \sum_{\overline{\mathcal{J}^{*} \subset I_{+1}^{*}}} V_{\overline{\mathcal{J}^{*}}}^{\underline{\mathcal{I}}}) P_{I_{+1}^{*}}^{(+)}$$

$$= P_{I_{+1}^{*}}^{(+)} (H_{I_{+1}^{*}}^{0} + \sqrt{t} \sum_{\overline{\mathcal{J}^{*} \subset I_{+1}^{*}}} P_{\overline{\mathcal{J}^{*}}}^{(+)} V_{\overline{\mathcal{J}^{*}}}^{\underline{\mathcal{I}}} P_{\overline{\mathcal{J}^{*}}}^{(+)}) P_{I_{+1}^{*}}^{(+)}$$

$$(4.10)$$

$$+P_{I_{+1}^{*}}^{(+)} \sqrt{t} \sum_{\overline{\mathcal{J}^{*}} \subset I_{+1}^{*}} \langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} (\mathbb{1} - P_{\overline{\mathcal{J}^{*}}}^{(+)}) P_{I_{+1}^{*}}^{(+)}.$$
(4.11)

Now, using (4.6) above together with

$$H_{\mathcal{I}_{+1}^{*}}^{C} - \langle H_{\mathcal{I}_{+1}^{*}}^{C} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}} \leq H_{\mathcal{I}_{+1}^{*}}^{0} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}}, \qquad (4.12)$$

(which trivially follows from (1.10) and (1.12)) we get

$$\pm \sum_{\overline{\mathcal{J}^{*} \subset I_{+1}^{*}}} P_{\overline{\mathcal{J}^{*}}}^{(+)} V_{\overline{\mathcal{J}^{*}}}^{I} P_{\overline{\mathcal{J}^{*}}}^{(+)}$$

$$\leq \frac{C_{J,h}}{2J+h} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J})+1) (H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}).$$

$$(4.13)$$

Thus, with the help of (4.4), (4.12), and using the identity

$$H^{0}_{\mathcal{I}^{*}_{+1}} - E_{\mathcal{I}^{*}_{+1}} = H^{0}_{\mathcal{I}^{*}_{+1}} - \langle H^{0}_{\mathcal{I}^{*}_{+1}} \rangle_{\Psi_{\mathcal{I}^{*}_{+1}}} - \sqrt{t} \sum_{\overline{\mathcal{J}^{*}} \subset \mathcal{I}^{*}_{+1}} \langle V^{\underline{\mathcal{I}}}_{\overline{\mathcal{J}^{*}}} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}},$$

we can conclude that

$$P_{\mathcal{I}_{+1}^{*}}^{(+)} (G_{\mathcal{I}_{+1}^{*}} - E_{\mathcal{I}_{+1}^{*}}) P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

$$\geq P_{\mathcal{I}_{+1}^{*}}^{(+)} \left(1 - \frac{2C_{J,h}}{2J + h} \cdot \sqrt{t} \sum_{K'=1}^{\infty} t^{\frac{K'-1}{16}} (K'+1) (H_{\mathcal{I}_{+1}^{*}}^{0} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}}) \right) P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

$$\geq (2J+2h) \left(1 - \frac{2C_{J,h}}{2J + h} \cdot \sqrt{t} \sum_{K'=1}^{\infty} t^{\frac{K'-1}{16}} (K'+1) \right) P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

where for the last inequality we have used

$$P_{\mathcal{I}_{+1}^{*}}^{(+)}(H_{\mathcal{I}_{+1}^{*}}^{0} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}}) P_{\mathcal{I}_{+1}^{*}}^{(+)} \ge (2J + 2h) P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

which follows from Proposition 1.1. \Box

4.1.2 Low-lying spectrum of $G_{I_{\perp 1}^*}$ in the antiferromagnetic case

The discussion is carried out in three steps.

- I) In Lemma 4.3 we show that, due to the definition of $V_{\overline{\mathcal{J}^*}}^{\mathcal{J}}$ in point b) of the algorithm (see (3.20)-(3.21)), the operator $G_{I_{+1}^*}$ is not only block-diagonal with respect to $P_{I_{+1}^*}^{(-)}, P_{\overline{\mathcal{I}^*_{+1}}}^{(+)}$ but also w.r.t. $P_{I_{+1}^*}^{(-),A}, P_{I_{+1}^*}^{(-),B}$;
- II) The result is then used in Lemma 4.4 where we estimate the difference $|E_{I_{+1}^*}^B E_{I_{+1}^*}^A|$ of the two lowest eigenvalues of $G_{I_{+1}^*}$;
- III) In Lemma 4.6 we finally estimate the distance between the spectrum of $P_{I_{+1}^*}^{(+)}G_{I_{+1}^*}P_{I_{+1}^*}^{(+)}$ and the two lowest eigenvalues of $G_{I_{+1}^*}$, with the help of Lemma 4.5 which provides an intermediate result.

In the following, due to the structure of the two lowest-energy eigenvectors of the local unperturbed Hamiltonians, it will be useful to split the set consisting of intervals \mathcal{J}^* such that $\overline{\mathcal{J}^*} \subset I_{+1}^*$, i.e.,

$$(\mathcal{I}_{+1}^*)_{\text{int}} := \left\{ \mathcal{J}^* \in \mathfrak{I}^* : \overline{\mathcal{J}^*} \subset \mathcal{I}_{+1}^* \right\}, \qquad (4.14)$$

into two sets which are defined below, where $j_{\ell}(\mathcal{J})$ stands for the microscopic coordinate of the leftmost site of the interval \mathcal{J} :

$$(\mathcal{I}_{+1}^{*})_{\text{ev}} := \left\{ \mathcal{J}^{*} \in \mathfrak{I}^{*} : \overline{\mathcal{J}^{*}} \subset \mathcal{I}_{+1}^{*}, |j_{\ell}(\mathcal{J}^{*}) - j_{\ell}(\mathcal{I}_{+1}^{*})| \text{ is even } \right\},$$
(4.15)

$$(I_{+1}^*)_{\text{odd}} := \left\{ \mathcal{J}^* \in \mathfrak{I}^* : \overline{\mathcal{J}^*} \subset I_{+1}^*, |j_{\ell}(\mathcal{J}^*) - j_{\ell}(I_{+1}^*)| \text{ is odd } \right\}.$$
(4.16)

Remark 4.2. We observe that, by definition of the vectors Ψ_I^A and Ψ_I^B , and of the sets $(\mathcal{I}_{+1}^*)_{\text{ev}}$, $(\mathcal{I}_{+1}^*)_{\text{odd}}$, the following identities hold true: if $\mathcal{J}^* \in (\mathcal{I}_{+1}^*)_{\text{ev}}$

$$\langle V_{\mathcal{J}^*} \rangle_{\Psi^{A/B}_{\mathcal{I}^*_{\mathcal{I}^*}}} = \langle V_{\mathcal{J}^*} \rangle_{\Psi^{A/B}_{\mathcal{J}^*}} \tag{4.17}$$

if $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{odd}}$

$$\langle V_{\mathcal{J}^*} \rangle_{\Psi^{A/B}_{\mathcal{I}^*_{+1}}} = \langle V_{\mathcal{J}^*} \rangle_{\Psi^{B/A}_{\mathcal{J}^*}}.$$
(4.18)

In the lemma below we prove a property that can be seen as a weak form of the LTQO condition for the ground-state subspace of $H_{I_{+1}^*}^C$. It consists in the absence of off-diagonal terms of the effective potentials $V_{\overline{\mathcal{I}}^*}^I$ w.r.t. $P_{I_{+1}^*}^{(-),A}$ and $P_{I_{+1}^*}^{(-),B}$. This property results from using enlarged intervals in the block-diagonalization procedure.

Lemma 4.3. For any $I, \mathcal{J} \in \mathfrak{I}$ with $\overline{\mathcal{J}^*} \subset \mathcal{I}^*_{+1}$, we have the following:

(a) $V_{\overline{\mathcal{J}^*}}^{\underline{I}}$ is block-diagonal with respect to $P_{\overline{\mathcal{J}^*}}^{(-),A}$, $P_{\overline{\mathcal{J}^*}}^{(-),B}$, $P_{\overline{\mathcal{J}^*}}^{(+)}$ (see their definitions below (2.41)); (b) $V_{\overline{\mathcal{J}^*}}^{\underline{I}}$ is block-diagonal with respect to $P_{I_{+1}}^{(-),A}$, $P_{I_{+1}}^{(-),B}$, $P_{I_{+1}^*}^{(+)}$.

Consequently, $G_{\mathcal{I}_{+1}^*}$ is block-diagonal with respect to $P_{\mathcal{I}_{+1}^*}^{(-),A}$, $P_{\mathcal{I}_{+1}^*}^{(-),B}$, $P_{\mathcal{I}_{+1}^*}^{(+)}$.

Proof

The proofs of points (a) and (b) are identical; thus we show only (b), that is we prove that

$$P_{I_{+1}^{*}}^{(+)} V_{\overline{I}_{+}^{*}}^{\overline{I}} P_{I_{+1}^{*}}^{(-)} = 0, ag{4.19}$$

$$P_{I_{+1}^*}^{(-),A} V_{\overline{\mathcal{I}}^*}^{I} P_{I_{+1}^*}^{(-),B} = 0.$$
(4.20)

We recall that (see point b) in Definition 3.3) for $\mathcal{I} \geq \mathcal{J}$

$$V_{\overline{\mathcal{J}^{*}}}^{I} = V_{\overline{\mathcal{J}^{*}}}^{\mathcal{J}} := P_{\mathcal{J}^{*}}^{(+)} V_{\mathcal{J}}^{\mathcal{J}_{-1}} P_{\mathcal{J}^{*}}^{(+)} + P_{\mathcal{J}^{*}}^{(-)} V_{\mathcal{J}}^{\mathcal{J}_{-1}} P_{\mathcal{J}^{*}}^{(-)}$$
(4.21)

$$+P_{\overline{\mathcal{J}^{*}}}^{(+)} \Big(adZ_{\overline{\mathcal{J}^{*}}} \Big(\frac{\sigma_{i_{-}}^{\zeta_{*}} - 1}{\sqrt{t}} + \frac{\sigma_{i_{+}}^{\zeta_{*}} \sigma_{i_{+}}^{\zeta_{*}} + 1}{\sqrt{t}} \Big) \Big) P_{\overline{\mathcal{J}^{*}}}^{(+)}, \tag{4.22}$$

and we observe the following relations:

- (i) $P_{I_{+1}^*}^{(-)} P_{\mathcal{J}^*}^{(+)} = P_{I_{+1}^*}^{(-)} P_{\overline{\mathcal{J}^*}}^{(+)} = 0$ since $\overline{\mathcal{J}^*} \subset I_{+1}^*$ by assumption;
- (ii) $P_{\mathcal{J}^*}^{(-),A}V_{\mathcal{J}}^{\mathcal{J}_{-1}}P_{\mathcal{J}^*}^{(-),B} = 0$, due to the fact that $V_{\mathcal{J}}^{\mathcal{J}_{-1}}$ is localized in \mathcal{J} , which is strictly contained in \mathcal{J}^* ; hence

$$P_{\mathcal{J}^*}^{(-)}V_{\mathcal{J}}^{\mathcal{J}_{-1}}P_{\mathcal{J}^*}^{(-)} = \langle V_{\mathcal{J}}^{\mathcal{J}_{-1}}\rangle_{\Psi_{\mathcal{J}^*}^A}P_{\mathcal{J}^*}^{(-),A} + \langle V_{\mathcal{J}}^{\mathcal{J}_{-1}}\rangle_{\Psi_{\mathcal{J}^*}^B}P_{\mathcal{J}^*}^{(-),B};$$

(iii) If
$$\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{ev}}$$
, $P_{\mathcal{I}^*_{+1}}^{(-),A} P_{\mathcal{J}^*}^{(-),A} = P_{\mathcal{I}^*_{+1}}^{(-),A} \text{ and } P_{\mathcal{I}^*_{+1}}^{(-),B} P_{\mathcal{J}^*}^{(-),B} = P_{\mathcal{I}^*_{+1}}^{(-),B}$.
If $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{odd}}$, $P_{\mathcal{I}^*_{+1}}^{(-),A} P_{\mathcal{J}^*}^{(-),A} = P_{\mathcal{I}^*_{+1}}^{(-),B} \text{ and } P_{\mathcal{I}^*_{+1}}^{(-),B} P_{\mathcal{J}^*}^{(-),B} = P_{\mathcal{I}^*_{+1}}^{(-),A}$.
(iv) If $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{ev}}$, $P_{\mathcal{I}^*_{+1}}^{(-),A} P_{\mathcal{J}^*}^{(-),A} = P_{\mathcal{I}^*_{+1}}^{(-),A}$, $P_{\mathcal{I}^*_{+1}}^{(-),B} P_{\mathcal{J}^*}^{(-),B} = P_{\mathcal{I}^*_{+1}}^{(-),B}$, and $P_{\mathcal{I}^*_{+1}}^{(-),A} P_{\mathcal{J}^*}^{(-),B} = 0$.

$$\text{If } \mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{odd}}, \quad P_{\mathcal{I}^*_{+1}}^{(-),A} P_{\mathcal{J}^*}^{(-),A} = 0, \quad P_{\mathcal{I}^*_{+1}}^{(-),B} P_{\mathcal{J}^*}^{(-),B} = 0, \text{ and } P_{\mathcal{I}^*_{+1}}^{(-),A} P_{\mathcal{J}^*}^{(-),B} = P_{\mathcal{I}^*_{+1}}^{(-),A}.$$

Thus, using (4.21),

$$P_{I_{+1}^{*}}^{(+)} V_{\overline{\mathcal{J}}^{*}}^{\overline{\mathcal{I}}} P_{I_{+1}^{*}}^{(-)}$$

$$= P_{I_{+1}^{*}}^{(+)} P_{\mathcal{J}^{*}}^{(-)} V_{\mathcal{J}}^{\mathcal{J}_{-1}} P_{\mathcal{J}^{*}}^{(-)} P_{I_{+1}^{*}}^{(-)}$$

$$= P_{I_{+1}^{*}}^{(+)} \left(\langle V_{\mathcal{J}}^{\mathcal{J}_{-1}} \rangle_{\Psi_{\overline{\mathcal{J}}^{*}}} P_{\mathcal{J}^{*}}^{(-),A} + \langle V_{\mathcal{J}}^{\mathcal{J}_{-1}} \rangle_{\Psi_{\overline{\mathcal{J}}^{*}}} P_{\mathcal{J}^{*}}^{(-),B} \right) P_{I_{+1}^{*}}^{(-)} = 0$$

which proves (4.19), where the first equality is due to item (i), the second to item (ii), and the last one to item (iii).

Concerning (4.20),

$$\begin{split} & P_{I_{+1}^*}^{(-),A} V_{\overline{\mathcal{J}^*}}^{\mathcal{I}} P_{I_{+1}^*}^{(-),B} \\ &= P_{I_{+1}^*}^{(-),A} P_{\overline{\mathcal{J}^*}}^{(-)} V_{\overline{\mathcal{J}^*}}^{\mathcal{J}_{-1}} P_{\overline{\mathcal{J}^*}}^{(-),B} P_{I_{+1}^*}^{(-),B} \\ &= P_{I_{+1}^*}^{(-),A} (\langle V_{\mathcal{J}}^{\mathcal{J}_{-1}} \rangle_{\Psi_{\mathcal{J}^*}^A} P_{\overline{\mathcal{J}^*}}^{(-),A} + \langle V_{\mathcal{J}}^{\mathcal{J}_{-1}} \rangle_{\Psi_{\mathcal{J}^*}^B} P_{\overline{\mathcal{J}^*}}^{(-),B}) P_{I_{+1}^*}^{(-),B} = 0 \,, \end{split}$$

where the first equality follows from item (i), the second from item (ii), and the last one from item (iv). \Box

In the next lemma we make use of the argument (see Section 2.4) regarding what we refer to as *degeneracy of the bulk ground-state energy*.

Lemma 4.4. Let

$$E_{\mathcal{I}_{+1}^{*}}^{A} := \langle G_{\mathcal{I}_{+1}^{*}} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A}}, \qquad (4.23)$$

and

$$E^B_{\mathcal{I}^*_{+1}} := \langle G_{\mathcal{I}^*_{+1}} \rangle_{\Psi^B_{\mathcal{I}^*_{+1}}}.$$

If I^* has an odd number of sites, then

$$E^{B}_{\mathcal{I}^{*}_{+1}} - E^{A}_{\mathcal{I}^{*}_{+1}} = 2h + O(\sqrt{t}).$$

If I^* has an even number of sites, then

$$|E^B_{\mathcal{I}^*_{+1}} - E^A_{\mathcal{I}^*_{+1}}| \le O(\sqrt{t}).$$

Proof

If the set \mathcal{I}^* has odd cardinality, then

$$\langle H^0_{\mathcal{I}^*_{+1}} \rangle_{\Psi^B_{\mathcal{I}^*_{+1}}} - \langle H^0_{\mathcal{I}^*_{+1}} \rangle_{\Psi^A_{\mathcal{I}^*_{+1}}} = 2h,$$

whereas if \mathcal{I}^* has even cardinality

$$\langle H^0_{\mathcal{I}^*_{+1}} \rangle_{\Psi^B_{\mathcal{I}^*_{+1}}} - \langle H^0_{\mathcal{I}^*_{+1}} \rangle_{\Psi^A_{\mathcal{I}^*_{+1}}} = 0.$$

Next we consider two subsets of $(\mathcal{I}_{+1}^*)_{ev}$ and $(\mathcal{I}_{+1}^*)_{odd}$, respectively:

$$(\mathcal{I}_{+1}^{*})'_{\text{ev}} := \{\mathcal{J}^{*} \in (\mathcal{I}_{+1}^{*})_{\text{ev}} : \tau_{-1}(\overline{\mathcal{J}^{*}}) \subset \mathcal{I}_{+1}^{*}\}$$
(4.24)

and

$$(\mathcal{I}_{+1}^{*})'_{\text{odd}} := \{\mathcal{J}^{*} \in (\mathcal{I}_{+1}^{*})_{\text{odd}} : \tau_{1}(\overline{\mathcal{J}^{*}}) \subset \mathcal{I}_{+1}^{*}\}.$$
(4.25)

The set $(\mathcal{I}_{+1}^*)'_{ev}$, respectively $(\mathcal{I}_{+1}^*)'_{odd}$, consists of intervals that are still contained in \mathcal{I}_{+1}^* when shifted by τ_k with k = -1, respectively k = 1 (see Definition 3.6). We also observe that

$$\tau_{-1}((\mathcal{I}_{+1}^*)'_{\rm ev}) = (\mathcal{I}_{+1}^*)'_{\rm odd} \,. \tag{4.26}$$

If \mathcal{I}^* has an odd number of sites, by splitting $(\mathcal{I}^*_{+1})_{int}$ (see (4.14)) into $(\mathcal{I}^*_{+1})_{ev} \cup (\mathcal{I}^*_{+1})_{odd}$, and by using the properties in Remark 4.2, we can write

$$E_{I_{+1}^*}^B - E_{I_{+1}^*}^A \tag{4.27}$$

$$= \langle H^{0}_{\mathcal{I}^{*}_{+1}} \rangle_{\Psi^{B}_{\mathcal{I}^{*}_{+1}}} - \langle H^{0}_{\mathcal{I}^{*}_{+1}} \rangle_{\Psi^{A}_{\mathcal{I}^{*}_{+1}}} + \sqrt{t} \cdot \sum_{\overline{\mathcal{J}^{*}} \subset \mathcal{I}^{*}_{+1}} (\langle V^{\underline{I}}_{\overline{\mathcal{J}^{*}}} \rangle_{\Psi^{B}_{\mathcal{I}^{*}_{+1}}} - \langle V^{\underline{I}}_{\overline{\mathcal{J}^{*}}} \rangle_{\Psi^{A}_{\mathcal{I}^{*}_{+1}}})$$
(4.28)

$$= 2h + \sqrt{t} \cdot \left(\sum_{\mathcal{J}^* \in (\overline{I}^*_{+1})_{\text{ev}}} (\langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^B} - \langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^A}) + \sum_{\mathcal{J}^* \in (\overline{I}^*_{+1})_{\text{odd}}} (\langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^A} - \langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^B}) \right).$$

Next, by splitting $(\mathcal{I}_{+1}^*)_{ev/odd}$ into $(\mathcal{I}_{+1}^*)'_{ev/odd} \cup [(\mathcal{I}_{+1}^*)_{ev/odd} \setminus (\mathcal{I}_{+1}^*)'_{ev/odd}]$, we can estimate

$$\begin{aligned} |E_{I_{+1}^{*}}^{B} - E_{I_{+1}^{*}}^{A} - 2h| \\ \leq & \sqrt{t} \cdot \left| \left(\sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{ev}^{\prime}} (\langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}} - \langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}}) + \sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{odd}^{\prime}} (\langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}} - \langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}}) \right) \right| \\ & + \sqrt{t} \cdot \sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{ev} \setminus (I_{+1}^{*})_{ev}^{\prime}} |(\langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}} - \langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}})| + \sqrt{t} \cdot \sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{odd} \setminus (I_{+1}^{*})_{odd}^{\prime}} |(\langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}} - \langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}})| + \sqrt{t} \cdot \sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{odd} \setminus (I_{+1}^{*})_{odd}^{\prime}} |(\langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}} - \langle V_{\mathcal{J}^{*}}^{I} \rangle_{\Psi_{\mathcal{J}^{*}}})| . \end{aligned}$$

Now we claim that the terms in (4.29) cancel out. In order to show this we recall (4.26) and observe that, for each $\mathcal{J}^* \in (\mathcal{I}^*_{+1})'_{ev}$, we have:

• by invoking Proposition 3.7

$$\langle V_{\overline{\mathcal{J}^*}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^*}}^{A/B}} = \langle V_{\tau_{-1}(\overline{\mathcal{J}^*})}^{I_{-1}} \rangle_{\Psi_{\tau_{-1}(\overline{\mathcal{J}^*})}^{A/B}}; \qquad (4.30)$$

• due to the rules in Definition 3.3, a potential that has been block-diagonalized does not change along the flow; since $I > \mathcal{J}$, the potential $V_{\tau_{-1}(\overline{\mathcal{J}^*})}^{\tau_{-1}(I)}$ is already block-diagonalized and, consequently,

$$\langle V_{\tau_{-1}(\overline{\mathcal{J}^*})}^{\mathcal{I}_{-1}} \rangle_{\Psi^{A/B}_{\tau_{-1}(\overline{\mathcal{J}^*})}} = \langle V_{\tau_{-1}(\overline{\mathcal{J}^*})}^{\mathcal{I}} \rangle_{\Psi^{A/B}_{\tau_{-1}(\overline{\mathcal{J}^*})}}, \qquad (4.31)$$

where $\tau_{-1}(\overline{\mathcal{J}^*})$ is indeed an interval belonging to $(\mathcal{I}_{+1}^*)'_{odd}$.

The identities in (4.30) and (4.31) imply that with each term in the first sum of (4.29) we can associate another one in the second sum that is exactly its opposite.

Then we can write

$$|E_{I_{+1}^{*}}^{B} - E_{I_{+1}^{*}}^{A} - 2h|$$

$$\leq \sqrt{t} \cdot \sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{ev} \setminus (I_{+1}^{*})_{ev}^{L}} |(\langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}})| + \sqrt{t} \cdot \sum_{\mathcal{J}^{*} \in (I_{+1}^{*})_{odd} \setminus (I_{+1}^{*})_{odd}^{\prime}} |(\langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}})|$$

$$\leq 4\sqrt{t} \cdot \sum_{K=1}^{\infty} C_{J,h} \cdot \frac{t^{\frac{K-1}{16}}}{K^{2}},$$

$$(4.33)$$

where for the last inequality we use that there is at most one interval of length *K* for each sum and its norm is bounded by $C_{J,h} \cdot \frac{t \frac{K-1}{16}}{K^2}$ by (4.1). This proves the inequality in the statement of the lemma. An analogous argument applies if I_{+1}^* has an even number of sites. \Box

Similarly to the ferromagnetic case, we derive

Lemma 4.5. Assuming that the bound in (4.1) holds in step I of the block-diagonalization, the following holds true

1. If the number of sites of \mathcal{I}_{+1}^* is odd

$$\pm \sum_{\overline{\mathcal{J}^{*}} \subset \mathcal{I}^{*}_{+1}} P^{(+)}_{\overline{\mathcal{J}^{*}}} V^{\overline{\mathcal{I}}}_{\overline{\mathcal{J}^{*}}} P^{(+)}_{\overline{\mathcal{J}^{*}}} \leq \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C_{J,h}}{2|J|-h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(\ell(\mathcal{J})+1\right) \left(H^{0}_{\mathcal{I}^{*}_{+1}}-\langle H^{0}_{\mathcal{I}^{*}_{+1}} \rangle_{\Psi^{A}_{\mathcal{I}^{*}_{+1}}}\right), \quad (4.34)$$

and

$$P_{\overline{I_{+1}^*}}^{(+)} \le \frac{1}{2|J|} \left(H_{\overline{I_{+1}^*}}^0 - \langle H_{\overline{I_{+1}^*}}^0 \rangle_{\Psi_{\overline{I_{+1}^*}}^A} \right), \tag{4.35}$$

while, concerning $\Psi^{B}_{I^{*}_{+1}}$,

$$\pm \sum_{\overline{\mathcal{J}^{*}} \subset I_{+1}^{*}} P_{\overline{\mathcal{J}^{*}}}^{(+)} V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}} P_{\overline{\mathcal{J}^{*}}}^{(+)} \leq \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C_{J,h}}{2|J|-h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(\ell(\mathcal{J})+1\right) \left(H_{I_{+1}^{*}}^{0}-\langle H_{I_{+1}}^{0} \rangle_{\Psi_{I_{+1}}^{B}}+2h\right); (4.36)$$

2. *if the number of sites of* I_{+1}^* *is even*

$$\pm \sum_{\overline{\mathcal{J}^{*} \subset \mathcal{I}^{*}_{+1}}} P_{\overline{\mathcal{J}^{*}}}^{(+)} V_{\overline{\mathcal{J}^{*}}}^{\overline{I}} P_{\overline{\mathcal{J}^{*}}}^{(+)} \leq \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C_{J,h}}{2|J|-h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(\ell(\mathcal{J})+1\right) \left(H_{\mathcal{I}^{*}_{+1}}^{0}-\langle H_{\mathcal{I}^{*}_{+1}}^{0} \rangle_{\Psi_{\mathcal{I}^{*}_{+1}}}^{A/B}+h\right), (4.37)$$

and

$$P_{\overline{I}_{+1}^*}^{(+)} \le \frac{1}{2|J| - 2h} \left(H_{\overline{I}_{+1}^*}^0 - \langle H_{\overline{I}_{+1}^*}^0 \rangle_{\Psi_{\overline{I}_{+1}^*}} \right).$$
(4.38)

Proof

From the definitions of H_I^0 and H_I^C in (1.7) and (1.9), respectively, we easily deduce: if I_{+1}^* contains an even number of sites

$$H_{\mathcal{I}_{+1}^{*}}^{C} - \langle H_{\mathcal{I}_{+1}^{*}}^{C} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A}} \le H_{\mathcal{I}_{+1}^{*}}^{0} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A}} + h$$
(4.39)

and

$$H_{I_{+1}^*}^C - \langle H_{I_{+1}^*}^C \rangle_{\Psi_{I_{+1}^*}^B} \le H_{I_{+1}^*}^0 - \langle H_{I_{+1}^*}^0 \rangle_{\Psi_{I_{+1}^*}^B} + h; \qquad (4.40)$$

if \mathcal{I}_{+1}^* contains an odd number of sites

$$H_{\mathcal{I}_{+1}^{*}}^{C} - \langle H_{\mathcal{I}_{+1}^{*}}^{C} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A}} \le H_{\mathcal{I}_{+1}^{*}}^{0} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A}}$$
(4.41)

$$H_{\mathcal{I}_{+1}^{*}}^{C} - \langle H_{\mathcal{I}_{+1}^{*}}^{C} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{B}} \le H_{\mathcal{I}_{+1}^{*}}^{0} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{B}} + 2h.$$
(4.42)

Analogously to (4.6), we can show that

$$\pm \sum_{\mathcal{J} \in (\mathcal{I}_{+1}^{*})_{K}} P_{\mathcal{J}^{*}}^{(+)} V_{\mathcal{J}^{*}}^{\mathcal{I}} P_{\mathcal{J}^{*}}^{(+)} \leq \frac{C_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J}) - 1}{16}} (K+1) \Big(H_{\mathcal{I}_{+1}^{*}}^{C} - \langle H_{\mathcal{I}_{+1}^{*}}^{C} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A/B}} \Big), \tag{4.43}$$

where $(\mathcal{I}_{+1}^*)_K$ is defined in item (*ii*) of Subsection 4.1.1. Finally, by combining (4.39)-(4.40) and (4.41)-(4.42) with (4.43), we get the inequalities in (4.34), (4.36), and (4.37).

Depending on the odd/even case the spectral gap changes according to Proposition 1.1, and consequently the inequalities (4.35) and (4.38) follow.

In the lemma below we use all the crucial ingredients: 1) the translation covariance of the model combined with the antiferromagnetic structure of the two groundstates of the unperturbed local Hamiltonians H_{I}^{C} ; 2) the block-diagonalization of the effective potentials $V_{\overline{J}^{*}}^{I}$ with respect to the triples of projections displayed in (a) and (b) of Lemma 4.3; 3) the spectral gap of the unperturbed local Hamiltonians.

Lemma 4.6. Assuming that the bound in (4.1) holds in step I of the block-diagonalization, there exists $\overline{t} > 0$ small enough such that $\forall t \leq \overline{t}$

$$P_{I_{+1}^*}^{(+)}\left(G_{I_{+1}^*} - E_{I_{+1}^*}^{A/B}\right)P_{I_{+1}^*}^{(+)} \ge (2|J| - 2h) \cdot \left[1 - O(\sqrt{t})\right]P_{I_{+1}^*}^{(+)}.$$
(4.44)

Proof

We recall that thanks to point (a) of Lemma 4.3

$$P_{I_{+1}^*}^{(+)} G_{I_{+1}^*} P_{I_{+1}^*}^{(+)} = P_{I_{+1}^*}^{(+)} (H_{I_{+1}^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I_{+1}^*} V_{\overline{\mathcal{J}^*}}^{\underline{\mathcal{I}}}) P_{I_{+1}^*}^{(+)}$$
(4.45)

$$= P_{I_{+1}^*}^{(+)} (H_{I_{+1}^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I_{+1}^*} P_{\overline{\mathcal{J}^*}}^{(+)} V_{\overline{\mathcal{J}^*}}^{I} P_{\overline{\mathcal{J}^*}}^{(+)}) P_{I_{+1}^*}^{(+)}$$
(4.46)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\overline{\mathcal{J}^{*}}\subset I_{+1}^{*}}\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}P_{\overline{\mathcal{J}^{*}}}^{(-),B}\right)P_{I_{+1}^{*}}^{(+)}$$
(4.47)

$$+P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\overline{\mathcal{J}^{*}}\subset\mathcal{I}_{+1}^{*}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{\mathcal{A}}}P_{\overline{\mathcal{J}^{*}}}^{(-),\mathcal{A}}\right)P_{\mathcal{I}_{+1}^{*}}^{(+)}.$$
(4.48)

Using the definitions in (4.15) and (4.16), we can write:

$$P_{\mathcal{I}_{+1}^{*}}^{(+)} G_{\mathcal{I}_{+1}^{*}} P_{\mathcal{I}_{+1}^{*}}^{(+)}$$
(4.49)

$$= P_{I_{+1}^*}^{(+)} (H_{I_{+1}^*}^0 + \sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I_{+1}^*} P_{\overline{\mathcal{J}^*}}^{(+)} V_{\overline{\mathcal{J}^*}}^{I} P_{\overline{\mathcal{J}^*}}^{(+)}) P_{I_{+1}^*}^{(+)}$$
(4.50)

$$+P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{ev}}\langle V_{\mathcal{J}^{*}}^{\mathcal{I}}\rangle_{\Psi_{\mathcal{J}^{*}}^{A}}P_{\mathcal{J}^{*}}^{(-),A}\right)P_{\mathcal{I}_{+1}^{*}}^{(+)}+P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{odd}}\langle V_{\mathcal{J}^{*}}^{\mathcal{I}}\rangle_{\Psi_{\mathcal{J}^{*}}^{A}}P_{\mathcal{J}^{*}}^{(-),A}\right)P_{\mathcal{I}_{+1}^{*}}^{(+)}$$
(4.51)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\mathcal{J}^{*}\in(\bar{I}_{+1}^{*})_{\text{odd}}}\langle V_{\mathcal{J}^{*}}^{\bar{I}}\rangle_{\Psi_{\mathcal{J}^{*}}^{B}}P_{\mathcal{J}^{*}}^{(-),B}\right)P_{I_{+1}^{*}}^{(+)}+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\mathcal{J}^{*}\in(\bar{I}_{+1}^{*})_{\text{ev}}}\langle V_{\mathcal{J}^{*}}^{\bar{I}}\rangle_{\Psi_{\mathcal{J}^{*}}^{B}}P_{\mathcal{J}^{*}}^{(-),B}\right)P_{I_{+1}^{*}}^{(+)}.$$
(4.52)

Next, for each $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{ev}}$ and $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{odd}}$, we add and subtract

$$\langle V^{I}_{\overline{\mathcal{J}^{*}}} \rangle_{\Psi^{A}_{\overline{\mathcal{J}^{*}}}} P^{(-),B}_{\overline{\mathcal{J}^{*}}} \quad , \quad \langle V^{I}_{\overline{\mathcal{J}^{*}}} \rangle_{\Psi^{B}_{\overline{\mathcal{J}^{*}}}} P^{(-),A}_{\overline{\mathcal{J}^{*}}} \, ,$$

respectively. Thus we obtain

$$P_{I_{+1}^*}^{(+)} G_{I_{+1}^*} P_{I_{+1}^*}^{(+)}$$
(4.53)

$$= P_{I_{+1}^{*}}^{(+)} (H_{I_{+1}^{*}}^{0} + \sqrt{t} \sum_{\overline{\mathcal{J}^{*}} \subset I_{+1}^{*}} P_{\overline{\mathcal{J}^{*}}}^{(+)} V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}} P_{\overline{\mathcal{J}^{*}}}^{(+)}) P_{I_{+1}^{*}}^{(+)}$$
(4.54)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(\tilde{I}_{+1}^{*})_{ev}}(\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}-\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}})P_{\overline{\mathcal{J}^{*}}}^{(-),B}\Big]P_{I_{+1}^{*}}^{(+)}$$
(4.55)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\mathcal{J}^{*}\in(I_{+1}^{*})_{\text{odd}}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}P_{\overline{\mathcal{J}^{*}}}^{(-)}\right)P_{I_{+1}^{*}}^{(+)}$$
(4.56)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\left[\sum_{\mathcal{J}^{*}\in(I_{+1}^{*})_{\text{odd}}}\left(\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}}-\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}}\right)P_{\overline{\mathcal{J}^{*}}}^{(-),A}\right]P_{I_{+1}^{*}}^{(+)}$$
(4.57)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\left(\sum_{\mathcal{J}^{*}\in(I_{+1}^{*})_{\text{ev}}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{\mathcal{A}}}P_{\overline{\mathcal{J}^{*}}}^{(-)}\right)P_{I_{+1}^{*}}^{(+)},$$
(4.58)

where we have used

$$\langle V_{\overline{\mathcal{J}^*}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}^*}}^{A/B}} \Big\{ P_{\overline{\mathcal{J}^*}}^{(-),B} + P_{\overline{\mathcal{J}^*}}^{(-),A} \Big\} = \langle V_{\overline{\mathcal{J}^*}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}^*}}^{A/B}} P_{\overline{\mathcal{J}^*}}^{(-)}.$$

The rest of the proof is separated into two parts, namely the study of (4.55) + (4.57) and of (4.56) + (4.58), respectively, and in a conclusion where we collect our partial estimates and finally prove the result stated in (4.44).

Study of the terms (4.55) + (4.57)

We intend to show some cancellations in the expression above. For this purpose we observe that if $\mathcal{J}^* \in (\mathcal{I}^*_{+1})'_{odd} \subset (\mathcal{I}^*_{+1})_{odd}$ (see the definition of $(\mathcal{I}^*_{+1})'_{odd}$ in (4.25)) then (i).a

$$P_{\overline{\mathcal{J}^*}}^{(-),A}P_{\overline{\mathcal{J}^*}\cup\tau_1(\overline{\mathcal{J}^*})}^{(-),A} = P_{\overline{\mathcal{J}^*}\cup\tau_1(\overline{\mathcal{J}^*})}^{(-),A},$$

(ii).a

$$P_{\overline{\mathcal{J}^*}}^{(-),A} P_{\overline{\mathcal{J}^*} \cup \tau_1(\overline{\mathcal{J}^*})}^{(-),B} = 0$$

Analogously, for $\mathcal{J}^* \in (\mathcal{I}^*_{+1})'_{ev} \subset (\mathcal{I}^*_{+1})_{ev}$, the following hold (i).b

(1).0

$$P_{\overline{\mathcal{J}^*}}^{(-),B} P_{\overline{\mathcal{J}^*} \cup \tau_{-1}(\overline{\mathcal{J}^*})}^{(-),B} = 0,$$

(ii).b

$$P_{\overline{\mathcal{J}^*}}^{(-),B}P_{\overline{\mathcal{J}^*}\cup\tau_{-1}(\overline{\mathcal{J}^*})}^{(-),A} = P_{\overline{\mathcal{J}^*}\cup\tau_{-1}(\overline{\mathcal{J}^*})}^{(-),A}$$

Therefore, regarding each summand in (4.57) associated with an interval $\mathcal{J}^* \in (\mathcal{I}^*_{+1})'_{odd}$, we decompose the identity into

$$\mathbb{1} = P_{\overline{\mathcal{J}^*} \cup \tau_1(\overline{\mathcal{J}^*})}^{(-),A} + P_{\overline{\mathcal{J}^*} \cup \tau_1(\overline{\mathcal{J}^*})}^{(-),B} + P_{\overline{\mathcal{J}^*} \cup \tau_1(\overline{\mathcal{J}^*})}^{(+)}; \qquad (4.59)$$

analogously, for an interval $\mathcal{J}^* \in (\mathcal{I}^*_{+1})'_{ev}$ in (4.55) we use

$$\mathbb{1} = P_{\overline{\mathcal{J}^* \cup \tau_{-1}(\overline{\mathcal{J}^*})}}^{(-),A} + P_{\overline{\mathcal{J}^* \cup \tau_{-1}(\overline{\mathcal{J}^*})}}^{(-),B} + P_{\overline{\mathcal{J}^* \cup \tau_{-1}(\overline{\mathcal{J}^*})}}^{(+)}.$$
(4.60)

Then we split $(\mathcal{I}_{+1}^*)_{\text{ev/odd}}$ into $(\mathcal{I}_{+1}^*)'_{\text{ev/odd}} \cup [(\mathcal{I}_{+1}^*)_{\text{ev/odd}} \setminus (\mathcal{I}_{+1}^*)'_{\text{ev/odd}}]$, and we substitute (4.60) into (4.55) and (4.59) into (4.57). For each $\mathcal{J} \in (\mathcal{I}_{+1}^*)'_{\text{ev}}$, the corresponding term in (4.55) is

$$P_{\mathcal{I}_{+1}^{*}}^{(+)} \left(\langle V_{\overline{\mathcal{J}}^{*}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}}^{*}}^{B}} - \langle V_{\overline{\mathcal{J}}^{*}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}}^{*}}^{A}} \right) P_{\overline{\mathcal{J}}^{*}}^{(-),B} P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

$$(4.61)$$

$$= P_{I_{+1}^*}^{(+)} \left(\langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^B} - \langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^A} \right) P_{\overline{\mathcal{J}^*}}^{(-),B} \mathbb{1} P_{I_{+1}^*}^{(+)}$$

$$(4.62)$$

$$= P_{\mathcal{I}_{+1}^{*}}^{(+)} \left(\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}} - \langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}} \right) P_{\overline{\mathcal{J}^{*}}}^{(-),B} P_{\overline{\mathcal{J}^{*}}}^{(-),A} P_{\mathcal{I}_{+1}}^{(+)}$$

$$(4.63)$$

$$+P_{I_{+1}^{*}}^{(+)}\left(\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}-\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}}\right)P_{\overline{\mathcal{J}^{*}}}^{(-),B}P_{\overline{\mathcal{J}^{*}}\cup\tau_{-1}(\overline{\mathcal{J}^{*}})}^{(-),B}P_{I_{+1}}^{(+)}$$

$$(4.64)$$

$$+P_{I_{+1}^{*}}^{(+)}\left(\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}-\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}}\right)P_{\overline{\mathcal{J}^{*}}}^{(-),B}P_{\overline{\mathcal{J}^{*}}\cup\tau_{-1}(\overline{\mathcal{J}^{*}})}^{(+)}P_{I_{+1}^{*}}^{(+)}.$$
(4.65)

Using property (ii).b above, (4.63) is equal to

$$P_{\mathcal{I}_{+1}^*}^{(+)}\left(\langle V_{\overline{\mathcal{J}^*}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^*}}^{\mathcal{B}}} - \langle V_{\overline{\mathcal{J}^*}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^*}}^{\mathcal{A}}}\right)P_{\overline{\mathcal{J}^*}\cup\tau_{-1}(\overline{\mathcal{J}^*})}^{(-),\mathcal{A}}P_{\mathcal{I}_{+1}^*}^{(+)},$$

while, using property (i).b above, (4.64) is equal to 0. Thus, we get

$$P_{\mathcal{I}_{+1}^{*}}^{(+)} \left(\langle V_{\overline{\mathcal{J}}^{*}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}}^{*}}} - \langle V_{\overline{\mathcal{J}}^{*}}^{\mathcal{I}} \rangle_{\Psi_{\overline{\mathcal{J}}^{*}}} \right) P_{\overline{\mathcal{J}}^{*}}^{(-),B} P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

$$(4.66)$$

$$= P_{I_{+1}^{*}}^{(+)} \Big(\langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} \Big) P_{\overline{\mathcal{J}^{*}\cup\tau_{-1}}(\overline{\mathcal{J}^{*}})}^{(-),A} P_{I_{+1}^{*}}^{(+)}$$

$$+ P_{I_{+1}^{*}}^{(+)} \Big(\langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \langle V_{\overline{\mathcal{J}^{*}}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} \Big) P_{\overline{\mathcal{J}^{*}}}^{(-),B} P_{\overline{\mathcal{J}^{*}}}^{(+)} P_{I_{+1}^{*}}^{(+)}.$$

$$(4.67)$$

Analogously, from the properties in (i).a and (ii).a above, we deduce that, for each $\mathcal{J} \in (\mathcal{I}_{+1}^*)'_{odd}$, the corresponding term in (4.57) is

$$P_{I_{+1}^{*}}^{(+)} \left(\left\langle V_{\overline{\mathcal{J}^{*}}}^{I} \right\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \left\langle V_{\overline{\mathcal{J}^{*}}}^{I} \right\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} \right) P_{\overline{\mathcal{J}^{*}}}^{(-),A} P_{I_{+1}^{*}}^{(+)}$$

$$= P_{I_{+1}^{*}}^{(+)} \left(\left\langle V_{\overline{\mathcal{J}^{*}}}^{I} \right\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \left\langle V_{\overline{\mathcal{J}^{*}}}^{I} \right\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} \right) P_{\overline{\mathcal{J}^{*}}\cup\tau_{1}(\overline{\mathcal{J}^{*}})}^{(-),A} P_{I_{+1}^{*}}^{(+)}$$

$$+ P_{I_{+1}^{*}}^{(+)} \left(\left\langle V_{\overline{\mathcal{J}^{*}}}^{I} \right\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} - \left\langle V_{\overline{\mathcal{J}^{*}}}^{I} \right\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}} \right) P_{\overline{\mathcal{J}^{*}}}^{(-),A} P_{\overline{\mathcal{J}^{*}}}^{(+)} P_{I_{+1}^{*}}^{(+)} .$$

$$(4.68)$$

For the terms corresponding to intervals in $(\mathcal{I}_{+1}^*)_{ev/odd} \setminus (\mathcal{I}_{+1}^*)'_{ev/odd}$ we do nothing. Thus, we have

$$(4.55) + (4.57) = P_{I_{+1}^*}^{(+)} \sqrt{t} \Big[\sum_{\mathcal{J}^* \in (\overline{I_{+1}^*})_{\text{odd}}'} \Big(\langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^A} - \langle V_{\overline{\mathcal{J}^*}}^I \rangle_{\Psi_{\overline{\mathcal{J}^*}}^B} \Big) P_{\overline{\mathcal{J}^*} \cup \tau_1(\overline{\mathcal{J}^*})}^{(-),A} \Big] P_{I_{+1}^*}^{(+)}$$

$$(4.69)$$

$$+P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{\text{odd}}^{\prime}}\Big(\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}}-\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}\Big)P_{\overline{\mathcal{J}^{*}}}^{(-),A}P_{\overline{\mathcal{J}^{*}}}^{(+)}|_{\mathcal{I}_{+1}^{*}}\Big]P_{\mathcal{I}_{+1}^{*}}^{(+)}$$
(4.70)

$$+P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{ev}^{\prime}}\left(\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}-\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}}\right)P_{\overline{\mathcal{J}^{*}\cup\tau_{-1}(\overline{\mathcal{J}^{*}})}^{(-),A}\Big]P_{\mathcal{I}_{+1}^{*}}^{(+)}$$
(4.71)

$$+P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{ev}^{\prime}}(\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}-\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}})P_{\overline{\mathcal{J}^{*}}}^{(-),B}P_{\overline{\mathcal{J}^{*}}\cup\tau_{-1}(\overline{\mathcal{J}^{*}})}^{(+)}\Big]P_{\mathcal{I}_{+1}^{*}}^{(+)}$$
(4.72)

$$+P_{I_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(I_{+1}^{*})_{ev}\setminus(I_{+1}^{*})_{ev}^{\prime}}\left(\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}}-\langle V_{\overline{\mathcal{J}^{*}}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}}\right)P_{\overline{\mathcal{J}^{*}}}^{(-),B}\Big]P_{I_{+1}^{*}}^{(+)}$$

$$(4.73)$$

$$+P_{\mathcal{I}_{+1}^{*}}^{(+)} \sqrt{t} \Big[\sum_{\mathcal{J}^{*} \in (\mathcal{I}_{+1}^{*})_{\text{odd}} \setminus (\mathcal{I}_{+1}^{*})_{\text{odd}}'} \Big(\langle V_{\mathcal{J}^{*}}^{\mathcal{I}} \rangle_{\Psi_{\mathcal{J}^{*}}^{A}} - \langle V_{\mathcal{J}^{*}}^{\mathcal{I}} \rangle_{\Psi_{\mathcal{J}^{*}}^{B}} \Big) P_{\mathcal{J}^{*}}^{(-),A} \Big] P_{\mathcal{I}_{+1}^{*}}^{(+)} .$$

$$(4.74)$$

We proceed our study of (4.55) + (4.57) by combining different terms in (4.69)-(4.74). We start with the estimate of the following terms

• (4.69) + (4.71).

From (4.26) we have that for each $\mathcal{J}^* \in (\mathcal{I}^*_{+1})'_{odd}$, the projection $P^{(-),A}_{\overline{\mathcal{J}^*} \cup \tau_1(\overline{\mathcal{J}^*})}$ is equal to the projection $P^{(-),A}_{\overline{\mathcal{K}^*} \cup \tau_{-1}(\overline{\mathcal{K}^*})}$ where $\mathcal{K} = \tau_1(\mathcal{J}) \in (\mathcal{I}^*_{+1})'_{even}$. Thus each term in (4.69) is paired with a term in (4.71); their combination gives

$$\left(\langle V_{\overline{\mathcal{J}^*}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^*}}^{A}} - \langle V_{\overline{\mathcal{J}^*}}^{I}\rangle_{\Psi_{\overline{\mathcal{J}^*}}^{B}} + \langle V_{\tau_1(\overline{\mathcal{J}^*})}^{I}\rangle_{\Psi_{\tau_1(\overline{\mathcal{J}^*})}^{B}} - \langle V_{\tau_1(\overline{\mathcal{J}^*})}^{I}\rangle_{\Psi_{\tau_1(\overline{\mathcal{J}^*})}^{A}}\right)P_{\overline{\mathcal{J}^*}\cup\tau_1(\overline{\mathcal{J}^*})}^{(-),A}$$
(4.75)

Now we recall that (see an analogous identity in (4.30)-(4.31))

$$\langle V_{\overline{\mathcal{J}^*}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^*}}^{B}} = \langle V_{\tau_1(\overline{\mathcal{J}^*})}^{I} \rangle_{\Psi_{\tau_1(\overline{\mathcal{J}^*})}^{B}}, \qquad \langle V_{\overline{\mathcal{J}^*}}^{I} \rangle_{\Psi_{\overline{\mathcal{J}^*}}^{A}} = \langle V_{\tau_1(\overline{\mathcal{J}^*})}^{I} \rangle_{\Psi_{\tau_1(\overline{\mathcal{J}^*})}^{A}}, \qquad (4.76)$$

which follow from the translation covariance stated in Proposition 3.7. Thus we conclude that

$$(4.69) + (4.71) = 0. (4.77)$$

• (4.73) + (4.74).

For each $K \in \mathbb{N}$, there is at most one $\mathcal{J} \in \mathfrak{I}$ with $\ell(\mathcal{J}) = K$ and $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{odd} \setminus (\mathcal{I}^*_{+1})'_{odd}$; analogously, there is at most one $\mathcal{K} \in \mathfrak{I}$ with $\ell(\mathcal{K}) = K$ such that $\mathcal{K}^* \in (\mathcal{I}^*_{+1})_{ev} \setminus (\mathcal{I}^*_{+1})'_{ev}$. Thanks to (4.1), this implies

$$\pm ((4.73) + (4.74)) \le 4 \sqrt{t} \cdot \sum_{K=1}^{\ell(I)-1} C_{J,h} \cdot \frac{t^{\frac{K-1}{16}}}{K^2} P_{I_{+1}^*}^{(+)}.$$
(4.78)

• (4.70) + (4.72).

We use Lemma 4.5 and estimate (the factor 4 in (4.79) and (4.80) comes from applying the estimates in (4.34) (odd case) and in (4.37) (even case) to each of the two terms contained in (4.70) and (4.72)):

- if \mathcal{I}^* has an odd number of sites,

$$\pm ((4.70) + (4.72))$$

$$\leq P_{I_{+1}^{*}}^{(+)} \sqrt{t} \Big[\sum_{K=2}^{\ell(I)-1} \sum_{\mathcal{K} \in (I_{+1}^{*})_{K}} (4 \cdot C_{J,h} \cdot \frac{t^{\frac{K-2}{16}}}{(K-1)^{2}}) P_{\overline{\mathcal{K}^{*}}}^{(+)} \Big] P_{I_{+1}^{*}}^{(+)}$$

$$\leq P_{I_{+1}^{*}}^{(+)} \sqrt{t} \Big[\sum_{K=2}^{\ell(I)-1} (4 \cdot C_{J,h} \cdot \frac{t^{\frac{K-2}{16}}}{(K-1)^{2}}) (K+1) \frac{1}{2|J|-h} (H_{I_{+1}}^{0} - \langle H_{I_{+1}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}) \Big] P_{I_{+1}^{*}}^{(+)}; \quad (4.79)$$

- if \mathcal{I}^* has an even number of sites,

$$\pm ((4.70) + (4.72)) \\
\leq P_{I_{+1}^{*}}^{(+)} \sqrt{t} \Big[\sum_{K=2}^{\ell(J)-1} \sum_{\mathcal{K} \in (I_{+1}^{*})K} (4 \cdot C_{J,h} \cdot \frac{t^{\frac{K-2}{16}}}{(K-1)^{2}}) P_{\overline{\mathcal{K}^{*}}}^{(+)} \Big] P_{I_{+1}^{*}}^{(+)} \\
\leq P_{I_{+1}^{*}}^{(+)} \sqrt{t} \Big[\sum_{K=2}^{\ell(J)-1} (4 \cdot C_{J,h} \cdot \frac{t^{\frac{K-2}{16}}}{(K-1)^{2}}) (K+1) \frac{1}{2|J|-h} (H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{A} + h) \Big] P_{I_{+1}^{*}}^{(+)} . \quad (4.80)$$

where $(\mathcal{I}_{+1}^*)_K$ is defined in item (*ii*) of Subsection 4.1.1.

Study of the terms (4.54) + (4.56) + (4.58)

By using $P_{\overline{\mathcal{J}^*}}^{(-)} = \mathbb{1} - P_{\overline{\mathcal{J}^*}}^{(+)}$, we can write

$$(4.56) + (4.58) \tag{4.81}$$

$$= P_{I_{+1}^*}^{(+)} \sqrt{t} \Big[\sum_{\mathcal{J}^* \in (I_{+1}^*)_{\text{ev}}} \langle V_{\mathcal{J}^*}^I \rangle_{\Psi_{\mathcal{J}^*}^A} + \sum_{\mathcal{J}^* \in (I_{+1}^*)_{\text{odd}}} \langle V_{\mathcal{J}^*}^I \rangle_{\Psi_{\mathcal{J}^*}^B} \Big] P_{I_{+1}^*}^{(+)}$$
(4.82)

$$-P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{\text{ev}}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{A}}P_{\overline{\mathcal{J}^{*}}}^{(+)} + \sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{\text{odd}}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{B}}P_{\overline{\mathcal{J}^{*}}}^{(+)}\Big]P_{\mathcal{I}_{+1}^{*}}^{(+)}.$$
(4.83)

We recall that

(i) for $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{ev}}$, $\langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^A_{\mathcal{J}^*}} = \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^A_{\mathcal{I}^*_{+1}}}$ and $\langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^B_{\mathcal{J}^*}} = \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^B_{\mathcal{I}^*_{+1}}}$, (ii) for $\mathcal{J}^* \in (\mathcal{I}^*)$ is $\langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\mathcal{I}^*} = \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\mathcal{I}^*_{+1}}$ and $\langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\mathcal{I}^*_{+1}} = \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\mathcal{I}^*_{+1}}$,

(ii) for $\mathcal{J}^* \in (\mathcal{I}^*_{+1})_{\text{odd}}$, $\langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^A_{\mathcal{J}^*}} = \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^B_{I^*_{+1}}} \text{ and } \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^B_{\mathcal{J}^*}} = \langle V^{\mathcal{I}}_{\mathcal{J}^*} \rangle_{\Psi^A_{I^*_{+1}}}$. Consequently, we get

 $(4.82) = (E_{\mathcal{I}_{+1}^*}^A - \langle H_{\mathcal{I}_{+1}^*}^0 \rangle_{\Psi_{\mathcal{I}_{+1}^*}^A}) P_{\mathcal{I}_{+1}^*}^{(+)} .$ (4.84)

Hence, we have

$$(4.54) + (4.56) + (4.58) \tag{4.85}$$

$$= P_{\mathcal{I}_{+1}^{*}}^{(+)} (H_{\mathcal{I}_{+1}^{*}}^{0} + E_{\mathcal{I}_{+1}^{*}}^{A} - \langle H_{\mathcal{I}_{+1}^{*}}^{0} \rangle_{\Psi_{\mathcal{I}_{+1}^{*}}^{A}}) P_{\mathcal{I}_{+1}^{*}}^{(+)}$$

$$(4.86)$$

$$+P_{I_{+1}^{*}}^{(+)}\left(\sqrt{t}\sum_{\overline{\mathcal{J}^{*}}\subset I_{+1}^{*}}P_{\overline{\mathcal{J}^{*}}}^{(+)}V_{\overline{\mathcal{J}^{*}}}^{I}P_{\overline{\mathcal{J}^{*}}}^{(+)}\right)P_{I_{+1}^{*}}^{(+)}$$
(4.87)

$$-P_{\mathcal{I}_{+1}^{*}}^{(+)}\sqrt{t}\Big[\sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{ev}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{\mathcal{A}}}P_{\overline{\mathcal{J}^{*}}}^{(+)} + \sum_{\mathcal{J}^{*}\in(\mathcal{I}_{+1}^{*})_{odd}}\langle V_{\overline{\mathcal{J}^{*}}}^{\mathcal{I}}\rangle_{\Psi_{\overline{\mathcal{J}^{*}}}^{\mathcal{B}}}P_{\overline{\mathcal{I}^{*}}}^{(+)}\Big]P_{\mathcal{I}_{+1}^{*}}^{(+)}.$$
(4.88)

Concerning the terms in (4.87)-(4.88), we must distinguish between odd/even number of sites so as to apply Lemma 4.5 similarly to what we have done for (4.70)+(4.72).

Conclusion

We now collect our estimates obtained so far and, depending on the parity of the number of sites of I_{+1}^* , we show how to derive the inequalities in the statement. In the following $C'_{J,h}$ stands for a quantity which depends on J and h, and may vary line by line.

For an odd number of sites, we use point 1 in Lemma 4.5, to get

$$P_{I_{+1}^{(+)}}^{(+)} \left(G_{I_{+1}^{*}} - E_{I_{+1}^{*}}^{A}\right) P_{I_{+1}^{(+)}}^{(+)} \\ \geq \left(1 - \sqrt{t} \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(\ell(\mathcal{J}) + 1\right)\right) \left(H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{A}\right) P_{I_{+1}^{*}}^{(+)} \quad (4.89) \\ - \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{J})-1} C'_{J,h} \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{(\ell(\mathcal{J}))^{2}} P_{I_{+1}^{*}}^{(+)} \\ \geq \left(1 - \sqrt{t} \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} \left(\ell(\mathcal{J}) + 1\right)\right) 2|J| P_{I_{+1}^{*}}^{(+)} \quad (4.90) \\ - \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} C'_{J,h} \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{(\ell(\mathcal{J}))^{2}} P_{I_{+1}^{*}}^{(+)} \\ \end{cases}$$

where we have used (4.35) to estimate the term proportional to $H^0_{\mathcal{I}^*_{+1}} - \langle H^0_{\mathcal{I}^*_{+1}} \rangle_{\Psi^A_{\mathcal{I}^*_{+1}}}$ in (4.89).

For an even number of sites, we use point 2. in Lemma 4.5, to get

$$P_{I_{+1}^{*}}^{(+)} (G_{I_{+1}^{*}} - E_{I_{+1}^{*}}^{A}) P_{I_{+1}^{*}}^{(+)}$$

$$\geq \left[1 - \sqrt{t} \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J}) + 1) \right] (H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{A}) P_{I_{+1}^{*}}^{(+)}$$

$$= \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} C'_{J,h} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J}) + 1) \right] P_{I_{+1}^{*}}^{(+)}$$

$$\geq \left[1 - \sqrt{t} \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J}) + 1) \right] (2|J| - 2h) P_{I_{+1}^{*}}^{(+)}$$

$$= \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{I})-1}{16}} (\ell(\mathcal{J}) + 1) \right] (2|J| - 2h) P_{I_{+1}^{*}}^{(+)}$$

$$+ h \sqrt{t} \cdot \left[- \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{I})-1}{16}} (\ell(\mathcal{J}) + 1) \right] P_{I_{+1}^{*}}^{(+)}$$

$$+ h \sqrt{t} \cdot \left[- \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{I})-1}{16}} (\ell(\mathcal{J}) + 1) \right] P_{I_{+1}^{*}}^{(+)}$$

$$+ h \sqrt{t} \cdot \left[- \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{I})-1}{16}} (\ell(\mathcal{I}) + 1) \right] P_{I_{+1}^{*}}^{(+)}$$

where we have used (4.38) to estimate the term proportional to $H^0_{I^*_{+1}} - \langle H^0_{I^*_{+1}} \rangle_{\Psi^A_{I^*_{+1}}}$ in (4.91). From the inequalities in (4.90) and (4.92) we easily deduce the statement in (4.44) regarding $G_{I^*_{+1}} - E^A_{I^*_{+1}}$.

We can prove the estimate concerning $P_{I_{+1}^*}^{(+)}(G_{I_{+1}^*} - E_{I_{+1}^*}^B)P_{I_{+1}^*}^{(+)}$ by repeating almost verbatim the proof used for $P_{I_{+1}^*}^{(+)}(G_{I_{+1}^*} - E_{I_{+1}^*}^A)P_{I_{+1}^*}^{(+)}$. Namely, in order to get inequalities analogous to (4.90) and (4.92) and finally the statement in (4.44), we have to use (4.36) (in point *I*) or (4.37) (in point 2) of Lemma 4.5 for a number of sites of I_{+1}^* odd or even, respectively:

odd number of sites

$$P_{I_{+1}^{*}}^{(+)} (G_{I_{+1}^{*}} - E_{I_{+1}^{*}}^{B}) P_{I_{+1}^{*}}^{(+)}$$

$$\geq P_{I_{+1}^{*}}^{(+)} (H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{B}) P_{I_{+1}^{*}}^{(+)}$$

$$+ \sqrt{t} P_{I_{+1}^{*}}^{(+)} \Big[-\sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J|-h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J})+1) \Big] (H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{B} + 2h) P_{I_{+1}^{*}}^{(+)}$$

$$- \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{J})-1} C'_{J,h} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{(\ell(\mathcal{J}))^{2}} P_{I_{+1}^{*}}^{(+)}$$

$$\geq \Big[1 - \sqrt{t} \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J|-h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J})+1) \Big] (2|J|-2h) P_{I_{+1}^{*}}^{(+)}$$

$$- \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} C'_{J,h} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{(\ell(\mathcal{J}))^{2}} P_{I_{+1}^{*}}^{(+)}$$

$$+ 2h \sqrt{t} \cdot \Big[-\sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J|-h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J})+1) \Big] P_{I_{+1}^{*}}^{(+)} ;$$

even number of sites

$$P_{I_{+1}^{*}}^{(+)} (G_{I_{+1}^{*}} - E_{I_{+1}^{*}}^{B}) P_{I_{+1}^{*}}^{(+)}$$

$$\geq P_{I_{+1}^{*}}^{(+)} (H_{I_{+1}^{*}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{B}) P_{I_{+1}^{*}}^{(+)}$$

$$+ \sqrt{t} P_{I_{+1}^{*}}^{(+)} \Big[-\sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J}) + 1) \Big] (H_{I_{+1}}^{0} - \langle H_{I_{+1}^{*}}^{0} \rangle_{\Psi_{I_{+1}^{*}}}^{B} + h) P_{I_{+1}^{*}}^{(+)}$$

$$- \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} C'_{J,h} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{(\ell(\mathcal{J}))^{2}} P_{I_{+1}^{*}}^{(+)}$$

$$\geq \Big[1 - \sqrt{t} \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{J})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J}) + 1) \Big] (2|J| - 2h) P_{I_{+1}^{*}}^{(+)}$$

$$- \sqrt{t} \cdot \sum_{\ell(\mathcal{J})=1}^{\ell(\mathcal{I})-1} \frac{C'_{J,h}}{2|J| - h} \cdot t^{\frac{\ell(\mathcal{J})-1}{16}} (\ell(\mathcal{J}) + 1) \Big] P_{I_{+1}^{*}}^{(+)} .$$

4.2 Estimates of operator norms of potentials and main theorem

In this section we prove our main results. Namely, in Theorem 4.9 we collect the preliminary ingredients and show by induction the crucial bounds on the operator norms of the potentials assumed in some of the previous arguments. This proves that the block-diagonalization can be

implemented up to the last step. In Theorem 4.11 we draw the conclusions about the low lying spectrum of the XXZ Hamiltonian in (1.3).

In order to make Theorem 4.9 as straightforward as possible, we prepare the ground in Sections 4.2.1 and 4.2.2 below, where we defer parts of the proof by induction (of Theorem 4.9); namely we study some of the expressions entering the algorithm, in order to estimate their operator norms in terms of the norms of the effective potentials at a given step. Similarly, the control of the Lie-Schwinger series (which is part of the induction) is deferred to Lemma 4.10.

4.2.1 Hooked potentials

Assuming the inductive hypothesis in (4.1) and the bounds in (4.184), (4.182), and (4.183) proven in Lemma 4.10, for *t* sufficiently small we readily derive the following relations.

• Ferromagnetic case

$$\left\|\sum_{n=1}^{\infty} \frac{1}{n!} a d^{n} Z_{I^{*}}(V_{\overline{\mathcal{K}^{*}}}^{I_{-1}})\right\|$$
(4.95)

$$\leq C \cdot \frac{A_1}{J+h} \cdot \sqrt{t} \cdot \|V_{\overline{I}}^{I_{-1}}\| \cdot \|V_{\overline{K}^*}^{I_{-1}}\|$$

$$(4.96)$$

$$\leq C_{J,h} \cdot C \cdot \frac{A_1}{J+h} \cdot \sqrt{t} \cdot \|V_{I}^{I_{-1}}\| \cdot \|V_{\mathcal{K}}^{I_{-1}}\|.$$
(4.97)

where *C* and A_1 are universal constants, and we recall that $C_{J,h}$ is defined in (4.1). By similar steps we can prove that

$$\left\|\sum_{n=1}^{\infty} \frac{1}{n!} a d^{n} Z_{\mathcal{I}^{*}}(V_{\mathcal{K}}^{\mathcal{I}_{-1}})\right\| \leq C \cdot \frac{A_{1}}{J+h} \cdot \sqrt{t} \cdot \|V_{\mathcal{I}}^{\mathcal{I}_{-1}}\| \cdot \|V_{\mathcal{K}}^{\mathcal{I}_{-1}}\|.$$
(4.98)

• Antiferromagnetic case

$$\left\|\sum_{n=1}^{\infty} \frac{1}{n!} a d^{n} Z_{I^{*}}(V_{\overline{\mathcal{K}^{*}}}^{I_{-1}})\right\|$$
(4.99)

$$\leq C \cdot \frac{A_1}{|J| - h} \cdot \sqrt{t} \cdot \|V_I^{I_{-1}}\| \cdot \|V_{\overline{\mathcal{K}^*}}^{I_{-1}}\|$$
(4.100)

$$\leq C_{J,h} \cdot C \cdot \frac{A_1}{|J| - h} \cdot \sqrt{t} \cdot \|V_I^{I_{-1}}\| \cdot \|V_{\mathcal{K}}^{I_{-1}}\|$$
(4.101)

and

$$\left\|\sum_{n=1}^{\infty} \frac{1}{n!} a d^{n} Z_{\mathcal{I}^{*}}(V_{\mathcal{K}}^{\mathcal{I}_{-1}})\right\| \leq C \cdot \frac{A_{1}}{|\mathcal{J}| - h} \cdot \sqrt{t} \cdot \|V_{\mathcal{I}}^{\mathcal{I}_{-1}}\| \cdot \|V_{\mathcal{K}}^{\mathcal{I}_{-1}}\|.$$
(4.102)

4.2.2 Off-diagonal part of the hooked Ising terms and Lieb-Robinson bounds

For an Hamiltonian like $H_{\mathcal{J}}^0$, defined in (1.7), a Lieb-Robinson bound (see [LR]) on the speed of propagation of observables holds as shown in [NS]. We follow the notation used in [NS], where the authors introduce a family of functions labelled by a parameter a > 0

$$F_a: [0,\infty) \to (0,\infty)$$
 , $F_a(r) := e^{-ar} e^{-\sqrt{r}} \frac{1}{(1+r)^3}$, (4.103)

which belong to the class of \mathcal{F} -functions [NS], namely they satisfy the properties

- $||F_a|| := \sum_{i \in \mathbb{Z}^+} F_a(i) < \infty$,
- there exists $C_a > 0$ such that for all $i, j \in \mathbb{Z}$,

$$\sum_{z \in \mathbb{Z}} F_a(|i-z|) F_a(|z-j|) \le C_a F_a(|i-j|);$$
(4.104)

see Section 6.1 of [MN]. Now, let $\{\exp(isH_{\mathcal{J}}^0), s \in \mathbb{R}\}\$ be the one-parameter group generated by $H_{\mathcal{J}}^0, \mathcal{J} \subseteq \Lambda$, then, for given observables *A*, *B* localized in intervals $\mathcal{I}_1, \mathcal{I}_2$ respectively, by Eq. (16) in [NS] we have

$$\|[\exp(isH_{\mathcal{J}}^{0})A \exp(-isH_{\mathcal{J}}^{0}), B]\| \le \frac{4 \|A\| \|B\| \|F_{0}\|}{C_{a}} \cdot e^{-a \cdot [d(\mathcal{I}_{1}, \mathcal{I}_{2}) - \frac{2\|\Phi\|_{a}C_{a}|s|}{a}]},$$
(4.105)

where $d(I_1, I_2)$ is the distance between the intervals I_1, I_2 , e.g., referring to the notation introduced in (4.15), in case I_1 sits on the left of I_2 and $I_1 \cap I_2 = \emptyset$ then the distance is given by $j_\ell(I_2) - j_r(I_1)$, where $j_r(I_1)$ is the right-most site of I_1 , and

$$\|\Phi\|_{a} := \max_{i \in \Lambda} \left\{ \frac{\|-J\sigma_{i}^{z}\sigma_{i+1}^{z} - \frac{h}{2}\sigma_{i}^{z} - \frac{h}{2}\sigma_{i+1}^{z}\|}{F_{a}(1)}; \frac{\|\frac{h}{2}\sigma_{i}^{z}\|}{F_{a}(0)} \right\}$$
(4.106)

which in our setting is trivially uniformly bounded in a. In the sequel we will set a = 1.

Lemma 4.7. Assuming S1) and S2) of Theorem 4.9 in step I_{-1} , the following inequality holds true

$$\left\| P_{\overline{I^*}}^{(+)} \left(ad \, Z_{\overline{I^*}} \left(\frac{\sigma_{i_-^* - 1}^z \sigma_{i_-^*}^z}{t^{1/2}} \right) \right) P_{\overline{I^*}}^{(-)} \right\| \le O(t^{\frac{1}{4}} \cdot \|V_{\overline{I}}^{\overline{I}_{-1}}\|) \,. \tag{4.107}$$

Proof

We treat the antiferromagnetic case first and then explain how to recover the ferromagnetic case from it.

Recall the formulae introduced in Section 3.1 and write

$$P_{\overline{I^*}}^{(+)} \left(ad Z_{\overline{I^*}} \left(\frac{\sigma_{i_-}^{\zeta_*} - 1 \sigma_{i_-}^{\zeta_*}}{t^{1/2}} \right) \right) P_{\overline{I^*}}^{(-)} = P_{\overline{I^*}}^{(+)} \left[Z_{\overline{I^*}} , \frac{\sigma_{i_-}^{\zeta_*} - 1 \sigma_{i_-}^{\zeta_*}}{t^{1/2}} \right] P_{\overline{I^*}}^{(-)}$$

$$(4.108)$$

$$= P_{\overline{I^*}}^{(+)} \left[Z_{\overline{I^*}}, \frac{\sigma_{i_-^*-1}^* \sigma_{i_-^*}^*}{t^{1/2}} + \frac{1}{t^{1/2}} \right] P_{\overline{I^*}}^{(-)}$$
(4.109)

$$= -P_{\overline{I^*}}^{(+)} \left(\frac{\sigma_{i_-^*-1}^{\zeta} \sigma_{i_-^*}^{\zeta}}{t^{1/2}} + \frac{1}{t^{1/2}} \right) Z_{\overline{I^*}} P_{\overline{I^*}}^{(-)}$$
(4.110)

$$= -\sum_{j=1}^{\infty} t^{\frac{j}{2}} P_{\overline{I^*}}^{(+)} \left(\frac{\sigma_{i^*_{-}-1}^{z} \sigma_{i^*_{-}}^{z}}{t^{1/2}} + \frac{1}{t^{1/2}} \right) (Z_{I^*})_j P_{\overline{I^*}}^{(-)}, \quad (4.111)$$

where in the step from (4.108) to (4.110) we have used

$$(\frac{\sigma_{i_--1}^z\sigma_{i_-}^z}{t^{1/2}}+\frac{1}{t^{1/2}})P_{\overline{I^*}}^{(-)}=0\,.$$

In expression (4.111) above we can discard all the terms of the series starting from j = 2, i.e.,

$$-\sum_{j=2}^{\infty} t^{\frac{j}{2}} P_{\overline{I^*}}^{(+)} \left(\frac{\sigma_{i^*_*-1}^z \sigma_{i^*_*}^z}{t^{1/2}} + \frac{1}{t^{1/2}} \right) (Z_{\overline{I^*}})_j P_{\overline{I^*}}^{(-)}, \qquad (4.112)$$

since ||(4.112)|| is bounded by $O(t^{1/2} \cdot ||V_I^{\mathcal{I}_{-1}}||^2)$ and, consequently, the bound in (4.107) is fulfilled thanks to S1) of Theorem 4.9 in step \mathcal{I}_{-1} . Now we focus on the remaining quantity

$$-t^{\frac{1}{2}}P_{\overline{I^*}}^{(+)}\left(\frac{\sigma_{i^*_{-}-1}^{z}\sigma_{i^*_{-}}^{z}}{t^{1/2}}+\frac{1}{t^{1/2}}\right)(Z_{I^*})_1P_{\overline{I^*}}^{(-)}$$
(4.113)

which, in the antiferromagnetic case, corresponds to

$$-P_{\overline{I^*}}^{(+)}(\sigma_{i^*_{-}-1}^z\sigma_{i^*_{-}}^z+1)\frac{1}{G_{I^*}-E_{I^*}^A}P_{I^*}^{(+)}V_{I}^{\overline{I}_{-}1}P_{I^*}^{(-),A}P_{\overline{I^*}}^{(-)}$$
(4.114)

$$-P_{\overline{I^*}}^{(+)} \left(\sigma_{i^*_{-}-1}^z \sigma_{i^*_{-}}^z + 1\right) \frac{1}{G_{I^*} - E_{I^*}^B} P_{I^*}^{(+)} V_I^{\mathcal{I}_{-1}} P_{I^*}^{(-),B} P_{\overline{I^*}}^{(-)}.$$
(4.115)

We make use of the identity

$$\frac{1}{G_{\mathcal{I}^*} - E_{\mathcal{I}^*}^{A/B}} P_{\mathcal{I}^*}^{(+)} = \frac{1}{G_{\mathcal{I}^*} - E_{\mathcal{I}^*}^{A/B} + i\delta_t} P_{\mathcal{I}^*}^{(+)} + \frac{i\delta_t}{G_{\mathcal{I}^*} - E_{\mathcal{I}^*}^{A/B}} \frac{1}{G_{\mathcal{I}^*} - E_{\mathcal{I}^*}^{A/B} + i\delta_t} P_{\mathcal{I}^*}^{(+)}$$
(4.116)

where δ_t is set equal to $t^{\frac{1}{4}}$. Then, thanks to the gap bound (4.44) proven in Lemma 4.6 (in step I_{-1}), we can write

$$(4.114) + (4.115) = -P_{\overline{I^*}}^{(+)} (\sigma_{i^*_{-1}}^z \sigma_{i^*_{-}}^z + 1) \frac{1}{G_{I^*} - E_{\overline{I^*}}^A + i\delta_t} P_{I^*}^{(+)} V_{I}^{I_{-1}} P_{\overline{I^*}}^{(-),A} P_{\overline{I^*}}^{(-)} (4.117) -P_{\overline{I^*}}^{(+)} (\sigma_{i^*_{-1}}^z \sigma_{i^*_{-}}^z + 1) \frac{1}{G_{I^*} - E_{\overline{I^*}}^B + i\delta_t} P_{I^*}^{(+)} V_{I}^{I_{-1}} P_{\overline{I^*}}^{(-),B} P_{\overline{I^*}}^{(-)} (4.118) +R_1$$

$$(4.119)$$

with $||R_1|| \le O(\frac{\delta_t}{2|J|-2h} \cdot ||V_I^{I_{-1}}||)$. The remainder term, R_1 , appearing in (4.119) fulfills the bound in (4.107), whereas the first and second terms require some further manipulation explained below. Since the latter ones are estimated in the same way, we will proceed by analyzing the right hand side of (4.117) only. For this purpose, we use the Neumann expansion displayed below

$$\frac{1}{G_{I^{*}} - E_{I^{*}}^{A} + i\delta_{t}}P_{I^{*}}^{(+)} = \frac{1}{P_{I^{*}}^{(+)}(G_{I^{*}} - E_{I^{*}}^{A} + i\delta_{t})P_{I^{*}}^{(+)}}P_{I^{*}}^{(+)} = \frac{1}{H_{I^{*}}^{0} - \langle H_{I^{*}}^{0} \rangle_{\Psi_{I^{*}}^{A}} + i\delta_{t}}P_{I^{*}}^{(+)} + (4.120) + \frac{1}{H_{I^{*}}^{0} - \langle H_{I^{*}}^{0} \rangle_{\Psi_{I^{*}}^{A}} + i\delta_{t}}\sum_{j=1}^{\infty} \left[\left(-\sqrt{t} \sum_{\overline{\mathcal{J}^{*} \subset I^{*}}} P_{I^{*}}^{(+)} (V_{\overline{\mathcal{J}^{*}}}^{I_{-1}} - \langle V_{\overline{\mathcal{J}^{*}}}^{I_{-1}} \rangle_{\Psi_{I^{*}}^{A}}) P_{I^{*}}^{(+)} \right]^{j} P_{I^{*}}^{(+)}$$

which is well defined due to arguments as in the proof of Lemma 4.6 (see in particular (4.43)), and we can estimate

$$\left\| \frac{1}{H_{I^*}^0 - \langle H_{I^*}^0 \rangle_{\Psi_{I^*}^A} + i\delta_t} \sum_{j=1}^{\infty} \left[\left(-\sqrt{t} \sum_{\overline{\mathcal{J}^*} \subset I^*} P_{I^*}^{(+)} (V_{\overline{\mathcal{J}^*}}^{I_{-1}} - \langle V_{\overline{\mathcal{J}^*}}^{I_{-1}} \rangle_{\Psi_{I^*}^A}) P_{I^*}^{(+)} \right] \frac{1}{H_{I^*}^0 - \langle H_{I^*}^0 \rangle_{\Psi_{I^*}^A} + i\delta_t} \right]^j P_{I^*}^{(+)} \right|$$

$$\leq O(\sqrt{t}).$$

$$(4.121)$$

Hence we write

$$-P_{\overline{I^*}}^{(+)}(\sigma_{i^*_{-}-1}^z\sigma_{i^*_{-}}^z+1)\frac{1}{G_{I^*}-E_{I^*}^A+i\delta_t}P_{I^*}^{(+)}V_I^{\mathcal{I}_{-1}}P_{I^*}^{(-),A}P_{\overline{I^*}}^{(-)}$$
(4.122)

$$= -P_{\overline{I^*}}^{(+)} (\sigma_{i^*_{-}-1}^z \sigma_{i^*_{-}}^z + 1) \frac{1}{H^0_{I^*} - \langle H^0_{I^*} \rangle_{\Psi^A_{I^*}} + i\delta_t} P_{I^*}^{(+)} V_I^{\mathcal{I}_{-1}} P_{I^*}^{(-),A} P_{\overline{I^*}}^{(-)}$$
(4.123)

$$+R_2$$
, (4.124)

where $||R_2|| \le O(\sqrt{t} \cdot ||V_I^{I_{-1}}||).$

In expression (4.123), first we substitute $P_{I^*}^{(+)} = \mathbb{1} - P_{I^*}^{(-)}$, then we exploit that $V_I^{I_{-1}}$ is block diagonal w.r.t. $P_{I^*}^{(-),A/B}$ and write

$$-P_{\overline{I^*}}^{(+)}(\sigma_{i^*_{-}-1}^{z}\sigma_{i^*_{-}}^{z}+1)\frac{1}{H^0_{I^*}-\langle H^0_{I^*}\rangle_{\Psi^A_{I^*}}+i\delta_t}P_{I^*}^{(+)}V_I^{I_{-1}}P_{I^*}^{(-),A}P_{\overline{I^*}}^{(-)}$$
(4.125)

$$= -P_{\overline{I^*}}^{(+)} (\sigma_{i^*_{-}-1}^z \sigma_{i^*_{-}}^z + 1) \frac{1}{H^0_{I^*} - \langle H^0_{I^*} \rangle_{\Psi^A_{I^*}} + i\delta_t} V_I^{I_{-1}} P_{I^*}^{(-),A} P_{\overline{I^*}}^{(-)}$$
(4.126)

$$+P_{\overline{I^{*}}}^{(+)}(\sigma_{i^{*}_{-}-1}^{z}\sigma_{i^{*}_{-}}^{z}+1)\frac{1}{H^{0}_{I^{*}}-\langle H^{0}_{I^{*}}\rangle_{\Psi^{A}_{I^{*}}}+i\delta_{t}}P_{I^{*}}^{(-),A}V_{I}^{I_{-1}}P_{I^{*}}^{(-),A}P_{\overline{I^{*}}}^{(-),A}P_{\overline{I^{*}}}^{(-),A}$$
(4.127)

Next, we recall the following three identities:

1)

$$\frac{1}{H_{I^*}^0 - \langle H_{I^*}^0 \rangle_{\Psi_{I^*}^A} + i\delta_t} P_{I^*}^{(-),A} V_I^{I_{-1}} P_{I^*}^{(-),A} P_{\overline{I^*}}^{(-)} = \frac{1}{i\delta_t} P_{I^*}^{(-),A} V_I^{I_{-1}} P_{I^*}^{(-),A} P_{\overline{I^*}}^{(-)}$$
(4.128)

which holds since $(H^0_{I^*} - \langle H^0_{I^*} \rangle_{\Psi^A_{I^*}}) P^{(-),A}_{I^*} = 0;$ 2)

$$P_{\mathcal{I}^{*}}^{(-),A} V_{\mathcal{I}}^{\mathcal{I}_{-1}} P_{\mathcal{I}^{*}}^{(-),A} P_{\overline{\mathcal{I}^{*}}}^{(-)} = \langle V_{\mathcal{I}}^{\mathcal{I}_{-1}} \rangle_{\Psi_{\mathcal{I}^{*}}^{A}} P_{\overline{\mathcal{I}^{*}}}^{(-),A} P_{\overline{\mathcal{I}^{*}}}^{(-)} = \langle V_{\mathcal{I}}^{\mathcal{I}_{-1}} \rangle_{\Psi_{\mathcal{I}^{*}}^{A}} P_{\overline{\mathcal{I}^{*}}}^{(-),A}$$

$$(4.129)$$

which holds since $\overline{I^*}$ is an extension of I^* by two sites both on the left and on the right for bulk-intervals, and by two sites either on the left or on the right for boundary-intervals; 3)

$$(\sigma_{i_{-}^{*}-1}^{z}\sigma_{i_{-}^{*}}^{z}+1)P_{\overline{I^{*}}}^{(-)}=0, \qquad (4.130)$$

see Remark 4.8. From 1), 2), and 3) above we deduce

$$(4.127) = 0. (4.131)$$

Finally, we show how to control

$$-P_{\overline{I^*}}^{(+)}(\sigma_{i_-^{z-1}}^{z}\sigma_{i_-^{z}}^{z}+1)\frac{1}{H_{\overline{I^*}}^{0}-\langle H_{\overline{I^*}}^{0}\rangle_{\Psi_{\overline{I^*}}^{A}}+i\delta_{t}}V_{I}^{\overline{I}_{-1}}P_{\overline{I^*}}^{(-),A}P_{\overline{I^*}}^{(-)}.$$
(4.132)

We re-write

$$\frac{1}{H^{0}_{I^{*}} - \langle H^{0}_{I^{*}} \rangle_{\Psi^{A}_{I^{*}}} + i\delta_{t}} = -i \int_{0}^{t^{-\frac{1}{3}}} e^{i \left(H^{0}_{I^{*}} - \langle H^{0}_{I^{*}} \rangle_{\Psi^{A}_{I^{*}}} + i\delta_{t}\right)s} ds - i \int_{t^{-\frac{1}{3}}}^{+\infty} e^{i \left(H^{0}_{I^{*}} - \langle H^{0}_{I^{*}} \rangle_{\Psi^{A}_{I^{*}}} + i\delta_{t}\right)s} ds$$

$$(4.133)$$

and define

$$R_4 := i P_{\overline{I^*}}^{(+)} \left(\sigma_{i_-^*-1}^z \sigma_{i_-^*}^z + 1 \right) \int_{t^{-1/3}}^{+\infty} e^{i \left(H_{I^*}^0 - \langle H_{I^*}^0 \rangle_{\Psi_{I^*}}^A + i\delta_t \right) s} \, ds \, V_I^{\mathcal{I}_{-1}} \, P_{I^*}^{(-),A} \, P_{\overline{I^*}}^{(-)} \tag{4.134}$$

with $||R_4|| \le O(\frac{e^{-\delta_t \cdot t^{-\frac{1}{3}}}}{\delta_t} ||V_I^{I_{-1}}||)$. By using the identities in (4.133) and in (4.130), we write

$$-P_{\overline{I^*}}^{(+)}\left(\sigma_{i^*_{-}-1}^{z}\sigma_{i^*_{-}}^{z}+1\right)\frac{1}{H^0_{I^*}-\langle H^0_{I^*}\rangle_{\Psi^A_{I^*}}+i\delta_t}V_I^{\mathcal{I}_{-1}}P_{I^*}^{(-),A}P_{\overline{I^*}}^{(-)}-R_4$$
(4.135)

$$= i \cdot \int_{0}^{t^{-1/3}} P_{\overline{I^{*}}}^{(+)} \left(\sigma_{i_{-}^{*}-1}^{z} \sigma_{i_{-}^{*}}^{z} + 1 \right) e^{i \left(H_{I^{*}}^{0} - \langle H_{I^{*}}^{0} \rangle_{\Psi_{I^{*}}}^{A} + i\delta_{t} \right) s} V_{I}^{\mathcal{I}_{-1}} P_{\overline{I^{*}}}^{(-),A} ds$$

$$(4.136)$$

$$= i \cdot \int_{0}^{t^{-1/3}} P_{\overline{I^{*}}}^{(+)} (\sigma_{i_{-}^{z}1}^{z} \sigma_{i_{-}^{z}}^{z} + 1) e^{-\delta_{t} \cdot s} e^{i \left(H_{I^{*}}^{0} - \langle H_{I^{*}}^{0} \rangle_{\Psi_{I^{*}}}\right) s} V_{I}^{\overline{I}_{-1}} e^{-i \left(H_{I^{*}}^{0} - \langle H_{I^{*}}^{0} \rangle_{\Psi_{I^{*}}}\right) s} P_{\overline{I^{*}}}^{(-),A} ds$$

$$= i \cdot \int_{0}^{t^{-1/3}} P_{\overline{I^{*}}}^{(+)} e^{-\delta_{t} \cdot s} \left[(\sigma_{i_{-}^{z}1}^{z} \sigma_{i_{-}^{z}}^{z} + 1), e^{i H_{I^{*}}^{0} s} V_{I}^{\overline{I}_{-1}} e^{-i H_{I^{*}}^{0} s} \right] P_{\overline{I^{*}}}^{(-),A} ds.$$

$$(4.137)$$

At this point, we can make use of the Lieb-Robinson bound displayed in (4.105); hence we get that for *t* sufficiently small

$$\|(4.135)\| \leq \delta_t^{-1} \cdot \sup_{0 \leq s \leq t^{-1/3}} \left\| \left[(\sigma_{i_{-1}^s}^z \sigma_{i_{-}^s}^z + 1), e^{i \cdot H_{I^*}^0 \cdot s} V_I^{I_{-1}} e^{-i \cdot H_{I^*}^0 \cdot s} \right] \right\|$$
(4.138)

$$\leq t^{-\frac{1}{4}} \cdot \frac{4 \left\| (\sigma_{i_{-}^{*}}^{z} \sigma_{i_{-}^{*}}^{z} + 1) \right\| \|V_{I}^{I_{-1}}\| \|F_{0}\|}{C_{1}} \cdot e^{-\left[d(i_{-}^{*}, I) - 2 \|\Phi\|_{1}C_{1}t^{-1/3}\right]}$$
(4.139)

$$\leq e^{-\frac{\sqrt{I^{-1}}}{4}} \|V_{I}^{I_{-1}}\| \tag{4.140}$$

where $d(i_{-}^*, I) = \frac{\sqrt{t^{-1}}}{3}$, and C_1 , $\|\Phi\|_1$ and F_0 are positive constants defined in (4.104), (4.106), and (4.103), respectively, with *a* set equal to 1.

By collecting all the estimates the bound in (4.107) is proven.

Remark 4.8. Concerning the relation between a single Ising term and the ground-state vectors of the unperturbed Hamiltonian of the entire chain, we observe that in the estimate of the commutator in (4.107) we only use that they are eigenvectors of each Ising term.

Regarding the ferromagnetic case, in the analogous proof starting from (4.109) we must replace

$$\frac{\sigma_{i_{-}^{*}-1}^{z}\sigma_{i_{-}^{*}}^{z}}{t^{1/2}} + \frac{1}{t^{1/2}} \quad \text{with} \quad \frac{\sigma_{i_{-}^{*}-1}^{z}\sigma_{i_{-}^{*}}^{z}}{t^{1/2}} - \frac{1}{t^{1/2}};$$

the rest of the proof is indeed simpler due to the one-dimensionality of the ground-state subspace of $H^0_{T^*}$.

4.2.3 Main theorems

We are now ready to collect all the ingredients and prove our main results.

Theorem 4.9. There exists $\overline{t} > 0$ independent of N, such that for all $|t| < \overline{t}$, for any $\Im \ni \mathcal{L} \leq \Lambda_{-1}$, the Hamiltonians $G_{\mathcal{L}^*}$ are well defined, and the properties below hold true:

S1) for any interval $\mathcal{J} \in \mathfrak{I}$, the following operator norms estimates hold

$$\|V_{\mathcal{J}}^{\mathcal{L}}\| \le \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{\ell(\mathcal{J})^2}$$

for $\mathcal{J} > \mathcal{L}$;

S2) in the antiferromagnetic case, the spectrum of the operators $(G_{\mathcal{L}_{+1}^*} - E_{\mathcal{L}_{+1}^*}^{A/B})$ restricted to $P_{\mathcal{L}_{+1}^*}^{(+)} \mathcal{H}^{\Lambda}$ is bounded below by $\Delta_{\mathcal{L}_{+1}^*} \geq \frac{2|J|-2h}{2}$, where $G_{\mathcal{L}_{+1}^*}$ is defined in formula (3.5). Analogously, in the ferromagnetic case, the spectral gap of $G_{\mathcal{L}_{+1}^*}$ above the ground-state energy is bounded below by $\Delta_{\mathcal{L}_{+1}^*} \geq \frac{2J+2h}{2}$.

Proof

The proof is by induction in the interval I that labels the block-diagonalization step. Hence for each operator support $\mathcal{J} \in \mathfrak{I}$ we shall prove S1) and S2) from step $I = I_0$ up to step $I = \Lambda_{-1}$. That is we assume that S1) holds for all $V_{\mathcal{J}}^{\mathcal{K}}$ with $\mathcal{K} < I$ and S2) for all $\mathcal{K} < I$. Then we show that they hold for all $V_{\mathcal{J}}^{I}$ and for $G_{(I^*)_{+1}}$, respectively. This implies that K_{Λ}^{I} and (by Lemma 4.10) Z_{I^*} are well defined operators and, consequently, (3.30) holds true.

For $I = I_0$, S1) can be verified by direct computation, indeed $||V_{\mathcal{J}}^{I_0}|| \le 1$ for \mathcal{J} with $\ell(\mathcal{J}) = 1$ due to (3.16) and (2.20), and $||V_{\mathcal{J}}^{I_0}|| = 0$ (see (3.17)) for \mathcal{J} with $\ell(\mathcal{J}) > 1$; S2) holds trivially since, by definition, $G_{(I_0)_{+1}} = H_{(I_0)_{+1}}^0$ where $(I_0)_{+1}$ is the interval with $Q((I_0)_{+1}) = \ell((I_0)_{+1}) = 1$.

The induction step consists of several parts and for each of them we choose $t \ge 0$ in an interval such that the previous parts and Lemma 4.10 are verified. By this procedure we may progressively restrict such interval until we determine a $\overline{t} > 0$ for which all the parts hold true for $0 \le t < \overline{t}$.

Induction step in the proof of S1)

Starting from \mathcal{L} down to $(\mathcal{I}_0)_{+1}$, step by step, we relate the norm of $V_{\mathcal{J}}^I$ to the ones of the operators, in step \mathcal{I}_{-1} , in terms of which $V_{\mathcal{J}}^I$ is expressed according to the algorithm. It is then clear that, at fixed \mathcal{J} , for most of the steps the norm is preserved, i.e., $||V_{\mathcal{J}}^I|| = ||V_{\mathcal{J}}^{I-1}||$, and only for special steps we have nontrivial relations.

By the rules of the algorithm displayed in Definition 3.3, $V_{\mathcal{J}}^{\mathcal{L}}$ is defined only for $\mathcal{J} > \mathcal{L}$, and this is possible if $\ell(\mathcal{J}) > \ell(\mathcal{L})$ or if $\ell(\mathcal{J}) = \ell(\mathcal{L})$ and $Q(\mathcal{J}) > Q(\mathcal{L})$.

Case $\ell(\mathcal{J}) = 1$

In this case statement S1) holds since, due to a-1) of Definition 3.3, by repeated steps back we readily obtain

$$\|V_{\mathcal{J}}^{\mathcal{L}}\| = \|V_{\mathcal{J}}^{I_0}\| = \|V_{\mathcal{J}}\| \le 1.$$
(4.141)

Case $\ell(\mathcal{J}) \geq 2$

We recall that by construction $V_{\mathcal{J}}^{I}$ is defined only for $\ell(\mathcal{J}) \geq \ell(I)$. For the re-expansion from step I to I_{-1} , we distinguish the following situations related to cases a), b), and c) of Definition 3.3:

i) In case a-1), and in case c) with the additional requirement that $\widetilde{I^*}$ does not contain endpoints of \mathcal{J} , we have that

$$\|V_{\mathcal{J}}^{I}\| = \|V_{\mathcal{J}}^{I_{-1}}\|; \qquad (4.142)$$

indeed, in case a-1) the statement is trivial, while in case c) only the contribution in (3.23) is nonzero due to requirement that $\widetilde{I^*}$ does not contain endpoints of \mathcal{J} .

ii) In case c), if $\widetilde{I^*}$ contains only one of the endpoints of \mathcal{J} , the contributions are those in (3.23), (3.24), and (3.25), hence we can estimate

$$\|V_{\mathcal{J}}^{I}\| \leq \|V_{\mathcal{J}}^{I_{-1}}\| \tag{4.143}$$

$$+\sum_{\mathcal{K}\in[\mathcal{G}_{\mathcal{T}}^{I}]_{1}}\left\|\sum_{n=1}^{\infty}\frac{1}{n!}ad^{n}Z_{I^{*}}(V_{\mathcal{K}}^{I_{-1}})\right\|$$
(4.144)

$$+\sum_{\mathcal{K}^* \in [\mathcal{G}_{\mathcal{J}}^{\mathcal{I}}]_2} \left\| \sum_{n=1}^{\infty} \frac{1}{n!} a d^n Z_{\mathcal{I}^*}(V_{\overline{\mathcal{K}^*}}^{\mathcal{I}_{-1}}) \right\|$$
(4.145)

where for the convenience of the reader we recall that

$$[\mathcal{G}_{\mathcal{J}}^{I}]_{1} := \left\{ \mathcal{K} \in \mathfrak{I} \mid \mathcal{K} \succ I, \, \mathcal{K} \cap I^{*} \neq \emptyset, \qquad (4.146) \\ \mathcal{K} \neq \mathcal{J}, \, \text{and} \, \widetilde{I^{*}} \cup \mathcal{K} = \mathcal{J} \right\}$$

$$[\mathcal{G}_{\mathcal{J}}^{I}]_{2} := \left\{ \mathcal{K}^{*} \in \mathfrak{I}^{*} \mid I \succ \mathcal{K}, \, \mathcal{K}^{*} \cap I^{*} \neq \emptyset, \, \mathcal{K}^{*} \notin I^{*} \right.$$

and $\widetilde{I^{*}} \cup \widetilde{\mathcal{K}^{*}} = \mathcal{J} \left. \right\}.$

For the estimate of (4.144) + (4.145), we invoke the computations in (4.95)-(4.102) of Section 4.2.1, the inductive hypothesis S1), and we split our study into three parts corresponding to the following subcases (see Remark 2.7): ii-1) \mathcal{J} is a bulk-interval (hence also \mathcal{I} is a bulk-interval since $\mathcal{I}^* \subset \mathcal{J}$); ii-2) \mathcal{J} and \mathcal{I} are both boundary intervals; ii-3) \mathcal{J} is a boundary interval but \mathcal{I} is a bulk-interval.

We also use the symbol $C'_{J,h}$ for any positive constants that depend on J and h. They may change from line to line.

ii-1) If \mathcal{J} is a bulk-interval then $\ell(\mathcal{J}) - \ell(\mathcal{I}) \ge 3$ and $\ell(\mathcal{K}) \ge \ell(\mathcal{J}) - \ell(\mathcal{I}) - 2$ in the summation concerning (4.145). Hence, we can estimate (see (4.95)-(4.102))

$$(4.144) + (4.145) \leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{\ell(\mathcal{K}) = \ell(\mathcal{J}) - \ell(I) - 2}^{\ell(\mathcal{J}) - 1} \frac{t^{\frac{\ell(I) - 1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(\mathcal{K}) - 1}{16}}}{\ell(\mathcal{K})^2}$$
(4.147)

$$\leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{m=0}^{\ell(J)+1} \frac{t^{\frac{\ell(J)-1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(J)-\ell(J)+m-3}{16}}}{(\ell(J)-\ell(I)-2+m)^2}$$
(4.148)

$$= C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot t^{\frac{\ell(\mathcal{J})-4}{16}} \sum_{m=0}^{\ell(\mathcal{J})+1} \frac{t^{\frac{m}{16}}}{\ell(\mathcal{I})^2 \cdot (\ell(\mathcal{J}) - \ell(\mathcal{I}) - 2 + m)^2} (4.149)$$

$$\leq C'_{J,h} \cdot t^{\frac{5}{16}} \cdot \frac{t^{\frac{5}{16}}}{\ell(I)^2 \cdot (\ell(\mathcal{J}) - \ell(I) - 2)^2} \,. \tag{4.150}$$

ii-2) If \mathcal{J} is a boundary-interval but I is a bulk-interval then $\ell(\mathcal{J}) - \ell(I) \ge 3$ and $\ell(\mathcal{K}) \ge \ell(\mathcal{J}) - \ell(I) - 2$ in the summation concerning (4.145). In this case we get an analogous estimate

$$(4.144) + (4.145) \leq C'_{J,h} \cdot t^{\frac{5}{16}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{\ell(\mathcal{I})^2 \cdot (\ell(\mathcal{J}) - \ell(\mathcal{I}) - 2)^2}.$$
(4.151)

ii-3) If both \mathcal{J} and \mathcal{I} are boundary-intervals then $\ell(\mathcal{J}) - \ell(\mathcal{I}) \ge 2$ and $\ell(\mathcal{K}) \ge \ell(\mathcal{J}) - \ell(\mathcal{I}) - 1$ in the summation concerning (4.145). In this case we estimate

$$(4.144) + (4.145) \leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{\ell(\mathcal{K})=\ell(\mathcal{J})-\ell(I)-1}^{\ell(\mathcal{J})-1} \frac{t^{\frac{\ell(I)-1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(\mathcal{K})-1}{16}}}{\ell(\mathcal{K})^2}$$
(4.152)

$$\leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{m=0}^{\ell(I)} \frac{t^{\frac{\ell(J)-1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(J)-\ell(I)+m-2}{16}}}{(\ell(J)-\ell(I)-1+m)^2}$$
(4.153)

$$= C'_{J,h} \cdot C \cdot A_{1} \cdot \sqrt{t} \cdot t^{\frac{\ell(\mathcal{J})-3}{16}} \sum_{m=0}^{\ell(I)} \frac{t^{\frac{m}{16}}}{\ell(I)^{2} \cdot (\ell(\mathcal{J}) - \ell(I) - 1 + m)^{2}} (4.154)$$

$$\leq C'_{J,h} \cdot t^{\frac{3}{8}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{\ell(I)^{2} \cdot (\ell(\mathcal{J}) - \ell(I) - 1)^{2}} .$$

$$(4.155)$$

iii) In case c), if $\widetilde{I^*}$ contains both the two endpoints of \mathcal{J} , i.e., $\widetilde{I^*} = \mathcal{J}$, the possible contributions are all those in (3.23)-(3.29) of Definition 3.3, and we can estimate

$$\|V_{\mathcal{J}}^{I}\| \leq \|V_{\mathcal{J}}^{I_{-1}}\| \tag{4.156}$$

$$+ \|(3.24)\| + \|(3.25)\| \tag{4.157}$$

$$+ \|(3.26)\| + \|(3.27)\| + \|(3.28)\| + \|(3.29)\|.$$
(4.158)

Notice that the terms in (4.157) are those in (4.144) and (4.145). Concerning (4.158), together with the inductive hypothesis S1) we invoke:

a) either (4.182) or (4.183) (depending on the sign of J) and (4.185) in Lemma 4.10 for the estimate of ||(3.26)||, ||(3.27)||, and ||(3.28)||;

b) Lemma 4.7 for the estimate of ||(3.29)||.

In order to study the terms above we split the study into three parts.

iii-1) If \mathcal{J} is a bulk-interval then $\ell(\mathcal{J}) - \ell(\mathcal{I}) = 2$, the contribution of (4.145) is absent (i.e., the corresponding set $[\mathcal{G}_{\mathcal{I}}^{\mathcal{I}}]_2$ is empty), and $\ell(\mathcal{K}) \ge \ell(\mathcal{J}) - \ell(\mathcal{I}) - 1$ in the sum

in (4.144). Hence, we estimate

$$(4.157) + (4.158) \tag{4.159}$$

$$\leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{\ell(\mathcal{K}) = \ell(\mathcal{J}) - \ell(I) - 1}^{\ell(\mathcal{J}) - 1} \frac{t^{\frac{\ell(I) - 1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(\mathcal{K}) - 1}{16}}}{\ell(\mathcal{K})^2}$$
(4.160)

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-3}{16}}}{(\ell(\mathcal{J})-2)^2}$$
(4.161)

$$\leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{m=0}^{\ell(I)} \frac{t^{\frac{\ell(J)-1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(J)-\ell(J)+m-2}{16}}}{(\ell(\mathcal{J})-\ell(I)-1+m)^2}$$
(4.162)

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{1}{(J-J)}}}{(\ell(\mathcal{J})-2)^2}$$
(4.163)

$$= C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot t^{\frac{\ell(\mathcal{J}) - 3}{16}} \sum_{m=0}^{\ell(\mathcal{I})} \frac{t^{\frac{m}{16}}}{\ell(\mathcal{I})^2 \cdot (\ell(\mathcal{J}) - \ell(\mathcal{I}) - 1 + m)^2}$$
(4.164)

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(J)-3}{16}}}{(\ell(\mathcal{J})-2)^2}$$
(4.165)

$$\leq C'_{J,h} \cdot t^{\frac{3}{8}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{\ell(\mathcal{I})^2 \cdot (\ell(\mathcal{J}) - \ell(\mathcal{I}) - 1)^2} + C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-3}{16}}}{(\ell(\mathcal{J}) - 2)^2}$$
(4.166)

where for the origin of the summand $C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-3}{16}}}{(\ell(\mathcal{J})-2)^2}$ see the explanations in a) and b) above.

iii-2) If both \mathcal{J} and \mathcal{I} are boundary-intervals then $\ell(\mathcal{J}) - \ell(\mathcal{I}) = 1$, the contribution of (4.145) is absent and $\ell(\mathcal{K}) \ge \ell(\mathcal{J}) - \ell(\mathcal{I})$ in the sum in (4.144). Hence, similarly to subcase iii-1), we can estimate

$$\|(4.158)\| \leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{\ell(\mathcal{K})=\ell(\mathcal{J})-\ell(I)}^{\ell(\mathcal{J})-1} \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{\ell(I)^2} \cdot \frac{t^{\frac{\ell(\mathcal{K})-1}{16}}}{\ell(\mathcal{K})^2}$$
(4.167)

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-2}{16}}}{(\ell(\mathcal{J})-1)^2}$$
(4.168)

$$\leq C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot \sum_{\substack{m=0\\\ell(\mathcal{J})-2}}^{\ell(\mathcal{I})-1} \frac{t^{\frac{\ell(\mathcal{I})-1}{16}}}{\ell(\mathcal{I})^2} \cdot \frac{t^{\frac{\ell(\mathcal{J})-\ell(\mathcal{I})+m-1}{16}}}{(\ell(\mathcal{J})-\ell(\mathcal{I})+m)^2}$$
(4.169)

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{(J)^{-2}}{16}}}{(\ell(\mathcal{J}) - 1)^2}$$
(4.170)

$$= C'_{J,h} \cdot C \cdot A_1 \cdot \sqrt{t} \cdot t^{\frac{\ell(\mathcal{J})-2}{16}} \sum_{m=0}^{\ell(\mathcal{I})-1} \frac{t^{\frac{m}{16}}}{\ell(\mathcal{I})^2 \cdot (\ell(\mathcal{J}) - \ell(\mathcal{I}) + m)^2} (4.171)$$

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{-16}{16}}}{(\ell(\mathcal{J}) - 1)^2}$$
(4.172)

$$\leq C'_{J,h} \cdot t^{\frac{7}{16}} \cdot \frac{t^{\frac{\ell(J)^{-1}}{16}}}{\ell(I)^2 \cdot (\ell(\mathcal{J}) - \ell(I))^2} + C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(J)^{-2}}{16}}}{(\ell(\mathcal{J}) - 1)^2}$$
(4.173)

iii-3) If \mathcal{J} is a boundary-interval but I is a bulk-interval then $\ell(\mathcal{J}) - \ell(I) = 2$ and the contribution of (4.145) is not absent and in the corresponding sum we have $\ell(\mathcal{K}) \ge 1$

 $\ell(\mathcal{J}) - \ell(\mathcal{I}) - 1$. Thus we estimate

$$\|(4.158)\| \leq C'_{J,h} \cdot t^{\frac{3}{8}} \cdot \frac{t^{\frac{\ell(J)-1}{16}}}{\ell(I)^2 \cdot (\ell(J) - \ell(I) - 1)^2}$$
(4.174)

$$+C'_{J,h} \cdot t^{\frac{1}{4}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-3}{16}}}{(\ell(\mathcal{J})-2)^2} \,. \tag{4.175}$$

At fixed \mathcal{J} and fixed $\ell(\mathcal{I})$, the situation described in ii) happens only three times at most, namely it happens three times when \mathcal{J} is a boundary-interval. At fixed \mathcal{J} , the situation described in iii) happens only for two intervals \mathcal{I} at most, namely it happens twice when \mathcal{J} is a boundary-interval. Hence, for fixed \mathcal{L} , by re-expanding back down to level \mathcal{I}_0 we can estimate for *t* sufficiently small

$$\|V_{\mathcal{J}}^{\mathcal{L}}\| \leq \|V_{\mathcal{J}}^{\mathcal{I}_0}\| \tag{4.176}$$

$$+\mathbb{1}(\ell(\mathcal{J})-2)\sum_{K=1}^{\ell(\mathcal{J})-1} 3 \cdot C'_{J,h} \cdot t^{\frac{7}{16}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{K^2 \cdot (\ell(\mathcal{J})-K)^2}$$
(4.177)

$$+\mathbb{1}(\ell(\mathcal{J})-3)\sum_{K=1}^{\ell(\mathcal{J})-1} 3 \cdot C'_{J,h} \cdot t^{\frac{3}{8}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{K^2 \cdot (\ell(\mathcal{J})-K-1)^2}$$
(4.178)

$$+\mathbb{1}(\ell(\mathcal{J})-4)\sum_{K=1}^{\ell(\mathcal{J})-1} 3 \cdot C'_{J,h} \cdot t^{\frac{5}{16}} \cdot \frac{t^{\frac{\ell(\mathcal{J})-1}{16}}}{K^2 \cdot (\ell(\mathcal{J})-K-2)^2}$$
(4.179)

$$+C'_{J,h} \cdot t^{\frac{1}{8}} \cdot \frac{t^{\frac{1}{(U)-1}}}{(\ell(\mathcal{J})-1)^2}$$
(4.180)

$$\leq \frac{t^{\frac{\ell(f)-1}{16}}}{\ell(f)^2} \tag{4.181}$$

where $\mathbb{1}(z)$ is the characteristic function of $[0, +\infty]$ and in the last step we have used that $\|V_{\mathcal{T}}^{\mathcal{I}_0}\| = 0$ for $\ell(\mathcal{J}) > 1$.

Induction step in the proof of S2)

The statement follows from Lemmata 4.1 and 4.6 where we assume S2) in step \mathcal{L}_{-1} and exploit the result just proven for S1) in step \mathcal{L} . \Box

In the next lemma, which is part of the induction, we (also) estimate the operator norm of a potential after its block-diagonalization. This estimate applies to the subsequent steps as well, since by construction the block-diagonalized potential does not change along the flow; see the algorithm in Definition 3.3.

Lemma 4.10. Assume that t > 0 is sufficiently small independently of N, and such that S1) and S2) of Theorem 4.9 hold true in step I_{-1} . Then the inequalities

(a)

$$||Z_{I^*}|| \le A_1 \cdot \frac{\sqrt{t}}{J+h} \cdot ||V_I^{I_{-1}}||, \qquad (4.182)$$

in the ferromagnetic case,

(b)

$$||Z_{I^*}|| \le A_1 \cdot \frac{\sqrt{t}}{|J| - h} \cdot ||V_I^{I_{-1}}||, \qquad (4.183)$$

in the antiferromagnetic case,



Figure 5: On the left, we display the relation between Theorem 4.9 and Lemma 4.10 in the inductive step of the block-diagonalization procedure. On the right, we show how Lemma 4.10 implies *S2*) in Theorem 4.9 by means of Lemmata 4.1 and 4.6.

and

$$\|V_{\overline{I^*}}^{I}\| \le C_{J,h} \cdot \|V_{\overline{I}}^{I_{-1}}\|, \qquad (4.184)$$

$$\sum_{j=2}^{\infty} t^{\frac{j-1}{2}} \| (V_{I^*}^{\mathcal{I}_{-1}})_j^{diag} \| \le \frac{C}{|J| \pm h} \cdot \sqrt{t} \cdot \| V_{I}^{\mathcal{I}_{-1}} \|,$$
(4.185)

hold true where \pm are referred to the ferromagnetic and the antiferromagnetic case, respectively, and

$$C_{J,h} := 1 + 2 \cdot \frac{A_1}{J+h}$$

in the ferromagnetic case,

$$C_{J,h} := 1 + 2 \cdot \frac{A_1}{|J| - h}$$

in the antiferromagnetic case; A₁ and C are universal constants.

Proof

Concerning (4.182), (4.183), and (4.185) the argument is the same as in [FP, Lemma A.3]. The inequality in (4.184) can be obtained directly from (3.20)-(3.21). \Box

We can now prove the main result of the paper.

Theorem 4.11.

(a) If J > 0, there exists a $\bar{t} > 0$ dependent on J and h, but independent of $N > \frac{J}{h} + \frac{3}{2}$ such that for all $|t| < \bar{t}$ the ground-state energy E_{Λ} of the Hamiltonian K_{Λ} in (1.3) is non-degenerate and the spectral gap is bounded below by $2J + 2h - O(\sqrt{|t|})$.

(b) If J < 0, for |J| > 2h there exists a $\overline{t} > 0$ dependent on J and h, but independent of N such that for all $|t| < \overline{t}$:

- if Λ has an odd number of sites, the set $\mathfrak{S} := \sigma(K_{\Lambda}) \cap [E_{\Lambda}, E_{\Lambda} + 2|J| - O(\sqrt{|t|})]$, where $\sigma(K_{\Lambda})$ is the spectrum of K_{Λ} and E_{Λ} its ground-state energy, consists of two points, E_{Λ} and E'_{Λ} , with $E'_{\Lambda} - E_{\Lambda} = 2h - O(\sqrt{|t|})$, and the spectral projection associated with \mathfrak{S} is of rank 2:

- if Λ has an even number of sites, the set $\mathfrak{S} := \sigma(K_{\Lambda}) \cap [E_{\Lambda}, E_{\Lambda} + 2|J| - 2h - O(\sqrt{|t|})]$ consists of at most two points, E_{Λ} and E'_{Λ} , with $|E'_{\Lambda} - E_{\Lambda}| \leq O(\sqrt{|t|})$, and the spectral projection associated with \mathfrak{S} is of rank 2.

Proof

By means of the conjugation $e^{Z_{\Lambda_{-1}}}$, we get the transformed Hamiltonian

$$e^{Z_{\Lambda_{-1}}} K_{\Lambda}(t) e^{-Z_{\Lambda_{-1}}} = G_{\Lambda} + \sqrt{t} V_{\Lambda}^{\Lambda_{-1}}.$$
(4.186)

Next, we implement a standard Lie-Schwinger block-diagonalization, w.r.t. to the pair of projections $P_{\Lambda}^{(-)}, P_{\Lambda}^{(+)}$; hence we have that $K_{\Lambda}(t)$ is unitarily equivalent to

$$\check{K}_{\Lambda}(t) := G_{\Lambda} + \sqrt{t} \check{V}_{\Lambda}^{\Lambda},$$

where $\check{V}^{\Lambda}_{\Lambda}$ is block diagonal w.r.t. $P^{(-)}_{\Lambda}, P^{(+)}_{\Lambda}$. By using the results of Theorem 4.9 (combined with Lemma 4.10), item (a) follows from the argument leading to Lemma 4.1 by also including the term $\sqrt{t}\check{V}^{\Lambda}_{\Lambda}$. Similarly, item (b) follows from the arguments leading to Lemmata 4.4 and 4.6 by also including the term $\sqrt{t}\check{V}^{\Lambda}_{\Lambda}$, which, despite not being block-diagonal w.r.t. $P^{(-),A}_{\Lambda}$, $P^{(-),B}_{\Lambda}$, is easily controlled since its norm is estimated much smaller than \sqrt{t} by the procedure of Lemma 4.10. In order to get the result in item (b) when Λ has an odd number of sites, we also exploit that with regard to $E_{I^*}^A$ the statement of Lemma 4.6 can be replaced by $P_{I_{+1}^*}^{(+)}(G_{I_{+1}^*} - E_{I_{+1}^*}^A)P_{I_{+1}^*}^{(+)} \ge 2|J| \cdot \left[1 - O(\sqrt{t})\right]P_{I_{+1}^*}^{(+)}$ as it is evident from the details of the proof. \Box

Remark 4.12. We can treat the ferromagnetic XXZ model without magnetic field similarly to the antiferromagnetic one, in the sense that we use the algorithm in Definition 3.3 and define for any \mathcal{J} the two spectral projections $P_{\overline{\mathcal{J}^*}}^{(-)A'}$ and $P_{\overline{\mathcal{J}^*}}^{(-)B'}$ which, in this case, are associated with the two vectors

$$\Psi_{\overline{\mathcal{J}^*}}^{A'} := |\uparrow\uparrow \cdots \uparrow\rangle \quad \text{and} \quad \Psi_{\overline{\mathcal{J}^*}}^{B'} := |\downarrow\downarrow \cdots \downarrow\rangle,$$

respectively. As for the antiferromagnetic case, we observe that the block-diagonalized potentials (defined in b) of Definition 3.3) are also block-diagonal w.r.t. to $P_{\overline{\mathcal{J}^*}}^{(-)A'}$ and $P_{\overline{\mathcal{J}^*}}^{(-)B'}$. Furthermore, the two energy levels $E_{I_{+1}^*}^{A'}$ and $E_{I_{+1}^*}^{B'}$ of the Hamiltonian $G_{I_{+1}^*}$ (see Section 4.1.1), corresponding to the two eigenvectors $\Psi_{I_{+1}^*}^{A'}$ and $\Psi_{I_{+1}^*}^{B'}$, are defined as in (3.12) and (3.13). Next, we fix a perpendicular direction and consider a rotation by π around it of the spin variables, for each site of the chain. Since all local terms of K_{Λ} are invariant under such rotation, and since $\Psi_{I_{+1}^*}^{A'}$ and $\Psi_{I_{+1}^*}^{B'}$ are mapped one to each other, we deduce that $E_{I_{+1}^*}^{A'}$ and $E_{I_{+1}^*}^{B'}$ coincide. Hence the statement on the spectral gap follows by essentially the same procedure used in Lemma 4.6, and the control of the block-diagonalization flow can be carried out as shown in this section.

References

- [AKLT] I. Affleck, T. Kennedy, E.H. Lieb, and H. Tasaki *Rigorous results on valence-bond* ground states in antiferromagnets Physical Review Letters. 59(7): 799-802 (1987).
- [BN] S. Bachmann, B. Nachtergaele. *On gapped phases with a continuous symmetry and boundary operators* J. Stat. Phys. 154(1-2): 91-112 (2014)
- [BH] S. Bravyi and M.B. Hastings. *A short proof of stability of topological order under local perturbations* Commun. Math. Physics. **307**, 609 (2011).
- [BHM] S. Bravyi, M.B. Hastings and S. Michalakis. *Topological quantum order: stability under local perturbations* J. Math. Phys. **51**, 093512, (2010).
- [DFFR] N. Datta, R. Fernandez, J. Fröhlich, L. Rey-Bellet. Low-Temperature Phase Diagrams of Quantum Lattice Systems. II. Convergent Perturbation Expansions and Stability in Systems with Infinite Degeneracy Helvetica Physica Acta 69, 752–820 (1996)
- [DFPRa] S. Del Vecchio, J. Fröhlich, A. Pizzo, A. Ranallo. *Boundary effects and the stability* of the low energy spectrum of the AKLT model (in preparation).
- [DFPR1] S. Del Vecchio, J. Fröhlich, A. Pizzo, S. Rossi. Lie-Schwinger block-diagonalization and gapped quantum chains with unbounded interactions Comm. Math. Phys. https://link.springer.com/article/10.1007/s00220-020-03878-y
- [DFPR2] S. Del Vecchio, J. Fröhlich, A. S. Rossi. Lie-Schwinger Pizzo, block-diagonalization chains: analyticand gapped quantum ity of the ground-state V. Issue 8 energy J. Funct. Anal. 279 https://www.sciencedirect.com/science/article/abs/pii/S0022123620302469
- [DFPR3] S. Del Vecchio, J. Fröhlich, A. Pizzo, S. Rossi. *Local* iterative block-diagonalization of gapped Hamiltonians: a new tool in singular perturbation theory J. Math. Phys. 63 (2022) 073503, https://doi.org/10.1063/5.0084552.
- [DFP] S. Del Vecchio, J. Fröhlich, A. Pizzo. Block-diagonalization of infinite-volume lattice Hamiltonians with unbounded interactions J. Funct. Anal. V. 284, Issue 1, 1 January 2023, 109734
- [DS] W. De Roeck, M. Salmhofer. Persistence of Exponential Decay and Spectral Gaps for Interacting Fermions Comm. Math. Phys. https://doi.org/10.1007/s00220-018-3211z
- [FS] C. Fishbacher, G. Stoltz The infinite XXZ quantum spin chain revisited: structure of low lying spectral bands and gaps Math. Model. Nat. Phenom. Vol. 9, No. 5, 2014, pp. 44-72
- [F] F. Franchini. An Introduction to Integrable Techniques for One-Dimensional Quantum Systems Lecture Notes in Physics (LNP, volume 940)
- [FP] J. Fröhlich, A. Pizzo. *Lie-Schwinger block-diagonalization and gapped quantum chains* Comm. Math. Phys. 375, 2039-2069 (2020)
- [H] M.B. Hastings. *The Stability of Free Fermi Hamiltonians* J. Math. Phys. 60, 042201 (2019)
- [K] T. Koma. Stability of the Spectral Gap for Lattice Fermions arXiv:2005.04548
- [KN] T. Koma, B. Nachtergaele. *The Spectral Gap of the Ferromagnetic XXZ-Chain* Letters in Mathematical Physics (40), 1-16 (1997)

- [KNS] T. Koma, B. Nachtergaele., S. Starr. The Spectral Gap for the Ferromagnetic Spin-J XXZ Chain Adv. Theor. Math. Phys, 5, 1047-1090 (2001)
- [KT] T. Kennedy, H. Tasaki. *Hidden symmetry breaking and the Haldane phase in* S = 1 *quantum spin chains* Comm.. Math. Phys. 147, 431-484 (1992)
- [LMY] A. Lucia, A. Moon, A. Young. Stability of the spectral gap and grounds state indistinguishability for a decorated AKLT model. preprint arXiv:2209.01141
- [LR] E. H. Lieb, D. W. Robinson. *The finite group velocity of quantum spin systems* Communications in Mathematical Physics. 28 (3) 251-257 (1972)
- [MN] A. Moon, B. Nachtergaele. *Stability of Gapped Ground State Phases of Spins and Fermions in One Dimension* J. Math. Phys. 59, 091415 (2018)
- [NS] B. Nachtergaele, R. Sims. Locality Estimates for Quantum Spin Systems in New Trends in Mathematical Physics 591-614 (2009). Spinger Netherlands.
- [NSY] B. Nachtergaele, R. Sims, A. Young. *Lieb-Robinson bounds, the spectral flow, and stability of the spectral gap for lattice fermion systems* Mathematical Problems in Quantum Physics, pp.93-115
- [NSY2] B. Nachtergaele, R. Sims, A. Young. *Stability of the bulk gap for frustration-free topologically ordered quantum lattice systems* preprint arXiv:2102.07209
- [NSY3] B. Nachtergaele, R. Sims, A. Young. Quasi-Locality Bounds for Quantum Lattice Systems. Part II. Perturbations of Frustration-Free Spin Models with Gapped Ground States Ann. Henri Poincaré, 23, 393-511 (2022)
- [NSt] B. Nachtergaele, S. Starr Droplet States in the XXZ Heisenberg Chain Comm. Math. Phys. 218 (2001), 569-607
- [NSS] B. Nachtergaele, W. Spitzer, and S. Starr *Droplet excitations for the spin-1/2 XXZ chain with kink boundary conditions* Ann. Henri Poincaré 8 (2007), 165-201
- [O] Y. Ogata. An $H^3(G, \mathbb{T})$ -valued index of symmetry-protected topological phases with on-site finite group symmetry for two-dimensional quantum spin systems Forum of Mathematics, Pi, Vol. 9:e13 1-62 (2021)
- [RW] Rakov, Mykhailo V., Weyrauch Michael. *Spin-*¹/₂ *XXZ Heisenberg chain in a longitudinal magnetic field* Physical Review B 100.13 (2019): 134434
- [R] S. B. Rutkevich *Kink confinement in the antiferromagnetic XXZ spin-(1/2) chain in a weak staggered magnetic field* Europhysics Letters, Volume 121, Number 3
- [S] N. Sopenko. An index for two-dimensional SPT states J. Math. Phys. 62, 111901 (2021)