

TOWARDS A THEORY OF NATURAL DIRECTED PATHS

PHILIPPE GAUCHER

ABSTRACT. We introduce the abstract setting of presheaf category on a thick category of cubes. Precubical sets, symmetric transverse sets, symmetric precubical sets and the new category of (non-symmetric) transverse sets are examples of this structure. All these presheaf categories share the same metric and homotopical properties from a directed homotopy point of view. This enables us to extend Raussen’s notion of natural d -path for each of them. Finally, we adapt Ziemiański’s notion of cube chain to this abstract setting and we prove that it has the expected behavior on precubical sets. As an application, we verify that the formalization of the parallel composition with synchronization of process algebra using the coskeleton functor of the category of symmetric transverse sets has a category of cube chains with the correct homotopy type.

CONTENTS

1. Introduction	1
2. Thick category of cubes	2
3. Metric and homotopical study	7
4. Ziemiański’s cube chain	10
5. Application	14
References	15

1. INTRODUCTION

Precubical sets are *de facto* the standard geometric model for directed homotopy for concurrency [2]. In fact, most of them are even non-positively curved in the sense of [12, Definition 1.28 and Proposition 1.29], or in the worst case proper in the sense of [19, page 499]. The motivation for introducing symmetric transverse sets in [6] is to formalize the parallel product with synchronization for process algebra using the associated coskeleton functor [6, Theorem 4.1.8]. Indeed, it is impossible to use the coskeleton functor associated with the category of precubical sets because of its pathological behavior (see [5, Proposition 3.15] and [6, Definition 3.1.3]). However, precubical sets still remain sufficient to model this parallel product by tweaking the coskeleton functor of this category (see [5, Section 3.3]).

Symmetric transverse sets share with precubical sets similar metric and homotopical properties by [9]. Indeed, their geometric realization carries a Lawvere metric structure which enables us to extend Raussen’s notion of (tame) natural d -path originally defined for precubical sets [17, Definition 2.14] [19, Definition 5.3] [20, Section 2.9]. Moreover, the

2020 *Mathematics Subject Classification.* 55U35,68Q85.

Key words and phrases. precubical set, directed path, presheaf, Reedy category, Lawvere metric space, process algebra.

full subcategory of representable objects of this presheaf category is c-Reedy in the sense of [18, Definition 8.25], like the box category (see Definition 2.3) for the precubical sets (the box category is even direct Reedy in the sense of [14, Definition 15.1.2]). This makes possible to compare in [9, Theorem 7.4] the natural realization of a symmetric transverse set with other realization functors and to generalize homotopical results proved in [4] and [10] for precubical sets.

The technical contribution of this note is threefold. Firstly, we explain why precubical sets and symmetric transverse sets belong to a larger family of presheaf categories on a *thick category of cubes* (see Definition 2.10). This family of presheaf categories contains also the symmetric precubical sets of [13] and a new category of non-symmetric transverse sets. Symmetric transverse sets are presheaves on a thick category of cubes which turns out to be, for tautological reasons, the greatest one for the inclusion.

Theorem. (*Proposition 2.13, Theorem 2.16 and Theorem 2.17*) *There exists a greatest thick category of cubes for the inclusion not containing the symmetry maps.*

Secondly, we prove that all results of [9] are valid for all presheaf categories on a thick category of cubes.

Theorem. (*Section 3 and more specifically Theorem 3.14*) *All metric and homotopical results of [9] are valid for the category $\mathcal{A}^{op}\mathbf{Set}$ of \mathcal{A} -sets when \mathcal{A} is a thick category of cubes.*

Thirdly, we obtain a statement which coincides with (a part of) [20, Theorem 7.5] when \mathcal{A} is the box category used to define the precubical sets:

Theorem. (*Corollary 4.8*) *Let K be a precubical set. Let \mathcal{A} be a thick category of cubes. The space of tame natural d -paths of the free \mathcal{A} -set $\mathcal{L}_{\mathcal{A}}(K)$ generated by K is homotopy equivalent to the classifying space of the small category of Ziemiański's cube chains of the free \mathcal{A} -set $\mathcal{L}_{\mathcal{A}}(K)$ generated by K .*

This leads to the following application:

Theorem. (*Section 5*) *The formalization of the parallel composition with synchronisation of [6] using the coskeleton functor of the category of symmetric transverse sets has a category of cube chains which gives the correct space of tame natural d -paths up to homotopy.*

The formal setting of presheaf category on a thick category of cubes is a first step towards an axiomatization of the notion of tame natural d -path. The next step would be to find a way of taking into account the globular version of this notion as it is introduced in [11]. This could lead to a general framework unifying all geometric approaches of directed homotopy for concurrency ¹.

2. THICK CATEGORY OF CUBES

2.1. Definition. A small category enriched over $([0, \infty], \geq, +, 0)$ [15] is called a *Lawvere metric space*. The category of Lawvere metric spaces is denoted by **LvMet**.

¹The expression “directed homotopy” has several quite distinct meanings. It is the reason why I add “for concurrency” on purpose.

A *Lawvere metric space* (X, d) is a set X equipped with a map $d : X \times X \rightarrow [0, \infty]$ called a *Lawvere metric* such that:

- $\forall x \in X, d(x, x) = 0$
- $\forall (x, y, z) \in X \times X \times X, d(x, y) \leq d(x, z) + d(z, y)$.

A map $f : (X, d) \rightarrow (Y, d)$ of Lawvere metric spaces is a set map $f : X \rightarrow Y$ which is *non-expansive*, i.e. $\forall (x, y) \in X \times X, d(f(x), f(y)) \leq d(x, y)$.

2.2. Notation. The category of partially ordered sets or posets together with the *strictly increasing maps* is denoted by \mathbf{PoSet}^+ .

Let $[0] = \{()\}$ and $[n] = \{0 < 1\}^n$ for $n \geq 1$ equipped with the product order. By convention, one has $\{0 < 1\}^0 = [0] = \{()\}$. In the sequel, for all $n \geq 1$, both the sets $[n]$ and $[0, 1]^n$ are equipped with the product order. By convention, $[0, 1]^0$ is a singleton. Let $\delta_i^\alpha : [n-1] \rightarrow [n]$ be the *coface map* defined for $1 \leq i \leq n$ and $\alpha \in \{0, 1\}$ by

$$\delta_i^\alpha(x_1, \dots, x_{n-1}) = (x_1, \dots, x_{i-1}, \alpha, x_i, \dots, x_{n-1}).$$

2.3. Definition. The *box category* \square is the subcategory of \mathbf{PoSet}^+ generated by the coface maps δ_i^α .

Let $x = (x_1, \dots, x_n)$ and $x' = (x'_1, \dots, x'_n)$ be two elements of $[0, 1]^n$ with $n \geq 1$. Let $\vec{d}_1 : [0, 1]^n \times [0, 1]^n \rightarrow [0, \infty]$ be the set map defined by

$$\vec{d}_1(x, x') = \begin{cases} \sum_{i=1}^n |x_i - x'_i| & \text{if } x \leq x' \\ \infty & \text{otherwise.} \end{cases}$$

Let $n \geq 0$. The set map $\vec{d}_1 : [0, 1]^n \times [0, 1]^n \rightarrow [0, \infty]$ is a Lawvere metric by [9, Proposition 1.5]. It restricts to a Lawvere metric on $\{0 < 1\}^n$.

2.4. Definition. [6, Definition 2.1.5] A map $f : [m] \rightarrow [n]$ of \mathbf{PoSet}^+ is *cotransverse* if

$$\text{For all } x, y \in [m], \vec{d}_1(x, y) = 1 \text{ implies } \vec{d}_1(f(x), f(y)) = 1.$$

Denote by $\widehat{\square}_S$ the subcategory of \mathbf{PoSet}^+ generated by the cotransverse maps.

A *cotransverse degeneracy map* is a cotransverse map $[n] \rightarrow [n]$ for $n \geq 2$ which is not one-to-one. Proposition 2.5 and Proposition 2.6 are important for the sequel.

2.5. Proposition. [6, Proposition 3.1.14] *Let $0 \leq m \leq n$. Every cotransverse map $f : [m] \rightarrow [n]$ factors uniquely as a composite $[m] \xrightarrow{\psi} [m] \xrightarrow{\phi} [n]$ with $\phi \in \square$ and ψ cotransverse.*

2.6. Proposition. *Let $f : [m] \rightarrow [n]$ be a cotransverse map. Let $\delta : [n] \rightarrow [p]$ be a map of \square . Suppose that $\delta f \in \square$. Then $f \in \square$.*

Proof. There exists a set map $s : [p] \rightarrow [n]$ obtained by removing some coordinates such that $s\delta = \text{Id}_{[n]}$. We deduce that $f = s(\delta f)$. From $\delta f \in \square$, we then deduce that $f \in \square$. \square

Let $\sigma_i : [n] \rightarrow [n]$ be the set map defined for $1 \leq i \leq n-1$ and $n \geq 2$ by

$$\sigma_i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, x_{i+1}, x_i, x_{i+2}, \dots, x_n).$$

These maps are called the *symmetry maps* [13]. The symmetry maps are clearly cotransverse. This is the reason of the S in the notation $\widehat{\square}_S$.

2.7. Notation. The subcategory of \mathbf{PoSet}^+ generated by the coface maps and the symmetry maps is denoted by \square_S .

2.8. Definition. [6, Definition 2.1.7 and Definition 2.1.12] A *category of cubes* is a small category \mathcal{A} satisfying the inclusions

$$\square \subset \mathcal{A} \subset \widehat{\square}_S.$$

A presheaf on \mathcal{A} is called an \mathcal{A} -set. The category of \mathcal{A} -sets is denoted by $\mathcal{A}^{op}\mathbf{Set}$.

The category \square , \square_S and $\widehat{\square}_S$ are examples of categories of cubes. The \square -sets are the *precubical sets* [1]. The \square_S -sets are the *symmetric precubical sets* (see [13]). The $\widehat{\square}_S$ -sets are the *symmetric transverse sets* introduced in [9].

The inclusion of small categories $j_{\mathcal{A}} : \square \subset \mathcal{A}$ induces by precomposition a forgetful functor

$$\omega_{\mathcal{A}} : \mathcal{A}^{op}\mathbf{Set} \longrightarrow \square^{op}\mathbf{Set}$$

which has a left adjoint

$$\mathcal{L}_{\mathcal{A}} : \square^{op}\mathbf{Set} \longrightarrow \mathcal{A}^{op}\mathbf{Set}$$

given by the left Kan extension along $j_{\mathcal{A}}$. For a precubical set K , the \mathcal{A} -set $\mathcal{L}_{\mathcal{A}}(K)$ is called the *free \mathcal{A} -set* generated by K .

2.9. Proposition. *The functor $\mathcal{L}_{\mathcal{A}} : \square^{op}\mathbf{Set} \rightarrow \mathcal{A}^{op}\mathbf{Set}$ is faithful.*

Proof. By [6, Proposition 2.1.15], the identity map $\text{Id}_{\mathcal{L}_{\mathcal{A}}(K)}$ induces for all precubical sets K a natural inclusion of precubical sets $K \subset \omega_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(K)$. Let $f, g : K \rightarrow L$ be two maps of precubical sets such that $\mathcal{L}_{\mathcal{A}}(f) = \mathcal{L}_{\mathcal{A}}(g)$. Then $\omega_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(f) = \omega_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(g)$. Thus

$$f = \omega_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(f) \upharpoonright_K = \omega_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(g) \upharpoonright_K = g.$$

□

Let K be an object of $\mathcal{A}^{op}\mathbf{Set}$. The set $K([n])$ is denoted by K_n . For any map $k : [m] \rightarrow [n]$ of \mathcal{A} and any \mathcal{A} -set K , denote by $k^* : K_n \rightarrow K_m$ the set map induced by k . Let $p \geq 0$. The *p-cube* $\mathcal{A}[p]$ is by definition the presheaf $\mathcal{A}(-, [p])$. For any \mathcal{A} -set K , the data

$$(K_{\leq n})_p = \begin{cases} K_p & \text{if } p \leq n \\ \emptyset & \text{if } p > n. \end{cases}$$

assemble into an \mathcal{A} -set denoted by $K_{\leq n}$ because $\mathcal{A}([m], [n]) = \emptyset$ when $m > n$. Let

$$\partial\mathcal{A}[n] = \mathcal{A}_{\leq n-1}$$

for all $n \geq 0$.

2.10. Definition. A category of cubes \mathcal{A} is *thick* if the factorization of Proposition 2.5 is a factorization in \mathcal{A} , i.e $f \in \mathcal{A}$ implies $\psi \in \mathcal{A}$.

The category of cubes $\widehat{\square}_S$ is thick for tautological reasons. The terminology must be understood as follows. A thick category of cubes \mathcal{A} is morally a thick subcategory of the category of cubes $\widehat{\square}_S$: it is an analogy with the notion of thick subcategory of a triangulated category. Theorem 2.16 provides other examples of thick categories of cubes.

2.11. **Proposition.** *Let \mathcal{A} be a thick category of cubes. For $n \geq 0$, let*

$$j_{\mathcal{A}}^n : (\square \downarrow [n]) \xrightarrow{\subset} (\mathcal{A} \downarrow [n])$$

be the functor between comma categories induced by the inclusion $j_{\mathcal{A}} : \square \subset \mathcal{A}$. Then for all $n \geq 0$ and for all objects k of $(\mathcal{A} \downarrow [n])$, the comma category $(k \downarrow j_{\mathcal{A}}^n)$ has an initial object.

Proof. Let $k : [p] \rightarrow [n]$. Using Proposition 2.5 and since \mathcal{A} is thick by hypothesis, we obtain the commutative diagram of \mathcal{A}

$$\begin{array}{ccc} [p] & \xrightarrow{f} & [p] \\ k \downarrow & & \downarrow \delta \in \square \\ [n] & \xlongequal{\quad} & [n] \end{array}$$

which is an element of the comma category $(k \downarrow j_{\mathcal{A}}^n)$. Consider another element of $(k \downarrow j_{\mathcal{A}}^n)$ depicted by the following commutative diagram of solid arrows of \mathcal{A} :

$$\begin{array}{ccc} [p] & \xrightarrow{g} & [p] \\ \parallel & & \downarrow \delta'' \in \square \\ [p] & \xrightarrow{f'} & [q] \\ k \downarrow & & \downarrow \delta' \in \square \\ [n] & \xlongequal{\quad} & [n] \end{array}$$

Since \mathcal{A} is thick, write $f' = \delta''g$ with $\delta'' \in \square$. We obtain $\delta f = k = \delta'f' = \delta'\delta''g$. By uniqueness of the factorization of Proposition 2.5, we obtain $\delta = \delta'\delta''$ and $f = g$. We have obtain the map of $(k \downarrow j_{\mathcal{A}}^n)$

$$\begin{array}{ccccc} [p] & \xrightarrow{f} & [p] & & \\ & \searrow & & \searrow \delta'' & \\ & & [p] & \xrightarrow{f'} & [q] \\ k \downarrow & & \downarrow \delta & & \downarrow \delta' \\ [n] & \xlongequal{\quad} & [n] & & [n] \\ & \searrow & & \searrow & \\ & & [n] & \xlongequal{\quad} & [n] \end{array}$$

Moreover, the map $\delta'' : [p] \rightarrow [q]$ is unique because it is given by the factorization of f' using Proposition 2.5. Hence the proof is complete. \square

2.12. **Proposition.** *Let \mathcal{A} be a category of cubes. For all $n \geq 0$, one has the isomorphism of \mathcal{A} -sets*

$$\mathcal{L}_{\mathcal{A}}(\square[n]) \cong \mathcal{A}[n].$$

If moreover \mathcal{A} is thick, then there is the isomorphism of \mathcal{A} -sets

$$\mathcal{L}_{\mathcal{A}}(\partial \square[n]) \cong \partial \mathcal{A}[n]$$

for all $n \geq 0$.

Proof. The first statement is [6, Proposition 2.1.14]. Let $n \geq 0$. Since $\mathcal{L}_{\mathcal{A}}$ is colimit-preserving, there is a natural map of \mathcal{A} -sets ($\square_{<n}$ and $\mathcal{A}_{<n}$ are the full subcategory of \square and \mathcal{A} respectively containing only $[0], \dots, [n-1]$):

$$\mathcal{L}_{\mathcal{A}}(\partial \square[n]) \cong \varinjlim_{(\square_{<n} \downarrow [n])} \mathcal{L}_{\mathcal{A}}(\square[p]) \cong \varinjlim_{(\square_{<n} \downarrow [n])} \mathcal{A}[p] \rightarrow \varinjlim_{(\mathcal{A}_{<n} \downarrow [n])} \mathcal{A}[p] \cong \partial \mathcal{A}[n].$$

The above arrow is an isomorphism by Proposition 2.11 and [16, Theorem 1 p. 213]. \square

2.13. Proposition. *The set of maps*

$$\widehat{\square} = \{\phi : [m] \rightarrow [n] \in \widehat{\square}_S \mid \forall \delta : [p] \rightarrow [m] \in \square, \phi\delta \text{ one-to-one} \Rightarrow \phi\delta \in \square\}$$

is closed under composition and contains all identities of $\widehat{\square}_S$. There are the inclusions $\square \subset \widehat{\square} \subset \widehat{\square}_S$. In other terms, the set of maps $\widehat{\square}$ yields a well-defined category of cubes. The only one-to-one set maps of $\widehat{\square}$ are the maps of \square .

Proof. Let $\phi_1, \phi_2 \in \widehat{\square}$ such that $\phi_1\phi_2$ exists. Let $\delta \in \square$ such that $\phi_1\phi_2\delta$ exists and is one-to-one. Then $\phi_2\delta$ is a one-to-one set map. Thus $\phi_2\delta \in \square$, ϕ_2 belonging to $\widehat{\square}$. We deduce that $\phi_1\phi_2\delta = \phi_1(\phi_2\delta) \in \square$ since $\phi_1 \in \widehat{\square}$. This means that $\phi_1\phi_2 \in \widehat{\square}$. For all $\phi = \text{Id}$, one has $\phi\delta = \delta \in \square$. Hence $\widehat{\square}$ contains all identity maps. Finally suppose that $f : [m] \rightarrow [n] \in \widehat{\square}$ is one-to-one. Then $f \text{Id}_{[m]}$ is one-to-one, which implies that $f \in \square$. \square

2.14. Definition. The $\widehat{\square}$ -sets are called *transverse sets*.

The following maps, introduced in [6, Definition 3.1.11], are examples of cotransverse degeneracy maps. Let $\gamma_i : [n] \rightarrow [n]$ be the set map defined for $1 \leq i \leq n-1$ and $n \geq 2$ by

$$\gamma_i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, \max(x_i, x_{i+1}), \min(x_i, x_{i+1}), x_{i+2}, \dots, x_n).$$

2.15. Notation. [6, Theorem 3.1.16] The category of cubes generated by the δ_i^α , σ_i and γ_i is denoted by $\overline{\square}$.

Unlike $\widehat{\square}$, the category of cubes $\overline{\square}$ has a conjectural presentation by generators and relations [6, Proposition 3.1.20 and Conjecture 3.1.21].

2.16. Theorem. *The categories of cubes \square , \square_S , $\widehat{\square}$ and $\overline{\square}$ are thick.*

Proof. Every map $f : [m] \rightarrow [n]$ of \square factors uniquely in $\widehat{\square}_S$ as a composite $[m] \rightarrow [m] \rightarrow [n]$ such that the right-hand map $[m] \rightarrow [n]$ belongs to \square . Since $f = f \text{Id}_{[m]}$, we deduce by uniqueness that the left-hand map belongs to \square . Hence the category \square is thick.

Every map $f : [m] \rightarrow [n]$ of \square_S factors uniquely in $\widehat{\square}_S$ as a composite $[m] \rightarrow [m] \rightarrow [n]$ such that the right-hand map $[m] \rightarrow [n]$ belongs to \square . Since all maps of \square_S are one-to-one, the left-hand map $[m] \rightarrow [m]$ is one-to-one, hence bijective for cardinality reason. We deduce that the left-hand map is a composite of symmetry maps, which means that it belongs to \square_S . Thus the category \square_S is thick.

Consider a map $f : [m] \rightarrow [n]$ of $\widehat{\square}$. It factors in $\widehat{\square}_S$ as a composite $[m] \rightarrow [m] \rightarrow [n]$ such that the right-hand map $\delta : [m] \rightarrow [n]$ belongs to the box category. Denote by $g : [m] \rightarrow [m]$ the left-hand map. Let $\delta' : [p] \rightarrow [m] \in \square$ such that $g\delta'$ exists and is

one-to-one. Then $f\delta' = \delta(g\delta')$ is one-to-one, being the composite of two one-to-one set maps. Since $f \in \widehat{\square}$, we deduce that $\delta g\delta' \in \square$. Using Proposition 2.6, we deduce that $g\delta' \in \square$. We have proved that $g \in \widehat{\square}$. This means that the category of cubes $\widehat{\square}$ is thick.

Finally, one has $\gamma_i \delta_j^\alpha = \delta_j^\alpha \gamma_i$ for $j \leq i - 1$ and $j \geq i + 2$, $\gamma_i \delta_i^0 = \delta_{i+1}^0$, $\gamma_i \delta_i^1 = \delta_i^1$, $\gamma_i \delta_{i+1}^0 = \delta_{i+1}^0$ and $\gamma_i \delta_{i+1}^1 = \delta_i^1$. Thus, $\overline{\square}$ is thick. \square

It is possible to obtain even more examples of thick categories of cubes by removing the symmetry maps from $\overline{\square}$, and also by adding one by one other cotransverse degeneracy maps in the set of generators. The only interest of the category of cubes $\overline{\square}$ for this note is to show that it is really easy to construct other thick categories of cubes. Proposition 2.17 gives a characterization of $\widehat{\square}$.

2.17. Theorem. *Let \mathcal{A} be a thick category of cubes which does not contain the symmetry maps. Then $\mathcal{A} \subset \widehat{\square}$. In other terms, $\widehat{\square}$ is the greatest thick category of cubes for the inclusion which does not contain the symmetry maps.*

Proof. Let $\phi : [m] \rightarrow [n]$ be a map of \mathcal{A} . Let $\delta : [p] \rightarrow [m] \in \square$ such that $\phi\delta$ is one-to-one. Write $\phi\delta = \delta'\phi'$ with $\delta' : [p] \rightarrow [n] \in \square$ and $\phi' : [p] \rightarrow [p] \in \mathcal{A}$, the category of cubes \mathcal{A} being thick by hypothesis. Since $\phi\delta$ is one-to-one, the set map $\phi' : [p] \rightarrow [p]$ is one-to-one. By hypothesis, this implies that ϕ' is the identity map. Thus $\phi\delta = \delta' \in \square$. We have proved that $\phi \in \widehat{\square}$. \square

To summarize, $\widehat{\square}$ is the greatest thick category of cubes for the inclusion not containing the symmetry maps and $\widehat{\square}_S$ is the greatest thick category of cubes for the inclusion.

For the sequel, \mathcal{A} denotes a fixed thick category of cubes. Note that with the choice $\mathcal{A} = \square$ (the least thick category of cubes), all following results remain valid. However, their formulation is not necessarily the best one.

3. METRIC AND HOMOTOPICAL STUDY

3.1. Notation. Let

$$\begin{aligned} \overrightarrow{\mathcal{A}} &= \mathcal{A}, \\ \overleftarrow{\mathcal{A}} &= \overleftarrow{\mathcal{A}} = \coprod_{n \geq 0} \{f : [n] \rightarrow [n] \mid f \in \mathcal{A}\}. \end{aligned}$$

We consider the degree function $d([n]) = n$ for all $n \geq 0$.

Every morphism $f : [m] \rightarrow [n]$ of \mathcal{A} is basic in the sense of [18, Definition 6.12] since every factorization of f as a composite $[m] \rightarrow [p] \rightarrow [n]$ implies that $m \leq p \leq n$, and therefore that every factorization is not fundamental in the sense of [18, Definition 6.12]: $p < \min(m, n) = m$ is impossible indeed. Hence, for all $n \geq 0$, the subcategory $\mathcal{A}_{=n}$ of \mathcal{A} generated by the objects of degree n and the basic morphisms in the sense of [18, Definition 6.12] between them is the full subcategory of \mathcal{A} having one object $[n]$. In particular, one has $\mathcal{A}_{=n}([n], [n]) = \mathcal{A}([n], [n])$.

3.2. Proposition. *The small category \mathcal{A} is c-Reedy in the sense of [18, Definition 8.25].*

Proof. Let f be a map of \mathcal{A} . Consider the category of factorizations of f : its objects are the pairs of morphisms (h, g) such that $hg = f$ with $h \in \overrightarrow{\mathcal{A}}$ and $g \in \overleftarrow{\mathcal{A}}$ and its morphisms

$k : (h, g) \rightarrow (h', g')$ are morphisms k (which are called connecting morphisms) such that there is a commutative diagram

$$\begin{array}{ccccc}
 \bullet & \xrightarrow{g} & \bullet & \xrightarrow{h} & \bullet \\
 \parallel & & \uparrow k & & \parallel \\
 \bullet & \xrightarrow{g'} & \bullet & \xrightarrow{h'} & \bullet
 \end{array}$$

By [9, Proposition 5.3] and since the factorization of Proposition 2.5 restricts to a factorization in \mathcal{A} , the category of factorization of f has a final object. The rest of the proof goes like the proof of [9, Proposition 5.4]. Let us summarize the argument. One has $\overleftarrow{\mathcal{A}} \subset \overrightarrow{\mathcal{A}} \cap \overleftarrow{\mathcal{A}}$ (first axiom). Every morphism of $\overleftarrow{\mathcal{A}}$ is degree-preserving (second axiom). Every morphism of $\overrightarrow{\mathcal{A}} \setminus \overleftarrow{\mathcal{A}}$ strictly raises degree and every morphism of $\overleftarrow{\mathcal{A}} \setminus \overrightarrow{\mathcal{A}} = \emptyset$ strictly lowers degree (third axiom). The category of factorizations of f with connecting maps in $\overleftarrow{\mathcal{A}}$ is connected since it has a final object (fourth axiom). For every $n \geq 0$, and any degree $m < n$, the functor $\overleftarrow{\mathcal{A}}([n], -) : \mathcal{A}_{=m} \rightarrow \mathbf{Set}$ is an (empty) coproduct of retracts of representables because $\mathcal{A}([n], [m]) = \emptyset$ (fifth axiom). \square

3.3. Notation. Let \mathcal{C} be a small category. Let \mathcal{M} be a locally small category. The category of functors from \mathcal{C} to \mathcal{M} together with the natural transformations is denoted by $\mathcal{M}^{\mathcal{C}}$.

3.4. Notation. Let $n \geq 0$. Following the notations of [18, page 37], let

$$\partial_n \mathcal{A}([p], [q]) = \int^{[m] \in \mathcal{A}^{<n}} \mathcal{A}([m], [q]) \times \mathcal{A}([p], [m])$$

The latching and matching object functors $L_n, M_n : \mathcal{M}^{\mathcal{A}} \rightarrow \mathcal{M}^{\mathcal{A}^{=n}}$ are given by

$$\begin{aligned}
 (M_n A)_{[n]} &= \int_{[m] \in \mathcal{A}} A([m])^{\partial_n \mathcal{A}([n], [m])} \\
 (L_n A)_{[n]} &= \int^{[p] \in \mathcal{A}} \partial_n \mathcal{A}([p], [n]) \cdot A([p])
 \end{aligned}$$

We obtain:

3.5. Theorem. *Let \mathcal{M} be a model category. Suppose that the projective model structure on $\mathcal{M}^{\mathcal{A}^{=n}}$ exists for all $n \geq 0$. There exists a unique model structure on $\mathcal{M}^{\mathcal{A}}$ such that*

- *The weak equivalences are objectwise.*
- *A map $A \rightarrow B$ of $\mathcal{M}^{\mathcal{A}}$ is a fibration (trivial fibration resp.) if for all $n \geq 0$, the map $A([n]) \rightarrow (M_n A)_{[n]} \times_{(M_n B)_{[n]}} B([n])$ is a fibration (trivial fibration resp.) of \mathcal{M} .*
- *A map $A \rightarrow B$ of $\mathcal{M}^{\mathcal{A}}$ is a cofibration (trivial cofibration resp.) if for all $n \geq 0$, $L_n B \sqcup_{L_n A} A \rightarrow B$ is a projective cofibration (trivial cofibration resp.) of the projective model structure of $\mathcal{M}^{\mathcal{A}^{=n}}$.*

This model structure is called the c-Reedy model structure of $\mathcal{M}^{\mathcal{A}}$.

Proof. By Proposition 3.2 and [18, Theorem 8.26], the small category \mathcal{A} is almost c-Reedy in the sense of [18, Definition 8.8]. The proof is complete thanks to [18, Theorem 8.9]. \square

3.6. Proposition. *One has*

$$\partial_n \mathcal{A}([p], [q]) = \begin{cases} \emptyset & \text{if } p > q \text{ or } n \leq p \\ \mathcal{A}([p], [q]) & \text{if } p \leq q \text{ and } p < n \end{cases}$$

Proof. It is mutatis mutandis the proof of [9, Proposition 5.8]: it suffices to change the category of cubes in the proof and to use Proposition 2.5 which restricts to a factorization in \mathcal{A} by definition of a thick category of cubes. \square

3.7. Theorem. *Let \mathcal{M} be a model category. Suppose that the projective model structure on $\mathcal{M}^{\mathcal{A}=n}$ exists for all $n \geq 0$. Then the projective model structure on $\mathcal{M}^{\mathcal{A}}$ exists and coincides with the *c-Reedy* model structure.*

Proof. The proof follows the road map of the proof of [9, Theorem 5.12] and makes use of Theorem 3.5 and Proposition 3.6. \square

3.8. Notation. The category of Δ -generated spaces or of Δ -Hausdorff Δ -generated spaces (cf. [7, Section 2 and Appendix B]) is denoted by **Top**.

3.9. Definition. [9, Definition 3.2] Let $f = (f_1, \dots, f_n) : [n] \rightarrow [n]$ be a cotransverse map. Let $T(f) : [0, 1]^n \rightarrow [0, 1]^n$ be the map defined by

$$T(f)(x_1, \dots, x_n) = (T(f)_1(x_1, \dots, x_n), \dots, T(f)_n(x_1, \dots, x_n))$$

with

$$T(f)_i(x_1, \dots, x_n) = \max_{(\epsilon_1, \dots, \epsilon_n) \in f_i^{-1}(1)} \min\{x_k \mid \epsilon_k = 1\}$$

for all $1 \leq i \leq n$.

3.10. Notation. For $\delta_i^\alpha : [n-1] \rightarrow [n] \in \square$, let

$$T(\delta_i^\alpha) = \begin{cases} [0, 1]^{n-1} \rightarrow [0, 1]^n \\ (\epsilon_1, \dots, \epsilon_{n-1}) \mapsto (\epsilon_1, \dots, \epsilon_{i-1}, \alpha, \epsilon_i, \dots, \epsilon_{n-1}) \end{cases}$$

for all $n \geq 1$ and $\alpha \in \{0, 1\}$.

The three mappings $[n] \mapsto [0, 1]^n$ for $n \geq 0$, $f : [n] \rightarrow [n] \in \mathcal{A} \mapsto T(f)$ and $\delta_i^\alpha : [n-1] \rightarrow [n] \mapsto T(\delta_i^\alpha)$ for $n \geq 1$ give rise to a functor from $\mathcal{A} \subset \widehat{\square}_S$ to **Top** denoted by $|\mathcal{A}[*]|_{geom}$ by [9, Theorem 3.9] and to a functor from $\mathcal{A} \subset \widehat{\square}_S$ to **LvMet** denoted by $|\mathcal{A}[*]|_{\vec{d}_1}$ by [9, Theorem 3.15]. Let K be an \mathcal{A} -set. Let

$$|K|_{geom} = \int^{[n] \in \mathcal{A}} K_n \cdot |\mathcal{A}[n]|_{geom} \text{ and } |K|_{\vec{d}_1} = \int^{[n] \in \mathcal{A}} K_n \cdot |\mathcal{A}[n]|_{\vec{d}_1}.$$

These give rise to two colimit-preserving functors from \mathcal{A} -sets to topological spaces and Lawvere metric spaces respectively. The latter functor factoring as a composite $\mathcal{A}^{op} \mathbf{Set} \rightarrow \widehat{\square}_S^{op} \mathbf{Set} \rightarrow \mathbf{LvMet}$, we define the notion of (tame or not) natural d -path in the geometric realization $|K|_{geom}$ like in [9, Section 4]. In fact, by using the inclusion $\mathcal{A} \subset \widehat{\square}_S$ and the fact that \mathcal{A} is thick, we can mimick all constructions of [9] and recover, thanks to Theorem 3.7, the results already proved for $\widehat{\square}_S$. More precisely, we obtain what follows.

3.11. Definition. [3, Definition 4.11] A *flow* is a small semicategory enriched over the closed monoidal category (\mathbf{Top}, \times) . The corresponding category is denoted by **Flow**. The objects are called *states* and the morphisms *execution paths*.

There is an inclusion functor $\mathbf{PoSet}^+ \subset \mathbf{Flow}$ such that there is an execution path from α to β if and only if $\alpha < \beta$. Let $r \in \{q, m, h\}$. The category \mathbf{Flow} is equipped with its r -model structure [8, Theorem 7.4]. The weak equivalences of the r -model structure are the maps of flows $f : X \rightarrow Y$ inducing a bijection on states and a weak equivalence of the r -model structure of \mathbf{Top} on execution paths.

3.12. Definition. Let $r \in \{q, m, h\}$. A functor $F : \mathcal{A}^{op}\mathbf{Set} \rightarrow \mathbf{Flow}$ is a r -realization functor if it satisfies the following properties:

- F is colimit-preserving.
- For all $n \geq 0$, the map $F(\partial\mathcal{A}[n]) \rightarrow F(\mathcal{A}[n])$ is a r -cofibration of \mathbf{Flow} .
- There is a map $F(\mathcal{A}[*]) \rightarrow \{0 < 1\}^*$ in $\mathbf{Flow}^{\mathcal{A}}$ which is an objectwise weak equivalence of the r -model structure of \mathbf{Flow} .

3.13. Definition. An \mathcal{A} -set K is *cellular* if the canonical map $\emptyset \rightarrow K$ is a transfinite composition of pushouts of the maps $\partial\mathcal{A}[n] \rightarrow \mathcal{A}[n]$ for $n \geq 0$. An \mathcal{A} -set K is *cofibrant* if it is a retract of a cellular \mathcal{A} -set.

The composite functor $\mathcal{A} \subset \widehat{\square}_S \rightarrow \mathbf{Flow}$ taking $[n]$ to the flow $|\square_S[n]|_{nat}$ defined in [9, Proposition 7.1] induces a colimit-preserving functor

$$|K|_{nat} = \int^{[n] \in \mathcal{A}} K_n \cdot |\square_S[n]|_{nat}$$

from $\mathcal{A}^{op}\mathbf{Set}$ to \mathbf{Flow} which is a m -realization functor by [9, Theorem 7.4]. The composite functor $\mathcal{A} \subset \widehat{\square}_S \rightarrow \mathbf{Flow}$ taking $[n]$ to the flow $(\{0 < 1\}^n)^{cof}$, where $(-)^{cof}$ is a q -cofibrant replacement of \mathbf{Flow} , induces a colimit-preserving functor

$$|K|_q = \int^{[n] \in \mathcal{A}} K_n \cdot (\{0 < 1\}^n)^{cof}$$

from $\mathcal{A}^{op}\mathbf{Set}$ to \mathbf{Flow} which is a q -realization functor by [9, Theorem 6.7].

3.14. Theorem. *There exists an m -realization functor $F : \mathcal{A}^{op}\mathbf{Set} \rightarrow \mathbf{Flow}$ and two natural transformations inducing bijections on the sets of states*

$$|-|_q \longleftarrow F(-) \Longrightarrow |-|_{nat}$$

such that for all cofibrant \mathcal{A} -sets K and all $(\alpha, \beta) \in K_0 \times K_0$, there is the zigzag of natural homotopy equivalences between m -cofibrant topological spaces

$$\mathbb{P}_{\alpha, \beta} |K|_q \xleftarrow{\simeq} \mathbb{P}_{\alpha, \beta} F(K) \xrightarrow{\simeq} \mathbb{P}_{\alpha, \beta} |K|_{nat} .$$

Proof. We follow the proof of [9, Theorem 7.4]. Details are left to the reader. \square

4. ZIEMIAŃSKI'S CUBE CHAIN

Cube chains are introduced in [19, Definition 1.1]. We use the presentation given in [20, Section 7] instead. Let $\text{Seq}(n)$ be the set of sequences of positive integers $\underline{n} = (n_1, \dots, n_p)$ with $n_1 + \dots + n_p = n$. Let $\underline{n} = (n_1, \dots, n_p) \in \text{Seq}(n)$. Then $|\underline{n}| = n$ is the length of \underline{n} and $\ell(\underline{n}) = p$ is the number of elements of \underline{n} . Let K be an \mathcal{A} -set and $A = a_1 < \dots < a_k \subset \{1, \dots, n\}$ and $\epsilon \in \{0, 1\}$. The *iterated face map* is defined by

$$\partial_A^\epsilon = \partial_{a_1}^\epsilon \partial_{a_2}^\epsilon \dots \partial_{a_k}^\epsilon$$

with $\partial_p^\epsilon = (\delta_p^\epsilon)^*$.

4.1. **Definition.** Let $\underline{n} \in \text{Seq}(n)$. The \underline{n} -cube is the \mathcal{A} -set

$$\mathcal{A}[\underline{n}] = \mathcal{A}[n_1] * \cdots * \mathcal{A}[n_p]$$

where the notation $*$ means that the final state 1_{n_i} of the \mathcal{A} -set $\mathcal{A}[n_i]$ is identified with the initial state $0_{n_{i+1}}$ of the \mathcal{A} -set $\mathcal{A}[n_{i+1}]$ for $1 \leq i \leq p-1$.

Let K be an \mathcal{A} -set. Let $\alpha, \beta \in K_0$. Let $n \geq 1$. The small category $\text{Ch}_{\alpha, \beta}(K, n)$ is defined as follows. The objects are the maps of \mathcal{A} -sets $\mathcal{A}[\underline{n}] \rightarrow K$ with $|\underline{n}| = n$ where the initial state of $\mathcal{A}[n_1]$ is mapped to α and the final state of $\mathcal{A}[n_p]$ is mapped to β . Let $A \sqcup B = \{1, \dots, m_1 + m_2\}$ be a partition with the cardinal of A equal to $m_1 > 0$ and the cardinal of B equal to $m_2 > 0$. Let

$$\phi_{A,B} : \mathcal{A}[m_1] * \mathcal{A}[m_2] \longrightarrow \mathcal{A}[m_1 + m_2]$$

be the unique map of \mathcal{A} -sets such that

$$\begin{aligned} \phi_{A,B}(\text{Id}_{[m_1]}) &= \partial_B^0(\text{Id}_{[m_1+m_2]}), \\ \phi_{A,B}(\text{Id}_{[m_2]}) &= \partial_A^1(\text{Id}_{[m_1+m_2]}). \end{aligned}$$

For $i \in \{1, \dots, \ell(\underline{n})\}$ and a partition $A \sqcup B = \{1, \dots, n_i\}$, let

$$\delta_{i,A,B} = \text{Id}_{\mathcal{A}[n_1]} * \cdots * \text{Id}_{\mathcal{A}[n_{i-1}]} * \phi_{A,B} * \text{Id}_{\mathcal{A}[n_{i+1}]} * \cdots * \text{Id}_{\mathcal{A}[n_{\ell(\underline{n})}]}$$

The morphisms are the commutative diagrams

$$\begin{array}{ccc} \mathcal{A}[\underline{n}_a] & \xrightarrow{a} & K \\ \downarrow & & \parallel \\ \mathcal{A}[\underline{n}_b] & \xrightarrow{b} & K \end{array}$$

where the left vertical map is a composite of maps of \mathcal{A} -sets of the form $\delta_{i,A,B}$. Let

$$\text{Ch}(K) = \coprod_{(\alpha, \beta) \in K_0 \times K_0} \text{Ch}_{\alpha, \beta}(K).$$

4.2. **Proposition.** *Suppose for this proposition only that \mathcal{A} is not necessarily thick. Let K be an \mathcal{A} -set. Then there is the isomorphism of categories $\text{Ch}(\omega_{\mathcal{A}}(K)) \cong \text{Ch}(K)$.*

Proof. It is an immediate consequence of the adjunction and $\mathcal{L}_{\mathcal{A}} \dashv \omega_{\mathcal{A}}$ and of the first part of Proposition 2.12 which does not require \mathcal{A} to be thick. \square

4.3. **Notation.** Since $\mathcal{L}_{\mathcal{A}}(\square[\underline{n}]) = \mathcal{A}[\underline{n}]$ for all $\underline{n} \in \text{Seq}(n)$, the functor

$$\mathcal{L}_{\mathcal{A}} : \square^{op}\text{Set} \longrightarrow \mathcal{A}^{op}\text{Set}$$

induces a functor

$$\mathcal{L}_{\mathcal{A}}^K : \text{Ch}(K) \longrightarrow \text{Ch}(\mathcal{L}_{\mathcal{A}}(K))$$

for all precubical sets K .

4.4. **Proposition.** *Let $0 \leq m \leq n$. Every map of \mathcal{A} -sets $f : \mathcal{A}[m] \rightarrow \mathcal{A}[n]$ factors uniquely as a composite $f = \mathcal{L}_{\mathcal{A}}(h)g$ with $g : \mathcal{A}[m] \rightarrow \mathcal{A}[m]$ and $h : \square[m] \rightarrow \square[n]$.*

Proof. It is a rephrasing of Proposition 2.5 obtained with the Yoneda lemma. \square

4.5. Proposition. *Let $m \geq 1$. Let K be a precubical set. Every map of \mathcal{A} -sets*

$$f : \mathcal{A}[m] \longrightarrow \mathcal{L}_{\mathcal{A}}(K)$$

factors uniquely as a composite of maps of \mathcal{A} -sets

$$\mathcal{A}[m] \xrightarrow{g} \mathcal{A}[m] \xrightarrow{\mathcal{L}_{\mathcal{A}}(h)} \mathcal{L}_{\mathcal{A}}(K)$$

where $h : \square[m] \rightarrow K$ is a map of precubical sets.

Proof. The functor $L \mapsto L_m$ from \mathcal{A} -sets to sets is colimit-preserving. Thus, there are the bijections

$$(\mathcal{L}_{\mathcal{A}}(K))_m \cong \varinjlim_{\square[p] \rightarrow K} (\mathcal{L}_{\mathcal{A}}(\square[p]))_m \cong \varinjlim_{\square[p] \rightarrow K} (\mathcal{A}([p]))_m \cong \varinjlim_{\square[p] \rightarrow K} \mathcal{A}([m], [p]).$$

The map of \mathcal{A} -sets $f : \mathcal{A}[m] \rightarrow \mathcal{L}_{\mathcal{A}}(K)$ gives rise to an element of the set $(\mathcal{L}_{\mathcal{A}}(K))_m$ by Yoneda. Therefore there exists a map $g : \mathcal{A}[m] \rightarrow \mathcal{A}[p]$ and a map of precubical sets $h : \square[p] \rightarrow K$ such that $f = \mathcal{L}_{\mathcal{A}}(h)g$. From Proposition 2.5 applied to g , we obtain $g = \mathcal{L}_{\mathcal{A}}(h')g'$ with $g' \in \mathcal{A}([m], [m])$ and we deduce $f = \mathcal{L}_{\mathcal{A}}(hh')g'$, which is the desired factorization. Consider two factorizations $f = \mathcal{L}_{\mathcal{A}}(h_i)g_i$ with $h_i : \square[m] \rightarrow K$ and $g_i : \mathcal{A}[m] \rightarrow \mathcal{A}[m]$ with $i = 1, 2$. They correspond to two representatives of the map f in the colimit of sets $\varinjlim_{\square[p] \rightarrow K} \mathcal{A}([m], [p])$. It means that there is a commutative diagram of \mathcal{A} -sets of the form

$$\begin{array}{ccccc} \mathcal{A}[m] & \xrightarrow{g_1} & \mathcal{A}[m] & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h_1)} & \mathcal{L}_{\mathcal{A}}(K) \\ \parallel & & \downarrow \mathcal{L}_{\mathcal{A}}(k_1) & & \parallel \\ \mathcal{A}[m] & \xrightarrow{g_3} & \mathcal{A}[p] & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h_3)} & \mathcal{L}_{\mathcal{A}}(K) \\ \parallel & & \uparrow \mathcal{L}_{\mathcal{A}}(k_2) & & \parallel \\ \mathcal{A}[m] & \xrightarrow{g_2} & \mathcal{A}[m] & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h_2)} & \mathcal{L}_{\mathcal{A}}(K) \end{array}$$

for some $g_3 \in \mathcal{A}([m], [p])$ indexed by the map of precubical sets $h_3 : \square[p] \rightarrow K$ in the diagram of sets. We obtain the equality $g_3 = \mathcal{L}_{\mathcal{A}}(k_1)g_1 = \mathcal{L}_{\mathcal{A}}(k_2)g_2$. From Proposition 4.4, we deduce $g_1 = g_2$ and $k_1 = k_2$. From the last equality, we deduce $\mathcal{L}_{\mathcal{A}}(h_1) = \mathcal{L}_{\mathcal{A}}(h_3)\mathcal{L}_{\mathcal{A}}(k_1) = \mathcal{L}_{\mathcal{A}}(h_3)\mathcal{L}_{\mathcal{A}}(k_2) = \mathcal{L}_{\mathcal{A}}(h_2)$. Since the functor $\mathcal{L}_{\mathcal{A}} : \square^{op}\mathbf{Set} \rightarrow \mathcal{A}^{op}\mathbf{Set}$ is faithful, we obtain $h_1 = h_2$. Hence the proof is complete. \square

4.6. Proposition. *Let $\underline{n} \in \text{Seq}(n)$ with $n \geq 1$. Let K be a precubical set. Every map of \mathcal{A} -sets $f : \mathcal{A}[\underline{n}] \longrightarrow \mathcal{L}_{\mathcal{A}}(K)$ factors uniquely as a composite of maps of \mathcal{A} -sets*

$$\mathcal{A}[\underline{n}] \xrightarrow{g} \mathcal{A}[\underline{n}] \xrightarrow{\mathcal{L}_{\mathcal{A}}(h)} \mathcal{L}_{\mathcal{A}}(K)$$

where $h : \square[\underline{n}] \rightarrow K$ is a map of precubical sets.

Proof. Let $\underline{n} = (n_1, \dots, n_p) \in \text{Seq}(n)$. Giving f is equivalent to giving p maps of \mathcal{A} -sets $f_i : \mathcal{A}[n_i] \rightarrow \mathcal{L}_{\mathcal{A}}(K)$ for $1 \leq i \leq p$ satisfying $f_i(1_{n_i}) = f_{i+1}(0_{n_{i+1}})$ for $1 \leq i \leq p-1$. Write

$$f = f_1 * \dots * f_p.$$

In the same way, given h is equivalent to giving p maps of precubical sets $h_i : \square[n_i] \rightarrow K$ for $1 \leq i \leq p$ satisfying $h_i(1_{n_i}) = h_{i+1}(0_{n_{i+1}})$ for $1 \leq i \leq p-1$. Write

$$h = h_1 * \cdots * h_p \text{ and } \mathcal{L}_{\mathcal{A}}(h) = \mathcal{L}_{\mathcal{A}}(h_1) * \cdots * \mathcal{L}_{\mathcal{A}}(h_p).$$

Then $f = \mathcal{L}_{\mathcal{A}}(h)g$ if and only if $f_i = \mathcal{L}_{\mathcal{A}}(h_i)g_i$ for all $1 \leq i \leq p$ where $g_i : [n] \rightarrow [n]$ is a map of \mathcal{A} . The proof is complete thanks to Proposition 4.5. \square

Proposition 4.2 shows that, for all precubical sets K , there is the isomorphism of small categories $\text{Ch}(\mathcal{L}_{\mathcal{A}}(K)) \cong \text{Ch}(\omega_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(K))$. Thus the small category $\text{Ch}(\mathcal{L}_{\mathcal{A}}(K))$ contains, in general, much more morphisms than $\text{Ch}(K)$.

4.7. Theorem. *Let K be a precubical set. The functor $\mathcal{L}_{\mathcal{A}}^K : \text{Ch}(K) \rightarrow \text{Ch}(\mathcal{L}_{\mathcal{A}}(K))$ induces a homotopy equivalence*

$$|\text{Ch}_{\alpha,\beta}(K)| \simeq |\text{Ch}_{\alpha,\beta}(\mathcal{L}_{\mathcal{A}}(K))|$$

for all $(\alpha, \beta) \in K_0 \times K_0$ between the classifying spaces where $|\mathcal{C}|$ means the classifying space of \mathcal{C} , i.e. the geometric realization of the simplicial nerve of \mathcal{C} .

Proof. By Quillen's Theorem A [14, Theorem 19.6.14], it suffices to prove that the comma category $(\underline{c} \downarrow \mathcal{L}_{\mathcal{A}}^K)$ is contractible for all cube chains \underline{c} of $\mathcal{L}_{\mathcal{A}}(K)$. By [14, Proposition 14.3.14], it suffices to prove that the comma category $(\underline{c} \downarrow \mathcal{L}_{\mathcal{A}}^K)$ has an initial object for all cube chains \underline{c} of $\mathcal{L}_{\mathcal{A}}(K)$. An object of the comma category $(\underline{c} \downarrow \mathcal{L}_{\mathcal{A}}^K)$ is a commutative square of \mathcal{A} -sets of the form

$$\begin{array}{ccc} \mathcal{A}[\underline{m}] & \xrightarrow{g} & \mathcal{L}_{\mathcal{A}}(\square[\underline{n}]) \\ \underline{c} \downarrow & & \downarrow \mathcal{L}_{\mathcal{A}}(h) \\ \mathcal{L}_{\mathcal{A}}(K) & \xlongequal{\quad} & \mathcal{L}_{\mathcal{A}}(K) \end{array}$$

where $h : \square[\underline{n}] \rightarrow K$ is a map of precubical sets. Proposition 4.6 provides such a commutative square

$$\begin{array}{ccc} \mathcal{A}[\underline{m}] & \xrightarrow{g_0} & \mathcal{L}_{\mathcal{A}}(\square[\underline{m}]) \\ \underline{c} \downarrow & & \downarrow \mathcal{L}_{\mathcal{A}}(h_0) \\ \mathcal{L}_{\mathcal{A}}(K) & \xlongequal{\quad} & \mathcal{L}_{\mathcal{A}}(K) \end{array}$$

Consider the commutative diagram of solid arrows of \mathcal{A} -sets

$$\begin{array}{ccccc} \mathcal{A}[\underline{m}] & \xrightarrow{g_0} & \mathcal{L}_{\mathcal{A}}(\square[\underline{m}]) & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h_0)} & \mathcal{L}_{\mathcal{A}}(K) \\ \parallel & \dashrightarrow^{g_1} & \downarrow \mathcal{L}_{\mathcal{A}}(h_1) & & \parallel \\ \mathcal{A}[\underline{m}] & \xrightarrow{g} & \mathcal{L}_{\mathcal{A}}(\square[\underline{n}]) & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h)} & \mathcal{L}_{\mathcal{A}}(K) \end{array}$$

with $\underline{c} = \mathcal{L}_{\mathcal{A}}(h_0)g_0 = \mathcal{L}_{\mathcal{A}}(h)g$. Using Proposition 4.6, write $g = \mathcal{L}_{\mathcal{A}}(h_1)g_1$. We obtain the equalities

$$\mathcal{L}_{\mathcal{A}}(h_0)g_0 = \underline{c} = \mathcal{L}_{\mathcal{A}}(h)g = \mathcal{L}_{\mathcal{A}}(h)\mathcal{L}_{\mathcal{A}}(h_1)g_1 = \mathcal{L}_{\mathcal{A}}(hh_1)g_1.$$

By Proposition 4.6 and since $\mathcal{L}_{\mathcal{A}}$ is faithful by Proposition 2.9, we obtain $hh_1 = h_0$ and $g_1 = g_0$. We have proved that there exists a diagram of solid arrows of \mathcal{A} -sets

$$\begin{array}{ccccc} \mathcal{A}[\underline{m}] & \xrightarrow{g_0} & \mathcal{L}_{\mathcal{A}}(\square[\underline{m}]) & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h_0)} & \mathcal{L}_{\mathcal{A}}(K) \\ \parallel & & \downarrow \mathcal{L}_{\mathcal{A}}(h_1) & & \parallel \\ \mathcal{A}[\underline{m}] & \xrightarrow{g} & \mathcal{L}_{\mathcal{A}}(\square[\underline{n}]) & \xrightarrow{\mathcal{L}_{\mathcal{A}}(h)} & \mathcal{L}_{\mathcal{A}}(K) \end{array}$$

From $g = \mathcal{L}_{\mathcal{A}}(h_1)g_0$ and Proposition 4.6, we deduce that there exists a unique map $\mathcal{L}_{\mathcal{A}}(h_1)$ making the above diagram commutative. We have proved that the commutative square

$$\begin{array}{ccc} \mathcal{A}[\underline{m}] & \xrightarrow{g_0} & \mathcal{L}_{\mathcal{A}}(\square[\underline{m}]) \\ \downarrow \underline{c} & & \downarrow \mathcal{L}_{\mathcal{A}}(h_0) \\ \mathcal{L}_{\mathcal{A}}(K) & \xlongequal{\quad} & \mathcal{L}_{\mathcal{A}}(K) \end{array}$$

is an initial object of the comma category $(\underline{c} \downarrow \mathcal{L}_{\mathcal{A}}^K)$ and the proof is complete. \square

4.8. Corollary. *Let K be a precubical set. The space of tame natural d -paths of $\mathcal{L}_{\mathcal{A}}(K)$ is homotopy equivalent to $|\mathrm{Ch}(\mathcal{L}_{\mathcal{A}}(K))|$.*

Proof. There are the homeomorphisms (natural with respect to $[n] \in \square$)

$$|\square[n]|_{\mathrm{geom}} \cong [0, 1]^n \cong |\mathcal{A}[n]|_{\mathrm{geom}} \cong |\mathcal{L}_{\mathcal{A}}(\square[n])|_{\mathrm{geom}},$$

the two homeomorphisms on the left by definition of the geometric realization, the right-hand homeomorphism by Proposition 2.12. Since all involved functors are colimit-preserving, we obtain for all precubical sets K the natural homeomorphism

$$|K|_{\mathrm{geom}} \cong |\mathcal{L}_{\mathcal{A}}(K)|_{\mathrm{geom}}.$$

Hence the space of tame natural d -paths of K is equal to the one of $\mathcal{L}_{\mathcal{A}}(K)$. By [20, Theorem 7.5], we deduce that the space of tame natural d -paths of $\mathcal{L}_{\mathcal{A}}(K)$ is homotopy equivalent to $|\mathrm{Ch}(K)|$, and therefore homotopy equivalent to $|\mathrm{Ch}(\mathcal{L}_{\mathcal{A}}(K))|$ by Theorem 4.7. \square

5. APPLICATION

We refer to [5, 6] for further details. [6, Theorem 4.1.8] states that, for two labelled cubes $\square[a_1, \dots, a_m]$ and $\square[a_{m+1}, \dots, a_{m+n}]$ with $m \geq 0$ and $n \geq 0$, there is an isomorphism of σ -labelled symmetric transverse sets

$$\begin{aligned} \mathcal{L}_{\widehat{\square}_S} \left(\overrightarrow{\mathrm{cosk}}^{\Sigma} \left(\square[a_1, \dots, a_m]_{\leq 1} \times_{\Sigma} \square[a_{m+1}, \dots, a_{m+n}]_{\leq 1} \right) \right) \\ \cong \mathrm{cosk}_1^{\widehat{\square}_S, \overline{\Sigma}} \left(\widehat{\square}_S[a_1, \dots, a_m]_{\leq 1} \overline{\times}_{\Sigma} \widehat{\square}_S[a_{m+1}, \dots, a_{m+n}]_{\leq 1} \right) \end{aligned}$$

where

- $\overrightarrow{\text{cosk}}^\Sigma$ is the σ -labelled directed coskeleton of [5, Section 3.3] which is a tweak of the σ -labelled coskeleton functor of the category of precubical sets, the latter being badly behaved by [5, Proposition 3.15]: it contains too many cubes and some of them have to be identified;
- $\text{cosk}_1^{\widehat{\square}_S, \overline{\Sigma}}$ is the σ -labelled coskeleton functor of the category of symmetric transverse sets which is well behaved by [6, Theorem 3.1.15].

By Theorem 4.7, the underlying precubical set of

$$\overrightarrow{\text{cosk}}^\Sigma (\square_{[a_1, \dots, a_m]_{\leq 1}} \times_\Sigma \square_{[a_{m+1}, \dots, a_{m+n}]_{\leq 1}})$$

and the underlying symmetric transverse set of

$$\text{cosk}_1^{\widehat{\square}_S, \overline{\Sigma}} (\widehat{\square}_S[a_1, \dots, a_m]_{\leq 1} \overline{\times}_\Sigma \widehat{\square}_S[a_{m+1}, \dots, a_{m+n}]_{\leq 1})$$

have homotopy equivalent categories of cube chains.

This implies that the parallel composition with synchronization for process algebra, as it is formalized in [6] using the labelled coskeleton functor of the category of symmetric transverse sets, has a category of cube chains which gives the correct homotopy type of tame natural d -paths.

REFERENCES

- [1] R. Brown and P. J. Higgins. On the algebra of cubes. *J. Pure Appl. Algebra*, 21(3):233–260, 1981. [https://doi.org/10.1016/0022-4049\(81\)90018-9](https://doi.org/10.1016/0022-4049(81)90018-9). 4
- [2] L. Fajstrup, E. Goubault, E. Haucourt, S. Mimram, and M. Raussen. *Directed algebraic topology and concurrency. With a foreword by Maurice Herlihy and a preface by Samuel Mimram*. SpringerBriefs Appl. Sci. Technol. Springer, 2016. <https://doi.org/10.1007/978-3-319-15398-8>. 1
- [3] P. Gaucher. A model category for the homotopy theory of concurrency. *Homology Homotopy Appl.*, 5(1):p.549–599, 2003. <https://doi.org/10.4310/hha.2003.v5.n1.a20>. 9
- [4] P. Gaucher. Globular realization and cubical underlying homotopy type of time flow of process algebra. *New York J. Math.*, 14:101–137, 2008. 2
- [5] P. Gaucher. Towards a homotopy theory of process algebra. *Homology Homotopy Appl.*, 10(1):353–388, 2008. <https://doi.org/10.4310/HHA.2008.v10.n1.a16>. 1, 14, 15
- [6] P. Gaucher. Combinatorics of labelling in higher-dimensional automata. *Theor. Comput. Sci.*, 411(11-13):1452–1483, March 2010. <https://doi.org/10.1016/j.tcs.2009.11.013>. 1, 2, 3, 4, 6, 14, 15
- [7] P. Gaucher. Left properness of flows. *Theory Appl. Categ.*, 37(19):562–612, 2021. 9
- [8] P. Gaucher. Six model categories for directed homotopy. *Categ. Gen. Algebr. Struct. Appl.*, 15(1):145–181, 2021. <https://doi.org/10.52547/cgasa.15.1.145>. 10
- [9] P. Gaucher. Directed degeneracy maps for precubical sets, 2022. <https://doi.org/10.48550/arXiv.2209.02667>. 1, 2, 3, 4, 8, 9, 10
- [10] P. Gaucher. Comparing cubical and globular directed paths. *Fundamenta Mathematicae*, 2023. <https://doi.org/10.4064/fm219-3-2023>. 2

- [11] P. Gaucher. Homotopy theory of Moore flows (III), 2023. <https://doi.org/10.48550/arXiv.2303.16174>. 2
- [12] E. Goubault and S. Mimram. Directed homotopy in non-positively curved spaces. *Log. Methods Comput. Sci.*, 16(3):55, 2020. Id/No 4. [https://doi.org/10.23638/LMCS-16\(3:4\)2020](https://doi.org/10.23638/LMCS-16(3:4)2020). 1
- [13] M. Grandis and L. Mauri. Cubical sets and their site. *Theory Appl. Categ.*, 11(8):185–211, 2003. 2, 3, 4
- [14] P. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003. <https://doi.org/10.1090/surv/099>. 2, 13
- [15] F. W. Lawvere. Metric spaces, generalized logic, and closed categories. *Repr. Theory Appl. Categ*, 2002(1):1–37, 2002. 2
- [16] S. Mac Lane. *Categories for the working mathematician*. Springer-Verlag, New York, second edition, 1998. <https://doi.org/10.1007/978-1-4757-4721-8>. 6
- [17] M. Raussen. Trace spaces in a pre-cubical complex. *Topology Appl.*, 156(9):1718–1728, 2009. <https://doi.org/10.1016/j.topol.2009.02.003>. 1
- [18] M. Shulman. Reedy categories and their generalizations, 2015. <https://doi.org/10.48550/arXiv.1507.01065>. 2, 7, 8
- [19] K. Ziemiański. Spaces of directed paths on pre-cubical sets. *Appl. Algebra Engrg. Comm. Comput.*, 28(6):497–525, 2017. <https://doi.org/10.1007/s00200-017-0316-0>. 1, 10
- [20] K. Ziemiański. Spaces of directed paths on pre-cubical sets II. *J. Appl. Comput. Topol.*, 4(1):45–78, 2020. <https://doi.org/10.1007/s41468-019-00040-z>. 1, 2, 10, 14

UNIVERSITÉ PARIS CITÉ, CNRS, IRIF, F-75013, PARIS, FRANCE
 URL: <http://www.irif.fr/~gaucher>