

A formula for the upper box-counting dimension of self-projective sets

Benedict Sewell*

November 8, 2023

Abstract

We prove a packing exponent–style formula for the upper box-counting dimension of attractors of certain projective iterated function systems. This partially affirms a conjecture of De Leo [11], and gives that the box-counting dimension of the Rauzy gasket \mathcal{R} , $\dim_B(\mathcal{R})$ exists and satisfies

$$\dim_B(\mathcal{R}) = \dim_H(\mathcal{R}) \in [1.6196, 1.7415].$$

1 Introduction

The Rauzy gasket is an interesting set. Discovered independently no fewer than three times, it is an important subset of parameter space for numerous problems in dynamics and topology, in fields as diverse as crystalline conductivity [24], triangular tiling billiards [20], and asymptotics on Markov–Hurwitz varieties [18]. Outside of these contexts, it is interesting in its own right as a fractal object, specifically for the challenge it poses: concrete bounds on its fractal dimensions (specifically Hausdorff and box-counting) have been hard to come by, despite a lot of recent attention. The gasket is a prototypical example of a self-projective attractor. The fractal geometric theory of such sets has so far only been undertaken in one dimension [7, 9, 32], with the recent exception of [21].

As suggested by the title, our main result is an explicit formula for the upper box-counting dimension of self-projective attractors with simplicial holes, which we prove using tools from classical analytic number theory. This allows us to derive a lower bound supporting a general conjecture of De Leo [11], and we can also give an upper bound in the more general case of convex holes. Concerning the Rauzy gasket \mathcal{R} , the main result applies, together with a corresponding formula of [21], to show that its box-counting dimension $\dim_B(\mathcal{R})$ exists and equals its Hausdorff dimension $\dim_H(\mathcal{R})$, and to give the strongest numerical lower bound for this (Hausdorff) dimension to date.

Before we state our results, we introduce the Rauzy gasket, illustrating some of the difficulty concerned by comparing it to two other well-known gaskets, and listing what is known about its dimension.

*Supported generously by the Alfréd Rényi Young Researcher Fund.

1.1 Good, bad and ugly gaskets

The three gaskets we touch upon in this section are mutually homeomorphic [2] but extremely different in the fractal geometric sense. Such differences can even be seen visually in the figures given below.

The word gasket¹ in each case refers to the attractor \mathcal{G} of three distance-decreasing maps f_k of some triangular subset of the plane, i.e., \mathcal{G} is the unique non-empty compact set such that $\mathcal{G} = f_1(\mathcal{G}) \cup f_2(\mathcal{G}) \cup f_3(\mathcal{G})$.

Certain properties of the f_j , specifically whether they are weakly or strongly contracting, and how distorting they are, are reflected in the relative ease of determining, or estimating, the fractal dimensions of the corresponding \mathcal{G} , as we shall see.

1.1.1 The Sierpiński gasket




Figure 1: The Sierpiński gasket, \mathcal{S} .

The Sierpiński gasket, \mathcal{S} , introduced by Waław Sierpiński in 1915 [29] and shown in Figure 1, is the example seen in first courses in fractal geometry.


Up to linear deformation, it is the limit set of three contracting similarities $x \mapsto \frac{1}{2}(x + e_k)$, where $(e_k)_{k=1}^3 \subset \mathbb{R}^2$ are fixed, non-colinear points. The rigidity and strong contractiveness of these maps makes the Hausdorff and box-counting dimensions easy to explicitly calculate:

$$\dim_H(\mathcal{S}) = \dim_B(\mathcal{S}) = \frac{\log(3)}{\log(2)} = 1.5849\dots$$

Hence it is not particularly interesting from the fractal geometric viewpoint. We however note some interesting connections to, e.g., cellular automata and Pascal's triangle, which can be found on Wikipedia.

¹We much prefer the French term, *baderne*—thanks to P. Arnoux for the introduction.

1.1.2 The Apollonian gasket

Figure 2: The Apollonian gasket, \mathcal{A}

The Apollonian gasket \mathcal{A} has perhaps the longest and richest history of any fractal, dating back to a theorem of Apollonius of Perga (ca. 200BC), which even features a formula of Descartes. We defer the reader to [26] for a readable historical account.

Up to a conformal change of coordinates, \mathcal{A} is the attractor of three Möbius maps:

$$z \mapsto \frac{z-1}{z+3}, \quad z \mapsto \frac{z+1}{3-z}, \quad \text{and} \quad z \mapsto \frac{1}{z-2i},$$

and is depicted in Figure 2.

The key difference between \mathcal{S} and \mathcal{A} is that the former is a hyperbolic attractor whereas the latter is parabolic: whereas the previous maps were contracting uniformly, here they are weakly attracting, having an indifferent fixed point. Visually, this corresponds to the fact that the largest circular holes approaching a vertex decay polynomially in size, not exponentially as in the previous example.

Correspondingly, the precise value for the Hausdorff and box-counting dimensions is unknown, but they do coincide. This is due to the conformality of the maps, as is the following formula of Boyd, which equates the so-called packing exponent to both Hausdorff and box-counting dimensions.

Proposition 1.1 (Boyd [8]). *Enumerating the radii of all the circles excised from \mathcal{A} as $(r_k)_{k=1}^{\infty}$,*

$$\dim_B(\mathcal{A}) = \dim_H(\mathcal{A}) = \inf \left\{ t > 0 : \sum_{k=1}^{\infty} r_k^t < \infty \right\}.$$

The dimension has other useful formulations, for example thermodynamically in terms of the root of the topological pressure function [17, 22]. As for the value, Boyd [8] showed

$\dim_H(\mathcal{A}) \in [1.300197, 1.314534]$ and McMullen [25] estimated $\dim_H(\mathcal{A}) \approx 1.30568$, agreeing with more recent empirical estimates of Bai–Finch [5] and others.

1.1.3 The Rauzy gasket




Figure 3: The Rauzy gasket, \mathcal{R} .

The Rauzy gasket, \mathcal{R} , in the fractal geometric sense represents the worst of all worlds. Not only is it a parabolic attractor like the Apollonian gasket, but the maps involved are non-conformal and highly distorting.

Here, the three attracting maps T_1, T_2, T_3 , acting on the standard 2-simplex, are the projective action of the matrices

$$N_1 = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad N_2 = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad N_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$$

(general definitions are given below), and are explicitly given by

$$\begin{aligned} T_1(x, y, z) &= \left(\frac{1}{2-x}, \frac{y}{2-x}, \frac{z}{2-x} \right), \\ T_2(x, y, z) &= \left(\frac{x}{2-y}, \frac{1}{2-y}, \frac{z}{2-y} \right), \\ T_3(x, y, z) &= \left(\frac{x}{2-z}, \frac{y}{2-z}, \frac{1}{2-z} \right). \end{aligned}$$

Visually, the triangles in Figure 3 decay polynomially in size as they approach the corners, and are often quite elongated. Correspondingly, much less is known in comparison to the other two examples, as we now describe.

1.2 Previous results about the area and dimension of \mathcal{R}

The Rauzy gasket has an interesting history, appearing for the first time in 1991 in the work of Arnoux and Rauzy [1] in the context of interval exchange transformations, where it was conjectured that $\text{area}(\mathcal{R}) = 0$. The gasket was rediscovered by Levitt in 1993, in a paper [23] which also included a proof that $\text{area}(\mathcal{R}) = 0$, due to Yoccoz. The gasket emerged for a third time in the work of De Leo and Dynnikov [13], this time in the context of Novikov's theory of magnetic induction on monocrystals.² They gave an alternative proof that $\text{area}(\mathcal{R}) = 0$ and proposed the stronger result $\dim_H(\mathcal{R}) < 2$. Furthermore, a conjecture of Maltsev and Novikov [24] implies $\dim_H(\mathcal{R}) \in (1, 2)$. But what is actually known?

- De Leo and Dynnikov in [13] empirically estimate $\dim_B(\mathcal{R}) \approx 1.72$.
- Avila, Hubert and Skripchenko in [4] prove $\dim_H(\mathcal{R}) < 2$.
- Gutiérrez-Romo and Matheus in [19] show $\dim_H(\mathcal{R}) \geq 1.19$.
- Fougeron in [16] shows $\dim_H(\mathcal{R}) \leq 1.825$, using estimates of Baragar [6].
- Pollicott and the author in [28] prove $\dim_H(\mathcal{R}) \leq 1.7415$.

Most recently, Jiao, Li, Pan and Xu in [21] proved a formula expressing $\dim_H(\mathcal{R})$ as the critical exponent of a certain Poincaré series, as we will present and use explicitly later.

1.3 Main results

1.3.1 Set-up

We first set-up the context and notation needed to state our general results.

Definition 1.2. *Given an $(n + 1) \times (n + 1)$ matrix $N \in \mathbb{R}^{n+1, n+1}$ such that all its entries are non-negative and $\det(N) = \pm 1$, its **projectivisation** is the normalised map*

$$T : \Delta^{(n)} \rightarrow \Delta^{(n)}, \quad T(x) = \frac{N \cdot x}{|N \cdot x|},$$

where $|\cdot|$ denotes the l^1 norm and $\Delta^{(n)} = \{x \in [0, 1]^{n+1} : |x| = 1\}$ the standard n -simplex. (Note that T is injective because N is invertible.)

²See [14] for the dichotomy between [13] and [1].

Fix a dimension $d \geq 2$, and a collection of $(d+1) \times (d+1)$ matrices N_1, \dots, N_m as in the above definition. Denoting their respective projectivisations by T_1, \dots, T_m and Δ as $\Delta^{(d)}$, we take \mathcal{G} to be the limit set of these maps, i.e., the unique non-empty subset of Δ such that

$$\mathcal{G} = \bigcup_{j=1}^m T_j(\mathcal{G}).$$

This \mathcal{G} we call a **self-projective attractor**.

Definition 1.3. We refer to a connected component of $\Delta \setminus \mathcal{G}$ as a **hole** (of \mathcal{G}), and a connected component of $\Delta \setminus \bigcup_{j=1}^m T_j(\Delta)$ as a **main hole**.

Finally note that, when we refer to the interior or boundary of a set, we mean as a subset of the d -dimensional hyperplane $\{x \in \mathbb{R}^{d+1} : |x| = 1\} \supset \Delta$.

1.3.2 General results

Our main theorem is the following.

Theorem 1.4. Suppose that the images $T_1(\Delta), \dots, T_m(\Delta)$ have disjoint interiors, that \mathcal{G} has zero d -dimensional Lebesgue measure, and that the main holes of \mathcal{G} are the interiors of d -simplices. Then, enumerating the holes as $\{\nabla_n\}_{n=1}^\infty$,

$$\overline{\dim}_B(\mathcal{G}) = d + \inf \left\{ t \geq -1 : \sum_{n=1}^\infty \text{vol}(\nabla_n) \text{in}(\nabla_n)^t < \infty \right\}, \quad (1)$$

where $\text{in}(\nabla_n)$ denotes the **inradius** of ∇_n , the radius of the largest ball contained in ∇_n , and $\text{vol}(\nabla_n)$ its d -dimensional volume.

Example 1.5. In the case of the Rauzy gasket, the main hole is the interior of the triangle with vertices $(\frac{1}{2}, \frac{1}{2}, 0)$, $(0, \frac{1}{2}, \frac{1}{2})$ and $(\frac{1}{2}, 0, \frac{1}{2})$, the $T_j(\Delta)$ meet each other only at their vertices, and $\text{area}(\mathcal{R}) = 0$, as cited above. Thus Theorem 1.4 applies.




Figure 4: Illustrating the geometric hypotheses of Theorem 1.4.

Example 1.6. To illustrate the geometric hypotheses of the theorem, Figure 4 shows four examples for $d = 2$ (the images $T_j(\Delta)$ are shown in yellow and the main holes in white). Considering the hypotheses of Theorem 1.4, only in the first does the theorem apply:

1. The geometric hypotheses are satisfied here.
2. The main hole contains part of the boundary of Δ , so is not the interior of a triangle.
3. The images do not have disjoint interiors.
4. The main hole is the interior of a convex quadrilateral, not a triangle.

Note that, in the last case, Theorem 1.7 below would give an upper bound on the dimension.

As mentioned in the example, the method of proof generalises to give an upper bound when the holes are merely convex, as follows.

Theorem 1.7. *Suppose that $T_1(\Delta), \dots, T_m(\Delta)$ have disjoint interiors, that $\text{vol}(\mathcal{G}) = 0$, and that the main holes are the interiors of convex d -dimensional polytopes. Then, enumerating the holes by $\{\nabla_n\}_{n=1}^\infty$,*

$$\overline{\dim}_B(\mathcal{G}) \leq d + \inf \left\{ t \geq -1 : \sum_{n=1}^{\infty} \text{vol}(\nabla_n) \text{in}(\nabla_n)^t < \infty \right\}. \quad (2)$$

Returning to Theorem 1.4, one of the main corollaries is the following result which partially confirms a conjecture of De Leo [11]. It will be written in a more convenient way later, once some notation is established.

Theorem 1.8. *Under the assumptions of Theorem 1.4,*

$$\overline{\dim}_B(\mathcal{G}) \geq \frac{d-1}{d} \inf \left\{ t > 0 : \sum_{n=1}^{\infty} \sum_{|i|=n} \|N_{i_1} N_{i_2} \cdots N_{i_n}\|^{-t} < \infty \right\} \quad (3)$$

$$= \frac{d-1}{d} \lim_{t \rightarrow \infty} \left(\frac{1}{t} \log \sum_{n=0}^{\infty} \# \left\{ |i| = n : \|N_{i_1} N_{i_2} \cdots N_{i_n}\| < e^t \right\} \right), \quad (4)$$

where $|i| = n$ is shorthand for $i \in \{1, 2, \dots, m\}^n$ and $\|\cdot\|$ is any matrix norm.

1.3.3 Application to the Rauzy gasket

In the context of the Rauzy gasket, we combine Theorem 1.4 with the main result of [21] to show that the box-counting dimension of the Rauzy gasket exists and equals its Hausdorff dimension:

Theorem 1.9. $\dim_B(\mathcal{R}) = \dim_H(\mathcal{R})$.

Regarding its value, along with the upper bound $\dim_H(\mathcal{R}) \leq 1.7415$, we give an effective lower bound via Theorem 1.8, using estimates of [30] and the Mathematica code available at [31]:

Theorem 1.10.

$$\dim_H(\mathcal{R}) \geq 1.6196.$$

1.4 Contents of this note

In section 2, we introduce some necessary notation and definitions. In section 3, we relate the ε -neighbourhood of \mathcal{G} to the interior neighbourhoods of the holes, and in section 4, we use the fact that the holes are simplicial to derive an explicit formula. In section 5 we deduce an expression for the Mellin–Laplace transform, which we use to complete the proof of Theorem 1.4 in section 6. Section 7 is dedicated to proving the lower bound in Theorem 1.8, and section 8 the upper bound in the case of convex holes. The final section 9 contains the applications to the Rauzy gasket, and the appendix contains the proofs of all our geometric lemmas.

2 Definitions

2.1 Notation

From now on, unless otherwise stated we fix the above notation and the assumptions of Theorem 1.4, to which we add the following convenient notation.

Definition 2.1. *Let*

$$\mathcal{I} := \{\emptyset\} \cup \bigcup_{n \in \mathbb{N}} \{1, 2, \dots, m\}^n$$

denote the set of finite words in $\{1, 2, \dots, m\}$. Given $i = (i_1, \dots, i_n) \in \mathcal{I}$, write

- $N_i = N_{i_1} \cdot N_{i_2} \cdot \dots \cdot N_{i_n}$,
- $T_i = T_{i_1} \circ T_{i_2} \circ \dots \circ T_{i_n}$, and
- $\Delta_i = T_i(\Delta)$.

Conventionally, N_\emptyset refers to the identity matrix, T_\emptyset to the identity map, and Δ_\emptyset to Δ .

Furthermore, since by assumption the main holes $\nabla_1, \dots, \nabla_K$ are the interiors of d -simplices, one can construct matrices M_1, \dots, M_K such that the closure of each ∇_k is the image of the projectivisation, S_k say, of M_k (in the case of the Rauzy gasket, we omit the subscripts). Fixing these, we naturally extend our alphabet \mathcal{I} to $\mathcal{I}^* = \mathcal{I} \times \{1, 2, \dots, K\}$ and our notation as follows: if $i' = (i, k) \in \mathcal{I}^*$, write

- $N_{i'} := N_i M_k$,
- $T_{i'} := T_i S_k$, and
- $\nabla_{i'} = T_i(\nabla_k)$.

Note that our use of ∇ is to distinguish holes from images.

2.2 Minkowski box-counting dimension

It is convenient here to use the following version of the box-counting dimension. All definitions and statements shall be made in the context of $\Delta = \Delta^{(d)}$ for simplicity.

Definition 2.2. Given $\Omega \subset \Delta$ closed and $\varepsilon > 0$, let

$$\Omega_\varepsilon = \{x \in \Delta : \|x - y\| \leq \varepsilon \text{ for some } y \in \Omega\}$$

denote the ε -neighbourhood of Ω , where $\|\cdot\|$ is the standard Euclidean metric.

Definition 2.3. If $\mathcal{G} \subset \Delta$ is closed, we define its upper box-counting dimension $\overline{\dim}_B(\mathcal{G})$ by

$$\overline{\dim}_B(\mathcal{G}) = d - \liminf_{\varepsilon \rightarrow 0^+} \left(\frac{\log(\text{vol}(\mathcal{G}_\varepsilon))}{\log(\varepsilon)} \right),$$

and its lower box-counting dimension $\underline{\dim}_B(\mathcal{G})$ by

$$\underline{\dim}_B(\mathcal{G}) = d - \limsup_{\varepsilon \rightarrow 0^+} \left(\frac{\log(\text{vol}(\mathcal{G}_\varepsilon))}{\log(\varepsilon)} \right).$$

If the two limits coincide, its box-counting dimension $\dim_B(\mathcal{G})$ exists and takes this value.

3 Excised holes

In this short section, we relate the ε -neighbourhood of \mathcal{G} to the so-called ε -inner neighbourhoods of the holes. This we do with two simple lemmas whose proofs can be safely omitted.

The first lemma formalises the intuitive, geometric description of \mathcal{G} as the limiting set of a process of excising holes, as fractals are often described informally. More precisely, it asserts that any hole is the image of one of the main holes under the T_i , which follows simply from the standard definition of \mathcal{G} as a limit set:

$$\mathcal{G} = \bigcap_{n=1}^{\infty} \bigcup_{|i|=n} \Delta_i,$$

and also that T_1, \dots, T_m are injective and their images $\Delta_1, \dots, \Delta_m$ have disjoint interiors.

Lemma 3.1. Every hole of \mathcal{G} is of the form ∇_i for some $i \in \mathcal{I}^*$. I.e.,

$$\Delta = \mathcal{G} \cup \bigcup_{i \in \mathcal{I}^*} \text{int}(\nabla_i), \tag{5}$$

where the unions are disjoint.

The second lemma requires the following simple definition.

Definition 3.2. For any $\Omega \subset \Delta$ and $\varepsilon > 0$, define its *inner ε -neighbourhood* $L_\varepsilon(\Omega)$ by

$$L_\varepsilon(\Omega) = \{x \in \Omega : \|x - y\| \leq \varepsilon \text{ for some } y \in \partial\Omega\},$$

the set of points in Ω at most ε away from the boundary $\partial\Omega$.

This second lemma is a natural consequence of the first.

Lemma 3.3. For all $\varepsilon > 0$,

$$\mathcal{G}_\varepsilon = \mathcal{G} \cup \bigcup_{i \in \mathcal{I}^*} L_\varepsilon(\nabla_i). \quad (6)$$

4 Simplices, inner neighbourhoods and inradii




Figure 5: The inner ε -neighbourhood $L_\varepsilon(\Omega)$ of triangle Ω , when $\varepsilon < \text{in}(\Omega)$.

In this section we make the inner neighbourhoods $L_\varepsilon(\nabla_i)$ explicit. This exploits the fact that an (injective) projective map on Δ maps any k -simplex onto another k -simplex. Hence, our assumption that the main holes are the interiors of d -simplices extends immediately to all holes, and we can apply the following lemmas. Their proofs, as with all geometric results, can be found in the appendix. In any case, we need a notion of inradius and incentre

The properties of the incentre of a simplex, described in the next lemma, are essential in the proofs of the two lemmas that follow.

Lemma 4.1. For any simplex Ω , there is a unique point $x \in \Omega$ furthest from $\partial\Omega$, which we call the *incentre* of Ω .

Our next result explicitly describes the inner neighbourhood of a simplex in terms of its incentre and inradius.

Definition 4.2. For any convex body $\Omega \subset \Delta$, let $\text{in}(\Omega)$ denote its *inradius*

$$\text{in}(\Omega) = \max \{ \|x - y\| : x \in \Omega, y \in \partial\Omega \}, \quad (7)$$

i.e., the radius of the largest ball contained within Ω .

Lemma 4.3. For any simplex Ω , $L_\varepsilon(\Omega)$ either equals Ω or is obtained by excising a similar copy of it:

$$L_\varepsilon(\Omega) = \begin{cases} \Omega, & \text{if } \varepsilon \geq \text{in}(\Omega), \\ \Omega \setminus \text{int} \left[\left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot \Omega \right], & \text{if } \varepsilon < \text{in}(\Omega), \end{cases} \quad (8)$$

where, for $\lambda > 0$, $\lambda \cdot \Omega$ denotes the similar copy to Ω obtained by enlarging Ω by a factor of λ about its incentre.

Remark 4.4. The second case of (8) is illustrated in Figure 5 when Ω is a triangle.

Our final result relates the inradius of a simplex to its volume and surface area as follows, generalising Heron's formula for the inradius of a triangle.

Lemma 4.5. For any non-degenerate d -simplex $\Omega \subset \Delta$,

$$\text{in}(\Omega) = d \cdot \frac{\text{vol}(\Omega)}{\text{per}(\Omega)}.$$

where $\text{per}(\Omega)$ denotes the $(d-1)$ -dimensional Lebesgue measure of the boundary $\partial\Omega$.

5 Laplace transform of $\text{vol}(\mathcal{G}_\varepsilon)$

Our analysis begins after the following simple consequence of Lemmas 3.3 and 4.3.

Proposition 5.1. For any d -simplex $\Omega \subset \Delta$,

$$\text{vol}(L_\varepsilon(\Omega)) = \text{vol}(\Omega) \left(1 - \max \left(0, 1 - \frac{\varepsilon}{\text{in}(\Omega)} \right)^d \right).$$

Consequently, since $\text{vol}(\mathcal{G}) = 0$,

$$\text{vol}(\mathcal{G}_\varepsilon) = \sum_{i \in \mathcal{I}^*} \text{vol}(L_\varepsilon(\nabla_i)) = \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \left(1 - \max \left(0, 1 - \frac{\varepsilon}{\text{in}(\nabla_i)} \right)^d \right).$$

The following lemma gives an explicit form for the Laplace transform

$$f(t) := \int_0^\infty e^{-tx} \text{vol}(\mathcal{G}_{\exp(-x)}) \, dx = \int_0^1 \varepsilon^{t-1} \text{vol}(\mathcal{G}_\varepsilon) \, d\varepsilon.$$

Lemma 5.2. For all $t > 0$,

$$f(t) = \frac{\text{vol}(\Delta)}{t} - \frac{d!}{t(t+1)(t+2)\cdots(t+d)} \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t.$$

In particular, f is analytic on the half-plane $\{z \in \mathbb{C} : \Re(z) > \sigma\}$, where

$$\sigma := \inf \left\{ t \geq -1 : \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t < \infty \right\}.$$




Figure 6: Sketch of $\varepsilon \mapsto \text{vol}(\nabla_i) \left(1 - \max\left(0, 1 - \frac{\varepsilon}{\text{in}(\nabla_i)}\right)^d\right)$

Proof. We prove the formula by direct computation, which we justify line-by-line afterwards.

$$f(t) = \int_0^1 \varepsilon^{t-1} \text{vol}(\mathcal{G}_\varepsilon) \, d\varepsilon \quad (9)$$

$$= \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \int_0^1 \varepsilon^{t-1} \left(1 - \max\left(0, 1 - \frac{\varepsilon}{\text{in}(\nabla_i)}\right)^d\right) \, d\varepsilon \quad (10)$$

$$= \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \left(\int_0^1 \varepsilon^{t-1} \, d\varepsilon - \int_0^{\text{in}(\nabla_i)} \varepsilon^{t-1} \left(1 - \frac{\varepsilon}{\text{in}(\nabla_i)}\right)^d \, d\varepsilon \right) \quad (11)$$

$$= \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \left(\frac{1}{t} - \text{in}(\nabla_i)^t \int_0^1 y^{t-1} (1-y)^d \, dy \right) \quad (12)$$

$$= \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \left(\frac{1}{t} - \frac{d!}{t(t+1)(t+2)\cdots(t+d)} \text{in}(\nabla_i)^t \right) \quad (13)$$

$$= \frac{\text{vol}(\Delta)}{t} - \frac{d!}{t(t+1)(t+2)\cdots(t+d)} \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t, \quad (14)$$

as required. As for the explanation:

- The first equality is the definition of f given above.
- The second uses Proposition 5.1 and exchanges integral and sum, using positivity.
- The third separates the two terms of the integral.
- The fourth uses the substitution $y = \varepsilon/\text{in}(\nabla_i)$.
- The fifth evaluates the Bernoulli integral

$$\int_0^1 y^{t-1} (1-y)^d \, dy = \frac{d!}{t(t+1)(t+2)\cdots(t+d)}.$$

- The final equality simply expands the bracket, noting that

$$\sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) = \text{vol}(\Delta),$$

which is equivalent to $\text{vol}(\mathcal{G}) = 0$.

As for the analytic continuation, we note that, decreasing along the real line, the singularity we first meet can only occur at $z = 0, -1, \dots, -d$, or at σ . It therefore suffices to show that there is no pole at 0 itself, assuming $\sigma \neq 0$: indeed, it is easy to see that the residues of the two terms at zero cancel perfectly in this case. \square

6 Proof of main theorem

We are now ready to complete the prove of our main theorem, Theorem 1.4, which we restate in our new notation below.

Theorem 6.1. *Under the assumptions of Theorem 1.4, $\overline{\dim}_B(\mathcal{G}) = d + \sigma$, i.e.,*

$$\overline{\dim}_B(\mathcal{G}) = d + \inf \left\{ t \geq -1 : \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t < \infty \right\}.$$

Proof. We prove it in two inequalities:

\otimes Writing $\overline{\dim}_B(\mathcal{G}) = r$, by definition,

$$\liminf_{\varepsilon \rightarrow 0^+} \frac{\log(\text{vol}(\mathcal{G}_\varepsilon))}{\log(\varepsilon)} = d - r.$$

I.e., for any $\delta > 0$, there exists $\varepsilon' > 0$ such that whenever $\varepsilon < \varepsilon'$,

$$\frac{\log(\text{vol}(\mathcal{G}_\varepsilon))}{\log(\varepsilon)} \geq d - r - \delta \quad \iff \quad \text{vol}(\mathcal{G}_\varepsilon) \leq \varepsilon^{d-r-\delta}.$$

Thus, whenever $t > r + \delta - d$, trivial splitting and estimation shows $f(t)$ is finite:

$$\begin{aligned} f(t) &= \int_0^{\varepsilon'} \varepsilon^{t-1} \text{vol}(\mathcal{G}_\varepsilon) \, d\varepsilon + \int_{\varepsilon'}^1 \varepsilon^{t-1} \text{vol}(\mathcal{G}_\varepsilon) \, d\varepsilon \\ &\leq \int_0^{\varepsilon'} \varepsilon^{-1+(t+d-r-\delta)} \, d\varepsilon + \text{vol}(\Delta) \int_{\varepsilon'}^1 \varepsilon^{t-1} \, d\varepsilon < \infty. \end{aligned}$$

Thus $t \geq \sigma$. The required inequality $\overline{\dim}_B(\mathcal{G}) \geq d + \sigma$ follows since t and δ are arbitrary.

\ominus We may assume $\sigma < 0$, since otherwise $\mathcal{G} \subset \Delta \implies \overline{\dim}_B(\mathcal{G}) \leq d$ trivially suffices. Given any $\tau \in (\sigma, 0)$, the sum

$$\sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^\tau$$

is finite for $z = \tau$, and hence uniformly bounded on the line $\{\Re(z) = \tau\}$. Consider the Laplace transform

$$\int_0^\infty \frac{d}{dx} \text{vol}(\mathcal{G}_{\exp(-x)}) \cdot e^{-tx} dx.$$

Integrating by parts [10, p.30], this transform equals

$$t \cdot (f(0^+) - f(t)) = \frac{d!}{(t+1) \cdots (t+d)} \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t.$$

Since the right hand side is integrable on the line $\{\Re(z) = \tau\}$ and has no singularities on or to the right of it, the Laplace inversion theorem [10, p.42ff] applies to give, for all $x > 0$,

$$\frac{d}{dx} \text{vol}(\mathcal{G}_{\exp(-x)}) = \frac{1}{2\pi i} \int_{\tau-i\infty}^{\tau+i\infty} \frac{d!}{(t+1) \cdots (t+d)} \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t e^{tx} dt \quad (15)$$

$$= e^{\tau x} \int_{\tau-i\infty}^{\tau+i\infty} \mathcal{O}\left(\frac{1}{(t+1)(t+2) \cdots (t+d)}\right) dt \quad (16)$$

$$= \mathcal{O}(e^{\tau x}). \quad (17)$$

In particular, since $\tau < 0$, integrating this inequality gives

$$\text{vol}(\mathcal{G}_{\exp(-x)}) = \mathcal{O}\left(\int_x^\infty e^{\tau y} dy\right) = \mathcal{O}(e^{\tau x}),$$

i.e., $\text{vol}(\mathcal{G}_\varepsilon) = \mathcal{O}(\varepsilon^{-\tau})$. Thus, by definition, $\overline{\dim}_B(\mathcal{G}) \leq d + \tau$, and we may take $\tau \rightarrow \sigma$. \square

7 A lower bound for $\overline{\dim}_B(\mathcal{G})$

In this section, we prove Theorem 1.8, affirming the conjecture of De Leo [11] under the geometric assumptions of our main theorem if one further assumes that $\dim_B \mathcal{G}$ exists.

The new notation makes the definition of ρ much easier to state, as we do below.

Theorem 7.1. *Under the assumptions of Theorem 1.4,*

$$\overline{\dim}_B(\mathcal{G}) \geq \max\left(d - 1, \frac{d \cdot \rho}{d + 1}\right),$$

where

$$\rho := \inf \left\{ r \geq 0 : \sum_{i \in \mathcal{I}} \|N_i\|^{-r} < \infty \right\} = \lim_{t \rightarrow \infty} \left(\frac{\log \#\{i \in \mathcal{I} : \|N_i\| \leq t\}}{\log t} \right).$$

The proof of this theorem uses the following useful formula for the action of projective maps on the volume of simplices.

Proposition 7.2. *Let $T : \Delta \rightarrow \Delta$ projectivise the matrix N with $\det(N) = \pm 1$. Then, for any d -simplex $\Omega \subset \Delta$,*

$$\frac{\text{vol}(T(\Omega))}{\text{vol}(\Omega)} = \prod_{e: \text{vertex of } \Omega} |N \cdot e|^{-1},$$

where $|\cdot|$ denotes the l^1 norm.

Proof. The case $\Omega = \Delta$ was proven in [27, Lem. 4.1], and we show how this more general case follows: Let $M \in \text{SL}_d(\mathbb{R})$ be such that its projectivisation maps the vertices of Δ onto those of Ω . Then by linearity, the $\Omega = \Delta$ case implies

$$\frac{\text{vol}(T(\Omega_i))}{\text{vol}(\Omega)} = \prod_{e: \text{vertex of } \Delta} \frac{|M \cdot e|}{|N \cdot M \cdot e|} = \prod_{e: \text{vertex of } \Delta} \left| N \cdot \left(\frac{M \cdot e}{|M \cdot e|} \right) \right|^{-1} = \prod_{e: \text{vertex of } \Omega} |N \cdot e|^{-1},$$

as required. \square

Proof of Theorem 1.7. We first prove the inequality involving the definition of ρ . For simplicity, we will use the symbol \lesssim to denote that an inequality holds up to multiplication by a fixed constant, depending at most on d and s . First observe that, for any d -simplex Ω , the volume of the largest contained ball cannot exceed the total volume: $\text{in}(\nabla_i) \lesssim \text{vol}(\nabla_i)^{1/d}$. Therefore, for any $t \in (-1, 0)$,

$$\begin{aligned} \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t &\gtrsim \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i)^{1+t/d} \\ &= \sum_{i \in \mathcal{I}} \sum_{k=1}^K \text{vol}(\nabla_k) \prod_{e: \text{vertex of } \nabla_k} |N_i \cdot e|^{-(1+t/d)} \\ &\geq \sum_{k=1}^K \text{vol}(\nabla_k) \cdot \sum_{i \in \mathcal{I}} \|N_i\|^{-(d+1)(1+t/d)}. \end{aligned}$$

Thus, recalling σ , the infimal value of $t \in (-1, 0)$ for which the left hand side converges, it satisfies

$$0 \geq \sigma \geq \inf \left\{ t \geq -1 : \sum_{i \in \mathcal{I}} \|N_i\|^{-(d+1)(1+t/d)} < \infty \right\} = \max \left(-1, \frac{d\rho}{d+1} - d \right),$$

and the inequality follows from Theorem 1.4.

To show the equivalence of the two definitions of ρ , let ρ denote the left hand side and λ the right hand side, whose limit exists by submultiplicativity. For any $\varepsilon > 0$,

$$\begin{aligned} \sum_{i \in \mathcal{I}} \|N_i\|^{-(\lambda+2\varepsilon)} &= \sum_{n=1}^{\infty} \sum_{\substack{i \in \mathcal{I}: \\ 2^{n-1} \leq \|N_i\| < 2^n}} \|N_i\|^{-(\lambda+2\varepsilon)} \\ &\leq \sum_{n=1}^{\infty} 2^{-n(\lambda+2\varepsilon)} \#\{i \in \mathcal{I} : \|N_i\| \leq 2^{-n}\} \\ &\lesssim \sum_{n=1}^{\infty} 2^{-n(\lambda+2\varepsilon)} 2^{n(\lambda+\varepsilon)} < \infty. \end{aligned}$$

Hence $\lambda \geq \rho$. The reverse inequality is similar, and completes the proof. \square

8 An upper bound for $\overline{\dim}_B(\mathcal{G})$ in the convex hole case

In this penultimate section we outline how, relaxing the assumptions so that the main holes may be convex polytopes, the proof of Theorem 1.4 can be extended to give an upper bound. Previously given as Theorem 1.7, it can be restated in our new notation as follows.

Theorem 8.1. *Under the assumptions of Theorem 1.7,*

$$\overline{\dim}_B(\mathcal{G}) \leq d + \inf \left\{ t \geq -1 : \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \text{in}(\nabla_i)^t < \infty \right\}.$$

The fundamental observation in the proof is that projectivised linear maps preserve convexity (so that the image of a convex polytope is a convex polytope), which can be shown directly from the definition. The main obstacle is that inner neighbourhoods of general convex bodies are more difficult to work with; we nevertheless obtain a general inclusion and inequality.

Note that, since there may be multiple points furthest from the boundary of a convex body Ω (e.g., when Ω the rectangle), we will denote any one of these points as an incentre.

Lemma 8.2. *For any convex body $\Omega \subset \Delta$,*

$$L_\varepsilon(\Omega) \subset \begin{cases} \Omega, & \text{if } \varepsilon \geq \text{in}(\Omega), \\ \Omega \setminus \text{int} \left[\left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot \Omega \right], & \text{if } \varepsilon < \text{in}(\Omega); \end{cases} \quad (18)$$

where, for $\lambda > 0$, $\lambda \cdot \Omega$ denotes the similar copy to Ω obtained by enlarging Ω by a factor of λ about any fixed incentre. Consequently,

$$\text{vol}(L_\varepsilon(\Omega)) \leq \text{vol}(\Omega) \left[1 - \max \left(0, 1 - \frac{\varepsilon}{\text{in}(\Omega)} \right) \right]^d$$

and

$$\text{vol}(\mathcal{G}_\varepsilon) \leq \sum_{i \in \mathcal{I}^*} \text{vol}(\nabla_i) \left[1 - \max \left(0, 1 - \frac{\varepsilon}{\text{in}(\nabla_i)} \right) \right]^d. \quad (19)$$

Proof. We defer the proof of the first inclusion to the appendix, and simply note that the inequalities are immediate, the second following by combining the first with 3.3 and the knowledge that all holes ∇_i are convex and the assumption that $\text{vol}(\mathcal{G}) = 0$. \square

From this point, the proof of the theorem continues analogously to that of Theorem 1.4, applying the results of section 5 and onwards to the right hand side of (19).

Finally, for completeness, we generalise the formula for the inradius of a simplex to an inequality. It is proven in the appendix.

Proposition 8.3. *For any non-empty convex set $\Omega \subset \Delta$,*

$$\frac{\text{vol}(\Omega)}{\text{per}(\Omega)} \leq \text{in}(\Omega) \leq (d-1) \frac{\text{vol}(\Omega)}{\text{per}(\Omega)}.$$

9 Application to the Rauzy gasket

In this final section, we prove the important results on the equality of Hausdorff and box-counting dimensions of the Rauzy gasket and go on to outline the proof of the numerical lower bound provided by Theorem 1.8

In this section, given a matrix N , we will denote by σ_j its j th largest singular value. The following result of [21] is reminiscent of singularity dimension, first introduced by Falconer.

Theorem 9.1 ([21]). *The Hausdorff dimension of the Rauzy gasket $\dim_H(\mathcal{R})$ is equal to the infimal value of $s \in (1, 2)$ for which the Poincaré series*

$$\sum_{i \in \mathcal{I}} \left(\frac{\sigma_2}{\sigma_1} \right) \left(\frac{\sigma_3}{\sigma_1} \right)^{s-2} (N_i) \quad (20)$$

converges.

Note that we will sometimes, for convenience, denote the summand of (20) by $\phi_s(N_i)$.

We will apply this result by showing the following.

Lemma 9.2. *For any $s \in (1, 2)$,*

$$\sum_{i \in \mathcal{I}^*} \text{area}(\nabla_i) \text{in}(\nabla_i)^{s-2} \lesssim \sum_{i \in \mathcal{I}} \left(\frac{\sigma_2}{\sigma_1} \right) \left(\frac{\sigma_3}{\sigma_1} \right)^{s-2} (N_i).$$

Thus combining this lemma with Theorem 1.4 and the above result, we see that $\overline{\dim}_B(\mathcal{R}) \leq \dim_H(\mathcal{R})$, immediately giving Theorem 1.9.

Proof of Lemma 9.2. Following [21], we note that the Euclidean distance between $x, y \in \Delta$, $\|x - y\|$ equals, up to a constant factor, the area of the triangle formed by x, y and the origin:

$$\|x - y\| \asymp \|x \wedge y\|,$$

where \asymp denotes that the two sides are comparable, i.e. their ratio is bounded away from 0 and ∞ , by constants depending at most on d and s . Heron's formula (Lemma 4.5) gives, for $s \in (1, 2)$,

$$\sum_{i \in \mathcal{I}^*} \text{area}(\nabla_i) \text{in}(\nabla_i)^{s-2} \asymp \sum_{i \in \mathcal{I}^*} \text{area}(\nabla_i) \left(\frac{\text{per}(\nabla_i)}{\text{area}(\nabla_i)} \right)^{2-s}. \quad (21)$$

Now, given $i \in \mathcal{I}^*$, applying the area formula given in Proposition 7.2,

$$\begin{aligned} \frac{\text{per}(\nabla_i)}{\text{area}(\nabla_i)} &\asymp \sum_{\{j,k,l\}=\{1,2,3\}} \|T_i(e_j) \wedge T_i(e_k)\| \cdot |N_i \cdot e_1| |N_i \cdot e_2| |N_i \cdot e_3| \\ &= \sum_{\{j,k,l\}=\{1,2,3\}} \|N_i \cdot e_j \wedge N_i \cdot e_k\| \cdot |N_i \cdot e_l| \\ &\lesssim 3 \sigma_1 \sigma_2 \cdot \sigma_1(N_i) = \sigma_1^2 \sigma_2(N_i). \end{aligned}$$

Hence, substituting into (21),

$$\sum_{i \in \mathcal{I}^*} \text{area}(\nabla_i) \text{in}(\nabla_i)^{s-2} \lesssim \sum_{i \in \mathcal{I}^*} \text{area}(\nabla_i) \cdot \sigma_1^{4-2s} \sigma_2^{2-s}(N_i). \quad (22)$$

To complete the proof, it will suffice to show $\text{area}(\nabla_i) \lesssim \sigma_1^{-3}(N_i)$ for all $i \in \mathcal{I}^*$, since then the right hand side of (22) will be dominated by

$$\sum_{i \in \mathcal{I}^*} \sigma_1^{-3} \sigma_1^{4-2s} \sigma_2^{2-s}(N_i) = \sum_{i \in \mathcal{I}^*} \left(\frac{\sigma_2}{\sigma_1} \right) \left(\frac{\sigma_3}{\sigma_1} \right)^{s-1} (N_i) \asymp \sum_{i' \in \mathcal{I}} \left(\frac{\sigma_2}{\sigma_1} \right) \left(\frac{\sigma_3}{\sigma_1} \right)^{s-1} (N_{i'}), \quad (23)$$

where the first inequality uses that $\det(N_i) = 1 = \sigma_1 \sigma_2 \sigma_3(N_i)$ for any $i \in \mathcal{I}^*$ and the second relation is justified by term-by-term comparison: for any $i' \in \mathcal{I}$, we have, for its corresponding $i \in \mathcal{I}^*$, $N_i = N_{i'} \cdot M$, where M is a non-singular matrix to be made explicit below. Since M is non-singular and $\det(N_i) = \det(N_{i'}) = 1$,

$$\sigma_1(N_i) \asymp \sigma_1(N_{i'}), \quad \sigma_3(N_i) \asymp \sigma_3(N_{i'}) \implies \sigma_2(N_i) = \frac{1}{\sigma_1 \sigma_3}(N_i) \asymp \frac{1}{\sigma_1 \sigma_3}(N_{i'}) = \sigma_2(N_{i'})$$

and hence $\phi(N_i) \asymp \phi(N_{i'})$, showing that the two series in (23) are comparable.

We prove the assertion $\text{area}(\nabla_i) \lesssim \sigma_1^{-3}(N_i)$ in three cases, writing $N = N_i$ for simplicity:

Case 1: $i = (i', \cdot) \in \mathcal{I}^*$, where $i' \in \mathcal{I}$ contains all three symbols.

Without loss of generality, write $N = A \cdot B$, where

$$B = N_1 \cdot N_2^a \cdot N_3^b \cdot M$$

for some $a, b \in \mathbb{N}$, fixing M to be the matrix with the following explicit form:

$$M = \begin{pmatrix} 0 & \alpha & \alpha \\ \alpha & 0 & \alpha \\ \alpha & \alpha & 0 \end{pmatrix},$$

where $\alpha = 2^{-1/3}$, so that $\det(M) = 1$. More explicitly,

$$B = \alpha \begin{pmatrix} ab + a + b + 2 & ab + 2a + b + 2 & 2ab + a + 2b + 2 \\ ab + a + 1 & ab + 2a & 2ab + a + 1 \\ b + 1 & b + 1 & 2b \end{pmatrix}.$$

We can see that the ratio of any two elements in a row of B is at most two:

$$\max_{j,k,l} \left(\frac{B_{j,k}}{B_{j,l}} \right) \leq 2,$$

independently of a and b . Applying this directly gives, for $\lambda = (1, 1, 1) \cdot A$ gives

$$\frac{|N \cdot e_k|}{|N \cdot e_l|} = \frac{\lambda \cdot B \cdot e_l}{\lambda \cdot B \cdot e_k} = \frac{\sum_{j=1}^3 \lambda_j B_{j,l}}{\sum_{j=1}^3 \lambda_j B_{j,k}} \leq \frac{\sum_{j=1}^3 \lambda_j (2B_{j,k})}{\sum_{j=1}^3 \lambda_j B_{j,k}} \leq 2.$$

Applying this in the area formula (Proposition 7.2) finishes the proof in this case:

$$\text{area}(\nabla_i) \asymp \prod_{j=1}^3 |N \cdot e_j|^{-1} \leq 4 \left(\max_j |N \cdot e_j| \right)^{-3} = 4 \cdot \|N\|_{l^1 \rightarrow l^1}^{-3} \asymp \sigma_1^{-3}(N), \quad (24)$$

where $\|N_i\|_{l^1 \rightarrow l^1}$ denotes the l^1 operator norm of N_i , using equivalence of norms at the end.

Case 2: $i = (i', \cdot) \in \mathcal{I}^*$; $i' \in \mathcal{I}$ contains precisely two symbols.

We assume symbols 1 and 2 appear in i' (in that order), and claim that N is of the form

$$\alpha \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ 1 & 1 & 0 \end{pmatrix},$$

where x_j, y_j are positive integers such that

$$\max_{j,k} \left(\frac{x_j}{x_k} \right) < 3 \quad \text{and} \quad \max_{j,k} \left(\frac{y_j}{y_k} \right) < 3.$$

This follows by a simple induction. As a base case, for $b \in \mathbb{N}$ consider the matrix

$$N_1 N_2^b M = \alpha \begin{pmatrix} b+2 & 2b+1 & b+1 \\ b+1 & 2b & b+1 \\ 1 & 1 & 0 \end{pmatrix},$$

which is of the required form. Inductively assuming that N is of the above form,

$$\begin{aligned} N_1 N &= \alpha \begin{pmatrix} x_1 + y_1 + 1 & x_2 + y_2 + 1 & x_3 + y_3 \\ y_1 & y_2 & y_3 \\ 1 & 1 & 0 \end{pmatrix}, \\ N_2 N &= \alpha \begin{pmatrix} x_1 & x_2 & x_3 \\ x_1 + y_1 + 1 & x_2 + y_2 + 1 & x_3 + y_3 \\ 1 & 1 & 0 \end{pmatrix}, \end{aligned}$$

which again are easily shown to be of the required form, proving the claim.

Thus we may write, for any column indices l and k ,

$$\frac{|N \cdot e_l|}{|N \cdot e_k|} = \frac{x_l + y_l + \delta_{\{l \neq 3\}}}{x_k + y_k + \delta_{\{k \neq 3\}}} \leq \frac{(3x_k - 1) + (3y_k - 1) + 1}{x_k + y_k} < 3.$$

Hence an analogue of (24) applies to show that $\text{area}(\nabla_i) \lesssim \sigma_1^{-3}(N)$ (with a larger implied constant).

Case 3: $i = (i', \cdot) \in \mathcal{I}^*$; $i' \in \mathcal{I}$ contains only one symbol. Assuming without loss that $i' = (1, \dots, 1) \in \{1\}^b$ for $b \in \mathbb{N}_0$, we explicitly write

$$N = N_1^b M = \alpha \begin{pmatrix} 2b & b+1 & b+1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

Hence $\max_{k,l} \left(\frac{|N \cdot e_k|}{|N \cdot e_l|} \right) = \frac{2b+2}{b+2} \leq 2$ and (24) applies as before to complete the proof. \square




Figure 7: Enlarging ball $B \subset \Omega$ to meet side S .

In order to outline the proof of Theorem 1.10, note that the above proof, particularly (24), can be altered to show more strongly that $\text{area}(\Delta_i) \asymp \text{area}(\nabla_{i'}) \asymp \|N_i\|^{-3}$ in the above notation.

This observation allows us to relate the theoretical lower bound in Theorem 1.8 to a numerical lower bound as per the following result. In particular, the author in [30] uses renewal theory and Mathematica to give the upper and lower bounds. See [31] for the code used and an in-depth explanation thereof. This altogether gives the following.

Theorem 9.3 ([30]).

$$\inf \left\{ \delta > 0 : \sum_{i \in \mathcal{I}} \|N_i\|^{-3\delta} < \infty \right\} = \inf \left\{ \delta > 0 : \sum_{i \in \mathcal{I}} \text{area}(\Delta_i)^\delta < \infty \right\} \in [0.8098, 0.8204].$$

Hence, by Theorems 1.8 and 1.9, $\dim_H(\mathcal{R}) \geq 2 \times 0.8098 = 1.6196$, as claimed.

A Geometric results

In this appendix, we give elementary and complete proofs of our geometric results in two parts. The first concerns simplices, and the second some generalisations to complex bodies. We will state and prove them in the context of \mathbb{R}^n for simplicity.

A.1 Inradii and incentres of simplices

In this subsection, let $\Omega \subset \mathbb{R}^n$ denote a given non-degenerate n -simplex.

One key property of simplices (compared to general convex bodies) is that the incentre is uniquely defined, as per the following simple but important lemma.

Lemma A.1. *There is a unique point in Ω furthest from the boundary $\partial\Omega$ (the incentre), which is equidistant from all faces of Ω .*

Proof. Let B be a ball contained in Ω , and suppose it does not meet side $S \subset \partial\Omega$. Then, letting e denote the opposing vertex, one can show the enlargement of B about e with scale

$$\frac{d(e, S)}{d(e, S) - d(B, S)} :$$

see Figure 7, is larger and contained in Ω . Thus, a maximal ball in Ω touches all of its sides, i.e., any point which is furthest from $\partial\Omega$ is equidistant from all sides.

We further claim there can be only one point equidistant from all sides: Since Ω is non-degenerate, the normal vectors to each face are linearly independent: thus, if there were two such distinct points, moving from one to the other would increase distance to some side and decrease distance to another, a contradiction. \square

The incentre provides a convenient decomposition of Ω which we use in the next two proofs. We begin with the simpler of the two; the formula for the inradius.

Lemma A.2.

$$\text{in}(\Omega) = n \frac{\text{vol}(\Omega)}{\text{per}(\Omega)}.$$

Thanks are due to András Sandor, who produced the proof idea in about a minute!

Proof. For a given side $S \subset \partial\Omega$, let Ω_S denote the set of points in Ω whose closest side is S . We know that Ω_S is bounded between S itself and each of the hyperplanes bisecting S and one of the other faces — i.e., the set of points equidistant to these two sides. By the previous lemma, these hyperplanes must all meet at the incentre of Ω , thus Ω_S is also an n -simplex. Taking S as the base of Ω_S , it has height $\text{in}(\Omega)$, and hence volume

$$\text{vol}(\Omega_S) = \frac{1}{n} \text{vol}_{n-1}(S) \cdot \text{in}(\Omega).$$

The formula follows by summing over S . \square

Next we prove the important expression for the inner ε -neighbourhood.


Lemma A.3. For all $\varepsilon > 0$,

$$L_\varepsilon(\Omega) = \begin{cases} \Omega, & \text{if } \varepsilon \geq \text{in}(\Omega), \\ \Omega \setminus \text{int} \left[\left(1 - \frac{\varepsilon}{\text{in}(\Omega)} \right) \cdot \Omega \right], & \text{if } \varepsilon \leq \text{in}(\Omega); \end{cases}$$

where $x \mapsto \lambda \cdot x$ denotes enlargement, scale factor λ , about the incentre of Ω .

Proof. It suffices to show, given $\varepsilon < \text{in}(\Omega)$, that

$$\{x \in \Omega \mid d(x, \partial\Omega) \geq \varepsilon\} = \left(1 - \frac{\varepsilon}{\text{in}(\Omega)} \right) \cdot \Omega \quad (25)$$

Figure 8: The action of $x \mapsto \lambda \cdot x$ on Ω_S , for $\lambda \in (0, 1)$.

and

$$\{x \in \Omega \mid d(x, \partial\Omega) = \varepsilon\} = \left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot \partial\Omega. \quad (26)$$

Moreover, recalling the notation from the previous proof, $x \mapsto \left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot x$ maps each Ω_S into itself, since it is a linear contraction towards one of the vertices of Ω_S . Hence, we only need to prove (25) and (26) with Ω_S in place of Ω and S in place of $\partial\Omega$: i.e.,

$$\{x \in \Omega_S \mid d_S(x) \geq \varepsilon\} = \left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot \Omega_S,$$

where $d_S := d(x, S)$ for $x \in \Omega$. Indeed, by linearity, for any $x \in \Omega_S$,

$$d_S \left(\left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot x \right) - \text{in}(\Omega) = \left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) (d_S(x) - \text{in}(\Omega))$$

(see Figure 8), which rearranges to

$$d_S \left(\left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot x \right) = \left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) d_S(x) + \varepsilon \geq \varepsilon, \quad (27)$$

with equality iff $x \in S$. Moreover, we may extend d_S linearly to the cone C_S based at the incentre and containing S ,³ so that $d_S(x) < 0$ for $x \in C_S \setminus \Omega_S$. The equality in (27) extends to $x \in C_S$ by linearity, and thus shows, for $x \in C_S$,

$$d_S \left(\left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot x \right) \geq \varepsilon \quad \text{if and only if} \quad d_S(x) \geq 0 \iff x \in \Omega.$$

Taking the union over sides S gives (25) and (26); thus the lemma follows. \square

³Equivalently, C_S can be defined as those points $x \in \mathbb{R}^n$ whose closest point in $\partial\Omega$ lies on S .

A.2 Inradii and incentres of convex bodies

We now fix $\Omega \in \mathbb{R}^n$ to be a convex body (i.e., a closed convex set with non-empty interior).

We first give the proof of the inclusion (18), which is surprisingly simple.

Lemma A.4.

$$L_\varepsilon(\Omega) \subset \begin{cases} \Omega, & \text{if } \varepsilon \geq \text{in}(\Omega), \\ \Omega \setminus \text{int} \left[\left(1 - \frac{\varepsilon}{\text{in}(\Omega)}\right) \cdot \Omega \right], & \text{if } \varepsilon < \text{in}(\Omega); \end{cases}$$

where $x \mapsto \lambda \cdot x$ denotes enlargement, scale factor λ , about a fixed incentre of Ω .

Proof. We first show $f : x \mapsto d(x, \partial\Omega)$ is concave on Ω , as follows. Suppose that B_0, B_1 are the largest balls in Ω based respectively at $x_0, x_1 \in \Omega$ (i.e., with radii $f(x_0)$ and $f(x_1)$). Then, for any $\lambda \in (0, 1)$,

$$B_\lambda := \lambda B_1 + (1 - \lambda)B_0$$

is a ball based at $x_\lambda := \lambda x_1 + (1 - \lambda)x_0$, with radius $\lambda f(x_1) + (1 - \lambda)f(x_0)$, contained wholly in Ω by convexity. Hence,

$$f(x_\lambda) \geq \lambda f(x_1) + (1 - \lambda)f(x_0),$$

as claimed. Therefore, given $y \in \text{int}(\Omega)$ and $\varepsilon < f(y)$, contracting $x \in \Omega$ towards y by a factor $\left(1 - \frac{\varepsilon}{f(y)}\right)$ takes it outside of the ε -inner neighbourhood:

$$f\left(\left(\frac{\varepsilon}{f(y)}\right)y + \left(1 - \frac{\varepsilon}{f(y)}\right)x\right) \geq \left(\frac{\varepsilon}{f(y)}\right)f(y) + \left(1 - \frac{\varepsilon}{f(y)}\right)f(x) \geq \varepsilon.$$

Thus taking y such that $f(y) = \text{in}(\Omega)$ gives the second case of (18). (The first is trivial.) \square




Figure 9: The decompositions of convex polygon Ω into $\{\Omega_S\}$ and $\{\Delta_S\}$. The largest inscribed ball is depicted in pastel yellow.

We finally prove the generalisation of Heron's formula, as below. Both bounds are in fact tight.

Proposition A.5.

$$\frac{\text{vol}(\Omega)}{\text{per}(\Omega)} \leq \text{in}(\Omega) \leq n \frac{\text{vol}(\Omega)}{\text{per}(\Omega)}.$$

Since any convex body can be approximated arbitrarily well by convex polytopes, we will prove the result assuming Ω is a polytope. The proof then relies on decomposing Ω in two different ways, illustrated in Figure 9, which coincide when Ω is a simplex.

Proof of lower bound. Consider one side $S \subset \partial\Omega$ of Ω , and recall the locus Ω_S of points in Ω whose closest side of Ω is S . Treating $S \subset \mathbb{R}^{n-1} \times \{0\}$ and $\Omega \subset \mathbb{R}^{n-1} \times [0, \infty)$, we see that

- every point in Ω_S can be at most $\text{in}(\Omega)$ distance from S , and
- the locus is bound by the hyperplanes bisecting S and each of the neighbouring sides:

by convexity, the interior angle between any two sides lies in $(0, \pi)$, and thus any bisector with S makes an interior angle at most $\pi/2$ with S : i.e., is either vertical or sloping inwards towards S . Hence $\Omega_S \subset S \times [0, \text{in}(\Omega)]$ and thus

$$\text{vol}(\Omega_S) \leq \text{vol}_{n-1}(S) \cdot \text{in}(\Omega).$$

Summing over S gives the required inequality. □

Proof of upper bound. Choose an incentre $x \in \Omega$. Fix side S , and let Δ_S denote the cone with base S and vertex x . Again take $S \subset \mathbb{R}^{n-1} \times \{0\}$ and $\Omega \subset \mathbb{R}^{n-1} \times [0, \infty)$ for simplicity. Since the ball of radius $\text{in}(\Omega)$ based at x is contained in $\Omega \subset \mathbb{R}^{n-1} \times [0, \infty)$, we must have $x \in \mathbb{R}^{n-1} \times [\text{in}(\Omega), \infty)$. Therefore, since Δ_S is a cone with base S and height at least $\text{in}(\Omega)$,

$$\frac{1}{n} \text{vol}_{n-1}(S) \cdot \text{in}(\Omega) \leq \text{vol}(\Delta_S).$$

The proof is finished by summing over S . □

References

- [1] P. Arnoux, and G. Rauzy. Représentation géométrique de suites de complexité $2n + 1$. *Bull. Soc. Math. France* **119** (1991), no. 2, 199–215. doi:10.24033/bsmf.2164
- [2] P. Arnoux and Š. Starosta (2013). *The Rauzy gasket*. In: J. Barral and S. Seuret (eds). Further developments in fractals and related fields, 1–23. Birkhäuser/Springer: New York. doi:10.1007/978-0-8176-8400-6
- [3] P. Arnoux, and J.-C. Yoccoz. Construction de difféomorphismes pseudo-Anosov. *C. R. Acad. Sci. Paris Sér. I Math.* **292** (1981), no. 1, 75–78.
- [4] A. Avila, P. Hubert and A. Skripchenko. On the Hausdorff dimension of the Rauzy gasket. *Bull. Soc. Math. France* **144** (2016), no. 3, 539–568. doi:10.24033/bsmf.2722

- [5] Z.-Q. Bai and S. R. Finch. Precise calculation of Hausdorff dimension of Apollonian gasket. *Fractals* **26** (2018), no. 4, 1850050–1850059. doi:10.1142/S0218348X18500500
- [6] A. Baragar. The exponent for the Markoff-Hurwitz equations. *Pacific J. Math.* **182** (1998), no.1, 1–21. doi:10.2140/pjm.1998.182.1
- [7] M. F. Barnsley and A. Vince. Real projective iterated function systems. *J. Geom. Anal.* **22** (2012), no. 4, 1137–1172.
- [8] D. W. Boyd. The residual set dimension of the Apollonian packing. *Mathematika* **20** (1973), 170–174. doi:10.1112/S0025579300004745
- [9] A. Christodoulou and N. Jurga. The Hausdorff dimension of self-projective sets. To appear: *Trans. Am. Math. Soc.*
- [10] B. Davies (2002). *Integral transforms and their applications*. 3rd edition. New York: Springer.
- [11] R. De Leo. Exponential growth of norms in semigroups of linear automorphisms and Hausdorff dimension of self-projective IFS. *J. of Geom. Anal.* **25** (2015) no. 3, 1798–1827. doi:10.1007/s12220-014-9494-1
- [12] R. De Leo. A conjecture on the Hausdorff dimension of attractors of real self-projective iterated function systems. *Exp. Math.* **24** (2015), no. 3, 270–288. doi:10.1080/10586458.2014.987884
- [13] R. De Leo and I. A. Dynnikov. Geometry of plane sections of the infinite regular skew polyhedron $\{4, 6 \mid 4\}$. *Geom. Dedicata* **138** (2009), 51–67. doi:10.1007/s10711-008-9298-1
- [14] I. A. Dynnikov, P. Hubert and A. Skripchenko. Dynamical systems around the Rauzy gasket and their ergodic properties. *Int. Math. Research Notices* **rnac440** (2022). arXiv:2011.15043
- [15] K. J. Falconer (1990). *Fractal Geometry: Mathematical Foundations and Applications*. 1st edition. John Wiley & Sons: Chichester.
- [16] C. Fougeron. *Dynamical properties of simplicial systems and continued fraction algorithms* (2020). arXiv:2001.01367v3
- [17] E. Fuchs. Counting problems in Apollonian packings. *Bull. Amer. Math. Soc.* **50** (2013), no. 2, 229–266. doi:10.1090/S0273-0979-2013-01401-0
- [18] A. Gamburd, M. Magee and R. Ronan. An asymptotic formula for integer points on Markoff-Hurwitz varieties. *Ann. Math. (2)* **190** (2019), no. 3, 751–809. doi:10.4007/annals.2019.190.3.2
- [19] R. Gutiérrez-Romo and C. Matheus. Lower bounds on the dimension of the Rauzy gasket. *Bull. Soc. Math. France* **148** (2020), no. 2, 321–327. doi:10.24033/bsmf.2807

- [20] P. Hubert and O. Paris-Romaskevich. Triangle tiling billiards and the exceptional family of their escaping trajectories: circumcenters and Rauzy gasket. *Exp. Math.* (2019). doi:10.1080/10586458.2019.1583615
- [21] Y. Jiao, J. Li, W. Pan and D. Xu. *On the dimension of limit sets on $\mathbb{P}(\mathbb{R}^3)$ via stationary measures: the theory and applications*. Preprint, available at <https://sites.google.com/view/jialunli> [accessed 18/09/2023].
- [22] A. Kontorovich and H. Oh. Apollonian circle packings and closed horospheres on hyperbolic 3-manifolds. *J. Am. Math. Soc.* **24** (2011), 603–648. doi:10.1090/S0894-0347-2011-00691-7
- [23] G. Levitt. La dynamique des pseudogroupes de rotations. *Invent. Math.* **113** (1993), no. 3, 633–670. doi:10.1007/BF01244321
- [24] A. Maltsev and S. P. Novikov. Dynamical systems, topology, and conductivity in normal metals. *J. Stat. Phys.* **115** (2004), no. 1–2, 31–46.
- [25] C. T. McMullen. Hausdorff Dimension and Conformal Dynamics, III: Computation of Dimension. *Am. J. Math.* **120** (1998), 691–721. doi:10.1353/ajm.1998.0031
- [26] M. Pollicott. *Apollonian Circle Packings*. In: Bandt, C., Falconer, K., Zähle, M. (eds) *Fractal Geometry and Stochastics V*, 121–142. Birkhäuser: Cham, 2015. doi:10.1007/978-3-319-18660-3_8
- [27] M. Pollicott and B. Sewell. An elementary proof that the Rauzy gasket is fractal. To appear: *Ergodic Theory and Dynamical Systems*. doi:10.1017/etds.2023.66
- [28] M. Pollicott and B. Sewell. An upper bound on the dimension of the Rauzy gasket. To appear: *Bull. Soc. Math. France*.
- [29] W. Sierpiński. Sur une courbe dont tout point est un point de ramification. *C. R. Acad. Sci. Paris* **160** (1915), 302–305.
- [30] B. Sewell. *Equidistribution of infinite interval substitution schemes, explicit resonances of Anosov toral maps, and the Hausdorff dimension of the Rauzy gasket*. PhD thesis: <https://wrap.warwick.ac.uk/164862/>
- [31] B. Sewell. Code repository: <https://github.com/sewell-math/rauzy-gasket-area>
- [32] B. Solomyak and Y. Takahashi. Diophantine property of matrices and attractors of projective iterated function systems in \mathbb{RP}^1 . *Int. Math. Res. Not. IMRN* (2021), no.16, 12639–12669.