A cyclic random motion in \mathbb{R}^3 driven by geometric counting processes

Antonella Iuliano
1*† and Gabriella Verasani
1†

^{1*}Department of Mathematics, Computer Science and Economics,, University of Basilicata, Viale dell'Ateneo Lucano, 10, Potenza, 85100, PZ, Italy.

*Corresponding author(s). E-mail(s): antonella.iuliano@unibas.it; Contributing authors: gabriella.verasani@unibas.it; †These authors contributed equally to this work.

Abstract

We consider the random motion of a particle that moves with constant velocity in \mathbb{R}^3 . The particle can move along four directions with different speeds that are attained cyclically. It follows that the support of the stochastic process describing the particle's position at time t is a tetrahedron. We assume that the sequence of sojourn times along each direction follows a geometric counting process (GCP). When the initial velocity is fixed, we obtain the explicit form of the probability law of the process $X(t) = (X_1(t); X_2(t); X_3(t)), t > 0$, for the particle's position. We also investigate the limiting behavior of the related probability density when the intensities of the four GCPs tend to infinity. Furthermore, we show that the process does not admit a stationary density. Finally, we introduce the first-passage-time problem for the first component of X(t) through a constant positive boundary $\beta > 0$ providing the bases for future developments.

 ${\bf Keywords:}$ Counting process, Finite speed, First-passage time, Random motion, Random evolution

 $\mathbf{MSC\ Classification:}\ 60\mathrm{K}99\ ,\ 60\mathrm{K}50$

ORCID: Antonella Iuliano: Orcid 0000-0001-8541-8120 Gabriella Verasani: Orcid 0009-0000-4994-1694

1 Introduction

In the last decades, finite-velocity random motions have been widely studied as a natural class of stochastic processes to model real phenomena on multi-dimensional Euclidean spaces. In the one-dimensional case, the model that describe these motions is the telegraph process, in which the changes of the two possible velocities are governed by the Poisson process. This process has been studied in (Orsingher, 1990), Kolesnik (1998), Beghin et al. (2001) and Kolesnik and Ratanov (2023). Later, the study of random motions in two or more dimensions has been performed by many authors over the years. In particular, finite-velocity planar random processes in continuous time have been investigated by (Orsingher, 1986) and Masoliver et al. (1993). Further investigations have been oriented to investigate the distribution for Markovian random motion in the plane (c.f. Kolesnik and Turbin (1998) and Kolesnik (2006)), and the random motion with possible reflections at Poisson-paced events (c.f. Kolesnik and Orsingher (2002)). The analysis of these random motions with finite speed is performed analytically by solving partial differential equations. Other methods use probabilistic approaches based on order statistics, or on more general renewal processes as in Di Crescenzo (2002) (with arbitrary random steps between successive switches).

Several generalizations of the basic model have been proposed especially in biology and physics, motivated by the need of describing a variety of random motions performed by cells, micro-organisms and animals. These problems have been analyzed in Martens *et al.* (2012), Hartmann *et al.* (2020), Santra *et al.* (2020) and Pogorui and Rodríguez-Dagnino (2021).

The main aim for the study of finite-velocity random motions in one or more dimensions is the determination of the probability distribution of the position vector X(t) of a particle at time t. For instance, cyclic planar motions with three and four directions have been treated by Orsingher (2002) and Orsingher and Sommella (2004) in which the changes of direction are governed by a homogenous Poisson process. Moreover, the authors have found a connection between the equations governing the probability distributions and the Bessel functions of higher order. The probability law of the motion of a particle performing a cyclic random motion is determined in Lachal (2006) where the particle can take a finite number of possible directions with different speeds. Here, the changes of direction occur at Poisson random times. For instance, see also Lachal et al. (2006) in which the probability distribution is obtained by using order statistics and is expressed in terms of hyper-Bessel functions of higher order. Recently, planar random motions with orthogonal directions switching at Poisson time have been examined in Orsingher et al. (2020), Cinque and Orsingher (2022) and Cinque and Orsingher (2023). Other remarkable results concerning random motions in high dimensional spaces with uniformly distributed directions are present in Orsingher and De Gregorio (2007), Pogorui (2012), and Pogorui and Rodríguez-Dagnino (2011).

Some modified versions of the telegraph process have been considered also by substituting the underlying Poisson process with different kinds of counting processes. For instance, in Di Crescenzo *et al.* (2023) the authors study a stochastic process which describes the dynamics of a particle performing a finite-velocity random motion in \mathbb{R} and \mathbb{R}^2 , whose velocities alternate cyclically along two and three different directions, respectively, with possibly unequal velocities. The novelty of this last paper is that the

number of displacements of the motion along each possible direction follows a Geometric Counting Process (GCP) (see Cha and Finkelstein (2013)). This new approach has the advantageous characteristic that consists in the possibility of describing phenomena whose interarrival times have heavy tails rather than the memoryless property, as observed often in real cases. Specifically, the properties of the interarrival times of GCP's are declined in several areas, such as in software reliability or actuarial theory, where such counting processes are often considered to describe occurrences of shocks or claims, and in earth sciences (climatology, hydrology, etc) to model the failures along times. For instance, some examples of applied fields where GCPs can be used are discussed in Section 6 and 7 of Di Crescenzo and Pellerey (2019).

Therefore, starting from the results obtained in Di Crescenzo et al. (2023), here, we propose an extension of the classical telegraph process in \mathbb{R}^3 with the aim of (i) determining the explicit general laws of the distribution of the current position, and (ii) presenting an approach of resolution based on the study of the intertimes between consecutive changes of speed. In particular, we assume that the motion proceeds along four velocities that alternate cyclically, where the intertimes between two subsequent velocity changes toward each direction are distributed as a GCP. The use of a limited number of speeds is justified by the results presented in Orsingher and Sommella (2004) where the authors consider a cyclic random motion with four directions forming a regular tetrahedron in \mathbb{R}^3 but where the changes of speed are governed by a homogenous Poisson process. Precisely, in our case the intertimes possess a heavy-tailed distribution, differently to the classical telegraph process in which the times between velocity changes have exponential distribution. The scheme of the present paper is useful in describing suitable dynamics where random occurrences between two events have infinite expectations, and are dependent. Thus, the proposed stochastic process provides new models for the description of phenomena that are no more governed by hyperbolic PDE's as the classical telegraph equation. Another benefit of the present study is the construction of new solvable models, whose probability laws are obtained in tractable and closed form (as in the special case with symmetry) using an approach based on the analysis of the intertimes between consecutive velocity changes. Note that the introduction of these finite-velocity random motions is also useful to describe the movement of a particle that chooses the new direction among all the possible ones in a cyclical manner.

Moreover, a further strength of this work concerns some interesting asymptotic results. In particular, we are able to obtain the limiting density of the process in a closed form when the parameters of the intertimes between velocity changes tend to infinity. In addition, we show that the process does not admit a stationary density as t goes to $+\infty$. All these motivations allow us to study the motion in \mathbb{R}^3 because it can be applied to concrete situations and the amount of calculations needed is comparatively acceptable. For instance, we mention that these models can approximate the random motion of some physical models linked to the movement of particles in the environment, birds and animals in the ocean, flying objects or users in a shopping mall.

In detail, we analyze the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ in \mathbb{R}^3 which describes the position of randomly moving particle performing a cyclic alternating motion with four specific and possible different directions \vec{v}_j , for $1 \leq j \leq 4$. The velocity-vectors form

a (possibly irregular) tetrahedron, say $\mathcal{T}(t)$, i.e., the set of all possible positions of the moving particle at time t>0 on the surface of the support in \mathbb{R}^3 . Note that the analysis of random motions in multidimensional Euclidean spaces is quite rare in the literature, since its analysis is rather difficult. Hence, to overcome the difficulties of the study we will refer in detail to the simpler case in which the region $\mathcal{T}(t)$ is regular, centered in the origin and with vertices placed on the coordinate axes. Once defined the geometry of the region $\mathcal{T}(t)$, the probability law of the process is determined when (i) the initial components are given by three terms, describing the situations in which the particle is found on the vertices, edges and faces of $\mathcal{T}(t)$, at the beginning of its motion, and (ii) the density concerning the absolutely continuous part is related to the motion of the particle in the interior of $\mathcal{T}(t)$. In particular, for the latter absolutely continuous component we exhibit an integral representation involving the probability density functions and the conditional survival function of the intertimes between two successive events.

This is the plan of the paper. In Section 2 we introduce the process $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$ describing the position of a particle performing a cyclic, minimal, random motion in \mathbb{R}^3 with four possible directions \vec{v}_j , for $1\leq j\leq 4$, and constant velocity c>0. Some preliminary results on the GCP are also briefly illustrated with reference to the distribution of the intertimes between two consecutive events. Then, in Section 3, we study the directional vectors and the analytic representation of the support in \mathbb{R}^3 identifying the tetrahedron $\mathcal{T}(t)$. In Section 4 we investigate the stochastic process and its probability laws with underlying GCP, and determine the explicit expression of the initial and absolutely continuous components. Section 5 illustrates a special case with four fixed cyclic directions forming a regular tetrahedron in \mathbb{R}^3 . Moreover, we examine the limiting distribution of the process when the parameters of the intertimes tend to infinity and when the time t goes to infinity. Finally, in Section 6 we provide some basic lines for the study of the first-passage-time problem of the first component of the process through a constant positive boundary $\beta>0$.

2 A random motion in \mathbb{R}^3 with cyclic velocities

Let $X(t) = (X_1(t), X_2(t), X_3(t))$ and V(t) be respectively the position and the velocity of the particle at an arbitrary time $t \geq 0$ in the space \mathbb{R}^3 . Assume that each point in \mathbb{R}^3 is represented by a triple $\boldsymbol{x} = (x_1, x_2, x_3)$. We consider a cyclic random motion $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ performed by a particle which can take four possible directions \vec{v}_j , for $1 \leq j \leq 4$, and moves with constant velocity c > 0. The motion is cyclic in the sense that at each event the particle switches from direction \vec{v}_j to \vec{v}_{j+1} , for j = 1, 2, 3, and then from \vec{v}_4 to \vec{v}_1 , and so on. Let $D_{j,k}$ be the random duration of the k-th time interval during which the motion proceeds with velocity c. For any $1 \leq j \leq 4$, the set $D_{j, \cdot} := \{D_{j,k}, k \in \mathbb{N}\}$ constitutes a sequence of non-negative and dependent absolutely continuous random variables such that

$$D_j^{(0)} = 0, \quad D_j^{(k)} = \sum_{i=1}^k D_{j,i}, \quad k \in \mathbb{N}.$$
 (1)

Moreover, the sets $D_{j,\cdot}$, for $1 \leq j \leq 4$, are mutually independent.

Now, let $\{N(t), t \geq 0\}$ be the alternating counting process having arrival times T_1, T_2, \ldots (i.e., the instants when the events occur), such that N(t) counts the total number of velocity reversals of the particle in [0, t], i.e.

$$N(0) = 0, \quad N(t) = \sum_{k=0}^{\infty} \mathbf{1}_{\{T_k \le t\}}, \quad t \ge 0.$$
 (2)

In order to obtain the probability law of the stochastic process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ introduced so far, we consider the stochastic equations for the position $\boldsymbol{X}(t)$ and the velocity V(t) of the particle at time $t \geq 0$. Specifically, we have

$$V(t) = \sum_{k=0}^{\infty} \mathbf{1}_{\{T_{k-1} \le t < T_k\}} \vec{v}_k,$$

with $V(0) \in \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4\}$, and

$$\boldsymbol{X}(t) = \boldsymbol{X}(0) + \int_0^t V(s)ds = \sum_{k=1}^{N(t)-1} (T_k - T_{k-1})\vec{v}_k + (t - T_{N(t)-1})\vec{v}_{N(t)}, \quad (3)$$

with $X(0) = \mathbf{0} \equiv (0,0,0)$ and $\sum_{k=1}^{N(t)-1} (T_k - T_{k-1}) = T_{N(t)-1}$, since $T_0 = 0$. The sum in Eq. (3) refers to the case where at least one change of direction has occurred before t, while the last term is related to the displacement along the current direction at time t.

We indicate with T_k the k-th random instant in which the motion modifies its direction, for $n \in \mathbb{N}_0$. Remembering Eq. (1) the following identity holds:

$$T_{4n+j} = \sum_{r=1}^{4} D_r^{(n+m_{j,r})},\tag{4}$$

where, for fixed $j \in \{1, 2, 3, 4\}$, $M = (m_{j,r})_{1 \le r \le 4} \in \mathbb{R}^{4 \times 4}$ is equal to

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

The relations obtained above will be used in Section 4 to study the probability law of the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$.

2.1 The distribution of the intertimes as a Geometric Counting Process

We introduce some preliminary results on the GCP (for more details, see Cha and Finkelstein (2013) and Di Crescenzo and Pellerey (2019)). Specifically, we assume that the random intertimes between consecutive changes of directions are governed by possible different GCPs (see, for instance, Di Crescenzo et al. (2023)).

For this purpose, we consider a mixed Poisson process $\{N(t), t \geq 0\}$ characterized by the following marginal distribution expressed as a mixture:

$$\mathbb{P}[N(t) = k] = \int_0^t \mathbb{P}[N^{(\alpha)}(t) = k] dU(\alpha), \quad t \ge 0, \quad k \in \mathbb{N}_0,$$

where $\{N^{(\alpha)}(t), t \geq 0\}$ is a Poisson process with intensity α and U is an exponential distribution with support \mathbb{R}^+ Here, we consider the special case when $U(\cdot) = U_{\lambda}(\cdot)$ is an exponential distribution with mean $\lambda \in \mathbb{R}^+$. Therefore, we refer to the process N(t) as a GCP with intensity λ , according to Cha and Finkelstein (2013), where the authors studied dependence properties of its increments in the general case of non-constant intensities. The probability distribution of the process N(t) satisfies the following properties:

- at time t = 0, one has N(0) = 0
- for all $s, t \geq 0$ and $k \in \mathbb{N}_0$ it holds:

$$\mathbb{P}\{N(t+s) - N(t) = k\} = \frac{1}{1+\lambda s} \left(\frac{\lambda s}{1+\lambda s}\right)^{k}.$$

Let T_k , with $k \in \mathbb{N}$, be the random times denoting the arrival instants of the process N(t) such that $T_0 = 0$. It is interesting to observe that the probability density function (p.d.f.) of T_k is expressed as follows:

$$f_{T_k}(t) = k \left(\frac{\lambda t}{1 + \lambda t}\right)^{k-1} \frac{\lambda}{(1 + \lambda t)^2}, \quad t \ge 0.$$

We define $D_{j,k} = T_k - T_{k-1}$, with $1 \le j \le 4$ and $k \in \mathbb{N}$, the increments between two consecutive events. Differently from the Poisson process, the GCP does not have the property of independent increments, hence, the random variables $D_{j,k}$ are dependent. Moreover, making use of Eq. (10) in Di Crescenzo and Pellerey (2019), we get the marginal density for all intertimes $D_{j,k}$, $1 \le j \le 4$ and $k \in \mathbb{N}$,

$$f_{D_{j,k}}(t) = \frac{\lambda}{(1+\lambda t)^2}, \quad t \ge 0.$$
 (5)

We remark that $D_{j,k}$ has a long right tail (see, for instance, Asmussen (2003)) since, for all t > 0,

$$\lim_{x \to \infty} \mathbb{P}\{D_{j,k} > x + t | D_{j,k} > x\} = 1.$$

From density (5) it is not hard to see that $D_{j,k}$ has a modified Pareto distribution which means that the intertimes of the GCP $\{N(t), t > 0\}$ have non-finite expectations. In addition, to obtain the explicit expressions of the process X(t), we recall the conditional survival function of $D_{j,k}$ conditional on $T_{k-1} = t$, for $k \in \mathbb{N}$, is given by

$$\overline{F}_{D_{j,k}|T_{k-1}}(s|t) = \left(\frac{1+\lambda t}{1+\lambda(t+s)}\right)^k, \quad s, t \ge 0,$$
(6)

and the corresponding probability density function (p.d.f.) of $D_{j,k}$ conditional on $T_{k-1} = t$, for $k \in \mathbb{N}$, i.e.,

$$f_{D_{j,k}|T_{k-1}}(s|t) = \frac{k\lambda(1+\lambda t)^k}{[1+\lambda(t+s)]^{k+1}}, \quad s,t \ge 0.$$
 (7)

In the following, we determine the initial and absolutely continuous components of the probability law of the stochastic process $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$, which describes a cyclic alternating random motion along four directions \vec{v}_j , for $1\leq j\leq 4$, driven by independent GCPs with intensity $?\lambda_j$ ($1\leq j\leq 4$. We remark that the intertimes of the GCPs are in practice more realistic since in real cases the counting processes under investigation do not have necessarily the independent increments property. Some applications may be found in geophysics (see Benson et al. (2007)), in climatology (see Lavergnat and Gole (1998)) or in modeling for internet traffic (see Clegg et al. (2010)). Moreover, in real applications the number of events in a fixed time interval does not always follow the Poisson distribution. Therefore, the class of GCPs is a possible alternative to Poisson processes. At least, the use of GCPs allows us to obtain the probability law of the process $\boldsymbol{X}(t)$ in a tractable way as shown in the remainder of the paper.

3 The velocity vectors and the support

In this section, we discuss the properties of the directional velocity vectors and the geometrical features of the region generated by them, i.e., the set of all possible positions of the moving particle at time t > 0.

3.1 The directional vectors

Let \vec{v}_j , $1 \leq j \leq 4$, be the vectors representing the possible directions of the cyclic motion in \mathbb{R}^3 (i.e., the minimal number of directions for a non trivial motion in \mathbb{R}^3). The velocity direction coordinates can be defined by using the spherical coordinates, given by

$$\vec{v}_j = (\vec{l}\cos\theta_j\sin\varphi_j + \vec{m}\sin\theta_j\sin\varphi_j + \vec{n}\cos\varphi_j),$$

where $\vec{l}, \vec{m}, \vec{n}$ are the unit vectors along the Cartesian coordinate axes in \mathbb{R}^3 , φ_j is the azimuthal angle in the xy-plane from the x-axis with $\theta_j \in [0, 2\pi]$ and θ_j is the polar angle from the positive z-axis with $\varphi_j \in [0, \pi]$, for $1 \leq j \leq 4$.

The particle starts from the origin $\mathbf{0} = (0,0,0) \in \mathbb{R}^3$ at time t = 0, running with constant velocity c > 0. Initially, it moves along the direction \vec{v}_1 . Then, after a random

duration $D_{1,1}$, the particle switches instantaneously its speed, moving along \vec{v}_2 for a random duration $D_{2,1}$. Subsequently, it goes along \vec{v}_3 and \vec{v}_4 for a length of time $D_{3,1}$ and $D_{4,1}$, respectively. Thus, the particle motion proceeds cyclically with velocities \vec{v}_j for the random periods $D_{j,2}, D_{j,3}, D_{j,4}, \ldots$, such that, for each $1 \leq j \leq 4$ and $k \in \mathbb{N}$, we have

$$D_{j,4k+i} \stackrel{d}{=} D_{j,i},\tag{8}$$

where $\stackrel{d}{=}$ means equality in distribution. Hence, during the *n*-th cycle the particle moves along directions \vec{v}_j in sequence for the random lengths $D_{j,n}$, with $1 \leq j \leq 4$ and $n \in \mathbb{N}$. In other words, the particle runs along direction \vec{v}_j and, after an intertime distributed as a CGP, it takes the direction \vec{v}_{j+1} , with $\vec{v}_{j+4n} = \vec{v}_j$, for $1 \leq j \leq 4$ and $n \in \mathbb{N}$. We note that the results concerning the cases of other initial direction, i.e., \vec{v}_2 , \vec{v}_3 , \vec{v}_4 , may be easily deduced. Now, in order to define the set of all possible positions occupied by the particle at an arbitrary time t > 0, we consider the following Remark.

Remark 1 Let be $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$ the stochastic process defined in Section 2. Given the direction vectors \vec{v}_j with $\vec{v}_{j+4n}=\vec{v}_j$, for $1\leq j\leq 4$ and $n\in\mathbb{N}$, we have that the particle motion reaches any state of \mathbb{R}^3 in a sufficiently large time t>0, if and only if,

- (i) the set of direction vectors $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ are linearly independent;
- (ii) the direction vector $\vec{v}_4 \in S$ with

$$S = \left\{ \xi(-\vec{v}_1) + \eta(-\vec{v}_2) + (1 - \xi - \eta)(-\vec{v}_3), 0 \le \xi, \eta \le 1, \ \xi + \eta \le 1 \right\}.$$

Hence, the set of all possible positions under the hypothesis of Remark 1 identifies as state space at time t a tetrahedron whose vertices coincide with the endpoints of the vectors $ct\vec{v}_j$. In other words, the tetrahedron is defined as a 3-dimensional simplex where the interior points are convex combinations of the four vertices such that $\sum_{j=1}^4 \alpha_j \vec{v}_j$ with $\sum_{j=1}^4 \alpha_j = 1$ and $\alpha_j \geq 0$.

3.2 Analytic representation of the support

Let us consider the points $A_j(t)$, $1 \le j \le 4$, defined by $\overrightarrow{\mathbf{0}}A_j(t) = \overrightarrow{v}_j t$, where $\mathbf{0} = (0,0,0)$ is the origin of the Cartesian coordinate system. The points $A_j(t)$, $1 \le j \le 4$, given by

$$A_{i}(t) = (ct\cos\theta_{i}\sin\varphi_{i}, ct\sin\theta_{i}\sin\varphi_{i}, ct\cos\varphi_{i}), \quad 1 \le j \le 4$$
 (9)

are the vertices of the support in \mathbb{R}^3 at time t>0, i.e., a time-dependent tetrahedron with edges $E_{ij}(t)$ and faces $F_{ijk}(t)$, $1 \le i, j, k \le 4$ and i < j < k. In particular, if the three different vertices $A_i(t)$, $A_j(t)$ and $A_k(t)$ are not aligned, then the particle reaches any point of the support in \mathbb{R}^3 . This is true since the matrix generated by two adjacent segments $A_i(t)A_k(t)$ and $A_k(t)A_j(t)$ has maximum rank, i.e. 2. Therefore, under the assumptions of Remark 1 the equations linking two adjacent vertices can expressed as

$$a_{ijk}x_1 + b_{ijk}x_2 + c_{ijk}x_3 - ctq_{ijk} = 0, \quad 1 \le i, j, k \le 4, \quad i < j < k,$$
 (10)

where

$$\begin{aligned} a_{ijk} &= (\sin\theta_j \sin\varphi_j - \sin\theta_i \sin\varphi_i)(\cos\varphi_k - \cos\varphi_i) \\ &- (\sin\varphi_k \sin\theta_k - \sin\varphi_i \sin\theta_i)(\cos\varphi_j - \cos\varphi_i), \\ b_{ijk} &= (\sin\varphi_k \cos\theta_k - \sin\varphi_i \cos\theta_i)(\cos\varphi_j - \cos\varphi_i) \\ &- (\sin\varphi_j \cos\theta_j - \sin\varphi_i \cos\theta_i)(\cos\varphi_k - \cos\varphi_i), \\ c_{ijk} &= \sin\varphi_i \cos\theta_i(\sin\varphi_j \sin\theta_j - \sin\varphi_k \sin\theta_k) \\ &+ \sin\varphi_j \cos\theta_j(\sin\varphi_k \sin\theta_k - \sin\varphi_i \sin\theta_i) \\ &+ \sin\varphi_k \cos\theta_k(\sin\varphi_i \sin\theta_i - \sin\varphi_j \sin\theta_j), \end{aligned}$$

and,

$$q_{ijk} = a_{ijk}\cos\theta_i\sin\varphi_i + b_{ijk}\sin\theta_i\sin\varphi_i + c_{ijk}\cos\varphi_i.$$

We recall that V(t) is the current direction of motion at time t. Assuming that V(0) takes value \vec{v}_1 with velocity c > 0, we are able to determine the analytic expression of the support in \mathbb{R}^3 , say $\mathcal{T}(t)$, i.e., the set of all positions allocated by the particle when it is confined in a tetrahedron, defined as

$$\mathcal{T}(t) = \left\{ (x_1, x_2, x_3) \in \mathbb{R}^3 : \begin{cases} a_{123}x_1 + b_{123}x_2 + c_{123}x_3 - ctq_{123} \ge 0 \\ a_{124}x_1 + b_{124}x_2 + c_{124}x_3 - ctq_{124} \ge 0 \\ a_{234}x_1 + b_{234}x_2 + c_{234}x_3 - ctq_{234} \ge 0 \\ a_{134}x_1 + b_{134}x_2 + c_{134}x_3 - ctq_{134} \le 0 \end{cases} \right\}, \quad (11)$$

where the given conditions derive from Eqs. (10). In general, the particle motion includes different mutually exclusive cases based on the initial assumptions $X(0) = \mathbf{0}$ and $V(0) = \vec{v_j}$, for a fixed $1 \leq j \leq 4$. More precisely, the particle is situated on the vertices $A_j(t)$ of $\mathcal{T}(t)$ if no velocity change occurs up to time t, while if one event occurs then, at time t, it will be located on some edge E_{ij} of $\mathcal{T}(t)$. Two events allow the particle to reach one of the faces F_{ijk} of $\mathcal{T}(t)$ and three or more changes of direction force the particle to be located inside $\mathcal{T}(t)$. Therefore, recalling Remark 1, the following holds.

Remark 2 For the probability law of the process $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$ we have:

- (i) the first component is related to the case when no event occurs in the interval (0, t), i.e., N(t) = 0, and the particle is concentrated on the vertices of $\mathcal{T}(t)$;
- (ii) the second component is concerning the situation in which one event happens at time t, i.e., N(t) = 1, and the particle is located on some edge of $\mathcal{T}(t)$;
- (iii) the third component corresponds to the instance when two events occur, i.e., N(t) = 2 and the particle reaches one of the faces of $\mathcal{T}(t)$;
- (iv) the forth component, which is absolutely continuous, refers to the case when $N(t) \ge 3$, i.e. when the particle is placed strictly in the interior of $\mathcal{T}(t)$, namely $\mathrm{Int}(\mathcal{T}(t))$.

4 The probability law

To define the conditional distributions of the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$, we denote by

$$C_i := \{ \mathbf{X}(0) = \mathbf{0}, V(0) = \vec{v}_i \},$$
 (12)

the event that the particle starts its motion from the origin $\mathbf{0}$ when the initial velocity is \vec{v}_i , $1 \leq i \leq 4$. Therefore, let \mathbb{P}_i be the probability conditional on C_i .

The conditional probability laws given C_i are composed by two contributions:

- (i) the initial components given by the terms related to the beginning of the motion. Specifically, they describe the cases in which the particle is over the border of the support $\mathcal{T}(t)$ in \mathbb{R}^3 .
- (ii) The absolutely continuous component related to the particle motion inside the support $\mathcal{T}(t)$ in \mathbb{R}^3 .

For the case (ii) we define

$$p_{ij}(\mathbf{x}, t)d\mathbf{x} = \mathbb{P}_i\{\mathbf{X}(t) \in d\mathbf{x}, V(t) = \vec{v}_j\}$$

$$= \mathbb{P}_i\{X_1(t) \in dx_1, X_2(t) \in dx_2, X_3(t) \in dx_3, V(t) = \vec{v}_j\},$$
(13)

with $1 \leq i, j \leq 4$ and where $d\boldsymbol{x}$ is the infinitesimal element in the space \mathbb{R}^3 with the Lebesgue measure $\mu(d\boldsymbol{x}) = dx_1 dx_2 dx_3$ (*i* refers to the initial velocity, and *j* to the current velocity at time *t*). The right-hand-side of Eq. (13) represents the probability that the particle at time t > 0 is located in a neighborhood of $\boldsymbol{x} \in \mathbb{R}^3$ and moves along direction \vec{v}_j , given the initial condition represented by C_i . Moreover, we can introduce the probability density function (p.d.f.) of the particle location $\boldsymbol{X}(t)$ at time t > 0, i.e.,

$$p_i(\boldsymbol{x}, t) = \mathbb{P}_i \{ \boldsymbol{X}(t) \in d\boldsymbol{x} \}, \quad 1 \le i \le 4. \tag{14}$$

Due to Eqs. (13) and (14), one immediately has

$$p_i(\mathbf{x}, t) = \sum_{j=1}^{4} p_{ij}(\mathbf{x}, t), \qquad 1 \le i \le 4.$$
 (15)

In order to determine the probability law of the stochastic process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$, we assume that the starting direction at time t is \vec{v}_1 , so that, the initial condition is C_1 as defined in Eq. (12). Therefore, the following results are obtained by taking into account the cases (i)-(iv) considered in Remark 2.

Theorem 1 (Initial components) Let be $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$ the stochastic process defined in Section 2. For $t\geq 0$ we have

$$\mathbb{P}_1\{\boldsymbol{X}(t) = (ct\cos\theta_j\sin\varphi_j, ct\sin\theta_j\sin\varphi_j, ct\cos\varphi_j), V(t) = \vec{v}_1\}
= \mathbb{P}\{D_{1,1} > t\},$$
(16)

$$\mathbb{P}_1\{X(t) \in E_{12}(t), V(t) = \vec{v}_2\} = \mathbb{P}\{D_{1,1} < t \le D_{1,1} + D_{2,1}\},\tag{17}$$

and

$$\mathbb{P}_1\{\boldsymbol{X}(t) \in F_{123}(t), V(t) = \vec{v}_3\} = \mathbb{P}\{D_{1,1} + D_{2,1} < t \le D_{1,1} + D_{2,1} + D_{3,1}\},\tag{18}$$

where $D_{j,k}$ is the random duration of the k-th time interval during which the particle moves with velocity \vec{v}_j , for $1 \leq j \leq 3$ and $k \in \mathbb{N}$.

Proof Eqs. (16), (17) and (18), when $V(0) = \vec{v}_1$, are easily obtained, under the conditions (i), (ii) and (iii) of Remark 2.

Using Eq. (13) and recalling (iv) of Remark 2, we determine the absolutely continuous components of the probability law of the X(t) conditioned on C_1 , for t > 0 and $1 \le j \le 4$, i.e., the densities

$$p_{1j}(\boldsymbol{x},t)d\boldsymbol{x} = \mathbb{P}_1\{\boldsymbol{X}(t) \in d\boldsymbol{x}, V(t) = \vec{v}_j\}. \tag{19}$$

In order to give the expression of $p_{1j}(\boldsymbol{x},t)$ we introduce the linear map $\zeta: \mathbb{R}^4 \longrightarrow \mathbb{R}^4$ defined by:

$$\zeta(\mathbf{t}) = \left(c\sum_{j=1}^{4} x_{1_{\vec{v}_j}} t_j, c\sum_{j=1}^{4} x_{2_{\vec{v}_j}} t_j, c\sum_{j=1}^{4} x_{3_{\vec{v}_j}} t_j, \sum_{j=1}^{4} t_j\right), \tag{20}$$

for $\mathbf{t} = (t_1, t_2, t_3, t_4)$, and where $x_{1\vec{v}_j}$, $x_{2\vec{v}_j}$ and $x_{3\vec{v}_j}$ are the components of the vectors \vec{v}_j ($1 \le j \le 4$) respectively along the \mathbf{x} -axes. When a cycle of the random motion ends after a period \mathbf{t} the function $\zeta(\mathbf{t})$, given in Eq. (20), provides a vector containing the displacements performed along the \mathbf{x} -axes during the cycle, as well as its whole duration.

Moreover, we consider the transformation matrix A of function $\zeta(t)$ expressed in terms of spherical coordinates, i.e.,

$$\mathbf{A} = \begin{pmatrix} c\cos\theta_1\sin\varphi_1 & c\cos\theta_2\sin\varphi_2 & c\cos\theta_3\sin\varphi_3 & c\cos\theta_4\sin\varphi_4 \\ c\sin\theta_1\sin\varphi_1 & c\sin\theta_2\sin\varphi_2 & c\sin\theta_3\sin\varphi_3 & c\sin\theta_4\sin\varphi_4 \\ c\cos\varphi_1 & c\cos\varphi_2 & c\cos\varphi_3 & c\cos\varphi_4, \\ 1 & 1 & 1 & 1 \end{pmatrix}$$
(21)

with

$$det \mathbf{A} = c^{3} \Big\{ \sin \varphi_{1} \sin \varphi_{2} [\sin(\theta_{2} - \theta_{1})] (\cos \varphi_{3} - \cos \varphi_{4})$$

$$+ \sin \varphi_{1} \sin \varphi_{3} [\sin(\theta_{3} - \theta_{1})] (\cos \varphi_{4} - \cos \varphi_{2})$$

$$+ \sin \varphi_{1} \sin \varphi_{4} [\sin(\theta_{4} - \theta_{1})] (\cos \varphi_{2} - \cos \varphi_{3})$$

$$+ \sin \varphi_{2} \sin \varphi_{3} [\sin(\theta_{3} - \theta_{2})] (\cos \varphi_{1} - \cos \varphi_{4})$$

$$+ \sin \varphi_{2} \sin \varphi_{4} [\sin(\theta_{4} - \theta_{2})] (\cos \varphi_{3} - \cos \varphi_{1})$$

$$+ \sin \varphi_{3} \sin \varphi_{4} [\sin(\theta_{4} - \theta_{3})] (\cos \varphi_{1} - \cos \varphi_{2}) \Big\}.$$

$$(22)$$

Remark 3 Recalling (9), we observe that the determinant $\det(t\mathbf{A})$ represents the volume $Vol(\mathcal{T}(t))$ of $\mathcal{T}(t)$ defined in (11).

Now, given a sample path of the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$, we denote by $\tau_j = \tau_j(\boldsymbol{x})$ the non negative random variables representing the dwelling times of the particle motion in each direction \vec{v}_j $(1 \leq j \leq 4)$ during [0, t] (i.e., the residence times for the process $\boldsymbol{X}(t)$ on the event that $\boldsymbol{X}(t) = \boldsymbol{x}, V(t) = \vec{v}_j$), so that

$$\sum_{j=1}^{4} \tau_j = t.$$

Therefore, recalling (21) and (9), we can express τ_j , $1 \le j \le 4$, as the 4-uple

$$\zeta^{-1}(\boldsymbol{x},t) = (\tau_1(\boldsymbol{x},t), \tau_2(\boldsymbol{x},t), \tau_3(\boldsymbol{x},t), \tau_4(\boldsymbol{x},t)),$$

with

$$\tau_j(\boldsymbol{x},t) = \sum_{k=1}^4 \sigma_{jk} x_k + \sigma_{j4} t, \qquad (23)$$

where $\sigma_{ij} = \frac{1}{\det \mathbf{A}} cof_{ij}(\mathbf{A})$ with $cof_{ij}(\mathbf{A}) = (-1)^{i+j} \det A_{ij}$, $1 \leq i, j \leq 4$. Moreover, since the function $\zeta(\mathbf{t})$, for all $\mathbf{x} \in \mathbb{R}^3$, is bijective (see, for more details, Lachal (2006)), it is easy to see that $\boldsymbol{\tau} = (\tau_1, \tau_2, \tau_3, \tau_4)^T$ is solution of the system $\mathbf{A}\boldsymbol{\tau} = (\mathbf{x}, t)^T$. More precisely, in the following proposition the explicit form of Eq. (23) can be obtained.

Proposition 2 Making use of Eq. (23), for all $\mathbf{x} \in \mathbb{R}^3$ the coordinates τ_j , for $1 \le j \le 4$ and c > 0, are expressed as

$$\tau_{j} = \frac{c^{2}}{\det \mathbf{A}} \left[L_{j} x_{1} + M_{j} x_{2} + N_{j} x_{3} - ct P_{j} \right], \quad 1 \le j \le 4, \tag{24}$$

where

$$\begin{split} L_j &= \sum_{i=1}^3 \sin \varphi_{j \hat{+} i} \sin \theta_{j \hat{+} i} \bigg(\cos \varphi_{j \hat{+} (i+1)} - \cos \varphi_{j \hat{+} (i+2)} \bigg), \\ M_j &= \sum_{i=1}^3 \sin \varphi_{j \hat{+} i} \cos \theta_{j \hat{+} i} \bigg(\cos \varphi_{j \hat{+} (i+2)} - \cos \varphi_{j \hat{+} (i+1)} \bigg), \\ N_j &= \sum_{i=1}^3 \sin \varphi_{j \hat{+} i} \cos \theta_{j \hat{+} i} \bigg(\sin \varphi_{j \hat{+} (i+1)} \sin \theta_{j \hat{+} (i+1)} - \sin \varphi_{j \hat{+} (i+2)} \sin \theta_{j \hat{+} (i+2)} \bigg), \\ P_j &= \sum_{i=1}^3 \sin \varphi_{j \hat{+} i} \cos \theta_{j \hat{+} i} \\ &\times \bigg(\sin \varphi_{j \hat{+} (i+1)} \sin \theta_{j \hat{+} (i+1)} \cos \varphi_{j \hat{+} (i+2)} - \sin \varphi_{j \hat{+} (i+2)} \sin \theta_{j \hat{+} (i+2)} \cos \varphi_{j \hat{+} (i+1)} \bigg), \end{split}$$

where a+b denotes (a+b) mod 4 and det **A** can be recovered from (21).

Proof Recalling Eqs. (9), (22) and (23), after straightforward calculations Eq. (24) is determined. \Box

Now, for the process $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$ we are able to formulate the following theorem about the absolutely continuous component concerning assumption (iv) of Remark 2. For this purpose, we indicate with $f_{D_j^{(n)}}$ the probability density function (p.d.f.) of $D_j^{(n)}$ given in Eq. (1), $1\leq j\leq 4$, and with $\overline{F}_{D_{j,k+1}|D_j^{(k)}}$ the conditional survival function of $D_{j,k+1}$ given $D_j^{(k)}$, for $1\leq j\leq 4$ and $k\in\mathbb{N}_0$.

Theorem 3 (Absolutely continuous component) For the stochastic process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ defined in Section 2, under the initial condition given in Eq. (12) with direction \vec{v}_1 , we have that, for all $\boldsymbol{x} \in \operatorname{Int}(\mathcal{T}(t))$ and t > 0, the absolutely continuous component of the probability law is given by

$$p_{1j}(\boldsymbol{x},t) = \frac{1}{\det \boldsymbol{A}} \sum_{k=0}^{\infty} \left\{ \prod_{i=1}^{j-1} f_{D_i^{(k+1)}}(\tau_i) \prod_{i=j+1}^{4} f_{D_i^{(k)}}(\tau_i) \right. \\ \left. \times \int_{t-\tau_j}^{t} f_{D_j^{(k)}}(s - (t - \tau_j)) \overline{F}_{D_{j,k+1}|D_j^{(k)}}(t - s|s - (t - \tau_j)) ds \right\},$$
(25)

where $\tau_j = \tau_j(\boldsymbol{x},t)$, for $1 \leq j \leq 4$, is defined in Eq. (24), and $\det \boldsymbol{A}$ is given in Eq. (22).

Proof By conditioning on the number of direction switches in (0, t), say k, and on the last time s previous t in which the particle changes its velocity from \vec{v}_4 to \vec{v}_1 in order to restart a new cycle, we reformulate Eq. (19), for $\mathbf{x} \in \mathbb{R}^3$ and t > 0, as follows:

$$p_{1j}(\mathbf{x}, t)d\mathbf{x} = \sum_{k=0}^{\infty} \int_{0}^{t} \mathbb{P}\{T_{4k+j-1} \in ds, X_{1}(s) + c(t-s)\cos\theta_{j}\sin\varphi_{j} \in dx_{1}, X_{2}(s) + c(t-s)\sin\theta_{j}\sin\varphi_{j} \in dx_{2}, X_{3}(s) + c(t-s)\cos\varphi_{j} \in dx_{3}, D_{j,k+1} > t-s\},$$
(26)

for $1 \leq j \leq 4$ and where T_{4k+j-1} is the instant occurring at time s in which the particle changes its direction to velocity \vec{v}_j . Thus, $\boldsymbol{X}(s) = (X_1(s), X_2(s), X_3(s))$ is the position of the particle when a change of direction occurs. Therefore, due to Eq.(4), we have

$$T_{4k+j-1} = \sum_{i=1}^{j-1} D_i^{(k+1)} + \sum_{i=j+1}^4 D_i^{(k)} + D_j^{(k)}.$$

Hence, for $s = T_{4k+j-1}$ and using Eq. (1), we can express the coordinates of the particle position X(s) as follows

$$X_{1}(s) = c \left[\sum_{i=1}^{j-1} D_{i}^{(k+1)} \cos \theta_{i} \sin \varphi_{i} + \sum_{i=j+1}^{4} D_{i}^{(k)} \cos \theta_{i} \sin \varphi_{i} + D_{j}^{(k)} \cos \theta_{j} \sin \varphi_{j} \right],$$

$$X_{2}(s) = c \left[\sum_{i=1}^{j-1} D_{i}^{(k+1)} \sin \theta_{i} \sin \varphi_{i} + \sum_{i=j+1}^{4} D_{i}^{(k)} \sin \theta_{i} \sin \varphi_{i} + D_{j}^{(k)} \sin \theta_{j} \sin \varphi_{j} \right],$$

$$X_{3}(s) = c \left[\sum_{i=1}^{j-1} D_{i}^{(k+1)} \cos \varphi_{i} + \sum_{i=j+1}^{4} D_{i}^{(k)} \cos \varphi_{i} + D_{j}^{(k)} \cos \varphi_{j} \right].$$
(27)

The conditions defined in (11) and $X(s) \in \mathcal{T}(s)$ implies that $s > t - \tau_j$, for $1 \leq j \leq 4$. Moreover, using (21) and substituting Eqs. (27) in Eq. (26). we have:

$$p_{1j}(\boldsymbol{x},t) = \frac{1}{\det \boldsymbol{A}} \sum_{k=0}^{\infty} \left\{ \int_{t-\tau_{j}}^{t} \Psi_{j,k} \left[s, x_{1} - c(t-s) \sin \varphi_{j} \cos \theta_{j}, x_{2} - c(t-s) \sin \varphi_{j} \cos \theta_{j}, x_{3} - c(t-s) \cos \varphi_{j} \right] \right.$$

$$\times \mathbb{P} \left[D_{j,k+1} > t - s | T_{4k+j-1} = s, X_{1}(T_{4k+j-1}) = x_{1} - x_{1_{\vec{v}_{j}}}(t-s), X_{2}(T_{4k+j-1}) = x_{2} - x_{2_{\vec{v}_{j}}}(t-s), X_{3}(T_{4k+j-1}) = x_{3} - x_{3_{\vec{v}_{j}}}(t-s) \right] ds \right\},$$

$$(28)$$

where $\Psi_{j,k}$ is the joint p.d.f. of $(T_{4k+j-1}, X_1(T_{4k+j-1}), X_2(T_{4k+j-1}), X_3(T_{4k+j-1}))$ with:

$$\begin{split} T_{4k+j-1} &= \sum_{i=1}^{j-1} D_i^{(k+1)} + \sum_{i=j+1}^4 D_i^{(k)} + D_j^{(k)}, \\ X_1(T_{4k+j-1}) &= \sum_{i=1}^{j-1} x_{1_{\vec{v}_i}} D_i^{(k+1)} + \sum_{i=j+1}^4 x_{1_{\vec{v}_i}} D_i^{(k)} + x_{1_{\vec{v}_j}} D_j^{(k)}, \\ X_2(T_{4k+j-1}) &= \sum_{i=1}^{j-1} x_{2_{\vec{v}_i}} D_i^{(k+1)} + \sum_{i=j+1}^4 x_{2_{\vec{v}_i}} D_i^{(k)} + x_{2_{\vec{v}_j}}^{(k)} D_j, \\ X_3(T_{4k+j-1}) &= \sum_{i=1}^{j-1} x_{3_{\vec{v}_i}} D_i^{(k+1)} + \sum_{i=j+1}^4 x_{3_{\vec{v}_i}} D_i^{(k)} + x_{3_{\vec{v}_j}} D_j^{(k)}. \end{split}$$

According to (20), for $1 \le j \le 4$, we have

$$x_{1_{\overrightarrow{v}_i}} = v_i \sin \varphi_i \cos \theta_i, \ x_{2_{\overrightarrow{v}_i}} = v_i \sin \varphi_i \sin \theta_i, \ x_{3_{\overrightarrow{v}_i}} = v_i \cos \varphi_i.$$

Moreover, given the mutual independence of the variables $\{D_{j,k}; k \in \mathbb{N}\}$, for $1 \leq j \leq 4$, we get

$$\Psi_{j,k} \left[s, x - c(t-s) \cos \theta_j \sin \varphi_j, y - c(t-s) \sin \theta_j \sin \varphi_j, z - c(t-s) \cos \varphi_j \right]$$

$$= \prod_{i=1}^{j-1} f_{D_i^{(k+1)}}(\tau_i) \prod_{i=j+1}^4 f_{D_i^{(k)}}(t-\tau_j) f_{D_j^{(k)}}[s-(t-\tau_j)].$$
(29)

Therefore, Eq. (25) is directly obtained replacing Eq. (29) in Eq. (28).

We point out that Eq. (25) is in analogy with Eq. (4.1) of Lachal (2006) in which a particular case of the minimal cyclic random motion in \mathbb{R}^d (n=d+1 directions, $n \leq d$) is investigated. Specifically, the integral formulation given in Eq. (25) is based on the application of Eqs. (6) and (7) where the assumption of dependent increments, instead of the typical exponential distribution, is taken into account. Differently, in Orsingher and Sommella (2004) and Lachal $et\ al.$ (2006) the explicit form of the absolutely continuous part of $\boldsymbol{X}(t)$ is determined but using an approach based on the Bessel functions of higher order.

Note that we analyze the motion in \mathbb{R}^3 since the amount of calculations needed is tractable and the obtained results are applicable for practical problems. Hence, we can easily conjecture that in \mathbb{R}^n the structure of the absolutely continuous component

of the distribution of a cyclic motion with n+1 directions (with directions forming a regular or possible irregular tetrahedron) is similar to Eq. (25). We remark that the n-dimensional case will be object of a future work.

Clearly, the results given in Theorems 1 and 3 for the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ can be extended similarly when the particle starts with velocity \vec{v}_l , for $2 \leq l \leq 4$. Using this new scheme, a fixed initial direction allows us to obtain tractable expressions for the probability law of the process as illustrated in the following section.

5 A special case

In this section we analyze a special case of the process $\{(X(t), V(t)), t \geq 0\}$, with initial condition defined by Eq. (12) when $V(0) = \vec{v}_1$ and for fixed directions.

Assumptions 1 The following conditions hold:

(i) The particle's motion is confined in a regular tetrahedron $\mathcal{T}(t)$ characterized by the vertices:

$$A_{1}(t) = ct(1,0,0), \quad A_{2}(t) = ct\left(-\frac{1}{3}, \frac{2\sqrt{2}}{3}, 0\right),$$

$$A_{3}(t) = ct\left(-\frac{1}{3}, -\frac{\sqrt{2}}{3}, \sqrt{\frac{2}{3}}\right), \quad A_{4}(t) = ct\left(-\frac{1}{3}, -\frac{\sqrt{2}}{3}, -\sqrt{\frac{2}{3}}\right).$$
(30)

- (ii) The vertices in Eq. (30), identified by the directions \vec{v}_j , for $1 \le j \le 4$, satisfy the conditions (i)-(iv) listed in Remark 2.
- (iii) The random times $D_{j,n}$, for $n \in \mathbb{N}$ and $1 \le j \le 4$, represent the j-th duration of the motion within the n-th cycle and constitute the intertimes of a GCP with intensity $\lambda_j \in \mathbb{R}^+$.

An example of projections onto the state-space \mathbb{R}^3 of suitable paths of the process $\{(\boldsymbol{X}(t),V(t)),t\geq 0\}$ under the Assumptions 1 is shown in Figure 1. It is easy to see that the particle can eventually reach any position of \mathbb{R}^3 , since $\mathcal{T}(t)\to\mathbb{R}^3$ as $t\to\infty$. Moreover, by taking into account the region defined in (11) and Assumptions 1 we have that at every time t>0 the set of possible positions $\boldsymbol{x}\in\mathbb{R}^3$ of the moving particle is the tetrahedron

$$\mathcal{T}(t) = \left\{ (x_1, x_2, x_3) \in \mathbb{R}^3 : -\frac{ct}{3} < x_1 < ct; -\frac{ct - x_1}{2\sqrt{2}} < x_2 < \frac{ct - x_1}{\sqrt{2}}, \right.$$

$$\left. |x_3| < \frac{\sqrt{6}}{6} (ct - x_1 - \sqrt{2}x_2) \right\}.$$

$$(31)$$

Remark 4 We observe that the set $\mathcal{T}(t)$ grows as time elapses and, recalling Remark 3, the volume is given by $Vol(\mathcal{T}(t)) = \left(\frac{4}{3}\right)^2 \sqrt{3}(ct)^3$.

Two sample paths of the region $\mathcal{T}(t)$, defined in (31), with directions \vec{v}_j , $1 \leq j \leq 4$, is illustrated in Figure 2. Under the hypothesis of Assumptions 1, we are now able to determine the explicit probability laws of the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ when the




Figure 1 A sample of the set $\mathcal{T}(t)$, with velocities \vec{v}_j , for $1 \leq j \leq 4$, and fixed vertices given in (30).




Figure 2 Two possible paths with the first four segments of the cyclic motion defined in Section 2 under the conditions of Remark 1 with c=t=1. The initial velocity is \vec{v}_1 .

random intertimes of the motion along the possible directions \vec{v}_j follow four possible independent GCPs with intensities $\lambda_j, 1 \leq j \leq 4$. Specifically, in the following theorem, in order to obtain a tractable form for the third component given in (iii) of Remark 2, we assume that the sojourn times along the directions \vec{v}_1, \vec{v}_2 , and \vec{v}_3 follow three independent GCPs with identical intensity λ (i.e., $\lambda_1 = \lambda_2 = \lambda_3 := \lambda$).

Theorem 4 (Initial components) Let $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ be the stochastic process defined in Section 2 under the initial condition defined in Eq. (12) with velocity \vec{v}_1 . If Assumptions 1 hold and the four sequences of intertimes are identically distributed and follow a GCP with intensity λ_j , $1 \leq j \leq 4$, then for all $t \geq 0$, we have

$$\eta_1(t) = \mathbb{P}_1 \left\{ \mathbf{X}(t) = (ct, 0, 0), V(t) = \vec{v}_1 \right\} = \frac{1}{1 + \lambda_1 t},$$
(32)

$$\eta_2(t) = \mathbb{P}_1 \left\{ \mathbf{X}(t) \in E_{12}, V(t) = \vec{v}_2 \right\} \\
= \frac{\lambda_1}{(\lambda_1 + \lambda_2 + \lambda_1 \lambda_2 t)^2} \left\{ \frac{\lambda_1 t (\lambda_1 + \lambda_2 + \lambda_1 \lambda_2 t)}{1 + \lambda_1 t} + \lambda_2 \log \left[(1 + \lambda_1 t)(1 + \lambda_2 t) \right] \right\}$$
(33)

and

$$\eta_3(t) = \mathbb{P}_1 \left\{ \mathbf{X}(t) \in F_{123}, V(t) = \vec{v}_3 \right\} = R_{\lambda}(t),$$
(34)

where for $\lambda_1 = \lambda_2 = \lambda_3 = \lambda$ one has

$$R_{\lambda}(t) = \frac{1}{(2+\lambda t)^2(3+\lambda t)^3}$$

$$\left\{ 2\log(1+\lambda t) \left[(3+\lambda t) \left(\lambda t (3+2\lambda t) \right) + 4(2+\lambda t)^2 \log(2+\lambda t) \right] + (2+\lambda t) \left[\pi^2 (2+\lambda t) + \lambda t (3+\lambda t)^2 - 4(2+\lambda t) \log \left(\frac{1}{2+\lambda t} \right) \log \left(\frac{1+\lambda t}{2+\lambda t} \right) \right] + 4(2+\lambda t)^2 \left[Li_2 \left(-(1+\lambda t) \right) + 2Li_2 \left(\frac{1+\lambda t}{2+\lambda t} \right) \right] \right\}$$

where $Li_2(\cdot)$ is the dilogarithm function.

Proof Recalling Theorem 1, under Assumption 1, Eq. (32) is obtained making use of Eq. (5) in the following expression

$$\mathbb{P}_1\{\boldsymbol{X}(t) \in A_1(t), V(t) = \vec{v}_1\} = 1 - \int_0^t f_{D_{1,1}}(s) ds.$$

Similarly, Eqs. (33) and (34), when $V(0) = \vec{v}_2$ and $V(0) = \vec{v}_3$, are given by

$$\mathbb{P}_1\{\boldsymbol{X}(t) \in E_{12}(t), V(t) = \vec{v}_2\} = \int_0^t f_{D_{1,1}}(s) \int_{t-s}^{\infty} f_{D_{2,1}}(u) du ds$$

and

$$\mathbb{P}_1\{\boldsymbol{X}(t) \in F_{123}(t), V(t) = \vec{v}_3\} = \int_0^t f_{D_{1,1} + D_{2,1}}(s) f_{D_{3,1}}(t - s) ds,$$

respectively, after some calculations.

In Figure 3 are shown some plots of the probabilities given in Eqs. (32), (33) and (34) for different choices of the intensities λ_j , $1 \le j \le 3$, respectively.




Figure 3 Left to right: plot of $\eta_1(t)$ with $\lambda_1 = 1$ (solid), 2 (dotted), 10 (dashed); plot of $\eta_2(t)$ with $\lambda_1 = 1$ and $\lambda_2 = 2$ (solid line), $\lambda_1 = 2$ and $\lambda_2 = 4$ (dotted line), $\lambda_1 = 5$ and $\lambda_2 = 10$ (dashed line); plot of $\eta_3(t)$ with $\lambda = 1$ (solid), 5 (dotted), 10 (dashed).

Theorem 5 (Absolutely continuous components) Let $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ be the stochastic process defined in Section 2 under the initial condition defined in Eq. (12) with velocity \vec{v}_1 . For all $\boldsymbol{x} \in \operatorname{Int}(\mathcal{T}(t))$, with $\mathcal{T}(t)$ given in (31), we have

$$p_{11}(\mathbf{x},t) = \lambda_{1}\lambda_{2}\lambda_{3}\lambda_{4}\tau_{1}$$

$$\times \frac{\left[1 + A(\tau) + B(\tau) + C(\tau)\right]^{2} + 6D(\tau)\left[1 + A(\tau) + B(\tau) + C(\tau) + D(\tau)\right]}{\det \mathbf{A}\left[1 + A(\tau) + B(\tau) + C(\tau)\right]^{4}},$$

$$p_{12}(\mathbf{x},t) = 2\lambda_{1}^{2}\lambda_{2}\lambda_{3}\lambda_{4}\tau_{1}\tau_{2}(1 + \lambda_{2}\tau_{2})(1 + \lambda_{3}\tau_{3})(1 + \lambda_{4}\tau_{4})$$

$$\times \frac{\left[1 + A(\tau) + B(\tau) + C(\tau) + 3D(\tau)\right]}{\det \mathbf{A}\left[1 + A(\tau) + B(\tau) + C(\tau)\right]^{4}},$$

$$p_{13}(\mathbf{x},t) = 2\lambda_{1}^{2}\lambda_{2}^{2}\lambda_{3}\lambda_{4}\tau_{1}\tau_{2}\tau_{3}(1 + \lambda_{3}\tau_{3})(1 + \lambda_{4}\tau_{4})$$

$$\times \frac{\left[1 + A(\tau) + B(\tau) + C(\tau)\right] + 3D(\tau)}{\det \mathbf{A}\left[1 + A(\tau) + B(\tau) + C(\tau)\right]^{4}},$$

$$p_{14}(\mathbf{x},t) = \lambda_{1}\lambda_{2}\lambda_{3}(1 + \lambda_{4}\tau_{4})$$

$$\times \frac{\left[1 + A(\tau) + B(\tau) + C(\tau)\right]^{2} + 6D(\tau)[1 + A(\tau) + B(\tau) + C(\tau) + D(\tau)]}{\det \mathbf{A}\left[1 + A(\tau) + B(\tau) + C(\tau)\right]^{4}},$$

with

$$det \mathbf{A} = \frac{16\sqrt{3}}{9}c^{3}, \quad A(\boldsymbol{\tau}) = \sum_{i=1}^{4} \lambda_{i}\tau_{i}, \quad B(\boldsymbol{\tau}) = \sum_{\substack{i,j=1\\i < j}}^{4} \lambda_{i}\lambda_{j}\tau_{i}\tau_{j},$$
$$C(\boldsymbol{\tau}) = \sum_{\substack{i,j,k=1\\i < i < k}}^{4} \lambda_{i}\lambda_{j}\lambda_{k}\tau_{i}\tau_{j}\tau_{k}, \quad D(\boldsymbol{\tau}) = \prod_{i=1}^{4} \lambda_{i}\tau_{i},$$

and where the terms $\tau_j = \tau_j(\boldsymbol{x}, t)$, for $1 \leq j \leq 4$, are given by

$$\tau_1 = \frac{ct + 3x_1}{4c}, \quad \tau_2 = \frac{ct - x_1 + 2\sqrt{2}x_2}{4c},
\tau_3 = \frac{ct - x_1 - \sqrt{2}x_2 + \sqrt{6}x_3}{4c}, \quad \tau_4 = \frac{ct - x_1 - \sqrt{2}x_2 - \sqrt{6}x_3}{4c}.$$
(36)

Proof Eqs. (35) are obtained in a closed form as an immediate consequence of Theorem 3 after straightforward calculations.

We observe that the values of τ_j in (36) are obtained directly from Eq. (24) in Proposition 2. Moreover, due to Eq. (15), under the assumptions of Theorem 5 the p.d.f. $p_1(\boldsymbol{x},t)$ can be immediately obtained from Eqs. (35).




Figure 4 Plot of $p_1(x,t)$ for c=t=1 and $x_1=1/2$ on the left-side and c=1, t=2 and $x_1=1$ on the right-side.

As example, some plots of $p_1(\boldsymbol{x},t)$ are illustrated in Figure 4 for fixed c>0 and t>0 and $x_1\in(-ct/3,ct)$. Note that the domain for (x_2,x_3) has a triangular form. Moreover, from Eq. (31) and Theorem 5 it is clear that the dependence on c and t is expressed through the product ct.

Hereafter, we study the asymptotic behaviour of the p.d.f. of the particle location X(t) defined in (15) when the intensities λ_j tend to infinity, for $1 \leq j \leq 4$. In particular, substituing Eqs. (35) in Eq. (15) we obtain the following corollary.

Corollary 1 Under the assumptions of Theorem 5, for t > 0 and $x \in Int(\mathcal{T}(t))$ one has

$$\lim_{\begin{subarray}{c} \forall j, \, \lambda_j \to +\infty \\ \forall i, \, \lambda_1/\lambda_i \to 1 \end{subarray}} p_1(\boldsymbol{x}, t) = \xi(\boldsymbol{x}, t),$$

where $\xi(\boldsymbol{x},t)$ is the following p.d.f.

$$\xi(\boldsymbol{x},t) = \frac{6t(\tau_1\tau_2\tau_3\tau_4)^2}{\det \boldsymbol{A}[(\tau_1\tau_2\tau_3)^4 + (\tau_1\tau_2\tau_4)^4 + (\tau_1\tau_3\tau_4)^4 + (\tau_2\tau_3\tau_4)^4]},$$
 (37)

with τ_j expressed in (24).

Some instances of $\xi(\boldsymbol{x},t)$ given in Eq. (37) are plotted in Figure 5 for fixed c>0 and t>0, with $x_1\in(-ct/3,ct)$.




Figure 5 Plot of $\xi(\boldsymbol{x},t)$ for c=t=1 and $x_1=1/2$ on the left-side and $c=1,\,t=2$ and $x_1=1$ on the right-side.

Remark 5 Under the assumptions of Theorem 5, we note that, for t > 0 and $1 \le j \le 4$, the limit of $p_1(\boldsymbol{x},t)$ for $(x_1,x_2,x_3) \to (ct\cos\theta_j\sin\varphi_j,ct\sin\theta_j\sin\varphi_j,ct\cos\varphi_j)$ can be computed in a closed form that we omit due to its complexity.

Similarly to the classical telegraph process driven by the Poisson process (see, e.g., Lemma 2 of Orsingher (1990)), it is not hard to see that the process X(t) does not admit a stationary state. In the next corollary making use of Eq. (15) we analyse the asymptotic behaviour of density $p_1(x,t)$ when the time t tends to $+\infty$.

Corollary 2 Under the assumptions of Theorem 5, for
$$t>0$$
 and $\boldsymbol{x}\in \mathrm{Int}(\mathcal{T}(t))$ one has
$$\lim_{t\to +\infty} p_1(\boldsymbol{x},t)=0.$$

Proof Recalling Eq. (15), the stated result follows by noting that each density $p_{1,j}(\boldsymbol{x},t)$ in (35), for $1 \le j \le 4$, behaves as t^{-3} as t tends to infinity.

The results expressed in this section for $V(0) = \vec{v}_1$ can be extended to the cases $V(0) = \vec{v}_j$, for $2 \le j \le 4$, by using a similar strategy.

6 Concluding remarks

In this paper we analyzed a finite random motion in \mathbb{R}^3 where the sojourn times along each direction form four independent GCPs, and where the possible directions alternate cyclically. This work has been inspired by Di Crescenzo *et al.* (2023) with the aim to de?ne similar processes in higher dimensions with possibly variable velocities. Potential applications in biomathematics, engineering, financial and actuarial sciences allow to investigate possible future developments also oriented to the study of the first-passage-time problem.

Here, in order to illustrate the basic issues of this problem, we introduce the (upward) first-passage time for the first component of the process $\{(\boldsymbol{X}(t), V(t)), t \geq 0\}$ through a constant barrier, say $\beta > 0$, conditional on C_1 (cf. Eq. (12)) given by

$$\tau_{\beta} = \inf\{t \ge 0 : X_1(t) \ge \beta\}, \quad \mathbf{X}(0) = \mathbf{0}, \quad V(0) = \vec{v}_1, \quad |\vec{v}_1| = c, \quad (38)$$

for c > 0. The probability distribution of (38) can be expressed in terms of the subdensity functions:

$$g_{\beta}(t,k):=\frac{\mathbb{P}_1\{\tau_{\beta}\in dt, N(t)=k\}}{dt}, \quad k\in\mathbb{N}.$$

In particular, by the law of total probability, we express the conditional distribution of τ_{β} in the form

$$\mathbb{P}_1\{\tau_\beta \in dt\} = \mathbb{P}\{D_{1,1} > t\}\delta_{\frac{\beta}{c}}(dt) + \sum_{k=1}^{+\infty} \mathbb{P}_1\{\tau_\beta \in dt, N(t) = k\},\tag{39}$$

where $\delta_{\frac{\beta}{c}}$ is the Dirac delta measure at $\frac{\beta}{c}$, and N(t) is the alternating counting process introduced in Eq. (2). The first term on the right-hand-side of (39) corresponds to the motion without any direction switching up to time t. The series on the right-hand side of (39) represents the absolutely continuous component of the first-passage-time distribution, which arises when at least one direction reversal occurs. We also recall that $D_{1,1}$, defined in Eq. (1), is the random duration of the first time interval during which the particle proceeds with velocity \vec{v}_1 . In order to describe the first-passage-time problem, we consider as threshold the plane $x_1 = \beta$ and project the vectors \vec{v}_j , for $1 \le j \le 4$, onto the x_1 -axes according to the following relation

$$\vec{v}_{j_{x_1}} = proj_{\vec{w}_1}(\vec{v}_j) = \frac{\vec{w}_1 \cdot \vec{v}_j}{||\vec{w}_1||^2} \cdot \vec{w}_1, \qquad 1 \le j \le 4.$$

The latter is the projection of the vector \vec{v}_j on \vec{w}_1 , which is the versor along the x_1 -axes. An illustration of the problem is shown in Figure 6. Clearly, under Assumption




Figure 6 Illustration the constant barrier $\beta > 0$ for the first-passage time problem introduced in (38).

1 of Section 5, we have:

$$\begin{split} \vec{v}_{1_{x_1}} &= c(1,0,0), \quad \vec{v}_{2_{x_1}} = c\Big(-\frac{1}{3},0,0\Big), \\ \vec{v}_{3_{x_1}} &= c\Big(-\frac{1}{3},0,0\Big), \quad \vec{v}_{4_{x_1}} = c\Big(-\frac{1}{3},0,0\Big). \end{split}$$

Due to the complexity of the problem, we discuss only the first cycle of particle motion concerning $D_{j,1}$ ($1 \le j \le 4$). We point out that the particle may reach the threshold $x_1 = \beta$ only when it runs with velocity $\vec{v}_{1_{x_1}}$, since the other directions $\vec{v}_{2_{x_1}}$, $\vec{v}_{3_{x_1}}$ and $\vec{v}_{4_{x_1}}$ refer to motion in the opposite direction. Moreover, with reference to the first term in the right-hand-side of (39), one has

$$\mathbb{P}(D_{1,1} > t) = \mathbb{P}\left\{X_1(\tau_\beta) = \beta, V(\tau_\beta) = \vec{v}_1\right\}$$
$$= \mathbb{P}\left\{\tau_\beta = \frac{\beta}{c}, N(\tau_\beta) = 0\right\}$$
$$= \overline{F}_{D_{1,1}}\left(\frac{\beta}{c}\right) = \frac{1}{1 + \lambda_1 \beta/c}.$$

Clearly, during the random periods $D_{2,1}, D_{3,1}, D_{4,1}$, when the particle moves with velocity $\vec{v}_{2_{x_1}} = \vec{v}_{3_{x_1}} = \vec{v}_{4_{x_1}} = -\frac{c}{3}$, it cannot reach the threshold β since it moves in the opposite direction. Therefore, the position occupied by the particle at the end of the first period of motion $D_1^{(1)}$ is given by

$$X_1(t) = cD_{1,1} - \frac{c}{3}(D_{2,1} + D_{3,1} + D_{4,1}).$$

It is worth mentioning that the determination of an explicit form for the terms in the series of (39) is in general very difficult even when k is small and when the intensities λ_i are equal. Hence, in view of possible future developments, in a forthcoming investigation we aim to apply computational methods to determine the related probabilities.

In conclusion, we stress that the analysis of finite-velocity random motions in multidimensional domains deserves interest in various applied fields. In particular, the motion of a particle in a three-dimensional space as studied in this paper provides possible applications also in chemistry, since the tetrahedral geometry characterizes the shape of many molecules. Indeed, the motion of a particle is isotropic, i.e., the direction of its movement is uniformly distributed on the unit sphere in \mathbb{R}^3 . Thus, according to this interpretation, the Eq. (14) can be employed to represent the wave function of the electron.

Further interesting real applications of finite-velocity random motions for modelling random occurrences of events in time and space are related, for instance, to biology (for the random motions of microorganisms), to geology (for alternating trends in volcanic areas), and to physics (for the vorticity motion in two or more dimensions, see Orsingher and Ratanov (2008)). Moreover, finite-velocity random motions in \mathbb{R}^3 are also useful to describe the movements of particles in gases, see, for instance, Reimberg and Abramo (2013), where the authors introduce an application to the study of photon propagation in the Cosmis Microwave Background (CMB) radiation. At least, using a similar approach, relevant real applications in higher dimensions concerning cyclic random motions under a GCP can be explored in future works even if the computation complexity in the resolution of the probability law is a very hard task. Hence, in our view the finite-velocity random motion discussed here is an extension of the telegraph process to the space \mathbb{R}^3 and it is probably one of the possible ways for which the explicit distribution of the position $\boldsymbol{X}(t)$, under the assumption introduced in Section 2.1, can be determined.

Acknowledgments. The authors are members of the group GNCS of INdAM (Istituto Nazionale di Alta Matematica).

Declarations

Competing Interest The authors have no other relevant financial or non-financial interests to disclose.

Funding This work is partially supported by INdAM-GNCS (project "Modelli di shock basati sul processo di conteggio geometrico e applicazioni alla sopravvivenza", CUP-E55F22000270001). This work is partially supported by PRIN 2022 PNRR, project P2022XSF5H "Stochastic Models in Biomathematics and Applications".

Author Contribution statement AI and GV contributed equally to this work.

Data Availability statement Not applicable.

References

- Asmussen, S. R.: Steady-State Properties of GI/G/1. Applied Probability and Queues. Stoch. Model. Appl. Probab., **51**, 266-301 (2003).
- Beghin, L., Nieddu, L. and Orsingher, E.: Probabilistic analysis of the telegrapher's process with drift by means of relativistic transformations. J. Appl. Math. Stoch. Anal., 14(1), 11-25 (2001).
- Benson D. A., Schumer R., Meerschaert M. M.: Recurrence of extreme events with power-law interarrival times. Geophys. Res. Lett., **34**:L16404 (2007).
- Cha, J. H. and Finkelstein, M.: A note on the class of geometric counting processes. Probab. Eng. Inf. Sci., **27**(2), 177-185 (2013).
- Cinque, F. and Orsingher, E.: Random motions in \mathbb{R}^3 with orthogonal directions. Stoch. Process. Their Appl., **161**, 173-200 (2023).
- Cinque, F. and Orsingher, E.: Stochastic dynamics of generalized planar random motions with orthogonal directions. J. Theor. Probab., 1-33 (2023).
- Clegg R. G., Di Cairano-Gilfedder C. and Shi Z.: A critical look at power law modelling of the Internet. Comput. Commun., **33**:259-268 (2010).
- Di Crescenzo, A. Exact transient analysis of a planar random motion with three directions. Stoch. Stoch. Rep., **72**(3-4), 175-189 (2002).
- Di Crescenzo, A., Iuliano, A. and Mustaro, V. On some finite-felocity random motions driven by the geometric counting process. J. Stat. Phys., **190**(3), 44 (2023).
- Di Crescenzo, A. and Pellerey, F. Some results and applications of geometric counting processes. Methodol. Comput. Appl. Probab., **21**(1), 203-233 (2019).
- Hartmann, A. K., Majumdar, S. N., hawe, H. and Schehr G.: The convex hull of the run-and-tumble particle in a plane. J. Stat. Mech. Theory Exp., **2020**(5), 053401 (2020).
- Kolesnik, A. D.: The equations of Markovian random evolution on the line. J. Appl. Probab., **35**(1), 27-35 (1998).
- Kolesnik, A. D.: Discontinuous term of the distribution for Markovian random evolution in \mathbb{R}^3 . Bul. Acad. de Stiinte Republicii Mold. Mat., 51(2), 62-68 (2006).
- Kolesnik, A. D. and Turbin, A. F.: The equation of symmetric Markovian random evolution in a plane. Stoch. Process. Their Appl., **75**(1), 67-87 (1998).
- Kolesnik, A. D. and Orsingher, E.: Analysis of a finite-velocity planar random motion with reflection. Theory Probab. its Appl., **46**(1), 132-140 (2002).

- Kolesnik, A. D. and Ratanov, N.: Telegraph Processes and Option Pricing. 2nd edn. Springer, Berlin (2023).
- Lavergnat J., Golé P.: A stochastic raindrop time distribution model. J. Appl. Meteorol., 37:805-818
- Lachal, A.: Cyclic random motions in-space with n directions. ESAIM Probab. Stat., **10**, 277-316 (2006).
- Lachal, A., Leorato, S. and Orsingher, E.: Minimal cyclic random motion in \mathbb{R}^n and hyper-Bessel functions. Ann. Inst. Henri Poincare (B) Probab., 42(6), 753-772 (2006).
- Martens, K., Angelani, L., Di Leonardo, R. and Bocquet, L.: Probability distributions for the run-and-tumble bacterial dynamics: An analogy to the Lorentz model. Eur. Phys. J. E., **35**, 1-6 (2012).
- Masoliver, J., Porra, J. M. and Weiss, G. H.: Some two and three-dimensional persistent random walks. Phys. A: Stat. Mech. Appl., 193(3-4), 469-482 (1993).
- Orsingher, E.: A planar random motion governed by the two-dimensional telegraph equation. J. Appl. Probab., 23(2), 385-397 (1986).
- Orsingher, E.: Probability law, flow function, maximum distribution of wave-governed random motions and their connections with Kirchoff's laws. Stoch. Process. Their Appl., **34**(1), 49-66 (1990).
- Orsingher, E.: Bessel functions of third order and the distribution of cyclic planar motions with three directions. Stoch. Stoch. Rep., **74**, 617-631 (2002).
- Orsingher, E. and De Gregorio, A.: Random flights in higher spaces. J. Theor. Probab., **20**, 769-806 (2007).
- Orsingher, E., Garra, R. and Zeifman, A. I.: Cyclic random motions with orthogonal directions. Markov Process. Relat. Fields., **26**(3), 381-402 (2020).
- Orsingher, E., and Ratanov, N.: Random motions in inhomogeneous media. Theory Probab. Math. Stat., **76**, 141-153 (2008).
- Orsingher, E. and Sommella, A. M.: A cyclic random motion in \mathbb{R}^3 with four directions and finite velocity. Stoch. Stoch. Rep., **76**(2), 113-133 (2004).
- Pogorui, A. A.: Evolution in multidimensional spaces. Random Oper. Stoch., **20**(2), 135-141 (2012).
- Pogorui, A. A. and Rodríguez-Dagnino, R. M.: Isotropic random motion at finite speed with K-Erlang distributed direction alternations. J. Stat. Phys., **145**, 102-112 (2011).

- Pogorui, A. A. and Rodríguez-Dagnino, R. M.: Distribution of random motion at renewal instants in three-dimensional space. J. Math. Sci., **254**, 416-424 (2021).
- Reimberg, P. H. and Abramo, L. R. CMB and random flights: temperature and polarization in position space. J. Cosmol. Astropart. Phys., **2013**(06), 043 (2013).
- Santra, I., Basu, U. and Sabhapandit, S.: Run-and-tumble particles in two dimensions: Marginal position distributions. Phys. Rev. E., 101(6), 062120 (2020).