

A three-dimensional cyclic random motion with finite velocities driven by geometric counting processes

Antonella Iuliano^{(1)*}, and Gabriella Verasani⁽¹⁾

(1) Dipartimento di Matematica, Informatica ed Economia
 Università degli studi della Basilicata, I-85100 Potenza, Italy
 Email: {antonella.iuliano, gabriella.verasani}@unibas.it

Abstract

We consider a stochastic process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ which describes a particle performing a minimal cyclic random motion with finite velocities in \mathbb{R}^3 . The particle can take four directions moving with different velocities \vec{v}_j , for $1 \leq j \leq 4$, so that the diffusion region is a tetrahedron $\mathcal{T}(t)$. Moreover, we assume that the sequence of sojourn times along each velocity \vec{v}_j follows a geometric counting process of intensity λ_j , $1 \leq j \leq 4$. We first describe the direction vectors \vec{v}_j and the domain $\mathcal{T}(t)$; then, we obtain the closed-form expressions of the initial and absolutely continuous components of the probability law of the process when the starting velocity is \vec{v}_1 . We also investigate the limiting behavior of the probability density of the process when the intensities λ_j tend to infinity. Finally, we introduce the first-passage time problem for the first component of $\mathbf{X}(t)$ through a constant positive boundary $\beta > 0$ providing the bases for future developments.

Keywords: random motion, Geometric counting process, Telegraph process, First-passage time

MSC Classification: 60K99, 60K50

Orcid: Antonella Iuliano: 0000-0001-8541-8120 ;
 Gabriella Verasani: 0009-0000-4994-1694

1 Introduction

In the last decades, finite-velocity random motions have been widely studied as a natural class of stochastic processes to model real phenomena on multi-dimensional Euclidean spaces. In the one-dimensional case, the model that describe these motions is the telegraph process, in which the changes of the two possible velocities are governed by the Poisson process (see, for instance, Orsingher [20], Kolesnik [9], Beghin et al. [1] and Kolesnik and Ratanov [13]).

*Corresponding author

Later, the study of random motions in two or more dimensions has been performed by many authors over the years. In particular, finite-velocity planar random processes in continuous time have been investigated by Orsingher [19] and Masoliver et al. [18]. Further investigations have been oriented to investigate the distribution for Markovian random motion in the plane (c.f. Kolesnik [11] and [10]), and the random motion with possible reflections at Poisson-paced events (c.f. Kolesnik and Orsingher [12]). The analysis of these random motions with finite speed is performed analytically using partial differential equations and a probabilistic approach based on order statistics, or on more general renewal processes as in Di Crescenzo [5] (with arbitrary random steps between successive switches).

Several generalizations of the basic model have been proposed especially in biology and physics, motivated by the need of describing a variety of random motions performed by cells, micro-organisms and animals (see, for instance, Martens et al. [17], Hartmann et al. [8], Santra et al. [28] and Pogorui and Rodríguez-Dagnino [27]).

The main aim for the study of n -dimensional random motions is the determination of the probability distribution of the position vector $(X_1(t), X_2(t), \dots, X_n(t)) \in \mathbb{R}^n$ of a particle at time t . For instance, cyclic planar motions with three and four directions have been treated by Orsingher [21], and Orsingher and Sommella [24] in which the changes of direction are governed by a homogenous Poisson process. Moreover, the authors have found a connection between the equations governing the probability distributions and the Bessel functions of higher order. The probability law of the motion of a particle performing a cyclic random motion in \mathbb{R}^n is also determined in Lachal [14] where the particle can take n possible directions with different speeds. Here, the changes of direction occur at Poisson random times forming a polyhedron. For instance, see also Lachal et al. [15] in which the probability distribution is obtained by using order statistics and is expressed in terms of hyper-Bessel functions of higher order (see, for instance, Leorato and Orsingher [16]). Recently, planar random motions with orthogonal directions switching at Poisson time have been examined in Orsingher et al. [23], Cinque and Orsingher [3, 4]. Other remarkable results concerning random motions in high dimensional spaces with uniformly distributed directions are present in Orsingher and De Gregorio [22], Pogorui [25] and Pogorui and Rodríguez-Dagnino [26].

Some modified versions of the telegraph process have been considered also by substituting the underlying Poisson process with different kinds of counting processes. For instance, Di Crescenzo et al. [6] study a stochastic process which describes the dynamics of a particle performing a finite-velocity random motion whose velocities alternate cyclically along two and three different directions with possibly unequal velocities. The novelty of this last paper is that the number of displacements of the motion along each possible direction follows a Geometric Counting Process (GCP) (see, Cha and Finkelstein [2]) in \mathbb{R} and \mathbb{R}^2 . The advantage of this new scheme allows in dealing with phenomena where the times separating consecutive events do not possess the memoryless property, since in many real cases their distributions are characterized by heavy tails.

Therefore, starting from the results obtained in [6], here, we propose an extension of the classical telegraph process in \mathbb{R}^3 with the aim of determining the explicit general laws of the distribution of the current position. We consider four speeds alternating cyclically where the intertimes between two subsequent changes of directions are distributed as a GCP. Specifically, the intertimes possess a heavy-tailed distribution, differently to the classical telegraph process in which the times between velocity changes have exponential distribution.

This scheme is useful in describing suitable dynamics where random occurrences between two events have infinite expectations, and are dependent. Thus, the proposed stochastic process provides new models for the description of phenomena that are no more governed by hyperbolic PDE's as the classical telegraph equation. Another benefit of the present study is the construction of new solvable models, whose probability laws are obtained in tractable and closed form (as in the special case with symmetry) using an approach based on the analysis of the intertimes between consecutive velocity changes.

In detail, we analyze the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ in \mathbb{R}^3 which describes the position of randomly moving particle performing a cyclic alternating motion with four specific and possible different directions \vec{v}_j , for $1 \leq j \leq 4$. The velocity-vectors form a (possible irregular) tetrahedron, say $\mathcal{T}(t)$, i.e., the set of all possible positions of the moving particle at time t . We remark that the analysis of random motion in multidimensional Euclidean spaces is quite rare in the literature. Once defined the geometry of the region $\mathcal{T}(t)$, the probability law of the process is determined by (i) the initial components given by three terms, describing the situations in which the particle is found on the vertices, edges and faces of $\mathcal{T}(t)$ at the beginning of its motion, and (ii) the density concerning the absolutely continuous part related to the motion of the particle in the interior of $\mathcal{T}(t)$. In particular, for the latter absolutely continuous component we exhibit an integral representation involving the probability density functions and the conditional survival function of the intertimes between two successive events.

This is the plan of the paper. In Section 2 we introduce the probability densities of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ describing the position of a particle performing a cyclic, minimal, random motion in \mathbb{R}^3 with four possible directions \vec{v}_j , for $1 \leq j \leq 4$, and constant velocity $c > 0$. Some preliminary results on the GCP are also briefly illustrated with reference to the distribution of the intertimes between two consecutive events. Then, in Section 3, we study the directional vectors and the analytic representation of the diffusion region identifying the tetrahedron $\mathcal{T}(t)$. In Section 4 we investigate the stochastic process and its probability laws with underlying GCP, and determine the explicit expression of the initial and absolutely continuous components. Section 5 illustrates a special case with four fixed cyclic directions forming a regular tetrahedron in \mathbb{R}^3 . Moreover, we examine the limiting distribution of the process when the parameters of the intertimes tend to infinity. Finally, in Section 6 we provide some basic lines for the study of the first-passage-time problem of the first component of the process through a constant positive boundary $\beta > 0$.

2 A finite cyclic random motion in \mathbb{R}^3

Let $\mathbf{X}(t) = (X_1(t), X_2(t), X_3(t))$ and $V(t)$ be respectively the position and the velocity of the particle at an arbitrary time $t \geq 0$ in the space \mathbb{R}^3 . Assume that each point in \mathbb{R}^3 is represented by a triple $\mathbf{x} = (x_1, x_2, x_3)$. We consider a cyclic random motion $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ performed by a particle which can take four possible directions \vec{v}_j , for $1 \leq j \leq 4$, and moves with constant velocity $c > 0$. The motion is cyclic in the sense that at each event the particle switches from direction \vec{v}_j to \vec{v}_{j+1} , for $j = 1, 2, 3$, and then from \vec{v}_4 to \vec{v}_1 , and so on. Let $D_{j,k}$ be the random duration of the k -th time interval during which the motion proceeds with velocity c . For any $1 \leq j \leq 4$, the set $D_{j\cdot} := \{D_{j,k}, k \in \mathbb{N}\}$ constitutes a sequence of

non-negative and dependent absolutely continuous random variables such that

$$D_j^{(0)} = 0, \quad D_j^{(k)} = \sum_{i=1}^k D_{j,i}, \quad k \in \mathbb{N}. \quad (1)$$

Moreover, the sets $D_{j,\cdot}$, for $1 \leq j \leq 4$, are mutually independent.

Now, let $\{N(t), t \geq 0\}$ be the alternating counting process having arrival times T_1, T_2, \dots (i.e., the instants when the events occur), such that $N(t)$ counts the total number of velocity reversals of the particle in $[0, t]$, i.e.

$$N(0) = 0, \quad N(t) = \sum_{k=0}^{\infty} \mathbf{1}_{\{T_k \leq t\}}, \quad t \geq 0. \quad (2)$$

In order to obtain the probability law of the stochastic process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ in \mathbb{R}^3 , describing the motion of a particle along four directions, we consider the position $\mathbf{X}(t)$ and the velocity $V(t)$ of a particle at time $t \geq 0$. Specifically, we have

$$V(t) = \sum_{k=0}^{\infty} \mathbf{1}_{\{T_{k-1} \leq t < T_k\}} \vec{v}_k, \quad (3)$$

with $V(0) \in \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4\}$, and

$$\mathbf{X}(t) = \mathbf{X}(0) + \int_0^t V(s) ds = \sum_{k=1}^{N(t)-1} (T_k - T_{k-1}) \vec{v}_k + (t - T_{N(t)-1}) \vec{v}_{N(t)}, \quad (4)$$

with $\mathbf{X}(0) = \mathbf{0} \equiv (0, 0, 0)$ and $\sum_{k=1}^{N(t)-1} (T_k - T_{k-1}) = T_{N(t)-1}$, since $T_0 = 0$. The sum in Eq. (4) refers to the case where at least one change of direction has occurred before t , while the last term is related to the displacement along the current direction at time t .

We indicate with T_k the k -th random instant in which the motion modifies its direction, for $n \in \mathbb{N}_0$. Remembering Eq. (1) the following identity holds:

$$T_{4n+j} = \sum_{r=1}^4 D_r^{(n+m_{j,r})}, \quad (5)$$

where, for fixed $j \in \{1, 2, 3, 4\}$, $M = (m_{j,r})_{1 \leq r \leq 4} \in \mathbb{R}^{4 \times 4}$ is equal to

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}.$$

To define the conditional distributions of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$, we denote by

$$C_i := \{\mathbf{X}(0) = \mathbf{0}, V(0) = \vec{v}_i\}, \quad (6)$$

the event that the particle starts its motion from the origin $\mathbf{0}$ when the initial velocity is \vec{v}_i , $1 \leq i \leq 4$. Therefore, let \mathbb{P}_i be the probability conditional on C_i .

Remark 2.1. *The conditional probability laws given C_i are composed by two contributions.*

- (i) *The initial components given by three terms related to the beginning of the motion. Specifically, they describe the cases in which the particle lies on the vertices, or it is located on some edge, or it reaches one of the faces of the diffusion region.*
- (ii) *The absolutely continuous component related to the particle motion in the internal part of the diffusion region.*

For the case (ii) of Remark 2.1 we define

$$\begin{aligned} p_{ij}(\mathbf{x}, t)d\mathbf{x} &= \mathbb{P}_i\{\mathbf{X}(t) \in d\mathbf{x}, V(t) = \vec{v}_j\} \\ &= \mathbb{P}_i\{X_1(t) \in dx_1, X_2(t) \in dx_2, X_3(t) \in dx_3, V(t) = \vec{v}_j\}, \end{aligned} \quad (7)$$

with $1 \leq i, j \leq 4$ and where $d\mathbf{x}$ is the infinitesimal element in the space \mathbb{R}^3 with the Lebesgue measure $\mu(d\mathbf{x}) = dx_1 dx_2 dx_3$ (i refers to the initial velocity, and j to current velocity at time t). The right-hand-side of Eq. (7) represents the probability that the particle at time $t > 0$ is located in a neighborhood of $\mathbf{x} \in \mathbb{R}^3$ and moves along direction \vec{v}_j , given the initial condition represented by C_i . Moreover, we can introduce the probability density function (p.d.f.) of the particle location $\mathbf{X}(t)$ at time $t > 0$, i.e.,

$$p_i(\mathbf{x}, t) = \mathbb{P}_i\{\mathbf{X}(t) \in d\mathbf{x}\}, \quad 1 \leq i \leq 4. \quad (8)$$

Due to Eqs. (7) and (8), one immediately has

$$p_i(\mathbf{x}, t) = \sum_{j=1}^4 p_{ij}(\mathbf{x}, t), \quad 1 \leq i \leq 4. \quad (9)$$

2.1 The distribution of the intertimes

We introduce some preliminary results on the GCP (for more details, see [2] and [7]). Specifically, we assume that the random intertimes between consecutive changes of directions are governed by possible different GCPs (see, for instance, [6]). Differently from the Poisson process, the GCP does not possess the property of independent increments, which usually complicates the probabilistic analysis. Moreover, the intertimes of the GCP have modified Pareto distributions, thus such process is suitable to be applied in those area where random occurrences between events have infinite expectations, and are not independent.

For this purpose, we consider a mixed Poisson process $\{\tilde{N}(t), t \geq 0\}$ characterized by the following marginal distribution expressed as a mixture:

$$\mathbb{P}[\tilde{N}_\lambda(t) = k] = \int_0^t \mathbb{P}[N^{(\alpha)}(t) = k] dU_\lambda(\alpha), \quad t \geq 0, \quad k \in \mathbb{N}_0, \quad (10)$$

where $\tilde{N}_\lambda(t)$ is a Poisson process with intensity α and where U_λ is an exponential distribution with mean $\lambda \in \mathbb{R}^+$. The process $\{\tilde{N}_\lambda(t)\}$ is called a GCP with intensity λ , according to [2]. In particular, its probability distribution satisfies the following properties:

- at time $t = 0$, one has the case $\tilde{N}_\lambda(0) = 0$;

- for all $s, t \geq 0$ and $k \in \mathbb{N}_0$ holds:

$$\mathbb{P}\{\tilde{N}_\lambda(t+s) - \tilde{N}_\lambda(t) = k\} = \frac{1}{1+\lambda s} \left(\frac{\lambda s}{1+\lambda s} \right)^k. \quad (11)$$

Let $\tilde{T}_{n,\lambda}$, with $n \in \mathbb{N}$, be the random times denoting the arrival instants of the process $\tilde{N}_\lambda(t)$ such that $\tilde{T}_{0,\lambda} = 0$. It is interesting to observe that the probability density function (p.d.f.) of $\tilde{T}_{n,\lambda}$ is expressed as follows:

$$f_{\tilde{T}_{n,\lambda}}(t) = n \left(\frac{\lambda t}{1+\lambda t} \right)^{n-1} \frac{\lambda}{(1+\lambda t)^2}, \quad t \geq 0. \quad (12)$$

Differently from the Poisson process, the GCP does not have the property of independent increments, therefore, $\tilde{N}_\lambda(t)$ possesses a dependence structure which means that the increments between two consecutive events $\tilde{D}_{n,\lambda} = \tilde{T}_{n,\lambda} - \tilde{T}_{n-1,\lambda}$, with $n \in \mathbb{N}$, are dependent. More precisely, the marginal p.d.f. of the random durations $D_{j,n}$, $n \in \mathbb{N}$, has a modified Pareto distribution of type I, i.e.

$$f_{D_{j,n}}(t) = \frac{\lambda_j}{(1+\lambda_j t)^2}, \quad t \geq 0. \quad (13)$$

In order to obtain the probability distribution of the process $\mathbf{X}(t)$, we consider the conditional survival function of $\tilde{D}_{n,\lambda}$ conditional on $\tilde{T}_{n-1,\lambda} = t$, for $n \in \mathbb{N}$, given by the following expression

$$\bar{F}_{\tilde{D}_{n,\lambda}|\tilde{T}_{n-1,\lambda}}(s|t) = \left(\frac{1+\lambda t}{1+\lambda(t+s)} \right)^n, \quad s, t \geq 0, \quad (14)$$

and the corresponding probability density function (p.d.f.) of $\tilde{D}_{n,\lambda}$ conditional on $\tilde{T}_{n-1,\lambda} = t$, for $n \in \mathbb{N}$, i.e.,

$$f_{\tilde{D}_{n,\lambda}|\tilde{T}_{n-1,\lambda}}(s|t) = \frac{n\lambda(1+\lambda t)^n}{[1+\lambda(t+s)]^{n+1}}, \quad s, t \geq 0. \quad (15)$$

For more details about Eq. (14) and Eq. (15) see, for instance, [7]. Therefore, considering the stochastic process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ and assuming the intertimes' dependence, we determine the initial and absolutely continuous component of a cyclic alternating random motion along four directions \vec{v}_j , for $1 \leq j \leq 4$ driven by independent GCPs with intensity λ_j ($1 \leq j \leq 4$).

3 The velocity vectors and the diffusion region

In this section, we discuss the properties of the directional velocity vectors and the geometrical features of the region generated by them, i.e., the set of all possible positions of the moving particle at time $t > 0$.

3.1 The directional vectors

Let \vec{v}_j , $1 \leq j \leq 4$, be the vectors representing the possible directions of the cyclic motion in \mathbb{R}^3 (i.e., the minimal number of directions for a non trivial motion in \mathbb{R}^3). The velocity direction coordinates can be defined by using the spherical coordinates, given by

$$\vec{v}_j = (\vec{l} \cos \theta_j \sin \varphi_j + \vec{m} \sin \theta_j \sin \varphi_j + \vec{n} \cos \varphi_j), \quad (16)$$

where $\vec{l}, \vec{m}, \vec{n}$ are the unit vectors along the Cartesian coordinate axes in a three-dimensional space \mathbb{R}^3 , φ_j is the azimuthal angle in the xy -plane from the x -axis with $\theta_j \in [0, 2\pi]$ and θ_j is the polar angle from the positive z -axis with $\varphi_j \in [0, \pi]$, for $1 \leq j \leq 4$.

The particle starts from the origin $\mathbf{0} = (0, 0, 0) \in \mathbb{R}^3$ at time $t = 0$, running with constant velocity $c > 0$. The particle at time $T_0 = 0$ begins its motion along the direction \vec{v}_1 for a random duration $D_{1,1}$. At instant T_1 the particle changes the velocity along \vec{v}_2 for a random period $D_{2,1}$. At time T_2 the particle switches the direction along \vec{v}_3 for a random time $D_{3,1}$. At instant T_3 the particle proceeds the running with speed \vec{v}_4 for a length of time $D_{4,1}$ and so on by achieving cyclically the directions \vec{v}_j for the random periods $D_{j,2}, D_{j,3}, D_{j,4}, \dots$, such that,

$$D_{j,4k+i} \stackrel{d}{=} D_{j,i}, \quad (17)$$

where $\stackrel{d}{=}$ means equality in distribution for each $1 \leq j \leq 4$ and $k \in \mathbb{N}$. Hence, during the n -th cycle the particle moves along directions \vec{v}_j in sequence for the random lengths $D_{j,n}$, with $1 \leq j \leq 4$ and $n \in \mathbb{N}$. In other words, the particle moves along direction \vec{v}_j and, after an intertime distributed as a CGP, it takes the direction \vec{v}_{j+1} , with $\vec{v}_{j+4n} = \vec{v}_j$, for $1 \leq j \leq 4$ and $n \in \mathbb{N}$. The results concerning the cases of other initial direction, i.e., $\vec{v}_2, \vec{v}_3, \vec{v}_4$, may be easily deduced. Now, in order to define the diffusion region, i.e., the set of all possible positions of the moving particle at time t , we consider the following Remark.

Remark 3.1. Let be $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ the stochastic process defined in Section 2. Given the direction vectors \vec{v}_j with $\vec{v}_{j+4n} = \vec{v}_j$, for $1 \leq j \leq 4$ and $n \in \mathbb{N}$, we have that the particle motion reaches any state of \mathbb{R}^3 in a sufficiently large time $t > 0$, if and only if, for a fixed j ,

(i) the set of direction vectors $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ are linearly independent;

(ii) the direction vector $\vec{v}_4 \in S$ with

$$S = \left\{ \xi(-\vec{v}_1) + \eta(-\vec{v}_2) + (1 - \xi - \eta)(-\vec{v}_3), 0 \leq \xi, \eta \leq 1, \xi + \eta \leq 1 \right\}. \quad (18)$$

Hence, the set of all possible positions under the hypothesis of Remark 3.1 identifies as state space at time t a tetrahedron whose vertices coincide with the endpoints of the vectors $ct\vec{v}_j$. In other words, the tetrahedron is defined as a 3-dimensional simplex where the interior points are convex combinations of the four vertices such that $\sum_{j=1}^4 \alpha_j \vec{v}_j$ with $\sum_{j=1}^4 \alpha_j = 1$ and $\alpha_j \geq 0$.

3.2 Analytic representation of the diffusion region

Let us consider the points $A_j(t)$, $1 \leq j \leq 4$, defined by $\overrightarrow{\mathbf{0}A_j(t)} = \vec{v}_j t$, where $\mathbf{0} = (0, 0, 0)$ is the origin of the Cartesian coordinate system. The points $A_j(t)$, $1 \leq j \leq 4$, given by

$$A_j(t) = (ct \cos \theta_j \sin \varphi_j, ct \sin \theta_j \sin \varphi_j, ct \cos \varphi_j), \quad 1 \leq j \leq 4 \quad (19)$$

are the vertices of the diffusion region of the particle at time t , i.e., a time-dependent tetrahedron with edges $E_{ij}(t)$ and faces $F_{ijk}(t)$, $1 \leq i, j, k \leq 4$ and $i < j < k$. In particular, given three different vertices $A_i(t)$, $A_j(t)$ e $A_k(t)$, for each $i \neq j \neq k$, the points are not aligned. This is true since the matrix generated by two adjacent segments $\overrightarrow{A_i(t)A_k(t)}$ and $\overrightarrow{A_k(t)A_j(t)}$ has maximum rank, i.e. 2. Therefore, under the assumptions of Remark 3.1 the equations linking two adjacent vertices can be expressed as

$$a_{ijk}x_1 + b_{ijk}x_2 + c_{ijk}x_3 - ctq_{ijk} = 0, \quad 1 \leq i, j, k \leq 4, \quad i < j < k, \quad (20)$$

where

$$a_{ijk} = (\sin \theta_j \sin \varphi_j - \sin \theta_i \sin \varphi_i)(\cos \varphi_k - \cos \varphi_i) - (\sin \varphi_k \sin \theta_k - \sin \varphi_i \sin \theta_i)(\cos \varphi_j - \cos \varphi_i), \quad (21)$$

$$b_{ijk} = (\sin \varphi_k \cos \theta_k - \sin \varphi_i \cos \theta_i)(\cos \varphi_j - \cos \varphi_i) - (\sin \varphi_j \cos \theta_j - \sin \varphi_i \cos \theta_i)(\cos \varphi_k - \cos \varphi_i), \quad (22)$$

$$c_{ijk} = \sin \varphi_i \cos \theta_i (\sin \varphi_j \sin \theta_j - \sin \varphi_k \sin \theta_k) + \sin \varphi_j \cos \theta_j (\sin \varphi_k \sin \theta_k - \sin \varphi_i \sin \theta_i) + \sin \varphi_k \cos \theta_k (\sin \varphi_i \sin \theta_i - \sin \varphi_j \sin \theta_j), \quad (23)$$

and,

$$q_{ijk} = a_{ijk} \cos \theta_i \sin \varphi_i + b_{ijk} \sin \theta_i \sin \varphi_i + c_{ijk} \cos \varphi_i. \quad (24)$$

We recall that $V(t)$ is the current direction of motion at time t . Assuming that $V(0)$ takes value \vec{v}_1 with velocity $c > 0$, we are able to determine the analytic expression of the diffusion region in which the particle is confined forming a tetrahedron defined as

$$\mathcal{T}(t) = \left\{ (x_1, x_2, x_3) \in \mathbb{R}^3 : \begin{cases} a_{123}x_1 + b_{123}x_2 + c_{123}x_3 - ctq_{123} \geq 0 \\ a_{124}x_1 + b_{124}x_2 + c_{124}x_3 - ctq_{124} \geq 0 \\ a_{234}x_1 + b_{234}x_2 + c_{234}x_3 - ctq_{234} \geq 0 \\ a_{134}x_1 + b_{134}x_2 + c_{134}x_3 - ctq_{134} \leq 0 \end{cases} \right\}, \quad (25)$$

where the given conditions derive from Eqs. (20). In general, the particle motion includes different mutually exclusive cases based on the initial assumptions $\mathbf{X}(0) = \mathbf{0}$ and $V(0) = \vec{v}_j$, for a fixed $1 \leq j \leq 4$. More precisely, the particle is situated on the vertices $A_j(t)$ of $\mathcal{T}(t)$ if no velocity change occurs up to time t , while if one event occurs then, at time t , it will be located on some edge E_{ij} of $\mathcal{T}(t)$. Two events allow the particle to reach one of the faces F_{ijk} of $\mathcal{T}(t)$ and three or more changes of direction force the particle to be located inside $\mathcal{T}(t)$. Therefore, recalling Remark 3.1, for every $\mathbf{x} \in \mathcal{T}(t)$, $t \geq 0$, we have:

- (i) the first component is related to the case when no event occurs in the interval $(0, t)$, i.e., $N(t) = 0$, and the particle is concentrated on the vertices of $\mathcal{T}(t)$;
- (ii) the second component is concerning the situation in which one event happens at time t , i.e., $N(t) = 1$, and the particle is located on some edge of $\mathcal{T}(t)$;
- (iii) the third component corresponds to the instance when two events occur, i.e., $N(t) = 2$ and the particle reaches one of the faces of $\mathcal{T}(t)$;
- (iv) When $N(t) \geq 3$, the particle is placed strictly in the interior of $\mathcal{T}(t)$, namely $\text{Int}(\mathcal{T}(t))$. Hence, for the absolutely continuous component of this case we refer to the density given in Eq. (8), i.e., $p_i(\mathbf{x}, t)$ for all $\mathbf{x} \in \text{Int}(\mathcal{T}(t))$, $t > 0$.

4 The probability law

In order to determine the probability law of the stochastic process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$, we assume that the starting direction at time t is \vec{v}_1 , so that initial condition is C_1 as defined in Eq. (6). Therefore, the following results are obtained by taking into account the cases (i)-(iv) considered above.

Theorem 4.1. *(Initial components) Let be $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ the stochastic process defined in Section 2. For $t \geq 0$ we have*

$$\begin{aligned} \mathbb{P}_1\{(X_1(t), X_2(t), X_3(t)) = (ct \cos \theta_j \sin \varphi_j, ct \sin \theta_j \sin \varphi_j, ct \cos \varphi_j), \\ V(t) = \vec{v}_1\} = \mathbb{P}\{D_{1,1} > t\}, \end{aligned} \quad (26)$$

$$\mathbb{P}_1\{\mathbf{X}(t) \in E_{12}(t), V(t) = \vec{v}_2\} = \mathbb{P}\{D_{1,1} < t \leq D_{1,1} + D_{2,1}\}, \quad (27)$$

and

$$\mathbb{P}_1\{\mathbf{X}(t) \in F_{123}(t), V(t) = \vec{v}_3\} = \mathbb{P}\{D_{1,1} + D_{2,1} < t \leq D_{1,1} + D_{2,1} + D_{3,1}\}, \quad (28)$$

where $D_{j,k}$ is the random duration of the k -th time interval during which the particle moves with velocity \vec{v}_j , for $1 \leq j \leq 3$ and $k \in \mathbb{N}$.

Proof. Eqs. (26), (27) and (28), when $V(0) = \vec{v}_1$, are easily obtained, under the assumptions (i), (ii) and (iii) of Section 3.2. \square

Using Eq. (7) and recalling (iv) of Section 3.2, we determine the absolutely continuous components of the probability law of the $X(t)$ conditioned on C_1 , for $t > 0$ and $1 \leq j \leq 4$, i.e.,

$$p_{1j}(\mathbf{x}, t)d\mathbf{x} = \mathbb{P}_1\{\mathbf{X}(t) \in d\mathbf{x}, V(t) = \vec{v}_j\}. \quad (29)$$

In order to give the expression of $p_{1j}(\mathbf{x}, t)$ we introduce the linear map $\zeta : \mathbb{R}^4 \longrightarrow \mathbb{R}^4$ defined by:

$$\zeta(\mathbf{t}) = \left(c \sum_{j=1}^4 x_{1\vec{v}_j} t_j, c \sum_{j=1}^4 x_{2\vec{v}_j} t_j, c \sum_{j=1}^4 x_{3\vec{v}_j} t_j, \sum_{j=1}^4 t_j \right), \quad (30)$$

for $\mathbf{t} = (t_1, t_2, t_3, t_4)$, and where $x_{1\vec{v}_j}$, $x_{2\vec{v}_j}$ and $x_{3\vec{v}_j}$ are the components of the vectors \vec{v}_j ($1 \leq j \leq 4$) respectively along the \mathbf{x} -axes. When a cycle of the random motion ends after a

period \mathbf{t} the function $\zeta(\mathbf{t})$, given in Eq. (30), provides a vector containing the displacements performed along the \mathbf{x} -axes during the cycle, as well as its whole duration.

Moreover, we consider the transformation matrix \mathbf{A} of function $\zeta(\mathbf{t})$ expressed in terms of spherical coordinates, i.e.,

$$\mathbf{A} = \begin{pmatrix} c \cos \theta_1 \sin \varphi_1 & c \cos \theta_2 \sin \varphi_2 & c \cos \theta_3 \sin \varphi_3 & c \cos \theta_4 \sin \varphi_4 \\ c \sin \theta_1 \sin \varphi_1 & c \sin \theta_2 \sin \varphi_2 & c \sin \theta_3 \sin \varphi_3 & c \sin \theta_4 \sin \varphi_4 \\ c \cos \varphi_1 & c \cos \varphi_2 & c \cos \varphi_3 & c \cos \varphi_4 \\ 1 & 1 & 1 & 1 \end{pmatrix} \quad (31)$$

with

$$\begin{aligned} \det \mathbf{A} = c^3 \{ & \sin \varphi_1 \sin \varphi_2 [\sin(\theta_2 - \theta_1)] (\cos \varphi_3 - \cos \varphi_4) \\ & + \sin \varphi_1 \sin \varphi_3 [\sin(\theta_3 - \theta_1)] (\cos \varphi_4 - \cos \varphi_2) \\ & + \sin \varphi_1 \sin \varphi_4 [\sin(\theta_4 - \theta_1)] (\cos \varphi_2 - \cos \varphi_3) \\ & + \sin \varphi_2 \sin \varphi_3 [\sin(\theta_3 - \theta_2)] (\cos \varphi_1 - \cos \varphi_4) \\ & + \sin \varphi_2 \sin \varphi_4 [\sin(\theta_4 - \theta_2)] (\cos \varphi_3 - \cos \varphi_1) \\ & + \sin \varphi_3 \sin \varphi_4 [\sin(\theta_4 - \theta_3)] (\cos \varphi_1 - \cos \varphi_2) \}. \end{aligned} \quad (32)$$

Remark 4.1. Recalling (19), we observe that the determinant $\det(t\mathbf{A})$ represents the volume $\text{Vol}(\mathcal{T}(t))$ of the diffusion region $\mathcal{T}(t)$ defined in (25).

Now, given a sample path of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$, we denote by $\tau_j = \tau_j(\mathbf{x})$ the non negative random variables representing the dwelling times of the particle motion in each direction \vec{v}_j ($1 \leq j \leq 4$) during $[0, t]$, so that

$$\sum_{j=1}^4 \tau_j = t. \quad (33)$$

Therefore, recalling (31) and (19), we can express τ_j , $1 \leq j \leq 4$, as the 4-uple

$$\zeta^{-1}(\mathbf{x}, t) = (\tau_1(\mathbf{x}, t), \tau_2(\mathbf{x}, t), \tau_3(\mathbf{x}, t), \tau_4(\mathbf{x}, t)), \quad (34)$$

with

$$\tau_j(\mathbf{x}, t) = \sum_{k=1}^4 \sigma_{jk} x_k + \sigma_{j4} t, \quad (35)$$

where $\sigma_{ij} = \frac{1}{\det \mathbf{A}} \text{cof}_{ij}(\mathbf{A})$ with $\text{cof}_{ij}(\mathbf{A}) = (-1)^{i+j} \det A_{ij}$, $1 \leq i, j \leq 4$. Moreover, since the function $\zeta(\mathbf{t})$, for all $\mathbf{x} \in \mathbb{R}^3$, is bijective (see, for more details, [14]), it is easy to see that $\boldsymbol{\tau} = (\tau_1, \tau_2, \tau_3, \tau_4)^T$ is solution of the system $\mathbf{A}\boldsymbol{\tau} = (\mathbf{x}, t)^T$. More precisely, in the following proposition the explicit form of Eq. (35) can be obtained.

Proposition 4.1. Making use of Eq. (35), for all $\mathbf{x} \in \mathbb{R}^3$ the coordinates τ_j , for $1 \leq j \leq 4$ and $c > 0$, are expressed as

$$\tau_j = \frac{c^2}{\det \mathbf{A}} \left[L_j x_1 + M_j x_2 + N_j x_3 - ct P_j \right], \quad 1 \leq j \leq 4, \quad (36)$$

where

$$\begin{aligned}
L_j &= \sum_{i=1}^3 \sin \varphi_{j\hat{+}i} \sin \theta_{j\hat{+}i} \left(\cos \varphi_{j\hat{+}(i+1)} - \cos \varphi_{j\hat{+}(i+2)} \right), \\
M_j &= \sum_{i=1}^3 \sin \varphi_{j\hat{+}i} \cos \theta_{j\hat{+}i} \left(\cos \varphi_{j\hat{+}(i+2)} - \cos \varphi_{j\hat{+}(i+1)} \right), \\
N_j &= \sum_{i=1}^3 \sin \varphi_{j\hat{+}i} \cos \theta_{j\hat{+}i} \left(\sin \varphi_{j\hat{+}(i+1)} \sin \theta_{j\hat{+}(i+1)} \right. \\
&\quad \left. - \sin \varphi_{j\hat{+}(i+2)} \sin \theta_{j\hat{+}(i+2)} \right), \\
P_j &= \sum_{i=1}^3 \sin \varphi_{j\hat{+}i} \cos \theta_{j\hat{+}i} \left(\sin \varphi_{j\hat{+}(i+1)} \sin \theta_{j\hat{+}(i+1)} \cos \varphi_{j\hat{+}(i+2)} \right. \\
&\quad \left. - \sin \varphi_{j\hat{+}(i+2)} \sin \theta_{j\hat{+}(i+2)} \cos \varphi_{j\hat{+}(i+1)} \right),
\end{aligned} \tag{37}$$

where $a\hat{+}b$ denotes $(a+b) \bmod 4$ and $\det \mathbf{A}$ can be recovered from (31).

Proof. Recalling Eqs. (19), (32) and (35), after straightforward calculations Eq. (36) is determined. \square

Now, for the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ we are able to formulate the following theorem about the absolutely continuous component concerning assumption (iv) of Section 3.2. For this purpose, we indicate with $f_{D_j^{(n)}}$ the probability density function (p.d.f.) of $D_j^{(n)}$ given in Eq. (1), $1 \leq j \leq 4$, and with $\bar{F}_{D_{j,k+1}|D_j^{(k)}}$ the conditional survival function of $D_{j,k+1}$ given $D_j^{(k)}$, for $1 \leq j \leq 4$ and $k \in \mathbb{N}_0$.

Theorem 4.2. (*Absolutely continuous component*) For the stochastic process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ defined in Section 2, under the initial condition given in Eq. (6) with direction \vec{v}_1 , we have that, for all $\mathbf{x} \in \text{Int}(\mathcal{T}(t))$ and $t > 0$, the absolutely continuous component of the probability law is given by

$$\begin{aligned}
p_{1j}(\mathbf{x}, t) &= \frac{1}{\det \mathbf{A}} \sum_{k=0}^{\infty} \left\{ \prod_{i=1}^{j-1} f_{D_i^{(k+1)}}(\tau_i) \prod_{i=j+1}^4 f_{D_i^{(k)}}(\tau_i) \right. \\
&\quad \left. \times \int_{t-\tau_j}^t f_{D_j^{(k)}}(s - (t - \tau_j)) \bar{F}_{D_{j,k+1}|D_j^{(k)}}(t - s | s - (t - \tau_j)) ds \right\},
\end{aligned} \tag{38}$$

where $\tau_j = \tau_j(\mathbf{x}, t)$, for $1 \leq j \leq 4$, is defined in Eq. (36), and $\det \mathbf{A}$ is given in Eq. (32).

Proof. By conditioning on the number of direction switches in $(0, t)$, say k , and on the last time s previous t in which the particle changes its velocity from \vec{v}_4 to \vec{v}_1 in order to restart

a new cycle, we reformulate Eq. (8), for $\mathbf{x} \in \mathbb{R}^3$ and $t > 0$, as follows:

$$\begin{aligned}
p_{1j}(\mathbf{x}, t) d\mathbf{x} &= \sum_{k=0}^{\infty} \int_0^t \mathbb{P}\{T_{4k+j-1} \in ds, X_1(s) + c(t-s) \cos \theta_j \sin \varphi_j \in dx_1, \\
&\quad X_2(s) + c(t-s) \sin \theta_j \sin \varphi_j \in dx_2, \\
&\quad X_3(s) + c(t-s) \cos \varphi_j \in dx_3, D_{j,k+1} > t-s\},
\end{aligned} \tag{39}$$

for $1 \leq j \leq 4$ and where T_{4k+j-1} is the instant occurring at time s in which the particle changes its direction to velocity \vec{v}_j . Thus, $\mathbf{X}(s) = (X_1(s), X_2(s), X_3(s))$ is the position of the particle when a change of direction occurs. Therefore, due to Eq.(5), we have

$$T_{4k+j-1} = \sum_{i=1}^{j-1} D_i^{(k+1)} + \sum_{i=j+1}^4 D_i^{(k)} + D_j^{(k)}. \tag{40}$$

Hence, for $s = T_{4k+j-1}$ and using Eq.(1), we can express the coordinates of the particle position $\mathbf{X}(s)$ as follows

$$\begin{aligned}
X_1(s) &= c \left[\sum_{i=1}^{j-1} D_i^{(k+1)} \cos \theta_i \sin \varphi_i + \sum_{i=j+1}^4 D_i^{(k)} \cos \theta_i \sin \varphi_i + D_j^{(k)} \cos \theta_j \sin \varphi_j \right], \\
X_2(s) &= c \left[\sum_{i=1}^{j-1} D_i^{(k+1)} \sin \theta_i \sin \varphi_i + \sum_{i=j+1}^4 D_i^{(k)} \sin \theta_i \sin \varphi_i + D_j^{(k)} \sin \theta_j \sin \varphi_j \right], \\
X_3(s) &= c \left[\sum_{i=1}^{j-1} D_i^{(k+1)} \cos \varphi_i + \sum_{i=j+1}^4 D_i^{(k)} \cos \varphi_i + D_j^{(k)} \cos \varphi_j \right].
\end{aligned} \tag{41}$$

The conditions defined in (25) and $\mathbf{X}(s) \in \mathcal{T}(s)$ implies that $s > t - \tau_j$, for $1 \leq j \leq 4$. Moreover, using (31) and substituting Eqs. (41) in Eq. (39). we have:

$$\begin{aligned}
p_{1j}(\mathbf{x}, t) &= \frac{1}{\det \mathbf{A}} \sum_{k=0}^{\infty} \left\{ \int_{t-\tau_j}^t \Psi_{j,k} \left[s, x_1 - c(t-s) \sin \varphi_j \cos \theta_j, \right. \right. \\
&\quad \left. \left. x_2 - c(t-s) \sin \varphi_j \cos \theta_j, x_3 - c(t-s) \cos \varphi_j \right] \right. \\
&\quad \times \mathbb{P} \left[D_{j,k+1} > t-s \mid T_{4k+j-1} = s, X_1(T_{4k+j-1}) = x_1 - x_{1\vec{v}_j}(t-s), \right. \\
&\quad \left. \left. X_2(T_{4k+j-1}) = x_2 - x_{2\vec{v}_j}(t-s), X_3(T_{4k+j-1}) = x_3 - x_{3\vec{v}_j}(t-s) \right] ds \right\},
\end{aligned} \tag{42}$$

where $\Psi_{j,k}$ is the joint p.d.f. of $(T_{4k+j-1}, X_1(T_{4k+j-1}), X_2(T_{4k+j-1}), X_3(T_{4k+j-1}))$ with:

$$\begin{aligned}
T_{4k+j-1} &= \sum_{i=1}^{j-1} D_i^{(k+1)} + \sum_{i=j+1}^4 D_i^{(k)} + D_j^{(k)}, \\
X_1(T_{4k+j-1}) &= \sum_{i=1}^{j-1} x_{1\vec{v}_i} D_i^{(k+1)} + \sum_{i=j+1}^4 x_{1\vec{v}_i} D_i^{(k)} + x_{1\vec{v}_j} D_j^{(k)}, \\
X_2(T_{4k+j-1}) &= \sum_{i=1}^{j-1} x_{2\vec{v}_i} D_i^{(k+1)} + \sum_{i=j+1}^4 x_{2\vec{v}_i} D_i^{(k)} + x_{2\vec{v}_j}^{(k)} D_j^{(k)}, \\
X_3(T_{4k+j-1}) &= \sum_{i=1}^{j-1} x_{3\vec{v}_i} D_i^{(k+1)} + \sum_{i=j+1}^4 x_{3\vec{v}_i} D_i^{(k)} + x_{3\vec{v}_j} D_j^{(k)}.
\end{aligned} \tag{43}$$

According to (30), for $1 \leq j \leq 4$, we have

$$x_{1\vec{v}_i} = v_i \sin \varphi_i \cos \theta_i, \quad x_{2\vec{v}_i} = v_i \sin \varphi_i \sin \theta_i, \quad x_{3\vec{v}_i} = v_i \cos \varphi_i. \tag{44}$$

Moreover, given the mutual independence of the variables $\{D_{j,k}; k \in \mathbb{N}\}$, for $1 \leq j \leq 4$, we get

$$\begin{aligned}
\Psi_{j,k} &\left[s, x - c(t-s) \cos \theta_j \sin \varphi_j, y - c(t-s) \sin \theta_j \sin \varphi_j, z - c(t-s) \cos \varphi_j \right] \\
&= \prod_{i=1}^{j-1} f_{D_i^{(k+1)}}(\tau_i) \prod_{i=j+1}^4 f_{D_i^{(k)}}(t - \tau_j) f_{D_j^{(k)}}[s - (t - \tau_j)].
\end{aligned} \tag{45}$$

Therefore, Eq. (38) is directly obtained replacing Eq. (45) in Eq. (42). \square

We point out that Eq. (38) is in analogy with Eq. (4.1) of [14] in which a particular case of the minimal cyclic random motion in \mathbb{R}^d ($n = d + 1$ directions, $n \leq d$) is investigated. Specifically, the integral formulation given in Eq. (38) is based on the application of Eqs. (13) and (15) where the assumption of dependent increments, instead of the typical exponential distribution, is taken into account. Differently, in [24, 15] the explicit form of the absolutely continuous part of $\mathbf{X}(t)$ is determined but using an approach based on the Bessel functions of higher order.

Clearly, the results given in Theorems 4.1 and 4.2 for the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ can be extended similarly when the particle starts with velocity \vec{v}_l , for $2 \leq l \leq 4$. Using this new scheme, a fixed initial direction allows us to obtain tractable expressions for the probability law of the process as illustrated in the following section.

5 A special case

In this section we analyze a special case of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$, with initial condition defined by Eq. (6) when $V(0) = \vec{v}_1$ and for fixed directions.

Assumptions 5.1. *The following conditions hold:*

(i) The particle's motion is confined in a regular tetrahedron $\mathcal{T}(t)$ characterized by the vertices:

$$\begin{aligned} A_1(t) &= ct(1, 0, 0), & A_2(t) &= ct\left(-\frac{1}{3}, \frac{2\sqrt{2}}{3}, 0\right), \\ A_3(t) &= ct\left(-\frac{1}{3}, -\frac{\sqrt{2}}{3}, \sqrt{\frac{2}{3}}\right), & A_4(t) &= ct\left(-\frac{1}{3}, -\frac{\sqrt{2}}{3}, -\sqrt{\frac{2}{3}}\right). \end{aligned} \quad (46)$$

(ii) The vertices in Eq. (46), identified by the directions \vec{v}_j , for $1 \leq j \leq 4$, satisfy the conditions (i)-(iv) listed in Section 3.2.

(iii) The random times $D_{j,n}$, for $n \in \mathbb{N}$ and $1 \leq j \leq 4$, represent the j -th duration of the motion within the n -th cycle and constitute the intertimes of a GCP with intensity $\lambda_j \in \mathbb{R}^+$.

An example of projections onto the state-space \mathbb{R}^3 of suitable paths of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ under the Assumptions 5.1 is shown in Figure 1. It is easy to see that the particle can eventually reach any position of \mathbb{R}^3 , since $\mathcal{T}(t) \rightarrow \mathbb{R}^3$ as $t \rightarrow \infty$. More-




Figure 1: A sample of the set $\mathcal{T}(t)$, with velocities \vec{v}_j , for $1 \leq j \leq 4$, and fixed vertices given in (46).

over, by taking into account the region defined in (25) and Assumptions 5.1 we have that at every time $t > 0$ the set of possible positions $\mathbf{x} \in \mathbb{R}^3$ of the moving particle is the tetrahedron

$$\mathcal{T}(t) = \left\{ (x_1, x_2, x_3) \in \mathbb{R}^3 : -\frac{ct}{3} < x_1 < ct; -\frac{ct - x_1}{2\sqrt{2}} < x_2 < \frac{ct - x_1}{\sqrt{2}}, \right. \\ \left. |x_3| < \frac{\sqrt{6}}{6}(ct - x_1 - \sqrt{2}x_2) \right\}. \quad (47)$$

Remark 5.1. We observe that the set $\mathcal{T}(t)$ grows as time elapses and, recalling Remark 4.1, the volume is give by $\text{Vol}(\mathcal{T}(t)) = \left(\frac{4}{3}\right)^2 \sqrt{3}(ct)^3$.

Two sample paths of the region $\mathcal{T}(t)$, defined in (47), with directions \vec{v}_j , $1 \leq j \leq 4$, is illustrated in Figure 2.




Figure 2: Two possible paths with the first four segments of the cyclic motion defined in Section 2 under the conditions of Remark 3.1 with $c = t = 1$. The initial velocity is \vec{v}_1 .

Under the hypothesis of Assumptions 5.1, we are now able to determine the explicit probability laws of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ when the random intertimes of the motion along the possible directions \vec{v}_j follow four possible independent GCPs with intensities λ_j , $1 \leq j \leq 4$.

Theorem 5.1. *(Initial components) Let $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ be the stochastic process defined in Section 2 under the initial condition defined in Eq. (6) with velocity \vec{v}_1 . If Assumptions 5.1 hold and the four sequences of intertimes are identically distributed and follow a GCP with intensity λ_j , $1 \leq j \leq 4$, then for all $t \geq 0$, we have*

$$\eta_1 = \mathbb{P}_1 \left\{ (X_1(t), X_2(t), X_3(t)) = (ct, 0, 0), V(t) = \vec{v}_1 \right\} = \frac{1}{1 + \lambda_1 t}, \quad (48)$$

$$\begin{aligned} \eta_2 &= \mathbb{P}_1 \left\{ \mathbf{X}(t) \in E_{12}, V(t) = \vec{v}_2 \right\} \\ &= \frac{\lambda_1}{(\lambda_1 + \lambda_2 + \lambda_1 \lambda_2 t)^2} \left\{ \frac{\lambda_1 t (\lambda_1 + \lambda_2 + \lambda_1 \lambda_2 t)}{1 + \lambda_1 t} + \lambda_2 \log [(1 + \lambda_1 t)(1 + \lambda_2 t)] \right\} \end{aligned} \quad (49)$$

and

$$\eta_3 = \mathbb{P}_1 \left\{ \mathbf{X}(t) \in F_{123}, V(t) = \vec{v}_3 \right\} = R_\lambda(t), \quad (50)$$

where for $\lambda_1 = \lambda_2 = \lambda_3 = \lambda$ one has

$$\begin{aligned}
R_\lambda(t) &= \frac{1}{(2 + \lambda t)^2(3 + \lambda t)^3} \\
&\left\{ 2 \log(1 + \lambda t) \left[(3 + \lambda t)(\lambda t(3 + 2\lambda t)) + 4(2 + \lambda t)^2 \log(2 + \lambda t) \right] \right. \\
&+ (2 + \lambda t) \left[\pi^2(2 + \lambda t) + \lambda t(3 + \lambda t)^2 - 4(2 + \lambda t) \log\left(\frac{1}{2 + \lambda t}\right) \log\left(\frac{1 + \lambda t}{2 + \lambda t}\right) \right] \\
&\left. + 4(2 + \lambda t)^2 \left[Li_2(- (1 + \lambda t)) + 2Li_2\left(\frac{1 + \lambda t}{2 + \lambda t}\right) \right] \right\} \quad (51)
\end{aligned}$$

where $Li_2(\cdot)$ is the dilogarithm function.

Proof. Recalling Theorem 4.1, under Assumption 5.1, Eq. (48) is obtained making use of Eq. (13) in the following expression

$$\mathbb{P}_1\{\mathbf{X}(t) \in A_1(t), V(t) = \vec{v}_1\} = 1 - \int_0^t f_{D_{1,1}}(s) ds. \quad (52)$$

Similarly, Eqs. (49) and (50), when $V(0) = \vec{v}_2$ and $V(0) = \vec{v}_3$, are given by

$$\mathbb{P}_1\{\mathbf{X}(t) \in E_{12}(t), V(t) = \vec{v}_2\} = \int_0^t f_{D_{1,1}}(s) \int_{t-s}^\infty f_{D_{2,1}}(u) du ds \quad (53)$$

and

$$\mathbb{P}_1\{\mathbf{X}(t) \in F_{123}(t), V(t) = \vec{v}_3\} = \int_0^t f_{D_{1,1}+D_{2,1}}(s) f_{D_{3,1}}(t-s) ds, \quad (54)$$

respectively. \square

In Figure 3 are shown some plots of Eq. (48), (49) and (50) for different choices of the intensities λ_j , $1 \leq j \leq 3$, respectively.

Theorem 5.2. (*Absolutely continuous components*) Let $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ be the stochastic process defined in Section 2 under the initial condition defined in Eq. (6) with velocity \vec{v}_1 .




Figure 3: Left to right: plot of Eq. (48) with $\lambda_1 = 1$ (solid), 2 (dotted), 10 (dashed); plot of Eq. (49) with $\lambda_1 = 1$ and $\lambda_2 = 2$ (solid line), $\lambda_1 = 2$ and $\lambda_2 = 4$ (dotted line), $\lambda_1 = 5$ and $\lambda_2 = 10$ (dashed line); plot of Eq. (50) with $\lambda = 1$ (solid), 5 (dotted), 10 (dashed).

For all $\mathbf{x} \in \text{Int}(\mathcal{T}(t))$, with $\mathcal{T}(t)$ given in (47), we have

$$\begin{aligned}
p_{11}(\mathbf{x}, t) &= \lambda_1 \lambda_2 \lambda_3 \lambda_4 \tau_1 \\
&\quad \times \frac{[1 + A(\tau) + B(\tau) + C(\tau)]^2 + 6D(\tau)[1 + A(\tau) + B(\tau) + C(\tau) + D(\tau)]}{\det \mathbf{A} [1 + A(\tau) + B(\tau) + C(\tau)]^4}, \\
p_{12}(\mathbf{x}, t) &= 2\lambda_1^2 \lambda_2 \lambda_3 \lambda_4 \tau_1 \tau_2 (1 + \lambda_2 \tau_2)(1 + \lambda_3 \tau_3)(1 + \lambda_4 \tau_4) \\
&\quad \times \frac{[1 + A(\tau) + B(\tau) + C(\tau) + 3D(\tau)]}{\det \mathbf{A} [1 + A(\tau) + B(\tau) + C(\tau)]^4}, \\
p_{13}(\mathbf{x}, t) &= 2\lambda_1^2 \lambda_2^2 \lambda_3 \lambda_4 \tau_1 \tau_2 \tau_3 (1 + \lambda_3 \tau_3)(1 + \lambda_4 \tau_4) \\
&\quad \times \frac{[1 + A(\tau) + B(\tau) + C(\tau)] + 3D(\tau)}{\det \mathbf{A} [1 + A(\tau) + B(\tau) + C(\tau)]^4}, \\
p_{14}(\mathbf{x}, t) &= \lambda_1 \lambda_2 \lambda_3 (1 + \lambda_4 \tau_4) \\
&\quad \times \frac{[1 + A(\tau) + B(\tau) + C(\tau)]^2 + 6D(\tau)[1 + A(\tau) + B(\tau) + C(\tau) + D(\tau)]}{\det \mathbf{A} [1 + A(\tau) + B(\tau) + C(\tau)]^4},
\end{aligned} \tag{55}$$

with

$$\begin{aligned}
\det \mathbf{A} &= \frac{16\sqrt{3}}{9}c^3, & A(\boldsymbol{\tau}) &= \sum_{i=1}^4 \lambda_i \tau_i, & B(\boldsymbol{\tau}) &= \sum_{\substack{i,j=1 \\ i < j}}^4 \lambda_i \lambda_j \tau_i \tau_j, \\
C(\boldsymbol{\tau}) &= \sum_{\substack{i,j,k=1 \\ i < j < k}}^4 \lambda_i \lambda_j \lambda_k \tau_i \tau_j \tau_k, & D(\boldsymbol{\tau}) &= \prod_{i=1}^4 \lambda_i \tau_i,
\end{aligned} \tag{56}$$

and where the terms $\tau_j = \tau_j(\mathbf{x}, t)$, for $1 \leq j \leq 4$, are given by

$$\begin{aligned}
\tau_1 &= \frac{ct + 3x_1}{4c}, & \tau_2 &= \frac{ct - x_1 + 2\sqrt{2}x_2}{4c}, \\
\tau_3 &= \frac{ct - x_1 - \sqrt{2}x_2 + \sqrt{6}x_3}{4c}, & \tau_4 &= \frac{ct - x_1 - \sqrt{2}x_2 - \sqrt{6}x_3}{4c}.
\end{aligned} \tag{57}$$

Proof. Eqs. (55) are obtained in a closed form as an immediate consequence of Theorem 4.2 after straightforward calculations. \square

We observe that the values of τ_j in (57) are obtained directly from Eq. (36) in Proposition 4.1. Moreover, due to Eq. (9), under the assumptions of Theorem 5.2 the p.d.f. $p_1(\mathbf{x}, t)$ can be immediately obtained from Eqs. (55).




Figure 4: Plot of $p_1(\mathbf{x}, t)$ for $c = t = 1$ and $x_1 = 1/2$ on the left-side and $c = 1$, $t = 2$ and $x_1 = 1$ on the right-side.

As example, some plots of $p_1(\mathbf{x}, t)$ are illustrated in Figure 4 for fixed $c > 0$ and $t > 0$ and $x_1 \in (-ct/3, ct)$. Note that the domain for (x_2, x_3) has a triangular form. Moreover, from Eq. (47) and Theorem 5.2 it is clear that the dependence on c and t is expressed through the product ct .

Hereafter, we study the asymptotic behaviour of the p.d.f. of the particle location $\mathbf{X}(t)$ defined in (9) when the intensities λ_j tend to infinity, for $1 \leq j \leq 4$. In particular, substituting Eqs. (55) in Eq. (9) we obtain the following corollary.

Corollary 5.1. *Under the assumptions of Theorem 5.2, for $t > 0$ and $\mathbf{x} \in \text{Int}(\mathcal{T}(t))$ one has*

$$\lim_{\substack{\forall j, \lambda_j \rightarrow +\infty \\ \forall i, \lambda_1/\lambda_i \rightarrow 1}} p_1(\mathbf{x}, t) = \xi(\mathbf{x}, t), \quad (58)$$

where $\xi(\mathbf{x}, t)$ is the following p.d.f.

$$\xi(\mathbf{x}, t) = \frac{6t(\tau_1\tau_2\tau_3\tau_4)^2}{\det \mathbf{A}[(\tau_1\tau_2\tau_3)^4 + (\tau_1\tau_2\tau_4)^4 + (\tau_1\tau_3\tau_4)^4 + (\tau_2\tau_3\tau_4)^4]}, \quad (59)$$

with τ_j expressed in (36).

Some instances of $\xi(\mathbf{x}, t)$ are plotted in Figure 5 for fixed $c > 0$ and $t > 0$, with $x_1 \in (-ct/3, ct)$.




Figure 5: Plot of $\xi(\mathbf{x}, t)$ for $c = t = 1$ and $x_1 = 1/2$ on the left-side and $c = 1, t = 2$ and $x_1 = 1$ on the right-side.

Remark 5.2. *Under the assumptions of Theorem 5.2, we note that, for $t > 0$ and $1 \leq j \leq 4$, the limit of $p_1(\mathbf{x}, t)$ for $(x_1, x_2, x_3) \rightarrow (ct \cos \theta_j \sin \varphi_j, ct \sin \theta_j \sin \varphi_j, ct \cos \varphi_j)$ can be computed in a closed form that we omit due to its complexity.*

The results expressed in this section for $V(0) = \vec{v}_1$ can be extended to the cases $V(0) = \vec{v}_j$, for $2 \leq j \leq 4$, by using a similar strategy.

6 Concluding remarks

In this paper we analyzed a finite random motion in \mathbb{R}^3 where the sojourn times along each direction form four independent GCPs, and where the possible directions alternate cyclically. This work has been inspired by a theory formulated in [6] with the aim to define similar processes in higher dimensions with possibly variable velocities. Potential applications in biomathematics, engineering, financial and actuarial sciences allow to investigate possible future developments also oriented to the study of the first-passage-time problem.

Here, in order to illustrate the basic issues of this problem, we introduce the (upward) first-passage time for the first component of the process $\{(\mathbf{X}(t), V(t)), t \geq 0\}$ through a constant barrier, say $\beta > 0$, conditional on C_1 (cf. Eq. (6)) given by

$$\tau_\beta = \inf\{t \geq 0 : X_1(t) \geq \beta\}, \quad \mathbf{X}(0) = \mathbf{0}, \quad V(0) = \vec{v}_1, \quad |\vec{v}_1| = c, \quad (60)$$

for $c > 0$. The probability distribution of (60) can be expressed in terms of the sub-density functions:

$$g_\beta(t, k) := \frac{\mathbb{P}_1(\tau_\beta \in dt, N(t) = k)}{dt}, \quad k \in \mathbb{N}. \quad (61)$$

In particular, by the law of total probability, we express the conditional distribution of τ_β in the form

$$\mathbb{P}_1(\tau_\beta \in dt) = \mathbb{P}(D_{1,1} > t)\delta_{\frac{\beta}{c}}(dt) + \sum_{k=1}^{+\infty} \mathbb{P}_1(\tau_\beta \in dt, N(t) = k), \quad (62)$$

where $\delta_{\frac{\beta}{c}}$ is the Dirac delta measure at $\frac{\beta}{c}$, and $N(t)$ is the alternating counting process introduced in Eq. (2). The first term on the right-hand-side of (62) corresponds to the motion without any direction switching up to time t . The series on the right-hand side of (62) represents the absolutely continuous component of the first-passage-time distribution, which arises when at least one direction reversal occurs. We also recall that $D_{1,1}$, defined in Eq. (1), is the random duration of the first time interval during which the particle proceeds with velocity \vec{v}_1 . In order to describe the first-passage-time problem, we consider as threshold the plane $x_1 = \beta$ and project the vectors \vec{v}_j , for $1 \leq j \leq 4$, onto the x_1 -axes according to the following relation

$$\vec{v}_{j_{x_1}} = \text{proj}_{\vec{w}_1}(\vec{v}_j) = \frac{\vec{w}_1 \cdot \vec{v}_j}{\|\vec{w}_1\|^2} \cdot \vec{w}_1, \quad 1 \leq j \leq 4.$$

The latter is the projection of the vector \vec{v}_j on \vec{w}_1 , which is the versor along the x_1 -axes. An illustration of the problem is shown in Figure 6. Clearly, under Assumption 5.1 of Section




Figure 6: Illustration the constant barrier $\beta > 0$ for the first-passage time problem introduced in (60).

5, we have:

$$\begin{aligned}\vec{v}_{1_{x_1}} &= c(1, 0, 0), & \vec{v}_{2_{x_1}} &= c\left(-\frac{1}{3}, 0, 0\right), \\ \vec{v}_{3_{x_1}} &= c\left(-\frac{1}{3}, 0, 0\right), & \vec{v}_{4_{x_1}} &= c\left(-\frac{1}{3}, 0, 0\right).\end{aligned}\tag{63}$$

Due to the complexity of the problem, we discuss only the first cycle of particle motion concerning $D_{j,1}$ ($1 \leq j \leq 4$). We point out that the particle may reach the threshold $x_1 = \beta$ only when it runs with velocity $\vec{v}_{1_{x_1}}$, since the other directions $\vec{v}_{2_{x_1}}$, $\vec{v}_{3_{x_1}}$ and $\vec{v}_{4_{x_1}}$ refer to motion in the opposite direction. Moreover, with reference to the first term in the right-hand-side of (62), one has

$$\begin{aligned}\mathbb{P}(D_{1,1} > t) &= \mathbb{P}\left\{X_1(\tau_\beta) = \beta, V(\tau_\beta) = \vec{v}_1\right\} \\ &= \mathbb{P}\left\{\tau_\beta = \frac{\beta}{c}, N(\tau_\beta) = 0\right\} \\ &= \bar{F}_{D_{1,1}}\left(\frac{\beta}{c}\right) = \frac{1}{1 + \lambda_1\beta/c}.\end{aligned}\tag{64}$$

Clearly, during the random periods $D_{2,1}, D_{3,1}, D_{4,1}$, when the particle moves with velocity $\vec{v}_{2_{x_1}} = \vec{v}_{3_{x_1}} = \vec{v}_{4_{x_1}} = -\frac{c}{3}$, it cannot reach the threshold β since it moves in the opposite direction. Therefore, the position occupied by the particle at the end of the first period of motion $D_1^{(1)}$ is given by

$$X_1(t) = cD_{1,1} - \frac{c}{3}(D_{2,1} + D_{3,1} + D_{4,1}).\tag{65}$$

It is worth mentioning that the determination of an explicit form for the terms in the series of (62) is in general very difficult even when k is small and when the intensities λ_i are equal. Hence, in view of possible future developments, in a forthcoming investigation we aim to apply computational methods to determine the related probabilities.

In conclusion, we stress that the analysis of finite-velocity random motions in multidimensional domains deserves interest in various applied fields. In particular, the motion of a particle in a three-dimensional space as studied in this paper provides possible applications also in chemistry, since the tetrahedral geometry characterizes the shape of many molecules. Indeed, the motion of a particle is isotropic, i.e., the direction of its movement is uniformly distributed on the unit sphere in \mathbb{R}^3 . Thus, according to this interpretation, the Eq. (8) can be employed to represent the wave function of the electron.

Further applications of finite-velocity random motions are related, for instance, to biology (for the random motions of microorganisms), to geology (for alternating trends in volcanic areas), and to physics (for the vorticity motion in two or more dimensions).

Acknowledgments

The authors are members of the group GNCS of INdAM (Istituto Nazionale di Alta Matematica). This work is partially supported by INdAM-GNCS (project ‘‘Modelli di shock basati sul processo di conteggio geometrico e applicazioni alla sopravvivenza’’, CUP-E55F22000270001).

Data Availability Statement

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

All authors contributed equally to this paper. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Beghin, L., Nietdu, L., Orsingher, E.: Probabilistic analysis of the telegrapher's process with drift by means of relativistic transformations. *J. appl. math. stoch. anal.* 14(1) 11-25 (2001)
- [2] Cha, J.H., Finkelstein, M.: A note on the class of geometric counting processes. *Probab. Eng. Inf. Sci.* 27(2), 177-185 (2013)
- [3] Cinque, F., Orsingher, E.: Random motions in \mathbb{R}^3 with orthogonal directions. arXiv preprint arXiv:2205.08829 (2022)
- [4] Cinque, F., Orsingher, E.: Stochastic dynamics of generalized planar random motions with orthogonal directions, *J. Theor. Probab.*1-33 (2023)
- [5] Di Crescenzo, A.: Exact transient analysis of a planar random motion with three directions. *Stochastics Stochastics Rep.* 72(3-4), 175-189 (2002)
- [6] Di Crescenzo, A., Iuliano, A., Mustaro, V.: On some finite-felocity random motions driven by the geometric counting process. *J. Stat. Phys.* 190(3), 44 (2023)
- [7] Di Crescenzo, A., Pellerey, F.: Some results and applications of geometric counting processes. *Methodol. Comput. Appl. Probab.* 21(1), 203-233 (2019)
- [8] Hartmann, A.K., Majumdar, S.N., Schawe, H., Schehr G.: The convex hull of the run-and-tumble particle in a plane. *J. Stat. Mech. Theory Exp.* 2020(5), 053401 (2020)
- [9] Kolesnik, A.D.: The equations of Markovian random evolution on the line. *J. Appl. Probab.* 35(1), 27-35 (1998)
- [10] Kolesnik, A.D.: Discontinuous term of the distribution for Markovian random evolution in \mathbb{R}^3 . *Bul. Acad. Ştiinţe Repub. Mold. Mat.* , 51(2), 62-68 (2006)
- [11] Kolesnik, A.D., Turbin, A.F.: The equation of symmetric Markovian random evolution in a plane, *Stoch. Process. Their Appl.* 75(1), 67-87 (1998)
- [12] Kolesnik, A.D., Orsingher, E.: Analysis of a finite-velocity planar random motion with reflection. *Theory Probab. Appl.* 46(1), 132-140 (2002)

- [13] Kolesnik, A.D., Ratanov, N.: Telegraph Processes and Option Pricing. Second Edition. Springer, Berlin (2023)
- [14] Lachal, A.: Cyclic random motions in-space with n directions. ESAIM Probab Stat . 10, 277-316 (2006)
- [15] Lachal, A., Leorato, S., Orsingher, E.: Minimal cyclic random motion in \mathbb{R}^n and hyper-Bessel functions. Ann. inst. Henri Poincare (B) Probab. Vol. 42, No. 6, pp. 753-772 (2006)
- [16] Leorato, S., Orsingher, E.: Bose-Einstein-type statistics, order statistics and planar random motions with three directions. Adv Appl Probab. 36(3), 937-970 (2004)
- [17] Martens, K., Angelani, L., Di Leonardo, R., Bocquet, L.: Probability distributions for the run-and-tumble bacterial dynamics: An analogy to the Lorentz model. Eur. Phys. J. E. 35, 1-6 (2012)
- [18] Masoliver, J., Porra, J.M., Weiss, G.H.: Some two and three-dimensional persistent random walks. Phys. A: Stat. Mech. Appl. 193(3-4), 469-482 (1993)
- [19] Orsingher, E.: A planar random motion governed by the two-dimensional telegraph equation. J. Appl. Probab. 23(2), 385-397 (1986)
- [20] Orsingher, E.: Probability law, flow function, maximum distribution of wave-governed random motions and their connections with Kirchoff's laws. Stoch. Process. Their Appl. 34(1), 49-66 (1990)
- [21] Orsingher, E.: Bessel functions of third order and the distribution of cyclic planar motions with three directions. Stochastics Stochastics Rep. 74, 617-631 (2002)
- [22] Orsingher, E., De Gregorio, A.: Random flights in higher spaces. J. Theor. Probab. 20, 769-806 (2007)
- [23] Orsingher, E.; Garra, R.; Zeifman, A. I.: Cyclic random motions with orthogonal directions. Markov Process. Related Fields 26 no. 3, 381-402 (2020)
- [24] Orsingher, E., Sommella, A. M.: A cyclic random motion in \mathbb{R}^3 with four directions and finite velocity. Stoch. stoch. rep. 76(2) 113-133 (2004)
- [25] Pogorui, A.A.: Evolution in multidimensional spaces. Random Oper. Stoch. 20(2), 135-141 (2012)
- [26] Pogorui, A.A., Rodríguez-Dagnino, R.M.: Isotropic random motion at finite speed with K-Erlang distributed direction alternations. J. Stat. Phys. 145, 102-112 (2011)
- [27] Pogorui, A.A., Rodríguez-Dagnino, R.M.: Distribution of random motion at renewal instants in three-dimensional space. Journal of Mathematical Sciences, 254, 416-424 (2021)
- [28] Santra, I., Basu, U., Sabhapandit, S.: Run-and-tumble particles in two dimensions: Marginal position distributions. Phys. Rev. E. 101(6), 062120 (2020)