

# Minimum intrinsic dimension scaling for entropic optimal transport

Austin J. Stromme\*

August 25, 2023

## Abstract

Motivated by the manifold hypothesis, which states that data with a high extrinsic dimension may yet have a low intrinsic dimension, we develop refined statistical bounds for entropic optimal transport that are sensitive to the intrinsic dimension of the data. Our bounds involve a robust notion of intrinsic dimension, measured at only a single distance scale depending on the regularization parameter, and show that it is only the minimum of these single-scale intrinsic dimensions which governs the rate of convergence. We call this the Minimum Intrinsic Dimension scaling (MID scaling) phenomenon, and establish MID scaling with no assumptions on the data distributions so long as the cost is bounded and Lipschitz, and for various entropic optimal transport quantities beyond just values, with stronger analogs when one distribution is supported on a manifold. Our results significantly advance the theoretical state of the art by showing that MID scaling is a generic phenomenon, and provide the first rigorous interpretation of the statistical effect of entropic regularization as a distance scale.

## 1 Introduction

Optimal transport (OT) is a powerful paradigm for comparing probability distributions based on a minimum-energy criterion, and has recently been employed throughout applied science, including statistics, computer science, biology and beyond [Vil08, San15, PC19, PZ19, SST<sup>+</sup>19]. Rather than comparing distributions pointwise, OT searches for the most efficient way to transform one into the other, and thus gives practitioners a geometrically meaningful method of contrasting and interpolating data that can be represented as probability distributions. For probability measures  $\mu, \nu$  on  $\mathbb{R}^d$  the OT problem, with respect to the squared Euclidean cost  $\|\cdot\|^2$ , is defined as

$$W_2^2(\mu, \nu) := \min_{\pi \in \Pi(\mu, \nu)} \int \|x - y\|^2 d\pi(x, y), \quad (1.1)$$

where  $\Pi(\mu, \nu)$  denotes the set of probability distributions  $\pi$  on  $\mathbb{R}^d \times \mathbb{R}^d$  which couple  $\mu$  to  $\nu$ , namely such that their marginal on the first  $d$  coordinates is  $\mu$  and their marginal on the last  $d$  coordinates is  $\nu$ . The fundamental theorem of OT guarantees that, so long as  $\mu, \nu$  have finite second moments, there is a solution to (1.1); moreover if  $\mu$  has a density

---

\*Department of EECS at the Massachusetts Institute of Technology, [austinjstromme.work@gmail.com](mailto:austinjstromme.work@gmail.com).

with respect to the Lebesgue measure, there is a unique minimizer which is supported on the graph of a deterministic map [Vi08]. For many applications, such as domain adaptation [FCTR16] or cellular biology [SST+19], it is the coupling minimizing (1.1), rather than the cost  $W_2^2(\mu, \nu)$  itself, which is of greatest interest.

**Entropic optimal transport.** When discretized, the OT problem (1.1) becomes a linear program (LP) with two equality constraints. Although this problem can be feasibly solved at moderate scale with specialized LP solvers which have computational complexity scaling cubically in the support size [DGP+20], in practice, OT is most often approximated with an entropic regularization term [Cut13]. This entropically regularized problem is then solved with a simple iterative rounding algorithm known as Sinkhorn’s algorithm [Sin64, SK67], and is preferred for its quadratic scaling in the support size, simplicity, and parallelizability. The entropically regularized OT (entropic OT) problem is defined, for a regularization parameter  $\varepsilon > 0$ , as

$$S_\varepsilon(\mu, \nu) := \min_{\pi \in \Pi(\mu, \nu)} \int \|x - y\|^2 d\pi(x, y) + \varepsilon \text{KL}(\pi \| \mu \otimes \nu), \quad (1.2)$$

where  $\mu \otimes \nu$  denotes the joint law of  $(x, y)$  when  $x \sim \mu$  and  $y \sim \nu$  are independent, and the Kullback–Leibler divergence is defined as

$$\text{KL}(P \| Q) := \begin{cases} \int \ln \left( \frac{dP}{dQ}(x) \right) dP(x) & P \ll Q, \\ \infty & P \not\ll Q, \end{cases}$$

and the notation  $P \ll Q$  means  $P$  is absolutely continuous with respect to  $Q$ .

## 1.1 Statistical aspects of (entropic) OT

Unfortunately, the un-regularized OT problem (1.1) is known to suffer from a statistical curse of dimensionality. To describe this barrier, consider the practical situation in which one does not have access to the entire distributions  $\mu, \nu$ , and instead only has access to iid samples of size  $n$  from each distribution, which we write as  $\mathcal{X} := (x_1, \dots, x_n) \sim \mu^{\otimes n}$  and  $\mathcal{Y} := (y_1, \dots, y_n) \sim \nu^{\otimes n}$ . Let  $\hat{\mu}$  and  $\hat{\nu}$  denote the empirical measures supported on  $\mathcal{X}$  and  $\mathcal{Y}$ , respectively, namely

$$\hat{\mu} := \frac{1}{n} \sum_{i=1}^n \delta_{x_i} \quad \text{and} \quad \hat{\nu} := \frac{1}{n} \sum_{j=1}^n \delta_{y_j}.$$

Given the empirical measures  $\hat{\mu}, \hat{\nu}$ , the most natural way to estimate the population quantity  $W_2^2(\mu, \nu)$  is with the plug-in estimator  $W_2^2(\hat{\mu}, \hat{\nu})$ . The discrepancy between the empirical OT value and its population counterpart is an old and well-studied area, and it has long been understood that rates like  $n^{-2/d}$  are typical – for instance, if  $d \geq 5$ , and  $\mu, \nu$  are absolutely continuous with respect to the Lebesgue measure on  $[0, 1]^d$  [Dud69, MNW21]. It is natural to wonder whether these rates can be improved with other estimators, but it was recently shown they are essentially un-improvable, both for OT value estimation [NWR22, MBNWW21] and OT map estimation [HR21].

**Statistical entropic OT.** Motivated by its computational benefits and corresponding prevalence in practice, as well as the curse of dimensionality for its un-regularized counterpart, a line of recent work has endeavored to understand the statistical consequences of entropic regularization [LSPC19, GCB<sup>+</sup>19, MNW19, RS22, dBSLNW23]. This work has shown that the entropic OT problem (1.2) offers significantly improved statistical performance, essentially transferring the curse of dimensionality from the sample size  $n$  to the regularization parameter  $\varepsilon$ ; the following result describes the current state of the art when the measures are bounded and  $\varepsilon$  is small.

**Theorem 1** ([GCB<sup>+</sup>19, MNW19]). *Suppose  $\mu, \nu$  are probability measures on  $\mathbb{R}^d$  with support contained in the unit ball  $B(0, 1)$ . Then for a constant  $C_d$  depending only on the dimension  $d$ ,*

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \leq C_d \cdot \frac{1 + \varepsilon^{-\lfloor d/2 \rfloor}}{\sqrt{n}},$$

where the expectation is taken over the iid samples  $\mathcal{X} \sim \mu^{\otimes n}, \mathcal{Y} \sim \nu^{\otimes n}$ .

By giving an explicit trade-off between regularization and statistical error, this result offers a strong and flexible description of the performance of entropic OT in practice. In this work, we build on Theorem 1 and develop a refined theory of the statistical behavior of entropic OT.

## 1.2 Effective statistical dimension of entropic OT

To describe our refinements, observe that the estimate in Theorem 1 is *extrinsic* in the sense that the dimension  $d$  is appearing because the ambient space is  $\mathbb{R}^d$ , rather than anything to do with the *intrinsic* dimensions of  $\mu$  and  $\nu$ . This extrinsic dimension dependence is fundamental to the proof, and can be very pessimistic. For a toy example, suppose  $\mu$  and  $\nu$  are supported on the same  $k$ -dimensional hyperplane in  $\mathbb{R}^d$ : then Theorem 1 itself implies that the statistical rate has the potentially much milder dependence  $\varepsilon^{-\lfloor k/2 \rfloor}$  on the regularization parameter  $\varepsilon$ . In such a case, we can say that  $\mu, \nu$  have extrinsic dimension  $d$  yet have intrinsic dimension (at most)  $k$ . Well beyond toy settings, the widely believed manifold hypothesis states that natural data with a high extrinsic dimension is typically near to or on a low-dimensional manifold embedded in Euclidean space, and so has low intrinsic dimension [FMN16, BGC17]. Given the ubiquity of entropic OT in practice, it is therefore of major interest to determine the effective statistical dimension of entropic OT, and identify how it relates to the extrinsic dimension  $d$  as well as the intrinsic dimensions of  $\mu$  and  $\nu$ . In this work, we thus study the following question:

*Suppose  $\mu$  and  $\nu$  are supported on  $\mathbb{R}^d$ , yet have (informally speaking) intrinsic dimensions  $d_\mu$  and  $d_\nu$ . How does the statistical rate of convergence of entropic OT depend on  $d, d_\mu$ , and  $d_\nu$ ?*

Our results will measure intrinsic dimension through the covering numbers of the supports of  $\mu$  and  $\nu$ . To introduce covering numbers, write the closed ball centered at  $z_0 \in \mathbb{R}^d$  with radius  $r > 0$  as

$$B^{\text{cl}}(z_0, r) := \{z \in \mathbb{R}^d: \|z - z_0\| \leq r\}.$$

Then the covering number of a set  $A \subset \mathbb{R}^d$  at scale  $\delta > 0$  is defined as

$$\mathcal{N}(A, \|\cdot\|, \delta) := \min \left\{ K \in \mathbb{N}_{>0} \mid \exists z_1, \dots, z_K \in \mathbb{R}^d : A \subset \bigcup_{k=1}^K B^{\text{cl}}(z_k, \delta) \right\}. \quad (1.3)$$

It will be convenient to define

$$\mathcal{N}(\mu, \delta) := \mathcal{N}(\text{supp}(\mu), \|\cdot\|, \delta), \quad \mathcal{N}(\nu, \delta) := \mathcal{N}(\text{supp}(\nu), \|\cdot\|, \delta), \quad \delta > 0.$$

**Contributions.** Using this notation, we can give our answer to the above question:

**MID scaling:** *the statistical rate of convergence of entropic OT can be upper bounded by quantities whose dimension-dependence is purely contained within the minimum covering number at scale  $\varepsilon$ , namely  $\mathcal{N}(\mu, \varepsilon) \wedge \mathcal{N}(\nu, \varepsilon)$ .*

We call this the minimum intrinsic dimension scaling (MID scaling) phenomenon. We emphasize that the MID scaling phenomenon encapsulates two related yet distinct phenomena:

1. **Minimum:** only the *minimum* of the intrinsic dimensions governs the convergence
2. **Intrinsic Dimension:** the dimension-dependence is intrinsic to  $\mu$  and  $\nu$ , and in fact is *intrinsic at a single distance scale depending on the regularization parameter*.

As we elaborate on in section 3, the MID scaling phenomenon provides a strong statistical justification for entropic OT in the context of the manifold hypothesis. It shows that distributions with support which may not be smooth or even locally low-dimensional can still have significantly improved dimension-dependence in their statistical rates, and clearly describes the statistical role of the regularization parameter  $\varepsilon$  as a distance scale. Moreover, it identifies the surprising fact that the dimension-dependence of entropic OT is driven by the lower-dimensional (in this single-scale sense) distribution.

The above statement of MID scaling is in the setting of Theorem 1, but our main results establish MID scaling more generally for bounded, Lipschitz costs, and for entropic OT quantities beyond values, including maps and densities. As our main results in this setting, we prove MID scaling for entropic OT values in Theorem 2, for entropic OT maps in empirical norms in Theorem 7, and for entropic OT densities in empirical norms in Theorem 8. Under the additional assumption that one distribution is supported on an embedded manifold, we show that MID scaling also holds for entropic dual potentials with fast, population norm convergence in Theorem 9, and apply this result to prove stronger analogs of the previous results on value, map, and density estimation.

**Organization of the paper.** We conclude this section with a discussion of related work and a summary of our notation. In section 2, we state our assumption on the cost function and give background and more notation for the entropic OT problem. In section 3, we state our main results and give examples and discussion. In section 4, we describe some preliminary observations that form the foundation of our approach. In section 5, we prove our main results on MID scaling with a thorough exposition of our proof strategy in the

case of value estimation. In section 6, we prove our main results in the case where one distribution is supported on an embedded manifold. In Appendix A, we collect some background on embedded manifolds that we use in section 6, and in Appendix B we include some deferred proofs.

**Related work.** Taken as a whole, our results provide strong evidence of the MID scaling phenomenon for entropic OT. In fact, there is additional evidence for MID scaling and related phenomena in the literature. Recent work studied the continuous to discrete case, and established that entropic OT maps achieve dimension-free rates of convergence, consistent with MID scaling [PDNW23]. And in the un-regularized setting, a similar phenomenon was recently established for value estimation, where it was dubbed “lower complexity adaptation” [HSM22]. The lower complexity adaptation phenomenon also states that the minimum of the intrinsic dimensions of  $\mu$  and  $\nu$  dictates the rate of convergence of un-regularized OT, but their notion of intrinsic dimension is different from ours. Indeed, as is common in the un-regularized literature, the intrinsic dimensions  $d_\mu$  and  $d_\nu$  in lower complexity adaptation are defined using the covering numbers at many distance scales, and so are distinct from the single-scale intrinsic dimensions in MID scaling. Also, their results require some structural assumptions, whereas our results show that MID scaling in the entropic setting is quite generic. In light of these results, it is natural to expect that the minimum part of MID scaling holds in greater generality, potentially including un-regularized maps and plans, and alternate forms of regularization.

To the best of our knowledge, there is only one prior work which considers the sample complexity of regularized OT and intrinsic dimension [BEZ22]. Those results apply to more general forms of regularization, but are only for value estimation and incur worse  $n$  dependence without identifying MID scaling. Concurrently with this work, [GH23] extended the lower complexity adaptation phenomenon to entropic OT values and Gromov-Wasserstein distances, yielding bounds which are similar to those of the un-regularized problem, with the curse of dimensionality residing primarily in the sample size, and which are mostly incomparable to our own.

As we discuss in detail in section 5, our technical approach is an intrinsic dimension-sensitive refinement of [RS22] where fully dimension-free bounds were established for entropic OT values, maps, and densities, but with exponential factors in  $1/\varepsilon$ . An alternative estimator of the entropic OT map which achieves sub-exponential dependence on  $1/\varepsilon$  was proposed in [Str23]. Other works have proven convergence of the dual potentials [LSPC19], used entropic OT quantities as computationally efficient estimators for their un-regularized counterparts [CRL<sup>+</sup>20, PNW21], considered the convergence of entropic OT maps [MWMA21, WMMA23], and studied the sample complexity of entropically regularized Gromov-Wasserstein distances [ZGMS22]. Significant recent effort has been devoted to developing central limit theorems for entropic OT [BCP19, MNW19, KTM20, dBSL<sup>+</sup>23, GKRS22, GSLNW22, GSH23]. Finally, a related form of minimum dimension-dependence appears in the study of asymptotics for entropic OT as  $\varepsilon \rightarrow 0^+$  [EN23].

Regarding the provenance of Theorem 1, we remark that it was originally proven for smooth, Lipschitz costs beyond  $\ell_2^2$ , but with an exponential dependence on  $1/\varepsilon$  [GCB<sup>+</sup>19]. The follow-up work [MNW19] showed how to remove this exponential factor in the com-

actly supported case that we study here, but primarily concentrated on extending Theorem 1 to un-bounded distributions.

Finally, our work can be seen as an entropic analog to the long line of work on the convergence of the un-regularized OT problem, which dates back to Dudley [Dud69] and encompasses precise asymptotics [DY95] as well as finer behavior in lower dimensions [AKT84, LS86]. In fact, the analysis of un-regularized OT is naturally sensitive to intrinsic dimension, at least when intrinsic dimension is measured through covering numbers at many scales [Dud69, BLG14, WB19]. The discrepancy between Theorem 1 and the natural appearance of intrinsic dimension in the un-regularized OT problem provides part of the motivation for this work. More discussion of the similarities and differences between this line of work and our results is provided in section 3.

**Notation.** Given  $a, b \in \mathbb{R}$ , we write the minimum  $a \wedge b := \min(a, b)$ , and the maximum  $a \vee b := \max(a, b)$ . For a non-negative integer  $K \in \mathbb{N}_{>0}$ , we write  $[K] = \{1 \dots, K\}$ . We always work with Borel probability distributions. Given probability distributions  $P, Q$  on  $\mathbb{R}^d$ , we denote their trivial coupling  $P \otimes Q$ , which is uniquely defined by taking Borel sets  $A, B \subset \mathbb{R}^d$  to  $(P \otimes Q)(A \times B) := P(A)Q(B)$ . The support of  $P$  is denoted  $\text{supp}(P)$ , and is defined to be the set of all points  $x \in \mathbb{R}^d$  such that  $P$  assigns positive value to all open sets containing  $x$ . The  $\ell_2$  Euclidean norm is always written  $\|\cdot\|$ , without a subscript. Given a Borel measurable  $\alpha: \mathbb{R}^d \rightarrow \mathbb{R}^k$  and  $p \in [1, \infty)$ , we define the  $L^p(P)$  norm as

$$\|\alpha\|_{L^p(P)} := \left( \int \|\alpha(x)\|^p dP(x) \right)^{1/p}.$$

The sup norm  $\|\alpha\|_{L^\infty(P)}$  is defined to be the essential supremum of  $\|\alpha\|$  with respect to  $P$ . For a Borel measurable  $\beta: \mathbb{R}^d \rightarrow \mathbb{R}$ , we will variously write

$$\int \beta(x) dP(x) = \mathbb{E}_P[\beta] = P(\beta).$$

We will also write

$$\text{Var}_P(\beta) := \mathbb{E}_P[(\beta - \mathbb{E}_P[\beta])^2].$$

The notation  $u \lesssim v$  indicates  $u \leq Cv$  for constant  $C$ ; whether the constant  $C$  is numerical or problem-dependent is a matter of context. The suppressed constants in our main results on MID scaling, described in sections 3.1 and 3.2 are numerical, while the suppressed constants for our results on embedded manifolds, described in section 3.3, depend on the low-dimensional distribution.

Given a metric  $\text{dist}_N$  on a set  $N \subset \mathbb{R}^d$ , we will write the  $\text{dist}_N$ -ball of radius  $r$  around  $z_0 \in N$  as

$$B_{\text{dist}_N}(z_0, r) := \{z \in N : \text{dist}_N(z, z_0) < r\}.$$

If  $\text{dist}_N = \|\cdot\|$  and  $N = \mathbb{R}^d$ , we simply write  $B(z_0, r)$ . The closed ball is written

$$B_{\text{dist}_N}^{\text{cl}}(z_0, r) := \{z \in N : \text{dist}_N(z, z_0) \leq r\},$$

and again, if  $\text{dist}_N = \|\cdot\|$  and  $N = \mathbb{R}^d$ , we simply write  $B^{\text{cl}}(z_0, r)$ .

Throughout, we will assume that  $\mu, \nu$  are probability measures on  $\mathbb{R}^d$ , and we have iid samples  $\mathcal{X} := (x_1, \dots, x_n) \sim \mu^{\otimes n}$  and  $\mathcal{Y} := (y_1, \dots, y_n) \sim \nu^{\otimes n}$ . Expectations  $\mathbb{E}$  without a subscript will always refer to integration over both of these samples.

## 2 Assumptions and background on entropic OT

In this section, we state our assumption on the cost function, and then establish important notation while reviewing duality for the entropic OT problem.

### 2.1 Assumptions on cost

Our results are most naturally stated for bounded, Lipschitz costs rather than the squared Euclidean cost  $\|\cdot\|^2$  and so we will work at this level of generality throughout the remainder of the paper. We make the following formal assumption on our cost function  $c$ .

**Assumption 1** (Bounded and uniformly Lipschitz cost). *We assume the cost  $c$  is Borel measurable and  $c \in L^\infty(\mu \otimes \nu)$ . By re-scaling the problem we assume without loss of generality that*

$$\|c\|_{L^\infty(\mu \otimes \nu)} \leq 1.$$

Also, we assume there exists  $L > 0$  such that for all  $y \in \text{supp}(\nu)$ ,

$$|c(x, y) - c(x', y)| \leq L\|x - x'\| \quad \forall x, x' \in \text{supp}(\mu),$$

and, similarly, for all  $x \in \text{supp}(\mu)$ ,

$$|c(x, y) - c(x, y')| \leq L\|y - y'\| \quad \forall y, y' \in \text{supp}(\nu).$$

We remark that in the case where  $c$  is the squared Euclidean cost  $\|\cdot\|^2$ , Assumption 1 just means the supports of  $\mu$  and  $\nu$  are compact.

### 2.2 Strong duality for entropic OT

Given the cost function  $c$ , we define the population entropic OT problem as

$$S_\varepsilon(\mu, \nu) := \inf_{\pi \in \Pi(\mu, \nu)} \{ \pi(c) + \varepsilon \text{KL}(\pi \| \mu \otimes \nu) \}. \quad (2.1)$$

The empirical entropic OT problem is then defined

$$S_\varepsilon(\hat{\mu}, \hat{\nu}) := \inf_{\pi \in \Pi(\hat{\mu}, \hat{\nu})} \{ \pi(c) + \varepsilon \text{KL}(\pi \| \hat{\mu} \otimes \hat{\nu}) \}. \quad (2.2)$$

There are unique optimal solutions to the primal problems (2.1) and (2.2), which we write as  $\pi_\varepsilon$  and  $\hat{\pi}_\varepsilon$ , respectively. We distinguish between  $c$  and  $S_\varepsilon(\mu, \nu)$  by referring to the former as the *cost function*, and the latter as the *entropic OT value*.

**Strong duality.** Under Assumption 1 on the cost  $c$ , a form of strong duality holds for both the population and empirical entropic OT problems. We refer to [DMG20] for these results as well as a thorough review of the literature on this topic. For the population entropic OT problem, strong duality is

$$S_\varepsilon(\mu, \nu) = \sup_{(f, g) \in L^\infty(\mu) \times L^\infty(\nu)} \Phi_\varepsilon(f, g) := \mu(f) + \nu(g) - \varepsilon(\mu \otimes \nu)(e^{-\varepsilon^{-1}(c-f-g)} - 1).$$



The function  $\Phi_\varepsilon: L^\infty(\mu) \times L^\infty(\nu) \rightarrow \mathbb{R}$  is the population entropic OT dual function. Similarly for the empirical entropic OT problem, strong duality is stated

$$S_\varepsilon(\hat{\mu}, \hat{\nu}) = \sup_{(f, g) \in L^\infty(\hat{\mu}) \times L^\infty(\hat{\nu})} \hat{\Phi}_\varepsilon(f, g) := \hat{\mu}(f) + \hat{\nu}(g) - \varepsilon(\hat{\mu} \otimes \hat{\nu})(e^{-\varepsilon^{-1}(c-f-g)} - 1).$$

And this  $\hat{\Phi}_\varepsilon: L^\infty(\hat{\mu}) \times L^\infty(\hat{\nu}) \rightarrow \mathbb{R}$  is the empirical entropic OT dual function.

**Population and empirical dual potentials.** The population and empirical dual problems have solutions  $(f_\varepsilon, g_\varepsilon) \in L^\infty(\mu) \times L^\infty(\nu)$  and  $(\hat{f}_\varepsilon, \hat{g}_\varepsilon) \in L^\infty(\hat{\mu}) \times L^\infty(\hat{\nu})$ , respectively. These solutions are unique up to the translation  $(f, g) \mapsto (f - a, g + a)$  for  $a \in \mathbb{R}$ , and we thus specify the solutions we consider by assuming  $\nu(g_\varepsilon) = 0$  and  $\hat{\nu}(\hat{g}_\varepsilon) = 0$ .

**Population and empirical densities.** For notational convenience, we also define the population density, for  $\mu$ -almost every  $x$  and  $\nu$ -almost every  $y$ , as

$$p_\varepsilon(x, y) := \frac{d\pi_\varepsilon}{d(\mu \otimes \nu)}(x, y) = e^{-\varepsilon^{-1}(c(x, y) - f_\varepsilon(x) - g_\varepsilon(y))}. \quad (2.3)$$

And we similarly define, for all  $x \in \mathcal{X}$  and  $y \in \mathcal{Y}$ , the empirical density

$$\hat{p}_\varepsilon(x, y) := \frac{d\hat{\pi}_\varepsilon}{d(\hat{\mu} \otimes \hat{\nu})}(x, y) = e^{-\varepsilon^{-1}(c(x, y) - \hat{f}_\varepsilon(x) - \hat{g}_\varepsilon(y))} \quad (2.4)$$

**Marginal constraints for the dual potentials.** The marginal constraints  $\pi_\varepsilon \in \Pi(\mu, \nu)$  and  $\hat{\pi}_\varepsilon \in \Pi(\hat{\mu}, \hat{\nu})$  in fact define necessary and sufficient optimality conditions for the corresponding dual potentials through equations (2.3) and (2.4), respectively. The resulting system of equations is sometimes known as the *Schrödinger system*, and is fundamental to entropic optimal transport. For the population potentials, the marginal constraints imply that for  $\mu$ -almost every  $x_0$  and  $\nu$ -almost every  $y_0$ ,

$$1 = \int e^{-\varepsilon^{-1}(c(x, y) - f_\varepsilon(x_0) - g_\varepsilon(y))} d\nu(y), \quad 1 = \int e^{-\varepsilon^{-1}(c(x, y) - f_\varepsilon(x) - g_\varepsilon(y_0))} d\mu(x). \quad (2.5)$$

The empirical potentials  $\hat{f}_\varepsilon, \hat{g}_\varepsilon$  satisfy the analogous marginal equations: for all  $x \in \mathcal{X}$  and  $y \in \mathcal{Y}$ ,

$$1 = \frac{1}{n} \sum_{j=1}^n e^{-\varepsilon^{-1}(c(x, y_j) - \hat{f}_\varepsilon(x) - \hat{g}_\varepsilon(y_j))}, \quad 1 = \frac{1}{n} \sum_{i=1}^n e^{-\varepsilon^{-1}(c(x_i, y) - \hat{f}_\varepsilon(x_i) - \hat{g}_\varepsilon(y))}. \quad (2.6)$$

**Canonical extensions of the dual potentials.** In fact, the marginal equations (2.6) yield a canonical means of extending the empirical entropic dual potentials to functions on all of  $\mathbb{R}^d$  [Ber20, PNW21]. We observe that solving for  $\hat{f}_\varepsilon$  in (2.6) yields, for  $x \in \mathcal{X}$ ,

$$\hat{f}_\varepsilon(x) = -\varepsilon \ln \left( \frac{1}{n} \sum_{j=1}^n e^{-\varepsilon^{-1}(c(x, y_j) - \hat{g}_\varepsilon(y_j))} \right). \quad (2.7)$$



Since  $c$  is defined everywhere, this equation actually makes sense for all  $x \in \text{supp}(\mu)$ , and so we thus *define*  $\hat{f}_\varepsilon$  on all of  $\text{supp} \mu$  with this formula. Similarly, we put,

$$\hat{g}_\varepsilon(y) := -\varepsilon \ln \left( \frac{1}{n} \sum_{i=1}^n e^{-\varepsilon^{-1}(c(x_i, y) - \hat{f}_\varepsilon(x_i))} \right) \quad y \in \text{supp}(\nu). \quad (2.8)$$

Using Equation (2.4), we extend the empirical density  $\hat{p}_\varepsilon$  to all of  $\text{supp}(\mu) \times \text{supp}(\nu)$  as well.

### 3 Main results

In this section we describe our main results on the effective statistical dimension of entropic optimal transport. In section 3.1, we introduce and discuss MID scaling in the context of entropic OT value estimation. In section 3.2, we state our results which establish MID scaling for entropic OT map and density estimation. And in section 3.3, we show how these results can be strengthened in the case where one of the measures is supported on an embedded manifold. Throughout, we work under Assumption 1 on the cost function.

#### 3.1 Introduction to MID scaling with value convergence

In this section, we introduce the reader to MID scaling in the context of the convergence of entropic OT values. We state our main result on MID scaling for values, Theorem 2, give several examples, and then conclude with discussion of its statistical significance and tightness.

**MID scaling for value estimation.** Recall the definition of covering numbers from (1.3), and that we define

$$\mathcal{N}(\mu, \delta) := \mathcal{N}(\text{supp}(\mu), \|\cdot\|, \delta), \quad \mathcal{N}(\nu, \delta) := \mathcal{N}(\text{supp}(\nu), \|\cdot\|, \delta), \quad \delta > 0.$$

Our main result on entropic OT value estimation follows.

**Theorem 2** (MID scaling for values). *For numerical constants independent of all problem parameters,*

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \sqrt{\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}}.$$

The only dimensional quantity in this estimate is contained in the minimum covering numbers at scale  $\varepsilon$ , demonstrating the MID scaling phenomenon. We emphasize that this result only requires that the cost  $c$  is bounded and Lipschitz (Assumption 1), in contrast to the smoothness assumptions in most of the previous literature.

**Examples of MID scaling.** To gain a feeling for Theorem 2, let us consider some examples.

**Example 3** (Generic distributions in  $B(0, 1)$ ). For generic probability distributions, we can apply Theorem 2 with standard upper bounds on covering numbers in Euclidean space [Ver18, Proposition 4.2.12]. We find that if  $\mu, \nu$  are supported in  $B(0, 1) \subset \mathbb{R}^d$ , then for numerical constants independent of all problem parameters, and for all  $\varepsilon > 0$ ,

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \cdot \left(1 + \frac{2L}{\varepsilon}\right)^{d/2} \cdot \frac{1}{\sqrt{n}}.$$

Specializing to the case of the  $\ell_2^2$  cost, this bound nearly recovers Theorem 1, being worse by a factor of  $\varepsilon^{-1/2}$  in the case where  $d$  is odd. In fact, Theorem 1 applies more generally to costs which are both Lipschitz and smooth to degree  $\lceil d/2 \rceil$  [GCB<sup>+</sup>19]. The authors of that work observed empirically that their smoothness assumption seemed unnecessary and left it as an open problem to remove that assumption. Theorem 2 therefore resolves this problem.<sup>1</sup>

In cases where just one of the measures is assumed to have low intrinsic dimension, we can obtain bounds which only depend on that measure.

**Example 4** (Semi-discrete). Suppose  $\nu$  is supported on  $K$  points. Then for numerical constants independent of all problem parameters, and for all  $\varepsilon > 0$ ,

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \cdot \sqrt{\frac{K}{n}}.$$

**Example 5** (Embedded manifold). When  $\nu$  is supported on a  $d_\nu$ -dimensional compact, smooth, embedded Riemannian manifold without boundary, we can apply Theorem 2 with the bound  $\mathcal{N}(\nu, \delta) \leq C_\nu \delta^{-d_\nu}$ , valid for some  $C_\nu > 0$  and  $\delta$  sufficiently small (Proposition 43 in Appendix A formally verifies this fact). In this case, Theorem 2 implies that for  $\varepsilon > 0$  sufficiently small and numerical constants independent of all problem parameters,

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim \sqrt{C_\nu} (1 + \varepsilon) \cdot \left(\frac{L}{\varepsilon}\right)^{d_\nu/2} \cdot \frac{1}{\sqrt{n}}.$$

Because MID scaling involves only a single distance scale, the above examples can be generalized to sets which are only low dimensional at some scales.

**Example 6** ( $\delta$ -fattening of sets [WB19]). For  $\delta \geq 0$  and  $A \subset \mathbb{R}^d$ , the  $\delta$ -fattening of  $A$  is

$$A_\delta := \bigcup_{a \in A} B^{\text{cl}}(a, \delta).$$

Suppose  $\text{supp}(\nu) \subset A_\delta$  for some  $\delta > 0$  and  $A$  such that  $\mathcal{N}(A, \|\cdot\|, \tau) \leq C_A \tau^{-k}$  for all  $\tau$  sufficiently small and some constants  $C_A, k \geq 0$ . Note that covering numbers on  $A_\delta$  can be compared to those on  $A$  itself, since for  $\tau \geq \delta$  we have  $\mathcal{N}(A_\delta, \|\cdot\|, \tau) \leq \mathcal{N}(A, \|\cdot\|, \tau - \delta)$ . Hence for  $\varepsilon > L\delta$  sufficiently small,

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim \sqrt{C_A} (1 + \varepsilon) \cdot \left(\frac{L}{\varepsilon - L\delta}\right)^{k/2} \cdot \frac{1}{\sqrt{n}}.$$

---

<sup>1</sup>For large  $\varepsilon$ , our bound diverges while Theorem 1 becomes  $O(1/\sqrt{n})$ , but it is straightforward to modify our proofs to fully recover Theorem 1 in this case.

The above example illustrates that, so long as the ratio  $\varepsilon/L$  is significantly larger than the fattening scale  $\delta$ , the rates are essentially the same as if  $\text{supp}(\nu)$  were actually contained in  $A$ , despite the fact that  $\text{supp}(\nu)$  may be full-dimensional at scale  $\delta$ . In this way, the dimension-dependence of MID scaling is oblivious to features below scale  $\varepsilon/L$ .

**Discussion of MID scaling.** We emphasize that MID scaling is similar to, but distinct from, the intrinsic dimension-dependence in the un-regularized OT literature [Dud69, BLG14, WB19, HSM22]. While in both settings the minimum of the intrinsic dimensions governs the rate of convergence, the notion of intrinsic dimension is different. In the un-regularized setting, intrinsic dimension is characterized through uniform covering number control at small scales, whereas in MID scaling the covering numbers play a role, but only at a single distance scale. The convergence of un-regularized OT is, in fact, adaptive to multi-scale behavior, but the relevant scales are determined by the sample size  $n$ , meaning that milder covering numbers at some scales only translate to improved rates while  $n$  is not too large; in fact, such rates are known to be tight [WB19]. In contrast, MID scaling shows that entropic OT benefits from better covering number control at some scales *for all sample sizes*  $n$ . Essentially, entropic regularization decouples the sample size and the distance scale of the problem, allowing for a flexible trade-off between the intrinsic dimension of the data distribution and the amount of regularization.<sup>2</sup>

MID scaling helps clarify the statistical role of entropic regularization, showing that beyond its well-known computational virtues, entropy also provides statistical regularization by specifying a distance scale, allowing the user to balance the intrinsic curse of dimensionality of the data with the statistical difficulty of the problem. Because of the manifold hypothesis, which states that natural data is typically supported on or near a low-dimensional embedded manifold, we expect data to have significantly smaller intrinsic dimension than extrinsic dimension. And so a means to flexibly adapt optimal transport to such intrinsic low-dimensional structure, particularly approximate low-dimensional structure, is of major interest. MID scaling demonstrates that entropic regularization provides this benefit for optimal transport.

**Remarks on tightness.** While the upper bounds in this work substantiate MID scaling and the statistical benefits of entropic regularization, a complete statistical account of entropic OT requires lower bounds. We leave a thorough study of this interesting direction to future work, but do give some indications on the tightness of Theorem 2 here. For this reason, we note that the CLT for entropic OT states that the asymptotic variance of  $\sqrt{n}(S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu))$  is  $\text{Var}_\mu(f_\varepsilon) + \text{Var}_\nu(g_\varepsilon)$  [GSH23]. In particular, it follows that the dependence on  $n$  in Theorem 2 is optimal, and moreover that Theorem 2 is tight in the case where  $\nu$  is supported on two points (see Example 4).

However, this example doesn't rule out the possibility that the dependence on the covering numbers of  $\mu$  and  $\nu$  can be improved. To reason about such issues, we can reduce to lower bounds for the un-regularized OT problem. For example, it is known that the approximation error  $|S_\varepsilon - S_0|$  is  $O(\varepsilon)$ , up to logarithmic factors [GCB<sup>+</sup>19]. Combining this approximation error with minimax lower bounds for estimating un-regularized OT

---

<sup>2</sup>Thanks to Yann Ollivier for stimulating comments on this point.

values then implies a firm speed limit on statistical bounds for entropic OT values. Arguing along these lines, we show in Appendix B.1 that Theorem 2 implies that entropic estimators can estimate  $W_1$  distances at the rate  $n^{-1/(d+2)}$ , close to the minimax optimal  $n^{-1/d}$  rate [NWR22], and that moreover the covering number dependence in the bound of Theorem 2 cannot be improved in general.

### 3.2 MID scaling for maps and densities

Theorem 2 gives a strong instance of the MID scaling phenomenon for entropic optimal transport, yet in many applications of entropic OT, such as trajectory reconstruction [SST<sup>+</sup>19] and domain adaptation [FCTR16, SDF<sup>+</sup>18], it is of greater interest to estimate the solution to the entropic OT problem than the value of the entropic OT objective alone. Motivated by this fact, we also study the performance of empirical plug-in estimators for the problems of entropic OT map estimation as well as density estimation, and develop results analogous to Theorem 2, showing that natural plug-in estimators also sport MID scaling.

We define the entropic OT map

$$T_\varepsilon(x) := \mathbb{E}_{\pi_\varepsilon}[y | x].$$

Note that the analogous map defined over  $y$  is symmetric with this one, and so we study only this case without loss of generality. The map  $T_\varepsilon$  is an entropic analog of the OT map, and has been the subject of much recent work as it offers greater computational and statistical efficiency than its un-regularized counterpart [PNW21, MWMA21, RS22, PDNW23]. We study the empirical analog of  $T_\varepsilon(x)$ :

$$\hat{T}_\varepsilon(x) := \mathbb{E}_{\hat{\pi}_\varepsilon}[y | x] = \frac{1}{n} \sum_{j=1}^n y_j \hat{p}_\varepsilon(x, y_j). \quad (3.1)$$

We consider  $\hat{T}_\varepsilon$  as an estimator for its population analog  $T_\varepsilon$  and show it enjoys a  $1/\sqrt{n}$  rate of convergence with MID scaling.

**Theorem 7.** *Suppose the diameter of  $\text{supp}(\nu)$  is at most  $R$ . Then for numerical constants independent of all problem parameters,*

$$\mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\hat{\mu})}^2] \lesssim R^2 \left(1 + \frac{1}{\varepsilon}\right) \sqrt{\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}}.$$

We note that this result is measured with respect to the empirical norm  $\|\cdot\|_{L^2(\hat{\mu})}$ , rather than the population norm  $\|\cdot\|_{L^2(\mu)}$ . In the next section we assume that one distribution is supported on an embedded manifold and show that this result can be strengthened to population norm convergence with fast  $1/n$  rates. As before, we may derive bounds from this result in cases where we have *a priori* control on the relevant covering numbers. Previous work on the convergence of  $\hat{T}_\varepsilon$  to  $T_\varepsilon$  for generic distributions incurred an exponential dependence on  $1/\varepsilon$  [MWMA21, RS22]. And an alternative estimator which achieves sub-exponential dependence on  $1/\varepsilon$  was considered in [Str23], but at the cost of worse rates in  $n$ . Theorem 2 avoids exponential factors and achieves MID scaling with a  $1/\sqrt{n}$  rate.

We finally remark that Theorem 7 only uses the boundedness of  $y$ , and can be extended to bound the convergence of other conditional expectations  $\mathbb{E}_{\pi_\varepsilon}[\alpha(x, y) | x]$  for bounded  $\alpha: \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}^k$ . Such generality is of interest, since some researchers study alternative definitions of entropic OT maps when the cost is not  $\ell_2^2$ . For example, when  $c(x, y) = h(x - y)$  for strictly convex  $h$ , a very recent work suggested a different definition of  $T_\varepsilon$  based on analogy to the un-regularized OT map in this case [CKA23]. If  $h$  satisfies Assumption 1 and is additionally strongly convex and differentiable everywhere, it is not hard to check that Theorem 7 also holds for the maps introduced in that work.

**MID scaling for entropic OT density estimation.** We can also consider estimating the full entropic OT density  $p_\varepsilon$ , defined in (2.3). A natural way to estimate  $p_\varepsilon$  from samples is with its plug-in counterpart,  $\hat{p}_\varepsilon$  from (2.4). We obtain the following result.

**Theorem 8.** *For numerical constants independent of all problem parameters,*

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \lesssim \left(1 + \frac{1}{\sqrt{\varepsilon}}\right) \left(\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}\right)^{1/4}.$$

This result shows that MID scaling even applies when estimating the full density  $p_\varepsilon$ , albeit in the empirical  $L^1(\hat{\mu} \otimes \hat{\nu})$  norm. Previously known bounds for the convergence of  $\hat{p}_\varepsilon$  to  $p_\varepsilon$  incurred an exponential dependence on  $1/\varepsilon$  [RS22], and Theorem 8 removes this dependence while only incurring MID scaling.

We remark that, unlike Theorem 2 on MID scaling for values, MID scaling for maps and densities doesn't yet have analogs in the un-regularized setting, to the best of our knowledge. Developing such analogs is an interesting direction for future work. In the next section, we demonstrate a stronger form of MID scaling for these problems when the low-dimensional distribution is supported on an embedded manifold.

### 3.3 Fast rates with MID scaling on embedded manifolds

In this section, we describe how we can strengthen the results from the previous section by assuming the low-dimensional measure is supported on an embedded manifold.

**Assumptions.** We work under the following assumptions, and for simplicity assume that  $\nu$  is the low-dimensional measure. See Appendix A for a review of embedded Riemannian manifolds, as well as a discussion of the tools from the theory of random geometric graphs on embedded manifolds that we use in our proofs.

**Assumption 2** ( $\nu$  is supported on an embedded manifold). *Assume  $\nu$  is supported on a compact, smooth, connected, Riemannian manifold  $(N, h)$  of dimension  $d_\nu \geq 3$  without boundary, where  $N$  is endowed with the submanifold geometry from its inclusion in  $\mathbb{R}^d$ .*

We also assume that  $\nu$  is compatible with  $N$  in the following sense.

**Assumption 3** ( $\nu$  is Lipschitz and non-vanishing). *Assume that  $\nu$  has a Lipschitz, non-vanishing density with respect to the Riemannian volume on  $N$ .*

These assumptions represent a natural instantiation of the manifold hypothesis while still furnishing enough analytical structure to yield stronger results than in the previous section. In this section and associated proofs, Assumptions 2 and 3 are made throughout, in addition to Assumption 1.

**Suppressing constants depending on  $N$  and  $\nu$ .** In contrast to the previous section, in this section and associated proofs we will generally suppress constants depending on  $N$  and  $\nu$ . The suppressed constants are strictly a function of the intrinsic dimension  $d_\nu$  (and not the extrinsic dimension  $d$ ), the geometry of  $N$ , and uniform bounds on the density of  $\nu$  and its Lipschitz constant. In Appendix A, where we collect all the background and necessary results on embedded manifolds and random geometric graphs, we describe the relevant geometric quantities from the manifold  $N$ .

**Main result on dual potential convergence.** The purpose of this section is to show that under these assumptions on  $\nu$  and  $N$ , we can derive significantly stronger instances of MID scaling than in the previous section. Our main result concerns the convergence of the empirical dual potentials  $\hat{f}_\varepsilon, \hat{g}_\varepsilon$  to their population counterparts  $f_\varepsilon, g_\varepsilon$ . Such convergence is essentially stronger than that of entropic OT values, maps, and densities, since each of those quantities is defined in terms of the dual potentials. Since this result measures convergence in population norms, we remind the reader that the out of sample extensions of  $\hat{f}_\varepsilon, \hat{g}_\varepsilon$  as defined in (2.7) and (2.8), respectively, are in full effect.

**Theorem 9.** *If  $\varepsilon/L$  is sufficiently small, then*

$$\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{13d_\nu+8} \cdot \frac{1}{n}.$$

*We also have convergence in empirical norms*

$$\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{9d_\nu+4} \cdot \frac{1}{n}.$$

Some remarks on this result are in order. First, as in all the results of this section, we require  $\varepsilon/L$  to be sufficiently small; the precise size is given in Equation (A.1). Second, the result concerns the sum of the squared norms, but this is merely for a convenient statement, and the proofs for each term are separate and different. Third, the empirical norm convergence is useful for some of our applications below, while also being an important preliminary step to the full population norm results. Fourth, we remark that, to the best of our knowledge, all previously known bounds on entropic OT dual potentials incurred an exponential dependence on  $\varepsilon$  [LSPC19, dBSLNW23, RS22]. The significance of Theorem 9 is that it avoids exponential dependence on  $\varepsilon$  while only incurring minimum intrinsic dimension-dependence, in accordance with MID scaling.

**MID scaling in the embedded manifold setting.** As do all the results in this section, Theorem 9 evinces a slightly modified form of MID scaling as compared to the previous sections. In our embedded manifold setting, the minimum part of MID scaling simply means that the intrinsic dimension of  $\mu$  doesn't appear in the bounds (remarkably, nothing about

$\mu$  appears in the bound at all); if  $\mu$  were also assumed to satisfy the above assumptions on a manifold of dimension  $d_\mu$ , we could write bounds with an appropriate minimum. In terms of the intrinsic dimension part of MID scaling, there is only a superficial difference, since although the rates in this section do not explicitly involve covering numbers at scale  $\varepsilon/L$  but instead the power  $(L/\varepsilon)^{d_\nu}$ , the two are, in fact, comparable (Proposition 43). And finally, as we mentioned above, there are hidden constants depending on  $N$  and  $\nu$ , and especially the intrinsic dimension  $d_\nu$ , as opposed to the previous results which had only numerical constants.

**Applications of dual convergence.** We demonstrate the power of Theorem 9 with a few applications. Our first two results improve upon the map and density estimation results from the previous section. For each of these applications, we again require  $\varepsilon/L$  to be sufficiently small, as specified in Equation (A.1).

Recall the setting of Theorem 7: we wish to estimate  $T_\varepsilon(x) := \mathbb{E}_{\pi_\varepsilon}[y | x]$  from samples, and we use the plug-in estimator  $\hat{T}_\varepsilon(x) := \mathbb{E}_{\hat{\pi}_\varepsilon}[y | x]$ . Using the canonical extensions of the empirical dual potentials  $\hat{f}_\varepsilon, \hat{g}_\varepsilon$  and correspondingly the empirical density  $\hat{p}_\varepsilon$  as in equations (2.7), (2.8), and (2.4), we can extend  $\hat{T}_\varepsilon$  to a map in  $L^2(\mu)$ , and consider convergence in this population norm. This extended map  $\hat{T}_\varepsilon$  can be evaluated in linear time once the empirical dual potentials are known, and was originally proposed as a computationally efficient estimator for the un-regularized OT map [PNW21]. We obtain the following corollary of Theorem 9.

**Corollary 10.** *If  $\varepsilon/L$  is sufficiently small, and the diameter of  $\text{supp}(\nu)$  is at most  $R$ , then*

$$\mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\mu)}^2] \lesssim R^2 \cdot \left(1 + \frac{1}{\varepsilon^4}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{11d_\nu+4} \cdot \frac{1}{n}$$

Corollary 10 improves over Theorem 7 in that it achieves a fast  $1/n$  rate of convergence in the full population norm  $L^2(\mu)$ .

We can also strengthen our result on density estimation from the previous section, Theorem 8. As in the case of map estimation, the canonical extensions of the empirical density  $\hat{p}_\varepsilon$  described in section 2, allow us to consider  $\hat{p}_\varepsilon$  as an out of sample estimator for  $p_\varepsilon$ .

**Corollary 11.** *If  $\varepsilon/L$  is sufficiently small, then*

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] \lesssim \left(1 + \frac{1}{\varepsilon^4}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{15d_\nu+8} \cdot \frac{1}{n}.$$

This result shows that, in the embedded manifold setting, Theorem 8 can be strengthened to full population norm convergence,  $L^2$  rather than  $L^1$ , and to faster rates.

Our final application concerns the *bias* of entropic OT, namely the quantity  $\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu})] - S_\varepsilon(\mu, \nu)$ . Recent work established that the bias converges at a fast  $1/n$  rate, yet with an exponential dependence on the regularization parameter  $\varepsilon$  [RS22, dBSLW23]. The following result removes this exponential dependence and achieves MID scaling. We state this result as a Theorem since, unlike the previous Corollaries, it doesn't directly follow from Theorem 9.



**Theorem 12.** *If  $\varepsilon/L$  is sufficiently small, then*

$$|\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu})] - S_\varepsilon(\mu, \nu)| \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{9d_\nu+4} \cdot \frac{1}{n}.$$

In sum, the results in this section demonstrate a strong form of the MID scaling phenomenon in the relatively general setting of Assumptions 2 and 3. These results underscore the main message of our work: MID scaling is a generic phenomenon for entropic OT. They demonstrate that MID scaling is certainly not limited to value estimation or map and density estimation in empirical norms with slow  $1/\sqrt{n}$  rates, but in the embedded manifold setting even holds for dual potentials, maps, densities, and biases, all with fast  $1/n$  rates in population norms.

## 4 Preliminary results

In this section, we fix final pieces of notation and describe some simple observations that form the foundation of our approach.

### 4.1 Concavity for the empirical dual

Given probability measures  $P, Q$  on  $\mathbb{R}^d$ , we define the entropic OT dual function  $\Phi_\varepsilon^{PQ} : L^\infty(P) \times L^\infty(Q) \rightarrow \mathbb{R}$  as

$$\Phi_\varepsilon^{PQ}(f, g) := P(f) + Q(g) - \varepsilon(P \otimes Q)(e^{-\varepsilon^{-1}(c-f-g)} - 1).$$

With this notation, we can express the population dual  $\Phi_\varepsilon = \Phi_\varepsilon^{\mu\nu}$  and the empirical dual  $\hat{\Phi}_\varepsilon = \Phi_\varepsilon^{\hat{\mu}\hat{\nu}}$ . We collect some basic facts about entropic dual functions below.

**Proposition 13.** *Let  $P, Q$  be probability measures on  $\mathbb{R}^d$ . The corresponding entropic OT dual function  $\Phi_\varepsilon^{PQ}$  has the following properties:*

1. **(Definition of gradient.)** *For every pair  $h_1 = (f_1, g_1) \in L^\infty(P) \times L^\infty(Q)$ , there exists an element of  $L^\infty(P) \times L^\infty(Q)$  which we denote by  $\nabla\Phi_\varepsilon^{PQ}(f_1, g_1)$  such that for all  $h_0 = (f_0, g_0) \in L^\infty(P) \times L^\infty(Q)$ ,*

$$\begin{aligned} \langle \nabla\Phi_\varepsilon^{PQ}(h_1), h_0 \rangle_{L^2(P) \times L^2(Q)} &= \int f_0(x) \left(1 - \int e^{-\varepsilon^{-1}(c(x,y)-f_1(x)-g_1(y))} dQ(y)\right) dP(x) \\ &\quad + \int g_0(y) \left(1 - \int e^{-\varepsilon^{-1}(c(x,y)-f_1(x)-g_1(y))} dP(x)\right) dQ(y). \end{aligned}$$

*In other words, the gradient of  $\Phi_\varepsilon^{PQ}$  at  $(f_1, g_1)$  is the marginal error corresponding to  $(f_1, g_1)$ .*

2. **(Concavity.)** *For any two pairs of dual potentials  $h_0, h_1 \in L^\infty(P) \times L^\infty(Q)$ , we have the inequalities*

$$\Phi_\varepsilon^{PQ}(h_0) - \Phi_\varepsilon^{PQ}(h_1) \leq \langle \nabla\Phi_\varepsilon^{PQ}(h_1), h_0 - h_1 \rangle_{L^2(P) \times L^2(Q)}, \quad (4.1)$$

and

$$\langle \nabla\Phi_\varepsilon^{PQ}(h_0), h_0 - h_1 \rangle_{L^2(P) \times L^2(Q)} \leq \Phi_\varepsilon^{PQ}(h_0) - \Phi_\varepsilon^{PQ}(h_1). \quad (4.2)$$

3. **(Marginal rounding improves the dual objective.)** Let  $(f, g) \in L^\infty(P) \times L^\infty(Q)$ , and set

$$\tilde{f}(x) := -\varepsilon \log \left( \int e^{-\varepsilon^{-1}(c(x,y)-g(y))} dQ(y) \right).$$

Then

$$\Phi_\varepsilon^{PQ}(f, g) \leq \Phi_\varepsilon^{PQ}(\tilde{f}, g).$$

And analogously when marginal rounding in the  $g$  variable.

*Proof of Proposition 13.* The first item is a definition. For the second item, write

$$\Phi_\varepsilon^{PQ}(f, g) = (P \otimes Q)(f(x) + g(y) - \varepsilon e^{-\varepsilon^{-1}(c(x,y)-f(x)-g(y))}) + \varepsilon,$$

and observe that the function  $t \mapsto t - \varepsilon e^{-\varepsilon^{-1}(c-t)}$  is concave. The result follows by applying the concavity of the integrand pointwise and collecting terms. The third item follows from the first two.  $\square$

## 4.2 Pointwise control and log-Lipschitz bounds

In this section we describe some simple yet powerful implications of Assumption 1.

The following pointwise bounds appear in a number of places [MNW19, DMG20, RS22], and so their proof is omitted.

**Proposition 14** (Pointwise control on dual potentials). *Under our normalization conventions that  $\nu(g_\varepsilon) = 0$  and  $\hat{\nu}(\hat{g}_\varepsilon) = 0$ , we have the uniform bounds*

$$\|\hat{f}_\varepsilon\|_{L^\infty(\mu)}, \|\hat{g}_\varepsilon\|_{L^\infty(\nu)} \leq 2, \quad \|f_\varepsilon\|_{L^\infty(\mu)}, \|g_\varepsilon\|_{L^\infty(\nu)} \leq 1.$$

The following Lipschitz bounds are also well-known [DMG20], but since they are the foundation of our approach we include their proof.

**Proposition 15** (Lipschitz bounds). *The population dual potentials  $f_\varepsilon$  and  $g_\varepsilon$  are  $L$ -Lipschitz over  $\text{supp}(\mu)$  and  $\text{supp}(\nu)$ , respectively. The extended empirical dual potentials  $\hat{f}_\varepsilon$  and  $\hat{g}_\varepsilon$  are also  $L$ -Lipschitz over  $\text{supp}(\mu)$  and  $\text{supp}(\nu)$ , respectively. In particular, the population density  $p_\varepsilon$  and the extended empirical density  $\hat{p}_\varepsilon$  are each  $\frac{2L}{\varepsilon}$ -log-Lipschitz in each of their variables over  $\text{supp}(\mu) \times \text{supp}(\nu)$ .*

*Proof.* By the marginal constraints (2.5), we have that for all  $x \in \text{supp}(\mu)$ ,

$$e^{-\frac{1}{\varepsilon}f_\varepsilon(x)} = \int e^{-\frac{1}{\varepsilon}(c(x,y)-g_\varepsilon(y))} d\nu(y).$$

Using Assumption 1 on the cost  $c$  being Lipschitz, we see that for  $x, x' \in \text{supp}(\mu)$

$$e^{-\frac{1}{\varepsilon}f_\varepsilon(x)} \leq e^{-\frac{1}{\varepsilon}f_\varepsilon(x') + \frac{L}{\varepsilon}\|x-x'\|}$$

By symmetry it follows that

$$|f_\varepsilon(x) - f_\varepsilon(x')| \leq L\|x - x'\|.$$

The analogous argument establishes the claim for  $g_\varepsilon$ , and also proves the claims for the extended empirical potentials  $\hat{f}_\varepsilon$  and  $\hat{g}_\varepsilon$ .

For the claims about the densities, note that

$$\log p_\varepsilon(x, y) = -\frac{1}{\varepsilon}(c(x, y) - f_\varepsilon(x) - g_\varepsilon(y)),$$

and so the claim follows from the previous ones for  $f_\varepsilon$  and  $g_\varepsilon$ , as well as Assumption 1. Similarly for  $\log \hat{p}_\varepsilon$ .  $\square$

## 5 Proofs of MID scaling with slow rates

In this section, we give our proofs of MID scaling for values, Theorem 2, MID scaling for maps in empirical norms, Theorem 7, and MID scaling for densities in empirical norms, Theorem 8. The proofs are organized in the following manner. In section 5.1, we give a detailed exposition of our approach in the case of value estimation. In section 5.2, we give the proof of one of our main technical lemmas, and the source of MID scaling, on sub-exponential bounds for  $\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2$ . In section 5.3, we show how to reduce map estimation to the estimates from the previous sections, and do the same for density estimation in section 5.4.

### 5.1 Proof of value rate and overview of technical approach

In this section, we illustrate the main ideas of our technical approach by describing our proof of Theorem 2 on the convergence of entropic OT values. To situate our approach, note that most existing statistical work on entropic OT studies smooth cost functions and shows that the smoothness of the cost functions implies smoothness for the dual potentials, leading to small function classes over which empirical processes can be well-controlled [GCB<sup>+</sup>19, LSPC19, MNW19, dBSLNW23, BEZ22]. The present work builds on a different approach, developed in [RS22], which entirely avoids empirical process theory by using strong concavity of the empirical dual objective. The benefit of this approach is that it is simple, requires no smoothness assumptions on the cost, leads to dimension-free bounds, and can be used to prove fast rates for most entropic OT quantities [RS22]. However, because it is dimension-free, it is not suited to a fine understanding of the dimension-dependence of entropic OT, and in particular it incurs exponential factors of  $1/\varepsilon$  at many points in the arguments. The main technical goal of this work is to refine this strong concavity of the empirical dual approach by replacing the exponential dependence on  $1/\varepsilon$  with a very fine dependence on the dimension of the problem: MID scaling.

To this end, note that the exponential dependence of [RS22] primarily arises from two separate sources: first, the strong concavity of the empirical dual objective, and second, pointwise bounds on the entropic density  $p_\varepsilon$ . For the results in this section, we avoid the exponential factors arising from strong concavity by using more delicate arguments which rely on only concavity, and remove the exponential factors from the pointwise bounds on the entropic density  $p_\varepsilon$  with a novel estimate (Lemma 16 below) which is itself the source of MID scaling.

To present the ideas as clearly as possible, we will suppress numerical constants with the notation  $u \lesssim v$ ; they can easily be extracted from the proof. Recall that  $\Phi_\varepsilon$  is

the population entropic OT dual function, and  $\hat{\Phi}_\varepsilon$  is its empirical counterpart. Then Theorem 2 concerns the quantity

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] = \mathbb{E}[|\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)|].$$

To bound this quantity, we first decompose it in the following manner:

$$\begin{aligned} \mathbb{E}[|\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)|] &\leq \mathbb{E}[|\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)|] + \mathbb{E}[|\hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)|] \\ &= \mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] + \mathbb{E}[|\hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)|]. \end{aligned}$$

For convenience, we refer to the first term as a bias term, and the second as a variance term. Note that our usage of “bias” and “variance” in this context is not intended to be standard, merely suggestive.

**Variance term.** If we just examine this variance term, we can compute:

$$\begin{aligned} &\mathbb{E}[|\hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)|] \\ &= \mathbb{E}[|(\hat{\mu} - \mu)(f_\varepsilon) + (\hat{\nu} - \nu)(g_\varepsilon) - \varepsilon(\hat{\mu} \otimes \hat{\nu} - \mu \otimes \nu)(p_\varepsilon)|] \\ &\leq \mathbb{E}[((\hat{\mu} - \mu)(f_\varepsilon) + (\hat{\nu} - \nu)(g_\varepsilon) - \varepsilon(\hat{\mu} \otimes \hat{\nu} - \mu \otimes \nu)(p_\varepsilon))^2]^{1/2} \\ &\lesssim (\mathbb{E}[(\hat{\mu} - \mu)(f_\varepsilon)^2] + \mathbb{E}[(\hat{\nu} - \nu)(g_\varepsilon)^2] + \varepsilon^2 \mathbb{E}[(\hat{\mu} \otimes \hat{\nu} - \mu \otimes \nu)(p_\varepsilon)^2])^{1/2} \\ &\leq \mathbb{E}[(\hat{\mu} - \mu)(f_\varepsilon)^2]^{1/2} + \mathbb{E}[(\hat{\nu} - \nu)(g_\varepsilon)^2]^{1/2} + \varepsilon \mathbb{E}[(\hat{\mu} \otimes \hat{\nu} - \mu \otimes \nu)(p_\varepsilon)^2]^{1/2}. \end{aligned}$$

For the first two terms, we use the pointwise boundedness of the dual potentials from Proposition 14 to yield

$$\mathbb{E}[(\hat{\mu} - \mu)(f_\varepsilon)^2]^{1/2} + \mathbb{E}[(\hat{\nu} - \nu)(g_\varepsilon)^2]^{1/2} = \sqrt{\frac{\text{Var}_\mu(f_\varepsilon)}{n}} + \sqrt{\frac{\text{Var}_\nu(g_\varepsilon)}{n}} \lesssim \frac{1}{\sqrt{n}}.$$

For the third term, the marginal constraints for  $p_\varepsilon$  from (2.5) allow us to cancel cross-terms and compute

$$\begin{aligned} \mathbb{E}[(\hat{\mu} \otimes \hat{\nu})(p_\varepsilon - (\mu \otimes \nu)(p_\varepsilon))^2]^{1/2} &= \mathbb{E}[(\hat{\mu} \otimes \hat{\nu})(p_\varepsilon - 1)^2]^{1/2} \\ &= \left( \frac{1}{n^4} \sum_{i,j,k,l=1}^n \mathbb{E}[(p_\varepsilon(x_i, y_j) - 1)(p_\varepsilon(x_k, y_l) - 1)] \right)^{1/2} \\ &= \frac{1}{n} \sqrt{\text{Var}_{\mu \otimes \nu}(p_\varepsilon)} \leq \frac{1}{n} \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}. \end{aligned}$$

We note that this kind of calculation is used repeatedly throughout the paper. We finally arrive at the bound

$$\begin{aligned} \mathbb{E}[|\hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)|] &\lesssim \frac{1}{\sqrt{n}} + \frac{\varepsilon \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}}{n} \\ &\lesssim (1 + \varepsilon) \frac{\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}}{\sqrt{n}}, \end{aligned} \tag{5.1}$$

where the last inequality follows because  $(\mu \otimes \nu)(p_\varepsilon) = 1$  implies  $1 \leq \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}$  by Cauchy-Schwarz.

**Bias term.** We use concavity of the empirical dual to control the bias term  $\mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)]$  by a quantity involving  $\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}$  again. Indeed, by Proposition 13 on concavity of  $\hat{\Phi}_\varepsilon$  and Proposition 14 on pointwise control for the dual potentials, we have

$$\begin{aligned} \mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] &\leq \mathbb{E}[\langle \nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}] \\ &\leq \mathbb{E}[\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})} \cdot \|(\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}] \\ &\lesssim \mathbb{E}[\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2]^{1/2}. \end{aligned}$$

Using the marginal constraints (2.5) to cancel cross terms once more, we find

$$\begin{aligned} \mathbb{E}[\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2] &= \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \left(\frac{1}{n} \sum_{j=1}^n p_\varepsilon(x_i, y_j) - 1\right)^2\right. \\ &\quad \left. + \frac{1}{n} \sum_{j=1}^n \left(\frac{1}{n} \sum_{i=1}^n p_\varepsilon(x_i, y_j) - 1\right)^2\right] \\ &= \frac{1}{n^3} \sum_{i, j_0, j_1=1}^n \mathbb{E}[(p_\varepsilon(x_i, y_{j_0}) - 1)(p_\varepsilon(x_i, y_{j_1}) - 1)] \\ &\quad + \frac{1}{n^3} \sum_{i_0, i_1, j=1}^n \mathbb{E}[(p_\varepsilon(x_{i_0}, y_j) - 1)(p_\varepsilon(x_{i_1}, y_j) - 1)] \\ &= \frac{2}{n} \text{Var}_{\mu \otimes \nu}(p_\varepsilon) \lesssim \frac{\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2}{n}. \end{aligned}$$

We arrive at an important, if elementary, bound that we will use frequently throughout this work,

$$\mathbb{E}[\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2] \lesssim \frac{\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2}{n}. \quad (5.2)$$

In particular, the bias is bounded as

$$\mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] \lesssim \frac{\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}}{\sqrt{n}}. \quad (5.3)$$

**Controlling**  $\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}$ . We can combine (5.1) and (5.3) to obtain

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \frac{\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}}{\sqrt{n}}. \quad (5.4)$$

We have thus reduced the problem to terms with a dimension-free dependence on the sample size  $n$ , and with the dimension and  $\varepsilon$ -dependence residing fully in the term  $(1 + \varepsilon)\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}$ . Up to this point, the differences between the techniques introduced in [RS22] and our approach are essentially contained in two steps, each taken to avoid factors of  $e^{1/\varepsilon}$ . First, we control the bias term with concavity rather than strong concavity. And second, we refrain from using the pointwise control of Proposition 14 to bound the  $p_\varepsilon$  terms, which would result in the bound  $\|p_\varepsilon\|_{L^\infty(\mu \otimes \nu)} \leq e^{C/\varepsilon}$  for a constant  $C$ . Such pointwise and exponential bounds were, to the best of our knowledge, all that was known about  $p_\varepsilon$  until this work.

One of the core technical innovations in this work, and indeed the source of MID scaling, is to provide the following sub-exponential, dimension-dependent bound on  $p_\varepsilon$ .

**Lemma 16** (MID scaling for the norm of entropic densities). *We have*

$$\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2 \lesssim \mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L}).$$

This result is proven, in the following section, by using the marginal constraints that  $p_\varepsilon$  satisfies in Equation (2.5) to gain pointwise control on  $p_\varepsilon$  that, when integrated, yields the covering numbers. In the embedded manifold setting of section 6, the  $L^2$  bound of Lemma 16 can be strengthened to pointwise  $L^\infty$  control, using the same technique (Lemma 26). The proof technique is used again in section 6.2, and we indeed expect that it will be of some broader utility. For example, it may be helpful for calculating quantities arising in the asymptotic distributions of entropic OT quantities [GSH23].

For our subsequent results, it is convenient to record the following Lemma, which results from combining Equation (5.3) with Lemma 16.

**Lemma 17** (MID scaling for the bias). *We have*

$$\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)] \lesssim \sqrt{\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}}.$$

**Summary of the technical approach.** In summary, we decomposed the difference  $|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|$  into a bias term and a variance term. For the variance term, an elementary calculation allows us to control it by something with the dimension of the problem residing purely in the  $\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}$  term. And for the bias term, a judicious use of concavity allows us to control it by another expression with the dimension of the problem purely in the  $\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}$  term. We then conclude with Lemma 16. The proofs of MID scaling with slow rates for maps and densities, Theorem 7 and Theorem 8, use additional estimates that also follow the pattern of splitting into bias and variance terms (loosely construed) to then reduce to Lemma 16, and are included in sections 5.3 and 5.4, respectively.

## 5.2 MID scaling for the norm of entropic densities

In this section, we will prove Lemma 16, which gives sub-exponential, MID scaling-type bounds on the entropic density  $p_\varepsilon$ . To this end, the following fact will be useful. Recall that we write the closed ball of size  $\delta$  at  $z \in \mathbb{R}^d$  as

$$B^{\text{cl}}(z, \delta) := \{z' \in \mathbb{R}^d : \|z - z'\| \leq \delta\}.$$

**Proposition 18** (Average inverse mass is bounded by the covering number). *Suppose  $P$  is a compactly supported probability measure on  $\mathbb{R}^d$ . Then for all  $\delta > 0$ ,*

$$\int P(B^{\text{cl}}(z, \delta))^{-1} dP(z) \leq \mathcal{N}(P, \delta/4).$$

To give the proof of this Proposition, recall that for  $\tau > 0$ , a *proper*  $\tau$ -covering of a set  $A \subset \mathbb{R}^d$ , is a covering of  $A$  at scale  $\tau$  with centers contained within  $A$ . We write the minimal size of a proper covering with respect to the Euclidean norm at scale  $\tau > 0$  as  $\mathcal{N}^{\text{pr}}(A, \|\cdot\|, \tau)$ . Note that proper covering numbers are comparable to vanilla covering numbers since  $\mathcal{N}^{\text{pr}}(A, \|\cdot\|, \tau) \leq \mathcal{N}(A, \|\cdot\|, \tau/2)$ .

*Proof of Proposition 18.* Let  $z_1, \dots, z_K \in \text{supp}(P)$  be a proper  $\delta/2$  covering of  $\text{supp}(P)$  achieving  $K = \mathcal{N}^{\text{pr}}(\text{supp}(P), \|\cdot\|, \delta/2)$ . Since this is a proper covering, we know that for all  $k \in [K]$ ,  $P(B^{\text{cl}}(z_k, \delta/2)) > 0$ . The triangle inequality then implies

$$z \in B^{\text{cl}}(z_k, \delta/2) \implies 0 < P(B^{\text{cl}}(z_k, \delta/2)) \leq P(B^{\text{cl}}(z, \delta)).$$

Thus

$$\begin{aligned} \int P(B^{\text{cl}}(z, \delta))^{-1} dP(z) &\leq \sum_{k=1}^K \int_{B^{\text{cl}}(z_k, \delta/2)} P(B^{\text{cl}}(z, \delta))^{-1} dP(z) \\ &\leq \sum_{k=1}^K \int_{B^{\text{cl}}(z_k, \delta/2)} P(B^{\text{cl}}(z_k, \delta/2))^{-1} dP(z) \\ &= K = \mathcal{N}^{\text{pr}}(P, \delta/2) \leq \mathcal{N}(P, \delta/4). \end{aligned}$$

□

*Proof of Lemma 16.* We show the bound for  $\mathcal{N}(\nu, \frac{\varepsilon}{L})$ , the bound for  $\mathcal{N}(\mu, \frac{\varepsilon}{L})$  follows by symmetry. We proceed via the marginal constraints on  $p_\varepsilon$  from (2.5): for all  $x \in \text{supp}(\mu)$  and  $y \in \text{supp}(\nu)$ ,

$$1 = p_\varepsilon(x, y) \int \frac{p_\varepsilon(x, y')}{p_\varepsilon(x, y)} d\nu(y'). \quad (5.5)$$

Applying Proposition 15 to Equation (5.5) yields,

$$\begin{aligned} 1 &\geq p_\varepsilon(x, y) \int_{B^{\text{cl}}(y, \frac{4\varepsilon}{L})} \frac{p_\varepsilon(x, y')}{p_\varepsilon(x, y)} d\nu(y') \\ &\geq p_\varepsilon(x, y) \int_{B^{\text{cl}}(y, \frac{4\varepsilon}{L})} e^{-\frac{2L}{\varepsilon} \|y' - y\|} d\nu(y') \\ &\geq e^{-8} \cdot p_\varepsilon(x, y) \cdot \nu(B^{\text{cl}}(y, \frac{4\varepsilon}{L})). \end{aligned}$$

Since  $y \in \text{supp}(\nu)$ ,  $\nu(B^{\text{cl}}(y, \frac{4\varepsilon}{L})) > 0$ , so we can rearrange this inequality and apply the marginal constraints (2.5) once more to yield

$$\begin{aligned} \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2 &= \int p_\varepsilon(x, y)^2 d(\mu \otimes \nu)(x, y) \\ &\lesssim \int \nu(B^{\text{cl}}(y, \frac{4\varepsilon}{L}))^{-1} p_\varepsilon(x, y) d(\mu \otimes \nu)(x, y) \\ &= \int \nu(B^{\text{cl}}(y, \frac{4\varepsilon}{L}))^{-1} d\nu(y). \end{aligned}$$

The result follows from Proposition 18. □



### 5.3 MID scaling for maps in empirical norm

As we discuss in the previous section, to prove Theorem 7 and Theorem 8, we reduce to the bound on the bias from Lemma 17. We emphasize that bounds on entropic maps and densities with exponential dependence on  $1/\varepsilon$  are known from [RS22]. To prove such bounds without incurring exponential factors requires further work beyond the previous sections. As in some previous works, we reduce from the error of map estimation to the error of value estimation; the difference in our approach is that we avoid factors of the KL-divergence [PDNW23] and it applies to the empirical entropic OT map rather than a modified estimator [Str23].

The idea is to consider the following rounded dual potential, defined for all  $x \in \text{supp}(\mu)$  as

$$\tilde{f}_\varepsilon(x) := -\varepsilon \log \left( \frac{1}{n} \sum_{j=1}^n e^{-\varepsilon^{-1}(c(x, y_j) - g_\varepsilon(y_j))} \right). \quad (5.6)$$

Let the corresponding density be defined for all  $x \in \text{supp}(\mu)$  and  $y \in \text{supp}(\nu)$  as  $\tilde{p}_\varepsilon(x, y) := e^{-\varepsilon^{-1}(c(x, y) - \tilde{f}_\varepsilon(x) - g_\varepsilon(y))}$ ; note that  $\tilde{p}_\varepsilon$  is such that for all  $x \in \text{supp}(\mu)$ ,

$$1 = \frac{1}{n} \sum_{j=1}^n \tilde{p}_\varepsilon(x, y_j).$$

This fact allows us to compare  $\frac{1}{n} \hat{p}_\varepsilon(x, \cdot)$  with  $\frac{1}{n} \tilde{p}_\varepsilon(x, \cdot)$  as probability distributions on  $\mathcal{Y}$ . The following Lemma is then a consequence of Pinsker's inequality, cancellation of some terms, and a crucial use of the monotonicity of the empirical dual under marginal rounding from Proposition 13 to replace  $\tilde{f}_\varepsilon$  with its un-rounded counterpart  $f_\varepsilon$ .

**Lemma 19.** *Let  $\tilde{f}_\varepsilon$  and  $\tilde{p}_\varepsilon$  be as above, and suppose the diameter of  $\text{supp}(\nu)$  is at most  $R$ . Then*

$$\mathbb{E} \left[ \left\| \hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j \tilde{p}_\varepsilon(x, y_j) \right\|_{L^2(\hat{\mu})}^2 \right] \lesssim \frac{R^2}{\varepsilon} \mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)].$$

However, this lemma does not yet yield Theorem 7 as it involves the rounded density  $\tilde{p}_\varepsilon$ . The following calculation addresses this issue.

**Lemma 20.** *Let  $\tilde{f}_\varepsilon$  and  $\tilde{p}_\varepsilon$  be as above. Then*

$$\mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\hat{\mu})}^2] \lesssim \mathbb{E} \left[ \left\| \hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j \tilde{p}_\varepsilon(x, y_j) \right\|_{L^2(\hat{\mu})}^2 \right] + R^2 \cdot \frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}.$$

We first dispatch Theorem 7, and then prove these lemmas.

*Proof of Theorem 7.* Combining Lemma 17 with the above two lemmas yields

$$\mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\hat{\mu})}^2] \lesssim R^2 \left\{ \frac{1}{\varepsilon} \cdot \sqrt{\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}} + \frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n} \right\}.$$

Now observe that the statement of Theorem 7 is trivial when  $\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L}) \geq n$ , since we always have the bound  $2R^2$ . Hence it suffices to consider the case  $\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L}) \leq n$ , in which case we may conclude from the above inequality.  $\square$

*Proof of Lemma 19.* To prove this result, let us first assume that  $0 \in \text{supp}(\nu)$  so that for each  $y \in \text{supp}(\nu)$ , we have  $\|y\| \leq R$ . Then for each  $x \in \mathcal{X}$ , we apply triangle inequality and Pinsker's inequality to see that

$$\begin{aligned} \left\| \hat{T}_\varepsilon(x) - \frac{1}{n} \sum_{j=1}^n y_j \tilde{p}_\varepsilon(x, y_j) \right\| &\leq \frac{R}{n} \sum_{j=1}^n |\tilde{p}_\varepsilon(x, y_j) - \hat{p}_\varepsilon(x, y_j)| \\ &\lesssim R \sqrt{\text{KL}\left(\frac{1}{n} \hat{p}_\varepsilon(x, \cdot) \parallel \frac{1}{n} \tilde{p}_\varepsilon(x, \cdot)\right)}. \end{aligned}$$

Therefore,

$$\begin{aligned} \left\| \hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j \tilde{p}_\varepsilon(x, y_j) \right\|_{L^2(\hat{\mu})}^2 &\lesssim R^2 \frac{1}{n} \sum_{i=1}^n \text{KL}\left(\frac{1}{n} \hat{p}_\varepsilon(x_i, \cdot) \parallel \frac{1}{n} \tilde{p}_\varepsilon(x_i, \cdot)\right) \\ &= \frac{R^2}{\varepsilon} \sum_{i,j=1}^n \frac{1}{n^2} \hat{p}_\varepsilon(x_i, y_j) (\hat{f}_\varepsilon(x_i) - \tilde{f}_\varepsilon(x_i) + \hat{g}_\varepsilon(y_j) - g_\varepsilon(y_j)) \\ &= \frac{R^2}{\varepsilon} (\hat{\mu}(\hat{f}_\varepsilon - \tilde{f}_\varepsilon) + \hat{\nu}(\hat{g}_\varepsilon - g_\varepsilon)), \end{aligned}$$

where we used the marginal constraints of  $\hat{p}_\varepsilon$  for the last step. Using the notation, as in the previous section,  $\hat{\Phi}_\varepsilon := \Phi_\varepsilon^{\hat{\mu}\hat{\nu}}$ , we recognize the result as a difference of  $\hat{\Phi}_\varepsilon$ , and apply Proposition 13 on marginal rounding improving dual objective value, to conclude that

$$\begin{aligned} \left\| \hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j \tilde{p}_\varepsilon(x, y_j) \right\|_{L^2(\hat{\mu})}^2 &\lesssim \frac{R^2}{\varepsilon} (\hat{\mu}(\hat{f}_\varepsilon - \tilde{f}_\varepsilon) + \hat{\nu}(\hat{g}_\varepsilon - g_\varepsilon)) \\ &= \frac{R^2}{\varepsilon} (\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(\tilde{f}_\varepsilon, g_\varepsilon)) \\ &\leq \frac{R^2}{\varepsilon} (\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)). \end{aligned}$$

Taking expectations,  $\mathbb{E}[\hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] = \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)$ , and we conclude the result. For the general case where 0 may not be in  $\text{supp}(\nu)$ , we can perform the above argument with suitably offset  $y$ .  $\square$

*Proof of Lemma 20.* As above, let us first assume that  $0 \in \text{supp}(\nu)$  so that for each  $y \in \text{supp}(\nu)$ , we have  $\|y\| \leq R$ . We apply Young's inequality twice, first comparing  $T_\varepsilon(x)$  to the empirical version  $\frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j)$ , and then the empirical version to the version involving  $\tilde{p}_\varepsilon$ :

$$\begin{aligned} \mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\hat{\mu})}^2] &\lesssim \mathbb{E}\left[\left\| \hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j \tilde{p}_\varepsilon(x, y_j) \right\|_{L^2(\hat{\mu})}^2\right] \\ &\quad + \mathbb{E}\left[\left\| \frac{1}{n} \sum_{j=1}^n y_j (\tilde{p}_\varepsilon(x, y_j) - p_\varepsilon(x, y_j)) \right\|_{L^2(\hat{\mu})}^2\right] \\ &\quad + \mathbb{E}\left[\left\| \frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j) - T_\varepsilon(x) \right\|_{L^2(\hat{\mu})}^2\right] \end{aligned}$$

The third term is controlled as

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j) - T_\varepsilon(x) \right\|_{L^2(\hat{\mu})}^2 \right] \\
&= \mathbb{E} \left[ \frac{1}{n^2} \sum_{j,k=1}^n \langle y_j p_\varepsilon(x, y_j) - T_\varepsilon(x), y_k p_\varepsilon(x, y_k) - T_\varepsilon(x) \rangle_{L^2(\hat{\mu})} \right] \\
&= \frac{1}{n} \|y p_\varepsilon(x, y) - T_\varepsilon(x)\|_{L^2(\mu \otimes \nu)}^2 \lesssim \frac{R^2}{n} \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2,
\end{aligned}$$

where the second equality follows because, for  $j \neq k$ ,  $y_j$  and  $y_k$  are iid draws from  $\nu$ , so that these terms cancel. For the second term, we begin by observing that for each  $y_j$ ,

$$\tilde{p}_\varepsilon(x, y_j) = \frac{p_\varepsilon(x, y_j)}{\frac{1}{n} \sum_{k=1}^n p_\varepsilon(x, y_k)}.$$

We thus apply triangle inequality and this equation to see that

$$\begin{aligned}
\left\| \frac{1}{n} \sum_{j=1}^n y_j (\tilde{p}_\varepsilon(x, y_j) - p_\varepsilon(x, y_j)) \right\| &\leq \frac{R}{n} \sum_{j=1}^n |\tilde{p}_\varepsilon(x, y_j) - p_\varepsilon(x, y_j)| \\
&= \frac{R}{n} \sum_{j=1}^n p_\varepsilon(x, y_j) \left| \frac{1}{\frac{1}{n} \sum_{k=1}^n p_\varepsilon(x, y_k)} - 1 \right| \\
&= R \left| \frac{1}{n} \sum_{j=1}^n p_\varepsilon(x, y_j) - 1 \right|.
\end{aligned}$$

Taking the  $\|\cdot\|_{L^2(\hat{\mu})}^2$  norm, we recognize this as the squared norm of the first component of  $\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)$ , and use the bound in Equation (5.2) to conclude

$$\begin{aligned}
\mathbb{E} \left[ \left\| \frac{1}{n} \sum_{j=1}^n y_j (\tilde{p}_\varepsilon(x, y_j) - p_\varepsilon(x, y_j)) \right\|_{L^2(\hat{\mu})}^2 \right] &\leq R^2 \mathbb{E} [\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2] \\
&\lesssim \frac{R^2}{n} \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2.
\end{aligned}$$

Applying Lemma 16 yields the result. For the general case where 0 may not be in  $\text{supp}(\nu)$ , we can perform the above argument with suitably offset  $y$ .  $\square$

#### 5.4 MID scaling for densities in empirical norm

We prove Theorem 8 on empirical norm convergence of the entropic OT density using the techniques introduced in the previous section. We prove the following two lemmas.

**Lemma 21.** *Let  $\tilde{f}_\varepsilon, \tilde{p}_\varepsilon$  be as in the previous section. Then*

$$\mathbb{E} [\|\hat{p}_\varepsilon - \tilde{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \lesssim \frac{1}{\sqrt{\varepsilon}} \mathbb{E} [S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)]^{1/2}$$

**Lemma 22.** Let  $\tilde{f}_\varepsilon, \tilde{p}_\varepsilon$  be as in the previous section. Then

$$\mathbb{E}[\|p_\varepsilon - \hat{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \lesssim \mathbb{E}[\|\hat{p}_\varepsilon - \tilde{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] + \sqrt{\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}}.$$

We first dispatch Theorem 8, and then prove these lemmas.

*Proof of Theorem 8.* Combining the above two lemmas with Lemma 17 yields

$$\mathbb{E}[\|p_\varepsilon - \hat{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \lesssim \frac{1}{\sqrt{\varepsilon}} \left( \frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n} \right)^{1/4} + \sqrt{\frac{\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L})}{n}}.$$

Now, observe that the statement of Theorem 8 is trivial when  $\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L}) \geq n$ , since we always have the bound  $\lesssim 1$  by triangle inequality. Hence it suffices to consider the case  $\mathcal{N}(\mu, \frac{\varepsilon}{L}) \wedge \mathcal{N}(\nu, \frac{\varepsilon}{L}) \leq n$ , in which case we may conclude from the above inequality.  $\square$

*Proof of Lemma 21.* Pinsker's and Jensen's imply

$$\begin{aligned} \|\tilde{p}_\varepsilon - \hat{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})} &\lesssim \frac{1}{n} \sum_{i=1}^n \sqrt{\text{KL}\left(\frac{1}{n} \hat{p}_\varepsilon(x_i, \cdot) \parallel \frac{1}{n} \tilde{p}_\varepsilon(x_i, \cdot)\right)} \\ &\leq \left( \frac{1}{n} \sum_{i=1}^n \text{KL}\left(\frac{1}{n} \hat{p}_\varepsilon(x_i, \cdot) \parallel \frac{1}{n} \tilde{p}_\varepsilon(x_i, \cdot)\right) \right)^{1/2}. \end{aligned}$$

Taking the expectation and applying Jensen's once more yields

$$\mathbb{E}[\|\tilde{p}_\varepsilon - \hat{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \lesssim \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^n \text{KL}\left(\frac{1}{n} \hat{p}_\varepsilon(x_i, \cdot) \parallel \frac{1}{n} \tilde{p}_\varepsilon(x_i, \cdot)\right) \right]^{1/2}.$$

The statement then follows as in Lemma 19.  $\square$

*Proof of Lemma 22.* Apply triangle inequality:

$$\mathbb{E}[\|p_\varepsilon - \hat{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \leq \mathbb{E}[\|\hat{p}_\varepsilon - \tilde{p}_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] + \mathbb{E}[\|\tilde{p}_\varepsilon - p_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}].$$

For the second term we use the same reasoning as in the proof of Lemma 20 to observe that

$$\|\tilde{p}_\varepsilon - p_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})} = \frac{1}{n^2} \sum_{i,j=1}^n |\tilde{p}_\varepsilon(x_i, y_j) - p_\varepsilon(x_i, y_j)| = \frac{1}{n} \sum_{i=1}^n \left| \frac{1}{n} \sum_{j=1}^n p_\varepsilon(x_i, y_j) - 1 \right|.$$

Applying Cauchy-Schwarz we find that

$$\|\tilde{p}_\varepsilon - p_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})} \leq \left( \frac{1}{n} \sum_{i=1}^n \left( \frac{1}{n} \sum_{j=1}^n p_\varepsilon(x_i, y_j) - 1 \right)^2 \right)^{1/2} \leq \|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}.$$

Taking expectations and applying Cauchy-Schwarz once more, we can apply Equation (5.2) to yield

$$\mathbb{E}[\|\tilde{p}_\varepsilon - p_\varepsilon\|_{L^1(\hat{\mu} \otimes \hat{\nu})}] \leq \mathbb{E}[\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2]^{1/2} \lesssim \frac{\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}}{\sqrt{n}}.$$

Finally, Lemma 16 yields the result.  $\square$

## 6 Proofs of MID scaling with fast rates on manifolds

In this section, we give the proofs of our stronger results when  $\nu$  is supported on an embedded Riemannian manifold. To introduce our approach, recall that fast rates of convergence for the entropic OT dual potentials, map, density, and bias are known, but at the price of exponential factors in  $1/\varepsilon$  [RS22]. As we mention in the previous section, those estimates are established through strong concavity of the empirical dual objective, but because the strong concavity parameter is exponentially small, such an approach is inadequate for establishing MID scaling. The techniques in the previous section represent a very weak form of this approach, using only concavity rather than strong concavity to replace exponential factors with MID scaling. In this section, we power our arguments with a *quadratic growth* condition [KNS16], which is a stronger condition than concavity alone, yet still weaker than true strong concavity.

In this regard, our approach is inspired by recent work [Mar21], which has found that a sufficient condition for a quadratic growth condition for the entropic OT dual is a Poincaré inequality. Indeed, [Mar21] shows that given probability measures  $P, Q$  on  $\mathbb{R}^d$ , if  $P$  satisfies a Poincaré inequality with constant  $C_P$ , then for any dual potentials  $f, g$ ,

$$\|f - f_\varepsilon^{PQ}\|_{L^2(P)}^2 \lesssim \frac{L^2}{\varepsilon C_P} (\Phi_\varepsilon^{PQ}(f_\varepsilon^{PQ}, g_\varepsilon^{PQ}) - \Phi_\varepsilon^{PQ}(f, g)), \quad (6.1)$$

where  $f_\varepsilon^{PQ}, g_\varepsilon^{PQ}$  are the optimal dual potentials from  $P$  to  $Q$  and we are suppressing numerical constants. However, this estimate cannot directly establish quadratic growth for the *empirical* dual  $\hat{\Phi}_\varepsilon$  since it relies on a Poincaré inequality for the source measure  $P$ . We emphasize that our approach is fundamentally about the empirical dual rather than the population dual, since working with the empirical dual is what allows us to bypass empirical processes by reducing to computing variances of population quantities (consider, for example, our proof of Lemma 17).

As our first major step in this section, we develop quadratic growth inequalities for the empirical dual. In particular, in section 6.1 we use different techniques than [Mar21] to show that under an empirical analog of a Poincaré inequality – a spectral gap of the random geometric graph (RGG) of  $\nu$  – an analog of (6.1) holds for the empirical dual  $\hat{\Phi}_\varepsilon$ . Using recent advances in the theory of RGGs on embedded manifolds [GTGHS20] which we describe in Appendix A.2, a spectral gap for the RGG of  $\nu$  then follows from a Poincaré inequality for  $\nu$  itself, which is entailed by Assumption 2 and Assumption 3. To summarize, the primary uses of our embedded manifold assumptions are that they imply a spectral gap for a weighted Laplacian associated to  $\nu$  (Proposition 37), which then implies a spectral gap for the RGG of  $\nu$  using spectral convergence theory [GTGHS20], which finally yields quadratic growth for the empirical dual; this argument is encapsulated in Lemma 23.

After we prove Lemma 23 in section 6.1, in section 6.2 we next bound the  $f$  dual potentials in terms of the  $g$  dual potentials, both in population and empirical norms, while only picking up MID scaling factors. These results are combined in section 6.3 to yield convergence of the  $g$  dual potentials in empirical norm, which then implies Theorem 12 and all of Theorem 9 on dual potential convergence except the convergence of the  $g$  dual potentials in population norm. In section 6.4, we complete the proof of Theorem 9 by

applying the previous results to establish the convergence of the  $g$  dual potentials in population norm. Finally, in section 6.5 we prove Corollary 10 and Corollary 11.

Throughout, we work under the assumption that  $n$  is large enough that

$$\left(\frac{\varepsilon}{L}\right)^{2d_\nu} \gtrsim \frac{1}{n} \left( \log \left( \frac{Ln}{\varepsilon} \right) + \frac{1}{\varepsilon} \right). \quad (6.2)$$

We may assume  $n$  is this large without loss of generality, since otherwise it is not hard to check that our bounds are worse than the trivial ones.

## 6.1 Quadratic growth for the empirical dual

In this section, we prove the following quadratic growth inequality for the empirical entropic OT dual function.

**Lemma 23** (Quadratic growth for the empirical dual). *Let  $\Psi_\varepsilon^{\hat{\mu}\hat{\nu}} : L^\infty(\hat{\nu}) \rightarrow \mathbb{R}$  denote the empirical dual objective with  $f$  rounded, namely*

$$\Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(g) := \hat{\mu} \left( -\varepsilon \log \left( \frac{1}{n} \sum_{j=1}^n e^{-\varepsilon^{-1}(c(x, y_j) - g(y_j))} \right) \right) + \hat{\nu}(g).$$

Then, for  $\varepsilon$  sufficiently small, we have

$$\varepsilon \left( \frac{\varepsilon}{L} \right)^{d_\nu + 2} \cdot \|g - \hat{g}_\varepsilon\|_{L^2(\hat{\nu})}^2 \lesssim \|g - \hat{g}_\varepsilon\|_{L^\infty(\hat{\nu})}^2 (\Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(\hat{g}_\varepsilon) - \Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(g)), \quad \forall g \in L^\infty(\hat{\nu}), \hat{\nu}(g) = 0.$$

This lemma is proven by reducing the claim to a spectral gap for the random geometric graph (RGG) of  $\nu$ . Using recent fundamental work on the spectral convergence of RGGs of embedded manifolds [GTGHS20], we conclude the result. We include this background material in Appendix A.2.

*Proof of Lemma 23.* Fix  $g \in L^\infty(\hat{\nu})$  with  $\hat{\nu}(g) = 0$ , and put  $\alpha := g - \hat{g}_\varepsilon$ . Let

$$\psi(t) := -\Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(\hat{g}_\varepsilon + t\alpha),$$

and also define

$$q_t(x_i, y_j) := \frac{e^{-\varepsilon^{-1}(c(x_i, y_j) - (\hat{g}_\varepsilon(y_j) + t\alpha(y_j)))}}{\frac{1}{n} \sum_{k=1}^n e^{-\varepsilon^{-1}(c(x_i, y_k) - (\hat{g}_\varepsilon(y_k) + t\alpha(y_k)))}}$$

Then it is straightforward to verify that  $\psi$  is a convex function with minimum attained at  $t = 0$  such that

$$\psi''(t) = \frac{1}{\varepsilon} \hat{\mu} \left( \text{Var}_{\frac{1}{n} q_t(x, \cdot)}(\alpha) \right).$$

Note that,

$$q_t(x_i, y_j) \geq e^{-2t\varepsilon^{-1}\|\alpha\|_{L^\infty(\hat{\nu})}} q_0(x_i, y_j) = e^{-2t\varepsilon^{-1}\|\alpha\|_{L^\infty(\hat{\nu})}} \hat{p}_\varepsilon(x_i, y_j).$$

Therefore,

$$\begin{aligned} \psi''(t) &= \frac{1}{2\varepsilon} \hat{\mu} \left( \mathbb{E}_{y, y' \sim \frac{1}{n} q_t(x, \cdot)} [(\alpha(y) - \alpha(y'))^2] \right) \\ &\geq \frac{e^{-4t\varepsilon^{-1}\|\alpha\|_{L^\infty(\hat{\nu})}}}{2\varepsilon} \hat{\mu} \left( \mathbb{E}_{y, y' \sim \frac{1}{n} \hat{p}_\varepsilon(x, \cdot)} [(\alpha(y) - \alpha(y'))^2] \right), \end{aligned}$$

It follows that, for  $t > 0$ ,

$$t^2 \frac{e^{-4t\varepsilon^{-1}\|\alpha\|_{L^\infty(\hat{\nu})}}}{4\varepsilon} \hat{\mu}(\mathbb{E}_{y,y' \sim \frac{1}{n}\hat{p}_\varepsilon(x,\cdot)}[(\alpha(y) - \alpha(y'))^2]) \leq \psi(t) - \psi(0) \leq \psi(1) - \psi(0),$$

where the first inequality is strong convexity for  $\psi$  on the interval  $(-\infty, t]$ , and the second inequality follows because  $\psi$  is a convex function with minimizer at  $t = 0$ . If  $\|\alpha\|_{L^\infty(\hat{\nu})} = 0$ , then the inequality is trivially true. Otherwise, we may set  $t = \varepsilon/4\|\alpha\|_{L^\infty(\hat{\nu})}$  to yield

$$\frac{\varepsilon}{\|\alpha\|_{L^\infty(\hat{\nu})}^2} \mathbb{E}_{y,y' \sim \frac{1}{n}\hat{p}_\varepsilon(x,\cdot)}[(\alpha(y) - \alpha(y'))^2] \lesssim \psi(1) - \psi(0).$$

It thus suffices to show that

$$\hat{\mu}(\mathbb{E}_{y,y' \sim \frac{1}{n}\hat{p}_\varepsilon(x,\cdot)}[(\alpha(y) - \alpha(y'))^2]) \gtrsim \left(\frac{\varepsilon}{L}\right)^{d_\nu+2} \hat{\nu}(\alpha^2), \quad (6.3)$$

Observe that

$$\hat{\mu}(\mathbb{E}_{y,y' \sim \frac{1}{n}\hat{p}_\varepsilon(x,\cdot)}[(\alpha(y) - \alpha(y'))^2]) = \frac{1}{n^2} \sum_{j,k=1}^n \left( \frac{1}{n} \sum_{i=1}^n \hat{p}_\varepsilon(x_i, y_j) \hat{p}_\varepsilon(x_i, y_k) \right) (\alpha(y_j) - \alpha(y_k))^2.$$

By Proposition 15,

$$\hat{p}_\varepsilon(x_i, y_j) \geq e^{-2\frac{L}{\varepsilon}\|y_j - y_k\|} \hat{p}_\varepsilon(x_i, y_k), \quad \forall j, k \in [n].$$

Hence

$$\begin{aligned} & \frac{1}{n^2} \sum_{j,k=1}^n \left( \frac{1}{n} \sum_{i=1}^n \hat{p}_\varepsilon(x_i, y_j) \hat{p}_\varepsilon(x_i, y_k) \right) (\alpha(y_j) - \alpha(y_k))^2 \geq \\ & \frac{1}{n^2} \sum_{j,k=1}^n e^{-2\frac{L}{\varepsilon}\|y_j - y_k\|} \left( \frac{1}{n} \sum_{i=1}^n \hat{p}_\varepsilon(x_i, y_k) \right)^2 (\alpha(y_j) - \alpha(y_k))^2 \\ & \geq \frac{1}{n^2} \sum_{j,k=1}^n e^{-2\frac{L}{\varepsilon}\|y_j - y_k\|} \left( \frac{1}{n} \sum_{i=1}^n \hat{p}_\varepsilon(x_i, y_k) \right)^2 (\alpha(y_j) - \alpha(y_k))^2 \\ & = \frac{1}{n^2} \sum_{j,k=1}^n e^{-2\frac{L}{\varepsilon}\|y_j - y_k\|} (\alpha(y_j) - \alpha(y_k))^2. \end{aligned}$$

Now, form the RGG with threshold  $\delta = \frac{\varepsilon}{L}$ , so that  $y_j \sim y_k$  if and only if  $\|y_j - y_k\| \leq \frac{\varepsilon}{L}$ . Then

$$\hat{\mu}(\mathbb{E}_{y,y' \sim \frac{1}{n}\hat{p}_\varepsilon(x,\cdot)}[(\alpha(y) - \alpha(y'))^2]) \gtrsim \frac{1}{n^2} \sum_{j \sim k} (\alpha(y_j) - \alpha(y_k))^2.$$

Applying Theorem 36 from Appendix A.2, we arrive at (6.3), and so the result follows.  $\square$



## 6.2 Reduction from $f$ dual potentials to $g$ dual potentials

In this section, we will prove the following Lemma, which shows that although the  $f$  dual potentials are defined on the support of  $\mu$  – a potentially higher dimensional set than the support of  $\nu$  – their convergence can be controlled by the convergence of the  $g$  dual potentials while only incurring MID scaling factors.

**Lemma 24** (Reduction from  $f$  dual potentials to  $g$  dual potentials). *We have*

$$\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] \lesssim \left(\frac{L}{\varepsilon}\right)^{3d_\nu} \left(\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] + \frac{1 + \varepsilon^2}{n}\right),$$

and similarly

$$\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2] \lesssim \left(\frac{L}{\varepsilon}\right)^{3d_\nu} \left(\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] + \frac{1 + \varepsilon^2}{n}\right).$$

To prove Lemma 24, as well as for the proofs in the following sections, the following helper lemmas are convenient (proven in Appendix B.2).

**Lemma 25** (Mass of empirical balls). *With probability at least  $1 - \frac{1}{n}e^{-10/\varepsilon}$ ,*

$$\inf_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L})) \gtrsim \left(\frac{\varepsilon}{L}\right)^{d_\nu}.$$

**Lemma 26** (Uniform control on entropic densities). *We have the uniform bound,*

$$\|p_\varepsilon\|_{L^\infty(\mu \otimes \nu)} \lesssim \left(\frac{L}{\varepsilon}\right)^{d_\nu}.$$

And, with probability at least  $1 - \frac{1}{n}e^{-10/\varepsilon}$ ,

$$\|\hat{p}_\varepsilon\|_{L^\infty(\mu \otimes \nu)} \lesssim \left(\frac{L}{\varepsilon}\right)^{d_\nu}.$$

*Proof of Lemma 24.* Let

$$\tilde{f}_\varepsilon(x) := -\varepsilon \log \left( \frac{1}{n} \sum_{j=1}^n e^{-\frac{1}{\varepsilon}(c(x, y_j) - g_\varepsilon(y_j))} \right).$$

Note that, for all  $x \in \text{supp } \mu$ ,

$$(\hat{f}_\varepsilon(x) - f_\varepsilon(x))^2 \lesssim (\hat{f}_\varepsilon(x) - \tilde{f}_\varepsilon(x))^2 + (\tilde{f}_\varepsilon(x) - f_\varepsilon(x))^2,$$

and we'll begin by studying each of these terms separately.

For the first term, put  $\alpha := g_\varepsilon - \hat{g}_\varepsilon$ , and let  $\varphi: [0, 1] \rightarrow \mathbb{R}$  be defined as

$$\varphi(t) := -\varepsilon \log \left( \frac{1}{n} \sum_{j=1}^n e^{-\varepsilon^{-1}(c(x, y_j) - \hat{g}_\varepsilon(y_j) - t\alpha(y_j))} \right),$$

and notice  $\varphi(0) = \hat{f}_\varepsilon(x)$  while  $\varphi(1) = \tilde{f}_\varepsilon(x)$ . For  $y \in \text{supp}(\nu)$ , put

$$q_t(x, y) := \frac{e^{-\varepsilon^{-1}(c(x, y) - \hat{g}_\varepsilon(y) - t\alpha(y))}}{\frac{1}{n} \sum_{k=1}^n e^{-\varepsilon^{-1}(c(x, y_k) - \hat{g}_\varepsilon(y_k) - t\alpha(y_k))}}$$

Then

$$|\varphi'(t)| = \left| \frac{1}{n} \sum_{j=1}^n \alpha(y_j) q_t(x, y_j) \right| \leq \|\alpha\|_{L^2(\hat{\nu})} \left( \frac{1}{n} \sum_{j=1}^n q_t(x, y_j)^2 \right)^{1/2}.$$

Using Proposition 15, we find that for any  $y, y' \in \text{supp}(\nu)$  and  $t \in [0, 1]$ ,

$$\begin{aligned} |\log q_t(x, y) - \log q_t(x, y')| &= \frac{1}{\varepsilon} |c(x, y) - \hat{g}_\varepsilon(y) - t\alpha(y) - (c(x, y') - \hat{g}_\varepsilon(y') - t\alpha(y'))| \\ &\leq \frac{2L}{\varepsilon} \|y - y'\|. \end{aligned}$$

Therefore, for any  $j \in [n]$ ,

$$1 = \frac{1}{n} \sum_{k=1}^n q_t(x, y_k) \gtrsim q_t(x, y_j) \hat{\nu}(B(y_j, \frac{\varepsilon}{L})) \geq q_t(x, y_j) \inf_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L})).$$

So that

$$\begin{aligned} |\hat{f}_\varepsilon(x) - \tilde{f}_\varepsilon(x)| &\leq \int_0^1 |\varphi'(t)| dt \leq \|\alpha\|_{L^2(\hat{\nu})} \int_0^1 \left( \frac{1}{n} \sum_{j=1}^n q_t(x, y_j)^2 \right)^{1/2} dt \\ &\lesssim \|\alpha\|_{L^2(\hat{\nu})} \sup_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L}))^{-1/2} \int_0^1 \left( \frac{1}{n} \sum_{j=1}^n q_t(x, y_j) \right)^{1/2} dt \\ &= \|\alpha\|_{L^2(\hat{\nu})} \sup_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L}))^{-1/2}. \end{aligned} \tag{6.4}$$

Now consider the  $(\tilde{f}_\varepsilon(x) - f_\varepsilon(x))^2$  term. Note that

$$|\tilde{f}_\varepsilon(x) - f_\varepsilon(x)| = \left| \varepsilon \log \left( \frac{1}{n} \sum_{j=1}^n p_\varepsilon(x, y_j) \right) \right|.$$

Since

$$1 = \int p_\varepsilon(x, y) d\nu(y),$$

there must exist some  $y' \in \text{supp}(\nu)$  such that  $p_\varepsilon(x, y') \geq 1$ . By Proposition 15, this implies

$$\frac{1}{n} \sum_{j=1}^n p_\varepsilon(x, y_j) \geq \frac{1}{n} \sum_{j=1}^n \frac{p_\varepsilon(x, y_j)}{p_\varepsilon(x, y')} \geq \frac{1}{n} \sum_{j=1}^n e^{-\frac{2L}{\varepsilon} \|y_j - y'\|} \gtrsim \hat{\nu}(B(y', \frac{\varepsilon}{L})).$$

Using Lipschitz-ness of log,

$$|\tilde{f}_\varepsilon(x) - f_\varepsilon(x)| \lesssim \varepsilon \hat{\nu}(B(y', \frac{\varepsilon}{L}))^{-1} \left| \frac{1}{n} \sum_{j=1}^n p_\varepsilon(x, y_j) - 1 \right|. \tag{6.5}$$

Now, let  $\mathcal{E}$  denote the event of Lemma 25. Observe that by the pointwise control in Proposition 14 and since  $\mathbb{P}[\mathcal{E}^c] \leq \frac{1}{n}$ ,

$$\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] \lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \frac{1}{n}.$$

Combining Equations (6.4) and (6.5) yields

$$\begin{aligned} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] &\lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \frac{1}{n} \\ &\lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}](\|\hat{f}_\varepsilon - \tilde{f}_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\tilde{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2)] + \frac{1}{n} \\ &\lesssim \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}\left[\mathbb{1}[\mathcal{E}]\left\{\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2 + \frac{\varepsilon^2}{n} \sum_{i=1}^n \left(\frac{1}{n} \sum_{j=1}^n p_\varepsilon(x_i, y_j) - 1\right)^2\right\}\right] + \frac{1}{n} \\ &\leq \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}\left[\mathbb{1}[\mathcal{E}]\left\{\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2 + \frac{\varepsilon^2}{n} \|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2\right\}\right] + \frac{1}{n} \\ &\lesssim \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \left\{\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] + \frac{\varepsilon^2}{n} \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2\right\} + \frac{1}{n}, \end{aligned}$$

where the final inequality follows from  $\mathbb{1}[\mathcal{E}] \leq 1$  and Equation (5.2). By Lemma 16 and Proposition 43 on the covering numbers of  $N$  (proved in Appendix A.3), we conclude the first inequality. The second follows in the same manner.  $\square$

### 6.3 MID scaling for the bias and $g$ dual potentials in empirical norm

In this section, we use our quadratic growth inequality from section 6.1 and our bound on the  $f$  dual potentials in terms of the  $g$  dual potentials from section 6.2 to prove the following result, on convergence of the  $g$  dual potentials in empirical norm.

**Lemma 27** (Convergence of  $g$  dual potentials in empirical norm). *We have*

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{6d_\nu+4} \cdot \frac{1}{n}.$$

By the previous section, this implies convergence of the  $f$  dual potentials in both empirical and population norms, and therefore yields all parts of Theorem 9 other than the population norm convergence of the  $g$  dual potentials, which is proved in the next section. As a byproduct of our proof, we will also obtain Theorem 12, on MID scaling with fast rates for the bias.

Notice that a  $1/\sqrt{n}$  rate follows by combining our result on quadratic growth for the empirical dual, Lemma 23, with our prior result on the bias, Lemma 17. To obtain a faster rate, we proceed by a self-bounding argument.

We first bound the bias  $\mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)]$  in terms of  $\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2$  and  $\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2$  plus a  $1/n$  term arising from the squared norm of  $\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)$ .

**Lemma 28.** *For all  $a > 0$ ,*

$$\mathbb{E}[\hat{\Phi}_\varepsilon(\hat{\mu}, \hat{\nu}) - \hat{\Phi}_\varepsilon(\mu, \nu)] \lesssim \frac{1}{an} \left(\frac{L}{\varepsilon}\right)^{d_\nu} + a\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2].$$

Using our result on the quadratic growth of  $\hat{\Phi}_\varepsilon$ , Lemma 23, the previous step then implies a bound for  $\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2$  in terms of itself,  $1/n$ , and  $\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2$ .

**Lemma 29.** *For all  $a > 0$ ,*

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] \lesssim \frac{1}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{d_\nu+2} \left\{ \frac{1}{an} \left(\frac{L}{\varepsilon}\right)^{d_\nu} + a \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] \right\} + \frac{1}{n}.$$

Combining this inequality with Lemma 24 from section 6.2 and taking the free parameter  $a$  sufficiently small, we can re-arrange and arrive at Lemma 27. Plugging our bounds on the dual potentials into Lemma 28 then yields Theorem 12. Let's give these proofs first, before we turn to the proofs of Lemma 28 and Lemma 29.

*Proof of Lemma 27.* By Lemma 29 and Lemma 24, for all  $a > 0$ ,

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] \lesssim \frac{1}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{d_\nu+2} \left\{ \frac{1}{an} \left(\frac{L}{\varepsilon}\right)^{d_\nu} + a \left(\frac{L}{\varepsilon}\right)^{3d_\nu} (\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] + \frac{1 + \varepsilon^2}{n}) \right\} + \frac{1}{n}.$$

If we take  $a = C\varepsilon(\frac{\varepsilon}{L})^{4d_\nu+2}$  for a sufficiently small constant  $C$ , we can re-arrange to yield

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] \lesssim (1 + \varepsilon^2 + \frac{1}{\varepsilon^2}) \cdot \left(\frac{L}{\varepsilon}\right)^{6d_\nu+4} \cdot \frac{1}{n} \lesssim (\varepsilon^2 + \frac{1}{\varepsilon^2}) \cdot \left(\frac{L}{\varepsilon}\right)^{6d_\nu+4} \cdot \frac{1}{n}.$$

□

*Proof of Theorem 12.* First notice that  $\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu})] - S_\varepsilon(\mu, \nu) = \mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] \geq 0$ , so that

$$|\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu})] - S_\varepsilon(\mu, \nu)| = \mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)].$$

Invoking Lemma 28 with  $a = 1$ , and applying Lemma 24 plus Lemma 27, we obtain

$$\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)] \lesssim (1 + \varepsilon^2 + \frac{1}{\varepsilon^2}) \cdot \left(\frac{L}{\varepsilon}\right)^{9d_\nu+4} \cdot \frac{1}{n} \leq (\varepsilon^2 + \frac{1}{\varepsilon^2}) \cdot \left(\frac{L}{\varepsilon}\right)^{9d_\nu+4} \cdot \frac{1}{n}.$$

and whence the result. □

The proofs of Lemma 28 and Lemma 29 follow.

*Proof of Lemma 28.* By Proposition 13, for all  $a > 0$ ,

$$\begin{aligned} \mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)] &= \mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)] \\ &= \mathbb{E}[\hat{\Phi}_\varepsilon(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] \\ &\leq \mathbb{E}[\langle \nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}] \\ &\leq \frac{1}{a} \mathbb{E}[\|\nabla \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\hat{\nu})}^2] + a \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2]. \end{aligned}$$

By Equation (5.2) and Lemma 16, we have

$$\mathbb{E}[S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)] \lesssim \frac{1}{an} \mathcal{N}(N, \frac{\varepsilon}{L}) + a \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2].$$

Applying Proposition 43 on the covering numbers of  $N$  (proved in Appendix A.3), we may conclude. □

*Proof of Lemma 29.* We begin by re-centering  $g_\varepsilon$  so that we can apply Lemma 23:

$$\begin{aligned} \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] &\leq \mathbb{E}[\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] + \mathbb{E}[(\hat{\nu}(g_\varepsilon))^2] \\ &\lesssim \mathbb{E}[\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] + \frac{1}{n}, \end{aligned} \quad (6.6)$$

where the last step follows by our convention that  $\nu(g_\varepsilon) = 0$  and the pointwise control on  $g_\varepsilon$  from Proposition 14. Let  $\mathcal{E}$  denote the event described in Lemma 23. Applying Proposition 14 once more we see that

$$\begin{aligned} \mathbb{E}[\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] &= \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] + \mathbb{E}[\mathbb{1}[\mathcal{E}^c]] \\ &\lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] + \frac{1}{n} \\ &\lesssim \mathbb{E}\left[\mathbb{1}[\mathcal{E}]\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^\infty(\hat{\nu})}^2\right. \\ &\quad \left. \times \frac{1}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{d_\nu+2} (\Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(\hat{g}_\varepsilon) - \Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(g_\varepsilon))\right] + \frac{1}{n}. \end{aligned}$$

Using  $\mathbb{1}[\mathcal{E}] \leq 1$  and Proposition 14 once more, we find

$$\mathbb{E}[\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] \lesssim \frac{1}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{d_\nu+2} \mathbb{E}[\Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(\hat{g}_\varepsilon) - \Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(g_\varepsilon)].$$

By Proposition 13 on marginal rounding

$$\Psi_\varepsilon^{\hat{\mu}\hat{\nu}}(g_\varepsilon) \geq \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon).$$

Hence

$$\mathbb{E}[\|\hat{g}_\varepsilon - (g_\varepsilon - \hat{\nu}(g_\varepsilon))\|_{L^2(\hat{\nu})}^2] \lesssim \frac{1}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{d_\nu+2} \mathbb{E}[\hat{\Phi}(f_\varepsilon, \hat{g}_\varepsilon) - \hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)].$$

Since  $\mathbb{E}[\hat{\Phi}_\varepsilon(f_\varepsilon, g_\varepsilon)] = \Phi_\varepsilon(f_\varepsilon, g_\varepsilon)$ , the result follows from Lemma 28.  $\square$

## 6.4 Fast rates for $g$ -dual potentials in population norms

In this section, we establish the remaining part of Theorem 9 by proving the following Lemma.

**Lemma 30** (Fast population norm convergence of  $g$ -dual potentials). *We have*

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{13d_\nu+8} \cdot \frac{1}{n}.$$

Lemma 30 is proved with another self-bounding argument that we outline here, ignoring  $\varepsilon/L$  dependence for clarity. Note that we will frequently use the canonical extension of  $\hat{g}_\varepsilon$  to consider  $(\hat{f}_\varepsilon, \hat{g}_\varepsilon)$  as a dual variable pair for the semi-empirical problem between  $\hat{\mu}$  and  $\nu$ . In particular, notation such as  $\nu(\hat{g}_\varepsilon)$  is not referring to integration over any part of the sample  $y_1, \dots, y_n \sim \nu^{\otimes n}$ , but rather over an *independent* draw from  $\nu$ .

1. Lemma 31: Use a Poincaré inequality on  $\nu$  (which follows from our assumptions and is described in Proposition 40 in Appendix A.3) to control  $\|\hat{g}_\varepsilon - g_\varepsilon - \nu(\hat{g}_\varepsilon)\|_{L^2(\nu)}^2$  in terms of entropic OT maps defined on  $N$

2. Lemma 32: Use the same analysis as in the proof of the empirical norm convergence of the entropic OT maps (section 5.3) to control the previous term in terms of  $1/n$ , the semi-empirical dual objective  $\Phi_\varepsilon^{\hat{\mu}\nu}$ , and  $\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2$ , with a free parameter from Young's inequality.
3. Lemma 33: Control the semi-empirical objective term with concavity to yield a bound in terms of  $1/n$  and  $\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2$ , with a free parameter from Young's inequality.
4. Lemma 34: Bound the shift  $\mathbb{E}[\nu(\hat{g}_\varepsilon)^2]$  by  $1/n$  using the semi-discrete objective once more
5. Use the free parameters from Young's inequality to conclude a  $1/n$  bound on  $\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2$ .

The formal statements of the lemmas outlined above are given below.

**Lemma 31.** *We have*

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon - \nu(\hat{g}_\varepsilon)\|_{L^2(\nu)}^2] \lesssim \mathbb{E}\left[\left\|\mathbb{E}_{\pi_\varepsilon}[\nabla_y c(x, y) | y] - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y)\right\|_{L^2(\nu)}^2\right].$$

**Lemma 32.** *For all  $a > 0$ , we have*

$$\begin{aligned} \mathbb{E}\left[\left\|\mathbb{E}_{\pi_\varepsilon}[\nabla_y c(x, y) | y] - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y)\right\|_{L^2(\nu)}^2\right] &\lesssim \frac{L^2}{\varepsilon} \mathbb{E}[\Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)] \\ &\quad + a \frac{1}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{2d_\nu+2} \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] \\ &\quad + L \left(\frac{a}{\varepsilon^2} + \frac{1}{a}\right) \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \left(\frac{L}{\varepsilon}\right)^{11d_\nu+5} \cdot \frac{1}{n}. \end{aligned}$$

**Lemma 33.** *For all  $a > 0$ , we have*

$$\mathbb{E}[\Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)] \lesssim \frac{1}{a} \left(\frac{L}{\varepsilon}\right)^{d_\nu} \cdot \frac{1}{n} + a \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2].$$

**Lemma 34.** *We have*

$$\mathbb{E}[\nu(\hat{g}_\varepsilon)^2] \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \left(\frac{L}{\varepsilon}\right)^{11d_\nu+4} \cdot \frac{1}{n}$$

*Proof of Lemma 30.* Note that

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] \lesssim \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon - \nu(\hat{g}_\varepsilon)\|_{L^2(\nu)}^2] + \mathbb{E}[\nu(\hat{g}_\varepsilon)^2].$$

We can thus apply each of the above lemmas to yield, for  $a, a' > 0$ ,

$$\begin{aligned} \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] &\lesssim \frac{L}{a} \left(\frac{L}{\varepsilon}\right)^{d_\nu+1} + a \frac{L^2}{\varepsilon} \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] \\ &\quad + \frac{a'}{\varepsilon} \left(\frac{L}{\varepsilon}\right)^{2d_\nu+2} \mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] + L \left(\frac{a'}{\varepsilon^2} + \frac{1}{a'}\right) \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \left(\frac{L}{\varepsilon}\right)^{11d_\nu+5} \cdot \frac{1}{n} \\ &\quad + \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \left(\frac{L}{\varepsilon}\right)^{11d_\nu+4} \cdot \frac{1}{n}. \end{aligned}$$

Taking  $a = C\varepsilon/L^2$  and  $a' = C\varepsilon(\varepsilon/L)^{2d_\nu+2}$  for  $C$  a sufficiently small constant and re-arranging yields

$$\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] \lesssim \left\{ L^2 \left( \frac{L}{\varepsilon} \right)^{d_\nu+2} + \left( \varepsilon^2 + \frac{1}{\varepsilon^2} \right) \left( \frac{L}{\varepsilon} \right)^{13d_\nu+8} + \left( \varepsilon^2 + \frac{1}{\varepsilon^2} \right) \left( \frac{L}{\varepsilon} \right)^{11d_\nu+4} \right\} \cdot \frac{1}{n}.$$

Using  $\varepsilon/L \lesssim 1$  yields the result.  $\square$

*Proof of Lemma 31.* Note that the term on the LHS is precisely  $\text{Var}_\nu(\hat{g}_\varepsilon - g_\varepsilon)$  since we have specified  $g_\varepsilon$  to be such that  $\nu(g_\varepsilon) = 0$ . We will thus apply the Poincaré inequality on  $\nu$  from Proposition 40 to bound this term; this is legitimate because  $\hat{g}_\varepsilon$  and  $g_\varepsilon$  are each Lipschitz with respect to the ambient Euclidean norm and so are Lipschitz with respect to  $(N, h)$ , since embedded manifold distances are always at least as large as extrinsic distances. We obtain

$$\|\hat{g}_\varepsilon - g_\varepsilon - \nu(\hat{g}_\varepsilon)\|_{L^2(\nu)}^2 \lesssim \int \|\nabla_N(\hat{g}_\varepsilon - g_\varepsilon)\|_h^2 d\nu(y).$$

Since  $N$  is an embedded manifold, the manifold norm of the manifold gradient is always upper bounded by the Euclidean norm of the Euclidean gradient (for the reader's convenience, this fact is formally stated and proved as Proposition 41 in Appendix A.3) so that

$$\|\hat{g}_\varepsilon - g_\varepsilon - \nu(\hat{g}_\varepsilon)\|_{L^2(\nu)}^2 \lesssim \|\nabla \hat{g}_\varepsilon - \nabla g_\varepsilon\|_{L^2(\nu)}^2.$$

To calculate  $\nabla g_\varepsilon$ , consider the marginal constraint

$$e^{-\frac{1}{\varepsilon}g_\varepsilon(y)} = \int e^{-\frac{1}{\varepsilon}(c(x,y)-f_\varepsilon(x))} d\mu(x).$$

Differentiating and re-arranging, we find that

$$\nabla g_\varepsilon(y) = \mathbb{E}_{\pi_\varepsilon}[\nabla_y c(x, y) | y].$$

Calculating in a similar fashion with the extended dual potential  $\hat{g}_\varepsilon$  we find

$$\nabla \hat{g}_\varepsilon(y) = \mathbb{E}_{\hat{\pi}_\varepsilon}[\nabla_y c(x, y) | y] = \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y).$$

The result follows.  $\square$

*Proof of Lemma 32.* For all  $y$  in  $\text{supp}(\nu)$ , put

$$\tilde{g}_\varepsilon(y) := -\varepsilon \log \left( \frac{1}{n} \sum_{i=1}^n e^{-\varepsilon^{-1}(c(x_i, y) - f_\varepsilon(x_i))} \right),$$

and let

$$\tilde{p}_\varepsilon(x, y) := e^{-\varepsilon^{-1}(c(x, y) - f_\varepsilon(x) - \tilde{g}_\varepsilon(y))}.$$

Then by Young's inequality,

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \mathbb{E}_{\pi_\varepsilon} [\nabla_y c(x, y) \mid y] - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right] \lesssim \\
& \mathbb{E} \left[ \left\| \mathbb{E}_{\pi_\varepsilon} [\nabla_y c(x, y) \mid y] - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) p_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right] \\
& + \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) p_\varepsilon(x_i, y) - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \tilde{p}_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right] \\
& + \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \tilde{p}_\varepsilon(x_i, y) - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right].
\end{aligned}$$

Using the same argument as in the proof of Lemma 20 just with the role of  $\mu$  and  $\nu$  swapped, the first two terms can each be bounded as  $\lesssim L^2 \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2 / n$ . By Lemma 16 and Proposition 43, we can bound  $\|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2 \lesssim (L/\varepsilon)^{d_\nu}$  and so arrive at

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \mathbb{E}_{\pi_\varepsilon} [\nabla_y c(x, y) \mid y] - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right] \lesssim \\
& \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \tilde{p}_\varepsilon(x_i, y) - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right] + L^2 \left( \frac{L}{\varepsilon} \right)^{d_\nu} \cdot \frac{1}{n}.
\end{aligned}$$

We proceed as in the proof of Lemma 19, by first applying triangle inequality and then Pinsker's. This yields

$$\begin{aligned}
& \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \tilde{p}_\varepsilon(x_i, y) - \frac{1}{n} \sum_{i=1}^n \nabla_y c(x_i, y) \hat{p}_\varepsilon(x_i, y) \right\|_{L^2(\nu)}^2 \right] \\
& \leq L^2 \mathbb{E} \left[ \left\| \frac{1}{n} \sum_{i=1}^n |\tilde{p}_\varepsilon(x_i, y) - \hat{p}_\varepsilon(x_i, y)| \right\|_{L^2(\nu)}^2 \right] \\
& \lesssim L^2 \mathbb{E} \left[ \int \text{KL} \left( \frac{1}{n} \hat{p}_\varepsilon(\cdot, y) \parallel \frac{1}{n} \tilde{p}_\varepsilon(\cdot, y) \right) d\nu(y) \right] \\
& = \frac{L^2}{\varepsilon} \mathbb{E} \left[ \int \left( \frac{1}{n} \sum_{i=1}^n \hat{p}_\varepsilon(x_i, y) (\hat{f}_\varepsilon(x_i) - f_\varepsilon(x_i) + \hat{g}_\varepsilon(y) - \tilde{g}_\varepsilon(y)) \right) d\nu(y) \right] \\
& = \frac{L^2}{\varepsilon} \mathbb{E} \left[ \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, \tilde{g}_\varepsilon) + \left\langle \int \hat{p}_\varepsilon(\cdot, y) d\nu(y) - 1, f_\varepsilon - f_\varepsilon \right\rangle_{L^2(\hat{\mu})} \right] \\
& \leq \frac{L^2}{\varepsilon} \mathbb{E} \left[ \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon) + \left\langle \int \hat{p}_\varepsilon(\cdot, y) d\nu(y) - 1, \hat{f}_\varepsilon - f_\varepsilon \right\rangle_{L^2(\hat{\mu})} \right],
\end{aligned}$$

where the second equality follows because  $(\hat{\mu} \otimes \nu)(\hat{p}_\varepsilon) = (\hat{\mu} \otimes \nu)(\tilde{p}_\varepsilon) = 1$ , and the final inequality is via Proposition 13. To conclude the argument, we apply Young's inequality



to the inner product term to obtain, for all  $a > 0$ ,

$$\begin{aligned}
\mathbb{E}\left[\left\langle \int \hat{p}_\varepsilon(\cdot, y) d\nu(y) - 1, \hat{f}_\varepsilon - f_\varepsilon \right\rangle_{L^2(\hat{\mu})}\right] &\leq a\mathbb{E}\left[\left\|\int (\hat{p}_\varepsilon(\cdot, y) - 1) d\nu(y)\right\|_{L^2(\hat{\mu})}^2\right] \\
&\quad + \frac{1}{a}\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\nu})}^2] \\
&= a\mathbb{E}\left[\left\|\int (\hat{p}_\varepsilon(\cdot, y) - p_\varepsilon(\cdot, y)) d\nu(y)\right\|_{L^2(\hat{\mu})}^2\right] \\
&\quad + \frac{1}{a}\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\nu})}^2] \\
&\leq a\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\hat{\mu}\otimes\nu)}^2] + \frac{1}{a}\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\nu})}^2].
\end{aligned}$$

By Lemma 35, which is stated and proved in the following section,

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\hat{\mu}\otimes\nu)}^2] \lesssim \frac{1}{\varepsilon^2} \cdot \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] + \frac{1}{n}.$$

Applying Lemma 24 and Lemma 27 we have

$$\mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] \lesssim \left(\varepsilon^2 + \frac{1}{\varepsilon^2}\right) \cdot \left(\frac{L}{\varepsilon}\right)^{9d_\nu+4} \cdot \frac{1}{n}.$$

The result follows by collecting terms.  $\square$

*Proof of Lemma 33.* Proposition 13 on concavity of the dual implies that

$$\mathbb{E}[\Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)] \leq \mathbb{E}[\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}].$$

Observe that because  $(f_\varepsilon, g_\varepsilon)$  attain the  $\nu$ -marginal, the first component of  $\nabla \Phi_\varepsilon^{\hat{\mu}\nu}$  vanishes, implying

$$\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)} = \left\langle 1 - \frac{1}{n} \sum_{i=1}^n p_\varepsilon(x_i, y), \hat{g}_\varepsilon - g_\varepsilon \right\rangle_{L^2(\nu)}.$$

Young's inequality yields, for all  $a > 0$ ,

$$\mathbb{E}[\Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)] \lesssim \frac{1}{a}\mathbb{E}\left[\left\|1 - \frac{1}{n} \sum_{i=1}^n p_\varepsilon(x_i, y)\right\|_{L^2(\nu)}^2\right] + a\mathbb{E}[\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2].$$

The first term can be calculated as in (5.2) to yield

$$\mathbb{E}\left[\left\|1 - \frac{1}{n} \sum_{i=1}^n p_\varepsilon(x_i, y)\right\|_{L^2(\nu)}^2\right] \lesssim \frac{\|p_\varepsilon\|_{L^2(\mu\otimes\nu)}^2}{n}.$$

Finally, applying Lemma 16 and Proposition 43, we obtain the result.  $\square$

*Proof of Lemma 34.* Note that, since  $(\hat{\mu} \otimes \nu)(\hat{p}_\varepsilon) = (\hat{\mu} \otimes \nu)(p_\varepsilon) = 1$  and we make the normalization assumption  $\nu(g_\varepsilon) = 0$ , we have

$$\nu(\hat{g}_\varepsilon) = \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon) + \hat{\mu}(f_\varepsilon - \hat{f}_\varepsilon).$$

By Jensen's inequality, we have that

$$|\nu(\hat{g}_\varepsilon)| \leq |\Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)| + \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}.$$

Observe that the two-sided concavity statements from Proposition 13 imply that

$$\begin{aligned} |\Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon) - \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)| &\leq |\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}| \\ &\quad + |\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}|. \end{aligned}$$

For the first term, Cauchy-Schwarz and Proposition 14 on uniform bounds for the dual potentials imply that

$$|\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}| \lesssim \|\nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\nu)}.$$

For the second term, since  $(\hat{f}_\varepsilon, \hat{g}_\varepsilon)$  attain the  $\hat{\mu}$  marginal, the second component of  $\nabla \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon)$  vanishes, so that

$$\begin{aligned} |\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}| &= |\langle 1 - \int \hat{p}_\varepsilon(x, y) d\nu(y), \hat{f}_\varepsilon - f_\varepsilon \rangle_{L^2(\hat{\mu})}| \\ &\leq \left\| \int \hat{p}_\varepsilon(x, y) d\nu(y) - 1 \right\|_{L^2(\hat{\mu})} \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})} \\ &\lesssim \|\hat{p}_\varepsilon\|_{L^\infty(\hat{\mu} \otimes \nu)} \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}. \end{aligned}$$

Now, let  $\mathcal{E}$  denote the event described in Lemma 26. Then by the pointwise control given in Proposition 14, we have

$$\begin{aligned} \mathbb{E}[\|\hat{p}_\varepsilon\|_{L^\infty(\hat{\mu} \otimes \nu)}^2 \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] &= \mathbb{E}[\mathbb{1}[\mathcal{E}] \|\hat{p}_\varepsilon\|_{L^\infty(\hat{\mu} \otimes \nu)}^2 \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] \\ &\quad + \mathbb{E}[\mathbb{1}[\mathcal{E}^c] \|\hat{p}_\varepsilon\|_{L^\infty(\hat{\mu} \otimes \nu)}^2 \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] \\ &\lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}] \|\hat{p}_\varepsilon\|_{L^\infty(\hat{\mu} \otimes \nu)}^2 \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \mathbb{E}[\mathbb{1}[\mathcal{E}^c]] e^{\frac{10}{\varepsilon}} \\ &\lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}] \|\hat{p}_\varepsilon\|_{L^\infty(\hat{\mu} \otimes \nu)}^2 \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \frac{1}{n} \\ &\lesssim \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\mathbb{1}[\mathcal{E}] \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \frac{1}{n} \\ &\leq \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \frac{1}{n} \end{aligned}$$

We thus conclude that

$$\begin{aligned} \mathbb{E}[\nu(\hat{g}_\varepsilon)^2] &\lesssim \mathbb{E}[\langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}^2 \\ &\quad + \langle \nabla \Phi_\varepsilon^{\hat{\mu}\nu}(\hat{f}_\varepsilon, \hat{g}_\varepsilon), (\hat{f}_\varepsilon - f_\varepsilon, \hat{g}_\varepsilon - g_\varepsilon) \rangle_{L^2(\hat{\mu}) \times L^2(\nu)}^2 + \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] \\ &\lesssim \mathbb{E}[\|\nabla \Phi_\varepsilon^{\hat{\mu}\nu}(f_\varepsilon, g_\varepsilon)\|_{L^2(\hat{\mu}) \times L^2(\nu)}^2 + \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2] + \frac{1}{n}. \end{aligned}$$

The squared gradient term can be calculated as in (5.2) and then bounded with Lemma 16 and Proposition 43, and the latter bounded with Lemma 24 and Lemma 27, to yield the result.  $\square$

## 6.5 Fast rates with MID scaling for maps and densities

In this section, we give the proof of Corollary 10 and Corollary 11. To this end, the following Lemma is convenient.

**Lemma 35.** *We have*

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] \lesssim \frac{1}{\varepsilon^2} \cdot \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] + \frac{1}{n}.$$

And we also have the analogous bounds

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \hat{\nu})}^2] \lesssim \frac{1}{\varepsilon^2} \cdot \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] + \frac{1}{n},$$

as well as,

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\hat{\mu} \otimes \nu)}^2] \lesssim \frac{1}{\varepsilon^2} \cdot \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\hat{\mu})}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] + \frac{1}{n}.$$

With this lemma in hand, the proofs of Corollary 10 and Corollary 11 are a simple application of the results from the previous section.

*Proof of Corollary 10.* To prove this result, let us first assume that  $0 \in \text{supp}(\nu)$  so that for each  $y \in \text{supp}(\nu)$ , we have  $\|y\| \leq R$ . By Young's inequality,

$$\begin{aligned} \mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\mu)}^2] &= \mathbb{E}[\|\hat{T}_\varepsilon - T_\varepsilon\|_{L^2(\mu)}^2] \\ &\lesssim \mathbb{E}\left[\left\|\hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j)\right\|_{L^2(\mu)}^2\right] + \mathbb{E}\left[\left\|\frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j) - T_\varepsilon\right\|_{L^2(\mu)}^2\right]. \end{aligned}$$

The second term is merely a variance, and can be bounded as

$$\begin{aligned} \mathbb{E}\left[\left\|\frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j) - T_\varepsilon\right\|_{L^2(\mu)}^2\right] &= \frac{1}{n} \|y p_\varepsilon(x, y) - T_\varepsilon(x)\|_{L^2(\mu \otimes \nu)}^2 \\ &\lesssim \frac{R^2}{n} \|p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2 \lesssim \frac{R^2}{n} \left(\frac{L}{\varepsilon}\right)^{d_\nu}, \end{aligned}$$

where the final inequality follows by applying Lemma 16 with Proposition 43. We now focus on the remaining expectation, and apply triangle inequality and Jensen's to yield, for any  $x \in \text{supp} \mu$ ,

$$\begin{aligned} \left\|\hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j)\right\|^2 &\lesssim \frac{1}{n} \sum_{j=1}^n \|y_j\|^2 (\hat{p}_\varepsilon(x, y_j) - p_\varepsilon(x, y_j))^2 \\ &\lesssim \frac{R^2}{n} \sum_{j=1}^n (\hat{p}_\varepsilon(x, y_j) - p_\varepsilon(x, y_j))^2. \end{aligned}$$

Taking expectations and applying Lemma 35 yields

$$\begin{aligned} \mathbb{E}\left[\left\|\hat{T}_\varepsilon - \frac{1}{n} \sum_{j=1}^n y_j p_\varepsilon(x, y_j)\right\|_{L^2(\mu)}^2\right] &\lesssim R^2 \mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \hat{\nu})}^2] \\ &\lesssim \frac{R^2}{\varepsilon^2} \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\hat{\nu})}^2] + \frac{R^2}{n}. \end{aligned}$$

The result follows from Lemma 24 and Lemma 27. For the general case where 0 may not be in  $\text{supp}(\nu)$ , we can perform the above argument with suitably offset  $y$ .  $\square$

*Proof of Corollary 11.* By Lemma 35,

$$\mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] \lesssim \frac{1}{\varepsilon^2} \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2] + \frac{1}{n}.$$

The result follows by applying Lemma 24 and Lemma 27 to control the  $\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2$  term, and Lemma 30 to control the  $\|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2$  term.  $\square$

**Acknowledgements.** The author gratefully acknowledges partial support from NSF awards IIS-1838071 and CCF-2106377, and is indebted to Enric Boix-Adsera, Sinho Chewi, Simone Di Marino, Augusto Gerolin, Dheeraj Nagaraj, Jonathan Niles-Weed, Aram-Alexandre Pooladian, Philippe Rigollet, George Stepaniants, and Stephanie Wu for enlightening conversations. Special thanks go to Sinho Chewi, Aram-Alexandre Pooladian, and Stephanie Wu for helpful comments on earlier drafts.

## A Background on embedded manifolds and RGGs

### A.1 Preliminaries on embedded manifolds

For a comprehensive introduction to Riemannian manifolds, we refer the reader to the books [DC16, Lee18]. In this section, we will establish notation while reviewing the definitions of some necessary geometric quantities.

To this end, suppose  $(N_0, h_0)$  is a compact, connected Riemannian manifold. Recall that, in coordinates, the canonical Riemannian volume has density

$$d \text{vol}_{N_0}(y) = \sqrt{\det h_0(y)}.$$

Given  $p \in N_0$  and  $u, v \in T_p N_0$ , we shall write  $\langle u, v \rangle_{h_0} := h_0(u, v)$ , and similarly  $\|u\|_{h_0}^2 := \langle u, u \rangle_{h_0}$ . Recall that for a point  $p \in N_0$ , the exponential map  $\exp_p: T_p N_0 \rightarrow N_0$  is defined as  $\exp_p(v) := \gamma_v(1)$  where  $\gamma_v$  is the constant-speed geodesic such that  $\gamma_v(0) = p$  and  $\dot{\gamma}_v(0) = v$ . For a point  $p \in N_0$ , the injectivity radius  $\text{inj}(p)$  is defined as

$$\text{inj}(p) := \sup \{R \geq 0 \mid \exp_p: B(0, R) \subset T_p N_0 \rightarrow N_0 \text{ is a diffeomorphism}\}.$$

The injectivity radius of  $N_0$  is then defined as

$$\text{inj}(N_0) := \inf_{p \in N_0} \text{inj}(p).$$

It is an elementary fact that when  $N_0$  is compact,  $\text{inj}(N_0) > 0$ , see e.g. [Lee18, Lemma 6.16].

The Riemannian curvature tensor  $\text{Rm}$  maps from 4-tuples of smooth vector fields to smooth functions  $\text{Rm}: \mathcal{X}(N_0)^4 \rightarrow C^\infty(N_0)$  and is defined as, for smooth vector fields  $W, X, Y, Z \in \mathcal{X}(N_0)$ ,

$$\text{Rm}(X, Y, Z, W) := \langle \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z, W \rangle_{h_0}.$$

Given  $p \in M$  and  $u, v \in T_p N_0$  linearly independent, the sectional curvature of  $u, v$  is defined as

$$\sec(u, v) := \frac{\text{Rm}(u, v, v, u)}{\|u\|_{h_0} \|v\|_{h_0} - \langle u, v \rangle_{h_0}}.$$

The sectional curvatures describe the geometry of the embedded surface generated by  $u, v$ , and when  $u, v$  range over  $T_p N_0$ , determine the full Riemannian curvature tensor  $\text{Rm}$ . It is, again, an elementary fact that when  $N_0$  is compact, the sectional curvatures are uniformly bounded in absolute value, see [BO69, Section 9.3].

Given a smooth manifold  $N_0$ , we say that  $N_0$  is embedded in  $\mathbb{R}^D$  if there is a smooth injection  $\iota: N_0 \hookrightarrow \mathbb{R}^D$  which is a homeomorphism onto its image and such that for all  $p \in N_0$ , the differential  $d\iota_p: T_p N_0 \rightarrow T_{\iota(p)} \mathbb{R}^D$  has trivial kernel. In such a case, we identify  $N_0$  with its image under  $\iota$ , and so write  $N_0 \subseteq \mathbb{R}^d$ . When  $N_0$  is endowed with a Riemannian structure  $(N_0, h_0)$ , we say that  $(N_0, h_0)$  is an embedded Riemannian manifold in  $\mathbb{R}^d$  if  $N_0$  is embedded in  $\mathbb{R}^d$  and  $h_0(u, v) = \langle d\iota_p(u), d\iota_p(v) \rangle$  for all  $u, v \in T_p N_0$  and the inner product is the standard Euclidean one.

The main quantitative property of subsets in  $\mathbb{R}^D$  we need is called the reach. To define this quantity, suppose  $S \subseteq \mathbb{R}^D$ , and for all  $x \in \mathbb{R}^D$  let  $d(x, S) := \inf_{y \in S} \|x - y\|$ . Then  $\text{reach}(S)$  is the supremum over all  $\varepsilon \geq 0$  such that for all  $x \in \mathbb{R}^D$  for which  $d(x, S) \leq \varepsilon$ , there is a unique  $y \in S$  so that  $\|x - y\| = d(x, S)$ . Once more, in our setting of compact smooth embedded manifolds  $N_0$ ,  $\text{reach}(N_0) > 0$  [Thä08, Prop. 14].

## A.2 Spectral gap for the RGG of $\nu$

The interface between our results and those relating to embedded Riemannian manifolds in Euclidean spaces is primarily contained in Theorem 36 below, which gives a spectral gap for the random geometric graph (RGG) of  $\nu$ , and is used to establish our main technical result in this embedded manifold setting, Lemma 23, which gives a weak form of strong concavity for the empirical dual function.

The RGG of  $\nu$  is defined, for a fixed threshold  $\delta > 0$ , as the graph on  $[n]$  with weights for  $j, k \in [n]$  defined as

$$w_{jk} := \begin{cases} \frac{C}{n\delta^{d_\nu+2}} & \text{if } \|y_j - y_k\| < \delta, \\ 0 & \text{else,} \end{cases}$$

where  $C = C(d_\nu)$  is constant only depending on the intrinsic dimension  $d_\nu$ . We write  $j \sim k$  when  $\|y_j - y_k\| < \delta$ , and the resulting weighted graph on  $[n]$  is written as  $\Gamma$ . For  $\alpha \in L^\infty(\hat{\nu})$ , define the associated Dirichlet form by

$$D(\alpha) := \frac{1}{n} \sum_{j \sim k} w_{jk} (\alpha(y_k) - \alpha(y_j))^2,$$

where  $C = C(d_\nu)$  is a constant only depending on the intrinsic dimension  $d_\nu$ . The unnormalized graph Laplacian on  $\Gamma$  with respect to  $L^2(\hat{\nu})$  is then

$$\Delta_\Gamma(\alpha)(y_j) := \sum_{k: k \sim j} w_{jk} (\alpha(y_k) - \alpha(y_j)).$$

We seek a spectral gap for  $\Delta_\Gamma$ , meaning that its second eigenvalue is bounded away from 0. To sketch how this is possible, note that under our assumptions on  $N$  and  $\nu$ , such a result holds for the continuous analog of  $\Delta_\Gamma$ . By employing recent work on the convergence of the spectrum of RGGs to their continuous analogs, we can transfer a continuous spectral gap to a (random) discrete one [GTGHS20]. The result we ultimately obtain is as follows.

**Theorem 36** (Spectral gap for the RGG of  $\nu$ ). *Suppose  $\delta \lesssim 1$  and  $n$  is large enough that  $\delta^{d_\nu} \gtrsim (\log n)/n$ . Then with probability at least  $1 - C/n$ ,  $\lambda_2(\Delta_\Gamma) \gtrsim 1$ . In this case, we have the inequality*

$$D(\alpha) \gtrsim \hat{\nu}((\alpha - \hat{\nu}(\alpha))^2) \quad \forall \alpha \in L^\infty(\hat{\nu}).$$

To derive this theorem from the results of [GTGHS20], we first introduce the continuous analog of the un-normalized graph Laplacian and subsequently discuss its spectrum.

**A certain weighted Laplacian.** For a thorough introduction to weighted Riemannian Laplacians we refer the reader to the book [Gri09]. Suppose  $\nu$  has density  $p: N \rightarrow \mathbb{R}_{\geq 0}$  with respect to  $\text{vol}_N$ . Consider the weighted Laplacian operator

$$\Delta_\nu(\cdot) := -\frac{1}{p} \text{div}(p^2 \nabla \cdot),$$

where  $\text{div}$  is the Riemannian divergence on  $N$ . In fact, this weighted Laplacian is the correct continuous analog of the un-normalized graph Laplacian  $\Delta_\Gamma$  [BCH03, HAL07], and we will thus consider its spectrum. Because  $p$  is bounded away from 0 and  $N$  is assumed to be compact,  $\Delta_\nu$  has a discrete, non-negative spectrum, written  $0 = \lambda_1 \leq \lambda_2 \leq \dots$  (with multiplicity), and the minimax principle holds [GTGHS20, Section 1.4]. Since  $N$  is connected and  $p$  is bounded above and away from 0, it follows that  $\lambda_2 > 0$ . Summarizing, we have the following Proposition.

**Proposition 37.** *The operator  $\Delta_\nu$  has a discrete, non-negative spectrum  $\{\lambda_k\}_{k \geq 1}$ , and  $\lambda_1 = 0$ , while  $\lambda_2 > 0$ .*

We will simply treat  $\lambda_2$  as a constant depending on  $N$  and  $\nu$ , but we remark that precise quantitative control on  $\lambda_2$  is intimately related to the Ricci curvature of the manifold  $N$  [Li12, Chapter 5].

**Spectral convergence of the RGG of  $\nu$ .** The results of [GTGHS20] quantify the convergence of the spectrum of  $\Delta_\Gamma$  to that of  $\Delta_\nu$ . For our purposes, it is enough to consider their result only in terms of lower bound control of the second eigenvalue of  $\Delta_\Gamma$ , namely  $\lambda_2(\Delta_\Gamma)$ , in terms of the second eigenvalue of  $\Delta_\nu$ , namely  $\lambda_2$ , but we emphasize that their results are far stronger than we represent here. To make the geometric constants involved in this bound explicit, let  $i_0 := \text{inj}(N)$  denote the injectivity radius of  $N$ ,  $K$  be a uniform upper bound on the absolute value of the sectional curvatures of  $N$ , and  $r := \text{reach}(N)$  the reach (these quantities are defined in the previous section). Since  $N$  is compact,  $i_0, r > 0$  and  $K < \infty$ .

**Theorem 38** (Theorem 4 (adapted) [GTGHS20]). *Suppose  $\delta > 0$  is such that*

$$\delta < \min \left\{ 1, \frac{i_0}{10}, \frac{1}{\sqrt{d_\nu K}}, \frac{r}{\sqrt{27 d_\nu}} \right\}.$$

Suppose  $W_\infty(\hat{\nu}, \nu)(d_\nu + 5) < \delta$ . Then, if

$$\sqrt{\lambda_2} \delta < C$$

we have

$$\lambda_2(\Delta_\Gamma) \geq \lambda_2(1 - C(\delta + \frac{1}{\delta}W_\infty(\hat{\nu}, \nu) + \sqrt{\lambda_2}\delta + K\delta^2)),$$

where the constants  $C$  only depend on  $d_\nu$  and uniform bounds on  $p$  and its Lipschitz constant.

This result combined with Proposition 37 shows that, so long as  $\delta$  is sufficiently small relative to constants depending on the geometry of  $N$ , the intrinsic dimension  $d_\nu$ , and uniform bounds on  $p$  and its Lipschitz constant, and  $W_\infty(\hat{\nu}, \nu) \lesssim \delta$ , we have  $\lambda_2(\Delta_\Gamma) \gtrsim 1$ .

To make this bound usable, we also need to get control on  $W_\infty(\hat{\nu}, \nu)$ .

**Theorem 39** (Theorem 2 (adapted) [GTGHS20]). *With probability at least  $1 - C/n$ ,*

$$W_\infty(\hat{\mu}, \mu) \lesssim \left(\frac{\log n}{n}\right)^{1/d_\nu},$$

where the constants only depend on  $d_\nu$ , upper and lower bounds on  $p$  and its Lipschitz constant, and the injectivity radius of  $N$ , uniform bound on the absolute value of the sectional curvature of  $N$ , and the reach of  $N$ .

Combining Proposition 37, Theorem 38, and Theorem 39, we obtain Theorem 36.

**Scale of our results.** The statement of Theorem 38 allows us to describe the required size of  $\delta$  in Theorem 36, and hence  $\varepsilon/L$  in our results from section 3.3, in terms of the injectivity radius  $i_0$ , uniform bound  $K$  on the absolute value of the sectional curvature, reach  $r$  (these geometric quantities are described in detail in section A.1) and second eigenvalue  $\lambda_2$ , as

$$\frac{\varepsilon}{L} < \min \left\{ 1, \frac{i_0}{10}, \frac{1}{\sqrt{d_\nu K}}, \frac{r}{\sqrt{27d_\nu}}, \frac{C}{\sqrt{\lambda_2}} \right\}, \quad (\text{A.1})$$

where  $C$  is a constant depending on  $d_\nu$ , uniform upper and lower bounds on  $p$  and its Lipschitz constant. In fact, we will use two more small-scale facts about the embedded manifold  $N$  beyond Theorem 38, namely Proposition 42 and Proposition 43 below, but this bound is enough to permit their use.

### A.3 Additional facts about embedded manifolds

We also use Assumptions 2 and 3 to derive a Poincaré inequality for  $\nu$  in Proposition 40, as well as to give convenient bounds on the covering numbers  $\mathcal{N}(\nu, \cdot)$  in Proposition 43.

Similar reasoning as in the previous section implies that the weighted Laplacian  $\Delta(\cdot) = -\frac{1}{p} \operatorname{div}(p\nabla\cdot)$  sports a spectral gap, and entails the following result.

**Proposition 40.** *For all locally Lipschitz functions  $\zeta: N \rightarrow \mathbb{R}$  we have*

$$\operatorname{Var}_\nu(\zeta) \lesssim \int \|\nabla_N \zeta(y)\|_h^2 d\nu(y).$$

The following Proposition is an elementary consequence of the fact that  $N$  is an embedded manifold of  $\mathbb{R}^d$ .

**Proposition 41.** *Let  $\zeta: N \rightarrow \mathbb{R}$  and  $p \in N$ . Suppose  $\zeta$  has a  $\nabla_N$  gradient at  $p$ , and admits a local extension  $\bar{\zeta}: U \rightarrow \mathbb{R}$  on some neighborhood  $U \subset \mathbb{R}^d$  of  $p$  such that  $\bar{\zeta}$  has a Euclidean gradient at  $p$ . Then*

$$\|\nabla_N \zeta(p)\|_h^2 \leq \|\nabla \bar{\zeta}(p)\|^2.$$

*Proof of Proposition 41.* Let  $\iota: N \hookrightarrow \mathbb{R}^d$  be the embedding of  $N$  into  $\mathbb{R}^d$ . Then for all  $v \in T_p N$ ,

$$d\zeta_p(v) = \langle \nabla_N \zeta(p), v \rangle_h = \langle d\iota_p(\nabla_N \zeta(p)), d\iota_p(v) \rangle,$$

and on the other hand,

$$d\zeta_p(v) = d\bar{\zeta}_{\iota(p)} \circ d\iota_p(v) = \langle \nabla \bar{\zeta}(\iota(p)), d\iota_p(v) \rangle.$$

Since this is true for all  $v \in T_p N$ , it follows that  $d\iota_p(\nabla_N \zeta(p))$  is the orthogonal projection of  $\nabla \bar{\zeta}(p)$  onto  $\text{im}(d\iota_p) \subset T_{\iota(p)} \mathbb{R}^d$ , and so

$$\|\nabla_N \zeta(p)\|_h^2 = \|d\iota_p(\nabla_N \zeta(p))\|^2 \leq \|\nabla \bar{\zeta}(p)\|^2.$$

□

**Proposition 42.** *Let  $i_0 := \text{inj}(N)$  be the injectivity radius of  $N$ . Let  $K$  be a uniform upper bound on the absolute value of the sectional curvatures of  $N$ , and  $r := \text{reach}(N)$ . Suppose  $\tau > 0$  is such that*

$$\tau \leq \min \left\{ \frac{1}{\sqrt{K}}, i_0, \frac{r}{2} \right\}.$$

Then for all  $y \in N$ ,

$$\tau^{d\nu} \lesssim \nu(B(y, \tau)) \lesssim \tau^{d\nu}.$$

*Proof of Proposition 42.* Fix some  $y \in N$ . By [GTGHS20, Proposition 2], for any  $y' \in N$  such that  $\|y - y'\| \leq r/2$ ,

$$\|y - y'\| \leq \text{dist}_N(y, y') \leq \|y - y'\| + \frac{8}{r^2} \|y - y'\|^3$$

Hence

$$B_{\text{dist}_N}(y, \tau) \subset B_{\|\cdot\|}(y, \tau) \cap N \subset B_{\text{dist}_N}\left(y, \tau + \frac{8}{r^2} \tau^3\right).$$

Whence

$$\nu(B(y, \tau)) = \nu(B_{\|\cdot\|}(y, \tau)) \geq \nu(B_{\text{dist}_N}(y, \tau)) \gtrsim \text{vol}_N(B_{\text{dist}_N}(y, \tau)),$$

where the final inequality follows because the density of  $\nu$  with respect to  $\text{vol}_N$  is bounded away from 0 under Assumptions 2 and 3. By [GTGHS20, Equation 1.35], when  $\tau \lesssim 1$ ,

$$\text{vol}_N(B_{\text{dist}_N}(y, \tau)) \gtrsim \tau^{d\nu},$$



yielding the lower bound. Similarly,

$$\nu(B(y, \tau)) = \nu(B_{\|\cdot\|}(y, \tau)) \leq \nu(B_{\text{dist}_N}(y, \tau + \frac{8}{r^2}\tau^3)) \lesssim \text{vol}_N(B_{\text{dist}_N}(y, \tau + \frac{8}{r^2}\tau^3)),$$

and we can conclude the upper bound with another application of [GTGHS20, Equation 1.35] where we have the opposite inequality

$$\text{vol}_N(B_{\text{dist}_N}(y, \tau + \frac{8}{r^2}\tau^3)) \lesssim (\tau + \frac{8}{r^2}\tau^3)^{d_\nu} \lesssim \tau^{d_\nu}.$$

□

The following control on the covering numbers of  $N$  will also be convenient.

**Proposition 43.** *Suppose  $\tau$  is as in Proposition 42. Then*

$$\tau^{-d_\nu} \lesssim \mathcal{N}(N, \|\cdot\|, \tau) \lesssim \tau^{-d_\nu}.$$

*Proof of Proposition 43.* For the upper bound, take a maximal  $\tau/2$  packing of  $N$  in the Euclidean norm,  $z_1, \dots, z_K$ . Then

$$1 \geq \sum_{k=1}^K \nu(B(z_k, \tau/2)).$$

For  $\tau \lesssim 1$ , by Proposition 42,

$$1 \gtrsim K\tau^{d_\nu}.$$

Observe that since the packing is maximal,  $K \geq \mathcal{N}(N, \|\cdot\|, \tau)$ , completing the proof of the upper bound.

For the lower bound, we argue similarly. Take any  $\tau$  cover of  $N$  in the Euclidean norm,  $z_1, \dots, z_K$ . Then by Proposition 42

$$1 \leq \sum_{k=1}^K \nu(B(z_k, \tau)) \lesssim K\tau^{d_\nu}.$$

Taking a minimal  $\tau$ -covering yields the result. □

## B Deferred proofs

### B.1 On the tightness of Theorem 2

In this section, we give our results concerning the tightness of Theorem 2 in terms of covering number dependence. Our main observation in this vein follows.

**Proposition 44.** *Suppose there exists  $r \geq 0$  such that for all  $\varepsilon > 0$ , probability measures  $\mu, \nu$  and cost functions  $c$  satisfying Assumption 1, there is a numerical constant such that*

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \sqrt{\frac{\mathcal{N}(\mu, (\frac{\varepsilon}{L})^r) \wedge \mathcal{N}(\nu, (\frac{\varepsilon}{L})^r)}{n}}.$$

*Then  $r \geq 1$ .*

To prove this result, we establish the following fact.

**Proposition 45** (Entropic estimation of  $W_1$  distances). *Suppose  $c(x, y) = \|x - y\|$ , and there exists  $r \geq 0$  such that for all  $\varepsilon > 0$ , and all  $\mu, \nu$  supported in  $B(0, 1/2)$ , there is a numerical constant for which*

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \sqrt{\frac{\mathcal{N}(\mu, (\frac{\varepsilon}{L})^r) \wedge \mathcal{N}(\nu, (\frac{\varepsilon}{L})^r)}{n}}.$$

Then for dimension-dependent constants  $C_d, C'_d$ ,

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - W_1(\mu, \nu)|] \lesssim C_d \left\{ \frac{1}{rd + 4} \log(C'_d n) + 1 \right\} \cdot n^{-1/(rd+2)}$$

In particular, Theorem 2 implies the rate  $n^{-1/(d+2)}$  for  $W_1$  distance estimation, up to logarithmic factors and dimension-dependent constants.

Proposition 44 follows from this result by recalling that for  $d \geq 3$ , the minimax rates for  $W_1$  distance estimation over all distributions supported in  $B(0, 1/2)$  are  $n^{-1/d}$  up to logarithmic factors [NWR22, Theorem 11].

*Proof of Proposition 45.* Using standard upper bounds on covering numbers in Euclidean spaces [Ver18, Proposition 4.2.12], the hypothesis implies

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - S_\varepsilon(\mu, \nu)|] \lesssim (1 + \varepsilon) \cdot \left(1 + \frac{1}{\varepsilon^r}\right)^{d/2} \cdot \frac{1}{\sqrt{n}}. \quad (\text{B.1})$$

Note that for  $\varepsilon = 0$  we recover the Wasserstein-1 distance, and the rate of approximation is known to be bounded as

$$|S_\varepsilon(\mu, \nu) - W_1(\mu, \nu)| \leq C_d \varepsilon \log\left(\frac{C'_d}{\varepsilon}\right), \quad (\text{B.2})$$

where  $C_d, C'_d$  are dimension-dependent constants [GCB<sup>+</sup>19, Theorem 1]. Using  $S_\varepsilon(\hat{\mu}, \hat{\nu})$  to estimate  $W_1(\mu, \nu)$ , suppose we take  $\varepsilon = n^{-1/(rd+2)}$ . We then find, for dimension-dependent constants  $C_d, C'_d$  (potentially different than before), that

$$\mathbb{E}[|S_\varepsilon(\hat{\mu}, \hat{\nu}) - W_1(\mu, \nu)|] \lesssim C_d \left\{ \frac{1}{rd + 4} \log(C'_d n) + 1 \right\} \cdot n^{-1/(rd+2)}$$

□

## B.2 Deferred proofs from section 6

*Proof of Lemma 25.* Let  $z_1, \dots, z_K$  be a proper  $\varepsilon/2L$ -net of  $N$  such that  $K = \mathcal{N}^{\text{pr}}(N, \frac{\varepsilon}{2L})$ . For each  $z \in N$ , there exists  $k \in [K]$  such that  $z \in B(z_k, \frac{\varepsilon}{2L})$ , so that  $B(z_k, \frac{\varepsilon}{2L}) \subset B(z, \frac{\varepsilon}{L})$ . Thus

$$\inf_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L})) \geq \min_{k \in [K]} \hat{\nu}(B(z_k, \frac{\varepsilon}{2L})).$$

Therefore, for all  $t > 0$ ,

$$\begin{aligned}
& \mathbb{P}[\inf_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L})) \leq \inf_{z \in N} \nu(B(z, \frac{\varepsilon}{2L})) - t] \\
& \leq \mathbb{P}[\min_{k \in [K]} \hat{\nu}(B(z_k, \frac{\varepsilon}{2L})) \leq \inf_{z \in N} \nu(B(z, \frac{\varepsilon}{2L})) - t] \\
& \leq \sum_{k=1}^K \mathbb{P}[\hat{\nu}(B(z_k, \frac{\varepsilon}{2L})) \leq \inf_{z \in N} \nu(B(z, \frac{\varepsilon}{2L})) - t] \\
& \leq \sum_{k=1}^K \mathbb{P}[\hat{\nu}(B(z_k, \frac{\varepsilon}{2L})) \leq \nu(B(z_k, \frac{\varepsilon}{2L})) - t].
\end{aligned}$$

By Hoeffding's inequality, for each  $k \in [K]$ ,

$$\mathbb{P}[\hat{\nu}(B(z_k, \frac{\varepsilon}{2L})) \leq \nu(B(z_k, \frac{\varepsilon}{2L})) - t] \leq e^{-2nt^2}.$$

Therefore, with probability at least  $1 - \frac{1}{n}e^{-\frac{10}{\varepsilon}}$ ,

$$\inf_{z \in N} \hat{\nu}(B(z, \frac{\varepsilon}{L})) \gtrsim \inf_{z \in N} \nu(B(z, \frac{\varepsilon}{2L})) - \sqrt{\frac{1/\varepsilon + \log(nK)}{n}}.$$

Note that  $K = \mathcal{N}^{\text{pr}}(N, \frac{\varepsilon}{2L}) \leq \mathcal{N}(N, \frac{\varepsilon}{4L})$ , and so we may conclude with Proposition 42 and Proposition 43 combined with our assumption on  $n$  in Equation (6.2).  $\square$

*Proof of Lemma 26.* For the first statement, observe that for all  $x \in \text{supp } \mu$

$$1 = \int p_\varepsilon(x, y') d\nu(y').$$

By Proposition 15 for any  $y \in \text{supp } \nu$ ,

$$1 \gtrsim p_\varepsilon(x, y) \nu(B(y, \frac{\varepsilon}{L})) \gtrsim p_\varepsilon(x, y) \left(\frac{\varepsilon}{L}\right)^{d_\nu},$$

where the second inequality follows by Proposition 42. Re-arranging yields the first statement since  $x \in \text{supp } \mu$  and  $y \in \text{supp } \nu$  were arbitrary.

For the second statement, it suffices to show the statement in the event described by Lemma 25. Use Proposition 15 to find that, for any  $x \in \text{supp } \mu, y \in \text{supp } \nu$ ,

$$\hat{p}_\varepsilon(x, y) \leq \min_{j \in [n]} \left\{ e^{\frac{2L}{\varepsilon} \|y - y_j\|} \hat{p}_\varepsilon(x, y_j) \right\}.$$

Reasoning as above,

$$\hat{p}_\varepsilon(x, y_j) \lesssim \hat{\nu}(B(y_j, \frac{\varepsilon}{L}))^{-1} \leq \sup_{y' \in N} \hat{\nu}(B(y', \frac{\varepsilon}{L}))^{-1}.$$

Since we are working in the event described by Lemma 25,

$$\sup_{y' \in N} \hat{\nu}(B(y', \frac{\varepsilon}{L}))^{-1} \lesssim \left(\frac{L}{\varepsilon}\right)^{d_\nu}.$$

It follows that there exists a  $y_j$  such that  $\|y - y_j\| \leq \frac{\varepsilon}{L}$ , in which case

$$\hat{p}_\varepsilon(x, y) \lesssim \left(\frac{L}{\varepsilon}\right)^{d_\nu}.$$

□

*Proof of Lemma 35.* We give the proof of the first claim, the remaining two follow in the same fashion. Let  $\mathcal{E}$  denote the event in Lemma 26. Observe that, by the pointwise bounds in Proposition 14,

$$\begin{aligned} \mathbb{E}[\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] &= \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] + \mathbb{E}[\mathbb{1}[\mathcal{E}^c]\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] \\ &\lesssim \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] + e^{\frac{10}{\varepsilon}} \mathbb{E}[\mathbb{1}[\mathcal{E}^c]] \\ &\leq \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] + \frac{1}{n}. \end{aligned}$$

Note that  $|e^a - e^b| \leq e^{a \vee b}|a - b| = (e^a \vee e^b)|a - b|$  for all  $a, b \in \mathbb{R}$ , and so

$$\begin{aligned} \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\hat{p}_\varepsilon - p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] &\leq \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\mathbb{1}[\mathcal{E}]\|\log \hat{p}_\varepsilon - \log p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] \\ &\leq \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\log \hat{p}_\varepsilon - \log p_\varepsilon\|_{L^2(\mu \otimes \nu)}^2] \\ &= \frac{1}{\varepsilon^2} \cdot \left(\frac{L}{\varepsilon}\right)^{2d_\nu} \mathbb{E}[\|\hat{f}_\varepsilon - f_\varepsilon\|_{L^2(\mu)}^2 + \|\hat{g}_\varepsilon - g_\varepsilon\|_{L^2(\nu)}^2]. \end{aligned}$$

The first inequality follows. □

## References

- [AKT84] Miklós Ajtai, János Komlós, and Gábor Tusnády. On optimal matchings. *Combinatorica*, 4(4):259–264, 1984.
- [BCH03] Olivier Bousquet, Olivier Chapelle, and Matthias Hein. Measure based regularization. *Advances in Neural Information Processing Systems*, 16, 2003.
- [BCP19] Jérémie Bigot, Elsa Cazelles, and Nicolas Papadakis. Central limit theorems for entropy-regularized optimal transport on finite spaces and statistical applications. *Electron. J. Stat.*, 13:5120–5150, 2019.
- [Ber20] Robert J Berman. The Sinkhorn algorithm, parabolic optimal transport and geometric Monge–Ampère equations. *Numer. Math.*, 145(4):771–836, 2020.
- [BEZ22] Erhan Bayraktar, Stephan Eckstein, and Xin Zhang. Stability and sample complexity of divergence regularized optimal transport. *arXiv preprint*, 2022.
- [BGC17] Yoshua Bengio, Ian Goodfellow, and Aaron Courville. *Deep learning*, volume 1. MIT press Cambridge, MA, USA, 2017.

- [BLG14] Emmanuel Boissard and Thibaut Le Gouic. On the mean speed of convergence of empirical and occupation measures in Wasserstein distance. In *Annales de l'IHP Probabilités et statistiques*, volume 50, pages 539–563, 2014.
- [BO69] Richard L Bishop and Barrett O’Neill. Manifolds of negative curvature. *Trans. Am. Math. Soc.*, 145:1–49, 1969.
- [CKA23] Marco Cuturi, Michal Klein, and Pierre Ablin. Monge, Bregman and Occam: Interpretable optimal transport in high-dimensions with feature-sparse maps. *arXiv preprint*, 2023.
- [CRL<sup>+</sup>20] Lénaïc Chizat, Pierre Roussillon, Flavien Léger, François-Xavier Vialard, and Gabriel Peyré. Faster Wasserstein distance estimation with the Sinkhorn divergence. *Advances in Neural Information Processing Systems*, 2020.
- [Cut13] Marco Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. In *Advances in Neural Information Processing Systems*, 2013.
- [dBSL<sup>+</sup>23] Eustasio del Barrio, Alberto González Sanz, Jean-Michel Loubes, and Jonathan Niles-Weed. An improved central limit theorem and fast convergence rates for entropic transportation costs. *SIAM J. Math. Data Sci.*, 5(3):639–669, 2023.
- [DC16] Manfredo P Do Carmo. *Differential geometry of curves and surfaces: revised and updated second edition*. Courier Dover Publications, 2016.
- [DGP<sup>+</sup>20] Yihe Dong, Yu Gao, Richard Peng, Ilya Razenshteyn, and Saurabh Sawlani. A study of performance of optimal transport. *arXiv preprint*, 2020.
- [DMG20] Simone Di Marino and Augusto Gerolin. An optimal transport approach for the Schrödinger bridge problem and convergence of Sinkhorn algorithm. *Journal of Scientific Computing*, 85(2):1–28, 2020.
- [Dud69] Richard Mansfield Dudley. The speed of mean Glivenko-Cantelli convergence. *Ann. Math. Stat.*, 40(1):40–50, 1969.
- [DY95] V Dobrić and Joseph E Yukich. Asymptotics for transportation cost in high dimensions. *J. Theor. Prob.*, 8:97–118, 1995.
- [EN23] Stephan Eckstein and Marcel Nutz. Convergence rates for regularized optimal transport via quantization. *Math. Oper. Res.*, 2023.
- [FCTR16] R Flamary, N Courty, D Tuia, and A Rakotomamonjy. Optimal transport for domain adaptation. *IEEE Trans. Pattern Anal. Mach. Intell.*, 1, 2016.
- [FMN16] Charles Fefferman, Sanjoy Mitter, and Hariharan Narayanan. Testing the manifold hypothesis. *J. Am. Math. Soc.*, 29(4):983–1049, 2016.

- [GCB<sup>+</sup>19] Aude Genevay, Lénaïc Chizat, Francis Bach, Marco Cuturi, and Gabriel Peyré. Sample complexity of Sinkhorn divergences. *International Conference on Artificial Intelligence and Statistics*, 2019.
- [GH23] Michel Groppe and Shayan Hundrieser. Lower complexity adaptation for empirical entropic optimal transport. *arXiv preprint*, 2023.
- [GKRS22] Ziv Goldfeld, Kengo Kato, Gabriel Rioux, and Ritwik Sadhu. Statistical inference with regularized optimal transport. *arXiv preprint*, 2022.
- [Gri09] Alexander Grigoryan. *Heat kernel and analysis on manifolds*. American Mathematical Society, 2009.
- [GSH23] Alberto González-Sanz and Shayan Hundrieser. Weak limits for empirical entropic optimal transport: Beyond smooth costs. *arXiv preprint*, 2023.
- [GSLNW22] Alberto Gonzalez-Sanz, Jean-Michel Loubes, and Jonathan Niles-Weed. Weak limits of entropy regularized optimal transport; potentials, plans and divergences. *arXiv preprint*, 2022.
- [GTGHS20] Nicolás García Trillos, Moritz Gerlach, Matthias Hein, and Dejan Slepčev. Error estimates for spectral convergence of the graph Laplacian on random geometric graphs toward the Laplace–Beltrami operator. *Found. Comput. Math.*, 20(4):827–887, 2020.
- [HAL07] Matthias Hein, Jean-Yves Audibert, and Ulrike von Luxburg. Graph Laplacians and their convergence on random neighborhood graphs. *J. Mach. Learn. Res.*, 8(6), 2007.
- [HR21] Jan-Christian Hütter and Philippe Rigollet. Minimax estimation of smooth optimal transport maps. *Ann. Stat.*, 49(2):1166–1194, 2021.
- [HSM22] Shayan Hundrieser, Thomas Staudt, and Axel Munk. Empirical optimal transport between different measures adapts to lower complexity. *arXiv preprint*, 2022.
- [KNS16] Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the Polyak–Lojasiewicz condition. *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, 2016.
- [KTM20] Marcel Klatt, Carla Taming, and Axel Munk. Empirical regularized optimal transport: statistical theory and applications. *SIAM J. Math. Data Sci.*, 2(2):419–443, 2020.
- [Lee18] John M Lee. *Introduction to Riemannian manifolds*, volume 2. Springer, 2018.
- [Li12] Peter Li. *Geometric analysis*. Cambridge University Press, 2012.

- [LS86] Frank Thomson Leighton and Peter Shor. Tight bounds for minimax grid matching, with applications to the average case analysis of algorithms. *ACM Symposium on Theory of Computing*, 1986.
- [LSPC19] Giulia Luise, Saverio Salzo, Massimiliano Pontil, and Carlo Ciliberto. Sinkhorn barycenters with free support via Frank-Wolfe algorithm. *Advances in Neural Information Processing Systems*, 32:9322–9333, 2019.
- [Mar21] Simone Di Marino. Linear convergence of Sinkhorn algorithm and asymptotic rate. Unpublished manuscript, 2021.
- [MBNWW21] Tudor Manole, Sivaraman Balakrishnan, Jonathan Niles-Weed, and Larry Wasserman. Plugin estimation of smooth optimal transport maps. *arXiv preprint*, 2021.
- [MNW19] Gonzalo Mena and Jonathan Niles-Weed. Statistical bounds for entropic optimal transport: Sample complexity and the central limit theorem. *Advances in Neural Information Processing Systems*, 2019.
- [MNW21] Tudor Manole and Jonathan Niles-Weed. Sharp convergence rates for empirical optimal transport with smooth costs. *arXiv preprint*, 2021.
- [MWMA21] Shoaib Bin Masud, Matthew Werenski, James M Murphy, and Shuchin Aeron. Multivariate rank via entropic optimal transport: Sample efficiency and generative modeling. *arXiv preprint*, 2021.
- [NWR22] Jonathan Niles-Weed and Philippe Rigollet. Estimation of Wasserstein distances in the spiked transport model. *Bernoulli*, 28(4):2663–2688, 2022.
- [PC19] Gabriel Peyré and Marco Cuturi. Computational optimal transport. *Found. Trends Mach. Learn.*, 11(5-6):355–607, 2019.
- [PDNW23] Aram-Alexandre Pooladian, Vincent Divol, and Jonathan Niles-Weed. Minimax estimation of discontinuous optimal transport maps: The semi-discrete case. *arXiv preprint*, 2023.
- [PNW21] Aram-Alexandre Pooladian and Jonathan Niles-Weed. Entropic estimation of optimal transport maps. *arXiv preprint*, 2021.
- [PZ19] Victor M. Panaretos and Yoav Zemel. Statistical aspects of Wasserstein distances. *Annu. Rev. Stat. Appl.*, 6:405–431, 2019.
- [RS22] Philippe Rigollet and Austin J. Stromme. On the sample complexity of entropic optimal transport. *arXiv preprint*, 2022.
- [San15] Filippo Santambrogio. *Optimal transport for applied mathematicians*, volume 87 of *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser/Springer, Cham, 2015. Calculus of variations, PDEs, and modeling.

- [SDF<sup>+</sup>18] Vivien Seguy, Bharath Bhushan Damodaran, Remi Flamary, Nicolas Courty, Antoine Rolet, and Mathieu Blondel. Large-scale optimal transport and mapping estimation. *International Conference on Learning Representations*, 2018.
- [Sin64] Richard Sinkhorn. A relationship between arbitrary positive matrices and doubly stochastic matrices. *Ann. Math. Stat.*, 35(2):876–879, 1964.
- [SK67] Richard Sinkhorn and Paul Knopp. Concerning nonnegative matrices and doubly stochastic matrices. *Pac. J. Math.*, 21(2):343–348, 1967.
- [SST<sup>+</sup>19] Geoffrey Schiebinger, Jian Shu, Marcin Tabaka, Brian Cleary, Vidya Subramanian, Aryeh Solomon, Joshua Gould, Siyan Liu, Stacie Lin, Peter Berube, et al. Optimal-transport analysis of single-cell gene expression identifies developmental trajectories in reprogramming. *Cell*, 176(4):928–943, 2019.
- [Str23] Austin J. Stromme. Sampling from a Schrödinger bridge. *International Conference on Artificial Intelligence and Statistics*, 2023.
- [Thä08] Christoph Thäle. 50 years sets with positive reach—a survey. *Surv. Math. Appl.*, 3:123–165, 2008.
- [Ver18] Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge University Press, 2018.
- [Vil08] Cédric Villani. *Optimal transport: Old and new*. Springer Science & Business Media, 2008.
- [WB19] Jonathan Weed and Francis Bach. Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. *Bernoulli*, 25(4 A):2620–2648, 2019.
- [WMMA23] Matthew Werenski, Shoaib Bin Masud, James M Murphy, and Shuchin Aeron. On rank energy statistics via optimal transport: Continuity, convergence, and change point detection. *arXiv preprint*, 2023.
- [ZGMS22] Zhengxin Zhang, Ziv Goldfeld, Youssef Mroueh, and Bharath K Sriperumbudur. Gromov-Wasserstein distances: Entropic regularization, duality, and sample complexity. *arXiv preprint*, 2022.