## BOUNDARY REGULARITY OF UNIFORMLY ROTATING VORTEX PATCHES AND AN UNSTABLE ELLIPTIC FREE BOUNDARY PROBLEM

YUCHEN WANG<sup>1,2</sup>, GUANGHUI ZHANG<sup>3</sup> AND MAOLIN ZHOU<sup>4</sup>

ABSTRACT. In this paper, we consider a sign-changing free boundary problem that comes from the boundary regularity of rotating vortex patches of the twodimensional incompressible Euler equations. The complete classification of singular points has been obtained through establishing a new Weiss-type monotonicity formula. Upon these results, we prove that only 90° corner type of singularity could happen at the boundary of a Lipschitz rotating vortex patch, while the other parts are  $C^{\infty}$  smooth.

**Keywords:** Vortex patches; Free boundary problem; Weiss-type monotonicity formula; Regularity of free boundary; Singularity

#### 1. INTRODUCTION

Consider the 2-dim incompressible Euler equations on the whole plane

(1.1) 
$$\begin{cases} \partial_t v + (v \cdot \nabla)v = -\nabla p, \quad x \in \mathbf{R}^2, \\ \nabla \cdot v = 0, \\ v \to 0, \quad |x| \to \infty, \end{cases}$$

where  $v = (v^1, v^2)^t \in \mathbf{R}^2$  is the velocity of fluids and p is the pressure. Throughout this paper, we shall mainly work with the vorticity formulation of the 2-dim incompressible Euler equations

(1.2) 
$$\begin{cases} \partial_t \omega + (v \cdot \nabla)\omega = 0, & x \in \mathbf{R}^2, \\ v = -\nabla^{\perp}(-\Delta)^{-1}\omega, & (a_1, a_2)^{\perp} := (-a_2, a_1)^t \end{cases}$$

with the vorticity field given by  $\omega := \partial_1 v^2 - \partial_2 v^1$ , where the velocity is uniquely determined by the vorticity due to the *stream-vorticity* formula. It is clearly a closed

Date: March 19, 2024.

 $<sup>^{1,2}</sup>$  partially supported by DFG ZH 605/1-2 and NSF of China No.11831009 .

<sup>&</sup>lt;sup>3</sup> partially supported by NSF of China (No. 12171176, 11971187).

<sup>&</sup>lt;sup>4</sup> supported by the National Key Research and Development Program of China (2021YFA1002400), Nankai Zhide Foundation and NSF of China (No. 12271437, 11971498).

PDEs system concerned  $\omega$  only and the global well-posedness of smooth solutions as well as the Yudovich-type weak solutions  $\omega \in L^1(\mathbf{R}^2) \cap L^{\infty}(\mathbf{R}^2)$  is well-known. We refer to [25, 33] for detailed discussions on the necessary backgrounds and reviews on the classical results for interested readers.

Among many other topics, special interest focuses on the vortex patch solution in the form of

(1.3) 
$$\omega(t,x) = \sum_{j=1}^{N} \kappa_j \mathbf{I}_{D_j(t)},$$

where  $\kappa_1, \ldots, \kappa_N$  are non-zero constants,  $D_1, \ldots, D_N$  are disjoint bounded domains and  $\mathbf{I}_D$  denotes the characteristic function of D. The global existence and the uniqueness of vortex patches follow from [33] straightforwardly. The global regularity of patches is proved by Chemin [3], Bertozzi and Constantin [7], Serfati [29] and Kiselev and Luo [23] if the initial patch is smooth. On the other hand, Chemin [8] studied the preserving of singularities, see also [10] where Elgindi and Jeong considered well/ill-posedness of specific singular structures. The global regularity of the patch solution of a 2-dim generalized transport equation was proved by Verdera in [31].

Moreover, Kiselev and Sverak [24] studied the small-scale formation of the 2-dim incompressible Euler equations, namely the infinite time blow-up solutions which satisfy  $\|\nabla \omega(t)\|_{\infty} \sim C_1 e^{e^{C_2 t}}$ . Kiselev and Li [22] proved that similar phenomenon that the curvature of the boundary tends to infinity happens for vortex patches. This is a long-standing conjecture that vortex patches would eventually weakly converge to some steady vortex patches. This conjecture seems beyond the current capability of PDEs, but also indicates that singular uniformly rotating vortex patches may be the global attractors of vortex patches.

Our main interest will focus on the steady vortex patches. Being aware of the rotational and translation invariances of the system, our consideration will mainly involve the uniformly rotating vortex patches

$$\omega(t, x) = \omega_0(e^{-i\Omega t}x),$$

which are relative equilibria of the Euler equations, i.e., the vorticity configuration is invariant up to rigid motions, where  $\Omega \in \mathbf{R}$  is the angular velocity and  $\omega_0 = \mathbf{I}_D$ is the initial patch. It is a natural setting due to the rotational invariance of  $\mathbf{R}^2$ and conservation of the center of vorticity. Since the incompressible condition implies that there exists a scalar function  $\Psi$ , referred to the stream function, such that  $v = -\nabla^{\perp}\Psi$ . It is indeed a nonlinear elliptic problem on the relative stream function  $\psi = \Psi + \frac{\Omega}{2}|x|^2$ . We call  $P \in \mathbf{R}^2$  is a stagnation point if v = 0, or equivalently  $\nabla\psi(P) = 0$ . We still reserve the name stream function for  $\psi$  for simplicity. Suppose D is a bounded domain that is enclosed by a rectifiable simple closed curve. It is not hard to verify that  $\omega = \mathbf{I}_D$  is a uniformly rotating vortex patch if and only if  $\psi$  solves the elliptic free boundary problem

(1.4) 
$$\begin{cases} -\Delta \psi = \mathbf{I}_D - 2\Omega, & x \in \mathbf{R}^2, \\ \nabla(\psi - \frac{\Omega}{2}|x|^2) \to 0, & |x| \to \infty, \\ \psi = 0, & x \in \partial D, \end{cases}$$

where the vortical domain D is the main unknown. Clearly circular patch  $\omega = \mathbf{I}_{B_1}$ (the Rankine vortex) is a uniformly rotating solution for any  $\Omega \in \mathbf{R}$ . The other explicit non-trivial uniformly rotating vortex patch is the Kirchhoff ellipse patch

$$\omega = \mathbf{I}_{E_{a,b}}, \quad E_{a,b} := \left\{ (x_1, x_2) \in \mathbf{R}^2 | \frac{x_1^2}{a^2} + \frac{x_2^2}{b^2} \le 1 \right\}$$

where the angular velocity satisfies  $0 < \Omega = \frac{ab}{(a+b)^2} \leq \frac{1}{4}$ . Of course the vortical domain may have a complicated topology and it is sufficient to slightly modify the equation (1.4) to

(1.5) 
$$\begin{cases} -\Delta \psi = \mathbf{I}_D - 2\Omega, & x \in \mathbf{R}^2, \\ \nabla(\psi - \frac{\Omega}{2}|x|^2) \to 0, & |x| \to \infty, \\ \psi = c_j, & x \in \Gamma_j, \end{cases}$$

in which each  $\Gamma_j$  is a connected component of  $\partial D$  and  $c_j$  are constants,  $1 \leq j \leq n$ , where *n* is the number of connected-components of  $\partial D$ . Clearly annular patches

$$\omega = \mathbf{I}_{C_{b,1}}, C_{b,1} := \{ x \in \mathbf{R}^2 | b \le |x| \le 1 \}$$

are trivial solutions of (1.5) holding for any  $\Omega \in \mathbf{R}$ . Beyond these explicit solutions, lots of uniformly rotating vortex patches are obtained via the local bifurcation approach [4, 19, 18, 9, 17, 6]. All of them are sufficiently close to explicit solutions for example the circular/annular/ellipse vortex patches in the sense of boundary perturbations.

On the other hand, Hassainia, Masmoudi and Wheeler [16] studied the global continuation of the *m*-fold local bifurcation curve  $\mathscr{S}_m$  emanating from the Rankine vortex for  $m \geq 3$ . They show that the end of bifurcation curves has the following alternatives:

- (1) Smooth vortex patch with overhanging profiles, i.e., the boundary can not be parameterized as a graph in the polar coordinates.
- (2) Limiting V-states, i.e., singular vortex patch where the boundary is parameterized as a non- $C^1$  graph in the polar coordinates.

Together with numerical observations [32, 27], they made the following conjecture.

# Conjecture: The singular solution with $90^{\circ}$ corners seen in numerics for $m \geq 3$ exists as the (weak) limits of patches along the *m*-fold branch $\mathscr{S}_m$ as the parameter $s \to \infty$ .

This conjecture is highly challenging since it essentially depends on the geometric property of special solutions that emanate from the disk. They are not easy to obtain explicitly. Some recent progress on the quantitative properties of the *m*-fold rotating vortex patch has been made in [28] for sufficiently large m via a variational argument based on the optimal transport. Inspired by the remarkable progress on the study of unstable obstacle problem [26, 2], it is an interesting problem on the boundary regularity of uniformly rotating vortex patches which may not be located on the bifurcation curves emanating from disk. This motivates the present paper.

Our consideration is based on the observations that uniformly rotating vortex patch is a particular case of the elliptic unstable free boundary problem

(1.6) 
$$\begin{cases} -\Delta u = \lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c} & \text{in } \mathbf{R}^2, \\ u = 0 & \text{on } \partial D, \end{cases}$$

in which  $\lambda_1, \lambda_2 > 0$ , D is a bounded domain. By the maximum principle u is positive in D, however u may change its sign in  $D^c$ . We shall focus on the regularity of free boundary (the regularity of  $\partial D \subset \{u = 0\}$ ) via tools developing in the study of obstacle-type problems since the pioneer work of Caffarelli [5], in particular, the Weiss-type monotonicity formula. See [13, 12] and references therein for more recent advantages. Up to the author's knowledge, there are only a few results on the regularity of uniformly rotating vortex patches except that are sufficiently close to explicit solutions. For example, in [19] Hmidi, Mateu and Verdera proved that the boundary of uniformly rotating vortex patches on the local branch emanating from the Rankine vortex is  $C^{\infty}$  smooth. The smoothness was improved to realanalytic very soon by Castro, Córdoba and Gómez-Serrano in [6]. On the other hand, regarding the unstable obstacle problem, results obtained in [26, 2] could not be implemented here straightforwardly since the level set  $\{u = 0\}$  does not completely coincide with free boundary and behaviors of the stream function are essentially different, nor the minimal/maximal solutions hardly exists.

The main result of this paper is given as follows:

**Theorem 1.1.** Suppose D is a bounded domain, u solves the 2-dim unstable elliptic free boundary problem and is positive in D

(1.7) 
$$\begin{cases} -\Delta u = \lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c} & \text{in } \mathbf{R}^2 \\ u = 0 & \text{on } \partial D \end{cases}, \quad \lambda_1, \lambda_2 > 0. \end{cases}$$

The free boundary  $\partial D$  is locally real-analytic up to the singular sets

$$\mathcal{S} := \{ x \in \partial D \mid \nabla u(x) = 0 \},\$$

which consist of

- (1) isolated singular points where the free boundary  $\partial D$  consists of two  $C^1$ -curves intersecting each other with a 90°-corner (Figure 1(a) or Figure 1(b)).
- (2) intersection points of finitely many local connected components of the boundary. The blow-up limits of the neighbourhood of the singular points must be one of the following alternatives:
  - The whole plane with a ray (Figure 1(c)) or a straight line (Figure 1(d)).
  - N-fold symmetric domain for  $N \ge 3$  (Figure 1(e)). It is made up of alternating regions of sub-harmonic domain and super-harmonic domain of u.
  - Axi-symmetric domain with even components (Figure 1(f)). There is exactly one sector with an opening angle greater than  $\pi$ , while the complement consists of odd components no less than 5 and is symmetric concerning the angular bisector.

For more accurate descriptions on singular points, please see Proposition 2.3.




FIGURE 1. Classification on the singular points

Remark 1.2. The free boundary near singular points may be very complicated hence it could not be written down as intersections of curves in general. We introduce the two dimensional **density** of domain E at some point x defined by

$$\lim_{r \to 0} \frac{\mathcal{L}^2(B_r(x) \cap E)}{\mathcal{L}^2(B_r(x))}.$$

therefore the two dimensional density of domain D would be 0 or constants depending on  $\lambda_1, \lambda_2$  only.

In Theorem 1.1, we assume that u is positive in the bounded domain D, following the background of vortex patches. Here we note that the complete classification on singular points of problem (1.6) can also be obtained by the analysis of our paper without assuming the positivity of u in D.

The rotating vortex patch problem is a special class of the sign-changing free boundary problem if and only if  $\Omega \in (0, \frac{1}{2})$ . Here the singularity problem is only meaning for vortex patch with Lipschitz boundary with  $\Omega \in (0, \frac{1}{2})$ . A domain is called a Lipschitz domain if its boundary is a Lipschitz curve. With the Lipschitz smoothness assumption on the boundary, we can give a complete description of the boundary regularity of uniformly rotating vortex patches.

**Theorem 1.3.** Suppose D is a Lipschitz domain and  $\omega = \mathbf{I}_D$  is a uniformly rotating vortex patch with angular velocity  $0 < \Omega < \frac{1}{2}$ . Then the singular set  $S \subset \partial D$  contains at most finitely many points, and  $\partial D \setminus S$  is smooth. Near each singular point  $x \in S$ , for sufficiently small r > 0,  $\partial D \cap B(x, r)$  consists of two  $C^1$  arcs meeting at the right angle. In particular, if there is no stagnation point on the boundary, the boundary is smooth.

Remark 1.4. The conclusion also holds for the symmetric rotating vortex pairs by the same argument. It holds regardless of solutions located on the bifurcation curves. The result recovers numerical observations obtained in [32]. We expect the analysis would be useful in solving the conjecture on the limiting V-states but a more delicate quantitative analysis on the geometry of the vortical domain seems necessary.

In contrast to Theorem 1.3, it seems another singularities even for Lipschitz domain when  $\Omega \notin (0, \frac{1}{2})$ . However, we recall the following remarkable result on the rigidity due to Gomes-Serrano, Park, Shi and Yao [15] hence another type of singularity is ruled out.

**Theorem A** Let  $D \subset \mathbf{R}^2$  be a bounded domain with Lipschitz boundary. Assume that  $\omega = \mathbf{I}_D$  is a stationary/uniformly rotating vortex patch of (1.5) for some  $\Omega \in \mathbf{R}$ . Then D must be radially symmetric if  $\Omega \in (-\infty, 0] \cup [\frac{1}{2}, +\infty)$  and radially symmetric up to a translation if  $\Omega = 0$ . It is also worth pointing out that the consideration here is a priori, whether a singular steady vortex patch exists remains an open problem up to our best knowledge. One could apply a variational method to study the existence of rotating vortex patches as [30], and the existence of uniformly rotating vortex patches could follow from maximizing the energy function penalized the angular momentum

(1.8) 
$$E(\omega) = \frac{1}{2} \int_{\mathbf{R}^2} \omega (-\Delta)^{-1} \omega d\mu - \frac{\Omega}{2} \int_{\mathbf{R}^2} |x|^2 \omega d\mu$$

among the rearrangement class of prescribed vorticity with prescribed angular impulse. But it seems highly non-trivial to determine the regularity of boundary except the vortex patch is highly concentrated. Indeed, for the steady vortex patches in the bounded domain, Turkington [30] also proposed a conjecture that whether the singular sets on steady vortex patch have Hausdorff dimension zero. So, if the conjecture in [16] holds, it verifies the existence of singular uniformly rotating patches, which may be of independent interest. We also note that Garcia and Hazioy [14] studied the global continuation of the local branch emanating from a pair of corotating point vortices in a very recent work. More limiting V-states occur and their geometry is much more complicated if  $D_1$  and  $D_2$  would touch with each other in several ways.

This paper is organized as follows. In Section 2 we consider a sign-changing unstable elliptic free boundary problem and derive the Weiss-type monotonicity formula. The classification of singular points is obtained due to their blow-up limits. In Section 3, we prove the uniform regularity near singular points where u have super-characteristic growth then finish the proof of Theorem 1.1.

### 2. A SIGN-CHANGING TWO PHASES UNSTABLE ELLIPTIC FREE BOUNDARY PROBLEM: WEISS-TYPE MONOTONICITY FORMULA AND CLASSIFICATION OF BLOW-UP LIMITS

The uniformly rotating vortex patch problem (1.4) is a special case of the twophase unstable elliptic free boundary problem

(2.1) 
$$\begin{cases} -\Delta u = \lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}, & \text{ in } \mathbf{R}^n, \ n \ge 2, \\ u = 0, & \text{ on } \partial D, \end{cases}$$

where  $\lambda_1, \lambda_2 > 0$  are prescribed constants and D is a bounded domain which is the main unknown.  $D^c := \mathbf{R}^n \setminus D$  denotes the complementary of D. Note that u may change its sign in  $\overline{D}^c$ ,  $\overline{D}^c$  might not be the set  $\{u < 0\}$ , thus (2.1) is different from the classical unstable obstacle problems which were studied in [26, 2]. The standard elliptic regularity theory implies  $u \in C^{1,\alpha}(\mathbf{R}^2)$  for all  $\alpha \in (0,1)$  and u is real-analytic in both D and  $\overline{D}^c$ . Moreover, for  $x \in \partial D$  where  $\nabla u(x) \neq 0$ , the implicit function

theorem implies that  $\partial D$  is a  $C^{1,\alpha}$  curve near x. It is indeed locally analytic. The singular set of  $\partial D$  is defined by

(2.2) 
$$\mathcal{S} := \left\{ x \in \partial D \middle| \nabla u(x) = 0 \right\}$$

**Lemma 2.1.** Suppose that u is a weak solution of (2.1). For any  $x_0 \in \mathbb{R}^n$ , let

$$\Phi_{x_0}(r) = r^{-n-2} \int_{B_r(x_0)} |\nabla u|^2 - 2u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^n - 2r^{-n-3} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1},$$

where  $\mathcal{H}^k$  is the k-dimensional Hausdorff measure. Then for any  $0 < \rho < \delta$ , the following Weiss-type monotonicity formula holds

(2.3) 
$$\Phi_{x_0}(\delta) - \Phi_{x_0}(\rho) = \int_{\rho}^{\delta} \int_{\partial B_r(x_0)} 2r^{-n-2} (\nabla u \cdot \nu - \frac{2u}{r})^2 d\mathcal{H}^{n-1} dr \ge 0.$$

We first prove the following lemma.

**Lemma 2.2.** If u is a weak solution of (2.1), then

(2.4) 
$$\int_{\mathbb{R}^n} |\nabla u|^2 \operatorname{div} \vec{X} - 2(\nabla u)^t \mathcal{D} \vec{X} \nabla u - 2u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) \operatorname{div} \vec{X} = 0$$

holds for all  $\vec{X} = (X_1, \dots, X_n) \in C_0^{\infty}(\mathbf{R}^n, \mathbf{R}^n).$ 

Proof. By a straightforward computation,

$$\begin{split} 0 &= \int_{\mathbf{R}^{n}} \operatorname{div}(|\nabla u|^{2}\vec{X}) \\ &= \int_{\mathbf{R}^{N}} 2\sum_{i=1}^{n} \sum_{j=1}^{n} \partial_{i} u \partial_{ij} u X_{j} + |\nabla u|^{2} \operatorname{div} \vec{X} \\ &= \int_{\mathbf{R}^{n}} -2\sum_{i=1}^{n} \sum_{j=1}^{n} \left( \partial_{ii} u \partial_{j} u X^{j} + \partial_{i} u \partial_{i} X^{j} \partial_{j} u \right) + |\nabla u|^{2} \operatorname{div} \vec{X} \\ &= \int_{\mathbf{R}^{n}} -2\Delta u (\nabla u \cdot \vec{X}) - 2(\nabla u)^{t} \mathcal{D} \vec{X} \nabla u + |\nabla u|^{2} \operatorname{div} \vec{X} \\ &= \int_{\mathbf{R}^{n}} 2\lambda_{1} \mathbf{I}_{D} \nabla u \cdot \vec{X} - 2\lambda_{2} \mathbf{I}_{D^{c}} \nabla u \cdot \vec{X} - 2(\nabla u)^{t} \mathcal{D} \vec{X} \nabla u + |\nabla u|^{2} \operatorname{div} \vec{X} \\ &= \int_{\mathbf{R}^{n}} |\nabla u|^{2} \operatorname{div} \vec{X} - 2(\nabla u)^{t} \mathcal{D} \vec{X} \nabla u - 2u(\lambda_{1} \mathbf{I}_{D} - \lambda_{2} \mathbf{I}_{D^{c}}) \operatorname{div} \vec{X}, \end{split}$$

the proof is complete.

Proof of Lemma 2.1. Suppose that r, k are positive constants and let

$$\eta_k(x) = \max\left\{0, \min\left\{1, (r - |x|)k\right\}\right\}, \vec{X}_k(x) = \eta_k(x)x = (\eta_k(x)x_1, \cdots, \eta_k(x)x_n).$$

A straightforward computation shows that

$$\operatorname{div} \vec{X}_k(x) = \sum_{i=1}^n \partial_i \left( \eta_k(x) x_i \right) = n \eta_k(x) + \nabla \eta_k(x) \cdot x,$$
$$(\nabla u)^t \mathcal{D} \vec{X}_k \nabla u = \sum_{i=1}^n \sum_{j=1}^n \partial_i u \partial_j (\eta_k(x) x_i) \partial_j u$$
$$= \sum_{i=1}^n \sum_{j=1}^n \partial_i u \eta_k \delta_{ij} \partial_j u + \partial_i u \partial_j (\eta_k) x_i \partial_j u.$$
$$= |\nabla u|^2 \eta_k + (\nabla u \cdot x) (\nabla u \cdot \nabla \eta_k).$$

According to (2.4), one has

$$0 = \int_{\mathbf{R}^n} n|\nabla u|^2 \eta_k + |\nabla u|^2 \nabla \eta_k \cdot x - 2|\nabla u|^2 \eta_k - 2(\nabla u \cdot x)(\nabla u \cdot \nabla \eta_k)$$
$$- \int_{\mathbf{R}^n} 2(\lambda_1 \mathbf{I}_D - 2\lambda_2 \mathbf{I}_{D^c})(n\eta_k + \nabla \eta_k \cdot x)u$$

Now letting  $k \to +\infty$ , we obtain

(2.5) 
$$0 = \int_{B_r(0)} (n-2) |\nabla u|^2 - 2nu(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^n - r \int_{\partial B_r(0)} |\nabla u|^2 - 2(\nabla u \cdot \nu)^2 - 2u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^{n-1},$$

for a.e. r > 0, where  $\nu$  is the unit outward normal vector of  $B_r(0)$ . Then one has

$$\frac{d}{dr}\Phi_{x_0}(r) = -(n+2)r^{-n-3}\int_{B_r(x_0)} |\nabla u|^2 - 2u(\lambda_1 \mathbf{I}_D - 2\lambda_2 \mathbf{I}_{D^c})d\mathcal{H}^n 
+ r^{-n-2}\int_{\partial B_r(x_0)} |\nabla u|^2 - 2u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c})d\mathcal{H}^{n-1} 
+ 8r^{-5}\int_{\partial B_1(0)} u^2(x_0 + rx)d\mathcal{H}^{n-1} - 4r^{-4}\int_{\partial B_1(0)} u(x_0 + rx)\left(\nabla u(x_0 + rx) \cdot x\right)d\mathcal{H}^{n-1}.$$

Inserting (2.5) into above equation, one has

$$\Phi_{x_0}'(r) = -r^{-n-3} \int_{B_r(x_0)} (n-2) |\nabla u|^2 - 2nu(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^n$$
  
$$- 4r^{-n-3} \int_{B_r(x_0)} |\nabla u|^2 - u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^n$$
  
$$+ r^{-n-2} \int_{\partial B_r(x_0)} |\nabla u|^2 - 2u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^{n-1}$$
  
$$+ 8r^{-n-4} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1} - 4r^{-n-3} \int_{\partial B_r(x_0)} u(\nabla u \cdot \nu) d\mathcal{H}^{n-1}$$
  
$$= 2r^{-n-2} \int_{\partial B_r(x_0)} (\nabla u \cdot \nu)^2 d\mathcal{H}^{n-1} + 8r^{-n-4} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1}$$
  
$$- 8r^{-n-3} \int_{\partial B_r(x_0)} u(\nabla u \cdot \nu) d\mathcal{H}^{n-1}$$
  
$$= 2r^{-n-2} \int_{\partial B_r(x_0)} \left(\nabla u \cdot \nu - \frac{2u}{r}\right)^2 d\mathcal{H}^{n-1},$$

in which we used the identity

$$\int_{\partial B_r(x_0)} u(\nabla u \cdot \nu) d\mathcal{H}^{n-1} = \int_{B_r(x_0)} |\nabla u|^2 + u\Delta u \ d\mathcal{H}^n$$
$$= \int_{B_r(x_0)} |\nabla u|^2 - u(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}) d\mathcal{H}^n.$$

(2.3) is obtained by integrating (2.6).

Lemma 2.1 implies a classification of the blow-up limits of u at singular points on the free boundary.

**Proposition 2.3** (Classification of blow-up limits). Suppose u is a solution of (2.1) and  $x_0 \in S^u$ . The following cases occur alternatively:

(1) In the case of  $\Phi_{x_0}(0+) = -\infty$ , one has  $\lim_{r \to 0} r^{-3-n} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1} = +\infty$ , and each limit of

$$v_r(x) = \frac{u(x_0 + rx)}{S(x_0, r)}$$

as  $r \to 0$  is a homogeneous harmonic polynomial of degree two, where

$$S(x_0, r) := \left( r^{1-n} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1} \right)^{1/2}.$$

(2) In the case of  $\Phi_{x_0}(0+) > -\infty$ , for any given R > 0, there exists  $\tau(R) > 0$  such that for  $r < \tau(R)$ ,

$$u_r(x) := \frac{u(x_0 + rx)}{r^2}$$

is bounded in  $W^{1,2}(B_R(0))$ . Let  $\{r_j\}_{j\in\mathbb{N}}$  be a sequence such that  $\{u_{r_j}\}_{j\in\mathbb{N}}$  converges to  $u_0$  weakly in  $W^{1,2}_{loc}(\mathbf{R}^n)$ , then  $u_0$  is a degree two homogeneous solution of

(2.7) 
$$\begin{cases} -\Delta u_0 = \lambda_1 \mathbf{I}_{D_0} - \lambda_2 \mathbf{I}_{D_0^c} & \text{in } \mathbf{R}^n, \\ u = 0 & \text{on } \partial D_0, \end{cases}$$

in which  $\mathbf{I}_{D_0} = \lim_{j \to \infty} \mathbf{I}_{D_{r_j}}$ , where  $D_r = \{x : x_0 + rx \in D\}$ .

Remark 2.4. Note that u may change its sign in  $D_0^c$ . We have

(2.8)  
$$\Psi_{x_0}(0+) = \int_{B_1(0)} |\nabla u_0|^2 - 2u_0(\lambda_1 \mathbf{I}_{D_0} - \lambda_2 \mathbf{I}_{D_0^c}) d\mathcal{H}^N - 2 \int_{\partial B_1(0)} u_0^2 d\mathcal{H}^{N_1}$$
$$= -2\lambda_1 \int_{D_0 \cap B_1(0)} u_0 d\mathcal{H}^N + 2\lambda_2 \int_{D_0^c \cap B_1(0)} u_0 d\mathcal{H}^N$$

may be non-negative for  $u_0 \neq 0$ .

Proof of Proposition 2.3. (1). If  $\Phi_{x_0}(0+) = -\infty$ , we suppose towards a contradiction that  $\limsup_{r\to 0} S(x_0, r) \leq Mr^2$  for some constant M, there exists an  $r_0 > 0$  such that, for  $r < r_0$ ,

$$\int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1} \le M^2 r^{n+3}.$$

Thus

$$\int_{B_r(x_0)} |u| d\mathcal{H}^n = \int_0^r \int_{\partial B_t(x_0)} |u| d\mathcal{H}^{n-1} dt$$

$$\leq C_1 \int_0^r \left( \int_{\partial B_t(x_0)} u^2 d\mathcal{H}^{n-1} \right)^{\frac{1}{2}} t^{\frac{n-1}{2}} dt$$

$$\leq C_2 \int_0^r t^{\frac{n+3}{2}} t^{\frac{n-1}{2}} dt$$

$$\leq C_3 r^{n+2}.$$

Consequently

$$\left|\frac{1}{r^{n+2}}\int_{B_r(x_0)} 2u\left(\lambda_1\mathbf{I}_D - \lambda_2\mathbf{I}_{D^c}\right) d\mathcal{H}^n\right| \le C_4.$$

We conclude that

$$\Phi_{x_0}(r) \ge - \left| r^{-n-2} \int_{B_r(x_0)} 22 \left( \lambda_1 \mathbf{I}_D u - \lambda_2 \mathbf{I}_{D^c} \right) d\mathcal{H}^n \right| - 2r^{-n-3} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1} \ge -C_4 - 2M^2,$$

for all  $r < r_0$ , which contradicts the assumption that  $\Phi_{x_0}(0+) = -\infty$ . Therefore  $\limsup_{r\to 0} \frac{S(x_0,r)}{r^2} = +\infty$ . We choose a sequence  $\{r_j\}_{j\in\mathbb{N}}$  such that  $\lim_{j\to\infty} r_j = 0$  and  $\lim_{j\to\infty} \frac{S(x_0,r_j)}{r_j^2} = +\infty$ . Let

$$v_k(x) = \frac{u(x_0 + r_k x)}{S(x_0, r_k)}.$$

Then  $||v_k||_{L^2(\partial B_1(0))} = 1$ . We may assume that  $\lim_{k\to\infty} v_k \to v_0$  weakly in  $L^2(\partial B_1(0))$ . Let  $K = \frac{\max\{\lambda_1, \lambda_2\}}{2n}$ . Then  $u_r^+ + K|x|^2$  and  $u_r^- + K|x|^2$  are subharmonic functions which imply

$$\frac{1}{t^{n-1}} \int_{\partial B_t(0)} u_r^+ + K|x|^2 d\mathcal{H}^{n-1},$$
$$\frac{1}{t^{n-1}} \int_{\partial B_t(0)} u_r^- + K|x|^2 d\mathcal{H}^{n-1},$$

are monotone increasing in t. Therefore

$$\int_{\partial B_t(0)} u_r^+ d\mathcal{H}^{n-1} \le t^{n-1} \int_{\partial B_1(0)} u_r^+ + K |x|^2 d\mathcal{H}^{n-1}$$
$$\le C_5 t^{n-1} \left( \|u_r\|_{L^2(\partial B_1(0))} + 1 \right),$$

for some constant  $C_5$ . Therefore

$$\int_{B_1(0)} u_r^+ d\mathcal{H}^n = \int_0^1 \int_{\partial B_t(0)} u_r^+ d\mathcal{H}^{n-1} dt \le \frac{C_5}{n} \left( \|u_r\|_{L^2(\partial B_1(0))} + 1 \right),$$

and similarly

$$\int_{B_1(0)} u_r^- d\mathcal{H}^n \le \frac{C_5}{n} \left( \|u_r\|_{L^2(\partial B_1(0))} + 1 \right).$$

By the monotonicity formula (2.3) we have

$$\int_{B_1(0)} |\nabla u_{r_k}|^2 + 2\lambda_1 \mathbf{I}_{D_{r_k}} u_{r_k} - 2\lambda_2 \mathbf{I}_{D_{r_k}^c} u_{r_k} d\mathcal{H}^n - 2\int_{\partial B_1(0)} u_{r_k}^2 d\mathcal{H}^{n-1} \le \Phi_{x_0}(r_1).$$

Let 
$$T_k = \frac{S(x_0, r_k)}{r_k^2}$$
 and  $v_k(x) = \frac{u(x_0 + r_k x)}{S(x_0, r_k)}$ . Then  

$$\int_{B_1(0)} |\nabla v_k|^2 d\mathcal{H}^n \le T_k^{-2} \Phi_{x_0}(r_1) + CT_k^{-2} \int_{B_1(0)} |u_{r_k}| d\mathcal{H}^n + 2 \int_{\partial B_1(0)} v_k^2 d\mathcal{H}^{n-1}.$$

12

Note that  $\lim_{k\to\infty} T_k = +\infty$  and

$$T_k^{-1} \int_{B_1(0)} |u_{r_k}| d\mathcal{H}^{n-1} \le CT_k^{-1} \left( \|u_{r_k}\|_{L^2(\partial B_1(0))} + 1 \right) \le C(1 + T_k^{-1}).$$

Thus

$$\sup_{k \in \mathbb{N}} \|\nabla v_k\|_{L^2(B_1(0))} \le C_6$$

for some constant  $C_6$ . By the Poincaré-Steklov inequality (see e.g. [11, Lemma 3.30]) we have

$$\|v_k\|_{L^2(B_1(0))} \le C_7 \left( \|\nabla v_k\|_{L^2(B_1(0))} + \|v_k\|_{L^2(\partial B_1(0))} \right) \le C_8,$$

where  $C_8$  does not depend on k. Thus we may assume that  $v_k$  converges to  $v_0$  weakly in  $W^{1,2}(B_1(0))$ . Letting  $k \to \infty$ , we obtain that  $v_0$  is harmonic and

$$\int_{B_1(0)} |\nabla v_0|^2 d\mathcal{H}^n \le 2 \int_{\partial B_1(0)} v_0^2 d\mathcal{H}^{n-1}$$

Since  $|\Delta v_k| \leq \max\{\lambda_1, \lambda_2\}$ , by the elliptic regularity and Sobolev embedding theorem, for any given  $\alpha \in (0, 1)$ , there exists a constant  $C_9$  such that

$$\sup_{k \in \mathbb{N}} \|v_k\|_{C^{1,\alpha}(B_1(0))} \le C_9.$$

We may assume that  $v_k$  converges to  $v_0$  in  $C^{1,\beta}(B_1(0))$  for some  $\beta \in (0, \alpha)$ . Notice that  $v_k(0) = 0$  and  $\nabla v_k(0) = 0$ , we have  $v_0(0) = 0$  and  $\nabla v_0(0) = 0$ . It follows from the frequency formula (see e.g. Lemma 4.2 of [26]) that  $v_0$  is a harmonic polynomial of degree 2.

(2). If  $\Phi_{x_0}(0+) > -\infty$ , there exists a constant  $M < \infty$  such that

$$\limsup_{r \to 0} r^{3-n} \int_{\partial B_r(x_0)} u^2 d\mathcal{H}^{n-1} < M.$$

Thus we can choose  $r_0 > 0$  small enough such that  $||u_r||_{L^2(\partial B_1(0))} \leq M + 1$  for  $r \in (0, r_0)$ . Without loss of generality, we set R = 1. Similar to case (1), we have  $\int_{B_1(0)} |u_r| d\mathcal{H}^n < C_{10}$  for some constant  $C_{10}$ . It follows from the monotonicity formula that there exists a constant  $C_{11}$  such that

$$\int_{B_1(0)} |\nabla u_r(x)|^2 \le \Phi_{x_0}(r) + 2 \max\{\lambda_1, \lambda_2\} \int_{B_1(0)} |u_r| d\mathcal{H}^2 + 2 \int_{\partial B_1(x)} u_r^2 d\mathcal{H}^{n-1} \le C_{11},$$

for all  $r \in (0, r_0)$ . Thus  $\{u_r\}$  is uniformly bounded in  $W^{1,2}(B_1(0))$ . We may choose a sequence  $\{r_k\}_{k\in\mathbb{N}}$  such that, for some  $u_0 \in W^{1,2}(B_1(0))$ ,  $u_{r_k} \to u_0$  weakly in  $W^{1,2}(B_1(0)), u_{r_k} \to u_0$  in  $L^2(B_1(0))$  and  $u_{r_k} \to u_0$  in  $L^2(\partial B_1(0))$ . Moreover, for any given  $\sigma > \rho > 0$ , we have

$$\int_{\rho}^{\delta} \int_{\partial B_r(x_0)} 2\left(\nabla u_{r_k} \cdot \nu - \frac{2u_{r_k}}{r}\right) d\mathcal{H}^{n-1} dr = \Phi_{x_0}(r_k\delta) - \Phi_{x_0}(r_k\rho) \to 0,$$

as  $k \to \infty$ . Letting  $k \to \infty$ , we have

$$\nabla u_0(x) \cdot x - 2u_0(x) = 0$$

for all  $x \in \mathbb{R}^n$ , and thus  $u_0$  is homogeneous of degree 2.

On the other hand, the exists a set  $D_0$  such that  $\mathbf{I}_{D_r} \to \mathbf{I}_{D_0}$  in  $L^1_{loc}(\mathbb{R}^N)$ , where  $D_r := \{x | x_0 + rx \in D\}$ . Thus  $u_0$  is a weak solution of degree 2 to

$$\Delta u_0 = \lambda_1 \mathbf{I}_{D_0} - \lambda_2 \mathbf{I}_{D_0^c},$$

It is clear that  $u_0 = 0$  on  $\partial D_0$ .

Proposition 2.3 implies the singular points on the free boundary would be classified into the following two classes:

(1) singular points where u has super-characteristic growth (the blow-up limit 1),

(2) singular points where u has characteristic growth (the blow-up limit 2).

 $P \in \partial D$  is called **degenerate** singular point if either the blow-up limit of domain D at P, or its complement, is a set of measure 0. All the singular points are isolated except the degenerate points provided n = 2.

# 3. Singularities of the two-dimensional unstable elliptic free boundary problem

The proof of our main results will be accomplished by the following lemmas. Firstly, we focus on the two-dimensional unstable elliptic free boundary problem

(3.1) 
$$\begin{cases} -\Delta \psi = \lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c} & \text{in } \mathbf{R}^2, \\ \nabla (\psi - \frac{\Omega}{2} |x|^2) \to 0, & |x| \to \infty, \\ \psi = 0 & \text{on } \partial D, \end{cases}$$

where D is a bounded domain. Due to the maximum principle, we have  $\psi > 0$  in D.

3.1. Blow-up analysis. Suppose  $x_0 \in S^{\psi}$ . It follows from the monotonicity formula

(3.2) 
$$\Phi_{x_0}^{\psi}(r) = \frac{1}{r^4} \int_{B_r(x_0)} |\nabla \psi|^2 - 2\psi \left(\lambda_1 \mathbf{I}_D - \lambda_2 \mathbf{I}_{D^c}\right) d\mathcal{H}^2 - \frac{2}{r^5} \int_{\partial B_r(x_0)} \psi^2 d\mathcal{H}^1_{\underline{\cdot}}$$

is monotone increasing. Proposition 2.3 implies the following alternatives occur hence we need to consider case by case.

**Blow-up limit I:** If  $\Phi_{x_0}^{\psi}(0+) = -\infty$ . One has

$$r^{-5} \int_{\partial B_r(x_0)} \psi^2 d\mathcal{H}^1 = +\infty.$$

Proposition 2.3 implies that, as  $r \to 0$ , any limit function of

(3.3) 
$$\frac{\psi(x_0 + rx)}{S(x_0, r)}$$

must be a homogeneous harmonic polynomial of the form

(3.4) 
$$\frac{x_1x_2 + A(x_1^2 - x_2^2)}{\|x_1x_2 + A(x_1^2 - x_2^2)\|_{L^2(\partial B_1)}}, \quad A \in \mathbf{R},$$

where  $S(x_0, r) = (\frac{1}{r} \int_{\partial B_r(x_0)} \psi^2 d\mathcal{H}^1)^{\frac{1}{2}}$ . Up to a rotation, we may assume that A = 0. Let  $D_0$  be the limit of  $D_r$ , then  $D_0 = \{(x_1, x_2) | x_1 x_2 > 0\}$ , or  $D_0 = \{(x_1, x_2) | x_1 > 0, x_2 > 0\}$ , or  $D_0 = \{(x_1, x_2) | x_1 < 0, x_2 < 0\}$ .

**Blow-up limit II:** If  $\Phi_{x_0}^{\psi}(0+) > -\infty$ , by Proposition 2.3 one has

(3.5) 
$$\psi_r(x) := \frac{\psi(x_0 + rx)}{r^2}$$

is bounded in  $W^{1,2}(B_R)$  for any given R > 0, and there exists a subsequence  $\{u_{r_j}\}$  weakly converging to degree two homogeneous function  $\psi_0$  satisfying

(3.6) 
$$\begin{cases} -\Delta\psi_0 = \lambda_1 \mathbf{I}_{D_0} - \lambda_2 \mathbf{I}_{D_0^c}, & x \in \mathbb{R}^2, \\ \psi_0 = 0, & x \in \partial D_0, \end{cases}$$

where  $D_r := \{x \mid x_0 + rx \in D\}$  and the blow-up limit  $I_{D_0} := \lim_{j \to \infty} I_{D_{r_j}}$  is dependent of extraction of the subsequence  $r_j \to 0$ . Let  $\psi_0 = r^2 f(\theta)$ . Due to the homogeneity, f solves the following ODE

(3.7) 
$$\begin{cases} -f''(\theta) - 4f(\theta) = \lambda_1, \ \theta_{2i} \le \theta \le \theta_{2i+1}, \\ -f''(\theta) - 4f(\theta) = -\lambda_2, \ \theta_{2i+1} \le \theta \le \theta_{2i+2}, \\ f(\theta_{2i}) = f(\theta_{2i+1}) = f(\theta_{2i+2}) = 0, \end{cases}$$

for i = 0, ..., N. With loss of generality we let  $\theta_0 = \theta_{2N+2} = 0$ . By the standard ODE technique, we can write down the solution in each sector explicitly as follows: (3.8)

$$f(\theta) = \begin{cases} -\frac{\lambda_1}{4} + \frac{\lambda_1}{4\sin 2(\theta_{2i+1} - \theta_{2i})} \left( (\sin 2\theta_{2i+1} - \sin 2\theta_{2i}) \cos 2\theta + (\cos 2\theta_{2i} - \cos 2\theta_{2i+1}) \sin 2\theta \right), \\ \text{for } \theta \in (\theta_{2i}, \theta_{2i+1}), \\ \frac{\lambda_2}{4} - \frac{\lambda_2}{4\sin 2(\theta_{2i+2} - \theta_{2i+1})} \left( (\sin 2\theta_{2i+2} - \sin 2\theta_{2i+1}) \cos 2\theta + (\cos 2\theta_{2i+1} - \cos 2\theta_{2i+2}) \sin 2\theta \right), \\ \text{for } \theta \in (\theta_{2i+1}, \theta_{2i+2}), \end{cases}$$

Note that  $\psi_0 \in C^{1,\alpha}$  implies  $f'(\theta)$  is continuous, which means that (3.9)

$$\frac{\lambda_1}{2\sin 2\alpha_i}(-1+\cos 2\alpha_i) = f'(\theta_{2i+1}) = -\frac{\lambda_2}{2\sin 2\beta_i}(1-\cos 2\beta_i), \quad \alpha_i := \theta_{2i+1} - \theta_{2i}, \ \beta_i := \theta_{2i+2} - \theta_{2i+1}.$$

provided  $\alpha_i, \beta_i \neq 0, \frac{\pi}{2}$  or  $\pi$ . It follows

(3.10) 
$$\frac{\tan \alpha_i}{\tan \beta_i} = \frac{\lambda_2}{\lambda_1} > 0, i = 0, \dots, N$$

which are independent of the index i. Similarly the continuity of  $f'(\theta)$  implies

(3.11) 
$$\frac{\tan \alpha_{i+1}}{\tan \beta_i} = \frac{\tan \alpha_i}{\tan \beta_{i-1}} = \frac{\lambda_2}{\lambda_1} > 0, i = 1, \dots, N-1.$$

The maximal principle implies  $\psi_0 > 0$  for  $(r, \theta) \in \mathbf{R}_{>0} \times (\theta_{2i}, \theta_{2i+1})$  hence  $\alpha_i \leq \pi$ . Moreover, due to the explicit formula (3.8),  $\psi_0$  does not change its sign on  $\theta \in (\theta_{2i+1}, \theta_{2i+2})$  unless  $\beta_i \geq \pi$ . Therefore, due to (3.10) and (3.11), we have the following alternatives:

(1) There exactly exists a super-harmonic sector of  $\psi_0$  with the opening angle denote by  $\beta > \pi$  without loss of generality. The complement consists of N sectors with opening angle  $\alpha$  and other N - 1 sectors with opening angle  $\beta - \pi$  which occur alternatively. Here  $\alpha$  and  $\beta$  satisfy

$$\alpha + \beta = \frac{N+1}{N}\pi$$
 and  $\frac{\tan \alpha}{\tan \beta} = \frac{\lambda_2}{\lambda_1}$ .

In particular, the pattern is  $Z_2$ -symmetric.

(2) The pattern is N-fold symmetric which super-harmonic and sub-harmonic domains of  $\psi_0$  occur alternatively.

Now let us consider the remaining cases  $\alpha_i, \beta_i = 0, \frac{\pi}{2}$  or  $\pi$ .

Firstly, we claim that  $\alpha_i \neq \frac{\pi}{2}$ . Otherwise, assume there are  $A, B \in \mathbf{R}$  such that the solution

$$f(\theta) = -\frac{\lambda_1}{4} + A\cos 2\theta + B\sin 2\theta, \quad \theta \in (\theta_{2i}, \theta_{2i+1}),$$

solving the ODE (3.7). Then one has

$$-\frac{\lambda_1}{4} + A\cos 2\theta_{2i} + B\sin 2\theta_{2i} = -\frac{\lambda_1}{4} + A\cos 2\left(\theta_{2i} + \frac{\pi}{2}\right) + B\sin 2\left(\theta_{2i} + \frac{\pi}{2}\right) = 0.$$

The first equality implies  $A \cos 2\theta_i = -B \sin 2\theta_i$  hence

$$-\frac{\lambda_1}{4} + A\cos 2\theta_{2i} + B\sin 2\theta_{2i} = -\frac{\lambda_1}{4} = 0.$$

It is a contradiction. The case  $\beta_i = \frac{\pi}{2}$  can be excluded similarly.

Secondly, we claim that  $\alpha_i \neq \pi$  hence  $\beta_i \neq 0$ . Otherwise, solving the ODE (3.7) and letting  $\theta_{2i} = 0$  without loss of generality, we have

$$f(\theta) = \frac{\lambda_1}{4} \left( \cos 2\theta - 1 \right) + B \sin 2\theta, \quad \theta \in (0, \pi), \quad B \in \mathbf{R}$$

Applying the strong maximal principle on (3.6), there must be  $f'(0)f'(\pi) \leq 0$  which implies B = 0. Therefore we have  $f(\theta_{2i}) = f(\theta_{2i+1}) = f'(\theta_{2i+1}) = 0$  and it implies  $\beta_i = 0$ . It is impossible since  $\psi_0 > 0$  in  $D_0$  inheriting from those solutions of (3.1) is positive in D.

Finaly, if  $\alpha_i = 0$  then one must have  $\beta_i = \pi$ . Otherwise, it would be  $f'(\theta_{2i+1}) = 0$ which is contradicted with the fact  $f'(\theta_{2i+1})f'(\theta_{2i+2}) < 0$  due to the equation (3.6) by implementing the strong maximal principle.

Last but not least, we show that  $N \ge 3$  in the alternative (2). Indeed, if N = 2, the opening angles  $\alpha$  and  $\beta$  satisfy

$$\alpha + \beta = \pi$$

thus

$$\frac{\tan \alpha}{\tan \beta} = \frac{\tan \alpha}{\tan(\pi - \alpha)} = \frac{\lambda_2}{\lambda_1} > 0.$$

It is impossible since  $\alpha \in (0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi)$ . Due to the discussion above, we have that

**Lemma 3.1.** Suppose  $x^0 \in S^{\psi}$  is a singular point. Regarding of the blow-up limit (2) in Proposition 2.3, for equation (3.7) we have the following conclusions:

- (1) There are finitely many super-harmonic and sub-harmonic domains of  $\psi_0$ .
- (2) The super-harmonic and sub-harmonic domains of  $\psi_0$  would occur as one of the following alternatives:
  - (a) The sub-harmonic domain of  $\psi_0$  is a ray or line on the plane where the super-harmonic domain is their complement.
  - (b) to assemble a N-fold symmetric pattern, where each sector consists of exactly one super-harmonic and sub-harmonic domain of  $\psi_0$  for  $N \ge 3$ . They occur alternatively.
  - (c) to assemble a  $Z_2$ -symmetric pattern consists of 2N sectors for  $N \ge 2$ , includes exactly one super-harmonic domain where  $\psi_0$  changes its sign.

Here N is determined by the constant  $\lambda_1, \lambda_2$ .

*Remark* 3.2. If  $\lambda_1 = 1, \lambda_2 = 0$ , one has  $\beta_i = \frac{\pi}{2}$  which indeed recovers the remarkable example given in [2].

Moreover, suppose the boundary of D is a Lipschitz curve. The boundary can be locally represented as the graph of a Lipschitz function near  $x_0 \in \partial D$ , namely for any  $x_0 \in \partial D$ , there exists  $f \in C^{0,1}(\mathbf{R}^{n-1}, \mathbf{R})$  such that

$$B_{\delta}(x_0) \cap D = \{ (\mathbf{x}, x_n) \big| \mathbf{x} \in \mathbf{R}^{n-1}, x_n \ge f(\mathbf{x}) \}.$$

Taking a blow-up sequence of  $D_{r_k} := \{x | x_0 + r_k x \in D\}$ . For each sequence  $D_{r_k}$ , the blow-up sequence of  $D_{r_k}$  is a Lipschitz domain thus there exists  $\delta_k > 0$ ,  $f_k \in C^{0,1}(\mathbf{R}^{n-1}, \mathbf{R})$  such that

$$B_{\delta}(x_0) \cap D_{r_k} = \{ (\mathbf{x}, x_n) | \mathbf{x} \in \mathbf{R}^{n-1}, x_n \ge f_k(\mathbf{x}) \}.$$

Note that the Lipschitz constant of  $f_k$  is uniformly. There is a subsequence still denoted by  $f_k$  converge to a Lipschitz function  $f_0$  and  $\delta_k \to \delta_0 > 0$  hence the blow-up limit is a Lipschitz domain

$$B_{\delta_0}(x_0) \cap D_0 = \{ (\mathbf{x}, x_n) \big| \mathbf{x} \in \mathbf{R}^{n-1}, x_n \ge f_0(\mathbf{x}) \}.$$

Therefore,  $D_0$  is locally represented as the graph of the Lipschitz function thus  $D_0$  is simply-connected which is a cone. Therefore consider the uniformly rotating vortex patch  $\lambda_1 = 1 - 2\Omega$ ,  $\lambda_2 = 2\Omega$ . Then (3.7) becomes

(3.12) 
$$\begin{cases} -f''(\theta) - 4f(\theta) = 1 - 2\Omega, \ 0 < \theta < \theta_1, \\ -f''(\theta) - 4f(\theta) = -2\Omega, \ \theta_1 < \theta \le 2\pi, \\ f(0) = f(\theta_1) = 0, \end{cases}$$

where  $D_0$  is a cone with an opening angle  $\theta_1$ .

We claim that  $\theta_1$  must be 0 or  $2\pi$ . Denote

$$\chi(\theta) = \begin{cases} 1 - 2\Omega, & 0 < \theta < \theta_1, \\ -2\Omega, & \theta_1 < \theta < 2\pi. \end{cases}$$

and consider the elliptic equation on

(3.13) 
$$-\partial_{\theta\theta}f - 4f = \chi(\theta) \quad \text{on } S^1$$

with the restrictions  $f(\theta_1) = f(0) = 0$ . It is a typical overdetermined problem that the equation has no solution if  $\chi(\theta) \in L^2(S^1)$  has a non-trivial projection onto the image of Ker $(-\partial_{\theta\theta} - 4I)$ . Therefore, let us consider the Fourier series of

$$f(\theta) = \sum_{k \in \mathbf{Z}} \hat{f}(k) e^{ik\theta} \in L^2(S^1)$$

Due to equation (3.13), we have the following computations on the Fourier series

(3.14) 
$$(k^2 - 4)\hat{f}(k) = \frac{1}{2\pi} \int_{S^1} \chi(\theta) e^{-ik\theta} d\theta := \begin{cases} \frac{1}{2\pi} \int_0^{\theta_1} e^{-ik\theta} d\theta, & k \neq 0, \\ \frac{\theta_1 - 4\Omega\pi}{2\pi}, & k = 0. \end{cases}$$

Note that  $\operatorname{Ker}(-\partial_{\theta\theta}-4\mathrm{I}) = \operatorname{span}\{e^{2i\theta}, e^{-2i\theta}\}$ . A necessary condition, such that (3.12) is solvable, is  $\hat{\chi}(\pm 2) = 0$ , namely

(3.15) 
$$\int_{0}^{\theta_{1}} e^{-2i\theta} d\theta = \frac{1 - e^{-2i\theta_{1}}}{2i} = 0.$$

It immediately implies  $\theta_1 = 0, \pi, 2\pi$ . By elliptic regularity theory, we have  $f \in C^{1,\alpha}(S^1)$  since  $\chi \in L^{\infty}(S^1)$ . Then the solution of (3.12) would be unique and the Fourier series of f would converge to f pointwisely if the overdetermined problem has a solution.

For  $\theta_1 = \pi$ , one has

(3.16) 
$$\hat{f}(k) = \frac{1 - e^{-ik\pi}}{2\pi i k (k^2 - 4)} = \begin{cases} \frac{1}{\pi i k (k^2 - 4)} & k \equiv 1 \pmod{2} \\ 0 & k \equiv 0 \pmod{2} \end{cases}$$

Therefore

$$(3.17)$$

$$f(\theta) = -\frac{\pi - 4\Omega\pi}{8\pi} + A_2 \cos 2\theta + B_2 \sin 2\theta + \sum_{n \ge 0} \frac{1}{\pi i (2n+1)((2n+1)^2 - 4)} (e^{i(2n+1)\theta} - e^{-i(2n+1)\theta})$$

$$= -\frac{\pi - 4\Omega\pi}{8\pi} + A_2 \cos 2\theta + B_2 \sin 2\theta + \sum_{n \ge 0} \frac{2}{\pi (2n+1)((2n+1)^2 - 4)} \sin(2n+1)\theta.$$

By a straightforward computation, we have

(3.18)

$$f'(0) = 2B_2 + \frac{2}{\pi} \sum_{n \ge 0} \frac{1}{(2n+1)^2 - 4} = 2B_2 + \frac{2}{\pi} \left( -\frac{1}{3} + \frac{1}{4} \left( \sum_{n \ge 1} \frac{1}{2n-1} - \frac{1}{2n+3} \right) \right)$$
  
= 2B<sub>2</sub>  
$$f'(\pi) = 2B_2 - \frac{2}{\pi} \sum_{n \ge 0} \frac{1}{(2n+1)^2 - 4} = 2B_2 - \frac{2}{\pi} \left( -\frac{1}{3} + \frac{1}{4} \left( \sum_{n \ge 1} \frac{1}{2n-1} - \frac{1}{2n+3} \right) \right)$$
  
= 2B<sub>2</sub>

Therefore, we have

 $f'(0)f'(\pi) = 4B_2^2 \ge 0.$ 

If  $B_2 \neq 0$ , f'(0) should have the same sign with  $f'(\pi)$ . Then  $f(0) = f(\pi) = 0$  immediately implies f is sign-changing in  $(0, \pi)$ , which is a contradiction with  $f \geq 0$  in  $(0, \theta_1)$ .

Otherwise,  $f'(0) = f'(\pi) = 0$ . Since f is  $C^2$  (actually  $C^{\infty}$ ) in the interval  $(0, \theta_1)$ , then there exists  $\xi \in (0, \theta_1)$  such that  $f''(\xi) = 0$ . Therefore, the equation (3.12)

$$-4f(\xi) = 1 - 2\Omega \ge 0$$

implies that  $f(\xi) \leq 0, \xi \in (0, \theta_1)$ , which is also a contradiction with f > 0 in  $(0, \theta_1)$  due to the maximum principle.

On the other hand, if  $\theta_1 = 0$ , we have

(3.19) 
$$f(\theta) = \frac{\Omega}{2} + A\cos 2\theta + B\sin 2\theta.$$

Being aware of the necessary condition f(0) = f'(0) = 0, one has  $f(\theta) = \frac{\Omega}{2}(1 - \cos 2\theta)$ . If  $\theta_1 = 2\pi$ , we have

(3.20) 
$$f(\theta) = -\frac{1-2\Omega}{4} + A\cos 2\theta + B\sin 2\theta,$$

then it implies  $f(\theta) = -\frac{1-2\Omega}{4} (1 - \cos 2\theta)$  due to  $f(2\pi) = f'(2\pi) = 0$ . For the case either  $D_0$  or  $(D^c)_0$  shrinks into a ray  $\{\psi_0 = 0\}$  forms a 0° cusp. Then we have

**Lemma 3.3.** Suppose the domain D is a Lipschitz domain. The singular points on the free boundary  $\partial D$  have two alternatives: (1) isolated singular points where the density of the singular point is density  $\frac{1}{4}$  for D and the free boundary around itforms a 90°-corner. (2) degenerate singular points that either the free boundary  $\partial D$  forms a 0° cusp while the density of D near the singular points is 0.

3.2. Uniform regularity. The Weiss-type monotionicity formula as well as the classification of the blow-up limits of singular points provide some geometric information concerned with the 0-level set of the blow-up limit function. In general the limits may not be unique, i.e., the limiting solution may depend on the choice of subsequence  $\{r_j\}_{j\in\mathbb{N}}$ . For the singular points where  $\psi$  has super characteristic growth (the blow-up limit (1)), we can prove that the blow-up limit is unique and the uniform regularity of the free boundary  $\partial D$  near the singularities. The situation is similar to Theorem A in [1] where the proof essentially depends on the uniqueness of generalized Newtonian potential of  $-\mathbf{I}_{\{x_1x_2>0\}}$ . It suffices for us to consider the specific equation

$$-\Delta \psi = \mathbf{I}_D - 2\Omega, \quad \psi = 0 \text{ on } \partial D$$

where D is simply-connected. Because the free boundary near the singular points of blow-up limit (1) is always cross-like, i.e., they consist of sectors with the opening angle  $\frac{\pi}{2}$  which is independent of the values of  $\lambda_1, \lambda_2 > 0$ .

**Lemma 3.4.** Let  $\psi$  be a solution of (3.1) and

$$S^{\psi}(x^0, r) \ge \frac{r^2}{\delta}$$

for some  $0 < \delta \leq \delta_0$ ,  $0 < r \leq r_0$ ,  $\sup_{B_3(x_0)} |\psi| \leq M$  and  $\psi(x^0) = |\nabla \psi(x^0)| = 0$  then:

(1) There exists a second order homogeneous harmonic polynomial  $p^{x^0,\psi} = p$  such that for each  $\alpha \in (0, \frac{1}{2})$  and each  $\beta \in (0, 1)$ ,

(3.21) 
$$\left\| \frac{\psi(x^0 + sx)}{\sup_{B_s(x^0)} |\psi|} - p \right\|_{C^{1,\beta}(\overline{B}_1)} \le C(M,\alpha,\beta) \left( \frac{\delta}{1 + \delta \log(\frac{r}{s})} \right)^{\alpha}$$

(2) The set  $\{\psi = 0\} \cap B_{r_0}(x^0)$  consists of two C<sup>1</sup>-curves intersecting each other at right angles at  $x^0$ .

It suffices for us to consider the generalized Newtonian potential

(3.22) 
$$-\Delta z = \mathbf{I}_{\mathbf{C}} - 2\Omega, \quad \mathbf{C} := \{(r,\theta) \subset \mathbf{R}^2 | r > 0, 0 < \theta < \frac{\pi}{2}\}$$

Note that the right-hand side of (3.22) is of class  $L^{\infty}$  and homogeneous of degree zero, by [20], z is uniquely representable in the form of

$$z(x) = p(x) \log |x| + |x|^2 \phi(\frac{x}{|x|}) + q_1(x), \quad x \in \mathbf{R}^2,$$

where p is a homogeneous harmonic polynomial of degree two,  $\phi \in C^1(S^1)$  and  $q_1(x) = a \cdot x + b$ . Suppose  $z(0) = \nabla z(0) = 0$ . It is sufficient to compute in the form of class  $p(x) \log |x| + |x|^2 \phi(\frac{x}{|x|})$ . Consider (3.22)

$$-(\partial_{rr} + \frac{1}{r}\partial_r + \frac{1}{r^2}\partial_{\theta\theta})z = \begin{cases} 1 - 2\Omega & 0 < \theta < \frac{\pi}{2} \\ -2\Omega & \frac{\pi}{2} < \theta < 2\pi \end{cases}$$

in the polar coordinate in which z is of the form

$$z(x_1, x_2) = \left(A(x_1^2 - x_2^2) + 2Bx_1x_2\right)\log|x| + |x|^2\phi(\frac{x}{|x|})$$
$$= r^2\log r \left(A\cos 2\theta + B\sin 2\theta\right) + r^2\phi(\theta)$$

with constants  $A, B \in \mathbf{R}$ . A straightforward computation shows that

$$-4(A\cos 2\theta + B\sin 2\theta) - \phi''(\theta) - 4\phi(\theta) = \begin{cases} 1 - 2\Omega & 0 < \theta < \frac{\pi}{2} \\ -2\Omega & \frac{\pi}{2} < \theta < 2\pi \end{cases}$$

Consider the Fourier series

$$\phi(\theta) = \frac{A_0}{2} + \sum_{n=1}^{\infty} A_n \cos n\theta + B_n \sin n\theta.$$

For  $n \neq 2$ , one has

$$A_0 = \frac{1 - 2\Omega}{2} - 3\Omega, \ A_n = \frac{1}{n\pi} \sin n\frac{\pi}{2}, \ B_n = \frac{1}{n\pi} \left(1 - \cos n\frac{\pi}{2}\right), \ n \ge 1, \ n \ne 2$$

and

$$A = 0, \quad B = \frac{1}{2\pi}.$$

Note that  $r^2(A_2 \cos 2\theta + B_2 \sin 2\theta)$  is a harmonic function. We let

(3.23) 
$$z(r,\theta) = r^2 \left( \frac{1-8\Omega}{4} + \frac{1}{\pi} \sum_{n \ge 1, n \ne 2} \frac{1}{n} \left( \sin \frac{n\pi}{2} \cos n\theta + (1-\cos \frac{n\pi}{2}) \sin n\theta \right) \right) + \frac{r^2 \log r}{2\pi} \sin 2\theta - \frac{1}{4\pi} r^2 \sin 2\theta$$

**Lemma 3.5.** The potential z given in (3.23) is the unique solution of equation (3.22) satisfying

(1)  $z(0) = |\nabla z(0)| = 0,$ (2)  $\lim_{x \to \infty} \frac{z(x)}{|x|^3} = 0,$ (3)  $\Pi(z) = 0,$ (4)  $\Pi(z_{\frac{1}{2}}) = \frac{\log 2}{\pi} x_1 x_2,$ (5)  $\tau(z_{\frac{1}{2}}) = \frac{\log 2}{\pi},$ 

where  $\mathbb{P}$  is the space of second-order homogeneous harmonic polynomials and  $\Pi$ :  $W^{2,2}(B_1) \to \mathbb{P}$  is a projection such that  $\Pi(u)$  is the unique minimizer of

$$p \to \int_{B_1} \left| D^2 v - D^2 p \right|^2$$

on  $\mathbb{P}$  and  $\tau(u) \ge 0$  is defined by  $\Pi(u) = \tau(u)p, \ p \in \mathbb{P}, \ \sup_{B_1} |p| = 1.$ 

*Proof.* Suppose  $h := \Pi(z)$ . Since h is harmonic we let

$$D^2h = \begin{pmatrix} a & b \\ b & -a \end{pmatrix}.$$

Since h is the minimizer of the mapping  $p \to \int_{B_1} |D^2 v - D^2 p|^2$ , we have

$$0 = \partial_b \int_{B_1} \left| D^2 v - D^2 h \right|^2 = 4b - 4 \int_{B_1} \partial_{12} z$$
  
=  $4b - \int_0^1 \int_{S^1} \left( \frac{\sin 2\theta}{2} \left( \partial_{rr} z + \frac{1}{r^2} \partial_{\theta\theta} \right) z(r,\theta) - \frac{1}{r} \partial_{r\theta} z(r,\theta) \right)$   
+  $\frac{1}{r^2} \partial_{\theta} z(r,\theta) + \frac{\sin 2\theta}{2r} \partial_r z(r,\theta) r dr d\theta$ 

22

Being aware of that only  $\frac{r^2 \log r}{2\pi} \sin 2\theta - \frac{1}{4\pi}r^2 \sin 2\theta$  provides non-zero integrals, we have b = 0 since

$$\int_{B_1} \partial_{12} \left( \frac{\log(x_1^2 + x_2^2)}{2\pi} x_1 x_2 - \frac{1}{2\pi} x_1 x_2 \right) dx_1 dx_2$$
  
=  $-\frac{1}{2} + \frac{1}{2\pi} \int_{B_1} \left( 2 + \log(x_1^2 + x_2^2) - \frac{4x_1^2 x_2^2}{(x_1^2 + x_2^2)^2} \right) dx_1 dx_2 = 0,$ 

while a = 0 is obtained similarly by considering

$$0 = \partial_a \int_{B_1} \left| D^2 v - D^2 h \right|^2 = 4a.$$

Therefore one has  $h := \Pi(z) = 0$  due to h is a second-order polynomial. Moreover, we have

$$\frac{z(sx_1, sx_2)}{s^2} = z(x_1, x_2) - \frac{x_1x_2}{\pi} \log s.$$

Then it follows

$$\Pi(z_{\frac{1}{2}}) = \Pi(z) - \Pi\left(\frac{x_1 x_2 \log\left(\frac{1}{2}\right)}{\pi}\right) = \frac{\log 2}{\pi} x_1 x_2.$$

If  $z_1$  and  $z_2$  are solutions of (3.23), the  $\eta := z_1 - z_2$  is a second-order harmonic polynomial then  $|\nabla \eta(0)| = \eta(0) = 0$  and  $\Pi(\eta) = 0$  imply  $z_1 = z_2$ , namely z is the unique solution.

**Lemma 3.6.** Suppose  $\psi$  solves (3.1),  $x_0 \in \mathcal{S}^u$  and  $\sup_{B_3(x_0)} |u| \leq M < +\infty$ . Then

$$\left(\int_{B_1} \left| D^2 \frac{u(x^0 + rx)}{r^2} - D^2 \Pi \left( \frac{u(x^0 + rx)}{r^2} \right) \right|^p \right)^{\frac{1}{p}} \le C(M, p)$$

and

$$\left\|\frac{u(x^0+rx)}{r^2} - \Pi\left(\frac{u(x^0+rx)}{r^2}\right)\right\|_{C^{1,\beta}(\overline{B}_1)} \le C(M,\beta).$$

**Lemma 3.7.** For each  $\epsilon > 0, M < +\infty, \alpha \in [1, +\infty)$  and  $\beta \in (0, 1)$ , there exist  $r_0, \delta > 0$  with the following property: suppose that  $0 < r \leq r_0$  and that  $\psi$  is a solution of (3.1) satisfying  $\sup_{B_2(x_0)} |u| \leq M$ ,  $u(x) = |\nabla u(x)| = 0$  and

$$\mathcal{L}^{2}((\{u(x+r\cdot)>0\}\Delta\{x_{1}x_{2}>0\})\cap B_{1})\leq\delta.$$

Then

(3.24) 
$$\left\|\frac{u(x+r\cdot)}{r^2} - \Pi\left(\frac{u(x+r\cdot)}{r^2}\right) - z\right\|_{C^{1,\beta}(\overline{B}_1)} \le \epsilon.$$

*Proof.* Suppose that  $r_j \to 0$  such that

$$\mathcal{L}^n\left(\{u(x+r\cdot)>0\}\Delta\{x_1x_2>0\}\right)\to 0 \quad \text{as } j\to\infty$$

and

$$\frac{u_j(x^j + r_j \cdot)}{r_j^2} - \prod \left( \frac{u_j(x^j + r_j \cdot)}{r_j^2} \right) \to \widetilde{z} \text{ strongly in } C_{loc}^{1,\beta}(\mathbf{R}^n) \text{ and weakly in } W_{loc}^{2,\alpha}(\mathbf{R}^n)$$

Now consider N be the Newtonian potential of  $\Delta u_j$ , i.e.,

$$\widetilde{N}(y) := \frac{1}{2\pi} \int_{\mathbf{R}^2} \log |y - \xi| \Delta u_j(\xi) d\xi$$

Let  $N(y) = \widetilde{N}(y) - \widetilde{N}(x^j) - \nabla \widetilde{N}(x^j)(y - x^j)$  then  $h(y) = u_j(y) - N(y)$  is a harmonic function. We have

$$|u_j(y) - N(y) - \mathcal{D}^2 h(x^j)(y - x^j)(y - x^j)| \le C |x^j - y|^3$$
 in  $B_{\frac{1}{2}}(x^j)$ .

For the scaled function  $v_j(y) = \frac{\psi_j(x^j + r_j y)}{r_j^2}$ ,  $N_j(y) = \frac{N(x^j + r_j y)}{r_j^2}$  and  $p_j(y) = \mathcal{D}^2 h(x^j) y \cdot y$ , we obtain

$$|v_j(y) - N_j(y) - p_j(y)| \le Cr_j |y|^3$$
 in  $B_{\frac{1}{2}}(x^j)$ .

Thus

$$v_j - \Pi(v_j) = N_j - \Pi(N_j) + o(1), \quad j \to \infty$$

Therefore  $N_j$  converges locally to  $N_0$ , where

 $-\Delta N_0 = \mathbf{I}_C - 2\Omega, \ N_0(0) = \nabla N_0(0) = 0, \text{ and } N_0 - \Pi(N_0) = \widetilde{z}.$ 

Then it suffices to show that  $N_0(y) = o(|y|^3)$  as  $|y| \to \infty$  since one would conclude  $\tilde{z} = N_0 - \Pi(N_0) = z$  due to the uniqueness result given in Lemma 3.5. Since  $\mathcal{D}^2 N_0 \in BMO$  so that

$$\int_{B_1} \left| \frac{\mathcal{D}^2\left(N_0(Ry)\right) - \overline{\mathcal{D}^2\left(N_0(R\cdot)\right)}}{\sup_{B_R} \left| \mathcal{D}^2 N_0 \right|} \right|^2 dy \le C \frac{R^4}{\sup_{B_R} \left| \mathcal{D}^2 N_0 \right|^2}$$

for all  $R \in (0, +\infty)$ , where  $\overline{\mathcal{D}^2(N_0(R\cdot))}$  denotes the mean value of  $\mathcal{D}^2(N_0(R\cdot))$  on  $B_1$ . Therefore we have a sun-sequence  $\frac{N_0(R_k\cdot)}{\sup_{B_{R_k}} |\mathcal{D}^2 N_0|}$  converges to a degree 2 homogeneous harmonic polynomial as  $R_k \to \infty$ . Assume

$$\limsup_{|y| \to \infty} \frac{|N_0(y)|}{|y|^3} > 0.$$

Since  $\Delta(N_0 - z) = 0$  and

$$\limsup_{|y| \to \infty} \frac{|N_0(y) - z(y)|}{|y|^3} > 0,$$

 $N_0 - z$  must be a harmonic polynomial of degree  $m \ge 3$  which is a contradiction.  $\Box$ 

Based on Lemma 3.7, Lemma 3.4 is proved by the same approach as Theorem A in [1], hence we omit it here.

Now we are in the position to prove the main theorems.

Proof of Theorem 1.1. The proof concerned with the regular points is standard hence we just outline it here. Due to the standard elliptic regularity theorem, we have  $u \in C^{1,\alpha}$  for  $\alpha < 1$  due to (1.6). For the free boundary  $\partial D$  near a regular point  $x_0 \in \partial D$ , i.e.,  $\nabla u(x_0) \neq 0$  and  $u(x_0) = 0$ , by applying the implicit function theorem we have  $\partial D$  is locally a  $C^{1,\alpha}$  curve near  $x_0$ , namely there exists r > 0 such that  $\partial D \cap B_r(x_0)$  is parameterized by the graph of a  $C^{1,\alpha}$  function. Following the idea in the proof of Theorem 3.1 of [21], we apply the hodograph transform and the local real-analyticity of the curve follows.

For the structure of singularity, the proof follows from Lemma 3.1 and Lemma 3.4.  $\Box$ 

Proof of Theorem 1.3. Let  $\lambda_1 = 1 - 2\Omega$  and  $\lambda_2 = 2\Omega$  in Theorem 1.1. Suppose that there is no stagnation point on the boundary, namely  $\nabla \psi \neq 0$  on  $\partial D$ . Since  $\partial D$  is a connected component of level set  $\{\psi = 0\}$  and  $\psi \in C^{1,\alpha}$  due to standard elliptic regularity theorem,  $\partial D$  is a  $C^{1,\alpha}$  curves, which is indeed locally real-analytic due to Theorem 1.1. Since the curve  $\partial D$  is real-analytic hence  $C^{\infty}$ ,  $\partial D$  is a  $C^{\infty}$  smooth manifold due to the fundamental theorem concerned partition of unity.

For the singular points, by Lemma 3.3 and Lemma 3.4, it suffices to exclude the case (2) provided  $\partial D$  is Lipschitz. Indeed, since  $\partial D$  is Lipschitz, for any  $x_0 \in \partial D$ , there exists r > 0 such that  $B_r(x_0) \cap \partial D$  can be parameterized by the graph of a Lipschitz function. Without loss of generality, we denote by the curve (s, f(s)) locally, where (0, f(0)) corresponds to the singular point. Note that the curve is  $C^1$  except s = 0, the tangential vector is given by T(s) = (1, f'(s)) therefore it is sufficient to show that T(0+) and T(0-) can not be co-linear. Otherwise, since the first component in the tangential vector is 1,  $f'(s) \to \pm \infty$  as  $s \to 0\pm$ , which is in contradict with f'(s) is uniformly bounded due to the Lipschitz continuity. There must be  $T(0^+) \cdot T(0^-) = 0$ , since the corner is 90°. The proof is complete.

#### References

<sup>[1]</sup> John Andersson, Henrik Shahgholian, and Georg S. Weiss. Uniform regularity close to cross singularities in an unstable free boundary problem. *Comm. Math. Phys.*, 296(1):251–270, 2010.

<sup>[2]</sup> John Andersson, Henrik Shahgholian, and Georg S. Weiss. On the singularities of a free boundary through Fourier expansion. *Inventiones Mathematicae*, 187(3):535–587, 2012.

- [3] A. L. Bertozzi and P. Constantin. Global regularity for vortex patches. Comm. Math. Phys., 152(1):19-28, 1993.
- [4] Jacob Burbea. Motions of vortex patches. Letters in Mathematical Physics. A Journal for the Rapid Dissemination of Short Contributions in the Field of Mathematical Physics, 6(1):1–16, 1982.
- [5] Luis A. Caffarelli. The regularity of free boundaries in higher dimensions. Acta Math., 139(3-4):155–184, 1977.
- [6] Angel Castro, Diego Córdoba, and Javier Gómez-Serrano. Uniformly rotating analytic global patch solutions for active scalars. Ann. PDE, 2(1), 2016.
- [7] Jean-Yves Chemin. Persistance de structures géométriques dans les fluides incompressibles bidimensionnels.(french) [[persistence of geometric structures in two-dimensional incompressible fluids]]. Ann. Sci. École Norm. Sup, (4):517–542, 1993.
- [8] Jean-Yves Chemin. Perfect incompressible fluids, volume 14 of Oxford Lecture Series in Mathematics and its Applications. The Clarendon Press, Oxford University Press, New York, 1998. Translated from the 1995 French original by Isabelle Gallagher and Dragos Iftimie.
- [9] Francisco de la Hoz, Taoufik Hmidi, Joan Mateu, and Joan Verdera. Doubly connected V-states for the planar Euler equations. SIAM Journal on Mathematical Analysis, 48(3):1892–1928, 2016.
- [10] Tarek M. Elgindi and In-Jee Jeong. On singular vortex patches, I: Well-posedness issues. Mem. Amer. Math. Soc., 283(1400):v+89, 2023.
- [11] Alexandre Ern and Jean-Luc Guermond. Theory and practice of finite elements, volume 159 of Applied Mathematical Sciences. Springer-Verlag, New York, 2004.
- [12] Alessio Figalli, Xavier Ros-Oton, and Joaquim Serra. Generic regularity of free boundaries for the obstacle problem. Publ. Math. Inst. Hautes Études Sci., 132:181–292, 2020.
- [13] Alessio Figalli and Joaquim Serra. On the fine structure of the free boundary for the classical obstacle problem. *Invent. Math.*, 215(1):311–366, 2019.
- [14] C Garcia and S Haziot. Global bifurcation for corotating and counter-rotating vortex pairs. arXiv:2204.11327, 2022.
- [15] Javier Gómez-Serrano, Jaemin Park, Jia Shi, and Yao Yao. Symmetry in stationary and uniformly rotating solutions of active scalar equations. *Duke Mathematical Journal*, 170(13):2957– 3038, 2021.
- [16] Zineb Hassainia, Nader Masmoudi, and Miles H. Wheeler. Global bifurcation of rotating vortex patches. Communications on Pure and Applied Mathematics, 73(9):1933–1980, 2020.
- [17] Taoufik Hmidi and Joan Mateu. Bifurcation of rotating patches from Kirchhoff vortices. Discrete and Continuous Dynamical Systems. Series A, 36(10):5401–5422, 2016.
- [18] Taoufik Hmidi and Joan Mateu. Degenerate bifurcation of the rotating patches. Advances in Mathematics, 302:799–850, 2016.
- [19] Taoufik Hmidi, Joan Mateu, and Joan Verdera. Boundary regularity of rotating vortex patches. Archive for Rational Mechanics and Analysis, 209(1):171–208, 2013.
- [20] Lavi Karp and Avmir S. Margulis. Newtonian potential theory for unbounded sources and applications to free boundary problems. J. Anal. Math., 70:1–63, 1996.
- [21] D. Kinderlehrer, L. Nirenberg, and J. Spruck. Regularity in elliptic free boundary problems. J. Analyse Math., 34:86–119 (1979), 1978.
- [22] Alexander Kiselev and Chao Li. Global regularity and fast small-scale formation for Euler patch equation in a smooth domain. *Comm. Partial Differential Equations*, 44(4):279–308, 2019.

- [23] Alexander Kiselev and Xiaoyutao Luo. Illposedness of  $C^2$  vortex patches. Arch. Ration. Mech. Anal., 247(3):Paper No. 57, 49, 2023.
- [24] Alexander Kiselev and Vladimir Šverák. Small scale creation for solutions of the incompressible two-dimensional Euler equation. Ann. of Math., 180(3):1205–1220, 2014.
- [25] Carlo Marchioro and Mario Pulvirenti. Mathematical theory of incompressible nonviscous fluids, volume 96 of Applied Mathematical Sciences. Springer-Verlag, New York, 1994.
- [26] R. Monneau and G. S. Weiss. An unstable elliptic free boundary problem arising in solid combustion. Duke Mathematical Journal, 136(2):321–341, 2007.
- [27] Edward A. Overman, II. Steady-state solutions of the Euler equations in two dimensions. II. Local analysis of limiting V-states. SIAM J. Appl. Math., 46(5):765–800, 1986.
- [28] Jaemin Park. Quantitative estimates for uniformly-rotating vortex patches. arXiv:2010.06754, 2020.
- [29] Philippe Serfati. Une preuve directe d'existence globale des vortex patches 2D. C. R. Acad. Sci. Paris Sér. I Math., 318(6):515–518, 1994.
- [30] Bruce Turkington. On steady vortex flow in two dimensions. I, II. Comm. Partial Differential Equations, 8(9):999–1030, 1031–1071, 1983.
- [31] Joan Verdera. The global regularity of vortex patches revisited. Adv. Math., 416:Paper No. 108917, 15, 2023.
- [32] H. M. Wu, E. A. Overman, II, and N. J. Zabusky. Steady-state solutions of the Euler equations in two dimensions: rotating and translating V-states with limiting cases. I. Numerical algorithms and results. J. Comput. Phys., 53(1):42–71, 1984.
- [33] V Yudovich. Non-stationary flows of an ideal incompressible fluid. Z. Vycisl. Mat. i. Mat. Fiz (Russian), 3:1032–1066, 1963.

<sup>1</sup> INSTITUTE OF MATHEMATICS, UNIVERSITY OF AUGSBURG, AUGSBURG, 86159, GERMANY,
 <sup>2</sup> SCHOOL OF MATHEMATICS SCIENCE, TIANJIN NORMAL UNIVERSITY, TIANJIN, 300387, CHINA

Email address: yuchen.wang@uni-a.de

<sup>3</sup> School of Mathematics and Statistics, Huazhong University of Science and Technology, Wuhan, 430074, China

Email address: guanghuizhang@hust.edu.cn

<sup>4</sup> Chern Institute of Mathematics and LPMC, Nankai University, Tianjin, 300071, China

Email address: zhouml123@nankai.edu.cn