

Lyapunov Exponents for Open Billiard Flows

Amal Al Dowais¹

Department of Mathematics and Statistics, University of Western Australia, Crawley 6009 WA, Australia
E-mail: amal.al dowais@research.uwa.edu.au

Luchozar Stoyanov

Department of Mathematics and Statistics, University of Western Australia, Crawley 6009 WA, Australia
E-mail: luchozar.stoyanov@uwa.edu.au

Abstract. In this paper we prove that the positive Lyapunov exponents for the billiard flow in an open billiard in \mathbb{R}^d ($d \geq 3$) are all equal. We should stress that we do not make any particular assumptions about the shape and size of the components of our obstacles – they are just assumed to be strictly convex and compact with C^3 boundaries and satisfy the so called no eclipse condition.

MSC: Primary: 37D20, 37D25; Secondary: 37D40

Keywords: Lyapunov exponents, open billiard flow, Oseledets subspaces

1 Introduction

Let K be a subset of \mathbb{R}^d ($d \geq 3$) of the form

$$K = K_1 \cup K_2 \cup \dots \cup K_{k_0},$$

where K_i are compact strictly convex disjoint domains in \mathbb{R}^d with C^3 boundaries $\Gamma_i = \partial K_i$ and $k_0 \geq 3$. Set $\Omega = \overline{\mathbb{R}^d} \setminus K$. We assume that K satisfies the following *no-eclipse condition*:

$$(H) \quad \begin{cases} \text{for every pair } K_i, K_j \text{ of different connected components of } K \\ \text{the convex hull of } K_i \cup K_j \text{ has no common points with any other} \\ \text{connected component of } K. \end{cases}$$

With this condition, the *billiard flow* ϕ_t defined on the *sphere bundle*

$$S(\Omega) = \{(q, v) : q \in \Omega, v \in \mathbb{S}^{d-1}\}$$

in the standard way is called an *open billiard flow*. It has singularities, however its restriction to the *non-wandering set* Λ has only simple discontinuities at reflection points.

In this paper we prove that all positive Lyapunov exponents of the billiard flow with respect to any invariant probability measure on Λ are equal. Equivalently, all positive Lyapunov exponents of the related billiard ball map are equal.

While this result may seem surprising, it has to be noted that conjectures of somethings similar happening, although in a more special and different situation, have been made before. E.g. Chernov conjectured in [Ch3] (see p.17 there; see also [Ch2]) that all positive Lyapunov exponents for a periodic Lorentz gas with spherical scatterers should be the same. We are not aware of any positive or negative results concerning this conjecture. However, as Chernov mentioned in [Ch3], it had been shown both theoretically and numerically that for 3D random Lorentz gases with a

¹Department of Mathematics, College of Science and Arts, Najran University, Najran, Saudi Arabia; E-mail: amalduas@nu.edu.sa

random configuration of scatterers the two positive Lyapunov exponents are different (see [LBS], [DP] and [BL]). There are probably some more recent results in this direction.

Here we do not assume that our scatterers K_i are spherical, in fact we do not assume anything about their shape and size except that they are strictly convex and compact with C^3 boundaries, however we do assume the no-eclipse condition (H) which is apparently significant. It is an interesting question what happens if we drop the latter assumption: are there examples of obstacles K in \mathbb{R}^d ($d \geq 3$) as above, however without assuming the condition (H), where the positive Lyapunov exponents are not all the same?

The *non-wandering set* Λ for the flow ϕ_t is the set of those $x \in S(\Omega)$ such that the trajectory $\{\phi_t(x) : t \in \mathbb{R}\}$ is bounded. Notice that the natural projection of ϕ_t on the quotient space $S(\Omega)/\sim$, where \sim is the equivalence relation $(q, v) \sim (p, w)$ iff $q = p$ and $v = w$ or $q = p \in \partial K$ and v and w are symmetric with respect to $T_q(\partial K)$, is continuous. Moreover whenever both x and $\phi_t(x)$ are in the interior of $S(\Omega)$ and sufficiently close to Λ , the map $y \mapsto \phi_t(y)$ is smooth on a neighbourhood of x . It follows from results of Sinai ([Si1], [Si2]) that Λ is a hyperbolic set for ϕ_t , and it is easily seen that there exists an open neighbourhood U of Λ in $S(\Omega)$ such that $\Lambda = \bigcap_{t \in \mathbb{R}} \phi_t(U)$. In fact, Λ is clearly the maximal compact ϕ_t -invariant subset of $S(\Omega)$. Thus, Λ is a basic set for ϕ_t and the classical theory of hyperbolic flows applies in the case under consideration (see e.g. Bowen [B] or part 4 in the monograph of Katok and Hasselblatt [KH]; see also [CP]).

The related *billiard ball map* B is defined in a natural way on

$$M = \{(q, v) \in \partial K \times \mathbb{S}^{d-1} : \langle \nu(q), v \rangle \geq 0\},$$

where $\nu(q)$ is the *outward unit normal* to ∂K at q . Namely, $B(q, v) = (p, w)$, where $p = q + tv \in \partial K$ for some $t > 0$, t is minimal with this property, and $w \in \mathbb{S}^{d-1}$ is the reflected direction of the corresponding billiard trajectory at p , namely $w = v - 2\langle v, \nu(p) \rangle \nu(p)$. The set $M_0 = M \cap \Lambda$ is the *non-wandering set* of the billiard ball map B . It is well-known (see below) that $B : M_0 \rightarrow M_0$ is naturally conjugate to a transitive sub-shift of finite type so the Multiplicative Ergodic Theorem (see Sect. 2) applies to it.

Given an invariant probability measure μ on M_0 , let

$$\lambda_k < \lambda_{k-1} < \dots < \lambda_2 < \lambda_1$$

be the positive Lyapunov exponents of B . Then $-\lambda_1 < -\lambda_2 < \dots < -\lambda_{k-1} < -\lambda_k$ are the negative Lyapunov exponents of B .

Here we prove the following

Theorem 1.1. *All positive Lyapunov exponents of the billiard ball map $B : M_0 \rightarrow M_0$ are equal. That is, $k = 1$ and λ_1 has multiplicity d .*

Equivalently, all positive Lyapunov exponents of the time-one map of the billiard flow ϕ_t on Λ with respect to any invariant probability measure are equal.

2 Preliminaries

2.1 Billiard ball map vs billiard flow

For $x \in \Lambda$ and a sufficiently small $\epsilon > 0$ be the (strong) *stable* and *unstable manifolds* of size ϵ **for the billiard flow** are defined by

$$\widetilde{W}_\epsilon^s(x) = \{y \in M : d(\phi_t(x), \phi_t(y)) \leq \epsilon \text{ for all } t \geq 0, d(\phi_t(x), \phi_t(y)) \rightarrow_{t \rightarrow \infty} 0\},$$

$$\widetilde{W}_\epsilon^u(x) = \{y \in M : d(\phi_t(x), \phi_t(y)) \leq \epsilon \text{ for all } t \leq 0, d(\phi_t(x), \phi_t(y)) \rightarrow_{t \rightarrow -\infty} 0\}.$$

Set $\widetilde{E}^u(x) = T_x \widetilde{W}_\epsilon^u(x)$ and $\widetilde{E}^s(x) = T_x \widetilde{W}_\epsilon^s(x)$. For any $x = (q, v) \in M_0$ the stable/unstable manifolds for the billiard ball map B are defined similarly:

$$W_\epsilon^s(x) = \{y \in M_0 : d(B^n(x), B^n(y)) \leq \epsilon \text{ for all } n \in \mathbb{N}, d(B^n(x), B^n(y)) \rightarrow_{n \rightarrow \infty} 0\},$$

$$W_\epsilon^u(x) = \{y \in M_0 : d(B^{-n}(x), B^{-n}(y)) \leq \epsilon \text{ for all } n \in \mathbb{N}, d(B^{-n}(x), B^{-n}(y)) \rightarrow_{n \rightarrow \infty} 0\},$$

where \mathbb{N} is the set of positive integers.

Given $x = (q, v) \in M_0$ and a small $t_0 > 0$, set $y = (q + t_0 v, v)$. For a small $\epsilon > 0$ and a sufficiently small $\delta > 0$ (depending on ϵ) the map

$$\Phi : W_\delta^s(x) \longrightarrow \widetilde{W}_\epsilon^s(y) \quad (2.1)$$

such that $\Phi(z, w) = (z + tw, w)$ for all $(z, w) \in W^u(x)$, where $t = t(z, w) > 0$, is a local diffeomorphism. In the same way there is a local diffeomorphism from $W_\delta^s(x)$ to $\widetilde{W}_\epsilon^s(y)$. Moreover $D\Phi(x) : T_x M_0 \longrightarrow T_y \Lambda$ is an isomorphism so that $D\Phi(x)(E^u(x)) = \widetilde{E}^u(y)$ and $D\Phi(x)(E^s(x)) = \widetilde{E}^s(y)$.

It is known (see [Si1], [Si2]) that $\widetilde{W}_\epsilon^s(y)$ has the form $\widetilde{W}_\epsilon^s(y) = \widetilde{Y}$, where

$$\widetilde{Y} = \{(p, \nu_Y(p)) : p \in Y\}$$

for some smooth $(d-1)$ -dimensional surface Y in \mathbb{R}^d containing the point y such that Y is strictly concave with respect to the unit normal field ν_Y , i.e. the curvature of Y is strictly negative. In a similar way one can describe $\widetilde{W}_\epsilon^u(y)$ using a strictly convex local surface.

Returning to the map (2.1), let $x_1 = (q_1, v_1) = B(x)$, where $q_1 = q + dv$ for some $d > 0$. Define $y_1 = (q_1 + t_1 v_1, v_1)$ for some small $t_1 > 0$ (e.g. $t_1 = t_0$). Then there is a local diffeomorphism (assuming again that $\epsilon > 0$ and $\delta > 0$ are sufficiently small) $\Phi_1 : W_\delta^s(x_1) \longrightarrow \widetilde{W}_\epsilon^s(y_1)$ defined as above. As above, $\widetilde{W}_\epsilon^s(y_1) = \widetilde{Y}_1$, where $\widetilde{Y}_1 = \{(p_1, \nu_Y(p_1)) : p_1 \in Y_1\}$ for some smooth $(d-1)$ -dimensional surface Y_1 in \mathbb{R}^d containing the point y_1 such that Y_1 is strictly concave with respect to the unit normal field ν_{Y_1} . Setting $t = d + t_1$, the following two diagrams are commutative:

$$\begin{array}{ccc} W^s(x) & \xrightarrow{B} & W^s(x_1) \\ \downarrow \Phi & & \downarrow \Phi_1 \\ \widetilde{W}^s(y) = \widetilde{Y} & \xrightarrow{\phi_t} & \widetilde{W}^s(y_1) = \widetilde{Y}_1 \end{array}, \quad \begin{array}{ccc} E^s(x) & \xrightarrow{DB(x)} & E^s(x_1) \\ \downarrow D\Phi & & \downarrow D\Phi_1 \\ E^s(y) & \xrightarrow{D\phi_t(y)} & E^s(y_1) \end{array}$$

For an open billiard, the billiard ball map $B : M_0 \longrightarrow M_0$ is naturally isomorphic to a transitive subshift of finite type. Here we briefly describe the natural conjugacy.

Let $k_0 \geq 2$ be as in Sect.1 and let $A = (A(i, j))_{i, j=1}^{k_0}$ be the $k_0 \times k_0$ matrix such that $A(i, j) = 1$ if $i \neq j$ and $A(i, i) = 0$ for all i, j . Consider the symbolic space

$$\Sigma_A = \{(i_j)_{j=-\infty}^{\infty} : 1 \leq i_j \leq k_0, A(i_j, i_{j+1}) = 1 \text{ for all } j\},$$

with the product topology and the *shift map* $\sigma : \Sigma_A \longrightarrow \Sigma_A$ given by $\sigma((i_j)) = ((i'_j))$, where $i'_j = i_{j+1}$ for all j . Given $0 < \theta < 1$, consider the *metric* d_θ on Σ_A defined by $d_\theta(\xi, \eta) = 0$ if $\xi = \eta$ and $d_\theta(\xi, \eta) = \theta^m$ if $\xi_i = \eta_i$ for $|i| < m$ and m is maximal with this property (see e.g. [B] or [PP] for general information on symbolic dynamics). Now define $R : M_0 \longrightarrow \Sigma_A$ by $R(x) = \{i_j\}_{j \in \mathbb{Z}}$ so that $\text{pr}_1(B^j(x)) \in K_{i_j}$ for all $j \in \mathbb{Z}$. It is well-known (see e.g. [I] or [PS]) that this is well-defined bijection which defines a conjugacy between $B : M_0 \longrightarrow M_0$ and $\sigma : \Sigma_A \longrightarrow \Sigma_A$, namely $R \circ B = \sigma \circ R$. Moreover, with an appropriate choice of $\theta \in (0, 1)$, R is a homeomorphism.

2.2 Multiplicative Ergodic Theorem

Let f be an invertible transitive subshift of finite type over a bilateral symbol space X and let μ be an invariant probability measure. We will assume in addition that μ is ergodic. Let L be an invertible continuous linear cocycle over f acting on a continuous \mathbb{R}^d -bundle E over X . Thus, $L(x) : E(x) \rightarrow E(f(x))$ is a linear map for each $x \in X$ and

$$L^n(x) = L(f^{n-1}(x)) \circ L(f^{n-2}(x)) \circ \dots \circ L(f(x)) \circ L(x) : E(x) \rightarrow E(f^n(x))$$

for every integer $n \geq 1$.

Given an integer $p = 1, \dots, d$, let $\text{Gr}_p(\mathbb{R}^d)$ be the *Grassman manifold* of the linear subspaces of \mathbb{R}^d of dimension p endowed with the usual distance $d(U, V)$ between subspaces $U, V \in \text{Gr}_p(\mathbb{R}^d)$ defined by

$$d(U, V) = \max\{|\langle u, w \rangle| : u \in U, w \in V^\perp, \|u\| = \|w\| = 1\}.$$

The following is Oseledets' Multiplicative Ergodic Theorem stated under the above assumptions (see e.g. [Ar], [BP] or [V] for related detailed exposition and proofs; see also [BPS]).

Theorem 2.1. (Multiplicative Ergodic Theorem). *There exists a subset \mathcal{L} of X with $\mu(\mathcal{L}) = 1$ such that:*

- (a) *For all $x \in \mathcal{L}$ there exists $N(x) = \lim_{n \rightarrow \infty} (L^n(x)^t L^n(x))^{1/2n}$.*
- (b) *There exist an integer $k \geq 1$ such that the operator (matrix) $N(x)$ has k distinct eigenvalues $t_k(x) < \dots < t_1(x)$ for all $x \in \mathcal{L}$.*
- (c) *There exist numbers $t_k < t_{k-1} < \dots < t_2 < t_1$ such that $(t_i^{(n)}(x))^{1/n} \rightarrow t_i$ for all $x \in \mathcal{L}$ and all $i = 1, \dots, k$.*
- (d) *For all $x \in \mathcal{L}$ and every $j = 1, \dots, k$ the dimension $\dim(E_j^{(n)}(x)) = m_j$ is constant and there exists $\lim_{n \rightarrow \infty} E_j^{(n)}(x) = E_j(x)$ in $\text{Gr}_{m_j}(\mathbb{R}^d)$.*
- (e) *For all $x \in \mathcal{L}$ and every $j = 1, \dots, k$ we have*

$$\lim_{n \rightarrow \pm\infty} \frac{1}{n} \log \|L^n(x) \cdot u\| = \log t_j \quad , \quad u \in E_j(x) \setminus \{0\}$$

for all $j = 1, \dots, k$.

The numbers

$$\lambda_k = \log t_k < \lambda_{k-1} = \log t_{k-1} < \dots < \lambda_2 = \log t_2 < \lambda_1 = \log t_1$$

are the (distinct) *Lyapunov exponents* of the cocycle L over f . The so called *Oseledets subspaces* $E_i(x)$ are invariant with respect to L .

Since the billiard ball map $B : M_0 \rightarrow M_0$ is conjugate to an invertible transitive subshift of finite type (see Sect. 2.1), the above theorem applies to it.

3 Derivative estimates for open billiards

Let $K \subset \mathbb{R}^d$ be as in Sect. 1. Set $d_0 = \min_{i \neq j} \text{dist}(K_i, K_j)$. Denote by $\text{pr}_1 : T(\mathbb{R}^d) \rightarrow \mathbb{R}^d$ the natural projection $\text{pr}_1(q, v) = q$. For any $x \in \Gamma = \partial K$ we will denote by $\nu(x)$ the *outward unit normal* to Γ at x .

Remark. Notice that the condition (H) implies the existence of a constant $\delta_0 > 0$ depending only on the obstacle K such that $\langle u, \nu(x) \rangle \geq \delta_0$ for any $(x, u) \in S^*(\Omega)$ whose backward and forward billiard trajectories both have common points with ∂K .

Since every K_i is strictly convex, the *curvature (shape) operator*

$$L_x : T_x(\partial K) \rightarrow T_x(\partial K) \quad , \quad L_x u = (\nabla_u \nu)(x),$$

defines a positive definite *second fundamental form* $\langle L_x u, u \rangle$. The *normal curvature* of ∂K at x in the direction of $u \in T_x(\partial K)$ ($\|u\| = 1$) is by definition $k(x, u) = \langle L_x u, u \rangle$. Clearly, $\|L_x u\| \geq k(x, u)$.

Given $\epsilon \in (0, d_0/2)$, set

$$S_\epsilon^*(\Omega) = \{x = (q, v) \in S^*(\Omega) : \text{dist}(q, \partial K) > \epsilon\} \quad , \quad \Lambda_\epsilon = \Lambda \cap S_\epsilon^*(\Omega).$$

In what follows in order to avoid ambiguity and unnecessary complications we will consider stable and unstable manifolds only for points x in $S_\epsilon^*(\Omega)$ or Λ_ϵ ; this will be enough for our purposes.

Here we derive a formula which is useful in getting estimates for $\|d\phi_t(x) \cdot u\|$ ($u \in E^u(x)$, $x \in \Lambda$), both from above and below. In a natural way this provides estimates for the derivatives of the billiard ball map B as well (see Sect. 2.1).

Fix for a moment a point $x_0 = (q_0, v_0) \in \Lambda_\epsilon$. If $\epsilon > 0$ is sufficiently small, as we remarked in Sect. 1.2, $\widetilde{W}_\epsilon^u(x_0)$ has the form (cf. [Si1],[Si2])

$$\widetilde{W}_\epsilon^u(x_0) = \{(x, \nu_X(x)) : x \in X\}$$

for some smooth hypersurface X in \mathbb{R}^d containing the point q_0 such that X is strictly convex with respect to the unit normal field ν_X . Let

$$D_q : T_q X \rightarrow T_q X \quad , \quad D_q(v) = (\nabla_v \nu_X)(q),$$

be the *curvature (shape) operator* of X at $q \in X$ defined by means of the normal field ν_X ; then the second fundamental form $\langle D_q(\xi), \xi \rangle$, $\xi \in T_q(X)$, is positive definite.

Given a point $q_0 \in X$, let $\gamma(x)$ be the *forward billiard trajectory* generated by $x = (q_0, \nu_X(q_0))$. Let $q_1(x), q_2(x), \dots$ be the reflection points of this trajectory and let $\xi_j(x) \in \mathbb{S}^{d-1}$ be the *reflected direction* of $\gamma(x)$ at $q_j(x)$.

Set $q_0(x) = q_0$, $t_0(x) = 0$ and denote by $t_1(x), t_2(x), \dots$ the times of the consecutive reflections of $\gamma(x)$ at ∂K . Then $t_j(x) = d_0(x) + d_1(x) + \dots + d_{j-1}(x)$, where $d_j(x) = \|q_{j+1}(x) - q_j(x)\|$, $0 \leq j$. Given $t \geq 0$, denote by $u_t(q_0)$ the *shift* of q_0 along the trajectory $\gamma(x)$ after time t . Set

$$X_t = \{u_t(q) : q \in X\}.$$

When $u_t(q_0)$ is not a reflection point of $\gamma(x)$, then locally near $u_t(q_0)$, X_t is a smooth convex $(d-1)$ -dimensional surface in \mathbb{R}^d with "outward" unit normal given by the *direction* $v_t(q_0)$ of $\gamma(x)$ at $u_t(q_0)$ (cf. [Si2]).

Fix for a moment $t > 0$ such that $t_m(x_0) < t < t_{m+1}(x_0)$ for some $m \geq 1$, and assume that $q_0(s)$, $0 \leq s \leq a$, is a C^3 curve on X with $q_0(0) = q_0$ such that for every $s \in [0, a]$ we have $t_m(x(s)) < t < t_{m+1}(x(s))$, where $x(s) = (q_0(s), \nu_X(q_0(s)))$. Assume also that $a > 0$ is so small that for all $j = 1, 2, \dots, m$ the reflection points $q_j(s) = q_j(x(s))$ belong to the same boundary component ∂K_{i_j} for every $s \in [0, a]$.

We will now estimate $\|d\phi_t(x_0) \cdot \dot{q}_0(0)\|$, where $\dot{q}(0) \in T_{q_0}X$.

Clearly $\phi_t(x(s)) = (p(s), v_m(x(s)))$, where $p(s)$, $0 \leq s \leq a$, is a C^3 curve on X_t . For brevity denote by $\gamma(s)$ the *forward billiard trajectory* generated by $(q_0(s), \nu(q_0(s)))$. Let $\xi_j(s) \in \mathbb{S}^{d-1}$ be the *reflected direction* of $\gamma(s)$ at $q_j(s)$ and let $\varphi_j(s)$ be the *angle* between $\xi_j(s)$ and the outward unit normal $\nu(q_j(s))$ to ∂K at $q_j(s)$. Let $\phi_t(x(s)) = (u_t(s), v_t(s))$, and let $t_j(s) = t_j(x(s))$ be the times of the consecutive reflections of the trajectory $\gamma(s)$ at ∂K . Set $d_j(s) = d_j(x(s)) = \|q_{j+1}(s) - q_j(s)\|$ ($0 \leq j \leq m-1$), $d_m(s) = \|p(s) - q_m(s)\|$, $t_0(s) = 0$ and $t_{m+1}(s) = t_m(s) + d_m(s)$.

Next, let $k_0(s)$ be the *normal curvature* of X at $q_0(s)$ in the direction of $\dot{q}_0(s)$. That is

$$k_0(s) = \frac{1}{\|\dot{q}_0(s)\|^2} \langle D_{q_0(s)}(\dot{q}_0(s)), \dot{q}_0(s) \rangle.$$

Similarly, for $j > 0$ let $k_j(s) > 0$ be the *normal curvature* of $X_{t_j(s)} = \lim_{t \searrow t_j(s)} X_t$ at $q_j(s)$ in the direction $e_j(s)$ ($\|e_j(s)\| = 1$) of $\lim_{t \searrow t_j(s)} \frac{d}{ds'} (u_t(s'))|_{s'=s}$, that is

$$k_j(s) = \langle D_{q_j(s)}(e_j(s)), e_j(s) \rangle.$$

Set $e_0(s) = \dot{q}_0(0)/\|\dot{q}_0(0)\|$ and for any $j \geq 0$ define $\ell_j(s) > 0$ by

$$[1 + d_j(s)\ell_j(s)]^2 = 1 + 2d_j(s)k_j(s) + (d_j(s))^2 \|D_{q_j(s)}(e_j(s))\|^2. \quad (3.1)$$

Finally, set

$$\delta_j(s) = \frac{1}{1 + d_j(s)\ell_j(s)}, \quad 0 \leq j \leq m. \quad (3.2)$$

Theorem 3.1. *For all $s \in [0, a]$ we have*

$$\|\dot{q}_0(s)\| = \|\dot{p}_0(s)\| \delta_0(s) \delta_1(s) \dots \delta_m(s). \quad (3.3)$$

This theorem was proved in [St1] in the case $d = 2$ and stated without proof in [St2] in the higher dimensional case. As we mentioned in [St2], the above formula can be derived from the more general study of the evolution of unstable fronts in multidimensional dispersing billiards in [BCST] (see Sect. 5 there). See also Ch. 3 in [ChM]. For completeness we provide a proof in the Appendix for all $d \geq 2$.

4 Proof of Theorem 1.1

Throughout we assume that μ is a B -invariant probability measure on M_0 . Fix a subset \mathcal{L} of M_0 of Lyapunov regular points with $\mu(\mathcal{L}) = 1$ and all the properties in Theorem 2.1. Let

$$\lambda_{2k} = -\lambda_1 < \lambda_{2k-1} = -\lambda_2 < \dots < \lambda_{k+1} = -\lambda_k < 0 < \lambda_k < \dots < \lambda_2 < \lambda_1$$

be the *Lyapunov spectrum* of dB with respect to μ . As before denote by $E_i(x)$ the *Oseledets subspace* corresponding to the Lyapunov exponent λ_i . The *stable bundle* $E^s(x)$ and the *unstable bundle* $E^u(x)$ ($x \in M_0$) are given by

$$E^s(x) = E_{2k}(x) \oplus E_{2k-1}(x) \oplus \dots \oplus E_{k+1}(x) \quad , \quad E^u(x) = E_k(x) \oplus E_{k-1}(x) \oplus \dots \oplus E_1(x).$$

Given a small $\epsilon > 0$, the *local stable manifold* $\widetilde{W}_\epsilon^s(x)$ and the *local unstable manifold* $\widetilde{W}_\epsilon^u(x)$ for the flow at $x \in \Lambda$ (see Sect. 2.1) have tangent spaces $T_x(\widetilde{W}_\epsilon^s(x)) = \widetilde{E}^s(x)$ and $T_x(\widetilde{W}_\epsilon^u(x)) = \widetilde{E}^u(x)$.

Consider the ‘stable’ linear cocycle

$$S(x) = dB(x) : E^s(x) \longrightarrow E^s(B(x)) \quad , \quad x \in M_0,$$

over the billiard ball map $B : M_0 \longrightarrow M_0$. Its Lyapunov exponents are

$$\lambda'_k = -\lambda_1 < \lambda'_{k-1} = -\lambda_2 < \dots < \lambda'_1 = -\lambda_k,$$

and its Oseledets subspaces are

$$E'_1(x) = E_{2k}(x), E'_2(x) = E_{2k-1}(x), \dots, E'_k(x) = E_{k+1}(x),$$

while its Oseledets flags are

$$F'_j(x) = E'_1(x) \oplus E'_2(x) \oplus \dots \oplus E'_j(x) = E_{2k}(x) \oplus E_{2k-1}(x) \oplus \dots \oplus E_{2k-j+1}(x)$$

for all $j = 1, \dots, k$ and $x \in \mathcal{L}$. Set $F'_0(x) = \{0\}$. The corresponding Oseledets subspaces and Oseledets flags for the flow will be denoted by $\widetilde{E}'_j(y)$ and $\widetilde{F}'_j(y)$, respectively.

Theorem 1.1 is an immediate consequence of the following

Theorem 4.1. *We have $k = 1$, i.e. B has only two non-zero Lyapunov exponents $\lambda_1 > 0$ and $\lambda_2 = -\lambda_1 < 0$, each with multiplicity d . That is, the linear cocycle S has only one Lyapunov exponent $\lambda_2 = -\lambda_1 < 0$ which has multiplicity d .*

Proof of Theorem 4.1. Assume that $k \geq 2$. Fix $\delta_0 \in (0, \epsilon/10)$ so small that

$$0 < \delta_0 < \frac{1}{10} \min\{\lambda_j - \lambda_{j+1} : j = 1, \dots, k-1\}.$$

We will use the notation and setup in Sect. 3, however backwards, since we will be dealing with local stable manifolds.

Fix a point $y_0 = (p_0, \eta_0) \in \Lambda_\epsilon$ for some small $\epsilon > 0$ so that $\widetilde{W}_\epsilon^s(y_0)$ has the form

$$\widetilde{W}_\epsilon^s(y_0) = \{(p, \nu_Y(p)) : p \in Y\}$$

for some smooth hypersurface Y in \mathbb{R}^d containing the point p_0 such that Y is strictly concave with respect to the *unit normal field* ν_Y . Then the *shape (curvature) operator*

$$\mathcal{K}_y : T_p Y \longrightarrow T_p Y$$

of Y at $p \in Y$ is negative definite at $y = (p, \nu_Y(p))$ with respect to the normal field ν_Y . Let $p_0(s)$, $0 \leq s \leq a$, be a C^3 curve on Y with $p_0(0) = p_0$ such that

$$\|\dot{p}_0(s)\| = 1 \quad , \quad s \in [0, a]. \tag{4.1}$$

Then $y_0(s) = (p_0(s), \nu_Y(p_0(s)))$ is a C^3 curve on $\widetilde{W}_\epsilon^s(y_0)$, so $\dot{y}_0(0) \in \widetilde{E}^s(y_0)$.

We will **assume that p_0 corresponds to a point in \mathcal{L}** via the local diffeomorphism (2.1) and that $\dot{y}_0(0) \in \widetilde{E}'_1(y_0)$, so

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \|d\phi_t(p_0) \cdot \dot{p}_0(0)\| = -\lambda_1. \tag{4.2}$$

Let $\tilde{p}_0(s)$, $0 \leq s \leq a$, be another C^3 curve on Y with $\tilde{p}_0(0) = p_0$ such that

$$\|\dot{\tilde{p}}_0(s)\| = 1 \quad , \quad s \in [0, a]. \quad (4.3)$$

Then $\tilde{y}_0(s) = (\tilde{p}_0(s), \nu_Y(\tilde{p}_0(s)))$ is a C^3 curve on $\widetilde{W}_\epsilon^s(\tilde{y}_0)$ with

$$\tilde{y}_0(0) = (\tilde{p}_0, \nu_Y(\tilde{p}_0)) = y_0(0)$$

and $\dot{\tilde{y}}_0(0) \in \widetilde{E}^s(\tilde{y})$.

Assume that $\dot{\tilde{y}}_0(0) \in \widetilde{F}'_{j_0}(y_0) \setminus \widetilde{F}'_{j_0-1}(y_0)$ **for some** $j_0 = 1, \dots, k$. Then we have

$$-\lambda_{j_0} = \lim_{t \rightarrow \infty} \frac{1}{t} \log \|d\phi_t(\tilde{p}_0) \cdot \dot{\tilde{p}}_0(0)\| \geq -\lambda_1.$$

Take $T > 0$ so large that

$$\frac{1}{t} \log \|d\phi_t(p_0) \cdot \dot{p}_0(0)\| < -\lambda_1 + \delta_0 \quad (4.4)$$

for all $t \geq T$.

Lemma 4.2. *Choosing $\delta_1 \in (0, \delta_0)$ sufficiently small, we can choose $T > 0$ so that if*

$$\|\dot{p}_0(0) - \dot{\tilde{p}}_0(0)\| < \delta_1, \quad (4.5)$$

then we have

$$\frac{1}{t} \log \|d\phi_t(\tilde{p}_0) \cdot \dot{\tilde{p}}_0(0)\| \leq -\lambda_1 + 2\delta_0 \quad (4.6)$$

for all $t \geq T$. Consequently, $j_0 = 1$, i.e. $\dot{\tilde{y}}_0(0) \in \widetilde{E}'_1(\tilde{y}_0)$.

Proof of Lemma 4.2. Denote by $\gamma(y_0(s))$ and $\gamma(\tilde{y}_0(s))$ the forward billiard trajectories in Ω_K determined by $y_0(s)$ and $\tilde{y}_0(s)$. Let $p_1(s), p_2(s), \dots$ and $\tilde{p}_1(s), \tilde{p}_2(s), \dots$, respectively, be the successive reflection points of these trajectories. We will denote by $\eta_j(s) \in \mathbb{S}^{d-1}$ the *reflected direction* of $\gamma(y_0(s))$ at $p_j(s)$ and by $\tilde{\eta}_j(s) \in \mathbb{S}^{d-1}$ the reflected direction of $\gamma(\tilde{y}_0(s))$ at $\tilde{p}_j(s)$. Set $t_0(s) = \tilde{t}_0(s) = 0$ and denote by $t_1(s), t_2(s), \dots$ the *times* of the consecutive reflections of the trajectory $\gamma(y_0(s))$ at ∂K . Then $t_j(s) = d_0(s) + d_1(s) + \dots + d_{j-1}(s)$, where $d_j(s) = \|p_{j+1}(s) - p_j(s)\|$. Similarly, denote by $\tilde{t}_1(s), \tilde{t}_2(s), \dots$ the *times* of the consecutive reflections of the trajectory $\gamma(\tilde{y}_0(s))$ at ∂K . Then $\tilde{t}_j(s) = \tilde{d}_0(s) + \tilde{d}_1(s) + \dots + \tilde{d}_{j-1}(s)$, where $\tilde{d}_j(s) = \|\tilde{p}_{j+1}(s) - \tilde{p}_j(s)\|$. Set $y_t(s) = \phi_t(y_0(s))$ and $\tilde{y}_t(s) = \phi_t(\tilde{y}_0(s))$ for all $t \geq 0$.

Given $t > 0$, denote by Y_t the *shift* of the surface Y along the flow ϕ_t after time t . That is

$$Y_t = \{\text{pr}_1(\phi_t(p, \nu_Y(p))) : p \in Y\},$$

where $\text{pr}_1(u, v) = u$ for all $(u, v) \in \mathbb{R}^d \times \mathbb{R}^d$. When $\phi_t(y_0)$ is not a reflection point of $\gamma(y_0)$, then locally near $\phi_t(y_0)$, Y_t is a smooth concave $(d-1)$ -dimensional surface in \mathbb{R}^d with "outward" unit normal given by the *direction* of $\gamma(y_0)$ at $\phi_t(y_0)$ (see [Si1], [Si2], or [Ch1], or [BCST]).

Fix a (large) integer m and $t > 0$ such that $t_m(0) > T$ and

$$t_m(0) < t < t_{m+1}(0). \quad (4.7)$$

Consider the surface $X = Y_t$ with the opposite normal $\nu_X = -\nu_{Y_t}$. Then X is strictly convex with respect to this normal and $(q, \nu_X(q)) \in W_\epsilon^u(x_0)$, where $x_0 = (q_0, v_0)$ with $q_0 = \text{pr}_1(y_t(0))$ and $v_0 = -\nu_{Y_t}(q_0)$. Consider the curve $q_0(s) = \text{pr}_1(y_t(s))$, $s \in [0, a]$, on X and let

$$x_\tau(s) = \phi_\tau(q(s), \nu_X(q(s))) = (u_\tau(s), v_\tau(s))$$

for $0 \leq \tau \leq t$. As in Sect. 3, let $q_1(s), q_2(s), \dots$ be the reflection points of the trajectory $\gamma(x_\tau(s))$ in Ω_K and let $\xi_j(s) \in \mathbb{S}^{d-1}$ be the *reflected direction* of $\gamma(x_\tau(s))$ at $q_j(s)$. Then $q_j(s) = p_{m-j}(s)$ for all $j = 1, \dots, m$, while $\xi_j(s)$ and $\eta_{m-j}(s)$ are symmetric with respect to the normal $\nu(q_j(s))$ to ∂K at $q_j(s)$. As in Sect. 3, let

$$D_{q_j(s)} : T_{q_j(s)}(X_{t_j(s)}) \longrightarrow T_{q_j(s)}(X_{t_j(s)})$$

be the *shape operator* of $X_{t_j(s)}$ at $q_j(s)$. Define $\ell_j(s) > 0$ by (3.1), where $\delta_j(s)$ is defined by (3.2). Then, by Theorem 3.1, formula (3.3) holds, where $\|\dot{p}_0(s)\| = 1$ by (4.1).

For the same m and t , we will need similar objects related to the trajectory $\gamma(\tilde{y}_0(s))$. Set $\tilde{x}_0 = (\tilde{q}_0, \tilde{v}_0)$ with $\tilde{q}_0 = \text{pr}_1(\tilde{y}_t(0))$ and $v_0 = -\nu_{Y_t}(\tilde{q}_0)$. Consider the curve $\tilde{q}_0(s) = \text{pr}_1(\tilde{y}_t(s))$, $s \in [0, a]$, on X and let

$$\tilde{x}_\tau(s) = \phi_\tau(\tilde{q}(s), \nu_X(\tilde{q}(s))) = (\tilde{u}_\tau(s), \tilde{v}_\tau(s)),$$

$0 \leq \tau \leq t$. As in Sect. 3, let $\tilde{q}_1(s), \tilde{q}_2(s), \dots$ be the reflection points of the trajectory $\gamma(\tilde{x}_\tau(s))$ and let $\tilde{\xi}_j(s) \in \mathbb{S}^{d-1}$ be the *reflected direction* of $\gamma(\tilde{x}_\tau(s))$ at $\tilde{q}_j(s)$. Then $\tilde{q}_j(s) = \tilde{p}_{m-j}(s)$ for all $j = 1, \dots, m$, while $\tilde{\xi}_j(s)$ and $\tilde{\eta}_{m-j}(s)$ are symmetric with respect to the normal $\nu(\tilde{q}_j(s))$ to ∂K at $\tilde{q}_j(s)$. Let

$$D_{\tilde{q}_j(s)} : T_{\tilde{q}_j(s)}(X_{t_j(s)}) \longrightarrow T_{\tilde{q}_j(s)}(X_{t_j(s)})$$

be the *shape operator* of $X_{t_j(s)}$ at $\tilde{q}_j(s)$. Define $\tilde{\ell}_j(s) > 0$ following (3.1), namely

$$[1 + \tilde{d}_j(s)\tilde{\ell}_j(s)]^2 = 1 + 2\tilde{d}_j(s)\tilde{k}_j(s) + (\tilde{d}_j(s))^2 \|D_{\tilde{q}_j(s)}(\tilde{e}_j(s))\|^2,$$

where $\tilde{k}_j(s)$ is defined analogously to $k_j(s)$. That is

$$\tilde{k}_0(s) = \frac{1}{\|\dot{\tilde{q}}_0(s)\|^2} \langle D_{\tilde{q}_0(s)}(\dot{\tilde{q}}_0(s)), \dot{\tilde{q}}_0(s) \rangle,$$

and for $j > 0$, $\tilde{k}_j(s) > 0$ denotes the *normal curvature* of $X_{\tilde{t}_j(s)} = \lim_{\tau \searrow \tilde{t}_j(s)} X_\tau$ at $\tilde{q}_j(s)$ in the direction $\tilde{e}_j(s)$ ($\|\tilde{e}_j(s)\| = 1$) of $\lim_{\tau \searrow \tilde{t}_j(s)} \frac{d}{ds'} (\tilde{u}_\tau(s'))|_{s'=s}$. Thus,

$$\tilde{k}_j(s) = \langle D_{\tilde{q}_j(s)}(\tilde{e}_j(s)), \tilde{e}_j(s) \rangle.$$

Set $\tilde{e}_0(s) = \dot{\tilde{q}}_0(0)/\|\dot{\tilde{q}}_0(0)\|$ and for any $j \geq 0$ define

$$\tilde{\delta}_j(s) = \frac{1}{1 + \tilde{d}_j(s)\tilde{\ell}_j(s)}. \quad (4.8)$$

Then, by Theorem 3.1, the analogue of (3.3) holds, namely

$$\|\dot{\tilde{q}}_0(s)\| = \|\dot{r}_0(s)\|\tilde{\delta}_0(s)\tilde{\delta}_1(s)\dots\tilde{\delta}_m(s) = \tilde{\delta}_0(s)\tilde{\delta}_1(s)\dots\tilde{\delta}_m(s), \quad (4.9)$$

taking into account (4.3).

Replacing $a > 0$ with a smaller constant we may assume that $d(p(s), \tilde{p}(s)) < \delta_1$ for all $s \in [0, a]$.

As is well-known (see e.g. [Si2], [I], [St1] or [PS]) it follows from the uniform hyperbolicity of the open billiard and general properties of (strong) stable manifolds that there exist global constants $C > 0$ and $\rho \in (0, 1)$, independent of the choice of y_0 such that

$$d(p_j(s), \tilde{p}_j(s)) \leq C\delta_1\rho^j \quad , \quad d(\eta_j(s), \tilde{\eta}_j(s)) \leq C\delta_1\rho^j \quad (4.10)$$

for all $j \geq 0$. It follows from the above that we have $d(q_j(s), \tilde{q}_j(s)) \leq C\delta_1\rho^{m-j}$ and $|d_j(s) - \tilde{d}_j(s)| \leq C\delta_1\rho^{m-j}$ for all $j = 0, 1, \dots, m$.

We will now estimate $\|e_j(s) - \tilde{e}_j(s)\|$ for all $j = 0, 1, \dots, m$. We have $\|\dot{p}_0(0) - \dot{\tilde{p}}_0(0)\| < \delta_1$ by (4.5). Assume $a > 0$ is so that² $\|\dot{p}_0(s) - \dot{\tilde{p}}_0(s)\| < \delta_1$ for all $s \in [0, a]$. Since

$$e_m(s) = \dot{p}_0(s)/\|\dot{p}_0(s)\| = \dot{p}_0(s) \quad , \quad \tilde{e}_m(s) = \dot{\tilde{p}}_0(s)/\|\dot{\tilde{p}}_0(s)\| = \dot{\tilde{p}}_0(s),$$

we have $\|e_m(s) - \tilde{e}_m(s)\| < \delta_1$ for all $s \in [0, a]$.

To get estimates for $\|e_j(s) - \tilde{e}_j(s)\|$ for $j < m$, notice that according to the definitions, the vectors $e_j(s)$ and $e_{j+1}(s)$ are symmetric with respect to the normal $\nu(q_j(s))$ to ∂K at $q_j(s)$. Therefore

$$e_j(s) = e_{j+1}(s) - 2\langle e_{j+1}(s), \nu(q_j(s)) \rangle \nu(q_j(s)).$$

Similarly,

$$\tilde{e}_j(s) = \tilde{e}_{j+1}(s) - 2\langle \tilde{e}_{j+1}(s), \nu(\tilde{q}_j(s)) \rangle \nu(\tilde{q}_j(s)).$$

Now using $d(q_j(s), \tilde{q}_j(s)) \leq C\delta_1\rho^{m-j}$, which implies $\|\nu(q_j(s)) - \nu(\tilde{q}_j(s))\| \leq C\delta_1\rho^{m-j}$ if the constant $C > 0$ is chosen sufficiently large, we derive the following³:

$$\begin{aligned} & \|e_j(s) - \tilde{e}_j(s)\| \\ &= \|[e_{j+1}(s) - 2\langle e_{j+1}(s), \nu(q_j(s)) \rangle \nu(q_j(s))] - [\tilde{e}_{j+1}(s) - 2\langle \tilde{e}_{j+1}(s), \nu(\tilde{q}_j(s)) \rangle \nu(\tilde{q}_j(s))]\| \\ &\leq \|(e_{j+1}(s) - \tilde{e}_{j+1}(s)) - 2\langle (e_{j+1}(s) - \tilde{e}_{j+1}(s)), \nu(q_j(s)) \rangle \nu(q_j(s))\| \\ &\quad + 2\|\langle \tilde{e}_{j+1}(s), \nu(q_j(s)) \rangle \nu(q_j(s)) - \langle \tilde{e}_{j+1}(s), \nu(\tilde{q}_j(s)) \rangle \nu(\tilde{q}_j(s))\| \\ &\leq \|e_{j+1}(s) - \tilde{e}_{j+1}(s)\| + 2\|\langle \tilde{e}_{j+1}(s), \nu(q_j(s)) - \nu(\tilde{q}_j(s)) \rangle \nu(q_j(s))\| \\ &\quad + 2\|\langle \tilde{e}_{j+1}(s), \nu(\tilde{q}_j(s)) \rangle (\nu(q_j(s)) - \nu(\tilde{q}_j(s)))\| \\ &\leq \|e_{j+1}(s) - \tilde{e}_{j+1}(s)\| + 4\|\nu(q_j(s)) - \nu(\tilde{q}_j(s))\| \\ &\leq \|e_{j+1}(s) - \tilde{e}_{j+1}(s)\| + 4C\delta_1\rho^{m-j}. \end{aligned}$$

A simple induction now gives

$$\begin{aligned} \|e_j(s) - \tilde{e}_j(s)\| &\leq \|e_{j+1}(s) - \tilde{e}_{j+1}(s)\| + 4C\delta_1\rho^{m-j} \\ &\leq \|e_{j+2}(s) - \tilde{e}_{j+2}(s)\| + 4C\delta_1\rho^{m-j-1} + 4C\delta_1\rho^{m-j} \\ &\leq \dots \\ &\leq \|e_m(s) - \tilde{e}_m(s)\| + 4C\delta_1\rho + \dots + 4C\delta_1\rho^{m-j-1} + 4C\delta_1\rho^{m-j} \\ &\leq \delta_1 + 4C\delta_1\frac{\rho}{1-\rho}. \end{aligned}$$

Therefore there exists a global constant $C_1 > 0$ such that

$$\|e_j(s) - \tilde{e}_j(s)\| < C_1\delta_1 \tag{4.11}$$

for all $j = 0, 1, \dots, m$ and all $s \in [0, a]$.

Next, it follows from (4.11) that there exists a global constant $C_2 > 0$ so that

$$|k_j(s) - \tilde{k}_j(s)| \leq C_2\delta_1 \quad , \quad \| \|D_{q_j(s)}(e_j(s))\| - \|D_{\tilde{q}_j(s)}(\tilde{e}_j(s))\| \| \leq C_2\delta_1,$$

²This is just for convenience – later on we only need this for $s = 0$.

³Using also the simple fact that if $w = v - 2\langle v, n \rangle n$ for a unit vector n , then $\|w\| = \|v\|$.

and thus $|\ell_j(s) - \tilde{\ell}_j(s)| \leq C_2\delta_1$ for all $s \in [0, a]$ and $j = 0, 1, \dots, m$. Now (3.2) and (4.8) yield that for some global constant $C_3 > 0$ we have

$$\begin{aligned} |\log \delta_j(s) - \log \tilde{\delta}_j(s)| &= |\log(1 + d_j(s)\ell_j(s)) - \log(1 + \tilde{d}_j(s)\tilde{\ell}_j(s))| \\ &\leq C |d_j(s)\ell_j(s) - \tilde{d}_j(s)\tilde{\ell}_j(s)| \leq C_3\delta_1, \end{aligned}$$

and so $\log \delta_j(s) \geq \log \tilde{\delta}_j(s) - C_3\delta_1$ for all $s \in [0, a]$ and $j = 0, 1, \dots, m$. From the latter, (3.3) and (4.9) it follows now that

$$\begin{aligned} \frac{1}{m+1} \log \|\dot{q}_0(s)\| &= \frac{1}{m+1} \sum_{j=0}^m \log \delta_j(s) \geq \frac{1}{m+1} \sum_{j=0}^m (\log \tilde{\delta}_j(s) - C_3\delta_1) \\ &\geq \frac{1}{m+1} \log \|\dot{\tilde{q}}_0(s)\| - C_3\delta_1. \end{aligned} \quad (4.12)$$

Assuming that $t > 0$ is sufficiently large, (4.4) holds and moreover for m with (4.7) we have

$$\frac{1}{m+1} \log \|\dot{q}_0(s)\| \leq \frac{1}{t} \log \|d\phi_t(p_0) \cdot \dot{p}_0(0)\| + \frac{C''}{m+1} < -\lambda_1 + \delta_0 + \frac{C''}{m+1}$$

for some global constant $C'' > 0$ independent of m and t . Thus, it follows from the above and (4.12) that

$$\frac{1}{m+1} \log \|\dot{\tilde{q}}_0(s)\| \leq \frac{1}{m+1} \log \|\dot{q}_0(s)\| + C_3\delta_1 \leq -\lambda_1 + \delta_0 + \frac{C''}{m+1} + C_3\delta_1.$$

However, again assuming that t and m with (4.7) are sufficiently large, we have

$$\frac{1}{m+1} \log \|\dot{\tilde{q}}_0(s)\| \geq \frac{1}{t} \log \|d\phi_t(\tilde{p}_0) \cdot \dot{\tilde{p}}_0(0)\| - \frac{C'''}{m+1}$$

for some global constant $C''' > 0$. Therefore

$$\frac{1}{t} \log \|d\phi_t(\tilde{p}_0) \cdot \dot{\tilde{p}}_0(0)\| \leq -\lambda_1 + \delta_0 + \frac{C''}{m+1} + \frac{C'''}{m+1} + C_3\delta_1$$

for $t \geq T$ for all sufficiently large T . This implies that there exists a global constant $C_4 > 0$ so that

$$\frac{1}{t} \log \|d\phi_t(\tilde{p}_0) \cdot \dot{\tilde{p}}_0(0)\| \leq -\lambda_1 + \delta_0 + C_4\delta_1$$

for all sufficiently large t . Assuming δ_1 is chosen with $C_4\delta_1 < \delta_0$, we see that (4.6) holds for $t \geq T$ for all sufficiently large T .

This proves Lemma 4.2. ■

We now continue with the proof of Theorem 4.1.

Assume as before that $y_0(s) = (p_0(s), \nu_Y(p_0(s)))$ is a C^3 curve on $\widetilde{W}_\epsilon^s(y_0)$ so that $\dot{y}_0(0) \in \widetilde{E}'_1(y_0)$. That is (4.2) holds. Then assume again that $T > 0$ is chosen so large that (4.4) is satisfied for all $t \geq T$.

Let $\delta_1 > 0$ and $T > 0$ be chosen as in Lemma 4.2. Since $k > 1$ by assumption, we can now choose a C^3 curve $\tilde{p}_0(s)$, $0 \leq s \leq a$, on Y with $\tilde{p}_0(0) = p_0$ and (4.3), and such that for $\tilde{y}_0(s) = (\tilde{p}_0(s), \nu_Y(\tilde{p}_0(s)))$ we have $\dot{\tilde{y}}_0(0) \in \widetilde{F}'_2(y_0) \setminus \widetilde{F}'_1(y_0) = (\widetilde{E}'_1(y_0) \oplus \widetilde{E}'_2(y_0)) \setminus \widetilde{E}'_1(y_0)$ and so that (4.5) holds as well. Now Lemma 4.2 implies that $\dot{\tilde{y}}_0(0) \in \widetilde{E}'_1(y_0)$, a contradiction. This proves Theorem 4.1. ■

5 Appendix

Here we prove Theorem 3.1.

To deal with Theorem 3.1 we will use the notation from the beginning of Sect. 3. We will need the following simple lemma which is a more precise version of Lemma 3.7 in [I].

Lemma 5.1. *Let Y and Z be the boundaries of two disjoint compact strictly convex domains in \mathbb{R}^d . Suppose Y and Z are C^3 -smooth and denote by ν_Y and ν_Z their outer unit normal fields. Given $\eta \in Y$, let η_t ($t \geq 0$) be the shift of η along the billiard trajectory issued from $(\eta, \nu_Y(\eta))$. Suppose $\eta(s)$, $0 \leq s \leq a$, is a C^3 (local) curve on Y such that for any $s \in [0, a]$ the trajectory $\{\eta_t(s) : t \geq 0\}$ hits transversally Z at $\eta_{\tau(s)}(s)$ and the angle $\varphi(s)$ between the reflected trajectory $\{\eta_t(s) : t \geq \tau(s)\}$ and the unit normal $\nu_Z(\eta_{\tau(s)}(s))$ is bounded from above by a constant $\varphi_0 \in (0, \pi/2)$. Let $L_\eta : T_\eta(Y) \rightarrow T_\eta(Y)$, $L_\eta u = (\nabla_u \nu_Y)(\eta)$, be the curvature (shape) operator of Y with respect to the normal field ν_Y . If $k_0 > 0$ is the normal curvature of Y at $\eta(0)$ in the direction of $u_0 = \dot{\eta}(0)$ and $\ell_0 > 0$ is such that*

$$(1 + \tau(0)\ell_0)^2 = 1 + 2\tau(0)k_0 + \tau^2(0) \|L_{\eta(0)}(u_0/\|u_0\|)\|^2,$$

then

$$\begin{aligned} (1 + \tau(0)\ell_0)\|\eta(s) - \eta(0)\| - C\|\eta(s) - \eta(0)\|^2 &\leq \|\eta_{\tau(0)}(s) - \eta_{\tau(0)}(0)\| \\ &\leq (1 + \tau(0)\ell_0)\|\eta(s) - \eta(0)\| + C\|\eta(s) - \eta(0)\|^2 \end{aligned}$$

for $0 \leq s \leq a$, where $C > 0$ is a constant depending only on the minimum and maximum of the curvatures of Y and Z , $M_i = \max_{s \in [0, a]} \|\frac{d^i \eta}{ds^i}(s)\|$ ($i = 1, 2, 3$), the constant φ_0 and $\sup_{0 \leq s \leq a} \tau(s)$. Moreover, there exist constants $0 < c < C$ such that

$$c\|\eta_{\tau(s)}(s) - \eta_{\tau(s)}(0)\| \leq \|\eta_{\tau(s)}(s) - \eta_{\tau(0)}(0)\| \leq C\|\eta_{\tau(s)}(s) - \eta_{\tau(s)}(0)\|$$

for all $s \in [0, a]$.

Proof of Lemma 5.1. We will first deal with the case when no reflection at Z is involved. Then we have to get estimates from below and above for

$$\lambda(s) = \|[\eta(s) + t\nu(s)] - [\eta(0) + t\nu(0)]\|^2,$$

for $t > 0$ and $s \rightarrow 0$, where $\nu(s) = \nu_Y(\eta(s))$.

We have $\lambda(s) = a(s) + 2tb(s) + t^2c(s)$, where

$$a(s) = \|\eta(s) - \eta(0)\|^2, \quad b(s) = \langle \eta(s) - \eta(0), \nu(s) - \nu(0) \rangle, \quad c(s) = \|\nu(s) - \nu(0)\|^2.$$

Since $\dot{a}(s) = 2\langle \eta(s) - \eta(0), \dot{\eta}(s) \rangle$ and $\ddot{a}(s) = 2\|\dot{\eta}(s)\|^2 + 2\langle \eta(s) - \eta(0), \ddot{\eta}(s) \rangle$, we have $\dot{a}(0) = 0$ and $\ddot{a}(0) = 2\|u_0\|^2$, so $a(s) = \|u_0\|^2 s^2 + O(s^3)$. (The estimate here obviously involves the constant M_3 .) Consequently, $\|\eta(s) - \eta(0)\| = \sqrt{a(s)} = \|u_0\|s + O(s^2)$.

Similarly, $\dot{b}(s) = \langle \dot{\eta}(s), \nu(s) - \nu(0) \rangle + \langle \eta(s) - \eta(0), \dot{\nu}(s) \rangle$ and $\ddot{b}(s) = O(s) + 2\langle \dot{\eta}(s), \dot{\nu}(s) \rangle$, so $b(0) = \dot{b}(0) = 0$ and $\ddot{b}(0) = 2\|u_0\|^2 k_0$. Hence $b(s) = \|u_0\|^2 k_0 s^2 + O(s^3)$.

Finally, $\dot{c}(s) = 2\langle \nu(s) - \nu(0), \dot{\nu}(s) \rangle$ and $\ddot{c}(s) = 2\|\dot{\nu}(s)\|^2 + O(s)$, so $c(0) = \dot{c}(0) = 0$ and $\ddot{c}(0) = 2\|\dot{\nu}(0)\|^2 = 2\|L_{\eta(0)}u_0\|^2$. Hence $c(s) = \|L_{\eta(0)}u_0\|^2 s^2 + O(s^3)$. This gives

$$\begin{aligned} \lambda(s) &= \|u_0\|^2 s^2 + 2t\|u_0\|^2 k_0 s^2 + t^2\|L_{\eta(0)}u_0\|^2 s^2 + O(s^3) \\ &= (\|u_0\|^2 + 2t\|u_0\|^2 k_0 + t^2\|L_{\eta(0)}u_0\|^2) s^2 + O(s^3). \end{aligned}$$

that $\max_{s \in [0, a]} \left\| \frac{d^i q_j(s)}{ds^i} \right\| \leq M$ for any $i = 1, 2, 3$ and any $j = 1, 2, \dots, m$. This implies similar estimates for $\left\| \frac{d^i u_{t_j(s)}(s)}{ds^i} \right\|$ for all $i = 1, 2, 3$ and $j = 1, 2, \dots, m$.

It is enough to estimate $\|\dot{q}(s)\|$ for $s = 0$. Next, fix for a moment an arbitrary $j = 1, \dots, m$. Given s close to 0, set $s_j^+ = 0$ if $t_j(0) \geq t_j(s)$ and $s_j^+ = s$ otherwise. Notice that $s_0^+ = 0$ and $s_{m+1}^+ = 0$. (In fact $t_0(s) = 0$ and $t_{m+1}(s) = t$ for all s .) Setting $\delta_j = \delta_j(0)$ for $j = 0, 1, \dots, m$ and $\epsilon_j = \|u_{t_j(s_j^+)}(s) - u_{t_j(s_j^+)}(0)\|$ for $j = 0, 1, \dots, m$ and $\epsilon_{m+1} = \|p(s) - p(0)\|$, Lemma 2 gives

$$[1 + d_j(0)\ell_j(0)] \epsilon_j - C \epsilon_j^2 \leq \epsilon_{j+1} \leq [1 + d_j(0)\ell_j(0)] \epsilon_j + C \epsilon_j^2$$

for some global constant $C > 0$. Thus, $\frac{\epsilon_j}{\delta_j} - C\epsilon_j^2 \leq \epsilon_{j+1} \leq \frac{\epsilon_j}{\delta_j} + C\epsilon_j^2$, which is equivalent to

$$\frac{\delta_j}{1 + C\delta_j\epsilon_j} \epsilon_{j+1} \leq \epsilon_j \leq \frac{\delta_j}{1 - C\delta_j\epsilon_j} \epsilon_{j+1}.$$

Applying these inequalities recursively, we get

$$\|p(s) - p(0)\| \prod_{i=j}^m \frac{\delta_i}{1 + C\delta_i\epsilon_i} \leq \epsilon_j \leq \|p(s) - p(0)\| \prod_{i=j}^m \frac{\delta_i}{1 - C\delta_i\epsilon_i}$$

for s close to 0. We have $\epsilon_0 = \|q_0(s) - q_0(0)\|$, so

$$\|p(s) - p(0)\| \prod_{i=0}^m \frac{\delta_i}{1 + C\delta_i\epsilon_i} \leq \|q_0(s) - q_0(0)\| \leq \|p(s) - p(0)\| \prod_{i=0}^m \frac{\delta_i}{1 - C\delta_i\epsilon_i}.$$

Dividing all sides of the last row by s and letting $s \rightarrow 0$, we get

$$\|\dot{p}(0)\| \prod_{i=0}^m \delta_i(0) \leq \|\dot{q}_0(0)\| \leq \|\dot{p}(0)\| \prod_{i=0}^m \delta_i(0),$$

which proves (3.3). ■

Acknowledgements. The work of the first author was supported by a scholarship from Najran University, Saudi Arabia.

References

- [Ar] Arnold, L.: Random Dynamical Systems. Springer Monographs in Mathematics, Springer, Berlin (1998).
- [BCST] Bálint, P., Chernov, N., Szász, D., Tóth, I.P.: Geometry of multi-dimensional dispersing billiards. *Asterisque* **286** 119-150 (2003)
- [BP] Barreira, L., Pesin, Ya.: Lyapunov exponents and smooth ergodic theory. Univ. Lect. Series 23, American Mathematical Society, Providence, R.I. (2002)
- [BPS] Barreira, L., Pesin, Ya., Sarig, O.: Smooth ergodic theory and nonuniformly hyperbolic dynamics. In: Handbook of Dynamical Systems; Vol. 1, No. PART B, Ch. 2 (2006)
- [BL] Bouchaud, J.-P., Le Doussal, P.: Numerical study of a d -dimensional periodic Lorentz gas with universal properties. *J. Statist. Phys.* **41**, 225-248 (1985)
- [B] Bowen, R.: Equilibrium states and the ergodic theory of Anosov diffeomorphisms. Lect. Notes in Maths. **470**, Springer-Verlag, Berlin (1975)
- [Burq] Burq, N.: Contrôle de l'équation des plaques en présence d'obstacles strictement convexes. *Mém. Soc. Math. France (N.S.)* **55** (1993)

- [Ch1] Chernov, N.: Structure of transversal leaves in multidimensional semidispersing billiards. *Funct. Anal. and Appl.* **16**, 35-46 (1982)
- [Ch2] Chernov, N.: A new proof of Sinai's formula for entropy of hyperbolic billiards. Its applications to Lorentz gas and stadium. *Funct. Anal. and Appl.* **25**, 204 - 219 (1991)
- [Ch3] Chernov, N.: Entropy, Lyapunov exponents, and Mean Free Path for Biliards. *J. Stat. Phys.* **88**, 1-29 (1997)
- [ChM] Chernov, N., Markarian, R.: Chaotic billiards. *Mathematical Surveys and Monographs Vol. 127*, Amer. Math. Soc. (2006)
- [CP] Chaubet, Y., Petkov, V.: Dynamical zeta functions for billiards. arXiv:2201.00683.
- [DP] Dellago, Ch., Posch, H.A.: Lyapunov spectrum and the conjugate pairing rule for a thermostatted random Lorentz gas: numerical simulations. *Phys. Rev. Lett* **78**, 211 (1997)
- [I] Ikawa, M.: Decay of solutions of the wave equation in the exterior of several strictly convex bodies. *Ann. Inst. Fourier* **38**, 113-146 (1988)
- [KH] Katok, A., Hasselblatt, B.: *Introduction to the Modern Theory of Dynamical System*. Cambridge Univ. Press, Cambridge (1995)
- [LBS] Latz, H., Van Beijeren, H., Dorfman, J.R.: Lyapunov spectrum and the conjugate pairing rule for a thermostatted random Lorentz gas: kinetic theory. *Phys. Rev. Lett* **78**, 207 (1997)
- [PP] Parry, W., Pollicott, M.: Zeta functions and the periodic orbit structure of hyperbolic dynamics. *Astérisque* **187-188**, (1990).
- [PS] Petkov, V., Stoyanov, L.: *Geometry of the generalized geodesic flow and inverse spectral problems*. 2nd edition. John Wiley & Sons, Chichester (2017)
- [Si1] Sinai, Ya.: Dynamical systems with elastic reflections. *Russian Math. Surveys* **25**, 137-190 (1970)
- [Si2] Sinai, Ya.: Development of Krylov's ideas. An addendum to: N.S.Krylov "Works on the foundations of statistical physics". Princeton Univ. Press, Princeton, pp. 239-281 (1979)
- [St1] Stoyanov, L.: Spectrum of the Ruelle operator and exponential decay of correlation for open billiard flows. *Amer. J. Math.* **123**, 715-759 (2001)
- [St2] Stoyanov, L.: Non-integrability of open billiard flows and Dolgopyat type estimates. *Ergod. Th.& Dynam. Sys.* **32**, 295-313 (2012)
- [V] Viana, M.: *Lectures on Lyapunov exponents*. Cambridge University Press, Cambridge (2014)