# Estimating Generalized Additive Conditional Quantiles for Absolutely Regular Processes

Yebin Cheng<sup>1</sup> and Jan G. De Gooijer<sup>2,\*</sup>

#### Abstract

We propose a nonparametric method for estimating the conditional quantile function that admits a generalized additive specification with an unknown link function. This model nests single-index, additive, and multiplicative quantile regression models. Based on a full local linear polynomial expansion, we first obtain the asymptotic representation for the proposed quantile estimator for each additive component. Then, the link function is estimated by noting that it corresponds to the conditional quantile function of a response variable given the sum of all additive components. The observations are supposed to be a sample from a strictly stationary and absolutely regular process. We provide results on (uniform) consistency rates, second order asymptotic expansions and point wise asymptotic normality of each proposed estimator.

**Key words and phrases:** Additive conditional quantiles, asymptotics, kernel estimation, unknown link function.

#### 1 Introduction

Suppose that Y is a response variable of interest which depends on a vector of random covariates  $X = (X_1, \ldots, X_d)^{\mathrm{T}} \in \mathbb{R}^d$ ,  $d \geq 2$ . We are interested in estimating the  $\alpha$ th  $(0 < \alpha < 1)$  conditional quantile q(x) of Y given X. For the ith subject, we assume that the sample  $\{(X_i, Y_i)\}_{i=1}^n$  is defined on a probability space  $(\Omega, \mathfrak{F}, \mathbb{P})$ , be a strictly stationary and absolutely regular stochastic process from the population (X, Y). It is well known that the  $\alpha$ th conditional quantile of Y given  $X = x = (x_1, \ldots, x_d)^{\mathrm{T}}$  is defined as the value q(x) such that  $q(x) = \inf\{t \colon F(t|x) \geq \alpha\}$ , where F(t|x) is the conditional distribution of Y given X = x. Equivalently,  $q(x) = \arg\inf_a \mathbb{E}\{\rho_\alpha(Y - a)|X = x\}$ , where  $\rho_\alpha(y) = |y| + (2\alpha - 1)y$  for any real y.

There is an extensive literature dealing with the estimation of q(x) when the functional relationship between Y and X is unknown. In particular, there have been many proposals using the additive quantile regression model  $q(x) = C + \sum_{u=1}^{d} q_u(x_u)$ , where C is a constant,  $q_u(x_u)$  (u = 1, 2, ..., d) are each additive components, and  $x_u$  is the uth component of x. For instance, Cheng, De Gooijer and Zerom (2011), De Gooijer and Zerom (2003), and Yu and Lu (2004) use this model to obtain estimates of additive conditional quantiles in a time series setting by nonparametric methods, and Horowitz and Lee (2005) and Noh and Lee (2014) for independent and identically distributed (i.i.d.) data by splines. In this paper,

<sup>&</sup>lt;sup>1</sup> Glorious Sun School of Business and Management, Donghua University, Shanghai, China

<sup>&</sup>lt;sup>2</sup> Amsterdam School of Economics, University of Amsterdam, the Netherlands

 $<sup>{\</sup>rm *Corresponding~author:~j.degooijer@contact.uva.nl}$ 

we consider estimating conditional quantiles in a more generalized setting. That is, we assume that the generalized additive model is of the form

$$q(x) = G\left(\sum_{u=1}^{d} q_u(x_u)\right),\tag{1.1}$$

where  $G(\cdot)$  is an *unknown* link function. It encompasses single-index, additive models and generalized additive models with *known* links as special cases. It also contains multiplicative models of the form  $q(x) = \widetilde{G}(\prod_{u=1}^d \widetilde{q}_u(x_u))$ , where  $\widetilde{G}(\cdot)$  and  $\widetilde{q}_u(\cdot)$  are unknown functions.

Building on the insight of Horowitz (2001) for the generalized additive conditional mean regression model, the main idea of this paper to estimate the components  $q_u(x_u)$ ,  $u=1,2,\ldots,d$ , is to write them as functionals of the distribution of the data, independent of  $G(\cdot)$ . Then, we estimate the unknown link function  $G(\cdot)$  by noting that it corresponds to the conditional quantile function of Y given  $q_0(X)$ , where  $q_0(x) = \sum_{u=1}^d q_u(x_u)$ . Also, we present theorems giving conditions under which the estimators of  $q_u(\cdot)$  and  $G(\cdot)$  are consistent and asymptotically normally distributed. Cheng (2007, Ch. 5) uses this latter result to formulate a test statistic for additivity of conditional quantile functions, under the assumption that the sample  $\{(X_i, Y_i)\}_{i=1}^n$  is still an absolutely regular process.

The rest of the paper is organized as follows. In Section 2, we provide a description of the estimator of the additive component and the estimator of the unknown link function, Section 3 gives the asymptotic representation of the nonparametric estimator of  $q_u(\cdot)$  for  $1 \le u \le d$ , in which the second order asymptotic representations are included, either. From this, it can be seen that the convergence rate of each estimator for each additive component is at the rate  $(nh)^{-\frac{1}{2}}$ , where h is the bandwidth. Compared to the rate of convergence  $(nh^d)^{-\frac{1}{2}}$  in the usual nonparametric setting, this kind of rate tends to 0 more quickly. In order to get the asymptotic representation for the estimator of the unknown link function  $G(\cdot)$ , we address in Section 4 the uniform convergence for the additive components. Then, in Section 5, we discuss the asymptotic representation of the estimation of the unknown link function  $G(\cdot)$  in (1.1) and subsequently we discuss the corresponding asymptotic normality. Concluding comments are presented in Section 6. The proofs of theorems and lemmas are provided in four supplementary appendices. This document also contains some useful lemmas and the proof of the Bahadur type linear representation for the local linear estimator of  $q_u(\cdot)$ .

Unless otherwise stated the symbol  $\stackrel{d}{\to}$  signifies convergence in distribution. The superscript T denotes matrix or vector transposition. For any a < b, we use the notation  $\mathfrak{M}_a^b$  to denote the sigma algebra generated by  $(Z_a,\ldots,Z_b)$ , where  $Z_i=(X_i,Y_i)$ . Given this notation, a process is called absolutely regular  $(\beta$ -mixing), if as  $\tau \to \infty$ ,  $\beta_{\tau}=\sup_{s\in N}\mathbb{E}\{\sup_{A\in\mathfrak{M}_{s+\tau}^{\infty}}\{\mathbb{P}(A|\mathfrak{M}_{-\infty}^s)-\mathbb{P}(A)\}\}\to 0$ ; see, e.g., Yoshihara (1978) and Arcones (1998).

# 2 Methodology

#### 2.1 Preamble

Following Horowitz (2001), we express  $q_k(\cdot)$ ,  $1 \le k \le d$ , as functionals of the population distribution of (X,Y). We first consider the case  $d \ge 3$ . The case d = 2 is briefly discussed at the end of this section. To simplify notation, for  $x \in \mathbb{R}^d$  and any fixed  $2 \le u \le d$ ,  $x_{\bar{u}}$  is a (d-2)-dimensional vector consisting of all components of x except  $x_1$  and  $x_u$ , and similar for the notations  $X_{\bar{u}}$  and  $X_{j,\bar{u}}$ , etc. For  $1 \le j \le n$ , let  $\widetilde{X}_j = \widetilde{X}_j^{t_1,u}$  be a d-dimensional vector with the first component  $t_1$ , the uth component  $t_u$  and the other components  $X_{j,\bar{u}}$ . Also, let  $\mathcal{X} = \prod_{k=1}^d [a_k, b_k] \subseteq \mathbb{R}^d$  be a subset for the support set of X and X does not contain its boundary. For identification of each component function and given some fixed points  $a_k < x_{k,0} < b_k$  for each  $k = 1, 2, \ldots, d$ , we assume that  $q_k(x_{k,0}) = 0$  and

$$\int_{a_1}^{b_1} \frac{w_1(x_1)}{q_1'(x_1)} dx_1 = 1, \tag{2.1}$$

where the weight function  $w_1(\cdot)$  defined on  $[a_1,b_1]$  is non-negative and integrates to one.

In order to make (2.1) hold, it is required that  $q'_1(x_1) \neq 0$  for  $a_1 \leq x_1 \leq b_1$ . Let  $\mathcal{X}_{1,u} = [a_1, b_1] \times [a_u, b_u]$ ,  $\mathcal{X}_{\bar{u}} = \prod_{1 \leq k \neq u \leq d} [a_k, b_k]$  and  $\partial_k q(x)$  be the first order partial derivative of q(x) with respect to the kth

component  $x_k$  of x. Define

$$D_{u}(x_{1}, x_{u}) = \int_{\mathcal{X}_{\bar{u}}} [\partial_{u} q(x)] p_{\bar{u}}(x_{\bar{u}}) dx_{\bar{u}} = \mathbb{E}[\partial_{u} q(x_{1}, x_{u}, X_{\bar{u}}) \mathbb{I}(X_{\bar{u}} \in \mathcal{X}_{\bar{u}})]$$

$$(2.2)$$

and

$$D_{1,u}(x_1, x_u) = \int_{\mathcal{X}_{\bar{u}}} [\partial_1 q(x)] p_{\bar{u}}(x_{\bar{u}}) \mathrm{d}x_{\bar{u}} = \mathbb{E}[\partial_1 q(x_1, x_u, X_{\bar{u}}) \mathbb{I}(X_{\bar{u}} \in \mathcal{X}_{\bar{u}})], \tag{2.3}$$

where  $p_{\bar{u}}(x_{\bar{u}})$  is the marginal density function of  $X_{\bar{u}}$ , and  $\mathbb{I}(\cdot)$  denotes the indicator function.

From (1.1) we have  $\partial_j q(x) = G'(\sum_{k=1}^d q_k(x_k))q'_j(x_j)$  for  $j = 1, \ldots, d$ . Then, from (2.2) and (2.3), we see that

$$D_{u}(x_{1}, x_{u}) = q'_{u}(x_{u}) \int_{\mathcal{X}_{\bar{u}}} G'\left(\sum_{k=1}^{d} q_{k}(x_{k})\right) p_{\bar{u}}(x_{\bar{u}}) dx_{\bar{u}}$$

and

$$D_{1,u}(x_1, x_u) = q_1'(x_1) \int_{\mathcal{X}_{\bar{u}}} G'(\sum_{k=1}^d q_k(x_k)) p_{\bar{u}}(x_{\bar{u}}) dx_{\bar{u}}.$$

Assume that  $D_{1,u}(x_1,x_u) \neq 0$  for any  $(x_1,x_u) \in \mathcal{X}_{1,u}$ . This implies that  $q_1'(x_1) \neq 0$  for  $x_1 \in [a_1,b_1]$ . Then it follows from (1.1) that

$$\frac{q_u'(x_u)}{q_1'(x_1)} = \frac{D_u(x_1, x_u)}{D_{1,u}(x_1, x_u)}.$$

Furthermore, by integrating both sides of the above expression, for any  $x_u \in [a_u, b_u]$ , we get

$$\int_{x_{u,0}}^{x_u} \int \frac{q'_u(t_u)}{q'_1(t_1)} w_1(t_1) dt_1 dt_u = \int_{x_{u,0}}^{x_u} \int \frac{D_u(t_1, t_u)}{D_{1,u}(t_1, t_u)} w_1(t_1) dt_u dt_1,$$

i.e.,

$$q_u(x_u) = \int_{x_u}^{x_u} \int \frac{D_u(t_1, t_u)}{D_{1,u}(t_1, t_u)} w_1(t_1) dt_u dt_1.$$
(2.4)

Observe that when  $x_u < x_{u,0}$ , (2.4) still holds because exchanging the location between  $x_u$  and  $x_{u,0}$  requires to add a minus notation before the integral simultaneously.

Analogously, for another non-negative weight function  $w_2(t_2)$  defined on  $[a_2, b_2]$ , which integrates to one, it follows that

$$c^{-1}q_1(x_1) = \int_{x_{1,0}}^{x_1} \int \frac{D_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} w_2(t_2) dt_2 dt_1,$$

with  $c^{-1} = \int (w_2(t_2)/q_2'(t_2)) dt_2$ . From this, we obtain that

$$\int w_1(t_1) \left[ \int \frac{D_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} w_2(t_2) dt_2 \right]^{-1} dt_1 = \int \frac{w_1(t_1)}{q'_1(t_1)} dt_1 \left[ \int \frac{w_2(t_2)}{q'_2(t_2)} dt_2 \right]^{-1} = c.$$
 (2.5)

Thus, we have

$$q_1(x_1) = c \int_{x_{1,0}}^{x_1} \int \frac{D_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} w_2(t_2) dt_2 dt_1.$$
(2.6)

Assume  $q_1'(\cdot) \neq 0$  and  $q_2'(\cdot) \neq 0$ . Then for the case d=2, going along the same lines as for the case  $d \geq 3$ , we can still establish (2.4), (2.5) and (2.6) if we set  $D_2(x_1, x_2) = \partial_2 q(x_1, x_2)$  and  $D_{1,2}(x_1, x_2) = \partial_1 q(x_1, x_2)$ .

### 2.2 Estimating the additive components $q_u(\cdot)$

Based on relationships (2.4)–(2.6), we construct the desired estimators by local polynomial fitting. To this end, we assume that q(x) is partially differentiable up to the order p + 1, which implies there are  $d^{p-1}$  parameters to estimate. By Taylor's expansion, for any w close to x, q(w) can be expressed as

$$q(w) = \sum_{\lambda \in \Lambda} \beta(\lambda, x) h^{-|\lambda|} (w - x)^{\lambda} + R_x(w) = \beta_x^{\mathrm{T}} A(\frac{w - x}{h}) + R_x(w),$$

where  $\Lambda = \{(\lambda_1, \ldots, \lambda_d)\}$ ,  $\lambda_i$  are non-negative integers and  $\sum_{i=1}^d \lambda_i \leq p-1\}$ ,  $|\lambda| = \sum_{i=1}^d \lambda_i$ ,  $x^{\lambda} = \prod_{i=1}^d x_i^{\lambda_i}$ . Here, h is a bandwidth specified below, and the two column vectors  $A(\frac{w-x}{h})$  and  $\beta_x$  are constructed from the elements  $h^{-|\lambda|}(w-x)^{\lambda}$  and  $\beta(\lambda,x)$  respectively, which are arranged in natural order with respect to  $\lambda \in \Lambda$ . Note that  $\beta(\lambda,x)$  is related to h and is of order  $h^{|\lambda|}$ .

To estimate  $\beta_x$ , using the local polynomial method, the corresponding estimator  $\widehat{\beta}_x$  can be obtained through minimizing the following objective function

$$\widehat{\beta}_x = \arg\min_{\beta} \sum_{i=1}^n w_{n,i} \rho_{\alpha} \left( Y_i - \beta^{\mathrm{T}} A \left( \frac{X_i - x}{h} \right) \right),$$

where the weight function  $w_{n,i}$  is equal to  $K\left(\frac{x-X_i}{h}\right)/\sum_{j=1}^n K\left(\frac{x-X_j}{h}\right)$  with the usual kernel function  $K(\cdot)$ . Then, the estimator of q(x) and its partial derivatives can be derived explicitly from  $\widehat{\beta}_x$ .

Based on this method, we can plug all the relevant estimators into (2.4) and obtain the estimator  $\widehat{q}_k(x_k)$  for  $q_k(x_k)$ ,  $1 \le k \le d$ . When  $d \ge 3$ ,  $\widehat{q}_u(x_u)$ ,  $2 \le u \le d$ , we have the representation

$$\widehat{q}_{u}(x_{u}) = \int_{x_{u,0}}^{x_{u}} \int \frac{\widehat{D}_{u}(t_{1}, t_{u})}{\widehat{D}_{1,u}(t_{1}, t_{u})} w_{1}(t_{1}) dt_{u} dt_{1},$$
(2.7)

where  $\widehat{D}_u(t_1,t_u)=\frac{1}{nh}\sum_{i=1}^n e_u^{\mathrm{T}}\widehat{\beta}_{\widetilde{X}_i}\mathbb{I}(X_{i,\bar{u}}\in\mathcal{X}_{\bar{u}})$  is the estimator of  $D_u(t_1,t_u), \frac{e_u^{\mathrm{T}}\widehat{\beta}_{\widetilde{X}_i}}{h}$  is the estimator of  $\partial_u q(\widetilde{X}_i)$  and  $e_u$ , which has the same dimension as  $\widehat{\beta}_x$ , denotes a unit vector such that its uth component is equal to 1 and all other components are equal to 0. Here, it should be noted that we adopt the leave-one-out rule to estimate  $\widehat{\beta}_{\widetilde{X}_i}$ . Similarly,  $\widehat{D}_{1,u}(t_1,t_u)=\frac{1}{nh}\sum_{i=1}^n e_1^{\mathrm{T}}\widehat{\beta}_{\widetilde{X}_i}\mathbb{I}(X_{i,\bar{u}}\in\mathcal{X}_{\bar{u}})$ . Analogously, for  $d\geq 3$ , the estimator of  $q_1(x_1)$  is given by

$$\widehat{q}_1(x_1) = \widehat{c} \int_{x_{1,0}}^{x_1} \int \frac{\widehat{D}_{1,2}(t_1, t_2)}{\widehat{D}_2(t_1, t_2)} w_2(t_2) dt_2 dt_1, \tag{2.8}$$

where

$$\widehat{c} = \int w_1(t_1) \left[ \int \frac{\widehat{D}_{1,2}(t_1, t_2)}{\widehat{D}_2(t_1, t_2)} w_2(t_2) dt_2 \right]^{-1} dt_1.$$
(2.9)

When d=2 in (2.7))–(2.9), we take the two estimators  $\widehat{D}_2(t_1,t_2)$  and  $\widehat{D}_{1,2}(t_1,t_2)$  as  $e_2^{\mathrm{T}}\widehat{\beta}_{(t_1,t_2)}$  and  $e_1^{\mathrm{T}}\widehat{\beta}_{(t_1,t_2)}$ . Then,  $\widehat{q}_2(x_2)$  and  $\widehat{q}_1(x_1)$  are of the same form as (2.7) and (2.8), respectively.

## 2.3 Estimating the link function $G(\cdot)$

To estimate  $G(\cdot)$ , we define  $q_0(x) = \sum_{k=1}^d q_k(x_k)$ ,  $\mathcal{V} = \{q_0(x) | x \in \mathcal{X}\}$  and  $\widehat{q}_0(X_i) = \sum_{k=1}^d \widehat{q}_k(X_{i,k})$ , where  $X_{i,k}$  is the kth component of  $X_i$ . Then,  $\mathcal{V}$  is also a compact set since the continuity of  $q_0(x)$ . From the definition of the conditional quantile, the property on the conditional expectation and the Borel measurability on the function  $q_0(\cdot)$ , we note that

$$1 - \alpha = \mathbb{E}(\mathbb{I}\{Y \le G(q_0(X))\}|X) = \mathbb{E}[\mathbb{E}(\mathbb{I}\{Y \le G(q_0(X))\}|X)|q_0(X)]$$
  
=  $\mathbb{E}(\mathbb{I}\{Y \le G(q_0(X))\}|q_0(X)).$ 

Thus the unknown function G(v) is also the conditional quantile function of Y given that  $q_0(X) = v$ . For any fixed  $v \in \mathcal{V}$ , the estimator of G(v) is given by the following empirical function

$$\widehat{G}_n(v) = \inf \{ y | \widehat{F}_n(y) | q_0(x) = v \} \ge 1 - \alpha \},$$

where

$$\widehat{F}_n(y|\ q_0(x) = v) = \sum_{i=1}^n \frac{K_G\left(\frac{v - \widehat{q}_0(X_i)}{h_G}\right) \mathbb{I}(Y_i \le y, X_i \in \mathcal{X})}{\sum_{j=1}^n K_G\left(\frac{v - \widehat{q}_0(X_j)}{h_G}\right) \mathbb{I}(X_j \in \mathcal{X})},\tag{2.10}$$

with  $K_G(\cdot)$  a kernel function of a scalar argument (in the sense of nonparametric density estimation), and  $h_G$  a bandwidth tending to 0 as  $n \to \infty$ .

# 3 Convergence of the additive components

For convenience of presentation, we introduce some notation. For  $2 \leq u \leq d$  and  $t_{1,u} = (t_1, t_u) \in \mathcal{X}_{1,u}$ , let  $\varepsilon_i = Y_i - q(X_i)$ ,  $Q = \int A(z)A^{\mathrm{T}}(z)K(z)\mathrm{d}z$ ,  $Q^* = \int K(x)A(x)A^{\mathrm{T}}(x)x^{\mathrm{T}}\mathrm{d}x$ ,  $\beta_{j,t_{1,u}} = \beta_{\widetilde{X}_j}$ ,  $K_{ij,t_{1,u}} = K\left(\frac{X_i - \widetilde{X}_j}{h}\right)$ ,  $A_{ij,t_{1,u}} = A\left(\frac{X_i - \widetilde{X}_j}{h}\right)$ ,  $r_{ij,t_{1,u}} = q(X_i) - \beta_{j,t_{1,u}}^{\mathrm{T}}A_{ij,t_{1,u}}$ ,  $P_{ij,t_{1,u}} = (\beta - \beta_{j,t_{1,u}})^{\mathrm{T}}A_{ij,t_{1,u}}$ ,  $\Delta_{1,u} = \widehat{D}_{1,u}(t_1,t_u) - D_{1,u}(t_1,t_u)$  and  $\Delta_u = \widehat{D}_u(t_1,t_u) - D_u(t_1,t_u)$ . Also, for any real y and  $1 \leq k \leq d$ , we assume that

$$f_k(y) = \int_{-1}^{y} \int A(t)K(t)dt_{\underline{k}}dt_k \, \mathbb{I}(|y| \le 1)$$

with  $t_{\underline{k}}$  a (d-1)-dimensional vector constructed from t by deleting the kth argument  $t_k$  of t. Let the set  $A_{(u)} = [x_{u,0} - h, x_u + h] \times \prod_{1 \le l \ne u \le d} [a_l - h, b_l + h]$  and  $\varepsilon > 0$  be an any sufficiently small constant. Also, let  $S_{t_1}$  be the support set of the distribution of  $X_j^{t_1,u}$  and  $S_{t_2}$  be the support set of the distribution of  $X_j^{t_2,u}$ ,  $t_1 \ne t_2$ . In addition, for ease of presentation, we introduce the notation  $\mathbb{E}_j$  defined as  $\mathbb{E}_{j0}g(\xi_i, \xi_0) = g_1(\xi_{j0})$ .

The asymptotic properties of  $\widehat{q}_n(\cdot)$  and  $\widehat{G}_n(\cdot)$  are established under the following regularity conditions:

- (B1) The density function  $p(\cdot)$  of X is bounded and continuous on its support set.
- (B2)  $K(\cdot)$  has bounded and continuous partial derivatives of order 1 and has the support set as unit sphere. Moreover,  $\int tK(t)dt = 0$ .
- (B3) The bandwidth  $h = O(n^{-\kappa})$  satisfies that  $\frac{1-\frac{1}{r}}{4p+d-\frac{d}{r^2}} < \kappa < \frac{1-\frac{1}{r}}{d+p+\frac{d+3p}{r}}$ . And the mixing coefficient  $\beta_k = O(k^{-r})$  with

$$r \geq \max \left\{ d + \frac{1}{2} - \frac{d}{p} + \frac{17p + d - 3}{2p(d - 1 - dp^{-1})}, d - 7 + \frac{2d^2 - 4 - 4d}{p} - \frac{22p + 6}{dp}, d, 11 \right\}.$$

- (B4) Let G(x, y) be the conditional distribution of  $\varepsilon_i$  given that  $X_i = x$ . Its conditional density function g(x, y) has first continuous derivative for y in the neighbourhood of 0 and  $x \in \mathcal{X}$ .
- (B5)  $g_1(x) = g(x,0)p(x)$  has bounded second derivatives and is bounded away from zero on  $\mathcal{X}$ .
- (B6) The bandwidth satisfies  $\frac{1}{2p+1} \le \kappa \le \frac{1-\frac{1}{r}}{3d+2+\frac{2d}{r}-\frac{d}{r^2}}$ .
- (B7) For any  $1 \leq l \leq n-1$ , the density function  $f(\cdot, \cdot)$  of  $(X_1, X_{1+l})$  exists and is continuous and bounded on its domain.
- (B8)  $D_{1,u}(t_1,t_u)$  has first continuous derivatives with respect to  $t_u$  for any  $t_1$ .
- (B9) There are m > 0 and compact intervals  $\overline{S}_1 \subset \operatorname{int}(S_{t_1})$  and  $\overline{S}_2 \subset \operatorname{int}(S_{t_2})$  such that  $|g_1'(t_1)| \geq m$  for all  $t_1 \in \overline{S}_1$  and  $g_2'(t_2)| \geq m$  for all  $t_2 \in \overline{S}_2$ .
- (B10)  $w_1(t_1) = O(t_1 a_1)$  as  $t_1 \to a_1$ ,  $w_1(t_1) = O(t_1 b_1)$  as  $t_1 \to b_1$ ,  $w_2(t_2) = O(t_2 a_2)$  as  $t_2 \to a_2$  and  $w_1(t_2) = O(t_2 b_2)$  as  $t_2 \to b_2$ .

**Lemma 3.1.** Under conditions (B1)–(B9), it holds uniformly for  $(t_1, t_u) \in \mathcal{X}_{1,u}$  with probability one that

$$\frac{1}{n^{2}h^{d+1}} \sum_{j=1}^{n} Q_{jn,t_{1,u}}^{-1} W_{j,n}(t_{1,u}) = B_{1,u}h^{p} (1 + o(1)), \qquad (3.1)$$

where  $Q_{jn,t_{1,u}} = \frac{1}{h^d} \mathbb{E}_j (K_{ij,t_{1,u}} A_{ij,t_{1,u}} A_{ij,t_{1,u}}^T g(X_i, 0)), B_{1,u} = \mathbb{E} \iint B_{2,u}(\widetilde{X}_j) dt_1 dt_u$ 

$$B_{2,u}(t) = e_u^{\mathrm{T}} Q^{-1} Q^* \frac{1}{p! q_1(t)} \frac{\partial g_1(t)}{\partial t} \int A(s) s^p K(s) ds \, q^{(p)}(t),$$

and

$$W_{j,n}(t_{1,u}) = \sum_{i=1}^{n} K_{ij,t_{1,u}} A_{ij,t_{1,u}} \left[ \mathbb{I}(\varepsilon_i \le 0) - \mathbb{I}(\varepsilon_i \le r_{ij,t_{1,u}}) \right] \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{\bar{u}}). \tag{3.2}$$

Lemma 3.2. Under conditions (B1)-(B9), it holds almost surely that

$$\sup_{(t_1,t_u)\in\mathcal{X}_{1,u}}\Delta_{1,u}=O\left(\left(nh^{4+\frac{1+\varepsilon}{r}}\right)^{-\frac{1}{2}}\right)\ and\ \sup_{(t_1,t_u)\in\mathcal{X}_{1,u}}\Delta_u=O\left(\left(nh^{4+\frac{1+\varepsilon}{r}}\right)^{-\frac{1}{2}}\right)$$

for  $2 \le u \le d$ .

**Theorem 3.1.** Let conditions (B1)–(B10) hold.

i) For  $2 \le u \le d$  and  $a_u \le x_u \le b_u$ , we have the following asymptotic representation

$$\sqrt{nh}(\widehat{q}_{u}(x_{u}) - q_{u}(x_{u})) = \sum_{i=1}^{n} \frac{\left((1-\alpha) - \mathbb{I}(\varepsilon_{i} \leq 0)\right) w_{1}(X_{i,1}) p_{\bar{u}}(X_{i,\bar{u}}) \mathbb{I}(X_{i} \in A_{(u)})}{\sqrt{nh} D_{1,u}(X_{i,1}, X_{i,u}) g_{1}(X_{i})} \\
\left(e_{u}^{T} - \frac{e_{1}^{T} D_{u}(X_{i,1}, X_{i,u})}{D_{1,u}(X_{i,1}, X_{i,u})}\right) Q^{-1} \left[f_{u}\left(\frac{X_{i,u} - x_{u}}{h}\right) - f_{u}\left(\frac{X_{i,u} - x_{u,0}}{h}\right) + \sum_{2 \leq k \leq d, k \neq u,} \left(f_{k}\left(\frac{X_{i,k} - b_{k}}{h}\right) - f_{k}\left(\frac{X_{i,k} - a_{k}}{h}\right)\right)\right] + o_{\mathbb{P}}(1). \tag{3.3}$$

ii) For  $a_1 \leq x_1 \leq b_1$ , it holds that

$$\sqrt{nh}(\widehat{q}_{1}(x_{1}) - q_{1}(x_{1})) = \sum_{i=1}^{n} \frac{\left((1-\alpha) - \mathbb{I}(\varepsilon_{i} \leq 0)\right) w_{2}(X_{i,2}) p_{\bar{2}}(X_{i,\bar{2}}) \mathbb{I}(X_{i} \in A_{(u)})}{\sqrt{nh} D_{2}(X_{i,1}, X_{i,2}) g_{1}(X_{i})} \cdot \left(e_{1}^{T} - \frac{e_{2}^{T} D_{1,2}(X_{i,1}, X_{i,2})}{D_{2}(X_{i,1}, X_{i,2})}\right) Q^{-1} \left[c\left(f_{1}\left(\frac{X_{i,1} - x_{1}}{h}\right) - f_{1}\left(\frac{X_{i,1} - x_{1,0}}{h}\right)\right) + \left(c - c_{1}(x_{1})w_{1}(X_{i,1})\right) \sum_{3 \leq k \leq d} \left(f_{k}\left(\frac{X_{i,k} - b_{k}}{h}\right) - f_{k}\left(\frac{X_{i,l} - a_{k}}{h}\right)\right)\right] + o_{\mathbb{P}}(1), \tag{3.4}$$

where  $c_1(x_1) = \int_{x_{1,0}}^{x_1} \int \frac{D_{1,2}(t_1,t_2)}{D_2(t_1,t_2)} w_2(t_2) dt_2 dt_1$ .

**Remark 3.1.** Moreover, if the more restrictive condition  $\frac{1}{2p-2-\frac{d}{r}} < \kappa < \frac{1-\frac{1}{r}}{3d+4+\frac{4d}{r}-\frac{d}{r^2}}$  is given compared to condition (B6), the second order representation in Theorem 3.1 can be specified explicitly as follows.

i) For  $2 \le u \le d$ , the remainder term in Theorem 3.1 is equal to

$$\xi_{n1}h^{\frac{1}{2}} + O_{\mathbb{P}}\left(h^{1-\frac{1+\varepsilon}{2r}} + \left(nh^{2p-1}\right)^{\frac{1}{2}} + \frac{n^{\frac{1}{2}}}{h^{\frac{1}{2}}}\left(n^{1-\frac{1}{3r}-\frac{2\varepsilon}{3}}h^{d\left(1+\frac{2}{3r}-\frac{1}{3r^2}\right)}\right)^{-\frac{3}{4}}\right),$$

where

$$\xi_{n1} = \sum_{i=1}^{n} \left\{ \frac{1}{\sqrt{n}} \left( a_{u}(x_{u}, i) - b_{u}(x_{u}, i) \right) + \left[ (1 - \alpha) - \mathbb{I}(\varepsilon_{i} \leq 0) \right] \mathbb{I}(X_{i} \in A_{(u)}) \right. \\ \left. \cdot \left[ \frac{e_{u}^{\mathsf{T}} Q^{-1}}{\sqrt{n}} \int \frac{\partial}{\partial x^{\mathsf{T}}} \left( \frac{w_{1}(x_{1}) p_{\bar{u}}(x_{\bar{u}})}{g_{1}(x) D_{1,u}(x_{1}, x_{u})} \right) \right|_{x = \widetilde{X}_{i}} t A(t) K(t) dt$$

$$\begin{split} & -\frac{e_1^{\mathrm{T}}Q^{-1}}{\sqrt{n}} \int \frac{\partial}{\partial x^{\mathrm{T}}} \left( \frac{w_1(x_1)p_{\bar{u}}(x_{\bar{u}})D_u(x_1,x_u)}{g_1(x)D_{1,u}^2(x_1,x_u)} \right) \bigg|_{x=\tilde{X}_i} tA(t)K(t)\mathrm{d}t \bigg] \\ & + \left( e_u^{\mathrm{T}} - e_1^{\mathrm{T}} \frac{D_u(X_{i,1},X_{i,u})}{D_{1,u}(X_{i,1},X_{i,u})} \right) \frac{w_1(X_{i,1})p_{\bar{u}}(X_{i,\bar{u}})Q^{-1}}{\sqrt{nh^2}g_1(X_i)D_{1,u}(X_{i,1},X_{i,u})} M_2^{(u)}(X_i) \bigg\} h^{\frac{1}{2}}. \end{split}$$

ii) For the first additive component, the remainder term in Theorem 3.1 is equal to

$$\xi_{n2}h^{\frac{1}{2}} + O_{\mathbb{P}}\left(h^{1 - \frac{1+\varepsilon}{2r}} + \frac{n^{\frac{1}{2}}}{h^{\frac{1}{2}}}\left(n^{1 - \frac{1}{3r} - \frac{2\varepsilon}{3}}h^{d\left(1 + \frac{2}{3r} - \frac{1}{3r^2}\right)}\right)^{-\frac{3}{4}} + \left(nh^{2p-1}\right)^{\frac{1}{2}}\right),$$

where

$$\xi_{n2} = \sum_{i=1}^{n} \left\{ \frac{b_{c}(i) - a_{c}(i) + a_{1}(x_{1}, i) - b_{1}(x_{1}, i)}{\sqrt{n}} + \left( (1 - \alpha) - \mathbb{I}(\varepsilon_{i} \leq 0) \right) \mathbb{I}(X_{i} \in A_{(1)}) \right. \\
\left. \cdot \left[ -c_{1}(x_{1})e_{1}^{\mathsf{T}}Q^{-1} \int \frac{\partial}{\partial x} \left( \frac{w_{1}(x_{1})w_{2}(x_{2})p_{\bar{2}}(x_{\bar{2}})}{g_{1}(x)D_{2}(x_{1}, x_{2})} \right) \Big|_{x=\tilde{X}_{i}^{1,2}} tA(t)K(t)dt \right. \\
\left. + c_{1}(x_{1})e_{2}^{\mathsf{T}}Q^{-1} \int \frac{\partial}{\partial x} \left( \frac{w_{1}(x_{1})w_{2}(x_{2})p_{\bar{2}}(x_{\bar{2}})D_{1,2}(x_{1}, x_{2})}{g_{1}(x)D_{2}^{2}(x_{1}, x_{2})} \right) \Big|_{x=\tilde{X}_{i}^{1,2}} tA(t)K(t)dt \right. \\
\left. + c e_{1}^{\mathsf{T}}Q^{-1} \int \frac{\partial}{\partial x} \left( \frac{w_{2}(x_{2})p_{\bar{2}}(x_{\bar{2}})}{g_{1}(x)D_{2}(x_{1}, x_{2})} \right) \Big|_{x=\tilde{X}_{i}^{1,2}} tA(t)K(t)dt \right. \\
\left. - c e_{2}^{\mathsf{T}}Q^{-1} \int \frac{\partial}{\partial x} \left( \frac{w_{2}(x_{2})p_{\bar{2}}(x_{\bar{2}})D_{1,2}(x_{1}, x_{2})}{g_{1}(x)D_{2}^{2}(x_{1}, x_{2})} \right) \Big|_{x=\tilde{X}_{i}^{1,2}} tA(t)K(t)dt \right. \\
\left. + \frac{w_{2}(X_{i,2})p_{\bar{2}}(X_{i,\bar{2}})\mathbb{I}(X_{i} \in A)}{\sqrt{nh^{2}}D_{2}(X_{i,1}, X_{i,2})g_{1}(X_{i})} \left( e_{1}^{\mathsf{T}} - \frac{e_{2}^{\mathsf{T}}D_{1,2}(X_{i,1}, X_{i,2})}{D_{2}(X_{i,1}, X_{i,2})} \right) Q^{-1}M_{1}^{(u)}(X_{i}) \right] \right\}. \tag{3.5}$$

Remark 3.2. Conditions (B3) and (B6) are about the restriction on the bandwidth. In order to get a chosen bandwidth, it should hold that  $p > \frac{d+1+\frac{d-1}{r}}{1-\frac{5}{r}}$ . Condition (B10) can be relaxed from Theorem 3.1. Otherwise, there are two extra similar terms which will be included in (3.3). The remaining conditions in Theorem 3.1 are standard; see, e.g., Chaudhuri (1991) and Honda (2004). Condition (B9) is used to identify the  $q'_u(\cdot)$ 's.

For convenience, let the set  $A_{(u)}^*$  be the limit of  $A_{(u)}$  for  $1 \le u \le d$ . From Theorem 3.1, the following Corollary 3.1 can be inferred from the standard Doob's large-block and small-block technique; see, e.g., Cai and Ould-Saïd (2003, Theorem 2).

**Corollary 3.1.** Under the conditions of Theorem 3.1, for  $1 \le u \le d$ , it holds that

$$\sqrt{nh}(\widehat{q}_u(x_u) - q_u(x_u) - B_{1,u}h^p) \stackrel{d}{\to} \mathcal{N}(0, \sigma_u^2),$$

where, for u = 1,  $\frac{\sigma_u^2}{\alpha(1-\alpha)}$  is defined as

$$\begin{split} &\int_{\left(t \in A_{(1)}^*\right)} \frac{c^2 w_2(t_2) p_{\bar{2}}(t_{\bar{2}})}{D_2(x_1, t_2) g_1(t)} \left( \left(e_1^{\mathrm{T}} - \frac{e_2^{\mathrm{T}} D_{1,2}(x_1, t_2)}{D_2(x_1, t_2)} \right) Q^{-1} f_1(t_1) \right)^2 p(x_1, t_2, t_{\bar{2}}) \mathrm{d}t \\ &+ \int_{\left(t \in A_{(1)}^*\right)} \frac{c^2 w_2(t_2) p_{\bar{2}}(t_{\bar{2}})}{D_2(x_{1,0}, t_2) g_1(t)} \left( \left(e_1^{\mathrm{T}} - \frac{e_2^{\mathrm{T}} D_{1,2}(x_{1,0}, t_2)}{D_2(x_{1,0}, t_2)} \right) Q^{-1} f_1(t_1) \right)^2 p(x_{1,0}, t_2, t_{\bar{2}}) \mathrm{d}t \\ &+ \sum_{3 \leq k \leq d} \int_{\left(t \in A_{(1)}^*\right)} \frac{\left(c - c(x_1, x_{1,0}) w_1(t_1)\right)^2 w_2(t_2) p_{\bar{2}}(t_{\bar{2}})}{D_2(t_1, t_2) g_1(t)} \\ &\cdot \left( \left(e_1^{\mathrm{T}} - \frac{e_2^{\mathrm{T}} D_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} \right) Q^{-1} f_k(t_k) \right)^2 \left(p(t_1, b_k, t_{\bar{k}}) + p(t_1, a_k, t_{\bar{k}})\right) \mathrm{d}t \end{split}$$

and, for  $2 \le u \le d$ ,  $\frac{\sigma_u^2}{\alpha(1-\alpha)}$  is defined as

$$\int_{\left(t \in A_{(u)}^{*}\right)} \frac{w_{1}^{2}(t_{1}) p_{\bar{u}}^{2}(t_{\bar{u}}) p(t_{1}, x_{u,0}, t_{\bar{u}})}{D_{1,u}^{2}(t_{1}, x_{u,0}) g_{1}^{2}(t_{1}, x_{u,0}, t_{\bar{u}})} \left(\left(e_{u}^{\mathsf{T}} - \frac{e_{1}^{\mathsf{T}} D_{u}(t_{1}, x_{u,0})}{D_{1,u}(t_{1}, x_{u,0})}\right) Q^{-1} f_{u}(t_{u})\right)^{2} dt 
+ \int_{\left(t \in A_{(u)}^{*}\right)} \frac{w_{1}^{2}(t_{1}) p_{\bar{u}}^{2}(t_{\bar{u}}) p(t_{1}, x_{u}, t_{\bar{u}})}{D_{1,u}^{2}(t_{1}, x_{u}, t_{\bar{u}})} \left(\left(e_{u}^{\mathsf{T}} - \frac{e_{1}^{\mathsf{T}} D_{u}(t_{1}, x_{u})}{D_{1,u}(t_{1}, x_{u})}\right) Q^{-1} f_{u}(t_{u})\right)^{2} dt 
+ \sum_{2 \leq k \leq d, k \neq u} \int_{\left(t \in A_{(u)}^{*}\right)} \frac{w_{1}^{2}(t_{1}) p_{\bar{u}}^{2}(t_{\bar{u}})}{D_{1,u}^{2}(t_{1}, t_{u}) g_{1}^{2}(t)} \left(\left(e_{u}^{\mathsf{T}} - \frac{e_{1}^{\mathsf{T}} D_{u}(t_{1}, t_{u})}{D_{1,u}(t_{1}, t_{u})}\right) Q^{-1} f_{k}(t_{k})\right)^{2} 
\cdot \left(p(t_{1}, b_{k}, t_{\bar{k}}) + p(t_{1}, a_{k}, t_{\bar{k}})\right) dt,$$

and where  $B_{1,u}$  is defined in Lemma 3.1.

**Remark 3.3.** The optimal bandwidth is equal to  $h_{opt} = Cn^{-\frac{1}{2p+1}}$ .

# 4 Uniform convergence of additive components

**Theorem 4.1.** Under conditions (B1)-(B8), it holds with probability one that

$$\sup_{x_u \in [a_u, b_u]} |\widehat{q}_u(x_u) - q_u(x_u)| = O\left(\left(nh^{1 + \frac{1 + \varepsilon}{r}}\right)^{-\frac{1}{2}}\right)$$
(3.6)

for  $1 \le u \le d$  and any sufficiently small constant  $\varepsilon > 0$ .

# 5 Convergence of the unknown link function

In this section, we address the asymptotic representation for the estimated link function  $\widehat{G}_n(v)$ . In particular, we show that the resulting representation holds uniformly for  $v \in \mathcal{V}$ . Then, the corresponding asymptotic normality with the bias will be illustrated. Furthermore, we discuss the choice of the optimal bandwidth  $h_{G,opt}$ . In the sequel, let  $F_n(t|v)$  be an empirical conditional distribution, which equals the right-hand side (RHS) of (2.10) with  $\widehat{q}_0(X_i)$  replaced by  $q_0(X_i)$ .

We impose the following conditions:

- (C1)  $K_G(x)$  is symmetric on the support set [-1,1].  $K_G'(1)=K_G'(-1)=0$ ,  $K_G(1)=K_G(-1)=0$  and  $K_G''\leq 0$ . There exists a constant C>0 such that  $|K_G''(x+t)-K_G''(x)|\leq C|t|$  for any x and t.
- (C2) The density function  $f_{q_0}(v)$  of  $q_0(X)$  has the second order continuous derivative for  $v \in \mathcal{V}$ , and  $f_{q_0}(v) > 0$ .
- (C3)  $\liminf_{n\to\infty} nh^{1+\frac{1+\varepsilon}{r}}h_G^3 > 0$ ,  $\sqrt{\frac{\log n}{nh_G^{5(1+\frac{1+\varepsilon}{r})}}} \ge 1$ ,  $h_G/h^{1+\frac{1+\varepsilon}{r}} \to 0$  hold.
- (C4) Let F(y|v) be the conditional distribution function of  $Y_i$  given  $q_0(X_i) = v$ . f(y|v) is the conditional density function of F(y|v) and has the first order continuous derivative at y = G(v). For any y in the neighbourhood of G(v), F(y|v) has the first order derivative with respect to  $v \in \mathcal{V}$ .
- (C5) Let  $f_{q_0}(y|z)$  be the conditional density function of  $\varepsilon_i$  given that  $q_0(X_i) = z$ . Furthermore,  $\frac{\partial^2 f_{q_0}(y|z)}{\partial z \partial y}$  exists in the neighbourhood of (0, v) for any  $v \in \mathcal{V}$ .

During the process of proving Theorem 5.1 in Appendix D, the following Lemma 5.1 is in fact proved. We list it here as an independent result.

Lemma 5.1. Under conditions of Theorem 5.1, it holds with probability one that

$$\frac{1}{nh_G^2} \sum_{m=1}^n K_G' \left( \frac{v - q_0(X_m)}{h_G} \right) \left( \widehat{q}_0(X_m) - q_0(X_m) \right) = o\left( \frac{1}{\sqrt{nh_G}} \right). \tag{5.1}$$

Lemma 5.2. i) Under conditions ii) of Theorem 5.1, it holds with probability one that

$$\left|\widehat{G}_n(v) - G(v)\right| = O\left(\left(\frac{1}{nh_G^{1 + \frac{1+\varepsilon}{r}}}\right)^{\frac{1}{2}}\right). \tag{5.2}$$

ii) Under conditions ii) of Theorem 5.1, with probability one, (5.2) holds uniformly with respect to  $v \in \mathcal{V}$ .

**Theorem 5.1.** Assume that the conditions of Theorem 4.1 and (C1)–(C4) hold. Then i) For any fixed  $v \in V$ , with probability one, we have the following asymptotic representation

$$\widehat{G}_n(v) - G(v) = \frac{1}{f(G(v)|v)} \left( (1 - \alpha) - F_n(G(v)|v) \right) + O\left(\frac{1}{\sqrt{nh^{1 + \frac{1+\varepsilon}{r}}}}\right).$$
 (5.3)

ii) Furthermore, if conditions (C1)-(C4) hold for any  $v \in \mathcal{V}$ , (5.3) holds uniformly for  $v \in \mathcal{V}$  with probability one.

**Corollary 5.1.** Under the conditions of Theorem 5.1 and condition (C5), we have that

$$\sqrt{\frac{nh_G f_{q_0}(v)}{\alpha (1-\alpha)}} \left( \widehat{G}_n(v) - G(v) - a(v) \left( 1 + o(1) \right) h_G^2 \right) \stackrel{d}{\to} \mathcal{N}(0,1), \tag{5.4}$$

where

$$a(v) = \frac{\int s^2 K_G(s) ds}{f_{q_0}(v)} \left[ \frac{\partial \left( f_{q_0}(0|v) f_{q_0}(v) G'(v) \right)}{\partial v} + f'_{q_0}(v) \left. \frac{\partial^2 f_{q_0}(y|v)}{\partial v \partial y} \right|_{v=0} \right].$$

**Remark 5.1.** From (5.4), the asymptotic mean squared error (AMSE) for  $\widehat{G}_n(v) - G(v)$  is equal to

$$\frac{\alpha (1 - \alpha)}{n h_G f_{q_0}(v)} + a^2(v) h_G^4 (1 + o(1)).$$

Hence, the optimal bandwidth of  $h_G$  in the sense of the AMSE is chosen as

$$h_{G,opt} = \left(\frac{\alpha (1 - \alpha)}{a^2(v) f_{q_0}(v)}\right)^{\frac{1}{5}} n^{-\frac{1}{5}}.$$

# 6 Concluding Remarks

This paper has been concerned with estimating the conditional quantile of a scalar random variable Y conditional on a vector of covariates X for a generalized additive model specification with an unknown link function. We have established various theoretical properties of the proposed estimators including consistency and asymptotic normality. This extension of estimating the generalized additive conditional mean regression model, is certainly non-trivial and demanding from a technical point of view for the large-sample properties of the proposed estimators. Furthermore, by allowing for a general form of serial dependence in the data, we enlarged the range of possible applications in practical situations.

#### References

Arcones, M.A. (1998). The law of large numbers for U-statistics under absolute regularity, *Elect. Comm. in Probab.*, 3, 13–19.

Cai, Z. and Ould-Saïd, E. (2003). Local M-estimator for nonparametric time series, Statist. Probab. Lett., 65(4), 433–449.

Chaudhuri, P. (1991). Global nonparametric estimation of conditional quantile functions and their derivatives, J. Multivariate Anal., 39(2), 246–269.

- Cheng, Y. (2007). Selected Topics on Nonparametric Conditional Quantiles and Risk Theory, Dissertation University of Amsterdam.
- Cheng, Y., De Gooijer, J.G. and Zerom, D. (2011). Efficient estimation of an additive quantile regression model, Scan. J. of Statistics, 38, 46–62.
- De Gooijer, J.G. and Zerom, D. (2003). On additive conditional quantiles with high-dimensional covariates, J. Amer. Statist. Assoc. 98, 135–146.
- Honda, T. (2004). Quantile regression in varying coefficient, J. Statist. Plann. Inference, 121, 113–125.
- Horowitz, J.L. (2001). Nonparametric estimation of a generalized additive model with an unknown link function, *Econometrica*, 69(2), 499–513.
- Horowitz , J.L. and Lee, S. (2005). Nonparametric estimation of an additive quantile regression model, J. Amer. Statist. Assoc., 100, 1238–1249.
- Lee, A.J. (1990). U-statistics: Theory and Practice, Marcel Dekker, Inc., New York.
- Lee, S. (2003). Efficient semiparametric estimation of a partially linear quantile regression model, *Econometric Theory*, 19, 1–31.
- Masry, E. (1996). Multivariate local polynomial regression for time series: Uniform strong consistency and rates, J. Time Ser. Anal., 17(6), 571–599.
- Mehra, K.L., Rama Krishnaiah, Y.S. and Rao, M.S. (1992). Bahadur representation of sample conditional quantiles based on smoothed conditional empirical distribution function, *Bull. Inform. Cybernet.*, 25(1-2), 99–107.
- Noh, H. and Lee, E.R. (2014). Component selection in additive quantile regression models. *J. Korean Statist. Soc.*, 43, 439–452.
- Stewart, G.W. and Sun J.G. (1990). Matrix Perturbation Theory, Academic Press: Boston, MA.
- Yoshihara, K. (1978). Probability inequalities for sums of absolutely regular processes and their applications, Z. Wahrsch. Verw. Gebiete, 43, 319–329.
- Yu, K. and Lu, Z. (2004). Local linear additive quantile regression, Scand. J. Statist., 31, 333–346.

# Supplementary Material for "Estimating Generalized Additive Conditional Quantiles by Absolutely Regular Processes"

By Yebin Cheng and Jan G. De Gooijer

# Appendices

#### A Preliminary Lemmas

In this Appendix we introduce two preliminary Lemmas A1 and A2 which are used in the proofs of Theorems 3.3, 4.1, and 5.3. Lemma A1 is a kind of Bernstein's inequality on absolutely regular processes. Lemma A2 is a moment inequality on degenerated U-statistics. This lemma extends Lemma 3 of Arcones (1998) to the case of higher moments. In the following two lemmas, we assume that  $\{\xi_i\}$ ,  $i=1,2,\ldots,n$ , is a sequence of strictly stationary random variables. Recall that  $\mathbb{E}_{j_0}g(\xi_i,\xi_{j_0})=g_1(\xi_{j_0})$ .

**Lemma A1.** Suppose that  $g(\cdot,\cdot)$  is a Borel measurable function with the bound M>0. Let  $g_2(\cdot)=\mathbb{E}g(\xi_i,\cdot)$ ,  $\sigma(\cdot)=Var(\sum_{i=1}^q g(\xi_i,\cdot))$  and  $U_{ij_0}=g(\xi_i,\xi_{j_0})-g_2(\xi_{j_0})$ , where  $1\leq j_0\leq n$  is fixed. Then, for any x>0,  $r_1>1$  and positive integer  $q\leq \frac{n}{4}$ , it holds that

$$\mathbb{P}\Big\{\Big|\sum_{1\leq i\leq n, i\neq j_0} U_{ij_0}\Big| \geq x\Big\} \leq 2\mathbb{E} \exp\Big\{\frac{-\left(\frac{x}{4}\right)^2}{\frac{n}{2q}\sigma(\xi_{j_0}) + \frac{2}{3}qM\frac{x}{4}}\Big\} + \frac{n}{q}\beta(q) + \frac{2^{r_1}q^{r_1-1}}{x^{r_1}} \sum_{|i-j_0|<2q} \mathbb{E}|U_{ij_0}|^{r_1}. \quad (A.1)^{r_1}$$

Proof. Assume that  $j_0 - 1 = m_1q + r_3$  and  $n - j_0 = m_2q + r_4$ , where  $m_1$  and  $m_2$  are two non-negative integers and  $0 \le r_3$ ,  $r_4 < q$ . The summation in the probability on the left-hand side (LHS) of (A.1) can be rewritten into a sum of two different summations according to whether the subscript i satisfies  $-(q+r_3) \le i - j_0 \le q + r_4$  or not. By using Markov's inequality, the last term on the RHS of the inequality (A.1) comes from the summation in which i satisfies the condition. Through exploring Berbee's lemma and Bernstein's inequality, the first two terms on the RHS of (A.1) can be derived from the summation in which i does not satisfy the condition.

**Remark 1.** The bound of  $\sigma(\cdot)$  in (A.1) can be obtained from Davydov's inequality, Hölder's inequality and  $C_r$ -inequality as follows

$$\sigma(\cdot) \le q \operatorname{Var}\left(g(\xi_1, \cdot)\right) + 2q \sum_{l=1}^{q} \left(\beta\left(l\right)\right)^{1-\frac{2}{r_2}} \left(\mathbb{E}\left|g(\xi_i, \cdot) - g_1(\cdot)\right|^{r_2}\right)^{\frac{2}{r_2}} \le Cq \left(\mathbb{E}\left|g(\xi_i, \cdot)\right|^{r_2}\right)^{\frac{2}{r_2}}, \tag{A.2}$$

where  $r_2$  satisfies that  $\sum_{l=1}^{\infty} (\beta(l))^{1-\frac{2}{r_2}} < \infty$  and  $1 - \frac{2}{r_2} = \frac{1+\varepsilon}{r}$  for a sufficiently small constant  $\varepsilon > 0$ . Without loss of generality (w.l.o.g.), q is assumed to be an integer throughout the next three appendices although it may not be sometimes according to its formula. In some cases, if  $\xi_{j_0}$  is replaced by a constant, then (A.1) still holds but its last term will be excluded.

**Lemma A2.** Let  $U_n = \sum_{1 \leq i < j \leq n} h_n(\xi_i, \xi_j)$  be a degenerated U-statistic with the symmetric kernel  $h_n(\cdot, \cdot)$ , i.e., for any  $t \in \mathbb{R}$ ,  $\mathbb{E}h(\xi_i, t) = 0$ . Then for  $k \in \mathbb{N}$ , there exists a universal constant C > 0 such that

$$\mathbb{E}U_n^k \le Cn^k \left(1 + \sum_{i=1}^{n-1} i^k \beta_i^{1-\frac{1}{s}}\right) M_{sk}^k, \tag{A.3}$$

where s > 1 and

$$M_{sk} = \sup_{(i_1, i_2), \mathbb{P}} \left( \int |h_n(\xi_{i_1}, \xi_{i_2})|^{sk} d\mathbb{P} \right)^{\frac{1}{sk}}$$

with  $\mathbb{P}$  being either the probability measure  $\mathbb{P}_{(\xi_{i_1},\xi_{i_2})}$  or  $\mathbb{P}_{\xi_{i_1}} \otimes \mathbb{P}_{\xi_{i_2}}$ .

*Proof.* Let  $J(i_1,\ldots,i_{2k})=\mathbb{E}[h_n(\xi_{i_1},\xi_{i_2})\cdots h_n(\xi_{i_{2k-1}},\xi_{i_{2k}})]$ . First, assume that  $i_1,\ldots,i_{2k}$  are different values. Rearrange the initial sequence  $i_1,\ldots,i_{2k}$  in natural order as  $j_1,\ldots,j_{2k}$ . Define  $d_1=j_2-j_1$ ,  $d_{2k} = j_{2k} - j_{2k-1}$  and  $d_i = \min\{j_i - j_{i-1}, j_{i+1} - j_i\}$  for  $i = 2, \dots, 2k-1$ . If,  $d_{i_0}$  is the largest number among  $\{d_i, i = 1, \ldots, 2k\}$ , then we compare the initial sequence  $(i_1, \ldots, i_{2k})$  with the one having the independent blocks  $(i_1, \ldots, i_{i_0-1})$ ,  $i_{i_0}$  and  $(i_{i_0+1}, \ldots, i_{2k})$  and the identical block distributions. Also, it can be inferred that there exist at least k numbers among  $(d_1, \ldots, d_{2k})$  which are in no excess of  $d_{i_0}$ . Thus, by part ii) of Arcones (1998, Lemma 2), monotonicity on  $\beta_i$  and Hölder's inequality, we obtain that

$$\sum_{i_1 \neq \dots \neq i_{2k}} J(i_1, \dots, i_{2k}) \le C n^k \sum_{i=1}^{n-1} i^k \beta_i^{1-\frac{1}{s}} M_{sk}^k.$$

The other cases of  $i_1, \ldots, i_{2k}$  can be dealt with similarly.

**Remark 2.** Usually, s is taken to be  $1 - \frac{k+1+\varepsilon}{r}$  with r > k+1.

#### Bahadur Representation

In this Appendix, and following an identical line as that of Honda (2004) and Chaudhuri (1991), we address the uniformly strong Bahadur representation for the conditional quantile and its derivatives at a random point. Let

$$V_{ij}(t_{1,u},\beta) = K_{ij,t_{1,u}} \left\{ \rho_{\alpha} \left( Y_i - \beta_{j,t_{1,u}}^{\mathsf{T}} A_{ij,t_{1,u}} \right) - \rho_{\alpha} \left( Y_i - \beta^{\mathsf{T}} A_{ij,t_{1,u}} \right) \right\}.$$

**Lemma B1.** If conditions (B1)-(B4) and  $r \ge d + \frac{1}{2} - \frac{d}{p} + \frac{17p+d-3}{2p(d-1-dp^{-1})}$  hold, then for any  $1 \le j \le n$ , it holds almost surely and uniformly both on  $\mathcal{X}_{1,u}$  and on  $|\beta - \beta_{j,t_{1,u}}| = \frac{B}{\sqrt{nh^{(1+\frac{1}{r})d+\varepsilon}}}$  that

$$\left| \sum_{i=1}^{n} (V_{ij}(t_{1,u}, \beta) - \mathbb{E}_{j} V_{ij}(t_{1,u}, \beta)) \right| \le Bh^{-\left(\frac{d}{r} + \varepsilon\right)}, \tag{B.1}$$

where B > 0 is a constant and may be chosen large.

*Proof.* By using Lemma A1 and taking  $q = \sqrt{nh^{\left(1-\frac{1}{r}\right)d}}$ , we have that

$$\mathbb{P}\left(\left|\sum_{i=1}^{n} \left(V_{ij}(t_{1,u},\beta) - \mathbb{E}V_{ij}(t_{1,u},\beta)\right)\right| > CBh^{-\left(\frac{d}{r}+\varepsilon\right)}\right) \le 2n^{-CB} + \frac{n}{q}\beta(q) + C\left(2h^{\frac{\varepsilon}{2}}\right)^{r_1}, \tag{B.2}$$

where the three terms on the RHS of (B.2) are derived from their counterparts in (A.1), respectively. More specifically, in the first term the relationship  $\sigma(Z_j) \leq Cqh^{\frac{2d}{r_2}}\delta^2$ , which follows from (A.2) and condition (B1), is used, where  $\delta = \frac{B}{\sqrt{nh^{(1+\frac{1}{r})d+\varepsilon}}}$  and  $1-\frac{2}{r_2}=\frac{1}{r}+\frac{\varepsilon}{2d}$ . And in the third term, the relationship  $\mathbb{E}|V_{ij}|^{r_1} \leq C\delta^{r_1}$  is used.

In order to prove uniformity related to  $t_{1,u}$  in (B.1), we divide  $\mathcal{X}_{1,u}$  into smaller squares with the side length  $\ell_n = \delta^2$ . For any point  $s = (s_1, s_u) \in \mathcal{X}_{1,u}$ , let  $t_{1,u} \in \mathcal{X}_{1,u}$  be the nearest grid points close to s. Then,  $|t_{1,u}-s| \leq \ell_n/\sqrt{2}$ . To prove uniformity related to  $\beta$  in (B.1), it is necessary to divide the two spheres  $|\beta - \beta_{j,t_{1,u}}| = B\delta$  and  $|\gamma - \beta_{j,s}| = B\delta$  into smaller cells with the radius  $d_1 = \ell_n$  such that the two divisions are location equivariant. It can be seen that the total number of such kind of cells related to each sphere is equal to  $O(\ell_n^{-\frac{d-1}{2}})$ . For any  $\alpha$  in the sphere  $|\gamma - \beta_{j,s}| = B\delta$ , let  $\beta$  in the sphere  $|\beta - \beta_{j,t_{1,u}}| = B\delta$  be the nearest grid point to the point which is in the sphere  $|\beta - \beta_{j,t_{1,u}}| = B\delta$  and is equivariant to  $\gamma$ . Then, it can be inferred that  $|\beta_{j,t_{1,u}} - \beta_{j,s}| \leq C\ell_n$  and  $|\gamma - \beta| \leq C(\ell_n + d_1) \leq C\ell_n$ . Next we prove that for s and  $t_{1,u}$  mentioned above, it holds that

$$|V_{ij}(s,\gamma) - V_{ij}(t_{1,u},\beta)| \le CB\ell_n \mathbb{I}_{\left(|X_i - \widetilde{X}_j| \le h\right)} + B\delta \mathbb{I}_{\left(h - \ell_n \le |X_i - \widetilde{X}_j| \le h + \ell_n\right)}. \tag{B.3}$$

In fact, if the two events  $|X_i - \widetilde{X}_{j,t_{1,u}}| \le h$  and  $|X_i - \widetilde{X}_{j,s}| \le h$  occur simultaneously, then from (B2), it

$$\left| (K_{ij,t_{1,u}} - K_{ij,s}) (\rho_{\alpha}(\varepsilon_i + r_{i,j,t_{1,u}}) - \rho_{\alpha}(\varepsilon_i + r_{i,j,t_{1,u}} + P_{i,j,t_{1,u}})) \right| \le C\ell_n \mathbb{I}_{(|X_i - \widetilde{X}_{j,t_{1,u}}| \le h)}$$
(B.4)

in view of  $\frac{\delta}{h} \to 0$ . Noting that

$$|r_{ij,t_{1,n}} - r_{ij,s}| \le |\beta_{j,t_{1,n}} - \beta_{j,s}| |A_{ij,t_{1,n}}| + |\beta_{j,s}| |A_{ij,t_{1,n}} - A_{ij,s}| \le C\ell_n, \tag{B.5}$$

we have

$$\left| K_{ij,s} \left\{ \left( \rho_{\alpha}(\varepsilon_{i} + r_{i,j,t_{1,u}}) - \rho_{\alpha}(\varepsilon_{i} + r_{i,j,t_{1,u}} + P_{i,j,t_{1,u}}) \right) - \left( \left( \rho_{\alpha}\varepsilon_{i} + r_{i,j,s} \right) - \rho_{\alpha}(\varepsilon_{i} + r_{i,j,s} + P_{i,j,s}^{\gamma}) \right) \right\} \right| \leq C \ell_{n} \mathbb{I}_{X_{i} - \widetilde{X}_{i,t_{1},u} \mid \leq h}.$$
(B.6)

From (B.4) and (B.6), it follows that

$$|V_{ij}(s,\gamma) - V_{ij}(t_{1,u},\beta)| \le C\ell_n \mathbb{I}_{(|X_i - \widetilde{X}_{i,t_{1,u}}| \le h)}$$

If  $|X_i - \widetilde{X}_{j,t_{1,u}}| \le h$  and  $|X_i - \widetilde{X}_{j,s}| > h$  occur simultaneously or  $|X_i - \widetilde{X}_{j,t_{1,u}}| > h$  and  $|X_i - \widetilde{X}_{j,s}| \le h$  occur simultaneously, then it can be inferred similarly that

$$|V_{ij}(s,\gamma) - V_{ij}(t_{1,u},\beta)| \le C\delta \mathbb{I}_{h-\ell_n \le |X_i - \widetilde{X}_{j,t_{1,u}}| \le h+\ell_n t}.$$

From the two cases above, we know that (B.3) holds. Thus, we consider the following two probabilities

$$\mathbb{P}\left(\sum_{i=1}^{n} \mathbb{I}_{\left(\left|X_{i}-\widetilde{X}_{j}\right| \leq h\right)} \geq \frac{CB}{\ell_{n}} h^{-\left(\frac{d}{r}+\varepsilon\right)}\right) \text{ and } \mathbb{P}\left(\sum_{i=1}^{n} \mathbb{I}_{\left(h-\ell_{n} \leq \left|X_{i}-\widetilde{X}_{j}\right| \leq h+\ell_{n}\right)} \geq CB\delta^{-1} h^{-\left(\frac{d}{r}+\varepsilon\right)}\right).$$

Similar as the proof of (B.2), the two probabilities above can also be bounded by the RHS of (B.2). For the first probability, to compare the two terms of the denominator in the first term on the RHS of (A.1), the fact that

$$\frac{n}{2q} \cdot q \left( \mathbb{E}_j \mathbb{I}_{\left( \left| X_i - \widetilde{X}_j \right| \le h \right)} \right)^{\frac{2}{r_2}} \le Cq \frac{h}{\ell_n} h^{-\left( \frac{d}{r} + \varepsilon \right)},$$

where  $r_2$  satisfies that  $1 - \frac{2}{r_2} = \frac{1}{r} + \frac{\varepsilon}{2d}$ , is used in view of  $nh^{\left(1 + \frac{1}{r}\right)d + \varepsilon} \to \infty$ . For the second probability, we use the fact that

$$\frac{n}{2q} \cdot q \left( \mathbb{E}_j \mathbb{I}_{\left(h-\ell_n \leq \left|X_i - \widetilde{X}_{j,t_{1,u}}\right| \leq h + \ell_n\right)} \right)^{\frac{2}{r_2}} \leq \frac{n}{2q} \cdot q \left( C\ell_n h^{d-1} \right)^{\frac{2}{r_2}} \leq Cq\delta^{-1} h^{-\left(\frac{d}{r} + \varepsilon\right)},$$

where  $r_2$  satisfies that  $1-\frac{2}{r_2}=\frac{1}{r}+\varepsilon_0$ , and the last inequality follows from  $\kappa<\frac{1-\frac{d}{r}}{d\left(1-\frac{1}{r}-\frac{d}{r^2}+\frac{1}{d}\right)}$ , and  $\varepsilon_0$  and  $\varepsilon$  are sufficiently small positive reals. Since the total number of small cells related to the divisions both on the domain  $\mathcal{X}_{1,u}$  and the sphere  $|\beta-\beta_j|=\delta$  is  $O(\ell_n^{-2}\cdot\ell_n^{\frac{d-1}{2}})=O(\ell_n^{-\frac{d+3}{2}})$ . Multiplying by  $O(\ell_n^{-\frac{d+3}{2}})$  on both sides of the inequality (B.2), then we can see that its RHS is controlled by  $\frac{C}{n(\log n)^2}$ , provided we choose B and  $r_1$  sufficiently large. Also, the inequality

$$\ell_n^{-\frac{d+3}{2}} \cdot \frac{n}{q} \cdot \beta(q) \le \frac{C}{n \left(\log n\right)^2} \tag{B.7}$$

follows from (B3) and the condition on r. Then, by the Borel-Cantelli Lemma, (B.1) holds.

**Lemma B2.** (Honda, 2004, Lemma 3.2) Under conditions (B1)–(B4), it holds uniformly on  $\mathcal{X}_{1,u}$  that

$$\left|\widehat{\beta}_{j,t_{1,u}} - \beta_{j,t_{1,u}}\right| = O\left(\frac{1}{\sqrt{nh^{(1+\frac{1}{r})d+\varepsilon}}}\right).$$

Let

$$\Delta_{ij}(t_{1,u},\beta) = K_{ij,t_{1,u}} A_{ij,t_{1,u}} \left[ \mathbb{I} \left( \varepsilon_i \le -r_{ij,t_{1,u}} - P_{ij,t_{1,u}} \right) - \mathbb{I} \left( \varepsilon_i \le -r_{ij,t_{1,u}} \right) \right]. \tag{B.8}$$

**Lemma B3.** Under conditions (B1)-(B4) and  $r \ge \max\left\{d-7+\frac{2d^2-4-4d}{p}-\frac{22p+6}{dp},d\right\}$ , with probability one, it holds uniformly on  $t_{1,u} \in \mathcal{X}_{1,u}$  and the sphere  $\left|\beta-\beta_{j,t_{1,u}}\right| = \frac{B}{\sqrt{nh^{(1+\frac{1}{n})d+\varepsilon}}}$  that

$$\left| \sum_{i=1}^{n} \left( \Delta_{ij}(t_{1,u}, \beta) - \mathbb{E}_{j} \Delta_{ij}(t_{1,u}, \beta) \right) \right| \le B \left( n^{1 + \frac{1}{r} + \varepsilon} h^{d\left(1 - \frac{1}{r}\right)^{2}} \right)^{\frac{1}{4}}. \tag{B.9}$$

**Theorem B1.** Under conditions (B1)–(B4), the following strong Bahadur representation

$$\widehat{\beta}_{j,t_{1,u}} - \beta_{j,t_{1,u}} = \frac{Q_{jn,t_{1,u}}^{-1}}{nh^d} \sum_{i=1}^{n} K_{ij,t_{1,u}} A_{ij,t_{1,u}} \left( (1-\alpha) - \mathbb{I}\left(\varepsilon_i \le -r_{i,j}\right) \right) + O\left( \left( n^{1-\frac{1}{3r} - \frac{2\varepsilon}{3}} h^{d\left(1 + \frac{2}{3r} - \frac{1}{3r^2}\right)} \right)^{-\frac{3}{4}} \right)$$
(B.10)

holds almost surely and uniformly for  $1 \le j \le n$ ,  $t_1$  and  $t_u$ .

*Proof.* We first note that

$$\sum_{i=1}^{n} K_{ij,t_{1,u}} A_{ij,t_{1,u}} \left[ (1-\alpha) - \mathbb{I} \left( \varepsilon_i \le -r_{i,j} \right) \right]$$

$$= n \mathbb{E}_j \Delta_{ij}(t_{1,u},\beta) + \sum_{i=1}^{n} \left( \Delta_{ij}(t_{1,u},\beta) - \mathbb{E}_j \Delta_{ij}(t_{1,u},\beta) \right) + R_n(\beta), \tag{B.11}$$

where

$$R_n(\beta) = \sum_{i=1}^n K_{ij,t_{1,u}} A_{ij,t_{1,u}} ((1-\alpha) - \mathbb{I}(\varepsilon_i \le -r_{ij,t_{1,u}} - P_{ij,t_{1,u}})).$$

Then, using condition (B4) and Taylor's expansion for  $g(X_i,\cdot)$ , it can be inferred that

$$n\mathbb{E}_{j}\Delta_{ij}(t_{1,u},\beta) = n\mathbb{E}_{j}K_{ij,t_{1,u}}A_{ij,t_{1,u}} \left[ G(X_{i}, -r_{ij,t_{1,u}} - P_{ij,t_{1,u}}) - G(X_{i}, -r_{ij,t_{1,u}}) \right]$$

$$= nh^{d}Q_{jn,t_{1,u}} \left( \left( \beta - \beta_{j,t_{1,u}} \right) + O(|\delta|^{2} + |\delta| h^{p}) \right). \tag{B.12}$$

Under condition (B3), we have that  $nh^d |\delta|^2 = O(\delta_n)$  and  $nh^d |\delta| h^p = O(\delta_n)$ . Also, there exists a constant  $\phi > 0$  such that  $|R_n(\widehat{\beta}_{j,t_{1,u}})| \le \phi$  holds almost surely. Thus, (B.10) holds.

#### C Proofs of Theorems

### Proof of Theorem 3.1

*Proof.* i) In considering (2.7), it can be seen that  $\widehat{q}_u(x_u) - q_u(x_u) = I_1 - I_2 - I_3$ , where  $I_1 = \int_{x_{u,0}}^{x_u} \int \frac{w_1(t_1)\Delta_u}{D_{1,u}(t_1,t_u)} dt_u dt_1$ ,  $I_2 = \int_{x_{u,0}}^{x_u} \int \frac{\Delta_{1,u}D_u(t_1,t_u)}{D_{1,u}^2(t_1,t_u)} w_1(t_1) dt_u dt_1$  and

$$I_3 = \int_{x_{u,0}}^{x_u} \int \frac{\left(\Delta_u D_{1,u}(t_1, t_u) - \Delta_{1,u} D_u(t_1, t_u)\right) \Delta_{1,u}}{\left(\Delta_{1,u} + D_{1,u}(t_1, t_u)\right) D_{1,u}^2(t_1, t_u)} w_1(t_1) dt_u dt_1.$$

From Lemma 3.2, we know that  $\sqrt{nh}I_3 = O\left(\left(nh^{7+\frac{2+2\varepsilon}{r}}\right)^{-\frac{1}{2}}\right)$ .

Next we deal with  $I_1$ . Note that  $\Delta_u$  can be decomposed into two terms as

$$\frac{1}{n}\sum_{j=1}^{n}\left(\partial_{u}q(\widetilde{X}_{j})\mathbb{I}(X_{j,\bar{u}}\in\mathcal{X}_{\bar{u}})-D_{u}(t_{1},t_{u})\right)+\frac{1}{n}\sum_{j=1}^{n}\left(\partial_{u}\widehat{q}(\widetilde{X}_{j})-\partial_{u}q(\widetilde{X}_{j})\right)\mathbb{I}\left(X_{j,\bar{u}}\in\mathcal{X}_{\bar{u}}\right).$$

Then, substituting this expression into  $I_1$ , we obtain two terms, say  $I_{11}$  and  $I_{12}$ , respectively. For  $I_{11}$ , it is included in  $\xi_{n1}$  given in Remark 3.1. This term, however, is not essential for understanding the results of this paper. In considering Theorem B1, we conclude that  $I_{12}$  is equal to

$$\frac{1}{n^{2}h^{d+1}} \sum_{j=1}^{n} \sum_{1 \leq i \neq j \leq n} \int_{x_{u}}^{x_{u,0}} \int e_{u}^{T} Q_{jn,t_{1,u}}^{-1} K_{ij,t_{1,u}} A_{ij,t_{1,u}} ((1-\alpha) - \mathbb{I}(\varepsilon_{i} \leq -r_{i,j})) dt_{1} dt_{u} \cdot \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{\bar{u}}) + O\left(\frac{1}{h} \left(n^{1-\frac{1}{3r} - \frac{2\varepsilon}{3}} h^{d\left(1+\frac{2}{3r} - \frac{1}{3r^{2}}\right)}\right)^{-\frac{3}{4}}\right).$$

Denote by  $I_{13}$  the first term of the expression above. Then, using Lemma 3.1, it holds with probability one that

$$I_{13} = \frac{1}{n^2 h^{d+1}} \sum_{1 \le i \ne j \le n} \eta(Z_i, Z_j) + B_{1,u} h^p (1 + o(1))$$
(C.1)

with

$$\eta(Z_i, Z_j) = e_u^{\mathrm{T}} \int_{x_{u,0}}^{x_u} \int Q_{jn,t_{1,u}}^{-1} K_{ij,t_{1,u}} A_{ij,t_{1,u}} \frac{((1-\alpha) - \mathbb{I}(\varepsilon_i \leq 0)) w_1(t_1)}{D_{1,u}(t_1, t_u)} dt_1 dt_u \cdot \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{\bar{u}}).$$

Let  $\psi(Z_i, Z_j) = \eta(Z_i, Z_j) + \eta(Z_j, Z_i)$ ,  $\psi_i = \mathbb{E}\psi(z, Z_j)|_{z=Z_i}$ ,  $\varphi_{ij} = \psi(Z_i, Z_j) - \psi_i - \psi_j$ ,  $U_n = \frac{1}{n^2h^{d+1}} \sum_{1 \leq i < j \leq n} \varphi_{ij}$  and  $I_4$  be the first term on the RHS of relationship (C.1). Taking into account the Hoeffding decomposition of an U-statistic (see, e.g., Lee (1990)), we rewrite  $I_4$  as  $I_4 = U_n + \frac{n-1}{n^2h^{d+1}} \sum_{i=1}^n \psi_i$ . Then, it can be inferred from Lemma A2 that

$$\mathbb{E}U_n^2 \le \frac{Cn^2M^2}{\left(n^2h^{d+1}\right)^2} \left(1 + \sum_{j=1}^{n-1} j^2 \beta_j^{(r_3-2)/r_3}\right),\,$$

where  $M = \sup_{i,j} (\mathbb{E} |\varphi_{ij}|^{r_3})^{\frac{1}{r_3}}$  for some  $r_3 > 2$  satisfying that  $1 - \frac{2}{r_3} = \frac{2+\varepsilon}{r}$ . It can be inferred from condition (B7) that  $\sup_{i,j} \mathbb{E} |\eta(Z_i,Z_j)|^{r_3} \leq Ch^d$ , and thus

$$\mathbb{E}U_n^2 \le \frac{C}{n^2 h^{d\left(1 + \frac{2+\varepsilon}{r}\right) + 2}},\tag{C.2}$$

i.e.,  $\sqrt{nh}U_n = O_{\mathbb{P}}\left(\left(nh^{d+1+\frac{d(2+\varepsilon)}{r}}\right)^{-\frac{1}{2}}\right)$ . Also, from the property of the conditional expectation, it follows that

$$\psi_i = ((1 - \alpha) - \mathbb{I}(\varepsilon_i \le 0))\zeta_i, \tag{C.3}$$

where

$$\zeta_i = \zeta(X_i) = \mathbb{E}_i \int_{x_u}^{x_u} \int \frac{w_1(t_1) e_u^{\mathrm{T}} Q_{jn,t_{1,u}}^{-1} K_{ij,t_{1,u}} A_{ij,t_{1,u}}}{D_{1,u}(t_1,t_u)} dt_1 dt_u \cdot \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{\bar{u}}).$$

Next, we consider the asymptotic expression of  $\sum_{i=1}^{n} \psi_i$ . Note that the domain of the covariates of  $\zeta(X_i)$  is  $A_{(u)}$ , which is defined at the beginning of Section 3. We then divide  $A_{(u)}$  into a sequence of subsets  $\{A_l\}$  and try to get the asymptotic representations  $\{M_l\}$  of  $\zeta(X_i)$  on these subsets, respectively. Let  $M(X_i)$  be the summation of all these  $\{M_l\}$ . Without loss of generality, we only consider some special cases of  $\{A_l\}$ , all other left cases of  $\{A_l\}$  can be settled similarly. By the inequality (A.2), we see that

$$\operatorname{Var}\left(\frac{(n-1)}{n^{2}h^{d+1}}\sum_{i=1}^{n}\left(1-\alpha-\mathbb{I}(\varepsilon_{i}\leq0)\right)\left(\zeta_{i}-M(X_{i})\right)\right)\leq\frac{C}{nh^{2d+2}}\left(\mathbb{E}\left|\zeta(X_{i})-M(X_{i})\right|^{r_{2}}\right)^{\frac{2}{r_{2}}}$$
(C.4)

for  $1 - \frac{2}{r_2} = \frac{1+\varepsilon}{r}$ . Let

$$A_{1} = \left[x_{u,0} - h, x_{u,0} + h\right] \times \Pi_{1 \leq l \neq u \leq d} \left[a_{l} - h, b_{l} + h\right], \qquad I_{u1} = \left(\int_{A_{1}} \left|\zeta\left(x\right) - M_{1}(x)\right|^{r_{2}} p(x) dx\right)^{\frac{2}{r_{2}}},$$

$$A_{2} = \left[a_{1} - h, a_{1} + h\right] \times \Pi_{2 \leq l \leq d} \left[a_{l} - h, b_{l} + h\right], \qquad I_{u2} = \left(\int_{A_{2}} \left|\zeta\left(x\right) - M_{2}(x)\right|^{r_{2}} p(x) dx\right)^{\frac{2}{r_{2}}},$$

$$A_{c} = \left[x_{u,0} + h, x_{u} - h\right] \times \Pi_{1 \leq l \neq u \leq d} \left[a_{l} + h, b_{l} - h\right], \qquad I_{c} = \left(\int_{A_{c}} \left|\zeta\left(x\right) - M_{c}(x)\right|^{r_{2}} p(x) dx\right)^{\frac{2}{r_{2}}}.$$

We now deal with each term mentioned above separately. First, we consider  $I_{u1}$ . By variable substitution, it can be seen from condition (B5) that

$$Q_{jn,t_{1,u}} = \int K(x)A(x)A^{\mathrm{T}}(x)g_1(\widetilde{X}_j + hx)\mathrm{d}x = g_1(\widetilde{X}_j)Q + hQ^* \left. \frac{\partial g_1(t)}{\partial t} \right|_{t=\widetilde{X}_j} + O(h^2).$$

Then, from Newman's expansion (see, e.g., Stewart and Sun, 1990) it can be established that

$$Q_{jn,t_{1,u}}^{-1} = \frac{Q^{-1}}{g_1(\widetilde{X}_j)} - \frac{hQ^{-1}Q^*}{g_1^2(\widetilde{X}_j)} \cdot \frac{\partial g_1(t)}{\partial t} \bigg|_{t=\widetilde{X}_j} Q^{-1} + O(h^2).$$
 (C.5)

Let  $B_k = \prod_{1 \le l \ne k \le d} [a_l + h, b_l - h] \subset \mathbb{R}^{d-1}$ ,

$$B_{k,i} = [a_i - h, a_i + h] \times \prod_{l=1, l \neq k \neq i}^d [a_l - h, b_l + h] \subset \mathbb{R}^{d-1},$$

$$M_{1}(z) = \int_{x_{u},0}^{x_{u}} \int \frac{e_{u}^{\mathrm{T}} Q^{-1} w_{1}(t_{1}) A\left(\frac{z-t}{h}\right) K\left(\frac{z-t}{h}\right) p_{t_{\bar{u}}}(t_{\bar{u}})}{g_{1}(t) D_{1,u}(t_{1},t_{u})} \mathbb{I}(t_{\bar{u}} \in \mathcal{X}_{\bar{u}}) dt \cdot \mathbb{I}(z \in A_{1})$$
(C.6)

and

$$M_u(z) = \frac{w_1(z_1)p_{\bar{u}}(z_{\bar{u}})e_u^{\mathrm{T}}Q^{-1}}{g_1(z)D_{1,u}(z_1, z_u)} \int_{-1}^{\frac{z_u - x_{u,0}}{h}} \int A(t)K(t)\mathrm{d}t \cdot \mathbb{I}(z \in A_1). \tag{C.7}$$

Then, by using (C.5) and variable substitution, we obtain

$$I_{u1} \leq \frac{C}{nh^{2d+2}} \left\{ \int_{x_{u,0}-h}^{x_{u,0}+h} \int \left| h^d \int_{-1}^{\frac{z_u-x_u}{h}} \int \left( -\frac{he_u^{\mathsf{T}} Q^{-1} Q^*}{g_1^2(x-ht)} \frac{\partial g_1(x-ht)}{\partial t} Q^{-1} + O(h^2) \right) \right. \\ \left. \frac{w_1(x_1-ht_1)A(t)K(t)}{D_{1,u}(x_1-ht_1,x_u-ht_u)} p_{\bar{u}}(x_{\bar{u}}-ht_{\bar{u}}) \mathrm{d}t \right|^{r_2} p(x) \mathrm{d}x \right\}^{\frac{2}{r_2}} \leq \frac{C}{nh} h^{\frac{2}{r_2}+1}.$$

In view of (C.6), (C.7) and variable substitution, it can be inferred that

$$\frac{1}{nh^{2+2d}} \left[ \int_{x_{u,0}-h}^{x_{u,0}+h} \int_{B_u} \left| M_1(z) - h^d M_u(z) \right|^{r_2} p(z) dz \right]^{\frac{2}{r_2}} \le \frac{C}{nh} h^{\frac{2}{r_2}+1}$$

and

$$\frac{1}{nh^{2+2d}} \left[ \sum_{k=1, k \neq u}^{d} \int_{x_{u,0}-h}^{x_{u,0}+h} \int_{B_{u,i}} \left| M_1(z) - h^d M_{u,k}(x_{u,0}, a_k, z) \right|^{r_2} p(z) dz \right]^{\frac{2}{r_2}} \leq \frac{C}{nh} h^{\frac{4}{r_2}+1}.$$

Hence, from the three inequalities above and the Cramér-Rao inequality, it can be seen that  $\frac{M_u(z)}{\sqrt{nh}}$  is included on the RHS of (3.3) with the remainder term  $O_{\mathbb{P}}(h^{\frac{1}{r_2}+\frac{1}{2}}+h^{\frac{d}{r_2}-\frac{1}{2}})$ .

Next, we consider  $I_{u2}$ . Let

$$I_{u2,1} = \left[ \int_{a_1 - h}^{a_1 + h} \int_{B_1} \left| \int_{x_{u,0}}^{x_u} \int \frac{w_1(t_1) e_u^{\mathrm{T}} Q_{jn}^{-1} A\left(\frac{z - t}{h}\right) K\left(\frac{z - t}{h}\right) p(t_{\bar{u}})}{D_{1,u}(t_1, t_u)} dt \right|^{r_2} p(z) dz \right]^{\frac{2}{r_2}},$$

$$I_{u2,2} = \left[ \int_{a_1 - h}^{a_1 + h} \int_{B_1^c} \left| \int_{x_{u,0}}^{x_u} \int \frac{w_1(t_1) e_u^{\mathrm{T}} Q_{jn}^{-1} A\left(\frac{z - t}{h}\right) K\left(\frac{z - t}{h}\right) p(t_{\bar{u}})}{D_{1,u}(t_1, t_u)} dt \right|^{r_2} p(z) dz \right]^{\frac{z}{r_2}}.$$

Obviously,  $I_{u2} \leq I_{u2,1} + I_{u2,2}$ . According to (C.5), variable substitution and the known condition w(t) = O(t) as  $t \to a$ , we obtain

$$\frac{I_{u2,1}}{nh^{2d+2}} \le \frac{C_p}{nh^2} \left\{ \int_{a-h}^{a+h} \int_{B_1} \left| \int_{-1}^{\frac{z_1-a}{h}} \int w_1 (z_1 - ht_1) \left( \frac{Q^{-1}}{g_1(z - ht)} + O(h) \right) \right. \\
\left. \cdot \frac{A(t)K(t)p (z_{\bar{u}} - ht_{\bar{u}})}{D_{1,u} (z_1 - ht_1, z_u - ht_u)} dt \right|^{r_2} f(z) dz \right\}^{\frac{2}{r_2}} \le \frac{C_{r_2}}{nh} h^{1 + \frac{2}{r_2}}.$$

Similarly, it can be shown that  $\frac{I_{u2,2}}{nh^{2d+2}} = O\left(\frac{1}{nh}h^{1+\frac{2}{r_2}}\right)$ . Therefore,  $I_{u2} = O\left(\frac{1}{nh}h^{1+\frac{2}{r_2}}\right)$ .

Finally, we consider  $I_c$ . Let

$$M_{c}(Z_{i}) = h^{d+1} \cdot e_{u}^{\mathsf{T}} Q^{-1} \int \left. \frac{\partial}{\partial x^{\mathsf{T}}} \left( \frac{w_{1}(x_{1}) p_{\bar{u}}(x_{\bar{u}})}{g_{1}(x) D_{1,u}(x_{1}, x_{u})} \right) \right|_{x = \widetilde{X}_{i}} t A(t) K(t) dt \cdot \mathbb{I}(X_{i} \in A_{(u)}).$$

From the fact that  $e_u^T Q^{-1} \int A(t)K(t)dt = 0$  and (C.5), by exploiting variable substitution and Taylor's expansion, we have

$$\frac{I_c}{nh^{2d+2}} \le \frac{Ch^2}{n}.$$

Thus,  $nh\frac{I_c}{n(h^{d+1})^2} = O(h^3)$  so that  $h^{\frac{3}{2}}$  appears in the remainder term. Finally, we note that the contribution of term  $I_2$  is similar to that of  $I_1$  with  $e_u^{\mathrm{T}}$  replaced by  $-D_u/D_1e_1^{\mathrm{T}}$ , which follows from comparing the expressions of  $\Delta_u$  and  $\Delta_{1,u}$ . This completes the proof of i).

ii) It can be seen from (2.8) and (2.9) that

$$\widehat{q}_1(x_1) - q_1(x_1) = c(x_1)(\widehat{c} - c) + (\widehat{c} - c) \int_{x_{1,0}}^{x_1} L_n(t_1) dt_1 + c \int_{x_{1,0}}^{x_1} L_n(t_1) dt_1,$$
 (C.8)

where

$$L_n(t_1) = \int \left( \frac{\widehat{D}_{1,2}(t_1, t_2)}{\widehat{D}_2(t_1, t_2)} - \frac{D_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} \right) w_2(t_2) dt_2.$$

For  $\hat{c} - c$ , it can be rewritten as  $I_{11} + I_{12}$ , where

$$I_{11} = -\int \frac{L_n(t_1)w_1(t_1)}{\left(\int \frac{D_{1,2}(t_1,t_2)}{D_2(t_1,t_2)}w_2(t_2)dt_2\right)^2} dt_1$$

and

$$I_{12} = \int \frac{w_1(t_1)L_n^2(t_1)}{\int \frac{\widehat{D}_{1,2}(t_1,t_2)}{\widehat{D}_2(t_1,t_2)} w_2(t_2) dt_2 \left(\int \frac{D_{1,2}(t_1,t_2)}{D_2(t_1,t_2)} w_2(t_2) dt_2\right)^2} dt_1.$$

From Lemma 3.2 it follows that

$$\sqrt{nh} \left| I_{12} \right| = \sqrt{nh} O\left( \sup_{t_1} L_n^2\left(t_1\right) \right) = O\left( \left( nh^{7 + \frac{2+2\varepsilon}{r}} \right)^{-\frac{1}{2}} \right).$$

As for  $I_{11}$ , it is equal to  $I_{11} = -I_{111} - I_{112} + I_{113} + I_{114}$ , where

$$I_{111} = \iint \frac{w_1(t_1)w_2(t_2)\Delta_{1,2}(t_1,t_2)}{D_2(t_1,t_2)} dt_2 dt_1,$$

$$I_{112} = \iint \frac{\Delta_{1,2}(t_1,t_2)\Delta_2(t_1,t_2)}{\widehat{D}_2(t_1,t_2)D_2(t_1,t_2)} w_1(t_1)w_2(t_2) dt_2 dt_1,$$

$$I_{113} = \iint \frac{\Delta_2(t_1,t_2)D_{1,2}(t_1,t_2)}{D_2^2(t_1,t_2)} w_1(t_1)w_2(t_2) dt_2 dt_1,$$

and

$$I_{114} = \iint \frac{\Delta_2^2(t_1, t_2) D_{1,2}(t_1, t_2)}{\widehat{D}_2(t_1, t_2) D_2^2(t_1, t_2)} w_1(t_1) w_2(t_2) dt_2 dt_1.$$

Using Lemma 3.2, we have

$$I_{112} = O\left(\left(nh^{4+\frac{1+\varepsilon}{r}}\right)^{-1}\right), \quad I_{114} = O\left(\left(nh^{4+\frac{1+\varepsilon}{r}}\right)^{-1}\right).$$

We now consider  $I_{111}$ . Similar as before, note that  $\Delta_{1,2}(t_1,t_2)$  can be rewritten as

$$\frac{1}{n}\sum_{j=1}^{n} \left(\partial_1 \widehat{q}(t_1, t_2, X_{j, \overline{2}}) - \partial_1 q(t_1, t_2, X_{j, \overline{2}})\right) \mathbb{I}(X_{j, \overline{2}} \in \mathcal{X}_{\overline{2}})$$

$$+\frac{1}{n}\sum_{j=1}^{n} \left(\partial_{1}q(t_{1},t_{2},X_{j,\bar{2}})\mathbb{I}(X_{j,\bar{2}}\in\mathcal{X}_{\bar{2}})-D_{1,2}(t_{1},t_{2})\right).$$

Substituting this into  $I_{111}$ , we derive two terms from this, say  $I_{115}$  and  $I_{116}$ . Then,  $I_{116}$  is included in (3.5). This term is not essential for understanding the proof in this case, and hence has been omitted. As for  $I_{115}$ , by using Theorem B1 and Lemma 3.1, it can be shown that

$$I_{115} = \frac{1}{n^2 h^{d+1}} \sum_{j=1}^n \iint e_1^{\mathrm{T}} Q_{jn}^{-1} \sum_{i=1}^n K_{ij,t_{1,2}} A_{ij,t_{1,2}} \left( (1-\alpha) - \mathbb{I}(\varepsilon_i \leq 0) \right) \frac{w_2(t_2) w_1(t_1)}{D_2(t_1, t_2)} dt_1 dt_2 + O\left(\frac{1}{h} \left( n^{1-\frac{1}{3r} - \frac{2\varepsilon}{3}} h^{d\left(1+\frac{2}{3r} - \frac{1}{3r^2}\right)} \right)^{-\frac{3}{4}} + h^{p-1} \right).$$

According to the previous analysis and by the Hoeffding decomposition the leading term of  $I_{115}$  is equal to

$$\frac{1}{nh^{d+1}} \sum_{i=1}^{n} \left( (1-\alpha) - \mathbb{I}(\varepsilon_i \leq 0) \right) \mathbb{E}_i \iint \frac{w_1(t_1)w_2(t_2)e_1^{\mathrm{T}} Q_{jn}^{-1} K_{ij,t_{1,2}} A_{ij,t_{1,2}} \mathbb{I}(X_{j,\bar{2}} \in \mathcal{X}_{\bar{2}})}{D_2(t_1,t_2)} \mathrm{d}t_1 \mathrm{d}t_2.$$

And then, by a similar method as the proof of part i), it can be shown that the leading term of  $I_{115}$  is equal to

$$\sum_{i=1}^{n} \frac{\left( (1-\alpha) - \mathbb{I}(\varepsilon_{i} \leq 0) \right) w_{1}(X_{i,1}) w_{2}(X_{i,2}) p(X_{i,\bar{2}}) e_{1}^{\mathsf{T}} Q^{-1}}{nh D_{2}(X_{i,1}, X_{i,2}) g_{1}(X_{i})} \sum_{3 \leq k \leq d} \left( f_{k} \left( \frac{X_{i,k} - b_{k}}{h} \right) - f_{k} \left( \frac{X_{i,l} - a_{l}}{h} \right) \right).$$

Analogously, we can deal with  $I_{113}$ . Its leading term is given by

$$\sum_{i=1}^{n} \frac{\left( (1-\alpha) - \mathbb{I}(\varepsilon_{i} \leq 0) \right) w_{1}(X_{i,1}) w_{2}(X_{i,2}) p(X_{i,\bar{2}}) D_{1,2}(X_{i,1}, X_{i,2}) e_{2}^{\mathsf{T}} Q^{-1}}{n h g_{1}(X_{i}) D_{2}^{2}(X_{i,1}, X_{i,2})} \cdot \sum_{3 \leq k \leq d} \left( f_{k} \left( \frac{X_{i,k} - b_{k}}{h} \right) - f_{k} \left( \frac{X_{i,l} - a_{l}}{h} \right) \right).$$

Let  $I_2 = c \int_{x_{1,0}}^{x_1} L_n(t_1) dt_1$ ,

$$I_{21} = \int_{x_{1.0}}^{x_1} \int \frac{w_2(t_2)\Delta_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} dt_2 dt_1, \quad I_{22} = \int_{x_{1.0}}^{x_1} \int \frac{w_2(t_2)\Delta_2(t_1, t_2)D_{1,2}(t_1, t_2)}{D_2^2(t_1, t_2)} dt_2 dt_1$$

and

$$I_{23} = \int_{x_1 \text{ o}}^{x_1} \!\! \int \!\! \left( \Delta_{12}(t_1, t_2) - \Delta_2(t_1, t_2) \frac{D_{1,2}(t_1, t_2)}{D_2(t_1, t_2)} \right) \frac{\Delta_2(t_1, t_2) w_2(t_2)}{\left( \Delta_2(t_1, t_2) + D_2(t_1, t_2) \right) D_2(t_1, t_2)} \mathrm{d}t_2 \mathrm{d}t_1.$$

Then,  $I_2 = (I_{21} - I_{22} - I_{23}) c$ . Similarly, the leading term of  $I_{21}$  is equal to

$$\frac{1}{nh^{d+1}} \sum_{i=1}^{n} \left( \left( 1 - \alpha \right) \mathbb{I}(\varepsilon_i \leq 0) \right) \mathbb{E}_i \int_{x_{1,0}}^{x_1} \int \frac{w_2(t_2) e_1^{\mathrm{T}} Q^{-1} K_{ij,t_{1,2}} A_{ij,t_{1,2}} \mathbb{I}(X_{j,\bar{2}} \in \mathcal{X}_{\bar{2}})}{D_2(t_1,t_2)} \mathrm{d}t_2 \mathrm{d}t_1.$$

By an analogous method, we can obtain the asymptotic representations of  $I_{21}$  and  $I_{22}$ , which are included in (3.4). This completes the proof of ii).

**Remark 3.** The argument of the proof of Theorem 3.1 uses  $x_u \ge u_{u,o}$  at some steps, for instance to derive the integration boundaries in (C.7). A similar argument is possible for  $x_u < x_{u,0}$ .

#### Proof of Theorem 4.1

Proof. We first consider the case  $2 \le u \le d$ . Let  $I_1$ ,  $I_2$  and  $I_3$  be the same notations as those in part i) of Theorem 3.1 and then  $\widehat{q}_u(x_u) - q_u(x_u)$  is equal to  $I_1 - I_2 - I_3$ . From Lemma 3.2, we know that  $\sqrt{nh} \sup_{x_u \in [a_u,b_u]} |I_3| = O\left(\left(nh^{7+\frac{2+2\varepsilon}{r}}\right)^{-\frac{1}{2}}\right)$ . For brevity, let  $\gamma_n = \sqrt{\frac{\log n}{nh^{1+\frac{1+\varepsilon}{r}}}}$ . Next, we only consider the uniform convergence rate of  $I_1$ , since the methodology to deal with  $I_1$  and  $I_2$  is almost completely the same. In view of Theorem B1 and Lemma 3.1, we know that  $I_1 = \phi(x_u) + o(\gamma_n)$ , where  $\phi(x_u) = \int_{x_{u,0}}^{x_u} \psi_n(t_u) \mathrm{d}t_u$ ,  $\psi_n(t_u) = \frac{1}{n^2h^{d+1}} \sum_{j=1}^n \sum_{i=1, i\neq j}^n \varphi_{ij}(t_u)$  and

$$\varphi_{ij}(t_u) = \left( (1 - \alpha) - \mathbb{I}(\varepsilon_i \le 0) \right) \int \frac{e_u^{\mathsf{T}} Q_{j,n}^{-1}(t_1, t_u) w_1(t_1) K_{ij, t_{1,u}} A_{ij, t_{1,u}}}{D_{1,u}(t_1, t_u)} dt_1.$$

In the sequel, we only need to prove that

$$\sup_{x_u \in [a_u, b_u]} \phi(x_u) = O(\gamma_n). \tag{C.9}$$

As usual, we divide the interval  $[a_u, b_u]$  into a sequence of disjoint subintervals, the length of which is equal to  $\ell_n$ . Without loss of generality, we assume that  $(b_u - a_u)\ell_n^{-1}$  is an integer, and  $\{x_{u,k}, k = 1, \dots, C\ell_n^{-1}\}$  are the corresponding grid points. Then,

$$\mathbb{P}\left(\sup_{x_{u}\in[a_{u},b_{u}]}\left|\phi\left(x_{u}\right)\right|\geq\gamma_{n}\right)\leq\sum_{1\leq k\leq\ell_{n}^{-1}}\mathbb{P}\left(\sup_{t_{u}\in[x_{u,k},x_{u,k+1}]}\left|\psi_{n}(t_{u})-\psi_{n}(x_{u,k})\right|\geq\frac{\gamma_{n}}{3\ell_{n}}\right) + \sum_{1\leq k\leq\ell_{n}^{-1}}\mathbb{P}\left(\left|\psi_{n}(x_{u,k})\right|\geq\frac{\gamma_{n}}{3\ell_{n}}\right) + \sum_{1\leq k\leq\ell_{n}^{-1}}\mathbb{P}\left(\left|\phi(x_{u,k})\right|\geq\frac{\gamma_{n}}{3}\right). \tag{C.10}$$

We now consider the super bound of the first term on the RHS of (C.10). From condition (B2), there exists another kernel function  $K_{-u}(\cdot)$  defined on  $\mathbb{R}^{d-1}$  such that

$$\left| \frac{\partial K(t)}{\partial t_u} \right| \le CK_{-u} \left( t_{-u} \right), \tag{C.11}$$

where  $t_{-u} \in \mathbb{R}^{d-1}$  is obtained from t by deleting its uth component. Similar as the proof of inequality (D.1) in Appendix D, we know that  $\|Q_{j,n}^{-1}(t_1,t_u)-Q_{j,n}^{-1}(t_1,x_{u,k})\| \leq C\ell_n$ . Also, from condition (B8), it can be seen that  $|D_{1,u}^{-1}(t_1,t_u)-D_{1,u}^{-1}(t_1,x_{u,k})| \leq C\ell_n$ . Therefore, if  $|X_{i,u}-t_u| \leq h$  and  $|X_{i,u}-x_{u,k}| \leq h$  occur simultaneously, then from (C.11), one of the leading terms for the bound of  $|\varphi_{ij}(t_u)-\varphi_{ij}(x_{u,k})|$  is  $\frac{C\ell_n}{h}W_{ij}$ , where

$$W_{ij} = \mathbb{I}(|X_{i,u} - x_{u,k}| \le h) \int K_{-u}\left(\frac{X_{i,1} - t_1}{h}, \frac{X_{i,\bar{u}} - X_{j,\bar{u}}}{h}\right) w_1(t_1) dt_1.$$

We have not specified the other terms since they can be dealt with analogously. If  $|X_{i,u} - t_u| \le h$  and  $|X_{i,u} - x_{u,k}| > h$  occur simultaneously, or  $|X_{i,u} - t_u| > h$  and  $|X_{i,u} - x_{u,k}| \le h$  occur simultaneously, one of the leading terms of the bound of

$$|\varphi_{ij}(t_u) - \varphi_{ij}(x_{u,k})|$$

is  $CU_{ij}$ , where

$$U_{ij} = \mathbb{I}(h - \ell_n \le |X_{i,u} - x_{u,k}| \le h + \ell_n)$$

$$\cdot \int \frac{\|Q_{j,n}^{-1}(t_1, x_{u,k})\|}{D_{1,u}(t_1, x_{u,k})} K_{-u}\left(\frac{X_{i,1} - t_1}{h}, \frac{X_{i,\bar{u}} - X_{j,\bar{u}}}{h}\right) w_1(t_1) dt_1.$$

Therefore, according to the two cases above, the following two summations, say  $I_{31}$  and  $I_{32}$ , respectively,

$$\frac{C}{n^2 h^{d+1}} \sum_{j=1}^{n} \sum_{i=1, i \neq j}^{n} \left( \frac{\ell_n}{h} W_{ij} + U_{ij} \right),$$

are one of the two leading terms of the bound of  $\sup_{t_u \in [x_{u,k}, x_{u,k+1}]} |\psi_n(t_u) - \psi_n(x_{u,k})|$ . Similar as the proof of Theorem 3.1,  $I_{31}$  can be rewritten as

$$L_{1} + \frac{(n-1)\ell_{n}}{n^{2}h^{d+2}} \sum_{j=1}^{n} \left( \mathbb{E}_{j}(W_{ij} + W_{ji}) - \mathbb{E}\left(\mathbb{E}_{j}(W_{ij} + W_{ji})\right) \right) - \frac{(n-1)\ell_{n}}{2nh^{d+2}} \mathbb{E}\left(\mathbb{E}_{j}(W_{ij} + W_{ji})\right),$$

where

$$L_1 = \frac{\ell_n}{n^2 h^{d+2}} \sum_{1 \le i \le j \le n} \left\{ (W_{ij} + W_{ji}) - \mathbb{E}_j (W_{ij} + W_{ji}) - \mathbb{E}_i (W_{ij} + W_{ji}) + \mathbb{E} \left( \mathbb{E}_i (W_{ij} + W_{ji}) \right) \right\}$$

is a degenerated U-statistic transformed from  $I_{31}$ . Next, we choose  $\ell_n = h\gamma_n^{\frac{1}{2}}$  and  $q_1 = \frac{nh}{\log n}$ . Noting that  $\mathbb{E}W_{ij}^{sk} \leq Ch^{sk+d-1}$ , by Lemma A2, we have that

$$\frac{1}{\ell_n} \cdot \frac{\ell_n^2}{\gamma_n^2} \cdot \mathbb{E} L_1^2 \leq \frac{C}{\ell_n} \cdot \frac{\ell_n^2}{\gamma_n^2} \cdot n^2 \left( \frac{\ell_n}{n^2 h^{d+2}} \right)^2 \cdot \left( h^{2s+d-1} \right)^{\frac{1}{s}} = \frac{C \ell_n^3}{\gamma_n^2 n^2 h^{d+3}} \cdot h^{-(d-1)\frac{3+\varepsilon}{r}} \leq \frac{1}{n (\log n)^2},$$

where  $\frac{1}{s} = 1 - \frac{k+1+\varepsilon}{r}$ . Since  $|\mathbb{E}_j W_{ji}| \leq Ch^{d-1}$  and  $\mathbb{E} |\mathbb{E}_j W_{ji}|^{r_1} \leq Ch^{r_1(d-1)+1}$ , by using Lemma A1, we have

$$\frac{1}{\ell_n} \mathbb{P}\left(\frac{\ell_n}{nh^{d+2}} \sum_{j=1}^n \left(\mathbb{E}_j W_{ji} - \mathbb{E}(\mathbb{E}_j W_{ji})\right) \ge B \frac{\gamma_n}{\ell_n}\right) \le \frac{1}{\ell_n} \cdot n^{-CB} + \frac{1}{\ell_n} \cdot \frac{n}{q_1} \cdot \beta\left(q_1\right) \le \frac{1}{n(\log n)^2}, \quad (C.12)$$

where we use the fact  $n\left(\mathbb{E}\left|\mathbb{E}_{j}W_{ji}\right|^{r_{1}}\right)^{\frac{2}{r_{1}}} \leq q_{1} \cdot Ch^{d-1} \cdot \frac{\gamma_{n}}{\ell_{n}} \cdot \frac{nh^{d+2}}{\ell_{n}}$  with  $\frac{2}{r_{1}} = 1 - \frac{1+\varepsilon}{r}$ . Also, it holds that  $\frac{\ell_{n}}{nh^{d+2}} \sum_{j=1}^{n} \mathbb{E}\left(\mathbb{E}_{j}(W_{ij} + W_{ji})\right) \leq \frac{C\gamma_{n}}{\ell_{n}}$ . Similarly,

$$\frac{1}{\ell_n} \mathbb{P}\left(\frac{\ell_n}{nh^{d+2}} \sum_{j=1}^n \left(\mathbb{E}_j W_{ij} - \mathbb{E}\left(\mathbb{E}_j W_{ij}\right)\right) \ge B \frac{\gamma_n}{\ell_n}\right) \le \frac{C}{n(\log n)^2}.$$

Next, we focus on  $I_{32}$ . Adopting the same method as that of  $I_{31}$ , let  $L_2$  be the corresponding degenerated U-statistic resulting from  $I_{32}$ . For k=4, since  $\mathbb{E}_{ij}^{ks} \leq C \frac{\ell_n}{h} h^{ks+d-2}$ , it follows that

$$\frac{1}{\ell_n} \cdot \frac{\ell_n^k}{\gamma_n^k} \cdot \mathbb{E}_2^k \le \frac{C}{\ell_n} \cdot \frac{\ell_n^k}{\gamma_n^k} \cdot n^k \cdot \left(\frac{1}{n^2 h^{d+1}}\right)^k \cdot \left(\frac{\ell_n}{h} h^{ks+d-2}\right)^{1-\frac{k+1+\varepsilon}{r}} \le \frac{C}{n(\log n)^2}.$$

Because  $|\mathbb{E}_i U_{ij}| \leq C h^{d-1}$  and  $\mathbb{E} |\mathbb{E}_i U_{ij}|^{r_1} \leq C h^{r_1(d-1)} \frac{\ell_n}{h}$ , it holds that

$$\frac{1}{\ell_n} \mathbb{P} \left\{ \left| \frac{1}{nh^{d+1}} \sum_{i=1}^n \left( \mathbb{E}_i U_{ij} - \mathbb{E} \left( \mathbb{E}_i U_{ij} \right) \right) \right| \ge \frac{B\gamma_n}{\ell_n} \right\} \le \frac{C}{n(\log n)^2},$$

where the fact

$$n \left( \mathbb{E} \left| \mathbb{E}_i U_{ij} \right|^{r_1} \right)^{\frac{2}{r_1}} \le C q_1 \cdot h^{d-1} \cdot \frac{B n h^{d+1} \gamma_n}{\ell_n}$$

is used. Also, it holds that  $\frac{1}{nh^{d+1}} \sum_{i=1}^n \mathbb{E}(\mathbb{E}_i(U_{ij} + U_{ji})) \leq \frac{C\gamma_n}{\ell_n}$ . Similarly,

$$\frac{1}{\ell_n} \mathbb{P}\left( \left| \frac{1}{nh^{d+1}} \sum_{i=1}^n \left( \mathbb{E}_i U_{ji} - \mathbb{E}(\mathbb{E}_i U_{ji}) \right) \right| \ge B \frac{\gamma_n}{\ell_n} \right) \le \frac{C}{n(\log n)^2}.$$

We now consider the last term on the RHS of (C.10). Here, we assume that the two notations  $U_n$  and  $\psi_i$  are the same as that appears in part i) of Theorem 3.1, with  $x_u$  replaced by  $x_{u,k}$ . Analogous to (C.2), for k = 8, it can be inferred that

$$\frac{1}{\gamma_n^k \ell_n} \cdot \mathbb{E} U_n^k \le \frac{1}{\gamma_n^k \ell_n} \cdot \frac{n^k h^{d\left(1 - \frac{k+1+\varepsilon}{r}\right)}}{(n^2 h^{d+1})^k} \le \frac{C}{n(\log n)^2}.$$

Noting that  $|\psi_i| \leq Ch^d$  and taking  $q = \sqrt{\frac{nh^{1-\frac{1+\varepsilon_0}{r}}}{\log n}}$ , it follows that

$$\frac{1}{\ell_n} \mathbb{P}\left( \left| \frac{1}{nh^{d+1}} \sum_{i=1}^n \psi_i \right| \ge \gamma_n \right) \le \frac{C}{n(\log n)^2}.$$

Finally, we consider the second term on the RHS of (C.10). As previously, let  $L_3$  be the degenerated U-statistic related to this term. Then, we have

$$\frac{1}{\ell_n} \cdot \frac{\ell_n^2}{\gamma_n^2} \cdot \mathbb{E}L_3^2 \leq \frac{C}{\ell_n} \cdot \frac{\ell_n^2}{\gamma_n^2} \cdot n^2 \left(\frac{1}{n^2 h^{d+1}}\right)^2 \cdot \left(h^d\right)^{1-\frac{2+\varepsilon}{r_3}} = \frac{C\ell_n}{\gamma_n^2} \cdot \frac{1}{n^2 h^{d+2+\frac{d(2+\varepsilon)}{r_3}}} \leq \frac{C}{n \left(\log n\right)^2}.$$

Also, since  $|\mathbb{E}_i \varphi_{ij}(x_{u,k})| \leq Ch^{d-1}$  and  $\mathbb{E} |\mathbb{E}_i \varphi_{ij}(x_{u,k})|^{r_2} \leq Ch^{r_2(d-1)+1}$ , it follows that

$$\left| \frac{1}{\ell_n} \mathbb{P}\left( \left| \frac{1}{nh^{d+1}} \sum_{i=1}^n \mathbb{E}_i \varphi_{ij}(x_{u,k}) \right| \ge \frac{B\gamma_n}{\ell_n} \right) \le \frac{C}{n \left(\log n\right)^2}.$$

Hence, (C.9) holds.

Next, we prove (3.6) for the u=1. Here, we adopt the same notations as in the proof of ii) of Theorem 3.1. From (C.8), it suffices to show that  $\widehat{c}-c=O(\gamma_n)$  and  $\sup_{x_1\in[a_1,b1]}\left|\int_{x_{1,0}}^{x_1}L_n(t_1)\mathrm{d}t_1\right|=O(\gamma_n)$ . The first one can be inferred from the proof of ii) of Theorem 3.1. Hence, we just consider the second one. Recalling that  $\frac{1}{c}I_2=I_{21}-I_{22}-I_{23}$ , we only need to prove that  $I_{21}=O(\gamma_n)$  and  $I_{22}=O(\gamma_n)$ . Now, analogous to the previous proof for the  $2\leq u\leq d$ , these two results follow immediately.  $\square$ 

#### Proof of Theorem 5.1

*Proof.* i) For convenience of notation, let  $\tau_n = \sqrt{\frac{\log n}{nh_G^{1+\frac{1+\varepsilon}{r}}}}$ ,  $\delta = \frac{1}{\sqrt{nh^{1+\frac{1+\varepsilon}{r}}}}$  and  $G_n(v)$  is the  $\alpha$ th conditional quantile of  $F_n(y|v)$ . First, we show that with probability one

$$\sup_{y \in \mathbb{R}} \left| \widehat{F}_n(y|v) - F_n(y|v) \right| = O\left(\frac{n^{-\varepsilon_0}}{\sqrt{nh_G}}\right). \tag{C.13}$$

In fact, this conclusion can be established from the following two relationships

$$\sup_{y \in \mathbb{R}} \left| \sum_{j=1}^{n} \left( K_G \left( \frac{v - \widehat{q}_0(X_j)}{h_G} \right) - K_G \left( \frac{v - q_0(X_j)}{h_G} \right) \right) \mathbb{I} \left( Y_i \le y \right) \right| = O\left( \sqrt{n^{1 - 2\varepsilon_0} h_G} \right) \tag{C.14}$$

and

$$I = \sum_{j=1}^{n} \left( K_G \left( \frac{v - \widehat{q}_0(X_j)}{h_G} \right) - K_G \left( \frac{v - q_0(X_j)}{h_G} \right) \right) = O\left( \sqrt{n^{1 - 2\varepsilon_0} h_G} \right). \tag{C.15}$$

Since the proofs of these two relationships are completely the same, we only verify the second relationship (C.15). According to Taylor's expansion, there exist  $0 \le \lambda_j \le 1, j = 1, ..., n$ , such that I can be written as

$$\sum_{j=1}^{n} K_{G}' \left( \frac{v - q_{0}(X_{j})}{h_{G}} \right) \frac{\widehat{q}_{0}(X_{j}) - q_{0}(X_{j})}{h_{G}} + \sum_{j=1}^{n} K_{G}'' \left( \frac{v - q_{0}(X_{j}) + \bar{\theta}_{j}}{h_{G}} \right) \frac{\left(\widehat{q}_{0}(X_{j}) - q_{0}(X_{j})\right)^{2}}{h_{G}^{2}},$$

where  $\bar{\theta}_j = \lambda_j (\hat{q}_0(X_j) - q_0(X_j))$ . Let  $I_1$  and  $I_2$  denote the two terms above, respectively. According to variable substitution, integration by parts, conditions (C1) and (C2), it can be shown subsequently that

$$-\mathbb{E}\left|K_{G}''\left(\frac{v-q_{0}(X_{j})}{h_{G}}\right)\right| = h_{G} \int_{-1}^{1} K_{G}''(t) f_{q_{0}}(v-th_{G}) dt = h_{G}^{2} \int_{-1}^{1} f_{q_{0}}'(v-th_{G}) K_{G}'(t) dt$$

$$= h_{G}^{3} \int_{-1}^{1} f_{q_{0}}''(v-th_{G}) K_{G}(t) dt = O(h_{G}^{3}). \tag{C.16}$$

Likewise, it follows that

$$\mathbb{E}\left|K_G'\left(\frac{v-q_0\left(X_j\right)}{h_G}\right)\right| = O(h_G^2). \tag{C.17}$$

By the standard method of the proof of Strong Law of Large Numbers (SLLN) we have

$$\sum_{j=1}^{n} \left( \left| K_G' \left( \frac{v - q_0(X_j)}{h_G} \right) \right| - \mathbb{E} \left| K_G' \left( \frac{v - q_0(X_j)}{h_G} \right) \right| \right) = O\left( \sqrt{nh_G^{1 - \frac{1 + \varepsilon}{r}}} \right). \tag{C.18}$$

Thus, from the two points above, Lemma 4.1 and condition (C3), it can be inferred that  $|I_1| = O(nh_G\delta)$ . By using conditions (C1) and (C3), for some fixed constant  $\varepsilon_0 > 0$ , we can see that

$$|I_2| \le \sum_{j=1}^n \left( \left| K_G'' \left( \frac{v - q_0(X_j)}{h_G} \right) \right| + \mathbb{I} \left( \frac{|v - q_0(X_j)|}{h_G} \le 1 + \varepsilon_0 \right) \frac{\delta}{h_G} \right) \left( \frac{\delta}{h_G} \right)^2 \equiv I_{21} + I_{22}$$
 (C.19)

holds for n sufficiently large. Analogous to the proof of relationship (C.18), by using Lemma A1, we have

$$\sum_{j=1}^{n} \left[ \mathbb{I}\left(\frac{|v - q_0(X_j)|}{h_G} \le 1 + \varepsilon_0\right) - \mathbb{E}\left(\frac{|v - q_0(X_j)|}{h_G} \le 1 + \varepsilon_0\right) \right] = O\left(\left(nh_G^{1 - \frac{1 + \varepsilon}{r}}\right)^{\frac{1}{2}}\right), \tag{C.20}$$

where we use the facts  $q = \left(nh_G^{1-\frac{1+\varepsilon}{r}}/\log n\right)^{\frac{1}{2}}$  and  $nq^{-(r+1)} \to 0$ . From this, condition (C3) and the fact

$$n\mathbb{E}\left(\frac{|v-q_0(X_j)|}{h_G} \le 1 + \varepsilon_0\right) = O(nh_G),$$

we know that  $I_{22} = o(nh_G\delta)$ . As for  $I_{21}$ , similar to the proof of (C.20), we can establish

$$\sum_{j=1}^{n} \left( \left| K_G'' \left( \frac{v - q_0(X_j)}{h_G} \right) \right| - \mathbb{E} \left| K_G'' \left( \frac{v - q_0\left(X_j\right)}{h_G} \right) \right| \right) = O\left( \sqrt{n h_G^{1 - \frac{1 + \varepsilon}{r}}} \right).$$

From this, (C.16) and condition (C3), we see that  $I_{21} = o(nh_G\delta)$ , where we use the fact  $\frac{\delta\sqrt{\log n}}{h_G^2\sqrt{nh_G^{1+\frac{1+\varepsilon}{r}}}} \to 0$ . Hence, (C.15) holds.

Next, in our of mixing processes, we adopt the standard procedure of the conditional quantiles (see, e.g., Mehra et al., 1992, Lemma 2.2), to prove that

$$\sup_{|y-G(v)| \le C\tau_n} |F_n(y|v) - F_n(G(v)|v) - F(y|v) + (1-\alpha)| = O\left(\tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}\right). \tag{C.21}$$

Divide the interval  $|y - G(v)| \leq C\tau_n$  into a sequence of subintervals with length  $\tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}$ . Let  $\eta_r$ ,  $r = 1, 2, \ldots, d_n$ , be the corresponding grid points. Then,  $d_n = C\tau_n^{-\frac{1}{2} + \frac{1+\varepsilon}{2r}}$ . Then, it holds that

$$\sup_{|y-G(v)| \le C\tau_n} |F_n(y|v) - F_n(G(v)|v) - F(y|v) + (1-\alpha)|$$

$$\le \max_{|r| \le Cd_n} |F_n(\eta_r|v) - F(\eta_r|v) - F_n(G(v)|v) + (1-\alpha)| + \max_{|r| \le Cd_n} |F(\eta_{r+1}|v) - F(\eta_r|v)|. \tag{C.22}$$

From condition (C4), it is easy to see that

$$\max_{|r| \le Cd_n} |F(\eta_{r+1}|v) - F(\eta_r|v)| = O\left(\tau n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}\right).$$
 (C.23)

Let  $\xi_{i,r,1} = K\left(\frac{v - q_0(X_i)}{h_G}\right) (\mathbb{I}(Y_i \leq \eta_r) - \mathbb{I}(Y_i \leq G(v)))$  and  $b_n = Bnh_G \tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}$ . By using Lemma A1, we have

$$d_{n}\mathbb{P}\left\{\left|\sum_{i=1}^{n}\left(\xi_{i,r,1} - \mathbb{E}\xi_{i,r,1}\right)\right| \ge b_{n}\right\} \le d_{n}\left(n^{-CB} + \frac{n}{q}\beta\left(q\right)\right) \le \frac{C}{n\left(\log n\right)^{2}},\tag{C.24}$$

where q is equal to  $\frac{b_n}{\log n}$ . In view of condition (C4), it follows that

$$n\mathbb{E}\xi_{i,r,1} = nh_G(\eta_r - G(v))f_{q_0}(v)(1 + O(h_G))$$

and

$$nh_G(F(\eta_r|v) - (1-\alpha)) = nh_G(\eta_r - G(v))f_{q_0}(v)(1 + O(h_G)).$$

We then have

$$n\mathbb{E}\xi_{i,r,1} - nh_G(F(\eta_r|v) - (1-\alpha)) = O(nh_G^2\tau_n).$$
 (C.25)

Also, it can be shown that  $\sum_{i=1}^{n} K_G\left(\frac{v-q_0(X_i)}{h_G}\right) = nh_G f_{q_0}(v) \left(1 + O(h_G^2)\right)$ . Hence, from this, (C.24), (C.25) and the fact  $b_n \geq nh_G^2 \tau_n$ , it can be inferred that

$$d_n \mathbb{P}\left\{ |F_n(\eta_r|v) - F_n(G(v)|v) - F(\eta_r|v) + (1 - \alpha)| \ge B\tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}} \right\} \le \frac{C}{n(\log n)^2}.$$
 (C.26)

Then (C.21) is implied by (C.22), (C.23) and (C.26). From (C.13) and (C.21), we have

$$\sup_{|y-G(v)| \le 2c\tau_n} \left| \widehat{F}_n(y|v) - F_n(G(v)|v) - F(y|v) + (1-\alpha) \right| = O\left(\tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}} + \delta\right). \tag{C.27}$$

Next, we prove (5.2). In order to do so, we first show that, with probability one,

$$|G_n(v) - G(v)| = O(\tau_n), \tag{C.28}$$

which is similar to Lemma 2.1 of Mehra et al. (1992). For any constant M > 0, let  $u_n^+ = G(v) + M\tau_n$  and  $u_n^- = G(v) - M\tau_n$  and then consider the following probability

$$\mathbb{P}\left\{\left|G_{n}(v)-G(v)\right|\geq M\tau_{n}\right\}=\mathbb{P}\left\{F_{n}\left(\left.u_{n}^{-}\right|v\right)\leq1-\alpha\right\}+\mathbb{P}\left\{F_{n}\left(\left.u_{n}^{+}\right|v\right)\geq1-\alpha\right\}.$$

Let  $\zeta_i = K\left(\frac{v - q_0(X_i)}{h_G}\right) \mathbb{I}(Y_i \leq u_n^+)$ . From Lemma A1, we have

$$\mathbb{P}\left\{ \left| \sum_{i=1}^{n} \left( \zeta_{i} - \mathbb{E}\zeta_{i} \right) \right| \geq M\tau_{n} \right\} \leq n^{-CM} + \frac{n}{q_{1}} \beta\left( q_{1} \right) \leq \frac{C}{n \left( \log n \right)^{2}},$$

where  $q_1$  is taken as  $\frac{1}{\tau_n h^{\frac{1+\varepsilon}{r}}}$ . From this and noting that  $n\left(\mathbb{E}\zeta_i - (1-\alpha)\mathbb{E}K\left(\frac{v-q_0(X_i)}{h_G}\right)\right) = O\left(nh_G\tau_n\right)$ , it can be inferred that  $\mathbb{P}\left\{F_n\left(u_n^+|v\right) \geq 1-\alpha\right\} \leq \frac{1}{n(\log n)^2}$ . Analogously, it can be shown that  $\mathbb{P}\left\{F_n\left(u_n^-|v\right) \leq 1-\alpha\right\} \leq \frac{1}{n(\log n)^2}$ . And thus, we know that (C.28) holds. Then there exists some constant C > 0 such that

$$1 - \alpha \in \left[ F_n \left( G(v) - C\tau_n | v \right), F_n \left( G(v) + C\tau_n | v \right) \right]$$
(C.29)

holds with probability one. Let

$$\xi_{i,r,2} = \xi_{i,r,2}(v) = K_G\left(\frac{v - q_0(X_i)}{h_G}\right) \mathbb{I}(G(v) + C\tau_n < Y_i \le G(v) + 2C\tau_n).$$

Then, from Lemma A1, it follows that

$$\mathbb{P}\left\{ \left| \sum_{i=1}^{n} \left( \xi_{i,r,2} - \mathbb{E}\xi_{i,r,2} \right) \right| \ge Bnh_{G}\tau_{n}^{\frac{3}{2} - \frac{1+\varepsilon}{2r}} \right\} \le n^{-CB} + \frac{n}{q_{2}}\beta(q_{2}) \le \frac{C}{n\left(\log n\right)^{2}}, \tag{C.30}$$

where  $q_2 = \frac{nh_G \tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}}{\log n}$ . From this and  $\sum_{i=1}^n K_G\left(\frac{v - q_0(X_i)}{h_G}\right) = nh_G f_{q_0}(v) + O(h_G^2)$ , it can be inferred that

$$F_n(G(v) + 2\tau_n|v) - F_n(G(v) + \tau_n|v) = c_{n1}(v)\tau_n + O\left(\tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}\right),$$
 (C.31)

where  $\liminf c_{n1}(v) > 0$  as  $n \to \infty$ . Similarly, it follows that

$$F_n(G(v) - 2\tau_n|v) - F_n(G(v) - \tau_n|v) = -c_{n2}(v)\tau_n + O\left(\tau_n^{\frac{3}{2} - \frac{1+\varepsilon}{2r}}\right),$$
(C.32)

where  $\liminf c_{n2}(v) > 0$ . Therefore, from  $\tau_n > \delta$  and monotonicity on the empirical conditional function, it can be inferred subsequently that

$$\widehat{F}_n(G(v) - 2C\tau_n|v) \le F_n(G(v) - 2C\tau_n|v) + \delta \le F_n(G(v) - C\tau_n|v)$$

$$\le 1 - \alpha \le F_n(G(v) + C\tau_n|v) \le F_n(G(v) + 2C\tau_n|v) - \delta \le \widehat{F}_n(G(v) + 2C\tau_n|v).$$

And this implies (5.2). In view of (5.2) and (C.27), it can be shown that (5.3) holds.

ii) To obtain the uniform convergence of (5.3), according to the previous proof, it suffices to show that (C.13), (C.21), (C.28), (C.31) and (C.32) hold uniformly for  $v \in V$ . We deal with each relationship separately. To prove the results on the uniform convergence for  $v \in V$ , it is necessary to divide V into smaller intervals with equal length  $d_n > 0$ . Assume that the corresponding grid points are  $v_l$ ,  $l = 1, 2, \ldots, O(d_n^{-1})$ , assuming l is an integer. Denote the lth smaller interval  $[v_l, v_{l+1}]$  by  $J_l$ . Note that  $d_n$  may take different values at different s, for simplicity of notation.

As for uniform convergence of (C.13), it can be inferred from the uniform convergence of (C.14) and (C.15). Here, since the proofs of the two relationships are the same, it suffices to prove that (C.15) holds uniformly for  $v \in \mathcal{V}$ . Then, going along the same lines as the proof of (C.15), we need to prove that  $\sup_{v \in \mathcal{V}} |I_1| = O(nh_G\delta_n)$ ,  $\sup_{v \in \mathcal{V}} |I_{21}| = o(nh_G\delta)$  and  $\sup_{v \in \mathcal{V}} |I_{22}| = o(nh_G\delta)$ . Since the proofs of these three relationships are analogous, we here only prove the first one. Divide  $\mathcal{V}$  as being stated, and let  $d_n = h_G^2$ . Then, for  $v \in J_l$ , it follows that

$$\left| K_G' \left( \frac{v - q_0\left(X_j\right)}{h_G} \right) - K_G' \left( \frac{v_l - q_0\left(X_j\right)}{h_G} \right) \right| \le \frac{Cd_n}{h_G} \mathbb{I}\left( |v_l - q_0\left(X_j\right)| \le h_G \right) + C\mathbb{I}\left( h_G - d_n \le |v_l - q_0\left(X_j\right)| \le h_G + d_n \right) \equiv \frac{Cd_n}{h_G} \xi_{i1} + C\xi_{i2}.$$

Thus, our objective is to show that

$$\mathbb{P}\left(\bigcup_{l} \left(\frac{d_n}{h_G} \sum_{j=1}^{n} \xi_{i1} \ge Cnh_G^2\right)\right) \le \frac{1}{n \left(\log n\right)^2} \text{ and } \mathbb{P}\left(\bigcup_{l} \left(\sum_{j=1}^{n} \xi_{i2} \ge Cnh_G^2\right)\right) \le \frac{1}{n \left(\log n\right)^2}. \quad (C.33)$$

We only prove the first one, because the proof of the other one is similar. Noting that  $f_{q^0}(\cdot)$  is bounded on  $\mathcal{V}$ , we know that  $n\mathbb{E}\xi_{i1} = O(nh_G)$ . Then, from Lemma A1, it can be derived that

$$\mathbb{P}\left(\frac{d_n}{h_G}\sum_{j=1}^n \xi_{i1} \ge CBnh_G^2\right) \le \mathbb{P}\left(\sum_{j=1}^n (\xi_{i1} - \mathbb{E}\xi_{i1}) \ge CBnh_G\right) \\
\le 2\exp\left\{\frac{-\left(\frac{Cnh_G}{4}\right)^2}{cn\left(\mathbb{E}\xi_{i1}^{r_2}\right)^{\frac{2}{r_2}} + \frac{2}{3}qM\frac{Cnh_G}{4}}\right\} + \frac{n}{q}\beta(q) \le n^{-CB} + \frac{n}{q}\beta(q) \le \frac{d_n}{n\left(\log n\right)^2},$$

where  $1 - \frac{2}{r_2} = \frac{1+\varepsilon_0}{r}$  and  $q = \frac{nh_G^{1+\frac{1+\varepsilon_0}{r}}}{\log n}$ . This implies that the first limit of (C.33) holds. In view of (C.18) and Lemma A1,

$$\left| \mathbb{P}\left( \bigcup_{l} \left( \left| \sum_{j=1}^{n} K_{G}' \left( \frac{v - q_{0}\left(X_{j}\right)}{h_{G}} \right) - \mathbb{E}K_{G}' \left( \frac{v - q_{0}\left(X_{j}\right)}{h_{G}} \right) \right| \geq CBnh_{G}^{2} \right) \right) \\
\leq 2 \sum_{l} \left( \exp\left\{ \frac{-\left(\frac{Cnh_{G}^{2}}{4}\right)^{2}}{cn\left(\mathbb{E}\xi_{i1}^{r_{2}}\right)^{\frac{2}{r_{2}}} + \frac{2}{3}q_{1}M\frac{Cnh_{G}^{2}}{4}} \right\} + \frac{n}{q_{1}}\beta\left(q_{1}\right) \right)$$

$$\leq d_n^{-1} \left( n^{-CB} + \frac{n}{q_1} \beta \left( q_1 \right) \right) \leq \frac{C}{n \left( \log n \right)^2},$$

where  $q_1 = \frac{nh_G^2}{\log n}$ . Therefore, from this, (C.33), Theorem 4.1 and the Borel-Cantelli lemma, we know that  $\sup_{n \in \mathcal{V}} |I_1| = O(nh_G\delta)$ .

We now begin to prove the uniform convergence of (C.21). For the division of this, we take  $d_n = \delta h_G$ . First, note that

$$I_{3} = \sup_{v \in J_{l}} \sup_{|y - G(v)| \le C\tau_{n}} |F_{n}(y|v) - F_{n}(G(v)|v) - F(y|v) + (1 - \alpha)| \le I_{31} + I_{32},$$

where

$$I_{31} = \sup_{v \in J_l} \sup_{|y - G(v)| \le C\tau_n} |F_n(y|v) - F_n(G(v)|v) - F(y|v) - (F_n(y|v_l) - F_n(G(v_l)|v_l) - F(y|v_l))|$$

and

$$I_{32} = \sup_{v \in J_l} \sup_{|y - G(v)| \le C\tau_n} |F_n(y|v_l) - F_n(G(v_l)|v_l) - F(y|v_l) + (1 - \alpha)|.$$

Since  $|G(v) - G(v_l)| \le C\delta$  for any  $v \in J_l$ , it then can be seen that

$$I_{32} \le \sup_{|y - G(v_l)| \le 2C\tau_n} |F_n(y|v_l) - F_n(G(v_l)|v_l) - F(y|v_l) + (1 - \alpha)|. \tag{C.34}$$

Let  $I_{311} = \sup_{v \in J_l} |F_n(G(v)|v) - F_n(G(v_l)|v_l)|$ ,  $I_{312} = \sup_{v \in J_l} \sup_y |F_n(y|v) - F_n(y|v_l)|$  and  $I_{313} = \sup_{v \in J_l} \sup_y |F(y|v) - F(y|v_l)|$ . Then,  $I_{31} \le I_{311} + I_{312} + I_{313}$ . Set

$$I_{3111} = \sup_{v \in J_l} |F_n(G(v_l \pm C\delta)|v) - F_n(G(v_l)|v)|$$

and  $I_{3112} = \sup_{v \in J_l} |F_n(G(v_l)|v) - F_n(G(v_l)|v_l)|$ . Then, we know that  $I_{311} \leq I_{3111} + I_{3112}$ . Note that

$$I_{3111} \leq 2I_{312} + \left| F_n \left( G \left( v_l \pm C \delta \right) | v_l \right) - F_n \left( G \left( v_l \right) | v_l \right) \right|$$

$$\leq 2I_{312} + I_{32} + \left| F \left( G \left( v_l \pm C \delta \right) | v_l \right) - F \left( G \left( v_l \right) | v_l \right) \right|.$$

Then, from the facts that  $I_{3112} \leq I_{312}$ ,  $\left| F\left(G\left(v_l \pm C\delta\right) | v_l\right) - F\left(G\left(v_l\right) | v_l\right) \right| \leq C\delta$  and  $I_{313} \leq C\delta$ , which is implied by condition (C6), it suffices to consider  $I_{312}$  and  $I_{32}$ . To get the result  $\mathbb{P}\left\{\left| \bigcup_l \left\{I_{312} \geq C\delta\right\} \right| \right\} \leq \frac{C}{n(\log n)^2}$ , it suffices to prove that

$$\mathbb{P}\left\{ \bigcup_{l} \left\{ \sup_{v \in J_{l}} \sup_{y} \left| \sum_{j=1}^{n} \left( K_{G}\left( \frac{v - q_{0}\left(X_{j}\right)}{h_{G}} \right) - K_{G}\left( \frac{v_{l} - q_{0}\left(X_{j}\right)}{h_{G}} \right) \right) \mathbb{I}\left(Y_{i} \leq y\right) \right| \geq Cnh_{G}\delta \right\} \right\}$$

and

$$\mathbb{P}\left\{ \bigcup_{l} \left\{ \sup_{v \in J_{l}} \left| \sum_{j=1}^{n} \left( K_{G} \left( \frac{v - q_{0}(X_{j})}{h_{G}} \right) - K_{G} \left( \frac{v_{l} - q_{0}(X_{j})}{h_{G}} \right) \right) \right| \geq Cnh_{G}\delta \right\} \right\}$$
 (C.35)

are less than  $\frac{C}{n(\log n)^2}$ . In considering that the proofs of two relationships above are the same, we only prove the later one. Since

$$\left| \sum_{j=1}^{n} \left( K_{G} \left( \frac{v - q_{0} \left( X_{j} \right)}{h_{G}} \right) - K_{G} \left( \frac{v_{l} - q_{0} \left( X_{j} \right)}{h_{G}} \right) \right) \right|$$

$$\leq C \frac{d_{n}}{h_{G}} \mathbb{I} \left( \left| v_{l} - q_{0} \left( X_{j} \right) \right| \leq h_{G} \right) + C \mathbb{I} \left( h_{G} - d_{n} \leq \left| v_{l} - q_{0} \left( X_{j} \right) \right| \leq h_{G} + d_{n} \right),$$

by using Lemma A1, we obtain

$$\mathbb{P}\left\{ \cup_{l} \left\{ \left| \sum_{j=1}^{n} \frac{d_{n}}{h_{G}} \mathbb{I}\left(\left| v_{l} - q_{0}\left(X_{j}\right)\right| \leq h_{G}\right) \right| \geq C n h_{G} \delta \right\} \right\} \leq \frac{C}{n \left(\log n\right)^{2}}$$

and

$$\mathbb{P}\left\{ \cup_{l} \left\{ \left| \sum_{j=1}^{n} \mathbb{I}\left(h_{G} - d_{n} \leq |v_{l} - q_{0}\left(X_{j}\right)| \leq h_{G} + d_{n}\right) \right| \geq Cnh_{G}\delta \right\} \right\} \leq \frac{C}{n\left(\log n\right)^{2}}.$$

In view of (C.34), by adopting the same proof as that of (C.21), we arrive at

$$\mathbb{P}\big\{\cup_{l} \{I_{32} \ge C\delta\}\big\} \le \frac{C}{n\left(\log n\right)^{2}}.$$

Thus, we know that (C.21) holds uniformly for  $v \in \mathcal{V}$  with the convergence rate  $\delta$ .

Next, we verify that (C.28) holds uniformly for  $v \in \mathcal{V}$ . Let  $d_n = \tau_n h_G a_n$ , where the positive reals  $a_n \to 0$  at a rather slow rate. For  $v \in J_l$ , since  $|G(v) - G(v_l)| \leq C\tau_n$ , it suffices to prove that

$$\sum_{l} \mathbb{P}\left\{ \sup_{v \in J_{l}} |G_{n}(v) - G_{n}(v_{l})| \ge C\tau_{n} \right\} \le \frac{C}{n \left(\log n\right)^{2}} \tag{C.36}$$

and

and

$$\mathbb{P}\left\{ \bigcup_{l} (|G_n(v_l) - G(v_l)| \ge C\tau_n) \right\} \le \frac{C}{n \left(\log n\right)^2}.$$

The second relationship above can be obtained similarly as (C.28). Analogous to the proof of  $I_{312}$ , it follows that

$$\sum_{l} \mathbb{P} \left\{ \sup_{v \in J_{l}} \sup_{y} |F_{n}(y|v) - F_{n}(y|v_{l})| \ge ca_{n}\tau_{n} \right\} \le \frac{C}{n \left(\log n\right)^{2}}.$$

Then, for any event  $w \in \{\sup_{v \in J_l} \sup_y |F_n(y|v) - F_n(y|v_l)| < ca_n \tau_n \}$  and  $v \in J_l$ , using relationship (C.31),

$$F_n(G(v_l) - 2C\tau_n|v) \le F_n(G(v_l) - 2C\tau_n|v_l) + Ca_n\tau_n \le F_n(G(v_l) - C\tau_n|v_l)$$

$$\le 1 - \alpha \le F_n(G(v_l) + C\tau_n|v_l) \le F_n(G(v_l) + 2C\tau_n|v_l) - Ca_n\tau_n \le F_n(G(v_l) + 2C\tau_n|v).$$

And this together with (C.36) implies (C.28).

Finally, we show that (C.31) and (C.32) hold uniformly for  $v \in \mathcal{V}$ . Let  $d_n = a_n \tau_n$ . First, note that

$$\begin{split} & \left| \sum_{i=1}^{n} \left( \xi_{i,r,2} \left( v \right) - \xi_{i,r,2} \left( v_{l} \right) \right) \right| \\ \leq & \left| \sum_{i=1}^{n} \left( K_{G} \left( \frac{v_{l} - q_{0}(X_{i})}{h_{G}} \right) + \left| K_{G} \left( \frac{v - q_{0}(X_{i})}{h_{G}} \right) - K_{G} \left( \frac{v_{l} - q_{0}(X_{i})}{h_{G}} \right) \right| \right) \\ & \cdot \left| \mathbb{I} \left( G(v) + C\tau_{n} < Y_{i} \leq G(v) + 2C\tau_{n} \right) - \mathbb{I} \left( G(v_{l}) + C\tau_{n} < Y_{i} \leq G(v_{l}) + 2C\tau_{n} \right) \right| \\ & + \sum_{i=1}^{n} \left| K_{G} \left( \frac{v - q_{0}(X_{i})}{h_{G}} \right) - K_{G} \left( \frac{v_{l} - q_{0}(X_{i})}{h_{G}} \right) \right| \mathbb{I} \left( G(v_{l}) + C\tau_{n} < Y_{i} \leq G(v_{l}) + 2C\tau_{n} \right). \end{split}$$

Also, since  $|G(v) - G(v_l)| \le ca_n \tau_n$  for  $v \in J_l$ , it holds that

$$\left| K_{G} \left( \frac{v - q_{0}(X_{i})}{h_{G}} \right) - K_{G} \left( \frac{v_{l} - q_{0}(X_{i})}{h_{G}} \right) \right|$$

$$\leq C\tau_{n} \mathbb{I} \left( |v_{l} - q_{0}(X_{i})| \leq h_{G} \right) + \mathbb{I} \left( h_{G} \left( 1 - C\tau_{n} \right) < v_{l} - q_{0}(X_{i}) \leq h_{G} \left( 1 + C\tau_{n} \right) \right)$$

$$\left| \mathbb{I} \left( G(v) + C\tau_{n} < Y_{i} \leq G(v) + 2C\tau_{n} \right) - \mathbb{I} \left( G(v_{l}) + C\tau_{n} < Y_{i} \leq G(v_{l}) + 2C\tau_{n} \right) \right|$$

$$\leq \mathbb{I} \left( G(v_{l}) + C\tau_{n} - Ch_{G}\tau_{n} < Y_{i} \leq G(v_{l}) + C\tau_{n} + Ch_{G}\tau_{n} \right)$$

$$+ \mathbb{I} \left( G(v_{l}) + 2C\tau_{n} - Ch_{G}\tau_{n} < Y_{i} \leq G(v_{l}) + 2C\tau_{n} + Ch_{G}\tau_{n} \right).$$

From the three relationships above, we see that  $\left|\sum_{i=1}^{n} (\xi_{i,r,2}(v) - \xi_{i,r,2}(v_l))\right|$  has a bound which is independent of v. Let

$$\zeta_{i1} = K_G \left( \frac{v_l - q_0(X_i)}{h_G} \right) \mathbb{I} \left( G(v_l) + c\tau_n - ca_n \tau_n < Y_i \le G(v_l) + c\tau_n + ca_n \tau_n \right)$$

and

$$\zeta_{i2} = c \frac{a_n \tau_n}{h_G} \mathbb{I} (|v_l - q_0(X_i)| \le h_G) \mathbb{I} (G(v_l) + c\tau_n < Y_i \le G(v_l) + 2c\tau_n).$$

Next, we focus on the convergence rate of this bound. Here, we only consider the two terms

$$\sum_{i=1}^{n} \zeta_{i1} \quad \text{and} \quad \sum_{i=1}^{n} \zeta_{i2}.$$

The other terms can be dealt with analogously. Noting that

$$n\mathbb{E}\zeta_{i1} = Cf_{q_0}(v_l)nh_G a_n \tau_n(1+o(1))$$
 and  $n\mathbb{E}\zeta_{i1} = Cna_n \tau_n^2 f_{q_0}(v_l)(1+o(1))$ ,

and then by using Lemma A1, we have

$$\mathbb{P}\left(\bigcup_{l} \left(\left|\sum_{i=1}^{n} \zeta_{i1}\right| \ge cnh_{G}a_{n}\tau_{n}\right)\right) \\
\le 2d_{n}^{-1} \exp\left\{\frac{-\left(\frac{Cnh_{G}a_{n}\tau_{n}}{4}\right)^{2}}{Cn\left(\mathbb{E}\zeta_{i1}^{r_{2}}\right)^{\frac{2}{r_{2}}} + \frac{2}{3}q\frac{Cnh_{G}a_{n}\tau_{n}}{4}}\right\} + d_{n}^{-1}\frac{n}{q}\beta(q) \le d_{n}^{-1}n^{-CB} + d_{n}^{-1}\frac{n}{q}\beta(q)$$

and, similarly,

$$\mathbb{P}\left(\bigcup_{l}\left(\left|\sum_{i=1}^{n}\zeta_{i2}\right| \geq Cnh_{G}a_{n}\tau_{n}\right)\right) \leq d_{n}^{-1}n^{-CB} + d_{n}^{-1}\frac{n}{q}\beta(q),$$

where  $q = \frac{a_n n h_G \tau_n}{\log n}$  and  $\frac{2}{r_2} = 1 - \frac{1+\varepsilon}{r}$ . By a similar method as that of (C.30), it follows that

$$\mathbb{P}\left\{ \bigcup_{l} \left( \left| \sum_{i=1}^{n} \left( \xi_{i,r,2}(v_l) - \mathbb{E}\xi_{i,r,2}(v_l) \right) \right| \ge Bnh_G a_n \tau_n \right) \right\} \le \frac{C}{n \left( \log n \right)^2}.$$

Therefore, from the analysis above, we have

$$\sup_{v \in \mathcal{V}} \left| \sum_{i=1}^{n} \left( \xi_{i,r,2}(v) - \mathbb{E}\xi_{i,r,2}(v) \right) \right| = O(nh_G a_n \tau_n).$$

Similar as the previous proof of (C.35), it follows that

$$\sup_{v \in \mathcal{V}} \left| \sum_{i=1}^{n} \left( K_G \left( \frac{v - q_0(X_i)}{h_G} \right) - \mathbb{E} K_G \left( \frac{v - q_0(X_i)}{h_G} \right) \right) \right| = O(nh_G a_n).$$

Then, from the two relationships above and the condition  $\inf_v f_{q_0(X_i)}(v) > 0$ , it follows that

$$\sup_{v \in \mathcal{V}} \left| \frac{\sum_{i=1}^{n} \xi_{i,r,2}(v)}{\sum_{i=1}^{n} K_G\left(\frac{v - q_0(X_i)}{h_G}\right)} - \frac{\mathbb{E}\xi_{i,r,2}(v)}{\mathbb{E}K_G\left(\frac{v - q_0(X_i)}{h_G}\right)} \right| = O(a_n \tau_n).$$

Since  $\mathbb{E}\xi_{i,r,2}(v) = cf_{q_0}(v)h_G\tau_n$ , we obtain

$$\sup_{v \in \mathcal{V}} |F_n(G(v) + 2C\tau_n|v) - F_n(G(v) + C\tau_n|v) - C(1 + o(1))\tau_n| = O(a_n\tau_n).$$

Similarly, we have

$$\sup_{v \in \mathcal{V}} |F_n(G(v) - 2C\tau_n|v) - F_n(G(v) - C\tau_n|v) + C(1 + o(1))\tau_n| = O(a_n\tau_n).$$

This completes the proof of part ii).

#### D Proofs of some Lemmas and a Corollary

#### Proof of Lemma 3.1

*Proof.* Denote  $W_n(t)$  by  $W_{j,n}(t_{1,u})$  with  $\widetilde{X}_j$  replaced by t. By the mean value of integration and by Taylor expansion, there exist  $0 < \theta_1 = \theta_1(X_i, t), \theta_2 = \theta_2(X_i, t) < 1$  such that

$$\mathbb{E}\left[\left.\mathbb{I}_{(\varepsilon_i \le 0)} - \mathbb{I}_{(\varepsilon_i \le r_{i,t})}\right| X_i\right] = \int_{r_{i,t}}^0 g(X_i, s) \mathrm{d}s = r_{i,t}g(X_i, \theta_2 r_{i,t}) = r_{i,t}g(X_i, 0) + O(r_{i,t}^2)$$

and

$$r_{i,t} = \frac{(X_i - t)^p}{p!} q^{(p)}(t) + \frac{(X_i - t)^{p+1}}{(p+1)!} q^{(p+1)} (t + \theta_1(X_i - t)).$$

From the two relationships above, variable substitution and the continuity of both the density function  $p(\cdot)$  and the (p+1)th derivative of  $q(\cdot)$ , it can be inferred that

$$\frac{1}{nh^{d+1}} \mathbb{E}W_n(t)h^{p-1} \int K(s)A(s)s^p ds \cdot \frac{q^{(p)}(t)g_1(t)}{p!} + h^p \left[ \int K(s)A(s)s^{p+1} ds \cdot \frac{q^{(p+1)}(t)g_1(t)}{(p+1)!} + \int K(s)A(s)s^p ds \cdot \frac{q^{(p)}(t)g_1'(t)}{p!} \right] + o(h^p).$$

According to this and the facts that (C.5),  $e_u^T Q^{-1} \int A(s) K(s) s^l ds = 0$  with l = p or p + 1, it can be inferred that

$$\frac{1}{nh^{d+1}}Q_{jn,t_{1,u}}^{-1}\mathbb{E}_{j}W_{j,n}(t_{1,u}) = h^{p}B_{2}(\widetilde{X}_{j}) + o(h^{p}).$$

By the uniformly SLLN for kernel regression function (see, e.g., Masry, 1996), with probability one, it holds uniformly for  $t \in \mathcal{X}$  that

$$\frac{1}{nh^{d+1}} (W_n(t) - \mathbb{E}W_n(t)) = O\left(\frac{1}{h} \left(\frac{\log n}{nh^d}\right)^{1/2}\right).$$

According to the SLLN, we have that

$$\frac{1}{n} \sum_{i=1}^{n} \left[ \iint B_2(\widetilde{X}_j) dt_1 dt_2 - \mathbb{E} \iint B_2(\widetilde{X}_j) dt_1 dt_2 \right] = O\left(\sqrt{\frac{\log n}{n}}\right).$$

Thus, from the three relationships above, it can be inferred that Lemma 3.1 holds.

### Proof of Lemma 3.2

*Proof.* Let  $\gamma_n = \frac{\sqrt{\log n}}{h^{2+\frac{1+\varepsilon}{r}}\sqrt{n}}$  and  $\ell_n = \varepsilon_n h^2$ . We only verify  $\Delta_{1,u}$  since the other cases can be dealt with analogously. As the usual way of getting the weak law of large number, it follows that

$$\frac{1}{n} \sum_{j=1}^{n} \left( \partial_1 q(\widetilde{X}_j) \right) \mathbb{E} \partial_1 q(\widetilde{X}_j)) = O\left( \left( \frac{\log n}{n} \right)^{\frac{1}{2}} \right)$$

holds uniformly for  $(t_1, t_u) \in \mathcal{X}_{1,u}$ . Then, from this, (3.1) and Theorem B1, we see that, with probability one,

$$\Delta_{1,u} = \frac{1}{n^2 h^{d+1}} \sum_{1 \le i \ne j \le n} \psi_{1,ij} + o(\gamma_n)$$

holds uniformly for  $t_{1,u} \in \mathcal{X}_{1,u}$ , where  $\psi_{1,ij} = e_1^T Q_{jn,t_{1,u}}^{-1} K_{ij,t_{1,u}} A_{ij,t_{1,u}} \left( \alpha \mathbb{I}(\varepsilon_i \leq 0) \right)$ . Let  $\xi_i = \mathbb{E}_i \psi_{1,ij}$ . Then,

$$\frac{1}{n^2h^{d+1}} \sum_{1 \le i \ne j \le n} \psi_{1,ij} = \frac{1}{n^2h^{d+1}} \sum_{1 \le i \le j \le n} (\psi_{1,ij} + \psi_{1,ji} - \xi_i - \xi_j) + \frac{(n-1)}{n^2h^{d+1}} \sum_{i=1}^n \xi_i.$$

Let  $U_n$  be the first term on the RHS of the relationship above. We now use Lemma A2 to get the convergence rate of  $U_n$ . And let  $M_{p_1k}$  be the corresponding quantity stated in Lemma A2 with  $k < r\left(1-\frac{1}{p_1}\right)-1$ . In view of condition (B7), it can be inferred that  $M_{p_1k} \leq Ch^{d/p_1}$ . Applying Lemma A2, in view of condition (B6), it follows that

$$\begin{split} \frac{\mathbb{E}(U_n^k)}{\gamma_n^k \ell_n^2} & \leq \frac{C h^{d/p_1}}{\left(\frac{\log n}{nh^{4+\frac{2+2\varepsilon}{r}}}\right)^{\frac{k+2}{2}} \left(nh^{d+1}\right)^k} \\ & \leq \frac{C}{\left(\log n\right)^{\frac{k+2}{2}} n^{\frac{k-2}{2}} h^{k(d+1)-2(k+2)-\frac{d}{p_1}-(k+2)\frac{1+\varepsilon}{r}}} \leq \frac{1}{n(\log n)^2} \end{split}$$

if k=10 and r>11. Then, it can be inferred that  $|\xi_i| \leq Ch^{d-2} \equiv M$  and  $(\mathbb{E}\,|\xi_i^{r_3}|)^{\frac{2}{r_3}} \leq Ch^{2(d-2)+\frac{4}{r_3}}$ , where  $1-\frac{2}{r_3}=\frac{1+\varepsilon}{r}$ . By Lemma A1 and taking  $q=\frac{h^{1+\frac{1}{r}+\varepsilon}\sqrt{n}}{\sqrt{\log n}}$ , we have

$$\mathbb{P}\left\{\left|\sum_{1\leq i\leq n}\xi_i\right|\geq nh^{d+1}\gamma_n\right\}\leq \exp\left\{\frac{-\left(\varepsilon_nnh^{d+1}\right)^2}{\frac{n}{2n}q\left(\mathbb{E}\left|\xi_i\right|^{r_3}\right)^{\frac{2}{r_3}}+qM\gamma_nnh^{d+1}}\right\}+\frac{n}{q}\beta(q)\leq n^{-CB}+\frac{n}{q}\beta(q).$$

For any  $s \in J_l$ , from condition (B5) and

$$Q_{jn,t_{1,u}} = \int K(x)A(x)A^{\mathrm{T}}(x)g_1(z-hx,0)\mathrm{d}x\bigg|_{z=\widetilde{X}_j},$$

we have  $\|Q_{jn,s} - Q_{jn,t_{1,u}}\| \le C\ell_n$ . Since  $Q_{jn,t_{1,u}} \to Q(t_{1,u})$  and  $Q(t_{1,u})$  is a positive definite matrix and is continuous on the compact set, it can be inferred that  $\|Q_{jn,t_{1,u}}^{-1}\|$  is bounded by a constant. Accordingly, we have

$$\left\| Q_{jn,s}^{-1} - Q_{jn,t_{1,u}}^{-1} \right\| \le \left\| Q_{jn,s}^{-1} \right\| \left\| Q_{jn,s} - Q_{jn,t_{1,u}} \right\| \left\| Q_{jn,t_{1,u}}^{-1} \right\| \le C\ell_n. \tag{D.1}$$

Similar as the previous lemma, it follows that

$$|L_1(s) - L_1(t_{1,u})| \le \frac{C}{n^2 h^{d+1}} \sum_{1 \le i \ne j \le n} \left\{ \mathbb{I}\left(\left|X_i - \widetilde{X}_j\right| \le h\right) \frac{\ell_n}{h} + \mathbb{I}\left(h - \ell_n \le X_i - \widetilde{X}_j \le h + \ell_n\right) \right\}.$$

Let  $L_2$  and  $L_3$  be the two terms on the RHS of the relationship above. We now consider the convergence rate of  $L_2$ . Let  $\psi_{ij} = \mathbb{I}(|X_i - \widetilde{X}_j| \le h) + \mathbb{I}(|X_j - \widetilde{X}_i| \le h)$ ,  $\psi_i = \mathbb{E}_i \psi_{ij}$  and  $\varphi_{ij} = \psi_{ij} - \psi_i - \psi_j + \mathbb{E}\psi_1$ . Then,

$$L_2 = \frac{2\ell_n}{n^2h^{d+2}} \sum_{1 \le i \le j \le n} \varphi_{ij} + \frac{2(n-1)\ell_n}{n^2h^{d+2}} \sum_{i=1}^n (\psi_i - \mathbb{E}\psi_i) - \frac{n(n-1)\ell_n}{n^2h^{d+2}} \mathbb{E}\psi_1 \equiv L_{21} + L_{22} + L_{23}.$$

By Lemma A2, condition (B7), for  $k < r\left(1 - \frac{1}{p_1}\right) - 1$ , we have

$$\mathbb{E}L_{21}^k \le Cn^k \left(\frac{\ell_n}{n^2h^{d+2}}\right)^k h^{d/p_1}.$$

Thus, if k is chosen to be 4, in view of condition (B6), then

$$\frac{\mathbb{E}_{21}^k}{\ell_n^2 \gamma_n^k} \le \frac{C}{n(\log n)^2}.$$

Since  $\psi_i$  includes two terms and the proofs of these two terms are the same, we just consider one of its terms  $\xi_i = \mathbb{E}_i \mathbb{I}(|X_j - \widetilde{X}_i| \leq h)$ . It is easy to see that  $|\xi_i| \leq Ch^{d-2} \mathbb{I}_{(t_{1,2} - h \leq X_{i,12} \leq t_{1,2} + h)}$  and  $(\mathbb{E} |\xi_i|^{r_3})^{\frac{2}{r_3}} \leq Ch^{2d-4} \cdot h^{\frac{4}{r_3}} = Ch^{2d-4} \cdot h^{2(1-\frac{1+\varepsilon}{r})} = Ch^{2d-2(1+\frac{1+\varepsilon}{r})}$ . Then, from Lemma A1, we have

$$\mathbb{P}\left\{ \left| \frac{\ell_n}{nh^{d+2}} \sum_{i=1}^n \left( \xi_i - \mathbb{E}\xi_i \right) \right| \ge \gamma_n \right\} \le \exp\left\{ \frac{-\left( \frac{\gamma_n nh^{d+2}}{\ell_n} \right)^2}{\frac{n}{2q} q \left( \mathbb{E} \left| \xi_i \right|^{r_3} \right)^{\frac{2}{r_3}} + qM \frac{\gamma_n nh^{d+2}}{\ell_n}} \right\} + \frac{n}{q} \beta(q) \le n^{-CB} + \frac{n}{q} \beta(q)$$

with  $1 - \frac{2}{r_3} = \frac{1+\varepsilon}{r}$  and  $q = \frac{nh^2}{\log n}$ . Then,  $\frac{1}{\ell_n^2} \frac{n}{q} \beta(q) \to 0$ . Since  $\mathbb{E}\psi_i = Ch^d$ ,  $|L_{23}| \leq \frac{1}{2} \gamma_n$ . Next, we will consider the convergence rate of  $L_3$ . Let

$$\psi_{ij} = \mathbb{I}(h - \ell_n \le X_i - \widetilde{X}_j \le h + \ell_n) + \mathbb{I}(h - \ell_n \le X_j - \widetilde{X}_i \le h + \ell_n),$$

 $\psi_i = \mathbb{E}\psi_{ij}$  and  $\varphi_{ij} = \psi_{ij} - \psi_i - \psi_j + \mathbb{E}\psi_1$ . Then,

$$L_{3} = \frac{2}{n^{2}h^{d+1}} \sum_{1 \leq i \leq j \leq n} \varphi_{ij} + \frac{2(n-1)}{n^{2}h^{d+1}} \sum_{i=1}^{n} (\psi_{i} - \mathbb{E}\psi_{i}) - \frac{n(n-1)}{n^{2}h^{d+1}} \mathbb{E}\psi_{1} = L_{31} + L_{32} + L_{33}.$$

Again by exploiting Lemma A2 and condition (B6), if k = 8 and  $r \ge 5.4$ , we have the following inequality

$$\frac{1}{\ell_n^2 \gamma_n^k} \mathbb{E} L_{31}^k \leq \frac{1}{\ell_n^2 \gamma_n^k} \cdot \frac{C}{n^k h^{dk+2k}} \left( h^d \frac{\ell_n}{h} \right)^{1-\frac{k+1}{r} - \varepsilon_1} \leq \frac{C}{n (\log n)^2}.$$

Similar to the case of  $L_{22}$ , we only consider one of the terms in  $L_{32}$ , i.e.,

$$\mathbb{E}_i \mathbb{I}(h - \ell_n \le |X_i - \widetilde{X}_j| \le h + \ell_n).$$

After some calculation, it is asymptotically equivalent to

$$h^{d-2} f(\bar{x}_{1,u}) \left( C_1 \mathbb{I} \left( h - \ell_n \le |x_{1,u} - t_{1,u}| \le h + \ell_n \right) + C_2 \mathbb{I} \left( |x_{1,u} - t_{1,u}| \le h - \ell_n \right) \frac{\ell_n}{h} \right).$$

Since the proofs of the two terms above are the same, we only consider the first one, say  $\xi_i$ . Thus,  $|\xi_i| \leq C h^{d-2}$  and  $(\mathbb{E} |\xi_i|^{r_3})^{\frac{2}{r_3}} \leq \left(h^{d-2}\right)^2 \left(h^2 \frac{\ell_n}{h}\right)^{1-\frac{1+\varepsilon}{r}}$  with  $\frac{2}{r_3} = 1 - \frac{1+\varepsilon}{r}$ .  $\frac{1}{h^{d+1}} \mathbb{E} \xi_i \leq \gamma_n$ . By using Lemma A1, we have

$$\mathbb{P}\left\{\left|\frac{1}{nh^{d+1}}\sum_{i=1}^{n}\left(\xi_{i}-\mathbb{E}\xi_{i}\right)\right| \geq \gamma_{n}\right\} \leq \exp\left\{\frac{-\left(\gamma_{n}nh^{d+1}\right)^{2}}{\frac{n}{2q}q\left(\mathbb{E}\left|\psi_{i}\right|^{r_{3}}\right)^{\frac{2}{r_{3}}}+qM\gamma_{n}nh^{d+1}}\right\} + \frac{n}{q}\beta(q),$$

where  $q = \frac{\gamma_n nh^3}{\log n}$ . Since  $\varepsilon$  is an arbitrary small constant,  $\varepsilon_n$  can be replaced by  $\left(nh^{4+\frac{1+\varepsilon}{r}}\right)^{-\frac{1}{2}}$ . This completes the proof of the lemma.

#### Proof of Lemma 5.1

*Proof.* In view of the definitions of  $\widehat{q}_0(X_m)$  and  $q_0(X_m)$ , it suffices to prove that, for  $1 \leq u \leq d$ ,

$$\frac{1}{nh_G^2} \sum_{m=1}^n K'_{G,m} (\widehat{q}_u(X_{m,u}) - q_u(X_{m,u})) = O\left(\frac{n^{-\varepsilon_0}}{\sqrt{nh_G}}\right).$$

We here only give the proof for the case  $2 \le u \le d$  while the proof of the case u=1 is similar. Noting from the proof of Theorem 3.1, we know that  $\widehat{q}_u(x_u) - q_u(x_u) = I_1 - I_2 - I_3$ . At a random point  $X_{m,u}$ , we write  $\widehat{q}_u(X_{m,u}) - q_u(X_{m,u}) = I_{1,m} - I_{2,m} - I_{3,m}$ , in which  $I_{i,m}$  is equal to  $I_m$  with  $x_u$  replaced by  $X_{m,u}$  for i=1,2,3. Denote by  $J_i = \frac{1}{nh_G^2} \sum_{m=1}^n K'_{G,m} I_{i,m}$ , i=1,2,3. Thus, it suffices to prove that  $J_i = O\left(\frac{n^{-\varepsilon_0}}{\sqrt{nh_G}}\right)$  for each i=1,2,3. Denote by

$$l(X_{m,u}, \widetilde{X}_j) = K'_{G,m} \int_{x_{u,0}}^{X_{m,u}} \int \frac{w_1(t_1) \left( \partial_u q(\widetilde{X}_j) \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{j,\bar{u}}) - D_u(t_1, t_u) \right)}{D_{1,u}(t_1, t_u)} dt_1 dt_u,$$

$$J_{11} = \frac{1}{n^2 h_G^2} \sum_{i=1}^n \sum_{j=1, j \neq m}^n l(X_{m,u}, \widetilde{X}_j)$$
 and

$$J_{12} = \frac{1}{n^2 h_G^2} \sum_{m=1}^n \sum_{i=1}^n \sum_{j \neq m}^n K_{G,m}^{'} \int_{x_{u,0}}^{X_{m,u}} \int \frac{w_1(t_1)}{D_{1,u}(t_1,t_u)} \left( \partial_u \widehat{q}_u(\widetilde{X}_j) - \partial_u q_u(\widetilde{X}_j) \right) \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{j,\bar{u}}) dt_1 dt_u.$$

Then,  $J_1 = J_{11} + J_{12}$ . Rewrite

$$J_{11} = \frac{1}{n^{2}h_{G}^{2}} \sum_{m=1}^{n} \sum_{j=1, j \neq m}^{n} [l(X_{m,u}, \widetilde{X}_{j}) - \mathbb{E}_{j}l(X_{m,u}, \widetilde{X}_{j})] + \frac{n-1}{n^{2}h_{G}^{2}} \sum_{j=1}^{n} \mathbb{E}_{j}l(X_{m,u}, \widetilde{X}_{j}) = J_{13} + J_{14}.$$

$$\left(\frac{\sqrt{nh_{G}}}{n^{-\varepsilon_{0}}}\right)^{k} \mathbb{E} |J_{13}|^{k} \leq Cn^{k} \left(\frac{\sqrt{nh_{G}}}{n^{2-\varepsilon_{0}}h_{G}^{2}}\right)^{k} M_{sk}^{k} = C \left(n^{1+3\varepsilon_{0}}h_{G}^{3}\right)^{-\frac{k}{2}} h_{G}^{\frac{1}{s}} \leq \frac{1}{n(\log n)^{2}}$$

$$\frac{\sqrt{nh_{G}}}{n^{-\varepsilon_{0}}} J_{14} = \frac{\sqrt{nh_{G}}}{n^{-\varepsilon_{0}}} \cdot \frac{n-1}{n^{2}h_{G}^{2}} \cdot \frac{Ch_{G}^{2}\sqrt{n}}{\log n} \to 0.$$

We now consider  $J_{12}$ . Noting from the proof of Theorem B1, to deal with  $J_{12}$ , it suffices to prove the following four expressions

$$\frac{1}{n^3 h^{d+1} h_G^2} \sum_{m=1}^n \sum_{j=1, j \neq m}^n \sum_{i=1, i \neq j}^n K_{G,m}^{'} \int_{x_{u,0}}^{X_{m,u}} \int \frac{w_1(t_1) \mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{j,\bar{u}})}{D_{1,u}(t_1, t_u)} Q_{ij,t_{1,u}}^{-1} K_{ij,t_{1,u}}$$

$$\cdot A_{ij,t_{1,u}}(\alpha - \mathbb{I}(\varepsilon_i \leq -r_{ij})) dt_1 dt_u$$

$$\frac{1}{n^3h^{d+1}h_G^2} \sum_{m=1}^n \sum_{i=1}^n \sum_{j\neq m}^n \sum_{i=1,i\neq m}^n K_{G,m}^{'} \int_{x_{u,0}}^{X_{m,u}} \int \frac{w_1(t_1)\mathbb{I}(X_{j,\bar{u}} \in \mathcal{X}_{j,\bar{u}})}{D_{1,u}(t_1,t_u)} \Delta_{ij}(t_{1,u},\beta) dt_1 dt_u,$$

 $\frac{1}{n^2hh_G^2}n\cdot nh_G\cdot (\delta^2+\delta h^p)$  and  $\frac{1}{n^2hh_G^2}n\cdot nh_G\cdot \frac{C}{nh^d}$  are all of the order  $\frac{n^{-\epsilon_0}}{\sqrt{nh_G}}$ , and moreover, the second expression is of order uniform for  $|\beta-\beta_{\tau}|\leq \frac{B}{\sqrt{nh^{(1+\frac{1}{\tau})^d+\epsilon}}}$ . In more details, the first three expressions come from the RHS of (B.11) and the last two terms on the LHS of (B.11), and the last expression comes from the last term of (B.12). For convenience, denote the first two terms by  $J_{13}$  and  $J_{14}$ , respectively. As for  $J_{13}$ , by using the SLLN three times, it can be obtained that

where the last step follows from the property of the conditional expectation.

$$\begin{split} \zeta_{m,i,j} &= K_{G,m}^{'} \int_{x_{u,0}}^{X_{m,u}} \int \frac{w_1(t_1)}{D_{1,u}(t_1,t_u)} Q_{jn,t_{1,u}}^{-1} \left( \Delta_{ij}(t_{1,u},\beta) - \mathbb{E} \Delta_{ij}(t_{1,u},\beta) \right) \mathrm{d}t_1 \mathrm{d}t_u \\ &\frac{1}{n^3 h^{d+1} h_G^2} \sum_{m=1}^n \sum_{j=1,j\neq m}^n \sum_{i=1,i\neq j}^n \zeta_{m,i,j} = \frac{1}{n^3 h^{d+1} h_G^2} \sum_{j=1}^n \sum_{i=1,i\neq j}^n \sum_{m=1,m\neq j}^n \zeta_{m,i,j} \\ &\eta_{i,j}(\beta) = \mathbb{E}_{i,j} \left[ K_{G,m}^{'} \int_{x_{u,0}}^{X_{m,u}} \int \frac{w_1(t_1)}{D_{1,u}(t_1,t_u)} Q_{jn,t_{1,u}}^{-1} \left( \Delta_{ij}(t_{1,u},\beta) - \mathbb{E} \Delta_{ij}(t_{1,u},\beta) \right) \mathrm{d}t_1 \mathrm{d}t_u \right] \\ &\frac{1}{n^3 h^{d+1} h_G^2} \sum_{j=1}^n \sum_{i=1,i\neq j}^n \sum_{m=1,m\neq j}^n \left( \zeta_{m,i,j} - \eta_{i,j}(\beta) \right) \\ &\xi_{m,i} = \zeta_{m,i,j} - \eta_{i,j}(\beta) \\ &\frac{(\sqrt{nh_G})^k \mathbb{E} |\xi_{m,i}|^k}{(n^2 h^{d+1} h_G^2)^k} \leq \frac{C}{n(\log n)^2}. \end{split}$$
 If  $k=6$ , 
$$\frac{Cnh_G h^{d+p}}{(\sqrt{nh_G}nh^{d+1}h_G)^k} = \frac{Cnh_G h^{d+p}}{(\sqrt{nh_G}nh^{d+1}h_G)^6} \leq \frac{C}{(\log n)^2}. \end{split}$$

This completes the proof.

#### Proof of Lemma A5

Proof. Let  $\delta_n$  denote the RHS of (B.9). In order to prove uniformity, we adopt a similar division on the domain  $\mathcal{X}_{1,u}$  and the sphere  $|\beta - \beta_{j,t_{1,u}}| = \delta$  as that of Lemma B3. And the same notations are used here. Let  $\ell_n = d_1 = \delta h$  and  $q = \delta_n n^{-\frac{\varepsilon}{2}}$ . For any real  $r_2 > 0$ , on the event  $\{|X_i - \widetilde{X}_j| \leq h\}$ , it can be inferred that

$$\mathbb{E}_{j}\left[\left|\mathbb{I}\left(\varepsilon_{i} \leq -r_{ij,t_{1,u}} - P_{ij,t_{1,u}}\right) - \mathbb{I}\left(\varepsilon_{i} \leq -r_{ij,t_{1,u}}\right)\right|^{r_{2}} \mid X_{i}\right] = \left|\int_{-r_{ij,t_{1,u}}}^{-r_{ij,t_{1,u}} - P_{ij,t_{1,u}}} g(X_{i}, u) du\right| \leq C\delta.$$

Furthermore, we have  $\mathbb{E}_j |\Delta_{ij}(t_{1,u},\beta)|^{r_2} \leq Ch^d\delta$ . Then, as previously proved, in view of the inequality (A.2), we know that

$$\frac{n}{q} \cdot q \left( h^d \delta \right)^{\frac{2}{r_2}} < Cq \delta_n,$$

where  $1 - \frac{2}{r_2} = \frac{1+\varepsilon}{r}$  for some sufficiently small constant  $\varepsilon > 0$ . Then, from Lemma A1, it follows that

$$\left\| \mathbb{P}\left( \left| \sum_{i=1}^{n} \left( \Delta_{ij}(t_{1,u}, \beta) - \mathbb{E}_{j} \Delta_{ij}(t_{1,u}, \beta) \right) \right| > \frac{\delta_{n}}{2} \right) \le n^{-CB} + \frac{n}{q} \beta(q) + C\left( \frac{q}{\delta_{n}} \right)^{r_{1}}.$$
(D.2)

In addition, note that

$$\left| \beta^{\mathrm{T}} A_{ij,t_{1,u}} - \gamma^{\mathrm{T}} A_{ij,s} \right| 
\leq \left| (\beta - \beta_{j,t_{1,u}})^{\mathrm{T}} A_{ij,t_{1,u}} - (\gamma - \beta_{j,s})^{\mathrm{T}} A_{ij,s} \right| + \left| \beta_{j,t_{1,u}}^{\mathrm{T}} A_{ij,t_{1,u}} - \beta_{j,s}^{\mathrm{T}} A_{ij,s} \right| \leq C \ell_{n}.$$
(D.3)

Analogous to the proof of Lemma A2, from (B.5) and (D.3), it follows that

$$\begin{aligned} & |[\Delta_{ij}(s,\gamma) - \mathbb{E}_{j}\Delta_{ij}(s,\gamma)] - [\Delta_{ij}(t_{1,u},\beta) - \mathbb{E}_{j}\Delta_{ij}(t_{1,u},\beta)]| \\ & \leq & C\mathbb{I}_{\left(\left|X_{i} - \widetilde{X}_{j}\right| \leq h\right)} \left\{ \frac{\ell_{n}}{h} \mathbb{I}_{\left(\left|\varepsilon_{i} + r_{ij,t_{1,u}}\right| \leq \delta\right)} + \mathbb{I}_{\left(\left|\varepsilon_{i} + r_{ij,t_{1,u}} + P_{ij,t_{1,u}}\right| \leq C\ell_{n}\right)} + \mathbb{I}_{\left(\left|\varepsilon_{i} + r_{ij,t_{1,u}}\right| \leq C\ell_{n}\right)} \right\} \\ & + C\mathbb{I}_{\left(h - \ell_{n} \leq \left|X_{i} - \widetilde{X}_{j}\right| \leq h + \ell_{n}\right)} \mathbb{I}_{\left(\left|\varepsilon_{i} + r_{ij,t_{1,u}}\right| \leq C(\ell_{n} + \delta)\right)}. \end{aligned}$$

Then, through using Lemma A1 and (A.2), we know that the following four probabilities

$$\mathbb{P}\left(\left|\sum_{i} \frac{\ell_{n}}{h} \mathbb{I}_{\left(\left|X_{i}-\widetilde{X}_{j}\right| \leq h\right)} \mathbb{I}_{\left(\left|\varepsilon_{i}+r_{ij,t_{1,u}}\right| \leq \delta\right)}\right| > \frac{\delta_{n}}{8}\right),$$

$$\mathbb{P}\left(\left|\sum_{i} \mathbb{I}_{\left(\left|X_{i}-\widetilde{X}_{j}\right| \leq h\right)} \mathbb{I}_{\left(\left|\varepsilon_{i}+r_{ij,t_{1,u}}+P_{ij,t_{1,u}}\right| \leq C\ell_{n}\right)}\right| > \frac{\delta_{n}}{8}\right),$$

$$\mathbb{P}\left(\left|\sum_{i} \mathbb{I}_{\left(\left|X_{i}-\widetilde{X}_{j_{n}}\right| \leq h\right)} \mathbb{I}_{\left(\left|\varepsilon_{i}+r_{ij,t_{1,u}}\right| \leq C\ell_{n}\right)}\right| > \frac{\delta_{n}}{8}\right)$$

and

$$\mathbb{P}\left(\left|\sum_{i} \mathbb{I}_{\left(h-\ell_{n} \leq \left|X_{i}-\tilde{X}_{j_{n}}^{u}\right| \leq h+\ell_{n}\right)} \mathbb{I}_{\left(\left|\varepsilon_{i}+r_{ij,t_{1,u}}\right| \leq C(\ell_{n}+\delta)\right)}\right| > \frac{\delta_{n}}{8}\right)$$

are bounded by the RHS of (D.2), where the relationships

$$\mathbb{E}_{j} \mathbb{I}_{\left(\left|X_{i}-\widetilde{X}_{j}\right| \leq h\right)} \mathbb{I}_{\left(\left|\varepsilon_{i}+r_{ij,t_{1,u}}+P_{ij,t_{1,u}}\right| \leq C\ell_{n}\right)} \leq Ch^{d} \ell_{n}$$

and

$$\mathbb{E}_{j}\left(\mathbb{I}_{\left(h-\ell_{n}\leq\left|X_{i}-\widetilde{X}_{j}\right|\leq h+\ell_{n}\right)}\mathbb{I}_{\left(\left|\varepsilon_{i}+r_{ij,t}\right|\leq C\left(\ell_{n}+\delta\right)\right)}\right)\leq C\left(\ell_{n}+\delta\right)\ell_{n}h^{d-1}$$

are used. Also, the inequality (B.7) follows from the condition on r and condition (B3). Similar as the proof of Lemma B1, (B.9) can be inferred from the Borel-Cantelli lemma.

#### Proof of Corollary 5.1 (Sketch)

*Proof.* From (5.3), we have

$$(1 - \alpha) - F_n(G(v)|v)$$

$$= \frac{\sum_{i=1}^n \left\{ \left[ K_{G,i,v} \left( (1 - \alpha) - \mathbb{I}(Y_i \leq G(v)) \right) \right] - \mathbb{E}\left[ K_{G,i,v} \left( (1 - \alpha) - \mathbb{I}(Y_i \leq G(v)) \right) \right] \right\}}{\sum_{i=1}^n K_{G,i}}$$

$$- \frac{n \left( \mathbb{E}\left[ K_{G,i,v} \mathbb{I}(Y_i \leq G(v)) \right] - \mathbb{E}\left[ K_{G,i,v} \mathbb{I}(\varepsilon_i \leq 0) \right] \right)}{\sum_{i=1}^n K_{G,i}}$$

$$= I_1 + I_2. \tag{D.4}$$

It can be proved that

$$\frac{\sum_{i=1}^{n} \left\{ \left[ K_{G,i,v} \left( \left( 1 - \alpha \right) - \mathbb{I} \left( Y_{i} \leq G(v) \right) \right) \right] - \mathbb{E} \left[ K_{G,i,v} \left( \left( 1 - \alpha \right) - \mathbb{I} \left( Y_{i} \leq G(v) \right) \right) \right] \right\}}{\sqrt{\alpha \left( 1 - \alpha \right) f_{q_{0}}(v) n h_{G}}} \stackrel{d}{\to} \mathcal{N}(0,1). \tag{D.5}$$

Moreover, it can be proved that

$$\sum_{i=1}^{n} K_{G,i,v} = n \mathbb{E} K_{G,i,v} + O\left(\frac{\sqrt{\log n}}{\sqrt{nh_G}}\right) = nh_G f_{q_0}(v) \left(1 + O(h_G^2)\right) + O\left(\frac{\sqrt{\log n}}{\sqrt{nh_G}}\right).$$
 (D.6)

By variable substitution, Taylor's expansion and  $\int sK_G(s) ds = 0$ , it follows that

$$\frac{\mathbb{E}\left[K_{G,i,v}\mathbb{I}\left(Y_{i} \leq G(v)\right)\right] - (1-\alpha)\mathbb{E}K_{G,i,v}}{h_{G}f_{q_{0}}(v)} = a(v)h_{G}^{2} + o\left(h_{G}^{2}\right). \tag{D.7}$$

Then, Corollary 5.1 can be inferred from part i) of Theorem 5.1, (D.4), (D.5), (D.6), (D.7) and the fact that  $\left(nh^{1+\frac{1+\epsilon}{r}}\right)^{-\frac{1}{2}} = o(h_G^2)$ .

### References

Arcones, M.A. (1998). The law of large numbers for U-statistics under absolute regularity, *Elect. Comm.* in *Probab.*, 3, 13–19.

Chaudhuri, P. (1991). Global nonparametric estimation of conditional quantile functions and their derivatives, J. Multivariate Anal., 39(2), 246–269.

Honda, T. (2004). Quantile regression in varying coefficient, J. Statist. Plann. Inference, 121, 113–125.

Lee, A.J. (1990). U-statistics: Theory and Practice, Marcel Dekker, Inc., New York.

Masry, E. (1996). Multivariate local polynomial regression for time series: Uniform strong consistency and rates, J. Time Ser. Anal., 17(6), 571–599.

Mehra, K.L., Rama Krishnaiah, Y.S. and Rao, M.S. (1992). Bahadur representation of sample conditional quantiles based on smoothed conditional empirical distribution function, *Bull. Inform. Cybernet.*, 25(1-2), 99–107.

Stewart, G.W. and Sun J.G. (1990). Matrix Perturbation Theory, Academic Press: Boston, MA.