

# Cohomology of $n$ -Lie algebras in Loday-Pirashvili category

Tao Zhang\*

College of Mathematics and Information Science, Henan Normal University, Xinxiang 453007, PR China

## Abstract

In this article, we present the concept of  $n$ -Lie algebras in the Loday-Pirashvili category. We examine the representation and cohomology theory of  $n$ -Lie algebras in this category, and also explore their applications in infinitesimal deformation and abelian extension theory.

## 1 Introduction

Filippov introduced the concept of  $n$ -Lie algebras in [13], which may also be referred to as Filippov algebras or Nambu-Lie algebras. An  $n$ -Lie algebra, denoted as  $\mathfrak{g}$ , is a vector space equipped with an  $n$ -ary totally skew-symmetric linear map ( $n$ -bracket)  $\mathfrak{g} \times \cdots \times \mathfrak{g} \rightarrow \mathfrak{g} : (x_1, \dots, x_n) \rightarrow [x_1, \dots, x_n]$  satisfying the following fundamental identity:

$$[x_1, \dots, x_{n-1}, [y_1, y_2, \dots, y_n]] = \sum_{i=1}^n [y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n], \quad (1)$$

for all  $x_i, y_i \in \mathfrak{g}$ . It dates back to Nambu's work [25] to generalize the classical Hamiltonian mechanics when  $n = 3$ :

$$\{h_1, h_2, \{x_1, x_2, x_3, z\}\} = \{\{h_1, h_2, x\}, y, z\} + \{x, \{h_1, h_2, y\}, z\} + \{x_1, x_2, x_3, \{h_1, h_2, z\}\},$$

where  $h_1, h_2$  are Hamiltonian and  $\{\cdot, \cdot, \cdot\}$  is a ternary product. When the  $n$ -ary bracket in (1) is not skew-symmetric, it is called a Leibniz  $n$ -algebra [8, 31, 4, 6].

The study of  $n$ -Lie algebra is closely related to many fields in mathematics and mathematical physics. For example, the foundations of the theory of Nambu-Poisson manifolds were developed by Takhtajan in [30]. The application of  $n$ -Lie algebras to the study of the supersymmetric gauge theory of multiple M2-branes and D-branes is given in [14, 27]. The investigation of 3-Lie algebras for integrable system is given in [7].

The algebraic theory of  $n$ -Lie algebras has been extensively studied by several authors. In particular, the (co)homology theory for  $n$ -Lie algebras was introduced by Takhtajan in [31, 8], Gautheron in [10], and Rotkiewicz in [28]. One compare to the homology theory introduced in [9]. This theory is specifically tailored to the study of formal differential geometry and formal deformations of Nambu structures. The representation and cohomology theory of Leibniz  $n$ -algebras is also investigated in [4, 6]. Additionally, for the special case when  $n = 3$ ,

---

\*E-mail: zhangtao@htu.edu.cn

<sup>0</sup>Keyword: Loday-Pirashvili category,  $n$ -Lie algebras, Leibniz algebras, cohomology, deformations

<sup>0</sup>Mathematics Subject Classification (2020): 17A32, 17A42, 17B56.

the research can be found in [2, 12, 22, 32, 33, 34]. Recently, non-abelian extensions of  $n$ -Lie algebras and perfect Leibniz  $n$ -algebras were studied in [1, 5].

In [24], Loday and Pirashvili introduced a tensor category  $\mathcal{LM}$  of linear maps. Roughly speaking,  $\mathcal{LM}$  consists of linear maps  $f : V \rightarrow W$  as objects. A morphism between two objects  $f : V \rightarrow W$  and  $f' : V' \rightarrow W'$  is a pair of linear maps  $(\phi, \psi)$ , where  $\phi : V \rightarrow V'$  and  $\psi : W \rightarrow W'$ , such that  $\psi \circ f = f' \circ \phi$ . In the same paper, they also introduced the concept of Lie algebra  $(M, \mathfrak{g}, f)$  in  $\mathcal{LM}$ . This Lie algebra is a representation  $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(M)$  of an ordinary Lie algebra  $\mathfrak{g}$  on a vector space  $M$ , such that  $f : M \rightarrow \mathfrak{g}$  is an equivariant map satisfying the equation:

$$f(\rho(x)(m)) = [x, f(m)]_{\mathfrak{g}}, \quad (2)$$

for all  $x \in \mathfrak{g}$  and  $m \in M$ . When  $f$  is a surjective map, then the above equation (24) is equivalent to

$$f(\rho(f(m))(n)) = [f(m), f(n)]_{\mathfrak{g}}, \quad (3)$$

for all  $m, n \in M$ . This is called an embedding tensor or tensor hierarchy algebra in [18, 20, 26]. Kurdiani constructed a novel cohomology theory for Lie algebras in  $\mathcal{LM}$  using the iterated cone constructions in [19], Rovi studied Lie algebroids in this category [29].

In this paper, we introduce the concept of  $n$ -Lie algebras in  $\mathcal{LM}$ . Let  $\mathfrak{g}$  be an  $n$ -Lie algebra,  $M$  be a vector space, and  $\rho$  a representation of  $\mathfrak{g}$  on  $M$ . We define an  $n$ -Lie algebra in  $\mathcal{LM}$  as an equivariant map  $f : M \rightarrow \mathfrak{g}$  satisfying the following condition:

$$f(\rho(x_1, \dots, x_{n-1})(m)) = [x_1, \dots, x_{n-1}, f(m)]_{\mathfrak{g}}, \quad (4)$$

for all  $x \in \mathfrak{g}$  and  $m \in M$ , where  $\rho : \wedge^{n-1}\mathfrak{g} \rightarrow \mathfrak{gl}(M)$  is a representation of  $\mathfrak{g}$ . When  $n = 2$ , we recover Loday and Pirashvili's Lie algebra  $(M, \mathfrak{g}, f)$  in  $\mathcal{LM}$  mentioned above. Therefore, the results of this paper are an  $n$ -order generalization of the results in [24, 19]. Additionally, we show that this new concept of  $n$ -Lie algebras in  $\mathcal{LM}$  gives rise to Leibniz  $n$ -algebras. Consequently, we investigate the representation and cohomology theory for this type of  $n$ -Lie algebras. Moreover, this paper provides a detailed exposition of the construction and properties of this cohomology theory. In a key observation, we find that the cohomology theory constructed by Kurdiani in [19] can be extended to the case of Leibniz algebras and  $n$ -Lie algebras in  $\mathcal{LM}$ . Finally, we study their infinitesimal deformations and abelian extension theory in detail as applications.

The organization of this paper is as follows. In Section 2, we review some basic facts about  $n$ -Lie algebras and Leibniz  $n$ -algebras. In Section 3, we introduce the concept of  $n$ -Lie algebras in  $\mathcal{LM}$  and study its elementary properties. We investigate the representation and cohomology theory of  $n$ -Lie algebras in  $\mathcal{LM}$ . Low-dimensional cohomologies are given in detail in this section. In Section 4, we prove that equivalent classes of abelian extensions are in one-to-one correspondence with the elements of the second cohomology groups. In Section 5, we study infinitesimal deformations of  $n$ -Lie algebras in  $\mathcal{LM}$ . The notion of Nijenhuis operators is introduced to describe trivial deformations.

## 2 Cohomology of $n$ -Lie algebras

First, we recall some facts about  $n$ -Lie algebras and Leibniz algebras, basic references are [10, 13, 30, 31, 23].

**Definition 2.1** ([13]). An  $n$ -Lie algebra is a vector space  $\mathfrak{g}$  with an  $n$ -ary totally skew-symmetric linear map ( $n$ -bracket)  $\wedge^n \mathfrak{g} \rightarrow \mathfrak{g} : (x_1, \dots, x_n) \rightarrow [x_1, \dots, x_n]$  satisfying the following fundamental identity:

$$[x_1, \dots, x_{n-1}, [y_1, y_2, \dots, y_n]] = \sum_{i=1}^n [y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n], \quad (5)$$

for all  $x_i, y_i \in \mathfrak{g}$ . If the  $n$ -bracket is not skew-symmetric, then it is called a Leibniz  $n$ -algebra.

Let  $\mathfrak{g}$  and  $\mathfrak{g}'$  be two  $n$ -Lie algebras. An  $n$ -Lie algebra homomorphism is a linear map  $\phi : \mathfrak{g} \rightarrow \mathfrak{g}'$  such that following condition holds

$$\phi([x_1, \dots, x_n]) = [\phi(x_1), \dots, \phi(x_n)]'. \quad (6)$$

Denote by  $x = (x_1, \dots, x_{n-1})$  and  $\text{ad}(x)y_i = [x_1, \dots, x_{n-1}, y_i]$ , then the above equality (5) can be rewritten in the form

$$\text{ad}(x)[y_1, y_2, \dots, y_n] = \sum_{i=1}^n [y_1, \dots, \text{ad}(x)y_i, \dots, y_n]. \quad (7)$$

Put  $L \triangleq \wedge^{n-1} \mathfrak{g}$ . The elements in  $L$  are called fundamental objects. Define an operation on fundamental objects by

$$x \circ y = \sum_{i=1}^{n-1} (y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1}). \quad (8)$$

It can be proved as in [8] that  $L$  is a Leibniz algebra satisfying the following Leibniz rule

$$x \circ (y \circ z) = (x \circ y) \circ z + y \circ (x \circ z),$$

and

$$\text{ad}(x)\text{ad}(y)w - \text{ad}(y)\text{ad}(x)w = \text{ad}(x \circ y)w,$$

for all  $x, y, z \in L, w \in \mathfrak{g}$ , i.e.  $\text{ad} : L \rightarrow \text{End}(\mathfrak{g})$  is a homomorphism of Leibniz algebras.

Recall that for a Leibniz algebra  $L$ , a representation is a vector space  $M$  together with two bilinear maps

$$[\cdot, \cdot]_L : L \otimes M \rightarrow M \quad \text{and} \quad [\cdot, \cdot]_R : M \otimes L \rightarrow M,$$

satisfying the following three axioms

- (LLM)  $[x \circ y, m]_L = [x, [y, m]_L]_L - [y, [x, m]_L]_L,$
- (LML)  $[m, x \circ y]_R = [[m, x]_R, y]_R + [x, [m, y]_R]_L,$
- (MLL)  $[m, x \circ y]_R = [x, [m, y]_R]_L - [[x, m]_L, y]_R.$

By (LML) and (MLL) we also have

- (MMM)  $[[m, x]_R, y]_R + [[x, m]_L, y]_R = 0.$

In fact, assume (LLM), one of (LML),(MLL),(MMM) can be derived from the other two.

The cohomology of Leibniz algebras was defined in [23] as follows. Let  $L$  be a Leibniz algebra,  $M$  be a representation of  $L$ . We denote the cochain complex by

$$CL^p(L, M) \triangleq \text{Hom}(\otimes^p L, M).$$

Define the coboundary operator

$$d_{p-1} : CL^{p-1}(L, M) \rightarrow CL^p(L, M)$$

by

$$\begin{aligned} & d_{p-1}\omega(x_1, x_2, \dots, x_p) \\ = & \sum_{i=1}^{p-1} (-1)^{i+1} [x_i, \omega(x_1, \dots, \widehat{x}_i, \dots, x_p)]_L + (-1)^p [\omega(x_1, \dots, x_{p-1}), x_p]_R \\ & + \sum_{1 \leq i < j \leq p} (-1)^i \omega(x_1, \dots, \widehat{x}_i, \dots, x_{j-1}, x_i \circ x_j, x_{j+1}, \dots, x_p), \end{aligned}$$

for all  $x_i \in L$ . It is proved that  $d \circ d = 0$ . For more details, see [23].

**Definition 2.2.** Let  $\mathfrak{g}$  be an  $n$ -Lie algebra and  $V$  be a vector space. Then  $(V, \rho)$  is called a representation of  $\mathfrak{g}$  if the following two conditions (R1) and (R2) are satisfied:

$$(R1) \quad [\rho(x_1, \dots, x_{n-1}), \rho(y_1, \dots, y_{n-1})] = \rho((x_1, \dots, x_{n-1}) \circ (y_1, \dots, y_{n-1})),$$

$$(R2) \quad \rho(x_1, \dots, x_{n-2}, [y_1, \dots, y_n]) = \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n) \rho(x_1, \dots, x_{n-2}, y_i).$$

**Remark 2.3.** The condition (R1) is equivalent to the following equation

$$\begin{aligned} & \rho(x_1, \dots, x_{n-1}) \rho(y_1, \dots, y_{n-1}) \\ = & \sum_{i=1}^{n-1} \rho(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1}) + \rho(y_1, \dots, y_{n-1}) \rho(x_1, \dots, x_{n-1}). \end{aligned} \quad (9)$$

When only this condition is satisfied, we call  $(V, \rho)$  a weak representation of  $\mathfrak{g}$ .

By direct computation, it is easy to obtain the following result.

**Proposition 2.4.** Given a representation  $\rho$  of the  $n$ -Lie algebra  $\mathfrak{g}$  on the vector space  $V$ . Define a skew-symmetric  $n$ -bracket on  $\mathfrak{g} \oplus V$  by

$$[x_1 + u_1, \dots, x_n + u_n] = [x_1, \dots, x_n] + \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n)(u_i), \quad (10)$$

then  $(\mathfrak{g} \oplus V, [,])$  is an  $n$ -Lie algebra, which is called the semidirect product of  $n$ -Lie algebra  $\mathfrak{g}$  with  $V$ . We denote this  $n$ -Lie algebra by  $\mathfrak{g} \ltimes V$ . Define a non-skew-symmetric  $n$ -bracket on  $\mathfrak{g} \oplus V$  by

$$[x_1 + u_1, \dots, x_n + u_n]_H = [x_1, \dots, x_n] + \rho(x_1, \dots, x_{n-1})(u_n), \quad (11)$$

then  $(\mathfrak{g} \oplus V, [, ]_H)$  is a Leibniz  $n$ -algebra, which is called the hemisemidirect product of  $\mathfrak{g}$  with  $V$ . We denote this Leibniz  $n$ -algebra by  $\mathfrak{g} \ltimes_H V$ . When  $n = 2$ , this hemisemidirect product coincides with the notion defined in [16].

For example, given an  $n$ -Lie algebra  $\mathfrak{g}$ , there is a natural adjoint representation on itself. The corresponding representation  $\text{ad}(x_1, \dots, x_{n-1})$  is given by

$$\text{ad}(x_1, \dots, x_{n-1})x_n = [x_1, \dots, x_{n-1}, x_n].$$

Thus, we obtain an  $n$ -Lie algebra  $\mathfrak{g} \times \mathfrak{g}$  by

$$[x_1 + y_1, \dots, x_n + y_n] = [x_1, \dots, x_n] + \sum_{i=1}^n (-1)^{n-i} [x_1, \dots, \widehat{x}_i, \dots, x_n, y_i] \quad (12)$$

and a Leibniz  $n$ -algebra  $\mathfrak{g} \times_H \mathfrak{g}$  by

$$[x_1 + y_1, \dots, x_n + y_n]_H = [x_1, \dots, x_n] + [x_1, \dots, x_{n-1}, y_n] \quad (13)$$

for all  $x_i, y_i \in \mathfrak{g}$ .

Let  $\mathfrak{g}$  be an  $n$ -Lie algebra and  $\rho$  be a representation of  $\mathfrak{g}$  on  $V$ . It can be proved similar as in [4, 6] that there is a representation of  $L = \wedge^{n-1} \mathfrak{g}$  as Leibniz algebra on  $\text{Hom}(\mathfrak{g}, V)$  under the following maps:

$$\begin{aligned} [\cdot, \cdot]_L &: L \otimes \text{Hom}(\mathfrak{g}, V) \rightarrow \text{Hom}(\mathfrak{g}, V), \\ [\cdot, \cdot]_R &: \text{Hom}(\mathfrak{g}, V) \otimes L \rightarrow \text{Hom}(\mathfrak{g}, V), \end{aligned}$$

by

$$[(x_1, \dots, x_{n-1}), \phi]_L(x_n) = \rho(x_1, \dots, x_{n-1})\phi(x_n) - \phi([x_1, \dots, x_{n-1}, x_n]), \quad (14)$$

$$[\phi, (x_1, \dots, x_{n-1})]_R(x_n) = \phi([x_1, \dots, x_n]) - \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n)\phi(x_i), \quad (15)$$

for all  $\phi \in \text{Hom}(\mathfrak{g}, V), x_i \in \mathfrak{g}$ . This observation is important because it provides a new characterization of the representation of  $n$ -Lie algebras in relation to the representation of Leibniz algebras. We will now establish that the converse is also valid.

**Proposition 2.5.** *Let  $\mathfrak{g}$  be an  $n$ -Lie algebra. Then  $\text{Hom}(\mathfrak{g}, V)$  equipped with the above two maps  $[\cdot, \cdot]_L$  and  $[\cdot, \cdot]_R$  is a representation of Leibniz algebra  $L = \wedge^{n-1} \mathfrak{g}$  if and only if the conditions (R1) and (R2) are satisfied, i.e.  $(V, \rho)$  is a representation of  $\mathfrak{g}$ .*

*Proof.* For  $x = (x_1, \dots, x_{n-1}), y = (y_1, \dots, y_{n-1}) \in L, y_n \in \mathfrak{g}$ , we compute the equality

$$[x \circ y, \phi]_L(y_n) = [x, [y, \phi]_L]_L(y_n) - [y, [x, \phi]_L]_L(y_n). \quad (16)$$

By definition (14), the left-hand side of (16) is equal to

$$[x \circ y, \phi]_L(y_n) = \rho(x \circ y)\phi(y_n) - \phi(\text{ad}(x \circ y)y_n),$$

and the right-hand side of (16) is equal to

$$\begin{aligned} & [x, [y, \phi]_L]_L(y_n) - [y, [x, \phi]_L]_L(y_n) \\ &= \rho(x)[y, \phi]_L(y_n) - [y, \phi]_L(\text{ad}(x)y_n) - \rho(y)[x, \phi]_L(y_n) + [x, \phi]_L(\text{ad}(y)y_n) \\ &= \rho(x)\rho(y)\phi(y_n) - \rho(x)\phi(\text{ad}(y)y_n) - \rho(y)\phi(\text{ad}(x)y_n) + \phi(\text{ad}(y)\text{ad}(x)y_n) \\ &\quad - \rho(y)\rho(x)\phi(y_n) + \rho(y)\phi(\text{ad}(x)y_n) + \rho(x)\phi(\text{ad}(y)y_n) - \phi(\text{ad}(x)\text{ad}(y)y_n) \end{aligned}$$

$$\begin{aligned}
&= \rho(x)\rho(y)\phi(y_n) + \phi(\text{ad}(y)\text{ad}(x)y_n) - \rho(y)\rho(x)\phi(y_n) - \phi(\text{ad}(x)\text{ad}(y)y_n) \\
&= [\rho(x), \rho(y)]\phi(y_n) - \phi([\text{ad}(x), \text{ad}(y)]y_n).
\end{aligned}$$

Since  $\text{ad} : L \rightarrow \text{End}(\mathfrak{g})$  is a homomorphism of Leibniz algebras, thus (LLM) holds for  $[\cdot, \cdot]_L$  if and only if (R1) is valid for  $\rho$ .

Next, we compute the equality

$$[[\phi, x]_R, y]_R(y_n) + [[x, \phi]_L, y]_R(y_n) = 0.$$

By (14) and (15) we have

$$\begin{aligned}
&[(x_1, \dots, x_{n-1}), \phi]_L(z) + [\phi, (x_1, \dots, x_{n-1})]_R(z) \\
&= -\sum_{i=1}^{n-1} (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, w) \phi(x_i),
\end{aligned}$$

thus

$$\begin{aligned}
&[(x_1, \dots, x_{n-1}), \phi]_L + [\phi, (x_1, \dots, x_{n-1})]_R \\
&= -\sum_{i=1}^{n-1} (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, \cdot) \phi(x_i),
\end{aligned}$$

where we denote  $\rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, \cdot) \phi(x_i) : \mathfrak{g} \rightarrow V$  by

$$\rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, \cdot) \phi(x_i)(z) = \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, w) \phi(x_i).$$

Now replace  $x_i$  by  $y_i$ ,  $\phi$  by  $-\sum_{i=1}^{n-1} (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, \cdot) \phi(x_i)$  in (15), then we obtain

$$\begin{aligned}
&[[\phi, x]_R + [x, \phi]_L, y]_R(y_n) \\
&= -\sum_{i=1}^{n-1} (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, \cdot) \phi(x_i)([y_1, \dots, y_n]) \\
&\quad + \sum_{i=1}^n (-1)^{n-j} \rho(y_1, \dots, \widehat{y}_j, \dots, y_n) \left\{ \sum_{i=1}^{n-1} (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, \cdot) \phi(x_i)(y_j) \right\} \\
&= -\sum_{i=1}^{n-1} (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, [y_1, \dots, y_n]) \phi(x_i) \\
&\quad + \sum_{i=1}^{n-1} (-1)^{n-i} \left( \sum_{j=1}^n (-1)^{n-j} \rho(y_1, \dots, \widehat{y}_j, \dots, y_n) \rho(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, y_j) \right) \phi(x_i).
\end{aligned}$$

Thus (MMM) is valid for  $[\cdot, \cdot]_L$  and  $[\cdot, \cdot]_R$  if and only if (R2) is valid for  $\rho$ .

At last, one can check that (LML) or (MLL) is valid for  $[\cdot, \cdot]_L$  and  $[\cdot, \cdot]_R$  if and only if (R1) and (R2) are valid for  $\rho$ .  $\square$

Next, we define the cohomology theory of an  $n$ -Lie algebra in virtue of the cohomology theory of Leibniz algebra as follows. A  $p$ -cochain on  $\mathfrak{g}$  with coefficients in a representation  $(V, \rho)$  is a multilinear map  $\omega : \wedge^{p(n-1)+1} \mathfrak{g} \rightarrow V$ . The space of  $p$ -cochains is denoted as  $C^p(\mathfrak{g}, V)$ .

**Theorem 2.6** ([4, 6]). *Let  $\mathfrak{g}$  be an  $n$ -Lie algebra and  $(V, \rho)$  be a representation of  $\mathfrak{g}$ . Then there exists a cochain complex  $\{C(\mathfrak{g}, V) = \bigoplus_{p \geq 0} C^p(\mathfrak{g}, V), \delta\}$  which is a subcomplex of the Leibniz algebra  $L = \wedge^{n-1} \mathfrak{g}$  with coefficients in  $\text{Hom}(\mathfrak{g}, V)$  such that the coboundary operator is given by*

$$\begin{aligned}
& \delta_p \omega(X_1, \dots, X_p, z) \\
= & \sum_{i=1}^p (-1)^i \omega(X_1, \dots, \widehat{X}_i, \dots, X_p, [X_i, z]) \\
& + \sum_{i=1}^p (-1)^{i+1} \rho(X_i) \omega(X_1, \dots, \widehat{X}_i, \dots, X_p, z) \\
& + \sum_{i=1}^{n-1} (-1)^{n+p-i+1} \rho(x_1^p, x_2^p, \dots, \hat{x}_i^p, \dots, x_{n-1}^p, z) \omega(X_1, \dots, X_{p-1}, x_i^p) \\
& + \sum_{1 \leq i < j} (-1)^i \omega(X_1, \dots, \widehat{X}_i, \dots, X_{j-1}, X_i \circ X_j, X_{j+1}, \dots, X_p, z) \tag{17}
\end{aligned}$$

for all  $X_i = (x_i^1, x_i^2, \dots, x_i^{n-1}) \in \wedge^{n-1} \mathfrak{g}$  and  $z \in \mathfrak{g}$  such that  $\delta \circ \delta = 0$ . Thus, we obtain the cohomology group  $H^\bullet(\mathfrak{g}, V) = Z^\bullet(\mathfrak{g}, V) / B^\bullet(\mathfrak{g}, V)$  where  $Z^\bullet(\mathfrak{g}, V)$  is the space of cocycles and  $B^\bullet(\mathfrak{g}, V)$  is the space of coboundaries.

*Proof.* For the convenience of the reader, we provide a detailed proof for the case of  $n$ -Lie algebras since the formula (17) in Theorem 2.6 is not explicitly given in [4, 6]. Now, to define the cochain complex for an  $n$ -Lie algebra  $\mathfrak{g}$  with coefficients in  $V$ , we consider a subcomplex of the Leibniz algebra  $\wedge^{n-1} \mathfrak{g}$  with coefficients in  $\text{Hom}(\mathfrak{g}, V)$ :

$$\begin{aligned}
C^{p+1}(\mathfrak{g}, V) & \triangleq \text{Hom}((\otimes^p(\wedge^{n-1} \mathfrak{g}) \otimes \mathfrak{g}, V) \cong \text{Hom}(\otimes^p(\wedge^{n-1} \mathfrak{g}), \text{Hom}(\mathfrak{g}, V))) \\
& = \text{Hom}(\otimes^p L, \text{Hom}(\mathfrak{g}, V)) = CL^p(L, \text{Hom}(\mathfrak{g}, V)).
\end{aligned}$$

By the definition of cohomology of Leibniz algebras, we get

$$d_{p-1} : CL^{p-1}(L, \text{Hom}(\mathfrak{g}, V)) \rightarrow CL^p(L, \text{Hom}(\mathfrak{g}, V)),$$

$$\begin{aligned}
& \delta_p \omega(X_1, X_2, \dots, X_p, z) = d_{p-1} \omega(X_1, X_2, \dots, X_p)(z) \\
= & \sum_{i=1}^{p-1} (-1)^{i+1} [X_i, \omega(X_1, \dots, \widehat{X}_i, \dots, X_p)]_L(z) + (-1)^p [\omega(X_1, \dots, X_{p-1}), X_p]_R(z) \\
& + \sum_{1 \leq i < j \leq p} (-1)^i \omega(X_1, \dots, \widehat{X}_i, \dots, X_{j-1}, X_i \circ X_j, X_{j+1}, \dots, X_p)(z) \\
= & \sum_{i=1}^{p-1} (-1)^{i+1} \{\rho(X_i) \omega(X_1, \dots, \widehat{X}_i, \dots, X_p, z) - \omega(X_1, \dots, \widehat{X}_i, \dots, X_p, [X_i, z])\} \\
& + (-1)^p \omega(X_1, \dots, X_{p-1}, [X_p, z]) \\
& - (-1)^p \sum_{i=1}^n (-1)^{n-i} \rho(x_1^p, x_2^p, \dots, \hat{x}_i^p, \dots, x_{n-1}^p, z) \omega(X_1, \dots, X_{p-1}, x_i^p) \\
& + \sum_{1 \leq i < j \leq p} (-1)^i \omega(X_1, \dots, \widehat{X}_i, \dots, X_{j-1}, X_i \circ X_j, X_{j+1}, \dots, X_p)(z)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^p (-1)^i \omega \left( X_1, \dots, \widehat{X}_i, \dots, X_p, [X_i, z] \right) \\
&\quad + \sum_{i=1}^p (-1)^{i+1} \rho(X_i) \omega \left( X_1, \dots, \widehat{X}_i, \dots, X_p, z \right) \\
&\quad + \sum_{i=1}^{n-1} (-1)^{n+p-i+1} \rho(x_1^p, x_2^p, \dots, \widehat{x}_i^p, \dots, x_{n-1}^p, z) \omega(X_1, \dots, X_{p-1}, x_i^p) \\
&\quad + \sum_{1 \leq i < j} (-1)^i \omega \left( X_1, \dots, \widehat{X}_i, \dots, X_{j-1}, X_i \circ X_j, X_{j+1}, \dots, X_p, z \right)
\end{aligned}$$

for all  $X_i = (x_1^i, x_2^i, \dots, x_{n-1}^i) \in L = \wedge^{n-1} \mathfrak{g}$ ,  $z \in \mathfrak{g}$ . In other words, the cohomology of an  $n$ -Lie algebra  $\mathfrak{g}$  with coefficients in  $V$  is defined to be the cohomology of Leibniz algebra  $L$  with coefficients in  $\text{Hom}(\mathfrak{g}, V)$ . Thus, we obtain that the cohomology group  $H^\bullet(\mathfrak{g}, V) = Z^\bullet(\mathfrak{g}, V)/B^\bullet(\mathfrak{g}, V)$  is well defined.  $\square$

**Remark 2.7.** L. Takhtajan in [31, 8], P. Gautheron in [10], M. Rotkiewicz in [28], and the authors in [9] have previously studied the cohomology theory for  $n$ -Lie algebras, each from a different perspective. However, we believe that the approach presented in Theorem 2.6 is the most direct and elegant. Specifically, for 3-Lie algebras, this method has been thoroughly examined in [33, 34].

According to the above definition, a 1-cochain is a map  $\nu \in \text{Hom}(\mathfrak{g}, V)$ , a 2-cochain is a map  $\omega \in \text{Hom}(\wedge^n \mathfrak{g}, V) \subseteq \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(\mathfrak{g}, V))$ , and the coboundary operator is given by

$$\delta_1 \nu(X_1, z) = d_0 \nu(X_1)(z) = -[\nu, X_1]_R(z), \quad (18)$$

$$\delta_2 \omega(X_1, X_2, z) = [X_1, \omega(X_2)]_L(z) + [\omega(X_1), X_2]_R(z) - \omega(X_1 \circ X_2)(z). \quad (19)$$

Put  $X_1 = (x_1, \dots, x_{n-1}) \in L$ ,  $z = x_n \in \mathfrak{g}$  in the equality (18), then by (15) we get

$$\delta_1 \nu(x_1, \dots, x_n) = \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n) \nu(x_i) - \nu([x_1, \dots, x_n]). \quad (20)$$

**Definition 2.8.** Let  $\mathfrak{g}$  be an  $n$ -Lie algebra and  $(V, \rho)$  be an  $\mathfrak{g}$ -module. Then a map  $\nu \in \text{Hom}(\mathfrak{g}, V)$  is called 1-cocycle if and only if

$$\nu([x_1, \dots, x_n]) = \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n) \nu(x_i), \quad (21)$$

and a map  $\omega : \wedge^n \mathfrak{g} \rightarrow V$  is called a 2-coboundary if there exists a map  $\nu \in \text{Hom}(\mathfrak{g}, V)$  such that  $\omega = \delta_1 \nu$ .

Put  $X_1 = (x_1, \dots, x_{n-1}) \in L$ ,  $X_2 = (y_1, \dots, y_{n-1}) \in L$ ,  $z = y_n \in \mathfrak{g}$  in the equality (19), then we get

$$\begin{aligned}
&\delta_2 \omega(x_1, \dots, x_{n-1}, y_1, \dots, y_{n-1}, y_n) \\
&= [x_1, \dots, x_{n-1}, \omega(y_1, \dots, y_{n-1})]_L(y_n) + [\omega(x_1, \dots, x_{n-1}), y_1, \dots, y_{n-1}]_R(y_n) \\
&\quad - \omega((x_1, \dots, x_{n-1}) \circ (y_1, \dots, y_{n-1}))(y_n).
\end{aligned}$$



We compute the right-hand side of the above equation as follows:

$$\begin{aligned}
& [x_1, \dots, x_{n-1}, \omega(y_1, \dots, y_{n-1})]_L(y_n) \\
= & \rho(x_1, \dots, x_{n-1})\omega(y_1, \dots, y_{n-1}, y_n) - \omega(y_1, \dots, y_{n-1}, [x_1, \dots, x_{n-1}, y_n]), \\
& [\omega(x_1, \dots, x_{n-1}), y_1, \dots, y_{n-1}]_R(y_n) \\
= & \omega(x_1, \dots, x_{n-1}, [y_1, \dots, y_n]) \\
& - \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n) \omega(x_1, \dots, x_{n-1}, y_i), \\
& \omega((x_1, \dots, x_{n-1}) \circ (y_1, \dots, y_{n-1}))(y_n) \\
= & \sum_{i=1}^{n-1} \omega(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1}, y_n).
\end{aligned}$$

Thus, we obtain the following Definition 2.9.

**Definition 2.9.** Let  $\mathfrak{g}$  be an  $n$ -Lie algebra and  $(V, \rho)$  be a representation. Then a map  $\omega : \wedge^n \mathfrak{g} \rightarrow V$  is called a 2-cocycle if  $\forall x_i, y_i \in \mathfrak{g}$ ,

$$\begin{aligned}
& \omega(x_1, \dots, x_{n-1}, [y_1, \dots, y_n]) + \rho(x_1, \dots, x_{n-1})\omega(y_1, \dots, y_n) \\
= & \sum_{i=1}^n \omega(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n) \\
& + \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n) \omega(x_1, \dots, x_{n-1}, y_i). \tag{22}
\end{aligned}$$

### 3 $n$ -Lie algebras in $\mathcal{LM}$

In this section, we will investigate the representation and cohomology theory of  $n$ -Lie algebras in  $\mathcal{LM}$ . We will begin by introducing the concept of  $n$ -Lie algebras in  $\mathcal{LM}$ , which recovers the concept of Lie algebras in  $\mathcal{LM}$  studied by Loday and Pirashvili in [23]. Furthermore, detailed information on low-dimensional cohomologies will be provided in this section.

**Definition 3.1.** An  $n$ -Lie algebra in  $\mathcal{LM}$  consists of a linear map  $f : M \rightarrow \mathfrak{g}$ , where  $\mathfrak{g}$  is an  $n$ -Lie algebra,  $(M, \rho)$  is a representation of  $\mathfrak{g}$ , and  $f$  is an equivariant map satisfying the following condition:

$$f(\rho(x_1, \dots, x_{n-1})(m)) = [x_1, \dots, x_{n-1}, f(m)], \tag{23}$$

for all  $x \in \mathfrak{g}$  and  $m \in M$ . We denote an  $n$ -Lie algebra in  $\mathcal{LM}$  by  $(M, \mathfrak{g}, f)$ .

When  $f$  is a surjective map, then the above equation (23) is equivalent to

$$f(\rho(f(m_1), \dots, f(m_{n-1}))(m_n)) = [f(m_1), \dots, f(m_n)]_{\mathfrak{g}}, \tag{24}$$

for all  $m_i \in M$ . This is called an embedding tensor of  $n$ -Lie algebras.

For any Leibniz  $n$ -algebra  $\mathfrak{h}$ , the quotient map  $\pi : \mathfrak{h} \rightarrow \mathfrak{h}_{\text{Lie}}$  is an  $n$ -Lie algebra in  $\mathcal{LM}$ , where  $\mathfrak{h}_{\text{Lie}}$  is  $\mathfrak{h}$  mod its Leibniz kernel

$$K = [\mathfrak{h}, \dots, \mathfrak{h}] = \text{span}\{[x_1, x_2, \dots, x_n] \mid \exists x_i, x_j \in \mathfrak{h}, x_i = x_j\}.$$

Conversely, given an  $n$ -Lie algebra  $(M, \mathfrak{g}, f)$  in  $\mathcal{LM}$ , a Leibniz  $n$ -algebra can be obtained on  $M$  using the following approach.

**Theorem 3.2.** For any  $n$ -Lie algebra  $(M, \mathfrak{g}, f)$  in  $\mathcal{LM}$ , we obtain a Leibniz  $n$ -algebra on  $M$  with the  $n$ -bracket given by:

$$[m_1, \dots, m_n] \triangleq \rho(f(m_1), \dots, f(m_{n-1}))(m_n). \quad (25)$$

*Proof.* First, we prove that  $f$  is an algebraic homomorphism from  $M$  to  $\mathfrak{g}$ . In fact, by equation (23) we get

$$f([m_1, \dots, m_n]) = f([f(m_1), \dots, f(m_{n-1}), m_n]) = [f(m_1), \dots, f(m_n)].$$

Next, we verify that the  $n$ -bracket defined by (25) on  $M$  satisfying the fundamental identity:

$$[m_1, \dots, m_{n-1}, [p_1, \dots, p_n]] = \sum_{i=1}^n [p_1, \dots, [m_1, \dots, m_{n-1}, p_i], \dots, p_n].$$

The left-hand side of the above equation is equal to

$$\begin{aligned} & [m_1, \dots, m_{n-1}, [p_1, \dots, p_{n-1}, p_n]] \\ &= [f(m_1), \dots, f(m_{n-1}), [f(p_1), \dots, f(p_{n-1}), p_n]] \\ &= \rho(f(m_1), \dots, f(m_{n-1}))\rho(f(p_1), \dots, f(p_{n-1}))(p_n), \end{aligned}$$

and the right-hand side is equal to

$$\begin{aligned} & \sum_{i=1}^n [p_1, \dots, [m_1, \dots, m_{n-1}, p_i], \dots, p_n] \\ &= \sum_{i=1}^n [f(p_1), [f(m_1), \dots, f(m_{n-1}), f(p_i)], \dots, f(p_{n-1}), p_n] \\ &= \sum_{i=1}^{n-1} \rho(f(p_1), [f(m_1), \dots, f(m_{n-1}), f(p_i)], \dots, f(p_{n-1}))(p_n) \\ & \quad + \rho(f(p_1), \dots, f(p_{n-1}))\rho(f(m_1), \dots, f(m_{n-1}))(p_n) \\ &= \sum_{i=1}^{n-1} \rho((f(m_1), \dots, f(m_{n-1})) \circ (f(p_1), \dots, f(p_{n-1})))(p_n) \\ & \quad + \rho(f(p_1), \dots, f(p_{n-1}))\rho(f(m_1), \dots, f(m_{n-1}))(p_n), \end{aligned}$$

Thus, the two sides are equal to each other because  $(M, \rho)$  is a representation of  $\mathfrak{g}$ . The proof is completed.  $\square$

Let  $(M, \mathfrak{g}, f)$  and  $(M', \mathfrak{g}', f')$  be two  $n$ -Lie algebras in  $\mathcal{LM}$ . A homomorphism between them is  $\phi = (\phi_0, \phi_1)$  where  $\phi_0 : \mathfrak{g} \rightarrow \mathfrak{g}'$  is an  $n$ -Lie algebra homomorphism and  $\phi_1 : M \rightarrow M'$  is an equivariant map such that

$$f' \circ \phi_1 = \phi_0 \circ f, \quad (26)$$

$$\phi_0([x_1, \dots, x_n]) = [\phi_0(x_1), \dots, \phi_0(x_n)]', \quad (27)$$

$$\phi_1(\rho(x_1, \dots, x_{n-1})(m)) = \rho'(\phi_0(x_1), \dots, \phi_0(x_{n-1}))(\phi_1(m)). \quad (28)$$

**Proposition 3.3.** Let  $\mathfrak{g}$  be an  $n$ -Lie algebra and  $(M, \rho)$  be a representation. Then  $f : M \rightarrow \mathfrak{g}$  is an  $n$ -Lie algebra in  $\mathcal{LM}$  if and only if the map  $\text{id} + f : \mathfrak{g} \rtimes M \rightarrow \mathfrak{g} \rtimes \mathfrak{g}$  is a homomorphism of  $n$ -Lie algebras.

*Proof.* We will verify when the map  $\text{id} + f : \mathfrak{g} \times M \rightarrow \mathfrak{g} \times \mathfrak{g}$  is a homomorphism of  $n$ -Lie algebras:

$$(\text{id} + f)([x_1 + m_1, \dots, x_n + m_n]) = [(\text{id} + f)(x_1 + m_1), \dots, (\text{id} + f)(x_n + m_n)]. \quad (29)$$

On one hand,

$$\begin{aligned} & (\text{id} + f)[x_1 + m_1, \dots, x_n + m_n] \\ &= (\text{id} + f)\left([x_1, \dots, x_n] + \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n)(m_i)\right) \\ &= [x_1, \dots, x_n] + \sum_{i=1}^n (-1)^{n-i} f(\rho(x_1, \dots, \widehat{x}_i, \dots, x_n)(m_i)). \end{aligned}$$

On the other hand,

$$\begin{aligned} & [(\text{id} + f)(x_1 + m_1), \dots, (\text{id} + f)(x_n + m_n)] \\ &= [x_1 + f(m_1), \dots, x_n + f(m_n)] \\ &= [x_1, \dots, x_n] + \sum_{i=1}^n (-1)^{n-i} [x_1, \dots, \widehat{x}_i, \dots, x_n, f(m_i)]. \end{aligned}$$

Thus, the two sides of above equation (29) are equal to each other if and only if

$$f(\rho(x_1, \dots, \widehat{x}_i, \dots, x_n)(m_i)) = [x_1, \dots, \widehat{x}_i, \dots, x_n, f(m_i)]$$

holds, which is the condition (23). Therefore  $(\text{id} + f)$  is a homomorphism of  $n$ -Lie algebras if and only if the conditions (23) hold. This completes the proof.  $\square$

**Example 3.4.** Consider the map  $\pi : \mathfrak{g} \times_H V \rightarrow \mathfrak{g}$  where  $\pi$  is the natural projection, and the action of  $\mathfrak{g}$  on  $\mathfrak{g} \times_H V$  given by

$$\rho(x_1, \dots, x_{n-1})(x + v) \triangleq [x_1, \dots, x_{n-1}, x] + \rho(x_1, \dots, x_{n-1})(v).$$

Then by Theorem 3.2,  $\mathfrak{g} \times_H V$  becomes a Leibniz  $n$ -algebra under the  $n$ -bracket

$$\begin{aligned} & [x_1 + u_1, \dots, x_n + u_n]_\pi \\ &= \rho(\pi(x_1 + u_1), \dots, \pi(x_{n-1} + u_{n-1}))(x_n + u_n) \\ &= [x_1, \dots, x_n] + \rho(x_1, \dots, x_{n-1})(u_n), \end{aligned}$$

which is exactly the  $n$ -bracket (11) in the hemisemidirect product  $\mathfrak{g} \times_H V$ .

**Example 3.5.** For an  $n$ -Lie algebra  $\mathfrak{g}$ , we define  $f = [\cdot, \dots, \cdot] : \wedge^n \mathfrak{g} \rightarrow \mathfrak{g}$  and the representation of  $\mathfrak{g}$  on  $\wedge^n \mathfrak{g}$  given by

$$\rho(x_1, \dots, x_{n-1})(y_1, \dots, y_n) \triangleq \sum_{i=1}^n (y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n)$$

for all  $x_i \in \mathfrak{g}, (y_1, \dots, y_n) \in \wedge^n \mathfrak{g}$ . In this case, by Theorem 3.2,  $\wedge^n \mathfrak{g}$  becomes a Leibniz  $n$ -algebra under the bracket

$$\begin{aligned} [Y_1, \dots, Y_n] &= \rho(f(Y_1), \dots, f(Y_{n-1}))(Y_n) \\ &= \sum_{i=1}^n (y_1, \dots, [[y_1^1, \dots, y_n^1], [y_1^{n-1}, \dots, y_n^{n-1}], y_i], \dots, y_n) \end{aligned}$$

for all  $Y_i = (y_1^i, \dots, y_n^i) \in \wedge^n \mathfrak{g}$ .

### 3.1 Representations

**Definition 3.6.** Let  $(M, \mathfrak{g}, f)$  be an  $n$ -Lie algebra in  $\mathcal{LM}$ . A representation  $\rho = (\rho_1, \rho_2, \rho_3)$  of  $(M, \mathfrak{g}, f)$  is an object  $(V, W, \varphi)$  in  $\mathcal{LM}$  such that the following conditions are satisfied:

- (i)  $(V, \rho_1)$  and  $(W, \rho_2)$  are representations of  $n$ -Lie algebra  $\mathfrak{g}$ ;
- (ii)  $\varphi$  is an equivariant map with respect to  $\rho_1$  and  $\rho_2$ :

$$\varphi \circ \rho_1(x_1, \dots, x_{n-1})(v) = \rho_2(x_1, \dots, x_{n-1}) \circ \varphi(v); \quad (30)$$

- (iii) there exists a map  $\rho_3 : \underbrace{\mathfrak{g} \otimes \dots \otimes \mathfrak{g}}_{n-2} \otimes M \longrightarrow \text{Hom}(W, V)$  such that:

$$\varphi \circ \rho_3(x_1, \dots, x_{n-2}, m)(w) = \rho_2(x_1, \dots, x_{n-2}, f(m))(w), \quad (31)$$

where  $x_1, \dots, x_{n-1} \in \mathfrak{g}$ ,  $m \in M$ ,  $v \in V$  and  $w \in W$ ;

- (iv) the following compatibility conditions hold:

$$\begin{aligned} & \rho_1(x_1, \dots, x_{n-1})\rho_3(y_1, \dots, y_{n-2}, m)(w) \\ = & \rho_3(y_1, \dots, y_{n-2}, \rho(x_1, \dots, x_{n-1})(m))(w) \\ & + \rho_3(y_1, \dots, y_{n-2}, m)\rho_2(x_1, \dots, x_{n-1})(w) \\ & + \sum_{i=1}^{n-2} \rho_3(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-2}, m)(w), \end{aligned} \quad (32)$$

$$\begin{aligned} & \rho_3(x_1, \dots, x_{n-2}, \rho(y_1, \dots, y_{n-1})(m))(w) \\ = & \rho_1(y_1, \dots, y_{n-1})\rho_3(x_1, \dots, x_{n-2}, m)(w) \\ & + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(y_1, \dots, \widehat{y}_i, \dots, y_{n-1}, m)\rho_2(x_1, \dots, x_{n-2}, y_i)(w), \end{aligned} \quad (33)$$

$$\begin{aligned} & \rho_3(x_1, \dots, x_{n-3}, [y_1, \dots, y_n], m)(w) \\ = & \sum_{i=1}^n (-1)^{n-i} \rho_1(y_1, \dots, \widehat{y}_i, \dots, y_n)\rho_3(x_1, \dots, x_{n-3}, y_i, m)(w), \end{aligned} \quad (34)$$

$$\begin{aligned} & \rho_3(x_1, \dots, x_{n-2}, m)\rho(y_1, \dots, y_{n-1})(w) \\ = & \rho_1(y_1, \dots, y_{n-1})\rho_3(x_1, \dots, x_{n-2}, m)(w) \\ & + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(y_1, \dots, \widehat{y}_i, \dots, y_{n-1}, \rho(x_1, \dots, x_{n-2}, y_i)(m))(w). \end{aligned} \quad (35)$$

From the map  $\rho_3$ , we define the  $n$ -linear map  $\{\cdot, \dots, \cdot\} : \underbrace{\mathfrak{g} \otimes \dots \otimes \mathfrak{g}}_{n-2} \otimes M \otimes W \rightarrow V$  by

$$\{x_1, \dots, x_{n-2}, m, w\} \triangleq \rho_3(x_1, \dots, x_{n-2}, m)(w). \quad (36)$$

Then the above equation (31) is equivalent to

$$\varphi(\{x_1, \dots, x_{n-2}, m, w\}) = \rho_2(x_1, \dots, x_{n-2}, f(m))(w). \quad (37)$$

The above equations (30) and (31) imply that the following two diagrams commute:

$$\begin{array}{ccc}
\mathfrak{g} \otimes \cdots \otimes \mathfrak{g} \otimes V & \xrightarrow{\rho_1} & V \\
\downarrow 1 \otimes \cdots \otimes 1 \otimes \varphi & & \downarrow \varphi \\
\mathfrak{g} \otimes \cdots \otimes \mathfrak{g} \otimes W & \xrightarrow{\rho_2} & W
\end{array}
\quad \text{and} \quad
\begin{array}{ccc}
\mathfrak{g} \otimes \cdots \otimes M \otimes W & \xrightarrow{\{\cdot, \dots, \cdot\}} & V \\
\downarrow 1_{\mathfrak{g}} \otimes \cdots \otimes f \otimes 1_W & & \downarrow \varphi \\
\mathfrak{g} \otimes \cdots \otimes \mathfrak{g} \otimes W & \xrightarrow{\rho_2} & W
\end{array}
\quad (38)$$

For  $(V, W, \varphi) = (M, \mathfrak{g}, f)$ , we get the adjoint representations of  $n$ -Lie algebras  $(M, \mathfrak{g}, f)$  on itself. This can be seen as following:

(i)  $M$  and  $\mathfrak{g}$  are representations of  $\mathfrak{g}$  given by  $\rho_1(x_1, \dots, x_{n-1})(y) = \text{ad}(x_1, \dots, x_{n-1})(y)$  and  $\rho_2(x_1, \dots, x_{n-1})(m) = \rho(x_1, \dots, x_{n-1})(m)$ ;

(ii)  $\varphi = f$  is an equivariant map:

$$f([x_1, \dots, x_{n-1}, m]) = [x_1, \dots, x_{n-1}, f(m)];$$

(iii) there exists a bilinear map  $\{\cdot, \dots, \cdot\} : \mathfrak{g} \otimes \cdots \otimes M \otimes W \rightarrow V$  defined by

$$\{x_1, \dots, x_{n-2}, m, x_{n-1}\} \triangleq -\rho(x_1, \dots, x_{n-1}, m)$$

such that

$$\begin{aligned}
f(\{x_1, \dots, x_{n-2}, m, x_{n-1}\}) &= -f(\rho(x_1, \dots, x_{n-1}, m)) \\
&= -[x_1, \dots, x_{n-1}, f(m)] \\
&= -[x_1, \dots, f(m), x_{n-1}] \\
&= \rho_2(x_1, \dots, x_{n-2}, f(m))(x_{n-1}).
\end{aligned}$$

(iv) the compatibility conditions (32)–(35) are satisfied by direct computations.

We construct semidirect products of an  $n$ -Lie algebra structure in  $\mathcal{LM}$  using its representations.

**Proposition 3.7.** *Given a representation of the  $n$ -Lie algebra  $(M, \mathfrak{g}, f)$  on  $(V, W, \varphi)$ . Define on  $(M \oplus V, \mathfrak{g} \oplus W, \widehat{f})$  an  $n$ -Lie algebra in  $\mathcal{LM}$  by*

$$\left\{ \begin{array}{l}
\widehat{f}(m + v) \triangleq f(m) + \varphi(v), \\
[x_1 + w_1, \dots, x_n + w_n] \triangleq [x_1, \dots, x_n] + \sum_{i=1}^n (-1)^{n-i} \rho_2(x_1, \dots, \widehat{x}_i, \dots, x_n)(w_i), \\
\widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})(m + v) \triangleq \rho(x_1, \dots, x_{n-1})(m) + \rho_1(x_1, \dots, x_{n-1})(v) \\
\quad + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, m)(w_i).
\end{array} \right. \quad (39)$$

Then  $(M \oplus V, \mathfrak{g} \oplus W, \widehat{f})$  is an  $n$ -Lie algebra in  $\mathcal{LM}$ , which is called the semidirect product of  $(M, \mathfrak{g}, f)$  and  $(V, W, \varphi)$ .

*Proof.* First, we verify that  $\widehat{f}$  is an equivariant map:

$$\widehat{f}(\widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})(m + v)) = [x_1 + w_1, \dots, x_{n-1} + w_{n-1}, \widehat{f}(m + v)]. \quad (40)$$

The left-hand side of (40) is

$$\begin{aligned}
&\widehat{f}(\widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})(m + v)) \\
&= \widehat{f}\rho(x_1, \dots, x_{n-1})(m) + \widehat{f}\rho_1(x_1, \dots, x_{n-1})(v)
\end{aligned}$$

$$\begin{aligned}
& -\widehat{f}\rho_3(w_1, \dots, w_{n-1})(m) \\
= & f(\rho(x_1, \dots, x_{n-1})(m)) + \varphi \circ \rho_1(x_1, \dots, x_{n-1})(v) \\
& + \sum_{i=1}^{n-1} (-1)^{n-i} \varphi \circ \rho_3(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, m)(w_i),
\end{aligned}$$

and the right-hand side of (40) is

$$\begin{aligned}
& [[x_1 + w_1, \dots, x_{n-1} + w_{n-1}, \widehat{f}(m + v)]] \\
= & [[x_1 + w_1, \dots, x_{n-1} + w_{n-1}, f(m) + \varphi(v)]] \\
= & [x_1, \dots, x_{n-1}, f(m)] + \rho_2(x_1, \dots, x_{n-1})(\varphi(v)) \\
& + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_2(x_1, \dots, \widehat{x}_i, \dots, f(m))(w_i).
\end{aligned}$$

Thus, the two sides of (40) are equal to each other by (31).

Next, we prove that  $\widehat{\rho}$  is a representation of  $\mathfrak{g} \oplus W$  on  $M \oplus V$  satisfying (R1):

$$\begin{aligned}
& \widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})\widehat{\rho}(y_1 + w'_1, \dots, y_{n-1} + w'_{n-1})(m + v) \\
& - \widehat{\rho}(y_1 + w'_1, \dots, y_{n-1} + w'_{n-1})\widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})(m + v) \\
= & \sum_{i=1}^{n-1} \widehat{\rho}(y_1 + w'_1, \dots, [x_1 + w_1, \dots, x_{n-1} + w_{n-1}, y_i + w'_i], \dots, y_{n-1} + w'_{n-1})(m + v)
\end{aligned} \tag{41}$$

The left-hand side of (41) is

$$\begin{aligned}
& \widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})\widehat{\rho}(y_1 + w'_1, \dots, y_{n-1} + w'_{n-1})(m + v) \\
& - \widehat{\rho}(y_1 + w'_1, \dots, y_{n-1} + w'_{n-1})\widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})(m + v) \\
= & \widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})\left(\rho(y_1, \dots, y_{n-1})(m) + \rho_1(y_1, \dots, y_{n-1})(v)\right) \\
& + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(y_1, \dots, \widehat{y}_i, \dots, y_{n-1}, m)(w'_i) - (x \leftrightarrow y, w_i \leftrightarrow w'_i) \\
= & [\rho(x_1, \dots, x_{n-1}), \rho(y_1, \dots, y_{n-1})](m) + [\rho_1(x_1, \dots, x_{n-1}), \rho_1(y_1, \dots, y_{n-1})](v) \\
& + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_1(x_1, \dots, x_{n-1})\rho_3(y_1, \dots, \widehat{y}_i, \dots, y_{n-1}, m)(w'_i) \\
& + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_1(y_1, \dots, y_{n-1})\rho_3(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, m)(w_i) \\
& + \sum_{j=1}^{n-1} (-1)^{n-j} \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(x_1, \dots, \widehat{x}_j, \dots, x_{n-1}, \rho(y_1, \dots, y_{n-1})(m))(w_j) \\
& + \sum_{j=1}^{n-1} (-1)^{n-j} \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(y_1, \dots, \widehat{y}_j, \dots, y_{n-1}, \rho(x_1, \dots, x_{n-1})(m))(w'_j),
\end{aligned}$$

and the right-hand side of (41) is

$$\sum_{i=1}^{n-1} \widehat{\rho}(y_1 + w'_1, \dots, [x_1 + w_1, \dots, x_{n-1} + w_{n-1}, y_i + w'_i], \dots, y_{n-1} + w'_{n-1})(m + v)$$

$$\begin{aligned}
&= \sum_{i=1}^{n-1} \widehat{\rho} \left( y_1 + w'_1, \dots, [x_1, \dots, x_{n-1}, y_i] + \rho_2(x_1, \dots, x_{n-1})(w'_i) \right. \\
&\quad \left. + \sum_{j=1}^{n-1} (-1)^{n-j} \rho_2(x_1, \dots, \widehat{x}_j, \dots, y_i)(w_j), \dots, y_{n-1} + w'_{n-1} \right) (m+v) \\
&= \sum_{i=1}^{n-1} \rho(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1})(m) \\
&\quad + \sum_{i=1}^{n-1} \rho_1(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1})(v) \\
&\quad + \sum_{i=1}^{n-1} \sum_{j=1}^{n-1} (-1)^{n-i} \rho_3(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, \widehat{y}_j, \dots, y_{n-1}, m)(w'_j) \\
&\quad + \sum_{i=1}^{n-1} \sum_{j=1}^{n-1} (-1)^{n-j} \rho_3(y_1, \dots, \widehat{y}_i, \dots, y_{n-1}, m) \\
&\quad \left( \rho_2(x_1, \dots, x_{n-1})(w'_i) + \sum_{j=1}^{n-1} (-1)^{n-j} \rho_2(x_1, \dots, \widehat{x}_j, \dots, y_i)(w_j) \right).
\end{aligned}$$

Thus, the two sides of (41) are equal to each other by  $\rho, \rho_1$  are representation of  $\mathfrak{g}$  on  $M, V$  and (32), (33) hold.

Finally, we prove that  $\widehat{\rho}$  is a representation of  $\mathfrak{g} \oplus W$  on  $M \oplus V$  satisfying (R2):

$$\begin{aligned}
&\widehat{\rho}(x_1 + w_1, \dots, x_{n-2} + w_{n-2}, [y_1 + w'_1, \dots, y_n + w'_n])(m+v) \\
&= \sum_{i=1}^n (-1)^{n-i} \widehat{\rho}(y_1 + w'_1, \dots, \widehat{y_i + w'_i}, \dots, y_n + w'_n) \\
&\quad \widehat{\rho}(x_1 + w_1, \dots, x_{n-2} + w_{n-2}, y_i + w'_i)(m+v). \tag{42}
\end{aligned}$$

The left-hand side of (42) is

$$\begin{aligned}
&\widehat{\rho}(x_1 + w_1, \dots, x_{n-2} + w_{n-2}, [y_1 + w'_1, \dots, y_n + w'_n])(m+v) \\
&= \widehat{\rho}(x_1 + w_1, \dots, x_{n-2} + w_{n-2}, [y_1, \dots, y_n]) \\
&\quad + \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n)(w'_i)(m+v) \\
&= \rho(x_1, \dots, x_{n-2}, [y_1, \dots, y_n])(m) + \rho(x_1, \dots, x_{n-2}, [y_1, \dots, y_n])(v) \\
&\quad + \sum_{i=1}^{n-2} (-1)^{n-i} \rho_3(x_1, \dots, \widehat{x}_i, \dots, x_{n-2}, [y_1, \dots, y_n], m)(w_i) \\
&\quad + \rho_3(x_1, \dots, x_{n-2}, m) \left( \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n)(w'_i) \right)
\end{aligned}$$

and the right-hand side of (42) is

$$\sum_{i=1}^n (-1)^{n-i} \widehat{\rho}(y_1 + w'_1, \dots, \widehat{y_i + w'_i}, \dots, y_n + w'_n)$$

$$\begin{aligned}
& \widehat{\rho}(x_1 + w_1, \dots, x_{n-2} + w_{n-2}, y_i + w'_i)(m + v) \\
= & \sum_{i=1}^n (-1)^{n-i} \widehat{\rho}(y_1 + w'_1, \dots, \widehat{y_i + w'_i}, \dots, y_n + w'_n) \\
& \left( \rho_1(x_1, \dots, x_{n-2}, y_i)(m) + \rho_1(x_1, \dots, x_{n-2}, y_i)(v) \right. \\
& \left. + \sum_{j=1}^{n-2} (-1)^{n-j} \rho_3(x_1, \dots, \widehat{x_j}, \dots, x_{n-2}, y_i, m)(w_j) + \rho_3(x_1, \dots, x_{n-2}, m)(w'_i) \right) \\
= & \sum_{i=1}^n (-1)^{n-i} \rho_1(y_1, \dots, \widehat{y_i}, \dots, y_n) \rho_1(x_1, \dots, x_{n-2}, y_i)(m) \\
& + \sum_{i=1}^n (-1)^{n-i} \rho_1(y_1, \dots, \widehat{y_i}, \dots, y_n) \rho_1(x_1, \dots, x_{n-2}, y_i)(v) \\
& + \sum_{i=1}^n (-1)^{n-i} \rho_1(y_1, \dots, \widehat{y_i}, \dots, y_n) \sum_{j=1}^{n-2} (-1)^{n-j} \rho_3(x_1, \dots, \widehat{x_j}, \dots, x_{n-2}, y_i, m)(w_j) \\
& + \sum_{i=1}^n (-1)^{n-i} \rho_1(y_1, \dots, \widehat{y_i}, \dots, y_n) \rho_3(x_1, \dots, x_{n-2}, m)(w'_i) \\
& + \sum_{i=1}^n (-1)^{n-i} \sum_{j=1}^{n-2} (-1)^{n-j} \rho_3(y_1, \dots, \widehat{y_j}, \dots, \widehat{y_i}, \dots, y_n, \rho_1(x_1, \dots, x_{n-2}, y_i)(m))(w'_j).
\end{aligned}$$

Thus, the two sides of (42) are equal to each other by  $\rho, \rho_1$  are representation of  $\mathfrak{g}$  on  $M, V$  and (34), (35) hold. This completes the proof.  $\square$

### 3.2 Cohomology

Let  $(M, \mathfrak{g}, f)$  be an  $n$ -Lie algebra in  $\mathcal{LM}$  and  $(V, W, \varphi)$  be a representation of  $(M, \mathfrak{g}, f)$ . We define the  $k$ -cochain  $C^k((M, \mathfrak{g}, f), (V, W, \varphi))$  to be the space

$$\text{Hom}(\wedge^{k(n-1)+1} \mathfrak{g}, W) \oplus \text{Hom}(\wedge^{k(n-1)} \mathfrak{g}, \text{Hom}(M, V)) \oplus \text{Hom}(\wedge^{(k-1)(n-1)} \mathfrak{g}, \text{Hom}(M, W))$$

which can be seen as the subspace of

$$\text{Hom}(\otimes^k L, \text{Hom}(\mathfrak{g}, W)) \oplus \text{Hom}(\otimes^k L, \text{Hom}(M, V)) \oplus \text{Hom}(\otimes^{k-1} L, \text{Hom}(M, W)),$$

and the coboundary map is given by

$$D(\omega, \nu, \theta) = (-\delta_1 \omega, -h_1 \omega + \delta_2 \nu, -h_2 \omega + \varphi_{\sharp} \nu - \delta_3 \theta) \quad (43)$$

where  $\delta_1, \delta_2, \delta_3$  are coboundary maps in  $n$ -Lie algebra cohomology of  $\mathfrak{g}$  with coefficient in  $W, \text{Hom}(M, V), \text{Hom}(M, W)$  respectively. The following maps:

$$\varphi_{\sharp} : \text{Hom}(\wedge^{k-1} L, \text{Hom}(M, V)) \rightarrow \text{Hom}(\wedge^{k-1} L, \text{Hom}(M, W)), \quad (44)$$

$$h_1 : \text{Hom}(\wedge^k L \otimes \mathfrak{g}, W) \rightarrow \text{Hom}(\wedge^{k+1} L, \text{Hom}(M, V)), \quad (45)$$

$$h_2 : \text{Hom}(\wedge^k L \otimes \mathfrak{g}, W) \rightarrow \text{Hom}(\wedge^k L, \text{Hom}(M, W)) \quad (46)$$

are defined by

$$(\varphi_{\sharp}(\nu)(X_1, \dots, X_{k-1}))(m) = \varphi(\nu(X_1, \dots, X_{k-1})(m)), \quad (47)$$



$$(h_1(\omega)(X_1, \dots, X_k))(m) = \rho_3(m)\omega(X_1, \dots, X_k), \quad (48)$$

$$(h_2(\omega)(X_1, \dots, X_{k-1}))(m) = \omega(f(m), X_1, \dots, X_{k-1}). \quad (49)$$

It can be verified by direct computations that  $D^2 = 0$ , thus we obtain a cochain complex  $C^k((M, \mathfrak{g}, f), (V, W, \varphi))$  whose cohomology group

$$H^k((M, \mathfrak{g}, f), (V, W, \varphi)) = Z^k((M, \mathfrak{g}, f), (V, W, \varphi))/B^k((M, \mathfrak{g}, f), (V, W, \varphi))$$

is defined as cohomology group of  $(M, \mathfrak{g}, f)$  with coefficients in  $(V, W, \varphi)$ .

The cochain complex is precisely given by

$$\begin{aligned} W &\xrightarrow{D_0} \\ \text{Hom}(\mathfrak{g}, W) \oplus \text{Hom}(M, V) &\xrightarrow{D_1} \\ \text{Hom}(\wedge^n \mathfrak{g}, W) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(M, V)) \oplus \text{Hom}(M, W) &\xrightarrow{D_2} \\ \text{Hom}(\wedge^{2n-1} \mathfrak{g}, W) \oplus \text{Hom}(\wedge^{2n-2} \mathfrak{g}, \text{Hom}(M, V)) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(M, W)) &\xrightarrow{D_3} \dots, \end{aligned} \quad (50)$$

where one writes the cochain complex in components as follows:

$$\begin{array}{ccccc} W & & & & \\ \downarrow -\delta_1 & \searrow & & & \\ \text{Hom}(\mathfrak{g}, W) \oplus & \text{Hom}(M, V) & & & \\ \downarrow -\delta_1 & \searrow -h_1 & \downarrow \delta_2 & \searrow -h_2 & \searrow \varphi_\# \\ \text{Hom}(\wedge^n \mathfrak{g}, W) \oplus & \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(M, V)) \oplus & & & \text{Hom}(M, W) \\ \downarrow -\delta_1 & \searrow -h_1 & \downarrow \delta_2 & \searrow -h_2 & \searrow \varphi_\# & \downarrow -\delta_3 \\ \text{Hom}(\wedge^{2n-1} \mathfrak{g}, W) \oplus & \text{Hom}(\wedge^{2n-2} \mathfrak{g}, \text{Hom}(M, V)) \oplus & & & \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(M, W)) \\ \downarrow -\delta_1 & \searrow -h_1 & \downarrow \delta_2 & \searrow -h_2 & \searrow \varphi_\# & \downarrow -\delta_3 \\ \text{Hom}(\wedge^{3n-2} \mathfrak{g}, W) \oplus & \text{Hom}(\wedge^{3n-3} \mathfrak{g}, \text{Hom}(M, V)) \oplus & & & \text{Hom}(\wedge^{2n-2} \mathfrak{g}, \text{Hom}(M, W)) \\ \vdots & & & & \end{array}$$

For 1-cochain  $(N_0, N_1) \in \text{Hom}(\mathfrak{g}, W) \oplus \text{Hom}(M, V)$ , the coboundary map is

$$D_1(N_0, N_1)(m) = (-h_2 N_0 + \varphi_\# N_1)(m) = \varphi \circ N_1(m) - N_0 \circ f(m),$$

$$\begin{aligned} D_1(N_0, N_1)(x_1, \dots, x_n) &= -\delta_1 N_0(x_1, \dots, x_n) \\ &= [N_0(x_1), \dots, x_n] + \dots + [x_1, \dots, N_0 x_n] - N_0[x_1, \dots, x_n], \end{aligned}$$

$$\begin{aligned} D_1(N_0, N_1)(x_1, \dots, x_{n-1}, m) &= (-h_1 N_0 + \delta_2 N_1)(x_1, x_2, m) \\ &= [N_0(x_1), \dots, x_{n-1}, m] + \dots + [x_1, \dots, N_0 x_{n-1}, m] \\ &\quad + [x_1, \dots, x_{n-1}, N_1 m] - N_1[x_1, \dots, x_{n-1}, m]. \end{aligned}$$

Thus, a 1-cocycle is  $(N_0, N_1) \in \text{Hom}(\mathfrak{g}, W) \oplus \text{Hom}(M, V)$ , such that

$$\varphi \circ N_1 = N_0 \circ f, \quad (51)$$

$$N_0[x_1, \dots, x_n] = \sum_{i=1}^n [x_1, \dots, N_0 x_i, \dots, x_n], \quad (52)$$

$$N_1[x_1, \dots, x_{n-1}, m] = \sum_{i=1}^{n-1} [x_1, \dots, N_0 x_i, \dots, x_{n-1}, m] + [x_1, \dots, x_{n-1}, N_1 m].$$

For a 2-cochain  $(\omega, \nu, \theta) \in \text{Hom}(\wedge^n \mathfrak{g}, W) \oplus \text{Hom}(\mathfrak{g}, \text{Hom}(M, V)) \oplus \text{Hom}(M, W)$ , we get that

$$D_2(\omega, \nu, \theta) \in \text{Hom}(\wedge^{2n-1} \mathfrak{g}, W) \oplus \text{Hom}(\wedge^{2n-2} \mathfrak{g}, \text{Hom}(M, V)) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(M, W))$$

where

$$\begin{aligned} & D_2(\omega, \nu, \theta)(x_1, \dots, x_{n-1}, y_1, \dots, y_n) \\ &= -\delta_1 \omega(x_1, \dots, x_{n-1}, y_1, \dots, y_n), \\ & D_2(\omega, \nu, \theta)(x_1, \dots, x_{n-1}, y_1, \dots, y_{n-1}, m) \\ &= (-h_1 \omega + \delta_2 \nu)(x_1, \dots, x_{n-1}, y_1, \dots, y_{n-1}, m), \\ & D_2(\omega, \nu, \theta)(x_1, \dots, x_{n-1}, m) \\ &= (-h_2 \omega + \varphi_{\#} \nu - \delta_3 \theta)(x_1, \dots, x_{n-1}, m). \end{aligned}$$

Thus, a 2-cocycle is  $(\omega, \nu, \theta) \in \text{Hom}(\wedge^n \mathfrak{g}, W) \oplus \text{Hom}(\wedge^{n-1} \mathfrak{g}, \text{Hom}(M, V)) \oplus \text{Hom}(M, W)$  such that the following conditions holds:

$$\begin{aligned} & \theta(\rho(x_1, \dots, x_{n-1})(m)) + \varphi(\nu(x_1, \dots, x_{n-1}, m)) \\ &= \omega(x_1, \dots, x_{n-1}, f(m)) + [x_1, \dots, x_{n-1}, \theta(m)], \end{aligned} \quad (53)$$

$$\begin{aligned} & \omega(x_1, \dots, x_{n-1}, [y_1, \dots, y_n]) + \rho_2(x_1, \dots, x_{n-1})\omega(y_1, \dots, y_n) \\ &= \sum_{i=1}^n \omega(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n) \\ & \quad + \sum_{i=1}^n (-1)^{n-i} \rho_2(y_1, \dots, \widehat{y}_i, \dots, y_n)\omega(x_1, \dots, x_{n-1}, y_i), \end{aligned} \quad (54)$$

$$\begin{aligned} & \nu(x_1, \dots, x_{n-1}, [y_1, \dots, y_{n-1}, m]) + \rho_1(x_1, \dots, x_{n-1})\nu(y_1, \dots, y_{n-1}, m) \\ &= \sum_{i=1}^{n-1} \nu(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1}, m) + \nu(y_1, \dots, y_{n-1}, [x_1, \dots, x_{n-1}, m]) \\ & \quad + \sum_{i=1}^{n-1} [y_1, \nu(x_1, \dots, x_{n-1}, y_i), \dots, y_{n-1}, m] + \rho_1(y_1, \dots, y_{n-1})\nu(x_1, \dots, x_{n-1}, m). \end{aligned} \quad (55)$$

## 4 Abelian extensions

**Definition 4.1.** Let  $(M, \mathfrak{g}, f)$  be an  $n$ -Lie algebra in  $\mathcal{LM}$ . An extension of  $(M, \mathfrak{g}, f)$  is a short exact sequence

$$\begin{array}{ccccccccc} 0 & \xrightarrow{0} & V & \xrightarrow{i_1} & \widehat{M} & \xrightarrow{p_1} & M & \xrightarrow{0} & 0 \\ 0 \downarrow & & \varphi \downarrow & & \widehat{f} \downarrow & & f \downarrow & & 0 \downarrow \\ 0 & \xrightarrow{0} & W & \xrightarrow{i_0} & \widehat{\mathfrak{g}} & \xrightarrow{p_0} & \mathfrak{g} & \xrightarrow{0} & 0 \end{array} \quad (56)$$

where  $(V, W, \varphi)$  is an  $n$ -Lie algebra in  $\mathcal{LM}$ .

We call  $(\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f})$  an extension of  $(M, \mathfrak{g}, f)$  by  $(V, W, \varphi)$ , and denote it by  $\widehat{E}$ . It is called an abelian extension if  $(V, W, \varphi)$  is an abelian  $n$ -Lie algebra in  $\mathcal{LM}$ , which means that  $\varphi : V \rightarrow W$  is a linear map between vector spaces  $V$  and  $W$  with zero bracket and trivial representation of  $V$  on  $W$ .

A splitting  $\sigma = (\sigma_0, \sigma_1) : (M, \mathfrak{g}, f) \rightarrow (\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f})$  consists of linear maps  $\sigma_0 : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$  and  $\sigma_1 : M \rightarrow \widehat{M}$  such that  $p_0 \circ \sigma_0 = \text{id}_{\mathfrak{g}}$  and  $p_1 \circ \sigma_1 = \text{id}_M$ .

Two extensions

$$\widehat{E} : 0 \longrightarrow (V, W, \varphi) \xrightarrow{i} (\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f}) \xrightarrow{p} (M, \mathfrak{g}, f) \longrightarrow 0$$

and

$$\widetilde{E} : 0 \longrightarrow (V, W, \varphi) \xrightarrow{j} (\widetilde{M}, \widetilde{\mathfrak{g}}, \widetilde{f}) \xrightarrow{q} (M, \mathfrak{g}, f) \longrightarrow 0$$

are equivalent, if there exists a homomorphism  $F : (\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f}) \rightarrow (\widetilde{M}, \widetilde{\mathfrak{g}}, \widetilde{f})$  such that  $F \circ i = j$ ,  $q \circ F = p$ .

Let  $(\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f})$  be an abelian extension of  $(M, \mathfrak{g}, f)$  by  $(V, W, \varphi)$  and  $\sigma : (M, \mathfrak{g}, f) \rightarrow (\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f})$  be a splitting. Define a representation of  $(M, \mathfrak{g}, f)$  over  $(V, W, \varphi)$  by

$$\begin{cases} \rho_1 : \wedge^{n-1} \mathfrak{g} \longrightarrow \mathfrak{gl}(V), \\ \rho_1(x_1, \dots, x_{n-1})(v) \triangleq \widehat{\rho}(\sigma_0(x_1), \dots, \sigma_0(x_{n-1}))(v), \\ \rho_2 : \wedge^{n-1} \mathfrak{g} \longrightarrow \mathfrak{gl}(W), \\ \rho_2(x_1, \dots, x_{n-1})(z) \triangleq [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), w]_{\widehat{\mathfrak{g}}}, \\ \rho_3 : \mathfrak{g} \otimes \dots \otimes \mathfrak{g} \otimes M \longrightarrow \text{Hom}(W, V), \\ \rho_3(x_1, \dots, x_{n-2}, m)(w) \triangleq -\widehat{\rho}(\sigma_0(x_1), \dots, \sigma_0(x_{n-2}), w)(\sigma_1(m)). \end{cases}$$

for all  $x_1, \dots, x_{n-1}, z \in \mathfrak{g}$ ,  $m \in M$ .

**Proposition 4.2.** *With the above notations,  $(\rho_1, \rho_2, \rho_3)$  is a representation of  $(M, \mathfrak{g}, f)$  on  $(V, W, \varphi)$ . Furthermore,  $(\rho_1, \rho_2, \rho_3)$  does not depend on the choice of the splitting  $\sigma$ . Moreover, equivalent abelian extensions give the same representation.*

*Proof.* Firstly, we show that  $\rho_1, \rho_2, \rho_3$  are well defined. Since  $\text{Ker } p_0 \cong W$ , then for  $w \in W$ , we have  $p_0(w) = 0$ . By the fact that  $(p_1, p_0)$  is a homomorphism between  $(\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f})$  and  $(M, \mathfrak{g}, f)$ , we get

$$\begin{aligned} p_0[\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), w]_{\widehat{\mathfrak{g}}} &= [p_0\sigma_0(x_1), \dots, p_0\sigma_0(x_{n-1}), p_0(w)]_{\widehat{\mathfrak{g}}} \\ &= [p_0\sigma_0(x_1), \dots, p_0\sigma_0(x_{n-1}), 0]_{\widehat{\mathfrak{g}}} = 0. \end{aligned}$$

Thus,  $\rho_2(x_1, \dots, x_{n-1})(w) = [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), w]_{\widehat{\mathfrak{g}}} \in \ker p_0 \cong W$ . Similar computations show that  $\rho_1(x_1, \dots, x_{n-1})(v) = \widehat{\rho}(\sigma_0(x_1), \dots, \sigma_0(x_{n-1}))(v)$ ,  $\rho_3(x_1, \dots, x_{n-2}, m)(w) = -\widehat{\rho}(\sigma_0(x_1), \dots, \sigma_0(x_{n-2}), w)(\sigma_1(m)) \in \text{Ker} p_0 = V$ .

Now we will show that  $\rho_i$  are independent of the choice of  $\sigma$ .

In fact, if we choose another section  $\sigma'_0 : \mathfrak{g} \rightarrow \widehat{\mathfrak{g}}$ , then

$$\begin{aligned} p_0(\sigma_0(x_i) - \sigma'_0(x_i)) &= x_i - x_i = 0 \\ \implies \sigma_0(x_i) - \sigma'_0(x_i) &\in \text{Ker} p_0 = W \\ \implies \sigma'_0(x_i) &= \sigma_0(x_i) + w_i \end{aligned}$$

for some  $w_i \in W$ .

Since we have  $[\dots, w, w']_{\widehat{\mathfrak{g}}} = 0$  for all  $w, w' \in W$ , which implies that

$$\begin{aligned} &[\sigma'_0(x_1), \dots, \sigma'_0(x_{n-1}), w]_{\widehat{\mathfrak{g}}} \\ &= [\sigma_0(x_1) + u_1, \dots, \sigma_0(x_{n-1}) + u_{n-1}, w]_{\widehat{\mathfrak{g}}} \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}) + u_{n-1}, w]_{\widehat{\mathfrak{g}}} + [u_1, \dots, \sigma_0(x_{n-1}) + u_{n-1}, w]_{\widehat{\mathfrak{g}}} \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), w]_{\widehat{\mathfrak{g}}} + \dots + [\sigma_0(x_1), \dots, u_{n-1}, w]_{\widehat{\mathfrak{g}}} \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), w]_{\widehat{\mathfrak{g}}}, \end{aligned}$$

thus,  $\rho_2$  is independent on the choice of  $\sigma$ . If we choose another splitting  $\sigma'_1 : M \rightarrow \widehat{M}$ , then  $p_1(\sigma_1(m) - \sigma'_1(m)) = m - m = 0$ , i.e.  $\sigma_1(m) - \sigma'_1(m) \in \text{Ker} p_1 = V$ . Similar argument shows that  $\rho_1, \rho_3$  are independent on the choice of  $\sigma$ . Thus  $\rho_1, \rho_2, \rho_3$  are well defined.

Secondly, we check that  $\rho = (\rho_1, \rho_2, \rho_3)$  is indeed a representation of  $(M, \mathfrak{g}, f)$  on  $(V, W, \varphi)$ . Since  $(V, W, \varphi)$  is an abelian  $n$ -Lie algebra in  $\mathcal{LM}$ , we have

$$\begin{aligned} &\rho_2(x_1, \dots, x_{n-1})\rho_2(y_1, \dots, y_{n-1})(w) \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), [\sigma_0(y_1), \dots, \sigma_0(y_{n-1}), w]]_{\widehat{\mathfrak{g}}} \\ &= [\sigma_0(y_1), \dots, [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), \sigma_0(y_i)] \dots, \sigma_0(y_{n-1}), w]_{\widehat{\mathfrak{g}}} \\ &\quad + [\sigma_0(y_1), \dots, \sigma_0(y_{n-1}), [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), w]]_{\widehat{\mathfrak{g}}} \\ &= \sum_{i=1}^{n-1} \rho_2(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1})(w) \\ &\quad + \rho_2(y_1, \dots, y_{n-1})\rho_2(x_1, \dots, x_{n-1})(w), \end{aligned}$$

which implies that  $\rho_2$  is a representation of  $\mathfrak{g}$  on  $W$ . Similarly, we get  $\rho_1$  is a representation of  $\mathfrak{g}$  on  $V$ . For the equivariant between  $\rho_1$  and  $\rho_2$ , we have

$$\begin{aligned} \varphi \circ \rho_1(x_1, \dots, x_{n-1})(v) &= \varphi \widehat{\rho}(\sigma_0(x_1), \dots, \sigma_0(x_{n-1}))(v) \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), \varphi(v)] \\ &= \rho_2(x_1, \dots, x_{n-1}) \circ \varphi(v). \end{aligned}$$

For  $\rho_3 : \mathfrak{g} \otimes \dots \otimes \mathfrak{g} \otimes M \rightarrow \text{Hom}(W, V)$ , we have

$$\begin{aligned} \varphi \circ \rho_3(x_1, \dots, x_{n-2}, m)(w) &= \varphi[\sigma_0(x_1), \dots, \sigma_0(x_{n-2}), w, \sigma_1(m)] \\ &= -[\sigma_0(x_1), \dots, \sigma_0(x_{n-2}), w, \widehat{f}\sigma_1(m)] \\ &= -[\sigma_0(x_1), \dots, \sigma_0(x_{n-2}), w, \sigma_0(f(m))] \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-2}), \sigma_0(f(m)), w] \end{aligned}$$

$$= \rho_2(x_1, \dots, x_{n-2}, f(m))(w).$$

Thus,  $(\rho_1, \rho_2, \rho_3)$  is a representation of  $(M, \mathfrak{g}, f)$  over  $(V, W, \varphi)$ .

Finally, suppose that  $\widehat{E}$  and  $\widetilde{E}$  are equivalent abelian extensions, and  $F : (\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f}) \longrightarrow (\widetilde{M}, \widetilde{\mathfrak{g}}, \widetilde{f})$  be the morphism. Choose linear sections  $\sigma$  and  $\sigma'$  of  $p$  and  $q$ . Then we have  $q_0 F_0 \sigma_0(x_i) = p_0 \sigma_0(x_i) = x_i = q_0 \sigma'_0(x_i)$ , thus  $F_0 \sigma_0(x_i) - \sigma'_0(x_i) \in \text{Ker} q_0 = W$ . Therefore, we obtain

$$\begin{aligned} [\sigma'_0(x_1), \dots, \sigma'_0(x_{n-1}), v + w]_{\widehat{\mathfrak{g}}} &= [F_0 \sigma_0(x_1), \dots, F_0 \sigma'_0(x_{n-1}), v + w]_{\widehat{\mathfrak{g}}} \\ &= F_0 [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), v + w]_{\widehat{\mathfrak{g}}} \\ &= [\sigma_0(x_1), \dots, \sigma_0(x_{n-1}), v + w]_{\widehat{\mathfrak{g}}}, \end{aligned}$$

which implies that equivalent abelian extensions give the same  $\rho_1$  and  $\rho_2$ . Similarly, we can show that equivalent abelian extensions also give the same  $\rho_3$ . Therefore, equivalent abelian extensions also give the same representation. The proof is finished.  $\square$

Let  $\sigma : (M, \mathfrak{g}, f) \rightarrow (\widehat{M}, \widehat{\mathfrak{g}}, \widehat{f})$  be a section of the abelian extension. Define the following linear maps:

$$\left\{ \begin{array}{l} \theta : M \longrightarrow W, \quad \theta(m) \triangleq \widehat{f} \sigma_1(m) - \sigma_0 f(m), \\ \omega : \wedge^n \mathfrak{g} \longrightarrow W, \quad \omega(x_1, \dots, x_n) \triangleq [\sigma_0(x_1), \dots, \sigma_0(x_n)]_{\widehat{\mathfrak{g}}} - \sigma_0([x_1, \dots, x_n]_{\mathfrak{g}}), \\ \nu : \wedge^{n-1} \mathfrak{g} \otimes M \rightarrow V, \\ \nu(x_1, \dots, x_{n-1}, m) \triangleq \widehat{\rho}(\sigma_0(x_1), \dots, \sigma_0(x_{n-1}))(\sigma_1(m)) - \sigma_1 \rho(x_1, \dots, x_{n-1})(m). \end{array} \right.$$

for all  $x_i \in \mathfrak{g}$ ,  $m \in M$ .

**Theorem 4.3.** *With the above notations,  $(\theta, \omega, \nu)$  is a 2-cocycle of  $(M, \mathfrak{g}, f)$  with coefficients in  $(V, W, \varphi)$ .*

*Proof.* By the equality

$$\widehat{f}(\widehat{\rho}(\sigma_0 x_1, \dots, \sigma_0 x_{n-1})(\sigma_1(m))) = [\sigma_0 x_1, \sigma_0 x_{n-1}, \widehat{f} \sigma_1(m)]_{\widehat{\mathfrak{g}}},$$

we obtain that

$$\begin{aligned} &\omega(x_1, \dots, x_{n-1}, f(m)) + \rho_1(x_1, \dots, x_{n-1}, \theta(m)) \\ &= \theta([x_1, \dots, x_{n-1}, m]) + \varphi(\nu(x_1, \dots, x_{n-1}, m)). \end{aligned} \tag{57}$$

By the equality

$$[\sigma_0 x_1, \dots, \sigma_0 x_{n-1}, [\sigma_0 y_1, \dots, \sigma_0 y_n]] = \sum_{i=1}^n [\sigma_0 y_1, \dots, [\sigma_0 x_1, \dots, \sigma_0 x_{n-1}, \sigma_0 y_i], \dots, \sigma_0 y_n],$$

we get that the left-hand side is equal to

$$\begin{aligned} &= [\sigma_0 x_1, \dots, \sigma_0 x_{n-1}, \omega(y_1, \dots, y_n) + \sigma_0([y_1, \dots, y_n])] \\ &= \rho(x_1, \dots, x_{n-1})\omega(y_1, \dots, y_n) + [\sigma_0 x_1, \dots, \sigma_0 x_{n-1}, \sigma_0([y_1, \dots, y_n])] \\ &= \rho(x_1, \dots, x_{n-1})\omega(y_1, \dots, y_n) + \omega(x_1, \dots, x_{n-1}, [y_1, \dots, y_n]) \\ &\quad + \sigma_0([x_1, \dots, x_{n-1}, [y_1, \dots, y_n]]), \end{aligned}$$

the right-hand side is equal to

$$\begin{aligned}
&= \sum_{i=1}^n (-1)^{n-i} [\sigma_0 y_1, \dots, \sigma_0 \widehat{y}_i, \dots, \sigma_0 y_n, [\sigma_0 x_1, \dots, \sigma_0 x_{n-1}, \sigma_0 y_i]] \\
&= \sum_{i=1}^n (-1)^{n-i} [\sigma_0 y_1, \dots, \sigma_0 \widehat{y}_i, \dots, \sigma_0 y_n, \omega(x_1, \dots, x_{n-1}, y_i) + \sigma_0([x_1, \dots, x_{n-1}, y_i])] \\
&= \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n) \omega(x_1, \dots, x_{n-1}, y_i) \\
&\quad + \omega\left(\sum_{i=1}^n [y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n]\right) \\
&\quad + \sigma_0\left(\sum_{i=1}^n [y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n]\right).
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
&\omega(x_1, \dots, x_{n-1}, [y_1, \dots, y_n]) + \rho(x_1, \dots, x_{n-1}) \omega(y_1, \dots, y_n) \\
&= \sum_{i=1}^n \omega([y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_n]) \\
&\quad + \sum_{i=1}^n (-1)^{n-i} \rho(y_1, \dots, \widehat{y}_i, \dots, y_n) \omega(x_1, \dots, x_{n-1}, y_i). \tag{58}
\end{aligned}$$

Finally, by the equality

$$\begin{aligned}
&\widehat{\rho}(\sigma_0 x_1, \dots, \sigma_0 x_{n-1}) \rho(\sigma_0 y_1, \dots, \sigma_0 y_{n-1})(\sigma_1 m) \\
&= \sum_{i=1}^{n-1} \rho(\sigma_0 y_1, \dots, [\sigma_0 x_1, \dots, \sigma_0 x_{n-1}, \sigma_0 y_i], \dots, \sigma_0 y_{n-1})(\sigma_1 m) \\
&\quad + \rho(\sigma_0 y_1, \dots, \sigma_0 y_{n-1}) \rho(\sigma_0 x_1, \dots, \sigma_0 x_{n-1})(\sigma_1 m),
\end{aligned}$$

we obtain that

$$\begin{aligned}
&\nu(x_1, \dots, x_{n-1}, [y_1, \dots, y_{n-1}, m]) + [x_1, \dots, x_{n-1}, \nu(y_1, \dots, y_{n-1}, m)] \\
&= \sum_{i=1}^{n-1} \nu(y_1, \dots, [x_1, \dots, x_{n-1}, y_i], \dots, y_{n-1}, m) + \nu(y_1, \dots, y_{n-1}, [x_1, \dots, x_{n-1}, m]) \\
&\quad + \sum_{i=1}^{n-1} [y_1, \nu(x_1, \dots, x_{n-1}, y_i), \dots, y_{n-1}, m] + [y_1, \dots, y_{n-1}, \nu(x_1, \dots, x_{n-1}, m)]. \tag{59}
\end{aligned}$$

Thus  $(\theta, \omega, \nu)$  is a 2-cocycle.  $\square$

Now, we define the  $n$ -Lie algebra structure in  $\mathcal{LM}$  using the 2-cocycle given above. More precisely, we have

$$\left\{ \widehat{f}(m + v) \triangleq f(m) + \theta(v) + \varphi(v), \right.$$

$$\left\{ \begin{array}{l} [x_1 + w_1, \dots, x_n + w_n] \\ \triangleq [x_1, \dots, x_n] + \omega(x_1, \dots, x_n) + \sum_{i=1}^n (-1)^{n-i} \rho_2(x_1, \dots, \widehat{x}_i, \dots, x_n)(w_i), \\ \widehat{\rho}(x_1 + w_1, \dots, x_{n-1} + w_{n-1})(m + v) \\ \triangleq \rho(x_1, \dots, x_{n-1})(m) + \nu(x_1, \dots, x_{n-1}, m) \\ \quad + \rho_1(x_1, \dots, x_{n-1})(v) + \sum_{i=1}^{n-1} (-1)^{n-i} \rho_3(x_1, \dots, \widehat{x}_i, \dots, x_{n-1}, f(m))(w_i), \end{array} \right.$$

for all  $x_i \in \mathfrak{g}$ ,  $m \in M$ ,  $v \in V$  and  $w_i \in W$ . Thus any extension  $E_{\widehat{\mathfrak{g}}}$  given by (56) is isomorphic to

$$\begin{array}{ccccccccc} 0 & \xrightarrow{0} & V & \xrightarrow{i_1} & M \oplus V & \xrightarrow{p_1} & M & \xrightarrow{0} & 0 \\ 0 \downarrow & & \varphi \downarrow & & \widehat{f} \downarrow & & f \downarrow & & 0 \downarrow \\ 0 & \xrightarrow{0} & W & \xrightarrow{i_0} & \mathfrak{g} \oplus W & \xrightarrow{p_0} & \mathfrak{g} & \xrightarrow{0} & 0, \end{array} \quad (60)$$

where the  $n$ -Lie algebra structure in  $\mathcal{LM}$  is given as above.

**Theorem 4.4.** *There is a one-to-one correspondence between equivalence classes of abelian extensions and elements of the second cohomology group  $\mathbf{H}^2((M, \mathfrak{g}, f), (V, W, \varphi))$ .*

**Proof.** Let  $\widehat{E}'$  be another abelian extension determined by the 2-cocycle  $(\theta', \omega', \nu')$ . We are going to show that  $\widehat{E}$  and  $\widehat{E}'$  are equivalent if and only if 2-cocycles  $(\theta, \omega, \nu)$  and  $(\theta', \omega', \nu')$  are in the same cohomology class.

Since  $F = (F_0, F_1)$  is an equivalence of extensions, there exist two linear maps  $b_0 : \mathfrak{g} \rightarrow W$  and  $b_1 : M \rightarrow V$  such that

$$F_0(x_i + w) = x_i + b_0(x_i) + w, \quad F_1(m + v) = m + b_1(m) + v.$$

First, by the equality

$$\widehat{f}' F_1(m) = F_0 \widehat{f}(m),$$

we have

$$\theta(m) - \theta'(m) = \varphi b_1(m) - b_0(f(m)). \quad (61)$$

Second, by direct computations we have

$$\begin{aligned} & F[x_1, \dots, x_n]_{\omega} \\ &= F_1([x_1, \dots, x_n] + \omega(x_1, \dots, x_n)) \\ &= [x_1, \dots, x_n] + \omega(x_1, \dots, x_n) + b_0([x_1, \dots, x_n]) \end{aligned}$$

and

$$\begin{aligned} & [F(x_1), \dots, F(x_n)]_{\omega'} \\ &= [x_1 + b_0(x_1), \dots, x_n + b_0(x_n)]_{\omega'} \\ &= [x_1, \dots, x_n] + \omega'(x_1, \dots, x_n) + \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n) b_0(x_i). \end{aligned}$$

Thus, we have

$$\begin{aligned} & (\omega - \omega')(x_1, \dots, x_n) \\ &= \sum_{i=1}^n (-1)^{n-i} \rho(x_1, \dots, \widehat{x}_i, \dots, x_n) b_0(x_i) - b_0([x_1, \dots, x_n]). \end{aligned} \quad (62)$$

Similarly, by the equality

$$\begin{aligned} & F_1([x_1, \dots, x_{n-1}, m] + \nu(x_1, \dots, x_{n-1}, m)) \\ = & [x_1, \dots, x_{n-1}, m] + \nu(x_1, \dots, x_{n-1}, m) - b_1(x_1, \dots, x_{n-1}, m) \end{aligned}$$

and

$$\begin{aligned} & \rho'(F_0(x_1), \dots, F_0(x_{n-1}))(F_1(m)) \\ = & [x_1, \dots, x_{n-1}, m] + \nu'(x_1, \dots, x_{n-1}, m) \\ & - \sum_{i=1}^{n-1} [x_1, \dots, b_0 x_i, \dots, x_{n-1}, m] - \rho_1(x_1, \dots, x_{n-1})(b_1 m). \end{aligned}$$

Thus we get

$$\begin{aligned} & (\nu - \nu')(x_1, \dots, x_{n-1}, m) \\ = & b_1 \rho(x_1, \dots, x_{n-1})(m) - \sum_{i=1}^{n-1} [x_1, \dots, b_0 x_i, \dots, x_{n-1}, m] - \rho_1(x_1, \dots, x_{n-1})(b_1 m). \end{aligned} \tag{63}$$

By (61)-(63), we deduce that  $(\psi, \omega, \nu) - (\psi', \omega', \nu') = D(b_0, b_1)$ . Thus, they are in the same cohomology class.

Conversely, if  $(\theta, \omega, \nu)$  and  $(\theta', \omega', \nu')$  are in the same cohomology class, assume that  $(\psi, \omega, \nu) - (\psi', \omega', \nu') = D(b_0, b_1)$ . Then define  $(F_0, F_1)$  by

$$F_0(x + w) = x + b_0(x) + w, \quad F_1(m + v) = m + b_1(m) + v.$$

Similar as the above proof, we can show that  $(F_0, F_1)$  is an equivalence. We omit the details.  $\square$

## 5 Infinitesimal deformations

In this section, we explore infinitesimal deformations of  $n$ -Lie algebras in  $\mathcal{LM}$  specifically focusing on the case of  $n = 3$ . The results presented in this section are applicable to the general  $n$  case. For more information on the theory of infinitesimal deformations for both Lie algebras and 3-Lie algebras, one refers to [11, 12, 17].

Let  $(M, \mathfrak{g}, f)$  be a 3-Lie algebra in  $\mathcal{LM}$  and  $\theta : M \rightarrow \mathfrak{g}$ ,  $\omega : \wedge^3 \mathfrak{g} \rightarrow \mathfrak{g}$ ,  $\nu : \wedge^2 \mathfrak{g} \otimes M \rightarrow M$  be linear maps. Consider a  $\lambda$ -parametrized family of linear operations:

$$\begin{aligned} f_\lambda(m) & \triangleq f(m) + \lambda \theta(m), \\ [x_1, x_2, x_3]_\lambda & \triangleq [x_1, x_2, x_3] + \lambda \omega(x_1, x_2, x_3), \\ [x_1, x_2, m]_\lambda & \triangleq [x_1, x_2, m] + \lambda \nu(x_1, x_2, m), \end{aligned}$$

where we denote  $[x_1, x_2, m] \triangleq \rho(x_1, x_2)(m)$  in the following of this section for simplicity.

If  $(M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  forms a 3-Lie algebra in  $\mathcal{LM}$ , then we say that  $(\theta, \omega, \nu)$  generates a 1-parameter infinitesimal deformation of  $(M, \mathfrak{g}, f)$ .

**Theorem 5.1.** *With the notations above,  $(\theta, \omega, \nu)$  generates a 1-parameter infinitesimal deformation of  $(M, \mathfrak{g}, f)$  is equivalent to the following conditions:*



- (i)  $(\theta, \omega, \nu)$  is a 2-cocycle of  $\mathfrak{g}$  with coefficients in the adjoint representation;
- (ii)  $(M, \mathfrak{g}, \theta)$  is a 3-Lie algebra in  $\mathcal{LM}$  with brackets  $\omega$  and  $\nu$ .

*Proof.* If  $(M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  is a 3-Lie algebra in  $\mathcal{LM}$ , then by Definition 3.1,  $f_\lambda$  is an equivariant map, thus we have

$$\begin{aligned}
& f_\lambda([x_1, x_2, m]_\lambda) - [x_1, x_2, f_\lambda(m)]_\lambda \\
= & (f + \lambda\theta)([x_1, x_2, m] + \lambda\nu(x_1, x_2, m)) \\
& - [x_1, x_2, f(m) + \lambda\theta(m)] - \lambda\omega(x_1, x_2, f(m) + \lambda\theta(m)) \\
= & f([x_1, x_2, m]) + \lambda(\theta[x_1, x_2, m] + f\nu(x_1, x_2, m)) + \lambda^2\theta\omega(x_1, x_2, m) \\
& - [x_1, x_2, f(m)] - \lambda(\omega(x_1, x_2, f(m)) + [x_1, x_2, \theta(m)]) - \lambda^2\nu(x_1, x_2, \theta(m)) \\
= & 0,
\end{aligned}$$

which implies that

$$\theta([x_1, x_2, m]) + f\nu(x_1, x_2, m) - \omega(x_1, x_2, f(m)) - [x_1, x_2, \theta(m)] = 0, \quad (64)$$

$$\theta\nu(x_1, x_2, m) - \omega(x_1, x_2, \theta(m)) = 0. \quad (65)$$

Since  $\mathfrak{g}_\lambda$  is a 3-Lie algebra, we have the equality

$$\begin{aligned}
& [x_1, x_2, [y_1, y_2, y_3]_\lambda]_\lambda \\
= & [[x_1, x_2, y_1]_\lambda, y_2, y_3]_\lambda + [y_1, [x_1, x_2, y_2]_\lambda, y_3]_\lambda + [y_1, y_2, [x_1, x_2, y_3]_\lambda]_\lambda,
\end{aligned}$$

the left-hand side is equal to

$$\begin{aligned}
& [x_1, x_2, [y_1, y_2, y_3] + \lambda\omega(y_1, y_2, y_3)]_\lambda \\
= & [x_1, x_2, [y_1, y_2, y_3]] + \lambda\omega(x_1, x_2, [y_1, y_2, y_3]) \\
& + [x_1, x_2, \lambda\omega(y_1, y_2, y_3)] + \lambda\omega(x_1, x_2, \lambda\omega(y_1, y_2, y_3)) \\
= & [x_1, x_2, [y_1, y_2, y_3]] + \lambda\{\omega(x_1, x_2, [y_1, y_2, y_3]) + [x_1, x_2, \omega(y_1, y_2, y_3)]\} \\
& + \lambda^2\omega(x_1, x_2, \omega(y_1, y_2, y_3)),
\end{aligned}$$

and the right-hand side is equal to

$$\begin{aligned}
& [[x_1, x_2, y_1] + \lambda\omega(x_1, x_2, y_1), y_2, y_3]_\lambda + [y_1, [x_1, x_2, y_2] + \lambda\omega(x_1, x_2, y_2), y_3]_\lambda \\
& + [y_1, y_2, [x_1, x_2, y_3] + \lambda\omega(x_1, x_2, y_3)]_\lambda \\
= & [[x_1, x_2, y_1], y_2, y_3] + [y_1, [x_1, x_2, y_2], y_3] + [y_1, y_2, [x_1, x_2, y_3]] \\
& + \lambda\{\omega([x_1, x_2, y_1], y_2, y_3) + [\omega(x_1, x_2, y_1), y_2, y_3] \\
& \quad + \omega(y_1, [x_1, x_2, y_2], y_3) + [y_1, \omega(x_1, x_2, y_2), y_3] \\
& \quad + \omega(y_1, y_2, [x_1, x_2, y_3]) + [y_1, y_2, \omega(x_1, x_2, y_3)]\} \\
& + \lambda^2\{\omega(\omega(x_1, x_2, y_1), y_2, y_3) + \omega(y_1, \omega(x_1, x_2, y_2), y_3) + \omega(y_1, y_2, \omega(x_1, x_2, y_3))\}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
& \omega(x_1, x_2, [y_1, y_2, y_3]) + [x_1, x_2, \omega(y_1, y_2, y_3)] \\
= & \omega([x_1, x_2, y_1], y_2, y_3) + \omega(y_1, [x_1, x_2, y_2], y_3) + \omega(y_1, y_2, [x_1, x_2, y_3]) \\
& + [\omega(x_1, x_2, y_1), y_2, y_3] + [y_1, \omega(x_1, x_2, y_2), y_3] + [y_1, y_2, \omega(x_1, x_2, y_3)], \quad (66)
\end{aligned}$$

$$\begin{aligned}
& \omega(x_1, x_2, \omega(y_1, y_2, y_3)) \\
= & \omega(\omega(x_1, x_2, y_1), y_2, y_3) + \omega(y_1, \omega(x_1, x_2, y_2), y_3) + \omega(y_1, y_2, \omega(x_1, x_2, y_3)). \quad (67)
\end{aligned}$$

Since  $M_\lambda$  is a  $\mathfrak{g}_\lambda$ -module, we get the equality

$$\begin{aligned}
& [x_1, x_2, [y_1, y_2, m]_\lambda]_\lambda \\
= & [[x_1, x_2, y_1]_\lambda, y_2, m]_\lambda + [y_1, [x_1, x_2, y_2]_\lambda, m]_\lambda + [y_1, y_2, [x_1, x_2, m]_\lambda]_\lambda,
\end{aligned}$$

the left-hand side is equal to

$$\begin{aligned}
& [x_1, x_2, [y_1, y_2, m] + \lambda\nu(y_1, y_2, m)]_\lambda \\
= & [x_1, x_2, [y_1, y_2, m]] + \lambda\nu(x_1, x_2, [y_1, y_2, m]) \\
& + [x_1, x_2, \lambda\nu(y_1, y_2, m)] + \lambda\nu(x_1, x_2, \lambda\nu(y_1, y_2, m)) \\
= & [x_1, x_2, [y_1, y_2, m]] + \lambda\{\nu(x_1, x_2, [y_1, y_2, m]) + [x_1, x_2, \nu(y_1, y_2, m)]\} \\
& + \lambda^2\nu(x_1, x_2, \nu(y_1, y_2, m)),
\end{aligned}$$

and the right-hand side is equal to

$$\begin{aligned}
& [[x_1, x_2, y_1] + \lambda\nu(x_1, x_2, y_1), y_2, m]_\lambda + [y_1, [x_1, x_2, y_2] + \lambda\nu(x_1, x_2, y_2), m]_\lambda \\
& + [y_1, y_2, [x_1, x_2, m] + \lambda\nu(x_1, x_2, m)]_\lambda \\
= & [[x_1, x_2, y_1], y_2, m] + [y_1, [x_1, x_2, y_2], m] + [y_1, y_2, [x_1, x_2, m]] \\
& + \lambda\{\nu([x_1, x_2, y_1], y_2, m) + [\nu(x_1, x_2, y_1), y_2, m] \\
& \quad + \nu(y_1, [x_1, x_2, y_2], m) + [y_1, \nu(x_1, x_2, y_2), m] \\
& \quad + \nu(y_1, y_2, [x_1, x_2, m]) + [y_1, y_2, \nu(x_1, x_2, m)]\} \\
& + \lambda^2\{\nu(\nu(x_1, x_2, y_1), y_2, m) + \nu(y_1, \omega(x_1, x_2, y_2), m) + \nu(y_1, y_2, \nu(x_1, x_2, m))\}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
& \nu(x_1, x_2, [y_1, y_2, m] + [x_1, x_2, \nu(y_1, y_2, m)]) \\
= & \nu([x_1, x_2, y_1], y_2, m) + \nu(y_1, [x_1, x_2, y_2], m) + \nu(y_1, y_2, [x_1, x_2, m]) \\
& + [\nu(x_1, x_2, y_1), y_2, m] + [y_1, \omega(x_1, x_2, y_2), m] + [y_1, y_2, \nu(x_1, x_2, m)], \quad (68) \\
& \nu(x_1, x_2, \nu(y_1, y_2, m))
\end{aligned}$$

$$= \nu(\omega(x_1, x_2, y_1), y_2, m) + \nu(y_1, \omega(x_1, x_2, y_2), m) + \nu(y_1, y_2, \nu(x_1, x_2, m)). \quad (69)$$

By (64), (66) and (68), we deduce that  $(\theta, \omega, \nu)$  is a 2-cocycle of  $(M, \mathfrak{g}, f)$  with the coefficients in the adjoint representation. Furthermore, by (65), (67) and (69),  $(M, \mathfrak{g}, \theta, \omega, \nu)$  is a 3-Lie algebra in  $\mathcal{LM}$ .  $\square$

Next, we introduce the notion of Nijenhuis operators which gives trivial deformations. In this case, we consider the second-order deformation  $(M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  where

$$\begin{aligned}
f_\lambda(m) & \triangleq f(m) + \lambda\theta_1(m), \\
[x_1, x_2, x_3]_\lambda & \triangleq [x_1, x_2, x_3] + \lambda\omega_1(x_1, x_2, x_3) + \lambda^2\omega_2(x_1, x_2, x_3), \\
[x_1, x_2, m]_\lambda & \triangleq [x_1, x_2, m] + \lambda\nu_1(x_1, x_2, m) + \lambda^2\nu_2(x_1, x_2, m).
\end{aligned}$$

**Definition 5.2.** A deformation is said to be trivial if there exist linear maps  $N_0 : \mathfrak{g} \rightarrow \mathfrak{g}, N_1 : M \rightarrow M$ , such that  $(T_0, T_1)$  is a homomorphism from  $(M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  to  $(M, \mathfrak{g}, f)$ , where  $T_0 = \text{id} + \lambda N_0, T_1 = \text{id} + \lambda N_1$ .

Note that  $(T_0, T_1)$  is a homomorphism means that

$$f \circ T_1(m) = T_0 \circ f_\lambda(m), \quad (70)$$

$$T_0([x_1, x_2, x_3]_\lambda) = [T_0x_1, T_0x_2, T_0x_3], \quad (71)$$

$$T_1([x_1, x_2, m]_\lambda) = [T_0x_1, T_0x_2, T_1m]. \quad (72)$$

From (70), we have that the left-hand is equal to

$$f \circ T_1(m) = f(m) + \lambda f N_1(m),$$

and the right-hand is equal to

$$\begin{aligned} & T_0 \circ f_\lambda(m) \\ &= (\text{id} + \lambda N_0)(f(m) + \lambda \theta(m)) \\ &= f(m) + \lambda N_0(f(m)) + \lambda \theta(m) + \lambda^2 N_0 \theta(m). \end{aligned}$$

Thus, we get

$$\theta(m) = (f N_1 - N_0 f)(m), \quad N_0 \theta(m) = 0.$$

It follows that  $N$  must satisfy the following condition:

$$N_0(f N_1 - N_0 f) = 0. \quad (73)$$

From the equation (71) we get

$$\begin{aligned} & \omega_1(x_1, x_2, x_3) + N_0[x_1, x_2, x_3] \\ &= [N_0(x_1), x_2, x_3] + [x_1, N_0(x_2), x_3] + [x_1, x_2, N_0(x_3)], \end{aligned} \quad (74)$$

$$\begin{aligned} & \omega_2(x_1, x_2, x_3) + N_0 \omega_1(x_1, x_2, x_3) \\ &= [N_0(x_1), N_0(x_2), x_3] + [N_0(x_1), x_2, N_0(x_3)] + [x_1, N_0(x_2), N_0(x_3)], \end{aligned} \quad (75)$$

$$N_0 \omega_2(x_1, x_2, x_3) = [N_0(x_1), N_0(x_2), N_0(x_3)]. \quad (76)$$

It follows from (74), (75) and (76) that  $N_0$  must satisfy the following condition

$$\begin{aligned} & [N_0(x_1), N_0(x_2), N_0(x_3)] \\ &= N_0([N_0(x_1), N_0(x_2), x_3] + [N_0(x_1), x_2, N_0(x_3)] + [x_1, N_0(x_2), N_0(x_3)]) \\ &\quad - N_0^2([N_0(x_1), x_2, x_3] + [x_1, N_0(x_2), x_3] + [x_1, x_2, N_0(x_3)]) \\ &\quad + N_0^3([x_1, x_2, x_3]). \end{aligned} \quad (77)$$

By definition the left-hand side of equation (72) is equal to

$$\begin{aligned} T_\lambda([x_1, x_2, m]_\lambda) &= [x_1, x_2, m] + \lambda(\nu_1(x_1, x_2, m) + N_1[x_1, x_2, m]) \\ &\quad + \lambda^2(\nu_2(x_1, x_2, m) + N_1\nu_1(x_1, x_2, m)) + \lambda^3 N_1 \nu_2(x_1, x_2, m), \end{aligned}$$

and the right-hand side equals to

$$\begin{aligned} & [T_\lambda(x_1), T_\lambda(x_2), T_\lambda(m)] \\ &= [x_1 + \lambda N_0(x_1), x_2 + \lambda N_0(x_2), m + \lambda N_1(m)] \\ &= [x_1, x_2, m] + \lambda([N_0(x_1), x_2, m] + [x_1, N_0(x_2), m] + [x_1, x_2, N_1(m)]) \\ &\quad + \lambda^2([N_0(x_1), N_0(x_2), m] + [N_0(x_1), x_2, N_1(m)] + [x_1, N_0(x_2), N_1(m)]) \end{aligned}$$

$$+\lambda^3[N_0(x_1), N_0(x_2), N_1(m)].$$

Thus, we have

$$\begin{aligned} \nu_1(x_1, x_2, m) + N_1[x_1, x_2, m] \\ = [N_0(x_1), x_2, m] + [x_1, N_0(x_2), m] + [x_1, x_2, N_1(m)], \end{aligned} \quad (78)$$

$$\begin{aligned} \nu_2(x_1, x_2, m) + N_1\nu_1(x_1, x_2, m) \\ = [N_0(x_1), N_0(x_2), m] + [N_0(x_1), x_2, N_1(m)] + [x_1, N_0(x_2), N_1(m)], \end{aligned} \quad (79)$$

$$N_1\nu_2(x_1, x_2, m) = [N_0(x_1), N_0(x_2), N_1(m)]. \quad (80)$$

It follows from (78), (79) and (80) that  $(N_0, N_1)$  must satisfy the following condition

$$\begin{aligned} & [N_0(x_1), N_0(x_2), N_1(m)] \\ & = N_1([N_0(x_1), N_0(x_2), m] + [N_0(x_1), x_2, N_1(m)] + [x_1, N_0(x_2), N_1(m)]) \\ & \quad - N_1^2([N_0(x_1), x_2, m] + [x_1, N_0(x_2), m] + [x_1, x_2, N_1(m)]) \\ & \quad + N_1^3([x_1, x_2, m])). \end{aligned} \quad (81)$$

**Definition 5.3.** A pair of linear maps  $N = (N_0, N_1)$  is called a Nijenhuis operator if the conditions (73), (77) and (81) hold.

We have seen that any second-order trivial deformation produces a Nijenhuis operator. Conversely, one proves that any Nijenhuis operator gives a second-order trivial deformation.

**Theorem 5.4.** Let  $N = (N_0, N_1)$  be a Nijenhuis operator. Then a second-order deformation can be obtained by putting

$$\left\{ \begin{array}{l} \theta(m) = (fN_1 - N_0f)(m), \\ \omega_1(x_1, x_2, x_3) = [N_0(x_1), x_2, x_3] + [x_1, N_0(x_2), x_3] + [x_1, x_2, N_0(x_3)] - N_0[x_1, x_2, x_3], \\ \omega_2(x_1, x_2, x_3) = [N_0(x_1), N_0(x_2), x_3] + [N_0(x_1), x_2, N_0(x_3)] + [x_1, N_0(x_2), N_0(x_3)] \\ \quad - N_1\omega_1(x_1, x_2, x_3), \\ \nu_1(x_1, x_2, m) = [N_0(x_1), x_2, m] + [x_1, N_0(x_2), m] + [x_1, x_2, N_1(m)] - N_1[x_1, x_2, m], \\ \nu_2(x_1, x_2, m) = [N_0(x_1), N_0(x_2), m] + [N_0(x_1), x_2, N_1(m)] + [x_1, N_0(x_2), N_1(m)] \\ \quad - N_1\nu_1(x_1, x_2, m). \end{array} \right. \quad (82)$$

Furthermore, this deformation is trivial.

Finally, we consider the formal deformations of any order. Let  $(M, \mathfrak{g}, f)$  be a 3-Lie algebra in  $\mathcal{LM}$  and  $\theta_i : M \rightarrow \mathfrak{g}$ ,  $\omega_i : \wedge^3 \mathfrak{g} \rightarrow \mathfrak{g}$ ,  $\nu_i : \wedge^2 \mathfrak{g} \otimes M \rightarrow M, i \geq 0$  be linear maps. Consider the  $\lambda$ -parametrized family of linear operations:

$$\begin{aligned} f_\lambda(m) & \triangleq f(m) + \lambda\theta_1(m) + \lambda^2\theta_2(m) + \dots, \\ [x_1, x_2, x_3]_\lambda & \triangleq [x_1, x_2, x_3] + \lambda\omega_1(x_1, x_2, x_3) + \lambda^2\omega_2(x_1, x_2, x_3) + \dots, \\ [x_1, x_2, m]_\lambda & \triangleq [x_1, x_2, m] + \lambda\nu_1(x_1, x_2, m) + \lambda^2\nu_2(x_1, x_2, m) + \dots \end{aligned}$$

In order that  $(M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  forms a 3-Lie algebra in  $\mathcal{LM}$ , we must have

$$\begin{aligned} \omega_\lambda(x_1, x_2, \omega_\lambda(y_1, y_2, y_3)) & = \omega_\lambda(\omega_\lambda(x_1, x_2, y_1), y_2, y_3) + \omega_\lambda(y_1, \omega_\lambda(x_1, x_2, y_2), y_3) \\ & \quad + \omega_\lambda(y_1, y_2, \omega_\lambda(x_1, x_2, y_3)), \\ \nu_\lambda(x_1, x_2, \nu_\lambda(y_1, y_2, m)) & = \nu_\lambda(\omega_\lambda(x_1, x_2, y_1), y_2, m) + \nu_\lambda(y_1, \omega_\lambda(x_1, x_2, y_2), m) \end{aligned}$$

$$f_\lambda \nu_\lambda(x_1, x_2, m) = \omega_\lambda(x_1, x_2, f_\lambda(m)) + \nu_\lambda(y_1, y_2, \nu_\lambda(x_1, x_2, m)),$$

which implies that

$$\begin{aligned} \sum_{i+j=k} \omega_i(x_1, x_2, \omega_j(y_1, y_2, y_3)) &= \sum_{i+j=k} \omega_i(\omega_j(x_1, x_2, y_1), y_2, y_3) + \omega_i(y_1, \omega_j(x_1, x_2, y_2), y_3) \\ &\quad + \omega_i(y_1, y_2, \omega_j(x_1, x_2, y_3)), \end{aligned} \quad (83)$$

$$\begin{aligned} \sum_{i+j=k} \nu_i(x_1, x_2, \nu_j(y_1, y_2, m)) &= \sum_{i+j=k} \nu_i(\omega_j(x_1, x_2, y_1), y_2, m) + \omega_i(y_1, \nu_j(x_1, x_2, m), y_3) \\ &\quad + \nu_i(y_1, y_2, \nu_j(x_1, x_2, m)), \end{aligned} \quad (84)$$

$$\sum_{i+j=k} f_i \nu_j(x_1, x_2, m) = \sum_{i+j=k} \omega_i(x_1, x_2, f_j(m)), \quad (85)$$

where we denote  $f_0(m) = f(m)$ ,  $\omega_0(x_1, x_2, x_3) = [x_1, x_2, x_3]$ ,  $\nu_0(x_1, x_2, m) = [x_1, x_2, m]$ .

For  $k = 0$ , conditions (83), (84) and (85) are equivalent to  $(\omega_0, \nu_0, f_0)$  is an  $n$ -Lie algebra in  $\mathcal{LM}$ . For  $k = 1$ , these conditions are equivalent to

$$\begin{aligned} &\omega_1(x_1, x_2, [y_1, y_2, y_3]) + [x_1, x_2, \omega_1(y_1, y_2, y_3)] \\ &= \omega_1([x_1, x_2, y_1], y_2, y_3) + \omega_1(y_1, [x_1, x_2, y_2], y_3) + \omega_1(y_1, y_2, [x_1, x_2, y_3]) \\ &\quad + [\omega_1(x_1, x_2, y_1), y_2, y_3] + [y_1, \omega_1(x_1, x_2, y_2), y_3] + [y_1, y_2, \omega_1(x_1, x_2, y_3)], \\ &\nu_1(x_1, x_2, [y_1, y_2, m]) + [x_1, x_2, \nu_1(y_1, y_2, m)] \\ &= \nu_1([x_1, x_2, y_1], y_2, m) + \nu_1(y_1, [x_1, x_2, y_2], m) + \nu_1(y_1, y_2, [x_1, x_2, m]) \\ &\quad + [\nu_1(x_1, x_2, y_1), y_2, m] + [y_1, \omega_1(x_1, x_2, y_2), m] + [y_1, y_2, \nu_1(x_1, x_2, m)], \\ &\theta_1([x_1, x_2, m]) + f \nu_1(x_1, x_2, m) = \omega_1(x_1, x_2, f(m)) + [x_1, x_2, \theta_1(m)]. \end{aligned}$$

Thus for  $k = 1$ ,  $(\omega_1, \nu_1, \theta_1) \in C^2((M, \mathfrak{g}, f), (M, \mathfrak{g}, f))$  is a 2-cocycle.

**Definition 5.5.** The 2-cochain  $(\omega_1, \nu_1, \theta_1)$  is called the infinitesimal of the deformation  $(\omega_\lambda, \nu_\lambda, f_\lambda)$ . More generally, if  $(\omega_i, \nu_i, f_i) = 0$  for  $1 \leq i \leq (n-1)$ , and  $(\omega_n, \nu_n, f_n)$  is a non-zero cochain in  $C^2((M, \mathfrak{g}, f), (M, \mathfrak{g}, f))$ , then  $(\omega_n, \nu_n, f_n)$  is called the  $n$ -infinitesimal of the deformation  $(\omega_\lambda, \nu_\lambda, f_\lambda)$ .

Let  $(\omega_\lambda, \nu_\lambda, f_\lambda)$  and  $(\omega'_\lambda, \nu'_\lambda, f'_\lambda)$  be two deformation. We say that they are equivalent if there exists a formal isomorphism  $(\Phi_\lambda, \Psi_\lambda) : (M'_\lambda, \mathfrak{g}'_\lambda, f'_\lambda) \rightarrow (M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  such that  $\omega'_\lambda(x_1, x_2, x_3) = \Psi_\lambda^{-1} \omega_\lambda(\Psi_\lambda(x_1), \Psi_\lambda(x_2), \Psi_\lambda(x_3))$ .

A deformation  $(\omega_\lambda, \nu_\lambda, f_\lambda)$  is said to be the trivial deformation if it is equivalent to  $(\omega_0, \nu_0, f)$ .

**Theorem 5.6.** Let  $(\omega_\lambda, \nu_\lambda, f_\lambda)$  and  $(\omega'_\lambda, \nu'_\lambda, f'_\lambda)$  be equivalent deformations of  $(M, \mathfrak{g}, f)$  be a 3-Lie algebra in  $\mathcal{LM}$ , then the first-order terms of them belong to the same cohomology class in the second cohomology group  $H^2((M, \mathfrak{g}, f), (M, \mathfrak{g}, f))$ .

**Proof.** Let  $(\Phi_\lambda, \Psi_\lambda) : (M'_\lambda, \mathfrak{g}'_\lambda, f'_\lambda) \rightarrow (M_\lambda, \mathfrak{g}_\lambda, f_\lambda)$  be an equivalence where  $\Phi_\lambda = \text{id}_M + \lambda\phi_1 + \lambda^2\phi_2 + \dots$  and  $\Psi_\lambda = \text{id}_M + \lambda\psi_1 + \lambda^2\psi_2 + \dots$ . Then we have  $\Psi_\lambda \omega'_\lambda(x_1, x_2, x_3) = \omega_\lambda(\Psi_\lambda(x_1), \Psi_\lambda(x_2), \Psi_\lambda(x_3))$   $\Psi_\lambda \nu'_\lambda(x_1, x_2, m) = \nu_\lambda(\Phi_\lambda(x_1), \Phi_\lambda(x_2), \Psi_\lambda(m))$   $\square$

A 3-Lie algebra  $(M, \mathfrak{g}, f)$  in  $\mathcal{LM}$  is called rigid if every deformation  $(\omega_\lambda, \nu_\lambda, f_\lambda)$  is equivalent to the trivial deformation.

**Theorem 5.7.** *If  $H^2((M, \mathfrak{g}, f), (M, \mathfrak{g}, f)) = 0$ , then  $(M, \mathfrak{g}, f)$  is rigid.*

*Proof.* Let  $(\omega_\lambda, \nu_\lambda, f_\lambda)$  be a deformation of  $(M, \mathfrak{g}, f)$ . It follows from the above results that  $D(\omega_\lambda, \nu_\lambda, f_\lambda) = 0$ , that is  $(\omega_\lambda, \nu_\lambda, f_\lambda) \in Z^2((M, \mathfrak{g}, f), (M, \mathfrak{g}, f))$ . Now assume  $H^2((M, \mathfrak{g}, f)) = 0$ , we can find  $(N_0, N_1)$  such that  $(\omega_\lambda, \nu_\lambda, f_\lambda) = D(N_0, N_1)$ . The proof is completed.  $\square$

## Acknowledgements

This research was supported by the National Natural Science Foundation of China (No. 11961049).

## Data availability

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

## Declarations

**Conflict of interest** On behalf of all authors, the corresponding author states that there is no conflict of interest.

## References

- [1] M. Afi and O. Basdouri, On the isomorphism of non-abelian extensions of  $n$ -Lie algebras, *J. Geom. Phys.* 173 (2022) 104427.
- [2] R. Bai, W. Wu, Y. Li and Z. Li, Module extensions of 3-Lie algebras, *Linear Multilinear Algebra* 60 (4) (2012) 583–594.
- [3] J. F. Cariñena, J. Grabowski and G. Marmo, Quantum bi-Hamiltonian systems, *Int. J. Mod. Phys. A* 15 (2000) 4797–4810.
- [4] J. M. Casas, E. Khmaladze and M. Ladra, Crossed modules for Leibniz  $n$ -algebras, *Forum Math.* 20 (2008) 841–858.
- [5] J. M. Casas, E. Khmaladze and M. Ladra, Notes on Leibniz  $n$ -algebras, *J. Geom. Phys.* 195 (2024) 105035.
- [6] J. M. Casas, J.-L. Loday and T. Pirashvili, Leibniz  $n$ -algebras, *Forum Math.* 14 (2002) 189–207.
- [7] M. Chen, S-K. Wang, K. Wu, W. Zhao, Infinite-dimensional 3-algebra and integrable system, *JHEP* 12 (2012) 030.
- [8] Y. I. Daletskii and L. A. Takhtajan, Leibniz and Lie algebra structures for Nambu algebra, *Lett. Math. Phys.* 39 (1997) 127–141.
- [9] X. García-Martínez, R. Turdibaev, T. Van der Linden, Do  $n$ -Lie algebras have universal enveloping algebras?, *J. Lie Theory* 28 (1) (2018), 43–55.

- [10] P. Gautheron, Some remarks concerning Nambu mechanics, *Lett. Math. Phys.* 37 (1996) 103–116.
- [11] I. Dorfman, Dirac Structures and Integrability of Nonlinear Evolution Equation, John Wiley & Sons, Ltd., Chichester, 1993.
- [12] J. Figueroa-O’Farrill, Deformations of 3-algebras, *J. Math. Phys.* 50 (2009) 113514.
- [13] V. T. Filippov,  $n$ -Lie algebras, *Sibirsk. Mat. Zh.* 26 (6) (1985) 126–140.
- [14] P. Ho, R. Hou and Y. Matsuo, Lie 3-algebra and multiple M2-branes, *JHEP* 06 (2008) 020.
- [15] Sh. M. Kasymov, On a theory of  $n$ -Lie algebras, *Algebra Logic* 26 (3) (1987) 277–297.
- [16] M. K. Kinyon and A. Weinstein, Leibniz algebras, Courant algebroids, and multiplications on reductive homogeneous spaces, *Amer. J. Math.* 123 (3) (2001), 525–550.
- [17] Y. Kosmann-Schwarzbach and F. Magri, Poisson-Nijenhuis structures, *Ann. Inst. Henri Poincaré* 53 (1) (1990) 35–81.
- [18] A. Kotov and T. Strobl, The embedding tensor, Leibniz-Loday algebras and their higher gauge theories, *Comm. Math. Phys.* 376 (2020) 235–258.
- [19] R. Kurdiani, Cohomology of Lie algebras in the tensor category of linear maps, *Comm. Algebra* 27 (10) (1999) 5033–5048.
- [20] S. Lavau and J. Palmkvist, Infinity-enhancing Leibniz algebras, *Lett. Math. Phys.* 110 (2020) 3121–3152.
- [21] W. X. Ling, On the structure of  $n$ -Lie algebras, Ph.D. Thesis, University-GHS-Siegen, Siegen, 1993.
- [22] W. Liu and Z. Zhang,  $T^*$ -extension of 3-Lie algebras, *Linear Multilinear Algebra* 60 (5) (2012) 583–594.
- [23] J.-L. Loday and T. Pirashvili, Universal enveloping algebras of Leibniz algebras and (co)-homology, *Math. Ann.* 296 (1993) 139–158.
- [24] J.-L. Loday and T. Pirashvili, The tensor category of linear maps and Leibniz algebras, *Georgian Math. J.* 5 (3) (1998) 263–276.
- [25] Y. Nambu, Generalized Hamiltonian dynamics, *Phys. Rev. D* 7 (1973) 2405–2412.
- [26] J. Palmkvist, The tensor hierarchy algebra, *J. Math. Phys.* 55 (2014) 011701.
- [27] C. Papageorgakis and C. Sämann, The 3-Lie algebra (2,0) tensor multiplet and equations of motion on loop space, *JHEP* 05 (2011) 099.
- [28] M. Rotkiewicz, Cohomology ring of  $n$ -Lie algebras, *Extracta Math.* 20 (2005) 219–232.
- [29] A. Rovi, Lie algebroids in the Loday-Pirashvili category, *SIGMA* 11 (2015) 079, 16 pages.
- [30] L. Takhtajan, On foundation of the generalized Nambu mechanics, *Comm. Math. Phys.* 160 (1994) 295–316.

- [31] L. Takhtajan, A higher order analog of Chevalley-Eilenberg complex and deformation theory of  $n$ -algebras, *St. Petersburg Math. J.* 6 (1995) 429–438.
- [32] T. Zhang, Notes on cohomologies of Lie triple systems, *J. Lie Theory* 24 (4) (2014) 909–929.
- [33] T. Zhang, Cohomology and deformations of 3-Lie colour algebras, *Linear Multilinear Algebra* 63 (2015) 651–671.
- [34] T. Zhang, Extending structures for 3-Lie algebras, *Comm. Algebra* 50 (4) (2022) 1469–1497.