

FINITISTIC SPACES WITH ORBIT SPACE A PRODUCT OF PROJECTIVE SPACES

ANJU KUMARI AND HEMANT KUMAR SINGH

ABSTRACT. Let $G = \mathbb{Z}_2$ act freely on a finitistic space X . If the mod 2 cohomology of X is isomorphic to the real projective space $\mathbb{R}P^{2n+1}$ (resp. complex projective space $\mathbb{C}P^{2n+1}$) then the mod 2 cohomology of orbit spaces of these free actions are $\mathbb{R}P^1 \times \mathbb{C}P^n$ (resp. $\mathbb{R}P^2 \times \mathbb{H}P^n$) [7]. In this paper, we have discussed converse of these result. We have showed that if the mod 2 cohomology of the orbit space X/G is $\mathbb{R}P^1 \times \mathbb{C}P^n$ (resp. $\mathbb{R}P^2 \times \mathbb{H}P^n$) then the mod 2 cohomology of X is $\mathbb{R}P^{2n+1}$ or $\mathbb{S}^1 \times \mathbb{C}P^n$ (resp. $\mathbb{C}P^{2n+1}$ or $\mathbb{S}^2 \times \mathbb{H}P^n$).

A partial converse of free involutions on the product of projective spaces $\mathbb{R}P^n \times \mathbb{R}P^{2m+1}$ (resp. $\mathbb{C}P^n \times \mathbb{C}P^{2m+1}$) are also discussed.

1. INTRODUCTION

Let G be a compact Lie group acting on a space X . An interesting problem is to classify X when G acts freely on X and the orbit space X/G is known. First, J. C. Su [9] worked in this direction. He proved that if $G = \mathbb{Z}_2$ or \mathbb{S}^1 , acts freely on a space X and the orbit space X/G is the mod 2 or integral cohomology projective space $\mathbb{F}P^n$, $\mathbb{F} = \mathbb{R}$ or \mathbb{C} , then the space X is the mod 2 cohomology n -sphere \mathbb{S}^n or integral cohomology $(2n + 1)$ -sphere \mathbb{S}^{2n+1} . Su also proved that if $G = \mathbb{Z}_q$, q an odd prime, and $X/G = L_q^{2n+1}$ (lens space) then X is the mod q cohomology $(2n + 1)$ -sphere \mathbb{S}^{2n+1} . It has also been proved that if $G = \mathbb{S}^3$ and $X/G = \mathbb{H}P^n$ (quaternion projective space), then X is either integral cohomology \mathbb{S}^{4n+3} or $\mathbb{S}^3 \times \mathbb{H}P^n$ [4]. Harvey [2] proved that for semi-free actions of $G = \mathbb{S}^1$ when the orbit space X/G is a simply connected 4-manifold, then the total space is a simply connected 5-manifold.

The orbit spaces of free actions of $G = \mathbb{Z}_2$ on projective spaces $X = \mathbb{F}P^{2n+1}$ and the product of projective spaces $X = \mathbb{F}P^n \times \mathbb{F}P^m$, where $\mathbb{F} = \mathbb{R}$ or \mathbb{C} , have been discussed in [7, 8]. For $G = \mathbb{Z}_2$ and $X = \mathbb{R}P^{2n+1}$ (resp. $\mathbb{C}P^{2n+1}$), it is proved that the mod 2 cohomology of the orbit space X/G is $\mathbb{R}P^1 \times \mathbb{C}P^n$ (resp. $\mathbb{R}P^2 \times \mathbb{H}P^n$) [7]. We have discussed converse

2010 *Mathematics Subject Classification*. Primary 55T10; Secondary 57S99 .

Key words and phrases. Free action; Finitistic space; Leray-Serre spectral sequence; Gysin sequence.

This paper is supported by the Science and Engineering Research Board (Department of Science and Technology, Government of India) with reference number- EMR/2017/002192.

of these results. We have determined the cohomology classification of a finitistic space X if the mod 2 cohomology of orbit space X/G is $\mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^n$ or $\mathbb{R}\mathbb{P}^2 \times \mathbb{H}\mathbb{P}^n$. For $G = \mathbb{Z}_2$ and $X = \mathbb{R}\mathbb{P}^n \times \mathbb{R}\mathbb{P}^{2m+1}$ (resp. $\mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{2m+1}$), one of the possibilities of the mod 2 cohomology of the orbit space is $\mathbb{R}\mathbb{P}^1 \times \mathbb{R}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^m$ (resp. $\mathbb{R}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m$) [8]. In this paper, we have also determined X equipped with a free action of G when the mod 2 cohomology of the orbit space X/G is $\mathbb{R}\mathbb{P}^1 \times \mathbb{R}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^m$ or $\mathbb{R}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m$. This gives a partial converse of results proved in [8].

All spaces are assumed to be finitistic connected spaces. Recall that a paracompact Hausdorff space X is said to be finitistic if every open cover of X has a finite dimensional open refinement. All compact spaces and all finite dimensional paracompact spaces are finitistic spaces. We have considered Čech cohomology with coefficients in \mathbb{Z}_2 . Throughout the paper, $X \sim_2 Y$ means $H^*(X; \mathbb{Z}_2) \cong H^*(Y; \mathbb{Z}_2)$. We have proved the following results:

Theorem 1.1. Let $G = \mathbb{Z}_2$ act freely on a finitistic connected space X with $X/G \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^n$. Then either $X \sim_{\mathbb{Z}_2} \mathbb{S}^1 \times \mathbb{C}\mathbb{P}^n$ or $X \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^{2n+1}$.

Theorem 1.2. Let $G = \mathbb{Z}_2$ act freely on a finitistic connected space X with $X/G \sim_2 \mathbb{R}\mathbb{P}^2 \times \mathbb{H}\mathbb{P}^n$. Then $X \sim_2 \mathbb{S}^2 \times \mathbb{H}\mathbb{P}^n$ or $X \sim_2 \mathbb{C}\mathbb{P}^{2n+1}$.

Theorem 1.3. Let $G = \mathbb{Z}_2$ act freely on a finitistic connected space X with $X/G \sim_2 \mathbb{R}\mathbb{P}^1 \times \mathbb{R}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^m$ and $\pi_1(B_G)$ act trivially on $H^*(X)$. Then the cohomology algebra of X is one of the following:

- (i) $\mathbb{Z}_2[x, y, z]/\langle x^{n+1}, y^{m+1}, z^2 + \alpha x^2 + \beta y \rangle$, where $\deg x = \deg z = 1$, $\deg y = 2$ and $\alpha, \beta \in \mathbb{Z}_2$;
- (ii) $\mathbb{Z}_2[x, y, z]/\langle x^2, y^{m+1}, z^2 + \alpha y^n \rangle$, where $\deg x = 1$, $\deg y = 2$, $\deg z = n$, $\alpha \in \mathbb{Z}_2$ and $\alpha = 0$ if $n > m$.

Theorem 1.4. Let $G = \mathbb{Z}_2$ act freely on a finitistic connected space X with $X/G \sim_2 \mathbb{R}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m$ and $\pi_1(B_G)$ acts trivially on $H^*(X)$. Then the cohomology algebra of X is isomorphic to

$$\mathbb{Z}_2[x, y, z]/\langle x^{n+1}, y^{m+1}, z^2 + \alpha x^2 + \beta y \rangle,$$

where $\deg x = 2$, $\deg y = 4$, $\deg z = 2$ and $\alpha, \beta \in \mathbb{Z}_2$.

2. PRELIMINARIES

In this section, we have recalled some results and definitions which are used to prove our main results.

Let G be a compact Lie group and $G \hookrightarrow E_G \rightarrow B_G$ be the universal principal G -bundle, where B_G is the classifying space. Suppose G acts freely on a finitistic space X . The associated bundle $X \hookrightarrow (X \times E_G)/G \rightarrow B_G$ is a fibre bundle with fibre X . Put $X_G = (X \times E_G)/G$. Then the bundle $X \hookrightarrow X_G \rightarrow B_G$ is called the Borel fibration. We consider the Leray-Serre spectral sequence for the Borel fibration. If $\pi_1(B_G)$ acts trivially on $H^*(X)$, then the system of local coefficients on B_G is simple and the E_2 -term of the Leray-Serre spectral sequence corresponding to the Borel fibration is given by

$$E_2^{k,l} = H^k(B_G) \otimes H^l(X).$$

This sequence converges to $H^*(X_G)$ as an algebra. We recall some results which are needed to prove our results:

Proposition 2.1. Let $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ be the Borel fibration. Suppose that the system of local coefficients on B_G is simple, then the edge homomorphisms

$$H^k(B_G) \cong E_2^{k,0} \longrightarrow E_3^{k,0} \longrightarrow \cdots \longrightarrow E_k^{k,0} \longrightarrow E_{k+1}^{k,0} = E_\infty^{k,0} \subset H^k(X_G)$$

and

$$H^l(X_G) \longrightarrow E_\infty^{0,l} = E_{l+2}^{0,l} \subset E_{l+1}^{0,l} \subset \cdots \subset E_2^{0,l} \cong H^l(X)$$

are the homomorphisms

$$\pi_k^* : H^k(B_G) \rightarrow H^k(X_G) \text{ and } i_l^* : H^l(X_G) \rightarrow H^l(X).$$

For details about spectral sequences, we refer [5]. Let G act freely on X and $h : X_G \rightarrow X/G$ be the map induced by G -equivariant projection $X \times E_G \rightarrow X$. Then h is a homotopy equivalence [1]. Now, we recall Gysin sequence of sphere bundles.

Proposition 2.2 ([3]). Let $G = \mathbb{Z}_2$ act freely on a finitistic space X . The Gysin sequence of the sphere bundle $G \hookrightarrow X \xrightarrow{p} X/G$ is:

$$\cdots \longrightarrow H^{i-1}(X/G) \xrightarrow{\cup} H^i(X/G) \xrightarrow{p_i^*} H^i(X) \xrightarrow{\rho_i} H^i(X/G) \xrightarrow{\cup} H^{i+1}(X/G) \longrightarrow \cdots$$

which start with

$$0 \longrightarrow H^0(X/G) \xrightarrow{p_0^*} H^0(X) \xrightarrow{\rho_0} H^0(X/G) \xrightarrow{\cup} H^1(X/G) \xrightarrow{p_1^*} H^1(X) \longrightarrow \cdots$$

where $\cup : H^i(X/G) \rightarrow H^{i+1}(X/G)$ maps $x \rightarrow xe$, $e \in H^1(X/G)$ denotes the image of 1 under the map $\cup : H^0(X/G) \rightarrow H^1(X/G)$ and e is called the Euler class of the sphere bundle.

Proposition 2.3 ([6]). Let $G = \mathbb{Z}_2$ act freely on a finitistic space X . Suppose that $H^j(X) = 0$ for all $j > n$, then $H^j(X/G) = 0$ for all $j > n$.

3. PROOF OF MAIN THEOREMS

Proof of Theorem 1.1. As $X/G \sim_2 \mathbb{R}P^1 \times \mathbb{C}P^n$, we have $H^*(X/G) = \mathbb{Z}_2[a, b]/\langle a^2, b^{n+1} \rangle$, where $\deg a = 1$ and $\deg b = 2$. By connectedness of X and the exactness of Gysin sequence of G -bundle $G \hookrightarrow X \xrightarrow{p} X/G$, we get p_0^* is an isomorphism and the Euler class $\cup(1)$ is a . This implies that for $0 \leq i < n$, $p_{2i+2}^* : H^{2i+2}(X/G) \rightarrow H^{2i+2}(X)$, $\rho_{2i+1} : H^{2i+1}(X) \rightarrow H^{2i+1}(X/G)$ and $\rho_{2n+1} : H^{2n+1}(X) \rightarrow H^{2n+1}(X/G)$ are isomorphisms. Consequently, $H^{2i+1}(X) \cong H^{2n+1}(X) \cong H^{2i+2}(X) \cong \mathbb{Z}_2$ with bases $\{u_{2i+1}\}$, $\{u_{2n+1}\}$ and $\{x^{i+1}\}$, respectively, where $x = p_2^*(b)$, $\rho_{2i+1}(u_{2i+1}) = ab^i$, $\rho_{2n+1}(u_{2n+1}) = ab^n$. By Proposition 2.3, $H^i(X) = 0$ for $i > 2n + 1$. Now, we compute the cohomology algebra of X . The Leray-Serre spectral sequence associated to the Borel fibration $X \hookrightarrow X_G \rightarrow B_G$ converges to $H^*(X_G)$. It is easy to see that G acts trivially on $H^*(X)$. Thus, the E_2 -term $E_2^{s,t}$ of the Leray-Serre spectral sequence associated to the Borel fibration $X \hookrightarrow X_G \xrightarrow{\pi} B_G$ is $H^s(B_G) \otimes H^t(X)$. By the edge homomorphism, we get $d_r(1 \otimes x) = 0$ for all $r > 0$ and $d_{r'}(1 \otimes u_1) \neq 0$ for some $r' > 0$. So, $d_2(1 \otimes u_1) = t^2 \otimes 1$. This gives that $x^i u_1 \neq 0$, and hence $u_{2i+1} = x^i u_1$ for all $0 \leq i \leq n$. Clearly, $x^{n+1} = 0$. If $u_1^2 = 0$ then $X \sim_2 \mathbb{S}^1 \times \mathbb{C}P^n$. If $u_1^2 \neq 0$ then $X \sim_2 \mathbb{R}P^{2n+1}$. \square

Proof of Theorem 1.2. As $X/G \sim_2 \mathbb{R}P^2 \times \mathbb{H}P^n$, we have $H^*(X/G) = \mathbb{Z}_2[a, b]/\langle a^3, b^{n+1} \rangle$, where $\deg a = 1$ and $\deg b = 4$. It is easy to see that $\cup : H^0(X/G) \rightarrow H^1(X/G)$ maps 1 to a . Since $H^j(X/G) = 0$ for $j \equiv 3 \pmod{4}$, we get $p_{4i}^* : H^{4i}(X/G) \rightarrow H^{4i}(X)$ and $\rho_{4i+2} : H^{4i+2}(X) \rightarrow H^{4i+2}(X/G)$ are isomorphisms for $1 \leq i \leq n$. Consequently, for $1 \leq i \leq n$, we get $H^{4i}(X) \cong H^{4i+2}(X) \cong \mathbb{Z}_2$ with bases $\{x^i\}$ and $\{u_{4i+2}\}$, respectively, where $x = p_4^*(b)$ and $\rho_{4i+2}(u_{4i+2}) = a^2 b^i$. By the exactness of Gysin sequence, $H^{4i+3}(X) = H^{4i+1}(X) = 0$ for all $0 \leq i \leq n$. Clearly, $H^i(X) = 0$ for all $i > 4n + 3$. Now, we compute cohomology algebra of X . As $\pi_1(B_G)$ acts trivially on $H^*(X)$, we get $E_2^{s,t} = H^s(B_G) \otimes H^t(X)$. By the edge homomorphism, we get $d_3(1 \otimes u_2) = t^3 \otimes 1$ and $d_3(1 \otimes x) = 0$. Thus, $x^i u_2 \neq 0$, and hence $u_{4i+2} = x^i u_2$ for all $0 \leq i \leq n$. Clearly, $x^{n+1} = 0$. If $u_2^2 = 0$ then $X \sim_2 \mathbb{S}^2 \times \mathbb{H}P^n$. Otherwise, we get $X \sim_2 \mathbb{C}P^{2n+1}$. \square

Proof of Theorem 1.3. It is given that $X/G \sim_2 \mathbb{R}P^1 \times \mathbb{R}P^n \times \mathbb{C}P^m$. So, we get $H^*(X/G) = \mathbb{Z}_2[a, b, c]/\langle a^{n+1}, b^{m+1}, c^2 \rangle$, where $\deg a = \deg c = 1$ and $\deg b = 2$. As X is connected, $H^0(X) \cong \mathbb{Z}_2$. First, we consider $1 \leq n < 2m$. By Proposition 2.3, $H^i(X) = 0$ for all $i > 2m + n + 1$. Consider the Gysin sequence of the sphere bundle $G \hookrightarrow X \xrightarrow{p} X/G$:

$$\dots \longrightarrow H^{i-1}(X/G) \xrightarrow{\cup} H^i(X/G) \xrightarrow{p_i^*} H^i(X) \xrightarrow{\rho_i} H^i(X/G) \xrightarrow{\cup} H^{i+1}(X/G) \longrightarrow \dots$$

which begins with

$$0 \longrightarrow H^0(X/G) \xrightarrow{p_0^*} H^0(X) \xrightarrow{\rho_0} H^0(X/G) \xrightarrow{\cup} H^1(X/G) \xrightarrow{p_1^*} H^1(X) \longrightarrow \dots$$

Clearly, the Euler class of the associated principal G -bundle $G \hookrightarrow X \rightarrow X/G$ must be nontrivial. We have considered the following three cases:

Case (1): When $\cup : H^0(X/G) \rightarrow H^1(X/G)$ maps 1 to c .

For $0 \leq i < \frac{n}{2}$, we have $H^{2i}(X/G) \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($2i + 1$ -copies) and $H^{2i+1}(X/G) \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($2i + 2$ -copies) with bases

$$\{b^i, ab^{i-1}c, a^2b^{i-1}, a^3b^{i-2}c, \dots, a^{2i-1}c, a^{2i}\}, \text{ and}$$

$$\{b^i c, ab^i, a^2b^{i-1}c, a^3b^{i-1}, \dots, a^{2i}c, a^{2i+1}\},$$

respectively. By the exactness of Gysin sequence, we get $\ker p_{2i}^* = \text{im } \rho_{2i} \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ (i -copies) with basis $\{a^{2j+1}b^{i-j-1}c \mid 0 \leq j \leq i-1\}$, and $\ker p_{2i+1}^* = \text{im } \rho_{2i+1} \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($i + 1$ -copies) with basis $\{a^{2j}b^{i-j}c \mid 0 \leq j \leq i\}$. Consequently, $\text{im } p_{2i}^* \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($i + 1$ -copies) with basis $\{x^{2j}y^{i-j} \mid 0 \leq j \leq i\}$, and $\text{im } p_{2i+1}^* \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($i + 1$ -copies) with basis $\{x^{2j+1}y^{i-j} \mid 0 \leq j \leq i\}$, where $p_1^*(a) = x, p_2^*(b) = y$. It is easy to see that

$$0 \rightarrow \text{im } p_j^* \hookrightarrow H^j(X) \xrightarrow{\rho_j} \text{im } \rho_j \rightarrow 0, j = 2i, 2i + 1$$

are split exact sequences. This gives that

$$H^{2i}(X) \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2 (2i + 1\text{-copies}), \text{ and}$$

$$H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2 (2i + 2\text{-copies})$$

with bases $\{x^{2j}y^{i-j}, u_{2i}^{2l+1} \mid 0 \leq j \leq i, 0 \leq l \leq i-1\}$ and $\{x^{2j+1}y^{i-j}, u_{2i+1}^{2j} \mid 0 \leq j \leq i\}$, respectively, where $\rho_{2i}(u_{2i}^{2l+1}) = a^{2l+1}b^{i-l-1}c, 0 \leq l \leq i-1$ and $\rho_{2i+1}(u_{2i+1}^{2j}) = a^{2j}b^{i-j}c, 0 \leq j \leq i$.

Firstly, suppose that $n = 2k$ for some k . Then for $\frac{n}{2} \leq i \leq m$, $H^{2i}(X/G) \cong H^{2i+1}(X/G) \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($n + 1$ -copies) with bases

$$\{b^i, ab^{i-1}c, a^2b^{i-1}, a^3b^{i-2}c, \dots, a^{n-1}b^{i-k}c, a^n b^{i-k}\}, \text{ and}$$

$$\{b^i c, ab^i, a^2b^{i-1}c, a^3b^{i-1}, \dots, a^{n-1}b^{i-k+1}, a^n b^{i-k}c\},$$

respectively. Again, by the exactness of Gysin sequence, $\ker p_{2i}^* = \text{im } \rho_{2i} \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ (k -copies) and $\ker p_{2i+1}^* = \text{im } \rho_{2i+1} \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with bases $\{a^{2j+1}b^{i-j-1}c \mid 0 \leq j \leq k-1\}$ and $\{a^{2j}b^{i-j}c \mid 0 \leq j \leq k\}$, respectively. This gives that $\text{im } p_{2i}^* \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) and $\text{im } p_{2i+1}^* \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ (k -copies) with bases $\{x^{2j}y^{i-j} \mid 0 \leq j \leq k\}$

and $\{x^{2j+1}y^{i-j} | 0 \leq j \leq k-1\}$, respectively. We observe that $H^j(X) \cong \text{im } p_j^* \oplus \text{im } \rho_j$, $j = 2i, 2i+1$. So, we get

$$H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+1\text{-copies})$$

with bases $\{x^{2j}y^{i-j}, u_{2i}^{2l+1} | 0 \leq j \leq k, 0 \leq l \leq k-1\}$ and $\{x^{2j+1}y^{i-j}, u_{2i+1}^{2l} | 0 \leq j \leq k-1, 0 \leq l \leq k\}$, respectively, where $\rho_{2i}(u_{2i}^{2l+1}) = a^{2l+1}b^{i-l-1}c$, $0 \leq l \leq k-1$ and $\rho_{2i+1}(u_{2i+1}^{2l}) = a^{2l}b^{i-l}c$, $0 \leq l \leq k$.

For $m < i \leq m + \frac{n}{2}$, we have $H^{2i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+2\text{-copies})$ and $H^{2i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+1\text{-copies})$ with bases

$$\{a^{2i-2m-1}b^m c, a^{2i-2m}b^m, a^{2i-2m+1}b^{m-1}c, a^{2i-2m+2}b^{m-1}, \dots, a^{n-1}b^{i-k}c, a^n b^{i-k}\}, \text{ and}$$

$$\{a^{2i-2m}b^m c, a^{2i-2m+1}b^m, a^{2i-2m+2}b^{m-1}c, a^{2i-2m+3}b^{m-1}, \dots, a^{n-1}b^{i-k+1}, a^n b^{i-k}c\},$$

respectively. Note that $\ker p_{2i}^* = \text{im } \rho_{2i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+1\text{-copies})$ and $\ker p_{2i+1}^* = \text{im } \rho_{2i+1} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+1\text{-copies})$ with bases $\{a^{2j+1}b^{i-j-1}c | i-m-1 \leq j \leq k-1\}$ and $\{a^{2j}b^{i-j}c | i-m \leq j \leq k\}$, respectively. This implies that $\text{im } p_{2i}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+1\text{-copies})$ and $\text{im } p_{2i+1}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i\text{-copies})$ with bases $\{x^{2j}y^{i-j} | i-m \leq j \leq k\}$ and $\{x^{2j+1}y^{i-j} | i-m \leq j \leq k-1\}$, respectively. So, we get

$$H^{2i}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+2\text{-copies}), \text{ and}$$

$$H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+1\text{-copies})$$

with bases $\{x^{2j}y^{i-j}, u_{2i}^{2l+1} | i-m \leq j \leq k, i-m-1 \leq l \leq k-1\}$ and $\{x^{2j+1}y^{i-j}, u_{2i+1}^{2l} | i-m \leq j \leq k-1, i-m \leq l \leq k\}$, respectively, where $\rho_{2i}(u_{2i}^{2l+1}) = a^{2l+1}b^{i-l-1}c$, $i-m-1 \leq l \leq k-1$ and $\rho_{2i+1}(u_{2i+1}^{2l}) = a^{2l}b^{i-l}c$, $i-m \leq l \leq k$.

Now, we compute the cohomology algebra of X . Consider the Leray Serre spectral sequence for the Borel fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ with $E_2^{s,t} = H^s(B_G) \otimes H^t(X)$. As $p_j^* = i_j^* \circ h^*$ for all $j \geq 0$, we get $d_r(1 \otimes x) = 0 = d_r(1 \otimes y)$ for all $r \geq 0$ and for all i, j ; $d_{r_{i,j}}(1 \otimes u_j^i) \neq 0$ for some $r_{i,j} > 0$. Put $u_1^0 = z$. So, $H^1(X)$ is generated by $\{x, z\}$. Trivially, $d_2(1 \otimes z) = t^2 \otimes 1$ and so $d_2(1 \otimes xz) \neq 0$. This implies that $xz = \alpha x^2 + \beta y + u_2^1$, $\alpha, \beta \in \mathbb{Z}_2$. So, $H^2(X)$ is generated by $\{x^2, y, xz\}$. By induction, we show that for $1 \leq i < \frac{n}{2}$, $H^{2i-1}(X)$ and $H^{2i}(X)$ are generated by $\{xy^{i-1}, x^3y^{i-2}, \dots, x^{2i-3}y, x^{2i-1}, y^{i-1}z, x^2y^{i-2}z, \dots, x^{2i-4}yz, x^{2i-2}z\}$ and $\{y^i, x^2y^{i-1}, \dots, x^{2i-2}y, x^{2i}, xy^{i-1}z, x^3y^{i-2}z, \dots, x^{2i-1}z\}$, respectively. Now, we prove it for $i+1$. We have $d_2(1 \otimes y^i z) = t^2 \otimes y^i$, $d_2(1 \otimes x^2 y^{i-1} z) = t^2 \otimes x^2 y^{i-1}$, \dots , $d_2(1 \otimes x^{2i-2} y z) = t^2 \otimes x^{2i-2} y$, $d_2(1 \otimes x^{2i} z) = t^2 \otimes x^{2i}$. This gives that $y^i z, x^2 y^{i-1} z, \dots, x^{2i-2} y z, x^{2i} z$ are nonzero

elements of $H^{2i+1}(X)$. Therefore,

$$\begin{aligned}
y^i z &= \alpha_1^0 x y^i + \alpha_3^0 x^3 y^{i-1} + \cdots + \alpha_{2i-1}^0 x^{2i-1} y + \alpha_{2i+1}^0 x^{2i+1} \\
&\quad + \beta_0^0 u_{2i+1}^0 + \beta_2^0 u_{2i+1}^2 + \cdots + \beta_{2i-2}^0 u_{2i+1}^{2i-2} + \beta_{2i}^0 u_{2i+1}^{2i} \\
x^2 y^{i-1} z &= \alpha_1^2 x y^i + \alpha_3^2 x^3 y^{i-1} + \cdots + \alpha_{2i-1}^2 x^{2i-1} y + \alpha_{2i+1}^2 x^{2i+1} \\
&\quad + \beta_0^2 u_{2i+1}^0 + \beta_2^2 u_{2i+1}^2 + \cdots + \beta_{2i-2}^2 u_{2i+1}^{2i-2} + \beta_{2i}^2 u_{2i+1}^{2i} \\
&\quad \vdots \\
x^{2i} z &= \alpha_1^{2i} x y^i + \alpha_3^{2i} x^3 y^{i-1} + \cdots + \alpha_{2i-1}^{2i} x^{2i-1} y + \alpha_{2i+1}^{2i} x^{2i+1} \\
&\quad + \beta_0^{2i} u_{2i+1}^0 + \beta_2^{2i} u_{2i+1}^2 + \cdots + \beta_{2i-2}^{2i} u_{2i+1}^{2i-2} + \beta_{2i}^{2i} u_{2i+1}^{2i}
\end{aligned}$$

where $\beta_j^k \neq 0$ for some $j \in \{0, 2, \dots, 2i\}$ and for all $k = 0, 2, \dots, 2i$. From this we get

$$\begin{aligned}
t^2 \otimes y^i &= \beta_0^0 d_2(1 \otimes u_{2i+1}^0) + \beta_2^0 d_2(1 \otimes u_{2i+1}^2) + \cdots + \beta_{2i}^0 d_2(1 \otimes u_{2i+1}^{2i}) \\
t^2 \otimes x^2 y^{i-1} &= \beta_0^2 d_2(1 \otimes u_{2i+1}^0) + \beta_2^2 d_2(1 \otimes u_{2i+1}^2) + \cdots + \beta_{2i}^2 d_2(1 \otimes u_{2i+1}^{2i}) \\
&\quad \vdots \\
t^2 \otimes x^{2i} &= \beta_0^{2i} d_2(1 \otimes u_{2i+1}^0) + \beta_2^{2i} d_2(1 \otimes u_{2i+1}^2) + \cdots + \beta_{2i}^{2i} d_2(1 \otimes u_{2i+1}^{2i})
\end{aligned}$$

As the set $\{y^i, x^2 y^{i-1}, \dots, x^{2i-2} y, x^{2i}\}$ is linearly independent, the coefficient matrix

$$\begin{bmatrix}
\beta_0^0 & \beta_2^0 & \cdots & \beta_{2i}^0 \\
\beta_0^2 & \beta_2^2 & \cdots & \beta_{2i}^2 \\
\vdots & \vdots & \vdots & \vdots \\
\beta_0^{2i} & \beta_2^{2i} & \cdots & \beta_{2i}^{2i}
\end{bmatrix}$$

is invertible. Then u_{2i+1}^{2k} is generated by $\{x y^i, x^3 y^{i-1}, \dots, x^{2i-1} y, x^{2i+1}, y^i z, x^2 y^{i-1} z, \dots, x^{2i-2} z, x^{2i} z\}$ for all $0 \leq k \leq i$. So, the basis for $H^{2i+1}(X)$ is $\{x y^i, x^3 y^{i-1}, \dots, x^{2i-3} y^2, x^{2i-1} y, x^{2i+1}, y^i z, x^2 y^{i-1} z, \dots, x^{2i-2} y z, x^{2i} z\}$. Also, we have $d_2(1 \otimes x y^i z) = t^2 \otimes x y^i, d_2(1 \otimes x^3 y^{i-1} z) = t^2 \otimes x^3 y^{i-1}, d_2(1 \otimes x^5 y^{i-2} z) = t^2 \otimes x^5 y^{i-2}, \dots, d_2(1 \otimes x^{2i+1} z) = t^2 \otimes x^{2i+1}$.

This implies that $x y^i z, x^3 y^{i-1} z, \dots, x^{2i+1} z$ are nonzero elements of $H^{2i+2}(X)$. So, we get

$$\begin{aligned}
x y^i z &= \alpha_0^1 y^{i+1} + \alpha_2^1 x^2 y^i + \cdots + \alpha_{2i-2}^1 x^{2i-2} y^2 + \alpha_{2i}^1 x^{2i} y \\
&\quad + \beta_1^1 u_{2i+2}^1 + \beta_3^1 u_{2i+2}^3 + \cdots + \beta_{2i+1}^1 u_{2i+2}^{2i+1} \\
x^3 y^{i-1} z &= \alpha_0^3 y^{i+1} + \alpha_2^3 x^2 y^i + \cdots + \alpha_{2i-2}^3 x^{2i-2} y^2 + \alpha_{2i}^3 x^{2i} y \\
&\quad + \beta_1^3 u_{2i+2}^1 + \beta_3^3 u_{2i+2}^3 + \cdots + \beta_{2i+1}^3 u_{2i+2}^{2i+1}
\end{aligned}$$

$$\begin{aligned} & \vdots \\ x^{2i+1}z &= \alpha_0^{2i+1}y^{i+1} + \alpha_2^{2i+1}x^2y^i + \cdots + \alpha_{2i-2}^{2i+1}x^{2i-2}y^2 + \alpha_{2i}^{2i+1}x^{2i}y \\ & \quad + \beta_1^{2i+1}u_{2i+2}^1 + \beta_3^{2i+1}u_{2i+2}^3 + \cdots + \beta_{2i+1}^{2i+1}u_{2i+2}^{2i+1} \end{aligned}$$

where $\beta_j^k \neq 0$ for some $j \in \{1, 3, \dots, 2i+1\}$ and for all $k = 1, 3, \dots, 2i+1$. Thus, we get

$$\begin{aligned} t^2 \otimes xy^i &= \beta_1^1 d_2(1 \otimes u_{2i+2}^1) + \beta_3^1 d_2(1 \otimes u_{2i+2}^3) + \cdots + \beta_{2i+1}^1 d_2(1 \otimes u_{2i+2}^{2i+1}) \\ t^2 \otimes x^3y^{i-1} &= \beta_1^3 d_2(1 \otimes u_{2i+2}^1) + \beta_3^3 d_2(1 \otimes u_{2i+2}^3) + \cdots + \beta_{2i+1}^3 d_2(1 \otimes u_{2i+2}^{2i+1}) \\ & \vdots \\ t^2 \otimes x^{2i+1} &= \beta_1^{2i+1} d_2(1 \otimes u_{2i+2}^1) + \beta_3^{2i+1} d_2(1 \otimes u_{2i+2}^3) + \cdots + \beta_{2i+1}^{2i+1} d_2(1 \otimes u_{2i+2}^{2i+1}) \end{aligned}$$

As the set $\{xy^i, x^3y^{i-1}, \dots, x^{2i+1}\}$ is linearly independent, u_{2i+2}^k is generated by $\{y^{i+1}, x^2y^i, \dots, x^{2i}y, xy^iz, x^3y^{i-1}z, \dots, x^{2i+1}z\}$ for all $k = 1, 3, \dots, 2i+1$. So, our assumption is true for $i+1$. Similarly, for $\frac{n}{2} \leq i \leq m$, bases for $H^{2i}(X)$ and $H^{2i+1}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z \mid 0 \leq j \leq k, 0 \leq l \leq k-1\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z \mid 0 \leq j \leq k-1, 0 \leq l \leq k\}$, respectively, and for $m < i \leq m + \frac{n}{2}$, bases for $H^{2i}(X)$ and $H^{2i+1}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z \mid i-m \leq j \leq k, i-m-1 \leq l \leq k-1\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z \mid i-m \leq j \leq k-1, i-m \leq l \leq k\}$, respectively.

Now, suppose that $n = 2k + 1$ for some k . For $n \leq 2i, 2i+1 \leq 2m$, $H^{n-1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2k+1$ -copies) and $H^{2i}(X/G) \cong H^{2i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2k+2$ -copies) with bases

$$\begin{aligned} & \{b^k, ab^{k-1}c, a^2b^{k-1}, a^3b^{k-2}c, \dots, a^{2k-1}c, a^{2k}\}, \\ & \{b^i, ab^{i-1}c, a^2b^{i-1}, a^3b^{i-2}c, \dots, a^{n-1}b^{i-k}, a^n b^{i-k-1}c\}, \text{ and} \\ & \{b^i c, ab^i, a^2b^{i-1}c, a^3b^{i-1}, \dots, a^{n-1}b^{i-k}c, a^n b^{i-k}\}, \end{aligned}$$

respectively. Then we have $\ker p_{n-1}^* = \text{im } \rho_{n-1} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ (k -copies), $\ker p_{2i}^* = \text{im } \rho_{2i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k+1$ -copies) and $\ker p_{2i+1}^* = \text{im } \rho_{2i+1} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k+1$ -copies) with bases $\{a^{2j+1}b^{k-j-1}c \mid 0 \leq j \leq k-1\}$, $\{a^{2j+1}b^{i-j-1}c \mid 0 \leq j \leq k\}$ and $\{a^{2j}b^{i-j}c \mid 0 \leq j \leq k\}$, respectively. So, $\text{im } p_{n-1}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k+1$ -copies), $\text{im } p_{2i}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k+1$ copies) and $\text{im } p_{2i+1}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k+1$ copies) with bases $\{x^{2j}y^{k-j} \mid 0 \leq j \leq k\}$, $\{x^{2j}y^{i-j} \mid 0 \leq j \leq k\}$ and $\{x^{2j+1}y^{i-j} \mid 0 \leq i \leq k\}$, respectively. As we have done above, we get

$$H^{n-1}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 \text{ (} n\text{-copies), and}$$

$$H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+1\text{-copies})$$

with bases $\{x^{2j}y^{k-j}, u_{2k}^{2l+1} | 0 \leq j \leq k, 0 \leq l \leq k-1\}$, $\{x^{2j}y^{i-j}, u_{2i}^{2j+1} | 0 \leq j \leq k\}$ and $\{x^{2j+1}y^{i-j}, u_{2i+1}^{2j} | 0 \leq j \leq k\}$, respectively, where $\rho_{2k}(u_{2k}^{2l+1}) = a^{2l+1}b^{k-l-1}c, 0 \leq l \leq k-1$, $\rho_{2i}(u_{2i}^{2j+1}) = a^{2j+1}b^{i-j-1}c, 0 \leq j \leq k$ and $\rho_{2i+1}(u_{2i+1}^{2j}) = a^{2j}b^{i-j}c, 0 \leq j \leq k$. Also, for $2m < 2i, 2i+1 \leq 2m+n+1$, we have $H^{2i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+2\text{-copies})$ and $H^{2i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+1\text{-copies})$ with bases

$$\{a^{2i-2m-1}b^m c, a^{2i-2m}b^m, a^{2i-2m+1}b^{m-1}c, a^{2i-2m+2}b^{m-1}, \dots, a^{n-1}b^{i-k}, a^n b^{i-k-1}c\}, \text{ and}$$

$$\{a^{2i-2m}b^m c, a^{2i-2m+1}b^m, a^{2i-2m+2}b^{m-1}c, a^{2i-2m+3}b^{m-1}, \dots, a^{n-1}b^{i-k}c, a^n b^{i-k}\},$$

respectively. Note that $\ker p_{2i}^* = \text{im } \rho_{2i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+2\text{-copies})$ and $\ker p_{2i+1}^* = \text{im } \rho_{2i+1} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+1\text{-copies})$ with bases $\{a^{2j+1}b^{i-j-1}c | i-m-1 \leq j \leq k\}$ and $\{a^{2j}b^{i-j}c | i-m \leq j \leq k\}$, respectively. This implies that $\text{im } p_{2i}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+1\text{-copies})$ and $\text{im } p_{2i+1}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (k+m-i+1\text{-copies})$ with bases $\{x^{2j}y^{i-j} | i-m \leq j \leq k\}$ and $\{x^{2j+1}y^{i-j} | i-m \leq j \leq k\}$, respectively. So, we get

$$H^{2i}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+2\text{-copies}), \text{ and}$$

$$H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n+2m-2i+1\text{-copies})$$

with bases $\{x^{2j}y^{i-j}, u_{2i}^{2l+1} | i-m \leq j \leq k, i-m-1 \leq l \leq k\}$ and $\{x^{2j+1}y^{i-j}, u_{2i+1}^{2j} | i-m \leq j \leq k\}$, respectively, where $\rho_{2i}(u_{2i}^{2l+1}) = a^{2l+1}b^{i-l-1}c, i-m-1 \leq l \leq k$ and $\rho_{2i+1}(u_{2i+1}^{2j}) = a^{2j}b^{i-j}c, i-m \leq j \leq k$.

For $1 \leq i < n$, $H^i(X)$ has the same basis as in the case when n is even. Again by induction, for $n \leq 2i, 2i+1 \leq 2m$, bases for $H^{n-1}(X)$, $H^{2i}(X)$ and $H^{2i+1}(X)$ are $\{x^{2j}y^{k-j}, x^{2l+1}y^{k-l-1}z | 0 \leq j \leq k, 0 \leq l \leq k-1\}$, $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z | 0 \leq j, l \leq k\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z | 0 \leq j, l \leq k\}$, respectively, and for $m < i \leq m + \frac{n}{2}$, bases for $H^{2i}(X)$ and $H^{2i+1}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z | i-m \leq j \leq k, i-m-1 \leq l \leq k\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z | i-m \leq j, l \leq k\}$, respectively.

As $d_2(1 \otimes z^2) = 0$, we get $z^2 = \alpha x^2 + \beta y$ for some $\alpha, \beta \in \mathbb{Z}_2$. Thus the cohomology algebra of X is given by

$$\mathbb{Z}_2[x, y, z] / \langle x^{n+1}, y^{m+1}, z^2 + \alpha x^2 + \beta y \rangle,$$

where $\deg x = \deg z = 1$, $\deg y = 2$ and $\alpha, \beta \in \mathbb{Z}_2$. This realizes possibility (i).

Case (2): When $\cup : H^0(X/G) \rightarrow H^1(X/G)$ maps 1 to a .

Clearly, $\rho_i : H^i(X) \rightarrow H^i(X/G)$ is trivial for all $1 \leq i < n$. For $0 < 2i, 2i+1 < n$, bases for $\ker p_{2i}^*$ and $\ker p_{2i+1}^*$ are $\{a^{2j+1}b^{i-j-1}c, a^{2l}b^{i-l} | 0 \leq j \leq i-1, 1 \leq l \leq i\}$

and $\{a^{2j+1}b^{i-j}, a^{2l}b^{i-l}c \mid 0 \leq j \leq i, 1 \leq l \leq i\}$, respectively. This implies that $\text{im } p_{2i}^* \cong \text{im } p_{2i+1}^* \cong \mathbb{Z}_2$ with bases $\{y^i\}$ and $\{xy^i\}$, respectively, where $p_1^*(c) = x$ and $p_2^*(b) = y$. So, $H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2$ with bases $\{y^i\}$ and $\{xy^i\}$, respectively.

If $n = 2k$ for some k , then for $n \leq 2i, 2i + 1 \leq 2m + 1$, $\{a^{2j+1}b^{i-j-1}c, a^{2l}b^{i-l} \mid 0 \leq j \leq k-1, 1 \leq l \leq k\}$ and $\{a^{2j+1}b^{i-j}, a^{2l}b^{i-l}c \mid 0 \leq j \leq k-1, 1 \leq l \leq k\}$ are bases for $\ker p_{2i}^*$ and $\ker p_{2i+1}^*$, respectively. This gives that $\text{im } p_{2i}^* \cong \text{im } p_{2i+1}^* \cong \mathbb{Z}_2$ with bases $\{y^i\}$ and $\{xy^i\}$, respectively, and $\text{im } \rho_{2i} \cong \text{im } \rho_{2i+1} \cong \mathbb{Z}_2$ with bases $\{a^n b^{i-k}\}$ and $\{a^n b^{i-k}c\}$, respectively. Consequently, $H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ with bases $\{y^i, u_{2i}^n\}$ and $\{xy^i, u_{2i+1}^n\}$, respectively, where $\rho_{2i}(u_{2i}^n) = a^n b^{i-k}$ and $\rho_{2i+1}(u_{2i+1}^n) = a^n b^{i-k}c$. If $2m + 1 < 2i, 2i + 1 \leq 2m + n + 1$ then bases for $\ker p_{2i}^*$ and $\ker p_{2i+1}^*$ are $\{a^{2j+1}b^{i-j-1}c, a^{2l}b^{i-l} \mid i-m-1 \leq j \leq k-1, i-m \leq l \leq k\}$ and $\{a^{2j+1}b^{i-j}, a^{2l}b^{i-l}c \mid i-m \leq j \leq k-1, i-m \leq l \leq k\}$, respectively. This implies that p_{2i}^*, p_{2i+1}^* are trivial and $\text{im } \rho_{2i} \cong \text{im } \rho_{2i+1} \cong \mathbb{Z}_2$ with bases $\{a^n b^{i-k}\}$ and $\{a^n b^{i-k}c\}$, respectively. Thus, $H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2$ with bases $\{u_{2i}^n\}$ and $\{u_{2i+1}^n\}$ respectively, where $\rho_{2i}(u_{2i}^n) = a^n b^{i-k}$ and $\rho_{2i+1}(u_{2i+1}^n) = a^n b^{i-k}c$. Trivially, $x^2 = y^{m+1} = 0$. Put $u_{2k}^n = z$. Then by the edge homomorphism, $d_r(1 \otimes x) = d_r(1 \otimes y) = 0$ for all $r > 0$ and $d_{r'}(1 \otimes z) \neq 0$ for some $r' > 0$. As $X/G \sim \mathbb{RP}^1 \times \mathbb{RP}^n \times \mathbb{CP}^m$, r' must be $n + 1$. Note that for all $0 \leq i \leq m$, $xy^i z$ and xy^i are nonzero elements of $H^{n+2i+1}(X)$ and $H^{n+2i}(X)$, respectively. Therefore, $xy^i z = \alpha xy^{k+m} + \beta u_{n+2i+1}^n$ and $xy^i = \alpha' y^{k+m} + \beta' u_{n+2i}^n$ for some $\alpha, \alpha', \beta, \beta' \in \mathbb{Z}_2$, where β and β' are nonzero. Note that for $n+2i > 2m$, we get $\alpha = \alpha' = 0$.

As $d_r(1 \otimes z^2) = 0$ for all $r \geq 0$, we get $z^2 = \alpha y^n$ for some $\alpha \in \mathbb{Z}_2$ and $\alpha = 0$ for $n > m$. Thus, the cohomology algebra of X is given by

$$\mathbb{Z}_2[x, y, z] / \langle x^2, y^{m+1}, z^2 + \alpha y^n \rangle,$$

where $\deg x = 1, \deg y = 2, \deg z = n$ and $\alpha = 0$ if $n > m$. This realizes possibility (ii).

Now, let $n = 2k + 1$ for some $k \geq 0$. For $n \leq 2i, 2i + 1 \leq 2m + 1$, we get $H^{2i}(X) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \cong H^{2i+1}(X)$ with bases $\{y^i, u_{2i}^n\}$ and $\{xy^i, u_{2i+1}^n\}$, respectively, where $\rho_{2i}(u_{2i}^n) = a^n b^{i-k-1}c$ and $\rho_{2i+1}(u_{2i+1}^n) = a^n b^{i-k}$. Also, for $2m + 1 < 2i, 2i + 1 \leq 2m + n + 1$, we have $H^{2i}(X) \cong \mathbb{Z}_2 \cong H^{2i+1}(X)$ with bases $\{u_{2i}^n\}$ and $\{u_{2i+1}^n\}$, respectively, where $\rho_{2i}(u_{2i}^n) = a^n b^{i-k-1}c$ and $\rho_{2i+1}(u_{2i+1}^n) = a^n b^{i-k}$. It is easy to observe that X has the same cohomology algebra as in the case when n is even.

Case (3): When $\cup : H^0(X/G) \rightarrow H^4(X/G)$ maps 1 to $a + c$.

Also, in this case $\rho_i : H^i(X) \rightarrow H^i(X/G)$ and p_j^* are trivial for all $1 \leq i < n$ and $2m + 1 < j \leq 2m + n + 1$. Similar calculations show that for $1 \leq i < n$, $H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2$ with bases $\{y^i\}$ and $\{xy^i\}$, respectively, where $x = p_1^*(a) = p_1^*(c), y = p_2^*(b)$. For $n \leq 2i, 2i +$

$1 \leq 2m+1$, $H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ with bases $\{y^i, u_{2i}\}$, $\{xy^i, u_{2i+1}\}$, respectively; and for $2m+1 < 2i, 2i+1 \leq 2m+n+1$, we have $H^{2i}(X) \cong H^{2i+1}(X) \cong \mathbb{Z}_2$ with bases $\{u_{2i}\}$, $\{u_{2i+1}\}$, respectively, where $\rho_{2i}(u_{2i}) = a^{n-1}b^{i-k}c + a^nb^{i-k}$ and $\rho_{2i+1}(u_{2i+1}) = a^nb^{i-k}c$ if $n = 2k$; and $\rho_{2i}(u_{2i}) = a^nb^{i-k-1}c$ and $\rho_{2i+1}(u_{2i+1}) = a^{n-1}b^{i-k}c + a^nb^{i-k}$ if $n = 2k + 1$. Consequently, the cohomology algebra of X is given by

$$\mathbb{Z}_2[x, y, z]/\langle x^2, y^{m+1}, z^2 + \alpha y^n \rangle,$$

where $\deg x = 1, \deg y = 2, \deg z = n$ and $\alpha = 0$ if $n > m$. This realizes possibility (ii).

Similarly, we get the same cohomology algebra for the case when $2m \leq n$. \square

Proof of Theorem 1.4. As $X/G \sim_2 \mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m \times \mathbb{R}\mathbb{P}^2$, we get $H^*(X/G) = \mathbb{Z}_2[a, b, c]/\langle a^{n+1}, b^{m+1}, c^3 \rangle$, where $\deg a = 2, \deg b = 4$ and $\deg c = 1$. Since X is connected, $H^0(X) \cong \mathbb{Z}_2$. By Proposition 2.3, $H^i(X) = 0$ for all $i > 4m + 2n + 2$. Firstly, assume that $1 \leq n < 2m$. Consider the Gysin sequence of the sphere bundle $G \hookrightarrow X \xrightarrow{p} X/G$:

$$\cdots \longrightarrow H^{i-1}(X/G) \xrightarrow{\cup} H^i(X/G) \xrightarrow{p_i^*} H^i(X) \xrightarrow{\rho_i} H^i(X/G) \xrightarrow{\cup} H^{i+1}(X/G) \longrightarrow \cdots$$

which begins with

$$0 \longrightarrow H^0(X/G) \xrightarrow{p_0^*} H^0(X) \xrightarrow{\rho_0} H^0(X/G) \xrightarrow{\cup} H^1(X/G) \xrightarrow{p_1^*} H^1(X) \longrightarrow \cdots$$

By the exactness of Gysin sequence, the Euler class of the associated principal G -bundle $G \hookrightarrow X \rightarrow X/G$ must be c .

For $0 \leq i < \frac{n}{2}$, we have $H^{4i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2i + 1$ -copies), $H^{4i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($i + 1$ -copies), $H^{4i+2}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2i + 2$ -copies) and $H^{4i+3}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($i + 1$ -copies) with bases

$$\begin{aligned} &\{b^i, ab^{i-1}c^2, a^2b^{i-1}, a^3b^{i-2}c^2, \dots, a^{2i-1}c^2, a^{2i}\}, \\ &\{b^i c, a^2b^{i-1}c, \dots, a^{2i-2}bc, a^{2i}c\}, \\ &\{b^i c^2, ab^i, a^2b^{i-1}c^2, a^3b^{i-1}, \dots, a^{2i}c^2, a^{2i+1}\}, \text{ and} \\ &\{ab^i c, a^3b^{i-1}c, \dots, a^{2i-1}bc, a^{2i+1}c\}, \end{aligned}$$

respectively. Again, by the exactness of Gysin sequence, we get $\ker p_{4i}^* = \text{im } \rho_{4i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ (i -copies) with basis $\{a^{2j+1}b^{i-j-1}c^2 \mid 0 \leq j \leq i-1\}$, $\ker p_{4i+1}^* = H^{4i+1}(X/G)$, $\ker p_{4i+2}^* = \text{im } \rho_{4i+2} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($i + 1$ -copies) with basis $\{a^{2j}b^{i-j}c^2 \mid 0 \leq j \leq i\}$ and $\ker p_{4i+3}^* = H^{4i+3}(X/G)$. This implies that $\text{im } p_{4i}^* \cong \text{im } p_{4i+2}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($i + 1$ -copies) with bases $\{x^{2j}y^{i-j} \mid 0 \leq j \leq i\}$ and $\{x^{2j+1}y^{i-j} \mid 0 \leq j \leq i\}$, respectively, where

$p_2^*(a) = x, p_4^*(b) = y$. Clearly, $\text{im } p_{4i+j}^* = \text{im } \rho_{4i+j} = 0, j = 1, 3$. Thus,

$$\begin{aligned} H^{4i+1}(X) &= H^{4i+3}(X) = 0 \\ H^{4i}(X) &\cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 \text{ (} 2i + 1\text{-copies), and} \\ H^{4i+2}(X) &\cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 \text{ (} 2i + 2\text{-copies)} \end{aligned}$$

with bases $\{x^{2j}y^{i-j}, u_{4i}^{2l+1} | 0 \leq j \leq i, 0 \leq l \leq i-1\}$ and $\{x^{2j+1}y^{i-j}, u_{4i+2}^{2j} | 0 \leq j \leq i\}$, respectively, where $\rho_{4i}(u_{4i}^{2l+1}) = a^{2l+1}b^{i-l-1}c^2, 0 \leq l \leq i-1$ and $\rho_{4i+2}(u_{4i+2}^{2j}) = a^{2j}b^{i-j}c^2, 0 \leq j \leq i$.

Let us assume that $n = 2k$ for some $k > 0$. In this case, for $2n \leq 4i+r \leq 4m, 0 \leq r \leq 3$, we have $H^{4i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2k + 1$ -copies), $H^{4i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies), $H^{4i+2}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2k + 1$ -copies) and $H^{4i+3}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ (k -copies) with bases

$$\begin{aligned} &\{b^i, ab^{i-1}c^2, a^2b^{i-1}, a^3b^{i-2}c^2, \dots, a^{2k-1}b^{i-k}c^2, a^{2k}b^{i-k}\}, \\ &\{b^i c, a^2b^{i-1}c, \dots, a^{2k-2}b^{i-k+1}c, a^{2k}b^{i-k}c\}, \\ &\{b^i c^2, ab^i, a^2b^{i-1}c^2, a^3b^{i-1}, \dots, a^{2k-1}b^{i-k+1}, a^{2k}b^{i-k}c^2\}, \text{ and} \\ &\{ab^i c, a^3b^{i-1}c, \dots, a^{2k-3}b^{i-k+2}c, a^{2k-1}b^{i-k+1}c\}, \end{aligned}$$

respectively. Then by the exactness of Gysin sequence, we get $\ker p_{4i}^* = \text{im } \rho_{4i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ (k -copies) with basis $\{a^{2j+1}b^{i-j-1}c^2 | 0 \leq j \leq k-1\}$, $\ker p_{4i+2}^* = \text{im } \rho_{4i+2} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with basis $\{a^{2j}b^{i-j}c^2 | 0 \leq j \leq k\}$ and $\ker p_{4i+j}^* = H^{4i+j}(X/G), j = 1, 3$. This gives that $\text{im } p_{4i}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with basis $\{x^{2j}y^{i-j} | 0 \leq j \leq k\}$ and $\text{im } p_{4i+2}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ (k -copies) with basis $\{x^{2j+1}y^{i-j} | 0 \leq j \leq k-1\}$. Also, $\text{im } p_{4i+j}^* = \text{im } \rho_{4i+j} = 0, j = 1$ and 3 , Therefore,

$$\begin{aligned} H^{4i+1}(X) &= H^{4i+3}(X) = 0 \\ H^{4i}(X) &\cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 \text{ (} n + 1\text{-copies), and} \\ H^{4i+2}(X) &\cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 \text{ (} n + 1\text{-copies)} \end{aligned}$$

with bases $\{x^{2j}y^{i-j}, u_{4i}^{2l+1} | 0 \leq j \leq k, 0 \leq l \leq k-1\}$ and $\{x^{2j+1}y^{i-j}, u_{4i+2}^{2l} | 0 \leq j \leq k-1, 0 \leq l \leq k\}$, respectively, where $\rho_{4i}(u_{4i}^{2l+1}) = a^{2l+1}b^{i-l-1}c^2, 0 \leq l \leq k-1$ and $\rho_{4i+2}(u_{4i+2}^{2l}) = a^{2l}b^{i-l}c^2, 0 \leq l \leq k$. Now, for $4m < 4i+r \leq 4m+2n+2, 0 \leq r \leq 3$; we have $H^{4i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2m+n-2i+2$ -copies), $H^{4i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($m+k-i+1$ -copies), $H^{4i+2}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2m+n-2i+1$ -copies) and $H^{4i+3}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$

$(m + k - i)$ -copies) with bases

$$\begin{aligned} & \{a^{2i-2m-1}b^m c^2, a^{2i-2m}b^m, a^{2i-2m+1}b^{m-1}c^2, a^{2i-2m+2}b^{m-1}, \dots, a^{2k-1}b^{i-k}c^2, a^{2k}b^{i-k}\}, \\ & \{a^{2i-2m}b^m c, a^{2i-2m+2}b^{m-1}c, \dots, a^{2k-2}b^{i-k+1}c, a^{2k}b^{i-k}c\}, \\ & \{a^{2i-2m}b^m c^2, a^{2i-2m+1}b^m, a^{2i-2m+2}b^{m-1}c^2, a^{2i-2m+3}b^{m-1}, \dots, a^{2k-1}b^{i-k+1}, a^{2k}b^{i-k}c^2\}, \text{ and} \\ & \{a^{2i-2m+1}b^m c, a^{2i-2m+3}b^{m-1}c, \dots, a^{2k-3}b^{i-k+2}c, a^{2k-1}b^{i-k+1}c\}, \end{aligned}$$

respectively. Then by the exactness of Gysin sequence, we get $\ker p_{4i}^* = \text{im } \rho_{4i} \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies) with basis $\{a^{2j+1}b^{i-j-1}c^2 | i - m - 1 \leq j \leq k - 1\}$, $\ker p_{4i+j}^* = H^{4i+j}(X/G)$, $j = 1, 3$; and $\ker p_{4i+2}^* = \text{im } \rho_{4i+2} \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies) with basis $\{a^{2j}b^{i-j}c^2 | i - m \leq j \leq k\}$. This gives that $\text{im } p_{4i}^* \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies) with basis $\{x^{2j}y^{i-j} | i - m \leq j \leq k\}$, $\text{im } p_{4i+j}^* = \text{im } \rho_{4i+j} = 0$, $j = 1, 3$; and $\text{im } p_{4i+2}^* \cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2$ ($m + k - i$ -copies) with basis $\{x^{2j+1}y^{i-j} | i - m \leq j \leq k - 1\}$. Therefore,

$$\begin{aligned} H^{4i+1}(X) &= H^{4i+3}(X) = 0 \\ H^{4i}(X) &\cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2 (2m + n - 2i + 2\text{-copies}), \text{ and} \\ H^{4i+2}(X) &\cong \mathbb{Z}_2 \oplus \dots \oplus \mathbb{Z}_2 (2m + n - 2i + 1\text{-copies}) \end{aligned}$$

with bases $\{x^{2j}y^{i-j}, u_{4i}^{2l+1} | i - m \leq j \leq k, i - m - 1 \leq l \leq k - 1\}$ and $\{x^{2j+1}y^{i-j}, u_{4i+2}^{2l} | i - m \leq j \leq k - 1, i - m \leq l \leq k\}$, respectively, where $\rho_{4i}(u_{4i}^{2l+1}) = a^{2l+1}b^{i-l-1}c^2$, $i - m - 1 \leq l \leq k - 1$ and $\rho_{4i+2}(u_{4i+2}^{2l}) = a^{2l}b^{i-l}c^2$, $i - m \leq l \leq k$.

Next, we compute the cohomology algebra of X . Consider the Leray-Serre spectral sequence for the Borel fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$. By the edge homomorphisms, we get $d_r(1 \otimes x) = 0 = d_r(1 \otimes y)$ for all $r > 0$ and for all i, j ; $d_{r_{i,j}}(1 \otimes u_j^i) \neq 0$ for some $r_{i,j} > 0$. Put $u_2^0 = z$. So, $H^2(X)$ is generated by $\{x, z\}$. Clearly, $d_3(1 \otimes z) = t^3 \otimes 1$ and so $d_3(1 \otimes xz) = t^3 \otimes x \neq 0$. This implies that $xz = \alpha x^2 + \beta y + u_4^1$, $\alpha, \beta \in \mathbb{Z}_2$. So, $H^4(X)$ is generated by $\{x^2, y, xz\}$. By the induction, as in Theorem 1.3, we get that for $1 \leq 4i + r < 2n$, $r = 0, 2$, $H^{4i}(X)$ and $H^{4i+2}(X)$ have bases $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z | 0 \leq j \leq i, 0 \leq l \leq i - 1\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z | 0 \leq j, l \leq i\}$, respectively. Similarly, for $2n \leq 4i + r \leq 4m$, $r = 0, 2$, bases for $H^{4i}(X)$ and $H^{4i+2}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z | 0 \leq j \leq k, 0 \leq l \leq k - 1\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z | 0 \leq j \leq k - 1, 0 \leq l \leq k\}$, respectively, and for $4m < 4i + r \leq 4m + 2n + 2$, $r = 0, 2$, bases for $H^{4i}(X)$ and $H^{4i+2}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z | i - m \leq j \leq k, i - m - 1 \leq l \leq k - 1\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z | i - m \leq j \leq k - 1, i - m \leq l \leq k\}$, respectively.

Now, we take the case when $n = 2k + 1$ for some $k > 0$. In this case, for $2n \leq 4i + r \leq 4m, 0 \leq r \leq 3$, we have $H^{4i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2k + 2$ -copies), $H^{4i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies), $H^{4i+2}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2k + 2$ -copies) and $H^{4i+3}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with bases

$$\begin{aligned} & \{b^i, ab^{i-1}c^2, a^2b^{i-1}, a^3b^{i-2}c^2, \dots, a^{2k}b^{i-k}, a^{2k+1}b^{i-k-1}c^2\}, \\ & \{b^i c, a^2b^{i-1}c, \dots, a^{2k-2}b^{i-k+1}c, a^{2k}b^{i-k}c\}, \\ & \{b^i c^2, ab^i, a^2b^{i-1}c^2, a^3b^{i-1}, \dots, a^{2k}b^{i-k}c^2, a^{2k+1}b^{i-k}\}, \text{ and} \\ & \{ab^i c, a^3b^{i-1}c, \dots, a^{2k-1}b^{i-k+1}c, a^{2k+1}b^{i-k}c\}, \end{aligned}$$

respectively. Again, by the exactness of Gysin sequence, we get $\ker p_{4i}^* = \text{im } \rho_{4i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with basis $\{a^{2j+1}b^{i-j-1}c^2 | 0 \leq j \leq k\}$, $\ker p_{4i+2}^* = \text{im } \rho_{4i+2} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with basis $\{a^{2j}b^{i-j}c^2 | 0 \leq j \leq k\}$ and $\ker p_{4i+j}^* = H^{4i+j}(X/G), j = 1, 3$. This gives that $\text{im } p_{4i}^* \cong \text{im } p_{4i+2}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($k + 1$ -copies) with bases $\{x^{2j}y^{i-j} | 0 \leq j \leq k\}$ and $\{x^{2j+1}y^{i-j} | 0 \leq j \leq k\}$, respectively. Note that $\text{im } p_{4i+j}^* = \text{im } \rho_{4i+j} = 0, j = 1$ and 3 . Thus,

$$\begin{aligned} H^{4i+1}(X) &= H^{4i+3}(X) = 0 \\ H^{4i}(X) &\cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n + 1\text{-copies}), \text{ and} \\ H^{4i+2}(X) &\cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (n + 1\text{-copies}) \end{aligned}$$

with bases $\{x^{2j}y^{i-j}, u_{4i}^{2l+1} | 0 \leq j, l \leq k\}$ and $\{x^{2j+1}y^{i-j}, u_{4i+2}^{2l} | 0 \leq j, l \leq k\}$, respectively, where $\rho_{4i}(u_{4i}^{2l+1}) = a^{2l+1}b^{i-l-1}c^2, 0 \leq l \leq k$ and $\rho_{4i+2}(u_{4i+2}^{2l}) = a^{2l}b^{i-l}c^2, 0 \leq l \leq k$. Now, for $4m < 4i + r \leq 4m + 2n + 2, 0 \leq r \leq 3$; we have $H^{4i}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2m + n - 2i + 2$ -copies), $H^{4i+1}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies), $H^{4i+2}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($2m + n - 2i + 1$ -copies) and $H^{4i+3}(X/G) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies) with bases

$$\begin{aligned} & \{a^{2i-2m-1}b^m c^2, a^{2i-2m}b^m, a^{2i-2m+1}b^{m-1}c^2, a^{2i-2m+2}b^{m-1}, \dots, a^{2k}b^{i-k}, a^{2k+1}b^{i-k-1}c^2\}, \\ & \{a^{2i-2m}b^m c, a^{2i-2m+2}b^{m-1}c, \dots, a^{2k-2}b^{i-k+1}c, a^{2k}b^{i-k}c\}, \\ & \{a^{2i-2m}b^m c^2, a^{2i-2m+1}b^m, a^{2i-2m+2}b^{m-1}c^2, a^{2i-2m+3}b^{m-1}, \dots, a^{2k}b^{i-k}c^2, a^{2k+1}b^{i-k}\}, \text{ and} \\ & \{a^{2i-2m+1}b^m c, a^{2i-2m+3}b^{m-1}c, \dots, a^{2k-1}b^{i-k+1}c, a^{2k+1}b^{i-k}c\}, \end{aligned}$$

respectively. Now, $\ker p_{4i}^* = \text{im } \rho_{4i} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($m + k - i + 2$ -copies) with basis $\{a^{2j+1}b^{i-j-1}c^2 | i - m - 1 \leq j \leq k\}$, $\ker p_{4i+j}^* = H^{4i+j}(X/G), j = 1, 3$; and $\ker p_{4i+2}^* = \text{im } \rho_{4i+2} \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies) with basis $\{a^{2j}b^{i-j}c^2 | i - m \leq j \leq k\}$. Consequently, $\text{im } p_{4i}^* \cong \text{im } p_{4i+2}^* \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2$ ($m + k - i + 1$ -copies) with bases

$\{x^{2j}y^{i-j}|i-m \leq j \leq k\}$ and $\{x^{2j+1}y^{i-j}|i-m \leq j \leq k\}$, respectively. Also, $\text{im } p_{4i+j}^* = \text{im } \rho_{4i+j} = 0, j = 1, 3$. Therefore,

$$H^{4i+1}(X) = H^{4i+3}(X) = 0$$

$$H^{4i}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (2m+n-2i+2\text{-copies}), \text{ and}$$

$$H^{4i+2}(X) \cong \mathbb{Z}_2 \oplus \cdots \oplus \mathbb{Z}_2 (2m+n-2i+1\text{-copies})$$

with bases $\{x^{2j}y^{i-j}, u_{4i}^{2l+1}|i-m \leq j \leq k, i-m-1 \leq l \leq k\}$ and $\{x^{2j+1}y^{i-j}, u_{4i+2}^{2l}|i-m \leq j, l \leq k\}$, respectively, where $\rho_{4i}(u_{4i}^{2l+1}) = a^{2l+1}b^{i-l-1}c^2, i-m-1 \leq l \leq k$ and $\rho_{4i+2}(u_{4i+2}^{2l}) = a^{2l}b^{i-l}c^2, i-m \leq l \leq k$. Now, we discuss the cohomology algebra of X . By the repeated use of induction, for $\frac{n}{2} \leq i \leq m$, bases for $H^{4i}(X)$ and $H^{4i+2}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z|0 \leq j \leq k, 0 \leq l \leq k\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z|0 \leq j \leq k, 0 \leq l \leq k\}$, respectively, and for $m < i \leq m + \frac{n}{2}$, bases for $H^{4i}(X)$ and $H^{4i+2}(X)$ are $\{x^{2j}y^{i-j}, x^{2l+1}y^{i-l-1}z|i-m \leq j \leq k, i-m-1 \leq l \leq k\}$ and $\{x^{2j+1}y^{i-j}, x^{2l}y^{i-l}z|i-m \leq j \leq k, i-m \leq l \leq k\}$, respectively.

Clearly, $x^{n+1} = y^{m+1} = 0$. As $d_2(1 \otimes z^2) = 0, y^2 = \alpha x^2 + \beta y$ for some $\alpha, \beta \in \mathbb{Z}_2$. Thus, the cohomology algebra of X is

$$\mathbb{Z}_2[x, y, z]/\langle x^{n+1}, y^{m+1}, z^2 + \alpha x^2 + \beta y \rangle,$$

where $\deg x = 2, \deg y = 4, \deg z = 2$ and $\alpha, \beta \in \mathbb{Z}_2$.

Similarly, we get the same cohomology algebra for the case when $n \geq 2m$. □

Now, we realize the possibilities of our main Theorems.

Example 3.1. Consider, the antipodal action of $G = \mathbb{Z}_2$ on \mathbb{S}^1 and trivial action of G on $\mathbb{C}\mathbb{P}^n$. Then the diagonal action of G on $X = \mathbb{S}^1 \times \mathbb{C}\mathbb{P}^n$ is a free action with the orbit space $X/G = \mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^n$. Also, the map $[x_0, x_1, \dots, x_{2n}, x_{2n+1}] \rightarrow [-x_1, x_0 \cdots, -x_{2n+1}, x_{2n}]$ gives a free action of G on $\mathbb{R}\mathbb{P}^{2n+1}$ with the orbit space $X/G \sim_2 \mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^n$ [7]. These examples realizes both possibilities of Theorem 1.1.

Example 3.2. Consider, the antipodal action of $G = \mathbb{Z}_2$ on \mathbb{S}^2 and trivial action of G on $\mathbb{H}\mathbb{P}^n$. Then the diagonal action of G on $X = \mathbb{S}^2 \times \mathbb{H}\mathbb{P}^n$ is a free action with the orbit space $X/G = \mathbb{R}\mathbb{P}^2 \times \mathbb{H}\mathbb{P}^n$. Also, the map $[z_0, z_1, \dots, z_{n-1}, z_n] \rightarrow [-\bar{z}_1, \bar{z}_0 \cdots, -\bar{z}_n, \bar{z}_{n-1}]$ gives a free action of G on $\mathbb{C}\mathbb{P}^{2n+1}$ with the orbit space $X/G \sim_2 \mathbb{R}\mathbb{P}^2 \times \mathbb{H}\mathbb{P}^n$ [7]. These examples realizes both possibilities of Theorem 1.2.

Example 3.3. Consider, the antipodal action of $G = \mathbb{Z}_2$ on \mathbb{S}^1 and trivial action on $\mathbb{R}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^m$. Then the diagonal action of G on $\mathbb{S}^1 \times \mathbb{R}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^m$ is a free action with

the orbit space $\mathbb{R}\mathbb{P}^1 \times \mathbb{R}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^m$. This realizes possibility (i) of Theorem 1.3 when $\alpha = \beta = 0$. Now, we consider trival action of G on $\mathbb{R}\mathbb{P}^n$ and free action of G on $\mathbb{R}\mathbb{P}^{2m+1}$ as defined in Example 3.1. Then G acts freely on $X = \mathbb{R}\mathbb{P}^n \times \mathbb{R}\mathbb{P}^{2m+1}$ with the orbit space $X/G \sim_2 \mathbb{R}\mathbb{P}^n \times \mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^m$. This realizes possibility (i) of Theorem 1.3 when $\alpha = 0, \beta = 1$. Also, if we take antipodal action of G on \mathbb{S}^n and trivial action on $\mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^m$ then G acts freely on $\mathbb{S}^n \times \mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^m$ with the orbit space $\mathbb{R}\mathbb{P}^n \times \mathbb{R}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^m$. This realizes possibility (ii) of Theorem 1.3 when $\alpha = 0$.

Example 3.4. Consider, the antipodal action of $G = \mathbb{Z}_2$ on \mathbb{S}^2 and trivial action on $\mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m$. Then the diagonal action of G on $\mathbb{S}^2 \times \mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m$ is a free action with the orbit space $\mathbb{R}\mathbb{P}^2 \times \mathbb{C}\mathbb{P}^n \times \mathbb{H}\mathbb{P}^m$. This realizes Theorem 1.4 when $\alpha = \beta = 0$. Take trival action of G on $\mathbb{C}\mathbb{P}^n$ and free action of G on $\mathbb{C}\mathbb{P}^{2m+1}$ as defined in Example 3.2. Then G acts freely on $X = \mathbb{C}\mathbb{P}^n \times \mathbb{C}\mathbb{P}^{2m+1}$ with the orbit space $X/G \sim_2 \mathbb{C}\mathbb{P}^n \times \mathbb{R}\mathbb{P}^2 \times \mathbb{H}\mathbb{P}^m$. This realizes Theorem 1.4 when $\alpha = 0, \beta = 1$.

ACKNOWLEDGEMENT

We are thankful to the referee for their suggestions to improve the paper, correcting typos and grammatical errors.

REFERENCES

- [1] T. Tom Dieck, *Transformation groups*, De Gruyter Studies in Mathematics, 8, Walter de Gruyter & Co., Berlin, 1987.
- [2] J. Harvey, M. Kerin and K. Shankar, *Semi-free actions with manifold orbit spaces*, Doc. Math. **25** (2020), 2085–21214.
- [3] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
- [4] A. Kumari and H. K. Singh, *Cohomology classification of spaces with free \mathbb{S}^1 and \mathbb{S}^3 -actions*, Filomat **36** (2022), no. 20, 7021–7026
- [5] J. McCleary, *A user's guide to spectral sequences*, second edition, Cambridge Studies in Advanced Mathematics, 58, Cambridge University Press, Cambridge, 2001.
- [6] P. L. Q. Pergher, H. K. Singh and T. B. Singh, *On \mathbb{Z}_2 and \mathbb{S}^1 free actions on spaces of cohomology type (a, b)* , Houston J. Math. **36** (2010), no. 1, 137–146. MR2610784
- [7] H. K. Singh and T. B. Singh, *Fixed point free involutions on cohomology projective spaces*, Indian J. Pure Appl. Math. **39** (2008), no. 3, 285–291.
- [8] M. Singh, *Orbit spaces of free involutions on the product of two projective spaces*, Results Math. **57** (2010), no. 1-2, 53–67.
- [9] J. C. Su, *Transformation groups on cohomology projective spaces*, Trans. Amer. Math. Soc. **106** (1963), 305–318.

Anju Kumari, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI – 110007, INDIA.
Email address: anjukumari0702@gmail.com

Hemant Kumar Singh, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI – 110007,
INDIA.
Email address: hemantksingh@maths.du.ac.in