

FINITISTIC SPACES WITH THE ORBIT SPACE $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$

ANJU KUMARI AND HEMANT KUMAR SINGH

ABSTRACT. Let $G = \mathbb{S}^d$, $d = 0, 1$ or 3 , act freely on a finitistic connected space X . This paper gives the cohomology classification of X if a mod 2 or rational cohomology of the orbit space X/G is isomorphic to the product of a projective space and sphere $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$, where $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , respectively. For a free involution on X , a lower bound of covering dimension of the coincidence set of a continuous map $f : X \rightarrow \mathbb{R}^k$ is also determined.

1. INTRODUCTION

Let G be a group acting on a compact Hausdorff space X . It has been an interesting problem to classify the orbit space X/G for a given space X when G acts freely on X . It is well known that the projective spaces $\mathbb{F}\mathbb{P}^n$, where $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , are the orbit spaces of the standard free actions of $G = \mathbb{S}^d$ on $\mathbb{S}^{(d+1)n+d}$, where $d = 0, 1$ and 3 , and the lens space L_p^{2n+1} is the orbit space of the standard free action of $G = \mathbb{Z}_p$, p an odd prime, on \mathbb{S}^{2n+1} . A lot of work have been done in the literature in this direction for various spaces X and different groups G . For example: (i) $X = L_p^{2n+1}$ or $\mathbb{F}\mathbb{P}^n$ or $\mathbb{F}\mathbb{P}^n \times \mathbb{F}\mathbb{P}^m$, where $\mathbb{F} = \mathbb{R}$ or \mathbb{C} and $G = \mathbb{Z}_2$ [16, 18, 19], (ii) $X = \mathbb{F}H_{r,s}$, $\mathbb{F} = \mathbb{R}$ or \mathbb{C} and $0 \leq s \leq r$, (the Real or complex Milnor manifold) and $G = \mathbb{Z}_2$ or \mathbb{S}^1 [3], (iii) $X = \mathbb{S}^n \times \mathbb{S}^m$, $1 \leq n \leq m$ and $G = \mathbb{Z}_p$, p a prime, or \mathbb{S}^1 or \mathbb{S}^3 [5, 10], and (iv) $X = P(1, n)$ (Dold manifold) and $G = \mathbb{Z}_2$ [13]. Another thread of research is to classify a space X for a given orbit space X/G , where G acts on X . For semi-free actions of $G = \mathbb{S}^1$ with the orbit space X/G is a simply connected 4-manifold, it has been proved that the total space is a simply connected 5-manifold [7]. Duan [6] shows that every 1-connected 4-manifold M is the quotient of some regular circle action on the space $\#_{r-1}\mathbb{S}^2 \times \mathbb{S}^3$, where $r = \text{rank } H^2(M, \mathbb{Z})$. If G acts freely on X , then the cohomology

2010 *Mathematics Subject Classification.* Primary 55T10; Secondary 57S99 .

Key words and phrases. Free action; Finitistic space; Leray-Serre spectral sequence; Gysin sequence; Euler class.

This paper is supported by the Science and Engineering Research Board (Department of Science and Technology, Government of India) with reference number- EMR/2017/002192.

classification of X is discussed for (i) $X/G = L_p^{2n+1}$ or $\mathbb{C}\mathbb{P}^n$ and $G = \mathbb{Z}_p, p$ a prime or \mathbb{S}^1 [20], and for $X/G = \mathbb{H}\mathbb{P}^n$ and $G = \mathbb{S}^3$ [11]. We contribute to this question by determining the cohomology classification of a finitistic space X equipped with free action of $G = \mathbb{Z}_2, \mathbb{S}^1$ or \mathbb{S}^3 and the orbit space X/G is a mod 2 or rational cohomology $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$, where $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , respectively. These results describes the converse of some results for $G = \mathbb{Z}_2$ or \mathbb{S}^1 actions proved by Dotzel et. al. [5], and for $G = \mathbb{S}^3$ actions proved in [10].

For free \mathbb{Z}_2 -space X and any space Y , the coincidence set of a continuous map $f : X \rightarrow Y$ is defined as $A(f) = \{x \in X | f(x) = f(gx) \text{ for each } g \in \mathbb{Z}_2\}$. The classical Borsuk-Ulam Theorem states that for a continuous map $f : \mathbb{S}^n \rightarrow \mathbb{R}^k$, the coincidence set $A(f) = \{x \in \mathbb{S}^n | f(x) = f(-x)\}$ is nonempty, where \mathbb{S}^n is equipped with antipodal action. Munkholm [14] shows that for a map $f : \mathbb{S}^n \rightarrow M$, topological dimension of coincidence set $A(f)$ is greater than or equal to $n - k$, where M is a compact k -dimensional topological manifold, $n > k$. Biasi et al. [1] extends this result for generalized manifolds. In this paper, we determine a lower bound of the covering dimension of coincidence set $A(f)$ of a continuous map $f : X \rightarrow \mathbb{R}^k$ if $H^*(X/G; \mathbb{Z}_2) = \mathbb{Z}_2[a, b] / \langle a^{n+1}, b^2 \rangle$, where $\deg a = 1$ and $\deg b = m$.

2. PRELIMINARIES

Let G be a compact Lie group and $G \rightarrow E_G \rightarrow B_G$ be the universal principal G -bundle, where B_G is the classifying space. Suppose G acts freely on a finitistic space X . The associated bundle $X \hookrightarrow (X \times E_G)/G \rightarrow B_G$ is a fibre bundle with fibre X . Put $X_G = (X \times E_G)/G$. The bundle $X \hookrightarrow X_G \rightarrow B_G$ is called the Borel fibration. Then there exists the Leray-Serre spectral sequence for the Borel fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ which converges to $H^*(X_G)$ as an algebra with $E_2^{k,l} = H^k(B_G; \mathcal{H}^l(X; R))$. If B_G is simply connected then the system of local coefficients on B_G is simple then the E_2 -term becomes

$$E_2^{k,l} = H^k(B_G; R) \otimes H^l(X; R).$$

We recall some results which are needed to prove our results:

Proposition 2.1. Let $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ be the Borel fibration. Suppose that the system of local coefficients on B_G is simple, then the edge homomorphisms

$$\begin{aligned} H^k(B_G) \cong E_2^{k,0} \longrightarrow E_3^{k,0} \longrightarrow \cdots \longrightarrow E_k^{k,0} \longrightarrow E_{k+1}^{k,0} = E_\infty^{k,0} \subset H^k(X_G), \text{ and} \\ H^l(X_G) \longrightarrow E_\infty^{0,l} = E_{l+1}^{0,l} \subset E_l^{0,l} \subset \cdots \subset E_2^{0,l} \cong H^l(X) \end{aligned}$$

are the homomorphisms

$$\pi^* : H^k(B_G) \rightarrow H^k(X_G) \text{ and } i^* : H^l(X_G) \rightarrow H^l(X).$$

For details about spectral sequences, we refer the reader to [12]. Let $h : X_G \rightarrow X/G$ be the map induced by the G -equivariant projection $X \times E_G \rightarrow X$. Then h is a homotopy equivalence [4]. All the cohomologies are Čech cohomology with coefficients in R , where $R = \mathbb{Q}$ or \mathbb{Z}_2 . Note that $X \sim_R Y$ means $H^*(X; R) \cong H^*(Y; R)$. For $R = \mathbb{Q}$ and for $G = \mathbb{S}^1$ or \mathbb{S}^3 , we assume that the associated sphere bundle $G \hookrightarrow X \rightarrow X/G$ is orientable.

Proposition 2.2. ([17, 9]) Let $G = \mathbb{S}^1$ or \mathbb{S}^3 , act freely on a finitistic space X . If $H^i(X; R) = 0$ for all $i > n$, then $H^i(X/G; R) = 0$ for all $i > n$.

Now, we recall Gysin sequence of sphere bundles.

Proposition 2.3. Let $G = \mathbb{S}^d$, $d = 0, 1$ or 3 , act freely on a finitistic space X . The Gysin sequence of the sphere bundle $G \hookrightarrow X \xrightarrow{p} X/G$ is:

$$\dots \rightarrow H^i(X/G) \xrightarrow{p_i^*} H^i(X) \xrightarrow{\rho_i} H^{i-d}(X/G) \xrightarrow{\cup} H^{i+1}(X/G) \xrightarrow{p_{i+1}^*} H^{i+1}(X) \rightarrow \dots$$

which start with

$$0 \longrightarrow H^d(X/G) \xrightarrow{p_d^*} H^d(X) \xrightarrow{\rho_d} H^0(X/G) \xrightarrow{\cup} H^{d+1}(X/G) \xrightarrow{p_{d+1}^*} H^{d+1}(X) \longrightarrow \dots$$

where $\cup : H^i(X/G) \rightarrow H^{i+d+1}(X/G)$ maps $x \rightarrow x \cup u$ and $u \in H^{d+1}(X/G)$ denotes the Euler class of the sphere bundle. It is easy to observe that p_i^* is an isomorphism for $0 \leq i \leq d-1$ for $d = 1$ or 2 .

In this paper, we have considered finitistic spaces. These spaces were introduced by R. G. Swan [21] and have been discovered as relevant spaces for the study of cohomological aspects of transformation groups [2]. Recall that a space is said to be finitistic if it is paracompact, Hausdorff and every open cover of it have a finite dimensional open refinement. All compact spaces, all finite dimensional and all finite-dimensional paracompact spaces are some examples of finitistic spaces. Note that the space $X = \prod_{n=1}^{\infty} \mathbb{S}^n \times \mathbb{R}^k$ is an example of finitistic space which is neither compact nor has finite covering dimension. Recall that $H^*(\mathbb{R}\mathbb{P}^n \times \mathbb{S}^m; R) = R[a, b]/\langle a^{n+1}, b^2 \rangle$, where $\deg a = 1$ and $\deg b = m$.

3. MAIN THEOREMS

Let $G = \mathbb{S}^d$, $d = 0, 1$ or 3 , act freely on a finitistic space X having the mod 2 or rational cohomology the product of spheres $\mathbb{S}^{(d+1)n+d} \times \mathbb{S}^m$. It has been proved that one of the possibilities of the orbit space X/G are the mod 2 or rational cohomology $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$, where $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , respectively. Using techniques of the Gysin sequence of sphere bundles and the Leray-Serre spectral sequence of Borel fibration, we discuss converse of these statements. These results also describes the cohomology classification of a finitistic connected free G -space X with the orbit space a product of projective space and sphere. First, we have discussed for free actions of $G = \mathbb{S}^3$.

Theorem 3.1. Let $G = \mathbb{S}^3$ act freely on a finitistic connected space X with $X/G \sim_R \mathbb{H}\mathbb{P}^n \times \mathbb{S}^m$, where R is \mathbb{Q} or \mathbb{Z}_2 . Then the cohomology algebra of X with coefficients in R is isomorphic to the cohomology algebra of one of the following:

- (i) $\mathbb{S}^m \times \mathbb{S}^{4n+3}$;
- (ii) $\mathbb{S}^3 \times \mathbb{S}^m \times \mathbb{H}\mathbb{P}^n$;
- (iii) $\mathbb{H}\mathbb{P}^n \times \mathbb{S}^7$ and $m = 4$;
- (iv) $R[x, y]/\langle x^{n+1}, y^4 \rangle$, where $\deg x = 4$, $\deg y = 3$, $m = 6$, $\beta(y) = x$ and $R = \mathbb{Z}_2$, where $\beta : H^3(X/G) \rightarrow H^4(X/G)$ denotes the Bockstein homomorphism associated with the coefficient sequence $0 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_4 \rightarrow \mathbb{Z}_2 \rightarrow 0$.

Proof. It is clear that $H^i(X/G) \cong H^i(X)$ for $i = 0, 1$ and 2 , and $H^i(X) = 0$ for all $i > m + 4n + 3$. We consider different possibilities of the Euler class in the Gysin sequence:

Case(a): When $\cup : H^0(X/G) \rightarrow H^4(X/G)$ is trivial.

First, assume that $m > 4n$. In this case, $\cup : H^i(X/G) \rightarrow H^{i+4}(X/G)$ is trivial for all $i \geq 0$. For $0 \leq i < n$ and $k = 0, m$, we get ρ_{k+4i+3} and p_{k+4i+4}^* are isomorphisms. This implies that $H^{k+4i+3}(X) \cong H^{k+4i+4}(X) \cong R$. Suppose that $\{u_{4i+3}\}$ and $\{x^{i+1}\}$ denotes bases for $H^{4i+3}(X)$ and $H^{4i+4}(X)$, respectively, where $\rho_{4i+3}(u_{4i+3}) = a^i$ and $p_4^*(a) = x$. Note that $H^i(X) = 0$ for $i \neq 4n + 3$, and $4n < i < m$. If $m \neq 4n + 3$, then by the exactness of the Gysin sequence, $H^{4n+3}(X) \cong H^m(X) \cong R$ with bases $\{u_{4n+3}\}$ and $\{y\}$, respectively; and if $m = 4n + 3$ then $H^{4n+3}(X) \cong R \oplus R$ with basis $\{u_{4n+3}, y\}$, where $\rho_{4n+3}(u_{4n+3}) = a^n$ and $p_m^*(b) = y$. Also, $H^{m+4n+3}(X) \cong R$. Let $\{v_{m+4i+3}\}$ be basis for $H^{m+4i+3}(X)$ with $\rho_{m+4i+3}(v_{m+4i+3}) = a^i b$ for all $0 \leq i \leq n$. Clearly, $H^j(X) = 0$ for all

$k \leq j \equiv k + 1$ or $k + 2 \pmod{4} < k + 4n$; and $k = 0, m$; and $j \neq 4n + 3$. Note that $u_3^2 = 0$.

Now we assume that $m \leq 4n$. We consider four cases $m \equiv j \pmod{4}, 0 \leq j \leq 3$.

$m \equiv 0 \pmod{4}$: Let $m = 4i_0$ for some $i_0 \leq n$. In this case, $H^i(X) = 0$ for $i \equiv 1 \pmod{4}$ or $i \equiv 2 \pmod{4}$. Also, ρ_{4i-1} and p_{4i}^* are isomorphisms for all $i \geq 0$. This implies that for each $0 \leq i < i_0$, we have $H^{4i}(X) \cong R$ and $H^{4i+3}(X) \cong R$ with bases $\{x^i\}$ and $\{u_{4i+3}\}$, respectively, and for $i_0 \leq i \leq n$, $H^{4i}(X) \cong R \oplus R$ and $H^{4i+3}(X) \cong R \oplus R$ with bases $\{x^i, x^{i-i_0}y\}$ and $\{u_{4i+3}, v_{4i+3}\}$, respectively, where $p_m^*(b) = y, p_4^*(a) = x, \rho_{4i+3}(u_{4i+3}) = a^i$ and $\rho_{4i+3}(v_{4i+3}) = a^{i-i_0}b$. Also, for $1 \leq i \leq i_0$, $H^{4n+4i}(X) \cong R$ and $H^{4n+4i+3}(X) \cong R$ with bases $\{x^{n+i-i_0}y\}$ and $\{v_{4n+4i+3}\}$, respectively, where $\rho_{4n+4i+3}(v_{4n+4i+3}) = a^{n+i-i_0}b$. Clearly, $u_3^2 = 0$. Thus,

$$H^i(X) = \begin{cases} R & \text{if } j \leq i \equiv 0 \text{ or } 3 \pmod{4} \leq m - 1 + j, j = 0 \text{ or } 4n + 4 \\ R \oplus R & \text{if } m \leq i \equiv 0 \text{ or } 3 \pmod{4} \leq 4n + m + 3 \\ 0 & \text{otherwise.} \end{cases}$$

$m \equiv 1 \pmod{4}$: Let $m = 4i_0 + 1$ for some $i_0 < n$. For $i \neq i_0 + j, 1 \leq j \leq n + 1$, we get p_{4i}^* are isomorphisms. This implies that $H^{4i}(X) \cong R$ with basis $\{x^i\}$ for $0 \leq i \leq i_0$, where $p_4^*(a) = x$. For $i_0 + 1 \leq i \leq n$, $H^{4i}(X) \cong R \oplus R$ with basis $\{x^i, v_{4i}\}$; and for $n + 1 \leq i \leq i_0 + n + 1$, $H^{4i}(X) \cong R$ with basis $\{v_{4i}\}$, where $\rho_{4i}(v_{4i}) = a^{i-i_0-1}b$. It is clear that $H^{4i+2}(X) = 0$ for all $i \geq 0$; $H^{4i+3}(X) \cong R \cong H^{m+4i}(X)$, for all $0 \leq i \leq n$, with bases $\{u_{4i+3}\}$ and $\{x^i y\}$, where $\rho_{4i+3}(u_{4i+3}) = a^i$ and $p_m^*(b) = y$; and $H^{4i+3}(X) = H^{4i+1}(X) = 0$ otherwise. Obviously, $u_3^2 = 0$. We have

$$H^i(X) = \begin{cases} R & \text{if } j \leq i \equiv 0 \pmod{4} \leq j + m - 1, j = 0 \text{ or } 4n + 4, \text{ or} \\ & j \leq i \equiv j \pmod{4} \leq j + 4n, j = 3 \text{ or } m \\ R \oplus R & \text{if } m < i \equiv 0 \pmod{4} \leq 4n \\ 0 & \text{otherwise.} \end{cases}$$

$m \equiv 2 \pmod{4}$: Let $m = 4i_0 + 2$ for some $i_0 < n$. We get ρ_{4i+3} and p_{4i}^* are isomorphisms, for all $i \geq 0$. Therefore, for $0 \leq i \leq n$, $H^{4i+3}(X) \cong H^{4i}(X) \cong R$ with bases $\{u_{4i+3}\}$ and $\{x^i\}$, respectively, and for $i > n$, $H^{4i+3}(X) = H^{4i}(X) = 0$, where $\rho_{4i+3}(u_{4i+3}) = a^i$ and $p_4^*(a) = x$. Also, we have $H^{4i+1}(X) = 0$, for $i \neq i_0 + j, 1 \leq j \leq n + 1$, and for $i_0 + 1 \leq i \leq i_0 + n + 1$, $H^{4i+1}(X) \cong H^{4i-2}(X) \cong R$ with bases $\{v_{4i+1}\}$ and $\{x^{i-i_0-1}y\}$,

where $\rho_{4i+1}(v_{4i+1}) = a^{i-i_0-1}b$ and $p_m^*(b) = y$. Consequently,

$$H^i(X) = \begin{cases} R & \text{if } j \leq i \equiv j \text{ or } j + 3 \pmod{4} \leq j + 4n + 3, j = 0 \text{ or } m \\ 0 & \text{otherwise.} \end{cases}$$

$m \equiv 3 \pmod{4}$: Let $m = 4i_0 + 3$ for some $i_0 < n$. Then $H^{4i}(X) = 0$ for $i > n$, and $H^{4i}(X) \cong R$ with basis $\{x^i\}$, for $0 \leq i \leq n$, where $p_4^*(a) = x$. Also, $H^{4i+1}(X) = 0$ for all $i \geq 0$. By the exactness of the Gysin sequence, for $i_0 + 1 \leq i \leq i_0 + n + 1$, $H^{4i+2}(X) \cong R$ with basis $\{v_{4i+2}\}$, where $\rho_{4i+2}(v_{4i+2}) = a^{i-i_0-1}b$, and $H^{4i+2}(X) = 0$, otherwise. Now, for $0 \leq i < i_0$, $H^{4i+3}(X) \cong R$ with basis $\{u_{4i+3}\}$; for $i_0 \leq i \leq n$, $H^{4i+3}(X) \cong R \oplus R$ with basis $\{u_{4i+3}, x^{i-i_0}y\}$; and for $n < i \leq i + i_0$, $H^{4i+3}(X) \cong R$ with basis $\{x^{i-i_0}y\}$, where $\rho_{4i+3}(u_{4i+3}) = a^i$ and $p_m^*(b) = y$. Clearly, $u_3^2 = 0$. We have

$$H^i(X) = \begin{cases} R & \text{if } j \leq i \equiv j \pmod{4} \leq j + 4n, j = 0 \text{ or } m + 3, \text{ or} \\ & j + 3 \leq i \equiv 3 \pmod{4} \leq j + m - 4, j = 0 \text{ or } 4n + 4 \\ R \oplus R & \text{if } m \leq i \equiv 3 \pmod{4} \leq 4n + 3 \\ 0 & \text{otherwise.} \end{cases}$$

Consider the Leray-Serre spectral sequence for the Borel fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ which converges to $H^*(X_G)$ as an algebra and $E_2^{k,l} = H^k(B_G; R) \otimes H^l(X; R)$ for all $k, l \geq 0$. The possible nontrivial differentials in the Leray-Serre spectral sequence for the Borel fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$ are $\{d_{4r}\}_{r \geq 1}$. By the edge homomorphisms and the fact that $p_j^* = i^* \circ h^*$ for all $j \geq 0$, we get $d_r(1 \otimes x) = 0 = d_r(1 \otimes y)$ for all $r \geq 0$. Note that u_{4i+3} and v_{m+4i+3} are not in image of p_{4i+3}^* and p_{m+4i+3}^* , respectively, so their image must be nonzero under some differential for all $0 \leq i \leq n$. Consequently, $x^i u_3 \neq 0$ and $x^i u_3 y \neq 0$, for all i . Obviously, $x^{n+1} = y^2 = 0$.

For $m > 4n$, it is clear that $u_{4i+3} = \alpha_i x^i u_3$ and $v_{m+4i+3} = \beta_i x^i u_3 y$ for some nonzero elements α_i, β_i in R and $0 \leq i \leq n$. In particular, for $m = 4n + 3$, $x^n u_3$ can not be equal to any multiple of y and so u_{4n+3} is generated by $\{y, x^n u_3\}$.

Now, suppose $m \leq 4n$. If $m = 4i_0$, for some $i_0 \leq n$ then $u_{4i+3} = \alpha_i x^i u_3$ for $0 \leq i < i_0$ and $v_{4i+3+m} = \beta_i x^i u_3 y$ for $n+1-i_0 \leq i \leq n$, where α_i 's and β_i 's are nonzero elements in R . We observe that $x^i u_3$ can not be equal to any multiple of $x^{i-i_0} u_3 y$, for all $i_0 \leq i \leq n$. Therefore, the elements u_{4i+3} and v_{4i+3} are generated by $\{x^i u_3, x^{i-i_0} u_3 y\}$ for each $i_0 \leq i \leq n$. If $m = 4i_0 + 1$ for some $i_0 < n$ then the elements $x^i u_3 y$ cannot be equal to any multiple of x^{4i+3+m} for each $0 \leq i \leq n - i_0 + 1$. Consequently, v_{4i+m+3} is generated by $\{x^i u_3 y, x^{4i+m+3}\}$ and u_{4i+3} is generated by $\{x^i u_3\}$ for all $0 \leq i \leq n$. If

$m = 4i_0 + 2$ for some $i_0 < n$ then we have $u_{4i+3} = \alpha_i x^i u_3$ and $v_{m+4i+3} = \beta_i x^i u_3 y$ for some nonzero elements α_i, β_i in R and $0 \leq i \leq n$. In this case, u_3^2 may be both zero or nonzero. If $m = 4i_0 + 3$ then $v_{m+4i+3} = \alpha_i x^i u_3 y$, where $\alpha_i \in R$, for all $0 \leq i \leq n$. As $x^i u_3$ can not be equal to $x^{i-i_0} y$, for all $i_0 \leq i \leq n$, therefore, u_{4i+3} is generated by $\{x^i u_3, x^{i-i_0} y\}$, for all $0 \leq i \leq n$. For $m \neq 3$, obviously $u_3^2 = 0$. If $u_3^2 \neq 0$, for $m = 3$, then $d_4(1 \otimes v_6) = 0$, a contradiction. Therefore, $u_3^2 = 0$. The cohomology algebra of X is given by

$$H^*(X) = R[x, u_3, y] / \langle x^{n+1}, u_3^2, y^2 \rangle,$$

where $\deg x = 4$, $\deg u_3 = 3$ and $\deg y = m$. This realizes the possibility (ii).

If $u_3^2 \neq 0$ then m must be 6 and $u_3^2 = \alpha y$ for some α nonzero in R . By the commutativity of cup product, we get $2u_3^2 = 0$. This implies that R cannot be \mathbb{Q} . Therefore,

$$H^*(X) \cong R[x, u_3] / \langle x^{n+1}, u_3^4 \rangle,$$

$\deg x = 4$ and $\deg u_3 = 3$ only when $R = \mathbb{Z}_2$. By the properties of Steenrod squares $Sq^3(y) = y^2 \neq 0$. As 3 is not a power of 2, we get $Sq^3 = Sq^2 \circ Sq^1 + Sq^1 \circ Sq^2$. This gives that $Sq^1(y) = x$. Note that Sq^1 is the Bockstein homomorphism associated with the coefficient sequence $0 \rightarrow \mathbb{Z}_2 \rightarrow \mathbb{Z}_4 \rightarrow \mathbb{Z}_2 \rightarrow 0$. This realizes the possibility (iv) of the Theorem.

Case(b): When $\cup : H^0(X/G) \rightarrow H^4(X/G)$ maps 1 to ca for some $c \neq 0$ in R .

First, we suppose that $m > 4n$. For $0 \leq i < n$ and $j = 0, m$, we get $\cup : H^{4i+j}(X/G) \rightarrow H^{4i+4+j}(X/G)$ are isomorphisms; ρ_{4i+3+j} and p_{4i+4+j}^* are trivial homomorphisms. Consequently, $H^{4i+3+j}(X) = H^{4i+4+j}(X) = 0$. Also, $H^{4i+1+j}(X) = H^{4i+2+j}(X) = 0$ except for $H^{4n+3}(X)$. Note that $H^i(X) = 0$ if $(4n < i < m$ and $i \neq 4n + 3)$ and $H^{m+4n+3}(X) \cong R$ with basis $\{u_{m+4n+3}\}$, where $\rho_{m+4n+3}(u_{m+4n+3}) = a^n b$. For $m \neq 4n + 3$, ρ_{4n+3} and p_m^* are isomorphisms. Consequently, $H^{4n+3}(X) \cong R \cong H^m(X)$ with bases $\{u_{4n+3}\}$ and $\{y\}$, respectively, where $p_m^*(b) = y$ and $\rho_{4n+3}(u_{4n+3}) = a^n$. If $m = 4n+3$ then $H^{4n+3}(X) \cong R \oplus R$ with basis $\{u_{4n+3}, y\}$, where $\rho_{4n+3}(u_{4n+3}) = a^n$ and $p_m^*(b) = y$. Also, $H^{m+4n+1}(X) = H^{m+4n+2}(X) = 0$. For $m \neq 4n + 3$, we get

$$H^i(X) = \begin{cases} R & \text{if } i = 0, 4n + 3, m, m + 4n + 3 \\ 0 & \text{otherwise} \end{cases}$$

and for $m = 4n + 3$, we get

$$H^i(X) = \begin{cases} R \oplus R & \text{if } i = 4n + 3 \\ R & \text{if } i = 0, m + 4n + 3 \\ 0 & \text{otherwise.} \end{cases}$$

For $m \leq 4n$, we get the same cohomology groups as in $m > 4n$.

Now, we compute the cohomology ring structure of X . In the Leray-Serre spectral sequence for the Borel fibration $X \hookrightarrow X_G \xrightarrow{\pi} B_G$, $d_{4r'}(1 \otimes u_{4n+3}) \neq 0$ for some $r' > 0$ and $d_{4r}(1 \otimes y) = 0$ for all $r \geq 0$. Firstly, suppose that $m = 4i_0$ and $r' = n - i_0 + 1$, where $1 \leq i_0 \leq n$, then $E_{4n-m+5}^{k,4n+3} = 0$, $E_{4n-m+5}^{k,q} = E_2^{k,q}$ for all $k \geq 0$ and $q = 0, m + 4n + 3$. Also, $E_{4n-m+5}^{k,m} = E_2^{k,m}$ for all $k \leq 4n - m$ and trivial otherwise. Since G acts freely on X , therefore, $d_{m+4n+4}(1 \otimes u_{m+4n+3}) = ct^{n+i_0+1} \otimes 1$ for some $c \neq 0$ in R . Then $E_\infty^{4k,q} = R$ for all $(4k \leq m + 4n, q = 0)$, $(4k \leq 4n - m, q = m)$ and trivial otherwise. This implies that $t \otimes 1 \in E_2^{4,0}$ and $1 \otimes y \in E_2^{0,m}$ are permanent cocycles. Then by edge homomorphism there exist $u \in E_\infty^{4,0}$ and $w \in E_\infty^{0,m}$ corresponding to $t \otimes 1$ and $1 \otimes y$ respectively with $\pi^*(t) = u$. We have $w^2 = u^{n+i_0+1} = u^{n-i_0+1}w = 0$. This implies that

$$\text{Tot}E_\infty^{*,*} \cong \mathbb{R}[u, w] / \langle w^2, u^{n+i_0+1}, u^{n-i_0+1}w \rangle,$$

where $\deg u = 4$ and $\deg w = m$. Then there exist an element $v \in H^m(X_G)$ corresponding to $w \in E_\infty^{0,m}$ such that $i^*(v) = y$. We have $u^{n-i_0+1}v = \alpha u^{n+1}$ and $v^2 = \beta u^{2i_0} + \gamma u^{i_0}v$, where $\alpha, \beta, \gamma \in R$ and $\beta = 0$ if $m > 4n$. So, the ring cohomology of X/G is given by $H^*(X_G) \cong R[u, v] / \langle u^{n-i_0+1}v - \alpha u^{n+1}, v^2 - \beta u^{2i_0} - \gamma u^{i_0}v, u^{n+i_0+1} \rangle$ where $\deg u = 4$, $\deg v = m$ and $\alpha, \beta, \gamma \in R$, $\beta = 0$ if $m > 4n$, which is a contradiction. So, r' must be $n + 1$. This implies that $d_{4n+4}(1 \otimes yu_{4n+3}) \neq 0$. Consequently, $u_{4n+m+3} = \alpha yu_{4n+3}$ for some $\alpha \neq 0$ in R . Obviously, $y^2 = 0$, and $u_{4n+3}^2 = 0$ for $m \notin \{4n + 3, 8n + 6\}$. If $m = 4n + 3$ then $u_{4n+3}^2 \neq \beta u_{4n+m+3}$ for any β in R and we get $u_{4n+3}^2 = 0$. If $m = 8n + 6$ then u_{4n+3}^2 may be both zero or nonzero. Thus,

$$H^*(X, R) = R[y, u_{4n+3}] / \langle y^2, u_{4n+3}^2 \rangle,$$

where $\deg y = m$ and $\deg u_{4n+3} = 4n + 3$. This realizes possibility (i). If $u_{4n+3}^2 \neq 0$ then $u_{4n+3} = \alpha y$ for some nonzero element $\alpha \in R$. Thus, the cohomology algebra of X is given by $R[u_{4n+3}] / \langle u_{4n+3}^4 \rangle$, where $\deg u_{4n+3} = 4n + 3$. By [8, Theorem 4L.9], this cohomology algebra is not possible for $R = \mathbb{Z}_2$ and by the commutativity of cup product this cohomology algebra is also not possible for $R = \mathbb{Q}$.

Case(c): When $\cup : H^0(X/G) \rightarrow H^4(X/G)$ maps 1 to cb , where $c \neq 0$ in R .

In this case m must be 4, and we get $H^{4i}(X) \cong R$ with basis $\{x^i\}$, where $p_4^*(a) = x$. Clearly, $H^3(X) = H^{4i+1}(X) = H^{4i+2}(X) = 0$ for all $i \geq 0$. As $\ker \cup : H^{4i}(X/G) \rightarrow H^{4i+4}(X/G)$ is generated by $a^{i-1}b$, we have $H^{4i+3}(X) \cong R$ with basis $\{u_{4i+3}\}$, where $\rho_{4i+3}(u_{4i+3}) = a^{i-1}b$ for all $1 \leq i \leq n+1$. We must have $d_4(1 \otimes u_7) = 0$ and $d_8(1 \otimes u_7) \neq 0$. Consequently, $u_{4i+7} = c_i x^i b_7$ for some nonzero $c_i \in R$, for all $1 \leq i \leq n$. Therefore, $H^*(X) \cong R[x, u_7]/\langle u_7^2, x^{n+1} \rangle$, where $\deg x = 4$ and $\deg u_7 = 7$. This realizes the possibility (iii).

Case(d): When $\cup : H^0(X/G) \rightarrow H^4(X/G)$ maps 1 to $ca + c'b$, where $c, c' \neq 0$ in R .

In this case also m must be 4 and $H^4(X) \cong R$ with basis $\{x\}$, where $p_4^*(a) = x$. By the exactness of the Gysin sequence, we get $H^3(X) = H^{4i+1}(X) = H^{4i+2}(X) = 0$ for all $i \geq 0$. As $\cup : H^{4i}(X/G) \rightarrow H^{4i+4}(X/G)$ is an isomorphism, we get $H^{4i+3}(X) = H^{4i+4}(X) = 0$ for all $0 < i < n$. Note that $H^{4n+4}(X) = 0$ and $\ker \cup : H^{4n}(X/G) \rightarrow H^{4n+4}(X/G)$ is generated by $\{a^{n-1}b - \frac{c}{c'}a^n\}$. This implies that $H^{4n+3}(X) \cong R$ with basis $\{u_{4n+3}\}$, where $\rho_{4n+3}(u_{4n+3}) = a^{n-1}b - \frac{c}{c'}a^n$. Obviously, $H^{4n+7}(X) \cong R$ with basis $\{u_{4n+7}\}$, where $\rho_{4n+7}(u_{4n+7}) = a^n b$. Using similar arguments as in case(b), $H^*(X) \cong R[x, u_{4n+3}]/\langle u_{4n+3}^2, x^2 \rangle$, where $\deg x = 4$ and $\deg u_{4n+3} = 4n+3$. This realizes the possibility (i). \square

Next, we discuss similar result for circle actions with the orbit space product of a complex projective space and sphere:

Theorem 3.2. Let $G = \mathbb{S}^1$ act freely on a finitistic connected space X with $X/G \sim_R \mathbb{C}\mathbb{P}^n \times \mathbb{S}^m$, where R is \mathbb{Q} or \mathbb{Z}_2 . The cohomology algebra of X with coefficients in R is isomorphic to the cohomology algebra of one of the following:

- (i) $\mathbb{S}^m \times \mathbb{S}^{2n+1}$;
- (ii) $\mathbb{S}^1 \times \mathbb{S}^m \times \mathbb{C}\mathbb{P}^n$;
- (iii) $\mathbb{R}\mathbb{P}^{2n+1} \times \mathbb{S}^m$;
- (iv) $R[x, y]/\langle x^{n+1}, y^2 + \alpha x^3 \rangle$, where $\deg x = 2, \deg y = 3, m = 2, \alpha \in R$ and $\alpha = 0$ for $R = \mathbb{Q}$.

Proof. As X is connected, $H^0(X/G) \cong H^0(X)$, and clearly, $H^i(X) = 0$ for all $i > m + 2n + 1$. We consider the following cases:

Case(a): When $\cup : H^0(X/G) \rightarrow H^2(X/G)$ is trivial.

First assume that $m > 2n$. In this case, $\cup : H^i(X/G) \rightarrow H^{i+2}(X/G)$ is trivial for all $i \geq 0$. Then ρ_{k+2i+1} and p_{k+2i+2}^* are isomorphisms for all $0 \leq i < n$ and $k = 0, m$.

This implies that $H^{k+2i+1}(X) \cong H^{k+2i+2}(X) \cong R$. Suppose that $\{u_{2i+1}\}$ and $\{x^{i+1}\}$ denotes the bases for $H^{2i+1}(X)$ and $H^{2i+2}(X)$, respectively, where $\rho_{2i+1}(u_{2i+1}) = a^i$ and $p_2^*(a) = x$. Also, $H^{m+2n+1}(X) \cong R$. Let $\{v_{m+2i+1}\}$ be basis for $H^{m+2i+1}(X)$ with $\rho_{m+2i+1}(v_{m+2i+1}) = a^i b$ for all $0 \leq i \leq n$. Note that $H^i(X) = 0$, for all $2n+1 < i < m$. If $m \neq 2n+1$, then by the exactness of Gysin sequence, $H^{2n+1}(X) \cong H^m(X) \cong R$ with bases $\{u_{2n+1}\}$ and $\{y\}$, respectively; and if $m = 2n+1$ then $H^{2n+1}(X) \cong R \oplus R$ with basis $\{u_{2n+1}, y\}$, where $\rho_{2n+1}(u_{2n+1}) = a^n$ and $p_m^*(b) = y$.

Now, suppose that $m \leq 2n$. We consider two cases when m is even or odd:

m is odd: Let $m = 2i_0 + 1$ for some $i_0 < n$. In this case, $\cup : H^{2i}(X/G) \rightarrow H^{2i+2}(X/G)$ is trivial for all $i \geq 0$. Then ρ_{2i-1} and p_{2i}^* are isomorphisms for all $0 \leq i \leq i_0$. This implies that $H^{2i-1}(X) \cong H^{2i}(X) \cong R$ with bases $\{u_{2i-1}\}$ and $\{x^i\}$, respectively, where $\rho_{2i-1}(u_{2i-1}) = a^{i-1}$ and $p_2^*(a) = x$. For $i_0 + 1 \leq i \leq n$, $H^{2i}(X) \cong H^{2i-1}(X) \cong H^{2n+1}(X) \cong R \oplus R$ with bases $\{x^i, v_{2i}\}$, $\{yx^{i-i_0-1}, u_{2i-1}\}$ and $\{yx^{n-i_0}, u_{2n+1}\}$, respectively; and for $n < i \leq i_0 + n$, we have $H^{2i}(X) \cong H^{2n+m+1}(X) \cong H^{2i+1}(X) \cong R$ with bases $\{v_{2i}\}$, $\{v_{2n+m+1}\}$ and $\{yx^{i-i_0}\}$, respectively, where $\rho_{2i}(v_{2i}) = a^{i-i_0-1}b$, $\rho_{2i+1}(u_{2i+1}) = a^i$ and $p_m^*(b) = y$.

m is even: Let $m = 2i_0$ for some $i_0 \leq n$. We get that ρ_{2i-1} and p_{2i}^* are isomorphisms for all $i \geq 0$. This implies that for each $0 \leq i < i_0$, $H^{2i}(X) \cong H^{2i+1}(X) \cong R$ with bases $\{x^i\}$ and $\{u_{2i+1}\}$, respectively, and for $i_0 \leq i \leq n$, we get $H^{2i}(X) \cong H^{2i+1}(X) \cong R \oplus R$ with bases $\{x^i, x^{i-i_0}y\}$ and $\{u_{2i+1}, v_{2i+1}\}$, respectively, where $p_2^*(a) = x, p_m^*(b) = y, \rho_{2i+1}(u_{2i+1}) = a^i$ and $\rho_{2i+1}(v_{2i+1}) = a^{i-i_0}b$. Also, for $1 \leq i \leq i_0$, $H^{2n+2i}(X) \cong H^{2n+2i+1}(X) \cong R$ with bases $\{x^{n+i-i_0}y\}$ and $\{v_{2n+2i+1}\}$, respectively, where $\rho_{2n+2i+1}(v_{2n+2i+1}) = a^{n+i-i_0}b$.

Finally, for all $m \leq 2n$, we have

$$H^i(X) = \begin{cases} R & \text{if } j \leq i \leq m + j - 1, j = 0 \text{ or } 2n + 2, \\ R \oplus R & \text{if } m \leq i \leq 2n + 1 \\ 0 & \text{otherwise.} \end{cases}$$

Now, we compute the cohomology ring structure of X . In the Leray Serre spectral sequence for the Borel fibration $X \xrightarrow{i} X_G \xrightarrow{\pi} B_G$, it is easy to observe that $d_r(1 \otimes x) = 0 = d_r(1 \otimes y)$ for all $r \geq 0$, and the images of u_{2i+1} and v_{m+2i+1} must be nonzero under some differential, for all $0 \leq i \leq n$. Consequently, $x^i u_1 \neq 0$ and $x^i u_1 y \neq 0$. Obviously, $x^{n+1} = y^2 = 0$.

For $m > 2n$, it is clear that for $0 \leq i \leq n$, $u_{2i+1} = \alpha_i x^i u_1$ and $v_{m+2i+1} = \beta_i x^i u_1 y$ for some nonzero elements α_i, β_i in R . In particular, for $m = 2n + 1$, $x^n u_1$ can not be equal to any multiple of y and so u_{2n+1} is generated by $\{y, x^n u_1\}$.

Now, suppose $m \leq 2n$. If $m = 2i_0 + 1$ for some $i_0 < n$ then $u_{2i+1} = \alpha_i x^i u_1$ for $0 \leq i < i_0$ and $v_{m+2i+1} = \beta_i x^i u_1 y$ for $n - i_0 \leq i \leq n$, where α_i 's and β_i 's are nonzero elements in R . As $x^i u_1$ can not be equal to $x^{i-i_0} y$, for all $i_0 \leq i \leq n$, therefore, u_{2i+1} is generated by $\{x^i u_1, x^{i-i_0} y\}$, for all $i_0 \leq i \leq n$. Also, $x^i u_1 y$ can not be equal to any multiple of x^{i+i_0+1} , so v_{m+2i+1} is generated by $\{x^{i+i_0+1}, x^i u_1 y\}$, for all $0 \leq i < n - i_0$. If $m = 2i_0$, for some $i_0 \leq n$ then $u_{2i+1} = \alpha_i x^i u_1$ for $0 \leq i < i_0$ and $v_{2i+m+1} = \beta_i x^i u_1 y$ for $n - i_0 + 1 \leq i \leq n$, where α_i 's and β_i 's are nonzero elements in R . We observe that $x^i u_1$ can not be equal to any multiple of $x^{i-i_0} u_1 y$, for all $i_0 \leq i \leq n$. Thus, the elements u_{2i+1} and v_{2i+1} are generated by $\{x^i u_1, x^{i-i_0} u_1 y\}$. Note that u_1^2 may be both zero or nonzero. If $u_1^2 = 0$ then $X \sim_R \mathbb{S}^1 \times \mathbb{C}\mathbb{P}^n \times \mathbb{S}^m$. If $u_1^2 \neq 0$, then by the commutativity of cup product, $R = \mathbb{Z}_2$ and $X \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^{2n+1} \times \mathbb{S}^m$. This realizes possibilities (ii) and (iii) of the theorem.

Case(b): When $\cup : H^0(X/G) \rightarrow H^2(X/G)$ maps 1 to ca for some $c \neq 0$ in R .

First, suppose that $m > 2n$. In this case, ρ_{2i+1+j} and p_{2i+2+j}^* are trivial for all $0 \leq i < n$ and $j = 0$ or m . Consequently, $H^{2i+1+j}(X) = H^{2i+2+j}(X) = 0$. Note that $H^i(X) = 0$ for $2n + 1 < i < m$ and $H^{m+2n+1}(X) \cong R$ with basis $\{u_{m+2n+1}\}$, where $\rho_{m+2n+1}(u_{m+2n+1}) = a^n b$. Thus, for $m \neq 2n + 1$, ρ_{2n+1} and p_m^* are isomorphisms. This implies that $H^{2n+1}(X) \cong H^m(X) \cong R$ with bases $\{u_{2n+1}\}$ and $\{y\}$ respectively. If $m = 2n + 1$ then $H^{2n+1}(X) \cong R \oplus R$ with basis $\{u_{2n+1}, y\}$, where $\rho_{2n+1}(u_{2n+1}) = a^n$ and $p_m^*(b) = y$.

It is easy to see that, for $m \leq 2n$, the cohomology groups and generators are the same as above.

In the Leray-Serre spectral sequence for the Borel fibration $X \hookrightarrow X_G \xrightarrow{\pi} B_G$, $d_{2r'}(1 \otimes u_{2n+1}) \neq 0$ for some $r' > 0$ and $d_{2r}(1 \otimes y) = 0$ for all $r \geq 0$. Let if possible $m = 2i_0$ and $r' = n - i_0 + 1$, where $1 \leq i_0 \leq n$. Then as done in previous theorem, we get $H^*(X_G) \cong R[u, v] / \langle u^{n-i_0+1} v - \alpha u^{n+1}, v^2 - \beta u^{2i_0} - \gamma u^{i_0} v, u^{n+i_0+1} \rangle$ where $\deg u = 2$, $\deg v = m$ and $\alpha, \beta, \gamma \in R$, $\beta = 0$ if $m > 2n$, which is a contradiction. Therefore, r' must be $n + 1$. This implies that $d_{2n+2}(1 \otimes y u_{2n+1}) \neq 0$. Consequently, $u_{m+2n+1} = \alpha y u_{2n+1}$ for some $\alpha \neq 0$ in R . Obviously, $y^2 = 0$ and $u_{2n+1}^2 = 0$ for $m \notin \{2n + 1, 4n + 2\}$. If $m = 2n + 1$ then $u_{2n+1}^2 \neq \alpha' u_{m+2n+1}$ for any α' in R and so $u_{2n+1}^2 = 0$. If $m = 4n + 2$ then u_{2n+1}^2 may be

both zero or nonzero. If $u_{2n+1}^2 = 0$ then $H^*(X) = R[y, u_{2n+1}]/\langle y^2, u_{2n+1}^2 \rangle$, where $\deg y = m$ and $\deg u_{2n+1} = 2n + 1$. This realizes possibility (i) of the theorem. If $u_{2n+1}^2 \neq 0$ then $u_{2n+1} = \beta y$ for some nonzero element β in R . So, we get $H^*(X) = R[u_{2n+1}]/\langle u_{2n+1}^4 \rangle$, where $\deg u_{2n+1} = 2n + 1$ and $m = 4n + 2$. By the commutativity of cup product this is not possible for $R = \mathbb{Q}$. Also, by [8, Theorem 4L.9], it is not possible for $R = \mathbb{Z}_2$.

Case(c): When $\cup : H^0(X/G) \rightarrow H^2(X/G)$ maps 1 to cb , where $c \neq 0$ in R .

In this case, m must be 2 and $H^1(X) = 0$. As $H^{2i-1}(X/G) = 0$ for all $i \geq 0$, therefore, we have $H^{2i-2}(X) \cong \text{im } p_{2i-2}^*$ and $H^{2i+1}(X) \cong \ker\{\cup : H^{2i}(X/G) \rightarrow H^{2i+2}(X/G)\}$ with bases $\{x^i\}$ and $\{u_{2i+1}\}$, respectively, where $\rho_{2i+1}(u_{2i+1}) = a^{i-1}b$ and $p_2^*(a) = x$ for all $1 \leq i \leq n + 1$. In the Leray-Serre spectral sequence, $d_{2r}(1 \otimes x) = 0$ for all $r \geq 0$ and $d_{2r'}(1 \otimes u_3) \neq 0$ for some $r' > 0$. If $r' = 1$ then $H^*(X_G) \cong R[u, v]/\langle u^{n+2}, v^{n+1}, uv \rangle$, where $\deg u = \deg v = 2$, a contradiction. Therefore, r' must be 2, and we get $x^i u_3 \neq 0$ for all $1 \leq i \leq n$. This implies that $u_{2i+3} = \alpha_i x^i u_3$ for some nonzero $\alpha_i \in R$. We have $u_3^2 = \alpha x^3$ for some $\alpha \in R$. By the commutativity of cup product, we get $2u_3^2 = 0$. Therefore, for $R = \mathbb{Q}$, α must be zero. Therefore, $H^*(X) \cong R[x, u_3]/\langle x^{n+1}, u_3^2 - \alpha x^3 \rangle$, where $\deg x = 2$, $\deg u_3 = 3$ and $\alpha = 0$ for $R = \mathbb{Q}$. This realizes possibility (iv).

Case(d): When $\cup : H^0(X/G) \rightarrow H^2(X/G)$ maps 1 to $ca + c'b$, where $c, c' \neq 0$ in R .

We have $m = 2$ and $\cup : H^{2i}(X/G) \rightarrow H^{2i+2}(X/G)$ is an isomorphism, for all $0 < i < n$. By the exactness of Gysin sequence, $H^{2i-1}(X) = H^{2i+2}(X) = 0$ for all $0 < i \leq n$; and $H^2(X) \cong H^{2n+1}(X) \cong H^{2n+3}(X) \cong R$ with bases $\{x\}$, $\{u_{2n+1}\}$ and $\{u_{2n+3}\}$, respectively, where $p_2^*(a) = x$, $\rho_{2n+1}(u_{2n+1}) = a^{n-1}b - \frac{c}{c'}a^n$ and $\rho_{2n+3}(u_{2n+3}) = a^n b$. Hence, $H^*(X) \cong R[x, u_{2n+1}]/\langle u_{2n+1}^2, x^2 \rangle$, where $\deg x = 2$ and $\deg u_{2n+1} = 2n + 1$. This realizes possibility (i). \square

Finally, we classify a finitistic space X equipped with a free involution and the orbit space a product of real projective space and sphere:

Theorem 3.3. Let $G = \mathbb{Z}_2$ act freely on a finitistic connected space X with $X/G \sim_{\mathbb{Z}_2} \mathbb{RP}^n \times \mathbb{S}^m$. Then the cohomology algebra of X is one of the following:

- (i) $\mathbb{S}^m \times \mathbb{S}^n$;
- (ii) $\mathbb{Z}_2[x, y]/\langle x^{n+1}, y^2 + \alpha x^2 \rangle$, where $\deg x = \deg y = 1$, $m = 1$ and $\alpha \in \mathbb{Z}_2$;
- (iii) $\mathbb{Z}_2[y]/\langle y^4 \rangle$, where $\deg y = n$, $m = 2n$ and $n = 1, 2$ or 4 .

Proof. Clearly, $H^0(X/G) \cong H^0(X)$ and $H^i(X) = 0$ for all $i > m + n$. It is easy to see that Euler class of the 0-sphere bundle $X \rightarrow X/G$ must be nontrivial. Now, we consider the following cases:

Case(a): When $\cup : H^0(X/G) \rightarrow H^1(X/G)$ maps 1 to a .

First, we suppose that $m \geq n$. For $m \neq n$, we get $\cup : H^{k+i}(X/G) \rightarrow H^{k+i+1}(X/G)$ is an isomorphism for all $1 \leq i < n$ and $k = 0$ or m . This implies that ρ_{k+i} and p_{k+i+1}^* are trivial homomorphism. Consequently, $H^{k+i}(X) = 0$ and $H^n(X) \cong H^m(X) \cong R$ with bases $\{u_n\}$ and $\{y\}$ respectively, where $\rho_n(u_n) = a^n$ and $p_m^*(b) = y$. Note that $H^i(X) = 0$ for $n < i < m$, $H^{m+n}(X) \cong R$ with basis $\{u_{m+n}\}$, where $\rho_{m+n}(u_{m+n}) = a^n b$. For $n = m$, $H^n(X) \cong R \oplus R$ with basis $\{u_n, y\}$, where $\rho_n(u_n) = a^n$ and $p_m^*(b) = y$.

Next, suppose that $m < n$. We get $\cup : H^i(X/G) \rightarrow H^{i+1}(X/G)$ is an isomorphism for all $1 < i < n+m$ and $i \neq m-1, n$. Note that the map $\cup : H^{m-1}(X/G) \rightarrow H^m(X/G)$ is injective with $\text{im} \cong R$ with basis $\{a^m\}$, and for $i = n$ it is surjective with $\ker \cong R$ with basis $\{a_n\}$, where $\rho_n(u_n) = a^n$. This implies that $H^i(X) = 0$ for $1 < i < n+m$ and $i \neq m, n$, and $H^m(X) \cong H^n(X) \cong R$ with basis $\{y\}$ and $\{u_n\}$, respectively, where $p_m^*(b) = y$ and $\rho_n(u_n) = a^n$. Also, $H^{n+m} \cong R$ with basis $\{u_{n+m}\}$, where $\rho_{n+m}(u_{n+m}) = a^n b$.

Now, we compute the cohomology ring structure of X . In the Leray-Serre spectral sequence for the Borel fibration $X \hookrightarrow X_G \xrightarrow{\pi} B_G$, $d_{r'}(1 \otimes u_n) \neq 0$ for some $r' > 0$ and $d_r(1 \otimes y) = 0$ for all $r \geq 0$. Let it possible for $n > m$, $r' = n - m + 1$. Then we get $H^*(X_G) \cong R[u, v]/\langle u^{n-m+1}v - \alpha u^{n+1}, v^2 - \beta u^{2m} - \gamma u^m v, u^{n+i_0+1} \rangle$ where $\deg u = 1, \deg v = m$ and $\alpha, \beta, \gamma \in R$, $\beta = 0$ if $m > n$, which is a contradiction. Therefore, r' must be $n + 1$. This implies that $d_{n+1}(1 \otimes y u_n) \neq 0$. Consequently, $u_{m+n} = y u_n$. Obviously, $y^2 = 0$, and $u_n^2 = 0$ for $m \notin \{n, 2n\}$. If $m = n$ then $u_n^2 \neq u_{m+n}$ and so $u_n^2 = 0$. If $m = 2n$ then u_n^2 may be both zero or nonzero. If $u_n^2 = 0$ then $H^*(X) = \mathbb{Z}_2[y, u_n]/\langle y^2, u_n^2 \rangle$, where $\deg y = m$ and $\deg u_n = n$. If $u_n^2 \neq 0$ then the cohomology algebra of X is $\mathbb{Z}_2[u_n]/\langle u_n^4 \rangle$, where $\deg u_n = n$ and $m = 2n$. By [8, Theorem 4L.9], $n = 1, 2, 4$ or 8 . This realizes possibility (i) and (iii).

Case(b): When $\cup : H^0(X/G) \rightarrow H^2(X/G)$ maps 1 to b .

In this case, m must be 1 and $\text{im } p_1^* = \mathbb{Z}_2$ with basis $\{p_1^*(a)\}$. Also, $\text{im } p_i^* \cong \text{im } \rho_i \cong \mathbb{Z}_2$ with basis $\{p_i^*(a^i)\}$ and $\{a^{i-1}b\}$, respectively, for all $0 < i \leq n$. Consequently, $H^i(X) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ with basis $\{x^i, u_i\}$, where $\rho_i(u_i) = a^{i-1}b$ and $p_1^*(a) = x$. Obviously, $H^{n+1}(X) \cong \mathbb{Z}_2$ with basis $\{u_{n+1}\}$, where $\rho_{n+1}(u_{n+1}) = a^n b$. In the Leray-Serre spectral

sequence, $d_r(1 \otimes x) = 0$ for all $r \geq 0$ and $d_2(1 \otimes u_1) \neq 0$. This implies that for $1 \leq i \leq n$, $x^i u_1 \neq 0$, and hence $u_{i+1} = x^i u_1 + \alpha_i x^{i+1}$ for some $\alpha_i \in \mathbb{Z}_2$. As $d_2(1 \otimes x u_1) \neq 0$, we have $u_1^2 = \alpha x^2$ for some $\alpha \in \mathbb{Z}_2$. Therefore, $H^*(X) \cong \mathbb{Z}_2[x, u_1]/\langle x^{n+1}, u_1^2 + \alpha x^2 \rangle$, where $\deg x = \deg u_1 = 1$. This realizes possibility (ii).

Case(c): When $\cup : H^0(X/G) \rightarrow H^2(X/G)$ maps 1 to $a + b$.

In this case also m must be 1 and $\cup : H^i(X/G) \rightarrow H^{i+1}(X/G)$ is an isomorphism for all $0 < i \leq n$. Consequently, $H^i(X) = 0$ for $1 < i < n$. Note that $\ker p_1^* \cong \text{im } \rho_n \cong \mathbb{Z}_2$ with bases $\{a + b\}$ and $\{a^n + a^{n-1}b\}$, respectively. This implies that for $n \neq 1$, $H^1(X) \cong H^n(X) \cong \mathbb{Z}_2$ with bases $\{x\}$ and $\{u_n\}$, respectively, where $p_1^*(a) = x$ and $\rho_n(u_n) = a^n + a^{n-1}b$. Obviously, $H^{n+1}(X) \cong \mathbb{Z}_2$ with basis $\{u_{n+1}\}$, where $\rho_{n+1}(u_{n+1}) = a^n b$. In particular for $n = 1$, $H^1(X) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$, $H^2(X) \cong \mathbb{Z}_2$ with bases $\{x, u_1\}$ and $\{u_2\}$, respectively, where $p_1^*(a) = x, \rho_1(u_1) = a + b$ and $\rho_2(u_2) = ab$. We must have $u_{n+1} = x u_n$ and $u_n^2 = 0$. Thus, $X \sim_{\mathbb{Z}_2} \mathbb{S}^1 \times \mathbb{S}^n$. This realizes possibility (i). \square

Now, we determine covering dimension of coincidence set $A(f)$ of continuous maps $f : X \rightarrow \mathbb{R}^k$, where X is a finitistic space equipped with free involution and $X/G \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^n \times \mathbb{S}^m$.

Theorem 3.4. Let $G = \mathbb{Z}_2$ act freely on a finitistic space X with $X/G \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^n \times \mathbb{S}^m$, where $n > 4, m > 1$. If $f : X \rightarrow \mathbb{R}^k$ be any continuous map, then $\text{cov.dim}(A(f)) \geq n - k$ for $k \leq n$.

Proof. By Theorem 3.3, the Volovikov's index $\text{in}(X)$ [22] is n . Note that $\text{in}(A(f)) \geq \text{in}(X) - k$ [23], so we have $\text{in}(A(f)) \geq n - k$. As G acts freely on X , it induces a free action on $A(f)$. By the definition of $\text{in}(X)$, $H^{n-k}(B_G) \rightarrow H^{n-k}(A(f)_G)$ is injective. This implies that $\text{cohom.dim}(A(f)_G) \geq n - k$, where cohom.dim denotes the cohomological dimension of a space. As $A(f)/G$ and $A(f)_G$ are homotopy equivalent, $\text{cohom.dim}(A(f)/G) \geq n - k$. Consequently, $\text{cohom.dim}(A(f)) \geq n - k$ [15, Proposition A.11]. The result follows from the fact that $\text{cov.dim}(A(f)) \geq \text{cohom.dim}(A(f))$. \square

4. EXAMPLES

Consider the standard free actions of $G = \mathbb{S}^d$ on $\mathbb{S}^{(d+1)n+d}$, for $d = 0, 1$ or 3 , and trivial action on \mathbb{S}^m , then under the diagonal action G acts freely on $\mathbb{S}^{(d+1)n+d} \times \mathbb{S}^m$ with the orbit space $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$, where $\mathbb{F} = \mathbb{R}, \mathbb{C}$ or \mathbb{H} , respectively. This realizes the possibility (i) of Theorems 3.1, 3.2 and 3.3. Similarly, if we take free actions of $G = \mathbb{S}^d$

on \mathbb{S}^{2d+1} and trivial action on $\mathbb{F}\mathbb{P}^n$ then G acts freely on $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^{2d+1}$ with the orbit space $\mathbb{F}\mathbb{P}^n \times \mathbb{F}\mathbb{P}^1$, respectively. This realizes the possibility (iii) of Theorem 3.1, when $\alpha = 0$, the possibility (iv) of Theorem 3.2 and the possibility (ii) of Theorem 3.3. If we take free action of $G = \mathbb{S}^d$ on itself, for $d = 1$ or 3 , and trivial action on $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$, where $\mathbb{F} = \mathbb{C}$ or \mathbb{H} , respectively then G acts freely on $\mathbb{S}^d \times \mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$ with the orbit space $\mathbb{F}\mathbb{P}^n \times \mathbb{S}^m$. This realizes the possibility (ii) of Theorems 3.1 and 3.2.

Now, consider free action of \mathbb{Z}_4 on $\mathbb{S}^3 \subseteq \mathbb{C}^2$ defined by $(z_1, z_2) \mapsto (z_1 e^{2\pi i/4}, z_2 e^{2\pi i/4})$. This induces a free involution on $\mathbb{R}\mathbb{P}^3$ with the orbit space $L^3(4, 1)$. We know that $L^3(4, 1) \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^1 \times \mathbb{S}^2$. Recall that if $G = \mathbb{Z}_2$ acts freely on a finitistic space X with the mod 2 cohomology $\mathbb{C}\mathbb{P}^3$, then the orbit spaces X/G is the mod 2 cohomology $\mathbb{R}\mathbb{P}^2 \times \mathbb{S}^4$ [16]. These examples realizes the possibility (iii) of Theorem 3.3 for $n = 1$ and $n = 2$, respectively.

Remark. Let X be a connected Hausdorff space equipped with free action of $G = \mathbb{Z}_2$. If the orbit space $X/G = \mathbb{R}\mathbb{P}^n \times \mathbb{S}^m$ then X is homeomorphic to $\mathbb{S}^n \times \mathbb{S}^m$. The significance of Theorem 3.3 lies with the fact that if $X/G \sim_{\mathbb{Z}_2} \mathbb{R}\mathbb{P}^n \times \mathbb{S}^m$ then X may have the mod 2 cohomology isomorphic to $\mathbb{R}\mathbb{P}^n \times \mathbb{S}^1$ or $\mathbb{S}^n \times \mathbb{S}^1$ for $m = 1$; and $\mathbb{S}^4 \times \mathbb{S}^2$ or $\mathbb{C}\mathbb{P}^3$ for $m = 4$ and $n = 2$.

REFERENCES

- [1] C. Biasi, D. De Mattos and E. L. Dos Santos, A Borsuk-Ulam theorem for maps from a sphere to a generalized manifold, *Geom. Dedicata* **107** (2004), 101–110. MR2110757
- [2] G. E. Bredon, Cohomological aspects of transformation groups, in *Proc. Conf. on Transformation Groups (New Orleans, La., 1967)*, 245–280, Springer, New York. MR0244990
- [3] P. Dey and M. Singh, Free actions of some compact groups on Milnor manifolds, *Glasg. Math. J.* **61** (2019), no. 3, 727–742. MR3991367
- [4] T. tom Dieck, *Transformation groups*, De Gruyter Studies in Mathematics, 8, Walter de Gruyter & Co., Berlin, 1987. MR0889050
- [5] R. M. Dotzel, T. B. Singh and S. P. Tripathi, The cohomology rings of the orbit spaces of free transformation groups of the product of two spheres, *Proc. Amer. Math. Soc.* **129** (2001), no. 3, 921–930. MR1712925
- [6] H. Duan and C. Liang, Circle bundles over 4-manifolds, *Arch. Math. (Basel)* **85** (2005), no. 3, 278–282. MR2172386
- [7] J. Harvey, M. Kerin and K. Shankar, Semi-free actions with manifold orbit spaces, *Doc. Math.* **25** (2020), 2085–2114. MR4187719
- [8] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. MR1867354
- [9] J. Kaur and H. K. Singh, On the existence of free action of \mathbb{S}^3 on certain finitistic mod P cohomology spaces, *J. Indian Math. Soc. (N.S.)* **82** (2015), no. 3-4, 97–106. MR3467619
- [10] A. Kumari and H. K. Singh, Fixed point free actions of spheres and equivariant maps, *Topology Appl.* **305** (2022), Paper No. 107886, 14 pp. MR4335843

- [11] A. Kumari and H. K. Singh, Cohomology classification of spaces with free S^3 -Actions, arXiv:2104.05377
- [12] J. McCleary, *A user's guide to spectral sequences*, second edition, Cambridge Studies in Advanced Mathematics, 58, Cambridge University Press, Cambridge, 2001. MR1793722
- [13] A. M. M. Morita, D. de Mattos and P. L. Q. Pergher, The cohomology ring of orbit spaces of free \mathbb{Z}_2 -actions on some Dold manifolds, Bull. Aust. Math. Soc. **97** (2018), no. 2, 340–348. MR3772666
- [14] H. J. Munkholm, A Borsuk-Ulam theorem for maps from a sphere to a compact topological manifold, Illinois J. Math. **13** (1969), 116–124. MR0238324
- [15] D. Quillen, The spectrum of an equivariant cohomology ring. I, II, Ann. of Math. (2) **94** (1971), 549–572; ibid. (2) **94** (1971), 573–602. MR0298694
- [16] H. K. Singh and T. B. Singh, Fixed point free involutions on cohomology projective spaces, Indian J. Pure Appl. Math. **39** (2008), no. 3, 285–291. MR2434244
- [17] H. K. Singh and T. B. Singh, The cohomology of orbit spaces of certain free circle group actions, Proc. Indian Acad. Sci. Math. Sci. **122** (2012), no. 1, 79–86. MR2909586
- [18] M. Singh, Orbit spaces of free involutions on the product of two projective spaces, Results Math. **57** (2010), no. 1-2, 53–67. MR2603012
- [19] M. Singh, Cohomology algebra of orbit spaces of free involutions on lens spaces, J. Math. Soc. Japan **65** (2013), no. 4, 1055–1078. MR3127816
- [20] J. C. Su, Transformation groups on cohomology projective spaces, Trans. Amer. Math. Soc. **106** (1963), 305–318. MR0143839
- [21] R. G. Swan, A new method in fixed point theory, Comment. Math. Helv. **34** (1960), 1–16. MR0115176
- [22] A. Yu. Volovikov, Russian Acad. Sci. Sb. Math. **76** (1993), no. 2, 361–387; translated from Mat. Sb. **183** (1992), no. 7, 115–144. MR1186977
- [23] A. Yu. Volovikov, Sb. Math. **191** (2000), no. 9-10, 1259–1277; translated from Mat. Sb. **191** (2000), no. 9, 3–22. MR1805595

Anju Kumari, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI – 110007, INDIA.

Email address: anjukumari0702@gmail.com

Hemant Kumar Singh, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI DELHI – 110007, INDIA.

Email address: hemantksingh@maths.du.ac.in