Semiparametric Discrete Choice Models for Bundles

Fu Ouyang*
University of Queensland

Thomas Tao Yang[†]
Australian National University

November 13, 2023

Abstract

We propose two approaches to estimate semiparametric discrete choice models for bundles. Our first approach is a kernel-weighted rank estimator based on a matching-based identification strategy. We establish its complete asymptotic properties and prove the validity of the nonparametric bootstrap for inference. We then introduce a new multi-index least absolute deviations (LAD) estimator as an alternative, of which the main advantage is its capacity to estimate preference parameters on both alternative- and agent-specific regressors. Both methods can account for arbitrary correlation in disturbances across choices, with the former also allowing for interpersonal heteroskedasticity. We also demonstrate that the identification strategy underlying these procedures can be extended naturally to panel data settings, producing an analogous localized maximum score estimator and a LAD estimator for estimating bundle choice models with fixed effects. We derive the limiting distribution of the former and verify the validity of the numerical bootstrap as an inference tool. All our proposed methods can be applied to general multi-index models. Monte Carlo experiments show that they perform well in finite samples.

JEL classification: C13, C14, C35.

Keywords: Multi-index models; Rank correlation; Maximum score; Least absolute deviations; Panel data.

^{*}Corresponding author. School of Economics, The University of Queensland, St Lucia, QLD 4072, Australia. Tel: +61 451 478 622. Email: f.ouyang@uq.edu.au.

[†]Research School of Economics, The Australian National University, Canberra, ACT 0200, Australia. Email: tao.yang@anu.edu.au.

1 Introduction

In many circumstances, consumers purchase bundles of goods (e.g., both chips and salsa) or services (e.g., a combination of mobile phone, home internet, and cable TV plans), instead of a single good or service. The literature on bundle choice is less well developed than that on ordinary multinomial choice models. An important but less extensively studied empirical question in industrial organization and marketing research relates to complementary or substitutive effects that may explain the bundle choice behavior of consumers. We refer readers to Berry et al. (2014) for a review of this literature. In an empirical study, Gentzkow (2007) estimated a parametric bundle choice model analyzing demand for print and online newspapers. Similarly, using aggregate data, Fan (2013) examined ownership consolidation in the newspaper market, where households on the demand side may purchase two newspapers as a bundle. A substantial literature focuses on bundle choice behavior, analysis of product demand, pricing strategy, customer subscriptions, and brand collaboration, among others. Examples include Manski and Sherman (1980), Train, McFadden, and Ben-Akiva (1987), Hendel (1999), Chung and Rao (2003), Dubé (2004), Nevo, Rubinfeld, and McCabe (2005), Augereau, Greenstein, and Rysman (2006), Song and Chintagunta (2006), Foubert and Gijsbrechts (2007), Liu, Chintagunta, and Zhu (2010), Gentzkow, Shapiro, and Sinkinson (2014), Lewbel and Nesheim (2019), Kim, Misra, and Shapiro (2020), Ershov, Laliberté, Marcoux, and Orr (2021), and Iaria and Wang (2021). The models in most of these applications are parametric.¹

There has been a growing interest in recent years in developing nonparametric and semiparametric methods for discrete choice models with bundles. Sher and Kim (2014) explored the identification of bundle choice models without stochastic components in the utility from the perspective of microeconomic theory, assuming a finite population of consumers. Dunker, Hoderlein, and Kaido (2022) studied nonparametric identification of market-level demand models in a general framework in which the demand for bundles is nested. Fox and Lazzati (2017) provided nonparametric identification results for both single-agent discrete choice bundle models and binary games of complete information, and established the mathematical equivalence between these models. Allen and Rehbeck (2019) studied nonparametric identification of a class of latent utility models with additively separable unobserved heterogeneity by exploiting asymmetries in cross-partial derivatives of conditional average demands. Ouyang, Yang, and Zhang (2020) used the "identification at infinity" for bundle choice models in cross-sectional settings. Allen and Rehbeck (2022) established partial identification results for complementarity (substitutability) in a latent utility model with binary

¹See Gentzkow (2007) pp. 722–723 for a brief review of pervasive parametric methods in the literature.

quantities of two goods. These methods assume cross-sectional data. Wang (2023) proposed a semiparametric framework for identifying and testing substitution and complementarity patterns in multinomial choice models with bundles, which can be applied to both cross-sectional and panel data settings.

Our paper differs from these studies in many respects. Using individual-level data, we study semiparametric point identification and estimation of preference coefficients for bundle choice models in both cross-sectional and panel data settings. Our methods do not impose distributional constraints on unobserved error terms and allow for arbitrary correlation among them. The robustness afforded by the distribution-free specification makes our approaches competitive alternatives to existing parametric methods. Our paper is most closely related to Wang (2023) in terms of model specification and the assumptions regarding the dependence of unobservables on observed covariates. However, our proposed estimation and inference methods are built on distinct identification inequalities and criterion functions.

For the cross-sectional model, our identification results rely on exogeneity conditions similar to Allen and Rehbeck (2019, 2022)) and Wang (2023), instead of either the exclusion restrictions ("special regressors") in Fox and Lazzati (2017) or aggregate data and valid instrumental variables required by Dunker et al. (2022). We first consider a maximum rank correlation (MRC) estimator in the spirit of Han (1987), using a matching insight. A recent study by Khan, Ouyang, and Tamer (2021) used a similar method to study multinomial choice models. We validate nonparametric bootstrapping for inference and introduce a criterion-based test for interaction effects. Additionally, we propose a novel least absolute deviations (LAD) procedure as an alternative to the matching-based MRC method. The LAD estimator offers the advantage of estimating coefficients for both choice- and individual-specific covariates. It is worth noting that while we introduce these methods in the context of discrete choice models with bundles, their fundamental principles can be readily adapted to estimate a wide range of "multi-index" models with minor modifications.

We then extend the insight gained from the identification of cross-sectional models to panel data models with fixed effects. In this context, our identification relies on a mild conditional homogeneity assumption, which is also employed by Pakes and Porter (2021), Shi, Shum, and Song (2018), and Gao and Li (2020), among others, for studying ordinary panel data multinomial choice models. Based on a series of moment inequalities, we propose a maximum-score-type estimation procedure in the spirit of Manski (1987) and derive its complete asymptotic properties. We also justify the validity of applying the numerical bootstrap for inference and propose a test for the presence of interaction effects. As in the

cross-sectional model, we also introduce a panel data LAD estimation method, which offers the same advantages of accommodating both choice- and individual-specific covariates. Our methods for constructing these estimations are general and can be applied to models with similar multi-index structures.

This paper contributes to the fast-growing literature on a broad class of multi-index models. Leading examples are the multinomial choice model and its variants. In addition to the previously mentioned studies, this literature includes works by Lee (1995), Lewbel (2000), Altonji and Matzkin (2005), Fox (2007), Berry and Haile (2009), Ahn, Ichimura, Powell, and Ruud (2018), Yan and Yoo (2019), and Chernozhukov, Fernández-Val, and Newey (2019), to name a few. These approaches, however, do not directly extend to bundle choice models because of the non-mutually exclusive choice set and distinctive random utilities associated with bundles. Another example of multi-index models involves the study of dyadic networks, which has gained increasing attention among researchers. Some of the notable works in this area include Graham (2017), Toth (2017), Candelaria (2020), and Gao, Li, and Xu (2023).

The remainder of this paper is organized as follows. Section 2 presents our MRC and LAD methods for the cross-sectional bundle choice model. Section 3 studies a panel data bundle choice model with fixed effects, introducing our localized MS and panel data LAD procedures. In Section 4, we assess the finite sample performance of these proposed procedures through Monte Carlo experiments. Section 5 concludes the paper. All tables, proofs, extensions, and further discussions are collected in the supplementary appendix (Appendixes A–F).

For ease of reference, the notations maintained throughout this paper are listed here.

Notation. All vectors are column vectors. \mathbb{R}^p is a p-dimensional Euclidean space equipped with the Euclidean norm $\|\cdot\|$, and $\mathbb{R}_+ \equiv \{x \in \mathbb{R} | x \geq 0\}$. We reserve letters i and m for indexing agents, j and l for indexing alternatives, and s and t for indexing time periods. The first element of a vector v is denoted by $v^{(1)}$ and the sub-vector comprising its remaining elements is denoted by \tilde{v} . $P(\cdot)$ and $\mathbb{E}[\cdot]$ denote probability and expectation, respectively. $1[\cdot]$ is an indicator function that equals 1 when the event in the brackets occurs, and 0 otherwise. For two random vectors U and V, the notation $U \stackrel{d}{=} V|\cdot$ means that U and V have identical distribution conditional on \cdot , and $U \perp V|\cdot$ means that U and V are independent conditional on \cdot . Symbols \setminus , ', \Leftrightarrow , \Rightarrow , \propto , $\stackrel{d}{\to}$, and $\stackrel{p}{\to}$ represent set difference, matrix transposition, if and only if, implication, proportionality, convergence in distribution, and convergence in probability, respectively. For any (random) positive sequences $\{a_N\}$ and $\{b_N\}$, $a_N = O(b_N)$ ($O_p(b_N)$) means that a_N/b_N is bounded (bounded in probability) and $a_N = o(b_N)$ ($o_p(b_N)$) means that $a_N/b_N \to 0$ ($a_N/b_N \stackrel{p}{\to} 0$).

2 Cross-Sectional Model

Throughout this paper, we focus on a choice model in which the choice set \mathcal{J} consists of three mutually exclusive alternatives (numbered 0–2) and a bundle of alternatives 1 and 2; that is, $\mathcal{J} = \{0, 1, 2, (1, 2)\}$. This simple model is sufficient to illustrate the main intuition that runs through both cross-sectional and panel data models.²

For ease of exposition, we re-number alternatives in \mathcal{J} with 2-dimensional vectors of binary indicators $d = (d_1, d_2) \in \{0, 1\} \times \{0, 1\}$, where d_1 and d_2 indicate if alternative 1 and 2 are chosen, respectively. In this way the choice set \mathcal{J} can be one-to-one mapped to the set $\mathcal{D} = \{(0, 0), (1, 0), (0, 1), (1, 1)\}$. An agent chooses the alternative in \mathcal{D} to maximize the latent utility

$$U_d = \sum_{j=1}^{2} F_j(X_j'\beta, \epsilon_j) \cdot d_j + \eta \cdot F_b(W'\gamma) \cdot d_1 \cdot d_2, \tag{2.1}$$

where $F_j(\cdot,\cdot)$'s and $F_b(\cdot)$ are unknown (to the econometrician) $\mathbb{R}^2 \to \mathbb{R}$ and $\mathbb{R} \to \mathbb{R}$, respectively, functions strictly monotonic in each of their arguments, $X_j \in \mathbb{R}^{k_1}$ collects alternative-specific covariates affecting the utility associated with stand-alone alternative $j, W \in \mathbb{R}^{k_2}$ is a vector of explanatory variables characterizing the interaction effects of the bundle (e.g., bundle discount), $(\epsilon_1, \epsilon_2, \eta) \in \mathbb{R}^2 \times \mathbb{R}_+$ captures unobserved (to the econometrician) heterogeneous effects, and $(\beta, \gamma) \in \mathbb{R}^{k_1+k_2}$ are unknown preference parameters to estimate.

We assume that $F_b(0) = 0$ so that the sign of $W'\gamma$ has its economic meaning (see below for details). For instance, $F_b(x)$ can be x or x^3 . The specification of model (2.1) takes the linear forms in Fox and Lazzati (2017) as a special case. We note that Fox and Lazzati (2017) does rely on the additive separability while ours does not. The utility of choosing (0,0) is normalized to 0. The utilities of choosing (1,0) and (0,1) are $F_1(X'_1\beta,\epsilon_1)$ and $F_2(X'_2\beta,\epsilon_2)$, respectively. The utility of choosing the bundle is the sum of the two standalone utilities plus the interaction term $\eta \cdot F_b(W'\gamma)$ which captures either complementary $(W'\gamma > 0)$ or substitution $(W'\gamma < 0)$ effects. (ϵ_1, ϵ_2) are idiosyncratic shocks associated with each stand-alone alternative as in common multinomial choice models, and η reflects unobserved heterogeneity for the bundle. In what follows, we assume $\eta \geq 0$ so that $W'\gamma > 0$ indicates interaction effect to be complementary, and otherwise substitutive.

²We discuss models with more alternatives in Appendix E.3.

Given the latent utility model (2.1), the observed dependent variable Y_d takes the form

$$Y_d = 1[U_d > U_{d'}, \forall d' \in \mathcal{D} \setminus d].^3 \tag{2.2}$$

Let $Z \equiv (X_1, X_2, W)$. The probabilities of choosing alternatives $d \in \mathcal{D}$ can be expressed as

$$(P(Y_{(0,0)} = 1|Z), P(Y_{(1,0)} = 1|Z), P(Y_{(0,1)} = 1|Z), P(Y_{(1,1)} = 1|Z))'$$

$$= \begin{bmatrix} P(\max\{F_1(X_1'\beta, \epsilon_1), F_2(X_2'\beta, \epsilon_2), F_1(X_1'\beta, \epsilon_1) + F_2(X_2'\beta, \epsilon_2) + \eta \cdot F_b(W'\gamma)\} < 0|Z) \\ P(\max\{0, F_2(X_2'\beta, \epsilon_2), F_1(X_1'\beta, \epsilon_1) + F_2(X_2'\beta, \epsilon_2) + \eta \cdot F_b(W'\gamma)\} < F_1(X_1'\beta, \epsilon_1)|Z) \\ P(\max\{0, F_1(X_1'\beta, \epsilon_1), F_1(X_1'\beta, \epsilon_1) + F_2(X_2'\beta, \epsilon_2) + \eta \cdot F_b(W'\gamma)\} < F_2(X_2'\beta, \epsilon_2)|Z) \\ P(\max\{0, F_1(X_1'\beta, \epsilon_1), F_2(X_2'\beta, \epsilon_2)\} < F_1(X_1'\beta, \epsilon_1) + F_2(X_2'\beta, \epsilon_2) + \eta \cdot F_b(W'\gamma)|Z) \end{bmatrix}.$$

2.1 Rank Correlation Method

Section 2.1.1 shows the identification result. Section 2.1.2 presents the MRC estimator and its asymptotic properties. Bootstrap inference procedure and a criterion-based test for interaction effects are discussed in Sections 2.1.3 and 2.1.4, respectively.

2.1.1 Identification

When $(\epsilon_1, \epsilon_2, \eta) \perp Z$, expression (2.3) implies that $P(Y_{(1,0)} = 1 | X_1 = x_1, X_2 = x_2, W = w)$ and $P(Y_{(1,1)} = 1 | X_1 = x_1, X_2 = x_2, W = w)$ are both increasing in $x'_1\beta$ for some constant vectors x_2 and w. That is, for $d_2 \in \{0, 1\}$,

$$x_1'\beta \ge \bar{x}_1'\beta \Leftrightarrow P(Y_{(1,d_2)} = 1|Z = (x_1, x_2, w)) \ge P(Y_{(1,d_2)} = 1|Z = (\bar{x}_1, x_2, w)).$$
 (2.4)

Similarly, for cases with $d_1 = 0$ and any d_2 , $x'_1\beta \geq \bar{x}'_1\beta$ is the "if-and-only-if" condition to the second inequality in (2.4) but with " \leq " instead.

Once β is identified, we can move on to identify γ using the following moment inequalities for $w'\gamma$ constructed by fixing $V(\beta) \equiv (X'_1\beta, X'_2\beta)$ at some constant vector (v_1, v_2) . For

Throughout this paper, we do not consider the case in which $U_d = U_{d'}$. As will be clear below, such utility ties occur with probability 0 under our identification consitions.

$$d = (1, 1),$$

$$w'\gamma \ge \bar{w}'\gamma$$

 $\Leftrightarrow P(Y_{(1,1)} = 1|V(\beta) = (v_1, v_2), W = w) \ge P(Y_{(1,1)} = 1|V(\beta) = (v_1, v_2), W = \bar{w}).$ (2.5)

For all $d \in \mathcal{D} \setminus (1,1)$, $w'\gamma \geq \bar{w}'\gamma$ is the "if-and-only-if" condition for the second inequality in (2.5) but with " \leq " instead.

To implement the above idea, we assume a random sample in Assumption C1 below. Thus, we can take two independent copies of (Y, Z), i.e., (Y_i, Z_i) and (Y_m, Z_m) from the sample. For (2.4), we can match X_2 and W for this two observations, so then we can rank their choice probabilities of $Y_{(1,d_2)}$ using the index $X'_1\beta$. This idea forms the basis of our MRC estimator, which we introduce in the next section. Note that γ can be alternatively identified by matching (X_1, X_2) across agents. However, some exploratory simulation studies have shown that matching the estimated index $V(\hat{\beta})$ (as suggested in (2.5)) can yield better finite sample performance than element-by-element matching.

To establish the identification of β and γ based on (2.4)–(2.5), the following conditions are sufficient:

- C1 (i) $\{(Y_i, Z_i)\}_{i=1}^N$ are i.i.d. across i, (ii) $(\epsilon_1, \epsilon_2, \eta) \perp Z$, and (iii) the joint distribution of $(\epsilon_1, \epsilon_2, \eta)$ is absolutely continuous on $\mathbb{R}^2 \times \mathbb{R}_+$.
- C2 For any pair of (i, m) and j = 1, 2, denote $X_{imj} = X_{ij} X_{mj}$, $W_{im} = W_i W_m$, and $V_{im}(\beta) = V_i(\beta) V_m(\beta)$. Then, (i) $X_{im1}^{(1)}(X_{im2}^{(1)})$ has almost everywhere (a.e.) positive Lebesgue density on \mathbb{R} conditional on $\tilde{X}_{im1}(\tilde{X}_{im2})$ and conditional on (X_{im2}, W_{im}) $((X_{im1}, W_{im}))$ in a neighborhood of (X_{im2}, W_{im}) $((X_{im1}, W_{im}))$ near zero, (ii) Elements in $X_{im1}(X_{im2})$, conditional on $(X_{im2}, W_{im})((X_{im1}, W_{im}))$ in a neighborhood of $(X_{im2}, W_{im})((X_{im1}, W_{im}))$ near zero, are linearly independent, (iii) $W_{im}^{(1)}$ has a.e. positive Lebesgue density on \mathbb{R} conditional on \tilde{W}_{im} and conditional on $V_{im}(\beta)$ in a neighborhood of $V_{im}(\beta)$ near zero, and (iv) Elements in W_{im} , conditional on $V_{im}(\beta)$ in a neighborhood of $V_{im}(\beta)$ near zero, are linearly independent.
- C3 $(\beta, \gamma) \in \mathcal{B} \times \mathcal{R}$, where $\mathcal{B} = \{b \in \mathbb{R}^{k_1} | b^{(1)} = 1, ||b|| \leq C_b\}$ and $\mathcal{R} = \{r \in \mathbb{R}^{k_2} | r^{(1)} = 1, ||r|| \leq C_r\}$ for some constants $C_b, C_r > 0$.
- C4 $F_1(\cdot,\cdot)$, $F_2(\cdot,\cdot)$, and $F_b(\cdot)$ are respectively $\mathbb{R}^2 \to \mathbb{R}$, $\mathbb{R}^2 \to \mathbb{R}$, and $\mathbb{R} \to \mathbb{R}$ functions strictly increasing in each of their arguments. $F_b(0) = 0$.

From the previous discussion, it can be seen that Assumptions C1 and C4 are the keys to establishing the moment inequalities (2.4)–(2.5). It is important to note that Assumption C1 allows for arbitrary correlation among $(\epsilon_1, \epsilon_2, \eta)$. Assumptions C2(i) and C2(iii) are standard requirements for MRC and MS types of estimators. Assumptions C2(ii) and C2(iv) are regular rank conditions. Furthermore, Assumption C3 defines a compact parameter space and normalizes the scale of coefficients, which is standard practice for discrete choice models. Essentially, we assume that $\beta^{(1)}$ and $\gamma^{(1)}$ are not equal to zero.

The following identification result for model (2.1)–(2.2) is proved in Appendix B.

Theorem 2.1. Suppose Assumptions C1–C4 hold. Then β and γ are identified.

Remark 2.1. The matching-based method discussed here is distribution-free, accommodating arbitrary correlation among $(\epsilon_1, \epsilon_2, \eta)$ and interpersonal heteroskedasticity. Furthermore, as demonstrated in Sections 2.1.2–2.1.3, the resulting estimator is computationally tractable and has an asymptotic normal distribution, rendering inference relatively straightforward, particularly when employing bootstrap methods. However, this method has two main limitations. Firstly, it becomes difficult to apply when the model has many covariates, as this decreases the number of potential matches. Secondly, in cases where "common regressors" are present, it cannot estimate coefficients related to these variables unless their coefficients are assumed to be not "alternative-specific".

2.1.2 Localized MRC Estimator

The local monotonic relations established in (2.4)–(2.5) naturally motivate a two-step localized MRC estimation procedure. Note that the probability of obtaining perfectly matched observations is zero for continuous regressors. Following the literature, we propose to use kernel weights as an approximation to the matching. These steps are described in turn below.

In the first step, we consider the localized MRC estimator $\hat{\beta}$ of β , analogous to the MRC estimator proposed in Han (1987) and Khan et al. (2021). Specifically, $\hat{\beta} = \arg \max_{\beta \in \mathcal{B}} \mathcal{L}_{N,\beta}^K(b)$. $\mathcal{L}_{N,\beta}^K(b)$ is defined as

$$\mathcal{L}_{N,\beta}^{K}(b) = \sum_{i=1}^{N-1} \sum_{m>i} \sum_{d \in \mathcal{D}} \{ \mathcal{K}_{h_N}(X_{im2}, W_{im}) (Y_{md} - Y_{id}) \operatorname{sgn}(X'_{im1}b) \cdot (-1)^{d_1} + \mathcal{K}_{h_N}(X_{im1}, W_{im}) (Y_{md} - Y_{id}) \operatorname{sgn}(X'_{im2}b) \cdot (-1)^{d_2} \}$$
(2.6)

with $\mathcal{K}_{h_N}(X_{imj}, W_{im}) \equiv h_N^{-(k_1+k_2)} \prod_{\iota=1}^{k_1} K(X_{imj,\iota}/h_N) \prod_{\iota=1}^{k_2} K(W_{im,\iota}/h_N)$, where $X_{imj,\iota}(W_{im,\iota})$ is the ι -th element of vector $X_{imj}(W_{im})$, $K(\cdot)$ is a standard kernel density function, and h_N is a bandwidth sequence that converges to 0 as $N \to \infty$. Obviously, $\mathcal{K}_{h_N}(X_{imj}, W_{im}) \approx [X_{imj} = 0, W_{im} = 0]$ for j = 1, 2, as $h_N \to 0$.

In the second step, we obtain $\hat{\gamma} = \arg \max_{\gamma \in \mathcal{R}} \mathcal{L}_{N,\gamma}^K(r; \hat{\beta})$ with

$$\mathcal{L}_{N,\gamma}^{K}(r;\hat{\beta}) = \sum_{i=1}^{N-1} \sum_{m>i} \mathcal{K}_{\sigma_N}(V_{im}(\hat{\beta}))(Y_{i(1,1)} - Y_{m(1,1)}) \operatorname{sgn}(W'_{im}r), \tag{2.7}$$

where $\mathcal{K}_{\sigma_N}(V_{im}(\hat{\beta})) = \sigma_N^{-2} K(X'_{im1}\hat{\beta}/\sigma_N) K(X'_{im2}\hat{\beta}/\sigma_N)$ and σ_N is a bandwidth sequence that converges to 0 as $N \to \infty$. Again, $\mathcal{K}_{\sigma_N}(V_{im}(\hat{\beta})) \approx 1[V_{im}(\hat{\beta}) = 0]$ as $\sigma_N \to 0$.

The rest of this section establishes the asymptotic properties of the estimators computed based on criterion functions (2.6) and (2.7). To do so, we need to place Assumptions C5–C7 in addition to Assumptions C1–C4 and introduce some new notations. Assumption C5 is a purely technical assumption and standard in the literature. To save space, we present it in Appendix B before the proof of Theorem 2.2. Here we only present Assumptions C6–C7 since they are of practical importance.

C6 $K(\cdot)$ is continuously differentiable and assumed to satisfy: (i) $\sup_{v} |K(v)| < \infty$, (ii) $\int K(v) dv = 1$, (iii) for any positive integer $\iota \leq \max\{\kappa_{\beta}, \kappa_{\gamma}\}$ where κ_{β} is an even integer greater than $k_1 + k_2$ and κ_{γ} is an even integer greater than $k_1 + k_2$ and k_{γ} is an even integer greater than $k_1 + k_2$ and $k_{\gamma} = 0$ if $k_{\gamma} = 0$

C7
$$h_N$$
 and σ_N are sequences of positive numbers such that as $N \to \infty$: (i) $h_N \to 0$, $\sigma_N \to 0$, (ii) $\sqrt{N} h_N^{k_1 + k_2} \to \infty$, $\sqrt{N} \sigma_N^2 \to \infty$, and (iii) $\sqrt{N} h_N^{\kappa_\beta} \to 0$, $\sqrt{N} \sigma_N^{\kappa_\gamma} \to 0$.

Assumptions C6 and C7 impose mild and standard restrictions on kernel functions and tuning parameters. These assumptions are essential to establish the consistency of the estimator and to ensure that the bias term becomes asymptotically negligible when deriving the limiting distribution. Assumption C7 requires the order of the kernel function to be at least $\max \{\kappa_{\beta}, \kappa_{\gamma}\}$. Using similar arguments employed in Han (1987), Sherman (1993, 1994a,b),

⁴In practice, kernel functions and bandwidths can be different for each univariate matching variable. Here we assume they are the same for all covariates just for notational convenience.

⁵Here we use the same kernel function $K(\cdot)$ for estimating β and γ to avoid additional notations. However, in practice, one can use kernel functions with different orders by replacing $\max\{\kappa_{\beta}, \kappa_{\gamma}\}$ with κ_{β} and κ_{γ} , respectively.

and Abrevaya, Hausman, and Khan (2010), we show the \sqrt{N} -consistency and asymptotic normality of the MRC estimators. The proof can be found in Appendix B.

Theorem 2.2. If Assumptions C1–C7 hold, then we have

(i) $\sqrt{N}(\hat{\beta} - \beta) \stackrel{d}{\to} N(0, 4 \{\mathbb{E}[\nabla^2 \varrho_i(\beta)]\}^{-1} \mathbb{E}[\nabla \varrho_i(\beta) \nabla \varrho_i(\beta)'] \{\mathbb{E}[\nabla^2 \varrho_i(\beta)]\}^{-1})$ or, alternatively, $\hat{\beta} - \beta$ has the linear representation:

$$\hat{\beta} - \beta = -\frac{2}{N} \left\{ \mathbb{E}[\nabla^2 \varrho_i(\beta)] \right\}^{-1} \sum_{i=1}^{N} \varrho_i(\beta) + o_p(N^{-1/2}),$$

with $\varrho_i(\cdot)$ defined in (B.1).

(ii)
$$\sqrt{N}(\hat{\gamma} - \gamma) \stackrel{d}{\to} N(0, \mathbb{E}[\nabla^2 \tau_i(\gamma)]^{-1} \mathbb{E}\left[\Delta_i \Delta_i'\right] \mathbb{E}[\nabla^2 \tau_i(\gamma)]^{-1}), \text{ where}$$

$$\Delta_i \equiv 2\nabla \tau_i(\gamma) + 2\mathbb{E}\left[\nabla_{13}^2 \mu\left(V_i(\beta), V_i(\beta), \gamma\right)\right] \left\{\mathbb{E}[\nabla^2 \varrho_i(\beta)]\right\}^{-1} \varrho_i(\beta),$$

with $\tau_i(\cdot)$ and $\mu(\cdot,\cdot,\cdot)$ defined in (B.2) and (B.3), respectively, and ∇^2_{13} denotes the second order derivative w.r.t. the first and third arguments.

We provide some intuition on the \sqrt{N} convergence rate in Appendix E.

2.1.3 Inference

Theorem 2.2 indicates that our localized MRC estimators are asymptotically normal and have asymptotic variances with the usual sandwich structure. The expressions for the asymptotic variances consist of first and second derivatives of the limit of the expectation of the maximands in (2.6) and (2.7). To use these results for statistical inference, Sherman (1993) proposed applying the numerical derivative method of Pakes and Pollard (1989). Hong, Mahajan, and Nekipelov (2015) investigated the application of the numerical derivative method in extremum estimators, including second-order U-statistics. As an alternative, Cavanagh and Sherman (1998) suggested nonparametrically estimating these quantities. These procedures, however, require selecting additional tuning parameters.

To avoid this complexity, we propose conducting inference using the classic nonparametric bootstrap for ease of implementation. Subbotin (2007) proved the consistency of the

nonparametric bootstrap for Han's (1987) MRC estimator. The structure of the criterion functions of our estimators is similar to that of the standard MRC estimator, but they do differ in two ways. First, our estimators require matching and thus contain kernel functions. Second, the criterion function for estimating $\hat{\gamma}$ contains a first-step (\sqrt{N} -consistent) $\hat{\beta}$ to approximate the true value of β . These two differences usually do not make the bootstrap inconsistent. For example, Horowitz (2001) demonstrated the consistency of the bootstrap for a range of estimators involved with kernel functions; Chen, Linton, and van Keilegom (2003) demonstrated the consistency of the bootstrap for estimators with a first-step estimation component and a well-behaved criterion function.

The bootstrap algorithm is standard. Draw $\{(Y_i^*, Z_i^*)\}_{i=1}^N$ independently from the sample $\{(Y_i, Z_i)\}_{i=1}^N$ with replacement. Obtain the bootstrap estimator $\hat{\beta}^*$ is obtained from

$$\hat{\beta}^* = \arg\max_{b \in \mathcal{B}} \mathcal{L}_{N,\beta}^{K*}(b) \equiv \arg\max_{b \in \mathcal{B}} \sum_{i=1}^{N-1} \sum_{m>i} \sum_{d \in \mathcal{D}} \{ \mathcal{K}_{h_N}(X_{im2}^*, W_{im}^*) Y_{mid}^* \operatorname{sgn}(X_{im1}^{*\prime}b) \cdot (-1)^{d_1} + \mathcal{K}_{h_N}(X_{im1}^*, W_{im}^*) Y_{mid}^* \operatorname{sgn}(X_{im2}^{*\prime}b) \cdot (-1)^{d_2} \}.$$
(2.8)

Then, obtain the bootstrap estimator $\hat{\gamma}^*$ through

$$\hat{\gamma}^* = \arg\max_{r \in \mathcal{R}} \mathcal{L}_{N,\gamma}^{K*}(r, \hat{\beta}^*) \equiv \arg\max_{r \in \mathcal{R}} \sum_{i=1}^{N-1} \sum_{m>i} \mathcal{K}_{\sigma_N}(V_{im}^*(\hat{\beta}^*)) Y_{im(1,1)}^* \operatorname{sgn}(W_{im}^{*\prime}r).$$
 (2.9)

Repeat the above steps for B times to yield a sequence of bootstrap estimates $\{(\hat{\beta}^*_{(b)}, \hat{\gamma}^*_{(b)})\}_{b=1}^B$.

Theorem 2.3. Suppose Assumptions C1–C7 hold. Then, we have

$$\sqrt{N}(\hat{\beta}^* - \hat{\beta}) \stackrel{d}{\to} N(0, 4 \left\{ \mathbb{E}[\nabla^2 \varrho_i(\beta)] \right\}^{-1} \mathbb{E}[\nabla \varrho_i(\beta) \nabla \varrho_i(\beta)'] \left\{ \mathbb{E}[\nabla^2 \varrho_i(\beta)] \right\}^{-1}) \text{ and}$$

$$\sqrt{N} \left(\hat{\gamma}^* - \hat{\gamma} \right) \stackrel{d}{\to} N \left(0, \mathbb{E}[\nabla^2 \tau_i(\gamma)]^{-1} \mathbb{E}\left[\Delta_i \Delta_i' \right] \mathbb{E}[\nabla^2 \tau_i(\gamma)]^{-1} \right)$$

conditional on the sample.

This result is proved in Appendix D. As a result, $\{(\hat{\beta}_{(b)}^*, \hat{\gamma}_{(b)}^*)\}_{b=1}^B$ can be used to compute the standard errors or confidence intervals of the estimators $\hat{\beta}$ and $\hat{\gamma}$.

⁶Jin, Ying, and Wei (2001) proposed an alternative resampling method for U-statistics by perturbing the criterion function repeatedly.

2.1.4 Testing η

Our identification requires $\gamma^{(1)} \neq 0$ and a scale normalization $\gamma^{(1)} = 1$. As a consequence, it is impossible to test for the presence of the interaction effect by simply looking at the significance of $\|\gamma\|$. This observation highlights the role of η -it determines the magnitude of the interaction effect between the two stand-alone alternatives. In this section, we introduce a criterion-based procedure to test whether η degenerates to zero. We formulate the null hypothesis as

$$\mathbb{H}_0: \eta > 0$$
 almost surely and $E(\eta) > 0$,

and the alternative hypothesis as

$$\mathbb{H}_1: \eta = 0$$
 almost surely.

Before presenting the theoretical result, we explain the idea underlying the test. We define $\hat{\mathcal{L}}_{N}^{K}\left(\hat{\gamma}\right)$ as

$$\hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) = \frac{1}{\sigma_{N}^{2} N\left(N-1\right)} \sum_{i \neq m} K_{\sigma_{N},\gamma}(V_{im}(\hat{\beta})) Y_{im(1,1)} \operatorname{sgn}\left(W_{im}'\hat{\gamma}\right).$$

Suppose we know the value of β . Define

$$\mathcal{L}_{N}^{K}\left(r\right) = \frac{1}{\sigma_{N}^{2}N\left(N-1\right)} \sum_{i \neq m} K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right) Y_{im(1,1)} \operatorname{sgn}\left(W'_{im}r\right).$$

Clearly, $\mathcal{L}_{N}^{K}(r)$ is a second-order U-statistic, and

$$\mathcal{L}_{N}^{K}(r) \stackrel{p}{\to} \bar{\mathcal{L}}(r) \equiv f_{V_{im}(\beta)}(0) \mathbb{E} \left[Y_{im(1,1)} \operatorname{sgn}(W'_{im}r) | V_{im}(\beta) = 0 \right].$$

If $\eta = 0$ almost surely,

$$\bar{\mathcal{L}}(r) = 0$$
 for any r

since $Y_{im(1,1)}$ is independent of W_{im} conditional on $V_{im}(\beta) = 0$ and $\mathbb{E}[Y_{im(1,1)}|V_{im}(\beta) = 0] = 0$. On the contrary, if $\eta > 0$ almost surely and $E(\eta) > 0$, then the signs of $W'_{im}\gamma$ and $P(Y_{i(1,1)}|V_i(\beta) = v, W_i) - P(Y_{m(1,1)}|V_m(\beta) = v, W_m)$ should be the same. As a result,

$$\bar{\mathcal{L}}(\gamma) > 0.$$

In light of this observation, testing \mathbb{H}_0 is equivalent to testing $\bar{\mathcal{L}}(\gamma) > 0$.

By some standard analysis for U-statistics, the limiting distribution of $\mathcal{L}_{N}^{K}(\gamma)$ is

$$\sqrt{N}\left(\mathcal{L}_{N}^{K}\left(\gamma\right)-\bar{\mathcal{L}}\left(\gamma\right)\right)\stackrel{d}{\to}N\left(0,\Delta_{\gamma}\right),$$

with

$$\Delta_{\gamma} = 4 \operatorname{Var} \left\{ f_{V_{im}(\beta)}(0) \mathbb{E} \left[Y_{im(1,1)} \operatorname{sgn} \left(W'_{im} \gamma \right) | Z_i, V_m \left(\beta \right) = V_i \left(\beta \right) \right] \right\}.$$

We can then test whether $\bar{\mathcal{L}}(\gamma) > 0$ using the above limiting distribution.

The analysis for our case is more complicated by the fact that we need to first estimate β and γ before running the test. In Theorem 2.4 below, we show that the plugged-in $\hat{\beta}$ and $\hat{\gamma}$ affect $\hat{\mathcal{L}}_N^K(\hat{\gamma})$ at the rates of $N^{-1}\sigma_N^{-2}$ and N^{-1} , respectively, around the true values (β, γ) . These rates are much faster than $N^{-1/2}$. Consequently, the asymptotic behavior of $\hat{\mathcal{L}}_N^K(\hat{\gamma})$ is the same as that of $\mathcal{L}_N^K(\gamma)$. The limiting distribution of $\hat{\mathcal{L}}_N^K(\hat{\gamma})$ under the null hypothesis is provided in the following theorem. The proof is included in Appendix B.

Theorem 2.4. Suppose Assumptions C1-C7 hold. Then, under \mathbb{H}_0 ,

$$\sqrt{N}(\hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) - \bar{\mathcal{L}}(\gamma)) \stackrel{d}{\to} N(0, \Delta).$$

To perform the test, we propose using a standard bootstrap procedure. Specifically, we independently draw $\{(Y_i^*, Z_i^*)\}_{i=1}^N$ with replacement from the sample $\{(Y_i, Z_i)\}_{i=1}^N$ for a total of B times. For each draw, we calculate

$$\hat{\mathcal{L}}_{N}^{K*}(\hat{\gamma}) = \frac{1}{\sigma_{N}^{2} N(N-1)} \sum_{i \neq m} K_{\sigma_{N}, \gamma}(V_{im}^{*}(\hat{\beta})) Y_{im(1,1)}^{*} \operatorname{sgn}(W_{im}^{*\prime} \hat{\gamma}).$$

Note that there is no need to re-compute $\hat{\beta}$ and $\hat{\gamma}$ using the bootstrap samples. We construct a 95% confidence interval for $\bar{\mathcal{L}}(\gamma)$ as a one-sided interval $[Q_{0.05}(\hat{\mathcal{L}}_N^{K*}(\hat{\gamma})), +\infty)$, where $Q_{0.05}(\cdot)$ denotes the 5% quantile of $\hat{\mathcal{L}}_N^{K*}(\hat{\gamma})$. By the same reason as we provided above, the plugged-in $\hat{\beta}$ and $\hat{\gamma}$ have no impact on the asymptotics of $\hat{\mathcal{L}}_N^{K*}(\hat{\gamma})$. The validity of the bootstrap for U-statistics has been established in the literature (see, e.g., Arcones and Gine (1992)). Since this analysis is standard, we omit it here for brevity.

2.2 Least Absolute Deviations Method

The main limitation of the MRC estimation proposed in Section 2.1 is its inability to estimate coefficients for regressors that are common across different alternatives, often referred to as common regressors. For example, individual-specific characteristics like gender may be present in multiple stand-alone utilities as well as the bundle utility. In such cases, the coefficients for these regressors are often alternative-specific. In this section, we propose an innovative least absolute deviations (LAD) estimator, offering the advantage of estimating coefficients for both common and alternative-specific regressors. The LAD estimator is also robust in that it does not impose any distributional assumptions on the error terms and allows for arbitrary correlations among them, aligning with the flexibility of the MRC estimator in these respects.

Consider the following model extended from the model specified in (2.1) and (2.2):

$$U_d = \sum_{j=1}^2 F_j(X_j'\beta + S'\rho_j, \epsilon_j) \cdot d_j + \eta \cdot F_b(W'\gamma + S'\rho_b) \cdot d_1 \cdot d_2,$$

$$Y_d = 1[U_d > U_{d'}, \forall d' \in \mathcal{D} \setminus d],$$
(2.10)

where $S \in \mathbb{R}^{k_3}$ is a vector of common regressors affecting the utilities of all stand-alone alternatives and the bundle. With a slight abuse of notation, we let $Z \equiv (X_1, X_2, W, S)$ in this section. Then, under Assumption C4 and $(\epsilon_1, \epsilon_2, \eta) \perp Z$, for two independent copies of (Y, Z), (Y_i, Z_i) and (Y_m, Z_m) , we can write the following inequalities from model (2.10) and equation (2.3) (with modified latent utilities specified in (2.10)):

$$\{X'_{im1}\beta + S'_{im}\rho_1 \ge 0, X'_{im2}\beta + S'_{im}\rho_2 \le 0, W'_{im}\gamma + S'_{im}\rho_b \le 0\} \Rightarrow \Delta p_{(1,0)}(Z_i, Z_m) \ge 0,$$
(2.11)

$$\{X'_{im1}\beta + S'_{im}\rho_1 \le 0, X'_{im2}\beta + S'_{im}\rho_2 \ge 0, W'_{im}\gamma + S'_{im}\rho_b \ge 0\} \Rightarrow \Delta p_{(1,0)}(Z_i, Z_m) \le 0,$$
(2.12)

where
$$S_{im} \equiv S_i - S_m$$
 and $\Delta p_{(1,0)}(Z_i, Z_m) \equiv P(Y_{i(1,0)} = 1 | Z_i) - P(Y_{m(1,0)} = 1 | Z_m)$.

We propose the following LAD criterion function to identify the unknown parameters $\theta \equiv (\beta, \gamma, \rho_1, \rho_2, \rho_b)$ in model (2.10):

$$Q_{(1,0)}(\vartheta) = \mathbb{E}\left[q_{(1,0)}(Z_i, Z_m, \vartheta)\right],\tag{2.13}$$

where $\vartheta = (b, r, \varrho_1, \varrho_2, \varrho_b)$ is an arbitrary vector in the parameter space Θ containing θ and

$$q_{(1,0)}(Z_i, Z_m, \vartheta) = [|I_{im(1,0)}^+(\vartheta) - 1[\Delta p_{(1,0)}(Z_i, Z_m) \ge 0]| + |I_{im(1,0)}^-(\vartheta) - 1[\Delta p_{(1,0)}(Z_i, Z_m) \le 0]|] \times [I_{im(1,0)}^+(\vartheta) + I_{im(1,0)}^-(\vartheta)]$$

with

$$I_{im(1,0)}^+(\vartheta) = 1[X_{im1}'b + S_{im}'\varrho_1 \ge 0, X_{im2}'b + S_{im}'\varrho_2 \le 0, W_{im}'r + S_{im}'\varrho_b \le 0]$$
 and (2.14)

$$I_{im(1,0)}^{-}(\vartheta) = 1[X'_{im1}b + S'_{im}\varrho_1 \le 0, X'_{im2}b + S'_{im}\varrho_2 \ge 0, W'_{im}r + S'_{im}\varrho_b \ge 0]$$
(2.15)

being indicator functions respectively for the events defined on the left hand sides of (2.11) and (2.12), but evaluated at ϑ .

The construction of the function $q_{(1,0)}(Z_i, Z_m, \vartheta)$ might seem complex at first glance, but it has an intuitive interpretation. Inequalities (2.11) and (2.12) imply that using the true parameter θ , we can flawlessly predict the occurrence of the event $\{\Delta p_{(1,0)}(Z_i, Z_m) \geq 0\}$ ($\{\Delta p_{(1,0)}(Z_i, Z_m) \leq 0\}$) when $I^+_{im(1,0)}(\theta) = 1$ ($I^-_{im(1,0)}(\theta) = 1$). However, for any $\vartheta \neq \theta$, such predictions may lead to errors, where $I^+_{im(1,0)}(\vartheta) = 1$ ($I^-_{im(1,0)}(\vartheta) = 1$) does not align with the observed data $\Delta p_{(1,0)}(Z_i, Z_m) \geq 0$ ($\Delta p_{(1,0)}(Z_i, Z_m) \leq 0$). Besides, when $I^+_{im(1,0)}(\vartheta) = I^-_{im(1,0)}(\vartheta) = 0$, no prediction can be made for the sign of $\Delta p_{(1,0)}(Z_i, Z_m)$. Then, treating $q_{(1,0)}(Z_i, Z_m, \vartheta)$ as a loss function, we can interpret it as the loss incurred from making incorrect predictions for the sign of $\Delta p_{(1,0)}(Z_i, Z_m)$ using ϑ :

- A wrong prediction incurs a loss of 2, i.e., $q_{(1,0)}(Z_i, Z_m, \vartheta) = 2$ when $\{I_{im(1,0)}^+(\vartheta) = 1, \Delta p_{(1,0)}(Z_i, Z_m) \leq 0\}$ or $\{I_{im(1,0)}^-(\vartheta) = 1, \Delta p_{(1,0)}(Z_i, Z_m) \geq 0\}$ occurs.
- A correct prediction or no prediction incurs 0 loss, i.e., $q_{(1,0)}(Z_i, Z_m, \vartheta) = 0$ when $\{I_{im(1,0)}^+(\vartheta) = 1, \Delta p_{(1,0)}(Z_i, Z_m) \geq 0\}$ or $\{I_{im(1,0)}^-(\vartheta) = 1, \Delta p_{(1,0)}(Z_i, Z_m) \leq 0\}$ or $\{I_{im(1,0)}^+(\vartheta) = I_{im(1,0)}^-(\vartheta) = 0\}$ occurs.

Therefore, $Q_{(1,0)}(\vartheta)$ represents the expected loss of making wrong predictions for the sign of $\Delta p_{(1,0)}(Z_i, Z_m)$ based on inequalities (2.11) and (2.12). It is evident that the true parameter θ minimizes $Q_{(1,0)}(\vartheta)$ since it never gives wrong predictions.

The following conditions are sufficient for establishing the identification of θ via (2.13), which are analogous to Assumptions C1–C4:

CL1 (i) $\{(Y_i, Z_i)\}_{i=1}^N$ are i.i.d. across i, (ii) $(\epsilon_1, \epsilon_2, \eta) \perp Z$, and (iii) the joint distribution of

 $(\epsilon_1, \epsilon_2, \eta)$ is absolutely continuous on $\mathbb{R}^2 \times \mathbb{R}_+$.

- CL2 For any pair of (i, m) and j = 1, 2, denote $X_{imj} = X_{ij} X_{mj}$, $W_{im} = W_i W_m$, and $S_{im} = S_i S_m$. Then, (i) $X_{im1}^{(1)}$ ($X_{im2}^{(1)}$) has a.e. positive Lebesgue density on \mathbb{R} conditional on $(\tilde{X}_{im1}, X_{im2}, W_{im}, S_{im})$ ($(\tilde{X}_{im2}, X_{im1}, W_{im}, S_{im})$), (ii) Elements in (X_{im1}, S_{im}) ((X_{im2}, S_{im})), conditional on (X_{im2}, W_{im}) ((X_{im1}, W_{im})), are linearly independent, (iii) $W_{im}^{(1)}$ has a.e. positive Lebesgue density on \mathbb{R} conditional on $(\tilde{W}_{im}, X_{im1}, X_{im2}, S_{im})$, and (iv) Elements in (W_{im}, S_{im}) , conditional on (X_{im1}, X_{im2}) , are linearly independent.
- **CL3** $\theta = (\beta, \gamma, \rho_1, \rho_2, \rho_b) \in \Theta \subset \mathbb{R}^{k_1 + k_2 + 3k_3}$, where $\Theta = \{\theta = (b, r, \varrho_1, \varrho_2, \varrho_b) | b^{(1)} = r^{(1)} = 1, \|\theta\| \le C\}$ for some constant C > 0.
- **CL4** $F_1(\cdot,\cdot)$, $F_2(\cdot,\cdot)$, and $F_b(\cdot)$ are respectively $\mathbb{R}^2 \to \mathbb{R}$, $\mathbb{R}^2 \to \mathbb{R}$, and $\mathbb{R} \to \mathbb{R}$ functions strictly increasing in each of their arguments. $F_b(0) = 0$.

The following theorem, proved in Appendix B, summarizes our identification result.

Theorem 2.5. Under Assumptions CL1-CL4, $Q_{(1,0)}(\theta) < Q_{(1,0)}(\vartheta)$ for all $\vartheta \in \Theta \setminus \{\theta\}$.

The true parameter θ can also be identified using an alternative population criterion function, replacing the $1[\Delta p_{(1,0)}(Z_i, Z_m) \geq 0]$ and $1[\Delta p_{(1,0)}(Z_i, Z_m) \leq 0]$ in (2.13) with $\Delta p_{(1,0)}(Z_i, Z_m)$ and $-\Delta p_{(1,0)}(Z_i, Z_m)$, respectively:

$$Q_{(1,0)}^D(\vartheta) = \mathbb{E}\left[q_{(1,0)}^D(Z_i, Z_m, \vartheta)\right],\tag{2.16}$$

where

$$q_{(1,0)}^{D}(Z_{i}, Z_{m}, \vartheta) = [|I_{im(1,0)}^{+}(\vartheta) - \Delta p_{(1,0)}(Z_{i}, Z_{m})| + |I_{im(1,0)}^{-}(\vartheta) + \Delta p_{(1,0)}(Z_{i}, Z_{m})|] \times [I_{im(1,0)}^{+}(\vartheta) + I_{im(1,0)}^{-}(\vartheta)] + [1 - (I_{im(1,0)}^{+}(\vartheta) + I_{im(1,0)}^{-}(\vartheta))].$$

In fact, minimizing criterion functions (2.16) and (2.13) is equivalent since $|\Delta p_{(1,0)}(Z_i, Z_m)| < 1$ and thus

$$\left\{ q_{(1,0)}(Z_i, Z_m, \vartheta) = 0 \right\} \Leftrightarrow \left\{ q_{(1,0)}^D(Z_i, Z_m, \vartheta) = 1 \right\} \text{ and }$$

$$\left\{ q_{(1,0)}(Z_i, Z_m, \vartheta) = 2 \right\} \Leftrightarrow \left\{ q_{(1,0)}^D(Z_i, Z_m, \vartheta) = 1 + 2|\Delta p_{(1,0)}(Z_i, Z_m)| \right\}.$$

Though criterion function (2.13) is more intuitive, in practice we recommend to construct estimation procedure based on (2.16) rather than (2.13). This is because $\Delta p_{(1,0)}(Z_i, Z_m)$

is unobservable, and hence a feasible estimation procedure will need to plug in its (uniformly) consistent estimate $\Delta \hat{p}_{(1,0)}(Z_i, Z_m)$. However, in finite samples $\Delta p_{(1,0)}(Z_i, Z_m)$ and $\Delta \hat{p}_{(1,0)}(Z_i, Z_m)$ can have different signs with positive probability, leading to finite-sample bias in the estimation. Using (2.16) can effectively reduce the impact of such problem in comparison to (2.13). To illustrate this, consider the situation where $\Delta p_{(1,0)}(Z_i, Z_m) > 0$ but $\Delta \hat{p}_{(1,0)}(Z_i, Z_m) < 0$. A simple calculation shows that this will generate an error of size 2 to $q_{(1,0)}(Z_i, Z_m, \vartheta)$ in (2.13) and an error of size $2|\Delta \hat{p}_{(1,0)}(Z_i, Z_m)|$ to $q_{(1,0)}^D(Z_i, Z_m, \vartheta)$ in (2.16). Since $\Delta \hat{p}_{(1,0)}(Z_i, Z_m) \stackrel{p}{\rightarrow} \Delta p_{(1,0)}(Z_i, Z_m)$, as the sample size increases, we expect not only $P(\Delta p_{(1,0)}(Z_i, Z_m) > 0, \Delta \hat{p}_{(1,0)}(Z_i, Z_m) < 0) \rightarrow 0$ but also $\Delta \hat{p}_{(1,0)}(Z_i, Z_m) \approx 0$ when its sign is different from that of $\Delta p_{(1,0)}(Z_i, Z_m)$. Consequently, the bias caused by the inconsistency in signs between $\Delta p_{(1,0)}(Z_i, Z_m)$ and $\Delta \hat{p}_{(1,0)}(Z_i, Z_m)$ decreases much faster when using (2.16) compared to (2.13). From this perspective, criterion function (2.16) can be considered a debiased version of (2.13).

Suppose there is a uniformly consistent estimator $\Delta \hat{p}_{(1,0)}(Z_i, Z_m)$ for $\Delta p_{(1,0)}(Z_i, Z_m)$. We propose the following LAD estimator $\hat{\theta}$ for θ :

$$\hat{\theta} = \arg\min_{\vartheta \in \Theta} \sum_{i=1}^{N-1} \sum_{m>i} \hat{q}_{im(1,0)}^{D}(\vartheta), \tag{2.17}$$

where

$$\hat{q}_{im(1,0)}^{D}(\vartheta) = [|I_{im(1,0)}^{+}(\vartheta) - \Delta \hat{p}_{(1,0)}(Z_i, Z_m)| + |I_{im(1,0)}^{-}(\vartheta) + \Delta \hat{p}_{(1,0)}(Z_i, Z_m)|] \times [I_{im(1,0)}^{+}(\vartheta) + I_{im(1,0)}^{-}(\vartheta)] + [1 - (I_{im(1,0)}^{+}(\vartheta) + I_{im(1,0)}^{-}(\vartheta))].$$

We need the following assumption on top of Assumptions CL1–CL4 to establish the consistency of the estimator $\hat{\theta}$ proposed in (2.17):

CL5
$$\sup_{(z_i,z_m)\in\mathcal{Z}^2} |\Delta \hat{p}_{(1,0)}(z_i,z_m) - \Delta p_{(1,0)}(z_i,z_m)| \stackrel{p}{\to} 0 \text{ as } n \to \infty, \text{ where } \mathcal{Z} \text{ is the support of } Z.$$

Assumption CL5, when combined with Assumption CL2, essentially requires a uniformly consistent estimator for $\Delta p_{(1,0)}(Z_i, Z_m)$ on a non-compact support.⁷ To our knowledge, the

⁷In empirical industrial organization and marketing research, choice models are often studied using aggregate data. Researchers can observe the aggregated choice probabilities (market shares) along with covariates of market-level. In this scenario, Assumption CL5 can be satisfied under mild conditions. See Section 6 of Shi et al. (2018) for a more detailed discussion.

nearest neighbor estimator (Devroye (1978)) and the kernel regression estimator (Hansen (2008)) can be shown to satisfy such uniform consistency condition under certain restrictions on the tail probability bounds for Z. For example, Hansen (2008) proved that when $\Delta \hat{p}_{(1,0)}(Z_i,Z_m)$ is obtained using kernel regression, Assumption CL5 is satisfied if elements in Z have distribution tails thinner than polynomial. Some recent advances in machine learning can also be used to estimate $\Delta p_{(1,0)}(Z_i,Z_m)$, especially for high-dimensional cases. However, similar results for uniform convergence with non-compact support are lacking in the literature. Kohler and Langer (2021) demonstrated that, with certain restrictions on the network architecture and smoothness assumptions on the regression function, the multi-layer neural network regression can circumvent the curse of dimensionality. They derive a uniform convergence rate of the multi-layer neural network estimator, assuming a compact support for covariates. In our simulation studies presented in Section 4, we use the neural network method to estimate the choice probabilities in a panel data bundle choice model (see Section 3.2). Our panel data LAD estimator using such plug-in estimates performs satisfactorily despite the lack of theoretical guarantees.

The following theorem, proved in Appendix B, shows the consistency of $\hat{\theta}$.

Theorem 2.6. Suppose Assumptions CL1–CL5 hold. We have $\hat{\theta} \stackrel{p}{\rightarrow} \theta$.

Using the plug-in nonparametric estimates $\Delta \hat{p}_{(1,0)}(Z_i, Z_m)$ complicates the derivation of the convergence rate and asymptotic distribution of the estimator, especially for its panel data counterpart which will be discussed in Section 3.2. We typically need $\Delta \hat{p}_{(1,0)}(Z_i, Z_m)$ converges to $\Delta p_{(1,0)}(Z_i, Z_m)$ sufficiently fast to prevent its prediction error from dominating the rate of the estimator. More importantly, we conjecture that applying certain transformations or smoothing techniques (e.g., Horowitz (1992)) to criterion function (2.13) (or (2.16)) may yield an estimator with a better convergence rate and facilitate the derivation of its asymptotic distribution. However, exploring these issues is a non-trivial task, and due to space constraints, we must defer a comprehensive investigation to future research. We conclude this section with a final remark.

Remark 2.2. In addition to (2.11) and (2.12), we can also construct population and sample

criterion functions based on the following inequalities:

$$\begin{aligned}
&\{X'_{im1}\beta + S'_{im}\rho_{1} \leq 0, X'_{im2}\beta + S'_{im}\rho_{2} \geq 0, W'_{im}\gamma + S'_{im}\rho_{b} \leq 0\} \Rightarrow \Delta p_{(0,1)}(Z_{i}, Z_{m}) \geq 0, \\
&(2.18)
\end{aligned}$$

$$\begin{aligned}
&\{X'_{im1}\beta + S'_{im}\rho_{1} \geq 0, X'_{im2}\beta + S'_{im}\rho_{2} \leq 0, W'_{im}\gamma + S'_{im}\rho_{b} \geq 0\} \Rightarrow \Delta p_{(0,1)}(Z_{i}, Z_{m}) \leq 0, \\
&(2.19)
\end{aligned}$$

$$\begin{aligned}
&\{X'_{im1}\beta + S'_{im}\rho_{1} \geq 0, X'_{im2}\beta + S'_{im}\rho_{2} \geq 0, W'_{im}\gamma + S'_{im}\rho_{b} \geq 0\} \Rightarrow \Delta p_{(0,1)}(Z_{i}, Z_{m}) \leq 0, \\
&(2.20)
\end{aligned}$$

$$\begin{aligned}
&\{X'_{im1}\beta + S'_{im}\rho_{1} \geq 0, X'_{im2}\beta + S'_{im}\rho_{2} \leq 0, W'_{im}\gamma + S'_{im}\rho_{b} \leq 0\} \Rightarrow \Delta p_{(1,1)}(Z_{i}, Z_{m}) \leq 0, \\
&(2.21)
\end{aligned}$$

$$\begin{aligned}
&\{X'_{im1}\beta + S'_{im}\rho_{1} \leq 0, X'_{im2}\beta + S'_{im}\rho_{2} \leq 0, W'_{im}\gamma + S'_{im}\rho_{b} \leq 0\} \Rightarrow \Delta p_{(0,0)}(Z_{i}, Z_{m}) \geq 0, \\
&(2.22)
\end{aligned}$$

$$\begin{aligned}
&\{X'_{im1}\beta + S'_{im}\rho_{1} \leq 0, X'_{im2}\beta + S'_{im}\rho_{2} \leq 0, W'_{im}\gamma + S'_{im}\rho_{b} \leq 0\} \Rightarrow \Delta p_{(0,0)}(Z_{i}, Z_{m}) \geq 0, \\
&(2.22)
\end{aligned}$$

where $\Delta p_d(Z_i, Z_m)$ for $d \in \{(0, 1), (1, 1), (0, 0)\}$ are similarly defined as $\Delta p_{(1,0)}(Z_i, Z_m)$ in (2.11)–(2.12). Note that the following two events are not informative in the sense that they are not sufficient to predict the sign of any $\Delta p_d(Z_i, Z_m)$:

$$\{X'_{im1}\beta + S'_{im}\rho_1 \le 0, X'_{im2}\beta + S'_{im}\rho_2 \le 0, W'_{im}\gamma + S'_{im}\rho_b \ge 0\},\$$
$$\{X'_{im1}\beta + S'_{im}\rho_1 \ge 0, X'_{im2}\beta + S'_{im}\rho_2 \ge 0, W'_{im}\gamma + S'_{im}\rho_b \le 0\}.$$

For any $d \in \{(0,1), (1,1), (0,0)\}$ corresponding to (2.18)–(2.19), (2.20)–(2.21), and (2.22)–(2.23), respectively, $(I_{imd}^+(\vartheta), I_{imd}^-(\vartheta))$ can be defined analogously to $(I_{im(1,0)}^+(\vartheta), I_{im(1,0)}^-(\vartheta))$ in (2.14)–(2.15). Then, any single criterion $Q_d(\vartheta) = \mathbb{E}[q_d(Z_i, Z_m, \vartheta)]$ $(Q_d^D(\vartheta)) = \mathbb{E}[q_d^D(Z_i, Z_m, \vartheta)]$ or their convex combination can be used to identify θ . For estimation purposes, we recommend to use a sample criterion constructed by combining the sample analogues of all available population criteria (e.g., summing them up), to achieve higher finite sample efficiency. For example, in the simulation studies presented in Section 4, we compute

$$\hat{\theta} = \arg\min_{\vartheta \in \Theta} \sum_{i=1}^{N-1} \sum_{m > i} \sum_{d \in \mathcal{D}} \hat{q}_{imd}^{D}(\vartheta),$$

where $\mathcal{D} = \{(0,0), (1,0), (0,1), (1,1)\}$ and $\hat{q}_{imd}^D(\vartheta)$ is similarly defined as in (2.17).

3 Panel Data Model

The structure of this section is similar to that of Section 2. Section 3.1 introduces a localized MS procedure, in which we establish the identification and the limiting distribution of the estimator (Section 3.1.1), show the validity of the numerical bootstrap inference procedure (Section 3.1.2), and propose a test for the interaction effects (Section 3.1.3). Section 3.2 develops a LAD estimation analogous to Section 2.2.

3.1 Maximum Score Estimation

3.1.1 Localized MS Estimator

The increasing availability of panel data sets provides new opportunities for the econometrician to control for unobserved heterogeneity across agents. This helps with the relaxing of the strict exogeneity restriction placed in the cross-sectional model. In this section we apply the matching-based identification strategy presented in Section 2.1 to a panel data bundle choice model. The latent utilities and observed choices are expressed as follows:

$$U_{dt} = \sum_{j=1}^{2} F_j(X'_{jt}\beta, \alpha_j, \epsilon_{jt}) \cdot d_j + \eta_t \cdot F_b(W'_t\gamma + \alpha_b) \cdot d_1 \cdot d_2, \tag{3.1}$$

and

$$Y_{dt} = 1[U_{dt} > U_{d't}, \forall d' \in \mathcal{D} \setminus d], \tag{3.2}$$

where we use t=1,...,T to denote time periods and suppress the agent subscript i to simplify notations. Again, we assume that functions $F_j(\cdot,\cdot,\cdot)$'s and $F_b(\cdot,\cdot)$ are strictly increasing in their arguments. The specification considered here is a natural extension of the cross-sectional model (2.1)–(2.2). Similar extensions are common in various discrete choice models. See, e.g., Pakes and Porter (2021), Shi et al. (2018), and Gao and Li (2020). Note that the random utility specified in (3.1) includes a set of unobserved (to the econometrician) agent-specific fixed effects $\alpha \equiv (\alpha_1, \alpha_2, \alpha_b)$ associated with the two stand-alone alternatives and the bundle, respectively. Denote $Z_t = (X_{1t}, X_{2t}, W_t)$ and $\xi_t = (\epsilon_{1t}, \epsilon_{2t}, \eta_t)$. In line with the literature on fixed effects methods, we place no restrictions on the distribution of α conditional on $Z^T \equiv (Z_1, ..., Z_T)$ and $\xi^T \equiv (\xi_1, ..., \xi_T)$.

Here we consider the identification and estimation of model (3.1)–(3.2) with $T < \infty$ and

 $N \to \infty$ (i.e., short panel). For any $T \ge 2$, our identification strategy relies on a similar conditional homogeneity assumption as those adopted by Manski (1987), Pakes and Porter (2021), Shi et al. (2018), and Gao and Li (2020) for binary and multinomial choice models. Specifically, we assume $\xi_s \stackrel{d}{=} \xi_t | (\alpha, Z_s, Z_t)$ for any two time periods s and t.

This restriction is much weaker than the strong exogeneity condition needed for the cross-sectional model. However, thanks to the panel data structure, it suffices to establish the following moment inequalities in the presence of the fixed effects α : for all d with $d_1 = 1$ and any fixed (x_2, w, c) ,

$$P(Y_{(1,d_2)t} = 1 | X_{1t} = x_1, X_{2t} = x_2, W_t = w, \alpha = c)$$

$$x_1'\beta \ge \tilde{x}_1'\beta \Leftrightarrow \ge$$

$$P(Y_{(1,d_2)s} = 1 | X_{1s} = \tilde{x}_1, X_{2s} = x_2, W_s = w, \alpha = c).$$
(3.3)

Similarly, for all d with $d_1 = 0$ and any fixed (x_2, w, c) , $x'_1\beta \geq \tilde{x}'_1\beta$ is the if-and-only-if condition to the second inequality in (3.3) but with a " \leq " instead.

For d = (1, 1) and any fixed (x_1, x_2, c) , we have

$$P(Y_{(1,1)t} = 1 | X_{1t} = x_1, X_{2t} = x_2, W_t = w, \alpha = c)$$

$$w'\gamma \ge \tilde{w}'\gamma \Leftrightarrow \ge$$

$$P(Y_{(1,1)s} = 1 | X_{1s} = x_1, X_{2s} = x_2, W_s = \tilde{w}, \alpha = c).$$
(3.4)

Similarly, for all $d \neq (1, 1)$ and any fixed (x_1, x_2, c) , $w'\gamma \geq \tilde{w}'\gamma$ is the if-and-only-if condition to the second inequality in (3.4) but with a " \leq " instead.

The moment inequalities (3.3)– $(3.4)^8$ are derived from the monotonicity conditions of agents' choices, analogous to (2.4)–(2.5) for the cross-sectional model. The main difference is that, in the presence of the fixed effects, we match and make our comparisons within agents over time, as opposed to pairs of agents. These moment inequalities are the foundation of our identification result for model (3.1)–(3.2).

We impose the following conditions for identification. For notational convenience, we let $X_{jts} \equiv X_{jt} - X_{js}$ and $W_{ts} \equiv W_t - W_s$ for all j = 1, 2 and (s, t).

⁸Once β is identified through (3.3), γ can be alternatively identified via matching indexes $(X'_{1t}\beta, X'_{2t}\beta)$; that is, for d=(1,1) and any fixed $(v_1,v_2), w'\gamma \geq \tilde{w}'\gamma \Leftrightarrow P(Y_{(1,1)t}=1|X'_{1t}\beta=v_1, X'_{2t}\beta=v_2, W_t=w, \alpha=c) \geq P(Y_{(1,1)s}=1|X'_{1s}\beta=v_1, X'_{2s}\beta=v_2, W_s=\tilde{w}, \alpha=c))$. However, due to some technical issues (see Appendix E), we do not adopt these moment inequalities to construct the MS estimator of γ .

- **P1** (i) $\{(Y_i^T, Z_i^T)\}_{i=1}^N$ are i.i.d, where $Y_i^T \equiv \{(Y_{i(0,0)t}, Y_{i(1,0)t}, Y_{i(0,1)t}, Y_{i(1,1)t})\}_{t=1}^T$, (ii) for almost all (α, Z^T) , $\xi_t \stackrel{d}{=} \xi_s | (\alpha, Z_t, Z_s)$, and (iii) the distribution of ξ_t conditional on α is absolutely continuous w.r.t. the Lebesgue measure on $\mathbb{R}^2 \times \mathbb{R}_+$ for all t.
- **P2** $(\beta, \gamma) \in \mathcal{B} \times \mathcal{R}$, where $\mathcal{B} = \{b \in \mathbb{R}^{k_1} | ||b|| = 1, b^{(1)} \neq 0\}$ and $\mathcal{R} = \{r \in \mathbb{R}^{k_2} | ||r|| = 1, r^{(1)} \neq 0\}$.
- P3 (i) $X_{1ts}^{(1)}$ ($X_{2ts}^{(1)}$) has a.e. positive Lebesgue density on \mathbb{R} conditional on \tilde{X}_{1ts} (\tilde{X}_{2ts}) and conditional on (X_{2ts}, W_{ts}) ((X_{1ts}, W_{ts})) in a neighborhood of (X_{2ts}, W_{ts}) ((X_{1ts}, W_{ts})) near zero, (ii) the support of X_{1ts} (X_{2ts}), conditional on (X_{2ts}, W_{ts}) ((X_{1ts}, W_{ts})) in a neighborhood of (X_{2ts}, W_{ts}) ((X_{1ts}, W_{ts})) near zero, is not contained in any proper linear subspace of \mathbb{R}^{k_1} , (iii) $W_{ts}^{(1)}$ has a.e. positive Lebesgue density on \mathbb{R} conditional on \tilde{W}_{ts} and conditional on (X_{1ts}, X_{2ts}) in a neighborhood of (X_{1ts}, X_{2ts}) near zero, and (iv) the support of W_{ts} , conditional on (X_{1ts}, X_{2ts}) in a neighborhood of (X_{1ts}, X_{2ts}) near zero, is not contained in any proper linear subspace of \mathbb{R}^{k_2} .
- **P4** $F_{j}(\cdot,\cdot,\cdot)$, j=1,2, and $F_{b}(\cdot)$ are strictly increasing in their arguments. $F_{b}(0)=0$.

Assumptions P1–P3 are analogous to the identification conditions used by Manski (1987). Note that Assumption P1 allows for arbitrary correlation between the fixed effects and the observed covariates, provided that the correlation is time stationary. This assumption substantially relaxes the strict exogeneity condition in the cross-sectional case, but the price to pay is a slower convergence rate, as shown in Theorem 3.2 below. A technical explanation for the slower convergence rate can be found in Appendix E.

We summarize our identification results in the following theorem. The proof is omitted for brevity, as it closely resembles that of the cross-sectional model.

Theorem 3.1. Suppose Assumptions P1-P4 hold. Then β and γ are identified.

The monotonic relations (3.3)–(3.4) motivate the following two-step localized MS estimation procedure. Given a random sample of N agents i = 1, ..., N, we obtain the estimator

⁹We adopt the same scale normalization as in Kim and Pollard (1990) and Seo and Otsu (2018). This simplifies the asymptotic analysis of our MS estimators, enabling us to use the results established in these works directly. It is easy to verify that the normalization used here is equivalent to that in Assumption C3.

 $\hat{\beta}$ of β through maximizing the criterion function

$$\mathcal{L}_{N,\beta}^{P,K}(b) = \sum_{i=1}^{N} \sum_{t>s} \sum_{d \in \mathcal{D}} \{ \mathcal{K}_{h_N}(X_{i2ts}, W_{its}) (Y_{ids} - Y_{idt}) \operatorname{sgn}(X'_{i1ts}b) \cdot (-1)^{d_1} + \mathcal{K}_{h_N}(X_{i1ts}, W_{its}) (Y_{ids} - Y_{idt}) \operatorname{sgn}(X'_{i2ts}b) \cdot (-1)^{d_2} \},$$
(3.5)

where $\mathcal{K}_{h_N}(X_{ijts}, W_{its}) \equiv \prod_{l=1}^{k_1} h_N^{-1} K\left(X_{ijts,l}/h_N\right) \prod_{l=1}^{k_2} h_N^{-1} K\left(W_{its,l}/h_N\right)$ for $j=1,2, K(\cdot)$ is a kernel density function, and h_N is a bandwidth sequence that converges to 0 as $N \to \infty$. Similarly, we compute the estimator $\hat{\gamma}$ of γ by maximizing the criterion function

$$\mathcal{L}_{N,\gamma}^{P,K}(r) = \sum_{i=1}^{N} \sum_{t>s} \mathcal{K}_{\sigma_N}(X_{i1ts}, X_{i2ts})(Y_{i(1,1)t} - Y_{i(1,1)s}) \operatorname{sgn}(W'_{its}r), \tag{3.6}$$

where $\mathcal{K}_{\sigma_N}(X_{i1ts}, X_{i2ts}) \equiv \prod_{l=1}^{k_1} \sigma_N^{-1} K\left(X_{i1ts,l}/\sigma_N\right) \prod_{l=1}^{k_1} \sigma_N^{-1} K\left(X_{i2ts,l}/\sigma_N\right)$ for j=1,2, and σ_N is a bandwidth sequence that converges to 0 as $N \to \infty$. Unlike the cross-sectional case, here we choose not to estimate γ by matching $X'_{jt}\hat{\beta}$ and $X'_{js}\hat{\beta}$, j=1,2. The reason is explained in Appendix E. Note that criterion functions (3.5) and (3.6) take value 0 for observations whose choice is time invariant; that is, our approach uses only data on "switchers" in a way similar to the estimator in Manski (1987).

To establish the asymptotic properties of $\hat{\beta}$ and $\hat{\gamma}$, we need the following conditions in addition to Assumptions P1–P4:

- **P5** Let $f_{X_{jt},W_t}(\cdot)$ denote the density of the random vector (X_{jt},W_t) for all j=1,2, and $f_{X_{1ts},X_{2ts}}(\cdot)$ denote the density of the random vector (X_{1ts},X_{2ts}) . $f_{X_{jt},W_t}(\cdot)$'s and $f_{X_{1ts},X_{2ts}}(\cdot)$ are absolutely continuous, bounded from above on their supports, strictly positive in a neighborhood of zero, and twice continuous differentiable a.e. with bounded derivatives.
- **P6** For all $d \in \mathcal{D}$, $\mathbb{E}[Y_{dts}\operatorname{sgn}(X'_{1ts}b)|X_{2ts},W_{ts}]$ and $\mathbb{E}[Y_{dts}\operatorname{sgn}(X'_{2ts}b)|X_{1ts},W_{ts}]$ are twice continuously differentiable w.r.t. b a.e. with bounded derivatives, and $\mathbb{E}[Y_{dts}\operatorname{sgn}(W'_{ts}r)|X_{1ts},X_{2ts}]$ is twice continuously differentiable w.r.t. r a.e. with bounded derivatives.
- **P7** $K(\cdot)$ is a function of bounded variation and has a compact support. It satisfies: (i) $K(v) \geq 0$ and $\sup_v |K(v)| < \infty$, (ii) $\int K(v) dv = 1$, (iii) $\int vK(v) dv = 0$, (iv) $\int v^2 K(v) dv < \infty$ and (v) $K(\cdot)$ is twice continuously differentiable with bounded first and second derivatives.

P8 $(\hat{\beta}, \hat{\gamma})$ satisfies

$$N^{-1}\mathcal{L}_{N,\beta}^{P,K}(\hat{\beta}) \ge \max_{b \in \mathcal{B}} N^{-1}\mathcal{L}_{N,\beta}^{P,K}(b) - o_p((Nh_N^{k_1 + k_2})^{-2/3})$$

and

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma}) \ge \max_{r \in \mathcal{R}} N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(r) - o_p((N\sigma_N^{2k_1})^{-2/3}).$$

P9 h_N and σ_N are sequences of positive numbers such that as $N \to \infty$: (i) $h_N \to 0$ and $\sigma_N \to 0$, (ii) $Nh_N^{k_1+k_2} \to \infty$ and $N\sigma_N^{2k_1} \to \infty$, and (iii) $(Nh_N^{k_1+k_2})^{2/3}h_N^2 \to 0$ and $(N\sigma_N^{2k_1})^{2/3}\sigma_N^2 \to 0$.

The boundedness and smoothness restrictions placed on Assumptions P5–P7 are regularity conditions needed for proving the uniform convergence of the criterion functions to their population analogues. Assumption P8 is a standard technical condition. Assumption P9 is also standard, where P9(iii) is included to ensure that the bias term from the kernel estimation is asymptotically negligible. Note that Assumption P6 implicitly assumes that the second moments of X_{jts} 's and W_{ts} exist.

For the asymptotic distribution, we focus on the case where all regressors are continuous, and introduce the following notations to ease the exposition. Let

$$\phi_{Ni}(b) \equiv \sum_{t>s} \sum_{d \in \mathcal{D}} \{ \mathcal{K}_{h_N}(X_{i2ts}, W_{its}) Y_{idst} (-1)^{d_1} \left(1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0] \right) + \mathcal{K}_{h_N}(X_{i1ts}, W_{its}) Y_{idst} (-1)^{d_2} \left(1[X'_{i2st}b > 0] - 1[X'_{i2st}\beta > 0] \right) \}$$

and

$$\varphi_{Ni}(r) \equiv \sum_{t>s} \mathcal{K}_{\sigma_N}(X_{i1ts}, X_{i2ts}) Y_{i(1,1)ts}(1[W'_{its}r > 0] - 1[W'_{its}\gamma > 0]).$$

Note that $\hat{\beta}$ and $\hat{\gamma}$ can be equivalently obtained from

$$\hat{\beta} = \arg\max_{b \in \mathcal{B}} N^{-1} \sum_{i=1}^{N} \phi_{Ni}(b) \text{ and } \hat{\gamma} = \arg\max_{r \in \mathcal{R}} N^{-1} \sum_{i=1}^{N} \varphi_{Ni}(r),$$

because $\operatorname{sgn}(\cdot) = 2 \times 1[\cdot] - 1$ and adding terms not related to b or r has no effects on the optimization.

Theorem 3.2. Suppose Assumptions P1–P9 hold. Then,

$$(Nh_N^{k_1+k_2})^{1/3}(\hat{\beta}-\beta) \stackrel{d}{\to} \arg\max_{\rho \in \mathbb{R}^{k_1}} \mathcal{Z}_1(\rho),$$

where \mathcal{Z}_1 is a Gaussian process taking values in $\ell^{\infty}\left(\mathbb{R}^{k_1}\right)$, with $\mathbb{E}\left(\mathcal{Z}_1\left(\rho\right)\right) = \frac{1}{2}\rho'\mathbb{V}\rho$, covariance kernel $\mathbb{H}_1\left(\rho_1,\rho_2\right)$, and $\rho,\rho_1,\rho_2\in\mathbb{R}^{k_1}$, and

$$(N\sigma_N^{2k_1})^{1/3} (\hat{\gamma} - \gamma) \stackrel{d}{\to} \arg\max_{\delta \in \mathbb{R}^{k_2}} \mathcal{Z}_2(\delta),$$

where \mathcal{Z}_2 is a Gaussian process taking values in $\ell^{\infty}\left(\mathbb{R}^{k_2}\right)$, with $\mathbb{E}\left(\mathcal{Z}_2\left(\delta\right)\right) = \frac{1}{2}\delta'\mathbb{W}\delta$, covariance kernel $\mathbb{H}_2\left(\delta_1,\delta_2\right)$, and $\delta,\delta_1,\delta_2\in\mathbb{R}^{k_2}$. \mathbb{V} , \mathbb{W} , \mathbb{H}_1 , and \mathbb{H}_2 are defined, respectively, by equations (C.1), (C.2), (C.3), and (C.4) in Appendix C.

The convergence rates of $\hat{\beta}$ and $\hat{\gamma}$ decline as the number of alternatives, J, increases due to the necessity of matching more covariates. We illustrate this in the model with J=3 in Section E.3.3. In line with existing results on the MS estimators (e.g., Kim and Pollard (1990) and Seo and Otsu (2018)), the limiting distributions of $\hat{\beta}$ and $\hat{\gamma}$ are non-Gaussian, and their rates of convergence are slower than $N^{-1/3}$. We refer interested readers to Appendix E for further insights into the convergence rates of both cross-sectional and panel data estimators. Inference using the limiting distribution derived in Theorem 3.2 directly is rather difficult. Therefore, we recommend a bootstrap-based procedure for the inference in the next section. One alternative is to adopt a smoothed MS approach (e.g., Horowitz (1992)), which may yield a faster and asymptotically normal estimator. We leave this topic for future research.

3.1.2 Inference

Abrevaya and Huang (2005) proved the inconsistency of the classic bootstrap for the ordinary MS estimator. Our panel data estimators are indeed of the MS type and we thus expect that the classic bootstrap does not work for them either. Valid inference can be made by the m-out-of-n bootstrap, according to Lee and Pun (2006). Recently, Hong and Li (2020) proposed the numerical bootstrap, and showed the superior performance of this procedure over the m-out-of-n bootstrap. For our panel data estimators, another advantage of the numerical bootstrap is that we do not need to choose another set of bandwidths (h_N and σ_N) for estimation using the bootstrap series as with the m-out-of-n bootstrap. Based on these considerations, we propose to conduct the inference using the numerical bootstrap.

The numerical bootstrap estimators $\hat{\beta}^*$ and $\hat{\gamma}^*$ are obtained as follows. First, draw $\{(Y_i^{T*}, Z_i^{T*})\}_{i=1}^N$ independently from the collection of the sample values $\{(Y_i^T, Z_i^T)\}_{i=1}^N$ with

replacement. Then, obtain $\hat{\beta}^*$ from

$$\hat{\beta}^* = \arg\max_{b \in \mathcal{B}} N^{-1} \sum_{i=1}^{N} \phi_{Ni}(b) + (N\varepsilon_{N1})^{1/2} \cdot [N^{-1} \sum_{i=1}^{N} \phi_{Ni}^*(b) - N^{-1} \sum_{i=1}^{N} \phi_{Ni}(b)], \quad (3.7)$$

where ε_{N1} is a tuning parameter to be discussed later, and $\phi_{Ni}^*(b)$ is the same as $\phi_{Ni}(b)$ except it uses the sampling series $\{(Y_i^{T*}, Z_i^{T*})\}_{i=1}^N$ as inputs. Similarly, compute $\hat{\gamma}^*$ from

$$\hat{\gamma}^* = \arg\max_{r \in \mathcal{R}} N^{-1} \sum_{i=1}^{N} \varphi_{Ni}(r) + (N\varepsilon_{N2})^{1/2} \cdot [N^{-1} \sum_{i=1}^{N} \varphi_{Ni}^*(r) - N^{-1} \sum_{i=1}^{N} \varphi_{Ni}(r)], \quad (3.8)$$

where ε_{N2} is a tuning parameter, and $\varphi_{Ni}^{*}(r)$ is similarly defined using bootstrap series.

When $\varepsilon_{N1}^{-1} = \varepsilon_{N2}^{-1} = N$, the numerical bootstrap reduces to the classic nonparametric bootstrap. The numerical bootstrap excludes this case and requires $N\varepsilon_{N1} \to \infty$ and $N\varepsilon_{N2} \to \infty$ as $N \to \infty$. We note that ε_{N1}^{-1} and ε_{N2}^{-1} play a similar role to the m in the m-out-of-n bootstrap (use only m observations for the estimation). The following conditions for the ε_N 's are required for the validity of the numerical bootstrap:

P10 ε_{N1} and ε_{N2} are sequences of positive numbers such that as $N \to \infty$: (i) $\varepsilon_{N1} \to 0$ and $\varepsilon_{N2} \to 0$, (ii) $N\varepsilon_{N1} \to \infty$ and $N\varepsilon_{N2} \to \infty$, (iii) $\varepsilon_{N1}^{-1}h_N^{k_1+k_2} \to \infty$ and $\varepsilon_{N2}^{-1}\sigma_N^{2k_1} \to \infty$, and (iv) $(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3}h_N^2 \to 0$ and $(\varepsilon_{N2}^{-1}\sigma_N^{2k_1})^{2/3}\sigma_N^2 \to 0$.

The following theorem shows the validity of the numerical bootstrap, whose proof is deferred to Appendix C. We note that our estimators do not directly satisfy all conditions required by Hong and Li (2020), for example, condition (vi) in their Theorem 4.1 is not the case here.

Theorem 3.3. Suppose Assumptions P1–P10 hold. Then

$$(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^*-\hat{\beta}) \xrightarrow{d} \arg\max_{\rho \in \mathbb{R}^{k_1}} \mathcal{Z}_1^*(\rho) \ \text{ conditional on the sample},$$

and

$$(\varepsilon_{N2}^{-1}\sigma_N^{2k_1})^{1/3}(\hat{\gamma}^* - \hat{\gamma}) \stackrel{d}{\to} \arg\max_{\delta \in \mathbb{R}^{k_2}} \mathcal{Z}_2^*(\delta)$$
 conditional on the sample,

where $\mathcal{Z}_{1}^{*}(\rho)$ and $\mathcal{Z}_{2}^{*}(\delta)$ are independent copies of $\mathcal{Z}_{1}(\rho)$ and $\mathcal{Z}_{2}(\delta)$ defined in Theorem 3.2, respectively.

Note that our ε_N was written as $\varepsilon_N^{1/2}$ in Hong and Li (2020).

We conclude this section by discussing the choice of the tuning parameters. Recall that $\hat{\beta}$ is of rate $(Nh_N^{k_1+k_2})^{-1/3}$, and additionally we need $(Nh_N^{k_1+k_2})^{2/3}h_N^2 \to 0$ (to handle the bias). To attain a fast rate of convergence, we tend to set h_N as large as possible. For example, we may set $(Nh_N^{k_1+k_2})^{2/3}h_N^2 \propto \log{(N)^{-1}}$. For the same reason, we may set $(N\sigma_N^{2k_1})^{2/3}\sigma_N^2 \propto \log{(N)^{-1}}$. These lead to $h_N \propto \log{(N)^{-\frac{3}{2k_1+2k_2+6}}}N^{-\frac{1}{k_1+k_2+3}}$ and $\sigma_N \propto \log{(N)^{-\frac{3}{4k_1+6}}}N^{-\frac{1}{2k_1+3}}$. As recommended by Hong and Li (2020), we can choose ε_{N1} and ε_{N2} such that $\varepsilon_{N1}^{-1}h_N^{k_1+k_2} \propto (Nh_N^{k_1+k_2})^{2/3}$ and $\varepsilon_{N2}^{-1}\sigma_N^{2k_1} \propto (N\sigma_N^{2k_1})^{2/3}$, which then implies that

$$\varepsilon_{N1} \propto N^{-\frac{k_1+k_2+2}{k_1+k_2+3}} \log(N)^{-\frac{k_1+k_2}{2k_1+2k_2+6}} \text{ and } \varepsilon_{N2} \propto N^{-\frac{2k_1+2}{2k_1+3}} \log(N)^{-\frac{k_1}{2k_1+3}}.$$
 (3.9)

3.1.3 Testing η

In this section, we use a similar approach as in Section 2.1.4 to propose a test for whether η_t tends to degenerate to 0. The hypotheses are formulated as

 $\mathbb{H}_0: \eta_t > 0$ almost surely and $\mathbb{E}(\eta_t) > 0$,

 $\mathbb{H}_1: \eta_t = 0$ almost surely.

We will just briefly outline how to construct this test since the approach is similar to that of Section 2.1.4.

The test is based on the criterion

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(r) = N^{-1} \sum_{i=1}^{N} \sum_{t>s} \mathcal{K}_{\sigma_N}(X_{i1ts}, X_{i2ts})(Y_{i(1,1)t} - Y_{i(1,1)s}) \operatorname{sgn}(W'_{its}r).$$

We also define

$$\bar{\mathcal{L}}^{P}(r) = \sum_{t>s} f_{X_{1ts}, X_{2ts}}(0, 0) \mathbb{E}\left[\left(Y_{i(1,1)t} - Y_{i(1,1)s}\right) \operatorname{sgn}(W'_{its}r) \middle| X_{1ts} = 0, X_{2ts} = 0\right].$$

Then, we have

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(r) \xrightarrow{p} \bar{\mathcal{L}}^{P}(r)$$

uniformly in a small neighborhood of γ . Using the same reasoning as before, we can conclude that under \mathbb{H}_0 , $\bar{\mathcal{L}}^P(\gamma) > 0$, and under \mathbb{H}_1 , $\bar{\mathcal{L}}^P(\gamma) = 0$. We show the limiting distribution of $N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma})$ in the following theorem. The proof is deferred to Appendix C.

Theorem 3.4. Suppose Assumptions P1–P9 hold. Then, under \mathbb{H}_0 ,

$$\sqrt{N\sigma_N^{2k_1}}(N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma}) - \bar{\mathcal{L}}^P(\gamma)) \stackrel{d}{\to} N(0,\Delta_{\gamma}^P),$$

where

$$\Delta_{\gamma}^{P} = \lim_{N \to \infty} \operatorname{Var} \left(\sigma_{N}^{k_{1}} \sum_{t > s} \mathcal{K}_{\sigma_{N}}(X_{i1ts}, X_{i2ts}) (Y_{i(1,1)t} - Y_{i(1,1)s}) \operatorname{sgn}(W'_{its}\gamma) \right).$$

Some standard calculations can verify that Δ_{γ}^{P} is well defined. Importantly, the plug-in estimator $\hat{\gamma}$ does not affect the asymptotics of $N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma})$. To understand this, we notice from the proof that

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma}) - N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\gamma) = O_p(\|\hat{\gamma} - \gamma\|^2) + O_p(\sigma_N^2) + O_p((N\sigma_N^{2k_1})^{-2/3}).$$

Since $\hat{\gamma} - \gamma = O_p((N\sigma_N^{2k_1})^{-1/3})$ by Theorem 3.2, we can show, by Assumption P9, that

$$\sqrt{N\sigma_N^{2k_1}}(N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma}) - N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\gamma)) = o_p(1).$$

For the implementation, one can employ the standard bootstrap with no need to re-estimate $\hat{\gamma}$. The validity of this approach can be justified in a similar way to what was discussed in Section 2.1.4.

3.2 LAD Estimator for Panel Data Models

The matching-based MS procedure proposed in Section 3.1 shares the same limitation as the MRC estimator in Section 2.1, as it cannot identify the coefficients for (time-varying) common regressors (e.g., income). In this section, we introduce a LAD estimation method for panel data bundle choice models, similar to the one discussed in Section 2.2 for cross-sectional models. This LAD estimator can estimate coefficients for both common and alternative-specific regressors. It offers the same level of robustness as the localized MS estimation (3.5)–(3.6), meaning it does not rely on distributional assumptions and can handle general correlations between fixed effects and regressors. The price to pay is that its implementation requires a first-step nonparametric estimation of choice probabilities.

We extend the model (3.1)–(3.2) to accommodate common regressors, similar to what

we did in Section 2.2 for the cross-sectional model. We consider the following panel data model:

$$U_{dt} = \sum_{j=1}^{2} F_{j}(X'_{jt}\beta + S'_{t}\rho_{j}, \alpha_{j}, \epsilon_{jt}) \cdot d_{j} + \eta_{t} \cdot F_{b}(W'_{t}\gamma + S'_{t}\rho_{b} + \alpha_{b}) \cdot d_{1} \cdot d_{2},$$
(3.10)

$$Y_{dt} = 1[U_{dt} > U_{d't}, \forall d' \in \mathcal{D} \setminus d], \tag{3.11}$$

where $S_t \in \mathbb{R}^{k_3}$ represents a vector of common regressors that enter more than one standalone alternative or the bundle. All other aspects of the model remain the same as in (3.1)–(3.2). We denote $Z_t = (X_{1t}, X_{2t}, W_t, S_t)$ and use $Z^T = (Z_1, ..., Z_T)$ to collect all observed covariates in (3.10)–(3.11).

Suppose $\xi_s \stackrel{d}{=} \xi_t | (\alpha, Z_t, Z_s)$ for any two time periods t and s. We can write the following identification inequalities:

$$\{X'_{1ts}\beta + S'_{ts}\rho_1 \ge 0, X'_{2ts}\beta + S'_{ts}\rho_2 \le 0, W'_{ts}\gamma + S'_{ts}\rho_b \le 0\} \Rightarrow \Delta p_{(1,0)}(Z_t, Z_s) \ge 0, \quad (3.12)$$

$$\{X'_{1ts}\beta + S'_{ts}\rho_1 \le 0, X'_{2ts}\beta + S'_{ts}\rho_2 \ge 0, W'_{ts}\gamma + S'_{ts}\rho_b \ge 0\} \Rightarrow \Delta p_{(1,0)}(Z_t, Z_s) \ge 0, \quad (3.13)$$

where $S_{ts} \equiv S_t - S_s$ and $\Delta p_{(1,0)}(Z_t, Z_s) \equiv P(Y_{(1,0)t} = 1|Z_t, Z_s) - P(Y_{(1,0)s} = 1|Z_t, Z_s)$. It is clear that (3.12) and (3.13) are analogous to (2.11) and (2.12). However, in the context of a panel data model, two differences from the cross-sectional model should be noted:

- We are comparing the same agent in different time periods, rather than different agents.
- $\Delta p_{(1,0)}(Z_t, Z_s)$ is defined as the difference between choice probabilities of (1,0) in time periods t and s, conditioning on all observable covariates from these two periods. This is because we do not impose any restriction on the way that α is dependent on (Z_t, Z_s) , and thus we must keep both Z_t and Z_s in the conditioning sets when integrating out the α in (3.12) and (3.13).¹¹

Inequalities (3.12) and (3.13) lead to the following criterion to identify model parameters $\theta \equiv (\beta, \gamma, \rho_1, \rho_2, \rho_b)$:

$$Q_{ts(1,0)}^{P}(\vartheta) = \mathbb{E}\left[q_{(1,0)}^{P}(Z_t, Z_s, \vartheta)\right], \tag{3.14}$$

The Recall that in cross-sectional model, $\Delta p_{(1,0)}(Z_i, Z_m) \equiv P(Y_{(1,0)i} = 1|Z_i) - P(Y_{(1,0)m} = 1|Z_m)$ since (Y_i, Z_i) and (Y_m, Z_m) are assumed to be independent.

where $\vartheta = (b, r, \varrho_1, \varrho_2, \varrho_b)$ is an arbitrary vector in the parameter space Θ containing θ and

$$q_{(1,0)}^{P}(Z_{t}, Z_{s}, \vartheta) = [|I_{ts(1,0)}^{+}(\vartheta) - 1[\Delta p_{(1,0)}(Z_{t}, Z_{s}) \ge 0]| + |I_{ts(1,0)}^{-}(\vartheta) - 1[\Delta p_{(1,0)}(Z_{t}, Z_{s}) \le 0]|] \times [I_{ts(1,0)}^{+}(\vartheta) + I_{ts(1,0)}^{-}(\vartheta)]$$

with

$$I_{ts(1,0)}^{+}(\vartheta) = 1[X_{ts1}'b + S_{ts}'\varrho_1 \ge 0, X_{ts2}'b + S_{ts}'\varrho_2 \le 0, W_{ts}'r + S_{ts}'\varrho_b \le 0] \text{ and}$$
(3.15)

$$I_{ts(1,0)}^{-}(\vartheta) = 1[X_{ts1}'b + S_{ts}'\varrho_1 \le 0, X_{ts2}'b + S_{ts}'\varrho_2 \ge 0, W_{ts}'r + S_{ts}'\varrho_b \ge 0].$$
(3.16)

Under the following assumptions, Theorem 3.5 establishes the identification of θ based on (3.14). The proof is omitted due to its similarity to that of Theorem 2.5.

- **PL1** (i) $\{(Y_i^T, Z_i^T)\}_{i=1}^N$ are i.i.d, where $Y_i^T \equiv \{(Y_{i(0,0)t}, Y_{i(1,0)t}, Y_{i(0,1)t}, Y_{i(1,1)t})\}_{t=1}^T$, (ii) for almost all (α, Z^T) , $\xi_t \stackrel{d}{=} \xi_s | (\alpha, Z_t, Z_s)$, and (iii) the distribution of ξ_t conditional on α is absolutely continuous w.r.t. the Lebesgue measure on $\mathbb{R}^2 \times \mathbb{R}_+$ for all t.
- **PL2** $\theta = (\beta, \gamma, \rho_1, \rho_2, \rho_b) \in \Theta \subset \mathbb{R}^{k_1 + k_2 + 3k_3}$, where $\Theta = \{ \vartheta = (b, r, \varrho_1, \varrho_2, \varrho_b) | b^{(1)} = r^{(1)} = 1, \|\vartheta\| \le C \}$ for some constant C > 0.
- **PL3** (i) $X_{1ts}^{(1)}(X_{2ts}^{(1)})$ has a.e. positive Lebesgue density on \mathbb{R} conditional on $(\tilde{X}_{1ts}, X_{2ts}, W_{ts}, S_{ts})$ $((\tilde{X}_{2ts}, X_{1ts}, W_{ts}, S_{ts}))$, (ii) the support of (X_{1ts}, S_{ts}) $((X_{2ts}, S_{ts}))$ is not contained in any proper linear subspace of $\mathbb{R}^{k_1+k_3}$ conditional on (X_{2ts}, W_{ts}) $((X_{1ts}, W_{ts}))$, (iii) $W_{ts}^{(1)}$ has a.e. positive Lebesgue density on \mathbb{R} conditional on $(\tilde{W}_{ts}, X_{1ts}, X_{2ts}, S_{ts})$, and (iv) the support of (W_{ts}, S_{ts}) is not contained in any proper linear subspace of $\mathbb{R}^{k_2+k_3}$ conditional on (X_{1ts}, X_{2ts}) .
- **PL4** $F_{j}(\cdot,\cdot,\cdot)$, j=1,2, and $F_{b}(\cdot)$ are strictly increasing in their arguments. $F_{b}(0)=0$.

Theorem 3.5. Under Assumptions PL1-PL4,
$$Q_{ts(1,0)}^P(\theta) < Q_{ts(1,0)}^P(\vartheta)$$
 for all $\vartheta \in \Theta \setminus \{\theta\}$.

Following similar arguments to (2.16), we can show that θ can be equivalently identified by the following criterion, which results in less bias in its empirical counterpart:

$$Q_{ts(1,0)}^{P,D}(\vartheta) = \mathbb{E}\left[q_{(1,0)}^{P,D}(Z_t, Z_s, \vartheta)\right],$$
(3.17)

where

$$q_{(1,0)}^{P,D}(Z_t, Z_s, \vartheta) = [|I_{ts(1,0)}^+(\vartheta) - \Delta p_{(1,0)}(Z_t, Z_s)| + |I_{ts(1,0)}^-(\vartheta) + \Delta p_{(1,0)}(Z_t, Z_s)|] \times [I_{ts(1,0)}^+(\vartheta) + I_{ts(1,0)}^-(\vartheta)] + [1 - (I_{ts(1,0)}^+(\vartheta) + I_{ts(1,0)}^-(\vartheta))].$$

Then, assuming a uniformly consistent estimator $\Delta \hat{p}_{(1,0)}(Z_t, Z_s)$ for $\Delta p_{(1,0)}(Z_t, Z_s)$, we propose the following LAD estimator $\hat{\theta}$ for θ :

$$\hat{\theta} = \arg\min_{\vartheta \in \Theta} \sum_{i=1}^{N-1} \sum_{t>s} \hat{q}_{its(1,0)}^{P,D}(\vartheta), \tag{3.18}$$

where

$$\hat{q}_{its(1,0)}^{P,D}(\vartheta) = [|I_{its(1,0)}^+(\vartheta) - \Delta \hat{p}_{(1,0)}(Z_{it}, Z_{is})| + |I_{its(1,0)}^-(\vartheta) + \Delta \hat{p}_{(1,0)}(Z_{it}, Z_{is})|] \times [I_{its(1,0)}^+(\vartheta) + I_{its(1,0)}^-(\vartheta)] + [1 - (I_{its(1,0)}^+(\vartheta) + I_{its(1,0)}^-(\vartheta))].$$

In practice, one can employ identification inequalities analogous to those presented in Remark 2.2 (i.e., (2.18)–(2.23)) and follow a procedure similar to (3.12)–(3.18) to compute $\hat{q}_{its(0,1)}^{P,D}(\vartheta)$, $\hat{q}_{its(0,0)}^{P,D}(\vartheta)$, and $\hat{q}_{its(1,1)}^{P,D}(\vartheta)$. Collectively, a more efficient $\hat{\theta}$ can be obtained by minimizing the sample criterion $\sum_{i=1}^{N} \sum_{t>s} \sum_{d \in \mathcal{D}} \hat{q}_{itsd}^{P,D}(\vartheta)$, where $\mathcal{D} = \{(0,0), (1,0), (0,1), (1,1)\}$.

The following assumption, together with Assumptions PL1–PL4, is sufficient for the consistency of the estimator $\hat{\theta}$ obtained from (3.18):

PL5 $\sup_{(z_t,z_s)\in\mathcal{Z}_t\times\mathcal{Z}_s} |\Delta \hat{p}_{(1,0)}(z_t,z_s) - \Delta p_{(1,0)}(z_t,z_s)| \stackrel{p}{\to} 0 \text{ as } n\to\infty \text{ for all } t\neq s, \text{ where } \mathcal{Z}_t \text{ is the support of } Z_t.$

Theorem 3.6. Suppose Assumptions PL1-PL5 hold. We have $\hat{\theta} \stackrel{p}{\rightarrow} \theta$.

The proof of Theorem 3.6 follows similar arguments to Theorem 2.6 and is omitted for brevity.

4 Monte Carlo Experiments

In this section, we investigate the finite sample performance of our proposed approaches, in both cross-sectional and panel data models, by means of Monte Carlo experiments. We

set sample sizes N=250,500,1000 for cross-sectional designs, and N=1000,2500,5000 for panel data designs, considering the slower convergence rates. All simulation results are obtained from 1000 independent replications. We report these results in tables collected in Appendix A.

We start from a benchmark cross-sectional design (Design 1) specified as follows:

$$U_{id} = \sum_{j=1}^{2} (X_{ij,1} + \beta X_{ij,2} + \rho_j S_i + \epsilon_{ij}) \cdot d_j + \eta_i \cdot (W_{i,1} + \gamma W_{i,2}) \cdot d_1 \cdot d_2,$$

$$Y_{id} = 1[U_{id} > U_{id'}, \forall d' \neq d],$$

where $\beta = \gamma = \rho_1 = \rho_2 = 1$ and $d = (d_1, d_2) \in \{(0, 0), (1, 0), (0, 1), (1, 1)\}$. We impose the scale normalization (Assumption C4) by setting the coefficients on $X_{ij,1}$ and $W_{i,1}$ to be 1, and thus $(\beta, \gamma, \rho_1, \rho_2)$ are free parameters to estimate. In this design, we let $\{\{X_{ij,\iota}\}_{j=1,2;\iota=1,2}, \{W_{i,\iota}\}_{\iota=1,2}, S_i, \{\epsilon_{ij}\}_{j=1,2}, \eta_i\}_{i=1,\ldots,N}$ be independent across i, j, and ι and from each other. $X_{ij,1}$'s and $W_{i,1}$ are Logistic(0,1), $X_{ij,2}$'s are Bernoulli(1/3), $W_{i,2}$, S_i , and ϵ_{ij} 's are N(0,1), and η_i is Beta(2,2).

To implement our MRC estimation, note that larger bandwidth means smaller variation. In light of this observation and requirements under Assumptions C6 and C7, we employ the sixth-order Gaussian kernel ($\kappa_{\beta}=6$) and tuning parameter (bandwidth) $h_N=c_1\hat{\sigma}\cdot N^{-1/8}\log(N)^{1/6}$ for "matching" the four continuous regressors ($k_1=k_2=2$) in the first-step estimation, where $\hat{\sigma}$ represents the standard deviation of the corresponding matching variable. Under this choice, $\sqrt{N}h_N^4\propto \log(N)^{2/3}\to\infty$ and $\sqrt{N}h_N^6\to0$. For similar reasons, we adopt the fourth-order Gaussian kernel ($\kappa_{\gamma}=4$) and bandwidth $\sigma_N=c_2\hat{\sigma}\cdot N^{-1/4}\log(N)^{1/4}$ for the second step, and thus $\sqrt{N}\sigma_N^2\propto \log(N)^{1/2}\to\infty$ and $\sqrt{N}\sigma_N^4\to0$. The discrete variable $X_{ij,2}$ is exactly matched across observations in the first step by using an indicator function. To test the sensitivity of our methods to the choice of bandwidths, we experiment with several values of $c_1\in\{0.6,1,1.4,1.8\}$ and several values of $c_2\in\{1.6,2,2.4,2.8\}$. It turns out that the results are not sensitive to the choice of (c_1,c_2) . Thus, to save space, we only report the results of $(c_1,c_2)=(1,2)$, for which the root mean squared errors (RMSE) are the smallest among all choices.

The implementation of our proposed LAD estimator requires a first-stage nonparametric estimation of $\Delta p_d(Z_i, Z_m)$. For the cross-sectional designs of this section, we use the standard Nadaraya-Watson estimator for conditional probability density (see, e.g., Li and Racine

(2007)). We employ the fourth-order Gaussian kernel¹² for continuous variables and the Aitchison-Aitken kernel for (unordered) discrete variables. To save computation time, we simply choose bandwidths for continuous variables using Silverman's "rule-of-thumb" and set N^{-1} as the smoothing parameter for discrete variables. When there is no need to repeat the computation thousands of times, we recommend considering more robust, data-driven methods, such as the cross-validation (CV) method proposed by Hall, Racine, and Li (2004), to select all the tuning parameters. The computation is conducted using the state-of-art R package np (Hayfield and Racine (2008)).

To compute the MRC and LAD estimators, we apply the differential evolution (DE) algorithm (Storn and Price (1997)) to globally search the optima for their corresponding sample criterion functions proposed in previous sections. The implementation uses the well-developed and tested R package DEoptim (Mullen, Ardia, Gil, Windover, and Cline (2011)).

Results for Design 1 are presented in tables numbered "1" and so on for other designs. The performance of the MRC estimator, its nonparametric bootstrap inference, and the LAD estimator are reported in tables labeled "A", "B", and "C", respectively. For example, Table 1A summarizes the performance of the MRC estimator for Design 1, in which we report the estimator's mean bias (MBIAS) and RMSE. Since these statistics are sensitive to outliers, we also present the median bias (MED) and the median absolute deviations (MAD). Table 1B reports the empirical coverage frequencies (COVERAGE) and lengths (LENGTH) of the 95% confidence intervals (CI) constructed using the standard bootstrap with B = 299 independent draws and estimations. Recall that the MRC procedure can only estimate (β, γ) , while the LAD method can also estimate the coefficients (ρ_1, ρ_2) on the common regressor S_i . Therefore, in Tables 1A and 1B, we only report results for (β, γ) , while in Table 1C we report results for $(\beta, \gamma, \rho_1, \rho_2)$.

The results for Design 1 are in line with our asymptotic theory. As the sample size increases, the RMSE of the MRC estimator shrinks at the parametric rate, indicating that these procedures are \sqrt{N} -consistent. We observe a similar root-n convergence rate for the LAD estimator, although we do not formally derive it in this paper. The MRC estimator has greater RMSE than the LAD estimator, as the latter makes full use of the data and can be more efficient. The nonparametric bootstrap for the MRC estimation also performs well and yields shrinking confidence intervals, with coverage rates approaching 95% as the

¹²We also try the second-order Gaussian kernel, but the simulation results exhibit larger bias than those obtained with higher-order kernels. This is in line with our discussion in Section 2.2, i.e., the bias in $\Delta \hat{p}_d(Z_i, Z_m)$, when making $\Delta \hat{p}_d(Z_i, Z_m)$ and $\Delta p_d(Z_i, Z_m)$ be of different signs, leads to the bias in the LAD estimation. Therefore, it is crucial to prioritize estimators that incur smaller bias to estimate $\Delta p_d(Z_i, Z_m)$.

sample size increases.

The outcome of Monte Carlo experiments can be significantly affected by the design of the experiment itself. For instance, estimators tend to perform better when all i.i.d. regressors and errors are used compared to cases where either the regressors or errors are correlated. In order to evaluate the performance of our estimators with correlated regressors and errors, we modify Design 1 by letting $Corr(X_{i1,1}, X_{i2,1}) = Corr(X_{i1,2}, X_{i2,2}) = Corr(\epsilon_{i1}, \epsilon_{i2}) = 0.5$. All other aspects of Design 1 remain the same. We refer to this modified version of the benchmark design as Design 2. Like Design 1, we use the same kernels and bandwidths in Design 2 and report the simulation results in Tables 2A–2C. The results are very similar to those for Design 1, though as expected, the performance slightly deteriorates. This confirms our theoretical results that our approaches allow for arbitrary correlation in errors.

We proceed to analyze the finite sample performance of the MS estimation, its numerical bootstrap, and the LAD estimator for panel data bundle choice models. We consider two designs (Designs 3 and 4) that have the same choice set as the cross-sectional designs $(\{(0,0),(1,0),(0,1),(1,1)\})$ and a panel of two time periods. We examine sample sizes of N=1000,2500,5000 for each panel data design. We construct 95% CI based on B=299 independent draws and estimations for inference. The first panel data design (Design 3) is specified as follows:

$$U_{idt} = \sum_{j=1}^{2} (X_{ijt,1} + \beta X_{ijt,2} + \rho_j S_{it} + \alpha_{ij} + \epsilon_{ijt}) \cdot d_j + \eta_{it} \cdot (W_{it,1} + \gamma W_{it,2} + \alpha_{ib}) \cdot d_1 \cdot d_2,$$

$$Y_{idt} = 1[U_{idt} > U_{id't}, \forall d' \neq d],$$

where $\beta = \gamma = \rho_1 = \rho_2 = 1$. $\{\{X_{ijt,\iota}\}_{j=1,2;\iota=1,2}, \{W_{it,\iota}\}_{\iota=1,2}, S_{it}, \{\epsilon_{ijt}\}_{j=1,2}, \eta_{it}\}_{i=1,...,N;t=1,2}$ are mutually independent random variables, where $X_{ijt,1}$'s and $W_{it,1}$'s are Logistic(0, 1), $X_{ijt,2}$'s are Bernoulli(1/3), S_{it} 's, $W_{it,2}$'s, and ϵ_{ijt} 's are N(0,1), and $\eta_{it} \sim \text{Beta}(2,2)$. We let $\alpha_{ij} = (X_{ij1,1} + X_{ij2,1})/4$ for j = 1, 2 and $\alpha_{ib} = (W_{i1,1} + W_{i2,1})/4$. For scale normalization (Assumption P3), we set the coefficients on $X_{ijt,1}$ and $W_{it,1}$ to be 1.¹³ In this design, the (unobserved) fixed effects are correlated with regressors, which invalidates the matching method in the cross-sectional case.

In the second panel data design (Design 4), the random utility model and coefficients are the same as in Design 3. However, similar to Design 2 for the cross-sectional setting, we

¹³Note that we employ the scale normalization in Assumption C3 rather than Assumption P2. The two normalization conventions are equivalent, but the former is often more computationally convenient and easier to interpret.

set $Corr(X_{ijt,1}, X_{ils,1}) = Corr(X_{ijt,2}, X_{ils,2}) = Corr(\epsilon_{ijt}, \epsilon_{ils}) = 0.25$ for all $j, l \in \{1, 2\}$ and $t, s \in \{1, 2\}$ with either $j \neq l$ or $t \neq s$, and $\{S_{it}, \{W_{it, l}\}_{l=1, 2}, \eta_{it}\}_{i=1, ..., N; t=1, 2}$ to have serial correlation 0.25. All other aspects remain the same.

To implement the MS estimation for these designs, we use the second-order Gaussian kernel and $h_N = \sigma_N = c_3 \hat{\sigma} \cdot N^{-1/7} \log(N)^{-1/14}$. We have two more tuning parameters to implement the numerical bootstrap: ε_{N1} and ε_{N2} . We set $\varepsilon_{N1} = \varepsilon_{N2} = c_4 \cdot N^{-5/7} \log(N)^{-5/14}$. This choice of $(h_N, \sigma_N, \varepsilon_{N1}, \varepsilon_{N2})$ meets Assumptions P9 and P10. We experiment with $c_3, c_4 \in \{1, 1.5, 2, 2.5\}$. It turns out that the results are not sensitive to the choice of (c_3, c_4) . Therefore, to save space, we report only the results with $(c_3, c_4) = (2, 2)$.

To implement our proposed LAD estimator, we need to estimate $\Delta p_d(Z_{it}, Z_{is})$ nonparametrically in the first stage. It is important to note that in order to estimate the probability of $Y_{idt} = 1$, we need to use observed (time-varying) regressors from both periods t and s due to the weak restrictions imposed on the unobservables of the model in Assumption P1. This makes the panel data LAD estimator more susceptible to the curse of dimensionality than the cross-sectional model. To address this issue, we employ nonparametric regression based on artificial neural networks with multiple hidden layers. Recent advances in statistics, such as Kohler and Langer (2021) and the references therein, establish the convergence rates of such estimators and demonstrate their ability to handle the curse of dimensionality under certain restrictions on the regression function and network (sparsity) architecture. Our simulation studies simply use neural networks with two hidden layers and three neurons per hidden layer for computational efficiency. The R package neuralnet (Fritsch and Guenther (2019)) is used for computation. In practice, one can consider using CV techniques to select the number of hidden layers and neurons.

Tables 3A–3C and Tables 4A–4C show the simulation results for Designs 3 and 4, respectively. According to the results, our MS and LAD estimators, calculated using the DE algorithm, perform reasonably well and yield similar results for these two designs. The MBIAS and RMSE of these estimators decrease as the sample size increases, indicating the consistency of our estimators. The MS estimator has a slower than cube root-N rate of convergence, while the LAD estimator appears to have a rate of convergence between \sqrt{N} and cube root-N. The inference results are also consistent with the theory. When the sample

¹⁴Though Hong and Li (2020) recommended ε_{N1} , $\varepsilon_{N2} \propto N^{-6/7} \log(N)^{-2/7}$ (see (3.9) in our Section 3.1.2), some tentative Monte Carlo simulations suggest that a slight modification like this leads to a better finite sample performance.

¹⁵Strictly speaking, the rate results of Kohler and Langer (2021) only apply to cases where the covariates have compact supports, which is not the case for our simulation designs. However, despite the lack of theoretical guarantees, our simulation results do not flag any problems.

size is small, the CI obtained by the numerical bootstrap tends to be too wide, resulting in coverage rates slightly higher than 95%. However, as the sample size increases, we observe that the CI shrink and their coverage rates approach 95%.

5 Conclusions

This paper develops estimation and inference procedures for semiparametric discrete choice models for bundles. We first propose a two-step maximum rank correlation estimator in the cross-sectional setting and establish its \sqrt{N} -consistency and asymptotic normality. Based on these results, we further show the validity of using the nonparametric bootstrap to carry out inference and propose a criterion-based test for the interaction effects. Additionally, we introduce a multi-index least absolute deviations (LAD) estimator to handle models with both alternative- and agent-specific covariates, which, to our knowledge, is new in the literature. We extend these approaches to the panel data setting, where the resulting maximum score (MS) and LAD estimators can consistently estimate bundle choice models with agent-, alternative-, and bundle-specific fixed effects. We also derive the limiting distribution of the MS estimator, justify the application of the numerical bootstrap for making inferences, and introduce an analogous test for interaction effects based on the MS criterion. It is important to note that although we propose these methods for estimating bundle choice models, they can be applied with minor modifications to many other multi-index models. A small-scale Monte Carlo study shows that our proposed methods perform adequately in finite samples.

This paper focuses on scenarios where the researcher can access individual-level data. We note that our proposed methods can also be adapted for models that use aggregate data. In these cases, the researcher can observe the aggregated choice probabilities (e.g., market shares) in many markets and the market-level covariates for each alternative. These types of models are commonly used in empirical industrial organizations (see, e.g., Fan (2013) and Shi et al. (2018)). A detailed exploration of this adaptation would require different estimators and asymptotics. Due to space limitations, we do not delve into this topic in this paper and reserve it for future study.

The work here leaves many other open questions for future research. One promising subsequent research is to establish the asymptotic properties of the proposed LAD estimators for both cross-sectional and panel data models, which can further lead to the development of easy-to-implement inference methods. Another potential area of research is to explore

the possibility of smoothing the criterion functions to achieve faster convergence rates and attain asymptotic normality for the MS and LAD estimators.

Supplementary Materials

Appendixes B–F are included in a supplementary appendix. Appendix B provides proofs for theorems presented in Section 2 for our cross-sectional estimators, except for Theorem 2.3, whose technical details are provided in Appendix D due to its length. Appendix C provides proofs for all the theorems for our panel data estimators introduced in Section 3. Appendix E provides additional discussions on the convergence rates of proposed procedures, the construction of our MS estimator, and the identification of models with more alternatives and bundles. Finally, Appendix F collects the proofs of all the technical lemmas used in Appendices B and C.

Acknowledgements

This work is partially supported by the Faculty of Business, Economics, and Law (BEL), University of Queensland, via the 2019 BEL New Staff Research Start-Up Grant. We thank Shakeeb Khan, Dong-Hyuk Kim, Firmin Tchatoka, Hanghui Zhang, and Yu Zhou for their valuable feedback. We are also grateful for the helpful comments and discussions from all seminar and conference participants at the University of Adelaide and the 31st Australia New Zealand Econometric Study Group Meeting. All errors are our responsibility.

Appendix

A Tables

Table 1A: Design 1, Performance of MRC Estimators

		β				γ		
	MBIAS	RMSE	MED	MAD	MBIAS	RMSE	MED	MAD
N = 250	0.068	0.534	0.075	0.424	0.101	0.452	0.074	0.318
N = 500	0.024	0.367	0.028	0.274	0.065	0.314	0.038	0.208
N = 1000	0.024	0.237	0.016	0.154	0.040	0.215	0.034	0.131

Table 1B: Design 1, Performance of Nonparametric Bootstrap

	β			γ				
	COVERAGE	LENGTH	С	OVERAGE	LENGTH			
N = 250	0.835	2.505		0.915	2.145			
N = 500	0.862	1.719		0.918	1.492			
N = 1000	0.926	1.164		0.938	1.002			

Table 1C: Design 1, Performance of LAD Estimators

		β					γ		
	MBIAS	RMSE	MED	MAD	•	MBIAS	RMSE	MED	MAD
N = 250	0.102	0.274	0.090	0.180	•	0.040	0.418	0.017	0.247
N = 500	0.112	0.213	0.114	0.144		0.032	0.274	0.004	0.173
N = 1000	0.106	0.164	0.100	0.110		0.030	0.189	0.022	0.124

		ρ_1					$ ho_2$		
	MBIAS	RMSE	MED	MAD	•	MBIAS	RMSE	MED	MAD
N = 250	0.022	0.208	0.011	0.133	•	0.019	0.200	0.011	0.128
N = 500	0.010	0.134	0.005	0.092		0.009	0.135	0.001	0.089
N = 1000	0.010	0.095	0.007	0.062		0.013	0.092	0.011	0.061

Table 2A: Design 2, Performance of MRC Estimators

		β				γ		
	MBIAS	RMSE	MED	MAD	MBIAS	RMSE	MED	MAD
N = 250 $N = 500$		0.585 0.433				0.471 0.328		
N = 300 $N = 1000$		0.453 0.283				0.328 0.219		

Table 2B: Design 2, Performance of Nonparametric Bootstrap

	β			γ			
	COVERAGE	LENGTH		COVERAGE	LENGTH		
N = 250	0.870	2.847	•	0.911	2.292		
N = 500	0.894	1.992		0.927	1.566		
N = 1000	0.925	1.358		0.933	1.067		

Table 2C: Design 2, Performance of LAD Estimators

		β					γ		
	MBIAS	RMSE	MED	MAD	-	MBIAS	RMSE	MED	MAD
N = 250	0.109	0.312	0.102	0.205	-	0.059	0.491	0.018	0.297
N = 500	0.107	0.222	0.103	0.147		0.048	0.308	0.023	0.189
N = 1000	0.112	0.172	0.113	0.124		0.028	0.209	0.013	0.143

		$ ho_1$					ρ_2		
	MBIAS	RMSE	MED	MAD		MBIAS	RMSE	MED	MAD
N = 250	-0.042	0.209	-0.050	0.139	-	-0.036	0.211	-0.047	0.154
N = 500	-0.034	0.146	-0.040	0.098		-0.037	0.147	-0.037	0.098
N = 1000	-0.041	0.105	-0.043	0.074		-0.037	0.100	-0.039	0.073

Table 3A: Design 3, Performance of MS Estimators

		β				γ		
	MBIAS	RMSE	MED	MAD	MBIAS	RMSE	MED	MAD
N = 1000	-0.008	0.371	0.003	0.244	-0.014	0.411	-0.057	0.276
N = 2500	0.006	0.332	0.003	0.236	-0.004	0.375	-0.040	0.246
N = 5000	-0.002	0.291	0.001	0.235	0.009	0.367	-0.016	0.252

Table 3B: Design 3, Performance of Numerical Bootstrap

	β			γ			
	COVERAGE	LENGTH	-	COVERAGE	LENGTH		
N = 1000	0.970	2.275		0.967	2.837		
N = 2500	0.967	1.973		0.971	2.527		
N = 5000	0.954	1.743		0.967	2.256		

Table 3C: Design 3, Performance of LAD Estimators

		β					γ		
	MBIAS	RMSE	MED	MAD		MBIAS	RMSE	MED	MAD
N = 1000	0.046	0.375	0.036	0.250	·	0.131	0.740	0.021	0.466
N = 2500	0.026	0.205	0.020	0.133		0.069	0.440	0.027	0.281
N = 5000	0.002	0.155	-0.003	0.107		0.067	0.327	0.039	0.214

		$ ho_1$					ρ_2		
	MBIAS	RMSE	MED	MAD		MBIAS	RMSE	MED	MAD
N = 1000	0.027	0.320	-0.006	0.198	-	0.031	0.318	0.000	0.203
N = 2500	0.012	0.172	0.011	0.112		0.009	0.167	0.012	0.114
N = 5000	0.002	0.118	-0.002	0.080		0.000	0.126	-0.006	0.086

Table 4A: Design 4, Performance of MS Estimators

	β				γ				
	MBIAS	RMSE	MED	MAD		MBIAS	RMSE	MED	MAD
N = 1000	-0.014	0.353	-0.012	0.241	-	-0.003	0.411	-0.034	0.280
N = 2500	0.000	0.0-0	0.00.	00		0.014		-0.006	
N = 5000	0.016	0.274	0.017	0.232		0.027	0.354	0.002	0.246

Table 4B: Design 4, Performance of Numerical Bootstrap

	β		γ			
	COVERAGE	LENGTH	COVERA	AGE LENGTH		
N = 1000	0.976	2.159	0.972	2.720		
N = 2500	0.959	1.856	0.969	2.383		
N = 5000	0.956	1.642	0.962	2.115		

Table 4C: Design 4, Performance of LAD Estimators

	β				γ				
	MBIAS	RMSE	MED	MAD	MBIAS	RMSE	MED	MAD	
N = 1000	0.060	0.337	0.045	0.215	0.132	0.666	0.025	0.406	
N = 2500	0.011	0.172	0.000	0.113	0.068	0.404	0.026	0.246	
N = 5000	0.002	0.128	-0.001	0.087	0.052	0.340	0.006	0.209	

	$ ho_1$					$ ho_2$				
	MBIAS	RMSE	MED	MAD		MBIAS	RMSE	MED	MAD	
N = 1000	0.069	0.328	0.029	0.197	_	0.044	0.314	0.011	0.204	
N = 2500	-0.015	0.163	-0.024	0.110		-0.009	0.167	-0.019	0.116	
N = 5000	-0.028	0.124	-0.035	0.087		-0.024	0.128	-0.024	0.087	

B Proofs for the Cross-Sectional Model

B.1 Proof of Theorem 2.1

Proof of Theorem 2.1. It suffices to show the identification of β based on (2.4) as the same arguments can be applied to the identification of β and γ based on similar moment inequalities. Denote $\Omega_{im} = \{X_{i2} = X_{m2}, W_i = W_m\}$. By Assumption C1, the monotonic relation (2.4) implies that for all $d \in \mathcal{D}$ with $d_1 = 1$, β maximizes

$$Q_1(b) \equiv \mathbb{E}[(P(Y_{i(1,d_2)} = 1|Z_i) - P(Y_{m(1,d_2)} = 1|Z_m))\operatorname{sgn}(X'_{im1}b)|\Omega_{im}]$$

for each pair of (i, m). To show that β attains a unique maximum, suppose that there is a $b \in \mathcal{B}$ such that $Q_1(b) = Q_1(\beta)$. In what follows, we show that $b = \beta$ must hold. To this end, we write $P[\operatorname{sgn}(X'_{im1}b) \neq \operatorname{sgn}(X'_{im1}\beta) | \Omega_{im}] = P[(\tilde{X}'_{im1}\beta < -X^{(1)}_{im1} < \tilde{X}'_{im1}b) \cup (\tilde{X}'_{im1}b < -X^{(1)}_{im1} < \tilde{X}'_{im1}\beta) | \Omega_{im}]$. Note that when $b \neq \beta$, $P(\tilde{X}'_{im1}\beta \neq \tilde{X}'_{im1}b | \Omega_{im}) > 0$ under Assumption C2(ii). Then by Assumption C2(i), we have $P[(\tilde{X}'_{im1}\beta < -X^{(1)}_{im1} < \tilde{X}'_{im1}b) \cup (\tilde{X}'_{im1}b < -X^{(1)}_{im1} < \tilde{X}'_{im1}\beta) | \Omega_{im}] > 0$. Combining with (2.4), this result implies that the event of $X'_{im1}b$ and $P(Y_{i(1,d_2)} = 1|Z_i) - P(Y_{m(1,d_2)} = 1|Z_m)$ having different signs occurs with strictly positive probability, and hence $Q_1(b) < Q_1(\beta)$, a contradiction. Since b is arbitrary, we conclude that $Q_1(b) \leq Q_1(\beta)$ and the equality holds only when $b = \beta$. This completes the proof. \square

B.2 Proof of Theorem 2.2

The proof of Theorem 2.2 involves utilizing several technical lemmas (i.e., Lemmas B.1–B.7). We will present these lemmas prior to the main proof. Before delving into the technical details, we will first introduce some new notations, state Assumption C5 that were omitted from the main text, and provide an outline of the proof process. In what follows, we assume that all regressors are continuous to streamline the exposition. ¹⁶

The following notations will be used for convenience: For all $d \in \mathcal{D}$ and j = 1, 2,

- $Y_{imd} \equiv Y_{id} Y_{md}$.
- $f_{X_{imj},W_{im}}(\cdot)$ and $f_{V_{im}(\beta)}(\cdot)$ denote the PDF of the random vectors (X_{imj},W_{im}) and $V_{im}(\beta)$, respectively. $f_{X_2,W|X_1}(\cdot)$ $(f_{X_1,W|X_2}(\cdot))$ denotes the PDF of (X_2,W) $((X_1,W))$ conditional on X_1 (X_2) . $f_{V(\beta)}(\cdot)$ $(f_{V(\beta)|W}(\cdot))$ is the PDF of $V(\beta)$ (conditional on W).
- For any function g, $\nabla g(v)$ and $\nabla^2 g(v)$ denote the gradient and Hessian matrix for $g(\cdot)$ evaluated at v, respectively.
- $V_i \equiv V_i(\beta)$, $\hat{V}_i \equiv V_i(\hat{\beta})$, $V_{im} \equiv V_i V_m$, and $\hat{V}_{im} \equiv \hat{V}_i \hat{V}_m$.
- For realized values, $v_i \equiv v_i(\beta)$, $\hat{v}_i \equiv v_i(\hat{\beta})$, $v_{im} \equiv v_i v_m$, and $\hat{v}_{im} \equiv \hat{v}_i \hat{v}_m$.
- $B(v_i, v_m, w_i, w_m) \equiv \mathbb{E}[Y_{im(1,1)}|V_i = v_i, V_m = v_m, W_i = w_i, W_m = w_m].$
- $S_{im}(r) \equiv \operatorname{sgn}(w'_{im}r) \operatorname{sgn}(w'_{im}\gamma)$.
- Define

$$\varrho_i(b) \equiv -\sum_{d \in \mathcal{D}} \{ \varrho_{i21d}(b) \cdot (-1)^{d_1} + \varrho_{i12d}(b) \cdot (-1)^{d_2} \}, \tag{B.1}$$

where

$$\varrho_{ijld}(b) \equiv \mathbb{E}\left[Y_{imd}\left[\operatorname{sgn}(X'_{iml}b) - \operatorname{sgn}(X'_{iml}\beta)\right] | Z_i, X_{mj} = X_{ij}, W_m = W_i\right]$$

for j, l = 1, 2 with $j \neq l$.

- Define

$$\tau_{i}(r) \equiv \mathbb{E}[Y_{im(1,1)}\left[\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma)\right] | V_{m}\left(\beta\right) = V_{i}\left(\beta\right), W_{i}]. \tag{B.2}$$

¹⁶This is a purely technical assumption, which is made to facilitate the derivation of the asymptotic variance of the proposed estimators. With discrete covariates, the asymptotic results summarized in Theorem 2.2 still valid but with more complex expressions.

- Define

$$\mu(v_{1}, v_{2}, r)$$

$$\equiv \mathbb{E}\left[Y_{im(1,1)}\left[\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma)\right] \begin{pmatrix} X'_{im1} \\ X'_{im2} \end{pmatrix} \middle| V_{i}\left(\beta\right) = v_{1}, V_{m}\left(\beta\right) = v_{2}\right] f_{V(\beta)}\left(v_{1}\right).$$
(B.3)

Let $\nabla_1 \mu(v_1, v_2, r)$ denote the partial derivative of $\mu(v_1, v_2, r)$ w.r.t. its first argument. Denote $\nabla_{1k}^2 \mu(v_1, v_2, r)$ as the partial derivative of $\nabla_1 \mu(v_1, v_2, r)$ w.r.t. its k-th argument and $\nabla_{33}^2 \nabla_1 \mu(v_1, v_2, r)$ as the Hessian matrix of $\nabla_1 \mu(v_1, v_2, r)$ w.r.t. its third argument.

- $K_{h_N,\beta}(\cdot) \equiv h_N^{k_1+k_2} \mathcal{K}_{h_N}(\cdot)$ and $K_{\sigma_N,\gamma}(\cdot) \equiv \sigma_N^2 \mathcal{K}_{\sigma_N}(\cdot)$. In proofs, we also write $K_{\beta}(\cdot/h_N)$ and $K_{\gamma}(\cdot/\sigma_N)$ for $K_{h_N,\beta}(\cdot)$ and $K_{\sigma_N,\gamma}(\cdot)$, respectively, when we need to apply changes of variables.

The following regularity conditions that were omitted in the main text will be used in proving Lemmas B.1–B.7 and Theorem 2.2:

C5 (i) $f_{V_{im}(\beta)}(\cdot)$ and $f_{X_{imj},W_{im}}(\cdot)$ for j=1,2 are bounded from above on their supports, strictly positive in a neighborhood of zero, and twice continuously differentiable with bounded second derivatives, (ii) For all $d \in \mathcal{D}$, $b \in \mathcal{B}$, and $r \in \mathcal{R}$, $\mathbb{E}[Y_{imd}[\operatorname{sgn}(X'_{im1}b) - \operatorname{sgn}(X'_{im1}\beta)]|X_{im2} = \cdot, W_{im} = \cdot]$, $\mathbb{E}[Y_{imd}[\operatorname{sgn}(X'_{im2}b) - \operatorname{sgn}(X'_{im2}\beta)]|X_{im1} = \cdot, W_{im} = \cdot]$, and $\mathbb{E}[Y_{im(1,1)}[\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma)]|V_{im}(\beta) = \cdot]$ are continuously differentiable with bounded first derivatives, (iii) $\mathbb{E}[Y_{imd}|Z_i = \cdot, Z_m = \cdot]$ is $\kappa_{1\beta}^{\text{th}}$ continuously differentiable with bounded $\kappa_{1\beta}^{\text{th}}$ derivatives. $f_{X_2,W|X_1}(\cdot)$ $(f_{X_1,W|X_2}(\cdot))$ is $\kappa_{2\beta}^{\text{th}}$ continuously differentiable with bounded $\kappa_{2\beta}^{\text{th}}$ derivatives. Denote $\kappa_{\beta} = \kappa_{1\beta} + \kappa_{2\beta}$. κ_{β} is an even integer greater than $k_1 + k_2$, (iv) $\mathbb{E}[Y_{im(1,1)}|V_i = \cdot, V_m = \cdot, W_i = \cdot, W_m = \cdot]$ is $\kappa_{1\gamma}^{\text{th}}$ continuously differentiable with bounded $\kappa_{1\gamma}^{\text{th}}$ derivatives, and $f_{V(\beta)|W}(\cdot)$ is $\kappa_{2\gamma}^{\text{th}}$ continuously differentiable with bounded $\kappa_{2\gamma}^{\text{th}}$ derivatives. Denote $\kappa_{\gamma} = \kappa_{1\gamma} + \kappa_{2\gamma}$. κ_{γ} is an even integer greater than 2, and (v) $\mathbb{E}[\|X_{imj}\|^2] < \infty$ for all j = 1, 2, and $\mathbb{E}[\|W_{im}\|^2] < \infty$. All derivatives in this assumption are with respect to \cdot .

The boundedness and smoothness restrictions under Assumption C5 are needed for proving the uniform convergence of the criterion functions to their population analogues.

In what follows, we will only show the asymptotics of $\hat{\gamma}$. The proof for $\hat{\beta}$ is omitted since one can derive the asymptotics of $\hat{\beta}$ by repeating the proof process for $\hat{\gamma}$ but skipping the

step handling the plug-in first step estimates (i.e., Lemma B.7). Throughout this section, we take it as a given that $\hat{\beta}$ is \sqrt{N} -consistent and asymptotically normal.

For ease of illustration, with a bit of abuse of notation, we will work with criterion function

$$\hat{\mathcal{L}}_{N}^{K}(r) \equiv \frac{1}{\sigma_{N}^{2}N(N-1)} \sum_{i \neq m} K_{\sigma_{N},\gamma}(V_{im}(\hat{\beta})) h_{im}(r),$$

where $K_{\sigma_N,\gamma}(\cdot) \equiv \sigma_N^2 \mathcal{K}_{\sigma_N}(\cdot)$ and $h_{im}(r) \equiv Y_{im(1,1)}[\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma)]$. Note that here we subtract the term $\operatorname{sgn}(W'_{im}\gamma)$ from criterion function (2.7), analogous to Sherman (1993). Doing this does not affect the value of the estimator, and will facilitate the proofs that follow. Besides, we define

$$\mathcal{L}_N^K(r) = \frac{1}{\sigma_N^2 N(N-1)} \sum_{i \neq m} K_{\sigma_N, \gamma}(V_{im}(\beta)) h_{im}(r)$$

and

$$\mathcal{L}(r) \equiv f_{V_{im}(\beta)}(0) \mathbb{E}[h_{im}(r)|V_{im}(\beta) = 0].$$

We establish the consistency of $\hat{\gamma}$ in Lemmas B.1–B.3 by applying Theorem 2.1 in Newey and McFadden (1994). The key step is to show the uniform convergence of $\hat{\mathcal{L}}_{N}^{K}(r)$ for $r \in \mathcal{R}$. To this end, we bound the differences among $\hat{\mathcal{L}}_{N}^{K}(r)$, $\mathcal{L}_{N}^{K}(r)$, and $\mathcal{L}(r)$ in Lemmas B.1 and B.2.

The next step is to show the asymptotic normality of $\hat{\gamma}$ by applying Theorem 2 of Sherman (1994a). Sufficient conditions for this theorem are that $\hat{\gamma} - \gamma = O_p(N^{-1/2})$ and uniformly over a neighborhood of γ with a radius proportional to $N^{-1/2}$,

$$\hat{\mathcal{L}}_{N}^{K}(r) = \frac{1}{2}(r - \gamma)' \mathbb{V}_{\gamma}(r - \gamma) + \frac{1}{\sqrt{N}}(r - \gamma)' \Psi_{N} + o_{p}(N^{-1}), \tag{B.4}$$

where V_{γ} is a negative definite matrix and Ψ_N is asymptotically normal, with mean zero and variance V_{Ψ} . To verify equation (B.4), we first show that uniformly over a neighborhood of γ , $\mathcal{R}_N \equiv \{r \in \mathcal{R} | ||r - \gamma|| \leq \delta_N \}$ with $\{\delta_N\} = O(N^{-\delta})$ for some $0 < \delta \leq 1/2$,

$$\hat{\mathcal{L}}_{N}^{K}(r) = \frac{1}{2}(r-\gamma)^{\prime} \mathbb{V}_{\gamma}(r-\gamma) + \frac{1}{\sqrt{N}}(r-\gamma)^{\prime} \Psi_{N} + o_{p}(\|r-\gamma\|^{2}) + O_{p}(\varepsilon_{N}), \tag{B.5}$$

which is the task of Lemmas B.4–B.7. The $O_p(\varepsilon_N)$ term in equation (B.5) can be shown to be of order $o_p(N^{-1})$ uniformly over \mathcal{R}_N . Further, by Theorem 1 of Sherman (1994b), equation (B.5) implies $\hat{\gamma} - \gamma = O_p(\sqrt{\varepsilon_N} \vee N^{-1/2}) = O_p(N^{-1/2})$. This rate result, together

with equation (B.5), further verifies equation (B.4).

To obtain equation (B.5), we will work with the following expansion

$$\hat{\mathcal{L}}_{N}^{K}(r) = \mathcal{L}_{N}^{K}(r) + \Delta \mathcal{L}_{N}^{K}(r) + R_{N}$$

$$= \mathbb{E}[\mathcal{L}_{N}^{K}(r)] + \frac{2}{N} \sum_{i} \left\{ \mathbb{E}[\mathcal{L}_{N}^{K}(r)|Z_{i}] - \mathbb{E}[\mathcal{L}_{N}^{K}(r)] \right\} + \rho_{N}(r) + \Delta \mathcal{L}_{N}^{K}(r) + R_{N}, \quad (B.6)$$

where

$$\Delta \mathcal{L}_{N}^{K}(r) \equiv \frac{1}{\sigma_{N}^{3} N(N-1)} \sum_{i \neq m} \nabla K_{\sigma_{N},\gamma}(V_{im}(\beta))'(V_{im}(\hat{\beta}) - V_{im}(\beta)) h_{im}(r), \qquad (B.7)$$

$$\rho_{N}(r) = \mathcal{L}_{N}^{K}(r) - \frac{2}{N} \sum_{i} \mathbb{E}[\mathcal{L}_{N}^{K}(r)|Z_{i}] + \mathbb{E}[\mathcal{L}_{N}^{K}(r)],$$

and R_N denotes the remainder term in the expansion of higher order (as $\sqrt{N}\sigma_N \to \infty$). The first three terms in (B.6) are the U-statistic decomposition for $\mathcal{L}_N^K(r)$ (see e.g., Sherman (1993) and Serfling (2009)). Lemmas B.4–B.6 establish asymptotic properties of these three terms, respectively. A linear representation for the fourth term in (B.6) is derived in Lemma B.7.

Here we present Lemmas B.1–B.7, whose proofs are relegated to Appendix F.

Lemma B.1. Under Assumptions C1–C7, $\sup_{r \in \mathcal{R}} |\hat{\mathcal{L}}_N^K(r) - \mathcal{L}_N^K(r)| = o_p(1)$.

Lemma B.2. Under Assumptions C1 and C5-C7, $\sup_{r \in \mathcal{R}} |\mathcal{L}_N^K(r) - \mathcal{L}(r)| = o_p(1)$.

Lemma B.3. Under Assumptions C1–C7,, $\hat{\gamma} \stackrel{p}{\rightarrow} \gamma$.

Lemma B.4. Under Assumptions C1–C7,, uniformly over \mathcal{R}_N , we have

$$\mathbb{E}[\mathcal{L}_N^K(r)] = \frac{1}{2}(r - \gamma)' \mathbb{V}_{\gamma}(r - \gamma) + o(\|r - \gamma\|^2),$$

where $\mathbb{V}_{\gamma} \equiv \mathbb{E}[\nabla^2 \tau_i(\gamma)]$.

Lemma B.5. Under Assumptions C1–C7,, uniformly over \mathcal{R}_N , we have

$$\frac{2}{N} \sum_{m} \mathbb{E}[\mathcal{L}_{N}^{K}(r)|Z_{m}] - 2\mathbb{E}[\mathcal{L}_{N}^{K}(r)] = \frac{1}{\sqrt{N}} (r - \gamma)' \Psi_{N,1} + o_{p}(\|r - \gamma\|^{2}),$$

where $\Psi_{N,1} \equiv N^{-1/2} \sum_{m} 2\nabla \tau_m(\gamma)$.

Lemma B.6. Under Assumptions C1 and C6–C7, uniformly over \mathcal{R}_N , $\rho_N(r) = O_p(N^{-1}\sigma_N^{-2})$.

Lemma B.7. Suppose

$$\hat{\beta} - \beta = \frac{1}{N} \sum_{i} \psi_{i,\beta} + o_p(N^{-1/2}),$$
 (B.8)

where $\psi_{i,\beta}$ is the influence function (of Z_i). $\psi_{i,\beta}$ is i.i.d across i with $\mathbb{E}(\psi_{i,\beta}) = 0.17$ Further, suppose Assumptions C1–C7 hold. Then uniformly over \mathcal{R}_N , we have

$$\Delta \mathcal{L}_{N}^{K}(r) = \frac{1}{\sqrt{N}} (r - \gamma)' \Psi_{N,2} + o_{p}(\|r - \gamma\|^{2}) + O_{p}(N^{-1}\sigma_{N}^{-2}),$$

with $\Psi_{N,2} = \frac{1}{\sqrt{N}} \sum_i \left(-\int \nabla_{13}^2 \mu(v_m, v_m, \gamma) f_V(v_m) dv_m \right) \psi_{i,\beta}$, where $\mu(v_i, v_m, r) \equiv G(v_i, v_m, r) f_V(v_i)$,

$$G(v_i, v_m, r) \equiv \mathbb{E}\left[B(V_i, V_m, W_i, W_m)S_{im}(r) \begin{pmatrix} x'_{im1} \\ x'_{im2} \end{pmatrix} | V_i = v_i, V_m = v_m \right],$$

 $\nabla_1 \mu(\cdot,\cdot,\cdot)$ denotes the partial derivative of $\mu(\cdot,\cdot,\cdot)$ w.r.t. its first argument, and $\nabla^2_{13}\mu(\cdot,\cdot,\cdot)$ denotes the partial derivative of $\nabla_1 \mu(\cdot,\cdot,\cdot)$ w.r.t. its third argument.

Proof of Theorem 2.2. Putting results in Lemmas B.4–B.7 together, we write equation (B.6) as

$$\hat{\mathcal{L}}_{N}^{K}(r) = \frac{1}{2}(r-\gamma)^{\prime} \mathbb{V}_{\gamma}(r-\gamma) + \frac{1}{\sqrt{N}}(r-\gamma)^{\prime} \Psi_{N} + o_{p}(\|r-\gamma\|^{2}) + O_{p}(\varepsilon_{N})$$
(B.9)

where $\Psi_N = \Psi_{N,1} + \Psi_{N,2}$ and $\varepsilon_N = N^{-1}\sigma_N^{-2}$. Theorem 1 of Sherman (1994b) then implies that $\hat{\gamma} - \gamma = O_p(\sqrt{\varepsilon_N}) = O_p(N^{-1/2}\sigma_N^{-1})$.

Next, take $\delta_N = O(\sqrt{\varepsilon_N})$ and $\mathcal{R}_N = \{r \in \mathcal{R} | ||r - \gamma|| \leq \delta_N \}$. We repeat the proof for Lemma B.6 and deduce from a Taylor expansion around γ that $\sup_{r \in \mathcal{R}_N} \mathbb{E}[\rho_{im}^*(r)^2] = O(\sigma_N^2 \delta_N^2)$. Apply Theorem 3 of Sherman (1994b) to see that uniformly over \mathcal{R}_N , $\rho_N(r) = O_p(N^{-1}\sigma_N^{\lambda-2}\delta_N^{\lambda})$ where $0 < \lambda < 1$. Then we have

$$\rho_N(r) = O_p(N^{-1}\sigma_N^{\lambda-2}\delta_N^{\lambda}) = O_p(N^{-1})O_p(N^{-\lambda/2}\sigma_N^{-2}) = o_p(N^{-1})$$

by invoking Assumption C7 and choosing λ sufficiently close to 1. This result in turn implies that the $O_p(\varepsilon_N)$ term in (B.9) has order $o_p(N^{-1})$, and hence $\hat{\gamma} - \gamma = O_p(N^{-1/2})$ by applying Theorem 1 of Sherman (1994b) once again.

¹⁷In fact, the linear representation of $\hat{\beta} - \beta$ in (B.8) can be obtained by applying the same arguments in Theorem 2 of Sherman (1993) to a representation for $\mathcal{L}_{N,\beta}^{K}(b)$ analogous to (B.4).

Now (B.9) can be expressed as

$$\hat{\mathcal{L}}_N^K(r) = \frac{1}{2}(r-\gamma)' \mathbb{V}_{\gamma}(r-\gamma) + \frac{1}{\sqrt{N}}(r-\gamma)' \Psi_N + o_p(N^{-1}).$$

Let $\Delta_i \equiv 2\nabla \tau_i(\gamma) - \left(\int \nabla_{13}^2 \mu(v_m, v_m, \gamma) f_V(v_m) dv_m\right) \psi_{i,\beta}$. Note that $\mathbb{E}[\Delta_i] = 0$ since $\mathbb{E}[\nabla \tau_i(\gamma)] = 0$ and $\mathbb{E}[\psi_{i,\beta}] = 0$. We deduce from Assumption C7 and Lindeberg-Lévy CLT that $\Psi_N \stackrel{d}{\to} N(0, \mathbb{V}_{\Psi})$ where $\mathbb{V}_{\Psi} = \mathbb{E}[\Delta_i \Delta_i']$. The asymptotic normality of $\hat{\gamma}$ then follows from Theorem 2 of Sherman (1994a), i.e., $\sqrt{N}(\hat{\gamma} - \gamma) \stackrel{d}{\to} N(0, \mathbb{V}_{\gamma}^{-1} \mathbb{V}_{\Psi} \mathbb{V}_{\gamma}^{-1})$.

B.3 Proof of Theorem 2.4

Recall that $\hat{\mathcal{L}}_{N}^{K}(r)$ was redefined in Appendix B.2. However, for the proof of Theorem 2.4, it is more convenient to still use the definition in the main body of the paper, specifically,

$$\hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) = \frac{1}{\sigma_{N}^{2} N\left(N-1\right)} \sum_{i \neq m} K_{\sigma_{N},\gamma}(V_{im}(\hat{\beta})) Y_{im(1,1)} \operatorname{sgn}\left(W_{im}'\hat{\gamma}\right).$$

Lemma B.8. Suppose Assumptions C1–C7 hold. Then, we have

$$\frac{1}{\sigma_N^2 N(N-1)} \sum_{i \neq m} [K_{\sigma_N,\gamma}(V_{im}(\hat{\beta})) - K_{\sigma_N,\gamma}(V_{im}(\beta))] Y_{im(1,1)} \operatorname{sgn}(W'_{im}\gamma) = O_p(N^{-1}\sigma_N^{-2}).$$

Lemma B.9. Suppose Assumptions C1–C7 hold. Then uniformly over $r \in \mathcal{R}$,

$$\hat{\mathcal{L}}_{N}^{K}(r) = \frac{1}{2} (r - \gamma)' \mathbb{V}_{\gamma} (r - \gamma) + o_{p} \left(N^{-1/2} \| r - \gamma \| \right) + o_{p} \left(\| r - \gamma \|^{2} \right) + \frac{1}{\sigma_{N}^{2} N (N - 1)} \sum_{i \neq m} K_{\sigma_{N}, \gamma} \left(V_{im} (\beta) \right) Y_{im(1, 1)} sgn \left(W'_{im} \gamma \right) + o_{p} (N^{-1/2}).$$

Lemma B.10. Suppose Assumptions C5–C7 hold. Then, we have

$$\mathbb{E}\left[\sigma_{N}^{-2}K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right)Y_{im\left(1,1\right)}sgn\left(W_{im}'\gamma\right)|Z_{i}\right]$$

$$= f_{V_{im}\left(\beta\right)}(0)\mathbb{E}\left[Y_{im\left(1,1\right)}sgn\left(W_{im}'\gamma\right)|Z_{i},V_{im}\left(\beta\right)=0\right] + o_{p}(N^{-1/2}).$$

Note that Lemma B.8 indicates that $\hat{\beta}$ affects $\hat{\mathcal{L}}_N^K(\hat{\gamma})$ at the rate of $N^{-1}\sigma_N^{-2}$, and Lemma B.9 shows that $\hat{\gamma}$ affects $\hat{\mathcal{L}}_N^K(\hat{\gamma})$ at the rate of N^{-1} .

Proof of Theorem 2.4. Note that $O_p(N^{-1}\sigma_N^{-2}) = o_p(N^{-1/2})$ by Assumption C7 and $\hat{\gamma} - \gamma =$

 $O_p(N^{-1/2})$ by Theorem 2.2. These results, together with Lemmas B.8 and B.9, imply that

$$\hat{\mathcal{L}}_{N}^{K}\left(\hat{\gamma}\right) = \frac{1}{\sigma_{N}^{2}N\left(N-1\right)} \sum_{i \neq m} K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right) Y_{im(1,1)} \operatorname{sgn}\left(W_{im}^{\prime}\gamma\right) + o_{p}(N^{-1/2}).$$

We can use the same arguments as those used for proving Lemmas B.1 and B.2 to show

$$\hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) \stackrel{p}{\to} \bar{\mathcal{L}}(\gamma) = f_{V_{im}(\beta)}(0) \mathbb{E}\left[Y_{im(1,1)}\operatorname{sgn}(W'_{im}\gamma) | V_{im}(\beta) = 0\right].$$

Then, we can write

$$\begin{split} & \hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) - \bar{\mathcal{L}}\left(\gamma\right) \\ & = \frac{1}{\sigma_{N}^{2}N\left(N-1\right)} \sum_{i \neq m} K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right) Y_{im(1,1)} \mathrm{sgn}\left(W_{im}'\gamma\right) - \bar{\mathcal{L}}\left(\gamma\right) + o_{p}(N^{-1/2}) \\ & = \frac{2}{N} \sum_{i=1}^{N} \left\{ \mathbb{E}\left[\sigma_{N}^{-2}K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right) Y_{im(1,1)} \mathrm{sgn}\left(W_{im}'\gamma\right) | Z_{i}\right] - \mathbb{E}\left[\sigma_{N}^{-2}K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right) Y_{im(1,1)} \mathrm{sgn}\left(W_{im}'\gamma\right)\right] \right\} \\ & + \mathbb{E}\left[\sigma_{N}^{-2}K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right) Y_{im(1,1)} \mathrm{sgn}\left(W_{im}'\gamma\right)\right] - \bar{\mathcal{L}}\left(\gamma\right) + o_{p}(N^{-1/2}) \\ & = \frac{2}{N} \sum_{i=1}^{N} f_{V_{im}(\beta)}(0) \left\{ \mathbb{E}\left[Y_{im(1,1)} \mathrm{sgn}\left(W_{im}'\gamma\right) | Z_{i}, V_{im}\left(\beta\right) = 0\right] - \mathbb{E}\left[Y_{im(1,1)} \mathrm{sgn}\left(W_{im}'\gamma\right) | V_{im}\left(\beta\right) = 0\right] \right\} \\ & + o_{p}(N^{-1/2}), \end{split}$$

where the second equality uses the decomposition (B.6) and Lemma B.6, and the last equality follows by Lemma B.10. Invoking the Lindeberg-Lévy CLT on the leading term of $\hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) - \bar{\mathcal{L}}(\gamma)$ in the above yields

$$\sqrt{N}(\hat{\mathcal{L}}_{N}^{K}(\hat{\gamma}) - \bar{\mathcal{L}}(\gamma)) \stackrel{d}{\to} N(0, \Delta_{\gamma}).$$

B.4 Proof of Theorem 2.5

Proof of Theorem 2.5. The proof is lengthy, but the underlying idea is straightforward. Using the (conditional) linear independence (equivalent to the regular full rank condition) and the free variation of the first regressor conditions in Assumption CL2, we show that there exists a positive probability of making an incorrect prediction for any $\vartheta \neq \theta$.

Some Definitions. We normalize the coefficients on $X_{im1}^{(1)}$, $X_{im2}^{(1)}$, and $W_i^{(1)}$ to 1 as indicated

in Assumption CL3. To simplify notation, we express the three utility indexes with a slight abuse of notation as $X_{im1}^{(1)} + u_{im1}(\vartheta)$, $X_{im2}^{(1)} + u_{im2}(\vartheta)$, and $W_{im}^{(1)} + u_{imb}(\vartheta)$, where $u_{im1}(\vartheta) \equiv \tilde{X}'_{im1}b + S'_{im}\varrho_1$, $u_{im2}(\vartheta) \equiv \tilde{X}'_{im2}b + S'_{im}\varrho_2$, and $u_{imb}(\vartheta) \equiv \tilde{W}'_{im}r + S'_{im}\varrho_b$, respectively. We use $a \vee b$ and $a \wedge b$ to denote $\max\{a,b\}$ and $\min\{a,b\}$, respectively. Finally, we define indicator functions for the events on the left-hand sides of (2.18)-(2.19), (2.20)-(2.21), and $(2.22)-(2.23)^{18}$ and the two uninformative events in Remark 2.2 in turn:

$$\begin{split} I^{+}_{im(0,1)}(\vartheta) &= 1[X^{(1)}_{im1} + u_{im1}(\vartheta) \leq 0, X^{(1)}_{im2} + u_{im2}(\vartheta) \geq 0, W^{(1)}_{im} + u_{imb}(\vartheta) \leq 0], \\ I^{-}_{im(0,1)}(\vartheta) &= 1[X^{(1)}_{im1} + u_{im1}(\vartheta) \geq 0, X^{(1)}_{im2} + u_{im2}(\vartheta) \leq 0, W^{(1)}_{im} + u_{imb}(\vartheta) \geq 0], \\ I^{+}_{im(1,1)}(\vartheta) &= 1[X^{(1)}_{im1} + u_{im1}(\vartheta) \geq 0, X^{(1)}_{im2} + u_{im2}(\vartheta) \geq 0, W^{(1)}_{im} + u_{imb}(\vartheta) \geq 0], \\ I^{-}_{im(1,1)}(\vartheta) &= 1[X^{(1)}_{im1} + u_{im1}(\vartheta) \leq 0, X^{(1)}_{im2} + u_{im2}(\vartheta) \leq 0, W^{(1)}_{im} + u_{imb}(\vartheta) \leq 0], \\ I^{+}_{imu}(\vartheta) &= 1[X^{(1)}_{im1} + u_{im1}(\vartheta) \leq 0, X^{(1)}_{im2} + u_{im2}(\vartheta) \leq 0, W^{(1)}_{im} + u_{imb}(\vartheta) \geq 0], \\ I^{-}_{imu}(\vartheta) &= 1[X^{(1)}_{im1} + u_{im1}(\vartheta) \geq 0, X^{(1)}_{im2} + u_{im2}(\vartheta) \geq 0, W^{(1)}_{im} + u_{imb}(\vartheta) \leq 0]. \end{split}$$

Recall that by construction $q_{(1,0)}(Z_i, Z_m, \theta) = 0 \le q_{(1,0)}(Z_i, Z_m, \vartheta)$ for all $\vartheta \in \Theta$ and (Z_i, Z_m) . In what follows, we show for any $\vartheta \in \Theta \setminus \{\theta\}$, $q_{(1,0)}(Z_i, Z_m, \vartheta) > 0$ occurs with positive probability, and hence $Q_{(1,0)}(\theta) < Q_{(1,0)}(\vartheta)$.

Identification of (β, ρ_1) . For any $\vartheta \in \Theta \setminus \{\theta\}$, we have

$$P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0)$$

$$\geq P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{im(1,1)}^{+}(\theta) + I_{im(1,1)}^{-}(\theta) = 1\})$$

$$= P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{im(1,1)}^{+}(\theta) = 1\}) + P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{im(1,1)}^{-}(\theta) = 1\})$$

$$\geq P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{im(1,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1\})$$

$$+ P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{im(1,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1\})$$

$$= P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0 | I_{im(1,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1) P(I_{im(1,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1)$$

$$+ P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0 | I_{im(1,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1) P(I_{im(1,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1).$$
(B.10)

There is no need to define $I^+_{im(0,0)}(\vartheta)$ and $I^-_{im(0,0)}(\vartheta)$ separately since $I^+_{im(0,0)}(\vartheta) = I^-_{im(1,1)}(\vartheta)$ and $I^-_{im(0,0)}(\vartheta) = I^+_{im(1,1)}(\vartheta)$.

By definitions, we can write

$$\begin{split} &P(I_{im(1,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1) \\ = &P(u_{im1}(\vartheta) \leq -X_{im1}^{(1)} \leq u_{im1}(\theta), -X_{im2}^{(1)} \leq u_{im2}(\vartheta) \wedge u_{im2}(\theta), -W_{im}^{(1)} \leq u_{imb}(\vartheta) \wedge u_{imb}(\theta)), \end{split}$$

and

$$\begin{split} &P(I_{im(1,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1) \\ = &P(u_{im1}(\theta) \leq -X_{im1}^{(1)} \leq u_{im1}(\vartheta), -X_{im2}^{(1)} \geq u_{im2}(\vartheta) \vee u_{im2}(\theta), -W_{im}^{(1)} \geq u_{imb}(\vartheta) \vee u_{imb}(\theta)). \end{split}$$

Assumptions CL2(i) and CL2(iii) imply that events $\{-X_{im2}^{(1)} \leq u_{im2}(\vartheta) \wedge u_{im2}(\theta), -W_{im}^{(1)} \leq u_{imb}(\vartheta) \wedge u_{imb}(\theta)\}$ and $\{-X_{im2}^{(1)} \geq u_{im2}(\vartheta) \vee u_{im2}(\theta), -W_{im}^{(1)} \geq u_{imb}(\vartheta) \vee u_{imb}(\theta)\}$ always have positive probabilities of occurring. Then, when conditioning on them, Assumption CL2(ii) implies that the event $u_{im1}(\vartheta) \neq u_{im1}(\theta)$ has a positive probability to occur when $(b, \varrho_1) \neq (\beta, \rho_1)$. Thus, under Assumption CL2(i), we conclude that either $u_{im1}(\vartheta) \leq -X_{im1}^{(1)} \leq u_{im1}(\theta)$ or $u_{im1}(\theta) \leq -X_{im1}^{(1)} \leq u_{im1}(\vartheta)$ occurs with positive probability. In summary, we can claim that for any ϑ with $(b, \varrho_1) \neq (\beta, \rho_1)$,

$$P(I_{im(1,1)}^+(\theta) = I_{im(1,0)}^-(\vartheta) = 1) > 0 \text{ or } P(I_{im(1,1)}^-(\theta) = I_{im(1,0)}^+(\vartheta) = 1) > 0.$$
 (B.11)

Then, if we can establish that

$$P(q_{(1,0)}(Z_i, Z_m, \vartheta) > 0 | I_{im(1,1)}^+(\theta) = I_{im(1,0)}^-(\vartheta) = 1) > 0 \text{ and}$$

$$P(q_{(1,0)}(Z_i, Z_m, \vartheta) > 0 | I_{im(1,1)}^-(\theta) = I_{im(1,0)}^+(\vartheta) = 1) > 0,$$
(B.12)

by substituting (B.11) into (B.10), we obtain $P(q_{(1,0)}(Z_i, Z_m, \vartheta) > 0) > 0$ for any $(b, \varrho_1) \neq (\beta, \rho_1)$. This establishes the identification of (β, ρ_1) in θ . We defer the justification of the inequalities in (B.12) to the end of this proof.

Identification of (β, ρ_2) . Similarly, we can write

$$P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0)$$

$$\geq P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{imu}^{+}(\theta) + I_{imu}^{-}(\theta) = 1\})$$

$$\geq P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0 | I_{imu}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1) P(I_{imu}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1)$$

$$+ P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0 | I_{imu}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1) P(I_{imu}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1), \quad (B.13)$$

where

$$\begin{split} &P(I_{imu}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1) \\ = &P(-X_{im1}^{(1)} \geq u_{im1}(\vartheta) \vee u_{im1}(\theta), u_{im2}(\theta) \leq -X_{im2}^{(1)} \leq u_{im2}(\vartheta), -W_{im}^{(1)} \leq u_{imb}(\vartheta) \wedge u_{imb}(\theta)), \end{split}$$

and

$$\begin{split} &P(I_{imu}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1) \\ =& P(-X_{im1}^{(1)} \leq u_{im1}(\vartheta) \wedge u_{im1}(\theta), u_{im2}(\vartheta) \leq -X_{im2}^{(1)} \leq u_{im2}(\theta), -W_{im}^{(1)} \geq u_{imb}(\vartheta) \vee u_{imb}(\theta)). \end{split}$$

Then, we can establish the identification of (β, ρ_2) in θ using similar arguments as above.

Identification of (γ, ρ_b) . The identification of (γ, ρ_b) can be similarly established based on the following inequality:

$$P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0)$$

$$\geq P(\{q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0\} \cap \{I_{im(0,1)}^{+}(\theta) + I_{im(0,1)}^{-}(\theta) = 1\})$$

$$\geq P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0 | I_{im(0,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1) P(I_{im(0,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1)$$

$$+ P(q_{(1,0)}(Z_{i}, Z_{m}, \vartheta) > 0 | I_{im(0,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1) P(I_{im(0,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1),$$
(B.14)

where

$$\begin{split} &P(I_{im(0,1)}^{+}(\theta) = I_{im(1,0)}^{-}(\vartheta) = 1) \\ = &P(-X_{im1}^{(1)} \leq u_{im1}(\vartheta) \wedge u_{im1}(\theta), -X_{im2}^{(1)} \geq u_{im2}(\vartheta) \vee u_{im2}(\theta), u_{imb}(\vartheta) \leq -W_{im}^{(1)} \leq u_{imb}(\theta)), \end{split}$$

and

$$\begin{split} &P(I_{im(0,1)}^{-}(\theta) = I_{im(1,0)}^{+}(\vartheta) = 1) \\ = &P(-X_{im1}^{(1)} \geq u_{im1}(\vartheta) \vee u_{im1}(\theta), -X_{im2}^{(1)} \leq u_{im2}(\vartheta) \wedge u_{im2}(\theta), u_{imb}(\theta) \leq -W_{im}^{(1)} \leq u_{imb}(\vartheta)). \end{split}$$

Finally, put all these results together to conclude that all elements in θ are identified.

On (B.12). To complete the proof, we need to justify the claim in (B.12) as well as similar claims for (B.13) and (B.14). Since these proofs follow similar arguments, we only present the proof for $P(q_{(1,0)}(Z_i, Z_m, \vartheta) > 0 | I_{im(1,1)}^+(\theta) = I_{im(1,0)}^-(\vartheta) = 1) > 0$ and omit the others for the sake of brevity. Note that by definition, when $I_{im(1,0)}^-(\vartheta) = 1$, $q_{(1,0)}(Z_i, Z_m, \vartheta) > 0$

if and only if $\Delta p_{(1,0)}(Z_i, Z_m) > 0$. Therefore, we show $P(\Delta p_{(1,0)}(Z_i, Z_m) > 0 | I_{im(1,1)}^+(\theta) = I_{im(1,0)}^-(\theta) = 1) > 0$ in the following, and the proof is done.

Under Assumption CL1(i), Z_{im} has distribution symmetric at 0, and hence we can pick $X_{im2}^{(1)} = W_{im}^{(1)} = 0$, $\tilde{X}_{im2} = 0$, $\tilde{W}_{im} = 0$, and $S_{im} = 0$ so that $\{-X_{im2}^{(1)} \leq u_{im2}(\vartheta) \land u_{im2}(\theta), -W_{im}^{(1)} \leq u_{imb}(\vartheta) \land u_{imb}(\theta)\}$ holds with equalities for any ϑ . Note that here we use the "matching" insight for developing the MRC procedure, that is, we match the utility indexes associated with the second stand-alone alternative and the bundle effect. Meanwhile, under Assumptions CL2(i) and CL2(ii), we can still have $\{u_{im1}(\vartheta) < -X_{im1}^{(1)} < u_{im1}(\theta)\}$ occur with positive probability conditional on $E \equiv \{X_{im2}^{(1)} = W_{im}^{(1)} = 0, \tilde{X}_{im2} = 0, \tilde{W}_{im} = 0, S_{im} = 0\}$. In this scenario, inequality (2.11) implies that $\Delta p_{(1,0)}(Z_i, Z_m) > 0$, and say $\Delta p_{(1,0)}(Z_i, Z_m) = \delta_1 > 0$ conditional on E. Due to the continuity, there exists a small $\delta_2 > 0$ such that conditional on

$$E^{\delta_2} = \{ |X_{im2}^{(1)}| + |W_{im}^{(1)}| + ||\tilde{X}_{im2}|| + ||\tilde{W}_{im}|| + ||S_{im}|| \le \delta_2 \},$$

a small neighborhood of E,

$$\Delta p_{(1,0)}(Z_i, Z_m) \ge \delta_1/2 > 0.$$

Again, due to the continuity, $P\left(E^{\delta_2}\right) > 0$. Since what we have discussed so far is only one possible scenario for $\Delta p_{(1,0)}(Z_i, Z_m) > 0$ to occur, we must have $P(q_{(1,0)}(Z_i, Z_m, \vartheta) > 0 | I_{im(1,1)}^+(\theta) = I_{im(1,0)}^-(\vartheta) = 1) > P\left(E^{\delta_2}\right) > 0$. This completes the proof.

B.5 Proof of Theorem 2.6

Proof of Theorem 2.6. Let

$$\hat{Q}_{N(1,0)}^{D}(\vartheta) := \frac{2}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} \hat{q}_{im(1,0)}^{D}(\vartheta),$$

where $\hat{q}_{im(1,0)}^D(\vartheta)$ is defined in (2.17). We prove the consistency of $\hat{\theta}$ by verifying the four sufficient conditions in Theorem 2.1 of Newey and McFadden (1994): (S1) Θ is a compact set, (S2) $\sup_{\vartheta \in \Theta} |\hat{Q}_{N(1,0)}^D(\vartheta) - Q_{(1,0)}^D(\vartheta)| = o_p(1)$, (S3) $Q_{(1,0)}^D(\vartheta)$ is continuous in ϑ , and (S4) $Q_{(1,0)}^D(\vartheta)$ is uniquely minimized at θ .

Condition (S1) is trivially satisfied under Assumption CL3. The identification condition (S4) is an immediate result of Theorem 2.5 and the discussion right after it. The continuity

condition (S3) is also satisfied since $Q_{(1,0)}^D(\vartheta)$, by definition, can be expressed as the sum of terms of the following type:

$$\int_{\Omega_{\theta}} (|1 - \Delta p_{(1,0)}(z_i, z_m)| - 1) f_Z(z_i) f_Z(z_m) dz_i dz_m,$$

where $\Omega_{\vartheta} := \{x'_{im1}b + s'_{im}\varrho_1 \geq 0, x'_{im2}b + s'_{im}\varrho_2 \leq 0, w'_{im}r + s'_{im}\varrho_b \leq 0\}$. Then it is easy to see that Assumptions CL1 and CL2 imply the continuity of $Q^D_{(1,0)}(\vartheta)$.

The remaining task is to show the uniform convergence condition (S2). To this end, we prove $\sup_{\vartheta \in \Theta} |\hat{Q}_{N(1,0)}^D(\vartheta) - Q_{N(1,0)}^D(\vartheta)| = o_p(1)$ and $\sup_{\vartheta \in \Theta} |Q_{N(1,0)}^D(\vartheta) - Q_{(1,0)}^D(\vartheta)| = o_p(1)$, where $Q_{N(1,0)}^D(\vartheta) := \frac{2}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} q_{(1,0)}^D(Z_i, Z_m, \vartheta)$.

Firstly, we have by definition

$$\sup_{\vartheta \in \Theta} |\hat{Q}_{N(1,0)}^{D}(\vartheta) - Q_{N(1,0)}^{D}(\vartheta)| \leq \frac{2}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} \sup_{\vartheta \in \Theta} |\hat{q}_{im(1,0)}^{D}(\vartheta) - q_{(1,0)}^{D}(Z_{i}, Z_{m}, \vartheta)|
\leq \frac{4}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} |\Delta \hat{p}_{(1,0)}(Z_{i}, Z_{m}) - \Delta p_{(1,0)}(Z_{i}, Z_{m})|
\leq \sup_{(z_{i}, z_{m}) \in \mathcal{Z}^{2}} 4|\Delta \hat{p}_{(1,0)}(z_{i}, z_{m}) - \Delta p_{(1,0)}(z_{i}, z_{m})| = o_{p}(1),$$

where the last equality follows by Assumption CL5.

Secondly, since $|\Delta p_{(1,0)}(Z_i, Z_m)| \leq 1$, we can write

$$\sup_{\vartheta \in \Theta} |Q_{N(1,0)}^{D}(\vartheta) - Q_{(1,0)}^{D}(\vartheta)|
\leq \sup_{\vartheta \in \Theta} |\frac{2}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} q_{(1,0)}^{D}(Z_{i}, Z_{m}, \vartheta) - \mathbb{E}[q_{(1,0)}^{D}(Z_{i}, Z_{m}, \vartheta)]|
\leq \sup_{\bar{\vartheta} \in \bar{\Theta}} |\frac{2}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} \psi_{(1,0)}^{D}(Z_{i}, Z_{m}, \bar{\vartheta}) - \mathbb{E}[\psi_{(1,0)}^{D}(Z_{i}, Z_{m}, \bar{\vartheta})]|,$$
(B.15)

where $\bar{\Theta} := {\bar{\vartheta} = (\vartheta, a) | \vartheta \in \Theta, a \in [-1, 1]}$ and

$$\psi_{(1,0)}^{D}(Z_{i}, Z_{m}, \bar{\vartheta}) := [|I_{im(1,0)}^{+}(\vartheta) - a| + |I_{im(1,0)}^{-}(\vartheta) + a|] \times [I_{im(1,0)}^{+}(\vartheta) + I_{im(1,0)}^{-}(\vartheta)] + [1 - (I_{im(1,0)}^{+}(\vartheta) + I_{im(1,0)}^{-}(\vartheta))].$$

Let $\mathcal{G} = \{\psi^D_{(1,0)}(\cdot,\cdot,\bar{\vartheta})|\bar{\vartheta} = (\vartheta,a), \vartheta \in \Theta, a \in [-1,1]\}$ define the class of bounded functions

 $\psi_{(1,0)}^D(\cdot,\cdot,\bar{\vartheta})$ indexed by $\bar{\vartheta}$. Using similar arguments in the proof of Lemma B.2, we can show that \mathcal{G} is a Euclidean class¹⁹ of functions. Hence, we have

$$\sup_{\bar{\vartheta}\in\bar{\Theta}} \left| \frac{2}{N(N-1)} \sum_{i=1}^{N-1} \sum_{m>i} \psi_{(1,0)}^{D}(Z_i, Z_m, \bar{\vartheta}) - \mathbb{E}[\psi_{(1,0)}^{D}(Z_i, Z_m, \bar{\vartheta})] \right| = o_p(1). \tag{B.16}$$

Plugging (B.16) into (B.15) gives $\sup_{\vartheta \in \Theta} |Q_{N(1,0)}^D(\vartheta) - Q_{(1,0)}^D(\vartheta)| = o_p(1)$. Then (S2) follows immediately from the triangle inequality, which completes the proof.

C Proofs for the Panel Data Model

C.1 Proof of Theorem 3.2

and

In this section, we provide the proof of Theorem 3.2, applying the asymptotic theory developed in Seo and Otsu (2018). Before the main proof, we first present two supporting lemmas, Lemmas C.1 and C.2, whose proofs are relegated to Appendix F. The outline of the proof process is as follows. Lemma C.1 verifies the technical conditions in Seo and Otsu (2018). Lemma C.2 obtains technical terms for the final asymptotics. Then we apply the results in Seo and Otsu (2018) and get the asymptotics of our estimators in the proof of Theorem 3.2.

Lemma C.1. Suppose Assumptions P1–P9 hold. Then $\phi_{Ni}(b)$ and $\varphi_{Ni}(r)$ satisfy Assumption M in Seo and Otsu (2018).

Lemma C.2. Suppose Assumptions P1-P9 hold. Then

$$\lim_{N \to \infty} (Nh_N^{k_1 + k_2})^{2/3} \mathbb{E}[\phi_{Ni}(\beta + \rho (Nh_N^{k_1 + k_2})^{-1/3})] = \frac{1}{2} \rho' \mathbb{V}\rho,$$

$$\lim_{N \to \infty} (N\sigma_N^{2k_1})^{2/3} \mathbb{E}[\varphi_{Ni}(\gamma + \delta(N\sigma_N^{2k_1})^{-1/3})] = \frac{1}{2} \delta' \mathbb{W} \delta,$$

 $\lim_{N\to\infty} (Nh_N^{k_1+k_2})^{1/3} \mathbb{E}[h_N^{k_1+k_2} \phi_{Ni}(\beta + \rho_1(Nh_N^{k_1+k_2})^{-1/3}) \phi_{Ni}(\beta + \rho_2(Nh_N^{k_1+k_2})^{-1/3})] = \mathbb{H}_1(\rho_1, \rho_2),$

$$\lim_{N \to \infty} (N\sigma_N^{2k_1})^{1/3} \mathbb{E}[\sigma_N^{2k_1} \varphi_{Ni} (\gamma + \delta_1 (N\sigma_N^{2k_1})^{-1/3}) \varphi_{Ni} (\gamma + \delta_2 (N\sigma_N^{2k_1})^{-1/3})] = \mathbb{H}_2 (\delta_1, \delta_2),$$

 $^{^{19}}$ Alternatively, one can also apply results in Section 2.6 of van der Vaart and Wellner (1996) to show that \mathcal{G} is a VC-class.

where ρ , ρ_1 , and ρ_2 are $k_1 \times 1$ vectors, δ , δ_1 , and δ_2 are $k_2 \times 1$ vectors, \mathbb{V} is a $k_1 \times k_1$ matrix defined as

$$\mathbb{V} = -\sum_{t>s} \sum_{d \in \mathcal{D}} \left\{ \int 1 \left[x'\beta = 0 \right] \left(\frac{\partial \kappa_{dts}^{(1)}(x)}{\partial x}'\beta \right) f_{X_{1ts} | \{X_{2ts} = 0, W_{ts} = 0\}}(x) x x' d\sigma_0^{(1)} \right. \\
+ \int 1 \left[x'\beta = 0 \right] \left(\frac{\partial \kappa_{dts}^{(2)}(x)}{\partial x}'\beta \right) f_{X_{2ts} | \{X_{1ts} = 0, W_{ts} = 0\}}(x) x x' d\sigma_0^{(2)} \right\},$$
(C.1)

W is a $k_2 \times k_2$ matrix defined as

$$W = -\sum_{t>s} \int 1 \left[w' \gamma = 0 \right] \left(\frac{\partial \kappa_{(1,1)ts}^{(3)}(w)}{\partial w} \gamma \right) f_{W_{ts} \mid \{X_{1ts} = 0, X_{2ts} = 0\}}(w) w w' d\sigma_0^{(3)}, \qquad (C.2)$$

and \mathbb{H}_1 and \mathbb{H}_2 are written respectively as

$$\mathbb{H}_{1}(\rho_{1}, \rho_{2}) = \frac{1}{2} \sum_{t>s} \sum_{d \in \mathcal{D}} \left\{ \int \kappa_{dts}^{(4)}(\bar{x}) \left[|\bar{x}'\rho_{1}| + |\bar{x}'\rho_{2}| - |\bar{x}'(\rho_{1} - \rho_{2})| \right] f_{X_{1ts}|\{X_{2ts} = 0, W_{ts} = 0\}}(0, \bar{x}) \, d\bar{x} f_{X_{2ts}, W_{ts}}(0, 0) \right. \\
\left. + \int \kappa_{dts}^{(5)}(\bar{x}) \left[|\bar{x}'\rho_{1}| + |\bar{x}'\rho_{2}| - |\bar{x}'(\rho_{1} - \rho_{2})| \right] f_{X_{2ts}|\{X_{1ts} = 0, W_{ts} = 0\}}(0, \bar{x}) \, d\bar{x} f_{X_{1ts}, W_{ts}}(0, 0) \right\} \bar{K}_{2}^{k_{1} + k_{2}},$$

and

$$\mathbb{H}_{2}(\delta_{1}, \delta_{2}) = \frac{1}{2} \sum_{t>s} \int \kappa_{(1,1)ts}^{(6)}(\bar{w}) \left[|\bar{w}'\delta_{1}| + |\bar{w}'\delta_{2}| - |\bar{w}'(\delta_{1} - \delta_{2})| \right]$$

$$\cdot f_{W_{ts}|\{X_{1ts}=0, X_{2ts}=0\}}(0, \bar{w}) \, d\bar{w} f_{X_{1ts}, X_{2ts}}(0, 0) \, \bar{K}_{2}^{2k_{1}}.$$
(C.4)

Technical terms that appear in \mathbb{V} , \mathbb{W} , \mathbb{H}_1 , \mathbb{H}_2 are defined as follows. $\sigma_0^{(1)}$, $\sigma_0^{(2)}$, and $\sigma_0^{(3)}$ are the surface measures of $\{X_{1ts}\beta=0\}$, $\{X_{2ts}\beta=0\}$, and $\{W_{ts}\gamma=0\}$, respectively. $\bar{K}_2 \equiv \int K(u)^2 du$. We decompose X_{1ts} , X_{2ts} and W_{ts} into $X_{jts}=a\beta+\bar{X}_{jts}$, j=1,2, and $W_{ts}=a\gamma+\bar{W}_{ts}$, where \bar{X}_{jts} , j=1,2, are orthogonal to β , and \bar{W}_{ts} is orthogonal to γ . The density for X_{1ts} is written as $f_{X_{1ts}}(a,\bar{X}_{1ts})$ under this decomposition. Densities for X_{2ts} and W_{ts} are written similarly. Further,

$$\kappa_{dts}^{(1)}(x) \equiv \mathbb{E}[Y_{idst}(-1)^{d_1} | X_{i1ts} = x, X_{i2ts} = 0, W_{its} = 0],$$

$$\kappa_{dts}^{(2)}(x) \equiv \mathbb{E}[Y_{idst}(-1)^{d_2} | X_{i2ts} = x, X_{i1ts} = 0, W_{its} = 0],$$

$$\kappa_{(1,1)ts}^{(3)}(w) \equiv \mathbb{E}[Y_{i(1,1)ts} | W_{its} = w, X_{i1ts} = 0, X_{i2ts} = 0],$$

$$\kappa_{dts}^{(4)}\left(x\right) \equiv \mathbb{E}\left[\left|Y_{idst}\right| \; \left|X_{i1ts}=x,X_{i2ts}=0,W_{its}=0\right],$$

$$\kappa_{dts}^{(5)}(x) \equiv \mathbb{E}\left[|Y_{idst}| \mid X_{i2ts} = x, X_{i1ts} = 0, W_{its} = 0\right],$$

and

$$\kappa_{(1,1)ts}^{(6)}(w) \equiv \mathbb{E}\left[\left|Y_{i(1,1)ts}\right| \mid W_{its} = w, X_{i1ts} = 0, X_{i2ts} = 0\right].$$

Proof of Theorem 3.2. Lemma C.1 verifies the technical conditions for applying the results in Seo and Otsu (2018), and shows that $\phi_{Ni}(b)$ and $\varphi_{Ni}(r)$ are manageable in the sense of Kim and Pollard (1990). By Assumption P9, Lemma 1 in Seo and Otsu (2018) and its subsequent analysis, we have

$$\hat{\beta} - \beta = O_p((Nh_N^{k_1+k_2})^{-1/3})$$
 and $\hat{\gamma} - \gamma = O_p((N\sigma_N^{2k_1})^{-1/3}).$

Notice that $\hat{\beta}$ can be equivalently obtained from

$$\arg\max_{b\in\mathcal{B}}(Nh_N^{k_1+k_2})^{2/3}\cdot N^{-1}\sum_{i=1}^N\phi_{Ni}(\beta+(Nh_N^{k_1+k_2})^{-1/3}\{(Nh_N^{k_1+k_2})^{1/3}(b-\beta)\}).$$

We get the asymptotics of $\hat{\beta}$ if we can get the asymptotics of

$$\arg \max_{\rho \in \mathbb{R}^{k_1}} (Nh_N^{k_1+k_2})^{2/3} \cdot N^{-1} \sum_{i=1}^N \phi_{Ni} (\beta + (Nh_N^{k_1+k_2})^{-1/3} \rho).$$

Since $\phi_{Ni}(b)$ is manageable in the sense of Kim and Pollard (1990), by Theorem 1 in Seo and Otsu (2018) and Lemma C.2, we have the uniform convergence of the above stochastic process and

$$(Nh_N^{k_1+k_2})^{2/3} \cdot N^{-1} \sum_{i=1}^N \phi_{Ni} (\beta + (Nh_N^{k_1+k_2})^{-1/3} \rho) \rightsquigarrow \mathcal{Z}_1(\rho) ,$$

where $\mathcal{Z}_1(\rho)$ is a Gaussian process with continuous sample path, expected value $\frac{1}{2}\rho'\mathbb{V}\rho$ and covariance kernel $\mathbb{H}_1(\rho_1, \rho_2)$. Apply the CMT to obtain

$$(Nh_N^{k_1+k_2})^{1/3}(\hat{\beta}-\beta) \xrightarrow{d} \arg\max_{\rho} \mathcal{Z}_1(\rho)$$
.

Apply similar arguments to $\hat{\gamma}$. By Theorem 1 in Seo and Otsu (2018) and Lemma C.2, we get

$$(N\sigma_N^{2k_1})^{1/3} (\hat{\gamma} - \gamma) \stackrel{d}{\to} \arg\max_{\delta \in \mathbb{R}^{k_2}} \mathcal{Z}_2(\delta),$$

where $\mathcal{Z}_2(\delta)$ is a Gaussian process with continuous sample path, expected value $\frac{1}{2}\delta'\mathbb{W}\delta$ and covariance kernel $\mathbb{H}_2(\delta_1, \delta_2)$.

C.2 Proof of Theorem 3.3

Proof of Theorem 3.3. We show that the numerical bootstrap works for β . The discussion for γ is omitted due to the similarity.

The key is to show the results in Hong and Li (2020) hold by modifying condition (vi) in Theorem 4.1 (using their notations), for example, for β , that

$$\Sigma_{1/2}(\rho_1, \rho_2) = \lim_{N \to \infty} (Nh_N^{k_1 + k_2})^{1/3} \mathbb{E}[h_N^{k_1 + k_2} \phi_{Ni} (\beta + \rho_1 (Nh_N^{k_1 + k_2})^{-1/3}) \phi_{Ni} (\beta + \rho_2 (Nh_N^{k_1 + k_2})^{-1/3})]$$

exists for all $\rho_1, \rho_2 \in \mathbb{R}^{k_1}$. This is indeed the case as shown by our Lemma C.2 in Appendix C. We now provide the details of the proof.

By some standard calculation as Seo and Otsu (2018), we can show that the convergence rate of $\hat{\beta}^*$ is $(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3}$. Thus, if we manage to show the limit of

$$(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3} \cdot N^{-1} \sum_{i=1}^N \phi_{Ni}(\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3}) + (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3} \cdot (N\varepsilon_{N1})^{1/2}$$

$$\cdot [N^{-1} \sum_{i=1}^N \phi_{Ni}^* (\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3}) - N^{-1} \sum_{i=1}^N \phi_{Ni}(\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3})],$$
(C.5)

then the limiting distribution of $\hat{\beta}^*$ can be established.

For the first term in equation (C.5),

$$(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3} \cdot N^{-1} \sum_{i=1}^N \phi_{Ni}(\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3})$$

$$= (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3} \mathbb{E}[\phi_{Ni}(\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3})]$$

$$+ (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3} \cdot N^{-1} \sum_{i=1}^N [\phi_{Ni}(\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3}) - \mathbb{E}[\phi_{Ni}(\beta + \rho(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/3})]]$$

$$= \frac{1}{2}\rho' \mathbb{V}\rho + O_p((\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3} \cdot N^{-1/2} \cdot h_N^{-(k_1+k_2)/2}(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{-1/6}) = \frac{1}{2}\rho' \mathbb{V}\rho + o_p(1),$$
(C.6)

where the second equality follows by Lemma C.2, the assumption on ε_{N1} such that the bias

term h_N^2 is a small order term compared to $(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3}$, and Markov inequality, and the last equality holds by $N\varepsilon_{N1} \to \infty$.

The second term in equation (C.5) is a sample average of mean 0 series under P^* multiplied by $(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{2/3}(N\varepsilon_{N1})^{1/2}$. The covariance kernel of the second term is

$$(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{4/3} \cdot (N\varepsilon_{N1}) \cdot N^{-1} \cdot \mathbb{E}^{*} [\phi_{Ni}^{*} (\beta + \rho_{1}(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{-1/3}) \phi_{Ni}^{*} (\beta + \rho_{2}(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{-1/3})]$$

$$= (\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{1/3} \mathbb{E}^{*} [h_{N}^{k_{1}+k_{2}} \phi_{Ni}^{*} (\beta + \rho_{1}(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{-1/3}) \phi_{Ni}^{*} (\beta + \rho_{2}(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{-1/3})]$$

$$= (\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{1/3} \mathbb{E} [h_{N}^{k_{1}+k_{2}} \phi_{Ni} (\beta + \rho_{1}(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{-1/3}) \phi_{Ni} (\beta + \rho_{2}(\varepsilon_{N1}^{-1}h_{N}^{k_{1}+k_{2}})^{-1/3})] + o_{p}(1)$$

$$= \mathbb{H}_{1}(\rho_{1}, \rho_{2}) + o_{p}(1),$$

$$(C.7)$$

by the i.i.d. sampling and Lemma C.2.

Equations (C.6) and (C.7) imply that the term in equation (C.5) converges to $\mathcal{Z}_{1}^{*}(\rho)$ with mean $\frac{1}{2}\rho'\mathbb{V}\rho$ and covariance kernel $\mathbb{H}_{1}(\rho_{1},\rho_{2})$. Thus, $\mathcal{Z}_{1}^{*}(\rho)$ is an independent copy of $\mathcal{Z}_{1}(\rho)$ and

$$(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^*-\beta) \stackrel{d}{\to} \arg\max_{\rho\in\mathbb{R}^{k_1}} \mathcal{Z}_1^*(\rho)$$
.

Finally, note, by $N\varepsilon_{N1} \to \infty$ and

$$\begin{split} (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^* - \hat{\beta}) &= (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^* - \beta) + (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta} - \beta) \\ &= (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^* - \beta) + O_p((\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3} \cdot (Nh_N^{k_1+k_2})^{-1/3}) \\ &= (\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^* - \beta) + o_p(1), \end{split}$$

we have

$$(\varepsilon_{N1}^{-1}h_N^{k_1+k_2})^{1/3}(\hat{\beta}^* - \hat{\beta}) \stackrel{d}{\to} \arg\max_{\rho \in \mathbb{R}^{k_1}} \mathcal{Z}_1^*(\rho).$$

C.3 Proof of Theorem 3.4

Proof of Theorem 3.4. Recall that under Assumptions P1–P9, we have shown that our estimator satisfies the technical conditions required by Seo and Otsu (2018) in Lemma C.1. Then applying Lemma 1 in Seo and Otsu (2018) to $N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(r)$, we get

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(r) - \bar{\mathcal{L}}^{P}\left(r\right) = N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\gamma) - \bar{\mathcal{L}}^{P}\left(\gamma\right) + \varepsilon \left\|r - \gamma\right\|^{2} + O_{p}(\sigma_{N}^{2}) + O_{p}((N\sigma_{N}^{2k_{1}})^{-2/3}) \quad (\text{C.8})$$

for any small $\varepsilon > 0$, where use some of the results in Lemma C.2.

Some standard calculations as in the proof of Lemma C.2 yield

$$\bar{\mathcal{L}}^{P}(r) - \bar{\mathcal{L}}^{P}(\gamma) = \frac{1}{2} (r - \gamma)' \mathbb{W}(r - \gamma) + o_p (\|r - \gamma\|^2)$$
 (C.9)

for any r in a small neighborhood of γ .

Recall that we assume $(N\sigma_N^{2k_1})^{2/3}\sigma_N^2 \to 0$ in Assumption P9 and that W is finite. Combining these two conditions with (C.8) and (C.9) gives

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(r) = N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\gamma) + O_p(\|r - \gamma\|^2) + O_p((N\sigma_N^{2k_1})^{-2/3}),$$

which further implies

$$N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma}) - N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\gamma) = O_p((N\sigma_N^{2k_1})^{-2/3})$$

since $\hat{\gamma} - \gamma = O_p((N\sigma_N^{2k_1})^{-1/3})$ as shown in Theorem 3.2. Therefore, we have

$$\sqrt{N\sigma_N^{2k_1}} \left(N^{-1} \mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma}) - N^{-1} \mathcal{L}_{N,\gamma}^{P,K}(\gamma) \right) = o_p(1). \tag{C.10}$$

Additionally, by applying the Lindeberg-Feller CLT, we have

$$\sqrt{N\sigma_N^{2k_1}}\left(N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\gamma) - \bar{\mathcal{L}}^P(\gamma)\right) \stackrel{d}{\to} N\left(0, \Delta_{\gamma}^P\right).$$

Combining this with (C.10), we arrive at the desired result

$$\sqrt{N\sigma_{N}^{2k_{1}}}\left(N^{-1}\mathcal{L}_{N,\gamma}^{P,K}(\hat{\gamma})-\bar{\mathcal{L}}^{P}\left(\gamma\right)\right)\overset{d}{\to}N\left(0,\Delta_{\gamma}^{P}\right).$$

D Technical Details for Theorem 2.3

In this section, we show the validity of the nonparametric bootstrap for the cross-sectional model (Section 2.1.3). A complete proof of the consistency of the bootstrap procedure (Theorem 2.3) is lengthy and tedious. Considering that this process is relatively standard

in the literature, we only present an outline of the proof. To fill in the gaps of the outline, one first needs to define a product probability space for both the original random series and the bootstrap series as in Wellner and Zhan (1996) (an application can be found in Abrevaya and Huang (2005)), define a suitable norm for functions, and then establish some uniform convergence results following Sherman (1993, 1994a,b) on this probability space with this norm. To apply Sherman's results, the criterion functions need to be manageable in the sense of Kim and Pollard (1990). This is indeed the case for our estimators. For example, equation (2.7) is a summation of

$$\mathcal{K}_{\sigma_N}(V_{im}(\hat{\beta}))Y_{im(1,1)}\operatorname{sgn}(W'_{im}r) = \mathcal{K}_{\sigma_N}(V_{im}(\hat{\beta}))Y_{im(1,1)}(2 \cdot 1[W'_{im}r > 0] - 1).$$

The collection of indicator functions $1[W'_{im}r > 0]$ for $r \in \mathcal{R}$ is well known to be a Vapnik–Chervonenkis (VC) class. The kernel function \mathcal{K} , when is chosen to satisfy some mild conditions (e.g., uniformly bounded with compact support and continuously differentiable), is also manageable. The complication of the bandwidth can be handled in the same way as in Seo and Otsu (2018).

Now suppose we have handled the technical details mentioned above. In what follows, we present the proof outline. To ease expositions, we assume $\hat{\beta}$ and $\hat{\gamma}$ are obtained from maximizing, respectively, $\mathbb{L}_{N,\beta}^K(b)$ and $\mathbb{L}_{N,\gamma}^K(r,\hat{\beta})$, which are defined as

$$\mathbb{L}_{N,\beta}^{K}(b) = \sum_{i=1}^{N-1} \sum_{m>i} \mathcal{K}_{h_{N}}(X_{im}) \chi_{1}(Z_{i}, Z_{m}, b)$$

and

$$\mathbb{L}_{N,\gamma}^{K}(r,\hat{\beta}) = \sum_{i=1}^{N-1} \sum_{m>i} \mathcal{K}_{\sigma_N}(X_{im}'\hat{\beta}) \chi_2(Z_i, Z_m, r),$$

where X_{im} is a sub-vector of $Z_i - Z_m$. Note that the criterion functions and notations defined here are meant for general U-processes, and thus are different from the ones used in the main text. However, the procedures that we study here share the same structure as those in Section 2.1 (i.e., (2.7)–(2.7)). Therefore, if we can demonstrate the consistency of the bootstrap here, we can easily apply the same method to our MRC estimators.

We do normalization such that $\chi_1(Z_m, Z_i, \beta) = 0$ and $\chi_2(Z_i, Z_m, \gamma) = 0$ for all Z^{20} Assume that $\chi_1(Z_i, Z_m, b) = \chi_1(Z_m, Z_i, b)$, β uniquely maximizes $\mathbb{E}[\chi_1(Z_i, Z_m, b) | X_{im} = 0]$, and \mathcal{K}_{h_N} is a p-th order kernel with h_N that satisfies $Nh_N^p = o(1)$. Similarly, we assume

²⁰Taking equation (2.7) as an example, this can be done by replacing $\operatorname{sgn}(W'_{im}r)$ with $\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma)$ in the criterion function.

that $\chi_2(Z_i, Z_m, r) = \chi_2(Z_m, Z_i, r)$, γ uniquely maximizes $\mathbb{E}\left[\chi_2(Z_i, Z_m, r) | X'_{im}\beta = 0\right]$, and \mathcal{K}_{σ_N} is a p-th order kernel with σ_N that satisfies $N\sigma_N^p = o(1)$. We similarly denote $\mathbb{L}_{N,\beta}^{K*}(b)$ and $\mathbb{L}_{N,\gamma}^{K*}(r, \hat{\beta}^*)$ as the corresponding criterion functions using the bootstrap series.

Before establishing the consistency of the bootstrap, it is helpful to do a recast of how we derive the asymptotics for $(\hat{\beta}, \hat{\gamma})$. $\mathbb{L}_{N,\beta}^{K}(b)$ can be decomposed into

$$2 [N (N-1)]^{-1} \mathbb{L}_{N,\beta}^{K} (b)$$

$$= \mathbb{E} [\mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b)]$$

$$+ \frac{2}{N} \sum_{i=1}^{N} \{ \mathbb{E} [\mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b) | Z_{i}] - \mathbb{E} [\mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b)] \}$$

$$+ \frac{1}{N (N-1)} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \{ \mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b) - \mathbb{E} [\mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b) | Z_{i}]$$

$$- \mathbb{E} [\mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b) | Z_{m}] + \mathbb{E} [\mathcal{K}_{h_{N}} (X_{im}) \chi_{1} (Z_{i}, Z_{m}, b)] \}$$

$$\equiv \mathcal{T}_{N1} + \mathcal{T}_{N2} + \mathcal{T}_{N3}.$$
(D.1)

Since $\chi_{1}\left(Z_{i},Z_{m},\beta\right)=0$ and β uniquely maximizes $\mathbb{E}\left[\chi_{1}\left(Z_{i},Z_{m},b\right)|X_{im}=0\right]$, we have

$$\mathcal{T}_{1N} = \mathbb{E} \left[\chi_1 \left(Z_i, Z_m, b \right) | X_{im} = 0 \right] + O_p \left(h_N^p \right)$$

= $\frac{1}{2} \left(b - \beta \right)' \mathbb{V}_{\beta} \left(b - \beta \right) + O_p \left(\| b - \beta \|^3 \right) + O_p \left(h_N^p \right),$

where the $O_p(h_N^p)$ is the bias term from matching and \mathbb{V}_{β} is a negative definite matrix. For the second term,

$$\mathcal{T}_{N2} = \frac{2}{N} \sum_{i=1}^{N} \left\{ \mathbb{E} \left[\chi_1 \left(Z_i, Z_m, b \right) | Z_i, X_{im} = 0 \right] f_{X_m} \left(X_i \right) - \mathbb{E} \left[\chi_1 \left(Z_i, Z_m, b \right) | X_{im} = 0 \right] \right\} + O_p \left(h_N^p \right)$$

$$= 2 \left(b - \beta \right)' \frac{W_{N1}}{\sqrt{N}} + o_p \left(\|b - \beta\|^2 \right) + O_p \left(h_N^p \right),$$
(D.2)

where W_{N1} converges to a normal distribution. For the last term,

$$\mathcal{T}_{N3} = o_p\left(N^{-1}\right),\,$$

by similar arguments in Sherman (1993). Put all these results together to obtain

$$2\left[N\left(N-1\right)\right]^{-1}\mathbb{L}_{N,\beta}^{K}\left(b\right) = \frac{1}{2}\left(b-\beta\right)'\mathbb{V}_{\beta}\left(b-\beta\right) + 2\left(b-\beta\right)'\frac{W_{N1}}{\sqrt{N}} + o_{p}\left(\|b-\beta\|^{2}\right) + o_{p}\left(N^{-1}\right).$$

 $\hat{\beta} - \beta$ can be shown to be $O_p(N^{-1/2})$. Using the rate result and invoking the CMT, we have

$$\sqrt{N}(\hat{\beta} - \beta) = -2\mathbb{V}_{\beta}^{-1}W_{N1} + o_p(1). \tag{D.3}$$

Let \mathbb{E}^* and P^* denote the expectation and the density for the bootstrap series, respectively. By definition, P^* puts N^{-1} probability on each of $\{Z_i\}_{i=1}^N$. We can do a similar decomposition of $\mathbb{L}_{N,\beta}^{K*}(b)$ as for $\mathbb{L}_{N,\beta}^K(b)$:

$$2 [N (N-1)]^{-1} \mathbb{L}_{N,\beta}^{K*}(b)$$

$$= [N (N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1,m\neq i}^{N} \mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b)$$

$$= \mathbb{E}^{*} [\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b)]$$

$$+ \frac{2}{N} \sum_{i=1}^{N} {\mathbb{E}^{*} [\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b) | Z_{i}^{*}] - \mathbb{E}^{*} [\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b)]}$$

$$+ \frac{1}{N(N-1)} \sum_{i=1}^{N} \sum_{m=1,m\neq i}^{N} {\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b) - \mathbb{E}^{*} [\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b) | Z_{i}^{*}]}$$

$$- \mathbb{E}^{*} [\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b) | Z_{m}^{*}] + \mathbb{E}^{*} [\mathcal{K}_{h_{N}}(X_{im}^{*}) \chi_{1}(Z_{i}^{*}, Z_{m}^{*}, b)]}$$

$$\equiv \mathcal{T}_{N1}^{*} + \mathcal{T}_{N2}^{*} + \mathcal{T}_{N3}^{*}. \tag{D.4}$$

For \mathcal{T}_{N1}^* , by the definition of of \mathbb{E}^* ,

$$\mathcal{T}_{N1}^{*} = \mathbb{E}^{*} \left[\mathcal{K}_{h_{N}} \left(X_{im}^{*} \right) \chi_{1} \left(Z_{i}^{*}, Z_{m}^{*}, b \right) \right] \\
= \frac{1}{N^{2}} \sum_{i=1}^{N} \sum_{m=1}^{N} \mathcal{K}_{h_{N}} \left(X_{im} \right) \chi_{1} \left(Z_{i}, Z_{m}, b \right) \\
= \frac{1}{2} \left(b - \beta \right)' \mathbb{V}_{\beta} \left(b - \beta \right) + 2 \left(b - \beta \right)' \frac{W_{N1}}{\sqrt{N}} + O_{p} \left(\|b - \beta\|^{3} \right) + o_{p} \left(N^{-1} \right), \tag{D.5}$$

where the last line holds because the second line is the same as $2 [N(N-1)]^{-1} \mathbb{L}_{N,\beta}^{K}(b)$ plus a small order term.

Both \mathcal{T}_{N2} and \mathcal{T}_{N2}^* are sample averages of mean zero series. Moreover, for $||b - \beta|| = O(N^{-1/2})$, $N\mathcal{T}_{N2} = 2\sqrt{N}(b-\beta)'W_{N1} + o_p(1)$ and W_{N1} converges to a normal distribution. Then, we can apply the results in Giné and Zinn (1990) that NT_{N2}^* approximates the distri-

bution of NT_{N2} for $||b - \beta|| = O(N^{-1/2})$. That is, for $||b - \beta|| = O(N^{-1/2})$,

$$\mathcal{T}_{N2}^{*} = 2 \left(b - \beta \right)' \frac{W_{N1}^{*}}{\sqrt{N}} + o_p \left(N^{-1} \right), \tag{D.6}$$

where W_{N1}^* is an independent copy of W_{N1} .

For \mathcal{T}_{N3}^* , terms across i and m are uncorrelated with each other under P^* . Therefore, $\left[\mathbb{E}^*\left(\mathcal{T}_{N3}^{*2}\right)\right]^{1/2}$ is of the same order as

$$N^{-1} \left\{ \mathbb{E}^* \left[\left[\mathcal{K}_{h_N} \left(X_{im}^* \right) \chi_1 \left(Z_i^*, Z_m^*, b \right) \right]^2 \right] \right\}^{1/2} = N^{-1} \left\{ \frac{1}{N^2} \sum_{i=1}^N \sum_{m=1}^N \mathcal{K}_{h_N} \left(X_{im} \right)^2 \chi_1 \left(Z_i, Z_m, b \right)^2 \right\}^{1/2}.$$

Note that $\chi_1(Z_i, Z_m, \beta) = 0$, so the above term is $o_p(N^{-1})$ for $||b - \beta|| = O(N^{-1/2})$. That means $\left[\mathbb{E}^* (\mathcal{T}_{N3}^{*2})\right]^{1/2} = o_p(N^{-1})$. By Markov inequality,

$$\mathcal{T}_{N3}^* = o_p\left(N^{-1}\right). \tag{D.7}$$

To summarize, equations (D.4)–(D.7) imply that for $\|b - \beta\| = O\left(N^{-1/2}\right)$,

$$2\left[N\left(N-1\right)\right]^{-1}\mathbb{L}_{N,\beta}^{K*}\left(b\right) = \frac{1}{2}\left(b-\beta\right)'\mathbb{V}_{\beta}\left(b-\beta\right) + 2\left(b-\beta\right)'\frac{W_{N1}}{\sqrt{N}} + 2\left(b-\beta\right)'\frac{W_{N1}^{*}}{\sqrt{N}} + o_{p}\left(N^{-1}\right).$$

We can similarly show that $\hat{\beta}^* - \beta = O_p(N^{-1/2})$. By the CMT again,

$$\sqrt{N}(\hat{\beta}^* - \beta) = -2\mathbb{V}_{\beta}^{-1}W_{N1} - 2\mathbb{V}_{\beta}^{-1}W_{N1}^* + o_p(1).$$

Substitute equation (D.3) into the equation above to get

$$\sqrt{N}(\hat{\beta}^* - \hat{\beta}) = -2V_{\beta}^{-1}W_{N1}^* + o_p(1).$$
 (D.8)

Since W_{N1}^* is an independent copy of W_{N1} , $\sqrt{N}(\hat{\beta}^* - \hat{\beta})$ converges to the same distribution as the one $\sqrt{N}(\hat{\beta} - \beta)$ converges to.

We turn to the inference for $\hat{\gamma}$. Again, we decompose $\mathbb{L}_{N,\gamma}^K(r,\hat{\beta})$ and analyze each term

in the decompositions one by one. We start with

$$2 [N (N-1)]^{-1} \mathbb{L}_{N,\gamma}^{K} \left(r, \hat{\beta}\right)$$

$$= [N (N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \mathcal{K}_{\sigma_{N}} \left(X'_{im} \hat{\beta}\right) \chi_{2} \left(Z_{i}, Z_{m}, r\right)$$

$$= [N (N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \mathcal{K}_{\sigma_{N}} \left(X'_{im} \beta\right) \chi_{2} \left(Z_{i}, Z_{m}, r\right)$$

$$+ [N (N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2} \left(Z_{i}, Z_{m}, r\right) \nabla \mathcal{K}_{\sigma_{N}} \left(X'_{im} \beta\right) X'_{im} \frac{\left(\hat{\beta} - \beta\right)}{\sigma_{N}}$$

$$+ [N (N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2} \left(Z_{i}, Z_{m}, r\right) \nabla^{2} \mathcal{K}_{\sigma_{N}} \left(X'_{im} \tilde{\beta}\right) \left[X'_{im} \frac{\left(\hat{\beta} - \beta\right)}{\sigma_{N}}\right]^{2}$$

$$= \mathcal{T}_{N4} + \mathcal{T}_{N5} + \mathcal{T}_{N6},$$
(D.9)

where $\tilde{\beta}$ is a vector that lies between β and $\hat{\beta}$ and makes the equality hold.

Note that \mathcal{T}_{N4} shares the same structure as $2\left[N\left(N-1\right)\right]^{-1}\mathbb{L}_{N,\beta}^{K}\left(b\right)$, and so

$$\mathcal{T}_{N4} = \frac{1}{2} (r - \gamma)' \mathbb{V}_{\gamma} (r - \gamma) + 2 (r - \gamma)' \frac{W_{N2}}{\sqrt{N}} + O_p (\|r - \gamma\|^3) + O_p (\sigma_N^p) + o_p (N^{-1}), \quad (D.10)$$

where \mathbb{V}_{γ} is a negative definite matrix, and W_{N2} converges to a normal distribution.

For term \mathcal{T}_{N5} ,

$$\mathcal{T}_{N5} = \left\{ [N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2}(Z_{i}, Z_{m}, r) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}}(X'_{im}\beta) X'_{im} \right\} \left(\hat{\beta} - \beta \right).$$

Note that

$$[N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_2(Z_i, Z_m, r) \, \sigma_N^{-1} \nabla \mathcal{K}_{\sigma_N}(X'_{im}\beta) \, X'_{im}$$
(D.11)

$$= \mathbb{E}\left[\chi_2\left(Z_i, Z_m, r\right) \sigma_N^{-1} \nabla \mathcal{K}_{\sigma_N}\left(X'_{im} \beta\right) X'_{im}\right]$$

$$+\frac{2}{N}\sum_{i=1}^{N}\left\{\mathbb{E}\left[\chi_{2}\left(Z_{i},Z_{m},r\right)\sigma_{N}^{-1}\nabla\mathcal{K}_{\sigma_{N}}\left(X_{im}'\beta\right)X_{im}'|Z_{i}\right]-\mathbb{E}\left[\chi_{2}\left(Z_{i},Z_{m},r\right)\sigma_{N}^{-1}\nabla\mathcal{K}_{\sigma_{N}}\left(X_{im}'\beta\right)X_{im}'\right]\right\}$$

$$+ \frac{1}{N(N-1)} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \left\{ \chi_{2}(Z_{i}, Z_{m}, r) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}}(X'_{im}\beta) X'_{im} - \mathbb{E} \left[\chi_{2}(Z_{i}, Z_{m}, r) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}}(X'_{im}\beta) X'_{im} \right] \right\}$$

$$-\mathbb{E}\left[\chi_{2}\left(Z_{i},Z_{m},r\right)\sigma_{N}^{-1}\nabla\mathcal{K}_{\sigma_{N}}\left(X_{im}'\beta\right)X_{im}'|Z_{m}\right]+\mathbb{E}\left[\chi_{2}\left(Z_{i},Z_{m},r\right)\sigma_{N}^{-1}\nabla\mathcal{K}_{\sigma_{N}}\left(X_{im}'\beta\right)X_{im}'\right]\right\}$$

with the lead term $\mathbb{E}[\chi_2(Z_i, Z_m, r) \, \sigma_N^{-1} \nabla \mathcal{K}_{\sigma_N}(X'_{im}\beta) \, X'_{im}]$. As $\int \nabla \mathcal{K}_{\sigma_N}(u) \, \mathrm{d}u = 0$, the lead term cancels one more σ_N^{-1} when calculating the expectation. Since $\chi_2(Z_i, Z_m, \gamma) = 0$, we can write

$$\mathbb{E}\left[\chi_{2}\left(Z_{i}, Z_{m}, r\right) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}}\left(X_{im}^{\prime} \beta\right) X_{im}\right] = \left(r - \gamma\right)^{\prime} \Gamma + O(\|r - \gamma\|^{2}) + O\left(\sigma_{N}^{p}\right), \quad (D.12)$$

for some matrix Γ . Collect all these result to obtain

$$\mathcal{T}_{N5} = (r - \gamma)' \Gamma\left(\hat{\beta} - \beta\right) + O_p\left(\|r - \gamma\|^2 \|\hat{\beta} - \beta\|\right) + O\left(\sigma_N^p\right)$$

$$= -2\left(r - \gamma\right)' \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}}{\sqrt{N}} + O_p\left(\|r - \gamma\|^2 \|\hat{\beta} - \beta\|\right) + o_p\left(\frac{\|r - \gamma\|}{\sqrt{N}}\right) + O\left(\sigma_N^p\right),$$
(D.13)

where the second line follows by substituting in equation (D.3).

For term \mathcal{T}_{N6} , it is not hard to see that

$$\mathcal{T}_{N6} = O_p \left(\left\| r - \gamma \right\| \left\| \hat{\beta} - \beta \right\|^2 / \sigma_N^2 \right) = o_p \left(N^{-1} \right), \tag{D.14}$$

for $||r - \gamma|| = O(N^{-1/2})$, and σ_N in Assumption C7.

To summarize, equations (D.9), (D.10), (D.13), and (D.14) imply that for $||r - \gamma|| = O(N^{-1/2})$,

$$2 [N (N - 1)]^{-1} \mathbb{L}_{N,\gamma}^{K} (r, \hat{\beta}) = \frac{1}{2} (r - \gamma)' \mathbb{V}_{\gamma} (r - \gamma) + 2 (r - \gamma)' \frac{W_{N2}}{\sqrt{N}} - 2 (r - \gamma)' \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}}{\sqrt{N}} + o_{p} (N^{-1}).$$

Once $\hat{\gamma} - \gamma = O_p\left(N^{-1/2}\right)$ is established, applying the CMT gives

$$\sqrt{N} (\hat{\gamma} - \gamma) = -2 \mathbb{V}_{\gamma}^{-1} W_{N2} + 2 \mathbb{V}_{\gamma}^{-1} \Gamma \mathbb{V}_{\beta}^{-1} W_{N1} + o_p (1).$$
 (D.15)

For $\mathbb{L}_{N,\gamma}^{K*}(r,b)$, we decompose it similarly:

$$2[N(N-1)]^{-1} \mathbb{L}_{N,\gamma}^{K*} \left(r, \hat{\beta}^{*}\right)$$

$$= [N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta}^{*}\right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r\right)$$

$$= [N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta}\right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r\right)$$

$$+ [N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r\right) \nabla \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta}\right) \left(\frac{X_{im}^{*\prime} \left(\hat{\beta}^{*} - \hat{\beta}\right)}{\sigma_{N}}\right)$$

$$+ [N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r\right) \nabla^{2} \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta}^{*}\right) \left(\frac{X_{im}^{*\prime} \left(\hat{\beta}^{*} - \hat{\beta}\right)}{\sigma_{N}}\right)^{2}$$

$$= \mathcal{T}_{N4}^{*} + \mathcal{T}_{N5}^{*} + \mathcal{T}_{N6}^{*},$$
(D.16)

where $\tilde{\beta}^*$ is some vector that lies between b and $\hat{\beta}$ and makes the equality hold.

For \mathcal{T}_{N4}^* , we have

$$\mathcal{T}_{N4}^{*} = [N(N-1)]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \\
= \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \right] \\
+ \frac{2}{N} \sum_{i=1}^{N} \left\{ \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) | Z_{i}^{*} \right] - \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \right] \right\} \\
+ \frac{1}{N(N-1)} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \left\{ \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) - \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) | Z_{i}^{*} \right] \\
- \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) | Z_{m}^{*} \right] + \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \right] \right\}. \\
\equiv \mathcal{T}_{N4.1}^{*} + \mathcal{T}_{N4.2}^{*} + \mathcal{T}_{N4.3}^{*}. \tag{D.17}$$

We analyze the three terms in (D.17) one by one.

For $\mathcal{T}_{N4,1}^*$, we have

$$\mathcal{T}_{N4,1}^{*} = \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*'} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \right] \\
= \frac{1}{N^{2}} \sum_{i=1}^{N} \sum_{m=1}^{N} \mathcal{K}_{\sigma_{N}} \left(X_{im}^{'} \hat{\beta} \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) \\
= \frac{1}{2} \left(r - \gamma \right)^{\prime} \mathbb{V}_{\gamma} \left(r - \gamma \right) + 2 \left(r - \gamma \right)^{\prime} \frac{W_{N2}}{\sqrt{N}} - 2 \left(r - \gamma \right)^{\prime} \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}}{\sqrt{N}} + o_{p} \left(N^{-1} \right),$$

where the last line holds because the term in the second line is $2[N(N-1)]^{-1}\mathbb{L}_{N,\gamma}^K(r,\hat{\beta})$ plus a small order term.

For $\mathcal{T}_{N4,2}^*$, we first note that

$$\sum_{i=1}^{N} \left\{ \mathbb{E} \left[\mathcal{K}_{\sigma_N} \left(X'_{im} \beta \right) \chi_2 \left(Z_i, Z_m, r \right) | Z_i \right] - \mathbb{E} \left[\mathcal{K}_{\sigma_N} \left(X'_{im} \beta \right) \chi_2 \left(Z_i, Z_m, r \right) \right] \right\}$$

$$= \sqrt{N} \left(r - \gamma \right)' W_{N2} + o_p \left(1 \right),$$
(D.19)

for $||r - \gamma|| = O(N^{-1/2})$, which holds for the same reason as for equation (D.2). Since $\hat{\beta} - \beta = O_p(N^{-1/2})$, by the equicontinuity we claimed,

$$\sum_{i=1}^{N} \left\{ \mathbb{E} \left[\mathcal{K}_{\sigma_{N}} \left(X'_{im} \beta \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) | Z_{i} \right] - \mathbb{E} \left[\mathcal{K}_{\sigma_{N}} \left(X'_{im} \beta \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) \right] \right\}$$

$$- \sum_{i=1}^{N} \left\{ \mathbb{E}_{N} \left[\mathcal{K}_{\sigma_{N}} \left(X'_{im} \hat{\beta} \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) | Z_{i} \right] - \mathbb{E}_{N} \left[\mathcal{K}_{\sigma_{N}} \left(X'_{im} \hat{\beta} \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) \right] \right\} = o_{p} \left(1 \right).$$

Then equations (D.19) and (D.20) imply that

$$2\sum_{i=1}^{N} \left\{ \mathbb{E}_{N} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}' \hat{\beta} \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) | Z_{i} \right] - \mathbb{E}_{N} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}' \hat{\beta} \right) \chi_{2} \left(Z_{i}, Z_{m}, r \right) \right] \right\}$$

$$= 2\sqrt{N} \left(r - \gamma \right)' W_{N2} + o_{p} \left(1 \right),$$

which approximates a normal distribution for $||r - \gamma|| = O(N^{-1/2})$. Thus, we are able to apply the results of Giné and Zinn (1990) and get that

$$N\mathcal{T}_{N4,2}^{*} = 2\sum_{i=1}^{N} \left\{ \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*'} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) | Z_{i}^{*} \right] - \mathbb{E}^{*} \left[\mathcal{K}_{\sigma_{N}} \left(X_{im}^{*'} \hat{\beta} \right) \chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \right] \right\}$$

approximates the distribution of the above term. That is

$$\mathcal{T}_{N4,2}^* = 2 \left(r - \gamma \right)' \frac{W_{N2}^*}{\sqrt{N}} + o_p \left(N^{-1} \right), \tag{D.21}$$

where W_{N2}^* is an independent copy of W_{N2} .

The same arguments as we used for \mathcal{T}_{N3}^* lead to

$$\mathcal{T}_{N4,3}^* = o_p\left(N^{-1}\right). \tag{D.22}$$

Then, equations (D.17), (D.18), (D.21), and (D.22) together imply that for $||r - \gamma|| = O(N^{-1/2})$,

$$\mathcal{T}_{N4}^{*} = \frac{1}{2} (r - \gamma)' \mathbb{V}_{\gamma} (r - \gamma) + 2 (r - \gamma)' \frac{W_{N2}}{\sqrt{N}} - 2 (r - \gamma)' \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}}{\sqrt{N}} + 2 (r - \gamma)' \frac{W_{N2}^{*}}{\sqrt{N}} + o_{p} (N^{-1}).$$
(D.23)

For \mathcal{T}_{N5}^* , we have

$$\mathcal{T}_{N5}^{*} = \left[N\left(N-1\right)\right]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2}\left(Z_{i}^{*}, Z_{m}^{*}, r\right) \nabla \mathcal{K}_{\sigma_{N}}\left(X_{im}^{*\prime}\hat{\beta}\right) \left(\frac{X_{im}^{*\prime}\left(\hat{\beta}^{*} - \hat{\beta}\right)}{\sigma_{N}}\right)$$

$$= \left\{\left[N\left(N-1\right)\right]^{-1} \sum_{i=1}^{N} \sum_{m=1, m \neq i}^{N} \chi_{2}\left(Z_{i}^{*}, Z_{m}^{*}, r\right) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}}\left(X_{im}^{*\prime}\hat{\beta}\right) X_{im}^{*\prime}\right\} \left(\hat{\beta}^{*} - \hat{\beta}\right).$$

Following the same analysis for \mathcal{T}_{N5} , the lead term for \mathcal{T}_{N5}^* is

$$\mathbb{E}^* \left[\chi_2 \left(Z_i^*, Z_m^*, r \right) \sigma_N^{-1} \nabla \mathcal{K}_{\sigma_N} \left(X_{im}^{*\prime} \hat{\beta} \right) X_{im}^{*\prime} \right] \left(\hat{\beta}^* - \hat{\beta} \right).$$

Note that

$$\begin{split} &\mathbb{E}^{*} \left[\chi_{2} \left(Z_{i}^{*}, Z_{m}^{*}, r \right) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}} \left(X_{im}^{*\prime} \hat{\beta} \right) X_{im}^{*\prime} \right] \left(\hat{\beta}^{*} - \hat{\beta} \right) \\ &= \left[\frac{1}{N^{2}} \sum_{i=1}^{N} \sum_{m=1}^{N} \chi_{2} \left(Z_{i}, Z_{m}, r \right) \sigma_{N}^{-1} \nabla \mathcal{K}_{\sigma_{N}} \left(X_{im}^{\prime} \beta \right) X_{im} \right] \left(\hat{\beta}^{*} - \hat{\beta} \right) \\ &= \left(r - \gamma \right)^{\prime} \Gamma \left(\hat{\beta}^{*} - \hat{\beta} \right) + O_{p} \left(\| r - \gamma \|^{2} \left\| \hat{\beta}^{*} - \hat{\beta} \right\| \right) + o_{p} \left(N^{-1} \right) \\ &= -2 \left(r - \gamma \right)^{\prime} \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}^{*}}{\sqrt{N}} + o_{p} \left(\frac{\| r - \gamma \|}{\sqrt{N}} \right) + O_{p} \left(\| r - \gamma \|^{2} \left\| \hat{\beta}^{*} - \hat{\beta} \right\| \right) + o_{p} \left(N^{-1} \right), \end{split}$$

where we use equations (D.11) and (D.12) to get the third line, and we substitute equation (D.8) in the last line. Then we have

$$\mathcal{T}_{N5}^{*} = -2 (r - \gamma)' \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}^{*}}{\sqrt{N}} + o_{p} \left(\frac{\|r - \gamma\|}{\sqrt{N}} \right) + O_{p} \left(\|r - \gamma\|^{2} \|\hat{\beta}^{*} - \hat{\beta}\| \right) + o_{p} \left(N^{-1} \right). \quad (D.24)$$

For \mathcal{T}_{N6}^* , it is not hard to see that

$$\mathcal{T}_{N6}^* = O_p \left(\frac{\left\| r - \gamma \right\| \left\| \hat{\beta}^* - \hat{\beta} \right\|^2}{\sigma_N^2} \right). \tag{D.25}$$

Then, equations (D.16), (D.23), (D.24), and (D.25) together imply that for $||r - \gamma|| = O(N^{-1/2})$,

$$2 \left[N \left(N - 1 \right) \right]^{-1} \mathbb{L}_{N,\gamma}^{K*} \left(r, \hat{\beta}^* \right)$$

$$= \frac{1}{2} \left(r - \gamma \right)' \mathbb{V}_{\gamma} \left(r - \gamma \right) + 2 \left(r - \gamma \right)' \frac{W_{N2}}{\sqrt{N}} - 2 \left(r - \gamma \right)' \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}}{\sqrt{N}} + 2 \left(r - \gamma \right)' \frac{W_{N2}^*}{\sqrt{N}} - 2 \left(r - \gamma \right)' \Gamma \mathbb{V}_{\beta}^{-1} \frac{W_{N1}^*}{\sqrt{N}} + o_p \left(N^{-1} \right),$$

from which we can show that $\hat{\gamma}^* - \gamma = O_p\left(N^{-1/2}\right)$. With this result and by the CMT, we get

$$\sqrt{N} \left(\hat{\gamma}^* - \gamma \right) = -2 \mathbb{V}_{\gamma}^{-1} W_{N2} + 2 \mathbb{V}_{\gamma}^{-1} \Gamma \mathbb{V}_{\beta}^{-1} W_{N1} - 2 \mathbb{V}_{\gamma}^{-1} W_{N2}^* + 2 \mathbb{V}_{\gamma}^{-1} \Gamma \mathbb{V}_{\beta}^{-1} W_{N1}^* + o_p \left(1 \right),$$

which implies

$$\sqrt{N} \left(\hat{\gamma}^* - \hat{\gamma} \right) = -2 \mathbb{V}_{\gamma}^{-1} W_{N2}^* + 2 \mathbb{V}_{\gamma}^{-1} \Gamma \mathbb{V}_{\beta}^{-1} W_{N1}^* + o_p \left(N^{-1} \right)$$

by substituting equation (D.15) in.

Since W_{N1}^* and W_{N2}^* are independent copies of W_{N1} and W_{N2} , respectively, $\sqrt{N}(\hat{\gamma}^* - \hat{\gamma})$ approximates the distribution of $\sqrt{N}(\hat{\gamma} - \gamma)$.

E Additional Discussions

We discuss the convergence rates of our cross-sectional MRC and panel data MS estimators in Section E.1. We explain why we choose not to construct the panel data MS estimator for γ via index matching in Section E.2.

E.1 On the Convergence Rates of MRC and MS Estimators

The rate of convergence of the MRC estimators for the cross-sectional model is $N^{-1/2}$. The intuition is that each observation can be matched with $Nh_N^{k_1+k_2}$ and $N\sigma_N^{2k_1}$ observations (in probability) for estimating β and γ , respectively. Thus, all observations are useful and the curse of dimensionality is not a concern for the cross-sectional estimators. In contrast, an observation is useful for the panel data MS estimators if all its covariates are very close to each other across two time periods. For a fixed T, the probability of an observation being useful is proportional to $h_N^{k_1+k_2}$ or $\sigma_N^{2k_1}$. Therefore, the number of "useful" observations is proportional to $Nh_N^{k_1+k_2}$ and $N\sigma_N^{2k_1}$, and the estimation suffers from the curse of dimensionality. We may remedy this problem if $T\to\infty$ such that $Th_N^{k_1+k_2}\to\infty$ and $T\sigma_N^{2k_1}\to\infty$ and each observation can be "matched" at certain two time periods with large probability. Since this paper focuses on models for short panel data, we leave the discussion on this issue to future research. Further, the criterion functions have a "sharp" edge as in Kim and Pollard (1990). As a result, the convergence rates for $\hat{\beta}$ and $\hat{\gamma}$ are expected to be $(Nh_N^{k_1+k_2})^{-1/3}$ and $(N\sigma_N^{2k_1})^{-1/3}$, respectively. The cross-sectional estimators, however, do not suffer this "sharp" edge effect because the criterion functions are U-statistics and the "sharp" edge effect vanishes after we decompose the criterion functions (see, e.g., the second line of equation (B.6)) when deriving the asymptotics.

E.2 On the Construction of Panel Data MS Estimator

Unlike the cross-section case, our panel data MS procedure does not estimate γ through matching $X'_{jt}\hat{\beta}$ and $X'_{js}\hat{\beta}$, j=1,2, because of the following observation. As demonstrated in Theorem 3.2, the convergence rate of $\hat{\beta}$ is $(Nh_N^{k_1+k_2})^{-1/3}$. If we knew the true value of β , $\hat{\gamma}$ could be obtained by matching $X'_{jt}\beta$ and $X'_{js}\beta$. Using the same arguments, we can show that this infeasible estimator has a rate of convergence $(N\sigma_N^2)^{-1/3}$. As a result, it is only possibly beneficial to match $X_{jt}\hat{\beta}$ and $X_{js}\hat{\beta}$ for estimating γ when $\sigma_N^2 = o(h_N^{k_1+k_2})$; otherwise the asymptotics of $\hat{\beta}$ would dominate the limiting distribution of $\hat{\gamma}$. However, this involves careful selection of tuning parameters h_N and σ_N . In more general cases, this selection relies not only on the dimension of the regressor space but also on the cardinality of the choice set, and thus should be made on a case-by-case basis. Because of the lack of a universally applicable treatment, we choose to simply match the covariates to avoid this complexity and the ensuing discussion.

E.3 Identification of the Case with Three Alternatives

This section discusses how to create identification inequalities for models with three standalone alternatives and four potential bundles. More complex cases can be handled similarly. Given these inequalities, one can prove the identification and construct estimation procedures for model coefficients using the same methods employed in Sections 2 and 3. The technical conditions required for identification are similar to those of the two-alternative model studied in Sections 2 and 3. We omit them for conciseness and only focus on the identification strategies.

Throughout this section, we consider choice set $\mathcal{J} = \{0, 1, 2, 3, (1, 2), (1, 3), (2, 3), (1, 2, 3)\}$ or equivalently $\mathcal{D} = \{d | d = (d_1, d_2, d_3) \in \{0, 1\}^3\}$, which is the set of all possible $d = (d_1, d_2, d_3) \in \{0, 1\}^3$.

E.3.1 Cross-Sectional MRC Method

In the cross-sectional model, assume that an agent chooses d that maximizes the latent utility

$$U_{d} = \sum_{j=1}^{3} F_{j}(X'_{j}\beta, \epsilon_{j}) \cdot d_{j} + \eta_{110} \cdot F_{110}(W'_{1}\gamma_{1}) \cdot d_{1} \cdot d_{2} + \eta_{101} \cdot F_{101}(W'_{2}\gamma_{2}) \cdot d_{1} \cdot d_{3}$$
$$+ \eta_{011} \cdot F_{011}(W'_{3}\gamma_{3}) \cdot d_{2} \cdot d_{3} + \eta_{111} \cdot F_{111}(W'_{4}\gamma_{4}) \cdot d_{1} \cdot d_{2} \cdot d_{3},$$
 (E.1)

where $X_1, X_2, X_3 \in \mathbb{R}^{k_1}$, $W_1 \in \mathbb{R}^{k_2}$, $W_2 \in \mathbb{R}^{k_3}$, $W_3 \in \mathbb{R}^{k_4}$, $W_4 \in \mathbb{R}^{k_5}$, $\eta \equiv (\eta_{110}, \eta_{101}, \eta_{011}, \eta_{111}) \in \mathbb{R}^4$ are the bundle-specific unobserved heterogeneity, and all $F_j(\cdot)$'s and $F_d(\cdot)$'s are strictly increasing in all arguments. Note that by properly re-organizing the covariate vector, expression (E.1) can accommodate both choice-specific and common regressors. See Cameron and Trivedi (2005) p.498 for a more detailed discussion. Thus, for notational convenience, here we do not distinguish common regressors form those choice-specific regressors as we did in Section 2.2. Let $Z \equiv (X_1, X_2, X_3, W_1, W_2, W_3, W_4)$ to collect all observed covariates. We assume $(\epsilon_1, \epsilon_2, \epsilon_3, \eta) \perp Z$.

The observed dependent variable Y_d takes the form

$$Y_d = 1[U_d > U_{d'}, \forall d' \in \mathcal{D} \setminus d].$$

Then, similar to the two-alternatives²¹ case, by $(\epsilon_1, \epsilon_2 \epsilon_3, \eta) \perp Z$,

$$P(Y_{(1,d_2,d_3)} = 1 | Z, X_2 = x_2, X_3 = x_3, W_1 = w_1, W_2 = w_2, W_3 = w_3, W_4 = w_4)$$

is increasing in $X'_1\beta$ for any constant vectors $(x_2, x_3, w_1, w_2, w_3, w_4)$ and any (d_2, d_3) . Let (Y_i, Z_i) and (Y_m, Z_m) be two independent copies of (Y, Z) with

$$Y \equiv (Y_{(0,0,0)}, Y_{(1,0,0)}, Y_{(0,1,0)}, Y_{(0,0,1)}, Y_{(1,1,0)}, Y_{(1,0,1)}, Y_{(0,1,1)}, Y_{(1,1,1)}).$$

Then we have the following (conditional) moment inequalities for all d_2 and d_3 :

$$\begin{aligned}
&\{X'_{i1}\beta \ge X'_{m1}\beta\} \\
\Leftrightarrow &\{P(Y_{i(1,d_2,d_3)} = 1 | Z_i, X_{i2} = x_2, X_{i3} = x_3, W_{i1} = w_1, W_{i2} = w_2, W_{i3} = w_3, W_{i4} = w_4) \\
&\ge P(Y_{m(1,d_2,d_3)} = 1 | Z_m, X_{m2} = x_2, X_{m3} = x_3, W_{m1} = w_1, W_{m2} = w_2, W_{m3} = w_3, W_{m4} = w_4) \}.
\end{aligned}$$

This refers to models with J=2 or equivalently $\mathcal{J}=\{0,1,2,(1,2)\}$ that we discussed in Sections 2 and 3.

Similar moment inequalities can be obtained for $X'_{ij}\beta \geq X'_{mj}\beta$ by fixing $\{X_1, X_2, X_3\} \setminus \{X_j\}$ and W_1, W_2, W_3, W_4 . Collectively, these moment inequalities establish the identification of β , with some standard technical conditions as for the two-alternatives case.

Once β is identified, we can proceed to identify γ 's by fixing $X'_j\beta$, j=1,2,3, or simply by fixing X_1, X_2, X_3 . We follow the former strategy. For γ_1 , we fix $(X'_1\beta, X'_2\beta, X'_3\beta)$ at some constant vector (v_1, v_2, v_3) and fix (W_{i2}, W_{i3}, W_{i4}) at some constant vector (w_2, w_3, w_4) . Then by $(\epsilon_1, \epsilon_2, \epsilon_3, \eta) \perp Z$,

$$P(Y_{(1,1,d_3)} = 1 | Z, X_1'\beta = v_1, X_2'\beta = v_2, X_3'\beta = v_3, W_2 = w_2, W_3 = w_3, W_4 = w_4)$$

is increasing in $W'_1\gamma_1$ for any d_3 . As a result, for any d_3 , (v_1, v_2, v_3) , and (w_2, w_3, w_4) , we have

$$\begin{aligned}
&\{W'_{i1}\gamma_{1} \geq W'_{m1}\gamma_{1}\} \\
\Leftrightarrow &\{P(Y_{i(1,1,d_{3})} = 1 | Z_{i}, X'_{i1}\beta = v_{1}, X'_{i2}\beta = v_{2}, X'_{i3}\beta = v_{3}, W_{i2} = w_{2}, W_{i3} = w_{3}, W_{i4} = w_{4}) \\
\geq &P(Y_{m(1,1,d_{3})} = 1 | Z_{m}, X'_{m1}\beta = v_{1}, X'_{m2}\beta = v_{2}, X'_{m3}\beta = v_{3}, W_{m2} = w_{2}, W_{m3} = w_{3}, W_{m4} = w_{4}) \}.
\end{aligned}$$

We can similarly identify γ_2 and γ_3 .

Once β , γ_1 , γ_2 , and γ_3 are identified, we can identify γ_4 by fixing either $(X_1, X_2, X_3, W_1, W_2, W_3)$ or $(X_1'\beta, X_2'\beta, X_3'\beta, W_1'\gamma_1, W_2'\gamma_2, W_3'\gamma_3)$. We consider the latter and fix $(X_1'\beta, X_2'\beta, X_3'\beta, W_1'\gamma_1, W_2'\gamma_2, W_3'\gamma_3)$ at some constant vector $(v_1, v_2, v_3, v_4, v_5, v_6)$. Then $(\epsilon_1, \epsilon_2, \epsilon_3, \eta) \perp Z$ implies that

$$P\left(Y_{(1,1,1)} = 1 | Z, X_1'\beta = v_1, X_2'\beta = v_2, X_3'\beta = v_3, W_1'\gamma_1 = v_4, W_2'\gamma_2 = v_5, W_3'\gamma_3 = v_6\right)$$

is increasing in $W'_4\gamma_4$ for any fixed $(v_1, v_2, v_3, v_4, v_5, v_6)$. Thus, the following inequality holds for any $(v_1, v_2, v_3, v_4, v_5, v_6)'$,

$$\{W'_{i4}\gamma_{4} \geq W'_{m4}\gamma_{4}\}$$

$$\Leftrightarrow \{P(Y_{i(1,1,1)} = 1 | Z_{i}, X'_{i1}\beta = v_{1}, X'_{i2}\beta = v_{2}, X'_{i3}\beta = v_{3}, W'_{i1}\gamma_{1} = v_{4}, W'_{i2}\gamma_{2} = v_{5}, W'_{i3}\gamma_{3} = v_{6})$$

$$\geq P(Y_{m(1,1,1)} = 1 | Z_{m}, X'_{m1}\beta = v_{1}, X'_{m2}\beta = v_{2}, X'_{m3}\beta = v_{3}, W'_{m1}\gamma_{1} = v_{4}, W'_{m2}\gamma_{2} = v_{5}, W'_{m3}\gamma_{3} = v_{6})\}.$$

The inequalities established in equations (E.2)–(E.4) can be used to construct localized MRC estimators as in Section 2.1. However, this matching-based identification strategy cannot identify coefficients on common regressors, i.e., covariates that appear in more than one index.

E.3.2 LAD Method

The LAD estimation for $\theta := (\beta, \gamma_1, \gamma_2, \gamma_3, \gamma_4)$ is analogous to the simple case presented in the main text. Here, we only outline the procedure for the cross-sectional model for illustration purposes. The panel data LAD estimation can be constructed similarly and is omitted for brevity.²²

1. By $(\epsilon_1, \epsilon_2, \epsilon_3, \eta) \perp Z$ and the monotonicity of $F_j(\cdot)$'s and $F_d(\cdot)$'s functions, we can write

$$\{X'_{im1}\beta \ge 0, X'_{im2}\beta \le 0, X'_{im3}\beta \le 0, W'_{im1}\gamma_1 \le 0, W'_{im2}\gamma_2 \le 0, W'_{im3}\gamma_3, \le 0, W'_{im4}\gamma_4 \le 0\}$$
(E.5)

 \Rightarrow

$$\Delta p_{(1,0,0)}(Z_i, Z_m) := P(Y_{i(1,0,0)} = 1|Z_i) - P(Y_{m(1,0,0)} = 1|Z_i) \ge 0$$
(E.6)

and

$$\{X'_{im1}\beta \le 0, X'_{im2}\beta \ge 0, X'_{im3}\beta \ge 0, W'_{im1}\gamma_1 \ge 0, W'_{im2}\gamma_2 \ge 0, W'_{im3}\gamma_3, \ge 0, W'_{im4}\gamma_4 \ge 0\}$$
(E.7)

 \Rightarrow

$$\Delta p_{(1,0,0)}(Z_i, Z_m) \le 0,$$
 (E.8)

i.e., the event (E.5) and (E.7) are respectively sufficient (but not necessary) conditions for inequality (E.6) and (E.8) to hold.

2. Define

$$Q_{(1,0,0)}(\vartheta) := \mathbb{E}[q_{(1,0,0)}(Z_i, Z_m, \vartheta)],$$

where

$$\begin{aligned} &q_{(1,0,0)}(Z_i,Z_m,\vartheta) \\ = &[|I_{im(1,0,0)}^+(\vartheta) - 1[\Delta p_{(1,0,0)}(Z_i,Z_m) \geq 0]| + |I_{im(1,0,0)}^-(\vartheta) - 1[\Delta p_{(1,0,0)}(Z_i,Z_m) \leq 0]|] \\ &\times [I_{im(1,0,0)}^+(\vartheta) + I_{im(1,0,0)}^-(\vartheta)] \end{aligned}$$

²²For the panel data setting, the identification relies on the conditional homogeneity condition $\xi_s \stackrel{d}{=} \xi_t | (\alpha, Z_s, Z_t)$.

with

$$\begin{split} &I^{+}_{im(1,0,0)}(\vartheta)\\ =&1[X'_{im1}\beta\geq 0,X'_{im2}\beta\leq 0,X'_{im3}\beta\leq 0,W'_{im1}\gamma_{1}\leq 0,W'_{im2}\gamma_{2}\leq 0,W'_{im3}\gamma_{3},\leq 0,W'_{im4}\gamma_{4}\leq 0]\\ &I^{-}_{im(1,0,0)}(\vartheta)\\ =&1[X'_{im1}\beta\leq 0,X'_{im2}\beta\geq 0,X'_{im3}\beta\geq 0,W'_{im1}\gamma_{1}\geq 0,W'_{im2}\gamma_{2}\geq 0,W'_{im3}\gamma_{3},\geq 0,W'_{im4}\gamma_{4}\geq 0]. \end{split}$$

The intuition underlying the construction of this population criterion function is the same as the discussion in Section 2.2 (below (2.13)–(2.15)). Then the identification of θ can be established using arguments similar to those for Theorem 2.5.

3. To compute the estimator $\hat{\theta}$ for θ , one can consider the "debiased" sample criterion analogous to (2.17), i.e.,

$$\hat{\theta} = \arg\min_{\vartheta \in \Theta} \sum_{i=1}^{N-1} \sum_{m>i} \hat{q}_{im(1,0,0)}^{D}(\vartheta), \tag{E.9}$$

where

$$\hat{q}_{im(1,0,0)}^{D}(\vartheta) = [|I_{im(1,0,0)}^{+}(\vartheta) - \Delta \hat{p}_{(1,0,0)}(Z_i, Z_m)| + |I_{im(1,0,0)}^{-}(\vartheta) + \Delta \hat{p}_{(1,0,0)}(Z_i, Z_m)|] \times [I_{im(1,0,0)}^{+}(\vartheta) + I_{im(1,0,0)}^{-}(\vartheta)] + [1 - (I_{im(1,0,0)}^{+}(\vartheta) + I_{im(1,0,0)}^{-}(\vartheta))].$$

4. Repeat Steps 1–3 to obtain sample criteria associated with $\Delta p_{(0,1,0)}(Z_i, Z_m)$, $\Delta p_{(0,0,1)}(Z_i, Z_m)$, $\Delta p_{(1,1,0)}(Z_i, Z_m)$, $\Delta p_{(1,1,0)}(Z_i, Z_m)$, $\Delta p_{(1,1,0)}(Z_i, Z_m)$, $\Delta p_{(0,1,1)}(Z_i, Z_m)$, $\Delta p_{(0,0,0)}(Z_i, Z_m)$, and $\Delta p_{(1,1,1)}(Z_i, Z_m)$. Collectively, use some convex combination of these criteria to obtain $\hat{\theta}$ with better finite sample performance.

E.3.3 Panel Data MS Method

We specify the latent utility of the panel data bundle choice model with J=3 as follows:

$$U_{dt} = \sum_{j=1}^{3} F_{j}(X'_{jt}\beta, \epsilon_{jt}, \alpha_{j}) \cdot d_{j} + \eta_{110t} \cdot F_{110} (W'_{1t}\gamma_{1} + \alpha_{b1}) \cdot d_{1} \cdot d_{2} + \eta_{101t} \cdot F_{101} (W'_{2t}\gamma_{2} + \alpha_{b2}) \cdot d_{1} \cdot d_{3}$$
$$+ \eta_{011t} \cdot F_{011} (W'_{3t}\gamma_{3} + \alpha_{b3}) \cdot d_{2} \cdot d_{3} + \eta_{111t} \cdot F_{111} (W'_{4t}\gamma_{4} + \alpha_{b4}) \cdot d_{1} \cdot d_{2} \cdot d_{3},$$

where $X_{1t}, X_{2t}, X_{3t} \in \mathbb{R}^{k_1}$, $W_{1t} \in \mathbb{R}^{k_2}, W_{2t} \in \mathbb{R}^{k_3}, W_{3t} \in \mathbb{R}^{k_4}$, $W_{4t} \in \mathbb{R}^{k_5}$, and $\eta_t \equiv (\eta_{110t}, \eta_{101t}, \eta_{011t}, \eta_{111t}) \in \mathbb{R}^4_+$. All $F_j(\cdot)$'s and $F_d(\cdot)$'s are strictly increasing in all arguments. An agent chooses d that maximizes the latent utility, i.e.,

$$Y_{dt} = 1[U_{dt} > U_{d't}, \forall d' \in \mathcal{D} \setminus d].$$

Denote $Z_t = (X_{1t}, X_{2t}, X_{3t}, W_{1t}, W_{2t}, W_{3t}, W_{4t})$, $\xi_t = (\epsilon_{1t}, \epsilon_{2t}, \epsilon_{3t}, \eta_{110t}, \eta_{101t}, \eta_{111t})$, and $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_{b1}, \alpha_{b2}, \alpha_{b3}, \alpha_{b4})$. The key identification assumption is that $\xi_s \stackrel{d}{=} \xi_t | (\alpha, Z_s, Z_t)$ holds for any two time periods s and t.

The identification of the panel data model is similar to the cross-sectional model discussed in Section E.3.1. The primary difference is that in a panel data setting, we need to match and compare the same agent in time periods s and t, rather than two independent agents i and m. As a result, we can construct the same set of identification inequalities as Section E.3.1, but replace the (i, m) therein with (s, t). For example, analogous to (E.2) and (E.3), we can write for any $d_2, d_3, x_1, x_2, x_3, w_1, w_2, w_3, w_4$, and c

$$\{X'_{1t}\beta \ge X'_{1s}\beta\}$$

$$\Leftrightarrow \{P(Y_{(1,d_2,d_3)t} = 1 | Z_t, X_{2t} = x_2, X_{3t} = x_3, W_{1t} = w_1, W_2 = w_{2t}, W_{3t} = w_3, W_{4t} = w_4, \alpha = c)$$

$$\ge P(Y_{(1,d_2,d_3)s} = 1 | Z_s, X_{2s} = x_2, X_{3s} = x_3, W_{1s} = w_1, W_{2s} = w_2, W_{3s} = w_3, W_{4s} = w_4, \alpha = c)\}$$

and

$$\{W'_{1t}\gamma_1 \ge W'_{1s}\gamma_1\}$$

$$\Leftrightarrow P(Y_{(1,1,d_3)t} = 1 | Z_t, X_{1t} = x_1, X_{2t} = x_2, X_{3t} = x_3, W_{2t} = w_2, W_{3t} = w_3, W_{4t} = w_4, \alpha = c)$$

$$\ge P(Y_{(1,1,d_3)s} = 1 | Z_s, X_{1s} = x_1, X_{2s} = x_2, X_{3s} = x_3, W_{2s} = w_2, W_{3s} = w_3, W_{4s} = w_4, \alpha = c).$$

Then, under similar regular conditions, we can identify and estimate (β, γ_1) based on inequalities (E.10) and (E.11). Identification inequalities for $(\gamma_2, \gamma_3, \gamma_4)$ can be similarly constructed and hence are omitted for brevity.

The panel data MS estimator differs from the cross-sectional MRC estimator in that its convergence rate decreases as the number of matching variables increases. For example, using similar analysis, we can show that for MS estimators based on inequalities (E.10) and (E.11), we have

$$\hat{\beta} - \beta = O_p \left((Nh_N^{2k_1 + k_2 + k_3 + k_4 + k_5})^{-1/3} \right) \text{ and } \hat{\gamma}_1 - \gamma_1 = O_p \left((N\sigma_N^{3k_1 + k_3 + k_4 + k_5})^{-1/3} \right),$$

where we assume we use the same type of estimator as in Section 3.1 and the same bandwidth $(h_N \text{ or } \sigma_N)$ for all covariates. The convergence rates of $\hat{\gamma}_2, \hat{\gamma}_3$, and $\hat{\gamma}_4$ can be similarly obtained. Generally speaking, we can show that the MS estimator has a convergence rate of $O_p((Nh_N^{\bar{k}})^{-1/3})$ $(O_p((N\sigma_N^{\bar{k}})^{-1/3}))$, where \bar{k} represents the number of matching variables.

F Proofs of Technical Lemmas

We prove Lemmas B.1–B.10 and Lemmas C.1–C.2 in order. For Lemmas B.1–B.7, we use the following results, which can be derived under Assumptions C2 and C5 by applying standard arguments (e.g., Example 6.4 of Kim and Pollard (1990)). We use $\|\cdot\|_F$ to denote the Frobenius norm; that is, for any matrix \mathbf{A} , $\|\mathbf{A}\|_F = \sqrt{\operatorname{trace}(\mathbf{A}\mathbf{A}')}$.

Let \mathcal{N}_{β} denote a neighborhood of β . Then, for $\varrho_i(b)$ defined in (B.1),

- All mixed second partial derivatives of $\varrho_i(b)$ exist on \mathcal{N}_{β} .
- There is an integrable function $M_1(Z_i)$ such that for all $b \in \mathcal{N}_{\beta}$, $\|\nabla^2 \varrho_i(b) \nabla^2 \varrho_i(\beta)\|_F \le M_1(Z_i)\|b \beta\|$.
- $\mathbb{E}[\|\nabla \varrho_i(\beta)\|^2] < \infty$, $\mathbb{E}[\|\nabla^2 \varrho_i(\beta)\|_F] < \infty$, and $\mathbb{E}[\nabla^2 \varrho_i(\beta)]$ is negative definite.

Let \mathcal{N}_{γ} denote a neighborhood of γ . Then, for $\tau_i(r)$ and $\mu(v_1, v_2, r)$ defined respectively in (B.2) and (B.3),

- All mixed second partial derivatives of $\tau_i(r)$ exist on \mathcal{N}_{γ} .
- There is an integrable function $M_2(Z_i)$ such that for all $r \in \mathcal{N}_{\gamma} \|\nabla^2 \tau_i(r) \nabla^2 \tau_i(\gamma)\|_F \le M_2(Z_i) \|r \gamma\|$.
- $\mathbb{E}[\|\nabla \tau_i(\gamma)\|^2] < \infty$, $\mathbb{E}[\|\nabla^2 \tau_i(\gamma)\|_F] < \infty$, and $\mathbb{E}[\nabla^2 \tau_i(\gamma)]$ is negative definite.
- All mixed second partial derivatives of $\mu(v_1, v_2, r)$ exist on \mathcal{N}_{γ} .
- There exists an continuous, integrable function $M_3(v_1, v_2)$ such that for all $r \in \mathcal{N}_{\gamma}$, $\|\nabla^2_{11}\mu(v_1, v_2, r) \nabla^2_{11}\mu(v_1, v_2, \gamma)\|_F \leq M_3(v_1, v_2)\|r \gamma\|$.
- $\mathbb{E}[\|\nabla_{13}^{2}\mu(V_{i}\left(\beta\right),V_{i}\left(\beta\right),\gamma)\|_{F}^{2}] < \infty \text{ and } \sup_{r \in \mathcal{N}_{\gamma}} \mathbb{E}[\|\nabla_{33}^{2}\nabla_{1}\mu(V_{i}\left(\beta\right),V_{i}\left(\beta\right),\gamma)\|_{F}] < \infty.$

Further, using the definition of $\tau_i(r)$, we can write $\tau_i(r)$ as

$$\tau_i(r) = f_{V_{im}}(0)\mathbb{E}[h_{im}(r)|V_i = v_i, W_i = w_i, V_{im} = 0] = \int B(v_i, v_i, w_i, w_m)S_{im}(r)f_{V,W}(v_i, w_m)dw_m.$$

Proof of Lemma B.1. First, note that

$$\frac{1}{2}\sigma_N^2 \sup_{r \in \mathcal{R}} |\hat{\mathcal{L}}_N^K(r) - \mathcal{L}_N^K(r)| \le \frac{1}{N(N-1)} \sum_{i \ne m} |K_{\sigma_N,\gamma}(\hat{V}_{im}) - K_{\sigma_N,\gamma}(V_{im})|.$$
 (F.1)

Applying a second-order Taylor expansion to the right-hand side of (F.1) yields the lead term of the form

$$\frac{1}{\sigma_N N(N-1)} \sum_{i \neq m} |\nabla_1 K_{\sigma_N,\gamma}(V_{im}) X'_{im1}(\hat{\beta} - \beta) + \nabla_2 K_{\sigma_N,\gamma}(V_{im}) X'_{im2}(\hat{\beta} - \beta)|$$

$$\leq \frac{1}{\sigma_N N(N-1)} \sum_{i \neq m} \sum_{j=1}^2 |\nabla_j K_{\gamma}(V_{im}/\sigma_N) X'_{imj}(\hat{\beta} - \beta)|,$$

where $\nabla_j K_{\gamma}(\cdot)$ ($\nabla_j K_{\sigma_N,\gamma}(\cdot)$) denotes the partial derivative of $K_{\gamma}(\cdot)$ ($K_{\sigma_N,\gamma}(\cdot)$) w.r.t. its j-th argument corresponding to alternative j. It suffices to focus on the term with j = 1. Then by Assumptions C1–C7, and the Cauchy-Schwarz inequality, we conclude that

$$\frac{1}{N(N-1)} \sum_{i \neq m} \left| \nabla_1 K_{\gamma}(V_{im}/\sigma_N) \frac{X'_{im1}(\hat{\beta} - \beta)}{\sigma_N} \right| \leq \frac{1}{N(N-1)} \sum_{i \neq m} \left| \nabla_1 K_{\gamma}(V_{im}/\sigma_N) \right| \|X_{im1}\| \frac{\|\hat{\beta} - \beta\|}{\sigma_N} \\
= [O_p(\sigma_N^2) + o_p(1)] O_p(\|\hat{\beta} - \beta\|/\sigma_N) = o_p(1),$$

where the penultimate equality follows from the SLLN of U-statistics (see e.g., Serfling (2009)) and $\mathbb{E}[|\nabla_1 K_{\gamma}(V_{im}/\sigma_N)|||X_{im1}||] = O_p(\sigma_N^2)$, and the last equality is due to the \sqrt{N} -consistency of $\hat{\beta}$. Then the desired convergence result follows.

Proof of Lemma B.2. Note that Lemma B.2 follows from $\sup_{r \in \mathcal{R}} |\mathcal{L}_N^K(r) - \mathbb{E}[\mathcal{L}_N^K(r)]| = o_p(1)$ and $\sup_{r \in \mathcal{R}} |\mathbb{E}[\mathcal{L}_N^K(r)] - \mathcal{L}(r)| = o(1)$.

Define $\mathcal{F}_N = \{K_{\sigma_N,\gamma}(v_{im})h_{im}(r)|r \in \mathcal{R}\}$ with $\sigma_N > 0$ and $\sigma_N \to 0$. \mathcal{F}_N is a sub-class of the fixed class $\mathcal{F} \equiv \{K_{\gamma}(v_{im}/\sigma)h_{im}(r)|r \in \mathcal{R}, \sigma > 0\} = \mathcal{F}_r\mathcal{F}_\sigma$, with $\mathcal{F}_\sigma \equiv \{K_{\gamma}(v_{im}/\sigma)|\sigma > 0\}$, which is Euclidean for the constant envelope $\sup_{v \in \mathbb{R}^2} |K_{\gamma}(v)|$ by Lemma 22 in Nolan and Pollard (1987) and $\mathcal{F}_r \equiv \{h_{im}(r)|r \in \mathcal{R}\}$. Noticing that $h_{im}(r)$ is uniformly bounded by 2, Example 2.11 and Lemma 2.15 in Pakes and Pollard (1989) then implies that \mathcal{F}_r is Euclidean for the constant envelope 2. Putting all these results together, we conclude using Lemma 2.14 in

Pakes and Pollard (1989) that \mathcal{F} is Euclidean for the constant envelope $2 \sup_{v \in \mathbb{R}^2} |K_{\gamma}(v)|$. Applying Corollary 4 in Sherman (1994a), we obtain that $\sup_{r \in \mathcal{R}} |\mathcal{L}_N^K(r) - \mathbb{E}[\mathcal{L}_N^K(r)]| = O_p(N^{-1}/\sigma_N^2) = o_p(1)$ by Assumption C7.

It remains to show that $\sup_{r \in \mathcal{R}} |\mathbb{E}[\mathcal{L}_N^K(r)] - \mathcal{L}(r)| = o(1)$. Let $\eta(\cdot) \equiv f_{V_{im}}(\cdot)\mathbb{E}[h_{im}(r)|V_{im} = \cdot]$. Then $\mathcal{L}(r) = \eta(0)$ by definition. We can write by Assumptions C6 and C7 that

$$\sup_{r \in \mathcal{R}} |\mathbb{E}[\mathcal{L}_{N}^{K}(r)] - \mathcal{L}(r)| = \sup_{r \in \mathcal{R}} |\sigma_{N}^{-2}\mathbb{E}[K_{\gamma}(V_{im}/\sigma_{N})h_{im}(r)] - \eta(0)|$$

$$= \sup_{r \in \mathcal{R}} |\int \sigma_{N}^{-2}K_{\gamma}(v/\sigma_{N})\eta(v)dv - \eta(0)| = \sup_{r \in \mathcal{R}} |\int \sigma_{N}^{-2}K_{\gamma}(v/\sigma_{N})[\eta(0) + v'\nabla\eta(\bar{v})]dv - \eta(0)|$$

$$= \sup_{r \in \mathcal{R}} |\int K_{\gamma}(u)[\eta(0) + \sigma_{N}u'\nabla\eta(u_{N})]du - \eta(0)| = \sup_{r \in \mathcal{R}} |\sigma_{N}\int K_{\gamma}(u)u'\nabla\eta(u_{N})du|$$

$$\leq \sigma_{N} \cdot \sup_{r \in \mathcal{R}} \int |K_{\gamma}(u)|||u|| ||\nabla\eta(u_{N})||du = O(\sigma_{N}) = o(1),$$

where the third equality applies a mean-value expansion and the fourth equality uses a change of variables. Then, the desired result follows. \Box

Proof of Lemma B.3. The proof proceeds by verifying the four sufficient conditions for Theorem 2.1 in Newey and McFadden (1994): (S1) \mathcal{R} is a compact set, (S2) $\sup_{r \in \mathcal{R}} |\hat{\mathcal{L}}_N^K(r) - \mathcal{L}(r)| = o_p(1)$, (S3) $\mathcal{L}(r)$ is continuous in r, and (S4) $\mathcal{L}(r)$ is uniquely maximized at γ .

Condition (S1) is satisfied by Assumption C4. Lemmas B.1 and B.2 together imply that condition (S2) holds. Note that

$$\mathbb{E}[h_{im}(r)|V_{im}=0] = \mathbb{E}\left\{\mathbb{E}[Y_{im(1,1)}(\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma))|Z_i, Z_m]|V_{im}=0\right\}$$

$$=\mathbb{E}\left[(P(Y_{i(1,1)}=1|Z_i) - P(Y_{m(1,1)}=1|Z_m))(\operatorname{sgn}(W'_{im}r) - \operatorname{sgn}(W'_{im}\gamma))|V_{im}=0\right].$$

Then the identification condition (S4) can be verified by similar arguments used in the proof of Theorem 2.1 given that $f_{V_{im}}(0) > 0$ by Assumption C5.

It remains to verify the continuity of $\mathcal{L}(r)$ in r. $\mathbb{E}[h_{im}(r)|V_{im}=0]$ can be expressed as the sum of terms like

$$P(Y_{im(1,1)} = d, W_{im}^{(1)} > -\tilde{W}'_{im}\tilde{r}|V_{im} = 0)$$

$$= \int \int_{-\tilde{W}'_{im}\tilde{r}/r^{(1)}}^{\infty} P(Y_{im(1,1)} = d|W_{im}^{(1)} = w, \tilde{W}_{im}, V_{im} = 0) f_{W_{im}^{(1)}|\tilde{W}_{im}, V_{im} = 0}(w) dw dF_{\tilde{W}_{im}|V_{im} = 0}.$$

for some $d \in \{-1, 0, 1\}$. Then $\mathcal{L}(r)$ is continuous in r if $f_{W_{im}^{(1)}|\tilde{W}_{im}, V_{im}=0}(\cdot)$ does not have any

mass points, which is guaranteed by Assumption C3. This completes the proof.

Proof of Lemma B.4. First, we express $\mathbb{E}[\mathcal{L}_N^K(r)]$ as the integral

$$\int \sigma_{N}^{-2} K_{\sigma_{N},\gamma}(v_{im}) B(v_{i}, v_{m}, w_{i}, w_{m}) S_{im}(r) dF_{V,W}(v_{i}, w_{i}) dF_{V,W}(v_{m}, w_{m})
= \int K_{\gamma}(u_{im}) B(v_{m} + u_{im}\sigma_{N}, v_{m}, w_{i}, w_{m}) S_{im}(r) f_{V,W}(v_{m} + u_{im}\sigma_{N}, w_{i}) dw_{i} du_{im} dF_{V,W}(v_{m}, w_{m}),$$
(F.2)

where we apply the change of variables $u_{im} = v_{im}/\sigma_N$ to obtain the equality.

Take the $\kappa_{\gamma}^{\text{th}}$ -order Taylor expansion inside the integral around v_m to obtain the lead term

$$\int K_{\gamma}(u_{im})B(v_{m}, v_{m}, w_{i}, w_{m})S_{im}(r)f_{V,W}(v_{m}, w_{i})dw_{i}du_{im}dF_{V,W}(v_{m}, w_{m})$$

$$= \int B(v_{m}, v_{m}, w_{i}, w_{m})S_{im}(r)f_{V,W}(v_{m}, w_{i})dw_{i}dF_{V,W}(v_{m}, w_{m}) = \mathbb{E}[\tau_{m}(r)], \qquad (F.3)$$

where the first equality follows by $\int K_{\gamma}(u)du = 1$.²³ All remaining terms are zero except the last one which is of order $O(\sigma_N^{\kappa_{\gamma}}\delta_N) = o(\|r - \gamma\|^2)$ by Assumptions C6 and C7.

Note that a second-order Taylor expansion around γ gives

$$\tau_m(r) - \tau_m(\gamma) = (r - \gamma)' \nabla \tau_m(\gamma) + \frac{1}{2} (r - \gamma)' \nabla^2 \tau_m(\bar{r}) (r - \gamma)$$
$$= (r - \gamma)' \nabla \tau_m(\gamma) + \frac{1}{2} (r - \gamma)' \nabla^2 \tau_m(\gamma) (r - \gamma) + \frac{1}{2} (r - \gamma)' \left[\nabla^2 \tau_m(\bar{r}) - \nabla^2 \tau_m(\gamma) \right] (r - \gamma),$$

and hence by Assumption C7

$$\mathbb{E}[\tau_m(r)] = (r - \gamma)' \mathbb{E}[\nabla \tau_m(\gamma)] + \frac{1}{2}(r - \gamma)' \mathbb{E}[\nabla^2 \tau_m(\gamma)](r - \gamma) + o(\|r - \gamma\|^2)$$

$$= \frac{1}{2}(r - \gamma)' \mathbb{V}_{\gamma}(r - \gamma) + o(\|r - \gamma\|^2), \tag{F.4}$$

where the second equality uses the fact that $\mathbb{E}[\nabla \tau_m(\gamma)] = 0$ since $\mathbb{E}[\tau_m(r)]$ is maximized at γ . Then, applying (F.2), (F.3), and (F.4) proves the lemma.

Proof of Lemma B.5. We can establish a representation for $\mathbb{E}[\mathcal{L}_N^K(r)|Z_m]$ using the same arguments as in proving Lemma B.4, but no longer integrating over Z_m . Specifically, a

²³More precisely, we apply $\kappa_{1\gamma}^{\text{th}}$ -order and $\kappa_{2\gamma}^{\text{th}}$ -order Taylor expansions to $B(\cdot, v_m, w_i, w_m)$ and $f_{V,W}(\cdot, w_i)$, respectively.

change of variables $u_{im} = v_{im}/\sigma_N$ gives

$$\mathbb{E}[\mathcal{L}_{N}^{K}(r)|Z_{m} = z_{m}] = \int \sigma_{N}^{-2} K_{\sigma_{N},\gamma}(v_{im}) B(v_{i}, v_{m}, w_{i}, w_{m}) S_{im}(r) dF_{V,W}(v_{i}, w_{i})$$

$$= \int K_{\gamma}(u_{im}) B(v_{m} + u_{im}\sigma_{N}, v_{m}, w_{i}, w_{m}) S_{im}(r) f_{V,W}(v_{m} + u_{im}\sigma_{N}, w_{i}) dw_{i} du_{im}.$$

The lead term of the $\kappa_{\gamma}^{\text{th}}$ -order Taylor expansion inside the integral around v_m is

$$\int K_{\gamma}(u_{im})B(v_m, v_m, w_i, w_m)S_{im}(r)f_{V,W}(v_m, w_i)dw_idu_{im}$$

$$= \int B(v_m, v_m, w_i, w_m)S_{im}(r)f_{V,W}(v_m, w_i)dw_i = \tau_m(r)$$

by $\int K_{\gamma}(u)du = 1$, and the sample average of the bias term is of order $o(||r - \gamma||^2)$.

Then, applying Lemma B.4 and Assumption C7, we can write

$$\frac{2}{N} \sum_{m} \mathbb{E}[\mathcal{L}_{N}^{K}(r)|Z_{m}] - 2\mathbb{E}[\mathcal{L}_{N}^{K}(r)]$$

$$= \frac{2}{N} \sum_{m} \tau_{m}(r) - (r - \gamma)' \mathbb{V}_{\gamma}(r - \gamma) + o_{p}(\|r - \gamma\|^{2})$$

$$= \frac{2}{N} \sum_{m} (r - \gamma)' \nabla \tau_{m}(\gamma) + \frac{1}{N} \sum_{m} (r - \gamma)' \left(\nabla^{2} \tau_{m}(\gamma) - \mathbb{V}_{\gamma}\right) (r - \gamma) + o_{p}(\|r - \gamma\|^{2}).$$

Then the desired result follows by noticing that $N^{-1}\sum_{m}\nabla^{2}\tau_{m}(\gamma)-\mathbb{V}_{\gamma}=o_{p}(1)$ by the SLLN.

Proof of Lemma B.6. By construction, we can write

$$\rho_N(r) = \frac{1}{N(N-1)} \sum_{i \neq m} \rho_{im}(r),$$

where $\rho_{im}(r) \equiv (K_{\sigma_N,\gamma}(V_{im})h_{im}(r) - 2N^{-1}\sum_i \mathbb{E}[K_{\sigma_N,\gamma}(V_{im})h_{im}(r)|Z_i] + \mathbb{E}[K_{\sigma_N,\gamma}(V_{im})h_{im}(r)])/\sigma_N^2$.

Note that $\rho_{im}(\gamma) = 0$ and $|\rho_{im}(r)|$ is bounded by a multiple of M/σ_N^2 where M is a positive constant. Define $\mathcal{F}_N^* = \{\rho_{im}^*(r)|r \in \mathcal{R}\}$ where $\rho_{im}^*(r) \equiv \sigma_N^2 \rho_{im}(r)/M$. Then the Euclidean properties of the (P-degenerate) class of functions \mathcal{F}_N^* are deduced using similar arguments for proving Lemma B.2 in combination with Corollary 17 and Corollary 21 in Nolan and Pollard (1987). As $\int K_{\sigma_N,\gamma}(v)^p dv = O(\sigma_N^2)$ for p = 1, 2, we have

 $\sup_{r\in\mathcal{R}_N} \mathbb{E}[\rho_{im}^*(r)^2] = O(\sigma_N^2)$. Then applying Theorem 3 of Sherman (1994b) gives

$$\frac{1}{N(N-1)} \sum_{i \neq m} \rho_{im}^*(r) = O_p(N^{-1}\sigma_N^{\alpha}),$$

where $0 < \alpha < 1$ and hence $q_N(r) = O_p(N^{-1}\sigma_N^{\alpha-2}) = O_p(N^{-1}\sigma_N^{-2})$.

Proof of Lemma B.7. The first step is to plug (B.8) into (B.7) so that $\Delta \mathcal{L}_N^K(r)$ can be expressed as

$$\frac{1}{\sigma_N^3 N(N-1)(N-2)} \sum_{i \neq m \neq l} [\nabla_1 K_{\sigma_N,\gamma}(V_{im}) X'_{im1} \psi_{l,\beta} + \nabla_2 K_{\sigma_N,\gamma}(V_{im}) X'_{im2} \psi_{l,\beta}] h_{im}(r) \quad (F.5)$$

plus a remainder term of higher order since $\sqrt{N}\sigma_N \to \infty$. (F.5) is a third order U-process. We use the U-statistic decomposition found in Sherman (1994b) (see also Serfling (2009)) to derive a linear representation. Note that (F.5) has unconditional mean zero, so as is its mean conditional on each of its first two arguments (Z_i, Z_m) . Furthermore, by Theorem 3 of Sherman (1994b) and similar arguments for proving Lemma B.6, the remainder term of the decomposition (projection) is of order $O_p(N^{-1}\sigma_N^{-2})$. Hence it suffices to derive a linear representation for its mean conditional on its third argument:

$$\frac{1}{\sigma_N^3 N} \sum_{l} \int B(v_i, v_m, w_i, w_m) S_{im}(r) \nabla K_{\sigma_N, \gamma}(v_{im})' \begin{pmatrix} x'_{im1} \psi_{l,\beta} \\ x'_{im2} \psi_{l,\beta} \end{pmatrix} dF_{X,W}(x_i, w_i) dF_{X,W}(x_m, w_m).$$

where $F_{X,W}(\cdot,\cdot)$ denotes the joint distribution function of (X_i,W_i) . While the above integral is expressed w.r.t. (x_i,w_i) and (x_m,w_m) , it will prove convenient to express the integral in terms of v_i and v_m . We do so as follows:

$$\frac{1}{\sigma_N^3 N} \sum_{l} \left(\int \nabla K_{\sigma_N,\gamma}(v_{im})' G(v_i, v_m, r) f_V(v_i) f_V(v_m) dv_i dv_m \right) \psi_{l,\beta}.$$
 (F.6)

Apply a change of variables in (F.6) with $u_{im} = v_{im}/\sigma_N$ to obtain

$$\frac{1}{\sigma_N^3 N} \sum_{l} \left(\int \nabla K_{\sigma_N,\gamma}(v_{im})' G(v_i, v_m, r) f_V(v_i) f_V(v_m) dv_i dv_m \right) \psi_{l,\beta}$$

$$= \frac{1}{\sigma_N N} \sum_{l} \left(\int \nabla K_{\gamma}(u_{im})' G(v_m + u_{im}\sigma_N, v_m, r) f_V(v_m + u_{im}\sigma_N) f_V(v_m) du_{im} dv_m \right) \psi_{l,\beta}.$$

As $\int \nabla K_{\gamma}(u)' du = 0$ and $\int u_j \nabla_j K_{\gamma}(u) du = -1$ for j = 1, 2, a second-order expansion inside

the integral around $u_{im}\sigma_N=0$ yields the lead term²⁴

$$-\frac{1}{N}\sum_{l}\left(\int \nabla_{1}\mu(v_{m},v_{m},r)f_{V}(v_{m})dv_{m}\right)\psi_{l,\beta}.$$

We next apply a second-order Taylor expansion of $\nabla_1 \mu(v_m, v_m, r)$ around γ to yield

$$(r-\gamma)'\frac{1}{N}\sum_{l}\left(-\int \nabla_{13}^{2}\mu(v_{m},v_{m},\gamma)f_{V}(v_{m})dv_{m}\right)\psi_{l,\beta}+o_{p}(\|r-\gamma\|^{2}),$$

which concludes the proof of this lemma.

Proof of Lemma B.8. Recall that $V_{im}(\beta) = (X'_{im1}\beta, X'_{im2}\beta)$. The following calculation is useful:

$$K_{\sigma_{N},\gamma}(V_{im}(\hat{\beta})) - K_{\sigma_{N},\gamma}(V_{im}(\beta))$$

$$= K\left(\frac{X'_{im1}\hat{\beta}}{\sigma_{N}}\right) K\left(\frac{X'_{im2}\hat{\beta}}{\sigma_{N}}\right) - K\left(\frac{X'_{im1}\beta}{\sigma_{N}}\right) K\left(\frac{X'_{im2}\beta}{\sigma_{N}}\right)$$

$$= K'\left(\frac{X'_{im1}\beta}{\sigma_{N}}\right) K\left(\frac{X'_{im2}\beta}{\sigma_{N}}\right) \frac{X'_{im1}(\hat{\beta} - \beta)}{\sigma_{N}}$$

$$+K\left(\frac{X'_{im1}\beta}{\sigma_{N}}\right) K'\left(\frac{X'_{im2}\beta}{\sigma_{N}}\right) \frac{X'_{im2}(\hat{\beta} - \beta)}{\sigma_{N}} + O_{p}(N^{-1}\sigma_{N}^{-2}), \tag{F.7}$$

where the last line applies the Taylor expansion. Suppose we have a random vector q_{im} such that $\mathbb{E}[q_{im}|V_{im}(\beta)=v]$ exists and is κ_{γ} -th differentiable in v, with derivatives up to κ_{γ} -th being uniformly bounded. We make the following claim

$$\frac{1}{N(N-1)} \sum_{i \neq m} K'\left(\frac{X'_{im1}\beta}{\sigma_N}\right) K\left(\frac{X'_{im2}\beta}{\sigma_N}\right) q_{im} = O_p(N^{-1/2}\sigma_N^2), \tag{F.8}$$

whose proof is deferred to the end. Using (F.8), we can write

$$\frac{1}{\sigma_N^2 N(N-1)} \sum_{i \neq m} K' \left(\frac{X'_{im1} \beta}{\sigma_N} \right) K \left(\frac{X'_{im2} \beta}{\sigma_N} \right) \frac{X'_{im1} (\hat{\beta} - \beta)}{\sigma_N} Y_{im(1,1)} \operatorname{sgn} \left(W'_{im} \gamma \right)$$

$$= \left[\frac{1}{\sigma_N^2 N(N-1)} \sum_{i \neq m} K' \left(\frac{X'_{im1} \beta}{\sigma_N} \right) K \left(\frac{X'_{im2} \beta}{\sigma_N} \right) X'_{im1} Y_{im(1,1)} \operatorname{sgn} \left(W'_{im} \gamma \right) \right] \frac{(\hat{\beta} - \beta)}{\sigma_N}$$

$$= O_p(N^{-1/2} (\hat{\beta} - \beta) \sigma_N^{-1}) = O_p(N^{-1} \sigma_N^{-1}), \tag{F.9}$$

The second-order term is of order $O_p(\delta_N \sigma_N/\sqrt{N})$, which will be $o_p(\|r-\gamma\|^2)$.

where we let $q_{im} = X'_{im1}Y_{im(1,1)}\operatorname{sgn}(W'_{im}\gamma)$ and use the result $\hat{\beta} - \beta = O_p(N^{-1/2})$. Clearly, this also holds if we replace $X'_{im1}(\hat{\beta} - \beta)$ with $X'_{im2}(\hat{\beta} - \beta)$.

Combine (F.7) and (F.9) to obtain

$$\frac{1}{\sigma_N^2 N(N-1)} \sum_{i \neq m} \left[K_{\sigma_N,\gamma}(V_{im}(\hat{\beta})) - K_{\sigma_N,\gamma}(V_{im}(\beta)) \right] Y_{im(1,1)} \operatorname{sgn}(W'_{im}\gamma)
= 2 \cdot O_p(N^{-1}\sigma_N^{-1}) + O_p(N^{-1}\sigma_N^{-2}) = O_p(N^{-1}\sigma_N^{-2}),$$

as desired.

The remaining task is to show the claim in (F.8). Some standard results on U-statistics imply that the variance of (F.8) is $O_p(N^{-1})$. So we only need show the expectation of the U-statistic is $O(N^{-1/2})$. Under Assumption C5, we can calculate the expectation of the summand of (F.8) as follows:

$$\mathbb{E}\left[K'\left(\frac{X'_{im1}\beta}{\sigma_N}\right)K\left(\frac{X'_{im2}\beta}{\sigma_N}\right)q_{im}\right] = \mathbb{E}\left[K'\left(\frac{X'_{im1}\beta}{\sigma_N}\right)K\left(\frac{X'_{im2}\beta}{\sigma_N}\right)\mathbb{E}\left(q_{im}|V_{im}(\beta)\right)\right]$$

$$= \int\int K'\left(\frac{v_1}{\sigma_N}\right)K\left(\frac{v_2}{\sigma_N}\right)\mathbb{E}\left[q_{im}|V_{im}(\beta) = (v_1, v_2)\right]f_{V_{im}(\beta)}(v_1, v_2)dv_1dv_2$$

$$= \sigma_N^2\int\int K'\left(u_1\right)K\left(u_2\right)\mathbb{E}\left[q_{im}|V_{im}(\beta) = (u_1\sigma_N, u_2\sigma_N)\right]f_{V_{im}(\beta)}(u_1\sigma_N, u_2\sigma_N)du_1du_2$$

$$= \sigma_N^2\mathbb{E}\left[q_{im}|V_{im}(\beta) = (0, 0)\right]f_{V_{im}(\beta)}(0, 0)\int\int K'\left(u_1\right)K\left(u_2\right)du_1du_2 + O_p(\sigma_N^4)$$

$$= 0 + O_p(N^{-1/2}) = O_p(N^{-1/2}),$$

where the last line holds by Assumptions C6 and C7. This shows the claim.

Proof of Lemma B.9. Applying similar arguments for obtaining (B.9), we can write

$$\hat{\mathcal{L}}_{N}^{K}(r) = \frac{1}{2} (r - \gamma)' \mathbb{V}_{\gamma} (r - \gamma) + O_{p}(N^{-1/2} \| r - \gamma \|) + o_{p}(\| r - \gamma \|^{2}) + \frac{1}{\sigma_{N}^{2} N(N - 1)} \sum_{i \neq m} K_{\sigma_{N}, \gamma}(V_{im}(\hat{\beta})) Y_{im(1, 1)} \operatorname{sgn}(W'_{im} \gamma) + O_{p}(N^{-1} \sigma_{N}^{-2}).$$

Then, using the result in Lemma B.8, we have

$$\hat{\mathcal{L}}_{N}^{K}(r) = \frac{1}{2} (r - \gamma)' \mathbb{V}_{\gamma} (r - \gamma) + O_{p}(N^{-1/2} \| r - \gamma \|) + o_{p}(\| r - \gamma \|^{2})
+ \frac{1}{\sigma_{N}^{2} N(N - 1)} \sum_{i \neq m} K_{\sigma_{N}, \gamma} (V_{im}(\beta)) Y_{im(1,1)} \operatorname{sgn}(W'_{im} \gamma) + o_{p}(N^{-1/2}) + O_{p}(N^{-1} \sigma_{N}^{-2}).$$

Since $N^{1/2}\sigma_N^2 \to \infty$ by Assumption C7, $O_p\left(N^{-1}\sigma_N^{-2}\right) = o_p\left(N^{-1/2}\right)$, which completes the proof.

Proof of Lemma B.10. Using the definition that $V_{im}(\beta) = (X'_{im1}\beta, X'_{im2}\beta)$, we can write

$$\mathbb{E}\left[\sigma_{N}^{-2}K_{\sigma_{N},\gamma}\left(V_{im}\left(\beta\right)\right)Y_{im(1,1)}\mathrm{sgn}\left(W_{im}'\gamma\right)|Z_{i}\right]$$

$$=\mathbb{E}\left[\sigma_{N}^{-2}K\left(\frac{X_{im1}'\beta}{\sigma_{N}}\right)K\left(\frac{X_{im2}'\beta}{\sigma_{N}}\right)Y_{im(1,1)}\mathrm{sgn}\left(W_{im}'\gamma\right)|Z_{i}\right]$$

$$=\mathbb{E}\left[\sigma_{N}^{-2}K\left(\frac{X_{im1}'\beta}{\sigma_{N}}\right)K\left(\frac{X_{im2}'\beta}{\sigma_{N}}\right)\mathbb{E}\left[Y_{im(1,1)}\mathrm{sgn}\left(W_{im}'\gamma\right)|Z_{i},X_{m1},X_{m2}\right]|Z_{i}\right]$$

$$=f_{V_{im}(\beta)}(0)\mathbb{E}\left[Y_{im(1,1)}\mathrm{sgn}\left(W_{im}'\gamma\right)|Z_{i},V_{im}\left(\beta\right)=0\right]+o_{p}(N^{-1/2}),$$

where the last line uses the same argument as in proving Lemma B.5.

Proof of Lemma C.1. We verify the conditions in Assumption M, specifically, M.i, M.ii, and M.iii, in Seo and Otsu (2018) one by one. Recall that

$$\phi_{Ni}(b) \equiv \sum_{t>s} \sum_{d \in \mathcal{D}} \{ \mathcal{K}_{h_N}(X_{i2ts}, W_{its}) Y_{idst} (-1)^{d_1} \left(1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0] \right) + \mathcal{K}_{h_N}(X_{i1ts}, W_{its}) Y_{idst} (-1)^{d_2} \left(1[X'_{i2st}b > 0] - 1[X'_{i2st}\beta > 0] \right) \},$$

The " h_n " in Seo and Otsu (2018) needs to be set as $h_N^{k_1+k_2}$ for the term $\phi_{Ni}(b)$. Similarly, the " h_n " in Seo and Otsu (2018) needs to be set as $\sigma_N^{2k_1}$ for the term $\varphi_{Ni}(r)$. We conduct the analysis for $\phi_{Ni}(b)$ first, and the results for $\varphi_{Ni}(r)$ follow similarly. For $\phi_{Ni}(b)$, we analyze the term

$$\mathcal{K}_{h_N}(X_{i2ts}, W_{its})Y_{idst}(-1)^{d_1}(1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0])$$

and the remaining terms can be analyzed similarly, thanks to the similar structure of those terms.

On Assumption M.i in Seo and Otsu (2018). Note that

$$\mathbb{E}[\mathcal{K}_{h_N}(X_{i2ts}, W_{its})Y_{idst}(-1)^{d_1} (1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0])]$$

$$= \int_{\mathbb{R}^{k_2}} \int_{\mathbb{R}^{k_1}} \mathbb{E}[Y_{idst}(-1)^{d_1} (1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0]) | X_{i2ts} = x_2, W_{its} = w]$$

$$\cdot \mathcal{K}_{h_N}(x_2, w) f_{X_{2ts}, W_{ts}}(x_2, w) dx_2 dw$$

$$= \mathbb{E}[Y_{idst}(-1)^{d_1} (1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0]) | X_{i2ts} = 0, W_{its} = 0] f_{X_{2ts}, W_{ts}}(0, 0) + O(h_N^2),$$

where the second equality holds by change of variables and Taylor expansion, and the small order term holds by the properties of the kernel function and the boundedness conditions imposed in Assumptions P5–P7. Further, by Assumption P8, the small order term satisfies $O(h_N^2) = o((Nh_N^{k_1+k_2})^{-2/3})$. As a result, we only need to focus on the lead term

$$\mathbb{E}[Y_{idst}(-1)^{d_1} (1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0]) | X_{i2ts} = 0, W_{its} = 0]$$

$$= \mathbb{E}[\mathbb{E}[Y_{idst}(-1)^{d_1} | X_{i1ts}, X_{i2ts} = 0, W_{its} = 0] (1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0]) | X_{i2ts} = 0, W_{its} = 0]$$

$$= \mathbb{E}[\kappa^{(1)}_{dts}(X_{i1ts}) (1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0]) | X_{i2ts} = 0, W_{its} = 0]$$

$$= \int_{\mathbb{R}^{k_1}} \kappa^{(1)}_{dts}(x) (1[x'b > 0] - 1[x'\beta > 0]) f_{X_{1ts}|\{X_{2ts} = 0, W_{ts} = 0\}}(x) dx$$

$$= \int_{\mathbb{R}^{k_1}} \kappa^{(1)}_{dts}(x) (1[x'b > 0] - 1[x'\beta > 0]) f_{X_{1ts}|\{X_{2ts} = 0, W_{ts} = 0\}}(x) dx, \tag{F.11}$$

where to ease notations we denote

$$\kappa_{dts}^{(1)}(x) = \mathbb{E}[Y_{idst}(-1)^{d_1} | X_{i1ts} = x, X_{i2ts} = 0, W_{its} = 0]$$

in the third line.

We now get the first and second derivatives of the above term w.r.t. b around β . Since the calculation is the classical differential geometry and very similar to that in Sections 5 and 6.4 of Kim and Pollard (1990) and Section B.1 of Seo and Otsu (2018), we only present key steps and omit some standard similar details. First, define the mapping

$$T_b = (I - ||b||_2^{-2}bb')(I - \beta\beta') + ||b||_2^{-2}b\beta'.$$

Then T_b maps $\{X'_{i1ts}b > 0\}$ onto $\{X'_{i1ts}\beta > 0\}$ and $\{X'_{i1ts}b = 0\}$ onto $\{X'_{i1ts}\beta = 0\}$. With equation (F.11), equations (5.2) and (5.3) in Kim and Pollard (1990) imply

$$\frac{\partial}{\partial b} \mathbb{E} \left[Y_{idst} (-1)^{d_1} \left(1 \left[X'_{i1ts} b > 0 \right] - 1 \left[X'_{i1ts} \beta > 0 \right] \right) \middle| X_{i2ts} = 0, W_{its} = 0 \right]
= \frac{\partial}{\partial b} \int_{\mathbb{R}^{k_1}} \kappa_{dts}^{(1)} (x) \left(1 \left[x'b > 0 \right] - 1 \left[x'\beta > 0 \right] \right) f_{X_{1ts} \mid \{X_{2ts} = 0, W_{ts} = 0\}} (x) \, \mathrm{d}x,
= \|b\|_2^{-2} b'\beta \left(I - \|b\|_2^{-2} bb' \right) \int 1 \left[x'\beta = 0 \right] \kappa_{dts}^{(1)} (T_b x) \, x f_{X_{1ts} \mid \{X_{2ts} = 0, W_{ts} = 0\}} (T_b x) \, \mathrm{d}\sigma_0^{(1)},$$

where $\sigma_0^{(1)}$ is the surface measure of $\{X'_{i1ts}\beta=0\}$. Note that $T_{\beta}x=x$, then

$$1 \left[x'\beta = 0 \right] \kappa_{dts}^{(1)} \left(T_{\beta} x \right) = 1 \left[x'\beta = 0 \right] \kappa_{dts}^{(1)} \left(x \right) = 0$$

because

$$\kappa_{dts}^{(1)}(X_{i1t})\Big|_{x'\beta=0} = \mathbb{E}\left[Y_{idst}(-1)^{d_1} \middle| X'_{i1ts}\beta = 0, X_{i2ts} = 0, W_{its} = 0\right]$$

$$= \mathbb{E}\left[\left(Y_{idt} - Y_{ids}\right)(-1)^{d_1} \middle| X'_{i1t}\beta = X'_{i1s}\beta, X_{i2t} = X_{i2s}, W_{it} = W_{is}\right]$$

$$= 0,$$

where the last line holds by Assumption P2 that $\xi_s \stackrel{d}{=} \xi_t | (\alpha, Z^T)$. This implies that

$$\frac{\partial}{\partial b} \mathbb{E} \left[\left. Y_{idst} \left(-1 \right)^{d_1} \left(1 \left[X_{i1ts}' b > 0 \right] - 1 \left[X_{i1ts}' \beta > 0 \right] \right) \right| X_{i2ts} = 0, W_{its} = 0 \right] \right|_{b=\beta} = 0.$$

For the same reason, the nonzero component of the second order derivative at $b = \beta$ only comes from the derivative of $\kappa_{dts}^{(1)}(X_{i1ts})$. Notice that $\frac{\partial \kappa_{dts}^{(1)}(T_b x)}{\partial b}\Big|_{b=\beta} = -\left(\frac{\partial \kappa_{dts}^{(1)}(x)}{\partial x}'\beta\right)x$, we have

$$\frac{\partial^{2}}{\partial b \partial b'} \mathbb{E} \left[Y_{idst} (-1)^{d_{1}} \left(1 \left[X'_{i1ts} b > 0 \right] - 1 \left[X'_{i1ts} \beta > 0 \right] \right) \middle| X_{i2ts} = 0, W_{its} = 0 \right] \bigg|_{b=\beta}$$

$$= - \int 1 \left[x' \beta = 0 \right] \left(\frac{\partial \kappa_{dts}^{(1)} (x)}{\partial x} \beta \right) f_{X_{1ts} | \{X_{2ts} = 0, W_{ts} = 0\}} (x) x x' d\sigma_{0}^{(1)} \equiv \mathbb{V}_{dts}^{(1)}.$$

With the first and second derivatives obtained, we can move on to apply the Taylor expansion to the expectation term in equation (F.11) around $b = \beta$ as

$$\mathbb{E}\left[Y_{idst}(-1)^{d_1}\left(1\left[X'_{i1ts}b > 0\right] - 1\left[X'_{i1ts}\beta > 0\right]\right)\middle|X_{i2ts} = 0, W_{its} = 0\right]$$

$$= \frac{1}{2}\left(b - \beta\right)' \mathbb{V}_{dts}^{(1)}\left(b - \beta\right) + o\left(\|b - \beta\|^2\right).$$

Substitute this back to equation (F.10), we have

$$\mathbb{E}\left[\mathcal{K}_{h_N}(X_{i2ts}, W_{its})Y_{idst}(-1)^{d_1}\left(1\left[X'_{i1ts}b > 0\right] - 1\left[X'_{i1ts}\beta > 0\right]\right)\right]$$

$$= \frac{1}{2}\left(b - \beta\right)' \mathbb{V}_{dts}^{(1)}\left(b - \beta\right) + o\left(\|b - \beta\|^2\right) + o\left(\left(Nh_N^{k_1 + k_2}\right)^{-2/3}\right).$$

We can similarly define

$$\mathbb{V}_{dts}^{(2)} \equiv -\int 1 \left[x'\beta = 0 \right] \left(\frac{\partial \kappa_{dts}^{(2)}(x)}{\partial x}'\beta \right) f_{X_{2ts} | \{X_{1ts} = 0, W_{ts} = 0\}}(x) xx' d\sigma_0^{(2)},$$

where

$$\kappa_{dts}^{(2)}(x) \equiv \mathbb{E}\left[Y_{idst}(-1)^{d_2} \middle| X_{i2ts} = x, X_{i1ts} = 0, W_{its} = 0\right],$$

and $\sigma_0^{(2)}$ is the surface measure of $\{X'_{i2ts}\beta=0\}$. A similar derivation yields

$$\mathbb{E}\left[\mathcal{K}_{h_N}(X_{i1ts}, W_{its})Y_{idst}(-1)^{d_2}\left(1\left[X'_{i2st}b > 0\right] - 1\left[X'_{i2st}\beta > 0\right]\right)\right]$$

$$= \frac{1}{2}\left(b - \beta\right)' \mathbb{V}_{dts}^{(2)}\left(b - \beta\right) + o\left(\|b - \beta\|^2\right) + o\left(\left(Nh_N^{k_1 + k_2}\right)^{-2/3}\right).$$

Put everything so far together, we have

$$\mathbb{E}\left[\phi_{Ni}(b)\right] = \frac{1}{2} (b - \beta)' \left[\sum_{t>s} \sum_{d \in \mathcal{D}} \left(\mathbb{V}_{dts}^{(1)} + \mathbb{V}_{dts}^{(2)} \right) \right] (b - \beta) + o(\|b - \beta\|^2) + o((Nh_N^{k_1 + k_2})^{-2/3}).$$
(F.12)

Assumption M.i in Seo and Otsu (2018) is then verified by

$$\mathbb{V} \equiv \sum_{t>s} \sum_{d \in \mathcal{D}} \left(\mathbb{V}_{dts}^{(1)} + \mathbb{V}_{dts}^{(2)} \right).$$

$$= -\sum_{t>s} \sum_{d \in \mathcal{D}} \left\{ \int 1 \left[x'\beta = 0 \right] \left(\frac{\partial \kappa_{dts}^{(1)}(x)'}{\partial x} \beta \right) f_{X_{1ts} | \{X_{2ts} = 0, W_{ts} = 0\}}(x) x x' d\sigma_0^{(1)} \right.$$

$$+ \int 1 \left[x'\beta = 0 \right] \left(\frac{\partial \kappa_{dts}^{(2)}(x)'}{\partial x} \beta \right) f_{X_{2ts} | \{X_{1ts} = 0, W_{ts} = 0\}}(x) x x' d\sigma_0^{(2)} \right\}.$$

We can obtain similar results for $\varphi_{Ni}(r)$ using exactly the same line of analysis, i.e.,

$$\mathbb{E}\left[\varphi_{Ni}(r)\right] = \frac{1}{2} (r - \gamma)' \mathbb{W}(r - \gamma) + o(\|r - \gamma\|^2) + o((N\sigma_N^{2k_1})^{-2/3}), \tag{F.13}$$

where

$$\mathbb{W} \equiv -\sum_{t>s} \int 1 \left[w' \gamma = 0 \right] \left(\frac{\partial \kappa_{(1,1)ts}^{(3)} \left(w \right)'}{\partial w} \gamma \right) f_{W_{ts} \mid \{X_{1ts} = 0, X_{2ts} = 0\}} \left(w \right) ww' d\sigma_0$$

with

$$\kappa_{(1,1)ts}^{(3)}(w) \equiv \mathbb{E}\left[Y_{i(1,1)ts}|W_{its}=w, X_{i1ts}=0, X_{i2ts}=0\right]$$

and $\sigma_0^{(3)}$ being the surface measure of $\{W'_{its}\gamma=0\}$.

On Assumption M.ii in Seo and Otsu (2018). Again, we first verify this condition for

$$\mathcal{K}_{h_N}(X_{i2ts}, W_{its})Y_{idst}(-1)^{d_1}(1[X'_{i1ts}b > 0] - 1[X'_{i1ts}\beta > 0]),$$

and other components in $\phi_{Ni}(b)$ follow similarly. We evaluate norm of the difference of the above term at $b=b_1$ and $b=b_2$ multiplied by $h_N^{(k_1+k_2)/2}$:

$$h_{N}^{(k_{1}+k_{2})/2} \left\| \mathcal{K}_{h_{N}}(X_{i2ts}, W_{its}) Y_{idst} \left(-1\right)^{d_{1}} \left(1 \left[X_{i1ts}' b_{1} > 0\right] - 1 \left[X_{i1ts}' b_{2} > 0\right]\right) \right\|$$

$$= \left\{ \mathbb{E} \left[\mathbb{E} \left[h_{N}^{k_{1}+k_{2}} \mathcal{K}_{h_{N}}^{2} (X_{i2ts}, W_{its}) Y_{idst}^{2} \middle| X_{i1ts} \right] \left(1 \left[X_{i1ts}' b_{1} > 0\right] - 1 \left[X_{i1ts}' b_{2} > 0\right]\right)^{2} \right] \right\}^{1/2}$$

$$= \left\{ \mathbb{E} \left[\mathbb{E} \left[h_{N}^{k_{1}+k_{2}} \mathcal{K}_{h_{N}}^{2} (X_{i2ts}, W_{its}) Y_{idst}^{2} \middle| X_{i1ts} \right] \left|1 \left[X_{i1ts}' b_{1} > 0\right] - 1 \left[X_{i1ts}' b_{2} > 0\right]\right| \right] \right\}^{1/2} \quad (F.14)$$

$$\geq C_{1} \mathbb{E} \left[\left|1 \left[X_{i1ts}' b_{1} > 0\right] - 1 \left[X_{i1ts}' b_{2} > 0\right]\right| \right] \geq C_{2} \left\| b - \beta \right\|_{2},$$

where the second equality holds by the fact that the difference of two indicator functions can only take values -1, 0, or 1. C_1 and C_2 are two positive constants. Applying the same analysis to all other terms in $\phi_{Ni}(b)$, we have

$$h_N^{(k_1+k_2)/2} \|\phi_{Ni}(b_1) - \phi_{Ni}(b_2)\| \ge C_3 \|b_1 - b_2\|$$

for some positive constant C_3 .

Applying similar analysis to $\varphi_{Ni}(r)$ leads to

$$\sigma_N^{k_1} \|\varphi_{Ni}(r_1) - \varphi_{Ni}(r_2)\| \ge C_4 \|r_1 - r_2\|$$

for some positive constant C_4 .

On Assumption M.iii in Seo and Otsu (2018). We still begin with the analysis on

$$\mathcal{K}_{h_N}(X_{i2ts}, W_{its})Y_{idst}(-1)^{d_1} (1 [X'_{i1ts}b > 0] - 1 [X'_{i1ts}\beta > 0]),$$

and other components in $\phi_{Ni}(b)$ follow similarly. We evaluate the square of difference of the

above term at $b=b_1$ and $b=b_2$, such that $||b_1-b_2||<\varepsilon$, multiplied by $h_N^{k_1+k_2}$:

$$\begin{split} & h_{N}^{k_{1}+k_{2}}\mathbb{E}\left[\sup_{b_{1},b_{2}\in\mathcal{B},\|b_{1}-b_{2}\|<\varepsilon}\left[\mathcal{K}_{h_{N}}(X_{i2ts},W_{its})Y_{idst}\left(-1\right)^{d_{1}}\left(1\left[X_{i1ts}^{\prime}b_{1}>0\right]-1\left[X_{i1ts}^{\prime}b_{2}>0\right]\right)\right]^{2}\right]\\ \leq& \mathbb{E}\left[\mathbb{E}\left[h_{N}^{k_{1}+k_{2}}\mathcal{K}_{h_{N}}^{2}(X_{i2ts},W_{its})Y_{idst}^{2}\left|X_{i1ts}\right]\sup_{b_{1},b_{2}\in\mathcal{B},\|b_{1}-b_{2}\|<\varepsilon}\left|1\left[X_{i1ts}^{\prime}b_{1}>0\right]-1\left[X_{i1ts}^{\prime}b_{2}>0\right]\right|\right]\right]\\ \leq& C_{5}\mathbb{E}\left[\sup_{b_{1},b_{2}\in\mathcal{B},\|b_{1}-b_{2}\|<\varepsilon}\left|1\left[X_{i1ts}^{\prime}b_{1}>0\right]-1\left[X_{i1ts}^{\prime}b_{2}>0\right]\right|\right]\leq C_{6}\varepsilon, \end{split}$$

where the second line follows by the fact that the maximum of absolute value of the difference of two indicator functions is 1, and the last line holds by Assumptions P4 and P5. Apply the same analysis to all other terms in $\phi_{Ni}(b)$ to obtain

$$h_N^{k_1+k_2}\mathbb{E}\left[\sup_{b_1,b_2\in\mathcal{B},\|b_1-b_2\|<\varepsilon}\left[\phi_{Ni}\left(b_1\right)-\phi_{Ni}\left(b_2\right)\right]^2\right]\leq C_7\varepsilon$$

for some positive constant C_7 .

Similar analysis on $\varphi_{Ni}(r)$ leads to

$$h_N^{2k_1} \mathbb{E}\left[\sup_{r_1, r_2 \in \mathcal{R}, \|r_1 - r_2\| < \varepsilon} \left[\varphi_{Ni}\left(r_1\right) - \varphi_{Ni}\left(r_2\right)\right]^2\right] \le C_8 \varepsilon$$

for some positive constant C_8 .

Proof of Lemma C.2. The first two equalities are direct results from Lemma B.1 by Taylor expansions. Taking $b = \beta + \rho (Nh_N^{k_1+k_2})^{-1/3}$ in equation (F.12) yields

$$(Nh_N^{k_1+k_2})^{2/3} \mathbb{E} \left[\phi_{Ni} \left(\beta + \rho \left(Nh_N^{k_1+k_2} \right)^{-1/3} \right) \right] = \frac{1}{2} \rho' \mathbb{V} \rho + o(1) \to \frac{1}{2} \rho' \mathbb{V} \rho.$$

Similarly, setting $r = \gamma + \delta(N\sigma_N^{2k_1})^{-1/3}$ in equation (F.13) gives

$$\left(N\sigma_N^{2k_1}\right)^{2/3} \mathbb{E}\left[\varphi_{Ni}\left(\gamma + \delta\left(N\sigma_N^{2k_1}\right)^{-1/3}\right)\right] = \frac{1}{2}\delta' \mathbb{W}\delta + o\left(1\right) \to \frac{1}{2}\delta' \mathbb{W}\delta.$$

 \mathbb{H}_1 and \mathbb{H}_2 can be obtained in the same way as in Kim and Pollard (1990). We omit some similar yet tedious details and refer interested readers to Section 6.4 in Kim and Pollard

(1990). To get \mathbb{H}_1 , we let $\upsilon_N \equiv (Nh_N^{k_1+k_2})^{1/3}$ and define

$$\mathbb{L}\left(\rho_{1}-\rho_{2}\right) \equiv \lim_{N\to\infty} \upsilon_{N} \mathbb{E}\left\{h_{N}^{k_{1}+k_{2}}\left[\phi_{Ni}\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)-\phi_{Ni}\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)\right]^{2}\right\},$$

$$\mathbb{L}\left(\rho_{1}\right) \equiv \lim_{N\to\infty} \upsilon_{N} \mathbb{E}\left\{h_{N}^{k_{1}+k_{2}}\left[\phi_{Ni}\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)-\phi_{Ni}\left(\beta\right)\right]^{2}\right\}, \text{ and }$$

$$\mathbb{L}\left(\rho_{2}\right) \equiv \lim_{N\to\infty} \upsilon_{N} \mathbb{E}\left\{h_{N}^{k_{1}+k_{2}}\left[\phi_{Ni}\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)-\phi_{Ni}\left(\beta\right)\right]^{2}\right\}.$$

Since $\phi_{Ni}(\beta) = 0$,

$$\mathbb{H}_{1}\left(\rho_{1}, \rho_{2}\right) = \frac{1}{2} \left[\mathbb{L}\left(\rho_{1}\right) + \mathbb{L}\left(\rho_{2}\right) - \mathbb{L}\left(\rho_{1} - \rho_{2}\right) \right]. \tag{F.15}$$

We calculate \mathbb{L} as follows. Notice that

$$\begin{split} &\mathbb{E}\left[h_{N}^{k_{1}+k_{2}}\mathcal{K}_{h_{N}}(X_{ijts},W_{its})^{2}\right] \\ &= \int \frac{1}{h_{N}^{k_{1}+k_{2}}} \left[\Pi_{\iota=1}^{k_{1}}K\left(\frac{X_{ijts,\iota}}{h_{N}}\right)\Pi_{\iota=1}^{k_{2}}K\left(\frac{W_{its,\iota}}{h_{N}}\right)\right]^{2}\mathrm{d}F_{X_{ijts},W_{its}} = O(1), \\ &\mathbb{E}\left[h_{N}^{k_{1}+k_{2}}\mathcal{K}_{h_{N}}(X_{ijts},W_{its})\mathcal{K}_{h_{N}}(X_{ij't's'},W_{it's'})\right] \\ &= \int \int \frac{1}{h_{N}^{k_{1}+k_{2}}}\Pi_{\iota=1}^{k_{1}}K\left(\frac{X_{ijts,\iota}}{h_{N}}\right)\Pi_{\iota=1}^{k_{2}}K\left(\frac{W_{its,\iota}}{h_{N}}\right) \\ &\cdot \Pi_{\iota=1}^{k_{1}}K\left(\frac{X_{ij't's',\iota}}{h_{N}}\right)\Pi_{\iota=1}^{k_{2}}K\left(\frac{W_{it's',\iota}}{h_{N}}\right)\mathrm{d}F_{X_{ijts},W_{its}}\mathrm{d}F_{X_{ij't's'},W_{it's'}} \\ &= o\left(1\right), \text{ for any } (j,t,s) \neq (j',t',s'), \end{split}$$

and

$$\begin{split} & \phi_{Ni} \left(\beta + \rho_1 v_N^{-1}\right) - \phi_{Ni} \left(\beta + \rho_2 v_N^{-1}\right) \\ &= \sum_{t>s} \sum_{d \in \mathcal{D}} \left\{ \mathcal{K}_{h_N} (X_{i2ts}, W_{its}) Y_{idst} \left(-1\right)^{d_1} \left(1 \left[X'_{i1ts} \left(\beta + \rho_1 v_N^{-1}\right) > 0\right] - 1 \left[X'_{i1ts} \left(\beta + \rho_2 v_N^{-1}\right) > 0\right]\right) \\ & + \mathcal{K}_{h_N} (X_{i1ts}, W_{its}) Y_{idst} \left(-1\right)^{d_2} \left(1 \left[X'_{i2st} \left(\beta + \rho_1 v_N^{-1}\right) > 0\right] - 1 \left[X'_{i2st} \left(\beta + \rho_2 v_N^{-1}\right) > 0\right]\right) \right\}. \end{split}$$

Thus, the interaction terms after expanding the square in L's are asymptotically negligible, and we only need to focus on the squares of each term above.

We now derive the square of the first term multiplied by $h_N^{k_1+k_2}$ and the rest follow

similarly.

$$\begin{split} &\mathbb{E}\left\{h_{N}^{k_{1}+k_{2}}\left[\mathcal{K}_{h_{N}}(X_{i2ts},W_{its})Y_{idst}\left(-1\right)^{d_{1}}\left(1\left[X_{i1ts}'\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)>0\right]-1\left[X_{i1ts}'\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)>0\right]\right)\right]^{2}\right\} \\ &=\mathbb{E}\left\{\mathbb{E}\left[|Y_{idst}|\left|1\left[X_{i1ts}'\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)>0\right]-1\left[X_{i1ts}'\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)>0\right]\right|\left|X_{i2ts},W_{its}\right|h_{N}^{k_{1}+k_{2}}\mathcal{K}_{h_{N}}^{2}(X_{i2ts},W_{its})\right\} \\ &=\mathbb{E}\left[|Y_{idst}|\left|1\left[X_{i1ts}'\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)>0\right]-1\left[X_{i1ts}'\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)>0\right]\right|\left|X_{i2ts}=0,W_{its}=0\right] \\ &\cdot f_{X_{2ts},W_{ts}}\left(0,0\right)\left[\int K^{2}\left(u\right)\mathrm{d}u\right]^{k_{1}+k_{2}}+O\left(h_{N}\right) \\ &=\mathbb{E}\left\{\mathbb{E}\left[|Y_{idst}|\left|X_{i1ts},X_{i2ts}=0,W_{its}=0\right]\right.\right. \\ &\cdot \left|1\left[X_{i1ts}'\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)>0\right]-1\left[X_{i1ts}'\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)>0\right]\right|\left|X_{i2ts}=0,W_{its}=0\right\} \\ &\cdot f_{X_{2ts},W_{ts}}\left(0,0\right)\left[\int K^{2}\left(u\right)\mathrm{d}u\right]^{k_{1}+k_{2}}+O\left(h_{N}\right) \\ &=\int \kappa_{dts}^{(4)}\left(x\right)\left|1\left[x'\left(\beta+\rho_{1}\upsilon_{N}^{-1}\right)>0\right]-1\left[x'\left(\beta+\rho_{2}\upsilon_{N}^{-1}\right)>0\right]\right|f_{X_{1ts}|\left\{X_{2ts}=0,W_{its}=0\right\}}\left(x\right)\mathrm{d}x \\ &\cdot f_{X_{2ts},W_{ts}}\left(0,0\right)\bar{K}_{2}^{k_{1}+k_{2}}+O\left(h_{N}\right), \end{split}$$

where $\kappa_{dts}^{(4)}(x) \equiv \mathbb{E}\left[|Y_{idst}| \mid X_{i1ts} = x, X_{i2ts} = 0, W_{its} = 0\right]$ and $\bar{K}_2 \equiv \int K^2(u) \, du$. By Assumption P9, $v_N h_N = (N h_N^{k_1 + k_2})^{1/3} h_N \to 0$, so the bias term is asymptotic negligible. To calculate the above integral, we follow Kim and Pollard (1990) and decompose $X_{1ts} = a\beta + \bar{X}_{1ts}$, with \bar{X}_{1ts} orthogonal to β . We use $f_{X_{1ts}}(a,\bar{x})$ to denote the density of X_{1ts} at $X_{1ts} = a\beta + \bar{x}$. Using the results in Kim and Pollard (1990),

$$\lim_{N \to \infty} v_N \cdot \int \kappa_{dts}^{(4)}(x) \left| 1 \left[x' \left(\beta + \rho_1 v_N^{-1} \right) > 0 \right] - 1 \left[x' \left(\beta + \rho_2 v_N^{-1} \right) > 0 \right] \right| f_{X_{1ts} \mid \{X_{2ts} = 0, W_{ts} = 0\}}(x) dx$$

$$= \int \kappa_{dts}^{(4)}(\bar{x}) \left| \bar{x}' \left(\rho_1 - \rho_2 \right) \right| f_{X_{1ts} \mid \{X_{2ts} = 0, W_{ts} = 0\}}(0, \bar{x}) d\bar{x}.$$

To summarize, we have

$$\lim_{N\to\infty} v_N \mathbb{E} \left\{ h_N^{k_1+k_2} \left[\mathcal{K}_{h_N}(X_{i2ts}, W_{its}) Y_{idst} \left(-1\right)^{d_1} \left(1 \left[X_{i1ts}' \left(\beta + \rho_1 v_N^{-1} \right) > 0 \right] - 1 \left[X_{i1ts}' \left(\beta + \rho_2 v_N^{-1} \right) > 0 \right] \right) \right]^2 \right\}$$

$$= \int \kappa_{dts}^{(4)} \left(\bar{x} \right) |\bar{x}' \left(\rho_1 - \rho_2 \right)| f_{X_{1ts} | \{X_{2ts} = 0, W_{ts} = 0\}} \left(0, \bar{x} \right) d\bar{x} f_{X_{2ts}, W_{ts}} \left(0, 0 \right) \bar{K}_2^{k_1 + k_2}.$$

Similarly, we have for the second term

$$\lim_{N\to\infty} v_N \mathbb{E} \left\{ h_N^{k_1+k_2} \left[\mathcal{K}_{h_N}(X_{i1ts}, W_{its}) Y_{idst} \left(-1\right)^{d_2} \left(1 \left[X'_{i2st} \left(\beta + \rho_1 v_N^{-1} \right) > 0 \right] - 1 \left[X'_{i2st} \left(\beta + \rho_2 v_N^{-1} \right) > 0 \right] \right) \right]^2$$

$$= \int \kappa_{dts}^{(5)} \left(\bar{x} \right) \left| \bar{x}' \left(\rho_1 - \rho_2 \right) \right| f_{X_{2ts} \mid \{X_{1ts} = 0, W_{ts} = 0\}} \left(0, \bar{x} \right) d\bar{x} f_{X_{1ts}, W_{ts}} \left(0, 0 \right) \bar{K}_2^{k_1 + k_2},$$

where

$$\kappa_{dts}^{(5)}(x) \equiv \mathbb{E}[|Y_{idst}| | X_{i2ts} = x, X_{i1ts} = 0, W_{its} = 0].$$

Therefore,

$$\mathbb{L}(\rho_{1} - \rho_{2}) = \lim_{N \to \infty} v_{N} \mathbb{E} \left\{ h_{N}^{k_{1}+k_{2}} \left[\phi_{Ni} \left(\beta + \rho_{1} v_{N}^{-1} \right) - \phi_{Ni} \left(\beta + \rho_{2} v_{N}^{-1} \right) \right]^{2} \right\}$$

$$= \sum_{t>s} \sum_{d \in \mathcal{D}} \left\{ \int \kappa_{dts}^{(4)} (\bar{x}) |\bar{x}' (\rho_{1} - \rho_{2})| f_{X_{1ts}|\{X_{2ts} = 0, W_{ts} = 0\}} (0, \bar{x}) d\bar{x} f_{X_{2ts}, W_{ts}} (0, 0) + \int \kappa_{dts}^{(5)} (\bar{x}) |\bar{x}' (\rho_{1} - \rho_{2})| f_{X_{2ts}|\{X_{1ts} = 0, W_{ts} = 0\}} (0, \bar{x}) d\bar{x} f_{X_{1ts}, W_{ts}} (0, 0) \right\} \bar{K}_{2}^{k_{1}+k_{2}}.$$

Finally, by equation (F.15),

$$\mathbb{H}_{1}(\rho_{1}, \rho_{2}) = \frac{1}{2} \sum_{t>s} \sum_{d \in \mathcal{D}} \left\{ \int \kappa_{dts}^{(4)}(\bar{x}) \left[|\bar{x}'\rho_{1}| + |\bar{x}'\rho_{2}| - |\bar{x}'(\rho_{1} - \rho_{2})| \right] f_{X_{1ts}|\{X_{2ts} = 0, W_{ts} = 0\}}(0, \bar{x}) \, d\bar{x} f_{X_{2ts}, W_{ts}}(0, 0) \right. \\
+ \int \kappa_{dts}^{(5)}(\bar{x}) \left[|\bar{x}'\rho_{1}| + |\bar{x}'\rho_{2}| - |\bar{x}'(\rho_{1} - \rho_{2})| \right] f_{X_{2ts}|\{X_{1ts} = 0, W_{ts} = 0\}}(0, \bar{x}) \, d\bar{x} f_{X_{1ts}, W_{ts}}(0, 0) \right\} \bar{K}_{2}^{k_{1} + k_{2}}.$$

The same arguments can be applied to obtain \mathbb{H}_2 as

$$\mathbb{H}_{2}\left(\delta_{1}, \delta_{2}\right) = \frac{1}{2} \sum_{t>s} \int \kappa_{(1,1)ts}^{(6)}\left(\bar{w}\right) \left[\left|\bar{w}'\delta_{1}\right| + \left|\bar{w}'\delta_{2}\right| - \left|\bar{w}'\left(\delta_{1} - \delta_{2}\right)\right|\right] f_{W_{ts}|\left\{X_{1ts} = 0, X_{2ts} = 0\right\}}\left(0, \bar{w}\right) d\bar{w}$$

$$\cdot f_{X_{1ts}, X_{2ts}}\left(0, 0\right) \bar{K}_{2}^{2k_{1}},$$

where $W_{ts} = a\gamma + \bar{W}_{ts}$ with \bar{W}_{ts} orthogonal to γ , $f_{W_{ts}}(a, \bar{w})$ denotes the density of W_{ts} at $W_{ts} = a\gamma + \bar{W}_{ts}$, and

$$\kappa_{(1,1)ts}^{(6)}\left(w\right) \equiv \mathbb{E}\left[\left|Y_{i(1,1)ts}\right| \; \left|W_{its}=w,X_{i1ts}=0,X_{i2ts}=0\right].$$

References

ABREVAYA, J., J. A. HAUSMAN, AND S. KHAN (2010): "Testing for causal effects in a generalized regression model with endogenous regressors," *Econometrica*, 78, 2043–2061.

ABREVAYA, J. AND J. HUANG (2005): "On the bootstrap of the maximum score estimator,"

- Econometrica, 73, 1175–1204.
- AHN, H., H. ICHIMURA, J. L. POWELL, AND P. A. RUUD (2018): "Simple estimators for invertible index models," *Journal of Business & Economic Statistics*, 36, 1–10.
- ALLEN, R. AND J. REHBECK (2019): "Identification with additively separable heterogeneity," *Econometrica*, 87, 1021–1054.
- ALTONJI, J. G. AND R. L. MATZKIN (2005): "Cross section and panel data estimators for nonseparable models with endogenous regressors," *Econometrica*, 73, 1053–1102.
- ARCONES, M. A. AND E. GINE (1992): "On the bootstrap of U and V statistics," *The Annals of Statistics*, 655–674.
- Augereau, A., S. Greenstein, and M. Rysman (2006): "Coordination versus differentiation in a standards war: 56K modems," *The RAND Journal of Economics*, 37, 887–909.
- BERRY, S., A. KHWAJA, V. KUMAR, A. MUSALEM, K. C. WILBUR, G. ALLENBY, B. ANAND, P. CHINTAGUNTA, W. M. HANEMANN, AND P. JEZIORSKI (2014): "Structural models of complementary choices," *Marketing Letters*, 25, 245–256.
- Berry, S. T. and P. A. Haile (2009): "Nonparametric Identification of Multinomial Choice Demand Models with Heterogeneous Consumers," Working Paper 15276, National Bureau of Economic Research.
- Cameron, A. C. and P. K. Trivedi (2005): *Microeconometrics: methods and applications*, Cambridge University Press.
- Candelaria, L. E. (2020): "A semiparametric network formation model with unobserved linear heterogeneity," arXiv preprint arXiv:2007.05403.
- CAVANAGH, C. AND R. P. SHERMAN (1998): "Rank estimators for monotonic index models," *Journal of Econometrics*, 84, 351–382.
- CHEN, X., O. LINTON, AND I. VAN KEILEGOM (2003): "Estimation of semiparametric models when the criterion function is not smooth," *Econometrica*, 71, 1591–1608.

- Chernozhukov, V., I. Fernández-Val, and W. K. Newey (2019): "Nonseparable multinomial choice models in cross-section and panel data," *Journal of Econometrics*, 211, 104–116.
- Chung, J. and V. R. Rao (2003): "A general choice model for bundles with multiple-category products: Application to market segmentation and optimal pricing for bundles," *Journal of Marketing Research*, 40, 115–130.
- DEVROYE, L. (1978): "The uniform convergence of nearest neighbor regression function estimators and their application in optimization," *IEEE Transactions on Information Theory*, 24, 142–151.
- Dubé, J.-P. (2004): "Multiple discreteness and product differentiation: Demand for carbonated soft drinks," *Marketing Science*, 23, 66–81.
- Dunker, F., S. Hoderlein, and H. Kaido (2022): "Nonparametric identification of random coefficients in endogenous and heterogeneous aggregate demand models," arXiv preprint arXiv:2201.06140.
- Ershov, D., J.-W. Laliberté, M. Marcoux, and S. Orr (2021): "Estimating complementarity with large choice sets: An application to mergers," *Available at SSRN:* https://ssrn.com/abstract=3802097.
- FAN, Y. (2013): "Ownership consolidation and product characteristics: A study of the US daily newspaper market," *American Economic Review*, 103, 1598–1628.
- FOUBERT, B. AND E. GIJSBRECHTS (2007): "Shopper response to bundle promotions for packaged goods," *Journal of Marketing Research*, 44, 647–662.
- Fox, J. T. (2007): "Semiparametric estimation of multinomial discrete-choice models using a subset of choices," *The RAND Journal of Economics*, 38, 1002–1019.
- Fox, J. T. And N. Lazzati (2017): "A note on identification of discrete choice models for bundles and binary games," *Quantitative Economics*, 8, 1021–1036.
- FRITSCH, S. AND F. GUENTHER (2019): "Package 'neuralnet'," Training of Neural Networks, 2, 30.
- GAO, W. Y. AND M. LI (2020): "Robust semiparametric estimation in panel multinomial choice models," arXiv preprint arXiv:2009.00085.

- GAO, W. Y., M. Li, and S. Xu (2023): "Logical differencing in dyadic network formation models with nontransferable utilities," *Journal of Econometrics*, 235, 302–324.
- Gentzkow, M. (2007): "Valuing new goods in a model with complementarity: Online newspapers," *American Economic Review*, 97, 713–744.
- Gentzkow, M., J. M. Shapiro, and M. Sinkinson (2014): "Competition and ideological diversity: Historical evidence from us newspapers," *American Economic Review*, 104, 3073–3114.
- GINÉ, E. AND J. ZINN (1990): "Bootstrapping general empirical measures," *The Annals of Probability*, 851–869.
- Graham, B. S. (2017): "An econometric model of network formation with degree heterogeneity," *Econometrica*, 85, 1033–1063.
- Hall, P., J. Racine, and Q. Li (2004): "Cross-validation and the estimation of conditional probability densities," *Journal of the American Statistical Association*, 99, 1015–1026.
- HAN, A. K. (1987): "Nonparametric analysis of a generalized regression model; the maximum rank correlation estimator," *Journal of Econometrics*, 35, 303–316.
- Hansen, B. E. (2008): "Uniform convergence rates for kernel estimation with dependent data," *Econometric Theory*, 24, 726–748.
- Hayfield, T. and J. S. Racine (2008): "Nonparametric econometrics: The np package," Journal of Statistical Software, 27, 1–32.
- HENDEL, I. (1999): "Estimating multiple-discrete choice models: An application to computerization returns," *The Review of Economic Studies*, 66, 423–446.
- Hong, H. and J. Li (2020): "The numerical bootstrap," *The Annals of Statistics*, 48, 397–412.
- Hong, H., A. Mahajan, and D. Nekipelov (2015): "Extremum estimation and numerical derivatives," *Journal of Econometrics*, 188, 250–263.
- HOROWITZ, J. L. (1992): "A smoothed maximum score estimator for the binary response model," *Econometrica*, 60, 505–531.
- ——— (2001): "The bootstrap," in *Handbook of Econometrics*, Elsevier, vol. 5, 3159–3228.

- IARIA, A. AND A. WANG (2021): "An empirical model of quantity discounts with large choice sets," CEPR Discussion Paper No. DP16666, Available at SSRN: https://ssrn.com/abstract=3960270.
- JIN, Z., Z. YING, AND L. J. WEI (2001): "A simple resampling method by perturbing the minimand," *Biometrika*, 88, 381–390.
- Khan, S., F. Ouyang, and E. Tamer (2021): "Inference on semiparametric multinomial response models," *Quantitative Economics*, 12, 743–777.
- Kim, J. and D. Pollard (1990): "Cube root asymptotics," *The Annals of Statistics*, 18, 191–219.
- Kim, Y., S. Misra, and B. Shapiro (2020): "Valuing brand collaboration: Evidence from a natural experiment," *Available at SSRN: https://ssrn.com/abstract=3335833*.
- Kohler, M. and S. Langer (2021): "On the rate of convergence of fully connected deep neural network regression estimates," *The Annals of Statistics*, 49, 2231–2249.
- Lee, L.-f. (1995): "Semiparametric maximum likelihood estimation of polychotomous and sequential choice models," *Journal of Econometrics*, 65, 381–428.
- LEE, S. M. S. AND M. C. Pun (2006): "On m out of n bootstrapping for nonstandard mestimation with nuisance parameters," *Journal of American Statistical Association*, 101, 1185–1197.
- Lewbel, A. (2000): "Semiparametric qualitative response model estimation with unknown heteroscedasticity or instrumental variables," *Journal of Econometrics*, 97, 145–177.
- Lewbel, A. and L. Nesheim (2019): "Sparse demand systems: corners and complements," Cemmap Working Paper CWP45/19, Institute for Fiscal Studies.
- LI, Q. AND J. S. RACINE (2007): Nonparametric econometrics: theory and practice, Princeton University Press.
- Liu, H., P. K. Chintagunta, and T. Zhu (2010): "Complementarities and the demand for home broadband internet services," *Marketing Science*, 29, 701–720.
- Manski, C. F. (1987): "Semiparametric analysis of random effects linear models from binary panel data," *Econometrica* (1986-1998), 55, 357–362.
- Manski, C. F. and L. Sherman (1980): "An empirical analysis of household choice among motor vehicles," *Transportation Research Part A: General*, 14, 349–366.

- Mullen, K., D. Ardia, D. L. Gil, D. Windover, and J. Cline (2011): "Deoptim: An R package for global optimization by differential evolution," *Journal of Statistical Software*, 40, 1–26.
- NEVO, A., D. L. RUBINFELD, AND M. McCabe (2005): "Academic journal pricing and the demand of libraries," *American Economic Review*, 95, 447–452.
- NEWEY, W. K. AND D. McFadden (1994): "Large sample estimation and hypothesis testing," *Handbook of Econometrics*, 4, 2111–2245.
- NOLAN, D. AND D. POLLARD (1987): "U-processes: rates of convergence," *The Annals of Statistics*, 15, 780–799.
- OUYANG, F., T. T. YANG, AND H. ZHANG (2020): "Semiparametric identification and estimation of discrete choice models for bundles," *Economics Letters*, 109321.
- Pakes, A. and D. Pollard (1989): "Simulation and the asymptotics of optimization estimators," *Econometrica*, 57, 1027–1057.
- PAKES, A. AND J. PORTER (2021): "Moment Inequalities for Multinomial Choice with Fixed Effects," Forthcoming at Quantitative Economics.
- SEO, M. H. AND T. OTSU (2018): "Local M-estimation with discontinuous criterion for dependent and limited observations," *The Annals of Statistics*, 46, 344–369.
- SERFLING, R. J. (2009): Approximation theorems of mathematical statistics, vol. 162, John Wiley & Sons.
- SHER, I. AND K. I. KIM (2014): "Identifying combinatorial valuations from aggregate demand," *Journal of Economic Theory*, 153, 428–458.
- SHERMAN, R. P. (1993): "The limiting distribution of the maximum rank correlation estimator," *Econometrica*, 61, 123–137.

- Shi, X., M. Shum, and W. Song (2018): "Estimating semiparametric panel multinomial choice models using cyclic monotonicity," *Econometrica*, 86, 737–761.

- Song, I. and P. K. Chintagunta (2006): "Measuring cross-category price effects with aggregate store data," *Management Science*, 52, 1594–1609.
- Storn, R. M. and K. V. Price (1997): "Differential evolution—a simple and efficient heuristic for global optimization over continuous spaces," *Journal of Global Optimization*, 11, 341–359.
- Subbotin, V. (2007): "Asymptotic and bootstrap properties of rank regressions," Available at SSRN: https://ssrn.com/abstract=1028548.
- TOTH, P. (2017): "Semiparametric estimation in network formation models with homophily and degree heterogeneity," Available at SSRN: https://ssrn.com/abstract=2988698.
- Train, K. E., D. L. McFadden, and M. Ben-Akiva (1987): "The demand for local telephone service: A fully discrete model of residential calling patterns and service choices," *The RAND Journal of Economics*, 109–123.
- VAN DER VAART, A. AND J. WELLNER (1996): Weak convergence and empirical processes, Springer, New York.
- Wang, R. (2023): "Testing and Identifying Substitution and Complementarity Patterns," arXiv preprint arXiv:2304.00678.
- Wellner, J. A. and Y. Zhan (1996): "Bootstrapping Z-estimators," *University of Washington Department of Statistics Technical Report*, 308.
- YAN, J. AND H. I. YOO (2019): "Semiparametric estimation of the random utility model with rank-ordered choice data," *Journal of Econometrics*, 211, 414–438.