Non-minimaxity of debiased shrinkage estimators

Yuzo Maruyama and Akimichi Takemura

Kobe University & Shiga University

e-mail: maruyama@port.kobe-u.ac.jp; a-takemura@biwako.shiga-u.ac.jp

Abstract: We consider the estimation of the *p*-variate normal mean of $X \sim N_p(\theta, I)$ under the quadratic loss function. We investigate the decision theoretic properties of debiased shrinkage estimator, the estimator which shrinks towards the origin for smaller $||x||^2$ and which is exactly equal to the unbiased estimator X for larger $||x||^2$. Such debiased shrinkage estimator seems superior to the unbiased estimator X, which implies minimaxity. However we show that it is not minimax under mild conditions.

MSC2020 subject classifications: Primary 62C20; secondary 62J07. Keywords and phrases: minimaxity, debiased shrinkage estimator, James-Stein estimator.

1. Introduction

Let X have a p-variate normal distribution $\mathcal{N}_p(\theta, I_p)$. We consider the problem of estimating the mean vector θ under the loss function

$$L(\theta, \hat{\theta}) = \|\hat{\theta} - \theta\|^2 = \sum_{i=1}^{p} (\hat{\theta}_i - \theta_i)^2.$$
(1.1)

The risk function of an estimator $\hat{\theta}(X)$ is

$$R(\theta, \hat{\theta}) = \mathbf{E} \left[\| \hat{\theta}(X) - \theta \|^2 \right] = \int_{R^p} \frac{\| \hat{\theta}(x) - \theta \|^2}{(2\pi)^{p/2}} \exp\left(-\frac{\|x - \theta\|^2}{2} \right) \mathrm{d}x.$$

The usual unbiased estimator X has the constant risk p and is minimax for $p \in \mathbb{N}$. Stein (1956) showed that there are orthogonally equivariant estimators of the form

$$\hat{\theta}_{\phi}(X) = \left(1 - \frac{\phi(\|X\|^2)}{\|X\|^2}\right) X$$
(1.2)

which dominate X when $p \ge 3$. James and Stein (1961) gave an explicit dominating procedure

$$\hat{\theta}_{\rm JS}(X) = \left(1 - \frac{p-2}{\|X\|^2}\right) X,$$
(1.3)

called the James-Stein estimator. Further, as shown in Baranchik (1964), the James-Stein estimator is inadmissible since the positive-part estimator

$$\hat{\theta}_{\rm JS}^+(X) = \max\left(0, 1 - \frac{p-2}{\|X\|^2}\right) X \tag{1.4}$$

dominates $\hat{\theta}_{JS}$. For a class of general shrinkage estimators $\hat{\theta}_{\phi}(X)$ given by (1.2), Baranchik (1970) proposed a sufficient condition for minimaxity, {**B.1** and **B.2**} where

B.1 $0 \le \phi(w) \le 2(p-2)$ for all $w \ge 0$, **B.2** $\phi'(w) \ge 0$ for all $w \ge 0$.

Further Stein (1974) expressed the risk of $\hat{\theta}_{\phi}(X)$ as

$$E[\|\hat{\theta}_{\phi} - \theta\|^{2}] = p + E[r_{\phi}(\|X\|^{2})], \qquad (1.5)$$

where

$$r_{\phi}(w) = \frac{\phi(w)}{w} \left\{ \phi(w) - 2(p-2) \right\} - 4\phi'(w).$$
(1.6)

Hence the shrinkage factor $\phi(w)$ with the inequality $r_{\phi}(w) \leq 0$ for all $w \geq 0$, implies minimaxity of $\hat{\theta}_{\phi}$. We see that **B.1** and **B.2** is a tractable sufficient condition for $r_{\phi}(w) \leq 0$ for all $w \geq 0$.

A series of papers, Efron and Morris (1971, 1972a,b, 1973), showed that the James-Stein estimator can be interpreted as an empirical Bayes estimator under $\theta \sim \mathcal{N}_p(0, \tau I_p)$. Hence the shrinkage estimator including the James-Stein estimator utilize the prior information that $\|\theta\|^2$ is relatively small. In fact, the risk function of the James-Stein estimator is

$$p - (p-2)^2 \mathbb{E}[1/||X||^2]$$
 (1.7)

which is increasing in $\|\theta\|^2$. On the other hand, the larger $\|x\|^2$ suggests that the prior information $(\|\theta\|^2$ is relatively small) is incorrect. Although the James-Stein estimator uniformly dominates X under the quadratic risk, for larger $\|x\|^2$, the unbiased estimator X seems superior to the shrinkage estimators with the bias given by

$$E\left[\left(1 - \frac{\phi(\|X\|^2)}{\|X\|^2}\right)X\right] - \theta = -E\left[\frac{\phi(\|X\|^2)}{\|X\|^2}X\right],$$
(1.8)

with $O(1/\|\theta\|)$ provided $\phi(w)$ is bounded. Note that many popular shrinkage estimators have ϕ with

$$\liminf_{w \to \infty} \phi(w) \ge p - 2. \tag{1.9}$$

See a sufficient condition for admissibility by Brown (1971).

In this paper, we define debiased shrinkage estimator by

DS.1 $\phi(w)$ is weakly differentiable with bounded $\phi'(w)$, **DS.2** For some a > 0, $0 < \phi(w) \le w$ on (0, a) and $\phi(w) = 0$ on $[a, \infty)$.

Hence the debiased shrinkage estimator shrinks towards the origin for smaller $||x||^2$ and is exactly equal to the unbiased estimator X for larger $||x||^2$. Such debiased shrinkage estimators seem superior to the unbiased estimator X, which

implies minimaxity. In this paper, we are interested in whether the debiased shrinkage estimators are minimax or not.

In the literature, there are some debiased estimators including SCAD (Smoothly Clipped Absolute Deviation) by Fan and Li (2001) and nearly unbiased estimators by MCP (Minimax Concave Penalty) by Zhang (2010), which have not necessarily aimed at enjoying the conventional minimaxity.

The organization of this paper is as follows. By (1.5), the risk difference between $\hat{\theta}$ and the minimax estimator X is given by

$$\mathbf{E}[\|\hat{\theta}_{\phi} - \theta\|^{2}] - p = \mathbf{E}[r_{\phi}(\|X\|^{2})] = \mathbf{E}\left[r_{\phi}(\|X\|^{2})I_{[0,a]}(\|X\|^{2})\right], \qquad (1.10)$$

where $r_{\phi}(w)$ is given by (1.6) and the second equality follows from **DS.2**. In Section 2, we give a useful result, Theorem 2.1, on the asymptotic behavior of this type of an expected value when $\|\theta\|^2 \to \infty$. In Section 3, we review SCAD and MCP as a solution of penalized least squares and investigate how the corresponding $\phi(w)$ approaches 0 as $w \nearrow a$. In Section 4, using Theorem 2.1, we show that the debiased shrinkage estimators with **DS.1** and **DS.2** as well as mild conditions on the way how $\phi(w)$ approaches 0 as $w \nearrow a$, are not minimax, which is not necessarily expected.

2. Asymptotic behavior of an expected value

For fixed a > 0, we investigate the asymptotic behavior of the expected value

$$G(\|\theta\|^2; a) = \mathbb{E}\left[g(\|X\|^2)I_{[0,a]}(\|X\|^2)\right] \text{ as } \|\theta\|^2 \to \infty$$
(2.1)

where g(w) satisfies **A.1** and **A.2**:

A.1 $w^{(p-1)/2}|g(w)|$ is bounded on [0, a]. **A.2** There exists a nonnegative real b such that

$$\lim_{w \neq a} \frac{g(w)}{(a-w)^b} = 1.$$
 (2.2)

Notice that, on **A.2**, we do not lose the generality even if we assume the limit of $g(w)/(a-w)^b$ is 1. If the limit is equal to $g_*(\neq 0)$, we have only to consider

$$g_* \mathbf{E}\left[\frac{g(\|X\|^2)}{g_*}I_{[0,a]}(\|X\|^2)\right].$$

Then we have the following result.

Theorem 2.1. Assume $p \ge 2$ and that g(w) satisfies **A.1** and **A.2**. Let $\nu = \|\theta\|^2$ and

$$c(a,b,p) = \frac{a^{(p-1)/4+b/2}2^b\Gamma(b+1)}{\sqrt{2\pi}\exp(a/2)}.$$
(2.3)

Then

$$\lim_{\nu \to \infty} \frac{\nu^{(p+1)/4 + b/2} e^{\nu/2}}{e^{\sqrt{\nu}\sqrt{a}}} G(\nu; a) = c(a, b, p).$$

Proof. We first prove the theorem under the proper subset of **A.1**;

A.1.1 |g(w)| is bounded on [0, a].

Note that $||X||^2$ can be decomposed as $U^2 + V$ where $U \sim N(\sqrt{\nu}, 1), V \sim \chi^2_{p-1}$ and U and V are mutually independent. Then we have

$$\begin{split} G(\nu; a) &= \mathbb{E} \left[g(U^2 + V) I_{[0,a]}(U^2 + V) \right] \\ &= \mathbb{E}_U \left[\mathbb{E}_V \left[g(U^2 + V) I_{[0,a]}(U^2 + V) \, | \, U = u \right] I_{[-\sqrt{a},\sqrt{a}]}(U) \right] \\ &= \mathbb{E}_U \left[\mathbb{E}_V \left[g(V + a - \{a - U^2\}) I_{[0,a - U^2]}(V) \, | \, U = u \right] I_{[-\sqrt{a},\sqrt{a}]}(U) \right] \\ &= \mathbb{E}_U \left[H(a - U^2)(a - U^2)^{b+q} I_{[-\sqrt{a},\sqrt{a}]}(U) \right], \end{split}$$

where q = (p-1)/2, $H(\cdot)$ is given by

$$H(y) = \frac{1}{y^{b+q}} \int_0^y g(v+a-y) f_{p-1}(v) \mathrm{d}v$$
 (2.4)

and $f_{p-1}(v)$ is the pdf of χ^2_{p-1} . Hence $G(\nu; a)$ is rewritten as

$$G(\nu; a) = \int_{-\sqrt{a}}^{\sqrt{a}} H(a - u^2)(a - u^2)^{b+q} \frac{1}{\sqrt{2\pi}} e^{-(u - \sqrt{\nu})^2/2} \mathrm{d}u.$$

Since the asymptotic behavior of $G(\nu; a)$ as $\nu \to \infty$ is of interest, $\nu > a$ is assumed in the following. For $G(\nu; a)$, apply the change of variables,

$$z = (\sqrt{\nu} - \sqrt{a})(-u + \sqrt{a})$$

which implies

$$u = \sqrt{a} - \frac{z}{\sqrt{\nu} - \sqrt{a}}, \ \mathrm{d}u = -\frac{1}{\sqrt{\nu} - \sqrt{a}}\mathrm{d}z.$$

Then we have

$$G(\nu; a)$$

$$= \frac{1}{\sqrt{2\pi}(\sqrt{\nu} - \sqrt{a})} \int_{0}^{2\sqrt{a}(\sqrt{\nu} - \sqrt{a})} \left(a - \left\{\sqrt{a} - \frac{z}{\sqrt{\nu} - \sqrt{a}}\right\}^{2}\right)^{b+q}$$

$$\times H\left(a - \left\{\sqrt{a} - \frac{z}{\sqrt{\nu} - \sqrt{a}}\right\}^{2}\right) \exp\left(-\frac{1}{2}\left\{\sqrt{a} - \frac{z}{\sqrt{\nu} - \sqrt{a}} - \sqrt{\nu}\right\}^{2}\right) \mathrm{d}z.$$

Further we rewrite it as C(x)

$$\begin{aligned} G(\nu;a) \\ &= \frac{\exp(-\{\sqrt{\nu} - \sqrt{a}\}^2/2)}{\sqrt{2\pi}(\sqrt{\nu} - \sqrt{a})} \int_0^{2\sqrt{a}(\sqrt{\nu} - \sqrt{a})} H\left(a - \left\{\sqrt{a} - \frac{z}{\sqrt{\nu} - \sqrt{a}}\right\}^2\right) \\ &\quad \times \left(\frac{2\sqrt{a}z}{\sqrt{\nu} - \sqrt{a}} - \frac{z^2}{(\sqrt{\nu} - \sqrt{a})^2}\right)^{b+q} \exp\left(-z - \frac{z^2}{2(\sqrt{\nu} - \sqrt{a})^2}\right) \mathrm{d}z \end{aligned}$$

Y. Maruyama & A. Takemura/Non-minimaxity of debiased estimators

$$=\frac{\exp(-\{\sqrt{\nu}-\sqrt{a}\}^2/2)2^{b+q}a^{(b+q)/2}}{\sqrt{2\pi}(\sqrt{\nu}-\sqrt{a})^{b+q+1}}\int_0^\infty z^{b+q}e^{-z}H_1(z;\nu)\mathrm{d}z,$$

where

$$H_{1}(z;\nu) = H\left(a - \left\{\sqrt{a} - \frac{z}{\sqrt{\nu} - \sqrt{a}}\right\}^{2}\right) \left(1 - \frac{z}{2\sqrt{a}(\sqrt{\nu} - \sqrt{a})}\right)^{b+q} \times \exp\left(-\frac{z^{2}}{2(\sqrt{\nu} - \sqrt{a})^{2}}\right) I_{(0,2\sqrt{a}(\sqrt{\nu} - \sqrt{a}))}(z).$$
(2.5)

From Part 2 of Lemma 2.1 below, H(y) on [0, a] is bounded under **A.1.1**. Hence, for any ν , we have

$$H_1(z;\nu) \begin{cases} \leq \max_{y \in [0,a]} |H(y)| & 0 \leq z \leq 2\sqrt{a}(\sqrt{\nu} - \sqrt{a}) \\ = 0 & z > 2\sqrt{a}(\sqrt{\nu} - \sqrt{a}). \end{cases}$$
(2.6)

Further, by (2.5) and Part 1 of Lemma 2.1, we have

$$\lim_{\nu \to \infty} H_1(z;\nu) = \lim_{y \to 0} H(y) = \frac{\Gamma(b+1)}{\Gamma(b+(p+1)/2)2^{(p-1)/2}}.$$
 (2.7)

By (2.6) and (2.7), the dominated convergence theorem, gives

$$\lim_{\nu \to \infty} \int_0^\infty z^{b+q} e^{-z} H_1(z;\nu) \mathrm{d}z = \int_0^\infty z^{b+q} e^{-z} \lim_{\nu \to \infty} H_1(z;\nu) \mathrm{d}z = \frac{\Gamma(b+1)}{2^{(p-1)/2}},$$

which completes the proof under **A.1.1**.

Now we assume **A.1**, that is, $w^{(p-1)/2}|g(w)|$ is bounded on [0, a] as

$$w^{(p-1)/2}|g(w)| < M.$$
 (2.8)

Let $f_p(w,\nu)$ be the density of $W = ||X||^2$. Note that $f_p(w,\nu)/f_p(w,0)$ for any fixed $\nu > 0$ is increasing in w and that $w^{-(p-1)/2}$ is decreasing in w. By the correlation inequality, we have

$$\begin{aligned} \left| \int_{0}^{a/2} g(w) f_{p}(w,\nu) \mathrm{d}w \right| &< \int_{0}^{a/2} |g(w)| f_{p}(w,\nu) \mathrm{d}w \\ &< M \int_{0}^{a/2} w^{-(p-1)/2} f_{p}(w,\nu) \mathrm{d}w \\ &= M \int_{0}^{a/2} w^{-(p-1)/2} \frac{f_{p}(w,\nu)}{f_{p}(w,0)} f_{p}(w,0) \mathrm{d}w \\ &\leq M \frac{\int_{0}^{a/2} w^{-(p-1)/2} f_{p}(w,0) \mathrm{d}w}{\int_{0}^{a/2} f_{p}(w,0) \mathrm{d}w} \int_{0}^{a/2} f_{p}(w,\nu) \mathrm{d}w \end{aligned}$$

Y. Maruyama & A. Takemura/Non-minimaxity of debiased estimators

$$\leq M_1 \int_0^{a/2} f_p(w,\nu) \mathrm{d}w$$

where

$$M_1 = M \frac{\int_0^{a/2} w^{-(p-1)/2} f_p(w,0) \mathrm{d}w}{\int_0^{a/2} f_p(w,0) \mathrm{d}w}.$$

Let

$$g_L(w) = \begin{cases} -M_1 & 0 \le w \le a/2\\ g(w) & a/2 < w \le a, \end{cases} \quad g_U(w) = \begin{cases} M_1 & 0 \le w \le a/2\\ g(w) & a/2 < w \le a, \end{cases}$$

which are both bounded. Then we have

$$\mathbf{E}\left[g_L(W)I_{[0,a]}(W)\right] < G(\nu;a) < \mathbf{E}\left[g_U(W)I_{[0,a]}(W)\right]$$

and, by the result under A.1.1,

$$\lim_{\nu \to \infty} \frac{\nu^{(p+1)/4+b/2} e^{\nu/2}}{e^{\sqrt{\nu}\sqrt{a}}} \mathbb{E}\left[g_L(W)I_{[0,a]}(W)\right]$$
$$= \lim_{\nu \to \infty} \frac{\nu^{(p+1)/4+b/2} e^{\nu/2}}{e^{\sqrt{\nu}\sqrt{a}}} \mathbb{E}\left[g_U(W)I_{[0,a]}(W)\right] = c(a,b,p),$$

where c(a, b, p) is given by (2.3). Hence Theorem 2.1 is valid for the case where $w^{(p-1)/2}|g(w)|$ is bounded.

The following lemma gives some properties on H(y) given by (2.4), needed in the proof of Theorem 2.1.

Lemma 2.1. We assume that |g(w)| is bounded on [0, a] as in A.1.1. Then we have the following results.

1.
$$\lim_{y \to 0} H(y) = \frac{\Gamma(b+1)}{\Gamma(b+(p+1)/2)2^{(p-1)/2}}.$$

2. $H(y)$ on $[0, a]$ is bounded.

Proof. By (2.2), for any $\epsilon > 0$, there exists $\delta_1(\epsilon) > 0$ such that

$$(1-\epsilon)(a-w)^b \le g(w) \le (1+\epsilon)(a-w)^b$$

for all $a - \delta_1 \leq w < a$ and hence

$$(1 - \epsilon)(y - v)^b \le g(v + a - y) \le (1 + \epsilon)(y - v)^b$$
(2.9)

for all $0 < v < y \le \delta_1(\epsilon)$. Further, for any $0 < \epsilon < 1$, we have $1 - \epsilon \le e^{-v/2}$ for all $0 < v \le \delta_2(\epsilon)$ where $\delta_2(\epsilon) = -2\log(1-\epsilon)$ and hence

$$(1-\epsilon)\frac{v^{q-1}}{\Gamma(q)2^q} \le f_{p-1}(v) \le \frac{v^{q-1}}{\Gamma(q)2^q}$$

 $\mathbf{6}$

with q = (p-1)/2 and for all $v \leq \delta_2(\epsilon)$. Then for any $\epsilon > 0$ and all $0 < y \leq \min(\delta_1(\epsilon), \delta_2(\epsilon))$, we have

$$(1-\epsilon)^2 \int_0^y \frac{(y-v)^b}{y^{b+q}} \frac{v^{q-1}}{\Gamma(q)2^q} \mathrm{d}v \le H(y) \le (1+\epsilon) \int_0^y \frac{(y-v)^b}{y^{b+q}} \frac{v^{q-1}}{\Gamma(q)2^q} \mathrm{d}v,$$

where

$$\int_0^y \frac{(y-v)^b}{y^{b+q}} v^{q-1} \mathrm{d}v = B(q,b+1) = B((p-1)/2,b+1).$$
(2.10)

Hence, for $0 < y \leq \min(\delta_1(\epsilon), \delta_2(\epsilon))$, we have

$$\frac{(1-\epsilon)^2 \Gamma(b+1)}{\Gamma(b+(p+1)/2) 2^{(p-1)/2}} \le H(y) \le \frac{(1+\epsilon) \Gamma(b+1)}{\Gamma(b+(p+1)/2) 2^{(p-1)/2}}.$$

and the part 1 follows.

By (2.4), we have

$$H(a) = \frac{1}{a^{b+(p-1)/2}} \int_0^a g(v) f_{p-1}(v) \mathrm{d}v, \qquad (2.11)$$

which is bounded under A.1.1. By the continuity of H(y) and Part 1 of this lemma, the part 2 follows.

3. Review of existing debiased shrinkage estimators

As we mentioned in Section 1, in the literature, there are some "debiased shrinkage" estimators including SCAD (Smoothly Clipped Absolute Deviation) by Fan and Li (2001) and nearly unbiased estimators by MCP (Minimax Concave Penalty) by Zhang (2010), although they do not necessarily aim at enjoying the conventional minimaxity. In this section, we assume p = 1 and review existing estimators as solutions of the penalized least squares problem;

$$\hat{\theta}(P;\lambda) = \underset{\theta}{\arg\min}\left\{(\theta - x)^2 + P(|\theta|;\lambda)\right\}.$$
(3.1)

Table 1 summarizes three popular penalty functions $P(|\theta|; \lambda)$, and the corresponding minimizers "ridge", "soft thresholding" and "hard thresholding". For the three estimators, the corresponding shrinkage factors, $\phi(x^2)$, from the form

$$\hat{\theta} = \left(1 - \frac{\phi(x^2)}{x^2}\right)x \tag{3.2}$$

are

$$\phi_{\mathrm{R}}(w) = \frac{w}{\lambda + 1}, \quad \phi_{\mathrm{ST}}(w) = \begin{cases} w & w \le \lambda^2 \\ \lambda w^{1/2} & w > \lambda^2 \end{cases}, \quad \phi_{\mathrm{HT}}(w) = \begin{cases} w & w \le \lambda^2 \\ 0 & w > \lambda^2 \end{cases}.$$

Y. Maruyama & A. Takemura/Non-minimaxity of debiased estimators

	$P(heta ;\lambda)$	$\hat{ heta}(P;\lambda)$	ϕ
ridge	$\lambda \theta^2$	$\hat{\theta}_{\mathrm{R}} = (1 - 1/\{\lambda + 1\})x$	DS.1
soft thresholding	$\lambda heta $	$\hat{\theta}_{\rm ST} = \begin{cases} 0 & x^2 \le \lambda^2 \\ (1 - \lambda/ x)x & x^2 > \lambda^2 \end{cases}$	DS.1
hard thresholding	$\lambda^2 - (\theta - \lambda)^2 I_{[0,\lambda]}(\theta)$	$\hat{ heta}_{ m HT} = egin{cases} 0 & x^2 \leq \lambda^2 \ x & x^2 > \lambda^2 \end{cases}$	DS.2

 TABLE 1

 Ridge, Soft-thresholding, Hard-thresholding

We see that **DS.2**, **DS.2** and **DS.1** are not satisfied by $\phi_{\rm R}(w)$, $\phi_{\rm ST}(w)$ and $\phi_{\rm HT}(w)$, respectively.

SCAD (Smoothly Clipped Absolute Deviation) by Fan and Li (2001) is the minimizer, (3.1), with the continuous differentiable penalty function defined by

$$P'(|\theta|;\lambda,\alpha) = \begin{cases} \lambda & |\theta| < \lambda \\ \frac{\alpha\lambda - |\theta|}{\alpha - 1} & \lambda \le |\theta| < \alpha\lambda \\ 0 & |\theta| \ge \alpha\lambda \end{cases}$$
(3.3)

where $\alpha > 2$. The resulting solution is

$$\hat{\theta}_{\text{SCAD}}(\lambda;\alpha) = \begin{cases} 0 & 0 < x^2 < \lambda^2 \\ (1-\lambda/|x|)x & \lambda^2 \le x^2 \le 4\lambda^2 \\ \left(1 - \frac{-x^2 + \alpha\lambda|x|}{(\alpha - 2)x^2}\right)x & 4\lambda^2 \le x^2 \le \alpha^2\lambda^2 \\ x & x^2 \ge \alpha^2\lambda^2, \end{cases}$$
(3.4)

where the corresponding shrinkage factor is

$$\phi_{\text{SCAD}}(w) = \begin{cases} w & 0 < w < \lambda^2 \\ \lambda w^{1/2} & \lambda^2 \le w \le 4\lambda^2 \\ \frac{-w + \alpha \lambda w^{1/2}}{a - 2} & 4\lambda^2 \le w \le \alpha^2 \lambda^2 \\ 0 & w \ge \alpha^2 \lambda^2. \end{cases}$$
(3.5)

We see that $\phi_{\text{SCAD}}(w)$ satisfies both **DS.1** and **DS.2**. Further, by (3.5), the derivative at $w = \alpha^2 \lambda^2$ is

$$\frac{\mathrm{d}}{\mathrm{d}w}\phi_{\mathrm{SCAD}}(w)|_{w=\alpha^2\lambda^2} = -\frac{1}{2(\alpha-2)} < 0. \tag{3.6}$$

As pointed in Strawderman and Wells (2012), the nearly unbiased estimator by MCP (Minimax Concave Penalty) considered in Zhang (2010) is equivalent to the minimizer of (3.1) with the continuous differentiable penalty function defined by

$$P'(|\theta|;\lambda,\alpha) = \begin{cases} 2\{\lambda - |\theta|/\alpha\} & |\theta| < \alpha\lambda\\ 0 & |\theta| \ge \alpha\lambda \end{cases}$$
(3.7)

where $\alpha > 1$. Then the resulting solution is given by

$$\hat{\theta}_{\mathrm{MCP}}(\lambda;\alpha) = \begin{cases} 0 & 0 < x^2 < \lambda^2 \\ \left(1 - \frac{-x^2 + \alpha\lambda|x|}{(\alpha - 1)x^2}\right) x & \lambda^2 \le x^2 \le \alpha^2\lambda^2 \\ x & x^2 \ge \alpha^2\lambda^2, \end{cases}$$
(3.8)

where the corresponding shrinkage factor is

$$\phi_{\rm MCP}(w) = \begin{cases} w & 0 < w < \lambda^2 \\ \frac{-w + \alpha \lambda w^{1/2}}{\alpha - 1} & \lambda^2 \le w \le \alpha^2 \lambda^2 \\ 0 & w \ge \alpha^2 \lambda^2. \end{cases}$$
(3.9)

We see that $\phi_{MCP}(w)$ satisfies both **DS.1** and **DS.2**. Further, by (3.9), the derivative at $w = \alpha^2 \lambda^2$ is

$$\frac{\mathrm{d}}{\mathrm{d}w}\phi_{\mathrm{MCP}}(w)|_{w=\alpha^{2}\lambda^{2}} = -\frac{1}{2(\alpha-1)} < 0.$$
(3.10)

By (3.6) and (3.10), both $\phi_{\text{SCAD}}(w)$ and $\phi_{\text{MCP}}(w)$ approach 0 as $w \nearrow \alpha^2 \lambda^2$ with the negative slope.

Aside from the justification as a solution of the penalized least squares problem (3.1), let us consider

$$\phi_{\mathbf{Q}}(w) = \begin{cases} w & 0 \le w < \frac{2a+1-\sqrt{4a+1}}{2} \\ (a-w)^2 & \frac{2a+1-\sqrt{4a+1}}{2} \le w \le a \\ 0 & w > a. \end{cases}$$
(3.11)

We see that $\phi_{\mathbf{Q}}(w)$ satisfies both **DS.1** and **DS.2** and

$$\lim_{w \to a} \frac{\mathrm{d}}{\mathrm{d}w} \phi_{\mathrm{Q}}(w) = 0 \quad \text{and} \quad \lim_{w \nearrow a} (a - w) \frac{(\mathrm{d}/\mathrm{d}w)\phi_{\mathrm{Q}}(w)}{\phi_{\mathrm{Q}}(w)} = -2.$$
(3.12)

When $\phi(w)$ of debiased shrinkage estimator approaches 0 from above as $w \to a$, it seems that both $\{(3.6) \text{ and } (3.10)\}$ and (3.12) are typical behaviors characterized by $\phi'(w)$.

4. Main result

In this section, we investigate the minimaxity of the shrinkage debiased estimators with **DS.1** and **DS.2**. Recall, as in (1.10), the risk difference between $\hat{\theta}$ and the minimax estimator X is

$$\mathbf{E}\left[\|\hat{\theta}_{\phi} - \theta\|^{2}\right] - p = \mathbf{E}\left[r_{\phi}(\|X\|^{2})I_{[0,a]}(\|X\|^{2})\right],\tag{4.1}$$

where $r_{\phi}(w)$ is given by (1.6). Under the assumptions on $\phi(w)$, **DS.1** and **DS.2**, $r_{\phi}(w)$ given by (1.6) is bounded, that is, there exists an M such that

$$|r_{\phi}(w)| < M \text{ on } [0, a].$$
 (4.2)

For $\phi(w)$ with $\lim_{w \nearrow a} \phi(w) = 0$ as wells as $\phi(w) > 0$ for w < a, we consider two cases as a generalization of $\{(3.6) \text{ and } (3.10)\}$ and (3.12):

Case 1 $\limsup_{w \nearrow a} \phi'(w) < 0.$

Case 2 $\lim_{w \nearrow a} \phi'(w) = 0$ and there exist $0 < \epsilon < 1$ and $\gamma > 1$ such that

$$-\gamma < (a-w)\frac{\phi'(w)}{\phi(w)} < -\frac{1}{\gamma},\tag{4.3}$$

for all $w \in (\epsilon a, a)$.

Under Case 1, there exist $\delta_1 > 0$ and $0 < \delta_2 < 1$ such that

$$\phi'(w) < -\delta_1 \text{ and } \frac{\phi(w)}{w} \{\phi(w) - 2(p-2)\} > -\delta_1,$$
 (4.4)

for all $w \in (\delta_2 a, a)$. Then, by (1.6) and (4.4), we have

$$r_{\phi}(w) = \frac{\phi(w)}{w} \left\{ \phi(w) - 2(p-2) \right\} - 4\phi'(w) \ge 3\delta_1 > 0, \tag{4.5}$$

for all $w \in (\delta_2 a, a)$. Hence, by Theorem 2.1 with (4.1), (4.2) and (4.5), we have

$$\liminf_{\nu \to \infty} \frac{\nu^{(p+1)/4} e^{\nu/2}}{e^{\sqrt{\nu}\sqrt{a}}} \left\{ \mathbf{E} \left[\|\hat{\theta}_{\phi} - \theta\|^2 \right] - p \right\}$$

$$\geq 3\delta_1 c(a, 0, p) > 0, \qquad (4.6)$$

which implies that the debiased shrinkage estimator is not minimax under **Case 1**.

Under Case 2, the inequality

$$-\gamma < (a-w)\frac{\phi'(w)}{\phi(w)}$$

for $w \in (\epsilon a, a)$ implies

$$\int_{\epsilon a}^{w} \frac{\phi'(t)}{\phi(t)} \mathrm{d}t > \int_{\epsilon a}^{w} \frac{-\gamma}{a-t} \mathrm{d}t$$

which is equivalent to

$$\phi(w) > \phi_*(a-w)^{\gamma}, \text{ where } \phi_* = \frac{\phi(\epsilon a)}{(a-\epsilon a)^{\gamma}},$$

$$(4.7)$$

for all $w \in (\epsilon a, a)$. Further let

$$\epsilon' = \frac{1}{1 + 1/\{(p-2)\gamma\}}.$$
(4.8)

Then, for $w \in (\epsilon' a, a)$, we have

$$-\frac{2(p-2)(a-w)}{w} \ge -\frac{2}{\gamma}.$$
(4.9)

Hence, for $w \in (\max(\epsilon, \epsilon')a, a)$, we have

$$r_{\phi}(w) = \frac{\{\phi(w)\}^2}{w} - 2(p-2)\frac{\phi(w)}{w} - 4\phi'(w)$$

$$\geq -2(p-2)\frac{\phi(w)}{w} - 4\phi'(w)$$

$$= \frac{\phi(w)}{a-w} \left\{ -\frac{2(p-2)(a-w)}{w} - 4(a-w)\frac{\phi'(w)}{\phi(w)} \right\}$$

$$\geq \frac{2}{\gamma} \frac{\phi(w)}{a-w},$$
(4.10)

where the second inequality follows from (4.3) and (4.9). Further, by (4.7) and (4.10), we have

$$r_{\phi}(w) \ge \frac{2\phi_*}{\gamma} (a-w)^{\gamma-1} \tag{4.11}$$

for $w \in (\max(\epsilon, \epsilon')a, a)$. Hence, by Theorem 2.1 with (4.1), (4.2) and (4.11), we have

$$\begin{split} \liminf_{\nu \to \infty} \frac{\nu^{(p+1)/4 + (\gamma - 1)/2} e^{\nu/2}}{e^{\sqrt{\nu}\sqrt{a}}} \left\{ \mathbf{E} \left[\|\hat{\theta}_{\phi} - \theta\|^2 \right] - p \right\} \\ \geq \frac{2\phi_*}{\gamma} c(a, \gamma - 1, p) > 0, \end{split}$$

which implies that the debiased shrinkage estimator not minimax under **Case 2**. In summary, we have the following theorem.

Theorem 4.1. The debiased shrinkage estimator with DS.1 and DS.2 is not minimax under either Case 1 or Case 2.

Remark 4.1. Yet another application of Theorem 2.1 is also related to Stein estimation, the gain of the positive-part estimator $\hat{\theta}_{JS}^+$ given by (1.4) over the naive James-Stein estimator $\hat{\theta}_{JS}$ given by (1.3). For these estimators, the corresponding $\phi(w)$ are given by

$$\phi_{\rm JS}^+(w) = \min(w, p-2), \quad \phi_{\rm JS}(w) = p-2.$$
 (4.12)

By the general expression of the risk, (1.5) and (1.6) with (4.12), we have

$$R(\theta, \hat{\theta}_{\rm JS}) - R(\theta, \hat{\theta}_{\rm JS}^+) = \mathrm{E}\Big[\Big\{-\frac{(p-2)^2}{\|X\|^2} + 2p - \|X\|^2\Big\}I_{[0,p-2]}(\|X\|^2)\Big].$$
(4.13)

Let

$$f_k(v) = \frac{v^{k/2-1}e^{-v/2}}{\Gamma(k/2)2^{k/2}}, \quad f_k(v;\nu) = \sum_{i=0}^{\infty} \frac{(\nu/2)^i \exp(-\nu/2)}{i!} f_{k+2i}(v),$$

11

Y. Maruyama & A. Takemura/Non-minimaxity of debiased estimators

$$F_k(v) = \int_0^v f_k(w) \mathrm{d}w, \quad F_k(v;\nu) = \int_0^v f_k(w;\nu) \mathrm{d}w.$$

Hansen (2022), in Theorem 15.7, expressed the risk difference (4.13) through $F_k(v)$ and $F_k(v; \nu)$, the distribution functions of the central chi-square with and non-central chi-square, as

$$R(\theta, \hat{\theta}_{\rm JS}) - R(\theta, \hat{\theta}_{\rm JS}^+) = 2pF_p(p-2;\nu) - pF_{p+2}(p-2;\nu) - \nu F_{p+4}(p-2;\nu) - (p-2)^2 \sum_{i=0}^{\infty} \frac{(\nu/2)^i \exp(-\nu/2)}{i!} \frac{F_{p+2i-2}(p-2)}{p+2i-2}.$$

Robert (1988) expressed the risk difference (4.13) through the Dawson integral given by

$$D(\lambda) = e^{-\lambda^2} \int_0^{\lambda} e^{t^2} \mathrm{d}t.$$

The results by Hansen (2022) and Robert (1988), do not seem to directly provide the exact asymptotic order of the major term of (4.13) with the exact coefficient.

Using Theorem 2.1, we can get it as follows. Since

$$\left\{-\frac{(p-2)^2}{w} + 2p - w\right\}\Big|_{w=p-2} = 4,$$

and

$$w^{(p-1)/2} \left| -\frac{(p-2)^2}{w} + 2p - w \right| = w^{(p-3)/2} \left| -(p-2)^2 + 2pw - w^2 \right|$$

is bounded for $w \in (0, p-2)$ and for $p \ge 3$, Theorem 2.1 gives

$$\begin{split} &\lim_{\nu \to \infty} \frac{\nu^{(p+1)/4} e^{\nu/2}}{e^{\sqrt{\nu}\sqrt{p-2}}} \big\{ R(\theta, \hat{\theta}_{\rm JS}) - R(\theta, \hat{\theta}_{\rm JS}^+) \big\} \\ &= 4c(p-2, 0, p) = 4 \frac{(p-2)^{(p-1)/4}}{\sqrt{2\pi} \exp(p/2 - 1)}. \end{split}$$

References

- BARANCHIK, A. J. (1964). Multiple regression and estimation of the mean of a multivariate normal distribution Technical Report No. 51, Department of Statistics, Stanford University.
- BARANCHIK, A. J. (1970). A family of minimax estimators of the mean of a multivariate normal distribution. Ann. Math. Statist. **41** 642–645. MR0253461
- BROWN, L. D. (1971). Admissible estimators, recurrent diffusions, and insoluble boundary value problems. Ann. Math. Statist. 42 855–903. MR0286209
- EFRON, B. and MORRIS, C. (1971). Limiting the risk of Bayes and empirical Bayes estimators. I. The Bayes case. J. Amer. Statist. Assoc. 66 807–815.

- EFRON, B. and MORRIS, C. (1972a). Limiting the risk of Bayes and empirical Bayes estimators. II. The empirical Bayes case. J. Amer. Statist. Assoc. 67 130–139.
- EFRON, B. and MORRIS, C. (1972b). Empirical Bayes on vector observations: an extension of Stein's method. *Biometrika* **59** 335–347.
- EFRON, B. and MORRIS, C. (1973). Stein's estimation rule and its competitors—an empirical Bayes approach. J. Amer. Statist. Assoc. 68 117–130.
- FAN, J. and LI, R. (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. J. Amer. Statist. Assoc. 96 1348–1360. MR1946581
- HANSEN, B. E. (2022). *Probability and Statistics for Economists*. Princeton Univ Press, Princeton, NJ.
- JAMES, W. and STEIN, C. (1961). Estimation with quadratic loss. In Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. I 361–379. Univ. California Press, Berkeley, Calif. MR0133191
- ROBERT, C. (1988). An explicit formula for the risk of the positive-part James-Stein estimator. Canad. J. Statist. 16 161–168. MR963730
- STEIN, C. (1956). Inadmissibility of the usual estimator for the mean of a multivariate normal distribution. In *Proceedings of the Third Berkeley Symposium* on Mathematical Statistics and Probability, 1954–1955, vol. I 197–206. University of California Press, Berkeley and Los Angeles. MR0084922
- STEIN, C. (1974). Estimation of the mean of a multivariate normal distribution. In Proceedings of the Prague Symposium on Asymptotic Statistics (Charles Univ., Prague, 1973), Vol. II 345–381. Charles Univ., Prague. MR0381062
- STRAWDERMAN, R. L. and WELLS, M. T. (2012). On hierarchical prior specifications and penalized likelihood. In Contemporary developments in Bayesian analysis and statistical decision theory: a Festschrift for William E. Strawderman. Inst. Math. Stat. (IMS) Collect. 8 154–180. Inst. Math. Statist., Beachwood, OH. MR3202509
- ZHANG, C.-H. (2010). Nearly unbiased variable selection under minimax concave penalty. Ann. Statist. 38 894–942. MR2604701