

GLOBAL MINIMIZERS OF THE TWO-PHASE BERNOULLI PROBLEM WITH THE p -LAPLACE OPERATOR

MASOUD BAYRAMI AND MORTEZA FOTOUHI

ABSTRACT. In this paper, we study the classification of Lipschitz global solutions for a two-phase p -Laplace Bernoulli problem, subject to a mild assumption. Specifically, we focus on the scenario where the *interior* two-phase points of the global solution are non-empty. Our results show that the expected $C^{1,\eta}$ regularity holds in a suitable neighborhood of certain two-phase points referred to as *regular* two-phase points.

1. INTRODUCTION AND MAIN RESULT

We study the regularity of the free boundary, obtaining from the minimization of the following two-phase functional

$$(1) \quad J_{\text{TP}}(v, D) := \int_D |\nabla v|^p + (p-1)\lambda_+^p \chi_{\{v>0\}} + (p-1)\lambda_-^p \chi_{\{v<0\}} dx, \quad v \in \mathcal{K}.$$

Here, D is a bounded and smooth domain in \mathbb{R}^n , χ_A denotes the characteristic function of the set A , the positive constants $\lambda_+, \lambda_- > 0$ are given, and $1 < p < \infty$. The class of admissible functions \mathcal{K} consists of all functions $v \in W^{1,p}(D)$ with boundary data $v = g$ on ∂D for a prescribed boundary datum $g \in W^{1,p}(D)$.

Before proceeding, we need to review some common terminologies and definitions:

- A function $u : D \rightarrow \mathbb{R}$ is said to be a *minimizer* of J_{TP} in D if and only if

$$J_{\text{TP}}(u, D) \leq J_{\text{TP}}(v, D),$$

for all $v \in \mathcal{K}$.

- The two sets of points $\Omega_u^+ = \{x \in D : u(x) > 0\}$ and $\Omega_u^- = \{x \in D : u(x) < 0\}$ are respectively referred to as the positivity sets of the functions $u^+ := \max\{u, 0\}$ and $u^- := \max\{-u, 0\}$.
- $F(u) := (\partial\Omega_u^+ \cup \partial\Omega_u^-) \cap D$ represents the free boundary of the minimizer u .
- The set $\Gamma_{\text{TP}}(u) := \partial\Omega_u^+ \cap \partial\Omega_u^- \cap D$ is the set of two-phase points of the free boundary $F(u)$, and for the simplicity, we denote it by Γ_{TP} .
- The boundary of positive and negative phases, i.e. $\partial\Omega_u^\pm \cap D$ can be decomposed as

$$\partial\Omega_u^\pm \cap D = \Gamma_{\text{OP}}^\pm \cup \Gamma_{\text{TP}},$$

Date: June 9, 2023.

2020 Mathematics Subject Classification. 35R35, 35B65, 35J60, 35J70.

Key words and phrases. Two-phase Bernoulli problem, singular/degenerate operator, viscosity solutions, Lipschitz global minimizers.

M. Bayrami and M. Fotouhi was supported by Iran National Science Foundation (INSF) under project No. 4001885.

where $\Gamma_{\text{OP}}^+ := (\partial\Omega_u^+ \setminus \partial\Omega_u^-) \cap D$ and $\Gamma_{\text{OP}}^- := (\partial\Omega_u^- \setminus \partial\Omega_u^+) \cap D$ are the one-phase parts of $F(u)$.

- We say that $x_0 \in \Gamma_{\text{TP}}$ is an *interior* two-phase point and denote it by $x_0 \in \Gamma_{\text{TP}}^{\text{int}}$, if

$$|B_r(x_0) \cap \{u = 0\}| = 0, \quad \text{for some } r > 0.$$

- We say that $x_0 \in \Gamma_{\text{TP}}$ is a *branch* point, denoted by $x_0 \in \Gamma_{\text{TP}}^{\text{br}}$, if

$$|B_r(x_0) \cap \{u = 0\}| > 0, \quad \text{for every } r > 0.$$

Any minimizer u satisfies, in a certain weak sense, the following problem

$$(2) \quad \begin{cases} \Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0, & \text{in } \Omega_u^+ \cup \Omega_u^-, \\ |\nabla u^+|^p - |\nabla u^-|^p = \lambda_+^p - \lambda_-^p, \quad |\nabla u^+| \geq \lambda_+, \quad |\nabla u^-| \geq \lambda_-, & \text{on } \Gamma_{\text{TP}}, \\ |\nabla u^+| = \lambda_+, & \text{on } \Gamma_{\text{OP}}^+, \\ |\nabla u^-| = \lambda_-, & \text{on } \Gamma_{\text{OP}}^-, \end{cases}$$

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is the p -Laplace operator; see [8, Lemma 3.1]. These types of problems are known as Bernoulli-type free boundary problems, which appear in various models of fluid mechanics or heat conduction; see e.g. [2, 4, 5, 7, 26].

For the admissible functions in $\mathcal{K}^+ := \{v \in \mathcal{K} : v \geq 0\}$, the analogous one-phase functional and the corresponding overdetermined problem called the one-phase Bernoulli problem, was first studied in [1] for the case $p = 2$. Also, the case of uniformly elliptic quasilinear equations has been treated in [3]. For the earlier known results of the one-phase Bernoulli's problem for the p -Laplace operator, see [15, 16, 25, 30]. See also [23] for the $p(x)$ -Laplacian Bernoulli problem with the right hand side. It is worth noting that, generally, in the one-phase Bernoulli problem, the $(n-1)$ -Hausdorff dimension of the singular points of the free boundary is zero and that the free boundary is analytic in a suitable neighborhood of the regular points, [15].

The two-phase problem, in the case $p = 2$, was first studied in [6], followed by a series of works by Luis Caffarelli, [10, 11, 12]; see also [14]. A comprehensive study and the review of the earlier known results for $p = 2$ can be found in [17, 31].

The challenge with problem (2) is that the governing operator, $\Delta_p u$, is not uniformly elliptic. However, the regularity of the free boundary implies non-degeneracy of $|\nabla u|$ near the free boundary, which makes the p -Laplacian uniformly elliptic. Without regularity, it is difficult to prove non-degeneracy up to the free boundary.

For the minimizers of the following functional instead of (1)

$$J(u, D) := \int_D |\nabla u|^p + \lambda_+^p \chi_{\{u>0\}} + \lambda_-^p \chi_{\{u \leq 0\}} dx = \int_D |\nabla u|^p + (\lambda_+^p - \lambda_-^p) \chi_{\{u>0\}} dx + \lambda_-^p |D|,$$

or with neglecting the constant term $\lambda_-^p |D|$, and denoting $\lambda_+^p - \lambda_-^p$ with Λ , i.e. considering the functional

$$(3) \quad u \mapsto \int_D |\nabla u|^p + \Lambda \chi_{\{u>0\}} dx,$$

the best known results comes from the stratification argument in [21]. The important fact about the minimizers of the functional (3) is that their free boundary does not contain any branch points. In [28, 29], the authors show that the Lipschitz

free boundaries of the minimizers of (3) are $C^{1,\eta}$ smooth for some $\eta \in (0, 1)$. Moreover, in [21], it is also showed that the $(n - 1)$ -Hausdorff dimension of the singular points of the free boundary is zero.

For the free boundary of the minimizers of the functional (1), which appearance of the branch point is expected, there is a complete regularity result in the case $p = 2$; see [17]. In this work, the authors proved that a suitable neighbourhood of the two-phase points (i.e. both interior and branch points) is $C^{1,\eta}$ regular. The generalization of this result, for any $p \in (1, \infty)$, is highly nontrivial due to the nonlinear and degenerate nature of the p -Laplace operator, as well as the lack of the monotonicity formula, for $p \neq 2$.

In [8], we extend the main result of [21] to functional (1), which covers the analysis of the behavior of the minimizers near the branch points as well. In fact, we prove the local Lipschitz regularity of the minimizers using a dichotomy argument. More precisely, we first establish $C^{1,\eta}$ regularity of the flat part of the free boundary. Then, we show that either the free boundary is flat or we have linear growth at the non-flat points of the free boundary. In order to prove that flatness implies $C^{1,\eta}$ regularity, the linearization technique of De Silva, [18], is used. It is vital to emphasize that in [8], we use a linear improvement of flatness to obtain the compactness of the linearizing sequences. However, in this paper we require a somewhat stronger, a quadratic improvement of flatness.

1.1. Main results. Our goal in this paper is to continue our previous research, [8], by classifying the Lipschitz global minimizers of (1) (which they are also Lipschitz solutions of (2) in $D = \mathbb{R}^n$). Our first main result is presented in the following classification theorem.

Theorem 1.1 (Classification of Lipschitz global two-phase solutions). *Let u be a Lipschitz global viscosity solution to (2), i.e. with $D = \mathbb{R}^n$. Assume that $\Gamma_{\text{TP}}^{\text{int}}(u) \neq \emptyset$ (without loss of generality, assume that $0 \in \Gamma_{\text{TP}}^{\text{int}}(u)$), then u is a two-plane solution*

$$u(x) = \alpha (x \cdot v)^+ - \beta (x \cdot v)^-,$$

for some $v \in \mathbb{S}^{n-1}$, with

$$\alpha \geq \lambda_+, \quad \beta \geq \lambda_-, \quad \alpha^p - \beta^p = \lambda_+^p - \lambda_-^p.$$

In order to prove Theorem 1.1, we will closely follow the work of De Silva and Savin, [20]. As a result of this classification of Lipschitz global solutions to (2), the regularity of the free boundary $F(u) = (\partial\Omega_u^+ \cup \partial\Omega_u^-) \cap D$, for any minimizers of J_{TP} , around the regular two-phase points, will be deduced.

Definition 1.1. *We say that $x_0 \in \Gamma_{\text{TP}}$ is a regular two-phase point if*

$$\liminf_{r \rightarrow 0} \frac{|B_r(x_0) \cap \{u = 0\}|}{|B_r(x_0)|} = 0.$$

More precisely, we prove that, in a suitable neighborhood of the regular two-phase points, the sets Ω_u^+ and Ω_u^- are two $C^{1,\eta}$ regular domains touching along the closed set of two-phase points Γ_{TP} .

Theorem 1.2 (Regularity of the regular part of the free boundary). *Let $u : D \rightarrow \mathbb{R}$ be a minimizer of J_{TP} in D . Then, for every regular two-phase point x_0 , there exists a radius $r_0 > 0$ (depending on x_0) such that $\partial\Omega_u^\pm \cap B_{r_0}(x_0)$ are $C^{1,\eta}$ graphs, for any $\eta \in (0, \frac{1}{3})$.*

Finally, notice that once we have $C^{1,\eta}$ regularity on two-phase points, it follows that $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is uniformly elliptic locally near the two-phase points (recall the boundary condition in (2)) and we obtain the analyticity by [27].

1.2. Notation. Some frequently used notations are gathered in the following table.

| | |
|--------------------------------------|---|
| c, c_0, C, C_0, \dots | generic constants |
| \bar{U} | the closure of a set U |
| ∂U | the boundary of a set U |
| x' | $x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}$ for $x = (x', x_n) \in \mathbb{R}^n$ |
| $B_r(x), B_r$ | the ball centered at x with radius $r > 0$, and $B_r := B_r(0)$ |
| a^\pm | $(a^+)^p - (a^-)^p = \lambda_+^p - \lambda_-^p$, $a^\pm \in [L_0, L_1]$ |
| $U_{a^+, \nu}(x)$ | $U_{a^+, \nu}(x) := a^+(x \cdot \nu)^+ - a^-(x \cdot \nu)^-$ two-plane solutions to (2) |
| $\mathcal{U}_{\lambda_+, \lambda_-}$ | the set of all two-plane solutions to (2) |
| ν, \mathbf{e}_n | $\nu \in \mathbb{S}^{n-1}$ and $\mathbf{e}_n = (0, \dots, 0, 1) \in \mathbb{R}^n$ |
| $M = (M_{ij})$ | $M = (M_{ij}) \in \mathcal{S}^{n \times n}$ a symmetric matrix |
| $\operatorname{tr}(M)$ | trace of matrix M |
| $P_{M, \nu}(x)$ | $P_{M, \nu}(x) := x \cdot \nu + \frac{1}{2} x^T M x$ |
| $V_{a^+, M, \nu}(x)$ | $V_{a^+, M, \nu}(x) := a^+ P_{M, \nu}^+(x) - a^- P_{M, \nu}^-(x)$ |
| ∂_n | partial derivative in the \mathbf{e}_n -direction |
| ∂_{nn} | second partial derivative in the \mathbf{e}_n -direction |
| ∂_n^\pm | partial derivative in the \mathbf{e}_n -direction from the side $\{\pm x_n > 0\}$ |
| $\mathcal{L}_p(u)$ | the linearized operator $\mathcal{L}_p(u) := \Delta u + (p-2)\partial_{nn}u$ |
| $u_{x_0, r}, u_r$ | $u_{x_0, r}(x) := \frac{u(x_0 + rx)}{r}$, and $u_r(x) = u_{0, r}(x)$ |
| $v_{\tau\tau}^h(x)$ | $v_{\tau\tau}^h(x) := \frac{v(x+h\tau) + v(x-h\tau) - 2v(x)}{h^2}$ |

1.3. Outline of the paper. The paper is organized as follows: In Section 2, we review some fundamental properties of minimizers of (1). Section 3 is devoted to a Harnack-type inequality (Proposition 3.4) and quadratic improvement of flatness (Proposition 3.1) for solutions of (2). In Section 4, we introduce a nonlinear dichotomy which is crucial for the proof of the main results. Section 5 contains the proof of the main theorems, namely Theorem 1.1 and Theorem 1.2. Finally, in Appendix A, we present a Harnack inequality for an auxiliary problem of the p -Laplacian type, while in Appendix B, we provide a classification of the global solutions for an auxiliary linearized operator.

2. BASIC PROPERTIES OF MINIMIZERS

In this section, we present some fundamental properties of minimizers of J_{TP} . We start with the following existence result.

Theorem 2.1 (Existence). *If the admissible set \mathcal{K} is nonempty, then there exists a minimizer u of J_{TP} over \mathcal{K} . Moreover, every minimizer satisfies*

$$\begin{cases} \Delta_p u = 0, & \text{in } \Omega_u^+ \cup \Omega_u^-, \\ \Delta_p u^\pm \geq 0, & \text{in } D, \\ \|u\|_{L^\infty(D)} \leq \|g\|_{L^\infty(D)}. \end{cases}$$

Proof. See [8, Theorem 2.1]. \square

In the following proposition, we present a non-degeneracy property for the minimizers. This property reveals the fact that each of the two phases Ω_u^+ and Ω_u^- are optimal with respect to one-sided inwards perturbations; see [9, 31].

Theorem 2.2 (Non-degeneracy). *Let $D \subset \mathbb{R}^n$ be an open set, and u be a minimizer of J_{TP} . Then, u is non-degenerate; i.e. there exists a constant $C = C(n, \lambda_\pm, p) > 0$ such that*

$$\int_{B_r(x_0)} (u^\pm)^p dx \geq Cr^p, \quad \text{for every } x_0 \in \overline{\Omega_u^\pm} \cap D \text{ and every } 0 < r < \text{dist}(x_0, \partial D).$$

Proof. See [8, Proposition 2.2]. \square

In [8], we have shown that the flat free boundary of the two-phase Bernoulli problem with the p -Laplace operator has $C^{1,\eta}$ regularity, and using this, we have established the Lipschitz regularity of the minimizers by a dichotomy argument.

Theorem 2.3 (Lipschitz regularity). *Let $u : D \rightarrow \mathbb{R}$ be a minimizer of J_{TP} in D . Then u is locally Lipschitz continuous; $u \in C_{\text{loc}}^{0,1}(D)$.*

Proof. See [8, Theorem 1.2]. \square

By Theorem 2.3 in hand, we would review the concept of blow-up, which is significant prior to studying the regularity of the free boundary. Let u be a minimizer of J_{TP} in the open set D . For every $x_0 \in F(u)$, and every $0 < r < \text{dist}(x_0, \partial D)$, we define the function

$$u_{x_0,r}(x) := \frac{u(x_0 + rx)}{r},$$

which is well-defined for $|x| < \frac{1}{r} \text{dist}(x_0, \partial D)$, and vanishes at the origin. When $x_0 = 0$, we denote $u_{0,r}$ by u_r for the sake of convenience. Note that, for every $R > 0$, and $r \ll 1$, the functions $u_{x_0,r}$ are defined on the ball B_R and are uniformly Lipschitz in B_R because of the local Lipschitz continuity of minimizer u ; Theorem 2.3. Given a sequence $r_k > 0$ such that $r_k \rightarrow 0$, we say that the sequence of functions u_{x_0,r_k} is a *blow-up sequence* of u at x_0 . Hence, there is a Lipschitz continuous function $v : \mathbb{R}^n \rightarrow \mathbb{R}$ and a subsequence of u_{x_0,r_k} converging to v uniformly on every ball $B_R \subset \mathbb{R}^n$. We say that v is a blow-up limit of u at x_0 . The blow-up limit v is a minimizer in any ball B_R (i.e. a global minimizer) and satisfies the following properties (see for instance [1, 3])

$$\begin{aligned} u_{x_0,r_k} &\rightarrow v && \text{in } C_{\text{loc}}^{0,\alpha}(\mathbb{R}^n) \text{ for every } 0 < \alpha < 1, \\ \nabla u_{x_0,r_k} &\rightarrow \nabla v && \text{a.e. in } \mathbb{R}^n, \\ \partial\Omega_{u_{x_0,r_k}}^\pm &\rightarrow \partial\Omega_v^\pm && \text{locally in the Hausdorff distance,} \\ \chi\Omega_{u_{x_0,r_k}}^\pm &\rightarrow \chi\Omega_v^\pm && \text{in } L_{\text{loc}}^1(\mathbb{R}^n). \end{aligned}$$

We remember that u is a weak solution (in the sense of distributions) to the p -Laplace equation $\Delta_p u = f$, in Ω , if $u \in W^{1,p}(\Omega)$, and for every $\varphi \in C_c^\infty(\Omega)$, there holds that

$$-\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} f \varphi \, dx.$$

Also, by viscosity solution, we will refer to the usual C -viscosity definition of solutions of elliptic PDEs, as presented in e.g. the book [13]. We emphasize on the equivalence of viscosity and weak solutions for the solutions to $\Delta_p u = f$, [22, Theorem 2.2 and Remark 2.3].

Now, we define the optimality conditions (2), in a weak (viscosity) sense on the free boundary.

Definition 2.1. We say that a function $Q : D \rightarrow \mathbb{R}$, touches a function $w : D \rightarrow \mathbb{R}$ from below (resp. from above) at a point $x_0 \in D$, if $Q(x_0) = w(x_0)$, and

$$Q(x) - w(x) \leq 0 \quad (\text{resp. } Q(x) - w(x) \geq 0),$$

for every x in a neighborhood of x_0 . We will say that Q touches w strictly from below (resp. above) if the above inequalities are strict for $x \neq x_0$.

A function Q is an (admissible) comparison function in D , if

- (a) $Q \in C^1(\overline{\{Q > 0\}} \cap D) \cap C^1(\overline{\{Q < 0\}} \cap D)$;
- (b) $Q \in C^2(\{Q > 0\} \cap D) \cap C^2(\{Q < 0\} \cap D)$;
- (c) $\partial\{Q > 0\}$ and $\partial\{Q < 0\}$ are smooth manifolds in D .

We should remark that if $\nabla Q \neq 0$ on $\partial\{Q > 0\} \cup \partial\{Q < 0\}$, then condition (c) above holds.

Definition 2.2. The optimality conditions on the free boundary $F(u)$ is as follows:

- (A) Suppose that Q is a comparison function that touches u from below at x_0 .
 - (A.1) If $x_0 \in \Gamma_{\text{OP}}^+$, then $|\nabla Q^+(x_0)| \leq \lambda_+$;
 - (A.2) if $x_0 \in \Gamma_{\text{OP}}^-$, then $Q^+ \equiv 0$ in a neighborhood of x_0 , and $|\nabla Q^-(x_0)| \geq \lambda_-$;
 - (A.3) if $x_0 \in \Gamma_{\text{TP}}$, then $|\nabla Q^-(x_0)| \geq \lambda_-$, and

$$|\nabla Q^+(x_0)|^p - |\nabla Q^-(x_0)|^p \leq \lambda_+^p - \lambda_-^p.$$

- (B) Suppose that Q is a comparison function that touches u from above at x_0 .
 - (B.1) If $x_0 \in \Gamma_{\text{OP}}^+$, then $Q^- \equiv 0$ in a neighborhood of x_0 , and $|\nabla Q^+(x_0)| \geq \lambda_+$;
 - (B.2) if $x_0 \in \Gamma_{\text{OP}}^-$, then $|\nabla Q^-(x_0)| \leq \lambda_-$;
 - (B.3) if $x_0 \in \Gamma_{\text{TP}}$, then $|\nabla Q^+(x_0)| \geq \lambda_+$, and

$$|\nabla Q^+(x_0)|^p - |\nabla Q^-(x_0)|^p \geq \lambda_+^p - \lambda_-^p.$$

Definition 2.3. If $u : D \rightarrow \mathbb{R}$ is a continuous function such that the claim (A) in Definition 2.2 holds for every comparison function Q , then we say that u satisfies (2) on the free boundary in the viscosity supersolution sense.

Similarly, we say that u satisfies (2) on the free boundary in the viscosity subsolution sense if the claim (B) holds for every comparison function Q .

Also, if both (A) and (B) hold, then we say that u satisfies (2) on the free boundary in the viscosity sense.

Remark 2.4. Notice that Definition 2.3 together with the usual definition of viscosity supersolution to $\Delta_p u = 0$ means that u is a supersolution to (2). Similarly, Definition 2.3 together with the definition of viscosity subsolution to $\Delta_p u = 0$ means that u is a subsolution to (2). Also, u is a solution to (2) if it is both a subsolution and a supersolution.

Now, we recall the following result, which states that the minimizers of J_{TP} in D are viscosity solution to (2).

Lemma 2.5. *Let u be a minimizer of J_{TP} in the open set $D \subset \mathbb{R}^n$. Then u satisfies the optimality conditions on the free boundary $F(u)$.*

Proof. See [8, Lemma 3.1]. \square

3. QUADRATIC IMPROVEMENT OF FLATNESS

In this section, we will provide a quadratic improvement of flatness. However, before we proceed, we need to introduce some definitions and notations.

First, we define a two-plane solution to the equation (2) as

$$U(x) = U_{a^+, \nu}(x) := a^+(x \cdot \nu)^+ - a^-(x \cdot \nu)^-,$$

for some $\nu \in \mathbb{S}^{n-1}$ and a^+, a^- that satisfy

$$(a^+)^p - (a^-)^p = \lambda_+^p - \lambda_-^p, \quad a^+ \geq \lambda_+, \quad a^- \geq \lambda_-.$$

We denote by $\mathcal{U}_{\lambda_+, \lambda_-}$ the set of all two-plane solutions.

Next, we introduce the following approximate quadratic polynomials

$$P_{M, \nu}(x) := x \cdot \nu + \frac{1}{2} x^T M x,$$

for some $\nu \in \mathbb{S}^{n-1}$ and a symmetric matrix $M = (M_{ij}) \in \mathbb{S}^{n \times n}$. In addition, they satisfy the conditions

$$(4) \quad M\nu = 0, \quad \text{tr}(M) = 0.$$

We also denote by $V_{a^+, a^-, M, \nu}$, or for simplicity by $V_{a^+, M, \nu}$, the two-phase quadratic polynomial of positive slopes a^+, a^-

$$(5) \quad V_{a^+, M, \nu}(x) := a^+ P_{M, \nu}^+(x) - a^- P_{M, \nu}^-(x), \quad (a^+)^p - (a^-)^p = \lambda_+^p - \lambda_-^p.$$

Notice that $V_{a^+, 0, \nu} = U_{a^+, \nu}$.

It is noteworthy that the quadratic polynomials $P_{M, \nu}^+$ are not p -harmonic when $p \neq 2$, despite the linear approximation $U_{a^+, \nu}$ satisfying (2). This poses a challenge when attempting to extend the linear improvement of flatness to the quadratic improvement of flatness.

Proposition 3.1 (Quadratic improvement of flatness). *For every $1 < p < \infty$, $\lambda_{\pm} > 0$ and $0 < L_0 \leq L_1$, there exists positive constants r_1 , ϵ_1 , and δ such that if u is a solution to (2) with the assumption $0 \in \Gamma_{\text{TP}}^{\text{int}}(u)$ and furthermore it satisfies*

$$(6) \quad |u - V_{a^+, M, \nu}| \leq \epsilon, \quad \text{in } B_1,$$

with $L_0 \leq a_+ \leq L_1$ and

$$(7) \quad \|M\| \leq \delta \epsilon^{\frac{1}{2}},$$

for some $0 < \epsilon \leq \epsilon_1$, then

$$(8) \quad |u - V_{\bar{a}^+, \bar{M}, \bar{\nu}}| \leq \epsilon r_1^{1+\alpha_1}, \quad \text{in } B_{r_1},$$

for some $\alpha_1 > 0$, with

$$|\bar{a}^+ - a^+|, |\bar{\nu} - \nu|, \|M - \bar{M}\| \leq C\epsilon.$$

Here, constants $r_1, \epsilon_1, \delta, \alpha_1$, and C depends only on n, p, λ_{\pm}, L_0 , and L_1 .

Remark 3.2. *The conclusion of Proposition 3.1 can be iterated indefinitely. Precisely, if u satisfies (8), then the rescaling*

$$u_r(x) := \frac{u(rx)}{r}, \quad r = r_1,$$

will satisfy

$$|u_r - V_{a^+, r\bar{M}, \bar{v}}| \leq \epsilon_r := \epsilon r^{\alpha_1}, \quad \text{in } B_1,$$

if we check that

$$\|r\bar{M}\| \leq \delta \epsilon_r^{\frac{1}{2}}, \quad \epsilon_r \leq \epsilon \leq \epsilon_1.$$

This is clearly possible since by choosing ϵ_1 small, we have

$$\|r\bar{M}\| \leq \|rM\| + C\epsilon r \leq r\delta\epsilon^{\frac{1}{2}} + C\epsilon r \leq \delta\epsilon_r^{\frac{1}{2}} = \delta(\epsilon r^{\alpha_1})^{\frac{1}{2}}.$$

A consequence of Proposition 3.1 is the corollary below. In fact, by invoking Proposition 3.1, similar to what we have done in [8, Section 5], when u is close enough to a two-plane solution, we can conclude that $\Gamma_{\text{TP}}^{\text{int}}(u)$ is a $C^{1,\eta}$ smooth manifold.

Corollary 3.3. *There exist universal constants $\epsilon_1, \delta > 0$ such that if (6) and (7) hold with $0 < \epsilon \leq \epsilon_1$, then $\Gamma_{\text{TP}}^{\text{int}}(u)$ has small $C^{1,\eta}$ norm in $B_{\frac{1}{2}}$. Moreover,*

$$|\nabla u^+(0)| \leq a^+ + C\epsilon,$$

and

$$\left| \frac{\nabla u^+(0)}{|\nabla u^+(0)|} - v \right| \leq C\epsilon,$$

with $C > 0$ universal.

3.1. Harnack-type inequality. We need the following Harnack-type inequality for u .

Lemma 3.4. *For every $1 < p < \infty$, $\lambda_{\pm} > 0$, and $0 < L_0 \leq L_1$, there exists positive constants $\bar{\epsilon}, \delta, c > 0$ such that the following holds. Assume that u is a solution to (2) which satisfies*

$$V(x + \sigma_1 \mathbf{e}_n) \leq u(x) \leq V(x + \sigma_2 \mathbf{e}_n), \quad \text{in } B_1,$$

where $0 \leq \epsilon = \sigma_2 - \sigma_1 \leq \bar{\epsilon}$, $V(x) := V_{a^+, M, \mathbf{e}_n}(x)$ with $a^+ \in [L_0, L_1]$ and M satisfying

$$(9) \quad \|M\| \leq \delta \epsilon^{\frac{1}{2}}, \quad 0 < \epsilon \leq \bar{\epsilon}.$$

Then, one can find new constants $\bar{\sigma}_1, \bar{\sigma}_2$ with

$$0 \leq \bar{\sigma}_2 - \bar{\sigma}_1 \leq c\epsilon$$

such that

$$V(x + \bar{\sigma}_1 \mathbf{e}_n) \leq u(x) \leq V(x + \bar{\sigma}_2 \mathbf{e}_n), \quad \text{in } \overline{B_{\frac{1}{2}}}.$$

Proof. Suppose that $\lambda_+ \geq \lambda_-$, which this is not a restrictive assumption. This assumption infer also that $a^+ \geq a^-$.

Let $\bar{x} = \frac{1}{2} \mathbf{e}_n$ and $\sigma = \frac{(\sigma_2 - \sigma_1)a^+}{2}$. We will show that

$$(10) \quad u(\bar{x}) \geq V(\bar{x} + \sigma_1 \mathbf{e}_n) + \sigma \implies u(x) \geq V(x + \bar{\sigma}_1 \mathbf{e}_n), \quad \text{in } \overline{B_{\frac{1}{2}}},$$

for $\bar{\sigma}_1 = \sigma_1 + c\epsilon$, where $0 < c < 1$ is a universal constant. Similarly,

$$u(\bar{x}) \leq V(\bar{x} + \sigma_2 \mathbf{e}_n) - \sigma \implies u(x) \leq V(x + \bar{\sigma}_2 \mathbf{e}_n), \quad \text{in } \overline{B_{\frac{1}{2}}},$$

for $\bar{\sigma}_2 = \sigma_2 - c\epsilon$.

We prove only the first implication. The second one can be obtained by the same arguments.

Step 1: We first prove that

$$(11) \quad u(x) \geq V(x + \sigma_1 \mathbf{e}_n) + c_0 \epsilon, \quad \text{for all } x \in B_{2r_0}(x_0),$$

for some universal constants c_0 , and r_0 such that $B_{4r_0}(x_0) \subset \{x_n > 0\}$. To do this, we distinguish two cases:

Case (i). Suppose $|\nabla u(\bar{x})| < \frac{1}{4}a^+$ for $\bar{x} = \frac{1}{5}\mathbf{e}_n$. Then, there exists $r_1 = r_1(n, p, L_0) > 0$ such that $|\nabla u(x)| < \frac{1}{2}a^+$ in $B_{4r_1}(\bar{x})$; note that u is universally bounded and p -harmonic in $B_{\frac{1}{10}}(\bar{x})$. Now, by invoking Lemma A.1 in $B_{4r_1}(\bar{x})$ for $h := u - V$, one can guarantee that

$$u(x) - V(x + \sigma_1 \mathbf{e}_n) \geq C^{-1}(u(\bar{x}) - V(\bar{x} + \sigma_1 \mathbf{e}_n)) - a^+ r_1, \quad \text{in } B_{r_1}(\bar{x}),$$

for an appropriate universal constant $C = C(n, p) > 0$. By the assumption (10), we obtain for all $x \in B_{r_1}(\bar{x})$

$$\begin{aligned} C^{-1}\sigma - a^+ r_1 &\leq u(x) - V(x + \sigma_1 \mathbf{e}_n) \\ &\leq u(x - 2r_1 \mathbf{e}_n) - V(x + (\sigma_1 - 2r_1) \mathbf{e}_n) \\ &\quad + (V(x + (\sigma_1 - 2r_1) \mathbf{e}_n) - V(x + \sigma_1 \mathbf{e}_n)) + 2r_1 \|\nabla u\|_{L^\infty(B_{4r_1}(\bar{x}))} \\ &\leq u(x - 2r_1 \mathbf{e}_n) - V(x + (\sigma_1 - 2r_1) \mathbf{e}_n) - 2a^+ r_1 + 2r_1 \|\nabla u\|_{L^\infty(B_{4r_1}(\bar{x}))} \\ &\leq u(x - 2r_1 \mathbf{e}_n) - V(x + (\sigma_1 - 2r_1) \mathbf{e}_n) - a^+ r_1. \end{aligned}$$

Thus, with $x^* = \bar{x} - 2r_1 \mathbf{e}_n$, we get

$$(12) \quad C^{-1}\sigma \leq u(x) - V(x + \sigma_1 \mathbf{e}_n), \quad \text{for all } x \in B_{r_1}(x^*).$$

Case (ii). Suppose $|\nabla u(\bar{x})| > \frac{1}{4}a^+$. By the interior gradient estimate, we know that $|\nabla u|$ is bounded in $B_{\frac{1}{10}}(\bar{x})$, and there exists a constant $0 < r_0 = r_0(n, p, L_0, L_1)$, with $8r_0 \leq \frac{1}{10}$ such that

$$\frac{1}{8}a^+ \leq |\nabla u(x)| \leq Ca^+, \quad \text{for all } x \in B_{8r_0}(\bar{x}),$$

for an appropriate universal constant $C = C(n, p) > 0$. Now, u will be the weak solution to the following uniformly elliptic equation

$$\sum_{i,j=1}^n \theta_{ij} \partial_{x_i x_j} u = 0, \quad \text{in } B_{4r_0}(\bar{x}),$$

with $\theta_{ij} = \delta_{ij} + (p-2)|\nabla u|^{-2} \partial_{x_i} u \partial_{x_j} u$. Also, in this way, we conclude that $h = u - V$ satisfies the following uniformly elliptic equation

$$\begin{aligned} \sum_{i,j=1}^n \theta_{ij} \partial_{x_i x_j} h &= -a^+ \sum_{i,j=1}^n \theta_{ij} M_{ij} = -a^+ \text{tr}(M) - a^+(p-2) \frac{(\nabla u)^T M \nabla u}{|\nabla u|^2} \\ &= -\frac{a^+(p-2)}{|\nabla u|^2} (\nabla u + Mx)^T M \nabla h - a^+(p-2) \frac{x^T M^3 x}{|\nabla u|^2}. \end{aligned}$$

Then, recalling (9), and applying Harnack's inequality (see e.g. [24, Chapter 9]), one can guarantee that

$$(13) \quad C^{-1}\sigma \leq u(x) - V(x + \sigma_1 \mathbf{e}_n), \quad \text{for all } x \in B_{r_0}(\bar{x}).$$

Thus, summarizing (12) and (13), we get (11) with $x_0 = x^*$, or $x_0 = \bar{x}$, and the universal constant c_0 .

Step 2: In this step, we will construct a barrier. For a given positive constant $\gamma > 0$, let us define

$$\psi(x) = c \left(r_0^{-\gamma} - |x - x_0|^{-\gamma} \right),$$

in the closure of the annulus $A := B_{\frac{3}{4}}(x_0) \setminus B_{r_0}(x_0)$. Also, the positive constant c is such that ψ satisfies the following boundary conditions:

$$\begin{cases} \psi = 1, & \text{on } \partial B_{\frac{3}{4}}(x_0), \\ \psi = 0, & \text{on } \partial B_{r_0}(x_0). \end{cases}$$

Extend ψ to be equal to zero on $B_{r_0}(x_0)$. We can choose $\gamma = \gamma(n, p)$ such that

$$(14) \quad \Delta_p(P - \epsilon\psi) \geq c(n, p)\epsilon > 0, \quad \text{in } A,$$

for sufficiently small $\epsilon > 0$, where $P(x) = \sigma_1 + x \cdot \mathbf{e}_n + \frac{1}{2}x^T Mx$. To see this, we have

$$\begin{aligned} \Delta_p(P - \epsilon\psi) &= \operatorname{div} \left(|\mathbf{e}_n + Mx - \epsilon\nabla\psi|^{p-2} (\mathbf{e}_n + Mx - \epsilon\nabla\psi) \right) \\ &= |\mathbf{e}_n + Mx - \epsilon\nabla\psi|^{p-4} \left(-\epsilon |\mathbf{e}_n + Mx - \epsilon\nabla\psi|^2 \Delta\psi \right. \\ &\quad \left. + (p-2)(\mathbf{e}_n + Mx - \epsilon\nabla\psi)^T (M - \epsilon D^2\psi) (\mathbf{e}_n + Mx - \epsilon\nabla\psi) \right) \\ &= -\epsilon (\Delta\psi + (p-2)\partial_{nn}\psi) + O(\epsilon^{\frac{3}{2}}). \end{aligned}$$

On the other hand,

$$\Delta\psi + (p-2)\partial_{nn}\psi = -\frac{c\gamma}{|x - x_0|^{\gamma+4}} \left((\gamma - n - p + 4)|x - x_0|^2 + (p-2)(\gamma+2)(x - x_0) \cdot \mathbf{e}_n \right)^2.$$

If we choose $\gamma > n + p - 4$, when $p \geq 2$, and $\gamma > \frac{n-p}{p-1}$, when $1 < p < 2$, then (14) holds for small $\epsilon > 0$.

Step 3: Now, define for $t \geq 0$

$$v_t(x) := (1 + \epsilon\ell)a^+ (P(x) + \epsilon\tilde{c}(t - \psi(x) - k))^+ - a^- (P(x) + \epsilon\tilde{c}(t - \psi(x) - k))^- ,$$

where $P(x) = \sigma_1 + x \cdot \mathbf{e}_n + \frac{1}{2}x^T Mx$, $k = \frac{1}{2}(1 - \|\psi\|_{L^\infty(B_{\frac{1}{2}}(0))})$, $\tilde{c} = \frac{c_0}{a^+}$ for c_0 in (11), and ℓ is small such that

$$\ell P(x) \leq \tilde{c} \min \left(\frac{k}{2}, cr_0^{-\gamma}(1 - 2^{-\gamma}) \right), \quad \text{in } B_{\frac{3}{4}}(x_0).$$

In this step, we show that

$$v_0(x) \leq u(x), \quad x \in \overline{B_{\frac{3}{4}}(x_0)}.$$

First, for $x \in B_{2r_0}(x_0)$ we apply (11)

$$v_0(x) \leq (1 + \epsilon\ell)a^+ P(x)^+ \leq V(x + \sigma_1 \mathbf{e}_n) + a^+ \tilde{c}\epsilon = V(x + \sigma_1 \mathbf{e}_n) + c_0\epsilon \leq u(x).$$

When $x \in B_{\frac{3}{4}}(x_0) \setminus B_{2r_0}(x_0)$, we have

$$\tilde{c}\psi(x) \geq \tilde{c}cr_0^{-\gamma}(1 - 2^{-\gamma}) \geq \ell P(x).$$

Now, assume that $P(x) - \epsilon\tilde{c}(\psi(x) + k) \geq 0$, hence

$$\begin{aligned} v_0(x) &= (1 + \epsilon\ell)a^+ (P(x) - \epsilon\tilde{c}(\psi(x) + k)) \\ &\leq V(x + \sigma_1\mathbf{e}_n) + \epsilon a^+ (\ell P(x) - (1 + \epsilon\ell)\tilde{c}\psi(x)) \\ &\leq V(x + \sigma_1\mathbf{e}_n) \leq u(x). \end{aligned}$$

Also, if $P(x) - \epsilon\tilde{c}(\psi(x) + k) \leq 0$, the result is obvious when $0 \leq P(x)$; because $v_0(x) \leq 0 \leq a^+P(x) = V(x + \sigma_1\mathbf{e}_n) \leq u(x)$. If $P(x) \leq 0$,

$$v_0(x) = a_-(P(x) - \epsilon\tilde{c}(\psi(x) + k)) \leq a_-P(x) = V(x + \sigma_1\mathbf{e}_n) \leq u(x).$$

Step 4: Let \bar{t} be the largest $t \geq 0$ such that

$$v_t(x) \leq u(x), \quad \text{in } \overline{B_{\frac{3}{4}}(x_0)}.$$

We want to show that $\bar{t} \geq 1$. Suppose this is true, then we will have

$$\begin{aligned} u(x) &\geq v_1(x) = (1 + \epsilon\ell)a^+ (P(x) + \epsilon\tilde{c}(1 - \psi(x) - k))^+ - a^- (P(x) + \epsilon\tilde{c}(1 - \psi(x) - k))^- \\ &\geq V(x) + c\epsilon, \quad \text{in } B_{\frac{1}{2}}(0) \Subset B_{\frac{3}{4}}(x_0), \end{aligned}$$

where c is a universal constant, and therefore we obtain (10). Note that in the last inequality, we have used that $\|\psi\|_{L^\infty(B_{\frac{1}{2}}(0))} = 1 - 2k$, and $a^+ \geq a^-$.

Suppose $\bar{t} < 1$. Then, at some $\tilde{x} \in \overline{B_{\frac{3}{4}}(x_0)}$, we have

$$(15) \quad v_{\bar{t}}(\tilde{x}) = u(\tilde{x}).$$

We show that such touching point can only occur on $\overline{B_{r_0}(x_0)}$. Indeed, since $\psi = 1$ on $\partial B_{\frac{3}{4}}(x_0)$, from the definition of v_t , we get that for $t < 1$,

$$v_t(x) = (1 + \epsilon\ell)a^+ (P(x) + \epsilon\tilde{c}(t - 1 - k))^+ - a^- (P(x) + \epsilon\tilde{c}(t - 1 - k))^- < V(x).$$

Also, \tilde{x} cannot belong to the annulus A . Indeed, in $\{x \in A : u(x) \neq 0\}$, we have $\Delta_p u(x) = 0$ and by the definition of viscosity solution for p -harmonic function, it follows that $\Delta_p(v_{\bar{t}})(\tilde{x}) \leq 0$ at touching point. This contradicts (14) for ϵ small enough.

Finally, if $\tilde{x} \in \{u = 0\}$, we will check the free boundary condition at \tilde{x} .

$$\begin{aligned} |\nabla v_{\bar{t}}^+(\tilde{x})|^p - |\nabla v_{\bar{t}}^-(\tilde{x})|^p &= (a_+^p(1 + \epsilon\ell)^p - a_-^p) |\mathbf{e}_n + M\tilde{x} - \epsilon\tilde{c}\nabla\psi(\tilde{x})|^p \\ &= (\lambda_+^p - \lambda_-^p + pa_+^p\epsilon\ell + O(\epsilon^2)) \left(1 + \frac{p}{2}|M\tilde{x}|^2 - p\epsilon\tilde{c}\partial_n\psi(\tilde{x}) + O(\epsilon^{\frac{3}{2}})\right). \end{aligned}$$

One can verify that

$$\partial_n\psi(\tilde{x}) = c\gamma|\tilde{x} - x_0|^{-\gamma-2}(\tilde{x} - x_0) \cdot \mathbf{e}_n < -C < 0,$$

for a universal constant C ; note that $\tilde{x}_n = O(\epsilon^{\frac{1}{2}})$. Taking into account $\|M\| \leq \delta\epsilon^{\frac{1}{2}}$ and $\lambda_+ \geq \lambda_-$, we get

$$|\nabla v_{\bar{t}}^+(\tilde{x})|^p - |\nabla v_{\bar{t}}^-(\tilde{x})|^p > \lambda_+^p - \lambda_-^p,$$

as long as δ, ϵ are small enough.

Therefore, $\tilde{x} \in \overline{B_{r_0}(x_0)}$, and for ϵ sufficiently small

$$\begin{aligned} v_{\tilde{t}}(\tilde{x}) &= (1 + \epsilon\ell)a^+ \left(P(\tilde{x}) + \epsilon\tilde{c}(\tilde{t} - k) \right) \\ &< V(\tilde{x}) + \epsilon a^+ (\ell P(\tilde{x}) + \tilde{c}(1 + \epsilon\ell)(1 - k)) \\ &\leq V(\tilde{x}) + \epsilon c_0 \left(\frac{k}{2} + (1 + \epsilon\ell)(1 - k) \right) \\ &< V(\tilde{x}) + \epsilon c_0 \leq u(\tilde{x}), \end{aligned}$$

contradicting (15); where in the last inequality, we have used (11). \square

3.2. Constructing comparison functions. In this subsection, our goal is to construct suitable strict subsolutions to the two-phase problem in B_1 that are ϵ -perturbations of V . Similarly, strict supersolutions can be also constructed.

Given $\epsilon, \delta > 0$, assume that

$$v = \mathbf{e}_n, \quad \|M\| \leq \delta\epsilon^{\frac{1}{2}},$$

and due to the condition (4) on v and M , we have:

$$M\mathbf{e}_n = 0, \quad \text{tr}(M) = 0.$$

In the sequel, we write $x = (x', x_n) \in \mathbb{R}^n$, where $x' = (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}$.

Let $s, t, A \in \mathbb{R}$, $\xi' \in \mathbb{R}^{n-1}$ and $N = (N_{ij})$ be an $n \times n$ symmetric matrix satisfying

$$(16) \quad \text{tr}(N) + (p-2)N_{nn} > 0, \quad \text{and} \quad N\mathbf{e}_n = N_{nn}\mathbf{e}_n.$$

Given $\epsilon > 0$, set

$$\bar{a}^+ := a^+(1 + \epsilon s), \quad \bar{a}^- := a^-(1 + \epsilon t), \quad \bar{M} := M + 2\epsilon N,$$

and

$$Q(x) := x \cdot \mathbf{e}_n + \frac{1}{2} x'^T \bar{M} x + A\epsilon + \epsilon \xi' \cdot x',$$

or

$$Q(x) = P_{\bar{M}, \mathbf{e}_n + \epsilon(\xi', 0)}(x) + A\epsilon,$$

and define

$$(17) \quad v(x) := \bar{a}^+ Q^+ - \bar{a}^- Q^-.$$

Lemma 3.5. *Suppose that the symmetric matrix N satisfies (16). Then, if s, t satisfy*

$$(18) \quad (a^+)^p s - (a^-)^p t > \delta^2 \left((a^+)^p + (a^-)^p \right),$$

then v defined in (17) is a strict subsolution to (2) in B_1 for all sufficiently small ϵ .

Proof. In $\Omega_v^+ \cap B_1$, we have

$$\begin{aligned}
\Delta_p Q^+ &= \operatorname{div} \left(|\nabla(Q^+)|^{p-2} \nabla Q^+ \right) \\
&= |\nabla Q^+|^{p-4} \left(|\nabla Q^+|^2 \Delta Q^+ + (p-2)(\nabla Q^+)^T D^2 Q^+ \nabla Q^+ \right) \\
&= |\nabla Q^+|^{p-4} \left(|\nabla Q^+|^2 \operatorname{tr}(M + 2\epsilon N) + (p-2)(\nabla Q^+)^T (M + 2\epsilon N) (\nabla Q^+) \right) \\
&= |\nabla Q^+|^{p-4} \left[(p-2)x^T M^3 x + 2\epsilon(|Mx + \mathbf{e}_n|^2 \operatorname{tr}(N) + (p-2)(Mx + \mathbf{e}_n)^T N(Mx + \mathbf{e}_n) \right. \\
&\quad \left. + (p-2)(2Nx + \xi')^T M^2 x) + O(\epsilon^2) \right] \\
&\geq |\nabla Q^+|^{p-4} \left[-(p-2)\delta^3 \epsilon^{\frac{3}{2}} + 2\epsilon(\operatorname{tr}(N) + (p-2)N_{nn} - 2\delta\epsilon^{\frac{1}{2}}\|N\|) + O(\epsilon^2) \right] \\
&= 2\epsilon |\nabla Q^+|^{p-4} \left(\operatorname{tr}(N) + (p-2)N_{nn} + O(\epsilon^{\frac{1}{2}}) \right),
\end{aligned}$$

where the implicit constant in $O(\epsilon^{\frac{1}{2}})$ can be controlled by δ, N, p, n . Hence, for sufficiently small ϵ , we have $\Delta_p(\bar{a}^+ Q^+) > 0$, in $\Omega_v^+ \cap B_1$. Similar computations show that $\Delta_p(-\bar{a}^- Q^-) > 0$, in $\Omega_v^- \cap B_1$. Thus v is a strict subsolution to (2) in B_1 .

For the free boundary conditions, we need to show that

$$|\nabla v^+|^p - |\nabla v^-|^p > \lambda_+^p - \lambda_-^p.$$

Let us for instance compute

$$\begin{aligned}
|\nabla v^+|^p &= (\bar{a}^+)^p \left(1 + |\bar{M}x|^2 + \epsilon^2 |\xi'|^2 + 4\epsilon(Nx) \cdot \mathbf{e}_n + 2\epsilon(\bar{M}x) \cdot \xi' \right)^{\frac{p}{2}} \\
&= (a^+)^p \left(1 + p\epsilon \left(s + \frac{1}{2}|Mx|^2 + 2(Nx) \cdot \mathbf{e}_n \right) + O(\epsilon^{\frac{3}{2}}) \right).
\end{aligned}$$

On $\{v = 0\}$, we have $x_n = O(\epsilon^{\frac{1}{2}})$, and by (16) we get $(Nx) \cdot \mathbf{e}_n = O(\epsilon^{\frac{1}{2}})$. This along with the assumption $\|M\| \leq \delta\epsilon^{\frac{1}{2}}$, gives that

$$|\nabla v^+|^p \geq (a^+)^p \left(1 + p\epsilon(s - \delta^2) \right),$$

as long as ϵ is small enough; depending on s, t, N, ξ' , and δ . Similarly,

$$|\nabla v^-|^p \leq (a^-)^p \left(1 + p\epsilon(t + \delta^2) \right).$$

Thus, according to (18) and the condition $(a^+)^p - (a^-)^p = \lambda_+^p - \lambda_-^p$, the boundary condition holds. \square

3.3. Proof of Proposition 3.1. Let $V_{a^+, M, v}$ be the quadratic polynomial defined in (5). Given a continuous function v in B_1 , we denote its ϵ -linearization around the function $V_{a^+, M, v}$, as

$$(19) \quad \tilde{v}_{a^+, M, v, \epsilon}(x) := \begin{cases} \frac{v(x) - a^+ P_{M, v}(x)}{a^+ \epsilon}, & x \in \overline{\Omega_v^+ \cap B_1}, \\ \frac{v(x) - a^- P_{M, v}(x)}{a^- \epsilon}, & x \in \overline{\Omega_v^- \cap B_1}. \end{cases}$$

In what follows, we will typically drop the indices from V, P, \tilde{v} whenever there is no possibility of confusion.

Proof of Proposition 3.1. We divide the proof into three steps.

Step 1. Compactness. Fix r_1 universal to be made precise later in step 3. Assume for a contradiction that we can find sequences $\epsilon_k \rightarrow 0$, and $\delta_k \rightarrow 0$, and also a sequence u_k of solutions to (2) in $D = B_1$ such that

$$(20) \quad |u_k - V_{a_k^+, M_k, \mathbf{e}_n}| \leq \epsilon_k, \quad \text{in } B_1, \quad \text{and} \quad 0 \in \Gamma_{\text{TP}}^{\text{int}}(u_k),$$

with

$$\|M_k\| \leq \delta_k \epsilon_k^{\frac{1}{2}}, \quad M_k \mathbf{e}_n = 0, \quad (a_k^+)^p - (a_k^-)^p = \lambda_+^p - \lambda_-^p, \quad a_k^+ \in [L_0, L_1],$$

but u_k does not satisfy the conclusion (8).

Let $\tilde{u}_k := \tilde{u}_{a_k^+, M_k, \mathbf{e}_n, \epsilon_k}$ be defined as in (19). Then, (20) gives us

$$|\tilde{u}_k| \leq C, \quad \text{for } x \in B_1.$$

From Lemma 3.4, we can easily obtain that the oscillation of \tilde{u} decreases by a factor $1-c$, as we restrict from B_1 to $B_{\frac{1}{2}}$ (see e.g. [31, proof of Lemma 7.13]). This result can be iterated m -times provided that $\epsilon_k(2(1-c))^m \leq \bar{\epsilon}$. By the Arzelá-Ascoli theorem it follows that, as $\epsilon_k \rightarrow 0$, \tilde{u}_k (up to a subsequence) converges uniformly in $B_{\frac{1}{2}}$ to a Hölder continuous function u^* . Also, up to a subsequence,

$$a_k^+ \rightarrow \bar{a}^+ \in [L_0, L_1],$$

and hence

$$a_k^- \rightarrow \bar{a}^- = \left((\bar{a}^+)^p + \lambda_-^p - \lambda_+^p \right)^{\frac{1}{p}}.$$

Notice that $0 \in \Gamma_{\text{TP}}^{\text{int}}(u_k)$ implies that $\tilde{u}_k(0) = 0$; hence $u^*(0) = 0$.

Step 2. Limiting solution. We now show that u^* solves

$$(21) \quad \begin{cases} \mathcal{L}_p(u) = 0, & \text{in } B_{\frac{1}{2}} \cap \{x_n \neq 0\}, \\ a \partial_n^+ u - b \partial_n^- u = 0, & \text{on } B_{\frac{1}{2}} \cap \{x_n = 0\}, \end{cases}$$

with $a = (\bar{a}^+)^p$ and $b = (\bar{a}^-)^p$, where

$$(22) \quad \mathcal{L}_p(u) := \Delta u + (p-2)\partial_{nn}u.$$

First, we prove that $\mathcal{L}_p(u^*) = 0$. Let $P(x)$ be a quadratic polynomial touching u^* at $\bar{x} \in B_{\frac{1}{2}} \cap \{x_n > 0\}$ strictly from below. We need to show that at this point

$$\mathcal{L}_p(P) = \Delta P + (p-2)\partial_{nn}P \leq 0.$$

Since $\tilde{u}_k \rightarrow u^*$, there exist points $x_k \in \Omega_{u_k}^+ \cap B_{\frac{1}{2}}$, $x_k \rightarrow \bar{x}$, and constants $c_k \rightarrow 0$ such that

$$(23) \quad \tilde{u}_k(x_k) = P(x_k) + c_k,$$

and

$$(24) \quad \tilde{u}_k \geq P + c_k, \quad \text{in a neighborhood of } x_k.$$

From the definition of \tilde{u}_k , (23) and (24) read

$$u_k(x_k) = Q_k(x_k),$$

and

$$u_k(x) \geq Q_k(x), \quad \text{in a neighborhood of } x_k,$$

respectively, where

$$Q_k(x) = \epsilon_k a_k^+ (P(x) + c_k) + a_k^+ P_{M_k, \mathbf{e}_n}(x).$$

Note that

$$\nabla Q_k(x) = \epsilon_k a_k^+ \nabla P(x) + a_k^+ (\mathbf{e}_n + M_k x),$$

thus by recalling $\|M_k\| = O(\epsilon_k^{\frac{1}{2}})$,

$$\nabla Q_k(x_k) \neq 0, \quad \text{for } k \text{ large.}$$

Since u_k is p -harmonic, and Q_k touches u_k from below at x_k , and $\nabla Q_k(x_k) \neq 0$, by the equivalence of weak and viscosity solutions of p -harmonic functions, we get

$$\begin{aligned} 0 &\geq \Delta_p Q_k(x_k) \\ &= \operatorname{div} \left(|\nabla Q_k(x_k)|^{p-2} \nabla Q_k(x_k) \right) \\ &= |\nabla Q_k(x_k)|^{p-2} \left(\Delta Q_k(x_k) + (p-2) \left(\frac{\nabla Q_k(x_k)}{|\nabla Q_k(x_k)|} \right)^T D^2 Q_k(x_k) \left(\frac{\nabla Q_k(x_k)}{|\nabla Q_k(x_k)|} \right) \right) \\ &= |\nabla Q_k(x_k)|^{p-2} \left(\epsilon_k a_k^+ \Delta P(x_k) + a_k^+ \operatorname{tr}(M_k) \right. \\ &\quad \left. + \frac{p-2}{|\nabla Q_k(x_k)|^2} (\nabla Q_k(x_k))^T (\epsilon_k a_k^+ D^2 P(x_k) + a_k^+ M_k) \nabla Q_k(x_k) \right) \\ &= a_k^+ |\nabla Q_k(x_k)|^{p-2} \left(\epsilon_k \Delta P(x_k) + \frac{p-2}{|\nabla Q_k(x_k)|^2} (\nabla Q_k(x_k))^T (\epsilon_k D^2 P(x_k) + M_k) \nabla Q_k(x_k) \right) \\ &= \epsilon_k a_k^+ |\nabla Q_k(x_k)|^{p-2} \left(\Delta P(x_k) + \frac{p-2}{|\nabla Q_k(x_k)|^2} (\nabla Q_k(x_k))^T D^2 P(x_k) \nabla Q_k(x_k) + O(\epsilon_k^{\frac{1}{2}}) \right). \end{aligned}$$

Now, dividing both sides by ϵ_k , and passing to the limit $k \rightarrow \infty$, and recalling that

$$\nabla Q_k(x_k) \rightarrow \bar{a}^+ \mathbf{e}_n,$$

we conclude that

$$\Delta P(\bar{x}) + (p-2) \partial_{nn} P(\bar{x}) \leq 0.$$

Touching from above and reaching the opposite inequality is similar. Also, the same reasoning can be applied to prove $\mathcal{L}_p(u^*) = 0$ in $B_{\frac{1}{2}} \cap \{x_n < 0\}$.

Now, we verify the boundary condition in the viscosity sense. Let

$$w(x) := A + s x_n^+ - t x_n^- + x^T N x + \xi' \cdot x',$$

with $A \in \mathbb{R}$ and N a matrix such that $\mathcal{L}_p(w) > 0$, and

$$as - bt > 0.$$

We can always assume, via a change of coordinate in \mathbb{R}^{n-1} , that N satisfies (16). Assume also that w touches u^* strictly from below at a point $x_0 = (x_0', 0) \in B_{\frac{1}{2}}$. Set

$$\bar{a}_k^+ = a_k^+ (1 + \epsilon_k s), \quad \bar{a}_k^- = a_k^- (1 + \epsilon_k t),$$

$$\bar{M}_k = M_k + 2\epsilon_k N, \quad Q_k := P_{\bar{M}_k, \mathbf{e}_n} + \epsilon_k \xi' \cdot x' + A\epsilon_k,$$

and

$$v_k := \bar{a}_k^+ Q_k^+ - \bar{a}_k^- Q_k^-.$$

Let \tilde{v}_k be the ϵ -linearization of v_k around the function $V_{a^+, M_k, \mathbf{e}_n}$; then, it converges uniformly to w on $B_{\frac{1}{2}}$. Since \tilde{u}_k converges uniformly to u^* , and w touches u^* strictly from below at x_0 , we conclude that for a sequence of constants $c_k \rightarrow 0$, and points $x_k \rightarrow x_0$, the function

$$w_k(x) := v_k(x + \epsilon_k c_k \mathbf{e}_n),$$

touches u_k from below at x_k . Thus, Lemma 3.5 gives that w_k is a strict subsolution to our free boundary problem provided that we first choose δ_k small enough, so as to guarantee that (18) holds. We reach a contradiction as we let $k \rightarrow \infty$; thus u^* is a solution to the linearized problem (21).

Step 3. Contradiction. Since \tilde{u}_k converges uniformly to u^* , and u^* enjoys the quadratic estimate of [19, Theorem 3.2] or [8, Lemma 4.12], we have

$$(25) \quad |\tilde{u}_k - (x' \cdot v' + \tilde{s}x_n^+ - \tilde{t}x_n^-)| \leq Cr^2, \quad \text{for } x \in B_r,$$

with

$$a\tilde{s} - b\tilde{t} = 0, \quad |v'| \leq C.$$

Define

$$\bar{a}_k^+ = a_k^+ (1 + \epsilon_k \tilde{s}), \quad \bar{a}_k^- = \left((\bar{a}_k^+)^p + \lambda_-^p - \lambda_+^p \right)^{\frac{1}{p}} = a_k^- (1 + \epsilon_k \tilde{t}) + o(\epsilon_k),$$

and

$$v_k = \frac{\mathbf{e}_n + \epsilon_k(v', 0)}{\sqrt{1 + \epsilon_k^2 |v'|^2}} = \mathbf{e}_n + \epsilon_k(v', 0) + \epsilon_k^2 \tau, \quad |\tau| \leq C.$$

We claim that we can decompose

$$(26) \quad M_k = \bar{M}_k + D_k,$$

as

$$\bar{M}_k v_k = 0, \quad \text{tr}(\bar{M}_k) = 0, \quad \|D_k\| \leq C \|M_k\| |v_k - \mathbf{e}_n|;$$

hence $\|D_k\| = O(\epsilon_k^{\frac{3}{2}})$.

To prove the claim, we first decompose M_k as

$$M_k = N_k + L_k,$$

such that $N_k v_k = 0$, and

$$\|L_k\| \leq C \|M_k\| |v_k - \mathbf{e}_n|.$$

Since $\text{tr}(M_k) = 0$, we obtain that

$$\text{tr}(N_k) = O(\|L_k\|).$$

Then, we can decompose N_k further (here v_k^\perp is a unit vector perpendicular to v_k):

$$N_k = \bar{M}_k + z_k (v_k^\perp \otimes v_k^\perp), \quad z_k = \text{tr}(N_k), \quad \bar{M}_k v_k = 0,$$

so that $\text{tr}(\bar{M}_k) = 0$, and the claim (26) is proved.

Let us now show that for $r = r_1$ universal (say, $\alpha_1 = \frac{1}{2}$)

$$|u_k - V_{\bar{a}_k^+, \bar{M}_k, v_k}| \leq \epsilon_k r^{1+\alpha_1},$$

which contradicts the fact that u_k does not satisfy (8). Using (25) and the definition of \tilde{u}_k , we get that in $\Omega_{u_k}^+ \cap B_r$ (we can argue similarly in the negative part)

$$|u_k - a_k^+ P_{M, \mathbf{e}_n} - \epsilon_k a_k^+ (x' \cdot v' + \tilde{s}x_n^+ - \tilde{t}x_n^-)| \leq 2\epsilon_k Cr^2.$$

Since in this set $x_n \geq -3\epsilon_k^{\frac{1}{2}}$, we conclude that for all large k

$$|u_k - \bar{a}_k^+ P_{M, \mathbf{e}_n} - \epsilon_k a_k^+ (x' \cdot v' + \tilde{s}x_n)| \leq 3\epsilon_k Cr^2,$$

which gives

$$\left| u_k - \bar{a}_k^+ \left(x \cdot v_k + \frac{1}{2} x^T M x \right) \right| \leq 4\epsilon_k Cr^2.$$

Finally, from (26) we infer that

$$|u_k - V_{\bar{a}_k, \bar{M}_k, \bar{v}_k}| \leq 5C\epsilon_k r^2,$$

from which the desired bound follows for r_1 small enough. \square

4. NONLINEAR DICHOTOMY

In this section, we prove the following proposition which provides a dichotomy result for the solutions of (2). This result is crucial for the proof of Theorem 1.1.

Proposition 4.1 (Nonlinear dichotomy). *Let u be a solution to (2) with $|\nabla u| \leq 1$ and assume that $0 \in \Gamma_{\text{TP}}^{\text{int}}(u)$. Then, there exist small universal positive constants $\epsilon_0, \delta_0, r_0, c_0, \alpha_0$ such that if*

$$|u(x) - P_{M, \bar{v}}(x)| \leq \epsilon, \quad \text{for } x \in B_1,$$

with $\epsilon \leq \epsilon_0$, and

$$\|M\| \leq \delta_0 \epsilon^{\frac{1}{2}},$$

then one of the following two alternatives holds:

- (i) $|\nabla u^+(0)| \leq 1 - c_0 \epsilon$, or
- (ii) $|u - P_{\bar{M}, \bar{v}}| \leq \epsilon r_0^{2+\alpha_0}$ in B_{r_0} for some \bar{M}, \bar{v} with $\|M - \bar{M}\| \leq C\epsilon$ and C universal.

Remark 4.2. *If u satisfies alternative (ii) of Proposition 4.1, then we can rescale and iterate one more time. Indeed, the rescaling*

$$u_r(x) := \frac{u(rx)}{r}, \quad r = r_0,$$

will satisfy

$$|u - P_{r\bar{M}, \bar{v}}| \leq \epsilon_r := \epsilon r_0^{1+\alpha_0}, \quad \text{in } B_1,$$

with

$$\|r\bar{M}\| \leq \delta_0 \epsilon_r^{\frac{1}{2}}, \quad \epsilon_r \leq \epsilon \leq \epsilon_0,$$

provided that ϵ_0 is chosen small enough; see Remark 3.2.

The proof of Nonlinear dichotomy, i.e. Proposition 4.1, is very similar to that of Proposition 3.1. However, before that, we need the following Linear dichotomy which is similar to the nonlinear one.

Proposition 4.3 (Linear dichotomy). *Let v solves*

$$(27) \quad \begin{cases} \mathcal{L}_p(v) = 0, & \text{in } B_1 \cap \{x_n \neq 0\}, \\ \partial_n^+ v - b\partial_n^- v = 0, & \text{on } B_1 \cap \{x_n = 0\}, \end{cases}$$

with $0 < b_0 \leq b \leq b_1$ and $\|v\|_{L^\infty} \leq 1$. Moreover, suppose that

$$\partial_n v \leq 0.$$

Then, there exist universal constants $c_0, r_0 > 0$ such that either

- (i) $\partial_n^+ v(0) \leq -c_0$, or
- (ii) $|v(x) - Q(x')| \leq \frac{1}{4}r_0^2$ in B_{r_0} ,

where $Q(x')$ is a quadratic polynomial in the x' -direction with $\|Q\| \leq C$ universal and

$$\mathcal{L}_p(Q) = \Delta Q = 0.$$

Proof. Let r_0 be given, to be specified later. Assume by contradiction that we can find a sequence $c_k \rightarrow 0$, and the sequences of constants $b_0 \leq b_k \leq b_1$, and bounded monotone solutions v_k to (27) that satisfy

$$\partial_n^+ v_k(0) > -c_k,$$

and for which condition (ii) does not hold. Then, up to extracting a sequence, v_k converges uniformly on compacts and in the $C^{1,\alpha}$ norm from either side of $\{x_n \neq 0\}$ to a solution \bar{v} for a limiting problem

$$(28) \quad \begin{cases} \mathcal{L}_p(\bar{v}) = 0, & \text{in } B_{\frac{3}{4}} \setminus \{x_n \neq 0\}, \\ \partial_n^+ \bar{v} - b \partial_n^- \bar{v} = 0, & \text{on } B_{\frac{3}{4}} \cap \{x_n = 0\}, \end{cases}$$

with $\|\bar{v}\|_{L^\infty} \leq 1$, and

$$(29) \quad \partial_n \bar{v} \leq 0, \quad \partial_n^+ \bar{v}(0) = 0.$$

After subtracting a linear function in the x' -variable, we can assume that (in view of the free boundary condition $\partial_n^- \bar{v}(0) = 0$)

$$(30) \quad \bar{v}(0) = 0, \quad \nabla \bar{v}(0) = 0.$$

Step 1. We prove that

$$D_{x'}^2 \bar{v} \leq C I_{x'}, \quad \text{in } B_{\frac{1}{4}},$$

or, in other words that \bar{v} is uniformly semiconcave in the x' -variable.

For this, it suffices to show that

$$(31) \quad \bar{v}_{\tau\tau}^h(x) := \frac{\bar{v}(x+h\tau) + \bar{v}(x-h\tau) - 2\bar{v}(x)}{h^2} \leq C, \quad \text{in } B_{\frac{1}{4}},$$

independently of h , where τ is a unit direction with $\tau \perp \mathbf{e}_n$, and $h > 0$. We claim that

$$(32) \quad \bar{v}_{\tau\tau}^h(x) \leq \frac{C}{x_n^2}, \quad \text{in } B_{\frac{1}{2}}.$$

This estimate and the fact that $\bar{v}_{\tau\tau}^h$ is a solution for the transmission problem gives (31); in fact it is enough to use a suitable version of L^p -Harnack inequality for small $p > 0$ (see e.g. [13, Theorem 4.8 (2)]). To prove (32), notice that by the $C^{2,\alpha}$ interior estimates in $B_{\frac{|x|}{2}}(x)$, we have

$$|D^2 \bar{v}(x)| \leq \frac{C}{x_n^2} \|\bar{v}\|_{L^\infty} \leq \frac{C}{x_n^2}, \quad \text{in } B_{\frac{3}{8}} \setminus \{x_n = 0\}.$$

Since we can write

$$\bar{v}_{\tau\tau}^h(x_0) = \int_{-1}^1 \bar{v}_{\tau\tau}(x_0 + th\tau)(1 - |t|) dt,$$

(32) follows.

Step 2. In this step, we wish to show that

$$(33) \quad \|\bar{v}\|_{L^\infty(B_r)} \leq Cr^2, \quad r \leq \frac{1}{4},$$

with C universal.

Let us consider

$$\tilde{v}(x) = \frac{\bar{v}(rx)}{r^2}, \quad x \in B_1.$$

Then, \tilde{v} satisfies (28) and (29). From (30), we also have

$$\tilde{v}(0) = 0, \quad \nabla \tilde{v}(0) = 0.$$

Then, the conclusion of Step 1 implies that

$$(34) \quad \tilde{v}(x', 0) \leq C|x'|^2, \quad \text{in } B_1 \cap \{x_n = 0\}.$$

Hence, by the monotonicity of \tilde{v} ,

$$(35) \quad \tilde{v} \leq C, \quad \text{in } B_1 \cap \{x_n > 0\}.$$

Next, we claim that

$$(36) \quad \tilde{v} \geq -MC, \quad \text{in } B_{\frac{1}{3}}\left(\frac{\mathbf{e}_n}{2}\right),$$

for some large constant M to be specified later. Suppose by contradiction that this does not hold. Then, for any arbitrary point on $x_0 \in \{x_n = d\} \cap \{|x'| = \frac{1}{3}\}$ for small d , choose a sequence of points x_0, \dots, x_k with $x_k \in B_{\frac{1}{3}}\left(\frac{\mathbf{e}_n}{2}\right)$ and

$$|x_{i+1} - x_i| \leq d2^{i-1}, \quad i = 0, \dots, k-1.$$

Note that k , i.e. the number of points, depends only on d and is less than $2|\log d|$. Then, by virtue of Harnack inequality for $C - \tilde{v} \geq 0$ in $B_{d2^i}(x_i)$, there is a constant $c_0 < 1$ such that

$$C - \tilde{v}(x_i) \geq c_0(C - \tilde{v}(x_{i+1})).$$

This implies that

$$C - \tilde{v} \geq (1 + M)Cd^K, \quad \text{on } x_n = d, \quad |x'| \leq \frac{1}{3},$$

with K universal. In particular, if

$$d^K \geq \frac{2}{1 + M},$$

then

$$(37) \quad \tilde{v} \leq -C, \quad \text{on } x_n = d, \quad |x'| \leq \frac{1}{3}.$$

We now compare \tilde{v} with the explicit barrier

$$\phi(x) := 10C|x'|^2 - Ax_n^2 - x_n,$$

where $A = A(C, p, n)$ is chosen so that $\mathcal{L}_p(\phi) \leq 0$. Define

$$R := \left\{0 < x_n < d, \quad |x'| < \frac{1}{3}\right\}.$$

We show that for d small enough and hence M large enough, we have

$$(38) \quad \tilde{v} \leq \phi, \quad \text{on } \partial R.$$

Thus, we conclude that the inequality holds in R and we reach a contradiction because

$$0 = \partial_n \tilde{v}(0) \leq \partial_n \phi(0) = -1.$$

Now, we check (38). On $\{x_n = d\}$, this follows from (37) if d is chosen small depending on C and A . Similarly, on $\{x_n = 0\}$, the desired bound follows immediately from (34). Finally, in the set

$$\left\{0 < x_n < d, \quad |x'| = \frac{1}{3}\right\},$$

we use (35) and again we obtain (38) for d sufficiently small. In conclusion, the claim (36) holds.

Since \tilde{v} is decreasing in the \mathbf{e}_n -direction, we obtain

$$\tilde{v} \geq -C, \quad \text{in } B_{\frac{1}{3}},$$

for some C . Now, recalling that $\tilde{v}(0) = 0$, and applying Harnack inequality for $\tilde{v} + C$, we get

$$|\tilde{v}| \leq C, \quad \text{in } B_{\frac{1}{4}},$$

which after rescaling gives the desired claim (33).

Step 3. We prove that if \bar{v} solves (28) and satisfies (29), (30), and (33), then there exists a universal r_0 such that alternative (ii) holds for \bar{v} with right-hand side $\frac{1}{8}r_0^2$ instead of $\frac{1}{4}r_0^2$.

This claim follows by compactness from the classification of the global solution of \mathcal{L}_p , i.e. Lemma B.1.

Assume by contradiction that there exists a sequence of $\delta_k \rightarrow 0$, and solutions \bar{v}_k to a sequence of problems (28)_k (satisfying the same properties as \bar{v}) for which the alternative (ii) fails in the ball of radius δ_k . Denote the quadratic rescalings by

$$w_k(x) = \frac{\bar{v}_k(\delta_k x)}{\delta_k^2}.$$

Then, up to extracting a subsequence, w_k converges uniformly on compacts to a global solution U to the limiting transmission problem

$$\begin{cases} \mathcal{L}_p(U) = 0, & \text{in } \mathbb{R}^n \cap \{x_n \neq 0\}, \\ \partial_n^+ U = g \partial_n^- U, & \text{on } \{x_n = 0\}, \end{cases}$$

with $0 < b_0 \leq g \leq b_1$. The convergence is in the $C^{1,\alpha}$ norm from either side of $\{x_n \neq 0\}$, up to $\{x_n = 0\}$, and in the $C^{2,\alpha}$ norm in the interior. Clearly the global solution U also satisfies (29), (30), and (33).

Then according to Lemma B.1, we conclude that $U = Q(x')$ with $Q(x')$ being a pure quadratic polynomial in the x' -direction and with $\mathcal{L}_p(Q) = \Delta Q = 0$. Therefore, for k large, \bar{v}_k satisfies the alternative (ii) in B_{δ_k} . The contradiction establishes Step 3.

By Step 3, \bar{v} satisfies alternative (ii) with right-hand side $\frac{1}{8}r_0^2$ for some quadratic polynomial $\bar{Q}(x')$ with $\mathcal{L}_p(\bar{Q}) = \Delta \bar{Q} = 0$. As above, this means that the v_k 's satisfy the alternative (ii) for all k large, which is a contradiction. \square

Proof of Proposition 4.1. We divide the proof into two steps.

Step 1. Compactness and the limiting solution. Fix r_0 universal to be made precise later. Toward a contradiction, assume that there is sequences $\epsilon_k \rightarrow 0$, $\delta_k \rightarrow 0$ and a sequence u_k of solutions to (2) in $D = B_1$ such that

$$|u_k - P_{M_k, \mathbf{e}_n}| \leq \epsilon_k, \quad \text{in } B_1, \quad \text{and} \quad 0 \in \Gamma_{\text{TP}}^{\text{int}}(u_k),$$

with

$$\|M_k\| \leq \delta_k \epsilon_k^{\frac{1}{2}}, \quad M_k \mathbf{e}_n = 0, \quad \text{tr}(M_k) = 0,$$

but u_k does not satisfy either alternative (i) (for some small constant c'_0 to be specified later) or alternative (ii).

Let $\tilde{u}_k := \tilde{u}_{1, M_k, \mathbf{e}_n, \epsilon_k}$ be defined as in (19). Using the Step 1 and 2 of the proof of Proposition 3.1, we know that as $\epsilon_k \rightarrow 0$, \tilde{u}_k converges uniformly (up to a subsequence) in $B_{\frac{1}{2}}$ to a Hölder continuous function u^* with $u^*(0) = 0$, $\|u^*\|_{L^\infty} \leq C$, and u^* solves the transmission problem

$$\begin{cases} \mathcal{L}_p(u^*) = 0, & \text{in } B_{\frac{1}{2}} \cap \{x_n \neq 0\}, \\ \partial_n^+ u^* - b \partial_n^- u^* = 0, & \text{on } B_{\frac{1}{2}} \cap \{x_n = 0\}, \end{cases}$$

with $b > 0$ bounded by a universal constant. (Indeed, in Proposition 3.1 we can choose $L_0 = \frac{1}{2}$ and $L_1 = 2$ due to the condition $|\nabla u| \leq 1$.)

Next, we claim that u^* is monotone decreasing in the \mathbf{e}_n -direction, i.e.

$$(39) \quad \partial_n u^* \leq 0.$$

To prove this claim, we consider the following nonlinear elliptic equation satisfied by \tilde{u}_k

$$(40) \quad \sum_{i,j=1}^n \theta_{ij} \partial_{x_i x_j} \tilde{u}_k + \sum_{i=1}^n \beta_i \partial_{x_i} \tilde{u}_k = \frac{2-p}{\epsilon_k |\nabla u_k|^2} x^T M_k^2 \nabla u_k, \quad \text{in } \Omega_{u_k}^+ \cap \{|\nabla u_k| \neq 0\}.$$

where $\theta_{ij} = \delta_{ij} + (p-2)|\nabla u_k|^{-2} \partial_{x_i} u_k \partial_{x_j} u_k$ and $\beta_i = (p-2)|\nabla u_k|^{-2} \sum_{j=1}^n M_{kij} \partial_{x_j} u_k$.

As $\Gamma_{\text{TP}}^{\text{int}}(u_k) \subset \{|x_n| \leq \epsilon_k^{\frac{1}{2}}\}$, if $\bar{x} \in \{x_n > 0\}$, we have for k large enough

$$B := B_{\frac{\bar{x}_n}{2}}(\bar{x}) \subset \Omega_{u_k}^+,$$

and we first show that

$$(41) \quad |\nabla \tilde{u}_k| \leq C(\bar{x}_n), \quad \text{in } B.$$

Inside $\{|\nabla u_k| \geq \frac{1}{4}\} \cap B$, we get $|\nabla \tilde{u}_k| \leq C$ from (40) for a universal constant C . Thus,

$$|\nabla u_k| \geq 1 - \delta_k \epsilon_k^{\frac{1}{2}} - C \epsilon_k \geq \frac{1}{2},$$

for sufficiently large k . This along with the continuity of ∇u inside $\Omega_{u_k}^+$ necessitates that $|\nabla u_k| \geq \frac{1}{2}$ for sufficiently large k . This proves (41)

Hence,

$$\nabla u_k(\bar{x}) = \nabla(P_{M_k, \mathbf{e}_n} + \epsilon_k \tilde{u}_k)(\bar{x}) = \mathbf{e}_n + M_k \bar{x} + \epsilon_k \nabla \tilde{u}_k(\bar{x}).$$

Using that $|\nabla u_k(\bar{x})|^2 \leq 1$ and $M_k \mathbf{e}_n = 0$, we get

$$1 \geq 1 + |M_k \bar{x}|^2 + 2\epsilon_k \partial_n \tilde{u}_k(\bar{x}) + O(\epsilon_k^{\frac{3}{2}}),$$

where the constant in $O(\epsilon_k^{\frac{3}{2}})$ depends on \bar{x}_n . By $C^{1,\alpha}$ estimates of (40) in the ball B , \tilde{u}_k converges to u^* in C^1 . Passing to the limit, as $k \rightarrow \infty$, we get that

$$\partial_n u^*(\bar{x}) \leq 0.$$

By continuity, since u^* is $C^{1,\alpha}$ up to $\{x_n = 0\}$, we conclude that

$$\partial_n^+ u^* \leq 0, \quad \text{in } B_1 \cap \{x_n \geq 0\}.$$

We can argue similarly to show that $\partial_n^- u^* \leq 0$.

Step 2. Contradiction. According to Proposition 4.3, i.e. Linear dichotomy (since (39) holds and $\|u^*\|_{L^\infty} \leq 1$), there exist universal constants $c_0 > 0$ and $r_0 > 0$ such that either of the following conditions is satisfied:

$$(i) \quad \partial_n^+ u^*(0) \leq -c_0, \text{ or}$$

(ii) $|u^*(x) - Q(x')| \leq \frac{1}{4}r_0^2$ in B_{r_0} , where $Q(x')$ is a quadratic polynomial in the x' -direction with $\|Q\| \leq C$ universal and $\mathcal{L}_p(Q) = \Delta Q = 0$.

If (i) above is satisfied, by the quadratic estimate for u^* (see [19, Theorem 3.2] or [8, Lemma 4.12]), we obtain that

$$|u^*(x) - (\xi' \cdot x' + sx_n^+ - tx_n^-)| \leq Cr^2, \quad \text{for } x \in B_r,$$

with $|s|, |t|, |\xi'| \leq C$, and

$$s - bt = 0, \quad s \leq -c_0.$$

Using the convergence of the \tilde{u}_k to u^* and the definition of \tilde{u}_k , we immediately get that for k large,

$$|u_k(x) - (P_{M_k, \mathbf{e}_n}(x) + \epsilon_k(\xi' \cdot x' + sx_n^+ - tx_n^-))| \leq C\epsilon_k r^2, \quad \text{for } x \in B_r.$$

Arguing as in Step 3 of the proof of Proposition 3.1, we conclude that for $r \leq r_1$,

$$|u_k - V_{\bar{a}_k^+, \bar{M}_k, \nu_k}| \leq \epsilon_k r^{1+\alpha_1}, \quad \text{with } \bar{a}^+ = 1 + \epsilon_k s.$$

Thus, after rescaling (see Remark 3.2), Corollary 3.3 gives that

$$|\nabla u_k^+(0)| \leq 1 + s\epsilon_k + C\epsilon_k r^{1+\alpha_1} \leq 1 - \frac{1}{2}c_0\epsilon_k,$$

provided that r is chosen small, universal. Thus, u_k satisfies alternative (i) in Proposition 4.1 for $c'_0 = \frac{c_0}{2}$ and we arrive at a contradiction.

If condition (ii) above is satisfied, since $u^*(0) = 0$, we have

$$Q(x') = \xi' \cdot x' + \frac{1}{2}x^T N x, \quad N \mathbf{e}_n = 0, \quad \text{tr}(N) = 0.$$

Now using the convergence of the \tilde{u}_k to u^* and the definition of \tilde{u}_k , we get

$$(42) \quad \left| u_k(x) - \left(x_n + \epsilon_k \xi' \cdot x' + \frac{1}{2} x^T N_k^* x \right) \right| \leq \frac{1}{4} \epsilon_k r_0^2,$$

with

$$N_k^* = M_k + \epsilon_k N, \quad N_k^* \mathbf{e}_n = 0, \quad \text{tr}(N_k^*) = 0,$$

where we have used $\text{tr}(M_k) = 0$ and $\text{tr}(N) = 0$. Finally, as before, denote

$$\nu_k = \frac{\mathbf{e}_n + \epsilon_k(\xi', 0)}{\sqrt{1 + \epsilon_k^2 |\xi'|^2}} = \mathbf{e}_n + \epsilon_k(\xi', 0) + \epsilon_k^2 \tau, \quad |\tau| \leq C.$$

We argue as in Step 3 of Proposition 3.1, and decompose

$$N_k^* = \bar{M}_k + D_k, \quad \bar{M}_k \nu_k = 0, \quad \|D_k\| = o(\epsilon_k),$$

with $\text{tr}(\bar{M}_k) = 0$. Thus, (42) yields for k large

$$|u_k - P_{\bar{M}_k, \nu_k}| \leq \frac{1}{2} \epsilon_k r_0^2 \leq \epsilon_k r_0^{2+\alpha_0},$$

for α_0 small enough, and again we have reached a contradiction. \square

5. CLASSIFICATION OF LIPSCHITZ GLOBAL SOLUTIONS

The objective of this section is to prove Theorem 1.1, a Liouville type theorem for the Lipschitz global viscosity solutions of the two-phase Bernoulli problem (2). Before that, we establish the following lemma which shows that a two-plane solution with maximum slopes at some scale (possibly small) can approximate the solution u well. Then, by using Proposition 4.1, we obtain a "reversed" improvement of the flatness, and hence conclude that the solution u is well-approximated by the same two-plane solution at all large scales.

Let u be a Lipschitz viscosity solution to (2) in $D = \mathbb{R}^n$, i.e. a global solution. Define

$$(43) \quad \alpha := \sup_{\Omega_u^+} |\nabla u|, \quad \beta := \sup_{\Omega_u^-} |\nabla u|.$$

An initial flatness condition for u can be obtained by the following lemma, at any scale.

Lemma 5.1. *Let u , α and β be as above. Then,*

$$\alpha^p - \beta^p = \lambda_+^p - \lambda_-^p \quad \text{and} \quad \alpha \geq \lambda_+, \quad \beta \geq \lambda_-.$$

Moreover, if $\Gamma_{\text{TP}}^{\text{int}}(u) \neq \emptyset$, then there exists a sequence $u_k(x) := \frac{u(x_k + d_k x)}{d_k}$ of rescalings of u in B_R (for any fixed $R > 1$) with

$$u_k \rightarrow \alpha x_n^+ - \beta x_n^-,$$

uniformly on B_R , as $k \rightarrow +\infty$.

Proof. Given $\epsilon > 0$, let $x_\epsilon \in \Omega_u^+ = \{x \in \mathbb{R}^n : u(x) > 0\}$ be such that

$$|\nabla u(x_\epsilon)| \geq \alpha - \epsilon.$$

Let d_ϵ be the distance from x_ϵ to $\partial\Omega_u^+$ and consider the rescaling

$$u_\epsilon(x) := \frac{u(x_\epsilon + d_\epsilon x)}{d_\epsilon}.$$

These rescalings will still satisfy (2), with $B_1 \subset \Omega_{u_\epsilon}^+$ being tangent to $\partial\Omega_{u_\epsilon}^+ = \partial\{x \in \mathbb{R}^n : u_\epsilon(x) > 0\}$. In fact, after a rotation, we can also assume that

$$\nabla u_\epsilon(0) = t_\epsilon \mathbf{e}_n, \quad \alpha \geq t_\epsilon \geq \alpha - \epsilon.$$

Since u_ϵ is uniformly Lipschitz, up to extracting a subsequence, we can conclude that

$$u_\epsilon \rightarrow \bar{u},$$

uniformly on compacts in \mathbb{R}^n , as ϵ goes to zero. In addition, the convergence is in $C^{1,\eta}(B_{\frac{1}{2}})$ due to $\Delta_p u_\epsilon = 0$ in B_1 . Thus \bar{u} solves (2) in $D = B_R$, for any fixed $R > 1$, and it satisfies

$$\partial_n \bar{u}(0) = \alpha, \quad \text{and} \quad |\nabla \bar{u}| \leq \alpha \quad \text{in} \quad \Omega_{\bar{u}}^+,$$

with $B_1 \subset \Omega_{\bar{u}}^+$ tangent to $\partial\Omega_{\bar{u}}^+$. Hence, we have $\Delta_p \bar{u} = 0$ in B_1 and $\bar{u} \geq 0$ in B_1 , but it is not identically zero because $\partial_n \bar{u}(0) = \alpha > 0$. Therefore $\bar{u} > 0$ in B_1 . Since \bar{u} is p -harmonic, it is locally $C^{1,\eta}$ and we can choose a small $\delta > 0$ such that $\partial_n \bar{u} > \frac{\alpha}{2}$ in B_δ . Thus, $v = \partial_n \bar{u}$ will satisfy the uniform elliptic equation

$$(44) \quad \sum_{i,j=1}^n \partial_i (a_{ij} \partial_j v) = 0, \quad a_{ij} = |\nabla \bar{u}|^{p-2} \left(\delta_{ij} + (p-2) \frac{\partial_i \bar{u} \partial_j \bar{u}}{|\nabla \bar{u}|^2} \right),$$

in B_δ . Hence by the strong maximum principle

$$\partial_n \bar{u} \equiv \alpha, \quad \text{in } \mathcal{O},$$

where \mathcal{O} is the connected component on Ω_u^+ containing B_1 . Therefore,

$$\bar{u}(x) = \alpha x_n + \text{const}, \quad \text{in } \mathcal{O}.$$

Moreover, $\bar{u} = 0$ at the point, say $-e_n$, where the unit ball is tangent to $\partial\Omega_u^+$; thus

$$\bar{u}(x) = \alpha(x_n + 1), \quad \text{in } \{x_n > -1\}.$$

On the other hand, \bar{u} is also a viscosity solution to (2) and we must have

$$\alpha \geq \lambda_+.$$

Now recall (43). Then, $|\nabla \bar{u}| \leq \beta$ in Ω_u^- shows that the function

$$H(x) := \alpha(x_n + 1)^+ - \beta(x_n + 1)^-,$$

touches \bar{u} from below on $\{x_n = -1\}$. We distinguish here two cases:

Case (i): $\{x_n = -1\} \cap \Gamma_{\text{OP}}(\bar{u}) \neq \emptyset$.

From the definition of viscosity solution, we obtain that

$$\alpha = |\nabla H^+| \leq \lambda_+.$$

So, we have $\alpha = \lambda_+$ in this case.

Case (ii): $\{x_n = -1\} \subseteq \Gamma_{\text{TP}}(\bar{u})$.

From the definition of viscosity solution, we must have

$$\alpha^p - \beta^p = |\nabla H^+|^p - |\nabla H^-|^p \leq \lambda_+^p - \lambda_-^p.$$

Similarly, we can argue as above, starting with points $x_\epsilon \in \{x \in \mathbb{R}^n : u(x) < 0\}$ where $\nabla u(x_\epsilon) > \beta - \epsilon$, to get $\beta \geq \lambda_-$, as well as one of the following cases hold:

$$\beta = \lambda_- \quad \text{or} \quad \alpha^p - \beta^p \geq \lambda_+^p - \lambda_-^p.$$

Thus, we have shown that

$$\alpha^p - \beta^p = \lambda_+^p - \lambda_-^p.$$

In order to prove the second part of the lemma, notice that if either $\alpha > \lambda_+$ or $\beta > \lambda_-$, we are in situation Case (ii), and hence $\Omega_u^- = \{x_n < -1\}$. Noting that $H(x)$ touches $\bar{u}(x)$ from below on $\{x_n = -1\}$, and invoking the Hopf boundary lemma, we infer that

$$\bar{u}(x) = \alpha(x_n + 1)^+ - \beta(x_n + 1)^-.$$

Now, consider the case $\alpha = \lambda_+$ and $\beta = \lambda_-$, and assume that $0 \in \Gamma_{\text{TP}}^{\text{int}}(u)$. Let

$$\ell = \limsup_{\Omega_u^+ \ni x \rightarrow 0} |\nabla u(x)| \leq \alpha = \lambda_+,$$

and choose a maximizing sequence $y_k \rightarrow 0$ such that $|\nabla u(y_k)| \rightarrow \ell$ and $u(y_k) > 0$. Repeating the argument in the first part of the proof, we conclude that $\ell \geq \lambda_+$ and $\frac{u(y_k + d_k x)}{d_k} \rightarrow \bar{u}(x)$, up to a subsequence. Also, $\bar{u}(x) = \lambda_+(x_n + 1)$ in $\{x_n > -1\}$. Since $0 \in \Gamma_{\text{TP}}^{\text{int}}(u)$ and $|B_r \cap \{u = 0\}| = 0$ for some $r > 0$, so $\{x_n = -1\} \subseteq \Gamma_{\text{TP}}(\bar{u})$ and

$$\bar{u}(x) = \lambda_+(x_n + 1)^+ - \lambda_-(x_n + 1)^-.$$

□

Remark 5.2. From now on, for the sake of convenience, we assume that

$$\alpha = \beta = 1.$$

This is not a restrictive assumption as one can replace u by $\alpha^{-1}u^+ - \beta^{-1}u^-$. In this case, we have $\lambda_+ = \lambda_- \leq 1$.

Lemma 5.3. Assume $|\nabla u| \leq 1$, $0 \in \Gamma_{\text{TP}}(u)$ and $te_n \in \Omega_u^+$ for all $t \in (0, 1]$. If

$$\partial_n u(te_n) \geq 1 - \eta, \quad \forall t \in (0, 1],$$

for some $\eta > 0$, then

$$|u(x) - x_n| \leq \epsilon(\eta), \quad \text{in } B_2,$$

with $\epsilon(\eta) \rightarrow 0$ as $\eta \rightarrow 0$.

Proof. This follows immediately by compactness, the same lines of arguments as the proof of Lemma 5.1. Assume by contradiction that there exists a sequence $\eta_k \rightarrow 0$, and a sequence of equi-Lipschitz solutions u_k such that

$$(45) \quad \partial_n u_k(te_n) \geq 1 - \eta_k, \quad \forall t \in (0, 1], te_n \in \Omega_{u_k}^+,$$

but

$$(46) \quad |u_k(x) - x_n| > \delta, \quad \text{at some point } x \in B_2 \text{ for some fixed } \delta > 0.$$

From $u_k(0) = 0$ and (45), we obtain

$$u_k(e_n) \geq 1 - \eta_k.$$

Since $|\nabla u_k| \leq 1$, it follows that

$$u_k > 0, \quad \text{in } B_{1-\eta_k}(e_n).$$

We can now argue as in the previous lemma and extract a subsequence that will converge uniformly on $\overline{B_2}$ to $\bar{u} = x_n^+ - x_n^- = x_n$ and contradict (46). \square

We are now ready to provide the proof of Theorem 1.1.

Proof of Theorem 1.1. According to Lemma 5.1, after a translation, a rotation, and a rescaling, we can assume that (recall also the assumption $\alpha = \beta = 1$)

$$0 \in \Gamma_{\text{TP}}^{\text{int}}(u), \quad \text{and} \quad |u - x_n| \leq \epsilon' \quad \text{in } B_1,$$

for ϵ' small, to be made precise later. Now, let ϵ_0 be a sufficiently small universal constant so that the conclusions of Proposition 4.1, Proposition 3.1, and Corollary 3.3 hold for all $\epsilon \leq \epsilon_0$. If $\epsilon' \leq \epsilon_0$, then Corollary 3.3 implies that $\Gamma_{\text{TP}}^{\text{int}}(u) \cap B_{\frac{1}{2}}$ is locally $C^{1,\eta}$ smooth and $u \in C^{1,\eta}(\overline{\Omega_u^+ \cap B_{\frac{1}{2}}})$.

By denoting $h := u - x_n \in C^{1,\eta}(\overline{\Omega_u^+ \cap B_{\frac{1}{2}}})$, we easily achieve the following non-linear uniformly elliptic equation

$$(47) \quad \sum_{i,j=1}^n \theta_{ij} \partial_{x_i x_j} h = 0, \quad \text{in } \{\nabla u \neq 0\},$$

with $C^{0,\eta}$ coefficients $\theta_{ij} = \delta_{ij} + (p-2)|\nabla u|^{-2} \partial_{x_i} u \partial_{x_j} u$. Now, choose the largest ball B_r (for $r \leq \frac{1}{2}$) such that $\nabla u \neq 0$ inside $\Omega_u^+ \cap B_r$; then, the flatness of the free boundary implies that $|u_r - x_n| \leq C\epsilon'$ for a universal constant C . So, $h_r = u_r - x_n$ satisfies (47) in $\Omega_{u_r}^+ \cap B_1$, and the $C^{1,\eta}$ estimate of h_r yields that

$$|\partial_n u - 1| \leq C\epsilon', \quad \text{in } \Omega_u^+ \cap B_r,$$

for a universal constant C . If ϵ_0 is sufficiently small, we can conclude that $\nabla u \neq 0$ in $\Omega_u^+ \cap B_{\frac{1}{2}}$. Using again $C^{1,\eta}$ estimate of h in (47), we deduce that

$$(48) \quad \partial_n u(t\mathbf{e}_n) \geq 1 - C\epsilon', \quad \forall t \in \left(0, \frac{1}{2}\right],$$

and also

$$(49) \quad \partial_n u^+(0) := \lim_{t \rightarrow 0^+} \frac{u(t\mathbf{e}_n)}{t} \geq 1 - C\epsilon'.$$

Now, let $\epsilon(\eta)$ be as defined in Lemma 5.3, and choose a universal constant $\bar{\eta}$ small enough such that $\epsilon(\bar{\eta}) \leq \epsilon_0$. We claim that the following inequality holds

$$(50) \quad \partial_n u(t\mathbf{e}_n) \geq 1 - \bar{\eta}, \quad \text{for all } t > 0,$$

provided that ϵ' is chosen sufficiently small. Suppose this is true, then according to Lemma 5.3,

$$|u - x_n| \leq \epsilon(\bar{\eta})R \leq \epsilon_0 R, \quad \text{in } B_R,$$

for all R large. This, along with the improvement of flatness [8, Theorem 4.1], implies that

$$|u - U_{a^+, \bar{v}}| \leq \epsilon_0 \rho^{1+\gamma} R, \quad \text{in } B_{\rho R},$$

for some $a^+ = a^+(R)$, $\bar{v} = \bar{v}(R)$, and universal constants $\rho, \gamma > 0$. Then, for all R

$$\text{dist}(u, \mathcal{U}_{\lambda_+, \lambda_-}) \leq \epsilon_0 \rho^\gamma R,$$

where the distance is measured in $L^\infty(B_R)$. Repeat this argument to find out $\text{dist}(u, \mathcal{U}_{\lambda_+, \lambda_-}) \rightarrow 0$, for a fixed R . So, u must be a two-plane solution which concludes the proof of Theorem 1.1.

Now, we should prove the claim (50). For this denote $C\epsilon'$ from (48) by $\epsilon'' := C\epsilon'$, and we will choose ϵ'' (and therefore ϵ') later, so that $\epsilon'' \ll \bar{\eta}$. Let \bar{t} be the first t for which (50) fails, and assume without loss of generality after a rescaling that $\bar{t} = 1$. In other words,

$$(51) \quad \partial_n u(\mathbf{e}_n) = 1 - \bar{\eta}, \quad \text{and} \quad \partial_n u(t\mathbf{e}_n) \geq 1 - \bar{\eta}, \quad \forall t \in [0, 1].$$

Notice that after rescaling, (49) is still satisfied, i.e.

$$\partial_n u^+(0) \geq 1 - \epsilon''.$$

From Lemma 5.3 and (51), we conclude that

$$|u - x_n| \leq \epsilon(\bar{\eta}) \leq \epsilon_0, \quad \text{in } B_2.$$

Now, Corollary 3.3 implies that $u \in C^{1,\eta}(\overline{\Omega_u^+ \cap B_1})$ and that $\Gamma_{\text{TP}}^{\text{int}}(u) \cap B_1$ is a $C^{1,\eta}$ graph with small norm.

Since $1 - \partial_n u$ satisfies the uniform elliptic equation (44), we are able to apply the Harnack inequality for $1 - \partial_n u \geq 0$ in $\{x_n > \epsilon_0\}$, and noting that $(1 - \partial_n u)(\mathbf{e}_n) = \bar{\eta}$, gives that

$$1 - \partial_n u \geq c\bar{\eta}, \quad \text{in } B_{\frac{1}{4}}\left(\frac{\mathbf{e}_n}{2}\right) \subset \Omega_u^+ \cap B_1.$$

In $(\Omega_u^+ \cap B_1) \setminus B_{\frac{1}{4}}\left(\frac{\mathbf{e}_n}{2}\right)$, we compare $1 - \partial_n u$ with a suitable barrier function w , which is a solution to (44) with the following boundary values

$$\begin{cases} w = 0, & \text{on } \partial\Omega_u^+ \cap B_1, \\ w = c\bar{\eta}, & \text{on } \partial B_{\frac{1}{4}}\left(\frac{\mathbf{e}_n}{2}\right), \\ w = 0, & \text{on } \partial B_1 \cap \Omega_u^+. \end{cases}$$

Using the Hopf lemma in $C^{1,\eta}$ domains, together with $C^{1,\eta}$ estimates up to the boundary, we conclude that w grows linearly away from $\partial\Omega_u^+ \cap B_1$; hence

$$\partial_n u(t\mathbf{e}_n) \leq 1 - c\bar{\eta}t, \quad \forall t \in \left[0, \frac{1}{4}\right],$$

with c independent of u . Integrating in the \mathbf{e}_n -direction, and using that $u(0) = 0$ and $c, \bar{\eta}$ are universal, we find

$$(52) \quad u(t\mathbf{e}_n) \leq t - c_1 t^2, \quad \forall t \in \left[0, \frac{1}{4}\right],$$

for some small universal constant $c_1 > 0$.

We now apply Proposition 4.1 and conclude that either alternative (i) or (ii) is satisfied. However, if ϵ'' is small enough, then alternative (i) cannot hold; otherwise, by (49)

$$1 - c_0\epsilon_0 \geq |\nabla u^+(0)| \geq \partial_n u^+(0) \geq 1 - \epsilon'',$$

which leads to a contradiction. Thus (ii) holds and we can rescale and iterate one more time; see Remark 4.2. Similarly, after applying the conclusion of Proposition 4.1 for m -times, we obtain that if ϵ'' is sufficiently small depending on m and ϵ_0, c_0, r_0 , then only alternative (ii) can hold for the m iterations and conclude that

$$(53) \quad |u - P_{M,v}| \leq \epsilon_0 r^{2+\alpha_0}, \quad \text{in } B_r, \quad r = r_0^m,$$

with

$$(54) \quad \left| \frac{\nabla u^+(0)}{|\nabla u^+(0)|} - v \right| \leq C\epsilon_0 r^{1+\alpha_0}, \quad \text{and} \quad Mv = 0.$$

Here we have also used Corollary 3.3. Now, notice that

$$1 - \epsilon'' \leq \partial_n u^+(0) = \mathbf{e}_n \cdot \nabla u^+(0) \leq |\nabla u^+(0)| \leq 1,$$

which implies

$$\left| \frac{\nabla u^+(0)}{|\nabla u^+(0)|} - \mathbf{e}_n \right| \leq (2\epsilon'')^{\frac{1}{2}}.$$

Thus, if ϵ'' is small enough depending on r , the inequality above together with (54) gives that

$$|v - \mathbf{e}_n| \leq 2C\epsilon_0 r^{1+\alpha_0}.$$

Thus, since $Mv = 0$ and $\|M\| \leq 1$,

$$\left| P_{M,v}\left(\frac{r}{2}\mathbf{e}_n\right) - \frac{r}{2} \right| = \left| \frac{r}{2}\mathbf{e}_n \cdot (v - \mathbf{e}_n) + \frac{r^2}{8}(\mathbf{e}_n - v)^T M(\mathbf{e}_n - v) \right| \leq C' r^{2+\alpha_0},$$

which combined with (53) gives that

$$\left| u\left(\frac{r}{2}\mathbf{e}_n\right) - \frac{r}{2} \right| \leq 2C' r^{2+\alpha_0}.$$

This contradicts (52) for $t = \frac{r}{2}$, as long as r is small enough universal, i.e. m is large enough, and ϵ'' is small enough. This completes the proof of the theorem. \square

The main consequence of Theorem 1.1 is the regularity of the free boundary at regular points, i.e. Theorem 1.2.

Proof of Theorem 1.2. Consider a regular two-phase point of the free boundary. Then, the Lipschitz regularity of the solution, i.e Theorem 2.3, and the nondegeneracy property, i.e. Theorem 2.2, proves that any blow-ups at this point will be a Lipschitz global solution with $0 \in \Gamma_{\text{TP}}^{\text{int}}(u)$. According to Theorem 1.1, we know that this global solution is a two-plane solution, and we get that the (initial) flatness assumption in [8, Theorem 1.1] is satisfied. Therefore, by [8, Theorem 1.1], the boundaries $\partial\Omega_u^\pm$ are $C^{1,\eta}$ smooth in a neighborhood of the regular two-phase point, for any $\eta \in (0, \frac{1}{3})$. \square

APPENDIX A. A HARNACK INEQUALITY

In the following lemma, for the convenience of the readers, we prove a Harnack inequality for an auxiliary problem of the p -Laplacian type.

Lemma A.1. *Let $1 < p < \infty$, and Ω be a smooth bounded domain in \mathbb{R}^n . Suppose $x_0 \in \Omega$, and $0 < R \leq 1$ such that $\overline{B_{4R}(x_0)} \subset \Omega$. Let $h \in W^{1,p}(\Omega) \cap L^\infty(\Omega)$ be a non-negative solution to*

$$(55) \quad \operatorname{div} \left(|\nabla h(x) + Fx + E|^{p-2} (\nabla h(x) + Fx + E) \right) = 0, \quad \text{in } \Omega,$$

with F being an $n \times n$ symmetric matrix, and $E \in \mathbb{R}^n$. Then, there exists C depending only on n, p, F, E , and $\|h\|_{L^\infty(B_{4R}(x_0))}$ such that

$$\sup_{B_R(x_0)} h \leq C \left(\inf_{B_R(x_0)} h + R \right).$$

Proof. Let $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be defined as

$$A(x, \xi) = |\xi + Fx + E|^{p-2} (\xi + Fx + E).$$

Then, (55) can be written in the form

$$\operatorname{div} A(x, \nabla u) = 0, \quad \text{in } \Omega.$$

We first observe that, for every $\xi \in \mathbb{R}^n$,

$$|A(x, \xi)| = |\xi + Fx + E|^{p-1} \leq C_1 \left(|\xi|^{p-1} + \max_{x \in \Omega} |Fx + E|^{p-1} \right),$$

where C_1 depends only on p . On the other hand, for every $\xi \in \mathbb{R}^n$,

$$(56) \quad \begin{aligned} \xi^T A(x, \xi) &= |\xi + Fx + E|^{p-2} \xi^T (\xi + Fx + E) \\ &= |\xi + Fx + E|^p - |\xi + Fx + E|^{p-2} (Fx + E)^T (\xi + Fx + E) \\ &\geq |\xi + Fx + E|^p - \left(\max_{x \in \Omega} |Fx + E| \right) |\xi + Fx + E|^{p-1}. \end{aligned}$$

Now, if $|\xi + Fx + E| \leq 2 \max_{x \in \Omega} |Fx + E|$, we get from (56)

$$(57) \quad \begin{aligned} \xi^T A(x, \xi) &\geq |\xi + Fx + E|^p - 2^{p-1} \left(\max_{x \in \Omega} |Fx + E| \right)^p \\ &\geq C_2 |\xi|^p - C_2 \max_{x \in \Omega} |Fx + E|^p - 2^{p-1} \max_{x \in \Omega} |Fx + E|^p, \end{aligned}$$

where C_2 depends only on p .

On the other hand, if $|\xi + Fx + E| > 2 \max_{x \in \Omega} |Fx + E|$, we obtain from (56)

$$\begin{aligned}
 \xi^T A(x, \xi) &\geq |\xi + Fx + E|^p - \left(\max_{x \in \Omega} |Fx + E| \right) |\xi + Fx + E|^{p-1} \\
 (58) \quad &= |\xi + Fx + E|^p \left(1 - \left(\max_{x \in \Omega} |Fx + E| \right) |\xi + Fx + E|^{-1} \right) \\
 &\geq \frac{1}{2} |\xi + Fx + E|^p \geq C_3 |\xi|^p - C_3 \max_{x \in \Omega} |Fx + E|^p,
 \end{aligned}$$

where C_3 depends only on p . Then, from (57) and (58), we deduce

$$\xi^T A(x, \xi) \geq C_4 |\xi|^p - C_5,$$

where C_4 depends only on p , and C_5 depends on F , E , and p . Now, the result follows from [32, Theorem 1.1]. \square

APPENDIX B. A LIOUVILLE TYPE RESULT FOR THE LINEARIZED OPERATOR

In the following lemma, we present a Liouville type result for the global solutions of the operator \mathcal{L}_p , introduced in (22).

Lemma B.1. *Let U be a global solution to*

$$(59) \quad \begin{cases} \mathcal{L}_p(U) = 0, & \text{in } \mathbb{R}^n \cap \{x_n \neq 0\}, \\ \partial_n^+ U = g \partial_n^- U, & \text{on } \{x_n = 0\}, \end{cases}$$

with $0 < b_0 \leq g \leq b_1$ and

$$(60) \quad \partial_n U \leq 0, \quad |U(x)| \leq C|x|^2,$$

for some constant C . Then, $U = Q(x')$ where $Q(x')$ is a pure quadratic polynomial in the x' -direction such that $\mathcal{L}_p(Q) = \Delta Q = 0$.

Proof. Let

$$U_k(x) := \frac{U(r_k x)}{r_k^2}, \quad \text{with } r_k \rightarrow \infty,$$

be a sequence of blow-downs that converges uniformly on compact sets to another global solution \bar{U} to (59) with (60).

Let τ be a unit tangential direction, i.e. $\tau \perp \mathbf{e}_n$, and denote

$$\gamma = \gamma(\tau) := \sup_{\mathbb{R}^n \setminus \{x_n=0\}} \partial_{\tau\tau} \bar{U}.$$

Notice that $\gamma \neq \infty$. Indeed, by a similar argument of the Step 1 in the proof of Proposition 4.3, we obtain that $(U_k)_{\tau\tau}^h$ is bounded above in B_1 independently of h , and this implies that γ is well-defined.

We claim that

$$(61) \quad \partial_{\tau\tau} \bar{U} \equiv \gamma, \quad \text{on } \{x_n \neq 0\}.$$

Let x_0 be a point, say for simplicity in $B_1 \setminus \{x_n = 0\}$, and let us show that

$$\partial_{\tau\tau} \bar{U}(x_0) = \gamma.$$

Since $U_k \rightarrow \bar{U}$ in $C^{2,\alpha}$ in a small ball $B \subset \{x_n \neq 0\}$ around x_0 , clearly,

$$\partial_{\tau\tau} \bar{U}(x_0) \leq \gamma.$$

Assume by contradiction that the inequality above is strict. Then, for all k large,

$$\partial_{\tau\tau}(U_k)(x_0) \leq \gamma - \delta,$$

for some small $\delta > 0$. By $C^{2,\alpha}$ regularity,

$$\partial_{\tau\tau}(U_k) \leq \gamma - \frac{\delta}{2}, \quad \text{in } B_c(x_0) \subset B.$$

Hence, for all $h > 0$ small,

$$(U_k^h)_{\tau\tau} \leq \gamma - \frac{\delta}{2}, \quad \text{in } B_{c'}(x_0).$$

On the other hand, from the definition of γ , we have

$$(U_k^h)_{\tau\tau} \leq \gamma, \quad \text{in } B_{\frac{3}{4}}.$$

Since $(U_k^h)_{\tau\tau}$ is a solution of (59), we can construct an explicit upper barrier in $B_{\frac{3}{4}} \setminus B_{c'}(x_0)$, and conclude that

$$(U_k^h)_{\tau\tau} \leq \gamma - c(\delta), \quad \text{in } B_{\frac{1}{2}},$$

for all k large. This contradicts the definition of γ , and the claim (61) is proved.

Next, we show that \bar{U} is equal to a quadratic polynomial Q_+ (resp. Q_-) in $\{x_n > 0\}$ (resp. $\{x_n < 0\}$) and

$$(62) \quad Q_+ = Q_- \quad \text{on } \{x_n = 0\}, \quad Q_{\pm}(0) = 0, \quad \nabla Q_{\pm}(0) = 0, \quad \partial_n Q_{\pm} \leq 0.$$

We know that

$$\partial_{\tau\tau} \bar{U} = \text{const}, \quad \forall \tau \perp \mathbf{e}_n;$$

hence

$$\bar{U}(x) = b(x_n) + x' \cdot a(x_n) + Q(x'), \quad a(x_n) = (a_1(x_n), \dots, a_{n-1}(x_n)),$$

for some quadratic polynomial Q and functions b and a_i depending on one variable.

Since \bar{U} satisfies (59), so does $\partial_i \bar{U} = a_i(x_n) + \partial_i Q(x')$ for $i < n$. On the other hand, $\partial_i Q$ is a linear function; hence $\mathcal{L}_p(a_i) = 0$. This means that a_i is linear in the set $\{x_n > 0\}$. Now, we use $\mathcal{L}_p(\partial_n \bar{U}) = 0$, and we argue as above to find that b' is linear in $\{x_n > 0\}$, and our claim (62) is proved.

Invoking condition $\partial_n Q_{\pm} \leq 0$, and $Q_+ = Q_-$ on $\{x_n = 0\}$, we have

$$\partial_{ni} Q_{\pm} = 0, \quad \text{for } i \neq n, \quad \text{and} \quad \partial_{ij} Q_+ = \partial_{ij} Q_-, \quad \text{for } i, j \neq n.$$

Finally, since Q_{\pm} both solve $\mathcal{L}_p(Q_{\pm}) = 0$, from either side of $\{x_n \neq 0\}$, we conclude that

$$\partial_{nn} Q_+ = \partial_{nn} Q_-,$$

and therefore $Q_+ \equiv Q_-$. Let us call this common polynomial Q . In particular, since $\partial_n Q \leq 0$, it follows that $\partial_n Q \equiv 0$. Thus, Q is a pure quadratic polynomial in the x' -direction.

Finally, we need to show that $U \equiv Q$. For all tangential directions τ , we have

$$\sup \partial_{\tau\tau} U = \partial_{\tau\tau} Q.$$

Thus, $U - Q$ is concave in the x' -direction. Moreover

$$(63) \quad (U - Q)(0) = 0, \quad \text{and} \quad \nabla(U - Q)(0) = 0.$$

Hence,

$$U(x', 0) \leq Q(x'),$$

and by the monotonicity of U , we get

$$U(x', x_n) \leq Q(x'), \quad \text{on } \{x_n \geq 0\}.$$

As both U and Q solve (59), we can use (63) and Hopf lemma to conclude that $U = Q$, in $\{x_n \geq 0\}$. By the monotonicity of U , we have $U \geq Q$, in the set $\{x_n \leq 0\}$; hence $U = Q$, again by the Hopf lemma. \square

Acknowledgements. Masoud Bayrami and Morteza Fotouhi was supported by Iran National Science Foundation (INSF) under project No. 4001885.

DECLARATIONS

Data availability statement: All data needed are contained in the manuscript.

Funding and/or Conflicts of interests/Competing interests: The authors declare that there are no financial, competing or conflict of interests.

REFERENCES

- [1] ALT, H. W., AND CAFFARELLI, L. A. Existence and regularity for a minimum problem with free boundary. *J. Reine Angew. Math.* **325** (1981), 105–144.
- [2] ALT, H. W., CAFFARELLI, L. A., AND FRIEDMAN, A. Axially symmetric jet flows. *Arch. Rational Mech. Anal.* **81**, 2 (1983), 97–149.
- [3] ALT, H. W., CAFFARELLI, L. A., AND FRIEDMAN, A. A free boundary problem for quasilinear elliptic equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **11**, 1 (1984), 1–44.
- [4] ALT, H. W., CAFFARELLI, L. A., AND FRIEDMAN, A. Jets with two fluids. I. One free boundary. *Indiana Univ. Math. J.* **33**, 2 (1984), 213–247.
- [5] ALT, H. W., CAFFARELLI, L. A., AND FRIEDMAN, A. Jets with two fluids. II. Two free boundaries. *Indiana Univ. Math. J.* **33**, 3 (1984), 367–391.
- [6] ALT, H. W., CAFFARELLI, L. A., AND FRIEDMAN, A. Variational problems with two phases and their free boundaries. *Trans. Amer. Math. Soc.* **282**, 2 (1984), 431–461.
- [7] ALT, H. W., CAFFARELLI, L. A., AND FRIEDMAN, A. Compressible flows of jets and cavities. *J. Differential Equations* **56**, 1 (1985), 82–141.
- [8] BAYRAMI, M., AND FOTOUHI, M. Regularity in the two-phase bernoulli problem for the p -laplace operator. (2023) [arxiv:2301.11775](https://arxiv.org/abs/2301.11775) (preprint).
- [9] BUCUR, D., AND VELICHKOV, B. Multiphase shape optimization problems. *SIAM J. Control Optim.* **52**, 6 (2014), 3556–3591.
- [10] CAFFARELLI, L. A. A Harnack inequality approach to the regularity of free boundaries. I. Lipschitz free boundaries are $C^{1,\alpha}$. *Rev. Mat. Iberoamericana* **3**, 2 (1987), 139–162.
- [11] CAFFARELLI, L. A. A Harnack inequality approach to the regularity of free boundaries. III. Existence theory, compactness, and dependence on X . *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **15**, 4 (1988), 583–602 (1989).
- [12] CAFFARELLI, L. A. A Harnack inequality approach to the regularity of free boundaries. II. Flat free boundaries are Lipschitz. *Comm. Pure Appl. Math.* **42**, 1 (1989), 55–78.
- [13] CAFFARELLI, L. A., AND CABRÉ, X. Fully nonlinear elliptic equations, vol. 43 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1995.
- [14] CAFFARELLI, L. A., AND SALSA, S. A geometric approach to free boundary problems, vol. 68 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2005.
- [15] DANIELLI, D., AND PETROSYAN, A. A minimum problem with free boundary for a degenerate quasilinear operator. *Calc. Var. Partial Differential Equations* **23**, 1 (2005), 97–124.
- [16] DANIELLI, D., AND PETROSYAN, A. Full regularity of the free boundary in a Bernoulli-type problem in two dimensions. *Math. Res. Lett.* **13**, 4 (2006), 667–681.
- [17] DE PHILIPPIS, G., SPOLAOR, L., AND VELICHKOV, B. Regularity of the free boundary for the two-phase Bernoulli problem. *Invent. Math.* **225**, 2 (2021), 347–394.
- [18] DE SILVA, D. Free boundary regularity for a problem with right hand side. *Interfaces Free Bound.* **13**, 2 (2011), 223–238.

- [19] DE SILVA, D., FERRARI, F., AND SALSA, S. Two-phase problems with distributed sources: regularity of the free boundary. *Analysis & PDE* 7, 2 (2014), 267–310.
- [20] DE SILVA, D., AND SAVIN, O. Global solutions to nonlinear two-phase free boundary problems. *Comm. Pure Appl. Math.* 72, 10 (2019), 2031–2062.
- [21] DIPIERRO, S., AND KARAKHANYAN, A. L. Stratification of free boundary points for a two-phase variational problem. *Adv. Math.* 328 (2018), 40–81.
- [22] FERRARI, F., AND LEDERMAN, C. Regularity of flat free boundaries for a $p(x)$ -Laplacian problem with right hand side. *Nonlinear Anal.* 212 (2021), Paper No. 112444, 25.
- [23] FERRARI, F., AND LEDERMAN, C. Regularity of lipschitz free boundaries for a $p(x)$ -laplacian problem with right hand side. *Journal de Mathématiques Pures et Appliquées* (2022).
- [24] GILBARG, D., AND TRUDINGER, N. S. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [25] KARAKHANYAN, A. Full and partial regularity for a class of nonlinear free boundary problems. *Annales de l'Institut Henri Poincaré C* 38, 4 (2021), 981–999.
- [26] KARAKHANYAN, A. Regularity for the two-phase singular perturbation problems. *Proceedings of the London Mathematical Society* 123, 5 (2021), 433–459.
- [27] KINDERLEHRER, D., NIRENBERG, L., AND SPRUCK, J. Regularity in elliptic free boundary problems. *J. Analyse Math.* 34 (1978), 86–119 (1979).
- [28] LEWIS, J. L., AND NYSTRÖM, K. Regularity of Lipschitz free boundaries in two-phase problems for the p -Laplace operator. *Adv. Math.* 225, 5 (2010), 2565–2597.
- [29] LEWIS, J. L., AND NYSTRÖM, K. Regularity of flat free boundaries in two-phase problems for the p -Laplace operator. *Ann. Inst. H. Poincaré C Anal. Non Linéaire* 29, 1 (2012), 83–108.
- [30] PETROSYAN, A. On the full regularity of the free boundary in a class of variational problems. *Proc. Amer. Math. Soc.* 136, 8 (2008), 2763–2769.
- [31] VELICHKOV, B. *Regularity of the one-phase free boundaries*. Springer Nature, 2023.
- [32] WOLANSKI, N. Local bounds, Harnack's inequality and Hölder continuity for divergence type elliptic equations with non-standard growth. *Rev. Un. Mat. Argentina* 56, 1 (2015), 73–105.

DEPARTMENT OF MATHEMATICAL SCIENCES, SHARIF UNIVERSITY OF TECHNOLOGY, TEHRAN, IRAN
Email address: masoud.bayrami1990@sharif.edu

Email address: fotouhi@sharif.edu