A ring structure on Tor

Jeffrey D. Carlson

June 9, 2023

Abstract

We prove that within a natural class of E_3 -algebras, the graded group $\bigoplus \operatorname{Tor}_A^i(X, Y)$ induced by a pair of E_3 -algebra maps $X \leftarrow A \rightarrow Y$ carries a functorial commutative graded algebra structure generalizing the classical structure when A, X, Y are genuine commutative differential graded algebras.

As a topological corollary, Munkholm's Eilenberg–Moore collapse result for pullbacks of spaces with polynomial cohomology is enhanced to a ring isomorphism. New applications compute the Borel cohomology rings of homogeneous spaces and singular cohomology rings of biquotients; existing results on the singular cohomology rings of free loop spaces and (generalized) homogeneous spaces are respectively recovered and strengthened.

Among the most fundamental objects of homological algebra are the derived functors Tor^{*i*} of the tensor product. These are individually merely modules, but if $M \leftarrow A \rightarrow N$ are maps of commutative differential graded algebras (CDGAS), then the graded group $\bigoplus_i \operatorname{Tor}_A^i(M, N)$ carries the structure of a bigraded ring denoted simply $\operatorname{Tor}_A(M, N)$, essentially because under the hypothesis of commutativity, the multiplications $A \otimes A \longrightarrow A$ and so on can themselves be seen as maps of differential graded algebras (DGAS). When the input rings are instead cochain algebras $C^*(X) \leftarrow C^*(B) \rightarrow C^*(E)$, it is again classical that there exists a ring structure on Tor, but this is because of the Eilenberg–Zilber theorem, on the face of it an entirely unrelated reason. It is thus natural to wonder under what general conditions on the input DGAS a ring structure on Tor should exist. The question, however, seems never to have been seriously considered.

This situation is the more surprising because there does exist a candidate for such a product. In 1974, Hans J. Munkholm realized the products on $\text{Tor}_{H^*B}(H^*X, H^*E)$ and $\text{Tor}_{C^*B}(C^*X, C^*E)$ could both be described in terms of a structure generalizing CDGAs, called *strongly homotopy commutative* (SHC-) *algebras*, of which both the cochain algebra $C^*(-)$ and cohomology ring $H^*(-)$ are examples.¹ He had used these SHC-algebra structures to obtain a collapse result for certain Eilenberg–Moore spectral sequences (EMSSS), and went on in the last section of his paper to note, almost as an afterthought, that with some additional homotopy data, this alternative construction produces a product on Tor more generally [Mun74, §9]. The construction is involved and rather speculative, and Munkholm was not sanguine about his product's prospects [Mun74, p. 49]:

¹ This language is now unfamiliar, but should not be intimidating: we will be able to state all the relevant facts about SHC-algebras in less than two pages in Section 5. Although they are not strictly comparable, the main examples show they should be thought of as somewhere between E_2 - and E_3 -algebras; see Theorem 5.7 and Remark 5.8.

The composition of (1), (2) and (3) now gives some sort of a product. We have no specific applications of this in mind, so we have not tried to investigate the properties of this product. Presumably they are relatively bad, because of the dependence of (1) on our choice of the homotopies.

But he was wrong to doubt.² It is the main task of the present paper to vindicate this product.³

Theorem 0.1. Let SHC-algebras A, X, Y, and SHC-algebra maps $\mathbf{B}X \leftarrow \mathbf{B}A \rightarrow \mathbf{B}Y$ be given. Then the product defined in Definition 6.2 is independent of the choice of homotopies used to define it and renders $\operatorname{Tor}_A(X, Y)$ a commutative graded algebra. Moreover, the product is functorial in triples of SHC-algebra maps making the two necessary squares commute up to homotopy.

There are other plausible approaches to defining a product on Tor running through highly structured ring spectra (see Remark 8.5.6 for sketches of how this might work), but the the advantage of our approach is that it obtains functoriality of the product under minimal hypotheses on maps of input DGAs. This is critical for us because these hypotheses are already known to hold in the motivating situation, Munkholm's original Eilenberg–Moore collapse result, whereas the stronger hypotheses required for the functoriality of other probable products are not.

Casting our minds back, the advent of the EMSS [EM065, Sm67] had made the cohomology of a wide range of fiber products accessible to computation, loop spaces and two-stage Postnikov systems being popular examples. Cartan [Car51] had famously shown that for coefficient ring $k = \mathbb{R}$ and K < G compact, connected Lie groups, one has

$$H^*(G/K) \cong \operatorname{Tor}_{H^*BG}(k, H^*BK), \tag{0.2}$$

and with a view especially toward generalizing this result to harder coefficient rings, authors including Baum, May, Gugenheim, Munkholm, Halperin, Stasheff, Husemoller, Moore, and Wolf with eventual success—set themselves the task of proving collapse results, if not for the EMSS of a general pullback, then at least for the one-sided variant applying to a fibration of the form $F \rightarrow E \rightarrow B$ with $H^*(E)$ and $H^*(B)$ polynomial and converging to $H^*(F)$.

Munkholm's 1974 result was the most far-reaching of these, not only showing that the sequence collapses, but resolving the additive extension problem. There was no general multiplicative result until 2019,⁴ when Franz [Fr19a] established multiplicativity in the "one-sided" case, meaning (0.2) holds as an isomorphism of graded rings when $H^*(BG;k)$ and $H^*(BK;k)$ are polynomial and 2 is a unit of k. While this represented the greatest progress on the problem in the forty-five years since Munkholm, in the two-sided case, Munkholm's additive isomorphism was not still known to be multiplicative.

It is our second central observation that it is (and one need not invert 2). Since many existing EMSS collapse results factor through Munkholm's, it follows that we have ring isomorphisms much more commonly than it had seemed reasonable to hope. This is Munkholm's result:

² Maddeningly, there is a 1976 reference [Mun76b] to an unpublished paper with the word "multiplicative" in its title [Mun76c]. It is hard not to wonder what Munkholm would have said or why this paper never appeared.

³ More accurately speaking, in this manuscript we construct and valorize our own product; to show it agrees with Munkholm's is in some ways more technical and can wait until a sequel.

⁴ excepting the special case $G/H \rightarrow BH \rightarrow BG$ when $\operatorname{rk} G = \operatorname{rk} H$ and k is a field, which Borel and Baum proved in different ways in their theses [Bor53, Prop. 3.2][Baum68, Cor. 7.5].

Theorem 0.3 (Munkholm). Let $X \to B \leftarrow E$ be a diagram of topological spaces with $E \longrightarrow B$ a Serre fibration such that $\pi_1(B)$ acts trivially on $H^*(E;k)$ and suppose that $H^*(X;k)$, $H^*(B;k)$, and $H^*(E;k)$ are polynomial rings on at most countably many generators. If the characteristic of the principal ideal domain k is 2, assume as well that the \smile_1 -square vanishes on some selection of polynomial generators for $H^*(X;k)$ and $H^*(E;k)$. Then there is a graded k-module isomorphism

$$\operatorname{Tor}_{H^*(B;k)}\left(H^*(X;k), H^*(E;k)\right) \xrightarrow{\sim} H^*(X \underset{B}{\times} E;k).$$
(0.4)

Corollary 0.5. In the situation of Theorem 0.3, the Eilenberg–Moore spectral sequence of $X \rightarrow B \leftarrow E$ collapses with no additive extension problem.

Our enhancement assumes only what Munkholm does, except in characteristic 2, and concludes the isomorphism is multiplicative:

Theorem o.6. Assume the hypotheses of Theorem 0.3 and moreover, if char k = 2, that the \smile_1 -square vanishes on some selection of polynomial generators for $H^*(B;k)$. Then (0.4) is a k-algebra isomorphism.

Corollary 0.7. In the situation of Theorem 0.6, the Eilenberg–Moore spectral sequence of $X \rightarrow B \leftarrow E$ collapses with no additive or multiplicative extension problem.

Applications of Theorem 0.6

The apparently restrictive hypothesis that the input spaces have polynomial cohomology holds for the motivating classical example of a homogeneous space G/K as in (0.2). In fact, we can substantially generalize this.

As noted by Singhof [Sin93], if H and K are subgroups of a topological group G, then the homotopy orbit space

$$EK \underset{K}{\otimes} G/H := \frac{EK \times G/H}{(ek, gH) \sim (e, kgH)}, \quad e \in EG, \ k \in K, \ g \in G$$

is the homotopy pullback of $BK \to BG \leftarrow BH$, realizable as the pullback of the G/H-bundle $EG/H \longrightarrow EG/G$ for a fixed model for EG. If the two-sided action of $K \times H$ on G by $(k,h) \cdot g := kgh^{-1}$ is free, this can be identified with the so-called *biquotient* $K \setminus G/H$, which if G is a Lie group and K, H < G are closed is naturally a smooth manifold. (For K = 1, of course, this reduces to the statement that G/H is the fiber of $EG/H \longrightarrow EG/G$.) Then if K and H are connected, so that after inverting finitely many primes $H^*(BK), H^*(BG)$, and $H^*(BH)$ are all polynomial rings, Theorem 0.6 immediately gives the Borel cohomology ring $H^*_K(G/H) := H^*(EK \otimes_K G/H)$ with suitable coefficients.⁵

Theorem o.8. Let G be a compact, connected Lie group, K and H closed subgroups, and k a principal ideal domain over which the cohomology of the classifying spaces BG, BH, and BK is polynomial. Then we have an isomorphism of graded k-algebras

$$H^*_K(G/H) \xrightarrow{\sim} \operatorname{Tor}_{H^*BG}(H^*BK, H^*BH).$$

⁵ It can happen that $H^*(BK)$ and $H^*(BH)$ are polynomial even if *K* or *H* is disconnected, for example if the component group K/K_0 acts as a reflection group on $H^*(BK_0)$ and *k* is a field of characteristic relatively prime to $|K/K_0|$.

In particular, this determines $H^*(K \setminus G/H)$ if $K \times H$ acts freely on G under the standard action and $H^*(G/H)$ if K = 1.

This improves even on the result of Franz [Fr19a] for G/H, in that 2 no longer need be a unit of *k*. A minimal example in which one recovers previously unavailable torsion is the following. *Example* o.9. Let *H* be the U(1) subgroup of SU(4) with diagonal entries diag(z^{-3} , z, z, z). One previously knew from Franz's result [Fr19a] that, indexing generators by degree,

$$H^*(\mathrm{SU}(4)/H;\mathbb{Z}[\frac{1}{2}]) \cong \frac{\mathbb{Z}[\frac{1}{2}][s_2] \otimes \Lambda[a_5, b_7]}{(3s^2, s^3, s^2a)}.$$

Now one sees that in fact

$$H^*(\mathrm{SU}(4)/H;\mathbb{Z}) \cong \frac{\mathbb{Z}[s_2] \otimes \Lambda[y_5, z_7]}{(6s^2, 2s^3, s^4, 2s^2y, 3s^2z)},$$
$$H^*(\mathrm{SU}(4)/H;\mathbb{F}_2) \cong \frac{\mathbb{F}_2[s_2]}{(s^4)} \otimes \Lambda[x_3, y_5].$$

Another obvious application is to the cohomology of a free loop space *LX*, immediately recovering the main result of a paper of Saneblidze [Sano9]:

Theorem 0.10 (Saneblidze). Let X be a space and k a principal ideal domain such that H^*X is polynomial on an at most countable set Q of generators whose \sim_1 -squares vanish. Then we have ring isomorphisms

$$H^*LX \cong \operatorname{Tor}_{H^*(X \times X)}(H^*X, H^*X) \cong H^*X \otimes \Lambda[s^{-1}Q] \cong H^*X \otimes H^*\Omega X,$$

where $s^{-1}Q$ is a set of generators $s^{-1}q$, for $q \in Q$, with degrees $|s^{-1}q| = |q| - 1$.

Proof. It is well known that the free loop space $LX = Map(S^1, X)$ can be identified with the homotopy pullback of the diagonal map $\Delta: X \longrightarrow X \times X$ along itself.

The third isomorphism in Theorem 0.10 implicitly used the following result on a based loop space ΩB , which also follows from Theorem 0.6:

Theorem 0.11 (Probably Borel). Let B be a space and k a principal ideal domain such that H^*B is polynomial on an at most countable set Q of generators whose \sim_1 -squares vanish. Then we have ring isomorphisms

$$H^*\Omega B \cong \operatorname{Tor}_{H^*(B)}(k,k) \cong \Lambda[s^{-1}Q],$$

where $s^{-1}Q$ is a the set of generators $s^{-1}q$, for $q \in Q$, with degrees $|s^{-1}q| = |q| - 1$.

Proof. In this case, we use the fact ΩB is the homotopy pullback of $* \to B \leftarrow *$.

Counterexample 0.12. We really need the added hypothesis on $H^*(B)$ for Theorem 0.6 to go through. To see that Theorem 0.11 fails without this hypothesis, note that $B^2(\mathbb{Z}/2) = K(\mathbb{Z}/2, 2)$ does *not* satisfy the hypotheses over $k = \mathbb{F}_2$, its cohomology being the polynomial ring generated by the iterated \smile_1 -squares $x_{1+2^{\ell}} = \operatorname{Sq}^{2^{\ell}} \operatorname{Sq}^{2^{\ell-1}} \cdots \operatorname{Sq}^2 \operatorname{Sq}^1 \iota_2 \in H^{1+2^{\ell}}K(\mathbb{Z}/2, 2)$ of the fundamental

class $\iota_2 \in H^2K(\mathbb{Z}/2, 2)$. Thus, although Munkholm's theorem holds for the homotopy pullback $\mathbb{R}P^{\infty} = K(\mathbb{Z}/2, 1) = \Omega K(\mathbb{Z}/2, 2)$ of $* \leftarrow K(\mathbb{Z}/2, 2) \rightarrow *$, affording us an isomorphism

 $\mathbb{F}_{2}[\iota_{1}] = H^{*}K(\mathbb{Z}/2, 1) \cong \operatorname{Tor}_{H^{*}K(\mathbb{Z}/2, 2)}(\mathbb{F}_{2}, \mathbb{F}_{2}) \cong \Lambda_{\mathbb{F}_{2}}[s^{-1}\iota_{2}, s^{-1}x_{3}, s^{-1}x_{5}, s^{-1}x_{9}, \ldots]$

of graded vector spaces, this isomorphism is not multiplicative. For more on loop spaces, see Remark 8.0.3. See Saneblidze [San17] for a detailed account of what can happen for $H^*(\Omega X; \mathbb{F}_2)$ when \smile_1 -squares do not vanish.

Theorem 0.6 also recovers the easiest cases of group cohomology:

Proposition 0.13 (Classical). Let a finitely-generated abelian group A and principal ideal domain k be given. Let $0 \rightarrow R \rightarrow F \rightarrow A \rightarrow 0$ be a presentation of A, which is to say a short exact sequence of groups with R and F free abelian. Then we have a ring isomorphism

$$H^*(BA;k) \cong \operatorname{Tor}_{H^*(B^2F;k)}(k, H^*(B^2R;k)).$$

Proof. There is an evident fiber sequence $BA \rightarrow B^2R \rightarrow B^2F$ of Eilenberg–Mac Lane spaces, and H^*B^2R and H^*B^2F are polynomial rings on generators of degree 2, hence Theorem 0.6 applies with X = * and $(E \rightarrow B) = (B^2R \rightarrow B^2F)$.

Outline

The plan of the work is as follows.

0.14.1. Section 1 defines algebras and coalgebras and the bar–cobar adjunction, as well as the intermediary notion of a *twisting cochain*.

0.14.2. Section 2 brings in the tensor product and some of its interactions with the adjunction.

0.14.3. Section 3 discusses how homotopies of algebra maps can themselves be realized by algebra maps. Particularly, Section 3.1 introduces the path object receiving such homotopies. This material is classical, dating back to Munkholm's work or earlier, until the critical new Proposition 3.1.4 and Corollary 3.1.5. Section 3.2, which is also new, then develops an array of categorical machinery allowing us to manipulate homotopies diagrammatically without leaving the category of algebras.

0.14.4. Section 4 discusses conditions under which maps on Tor of a span of DGAs can be defined, which are classical, and establishes their homotopy-invariance and functoriality, which are new.

0.14.5. Section 5 recalls the notion of an SHC-algebra.

0.14.6. Section 6 motivates and defines our reformulation of Munkholm's product.

0.14.7. Section 7 establishes a CGA structure on Tor, the first clause of Theorem 0.1.

0.14.8. Section 8 proves this algebra structure is functorial in the input data (the second clause of Theorem **0.1**) and homotopy-invariant. In particular, it proves Theorem **0.6**.

These two sections involve some micromanagement of DGA homotopies and a number of diagrams, but mostly rely on the formal properties of the homotopy categories of DGAs and DGCs discussed in the preliminary sections, without recourse to the cochain level. In particular, it proves possible to almost entirely black-box the SHC-algebra technology, and prior familiarity with notions other than DGAs and DGCs is not assumed.

Acknowledgments. The author's interest in this question originated in joint work with Matthias Franz. The author sketched a proof generalizing Franz's homogeneous space result [Fr19a] to biquotients and asked what was left to do. Franz said the missing ingredient was a natural product structure on a two-sided bar construction of homotopy Gerstenhaber algebras and within a month gave a mod-two reduction of what he was convinced were the correct formulas. The author then found the right signs and proved the formulas worked. A draft has by now appeared on the arXiv [CaF21] and a briefer version will appear in *Algebraic & Geometric Topology*.

The author would like to thank Omar Antolín Camarena for listening to the ongoing drama of this proof and for detailed explanations of several sections of Lurie's *Higher Algebra* back when it seemed that approach might be fruitful, Manuel Rivera for discussion of Husemoller–Stasheff–Moore, Jim Stasheff for real magnanimity in discussion of his paper, Peter May for discussing the history of his papers and requesting a sample computation, Bernhard Keller for directing the author to Jesse Burke, Jesse Burke for preventing him from spending any more time trying to find a model category structure on coalgebras over a general ring, Björn Eurenius and David White for showing him such a structure does exist, Markus Szymik for sending him a copy of Stasheff–Halperin [StH70], Larry Smith and Pedro Tamaroff for detailed comments on an earlier version of this paper, Richard Thomas, John Nicholson, and an anonymous, very thorough referee for advice on presentation, and Joanna Quigley for rigorous copyediting of an earlier draft.

1. Algebras, coalgebras, and twisting cochains

Fix for all time a commutative base ring k with unity with respect to which all tensor products and hom-modules are taken. We take as understood the notions of differential graded k-modules and quasi-isomorphisms, of differential graded k-algebras (henceforth DGAs) and differential graded k-coalgebras (DGCs) and maps between them, tensor products, and the Koszul sign convention. A commutative DGA is a CDGA.

All algebras we consider are graded and associative and all coalgebras graded and coassociative. *All algebras we consider are augmented*, with an important exception that we modify in short order to be augmented as well, *and coalgebras coaugmented*. All *differentials d increase degree* by 1, and we use the terms DG *k-module* and *cochain complex* interchangeably. Our DGAs and DGCs are *nonnegatively-graded*. The multiplication $A \otimes A \longrightarrow A$, unit $k \longrightarrow A$, and augmentation $A \longrightarrow k$ of a DGA A are respectively denoted μ , η , and ε , decorated with a subscript Awhen necessary, and the augmentation ideal ker $\varepsilon \cong \operatorname{coker} \eta$ is denoted \overline{A} . The comultiplication $C \longrightarrow C \otimes C$, counit $C \longrightarrow k$, and coaugmentation $k \longrightarrow C$ of a DGC C are respectively denoted Δ , ε , and η , and the coaugmentation coideal coker $\eta \cong \ker \varepsilon$ is denoted \overline{C} .

We write Mod for the category of graded *k*-modules (with no differential) and maps of fixed but arbitrary degree, DGA for the category of augmented *k*-DGAs and augmentation-preserving

7

DGA maps and DGC for the category of coaugmented, *cocomplete k*-DGCs (cocompleteness will be explained shortly) and coaugmentation-preserving DGC maps. All DGA and DGC maps will be of degree 0, but maps $C \longrightarrow A$ from a coalgebra to an algebra are allowed to be homogeneous of varying degrees, as will also be explained momentarily. The base ring *k* itself is considered a DG Hopf algebra concentrated in degree zero, the differential, multiplication, and comultiplication being what they must.

We briefly rehearse some well-known generalities, taking the opportunity to establish notation and conventions which will be leaned on throughout. General background resources include Munkholm [Mun74, §1], Husemoller–Moore–Stasheff [HuMS74, Pt. II], the thesis of Prouté [Pr11], and the book of Loday–Valette [LV, Chs. 1–2]. One must mind the direction of the differential: while our differentials *increase* degree, others' do not.

Given two graded *k*-modules *C* and *A*, we denote by $Mod_n(C, A)$ the *k*-module of *k*-linear maps *f* sending each C_j to A_{j+n} , and set the degree |f| to *n* for such a map. The hom-set $Mod(C, A) = \bigoplus_{n \in \mathbb{Z}} Mod_n(C, A)$ then becomes itself a graded *k*-module. If *C* and *A* are cochain complexes, then Mod(C, A) becomes a cochain complex under the differential $d = d_{Mod(C,A)}$ given by $d(f) \coloneqq d_A f - (-1)^{|f|} f d_C$ [Mun74, §1.1], cochain maps being described by the condition d(f) = 0.

If *C* is a DGC and *A* a DGA, then Mod(C, A) becomes a DGA when endowed with the *cup product* $f \smile g := \mu_A(f \otimes g)\Delta_C$ [Mun74, §1.8].⁶ An element $t \in Mod_1(C, A)$ satisfying the three conditions

$$\varepsilon_A t = 0 = t \eta_C, \qquad d(t) = t \smile t$$

is called a *twisting cochain* [HuMS74, §1.8][HuMS74, Prop. 3.5(1)][Pr11, §§1.5, 4]. Twisting cochains compose with DGC and DGA maps in the sense that given DGAs *A*', *A* and DGCs *C*, *C*' and maps

$$C' \xrightarrow{g} C \xrightarrow{t} A \xrightarrow{f} A',$$

g a DGC map, *t* a twisting cochain, and *f* a DGA map, the maps *ft*, *tg*, and hence *ftg* are all again twisting cochains. Given a DGA *A*, there is a *final* twisting cochain $t^A : \mathbf{B}A \longrightarrow A$ defined by the property that any twisting cochain $t: C \longrightarrow A$ factors uniquely through a DGC map $g_t: C \longrightarrow \mathbf{B}A$ such that $t = t^A \circ g_t$. We denote this conversion in the input–output "deduction rule" format borrowed from proof theory:

$$g_t \colon C \longrightarrow \mathbf{B}A$$
$$t \colon C \longrightarrow A.$$

The DGC **B***A* is referred to as the *bar construction*, and gives the object component of a functor **B**: DGA \longrightarrow DGC [Mun74, §1.6][Pr11, §2.5]. The tautological twisting cochain $t^{(-)}$: **B** \longrightarrow id is a natural transformation.

More explicitly, the bar construction is the tensor coalgebra on the desuspension $s^{-1}\overline{A}$ of \overline{A} , equipped with the sum of the tensor differential and the unique coderivation extending the "bar-deletion" map $(s^{-1}\overline{A})^{\otimes 2} \xrightarrow{\sim} \overline{A}^{\otimes 2} \xrightarrow{\mu} \overline{A} \xrightarrow{\sim} s^{-1}\overline{A}$. The tautological twisting cochain t^A is the composition of the projection $\mathbf{B}A \longrightarrow s^{-1}\overline{A}$ and the resuspension $s^{-1}\overline{A} \xrightarrow{\sim} \overline{A}$.

⁶ To explain the nomenclature, write out the definition of the product in the cohomology theory E^* represented by a ring spectrum E; the name *convolution* is also popular.

As we have defined DGC to contain only *cocomplete* DGCs, the foregoing assertions include the statement that **B***A* is cocomplete, in the following sense. Given a coaugmented DGC *C*, the comultiplication $\Delta: C \longrightarrow C \otimes C$ defines by reduction a map $\overline{\Delta} = (id - \eta \varepsilon)^{\otimes 2} \Delta: \overline{C} \longrightarrow \overline{C} \otimes \overline{C}$ on the coaugmentation coideal, and by coassociativity, the iterates $\overline{\Delta}^{[n]}: \overline{C} \longrightarrow \overline{C}^{\otimes n}$ starting with $\overline{\Delta}^{[1]} = id_{\overline{C}}, \ \overline{\Delta}^{[2]} = \overline{\Delta}, \ \overline{\Delta}^{[3]} = (\overline{\Delta} \otimes id)\overline{\Delta} = (id \otimes \overline{\Delta})\overline{\Delta}$ are well-defined. We say *C* is *cocomplete*⁷ if \overline{C} is exhausted by the increasing filtration by kernels ker $\overline{\Delta}^{[n]}$.

Cocompleteness is the condition needed to extend a twisting cochain $C \longrightarrow A$ to a DGC map $C \longrightarrow \mathbf{B}A$, and holds of any connected DGC and of the bar construction. Cocompleteness is also important for another reason, whose relevance will become clearer when we discuss homotopy [Mun74, §1.3]. If *C* is cocomplete and $h \in Mod_0(C, A)$ satisfies $h\eta_C = \eta_A$, then $(\eta_A \varepsilon_C - h)\eta_C = 0$. Since im $\eta_C = \ker(\operatorname{id} - \eta_C \varepsilon_C)$ and $\overline{\Delta}_C^{[\ell]} = (\operatorname{id} - \eta_C \varepsilon_C)^{\otimes \ell} \Delta_C^{[\ell]}$, the cup-power

$$(\eta_A \varepsilon_C - h)^{\smile \ell} = \mu_A^{[\ell]} (\eta_A \varepsilon_C - h)^{\otimes \ell} \Delta_C^{[\ell]}$$

annihilates the kernel of $\overline{\Delta}_C^{[\ell]}$. These kernels exhaust *C* by cocompleteness, so $\sum_{\ell=0}^{\infty} (\eta_A \varepsilon_C - h)^{-\ell}$ is a finite sum on any element of *C* and hence gives a sensible two-sided cup-inverse h^{-1} to *h*.

Given a DGC *C*, there is also a twisting cochain $t_C: C \longrightarrow \Omega C$ *initial* in the sense that any twisting cochain $t: C \longrightarrow A$ factors uniquely through a DGA map $f^t: \Omega C \longrightarrow A$ such that $t = f^t t_C$:

$$t\colon C\longrightarrow A$$
$$f^t\colon \mathbf{\Omega}C\longrightarrow A.$$

The DGA ΩC is referred to as the *cobar construction*, and gives the object component of a functor Ω : DGC \longrightarrow DGA [Mun74, §1.7]. The tautological twisting cochain $t_{(-)}$: id $\longrightarrow \Omega$ is a natural transformation. Thus the two functors $\Omega \dashv B$ form an adjoint pair [Mun74, §1.9–10]:

$$g_t \colon \mathbf{\Omega} C \longrightarrow A$$
$$f^t \colon C \longrightarrow \mathbf{B} A.$$

We will have frequent recourse to the unit and counit of the adjunction $\Omega \rightarrow B$,

$$\eta: \operatorname{id} \longrightarrow B\Omega$$
 and $\varepsilon: \Omega B \longrightarrow \operatorname{id}$

respectively. These are both natural quasi-isomorphisms and homotopy equivalences on the level of DG modules [HuMS74, Thm. II.4.4–5][Mun74, Cor. 2.15][LH, Lem. 1.3.2.3].

There are a few elementary, purely categorical properties of this adjunction we will use, here included for easy reference.

Lemma 1.1. Given a DGA A and a DGC C, one has $\varepsilon \circ t_{\mathbf{B}A} = t^A : \mathbf{B}A \longrightarrow A$ and $t^{\mathbf{\Omega}C} \circ \eta = t_C : C \longrightarrow \mathbf{\Omega}C$.

Lemma 1.2. Given a DGA map $f: \Omega C \longrightarrow A$, we have

$$\varepsilon_A \circ \mathbf{\Omega} \mathbf{B} f \circ \mathbf{\Omega} \eta_C = f;$$

⁷ Conilpotent is probably currently more popular.

in particular, $\varepsilon_{\Omega C} \circ \Omega \eta_C = \mathrm{id}_{\Omega C}$. Dually, given a DGC map $g \colon C \longrightarrow \mathbf{B}A$, we have

$$\mathbf{B}\varepsilon_A \circ \mathbf{B}\mathbf{\Omega}g \circ \eta_C = g;$$

in particular, $\mathbf{B}\varepsilon_A \circ \eta_{\mathbf{B}A} = \mathrm{id}_{\mathbf{B}A}$.

Corollary 1.3. If *C* is a DGC and *A* a DGA, then any DGA map $f: \Omega C \longrightarrow A$ naturally factors through $\varepsilon: \Omega BA \longrightarrow A$. We write $f^{\#} = \Omega Bf \circ \Omega \eta_C: \Omega C \longrightarrow \Omega BA$ for the first factor. The transformation $f \longmapsto f^{\#}$ is natural in that given another DGA map $\phi: A \longrightarrow B$, one has $(\phi \circ f)^{\#} = \Omega B\phi \circ f^{\#}$.

We will sometimes say $f^{\#}: \Omega C \longrightarrow \Omega BA$ is *induced up* from $f: \Omega C \longrightarrow A$.

Corollary 1.4. *If* C *is a* DGC *and* A *a* DGA*, and* $g: C \longrightarrow BA$ *is a* DGC *map, then* g *factors through* $B\Omega C$ *and* $\Omega g: \Omega C \longrightarrow \Omega BA$ *factors as*

```
\Omega C \xrightarrow{\Omega \eta} \Omega B \Omega C \xrightarrow{\Omega B \Omega g} \Omega B \Omega B A \xrightarrow{\Omega B \varepsilon} \Omega B A.
```

2. The tensor product

Much of the material in this section is not proven in the source literature, and in longer preliminary drafts of this document, proofs of each result were spelled out. For space reasons we have again suppressed these, but it is still convenient to at least gather the statements in one place.

The functor **B**: DGA \longrightarrow DGC is lax monoidal with respect to the monoidal structure given on both categories by the appropriate tensor products, and Ω : DGC \longrightarrow DGA is lax comonoidal.

Definition 2.1 (See Husemoller et al. [HuMS74, Def. IV.5.3]). There exist natural transformations

$$\nabla \colon \mathbf{B}A_1 \otimes \mathbf{B}A_2 \longrightarrow \mathbf{B}(A_1 \otimes A_2), \qquad \gamma \colon \mathbf{\Omega}(C_1 \otimes C_2) \longrightarrow \mathbf{\Omega}C_1 \otimes \mathbf{\Omega}C_2$$

of functors DGA × DGA \rightarrow DGC and DGC × DGC \rightarrow DGA, respectively, the *shuffle maps*, determined by the twisting cochains

$$t^{A_1 \otimes A_2} \nabla = t^{A_1} \otimes \eta_{A_2} \varepsilon_{\mathbf{B}A_2} + \eta_{A_1} \varepsilon_{\mathbf{B}A_1} \otimes t^{A_2}, \qquad \gamma t_{C_1 \otimes C_2} = t_{C_1} \otimes \eta_{\mathbf{\Omega}C_2} \varepsilon_{C_2} + \eta_{\mathbf{\Omega}C_1} \varepsilon_{C_1} \otimes t_{C_2}.$$

These are homotopy equivalences of cochain complexes and hence quasi-isomorphisms.

Remark 2.2. Although we manage to sidestep cochain-level computations completely in this paper with the exception of the easy check in Proposition 3.1.4, it may be psychologically helpful to have an idea of what some of these maps do. The bar shuffle ∇ on $\mathbf{B}_p A \otimes \mathbf{B}_p B$ and cobar shuffle γ on $\mathbf{\Omega}_{\ell}(C \otimes D)$ respectively take

$$\begin{aligned} & [a_1|\cdots|a_p] \otimes [b_1|\cdots|b_q] \longmapsto \sum_{\sigma} \sigma \big([a_1 \otimes 1|\cdots|a_p \otimes 1|1 \otimes b_1|\cdots|1 \otimes b_q] \big), \\ & \langle c_1 \otimes d_1; \cdots; c_\ell \otimes d_\ell \rangle \longmapsto \big(\langle c_1 \rangle \otimes \varepsilon(d_1) + \varepsilon(c_1) \otimes \langle d_1 \rangle \big) \cdots \big(\langle c_\ell \rangle \otimes \varepsilon(d_\ell) + \varepsilon(c_\ell) \otimes \langle d_\ell \rangle \big), \end{aligned}$$

where the shuffles σ are the (p + q)!/p!q! permutations of the bar-word interlacing the "*a*" and "*b*" letters while retaining the relative order of the a_i and that of the b_i , multiplied by a Koszul

sign⁸. Hence a typical term of the former sum for p = q = 2 is $\pm [1 \otimes b_1 | a_1 \otimes 1 | a_2 \otimes 1 | 1 \otimes b_2]$. Sample values of the latter are

$$\begin{split} &\gamma \langle c_1 \otimes d_1 \rangle = 0, & c_1 \in \overline{C}, \ d_1 \in \overline{D}, \\ &\gamma \langle c_1 \otimes 1; c_2 \otimes 1 \rangle = \langle c_1; c_2 \rangle \otimes 1, & c_i \in \overline{C}, \\ &\gamma \langle 1 \otimes d_1; c_2 \otimes 1 \rangle = (-1)^{(|c_2|+1)(|d_1|+1)} \langle c_2 \rangle \otimes \langle d_1 \rangle, & c_2 \in \overline{C}, \ d_1 \in \overline{D}. \end{split}$$

Of course any identity involving these maps needs to be checked for all p, q, ℓ simultaneously and thus involves infinitely many equations, so computations involving multiple such maps rapidly scale beyond tractability.

Fortunately, however, explicit formulae are actually beside the point for our purposes. All we really need to know about the natural transformations ∇ and γ (and also ψ and \otimes , to be introduced momentarily) is that they allow us to move tensor products in and out of **B** and Ω without changing homotopy type. That is, we may treat them as devices for packaging and formally manipulating homotopical information. We will not really need to look under the hood, and given the complexity of the machinery, most of the time it is actually safer not to.

There is another important natural transformation on DGA we will rely on.

Theorem 2.3 ([HuMS74, Prop. IV.5.5][Mun74, k_{A_1,A_2} , p. 21, via Prop. 2.14]). *There exists a natural transformation*

 $\psi \colon \mathbf{\Omega}\mathbf{B}(A_1 \otimes A_2) \longrightarrow \mathbf{\Omega}\mathbf{B}A_1 \otimes \mathbf{\Omega}\mathbf{B}A_2$

of functors $DGA \times DGA \longrightarrow DGA$. This transformation satisfies

$$(\varepsilon_{A_1} \otimes \varepsilon_{A_2}) \circ \psi = \varepsilon_{A_1 \otimes A_2} \colon \mathbf{\Omega} \mathbf{B}(A_1 \otimes A_2) \longrightarrow A_1 \otimes A_2$$

and reduces to the identity if A_1 or A_2 is k.

Remark 2.4. Husemoller–Moore–Stasheff's construction of such a map relies on a splitting arising from the notion of an *injective class* in their categorical reformulation of differential homological algebra, and is not very explicit. Munkholm's map arises from the fact $\varepsilon_{A_1 \otimes A_2}$: $\Omega B(A_1 \otimes A_2) \longrightarrow$ $A_1 \otimes A_2$ is the initial object in a category of *trivialized extensions* of $A_1 \otimes A_2$, of which $\varepsilon_{A_1} \otimes \varepsilon_{A_2}$ is another object; each is a DGA quasi-isomorphism with a DG-module quasi-inverse and a contracting homotopy. The initiality follows the homotopy transfer theorem for DGAs and the finality of the tautological twisting cochain $t^{A_1 \otimes A_2}$: $B(A_1 \otimes A_2) \longrightarrow A_1 \otimes A_2$, so that Munkholm's ψ is not given terribly explicitly either. However, the homotopy transfer theorem is proven via the inductive construction of a certain twisting cochain from a given first stage, and Munkholm gives an explicit contracting homotopy for ε , so that the construction is at least explicit enough that in a sequel paper, we are able to show by computation that $\psi \circ \Omega \nabla = \gamma$, allowing us to see the product on Tor defined in this paper agrees with that defined by Munkholm.

Definition 2.5. Let A and B be DGAS. A DGC map $\mathbf{B}A \longrightarrow \mathbf{B}B$ is called an A_{∞} -map from A to B.

Evidently for $f: A \longrightarrow B$ a DGA map, **B***f* is an A_{∞} -map, but most A_{∞} -maps are not of this type. The natural transformation ψ allows us to also take tensor products of A_{∞} -maps.

⁸ That is, -1 to the power of $\sum (|a_i| - 1)(|b_j| - 1)$, where the sum runs over a_i and b_j that the shuffle has moved past one another.

Definition 2.6 ([Mun74, Prop. 3.3]). Let A_1, A_2, B_1, B_2 be DGAs and $g_j: \mathbf{B}A_j \longrightarrow \mathbf{B}B_j$ be DGC maps for $j \in \{1, 2\}$. Then we define the *internal tensor product* $g_1 \otimes g_2: \mathbf{B}(A_1 \otimes A_2) \longrightarrow \mathbf{B}(B_1 \otimes B_2)$ as the composition

$$\mathbf{B}(A_1 \otimes A_2) \xrightarrow{\eta} \mathbf{B} \mathbf{\Omega} \mathbf{B}(A_1 \otimes A_2) \xrightarrow{\mathbf{B} \psi} \mathbf{B}(\mathbf{\Omega} \mathbf{B} A_1 \otimes \mathbf{\Omega} \mathbf{B} A_2) \xrightarrow{\mathbf{B}(\varepsilon \, \mathbf{\Omega} g_1 \otimes \varepsilon \, \mathbf{\Omega} g_2)} \mathbf{B}(B_1 \otimes B_2).$$

This construction exhibits some functoriality:

Lemma 2.7 ([Mun74, Prop. 3.3(ii)]). Given DGA maps $f_j: A_j \longrightarrow B_j$ for $j \in \{1, 2\}$, we have

$$\mathbf{B}f_1 \underline{\otimes} \mathbf{B}f_2 = \mathbf{B}(f_1 \otimes f_2) \colon \mathbf{B}(A_1 \otimes A_2) \longrightarrow \mathbf{B}(B_1 \otimes B_2)$$

If A'_i and B'_i are further DGAs and $g_j: \mathbf{B}A'_i \longrightarrow \mathbf{B}A_j$ and $\ell_j: \mathbf{B}B_j \longrightarrow \mathbf{B}B'_i$ DGC maps, then

$$(\ell_1 \underline{\otimes} \ell_2) \circ \mathbf{B}(f_1 \otimes f_2) = (\ell_1 \circ \mathbf{B}f_1) \underline{\otimes} (\ell_2 \circ \mathbf{B}f_2) \quad and \quad \mathbf{B}(f_1 \otimes f_2) \circ (g_1 \underline{\otimes} g_2) = (\mathbf{B}f_1 \circ g_1) \underline{\otimes} (\mathbf{B}f_2 \circ g_2).$$

The internal tensor product is related as one would hope with the classical:

Lemma 2.8 ([Fr19a, Lem. 4.4]). Let A_j and B_j be DGAs for $j \in \{1, 2\}$ and $g_j: \mathbf{B}A_j \longrightarrow \mathbf{B}B_j$ DGC maps. Then $\nabla \circ (g_1 \otimes g_2) = (g_1 \otimes g_2) \circ \nabla$.

3. Formal manipulation of homotopies

In this section we define the relevant notions of homotopy and discuss how to package homotopies into representing path (and path-allied) objects.

Definition 3.0.1 ([Mun74, §1.11][Mun78, §4.1]).

Given two DGA maps $f_0, f_1: A' \longrightarrow A$, a DGA *homotopy* $f_0 \simeq f_1$ is a *k*-linear map $h: A' \longrightarrow A$ of degree -1 such that

$$\varepsilon_A h = 0,$$
 $h\eta_{A'} = 0,^9$ $d(h) = f_0 - f_1,$ $h\mu_{A'} = \mu_A(f_0 \otimes h + h \otimes f_1).$

Given two DGC maps $g_0, g_1 \colon C \longrightarrow C'$, a DGC *homotopy* $g_0 \simeq g_1$ is a *k*-linear map $j \colon C \longrightarrow C'$ of degree -1 such that

$$\varepsilon_{C'}j=0, \qquad j\eta_C=0, \qquad d(j)=g_1-g_0, \qquad \Delta_{C'}j=(g_0\otimes j+j\otimes g_1)\Delta_C.$$

Given two twisting cochains $t_0, t_1: C \longrightarrow A$, a *twisting cochain homotopy* $t_0 \simeq t_1$ is a *k*-linear map $x: C \longrightarrow A$ of degree 0 such that

$$\varepsilon_A x = \varepsilon_C, \qquad x\eta_A = \eta_C, \qquad d(x) = t_0 \smile x - x \smile t_1.$$

These three notions evidently each compose well with maps in the appropriate categories.

⁹ In the definition from our main source [Mun74], the unit and counit conditions are omitted; in later work dealing more specifically with DGA as a category, Munkholm includes them [Mun78, 4.1]. These are actually critical for the adjunction to preserve homotopy and hence later to our verification of the path object.

Lemma 3.0.2. The postcomposition of a DGA map to a DGA or twisting cochain homotopy, the precomposition of a DGC map to a DGC or twisting cochain homotopy, the precomposition of a DGA map to a DGA homotopy, or the postcomposition of a DGC map to a DGC homotopy all result in another homotopy of the same type.

Moreover, the three notions are interchangeable under the adjunctions.

Lemma 3.0.3 ([Mun74, §1.11; Thm. 5.4, pf.]). Suppose given a DGC C and a DGA A. Then there are bijections of homotopies of maps

$$\begin{array}{c}
\Omega C \longrightarrow A \\
\hline
C \longrightarrow A \\
\hline
C \longrightarrow \mathbf{B}A
\end{array}$$
(3.0.4)

The adjoint functors **DGC**: $\Omega \rightarrow B$: DGA *thus also preserve the relation of homotopy*.¹⁰

Twisting cochain homotopies, despite being maps between different types of objects, are in a way more flexible than DGA or DGC homotopies, because they are composable.

Lemma 3.0.5 ([Mun74, §1.12]). Let a DGC C and a DGA A be given. A homotopy $h_{0,1}: t_0 \simeq t_1: C \longrightarrow A$ of twisting cochains admits a two-sided cup-inverse $h_{0,1}^{\sim -1}$ which is a homotopy $t_1 \simeq t_0$. Given another homotopy $h_{-1,0}: t_{-1} \simeq t_0: C \longrightarrow A$, the cup product $h_{-1,0} \smile h_{0,1}$ is a homotopy $t_{-1} \simeq t_1$.

Remark 3.o.6. A suggestive phrasing is that the twisting cochains in $Mod_0(C, A)$ are the objects of a groupoid whose morphisms are the homotopies. Particularly, homotopy is an equivalence relation. The same then holds for the equivalent hom-sets $DGA(\Omega C, A) \leftrightarrow DGC(C, \mathbf{B}A)$, which are thus privileged over generic hom-sets DGA(A', A) or DGC(C, C'), which lack this property. Note that cocompleteness is critical for the existence of inverses.

We can use Lemma 3.0.3 to exchange DGA homotopies $f_{-1} \simeq f_0 \simeq f_1$ we wish to concatenate for twisting cochains, take the cup product of these as in Lemma 3.0.5, and then move the resulting composite homotopy back to DGA using Lemma 3.0.3 to get a homotopy $f_{-1} \simeq f_1$. The next subsections attempt to describe this process internally to DGA.

3.1. Path objects

It is well known that the data of a homotopy $j: g_0 \simeq g_1: C \longrightarrow C'$ of maps of chain complexes can be realized as single map $C \otimes I \longrightarrow C'$, where I is the complex $k\{u_{[0,1]}\} \rightarrow k\{u_{[0]}, u_{[1]}\}$ of nondegenerate chains in the standard simplicial structure on the interval [0, 1]. Moreover, there is a natural coproduct endowing I with a DGC structure so that DGC homotopies can be realized in the same way. Dually [Mun74, Thm. 5.4, pf.], the algebra of normalized cochains on the simplicial interval, with the cup product, defines a DGA $I^* = k\{v_0, v_1, e\}$ such that the data of a DGA homotopy $h: f_0 \simeq f_1: A' \longrightarrow A$ can be realized by a DGA map

¹⁰ That the relation of homotopy is preserved is not to say that, for instance, if $h: f_0 \simeq f_1: A' \longrightarrow A$ is a DGA homotopy, then **B***h* is a DGC homotopy from **B***f*₀ to **B***f*₁, but that there exists a DGC homotopy. Early drafts of this document addressed this in much more detail, as some of it is never explained in the primary sources.

$$h^{P}: A' \longrightarrow I^{*} \otimes A,$$

$$a \longmapsto v_{0} \otimes f_{0}(a) - e \otimes h(a) + v_{1} \otimes f_{1}(a).$$
(3.1.1)

Explicitly, the grading on I^* is given by $|v_0| = |v_1| = 0$ and |e| = 1, the unit $\eta: k \longrightarrow I^*$ by $\eta(1) = v_0 + v_1$, the nonzero differentials by $dv_1 = e = -dv_0$, and the multiplication by

\smile	v_0	v_1	е
v_0	v_0	0	е
v_1	0	v_1	0
е	0	е	0.

It is easily seen that I^* has trivial cohomology $H^*(I^*) = H^0(I^*) \cong k$ generated by the class of $v_0 + v_1$, so by the Künneth theorem, the projections

$$\pi_i: I^* \otimes A \longrightarrow k\{v_i\} \otimes A \xrightarrow{\sim} A$$

are quasi-isomorphisms such that $\pi_j \circ h^P = f_j$. Thus tensoring with I^* functorially yields what we will call a *naive path object* for DGAS.

Unfortunately, none of the natural augmentations on $I^* \otimes A$ are such that both "endpoint" maps $I^* \otimes A \longrightarrow A$ are augmentation-preserving [Mun78, §4.3], which we need in order to apply the natural transformation ψ of Theorem 2.3 and to use the $\Omega \dashv \mathbf{B}$ adjunction. So we repair our path object by separating out $k := k\{v_0 + v_1\} \otimes \operatorname{im} \eta_A$ as the image of our unit and augmentation and defining the augmentation ideal to be $I^* \otimes \overline{A}$:

$$PA = k \oplus (I^* \otimes \overline{A})^{11}. \tag{3.1.2}$$

The inclusion into $I^* \otimes A$ is a quasi-isomorphism, and the condition $\varepsilon h = 0$ on homotopies h and unitality condition $f_j(1) = 1$ on DGA homomorphisms guarantee the map $h^P \colon A' \longrightarrow I^* \otimes A$ of (3.1.1) lands in *PA*.

Definition 3.1.3. Given a DGA *A*, we denote by *PA* the augmented DGA of (3.1.2), equipped with the projections $\pi_0, \pi_1: PA \longrightarrow A$ restricted from those of $I^* \otimes A$, and refer to it as the *standard path object* of *A*. Given a homotopy $h: f_0 \simeq f_1: A' \longrightarrow A$ of DGA maps, we refer to the associated DGA map $h^P: A' \longrightarrow PA$ of (3.1.1) as a *right homotopy* and the composites $\pi_j h^P = f_j$ as its *endpoint maps*. We write $\zeta: A \longrightarrow PA$ for unital map defined by $a \longmapsto (v_0 + v_1) \otimes a$ on \overline{A} .

In later proofs, we will encounter many right homotopies witnessed by *nonstandard* path objects, which is to say DGA quasi-isomorphisms $A' \rightarrow P'A \rightrightarrows A$ representing homotopies $h: f_0 \simeq f_1: A' \longrightarrow A$ through some other DGA P'A equipped with two surjective quasi-isomorphisms to A. Much of the material in this section is aimed at allowing us to convert these back to standard right homotopies when needed in Sections 7 and 8.

We now make an elementary observation about *PA* that looks like it should follow purely model-theoretically but seems to be a fact about the category DGA (see Remark 3.1.7).

¹¹ Munkholm's description of this substitution [Mun78, p. 229, last line] seems to suffer from a typo. What is there does not quite parse as written, and the most natural reading yields $k\{v_0 + v_1, e\} \oplus (I^* \otimes \overline{A})$, which is not quasi-isomorphic to A because e generates a new k summand in H^1 now that v_0 and v_1 have been removed.

Proposition 3.1.4. Given a DGA X, the standard path object PX is right-homotopy–equivalent to X with respect to the notion of right homotopy determined by PX itself. In particular, for any other DGA A, the mappings ζ and π_i induce bijections $[A, X] \longleftrightarrow [A, PX]$ of right-homotopy classes of DGA maps.

Proof. It would be enough to find a right homotopy between $\zeta \circ \pi_j$ and id_{PX} , but it is psychologically more convenient to maintain A. Since $\pi_j \circ \zeta = id_X$, the map $\zeta_* \colon [A, X] \longrightarrow [A, PX]$ is an injection, so it is enough to see it is surjective too. For this, we note an arbitrary map $A \longrightarrow PX$ is a right homotopy h^P between maps $f = \pi_0 \circ h^P \colon A \longrightarrow X$ and $g = \pi_1 \circ h^P$, and show h^P itself is right homotopic to the map $\zeta \circ f$ representing the constant homotopy $f \simeq f$. The intuition for why this should be is given by the square



where we think of the left edge as $\zeta \circ f$ and the right edge as h^p , the labels 0 and *h* on the edges representing the homotopies, which is to say the degree-(-1) maps $A \longrightarrow X$ which are the "*e*-components" of the right homotopies $\zeta \circ f$ and h^p . Explicitly, the degree-(-1) map $A \longrightarrow PX$ given by $a \longmapsto v_1 \otimes h(a)$ can be checked to be a DGA homotopy $\zeta \circ f \simeq h^p$ in the sense of Definition 3.0.1.

Corollary 3.1.5. Given a DGC C and a DGA X, any two right homotopies $\Omega C \longrightarrow PX$ both representing homotopies $f \simeq g$ of DGA maps $\Omega C \longrightarrow X$ are themselves homotopic as DGA maps.

Proof. Note that the two right homotopies are both homotopic to $\zeta \circ f$ by Proposition 3.1.4 and recall from Remark 3.0.6 that homotopy is an equivalence relation on DGA(ΩC , X).

Remark 3.1.6. There is a standard cofibrantly generated model structure on DGA, with weak equivalences quasi-isomorphisms and surjections fibrations, due when *k* is a field to later work of Munkholm [Mun78], and more generally to Jardine [Jar97]. The counit quasi-isomorphism $\varepsilon: \Omega BA \longrightarrow A$ is only a cofibrant replacement if *k* is a field, essentially because only projective modules will lift against surjections and ΩBA is a projective *k*-module if and only if *A* is. We will nevertheless be able to use ΩB much in the manner of a functorial cofibrant replacement.¹²

Remark 3.1.7. The standard model-categorical notion of right homotopy uses generic path objects, not specifically the standard path object of Definition 3.1.3 witnessing the classical notion of DGA homotopy which as we have pointed out in Remark 3.0.6 is not typically an equivalence relation on DGA(A, X) unless $A = \Omega C$ for some DGC C. The natural witness $PA \times_A PA$ for a composite of homotopies is not again PA, but *is* another path object, and model-theoretic right homotopy on DGA is the transitive closure of classical DGA homotopy. Because of the functoriality and

¹² The author at one point hoped to make real use of this model structure, but inexplicitly summoning liftings does not seem to give enough control over composition; one wants something as near functorial as possible to describe composition of DGA homotopies. Another thought was to use DGCs instead, but when k is not a field, the standard proof for the model structure breaks down and for some time the author was under the impression that for k an arbitrary ring, there is no model structure (in fact, there is [HeKRS17, Cor. 6.3.5]).

accessibility of the path objects and the cobar–bar adjunction, and to maintain back-compatibility with the classics, we prefer to keep our discussion in terms of the classical notion.

Proposition 3.1.4 looks, if we took RX = PX, a bit like the standard lemma that the righthomotopy class of a(ny choice of) fibrant replacement $RA \longrightarrow RX$ of a map $A \longrightarrow X$ is determined by the composition $A \rightarrow X \rightarrow RX$. But since all objects are fibrant, this statement is distinct and actually trivial, and $\zeta: X \longrightarrow PX$ is very rarely a cofibration anyway. The homotopy category of DGA is determined by right homotopy after cofibrant replacement of domains, $QA \longrightarrow RX = X$, so right-homotopy classes in DGA(ΩBA , X) give the correct homotopy notion for k a field, but in general the relation is not clear.

3.2. Double- and triple-path objects and concatenation

We have seen we can compose homotopies with compatible endpoint maps in DGA(ΩC , A), for A a DGA and C a DGC, but our procedure passes through twisting cochains $C \longrightarrow A$ and gives no explicit description of the result in terms internal to DGA. In this subsection, whose content we could find no other reference for, we rectify (or more honestly, circumnavigate) this shortcoming.

Definition 3.2.1. Given a DGA *A*, we write

$$DA := PA \times PA$$

for the pullback of the diagram $PA \xrightarrow{\pi_1} A \xleftarrow{\pi_0} PA$ and refer to it as the *double-path object*. By definition it comes equipped with two projections to *PA* and three maps

$$p_0 \coloneqq \pi_0 \times 0, \qquad \qquad \pi_1 \times 0 = 0 \times \pi_0, \qquad \qquad p_1 \coloneqq 0 \times \pi_1$$

to *A*, all quasi-isomorphisms. As *PA* is a subalgebra of $I^* \otimes A$, we may apply distributivity of \otimes over $\oplus = \times$ to identify *DA* with a subalgebra of $(I^* \times I^*) \otimes A$, and then, in terms of the naive, unaugmented double path-object $I^* \times_k I^* < I^* \times I^*$ of *k*, we have a decomposition

$$DA \cong k\{(v_0, 0) + (v_1, v_0) + (0, v_1)\} \oplus (I^* \times I^*) \otimes \overline{A}$$

identifying *DA* as an augmented DGA.

The *raison d'être* of the double-path object, of course, is to represent pairs of composable homotopies $f_{-1} \simeq f_0 \simeq f_1 \colon A' \longrightarrow A$ of DGA maps, which it achieves tautologically since a pair $h_{-1,0} \colon f_{-1} \simeq f_0$ and $h_{0,1} \colon f_0 \simeq f_1$ of homotopies induces representatives $h_{-1,0}^p \colon A' \longrightarrow PA$ such that $\pi_1 h_{-1,0}^p = f_0 = \pi_0 h_{0,1}^p \colon A' \longrightarrow A$, and thus the map $(h_{-1,0}^p, h_{0,1}^p) \colon A' \longrightarrow PA \times PA$ factors through the fiber product. Evidently $p_0(h_{-1,0}^p, h_{0,1}^p) = f_{-1}$ and $p_1(h_{-1,0}^p, h_{0,1}^p) = f_1$.

If the desired composition of homotopies were realized by a map $DA \longrightarrow PA$, then the concatenation of any pair of compatible homotopies $A' \longrightarrow A$, would be represented by the composite of the associated map $A' \longrightarrow DA$ and the unattested $DA \longrightarrow PA$, but we know this is only possible when A' is the cobar construction ΩC on some DGC C. The composition of homotopies is, nevertheless, a natural transformation $\underline{\Upsilon}$: DGA $(\Omega(-), DA) \longrightarrow$ DGA $(\Omega(-), PA)$.¹³ In particular, plugging **B**DA in as the variable, $\underline{\Upsilon}$ takes the counit ε : Ω **B**DA \longrightarrow DA to a map $\underline{\Upsilon} := \underline{\Upsilon}(\varepsilon)$: Ω **B**DA \longrightarrow PA, and a Yoneda-style argument yields the following.

¹³ The intended visual mnemonic is that Υ takes two things and combines them into one.

Lemma 3.2.2. Let a DGC C and DGA A and homotopies $f_{-1} \simeq f_0 \simeq f_1$ of DGA maps $\Omega C \longrightarrow A$ be given. If the DGA map $h^D : \Omega C \longrightarrow DA$ represents this pair of homotopies, then the composite homotopy $f_{-1} \simeq f_1$ is represented by

$$\Omega C \xrightarrow{\Omega \eta} \Omega B \Omega C \xrightarrow{\Omega B h^D} \Omega B D A \xrightarrow{\Upsilon} P A, \qquad (3.2.3)$$

where the map Υ implementing the concatenation is a quasi-isomorphism.

We note that $\Omega C \longrightarrow \Omega B D A$ is the map induced up from the original $\Omega C \longrightarrow D A$ as in Corollary 1.3, so composition of homotopies in DGA is implemented by a diagram of the shape

$$\Omega BDA \xrightarrow{\Upsilon} PA.$$

$$(3.2.4)$$

$$\Omega C \xrightarrow{h^{D}} DA$$

Proof. Since $h^D = \omega^* \varepsilon$ in (3.2.4), by naturality of $\underline{\Upsilon}$ we have $\underline{\Upsilon}(h^D) = \underline{\Upsilon}(\omega^* \varepsilon) = \omega^* \underline{\Upsilon}(\varepsilon) = \omega^* \Upsilon$.

To see Υ is a quasi-isomorphism, fix $C = \mathbf{B}A$ and let $h^D : \Omega \mathbf{B}A \longrightarrow DA$ represent a pair of constant (= trivial) homotopies of $f_{-1} = f_0 = f_1 := \varepsilon : \Omega \mathbf{B}A \longrightarrow A$, so that explicitly $h^D : x \longmapsto ((v_0, 0) + (v_1, v_0) + (0, v_1)) \otimes \varepsilon(x)$. Then h^D is a quasi-isomorphism by 2-of-3, as $p_0h^D = \varepsilon$ and p_0 itself both are quasi-isomorphisms. If we follow the composition in (3.2.3) with the quasi-isomorphism ε , so by 2-of-3 again, the composition in (3.2.3) is a quasi-isomorphism. But $\Omega\eta$ is always a quasi-isomorphism, and $\Omega \mathbf{B}h^D$ is a quasi-isomorphism since h^D is, so by 2-of-3 yet again, so too must be Υ .

The same trick works equally for composable triples of homotopies.

Definition 3.2.5. Given a DGA *A*, its *triple-path object* is the pullback

$$TA := PA \underset{A}{\times} PA \underset{A}{\times} PA,$$

equipped with the expected three projections $TA \longrightarrow PA$ and four projections $TA \longrightarrow A$.

The same proof as for *DA* yields the following.

Lemma 3.2.6. Let a DGC C and DGA A and homotopies $f_0 \simeq f_1 \simeq f_2 \simeq f_3$ of DGA maps $\Omega C \longrightarrow A$ be given. There is a natural map III: $\Omega BTA \longrightarrow PA$ such that if the DGA map $h^T \colon \Omega C \longrightarrow TA$ represents this triple of homotopies, then the composite homotopy $f_0 \simeq f_3$ is represented by

$$\mathbf{\Omega}C \xrightarrow{\mathbf{\Omega}\eta} \mathbf{\Omega}\mathbf{B}\mathbf{\Omega}C \xrightarrow{\mathbf{\Omega}\mathbf{B}h^T} \mathbf{\Omega}\mathbf{B}TA \xrightarrow{\mathrm{III}} PA.$$

The map III *representing composition is a quasi-isomorphism.*

In one instance we will encounter, the composable pair we apply Lemma 3.2.2 to comprises two adjacent sides of a square of homotopies $\Omega C^* \longrightarrow (I^*)^{\otimes 2} \otimes A^{\otimes 2}$.

Lemma 3.2.7. Let A be a DGA. Then there is a natural DGA map $r: PA \otimes PA \longrightarrow D(A \otimes A)$ such that $p_0r = \pi_0 \otimes \pi_0$ and $p_1r = \pi_1 \otimes \pi_1$.

Proof. First note there exists a natural quotient map

$$I^* \otimes I^* \longrightarrow k\{v_0 \otimes v_0, e \otimes v_0, v_1 \otimes v_0, v_1 \otimes e, v_1 \otimes v_1\}$$

modding out from the naive square object the ideal spanned by $v_0 \otimes e$, $v_0 \otimes v_1$, $e \otimes e$, and $e \otimes v_1$. It is not hard to see this quotient is isomorphic to the naive double-path object $I^* \times_k I^*$ under the assignment

$$v_0 \otimes v_0 \longmapsto (v_0, 0), \quad e \otimes v_0 \longmapsto (e, 0), \quad v_1 \otimes v_0 \longmapsto (v_1, v_0), \quad v_1 \otimes e \longmapsto (0, e), \quad v_1 \otimes v_1 \longmapsto (0, v_1), \quad v_1 \otimes v_1 \mapsto (0, v_1) \mapsto (0, v_$$

Write $r^I: I^* \otimes I^* \longrightarrow I^* \times_k I^*$ for the composition and note it sends the idempotent $(v_0 + v_1)^{\otimes 2}$ to the unity of $I^* \times_k I^*$.

Recalling from Definition 3.1.3 that *PA* is a subalgebra of the tensor product $I^* \otimes A$, we may permute tensor-factors to identify $PA \otimes PA$ with a subalgebra of $(I^*)^{\otimes 2} \otimes A^{\otimes 2}$. The unity of $PA \otimes PA$ is sent to $(v_0 + v_1)^{\otimes 2} \otimes 1^{\otimes 2}$ under this identification, and the augmentation ideal $\overline{PA \otimes PA}$ into $(I^*)^{\otimes 2} \otimes \overline{A^{\otimes 2}}$. Thus $r^I \otimes \operatorname{id}_A^{\otimes 2}$ sends the unity of $PA \otimes PA$ to the unity of DA and $\overline{PA \otimes PA}$ into $(I^* \times_k I^*) \otimes \overline{A \otimes A} = \overline{D(A \otimes A)}$, so we may define the intended r by corestriction.

The last formal trick we will perform with homotopies is to move *P* past ΩB . For this, starting with a right homotopy $h^P \colon \Omega C \longrightarrow PA$ representing a homotopy $h \colon f_0 \simeq f_1 \colon \Omega C \longrightarrow A$, note the string of conversions

$$h^{p}: \Omega C \longrightarrow PA$$

$$h: \Omega C \longrightarrow A$$

$$\tilde{h}: \Omega C \longrightarrow \Omega BA$$

$$\tilde{h}^{p}: \Omega C \longrightarrow P\Omega BA$$

afforded by Lemma 3.0.3, Corollary 1.3, and Definition 3.1.3, amounting to a natural transformation

$$\underline{Z}: \mathbf{DGA}(\mathbf{\Omega}(-), PA) \longrightarrow \mathbf{DGA}(\mathbf{\Omega}(-), P\mathbf{\Omega}BA).$$

Following through the construction, the induced \tilde{h} is a homotopy $f_0^{\#} \simeq f_1^{\#}$, so that $\pi_j \circ \tilde{h}^p = f_j^{\#}$ for $j \in \{0, 1\}$.

As with $\underline{\Upsilon}$, a Yoneda-esque argument shows the natural transformation \underline{Z} is represented by one map. Plugging **B***PA* into the hole and applying \underline{Z} to the counit ε : Ω **B***PA* \longrightarrow *PA* yields a DGA map $Z := \underline{Z}(\varepsilon)$: Ω **B***PA* \longrightarrow *P* Ω **B***A*.

Lemma 3.2.8. Given a DGA A, there is a natural DGA quasi-isomorphism $Z: \Omega BPA \longrightarrow P\Omega BA$ such that $\pi_j \circ Z = \Omega B \pi_j: \Omega BPA \longrightarrow \Omega BA$.

Proof. The counit ε : $\Omega BPA \longrightarrow PA$ itself represents a homotopy h between the two composites $\pi_j \circ \varepsilon$: $\Omega BPA \longrightarrow A$ for $j \in \{0,1\}$, and the induced \tilde{h} is a homotopy between the maps $(\pi_j \circ \varepsilon)^{\#}$: $\Omega BPA \longrightarrow \Omega BA$. By the naturality property Corollary 1.3 of induction, $(\pi_j \circ \varepsilon)^{\#} = \Omega B\pi_j \circ \varepsilon^{\#}$ and $\varepsilon^{\#} = \operatorname{id}_{\Omega BA}$, so \tilde{h} is a homotopy $\Omega B\pi_0 \simeq \Omega B\pi_1$, and thus by definition $Z = \tilde{h}^P$ satisfies $\pi_j \circ \tilde{h}^P = \Omega B\pi_j$ for $j \in \{0,1\}$. It follows Z is a quasi-isomorphism, for π_j and $\Omega B\pi_j$ both are.

Remark 3.2.9. Though we will not use these facts, we should mention that the transformations $\underline{\Upsilon}$, $\underline{\Pi}$, *r*, and \underline{Z} (from Lemmas 3.2.2 and 3.2.6 to 3.2.8 respectively) are also natural in the variable *A* and *r* is a quasi-isomorphism.

4. Maps on Tor

One of the goals of this work is to determine the cohomology ring of the homotopy pullback of a span $X \rightarrow B \leftarrow E$ of spaces from the cohomology of the input spaces. The tool of choice here is the EMSS, which is a special instance of a so-called *algebraic* EMSS, converging [Mac, XI.3.2] to differential Tor of a triple of DGAS from classical Tor of their cohomology and functorial in all three variables.

Lemma 4.1 ([GuM, Cor. 1.8][Mun74, Theorem 5.4]). Given a DGA map $f : R' \longrightarrow R$, a right R'-module M', a left R'-module N, a right R-module M, a left R-module N, and DG module maps $u : M' \longrightarrow M$ and $v : N' \longrightarrow N$ making the expected squares

commute, there is induced a map of algebraic EMSSS from that of (M', R', N') to that of (M, R, N), converging to the functorial map

$$\operatorname{Tor}_{f}(u, v) \colon \operatorname{Tor}_{R'}(M', N') \longrightarrow \operatorname{Tor}_{R}(M, N)$$

of graded modules. Moreover, if the maps f, u, v are quasi-isomorphisms, then the map of spectral sequences is an isomorphism from the E_2 page on and $\text{Tor}_f(u, v)$ is an isomorphism.

We will only apply these considerations in the special case that M and N are DGAs and the R-module structure maps are induced by DGA homomorphisms $M \leftarrow R \rightarrow N$, so that we have the condensed compatibility diagram

of DGA maps. In later sections, where we will produce diagrams of Tors comprising mainly isomorphisms, we will ceaselessly apply this result. We will also need to expand the notion of a map of Tors to include squares which commute only up to homotopy.

Lemma 4.4 ([Mun74, Thm. 5.4]). Let DGAs and DGA maps as in (4.3) be given such that the squares commute up to homotopies $h_M: u \circ \phi_{M'} \simeq \phi_M \circ f$ and $h_N: v \circ \phi_{N'} \simeq \phi_N \circ f$. Then there is induced a map

$$\operatorname{Tor}_f(u, v; h_M, h_N) \colon \operatorname{Tor}_{R'}(M', N') \longrightarrow \operatorname{Tor}_R(M, N)$$

of graded modules which is a quasi-isomorphism if each of u, f, and v is.

Proof. Letting h_M^p : $R \longrightarrow PM'$ and h_N^p : $R \longrightarrow PN'$ be the respective DGA representatives, as described in Definition 3.1.3, for the homotopies h_M , h_N , the following equivalent diagrams commute by definition:



Since the π_i are quasi-isomorphisms, three applications of Lemma 4.1 let us set

$$\operatorname{Tor}_{f}(u, v; h_{M}, h_{N}) \coloneqq \operatorname{Tor}_{f}(\pi_{1}, \pi_{1}) \circ \operatorname{Tor}_{\mathrm{id}}(\pi_{0}, \pi_{0})^{-1} \circ \operatorname{Tor}_{\mathrm{id}}(u, v).$$

This result accounts for our fixation on representing homotopies by DGA maps in Section 3. It seems to be a new observation that the choice of homotopies doesn't matter.

Proposition 4.6. The map $\operatorname{Tor}_{f}(u, v) := \operatorname{Tor}_{f}(u, v; h_{M}, h_{N})$ is independent of the homotopies h_{M}, h_{N} .

Proof. First assume $R' = \Omega C$ is the cobar construction of some DGC *C*. By Corollary 3.1.5, then the right homotopy $h_M^p: R' \longrightarrow PM$ and any other right homotopy $\tilde{h}_M^p: R' \longrightarrow PM$ representing some other homotopy $\tilde{h}_M: u \circ \phi_{M'} \simeq \phi_M \circ f$ are homotopic, and hence there is some right homotopy $H_M^p: A \longrightarrow PPM$ with $\pi_0 \circ H_M^p = h_M^p$ and $\pi_1 \circ H_M^p = \tilde{h}_M^p$. Similarly, given a second homotopy $\tilde{h}_N: v \circ \phi_{N'} \simeq \phi_N \circ f$, we have a homotopy $h_N^p \simeq \tilde{h}_N^p$ and witnessing right homotopy $H_N^p: A \longrightarrow PPN$. By construction, one has $\pi_0 \circ \pi_0 \circ H_M^p = \pi_0 \circ \pi_1 \circ H_M^p$ and $\pi_1 \circ \pi_0 \circ H_M^p = \pi_1 \circ \pi_1 \circ H_M^p$, and similarly for H_N^p . This gives us the commutative diagram of maps below, where the Tor in the central object is defined by H_M^p .

$$\operatorname{Tor}_{R'}(PM, PN)$$

$$\operatorname{Tor}_{id}(\pi_{0}, \pi_{0}) \xrightarrow{\operatorname{Tor}_{id}(\pi_{0}, \pi_{0})} \operatorname{Tor}_{R'}(M, N) \xrightarrow{\operatorname{Tor}_{id}(\pi_{0}, \pi_{0})} \operatorname{Tor}_{R'}(PPM, PPN) \xrightarrow{\operatorname{Tor}_{id}(\pi_{1}, \pi_{1})} \operatorname{Tor}_{R'}(M, N) \xrightarrow{\operatorname{Tor}_{R'}(M, N)} \xrightarrow{T$$

The horizontal composite along the top, $\neg \nearrow \neg$, is $\operatorname{Tor}_f(u, v; h_M, h_N)$ while the bottom composite $\neg \checkmark \nearrow$ is $\operatorname{Tor}_f(u, v; \tilde{h}_M, \tilde{h}_N)$. Both can be factored through $\operatorname{Tor}_{R'}(PPM, PPN)$ using the vertical isomorphisms, the top as $\neg \checkmark \uparrow \uparrow \checkmark \rightarrow$, the bottom as $\neg \land \uparrow \downarrow \nearrow \rightarrow$, but by commutativity of the diagram, one has $\land \downarrow = \checkmark \uparrow$ and $\uparrow \checkmark = \downarrow \nearrow$, so the composites are equal.

In the general case, we have the diagram (4.7) but with the central $\text{Tor}_{R'}(PPM, PPN)$ omitted, and we do not know *a priori* the top and bottom composites are the same. If we precompose all maps from R' in the discussion above with the counit ε : $\Omega B R' \longrightarrow R'$, then the preceding

discussion with $C = \mathbf{B}R'$ does apply, and we obtain the commutative diagram (4.7) but with R' everywhere replaced by $\mathbf{\Omega}\mathbf{B}R'$. In that diagram, we do know the top and bottom composites are equal by the preceding paragraph. Now, deleting the central $\operatorname{Tor}_{\mathbf{\Omega}\mathbf{B}R'}(PPM, PPN)$ from that diagram, the rest admits the map $\operatorname{Tor}_{\varepsilon}(\operatorname{id}, \operatorname{id})$ down to the first diagram, so that by the naturality of ε , all new squares commute. As ε is a quasi-isomorphism, we know $\operatorname{Tor}_{\varepsilon}(\operatorname{id}, \operatorname{id})$ is a natural isomorphism by Lemma 4.1, and thus equality of the two composites in the R' diagram follows from the equality of the corresponding composites in the $\mathbf{\Omega}\mathbf{B}R'$ diagram.

Not only do these maps not depend on the homotopies involved, they compose functorially. To make the proof more legible, we introduce a convention that saves on the mental space involved in repeating triples of operations. We will make increasing use of this convention as our diagrams grow unwieldy, eventually arriving at a point where the diagrams only fit on a page through its use. We beg the reader's patience and trust in adapting to this abbreviation: the mental effort of expanding things out does seem greater in the beginning, but as one learns to incorporate the parallelism, abbreviation greatly helps one hold the objects in mind simultaneously, and by the time one can understand the proofs, will make them much easier to process.

Notation 4.8. Given DGA maps $X \leftarrow A \rightarrow Y$, functors F, G, F', G': DGA \longrightarrow DGA, and natural transformations $F \longrightarrow G$, $F' \longrightarrow G'$, $\phi: F \longrightarrow F'$, and $\chi: G \longrightarrow G'$ such that the two composites $F \longrightarrow G'$ are equal, we make the abbreviations

$$\operatorname{Tor}_{FA} := \operatorname{Tor}_{FA}(FX, FY), \qquad \operatorname{Tor}_{FA}(GX) := \operatorname{Tor}_{FA}(GX, GY),$$
$$\operatorname{Tor}_{\phi} := \operatorname{Tor}_{\phi}(\phi, \phi) \colon \operatorname{Tor}_{FA} \longrightarrow \operatorname{Tor}_{F'A}, \qquad \operatorname{Tor}_{\phi}(\chi) \coloneqq \operatorname{Tor}_{\phi}(\chi, \chi) \colon \operatorname{Tor}_{FA}(GX) \longrightarrow \operatorname{Tor}_{F'A}(G'X)$$

The following, apparently original,¹⁴ shows the functoriality of Tor with respect to this extended mapping notion.

Theorem 4.9. Assume given a diagram of DGA maps



in which there is a DGA *homotopy making each square commute. Then* $\operatorname{Tor}_{ff'}(uu', vv')$ *is well defined and equals* $\operatorname{Tor}_{f}(u, v) \circ \operatorname{Tor}_{f'}(u', v')$.

Proof. If R'' is a cobar construction, we define $\text{Tor}_{ff'}(uu', vv')$ using the composite DGA homotopies guaranteed by Lemma 3.0.5 making the vertical rectangles R''M''MR and R''N''NR commute. We have seen in Proposition 4.6 that this choice of homotopies does not matter so long as any exist in the first place.

¹⁴ but *cf*. work of Gugenheim–Munkholm [GuM74] which achieves something similar for a version of Tor defined as the cohomology of the two-sided bar construction

If R'' is not a cobar construction, such composite homotopies do not necessarily exist. However, since Tor is functorial in its arguments and $\varepsilon \colon \Omega B R'' \longrightarrow R''$ is a quasi-isomorphism, the induced map $\operatorname{Tor}_{\varepsilon}(\operatorname{id}, \operatorname{id}) \colon \operatorname{Tor}_{\Omega B R''}(-, -) \longrightarrow \operatorname{Tor}_{R''}(-, -)$ is a natural isomorphism, and conjugating by these isomorphisms, we may define $\operatorname{Tor}_{ff'}(uu', vv')$ as $\operatorname{Tor}_{\varepsilon}(\operatorname{id}, \operatorname{id}) \circ \operatorname{Tor}_{\Omega B f \Omega B f'}(uu', vv') \circ$ $\operatorname{Tor}_{\varepsilon}(\operatorname{id}, \operatorname{id})^{-1}$. By naturality of the isomorphism $\operatorname{Tor}_{\varepsilon}(\operatorname{id}, \operatorname{id})$, the composite $\operatorname{Tor}_{f}(u, v) \circ \operatorname{Tor}_{f'}(u', v')$ can be written equally well as

$$\left[\operatorname{Tor}_{\varepsilon}(\operatorname{id},\operatorname{id})\circ\operatorname{Tor}_{\mathbf{\Omega}\mathbf{B}f}(u,v)\circ\operatorname{Tor}_{\varepsilon}(\operatorname{id},\operatorname{id})^{-1}\right]\circ\left[\operatorname{Tor}_{\varepsilon}(\operatorname{id},\operatorname{id})\circ\operatorname{Tor}_{\mathbf{\Omega}\mathbf{B}f'}(u',v')\circ\operatorname{Tor}_{\varepsilon}(\operatorname{id},\operatorname{id})^{-1}\right],$$

so it is enough to compare $\operatorname{Tor}_{\Omega B_f \Omega B_{f'}}(uu', vv')$ with $\operatorname{Tor}_{\Omega B_f}(u, v) \circ \operatorname{Tor}_{\Omega B_{f'}}(u, v)$, and thus we may assume $R'' = \Omega C$ for some DGC C to begin with.

Now, omitting the *N* arguments, taking for granted that the diagrams are symmetric in *M*'s and *N*'s, and doing a reflection through y = x, the four-square diagram $\hat{a} \, la$ (4.5) giving Tor_{*ff*}(uu', vv') can be rewritten as

while the diagram yielding $\text{Tor}_{f}(u, v) \circ \text{Tor}_{f'}(u', v')$ is

By Lemma 3.0.5, we can concatenate the right homotopies $R'' \to PM' \to PM$ and $R'' \to R' \to PM$ implied by (4.11), and by Lemma 3.2.2 this composite homotopy can be implemented via a map with codomain ΩBDM followed by Υ : $\Omega BDM \longrightarrow PM$. We can combine these ingredients into Figure 4.12, abusively labelling maps on Tor by the DGA maps inducing them. Here the maps induced by (4.10), along the left and bottom, are in red and those induced by (4.11), along the top, are in blue; the first and last arrows, labeled u' and f respectively, are violet because they are both. Using the commutativity of the various squares and triangles, one sees $\operatorname{Tor}_{ff'}(uu', vv')$ and $\operatorname{Tor}_f(u, v) \circ \operatorname{Tor}_{f'}(u', v')$ are indeed equal.

5. SHC-algebras

A commutative DGA *A* is one for which the multiplication $\mu: A \otimes A \longrightarrow A$ is itself a DGA homomorphism. Cohomology rings are of this sort, and a large part of why homotopy theory is so much more tractable over a field *k* of characteristic 0 is that there are functorial CDGA models for cochains. For other characteristics this is not the case [Bor51, Thm. 7.1], but we can weaken the requirement by asking only that μ extend to an A_{∞} -algebra map. Munkholm's product is defined in terms of such a structure, as first considered by Stasheff and Halperin.



Figure 4.12: The diagram for functoriality of Tor

To make sense of the following definition, recall from Section 1 that the canonical twisting cochain $t^A: \mathbf{B}A \longrightarrow A$ of a DGA A factors through the projection onto a DG direct summand $s^{-1}\overline{A}$, on which it restricts to a cochain isomorphism of degree 1, whose inverse we dub $s_A^{-1}: \overline{A} \longrightarrow \mathbf{B}A$ and call the *desuspension*.

Definition 5.1 (Stasheff-Halperin [StH70, Def. 8]). We refer to a DGA *A* equipped with a DGC map $\Phi_A: \mathbf{B}(A \otimes A) \longrightarrow \mathbf{B}A$ such that the composition $t_A \circ \Phi \circ s_{A \otimes A}^{-1}: \overline{A \otimes A} \longrightarrow \overline{A}$ is the multiplication $\mu_A: A \otimes A \longrightarrow A$ as a *weakly homotopy commutative* (whc-)*algebra*.¹⁵ Given two whc-algebras *A* and *Z*, a whc-*algebra map* from *A* to *Z* is a DGC map $g: \mathbf{B}A \longrightarrow \mathbf{B}Z$ such that there exists a DGC homotopy between the two paths around the square

The particular homotopy is not prescribed as part of the data of a whc-algebra map.

WHC-algebra structures thus enable us to upgrade the non-DGA map $\mu: A \otimes A \longrightarrow A$ to a legitimate DGA map $\Omega \Phi: \Omega B(A \otimes A) \longrightarrow \Omega BA$, which is more tractable categorically even if less intuitive on the point-set level, and which by the naturality of $\varepsilon: \Omega B \longrightarrow$ id carries the same information as μ up to quasi-isomorphism.

Munkholm's definition adds to Stasheff-Halperin's a weakening of the standard CGA axioms.

Definition 5.3. [Mun74, Def. 4.1] A *strongly homotopy commutative* (henceforth SHC-) *algebra* is a WHC-algebra *A* whose structure map Φ_A : **B**($A \otimes A$) \longrightarrow **B**A, satisfies the following conditions:

¹⁵ Stasheff–Halperin call this a *strongly homotopy commutative algebra* structure, but we will meet a stronger notion momentarily, so we rename their notion "weak."

- **1**. It is strictly unital: $\Phi \circ \mathbf{B}(\mathrm{id}_A \otimes \eta_A) = \mathrm{id}_{\mathbf{B}A} = \Phi \circ \mathbf{B}(\eta_A \otimes \mathrm{id}_A).$
- 2. It is homotopy-commutative: there is a DGC homotopy from Φ to $\Phi \circ \mathbf{B}\chi$: $\mathbf{B}(A \otimes A) \longrightarrow \mathbf{B}A$, where $\chi: A \otimes A \xrightarrow{\sim} A \otimes A$ is the factor transposition $a \otimes b \longmapsto (-1)^{|a||b|} b \otimes a$.
- 3. It is homotopy-associative: there is a DGC homotopy between the maps $\Phi(\Phi \otimes id_{BA})$ and $\Phi(id_{BA} \otimes \Phi)$: $\mathbf{B}(A \otimes A \otimes A) \longrightarrow \mathbf{B}A$.

The associativity and commutativity homotopies postulated are again not themselves specified in the data of an shc-algebra, only the fact of their existence. An shc-algebra map is a whc-algebra map between shc-algebras.

The canonical example is that of an authentically commutative algebra.

Example 5.4. If *A* is a CDGA, then the morphism $\Phi = \mathbf{B}\mu_A : \mathbf{B}(A \otimes A) \longrightarrow \mathbf{B}A$ makes *A* an sHC-algebra. The cohomology ring $H^*(X;k)$ of a simplicial set is of this type, and will always come considered with this sHC-algebra structure. If $\rho : A \longrightarrow B$ is a map of CDGAS, then $\mathbf{B}\rho$ is an sHC-algebra map.

Example 5.5. If *A* is any DGA, there is a unique DG-module section $i: A \longrightarrow \Omega BA$ of $\varepsilon: \Omega BA \longrightarrow A$ that is unital and restricts to $t_{BA} \circ s^{-1}$ on \overline{A} [Mun74, Prop. 2.14]. If *A* is an SHC-algebra, then the homotopy-commutativity of Φ implies μ and $\mu \circ \chi$ are homotopic cochain maps, and $\varepsilon \circ \Omega(\Phi \circ B\chi) \circ i = \mu \circ \chi$ and $\varepsilon \circ \Omega \Phi \circ i = \mu$, and so $H^*(A)$ is a CDGA (with trivial differential). Thus $\Phi = B\mu_{H^*(A)}$ gives an SHC-algebra structure on $H^*(A)$ by Example 5.4. The cohomology ring of an SHC-algebra will always be endowed with this SHC-algebra structure.

Theorem 5.6 ([Mun74, Prop. 4.7]). Let X be a simplicial set and k any ring. Then the normalized cochain algebra $C^*(X) = C^*(X;k)$, augmented by restriction to $C^*(*;k) \cong k$ for some basepoint $* \in X_0$, admits an SHC-algebra structure $\Phi_{C^*(X)}$, and this structure is strictly natural in the sense that given a basepoint-preserving map $f: Y \longrightarrow X$ of simplicial sets, the induced DGC map $BC^*(f): BC^*(X) \longrightarrow BC^*(Y)$ renders the square (5.2) commutative on the nose.

This natural shc-algebra structure on cochains is a reinterpretation of the classical Eilenberg–Zilber theorem, and only verifying the homotopy-associativity axiom requires substantial work.

The class of known shc-algebras has recently expanded significantly:

Theorem 5.7 (Franz [Fr19c]). A homotopy Gerstenhaber algebra A admits a strictly natural wHC-algebra structure Φ_A satifying the axioms Definition 5.3.1 and Definition 5.3.3. If A is an extended homotopy Gerstenhaber algebra, then Φ_A is in fact an SHC-algebra structure.

Remark 5.8. A homotopy Gerstenhaber algebra is an algebra over a certain E_2 -operad $F_2 \mathscr{X}$ and similarly, an extended homotopy Gerstenhaber algebra is an algebra over a certain suboperad of an E_3 -operad $F_3 \mathscr{X}$, accounting for the phrasing we employ in the abstract.¹⁶ Since E_2 is not very far along the road to E_{∞} , morally speaking we require *some* commutativity to obtain the product on Tor, but not very much.

On the other hand, since the readiest source of homotopy Gerstenhaber algebras is algebras over this particular operad, rather than just any E_2 -operad, an E_2 -algebra is not necessarily a wHC-algebra, so the notions are not strictly comparable.

¹⁶ Both of these are filtrands of the so-called *surjection operad* \mathscr{X} of interval-cut operations on cochains, which is a quotient of the pg-operad \mathscr{E} associated to the classical Barratt–Eccles simplicial operad [McSo₃, BeFo₄].

6. The product

Munkholm's product can be motivated as a sort of least common generalization of the classical products on $\text{Tor}_{C^*B}(C^*X, C^*E)$ and $\text{Tor}_{H^*B}(H^*X, H^*E)$, rephrased in terms of the canonical shc-algebra structures. We choose not to use his definition of the product, but an equivalent definition of our own.¹⁷

Given DGAS R_0 , R_1 and right and left DG R_i -modules M_i and N_i respectively, there is a classically defined exterior product [CarE, p. 206]

$$\operatorname{Tor}_{R_0}(M_0, N_0) \otimes \operatorname{Tor}_{R_1}(M_1, N_1) \longrightarrow \operatorname{Tor}_{R_0 \otimes R_1}(M_0 \otimes M_1, N_0 \otimes N_1)$$

functorial in all six variables in the sense that given similarly defined $R'_{i'} M'_{i'} N'_{i}$ such that the squares (4.2) commute, so does the square

and given further R_i'', M_i'', N_i'' , such squares glue. If $R = R_0 = R_1$ is a *commutative* DGA, then $\mu: R' = R \otimes R \longrightarrow R$ is a DGA map, and if $M = M_0 = M_1$ and $N = N_0 = N_1$ are themselves DGAs, then $\mu: M' = M \otimes M \longrightarrow M$ and $\mu: N' = N \otimes N \longrightarrow N$ make the diagrams (4.2) commute, so we may follow the external product with the map

$$\operatorname{Tor}_{\mu}(\mu,\mu)$$
: $\operatorname{Tor}_{R\otimes R}(M\otimes M,N\otimes N) \longrightarrow \operatorname{Tor}_{R}(M,N)$

to obtain the classical product on Tor. This particularly applies to $R = H^*A$, $M = H^*X$, $N = H^*Y$ for $X \leftarrow A \rightarrow Y$ maps of spaces.

If *A* fails to be commutative, this fails to give a product, but taking $C^*(X) \leftarrow C^*(B) \rightarrow C^*(E)$ as $X \leftarrow A \rightarrow Y$, one can use the natural DGA maps

$$C^*(B) \otimes C^*(B) \longrightarrow (C_*B \otimes C_*B)^* \xleftarrow{\nabla^*} C^*(B \times B) \xrightarrow{C^*(\Delta)} C^*(B)$$

inducing the cup product (the first two are quasi-isomorphisms, so the direction of ∇^* is not an issue) to obtain a map

$$\operatorname{Tor}_{C^*B\otimes C^*B}(C^*X\otimes C^*X, C^*E\otimes C^*E) \longrightarrow \operatorname{Tor}_{C^*B}(C^*X, C^*E)$$

which we apply after the exterior product, and this yields the product on Tor. In the situation of the Eilenberg–Moore theorem, this product can be shown to be preserved by the isomorphism with $H^*(X \times_B E)$ [McC, Corollary 7.18][GuM, Cor. 3.5][Sm67, Prop. 3.4][CaF21, Thm. A.27].¹⁸

¹⁷ In a sequel, we will elaborate on both definitions and prove their equivalence, but it is not necessary to show our product agrees with Munkholm's in order to use it.

¹⁸ No source the author knows actually shows the product is preserved, but McCleary at least reduces it to an exercise, and Carlson–Franz [CaF21, A.27] spell out some of the steps to this exercise.

Munkholm is able to describe both these products as instances of another product.¹⁹ Please assume the following homotopy-commutative squares of DGC maps.

Applying Ω , which preserves the relation of homotopy by Lemma 3.0.3, this induces a map $\operatorname{Tor}_{\Omega B(A \otimes A)} \longrightarrow \operatorname{Tor}_{\Omega BA}$ by Lemma 4.4, where we have utilized the abbreviation convention in Notation 4.8. The exterior product is a map $(\operatorname{Tor}_{\Omega BA})^{\otimes 2} \longrightarrow \operatorname{Tor}_{(\Omega BA)^{\otimes 2}}$, in this notation, so to define a candidate product we must connect $\operatorname{Tor}_{(\Omega BA)^{\otimes 2}}$ with $\operatorname{Tor}_{\Omega B(A^{\otimes 2})}$. Munkholm does this using ψ and ε , but we can do it in a somewhat simpler way applying the natural quasi-isomorphisms

$$\mathbf{\Omega}\mathbf{B}Z \otimes \mathbf{\Omega}\mathbf{B}Z \xleftarrow{\gamma} \mathbf{\Omega}(\mathbf{B}Z \otimes \mathbf{B}Z) \xrightarrow{\mathbf{\Omega} \vee} \mathbf{\Omega}\mathbf{B}(Z \otimes Z)$$

of Definition 2.1 to the span $X \leftarrow A \rightarrow Y$. The map Tor_{γ} goes in the wrong direction, but this is no issue by Lemma 4.1, since γ is a quasi-isomorphism. All told, one gets the following composite.

Definition 6.2. Given whc-algebra maps and homotopies as in (6.1), the *product on Tor* is

Compactifying notation as per Notation 4.8, this is

 $\left(\operatorname{Tor}_{\boldsymbol{\Omega}BA}\right)^{\otimes 2} \xrightarrow{\operatorname{ext}} \operatorname{Tor}_{(\boldsymbol{\Omega}BA)^{\otimes 2}} \xrightarrow{\sim} \operatorname{Tor}_{\boldsymbol{\Omega}(BA)^{\otimes 2}} \xrightarrow{\operatorname{Tor}} \operatorname{Tor}_{\boldsymbol{\Omega}BA} \xrightarrow{\operatorname{Tor}}_{\boldsymbol{\Omega}} \operatorname{Tor}_{\boldsymbol{\Omega}BA} \xrightarrow{\operatorname{Tor}}_{\boldsymbol{\Omega}BA} \xrightarrow{\operatorname{Tor}}_{\boldsymbol{\Omega}BA$

7. The algebra structure on Tor

As noted in the introduction, Munkholm's product depends for its definition on a choice of homotopies making (6.1) commute, and he conjectured that its properties might therefore be

¹⁹ It takes a little work to see these products as instances of Munkholm's, or that the product we give here agrees with Munkholm's, and we will spell out the details in a sequel article comparing definitions of products, but here we will take the specialization as given.

bad. We aim to show this is untrue, proving the CGA structure from the statement of Theorem 0.1 and discussing some consequences.

First, we show the product map does not depend on our choices.

Theorem 7.0.1. *Given* WHC-algebra maps and homotopies as in (6.1) the product of Definition 6.2 does not depend on the given homotopies of maps $\Omega B(A \otimes A) \longrightarrow \Omega BX$ and $\Omega B(A \otimes A) \longrightarrow \Omega BY$.

Proof. Recall that the use of the homotopy in Equation (6.3) is only in the last two steps, as an application of Lemma 4.4 with

 $R' = \mathbf{\Omega}\mathbf{B}(A \otimes A) \quad R = \mathbf{\Omega}\mathbf{B}A, \quad M' = \mathbf{\Omega}\mathbf{B}(X \otimes X), \quad M = \mathbf{\Omega}\mathbf{B}X, \quad N' = \mathbf{\Omega}\mathbf{B}(Y \otimes Y), \quad N = \mathbf{\Omega}\mathbf{B}Y.$

But then independence is immediate from Proposition 4.6.

The desired properties making Munkholm's product as described in Section 6 a CGA follow in bijection with the defining properties of an sHC-algebra in Definition 5.3. We subdivide the proof accordingly.

Theorem 7.0.2. Let wHC-algebras A, X, Y and wHC-algebra maps $\mathbf{B}X \stackrel{\tilde{\zeta}}{\leftarrow} \mathbf{B}A \rightarrow \mathbf{B}Y$ be given. Suppose each wHC-algebra structure satisfies

1. the unitality condition 5.3.1. Then the product (6.3) on $\text{Tor}_A(X, Y)$ is unital, with unit²⁰

$$k \xrightarrow{\sim} \operatorname{Tor}_{k}(k,k) \xrightarrow{\operatorname{Ior}_{\eta}(\eta,\eta)} \operatorname{Tor}_{\mathbf{\Omega}\mathbf{B}A}(\mathbf{\Omega}\mathbf{B}X,\mathbf{\Omega}\mathbf{B}Y).$$

- 2. the commutativity condition 5.3.2. Then the product is commutative.
- 3. the associativity condition 5.3.3. Then the product is associative.

7.1. Unitality

Unitality is easiest.

Proof of Theorem **7.0.2.1**. We prove that this map is a left unit, the right proof being symmetric. Using the identifications $k \cong \mathbf{B}k \cong \mathbf{\Omega}\mathbf{B}k$, and following through our modified definition of the product, we get the diagram



²⁰ The η here are the units of the *k*-algebra structures on ΩBZ for $Z \in \{A, X, Y\}$, unusually, not to be confused with the unit id $\longrightarrow B\Omega$ of the bar–cobar adjunction.

Commutativity of the first three squares follows from naturality, in brief. At length, the external product is functorial in all six of its entries, giving the first square [CarE, p. 206]. For the second and third, by Definition 2.1, note γ and $\Omega\nabla$ are respectively natural transformations $\Omega(-\otimes -) \longrightarrow \Omega(-) \otimes \Omega(-)$ and $\Omega(B-\otimes B-) \longrightarrow \Omega B(-\otimes -)$.

The last square obviously should commute as a result of the assumed condition $\Phi \circ \mathbf{B}(\eta \otimes \mathrm{id}) =$ id, but to formally verify it we require the six-square diagram (4.5), of which we display only the *A*-X portion, appending the vertical map Tor_{$\Omega B(\eta \otimes \mathrm{id})$}:

That the leftmost square commutes is the same as stating the vertical map we call $\operatorname{Tor}_{\Omega B(\eta \otimes id)}$ exists in the first place, and follows from Lemma 2.7, using the fact that $\mathbf{B}\eta_A = \eta_{\mathbf{B}A}$ and $\mathbf{B}\eta_X = \eta_{\mathbf{B}X}$ are the coaugmentations. Using $\Phi_X \circ \mathbf{B}(\eta_X \otimes id_X) = id_{\mathbf{B}X}$, we may merge the first two squares. Then we extend the commutative diagram to include the composite right homotopy $\Omega \mathbf{B}(k \otimes A) \to \Omega \mathbf{B}(A \otimes A) \to P \Omega \mathbf{B}X$:



where the map $\Omega \mathbf{B}(k \otimes A) \longrightarrow \Omega \mathbf{B}A$ is the one we have been calling $\mathrm{id}_{\Omega \mathbf{B}A}$ under the identification using the condition $\Phi_A(\eta_A \otimes \mathrm{id}_A) = \mathrm{id}_{\mathbf{B}A}$. Converting this to a commutative diagram of Tors, we get



where the isomorphisms along the top become the identity under the standard identifications. $\hfill \Box$

7.2. Commutativity

The proof of commutativity is more involved.

Proof of Theorem **7.0.2.2**. Assume also given whc-algebras A', X', Y' and whc-algebra maps $\mathbf{B}X' \leftarrow \mathbf{B}A' \rightarrow \mathbf{B}Y'$. We write $\chi: A \otimes A' \longrightarrow A' \otimes A$. As in the proof of Item **1**, we trace through the definition of the product; in the last square, we will assume A' = A, X' = X, and Y' = Y and finally use the homotopy-commutativity assumption on the whc-algebra structures.

The vertical maps in the first two squares make sense by the naturality of the external product and γ . The map we have written as Tor_{ΩB_{χ}} makes sense because Munkholm [Mun74, Prop. 3.5] shows the following square commutes up to homotopy:

$$\begin{array}{c|c}
\mathbf{\Omega}\mathbf{B}(A\otimes A') & \xrightarrow{\mathbf{\Omega}(\xi\otimes\xi')} \mathbf{\Omega}\mathbf{B}(X\otimes X') \\
\end{array} \\
\mathbf{\Omega}_{\mathbf{B}\chi} & & & & & \\
\mathbf{\Omega}_{\mathbf{B}\chi} & & & & \\
\mathbf{\Omega}_{\mathbf{B}(A'\otimes A)} & \xrightarrow{\mathbf{\Omega}(\xi'\otimes\xi)} \mathbf{\Omega}_{\mathbf{B}(X'\otimes X)}. \end{array}$$

$$(7.2.2)$$

Thus we are forced to induce the map using the technique of Lemma 4.4.

The commutativity of the external product square appears in Cartan–Eilenberg [CarE, Prop. X.2.1]; the extra sign in their expression is implicit in our definition of the interchange map. The next two squares say, roughly, that the shuffle maps γ and ∇ of Definition 2.1 are commutative. On prepending the tautological twisting cochain, for γ , and postpending it, for ∇ , this boils down to the equation chains

$$\begin{split} \chi \gamma t_{\mathbf{B}A \otimes \mathbf{B}A'} &= \chi (t_{\mathbf{B}A} \otimes \eta_{\mathbf{\Omega}\mathbf{B}A'} \varepsilon_{\mathbf{B}A'} + \eta_{\mathbf{\Omega}\mathbf{B}A} \varepsilon_{\mathbf{B}A} \otimes t_{\mathbf{B}A'}) \\ &= (t_{\mathbf{B}A'} \otimes \eta_{\mathbf{\Omega}\mathbf{B}A} \varepsilon_{\mathbf{B}A} + \eta_{\mathbf{\Omega}\mathbf{B}A'} \varepsilon_{\mathbf{B}A'} \otimes t_{\mathbf{B}A}) \chi = \gamma t_{\mathbf{B}A' \otimes \mathbf{B}A} \chi = \gamma \mathbf{\Omega} \chi t_{\mathbf{B}A \otimes \mathbf{B}A'}, \\ t^{A' \otimes A} \nabla \chi &= (t^{A'} \otimes \eta_{A} \varepsilon_{\mathbf{B}A} + \eta_{A'} \varepsilon_{\mathbf{B}A'} \otimes t^{A}) \chi \\ &= \chi (t^{A} \otimes \eta_{A'} \varepsilon_{\mathbf{B}A'} + \eta_{A} \varepsilon_{\mathbf{B}A} \otimes t^{A'}) = \chi t^{A \otimes A'} \nabla = t^{A' \otimes A} \mathbf{B} \chi \nabla, \end{split}$$

where we have used naturality of the tautological twisting cochains. This works fully for the γ square, but there is something to check for the $\Omega \nabla$ square because the Tor is defined in steps. Let $h: \Omega \mathbf{B}(A \otimes A') \longrightarrow P \Omega \mathbf{B}(X' \otimes X)$ be a right homotopy witnessing the homotopy making (7.2.2) commute. Note that this square commutes on the nose if we precompose $\Omega \nabla: \Omega(\mathbf{B}A \otimes \mathbf{B}A') \longrightarrow \Omega \mathbf{B}(A \otimes A')$, for by the preceding equations and Lemma 2.8, we have

$$\mathbf{B}\chi\left(\underline{\xi}\underline{\otimes}\underline{\xi}'\right)\nabla=\mathbf{B}\chi\nabla\left(\underline{\xi}\underline{\otimes}\underline{\xi}'\right)=\nabla\chi\left(\underline{\xi}\underline{\otimes}\underline{\xi}'\right)=\nabla\left(\underline{\xi}'\underline{\otimes}\underline{\xi}\right)\chi=\left(\underline{\xi}'\underline{\otimes}\underline{\xi}\right)\nabla\chi=\left(\underline{\xi}'\underline{\otimes}\underline{\xi}\right)\mathbf{B}\chi\nabla.$$

Thus the following diagram commutes:

There is less going on here than meets the eye, and the color-coding of objects by quasiisomorphism class is hoped to make the few changes somewhat easier to follow. Arrows defined as compositions are grey and dashed, other non-quasi-isomorphisms are red, and the one right homotopy that is not a composite is gold and wavy. The left face just expresses in a long-winded way that $\Omega \chi \Omega(\xi \otimes \xi') = \Omega(\xi' \otimes \xi) \Omega \chi$, and the next vertical face comes from postcomposing $\Omega \nabla$. The right face is the three-square diagram inducing the map we abusively called $\text{Tor}_{\Omega B\chi}$. The third vertical face is the only interesting one. Its top square is inherited from the second vertical face, its middle square expresses that $\pi_0 \eta \Omega \nabla$ is the function in the preceding long display, and the bottom square commutes since $\chi \circ \chi = \text{id}$. The map from the third face back to the second is trivial except for the bottom two maps in front, which are π_0 , which is possible because $\pi_0 \eta \Omega \nabla$ and $\pi_1 \eta \Omega \nabla$ are equal by the long display and again because $\chi \circ \chi = \text{id}$. The map from the third face to the right face makes sense entirely by definition in the top two cubes; in the bottom cube, the back commutes since $B\chi \circ \nabla = \nabla \circ \chi$, the front commutes trivially, and the bottom commutes from the long display and yet again because $\chi \circ \chi = \text{id}$.

We can finally consider the Φ triangle at the right of (7.2.1). The maps of Tors are induced by the homotopy-commutative squares of the prism in Section 7.2. There are five edge-paths from $\Omega B(A \otimes A')$ to $\Omega B X$, all of lengths one or two, and if we say neighboring paths together bound a face of the prism, then each path has two neighbors. The top and bottom face homotopies, which are the same, are prescribed by the fact $\xi \colon BA \longrightarrow BX$ is a wHC-algebra map, although we have seen we have some flexibility in which we use to define the product, and the left face homotopy is determined by the rest of the argument. The right homotopies representing these homotopies can be expanded to give the large cubical Figure 7.2.4. In this figure, the objects are color-coded by homotopy type, primitive arrows are red, right homotopies wavy gold, and composites defined so as to make squares commute grey. The right face will induce the identity isomorphism on Tor_{ΩBA}. The three small interior cubes and triangular prism on the left commute by definition.

By assumption, there is a DGC homotopy between Φ_A and $\Phi_A \circ \mathbf{B}\chi$, inducing a homotopy from $\Omega \Phi_A$ to $\Omega \Phi_A \circ \Omega \mathbf{B}\chi$, which is represented by a right homotopy $\Omega \mathbf{B}(A \otimes A') \longrightarrow P \Omega \mathbf{B}A$.



Figure 7.2.3: Auxiliary diagrams for the commutativity argument.



Figure 7.2.4: The cube diagram for commutativity.

Postcomposing $P\Omega\xi$ gives a right homotopy $\Omega B(A \otimes A') \longrightarrow P\Omega BX$. The composition of the homotopy represented by the top face in the large cube and this homotopy is a homotopy from $\Omega \Phi_X \circ \Omega(\xi \otimes \xi)$ to $\Omega\xi \circ \Omega \Phi_A \circ \Omega B\chi$, as is the composition of the three homotopies represented by the right homotopies along the lower left. The right homotopies representing these two compositions are themselves homotopic by Corollary 3.1.5, so one finds a commutative diagram of objects receiving maps from $\Omega B(A \otimes A')$ as in Figure 7.2.3b.

Noting that right homotopy $\Omega B(A \otimes A') \longrightarrow P\Omega BX$ on the right edge coming from $\Omega \Phi_A \simeq \Omega \Phi_A \circ \Omega B\chi$ factors through $P\Omega BA$ and the right homotopy on the bottom factors through $\Omega B(A' \otimes A)$, we may plug this diagram into the large rectangle in the front face of the previous cube and take Tor to obtain a large commutative diagram. The map induced on the right edge is the identity map of $\text{Tor}_{\Omega BA}(\Omega BX)$ since the projections π_1 and π_0 induce the same map in cohomology, so this completes the final square in (7.2.1) and with it the proof.

7.3. Associativity

The associativity proof is again more involved.²¹

Proof of Theorem **7.0.2.3**. The template is Figure **7.3.1**, in which we show each square commutes. We can be brief about the proofs the squares from the first three columns commute, which mostly involve only naturality and functoriality, and only deal with the unprimed labels.

- I: The associativity of the external product is classical; cf. Cartan–Eilenberg [CarE, p. 206].
- II: This is the naturality of the external product in the second three variables [CarE, XI.2.1].
- III: This too is the naturality of the external product in the second three variables.
- V: This follows from naturality of γ and the equation $(id \otimes \gamma)\gamma = (\gamma \otimes id)\gamma$. To see this, it is enough to precompose the tautological twisting cochain $t: (\mathbf{B}A)^{\otimes 3} \longrightarrow \mathbf{\Omega}(\mathbf{B}A)^{\otimes 3}$ and expand using the definition in Definition 2.1.
- VI: This follows because γ is a natural transformation $\Omega(-\otimes -) \longrightarrow \Omega(-) \otimes \Omega(-)$.
- VIII: This follows from functoriality of Ω , then naturality of ∇ and the equation $\nabla(id \otimes \nabla) = \nabla(\nabla \otimes id)$, whose proof is dual to that in [V].

The other squares require their own diagrams.

IV: Commutativity of the square follows from naturality of the external product in the last three variables (the first three variables fixed as $\Omega BX \leftarrow \Omega BA \rightarrow \Omega BY$), applied to the diagram

²¹ To follow this proof carefully, it may be helpful to first absorb Section 8, as broadly similar arguments are presented more expansively there.



Figure 7.3.1: The associativity diagram.

and the symmetric diagram with X replaced with Y. The map we call \tilde{h} can be obtained in various (equivalent) ways from the assumed DGC homotopy between DGC maps $\mathbf{B}(A \otimes A) \longrightarrow \mathbf{B}X$. For our purposes it will be most convenient to transpose this to a DGA homotopy between two DGA maps $\mathbf{B}(A \otimes A) \longrightarrow X$ using Lemma 3.0.3, then represent that as a right homotopy $h^P: \Omega \mathbf{B}(A \otimes A) \longrightarrow PX$. We write h^{\dagger} for the composite $\mathbf{B}h^P \circ \eta: \mathbf{B}(A \otimes A) \rightarrow \mathbf{B}\Omega\mathbf{B}(A \otimes A) \longrightarrow BPX$. Then Ωh^{\dagger} is the map $(h^P)^{\#}: \Omega \mathbf{B}(A \otimes A) \longrightarrow \Omega \mathbf{B}PX$ induced up from h^P by Corollary 1.3, and we finally define \tilde{h} to be the composite $Z \circ \Omega h^{\dagger}$ with the natural map $Z: \Omega \mathbf{B}PX \longrightarrow P\Omega \mathbf{B}X$ of Lemma 3.2.8.

We simplify life by referring only to the *A*-*X* side in the remaining squares.

A replacement. To proceed in the diagram, we will need to replace $\operatorname{Tor}_{\Omega BA \otimes \Omega B(A^{\otimes 2})}(\Omega BX \otimes P \Omega BX)$ with $\operatorname{Tor}_{\Omega BA \otimes \Omega B(A^{\otimes 2})}(\Omega BX \otimes \Omega BPX)$. Recall that we defined the right homotopy \tilde{h} as the composite of the DGA quasi-isomorphism $Z: \Omega BPX \longrightarrow P\Omega BX$ and a map $\Omega h^{\dagger}: \Omega B(A^{\otimes 2}) \longrightarrow \Omega BPX$. Factoring \tilde{h} in this way in (7.3.2), we get

(and symmetrically on the *A*-*Y* side). Tensoring this diagram with $\Omega BX \leftarrow \Omega BA \rightarrow \Omega BY$ yields another diagram inducing the map $\operatorname{Tor}_{\operatorname{id}\otimes\Omega\Phi}$ at the top of [VII], connected to the diagram inducing the same map $\operatorname{Tor}_{\operatorname{id}\otimes\Omega\Phi}$ at the bottom of [IV] by the triple $(\operatorname{id}_{\Omega BX}\otimes Z, \operatorname{id}_{\Omega BA\otimes^2}, \operatorname{id}_{\Omega BY}\otimes Z)$ in the third column, and by the identity elsewhere.

VII: This follows from the functoriality of Tor on applying γ , which is a natural transformation $\gamma: \Omega(-\otimes -) \longrightarrow \Omega(-) \otimes \Omega(-)$, to the diagram

once we observe that $\gamma \circ \mathbf{\Omega}(\xi \otimes h^{\dagger}) = (\mathbf{\Omega} \xi \otimes \mathbf{\Omega} h^{\dagger}) \circ \gamma$.

IX: This follows from functoriality of Tor on applying $\Omega \nabla$, which is a natural transformation $\Omega(\mathbf{B}(-) \otimes \mathbf{B}(-)) \longrightarrow \Omega \mathbf{B}(- \otimes -)$ to the diagram (7.3.3). The right homotopy is now witnessed by $\Omega(\xi \otimes h^{\dagger}) : \Omega \mathbf{B}(A \otimes A^{\otimes 2}) \longrightarrow \Omega \mathbf{B}(X \otimes PX)$.

Another replacement. We now want to free the *P* trapped inside the ΩB . To this end, we note that there is a quasi-isomorphic embedding $\iota: X \otimes PX \longrightarrow P(X \otimes X)$ probably most easily understood by identifying each as a subalgebra of $X \otimes I^* \otimes X \cong I^* \otimes X \otimes X$. Postcomposing this ι , we may replace $\operatorname{Tor}_{\Omega B(A^{\otimes 2})}(\Omega B(X \otimes PX))$ with $\operatorname{Tor}_{\Omega B(A^{\otimes 2})}(\Omega BP(X \otimes X))$ on the bottom of [IX]. We may now further postcompose *Z* and replace $\operatorname{Tor}_{\Omega B(A^{\otimes 2})}(\Omega BP(X \otimes X))$ with $\operatorname{Tor}_{\Omega B(A^{\otimes 2})}(P\Omega B(X \otimes X))$.



Figure 7.3.4: Auxiliary diagrams for the associativity argument.



Figure 7.3.5: A part of the system of DGAs underlying Square X from Figure 7.3.1.



Figure 7.3.6: Square X from Figure 7.3.1, filled.

X: Consider the cube in Figure 7.3.4a, giving six maps $\Omega B(A^{\otimes 3}) \longrightarrow \Omega BX$ and six homotopies between them, each the composite of a map and a homotopy across one of the faces.

We may use the associated right homotopies to fill out Figure 7.3.5. The edges from (7.3.4a) are red, the right homotopies corresponding to the faces are gold and wavy, composite maps are grey and dashed, and quasi-isomorphism classes of DGAs are color-coded. In particular all endpoint maps π_0 and π_1 are green. The top of Figure 7.3.5 comes from the replacement we have just made, and the left from its suppressed, symmetric twin following the parallel square IX'; the right and bottom edges are both from Lemma 4.4 and induce Tor_{$\Omega\Phi$}. The diagram as it stands now commutes by definition, and it remains to fill in the interior.

By Lemma 3.0.5, the homotopies from $\Omega B(A^{\otimes 3})$ can be composed, and by Lemma 3.2.6 the composite of two consecutive triples can be represented by a single right homotopy. By Corollary 3.1.5, these composite right homotopies $\Omega B(A^{\otimes 3}) \longrightarrow P\Omega BX$ are themselves homotopic, and this is witnessed by a right homotopy $\Omega B(A^{\otimes 3}) \longrightarrow PP\Omega BX$. We can combine all the codomains into the Figure 7.3.4b, to be thought of as the base of a cone under $\Omega B(A^{\otimes 3})$. Using the factorizations of the maps along the right and bottom edges through $\Omega B(A^{\otimes 2})$ and ΩBA , we may insert this cone into Figure 7.3.5, and taking Tor, obtain Figure 7.3.6, in which the black arrows are isomorphisms and the red are not. This is square X of Figure 7.3.1, and using the commutativity of all its constituent squares and triangles, we see it commutes.

8. Functoriality of the product

Now that we have a ring structure on Tor, we would like also to have ring maps. It is clear the product (6.3) is functorial with respect to spans of DGA maps, but our claims are more expansive.

Theorem 8.0.1. Given whc-algebras A', X', Y', A, X, Y and whc-algebra maps



such that the squares commute up to DGC homotopy, the k-linear map

$$\Xi := \operatorname{Tor}_{\mathbf{\Omega}\lambda_{A}}(\mathbf{\Omega}\lambda_{X}, \mathbf{\Omega}\lambda_{X}): \operatorname{Tor}_{\mathbf{\Omega}\mathbf{B}A'}(\mathbf{\Omega}\mathbf{B}X', \mathbf{\Omega}\mathbf{B}Y') \longrightarrow \operatorname{Tor}_{\mathbf{\Omega}\mathbf{B}A}(\mathbf{\Omega}\mathbf{B}X, \mathbf{\Omega}\mathbf{B}Y)$$

defined as in Lemma 4.4 is multiplicative with respect to the products

$$\Pi': \operatorname{Tor}_{\mathbf{\Omega}BA'}(\mathbf{\Omega}BX', \mathbf{\Omega}BY')^{\otimes 2} \longrightarrow \operatorname{Tor}_{\mathbf{\Omega}BA'}(\mathbf{\Omega}BX', \mathbf{\Omega}BY'),$$
$$\Pi: \operatorname{Tor}_{\mathbf{\Omega}BA}(\mathbf{\Omega}BX, \mathbf{\Omega}BY)^{\otimes 2} \longrightarrow \operatorname{Tor}_{\mathbf{\Omega}BA}(\mathbf{\Omega}BX, \mathbf{\Omega}BY)$$

described in (6.3). That is, $\Pi \circ (\Xi \otimes \Xi) = \Xi \circ \Pi'$.

These algebra homomorphisms are functorial in the following sense: given further wHC-algebra maps $\mathbf{B}X'' \leftarrow \mathbf{B}A'' \rightarrow \mathbf{B}Y''$ and a triple $(\lambda'_X, \lambda'_A, \lambda'_Y)$ of wHC-algebra-maps to $\mathbf{B}X' \leftarrow \mathbf{B}A'' \rightarrow \mathbf{B}Y'$ such that the resulting squares commute up to DGC homotopy, and hence an algebra map

$$\Xi' = \operatorname{Tor}_{\boldsymbol{\Omega}\lambda'_{A}}(\boldsymbol{\Omega}\lambda'_{X},\boldsymbol{\Omega}\lambda'_{Y}) \colon \operatorname{Tor}_{\boldsymbol{\Omega}BA''}(\boldsymbol{\Omega}BX'',\boldsymbol{\Omega}BY'') \longrightarrow \operatorname{Tor}_{\boldsymbol{\Omega}BA'}(\boldsymbol{\Omega}BX',\boldsymbol{\Omega}BY'),$$

the composite $\Xi \circ \Xi'$ is equal to $\operatorname{Tor}_{\Omega\lambda_A \circ \Omega\lambda'_A}(\Omega\lambda_X \circ \Omega\lambda'_X, \Omega\lambda_Y \circ \Omega\lambda'_Y)$.

Corollary 8.0.2. In the situation of Theorem 8.0.1, suppose that $A' = H^*A$, $X' = H^*X$, $Y' = H^*Y$, with $\xi_*: H^*A \longrightarrow H^*X$ and $v_*: H^*A \longrightarrow H^*X$ the DGA maps obtained by conjugating $H^*\Omega\xi$ and $H^*\Omega v$ respectively by $H^*\varepsilon$. Then the induced map $\operatorname{Tor}_{\Omega B H^*A}(\Omega B H^*X, \Omega B H^*Y) \xrightarrow{\sim} \operatorname{Tor}_A(\Omega B X, \Omega B Y)$ is multiplicative with respect to the products induced as in Section 6 by the homotopies witnessing that ξ , v, $B\xi_*$, and Bv_* are SHC-algebra maps.

Proof of Theorem o.6. Conjugating by Tor_{ε} , we may replace $\text{Tor}_{C^*(B)}$ with $\text{Tor}_{\Omega BC^*(B)}$ and $\text{Tor}_{H^*(B)}$ with $\text{Tor}_{\Omega BH^*(B)}$. Now, assuming the spaces in the span $X \leftarrow B \rightarrow E$ have polynomial cohomology, Munkholm [Mun74, 7.2] uses the shC-algebra structure on singular cochains from Theorem 5.6 to produce a trio of DGC quasi-isomorphisms

$$\begin{array}{c|c} \mathbf{B}H^*(X) \leftarrow \mathbf{B}H^*(B) \rightarrow \mathbf{B}H^*(E) \\ \lambda_X & \downarrow & \downarrow \\ \mathbf{B}C^*(X) \leftarrow \mathbf{B}C^*(B) \rightarrow \mathbf{B}C^*(E). \end{array}$$

If *k* has characteristic 2 then for each $Z \in \{X, B, E\}$, if the \smile_1 -squares of a set of polynomial generators of $H^*(Z)$ vanish, λ_Z is a wHC-algebra map; if *k* has characteristic $\neq 2$, they are wHC-algebra maps no matter what [Mun74, 7.3]. If λ_X and λ_E are wHC-algebra maps, the squares commute up to DGC homotopy [Mun74, 7.4], and since we assume additionally λ_B is a wHC-algebra map, Corollary 8.0.2 applies to show the isomorphism is multiplicative.

Remark 8.0.3. Munkholm's additive predecessor of Theorem 8.0.1 requires λ_A be a DGC map only; in his intended case, this is the only map doing the job anyway, but his proof does not require it to be an sHC-algebra map. Our proof, on the other hand, *does* require λ_A be a wHC-algebra map.

In the application to Theorem 0.6, when $A' = H^*(B)$ is a polynomial algebra, this only poses an additional restriction if the characteristic of k is 2, but the obstruction is genuine and not merely a defect of the proof. If Z is topological space with polynomial $H^*(Z; \mathbb{F}_2)$ but such that \smile_1 -squares of generators z are not all decomposable (*i.e.*, if it cannot be guaranteed that the $z \smile_1 z$ lie in $\tilde{H}^*(Z; \mathbb{F}_2) \cdot \tilde{H}^*(Z; \mathbb{F}_2)$), then Saneblidze showed $H^*(\Omega Z; \mathbb{F}_2)$ is not exterior [San17, Cor. 1]. In such cases, $A = C^*(Z; \mathbb{F}_2)$ and $X = Y = C^*(*; \mathbb{F}_2) = \mathbb{F}_2$ is a counterexample to the desired strengthening of Theorem 8.0.1.

Other work of Munkholm also analyzes this situation, and *exotic* shc-algebra structures on cochains.

Remark 8.0.4. The result is more sensitive than it might appear, and the author still does not know cases where the strategy of Theorem 0.6 yields an isomorphism if the cohomology of the inputs is not polynomial, even when DGA quasi-isomorphisms between cochains and cohomology are known. In what is arguably the next-best case, that of Davis–Januszkiewicz spaces, the strategy already fails. The failure of two separate published putative proofs of analogous results in this case was a major motivator for our decision to conduct the proof as much as possible at the level of diagrams of DGA maps.

Proof

The rest of the section constitutes the proof of Theorem 8.0.1. The functorial nature of the maps follows immediately from Theorem 4.9, so it remains to show multiplicativity. That is, we are to connect the products Π' and Π described in Definition 6.2 using the map Ξ of Theorem 8.0.1:

$$\begin{array}{c} \operatorname{Tor}_{\mathbf{\Omega}BA'}(\mathbf{\Omega}BX',\mathbf{\Omega}BY')^{\otimes 2} \xrightarrow{\Pi'} \operatorname{Tor}_{\mathbf{\Omega}BA'}(\mathbf{\Omega}BX',\mathbf{\Omega}BY') \\ \cong \otimes \Xi & \downarrow & \downarrow^{\Xi} \\ \operatorname{Tor}_{\mathbf{\Omega}BA}(\mathbf{\Omega}BX,\mathbf{\Omega}BY)^{\otimes 2} \xrightarrow{\Pi} \operatorname{Tor}_{\mathbf{\Omega}BA}(\mathbf{\Omega}BX,\mathbf{\Omega}BY). \end{array}$$

Expanding out definition of the product partially and employing the space-saving convention of Notation 4.8, we will fill in the following diagram in such a way that commutativity of each

square is manifest:



Expanding all squares simultaneously would result in a diagram more intimidating than illuminating, so we will consider each square separately in its own subsection.

Before we do that, we should give the reader some explanation as to why we might expect the thing to commute at all. The external product, γ , and $\Omega \nabla$ are all natural transformations, so one should expect the squares involving them to commute, and they do, transforming the objectwise tensor-square of the three-square diagram of (4.5) determining Ξ into another such three-square diagram determining the left edge of the Φ square. The only casualty in this process is the right homotopy, which is transmogrified from a standard right homotopy to one encumbered with an increasingly ornate witnessing path object. We then have to deploy the material developed in Section 3 to recover a right homotopy witnessed by the standard path object. The edges of the Φ square come from three- or four-square diagrams per (4.5) determined by the new homotopy we have transported over from the one giving $\Xi \otimes \Xi$ and three of the homotopies appear as hypotheses for Theorem 8.0.1. Filling in the Φ square amounts to constructing DGA maps making these homotopies coherent, and for this we will use the techniques developed in Section 3.

In what follows, we will continuously use the functoriality of Tor, viewed as a graded k-module, in triples of k-DGA maps making the two squares (4.3) commute. All homotopies will be expanded in terms of path objects so that nothing is swept under the rug. Because the squares involving A and Y are notationally symmetric with those involving A and X, we truncate to the A-X portions what will nevertheless be a crushing overburden of diagrams. Every argument will proceed symmetrically and silently on the A-Y side.

Notation 8.o.6. Recall from the statement of Theorem 8.o.1 that we assume a DGC homotopy between the two paths around the square

$$\begin{array}{ccc}
\mathbf{B}A' \xrightarrow{\xi'} \mathbf{B}X' \\
\lambda_A & & & \downarrow \lambda_X \\
\mathbf{B}A \xrightarrow{\xi} \mathbf{B}X.
\end{array}$$

Via the adjunction of Lemma 3.0.3, we obtain a DGA homotopy $h = h_X$ between the two transposed maps $\Omega BA' \longrightarrow X$. Write $h^P : \Omega BA' \longrightarrow PX$ for the DGA map representing this DGA homotopy per Definition 3.1.3. We write $(h^P)^{\#} : \Omega BA' \longrightarrow \Omega BPX$ for the map induced up as in Corollary 1.3, which encodes a homotopy between the induced maps $(\xi \circ \lambda_A)^{\#}$ and $(\lambda_X \circ \xi')^{\#} : \Omega BA' \longrightarrow \Omega BX$ in the sense that one gets these maps back from $(h^P)^{\#}$ by postcomposing $\Omega B\pi_0$ and $\Omega B\pi_1$ respectively.

8.1. The external product square

To express $\Xi \otimes \Xi$ on the left of (8.0.5) in terms of DGA maps, we will find it more convenient to describe Ξ using $(h^P)^{\#}$ rather than with the right homotopy $\Omega BA' \longrightarrow P\Omega BX$ with the same endpoints representing the DGA homotopy $\Omega(\xi \circ \lambda_A) \simeq \Omega(\lambda_X \circ \xi')$ induced from the assumed DGC homotopy. Fortunately, Lemma 3.2.8 gives us a quasi-isomorphism $Z: \Omega BPX \longrightarrow P\Omega BX$ such that $\pi_j \circ Z = \Omega B\pi_j$, so we can replace the expected diagram on the left below with that on the right and have the same induced map Ξ on Tor.

Then the external product square is the composite of subsquares

in which each horizontal map is the exterior product and the vertical maps in each file are determined functorially by applying Tor to the right diagram of (8.1.1) and its tensor-square. Here, as promised, we have suppressed the symmetric A-Y half of the diagram in a bid for comprehensibility, and the unlabeled map is the necessary composition rendering the diagram commutative.

In summary, the input diagram of DGAs commutes by definition and the output diagram of Tors commutes by the functoriality of the exterior product.

8.2. The γ square

What we call the γ square in (8.0.5) arises by applying the cobar shuffle $\gamma: \Omega(-\otimes -) \longrightarrow \Omega(-) \otimes \Omega(-)$ of Definition 2.1 across the board, landing in the right edge of the Tor diagram

(8.1.2) from the external product square of the preceding subsection:

This Tor diagram is induced by the DGA diagram



in which all horizontal arrows are γ and h^{\dagger} : **B** $A' \longrightarrow$ **B**PX is the transpose of h^{P} : **ΩB** $A' \longrightarrow PX$. The left face commutes by the previous step, and the top two and bottom horizontal faces commute by naturality of γ , but the right face and the third horizontal face remain to be explained.

Since the transpose is given by applying **B** and precomposing η : **B** $A' \longrightarrow$ **B** Ω **B**A', and h^P represents the transpose of the original homotopy $\xi \circ \lambda_A \simeq \lambda_X \otimes \xi'$ it follows that **B** $\pi_j \circ h^{\dagger} =$ **B** $(\pi_j \circ h^P) \circ \eta$ are respectively $\xi \circ \lambda_A$ and $\lambda_X \otimes \xi'$ again for $j \in \{0, 1\}$, giving commutativity of the right face. The remaining horizontal face commutes, since $\Omega h^{\dagger} = \Omega B h^P \circ \Omega \eta = (h^P)^{\#}$ by Corollary 1.3.

8.3. The $\Omega \nabla$ square

In what we called the $\Omega \nabla$ square in (8.0.5), all horizontal maps are id or $\Omega \nabla$. The expanded rectangle of Tors is

$$\operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega(\mathbf{B}X')^{\otimes 2} \right) \xrightarrow{\operatorname{Tor}_{\operatorname{id}}(\Omega\nabla)} \operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega\mathbf{B}(X')^{\otimes 2} \right) \xrightarrow{} \operatorname{Tor}_{\operatorname{id}}(\Omega\nabla) \xrightarrow{} \operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega(\mathbf{B}X)^{\otimes 2} \right) \xrightarrow{} \operatorname{Tor}_{\operatorname{id}}(\Omega\nabla) \xrightarrow{} \operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega\mathbf{B}(X^{\otimes 2}) \right) \xrightarrow{} \operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega(\mathbf{B}PX)^{\otimes 2} \right) \xrightarrow{} \operatorname{Tor}_{\operatorname{id}}(\Omega\nabla) \xrightarrow{} \operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega\mathbf{B}(PX)^{\otimes 2} \right) \xrightarrow{} \operatorname{Tor}_{\operatorname{id}}(\Omega\nabla) \xrightarrow{} \operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}} \left(\Omega\mathbf{B}(X^{\otimes 2}) \right) \xrightarrow{} \operatorname{Tor}_{\Omega(\mathbf{B}A)^{\otimes 2}} \left(\Omega(\mathbf{B}X)^{\otimes 2} \right) \xrightarrow{} \operatorname{Tor}_{\operatorname{id}}(\Omega\nabla) \xrightarrow{} \operatorname{Tor}_{\Omega\mathbf{B}(A^{\otimes 2})} \left(\Omega\mathbf{B}(X^{\otimes 2}) \right).$$

The prism of inducing DGA maps is morally (but not exactly) the following, in which all horizontal maps are $\Omega \nabla$:



The left face is the right face of the γ square of Section 8.2, and the top, bottom, front, and back, each containing two edges $\Omega \nabla$, commute because $\nabla (- \otimes -) = (- \otimes -) \nabla$ by Lemma 2.8.

The right face does not necessarily commute as stands. The issue is the limited functoriality of $\underline{\otimes}$ in DGC maps: we have $\mathbf{B}(\pi_0 \otimes \pi_0)(h^{\dagger} \underline{\otimes} h^{\dagger}) = \mathbf{B}\pi_0 h^{\dagger} \underline{\otimes} \mathbf{B}\pi_0 h^{\dagger} = \lambda_X \xi' \underline{\otimes} \lambda_X \xi'$, and similarly for π_1 , but no guarantee that $\lambda_X \xi' \underline{\otimes} \lambda_X \xi'$ should equal $(\lambda_X \underline{\otimes} \lambda_X)(\xi' \underline{\otimes} \xi')$. However, because the other five faces of each cube commute, when one prepends $\mathbf{\Omega} \nabla : \mathbf{\Omega}(\mathbf{B}A' \otimes \mathbf{B}A') \longrightarrow \mathbf{\Omega}\mathbf{B}(A' \otimes A')$ to the composites of the two maps around any square of the right face, the resulting maps *are* equal. Hence we replace the three copies of $\mathbf{\Omega}\mathbf{B}(A' \otimes A')$ along the back right edge with $\mathbf{\Omega}(\mathbf{B}A' \otimes \mathbf{B}A')$ and get commutative cubes. The right face we build off of in subsequent diagrams then becomes



8.4. Repackaging the homotopy

The format we require for representatives of homotopies in the upcoming Section 8.5 has only one *P*, whereas $\Omega B(PX \otimes PX)$ has two, so we need to reformat this map to fit into the diagram to come. Thus we will convert $\Omega(h^{\dagger} \otimes h^{\dagger}) : \Omega(BA' \otimes BA') \longrightarrow \Omega B(PX \otimes PX)$ and its associated endpoint maps $\Omega B(\pi_j^{\otimes 2}) \circ \Omega(h^{\dagger} \otimes h^{\dagger}) : \Omega(BA' \otimes BA')^{\otimes 2} \longrightarrow \Omega B(X \otimes X)$ into a standard right homotopy $\Omega(BA' \otimes BA') \rightarrow P\Omega B(X \otimes X) \rightrightarrows \Omega B(X \otimes X)$ in a number of steps.

- Recall the natural surjection *r*: (*P*(−))^{⊗2} → *D*(−) of Lemma 3.2.7, morally restricting a square of DGA maps to two adjacent edges. Postcomposing ΩB*r*, we obtain a homotopy representative Ω(B*r* ∘ (*h*[†] ⊗ *h*[†])): Ω(B*A*')^{⊗2} → ΩB*D*(*X*^{⊗2}). By construction, the new endpoint maps agree with the old: ΩB*p_j* ∘ Ω(B*r* ∘ (*h*[†] ⊗ *h*[†])) = ΩB(επ_j)^{⊗2} ∘ Ω(*h*[†] ⊗ *h*[†]).
- Recall also the composition operation $D(X^{\otimes 2}) \stackrel{\varepsilon}{\leftarrow} \Omega B D(X^{\otimes 2}) \stackrel{\Upsilon}{\longrightarrow} P(X^{\otimes 2})$ of Lemma 3.2.2. We want to attach our existing homotopy representative to ΩB of this operation, to wit,

$$\mathbf{\Omega}\mathbf{B}D(X^{\otimes 2}) \xleftarrow{\mathbf{\Omega}\mathbf{B}\varepsilon} \mathbf{\Omega}\mathbf{B}\mathbf{\Omega}\mathbf{B}D(X^{\otimes 2}) \xrightarrow{\mathbf{\Omega}\mathbf{B}\Upsilon} \mathbf{\Omega}\mathbf{B}P(X^{\otimes 2}).$$

In order to accomplish this, we need our map $\Omega(\mathbf{B}r \circ (h^{\dagger} \otimes h^{\dagger}))$: $\Omega \mathbf{B}(A')^{\otimes 2} \longrightarrow \Omega \mathbf{B}D(X^{\otimes 2})$ to factor through $\Omega \mathbf{B}\varepsilon$. Fortunately, our map is Ωg for the DGC map $g := \mathbf{B}r \circ (h^{\dagger} \otimes h^{\dagger})$, so by Corollary 1.4, it indeed factors, and we have the following commutative diagram:



Let \bar{h} denote the composition along the top, $\Omega B(\Upsilon \circ \Omega g) \circ \Omega \eta$. By the definition of Υ and functoriality of ΩB , the new representative has the same endpoint maps: $\Omega B \pi_j \circ \bar{h} = \Omega B p_j \circ \Omega g$ for $j \in \{0, 1\}$.

Now to free *P* from ΩB, we postcompose the natural map *Z*: ΩB*P*(*X*^{⊗2}) → *P*ΩB(*X*^{⊗2}) of Lemma 3.2.8 to get a map *h* := *Z* ∘ *h* satisfying π_j*h* = ΩBπ_j ∘ *h* for *j* ∈ {0,1}.

All told, we will be able to glue the left of the following diagram to the lower two squares on the existing right face (8.3.2) the ∇ diagram:



All of the horizontal maps are quasi-isomorphisms, so the diagrams induce isomorphisms in Tor. In terms of the Tor diagram (8.3.1) of the $\Omega \nabla$ square, this will replace $\operatorname{Tor}_{\Omega(\mathbf{B}A')^{\otimes 2}}(\Omega \mathbf{B}(PX)^{\otimes 2})$ with the isomorphic $\operatorname{Tor}_{\Omega \mathbf{B}(A')^{\otimes 2}}(P(\Omega \mathbf{B}X)^{\otimes 2})$ at the right of the third row and otherwise leave the diagram unchanged.

8.5. The Φ square

In filling in the Φ square of (8.0.5), we are subject to a few constraints. The isomorphisms of the previous large squares essentially carry $\Xi \otimes \Xi$ along to the left edge, and the top, bottom, and right of the square must respectively describe Π' , Π , and Ξ . Thus our pre-existing commitments amount to Figure 8.5.1. It is not yet obvious this should commute, but we will fill it in such a way as to make commutativity apparent. In terms of general strategy, counting *P*'s, it is visible that we already have four homotopies of maps present, to be accounted for as follows:

- Begin with the homotopies assumed in Theorem 8.0.1 witnessing that ξ' , ξ , v, v' are shcalgebra-maps and precompose the two $\Omega \nabla$ maps $\Omega(\mathbf{B}A' \otimes \mathbf{B}A') \longrightarrow \Omega \mathbf{B}(A' \otimes A')$ and $\Omega \nabla \colon \Omega(\mathbf{B}A \otimes \mathbf{B}A) \longrightarrow \Omega \mathbf{B}(A \otimes A)$. The maps along the top and bottom of Figure 8.5.1 then follow from the associated six-square diagrams as in Lemma 4.4
- The maps on the left are inherited from the previous squares of (8.0.5).



Figure 8.5.1: The constraints on filling the Φ square.

• The maps on the right come from the six-square diagram that Lemma 4.4 associates to the homotopy-commutativity of the squares of wHC-algebra maps in the statement of Theorem 8.0.1.

The two remaining homotopies are those we have assumed to make λ_A , λ_X , λ_Y sHC-algebra maps. The six of these on the *A*-*X* side (the *A*-*Y* argument as usual proceeds silently in parallel) together make up the cube of Figure 8.5.2a; note that we have precomposed $\Omega \nabla$ so that $\Omega(\mathbf{B}A' \otimes \mathbf{B}A')$ and $\Omega(\mathbf{B}A \otimes \mathbf{B}A)$ rather than $\Omega \mathbf{B}(A' \otimes A')$ and $\Omega \mathbf{B}(A \otimes A)$ appear.



Figure 8.5.2: Auxiliary diagrams for the functoriality argument.

The right homotopies witnessing these together fit into Figure 8.5.3. We have color-coded the DGAs by quasi-isomorphism type to match (8.5.2a) and colored the arrows coming from (8.5.2a) in red; we do not need to label them because they are uniquely determined by their source and target. Gold wavy arrows are right homotopies corresponding to the faces in (8.5.2a) and grey dashed arrows are the defined as the necessary composites making the diagram commutative. The projections from path objects are green, and arranged so that π_0 always points up or left, π_1

down or right. The reader should convince themself Figure 8.5.3 expresses only the existence of right homotopies representing the homotopies we have just discussed. We are not yet asserting anything about the front or back of the large prism on the lower right.



Figure 8.5.3: The assemblage of right homotopies implied by Figure 8.5.2a.

Again by Lemma 3.0.5, the homotopies from $\Omega(\mathbf{B}A')^{\otimes 2}$ can be composed, and by Lemma 3.2.6 the composite of two consecutive triples can be represented by a single right homotopy. By Corollary 3.1.5, these composite right homotopies $\Omega(\mathbf{B}A')^{\otimes 2} \longrightarrow P\Omega\mathbf{B}X$ are themselves homotopic, and this is witnessed by a right homotopy $\Omega(\mathbf{B}A')^{\otimes 2} \longrightarrow PP\Omega\mathbf{B}X$. We can combine all the codomains into Figure 8.5.2b, to be thought of as a cone under $\Omega(\mathbf{B}A')^{\otimes 2}$. Using the factorizations of the maps along the right and bottom edges through $\Omega\mathbf{B}(A^{\otimes 2})$ and $\Omega\mathbf{B}A$, we may insert this cone into Figure 8.5.3, and taking Tor, obtain Figure 8.5.4, in which the black arrows are isomorphisms and the red are not. This is the Φ square of (8.0.5), and using all subdividing commutative squares and triangles, we see it commutes.

This completes the Φ diagram and the proof Ξ is multiplicative with respect to Π' and Π .

Remark 8.5.5. It has long been known [Baues81, §2][GerV95, Cor. 6] that the bar construction of the normalized cochain algebra $BC^*(X)$ of a connected simplicial set carries a differential graded Hopf algebra structure functorial in X (also known as a homotopy Gerstenhaber algebra structure on $C^*(X)$, and related [Fr19c, Rmk. 4.2] to the traditional sHC-algebra structure on $C^*(X)$). As a result, there is an idea for an alternative construction of a product on Tor, starting with the exterior product and γ as we have done, then following not with $\Omega \nabla$ and squares involving Φ , but $\Omega \mu$ for μ : $BC^*(B) \otimes BC^*(B) \longrightarrow BC^*(B)$ the DG Hopf algebra multiplication, which is a DGC map. As far as we can tell, such a proof would require stronger hypotheses than the present one.



Figure 8.5.4: The completed Φ square.

Remark 8.5.6 (Sketches of other ways forward). It has been pointed out to us that results in E_n -spectra could likely also be used to prove an analogue to Theorem 0.1, with some interpretation. Basterra and Mandell show [BM11, Thm. 5.3] that the bar construction $B\tilde{A}$ of a so-called augmented partial \mathscr{C}_n -algebra \tilde{A} is an augmented partial \mathscr{C}_{n-1} -algebra, where \mathscr{C}_n is the little *n*-cubes operad. A variant of their proof likely establishes (but this should be checked) that given a span of partial \mathscr{C}_n -algebras $\tilde{X} \leftarrow \tilde{A} \rightarrow \tilde{Y}$, the two-sided bar construction $B(\tilde{X}, \tilde{A}, \tilde{Y})$ is also a partial \mathscr{C}_{n-1} -algebra. A suitably enhanced version of the Dold–Kan correspondence should take a span $X \rightarrow A \rightarrow Y$ of E_3 -algebras to a span of augmented partial \mathscr{C}_3 -algebras $\tilde{X} \leftarrow \tilde{A} \rightarrow \tilde{Y}$, so that $B_{\bullet}(\tilde{X}, \tilde{A}, \tilde{Y})$ becomes a partial \mathscr{C}_2 -algebra whose cohomology is $\operatorname{Tor}_A(X, Y)$, and this should also give a CGA structure on Tor functorial in triples of E_3 -algebra maps making the expected pair of squares commute.

There are minor attendant difficulties in formalizing this argument, which would involve generalizing the Basterra–Mandell proof rather than simply applying their result, and the connection of true and partial C_n -algebras is not direct, but passes through a certain zigzag of equivalences. An additional complication, for our intended topological application in Theorem 0.6, is that the existing sHC-algebra formality maps given by Munkholm inducing the additive isomorphism $\operatorname{Tor}_{H^*(B)}(H^*(X), H^*(E)) \longrightarrow \operatorname{Tor}_{C^*(B)}(C^*(X), C^*(E))$ have not been shown to be E_3 -algebra maps (and as far as this author can see may not be), and the squares only commute up to A_{∞} -homotopy anyway. It may of course be that sufficient functoriality properties can be recovered for this hypothetical other version of the product as well, but this is not obvious. A benefit of the existing argument leading to Theorem 0.6 is that the hypotheses seem to be minimal, and the homotopycommutative squares of sHC-algebra maps to be taken as input to the theorem are already known, so that from our current Theorem 0.1, Theorem 0.6 is automatic, whereas with another approach, as we have discussed, some additional massaging would be needed.

Another suggestion that has come to the author involves a result of Fresse [Fre10] that the bar construction of an E_{∞} -algebra A is another E_{∞} -algebra. This likely applies as well to the twosided bar construction $\mathbf{B}(X, A, Y)$ of a span of E_{∞} -algebras, hence inducing a CGA structure on its cohomology. If so, this would induce the expected CGA structures on $\operatorname{Tor}_{C^*(B)}(C^*(X), C^*(E))$ and $\operatorname{Tor}_{H^*(B)}(H^*(X), H^*(E))$ under mild flatness conditions. Again, however, there is not a reason to expect this construction of a CGA structure to be to be functorial under the sort of homotopycommutative squares of A_{∞} -algebra maps we already have, and thus this approach as well does not immediately yield our target application.

References

- [Ad56] J. Frank Adams. On the cobar construction. *Proc. Natl. Acad. Sci. USA*, 42(7):409, 1956. doi:10.1073/pnas.42.7.409.
- [AH56] J. Frank Adams and Peter J. Hilton. On the chain algebra of a loop space. Comment. Math. Helv., 30(1):305– 330, 1956. doi:10.1007/BF02564350.
- [At62] Michael F. Atiyah. Vector bundles and the Künneth formula. *Topology*, 1(3):245–248, Sept 1962. doi:10.1016/0040-9383(62)90107-6.
- [BaGM59] M. G. Barratt, Victor K.A.M. Gugenheim, and John C. Moore. On semisimplicial fibre-bundles. American J. Math., 81(3):639–657, 1959. doi:10.2307/2372920.

- [BM11] Maria Basterra and Michael A. Mandell. Homology of E_n ring spectra and iterated *THH*. *Algebr. Geom. Topol.*, 11(2):939–981, 2011. arXiv:1007.5315, doi:10.2140/agt.2011.11.939.
- [BaMR14] Tobias Barthel, J. Peter May, and Emily Riehl. Six model structures for DG-modules over DGAs: model category theory in homological action. New York J. Math, 20:1077–1159, 2014. http://nyjm.albany.edu/j/2014/20-53.html, arXiv:1310.1159, doi:10.2140/agt.2013.13.1089.
- [Baues81] Hans-Joachim Baues. The double bar and cobar constructions. *Comp. Math.*, 43(3):331-341, 1981. http://numdam.org/item/CM_1981_43_3_331_0.
- [Baum] Paul Frank Baum. Cohomology of Homogeneous Spaces. Ph.D. thesis, Princeton, December 1962. http://dropbox.com/s/u14kyz01id9vnet/cohomology_homogeneous_spaces%28Baum_dissertation_ORIGINAL%29.pdf?dl
- [Baum68] Paul F. Baum. On the cohomology of homogeneous spaces. *Topology*, 7(1):15–38, 1968. doi:10.1016/0040-9383(86)90012-1.
- [BaumS67] Paul Baum and Larry Smith. The real cohomology of differentiable fibre bundles. *Comment. Math. Helv.*, 42(1):171–179, 1967. doi:10.1007/BF02564416.
- [BeF04] Clemens Berger and Benoit Fresse. Combinatorial operad actions on cochains. *Math. Proc. Cambridge Philos. Soc.*, 137(1):135–174, 2004. arXiv:0109158, doi:10.1017/S0305004103007138.
- [Bor51] Armand Borel. *Cohomologie des espaces localement compacts d'aprés J. Leray,* volume 2 of *Lecture Notes in Math.* Springer-Verlag, 1951. Sém. de Top. alg., ETH.
- [Bor53] Armand Borel. Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts. Ann. of Math. (2), 57(1):115-207, 1953. http://web.math.rochester.edu/people/faculty/doug/otherpapers/Borel-Sur.pdf, doi:10.2307/1969728.
- [BousG] Aldridge Knight Bousfield and Victor K.A.M. Gugenheim. On PL De Rham Theory and Rational Homotopy Type, volume 179 of Mem. Amer. Math. Soc. Amer. Math. Soc., 1976. https://books.google.com/books?id=MAPUCQAAQBAJ.
- [Br59] Edgar H. Brown. Twisted tensor products, I. Ann. of Math. (2), 29(1):223–246, Jan 1959. doi:10.2307/1970101.
- [CaF21] Jeffrey D. Carlson (appendix joint with Matthias Franz). The cohomology of biquotients via a product on the two-sided bar construction (expository version). 2021. arXiv:2106.02986v1.
- [Car51] Henri Cartan. La transgression dans un groupe de Lie et dans un espace fibré principal. In Colloque de topologie (espace fibrés), Bruxelles 1950, pages 57–71, Liège/Paris, 1951. Centre belge de recherches mathématiques, Georges Thone/Masson et companie. http://eudml.org/doc/112227.
- [Car⁺54] Henri Cartan et al. *Séminaire Henri Cartan*, 7, 1954–5. http://numdam.org/volume/SHC_1954-1955__7.
- [Car55] Henri Cartan. DGA-modules (suite), notion de construction. *Séminaire Henri Cartan*, 7(1), 1954–1955. http://numdam.org/item/SHC_1954-1955__7_1_A3_0.
- Eilenberg. Algebra, [CarE] Henri Cartan and Samuel Homological volume 19 of Princeton Math. Ser. Princeton Univ. Press, 1999 (1956). http://www.math.stonybrook.edu/~mmovshev/BOOKS/homologicalalgeb033541mbp.pdf.
- [Cl65] Allan Clark. Homotopy commutativity and the Moore spectral sequence. *Pacific J. Math*, 15:65–74, 1965. doi:10.2140/pjm.1965.15.65.
- [DCH16] Gabriel C. Drummond-Cole and Joseph Hirsh. Model structures for coalgebras. *Proc. Amer. Math. Soc.*, 144(4):1467–1481, 2016. arXiv:1411.5526, doi:10.1090/proc/12823.
- [EM53] Samuel Eilenberg and Saunders Mac Lane. On the groups *H*(Π, *n*), I. *Ann. of Math.* (2), 58(1):55–106, 1953. doi:10.2307/1969820.
- [EM54] Samuel Eilenberg and Saunders Mac Lane. On the groups $H(\Pi, n)$, II: Methods of computation. *Ann. of Math.* (2), 60(1):49–139, 1954. doi:10.2307/1969702.
- [EM065] Samuel Eilenberg and John C. Moore. Homology and fibrations I: Coalgebras, cotensor product and its derived functors. *Comment. Math. Helv.*, 40(1):199–236, 1965. doi:10.1007/BF02564371.
- [EZ53] Samuel Eilenberg and Joseph A. Zilber. On products of complexes. *Amer. J. Math.*, 75(1):200–204, 1953. doi:10.2307/2372629.

- [Fr19a] Matthias Franz. The cohomology rings of homogeneous spaces. 2019. arXiv:1907.04777.
- [Fr19b] Matthias Franz. The cohomology rings of smooth toric varieties and quotients of moment-angle complexes. 2019. arXiv:1907.04791.
- [Fr19c] Matthias Franz. Homotopy Gerstenhaber algebras are strongly homotopy commutative. 2019. arXiv:1907.04778.
- [Fre10] Benoit Fresse. The bar complex of an E-infinity algebra. Adv. Math., 223(6):2049– 2096, 2010. https://www.sciencedirect.com/science/article/pii/S0001870809003430, doi:https://doi.org/10.1016/j.aim.2009.08.022.
- [GerV95] Alexander Murray Gerstenhaber and Α. Voronov. Homotopy G-algebras and moduli space operad. Int. Math. Res. Not., 1995(3):141-153, 02 1995. https://academic.oup.com/imrn/article-pdf/1995/3/141/6768479/1995-3-141.pdf, arXiv:hep-th/9409063,doi:10.1155/S1073792895000110.
- [GetG99] Ezra Getzler and Paul Goerss. A model category structure for differential graded coalgebras. 1999 manuscript. http://ncatlab.org/nlab/files/GetzlerGoerss99.pdf.
- [Gu6o] Victor K.A.M. Gugenheim. On a theorem of E. H. Brown. *Illinois J. Math.*, 4(2):292–311, 1960. doi:10.1215/ijm/1255455870.
- [GuM] Victor K.A.M. Gugenheim and J. Peter May. *On the Theory and Applications of Differential Torsion Products,* volume 142 of *Mem. Amer. Math. Soc.*, Amer. Math. Soc., 1974.
- [GuM74] Victor K.A.M. Gugenheim and Hans J. Munkholm. On the extended functoriality of Tor and Cotor. J. *Pure Appl. Algebra*, 4(1):9–29, 1974. doi:10.1016/0022-4049(74)90026-7.
- [HeKRS17] Kathryn Hess, Magdalena Kędziorek, Emily Riehl, and Brooke Shipley. A necessary and sufficient condition for induced model structures. J. Topology, 10(2):324–369, 2017. arXiv:1509.08154, doi:10.1112/topo.12011.
- [Hin97] Vladimir Hinich. Homological algebra of homotopy algebras. *Comm. Algebra*, 25(10):3291–3323, 1997. doi:10.1080/00927879708826055.
- [Hod75] Luke Hodgkin. The equivariant Künneth theorem in K-theory. In Topics in K-theory, pages 1–101. Springer, 1975. doi:10.1007/BFb0082285.
- Husemoller, John C. and James [HuMS74] Dale Moore, Stasheff. Differential homologihomogeneous spaces. J. Pure Appl. cal algebra and Algebra, 5(2):113-185, 1974. http://math.mit.edu/~hrm/18.917/husemoller-moore-stasheff.pdf.
- [Jar97] John Frederick Jardine. A closed model structure for differential graded algebras. In Joachim J.R. Cuntz and Masoud Khalkhali, editors, *Cyclic cohomology and noncommutative geometry* (*Waterloo*, *ON*, 1995), volume 17 of *Fields Inst. Commun.*, pages 55–58. Amer. Math. Soc., 1997. https://ncatlab.org/nlab/files/JardineModelDG.pdf.
- [KS05] Tornike Kadeishvili and Samson Saneblidze. A cubical model for a fibration. J. Pure Appl. Algebra, 196(2-3):203–228, 2005. arXiv:0210006, doi:10.1016/j.jpaa.2004.08.017.
- [LH] Kenji Lefèvre-Hasegawa. Sur les A_{∞} -catégories. Ph.D. thesis, Université de Paris VII—Denis Diderot, 2002. arXiv:CT/0310337.
- [LV] Jean-Louis Loday and Bruno Vallette. *Algebraic Operads*, volume 346 of *Grundlehren Math. Wiss.* Springer, 2012. https://pure.mpg.de/rest/items/item_3121746/component/file_3121747/content.
- [Mac] Saunders Mac Lane. *Homology*. Classics in Mathematics (Grundlehren Math. Wiss. vol. 114). Springer, 2012 (1963, 1975).
- [May68] J. Peter May. The cohomology of principal bundles, homogeneous spaces, and two-stage Postnikov systems. Bull. Amer. Math. Soc., 74(2):334–339, 1968. http://www.math.uchicago.edu/~may/PAPERS/6.pdf.
- [MayNo2] J. Peter May and Frank Neumann. On the cohomology of generalized homogeneous spaces. *Proc. Amer. Math. Soc.*, 130(1):267–270, 2002. doi:10.1090/S0002-9939-01-06372-9.
- [McC] John McCleary. A User's Guide to Spectral Sequences, volume 58 of Cambridge Stud. Adv. Math. Cambridge Univ. Press, Cambridge, 2001.
- [McS03] James McClure and Jeffrey Smith. Multivariable cochain operations and little *n*-cubes. J. Amer. Math. Soc., 16(3):681–704, 2003. arXiv:0106024, doi:10.1090/S0894-0347-03-00419-3.

- [Mil56] John W. Milnor. Construction of universal bundles, II. Ann. of Math. (2), 63(3):430-436, May 1956. http://math.mit.edu/~hrm/18.906/milnor-construction-universal-ii.pdf, doi:10.2307/1970012.
- [Mo60] John C. Moore. Algèbre homologique et homologie des espaces classifiants. *Séminaire Henri Cartan*, 12(1), 1959-1960. Talk 7. http://www.numdam.org/item/SHC_1959-1960__12_1_A7_0.
- [Mun74] Hans J. Munkholm. The Eilenberg–Moore spectral sequence and strongly homotopy multiplicative maps. *J. Pure Appl. Algebra*, 5(1):1–50, 1974. doi:10.1016/0022-4049(74)90002-4.
- [Mun76a] Hans J. Munkholm. shm maps of differential algebras, I. A characterization up to homotopy. J. Pure Appl. Algebra, 9(1):39–46, 1976. doi:10.1016/0022-4049(76)90004-9.
- [Mun76b] Hans J. Munkholm. shm maps of differential algebras, II. Applications to spaces with polynomial cohomology. J. Pure Appl. Algebra, 9(1):47–63, 1976. doi:10.1016/0022-4049(76)90005-0.
- [Mun76c] Hans J. Munkholm. shm maps of differential algebras, III. Multiplicative stuctures on torsion products. In preparation as of 1976, current whereabouts unknown.
- [Mun78] Hans J. Munkholm. DGA algebras as a Quillen model category: Relations to shm maps. J. Pure Appl. Algebra, 13(3):221–232, 1978. doi:10.1016/0022-4049(78)90009-9.
- [MuR019] Fernando Muro and Constanze Roitzheim. Homotopy theory of bicomplexes. J. Pure Appl. Algebra, 223(5):1913–1939, 2019.
- [On] Arkadi L. Onishchik. Topology of transitive transformation groups. Johann Ambrosius Barth, 1994.
- [Pan15] Taras Panov. On the cohomology of quotients of moment-angle complexes. Russian Math. Surveys, 70(4):779–781, aug 2015. arXiv:1506.06875, doi:10.1070/RM2015v070n04ABEH004961.
- [Pr11] Alain Prouté. A_∞-structures. Modèles minimaux de Baues-Lemaire et Kadeishvili et homologie des fibrations. *Repr. Theory Appl. Categ*, 21:1-99, 2011. http://tac.mta.ca/tac/reprints/articles/21/tr21abs.html.
- [Qui69] Daniel Quillen. Rational homotopy theory. Annals of Math., 90(2):205–295, Sept 1969. doi:10.2307/1970725.
- [Rec70] David L. Rector. Steenrod operations in the Eilenberg–Moore spectral sequence. *Comment. Math. Helv.*, 45(1):540–552, 1970.
- [Riv19] Manuel Rivera. Adams' cobar construction revisited. 2019. arXiv:1910.08455.
- [RZ18] Manuel Rivera and Mahmoud Zeinalian. Cubical rigidification, the cobar construction and the based loop space. Algebraic & Geometric Topology, 18(7):3789–3820, 2018. arXiv:1612.04801, doi:10.2140/agt.2018.18.3789.
- [Sano9] Samson Saneblidze. The bitwisted cartesian model for the free loop fibration. *Topology Appl.*, 156(5):897–910, 2009. arXiv:0707.0614, doi:10.1016/j.topol.2008.11.002.
- [San17] Samson Saneblidze. The loop cohomology of a space with the polynomial cohomology algebra. *Trans. A. Razmadze Math. Inst.*, 171(3):389–395, 2017. doi:10.1016/j.trmi.2017.07.002.
- [Sin93] Wilhelm Singhof. On the topology of double coset manifolds. *Math. Ann.*, 297(1):133–146, 1993. doi:10.1007/BF01459492.
- [Sm67] Larry Smith. Homological algebra and the Eilenberg–Moore spectral sequence. *Trans. Amer. Math. Soc.*, 129:58–93, 1967. doi:10.2307/1994364.
- [Smith] Larry Smith. Lectures on the Eilenberg–Moore spectral sequence, volume 134 of Lecture Notes in Math. Springer-Verlag New York, 1970.
- [St63a] James Dillon Stasheff. Homotopy associativity of *H*-spaces. I. *Trans. Amer. Math. Soc.*, 108(2):275–292, 1963. doi:10.2307/1993608.
- [St63b] James Dillon Stasheff. Homotopy associativity of *H*-spaces. II. *Trans. Amer. Math. Soc.*, 108(2):293–312, 1963. doi:10.2307/1993609.
- [StH70] James Stasheff and Steve Halperin [*sic*]. Differential algebra in its own rite [*sic*]. In *Proc. Adv. Study Inst. Alg. Top.(Aarhus 1970)*, volume 3, pages 567–577, 1970.
- [Su60] Masahiro Sugawara. On the homotopy-commutativity of groups and loop spaces. *Mem. College Sci. Univ. Kyoto Ser. A Math.*, 33(2):257–269, 1960. doi:10.1215/kjm/1250775911.

- [Wolf77] Joel L. Wolf. The cohomology of homogeneous spaces. *Amer. J. Math.*, pages 312–340, 1977. doi:10.2307/2373822.
- [Wolf78] Joel L. Wolf. The real and rational cohomology of differential fibre bundles. *Trans. Amer. Math. Soc.*, 245:211–220, 1978. doi:10.1090/S0002-9947-1978-0511406-7.

DEPARTMENT OF MATHEMATICS IMPERIAL COLLEGE LONDON j.carlson@imperial.ac.uk