DENIZALP GOKTAS, Brown University, USA JIAYI ZHAO, Pomona College, USA AMY GREENWALD, Brown University, USA

A prevalent theme in the economics and computation literature is to identify natural price-adjustment processes by which sellers and buyers in a market can discover equilibrium prices. An example of such a process is tâtonnement, an auction-like algorithm first proposed in 1874 by French economist Walras in which sellers adjust prices based on the Marshallian demands of buyers, i.e., budget-constrained utility-maximizing demands. A dual concept in consumer theory is a buyer's Hicksian demand, i.e., consumptions that minimize expenditure while achieving a desired utility level. In this paper, we identify the maximum of the absolute value of the elasticity of the Hicksian demand, i.e., the maximum percentage change in the Hicksian demand of any good w.r.t. the change in the price of some other good, as an economic parameter sufficient to capture and explain a range of convergent and non-convergent tâtonnement behaviors in a broad class of markets. In particular, we prove the convergence of *tâtonnement* at a rate of $O((1+e^2)/T)$, in homothetic Fisher markets with bounded price elasticity of Hicksian demand, i.e., Fisher markets in which consumers have preferences represented by homogeneous utility functions and the price elasticity of their Hicksian demand is bounded, where ε is the maximum absolute value of the price elasticity of Hicksian demand across all buyers. Our result not only generalizes known convergence results for CES Fisher markets, but extends them to mixed nested CES markets and Fisher markets with continuous, possibly non-concave, homogeneous utility functions. Our convergence rate covers the full spectrum of nested CES utilities, including Leontief and linear utilities, unifying previously existing disparate convergence and non-convergence results. In particular, for $\varepsilon = 0$, i.e., Leontief markets, we recover the best-known convergence rate of O(1/T), and as $\varepsilon \to \infty$, e.g., linear Fisher markets, we obtain non-convergent behavior, as expected.

CCS Concepts: • Theory of computation \rightarrow Exact and approximate computation of market equilibria.

ACM Reference Format:

Denizalp Goktas, Jiayi Zhao, and Amy Greenwald. 2023. Tâtonnement in Homothetic Fisher Markets. In Proceedings of the 24th ACM Conference on Economics and Computation (EC '23), July 9–12, 2023, London, United Kingdom. ACM, New York, NY, USA, 34 pages. https://doi.org/10.1145/3580507.3597746

1 INTRODUCTION

Competitive (or Walrasian or market) equilibrium [Arrow and Debreu 1954; Walras 1896], first studied by French economist Léon Walras in 1874, is the steady state of an economy—any system governed by supply and demand [Walras 1896]. Walras assumed that each producer in an economy would act so as to maximize its profit, while consumers would make decisions that maximize their preferences over their available consumption choices; all this, while perfect competition prevails, meaning producers and consumers are unable to influence the prices that emerge. Under these assumptions, the demand and supply of each commodity is a function of prices, as they are a consequence of the decisions made by the producers and consumers, having observed the prevailing

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

EC '23, July 9-12, 2023, London, United Kingdom

^{© 2023} Association for Computing Machinery.

ACM ISBN 979-8-4007-0104-7/23/07...\$15.00

https://doi.org/10.1145/3580507.3597746

prices. A competitive equilibrium then corresponds to prices that solve the system of simultaneous equations with demand on one side and supply on the other, i.e., prices at which supply meets demand. Unfortunately, Walras did not provide conditions that guarantee the existence of such a solution, and the question of whether such prices exist remained open until Arrow and Debreu's rigorous analysis of competitive equilibrium in their model of a competitive economy in the middle of last century [Arrow and Debreu 1954].

The Arrow-Debreu model comprises a set of commodities; a set of firms, each deciding what quantity of each commodity to supply; and a set of consumers, each choosing a quantity of each commodity to demand in exchange for their endowment [Arrow and Debreu 1954]. Arrow and Debreu define a *competitive equilibrium* as a collection of consumptions, one per consumer, a collection of productions, one per firm, and prices, one per commodity, such that fixing equilibrium prices: (1) no consumer can increase their utility by deviating to an alternative affordable consumption, (2) no firm can increase profit by deviating to another production in their production set, and (3) the *aggregate demand* for each commodity (i.e., the sum of the commodity's production and endowment across firms and consumers, respectively), while the total value of the aggregate demand is equal to the total value of the aggregate supply, i.e., *Walras' law* holds.

Arrow and Debreu proceeded to show that their competitive economy could be seen as an abstract economy, which today is better known as a pseudo-game [Arrow and Debreu 1954; Facchinei and Kanzow 2010]. A pseudo-game is a generalization of a game in which the actions taken by each player impact not only the other players' payoffs, as in games, but also their set of permissible actions. Arrow and Debreu proposed generalized Nash equilibrium as the solution concept for this model, an action profile from which no player can improve their payoff by unilaterally deviating to another action in the space of permissible actions determined by the actions of other players. Arrow and Debreu further showed that any competitive economy could be represented as a pseudogame inhabited by a fictional auctioneer, who sets prices so as to buy and resell commodities at a profit, as well as consumers and producers, who respectively, choose utility-maximizing consumptions of commodities in the budget sets determined by the prices set by the auctioneer, and profitmaximizing productions at the prices set by the auctioneer. The elegance of the reduction from competitive economies to pseudo-games is rooted in a simple observation: the set of competitive equilibria of a competitive economy is equal to the set of generalized Nash equilibria of the associated pseudo-game, implying the existence of competitive equilibrium in competitive economies as a corollary of the existence of generalized Nash equilibria in pseudo-games, whose proof is a straightforward generalization of Nash's proof for the existence of Nash equilibria [Nash 1950].¹

With the question of existence out of the way, this line of work on competitive equilibrium, which today is known as general equilibrium theory [McKenzie et al. 2005], turned its attention to questions of (1) efficiency, (under what assumptions are competitive equilibria Pareto-optimal?) (2) uniqueness (under what assumptions are competitive equilibria unique?), and (3) stability (under what conditions would a competitive economy settle into a competitive equilibrium?). The first two questions were answered between the 1950s and 1970s [Arrow 1951b; Arrow and Hurwicz 1958; Arrow and Nerlove 1958; Balasko 1975; Debreu 1951; Dierker 1982; Hahn 1958; Pearce and Wise 1973], showing that (1) under suitable assumptions (e.g., see Arrow [1951a]) competitive equilibrium demands are Pareto-optimal, and (2) competitive equilibria are unique in markets with an *excess demand function*, (i.e., the difference between the aggregate demand and supply functions),

¹McKenzie [1959] would prove the existence of competitive equilibrium independently, but concurrently. Much of his work, however, has gone unrecognized perhaps because his proof technique does not depend on this fundamental relationship between competitive and abstract economies.

which satisfies the *gross substitutes* (GS) condition (i.e., the excess demand of any commodity increases if the price of any other commodity increases, fixing all other prices). In regards to the question of stability, most relevant work is concerned with the convergence properties of a natural auction-like price-adjustment process, known as *tâtonnement*, which mimics the behavior of the *law of supply and demand*, updating prices at a rate equal to the excess demand [Arrow et al. 1971; Kaldor 1934]. Research on *tâtonnement* in the economics literature is motivated by the fact that it can be understood as a plausible explanation of how prices move in real-world markets. Hence, if one could prove convergence in all exchange economies, then perhaps it would be justifiable to claim real-world markets would also eventually settle at a competitive equilibrium.

Walras [1896] conjectured, albeit without conclusive evidence, that tâtonnement would converge to a competitive equilibrium. While a handful of results guarantee the convergence of $t\hat{a}$ tonnement under mathematical conditions without widely agreed-upon economic interpretations [Nikaidô and Uzawa 1960; Uzawa 1960], Arrow and Hurwicz [1958; 1960] were the first to formally establish the convergence of tâtonnement to unique competitive equilibrium prices in a class of economically well-motivated competitive economies, namely those that satisfy the GS assumption. Following this promising result, Scarf [1960] dashed all hope that tâtonnement would prove to be a universal price-adjustment process that converges in all economies, by showing that competitive equilibrium prices are unstable under tâtonnement dynamics in his eponymous competitive economy without firms, and with only three commodities and three consumers with Leontief preferences, i.e., the Scarf exchange economy. Scarf's negative result seems to have discouraged further research by economists on the stability of competitive equilibrium [Fisher 1975]. Despite research on this question coming to a near halt, one positive outcome was achieved, on the convergence of a non-tâtonnement update rule known as Smale's process [Herings 1997; Kamiya 1990; Smale 1976; van der Laan et al. 1987], which updates prices at the rate of the product of the excess demand and the inverse of its Jacobian, in most competitive economies, even beyond GS, again suggesting the possibility that real-world economies could indeed settle at a competitive equilibrium.

Nearly half a century after these seminal analyses of competitive economies, research on the stability of competitive equilibrium is once again coming to the fore, this time in computer science, perhaps motivated by applications of algorithms such as tâtonnement to load balancing over networks [Jain et al. 2013], or to pricing of transactions on crypotocurrency blockchains [Leonardos et al. 2021; Liu et al. 2022; Reijsbergen et al. 2021]. A detailed inquiry into the computational properties of market equilibria was initiated by Devanur et al. [2008], who studied a special case of the Arrow-Debreu competitive economy known as the Fisher market [Brainard et al. 2000]. This model, for which Irving Fisher computed equilibrium prices using a hydraulic machine in the 1890s, is essentially the Arrow-Debreu model of a competitive economy, but there are no firms, and buyers are endowed with only one type of commodity-hereafter good²-an artificial currency [Brainard et al. 2000; Codenotti et al. 2007]. Devanur et al. [2002] exploited a connection first made by Eisenberg [1961] between the Eisenberg-Gale program and competitive equilibrium to solve Fisher markets assuming buyers with linear utility functions, thereby providing a (centralized) polynomial-time algorithm for equilibrium computation in these markets [Devanur et al. 2002; Devanur et al. 2008]. Their work was built upon by Jain et al. [2005], who extended the Eisenberg-Gale program to all Fisher markets in which buyers have continuous, quasi-concave, and homogeneous utility functions, and proved that the equilibrium of Fisher markets with such buyers can be computed in polynomial time by interior point methods.

 $^{^{2}}$ In the context of Fisher markets, commodities are typically referred to as goods Cheung et al. [2013], as Fisher markets are often analyzed for a single time period only. More generally, in Arrow-Debreu markets, where commodities vary by time, location, or state of the world, "an apple today" may be different than "an apple tomorrow". For consistency with the literature, we refer to commodities as goods.

Concurrent with this line of work on computing competitive equilibrium using centralized methods, a line of work on devising and proving convergence guarantees for decentralized priceadjustment processes (i.e., iterative algorithms that update prices according to a predetermined update rule) developed. This literature has focused on devising *natural* price-adjustment processes, like *tâtonnement*, which might explain or imitate the movement of prices in real-world markets. In addition to imitating the law of supply and demand, *tâtonnement* has been observed to replicate the movement of prices in lab experiments, where participants are given endowments and asked to trade with one another [Gillen et al. 2020]. Perhaps more importantly, the main premise of research on the stability of competitive equilibrium in computer science is that for competitive equilibrium to be justified, not only should it be backed by a natural price-adjustment process as economists have long argued, but it should also be computationally efficient [Codenotti et al. 2007].

The first result on this question is due to Codenotti et al. [2005], who introduced a discrete-time version of *tâtonnement*, and showed that in exchange economies that satisfy *weak gross substitutes (WGS)*, the *tâtonnement* process converges to an approximate competitive equilibrium in a number of steps which is polynomial in the approximation factor and size of the problem. Unfortunately, soon after this positive result appeared, Papadimitriou and Yannakakis [2010] showed that it is impossible for a price-adjustment process based on the excess demand function to converge in polynomial time to a competitive equilibrium in general, ruling out the possibility of Smale's process (and many others) justifying the notion of competitive equilibrium in all competitive economies. Nevertheless, further study of the convergence of price-adjustment processes such as *tâtonnement* under stronger assumptions, or in simpler models than full-blown Arrow-Debreu competitive economies, remains worthwhile, as these processes are being deployed in practice [Jain et al. 2013; Leonardos et al. 2021; Liu et al. 2022; Reijsbergen et al. 2021].

Toward this end, in this paper we make strides towards analyzing the computational complexity of discrete-time *tâtonnement* in *homothetic Fisher markets*, i.e., Fisher markets in which consumers have continuous and homothetic preferences.³ An important concept in consumer theory is a buyer's Hicksian demand, i.e., consumptions that minimize expenditure while achieving a desired utility level. In this paper, we identify the maximum elasticity of the Hicksian demand, i.e., the maximum percentage change in the Hicksian of any good w.r.t. the change in the price of some other good, as an economic parameter sufficient to capture and explain a range of convergent and non-convergent *tâtonnement* behaviors in a broad class of markets. In particular, we prove the convergence of *tâtonnement* in homothetic Fisher markets with bounded elasticity of Hicksian demand, i.e., Fisher markets in which consumers have preferences represented by homogeneous utility functions for which the elasticity of their Hicksian demand is bounded.

Interpretation via Pseudo-Game Theory. Recall that in the pseudo-game associated with a competitive economy [Arrow and Debreu 1954], a (fictional) auctioneer sets prices for commodities, firms choose what quantity of each commodity to produce, and consumers choose what quantity of each commodity to consume in exchange for their endowment. Running *tâtonnement* in this pseudo-game amounts to the auctioneer running a first-order method, namely a gradient ascent dynamic on their profit function, while the consumers and firms reply with their best response. Interestingly, if the competitive economy satisfies WGS, then the excess demand function is monotone over all *tâtonnement* trajectories. This pseudo-game can then be understood as a monotone

³We refer to Fisher markets that comprise buyers with a certain utility function by the name of the utility function, e.g., we call a Fisher market that comprises buyers with Leontief utility functions a Leontief Fisher market. We omit the "continuous" qualifier as competitive equilibrium is not guaranteed to exist when preferences are not continuous.

Economy type	Pseudo-game type	Mathematical object
WGS Economy	Monotone pseudo-game	Monotone Variational Inequality
Homothetic Fisher market	Zero-sum game	Convex Potential

Table 1. Summary of the equivalences among economy types, pseudo-game types, and mathematical objects.

variational inequality [Facchinei and Kanzow 2010], whose solutions are competitive equilibrium prices.

As *tâtonnement* is intended to be an explanation of real-world market behavior, proofs of its convergence would ideally rely on justifiable economic assumptions. Via the connection between pseudo-games games and competitive economies, establishing convergence of *tâtonnement* can be reduced to discovering justifiable economic assumptions to impose so that the ensuing pseudo-game or variational inequality satisfies suitable mathematical conditions for convergence.

Unfortunately, without imposing significant additional assumptions (such as the excess demand function of the competitive economy being Lipschitz-smooth; see, for example, Golowich et al. [2020]) or relying on more complex update rules such as *extragradient descent* [Korpelevich 1976], first-order methods are not guaranteed to converge in last iterates⁴ in monotone pseudo-games or monotone variational inequalities. Undeterred, in addition to assuming WGS, Cole and Fleischer [2008] impose economic assumptions on the Marshallian own-price elasticity of demand and Marsallian income elasticity of demand, which imply Lipschitz-smoothness of the excess demand function over *tâtonnement* trajectories, and obtain convergence of *tâtonnement* in last iterates in Fisher markets with WGS and bounded price/income elasticity of Marshallian demand.

Paralleling the duality between WGS competitive economies and monotone pseudo-games, any homothetic Fisher market is equivalent to a zero-sum game. Moreover, this zero-sum game can be further reduced to a convex potential, i.e., the Eisenberg-Gale program's dual, whose solutions are competitive equilibrium prices [Cheung et al. 2013; Devanur et al. 2002]. With this equivalence in hand, we seek to identify economically justifiable assumptions that translate into mathematical conditions on the excess demand function that are sufficient for convergence, because first-order methods run on zero-sum games and convex potentials are otherwise not guaranteed to converge to an optimal solution in last iterates. One obvious candidate condition is Lipschitz-smoothness of the excess demand function, but this property does not hold in Leontief Fisher markets, a flavor of homothetic markets in which *tâtonnement* converges! Our research has led to the discovery that assuming bounded elasticity of Hicksian demand yields an excess demand function that is Lipschitz-continuous as well as Bregman-smooth w.r.t. to the KL divergence,⁵ two properties which are sufficient to guarantee the convergence of *tâtonnement* in homothetic Fisher markets. *Our contribution, then, is to identify the economic assumptions that imply the requisite mathematical properties that yield convergence of*


1.1 Technical Contributions

Earlier work [Cheung 2014; Cheung et al. 2013] has established a convergence rate of $(1 - \Theta(1))^T$ for CES Fisher markets excluding the linear and Leontief cases, and of O(1/T) for Leontief and


⁴We note that the standard convergence metric in the *tâtonnement* literature is convergence in last iterates (see, for example, Cheung et al. [2013]).

⁵Bregman-smoothness is a generalization of Lipschitz-smoothness introduced by Cheung et al. [2018] following work by Birnbaum et al. [2011]; see Section 2 for the mathematical definition.

Denizalp Goktas, Jiayi Zhao, and Amy Greenwald



(a) The convergence rates of *tâtonnement* for different Fisher markets. We color previous contributions in blue, and our contribution in red, i.e., we study homothetic Fisher markets where ε is the maximum absolute value of the price elasticity of Hicksian demand across all buyers. We note that the convergence rate for WGS markets does not apply to markets where the price elasticity of Marshallian demand is unbounded, e.g., linear Fisher markets; likewise, the convergence rate for nested CES Fisher markets does not apply to linear or Leontief Fisher markets.



(b) Cross-price elasticity taxonomy of wellknown homogeneous utility functions. There are no previously studied utility functions in the space of utility functions with negative Hicksian cross-price elasticity. Future work could investigate this space and prove faster convergence rates than those provided in this paper. We note that our convergence result covers the entire spectrum of this taxonomy (excluding limits of the *y*-axis).

Fig. 1. A summary of known results in Fisher markets.

nested⁶ CES Fisher markets, where $T \in \mathbb{N}_+$ is the number of iterations for which *tâtonnement* is run. In linear Fisher markets, however, *tâtonnement* does not converge. We generalize these results by proving a convergence rate of $O((1+\varepsilon^2)/T)$, where ε is the maximum absolute value of the price elasticity of Hicksian demand across all buyers. Our convergence rate covers the full spectrum of homothetic Fisher markets, including mixed CES markets, i.e., CES markets with linear, Leontief, and (nested) CES buyers, unifying previously existing disparate convergence and non-convergence results. In particular, for $\varepsilon = 0$, i.e., Leontief Fisher markets, we recover the best-known convergence rate of O(1/T), and as $\varepsilon \to \infty$, i.e., linear Fisher markets, we obtain the non-convergent behaviour of *tâtonnement* [Cole and Tao 2019]. We summarize known convergence results in light of our results in Figure 1a.

We observe that, in contrast to general competitive economies, in homothetic Fisher markets, concavity of the utility functions is not necessary for the existence of competitive equilibrium (Theorem 2.2). A computational analog of this result also holds, namely that *tâtonnement* converges in homothetic Fisher markets, even when buyers' utility functions are non-concave. Our results parallel known results on the convergence of *tâtonnement* in WGS markets, where concavity of utility functions is again not necessary for convergence [Codenotti et al. 2005].

1.2 Related Work

Following Codenotti et al.'s [2005] initial analysis of *tâtonnement* in competitive economies that satisfy WGS, Garg and Kapoor [2004] introduced an auction algorithm that also converges in polynomial time for linear exchange economies. More recently, Bei et al. [2015] established faster convergence bounds for *tâtonnement* in WGS exchange economies.

⁶See Chapter 10 of Cheung [2014].

Another line of work considers price-adjustment processes in variants of Fisher markets. Cole and Fleischer [2008] analyzed *tâtonnement* in a real-world-like model satisfying WGS called the ongoing market model. In this model, *tâtonnement* once-again converges in polynomial-time [Cole and Fleischer 2008; Cole et al. 2010], and it has the advantage that it can be seen as an abstraction for market processes. Cole and Fleischer's results were later extended by Cheung et al. [2012] to ongoing markets with *weak gross complements*, i.e., the excess demand of any commodity weakly increases if the price of any other commodity weakly decreases, fixing all other prices, and ongoing markets with a mix of WGC and WGS commodities. The ongoing market model these two papers study contains as a special case the Fisher market; however Cole and Fleischer [2008] assume bounded own-price elasticity of Marshallian demand, and bounded income elasticity of Marshallian demand, while Cheung et al. [2012] assume, in addition to Cole and Fleischer's assumptions, bounded adversarial market elasticity, which can be seen as a variant of bounded cross-price elasticity of Marshallian lian demand, from below. With these assumptions, these results cover Fisher markets with a small range of the well-known CES utilities, including CES Fisher markets with $\rho \in [0, 1)$ and WGC Fisher markets with $\rho \in (-1, 0]$.⁷

Cheung et al. [2013] built on this work by establishing the convergence of *tâtonnement* in polynomial time in nested CES Fisher markets, excluding the limiting cases of linear and Leontief markets, but nonetheless extending polynomial-time convergence guarantees for *tâtonnement* to Leontief Fisher markets as well. More recently, Cheung and Cole [2018] showed that Cheung et al.'s [2013] result extends to an asynchronous version of *tâtonnement*, in which good prices are updated during different time periods. In a similar vein, Cheung et al. [2019] analyzed *tâtonnement* in online Fisher markets, determining that *tâtonnement* tracks competitive equilibrium prices closely provided the market changes slowly.

Another price-adjustment process that has been shown to converge to market equilibria in Fisher markets is *proportional response dynamics*, first introduced by Wu and Zhang [2007] for linear utilities; then expanded upon and shown to converge by [Zhang 2011] for all CES utilities; and very recently shown to converge in Arrow-Debreu exchange economies with linear and CES ($\rho \in [0, 1)$) utilities by Brânzei et al.. The study of the proportional response process was proven fundamental when Cheung et al. noticed its relationship to gradient descent. This discovery opened up a new realm of possibilities in analyzing the convergence of market equilibrium processes. For example, it allowed Cheung et al. [2018] to generalize the convergence results of proportional response dynamics to Fisher markets for buyers with mixed CES utilities. This same idea was applied by Cheung et al. [2013] to prove the convergence of *tâtonnement* in Leontief Fisher markets, using the equivalence between mirror descent [Boyd et al. 2004] on the dual of the Eisenberg-Gale program and *tâtonnement*, first observed by Devanur et al. [2008]. More recently, Gao and Kroer [2020] developed methods to solve the Eisenberg-Gale convex program in the case of linear, quasi-linear, and Leontief Fisher markets.

An alternative to the (global) competitive economy model, in which an agent's trading partners are unconstrained, is the Kakade et al. [2004] model of a graphical economies. This model features local markets, in which each agent can set its own prices for purchase only by neighboring agents, and likewise can purchase only from neighboring agents. Auction-like price-adjustment processes have been shown to converge in variants of this model assuming WGS [Andrade et al. 2021].

2 PRELIMINARIES

Notation. We use caligraphic uppercase letters to denote sets (e.g., X); bold lowercase letters to denote vectors (e.g., p, π); bold uppercase letters to denote matrices (e.g., X) and lowercase letters

⁷We refer the reader to Section 2 for a definition of CES utilities in terms of the substitution parameter ρ .

to denote scalar quantities (e.g., x, δ). We denote the *i*th row vector of a matrix (e.g., X) by the corresponding bold lowercase letter with subscript *i* (e.g., x_i). Similarly, we denote the *j*th entry of a vector (e.g., p or x_i) by the corresponding Roman lowercase letter with subscript *j* (e.g., p_j or x_{ij}). If a correspondence $\mathcal{A} : X \rightrightarrows \mathcal{Y}$ is singleton valued for some $x \in X$, for notational convenience, we treat it as an element of X, i.e., $\mathcal{A}(x) \in X$, rather than a subset of it, i.e., $\mathcal{A}(x) \subset X$. We denote the set of numbers $\{1, \ldots, n\}$ by [n], the set of natural numbers by \mathbb{N} , the set of real numbers by \mathbb{R}_+ and the set of strictly positive real numbers by \mathbb{R}_{++} . We let $\Delta_n = \{x \in \mathbb{R}^n_+ \mid \sum_{i=1}^n x_i = 1\}$. We denote the interior of any set \mathcal{A} by $\operatorname{int}(\mathcal{A})$. We define $\mathcal{B}_{\varepsilon}(x) = \{z \in \mathcal{Z} \mid ||z - x|| \le \varepsilon\}$ to be the closed ε -ball centered at x, where \mathcal{Z} and $\|\cdot\|$ will be clear from context. We denote the gradient $\nabla_x : C^1(X) \to C^0(X)$ as the operator which takes as input a functional $f : X \to \mathbb{R}$, and outputs a vector-valued function consisting of the partial derivatives of f w.r.t. x. Finally, by notational overload, we define the subdifferential g of a function f at a point $a \in U$ by $\partial_x f(a) = \{g \mid f(x) \ge f(a) + g^T(x-a)\}$.

2.1 Mathematical Preliminaries

A function $f : \mathbb{R}^m \to \mathbb{R}$ is said to be *homogeneous of degree* $k \in \mathbb{N}_+$ if $\forall \mathbf{x} \in \mathbb{R}^m, \lambda > 0, f(\lambda \mathbf{x}) = \lambda^k f(\mathbf{x})$. Unless otherwise indicated, without loss of generality, a homogeneous function is assumed to be homogeneous of degree 1. Fix any norm $\|\cdot\|$. Given $\mathcal{A} \subset \mathbb{R}^d$, the function $f : \mathcal{A} \to \mathbb{R}$ is said to be λ_f -*Lipschitz-continuous* iff $\forall \mathbf{x}_1, \mathbf{x}_2 \in \mathcal{X}, \|f(\mathbf{x}_1) - f(\mathbf{x}_2)\| \le \lambda_f \|\mathbf{x}_1 - \mathbf{x}_2\|$. If the gradient of f is $\lambda_{\nabla f}$ -Lipschitz-continuous, we then refer to f as $\lambda_{\nabla f}$ -Lipschitz-smooth.

2.2 Mirror Descent

Consider the optimization problem $\min_{x \in V} f(x)$, where $f : \mathbb{R}^n \to \mathbb{R}$ is a differentiable convex function and *V* is the feasible set of solutions. A standard method for solving this problem is the *mirror descent algorithm* [Boyd et al. 2004]:

$$\mathbf{x}(t+1) = \underset{\mathbf{x}\in V}{\arg\min} \left\{ \ell_f(\mathbf{x}, \mathbf{x}(t)) + \gamma_t \delta_h(\mathbf{x}, \mathbf{x}(t)) \right\} \qquad \text{for } t = 0, 1, 2, \dots \qquad (1)$$
$$\mathbf{x}(0) \in \mathbb{R}^n \qquad (2)$$

Here, $\gamma_t > 0$ is the step size at time t, $\ell_f(\mathbf{x}, \mathbf{y})$ is the *linear approximation* of f at \mathbf{y} , that is $\ell_f(\mathbf{x}, \mathbf{y}) = f(\mathbf{y}) + \nabla f(\mathbf{y})^T (\mathbf{x} - \mathbf{y})$, and $\delta_h(\mathbf{x}, \mathbf{x}(t))$ is the *Bregman divergence* of a convex differentiable *kernel* function $h(\mathbf{x})$ defined as $\delta_h(\mathbf{x}, \mathbf{y}) = h(\mathbf{x}) - \ell_h(\mathbf{x}, \mathbf{y})$ [Bregman 1967]. In particular, when $h(\mathbf{x}) = \frac{1}{2}||\mathbf{x}||_2^2$, $\delta_h(\mathbf{x}, \mathbf{y}) = \frac{1}{2}||\mathbf{x} - \mathbf{y}||_2^2$. In this case, mirror descent reduces to projected gradient descent [Boyd et al. 2004]. If instead the kernel is the weighted entropy $h(\mathbf{x}) = \sum_{i \in [n]} (x_i \log(x_i) - x_i)$, the Bregman divergence reduces to the *generalized Kullback-Leibler (KL) divergence* [Joyce 2011]:

$$\delta_{\mathrm{KL}}(\boldsymbol{x}, \boldsymbol{y}) = \sum_{i \in [n]} \left[x_i \log \left(\frac{x_i}{y_i} \right) - x_i + y_i \right] \quad , \tag{3}$$

which, when $V = \Delta_n$, yields the following simplified *entropic descent* update rule:

A function f is said to be γ -Bregman-smooth [Cheung et al. 2018] w.r.t. a Bregman divergence with kernel function h if $f(\mathbf{x}) \leq \ell_f(\mathbf{x}, \mathbf{y}) + \gamma \delta_h(\mathbf{x}, \mathbf{y})$. Birnbaum et al. [2011] showed that if the objective function $f(\mathbf{x})$ of a convex optimization problem is γ -Bregman w.r.t. to some Bregman

divergence δ_h , then mirror descent with Bregman divergence δ_h converges to an optimal solution $f(\mathbf{x}^*)$ at a rate of O(1/t). We require a slightly modified version of this theorem, introduced by Cheung et al. [2013], where it suffices for the γ -Bregman-smoothness property to hold only for consecutive pairs of iterates.

THEOREM 2.1 (BIRNBAUM ET AL. [2011], CHEUNG ET AL. [2013]). Let $\{\mathbf{x}^t\}_t$ be the iterates generated by mirror descent with Bregman divergence δ_h . Suppose f and h are convex, and for all $t \in \mathbb{N}$ and for some $\gamma > 0$, it holds that $f(\mathbf{x}^{(t+1)}) \leq \ell_f(\mathbf{x}^{(t+1)}, \mathbf{x}^{(t)}) + \gamma \delta_h(\mathbf{x}^{(t+1)}, \mathbf{x}^{(t)})$. If \mathbf{x}^* is a minimizer of f, then the following holds for mirror descent with fixed step size γ : for all $t \in \mathbb{N}$, $f(\mathbf{x}^{(t)}) - f(\mathbf{x}^*) \leq$ $\gamma/t \delta_h(\mathbf{x}^*, \mathbf{x}^{(0)})$.

2.3 Consumer Theory

Let $X = \mathbb{R}_{+}^{m}$ be a set of possible consumptions over m goods s.t. for any $x \in \mathbb{R}_{+}^{m}$ and $j \in [m]$, $x_{j} \geq 0$ represents the amount of good $j \in [m]$ consumed by consumer (hereafter, buyer) i. The preferences of buyer i over d such that the buyer (resp. weakly) prefers a choice $x \in X$ to another choice $y \in X$ iff $x >_{i} y$ (resp. $x \ge_{i} y$). A preference relation is said to be *complete* iff for all $x, y \in X$, either $x \ge_{i} y$ or $y \ge_{i} x$, or both. A preference relation is said to be *transitive* if, for all $x, y, z \in X$, $x \ge_{i} z$ whenever $x \ge_{i} y$ and $y \ge_{i} z$. A preference relation is said to be *continuous* if for any sequence $\{x^{(n)}, y^{(n)}\}_{n \in \mathbb{N}_{+}} \subset X \times X$ $(x^{(n)}, y^{(n)}) \to (x, y)$ and $x^{(n)} \ge_{i} y^{(n)}$ for all $n \in \mathbb{N}_{+}$, it also holds that $x \ge_{i} y$. A preference relation \ge_{i} is said to be *locally non-satiated* iff for all $x \in X$ and $\epsilon > 0$, there exists $y \in \mathcal{B}_{\epsilon}(x)$ such that $y >_{i} x$. A utility function $u_{i} : X \to \mathbb{R}_{+}$ assigns a positive real value⁸ to elements of X, i.e., to all possible consumptions. Every continuous utility function represents some complete, transitive, and continuous preference relation \ge_{i} over goods s.t. if $u_{i}(x) \ge u_{i}(y)$ for two bundles of goods $x, y \in \mathbb{R}^{m}$, then $x \ge_{i} y$ [Arrow et al. 1971].

In this paper, we consider the general class of *homothetic* preferences \geq_i s.t. for any consumption $x, y \in X$ and $\lambda \in \mathbb{R}_+$, $x \geq_i y$ and $y \geq_i x$ implies $\lambda x \geq_i \lambda y$ and $\lambda y \geq_i \lambda x$, respectively. A preference relation \geq_i is complete, transitive, continuous, and homothetic iff it can be represented via a continuous and homogeneous utility function u_i of arbitrary degree [Arrow et al. 1971].⁹ We note that any homogeneous utility function u_i represents locally non-satiated preferences, since for all $\epsilon > 0$ and $x \in \mathcal{X}$, there exists an allocation $(1+\epsilon/||x||)x$ s.t. $u_i((1+\epsilon/||x||)x) = (1+\epsilon/||x||)u_i(x) > u_i(x)$, and $[x - (1 + \epsilon/||x||)x] \in \mathcal{B}_{\epsilon}(x)$.

The class of homogeneous utility functions includes the well-known *constant elasticity of sub*stitution (CES) utility function family, parameterized by a substitution parameter $-\infty \le \rho_i \le 1$, and given by $u_i(\mathbf{x}_i) = \sqrt[\rho_i]{\sum_{j \in [m]} a_{ij} \mathbf{x}_{ij}^{\rho_i}}$ with each utility function parameterized by the vector of valuations $\mathbf{a}_i \in \mathbb{R}^n_+$, where each a_{ij} quantifies the value of good *j* to buyer *i*. CES utilities are said to be gross substitutes (resp. gross complements) CES if $\rho_i > 0$ ($\rho_i < 0$). Linear utility functions are obtained when ρ is 1 (goods are perfect substitutes), while Cobb-Douglas and Leontief utility

⁸Without loss of generality, we assume that utility functions are positive real-valued functions, since any real-valued function can be made positive real-valued by passing it through the monotonic transformation $\mathbf{x} \mapsto e^{\mathbf{x}}$ without affecting the underlying preference relation.

⁹Throughout this work, without loss of generality, we assume that complete, transitive, continuous, and homothetic preference relations are represented via a homogeneous utility function of degree 1, since any homogeneous utility function of degree *k* can be made homogeneous of degree 1 without affecting the underlying preference relation by passing the utility function through the monotonic transformation $x \mapsto \sqrt[k]{x}$.

functions are obtained when $\rho \to 0$ and $\rho \to -\infty$ (goods are perfect complements), respectively:

Linear:

$$u_i(\mathbf{x}_i) = \sum_{j \in [m]} a_{ij} x_{ij}$$
Cobb-Doulas:

$$u_i(\mathbf{x}_i) = \prod_{j \in [m]} x_{ij}^{a_{ij}}$$
Leontief:

$$u_i(\mathbf{x}_i) = \min_{j:a_{ij} \neq 0} \frac{x_{ij}}{a_{ij}}$$

Associated with any consumption $\mathbf{x} \in X$ are prices $\mathbf{p} \in \mathbb{R}_+^m$ s.t. for all goods $j \in [m]$, $p_j \ge 0$ denotes the price of good j. A demand correspondence $\mathcal{F} : \mathbb{R}_+^m \to X$ takes as input prices $\mathbf{p} \in \mathbb{R}_+^m$ and outputs a set of consumptions $\mathcal{F}(\mathbf{p})$. If \mathcal{F} is singleton-valued for all $\mathbf{p} \in \mathbb{R}_+^m$, then it is called a demand function. Given a demand function f, we define the elasticity $\varepsilon_{f_i,x_j} : \mathbb{R}^m \to \mathbb{R}$ of output $f_i(\mathbf{x})$ w.r.t. the *j*th input x_j evaluated at $\mathbf{x} = \mathbf{y}$ as $\varepsilon_{f_i,x_j}(\mathbf{y}) = \partial_{x_j} f_i(\mathbf{y}) \frac{y_j}{f_i(\mathbf{y})}$.

A good $j \in [m]$ is said to be a substitute (resp. complement) w.r.t. a demand function f for a good $k \in [m] \setminus \{j\}$ if the demand $f_j(p)$ is increasing (resp. decreasing) in p_k . If a buyer's demand $f_j(p)$ for good j is instead weakly increasing (resp. decreasing), good j is said to be a weak substitute (resp. weak complement) for good k.

Next, we define the *consumer functions* [Jehle 2001; Mas-Colell et al. 1995]. The *indirect utility function* $v_i : \mathbb{R}^m_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ takes as input prices p and a budget b_i and outputs the maximum utility the buyer can achieve at that prices within that budget, i.e., $v_i(p, b_i) = \max_{\mathbf{x} \in X: p \cdot \mathbf{x} \le b_i} u_i(\mathbf{x})$.

The Marshallian demand is a correspondence $\mathcal{D}_i : \mathbb{R}^m_+ \times \mathbb{R}_+ \Rightarrow X$ that takes as input prices p and a budget b_i and outputs the utility-maximizing allocations of goods at that budget, i.e., $\mathcal{D}_i(p, b_i) = \arg \max_{\mathbf{x} \in X: p : \mathbf{x} \le b_i} u_i(\mathbf{x}).$

The expenditure function $e_i : \mathbb{R}^m_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ takes as input prices p and a utility level v_i and outputs the minimum amount the buyer must spend to achieve that utility level at those prices, i.e., $e_i(p, v_i) = \min_{x \in X: u_i(x) \ge v_i} p \cdot x$. If the utility function u_i is continuous, then the expenditure function is continuous and homogeneous of degree 1 in p and v_i jointly, non-decreasing in p, strictly increasing in v_i , and concave in p.

The *Hicksian demand* is a correspondence $\mathcal{H}_i : \mathbb{R}^m_+ \times \mathbb{R}_+ \rightrightarrows \mathbb{R}_+$ that takes as input prices \boldsymbol{p} and a utility level v_i and outputs the cost-minimizing allocations of goods at those prices and utility level, i.e., $\mathcal{H}_i(\boldsymbol{p}, v_i) = \arg \min_{\boldsymbol{x} \in \mathcal{X}: u_i(\boldsymbol{x}) \ge v_i} \boldsymbol{p} \cdot \boldsymbol{x}$.

In this paper, we study buyers whose utilities are s.t. Hicksian demand elasticity is well defined, i.e., buyers whose Hicksian demand is singleton-valued, since demands must be unique (and differentiable) for elasticity to be well-defined.¹⁰ As such, $\mathcal{D}_i(\boldsymbol{p}, b_i) = \{\boldsymbol{d}_i(\boldsymbol{p}, b_i)\}$ and $\mathcal{H}_i(\boldsymbol{p}, b_i) = \{\boldsymbol{h}_i(\boldsymbol{p}, b_i)\}$.

Fixing a buyer $i \in [n]$, a good $j \in [m]$ is said to be a gross (resp. net) substitute for a good $k \in [m] \setminus \{j\}$ if it is a substitute w.r.t. Marshallian (resp. Hicksian) demand. If j is instead a weak substitute w.r.t. Marshallian (resp. Hicksian) demand, then it is called a *weak gross (resp. net)* substitute. Gross and net complements, and their weak counterparts, are defined analogously.

Finally, the cross-price elasticity of Marshallian (resp. Hicksian) demand for good j w.r.t. the price of good $k \neq j$ at price p and budget b_i (resp. utility level v_i) is given by $\varepsilon_{d_{ij},p_k}(p, b_i)$ (resp. $\varepsilon_{h_{ij},p_k}(p, v_i)$). If k = j, then we instead have the Hicksian (resp. Marshallian) own-price elasticity of demand.

2.4 Fisher Markets

A *Fisher market* consists of *n* buyers and *m* divisible goods [Brainard et al. 2000]. Each buyer $i \in [n]$ has a budget $b_i \in \mathbb{R}_+$ and a utility function $u_i : \mathbb{R}_+^m \to \mathbb{R}$. As is standard in the literature,

¹⁰The definition of elasticity can be extended to non-unique and non-differentiable demands with additional care (see, for instance, Cheung [2014]). Nonetheless, we present our analysis in this restricted setting, as it is sufficient to capture all convergent and non-convergent behaviors of *tâtonnement*, and allows for a simpler analysis.

we assume there is one unit of each good, and one unit of currency available in the market, i.e. $\sum_{i \in [n]} b_i = 1$ [Codenotti et al. 2007]. An instance of a Fisher market is given by a tuple (n, m, u, b), where $u = (u_1, \ldots, u_n)$, and $b \in \mathbb{R}^n_+$ is the vector of buyer budgets. We abbreviate as (u, b), when n and m are clear from context.

An *allocation* X is a map from goods to buyers, represented as a matrix s.t. $x_{ij} \ge 0$ denotes the amount of good $j \in [m]$ allocated to buyer $i \in [n]$. Goods are assigned *prices* $p \in \mathbb{R}^m_+$. A tuple (X^*, p^*) is said to be a *competitive (or Walrasian) equilibrium* of a Fisher market (u, b) if 1. buyers are utility maximizing constrained by their budget, i.e., $\forall i \in [n], x_i^* \in d_i(p^*, b_i)$; and 2. the market clears, i.e., $\forall j \in [m], p_j^* > 0 \Rightarrow \sum_{i \in [n]} x_{ij}^* = 1$ and $p_j^* = 0 \Rightarrow \sum_{i \in [n]} x_{ij}^* \le 1$.

When the buyers' utility functions in a Fisher market are all of the same type, we qualify the market by the name of the utility function, e.g., a linear Fisher market. A *mixed CES Fisher market* is a Fisher market which comprises CES buyers with possibly different substitution parameters. Considering properties of goods, rather than buyers, a (Fisher) market satisfies *gross substitutes* (resp. *gross complements*) if all pairs of goods in the market are gross substitutes (resp. gross complements). We define net substitute Fisher markets and net complements Fisher markets similarly. We refer the reader to Figure 1a for a summary of the relationships among various Fisher markets. A Fisher market is called *homothetic* if the buyers' utility functions are continuous and homogeneous.

Given a Fisher market (u, b), we define the *aggregate demand* correspondence $q : \mathbb{R}_{+}^{m} \Rightarrow \mathbb{R}_{+}^{m}$ at prices p as the sum of the Marshallian demand at p, given budgets b, i.e., $q(p) = \sum_{i \in [n]} d_i(p, b_i)$. The *excess demand* correspondence $z : \mathbb{R}^{m} \Rightarrow \mathbb{R}^{m}$ of a Fisher market (u, b), which takes as input prices and outputs a set of excess demands at those prices, is defined as the difference between the aggregate demand for and the supply of each good: i.e., $z(p) = q(p) - 1_m$ where 1_m is the vector of ones of size m, and $q(p) - 1_m = \{x - 1_m \mid \forall x \in q(p)\}$.

The *discrete* tâtonnement *process* for Fisher markets is a decentralized, natural price adjustment, defined as:

$$p^{(t+1)} = p^{(t)} + G(g^{(t)}) \qquad \text{for } t = 0, 1, 2, \dots$$
$$g^{(t)} \in z(p^{(t)})$$
$$p^{(0)} \in \mathbb{R}^{m}_{+},$$

where $G : \mathbb{R}^m \to \mathbb{R}^m$ is a coordinate-wise monotonic function such that, for all $j \in [m], \mathbf{x}, \mathbf{y} \in \mathbb{R}^m$, if $x_j \ge y_j$, then $G_j(\mathbf{x}) \ge G_j(\mathbf{y})$. Intuitively, *tâtonnement* is an auction-like process in which the sellers increase (resp. decrease) the prices of goods whenever the demand (resp. supply) is greater than the supply (resp. demand).

Given a sequence of prices $\{\boldsymbol{p}^{(t)}\}_t$, for any $t \in \mathbb{N}_+$, we denote buyer *i*'s Marshallian demand for good *j*, i.e., $d_{ij}(\boldsymbol{p}^{(t)}, b_i)$, by $d_{ij}^{(t)}$; the aggregate demand for good *j*, i.e., $q_j(\boldsymbol{p}^{(t)})$, by $q_j^{(t)}$; and buyer *i*'s Hicksian demand for good *j* at utility level 1, i.e., $h_{ij}(\boldsymbol{p}^{(t)}, 1)$, by $h_{ij}^{(t)}$. Additionally, for any fixed $t \in \mathbb{N}_+$, which will be clear from context, we define $\Delta \boldsymbol{p} \doteq \boldsymbol{p}^{(t+1)} - \boldsymbol{p}^{(t)}$.

2.5 Homothetic Fisher Markets

Suppose that (u, b) is a continuous, concave, and homogeneous Fisher market. The optimal solutions (X^*, p^*) to the primal and dual of *Eisenberg-Gale program* constitute a competitive equilibrium of (u, b) [Devanur et al. 2002; Eisenberg and Gale 1959; Jain et al. 2005]:¹¹

$$\max_{X \in \mathbb{R}_{+}^{n \times m}} \sum_{i \in [n]} b_i \log (u_i(\mathbf{x}_i))$$

subject to
$$\sum_{i \in [n]} x_{ij} \le 1 \quad \forall j \in [m]$$

Dual
$$\min_{\mathbf{p} \in \Delta_m} \sum_{j \in [m]} p_j + \sum_{i \in [n]} [b_i \log (v_i(\mathbf{p}, b_i)) - b_i]$$

Recently, Goktas et al. [2022] proposed a convex program that is equivalent to the Eisenberg-Gale convex program, but whose optimal value differs from that of the Eisenberg-Gale convex program by an additive constant. The authors state their results only for continuous, concave, and homogeneous Fisher markets; however, their proof is valid for *all* homothetic Fisher markets, including those in which the buyers' utility functions are non-concave:

THEOREM 2.2 (GOKTAS ET AL. [2022]). The optimal solutions (X^*, p^*) to the following primal and dual convex programs correspond to competitive equilibrium allocations and prices, respectively, of the homothetic Fisher market (u, b):

I

Primal

Primal

$$\begin{array}{ll}
\max_{X \in \mathbb{R}_{+}^{n \times m}} & \sum_{i \in [n]} \left[b_{i} \log u_{i} \left(\frac{\boldsymbol{x}_{i}}{b_{i}} \right) + b_{i} \right] \\
\text{subject to} & \sum_{i \in [n]} x_{ij} \leq 1 \quad \forall j \in [m] \end{array} \qquad \begin{array}{l}
\text{Dual} \\
\min_{\boldsymbol{p} \in \Delta_{m}} \varphi(\boldsymbol{p}) \doteq \sum_{j \in [m]} p_{j} - \sum_{i \in [n]} b_{i} \log \left(e_{i}(\boldsymbol{p}, 1) \right) \\
\end{array}$$

Since the objective function of the primal in Theorem 2.2 is in general non-concave (i.e., if utilities *u* are not concave), strong duality need not hold; however, the dual is still guaranteed to be convex [Boyd et al. 2004]. This observation suggests that even if the problem of computing competitive equilibrium *allocations* is non-concave, the problem of computing competitive equilibrium *prices* can still be convex. Additionally, since this convex program differs from the Eisenberg-Gale program by an additive constant, we obtain as a corollary that solutions to the Einseberg-Gale program also correspond to competitive equilibria in *all* homothetic Fisher markets, including those in which the buyers' utility functions are non-concave.

An interesting property of this convex program is that its dual expresses competitive equilibrium prices via expenditure functions, and just like the Eisenberg-Gale program's dual objective [Cheung et al. 2013; Devanur et al. 2008], the gradient of its objective $\varphi(\mathbf{p})$ at any price \mathbf{p} is equal to the negative excess demand in the market at those prices.

THEOREM 2.3 (GOKTAS ET AL. [2022]). Given any homothetic Fisher market (u, b), the subdifferential of the dual of the program in Theorem 2.2 at any price p is equal to the negative excess demand in (u, b) at price p: i.e., $\partial_p \varphi(p) = -z(p)$.

Cheung et al. [2013] define a class of markets called *convex potential function (CPF)* markets. A market is a CPF market, if there exists a convex potential function φ such that $\partial_p \varphi(p) = -z(p)$. A corollary of Theorem 2.3 is that all homothetic Fisher markets are CPF markets. This in turn implies that mirror descent on φ over the unit simplex is equivalent to *tâtonnement* in all homothetic Fisher

¹¹The dual as presented here was formulated by Goktas et al. [2022].

markets, for some monotone function G. Using this equivalence, we can pick a particular kernel function h, and then potentially use Theorem 2.1 to establish convergence rates for *tâtonnement*.

Unfortunately, *tâtonnement* does not converge to equilibrium prices in all homothetic Fisher markets, e.g., linear Fisher markets [Cole and Tao 2019], which suggests the need for additional restrictions on the class of homothetic Fisher markets. Goktas et al. [2022] suggest the maximum absolute value of the Marshallian price demand elasticity, i.e., $c = \max_{j,k,i} \max_{(p,b) \in \Delta_m \times \Delta_n \times [n]} \|\varepsilon_{d_{ij},p_k}(p, b_i)\|$, as a possible market parameter to use to establish a convergence rate of $O((^{1+c})/_T)$. However, Cole and Fleischer's [2008] results suggest that it is unlikely that Marshallian demand elasticity could be enough, since the proof techniques used in work that makes this assumption require one to quantify the direction of the change in demand as a function of the change in the prices of the other goods, and hence only apply when one assumes WGS or WGC [Cole and Fleischer 2008].

3 MARKET PARAMETERS

One of the main contributions of this paper is the observation that the maximum absolute value of the price elasticity of Hicksian demand in a homothetic Fisher market is sufficient to analyze the convergence of *tâtonnement*. To this end, in this section, we anaylyze Hicksian demand price elasticity, exposit some of its properties in homothetic markets, and argue why it is a natural parameter to consider in the analysis of *tâtonnement*.

We first note that for Leontief utilities, the Hicksian cross-price elasticity of demand is equal to 0, while for linear utilities the Hicksian cross-price elasticity of demand is, by convention, ∞ .¹² For Cobb-Douglas utilities, the Hicksian cross-price elasticity of demand is strictly positive and upper bounded by 1, but it is not the same for all pairs of goods. Note that the behavior of the Hicksian cross-price elasticity of demand, for the elasticities of linear, Cobb-Douglas, and Leontief utilities are respectively given as ∞ , 0, and $-\infty$. A taxonomy of utility classes as a function of price elasticity of demand Hicksian) is shown in Figure 1b.

We start our analysis with following lemma, which shows that the Hicksian price elasticity of demand is constant across all utility levels in homothetic Fisher markets. This property implies that the Hicksian demand price elasticity at one unit of utility provides sufficient information about the market's reactivity to changes in prices, even without any information about the buyers' utility levels. This information is crucial when trying to bound the changes in Hicksian demand from one iteration of *tâtonnement* to another, since buyers' utilities can change.¹³

LEMMA 3.1. For any Hicksian demand h_i associated with a homogeneous utility function u_i , for all $j, k \in [m], p \in \mathbb{R}^m_+, v_i \in \mathbb{R}_+$, it holds that $\varepsilon_{h_{ij}, p_k}(p, v_i) = \varepsilon_{h_{ij}, p_k}(p, 1) = 1$.

With the above lemma in hand, we now explain why the Hicksian demand price elasticity¹⁴ is a better market parameter by which to analyze the convergence of *tâtonnement* than the Marshallian demand price elasticity. Cheung [2014]; Cheung et al. [2013] use the dual of the Eisenberg-Gale program as a potential to measure the progress that *tâtonnement* makes at each step, for (nested) CES and Leontief utilities. Under these functional forms, the authors are able to explain a change in the value of the buyers' indirect utilities as a function a change in prices, based on which they

¹³We include all omitted results and proofs in Appendix A.

¹²The limit of Hicksian price elasticity of demand as $\rho \to 1$ is not well defined, i.e., if $\rho \to 1^-$ the limit is + ∞ , while if $\rho \to 1^+$ the limit is - ∞ . However, for linear utilities, as the Hicksian demand for a good can only go up when the price of another good goes up, we set the elasticity of Hicksian price elasticity of demand for linear utilities to be + ∞ , by convention. We refer the reader to Ramskov and Munksgaard [2001] for a primer on elasticity of demand.

¹⁴Going forward we refer to the Hicksian price elasticity of demand, as simply Hicksian demand elasticity, because Hicksian price elasticity of demand w.r.t. utility level is always 1.

bound the change in the second term of the dual $\sum_{i \in [n]} b_i \log(v_i(\mathbf{p}, b_i)) - b_i$ from one time period to the next. Using this bound, they show that *tâtonnement* makes steady progress towards equilibrium.

However, in general homothetic Fisher markets, knowing how much the Marshallian demand for each good changes from one iteration of *tâtonnement* to another does not tell us how much the buyers' utilities change. More concretely, suppose that the Marshallian demand of a buyer *i* has changed by an additive vector Δd_i from time *t* to time *t* + 1, then the difference in indirect utilities from one period to another is given by $u_i(d_i^{(t+1)}) - u_i(d_i^{(t)}) = u_i(d_i^{(t)} + \Delta d_i) - u_i(d_i^{(t)})$. Without additional information about the utility functions, e.g., Lipschitz continuity, it is impossible to bound this difference, because utilities can change by an unbounded amount from one period to another. Hence, even if the Marshallian price elasticity of demand and the changes in prices from one period to another were known, it would only allow us to bound the difference in demands, and not the difference in indirect utilities. To get around this difficulty, one could consider making an assumption about the boundedness of the indirect utility function's price elasticity, or the utility function's Lipschitz-continuity, but such assumptions would not be economically justified, since utility functions are merely representations of preference orderings without any inherent meaning of their own.

We can circumvent this issue by instead looking at the dual of the convex program in Theorem 2.2. In this dual, the indirect utility term is replaced by the expenditure function. The advantage of this formulation is that if one knows the amount by which prices change from one iteration to the next, as well as the Hicksian demand elasticity, then we can easily bound the change in spending from one period to another.

The following lemma is crucial to proving the convergence of *tâtonnement*. This lemma allows us to bound the changes in buyer spending across all time periods, thereby allowing us to obtain a global convergence rate. In particular, it shows that the change in spending between two consecutive iterations of *tâtonnement* can be bounded as a function of the prices and the Hicksian demand elasticity.

More formally, suppose that we would like to bound the percentage change in expenditure at one unit of utility from one iteration to another, i.e., $\frac{e_i(\boldsymbol{p}^{(t+1)},1)-e_i(\boldsymbol{p}^{(t)},1)}{e_i(\boldsymbol{p}^{(t)},1)}$, using a first order Taylor expansion of $e_i(\boldsymbol{p}^{(t)} + \Delta \boldsymbol{p}, 1)$ around $\boldsymbol{p}^{(t)}$. By Taylor's theorem [Graves 1927], we have: $e_i(\boldsymbol{p}^{(t)} + \Delta \boldsymbol{p}, 1) = e_i(\boldsymbol{p}^{(t)}, 1) + \langle \nabla_{\boldsymbol{p}} e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \rangle + \frac{1}{2} \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle$ for some $c \in (0, 1)$. Re-organizing terms around, we get $e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1) = \langle \nabla_{\boldsymbol{p}} e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \rangle + \frac{1}{2} \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle$. Dividing both sides by $e_i(\boldsymbol{p}^{(t)}, 1)$ we obtain:

$$\frac{\left\langle \nabla_{\boldsymbol{p}} e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \right\rangle + \frac{1}{2} \left\langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1) \Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{e_i(\boldsymbol{p}^{(t)}, 1)}$$

We can now apply Shephard's lemma [Shephard 2015], a corollary of the envelope theorem [Afriat 1971; Milgrom and Segal 2002], to the numerator, which allows us to conclude that for all buyers $i \in [n], \nabla_{p} e_{i}(p, v_{i}) = h_{i}(p, v_{i})$. Next, using the definition of the expenditure function in the denominator, we obtain the following:

$$=\frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1),\Delta\boldsymbol{p}\right\rangle}{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1),\boldsymbol{p}^{(t)}\right\rangle}+\frac{\frac{1/2}{\left\langle \nabla_{\boldsymbol{p}}^{2}e_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\right\rangle}{e_{i}(\boldsymbol{p}^{(t)},1)}$$
(6)

If the change in prices is bounded, and the Hicksian demand elasticity is known, then one can bound the first term in Equation (6) with ease. It remains to be seen if the second term can be

bounded. The following lemma provides an affirmative answer to that question. In particular, we show that the second-order error term in the Taylor approximation above can be bounded as a function of the maximum absolute value of the Hicksian demand elasticity. We note that in the following lemma, by Lemma A.9, the Marshallian demand is unique, because the Hicksian demand is a singleton for bounded elasticity of Hicksian demand.

LEMMA 3.2. Fix $i \in [n]$ and $t \in \mathbb{N}_+$ and let $\Delta \mathbf{p} = \mathbf{p}^{(t+1)} - \mathbf{p}^{(t)}$. Suppose that $\frac{|\Delta p_j|}{p_j^{(t)}} \leq \frac{1}{4}$, then for all buyers $i \in [n]$, and for some $c \in (0, 1)$, it holds that:

$$\left|\frac{b_i}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{e_i(\boldsymbol{p}^{(t)}, 1)}\right| \le \frac{5\varepsilon}{6} \sum_j \frac{(\Delta p_j)^2}{p_j^{(t)}} d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) , \qquad (7)$$

where $\varepsilon \doteq \max_{\boldsymbol{p} \in \Delta_m, j, k \in [m]} |\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, 1)|.$

PROOF OF LEMMA 3.2. By Shephard's lemma [Shephard 2015] (Lemma A.1, Appendix A), it holds that $\langle \nabla_{p}^{2} e_{i}(p^{(t)} + c\Delta p, 1)\Delta p, \Delta p \rangle = \langle \nabla_{p} h_{i}(p^{(t)} + c\Delta p, 1)\Delta p, \Delta p \rangle$.

$$\begin{aligned} \left| \frac{b_{i}}{2} \frac{\left\langle \nabla_{p}^{2} e_{i}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1)\Delta\boldsymbol{p}, \Delta\boldsymbol{p} \right\rangle}{\left\langle \boldsymbol{h}_{i}^{t}, \boldsymbol{p}^{(t)} \right\rangle} \right| \\ &= \left| \frac{b_{i}}{2} \frac{\left\langle \nabla_{p} \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1)\Delta\boldsymbol{p}, \Delta\boldsymbol{p} \right\rangle}{\left\langle \boldsymbol{h}_{i}^{t}, \boldsymbol{p}^{(t)} \right\rangle} \right| \\ &\leq \frac{b_{i}}{2} \frac{\sum_{j,k} \left| \Delta p_{j} \right| \left| \partial_{p_{k}} \boldsymbol{h}_{ij}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1) \right| \left| \Delta p_{k} \right|}{\left\langle \boldsymbol{h}_{i}^{t}, \boldsymbol{p}^{(t)} \right\rangle} \\ &= \frac{b_{i}}{2} \frac{\sum_{j,k} \left| \Delta p_{j} \right| \sqrt{\left| \partial_{p_{k}} \boldsymbol{h}_{ij}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1) \right| \left\langle \sqrt{\left| \partial_{p_{k}} \boldsymbol{h}_{ij}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1) \right|} \left| \Delta p_{k} \right|}}{\left\langle \boldsymbol{h}_{i}^{t}, \boldsymbol{p}^{(t)} \right\rangle} \\ &= \frac{b_{i}}{2} \frac{\sum_{j,k} \left| \Delta p_{j} \right| \sqrt{\left| \partial_{p_{j}} \boldsymbol{h}_{ik}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1) \right| \left\langle \sqrt{\left| \partial_{p_{k}} \boldsymbol{h}_{ij}(\boldsymbol{p}^{(t)} + c\Delta\boldsymbol{p}, 1) \right|} \left| \Delta p_{k} \right|}}{\left\langle \boldsymbol{h}_{i}^{t}, \boldsymbol{p}^{(t)} \right\rangle} \end{aligned}$$

where the last was obtained from the symmetry of $\nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}, v_i) = \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}, v_i)^T$ for all $i \in [n], \boldsymbol{p} \in \mathbb{R}^m_+$, $v_i \in \mathbb{R}_+$ [Mas-Colell et al. 1995], which combined with Shepherd's lemma gives us $\nabla_{\boldsymbol{p}} \boldsymbol{h}_i(\boldsymbol{p}, v_i) = \nabla_{\boldsymbol{p}} \boldsymbol{h}_i(\boldsymbol{p}, v_i)^T$, i.e., for all $j, k \in [m], \partial_{p_j} h_{ik}(\boldsymbol{p}, v_i) = \partial_{p_k} h_{ij}(\boldsymbol{p}, v_i)$. Define the Hicksian demand elasticity of buyer *i* for good *j* w.r.t. the price of good *k* as $\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, v_i) = \varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, v_i)$.

Define the Hicksian demand elasticity of buyer *i* for good *j* w.r.t. the price of good *k* as $\varepsilon_{h_{ij},p_k}(\boldsymbol{p}, v_i) = \partial_{p_k} h_{ij}(\boldsymbol{p}, v_i) \frac{p_k}{h_{ij}(\boldsymbol{p}, v_i)}$. Since utility functions are homogeneous, by Lemma 3.1 we have for all $v_i \in \mathbb{R}_+$, $\varepsilon_{h_{ij},p_k}(\boldsymbol{p}, v_i) = \partial_{p_k} h_{ij}(\boldsymbol{p}, v_i) \frac{p_k}{h_{ij}(\boldsymbol{p}, v_i)} = \partial_{p_k} h_{ij}(\boldsymbol{p}, 1) \frac{p_k}{h_{ij}(\boldsymbol{p}, 1)}$. Re-organizing expressions, we get $\partial_{p_k} h_{ij}(\boldsymbol{p}, 1) = \varepsilon_{h_{ij},p_k}(\boldsymbol{p}, 1) \frac{h_{ij}(\boldsymbol{p}, 1)}{p_k}$. Going back to Equation (8), we get:

$$=\frac{b_{i}}{2}\frac{\sum_{j,k}\left|\Delta p_{j}\right|\sqrt{\left|\varepsilon_{h_{ik},p_{j}}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)\frac{h_{ik}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)}{p_{j}^{(t)}+c\Delta p_{j}}\right|}\sqrt{\left|\varepsilon_{h_{ij},p_{k}}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)\frac{h_{ij}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)}{p_{k}^{(t)}+c\Delta p_{k}}\right|}\left|\Delta p_{k}\right|}{\left\langle\boldsymbol{h}_{i}^{t},\boldsymbol{p}^{(t)}\right\rangle}$$

Denizalp Goktas, Jiayi Zhao, and Amy Greenwald

$$=\frac{b_{i}}{2}\frac{\sum_{j,k}\left|\Delta p_{j}\right|\sqrt{\left|\varepsilon_{h_{ik},p_{j}}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)\right|\frac{h_{ik}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)}{p_{j}^{(t)}+c\Delta p_{j}}}\sqrt{\left|\varepsilon_{h_{ij},p_{k}}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)\right|\frac{h_{ij}\left(\boldsymbol{p}^{(t)}+c\Delta \boldsymbol{p},1\right)}{p_{k}^{(t)}+c\Delta p_{k}}\left|\Delta p_{k}\right|}}{\left\langle h_{i}(\boldsymbol{p}^{(t)},1),\boldsymbol{p}^{(t)}\right\rangle}$$

Letting $\varepsilon = \max_{\boldsymbol{p} \in \mathbb{R}^m_+, v_i \in \mathbb{R}_+, j, k \in [m]} |\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, v_i)|$. Note that since utility functions are homogeneous, by Lemma 3.1 we have $\varepsilon = \max_{\boldsymbol{p} \in \mathbb{R}^m_+, v_i \in \mathbb{R}_+, j, k \in [m]} |\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, v_i)| = \max_{\boldsymbol{p} \in \mathbb{R}^m_+, j, k \in [m]} |\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, 1)|$, which gives us:

$$\leq \frac{\varepsilon b_i}{2} \frac{\sum_{j,k} \left| \Delta p_j \right| \sqrt{\frac{h_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)}{p_j^{(t)} + c\Delta p_j}} \sqrt{\frac{h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)}{p_k^{(t)} + c\Delta p_k}} \left| \Delta p_k \right|}{\left\langle \boldsymbol{h}_i(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle}$$

Since for all $j \in [m]$, $\frac{|\Delta p_j|}{p_j} \le \frac{1}{4}$, we have that for all $j \in [m]$ and for all $c \in [0, 1]$, $p_j^{(t)} + c\Delta p_j \ge \frac{3}{4}p_j^{(t)}$, which gives:

$$\leq \frac{\varepsilon b_{i}}{2} \frac{\sum_{j,k} \left| \Delta p_{j} \right| \sqrt{\frac{h_{ik}(\boldsymbol{p}^{(t)} + c\Delta p, 1)}{\frac{3}{4}p_{j}^{(t)}}} \sqrt{\frac{h_{ij}(\boldsymbol{p}^{(t)} + c\Delta p, 1)}{\frac{3}{4}p_{k}^{(t)}}} \left| \Delta p_{k} \right| }{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle} \\ = \frac{2\varepsilon b_{i}}{3} \frac{\sum_{j,k} \left| \Delta p_{j} \right| \sqrt{\frac{h_{ij}(\boldsymbol{p}^{(t)} + c\Delta p, 1)}{p_{j}^{(t)}}} \sqrt{\frac{h_{ik}(\boldsymbol{p}^{(t)} + c\Delta p, 1)}{p_{k}^{(t)}}} \left| \Delta p_{k} \right| }{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle} \\ = \frac{2\varepsilon b_{i}}{3} \frac{\sum_{j,k \in [m]} \sqrt{\frac{\left| \Delta p_{j} \right|^{2}}{p_{j}^{(t)}}} h_{ij}(\boldsymbol{p}^{(t)} + c\Delta p, 1) h_{ik}(\boldsymbol{p}^{(t)} + c\Delta p, 1) \frac{\left| \Delta p_{k} \right|^{2}}{p_{k}^{(t)}}}{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle}$$

Applying the AM-GM inequality, i.e., for all $x, y \in \mathbb{R}_+$, $\frac{x+y}{2} \ge \sqrt{xy}$, to the sum inside the numerator above, we obtain:

$$\leq \frac{2\varepsilon b_i}{3} \frac{\sum_{j,k\in[m]} \frac{1}{2} \left(\frac{|\Delta p_j|^2}{p_j^{(t)}} h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1) + h_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1) \frac{|\Delta p_k|^2}{p_k^{(t)}} \right)}{\left\langle h_i(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle} \\ \leq \frac{2\varepsilon b_i}{3} \frac{\sum_j \frac{(\Delta p_j)^2}{p_j^{(t)}} h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)}{\left\langle h_i(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle}$$

Since for all $j \in [m]$, $\frac{|\Delta p_j|}{p_j^{(t)}} \le \frac{1}{4}$, we have for all $c \in [0, 1]$ that $\frac{4}{5} \sum_j h_{ij}(\mathbf{p}^{(t)}, 1)(p_j^{(t)} + c\Delta p_j) \le \sum_j h_{ij}(\mathbf{p}^{(t)}, 1)p_j^{(t)}$:

$$\leq \frac{2\varepsilon b_i}{3} \frac{\sum_j \frac{(\Delta p_j)^2}{p_j^{(t)}} h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)}{\frac{4}{5} \sum_j h_{ij}(\boldsymbol{p}^{(t)}, 1)(p_j^{(t)} + c\Delta \boldsymbol{p}_j)}$$
$$= \frac{5\varepsilon b_i}{6} \frac{\sum_j \frac{(\Delta p_j)^2}{p_j^{(t)}} h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)}{\sum_j h_{ij}(\boldsymbol{p}^{(t)}, 1)(p_j^{(t)} + c\Delta \boldsymbol{p}_j)}$$

$$\leq \frac{5\varepsilon b_i}{6} \frac{\sum_j \frac{(\Delta p_j)^2}{p_j^{(t)}} h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)}{\sum_j h_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)(p_j^{(t)} + c\Delta p_j)}$$
$$= \frac{5\varepsilon}{6} \sum_j \frac{(\Delta p_j)^2}{p_j^{(t)}} d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i)$$

(Corollary A.7, Appendix A) (Lemma A.9, Appendix A)

Because we can bound the change in the expenditure function from one iteration of *tâtonnement* to the next, the Hicksian price elasticity of demand is a better tool with which to analyze the convergence of *tâtonnement* than Marshallian price elasticity of demand. Additionally, as shown previously by Cheung et al. [2013] (Lemma A.3, Appendix A), we can further upper bound the price terms in Lemma 3.2 by the KL divergence between the two prices In light of Theorem 2.1, this result suggests that running mirror descent with KL divergence as the Bregman divergence on the dual of the convex program in Theorem 2.2 could result in a *tâtonnement* update rule that converges to a competitive equilibrium.

4 CONVERGENCE BOUNDS FOR ENTROPIC TÂTONNEMENT

In this section, we analyze the rate of convergence of *entropic tâtonnement*, which corresponds to the *tâtonnement* process given by mirror descent with weighted entropy as the kernel function, i.e., entropic descent. This particular update rule reduces to Equations (4) to (5), and has been the focus of previous work [Cheung et al. 2013]. We provide a sketch of the proof used to obtain our convergence rate in this section. The omitted lemmas and proofs can be found in Appendix A.

At a high level, our proof follows Cheung et al.'s [2013] proof technique for Leontief Fisher markets [Cheung et al. 2013], although we encounter different lower-level technical challenges in generalizing to homothetic markets. This proof technique works as follows. First, we prove that under certain assumptions, the condition required by Theorem 2.1 holds when f is the convex potential function for homothetic Fisher markets, i.e., $f = \varphi$. For these assumptions to be valid, we need to set γ to be greater than a quadratic function of the maximum absolute value of the price elasticity of the Hicksian demand and the maximum Marshallian demand, for all goods throughout the *tâtonnment* process. Further, since γ needs to be set at the outset, we need to upper bound γ . To do so, we derive a bound on the maximum demand for any good during *tâtonnement* in all homothetic Fisher markets, which in turn allows us to derive an upper bound on γ . Finally, we use Theorem 2.1 to obtain the convergence rate of $O(1+\varepsilon^2/t)$.

The following lemma derives the conditions under which the antecedent of Theorem 2.1 holds for entropic *tâtonnement*.

LEMMA 4.1. Consider a homothetic Fisher market $(\boldsymbol{u}, \boldsymbol{b})$ and let $\varepsilon = \max_{\boldsymbol{p} \in \Delta_m, j, k \in [m]} |\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, 1)|$. Then, the following holds for entropic tâtonnement when run on $(\boldsymbol{u}, \boldsymbol{b})$: for all $t \in \mathbb{N}$,

$$\varphi(p^{(t+1)}) \le \ell_{\varphi}(p^{(t+1)}, p^{(t)}) + \gamma \delta_{\mathrm{KL}}(p^{(t+1)}, p^{(t)})$$

where $\gamma = \max_{j \in [m], t \in \mathbb{N}_+, c \in [0,1]} \left\{ 1 + q_j (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, \boldsymbol{b}) \right\} \left(6 + \frac{85\varepsilon}{12} + \frac{25\varepsilon^2}{72} \right).$

Proof of Lemma 4.1.

$$\varphi(\boldsymbol{p}^{(t+1)}) - \ell_{\varphi}(\boldsymbol{p}^{(t+1)}, \boldsymbol{p}^{(t)})$$

= $\varphi(\boldsymbol{p}^{(t+1)}) - \varphi(\boldsymbol{p}^{(t)}) + \boldsymbol{z}(\boldsymbol{p}^{(t)}) \cdot \left(\boldsymbol{p}^{(t+1)} - \boldsymbol{p}^{(t)}\right)$

Denizalp Goktas, Jiayi Zhao, and Amy Greenwald

$$\begin{split} &= \sum_{j \in [m]} \left(p_j^{(t)} + \Delta p_j \right) - \sum_{i \in [n]} b_i \log \left(e_i(\boldsymbol{p}^{(t+1)}, 1) \right) - \sum_{j \in [m]} p_j^{(t)} + \sum_{i \in [n]} b_i \log \left(e_i(\boldsymbol{p}^{(t)}, 1) \right) + \sum_{j \in [m]} z_j(\boldsymbol{p}^{(t)}) \Delta p_j \\ &= \sum_{j \in [m]} \left(p_j^{(t)} + \Delta p_j \right) - \sum_{i \in [n]} b_i \log \left(e_i(\boldsymbol{p}^{(t+1)}, 1) \right) - \sum_{j \in [m]} p_j^{(t)} + \sum_{i \in [n]} b_i \log \left(e_i(\boldsymbol{p}^{(t)}, 1) \right) + \sum_{j \in [m]} (q_j^{(t)} - 1) \Delta p_j \\ &= \sum_{j \in [m]} \Delta p_j q_j^{(t)} - \sum_{i \in [n]} b_i \log \left(e_i(\boldsymbol{p}^{(t+1)}, 1) \right) + \sum_{i \in [n]} b_i \log \left(e_i(\boldsymbol{p}^{(t)}, 1) \right) \\ &= \left\langle \Delta \boldsymbol{p}, \boldsymbol{q}^{(t)} \right\rangle + \sum_{i \in [n]} b_i \log \left(\frac{e_i(\boldsymbol{p}^{(t)}, 1)}{e_i(\boldsymbol{p}^{(t+1)}, 1)} \right) \\ &= \left\langle \Delta \boldsymbol{p}, \boldsymbol{q}^{(t)} \right\rangle + \sum_{i \in [n]} b_i \log \left(\frac{e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1)}{e_i(\boldsymbol{p}^{(t)}, 1) + e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1)} \right) \\ &= \left\langle \Delta \boldsymbol{p}, \boldsymbol{q}^{(t)} \right\rangle + \sum_{i \in [n]} b_i \log \left(1 - \frac{e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1)}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(1 + \frac{e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1)}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^{-1} \right) \end{split}$$

where the last line is obtained by simply noting that $\forall a, b \in \mathbb{R}, \frac{a}{a+b} = 1 - \frac{b}{a}(1 + \frac{b}{a})^{-1}$. Using Lemma A.12, we then obtain:

$$\begin{split} \varphi(\boldsymbol{p}^{(t+1)}) &= \ell_{\varphi}(\boldsymbol{p}^{(t+1)}, \boldsymbol{p}^{(t)}) \\ &\leq \left\langle \Delta \boldsymbol{p}, \boldsymbol{q}^{(t)} \right\rangle + \sum_{i \in [n]} \left[\left(\frac{4}{3} + \frac{20\varepsilon}{27} \right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} + \left(\frac{5\varepsilon}{6} + \frac{25\varepsilon^{2}}{324} \right) \sum_{l \in [m]} \frac{d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})}{p_{l}^{(t)}} (\Delta p_{l})^{2} - \left\langle \boldsymbol{d}_{i}^{(t)}, \Delta \boldsymbol{p} \right\rangle \right] \\ &= \left\langle \Delta \boldsymbol{p}, \boldsymbol{q}^{(t)} \right\rangle + \left(\frac{4}{3} + \frac{20\varepsilon}{27} \right) \sum_{l \in [m]} \frac{q_{l}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} + \left(\frac{5\varepsilon}{6} + \frac{25\varepsilon^{2}}{324} \right) \sum_{l \in [m]} \frac{q_{l}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, \boldsymbol{b})}{p_{l}^{(t)}} (\Delta p_{l})^{2} - \left\langle \boldsymbol{q}^{(t)}, \Delta \boldsymbol{p} \right\rangle \\ &= \left(\frac{4}{3} + \frac{20\varepsilon}{27} \right) \sum_{l \in [m]} \frac{q_{l}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} + \left(\frac{5\varepsilon}{6} + \frac{25\varepsilon^{2}}{324} \right) \sum_{l \in [m]} \frac{q_{l}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, \boldsymbol{b})}{p_{l}^{(t)}} (\Delta p_{l})^{2} \\ &\leq \left(\frac{4}{3} + \frac{20\varepsilon}{27} \right) \sum_{l \in [m]} q_{l}^{(t)} \left(\frac{9}{2} \right) \delta_{\mathrm{KL}}(p_{j}^{(t)} + \Delta p_{j}, p_{j}^{(t)}) + \left(\frac{5\varepsilon}{6} + \frac{25\varepsilon^{2}}{324} \right) \sum_{l \in [m]} q_{l}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, \boldsymbol{b}) \left(\frac{9}{2} \right) \delta_{\mathrm{KL}}(p_{j}^{(t)} + \Delta p_{j}, p_{j}^{(t)}) \end{split}$$

where the last line follows from Lemma A.3. Continuing,

$$= \left(6 + \frac{10\varepsilon}{3}\right) \sum_{l \in [m]} q_l^{(t)} \delta_{\mathrm{KL}}(p_l^{(t)} + \Delta p_l, p_l^{(t)}) + \left(\frac{15\varepsilon}{4} + \frac{25\varepsilon^2}{72}\right) \sum_{l \in [m]} q_l(\mathbf{p}^{(t)} + c\Delta \mathbf{p}, \mathbf{b}) \delta_{\mathrm{KL}}(p_l^{(t)} + \Delta p_l, p_l^{(t)})$$

$$\le \max_{\substack{j \in [m], \\ t \in \mathbb{N}_+ \\ c \in [0,1]}} \left\{q_j(\mathbf{p}^{(t)} + c\Delta \mathbf{p})\right\} \left(6 + \frac{10\varepsilon}{3}\right) \sum_{l \in [m]} \delta_{\mathrm{KL}}(p_l^{(t)} + \Delta p_l, p_l^{(t)}) + \sum_{\substack{l \in [m], \\ t \in \mathbb{N}_+ \\ c \in [0,1]}} \left\{q_j(\mathbf{p}^{(t)} + c\Delta \mathbf{p})\right\} \left(\frac{15\varepsilon}{4} + \frac{25\varepsilon^2}{72}\right) \sum_{l \in [m]} \delta_{\mathrm{KL}}(p_l^{(t)} + \Delta p_l, p_l^{(t)})$$

$$\le \max_{\substack{j \in [m], \\ t \in \mathbb{N}_+, \\ c \in [0,1]}} \left\{q_j(\mathbf{p}^{(t)} + c\Delta \mathbf{p})\right\} \left(6 + \frac{85\varepsilon}{12} + \frac{25\varepsilon^2}{72}\right) \delta_{\mathrm{KL}}(\mathbf{p}^{(t)} + \Delta \mathbf{p}, \mathbf{p}^{(t)})$$

Combining Lemma 4.1 with Theorem 2.1, we obtain our main result, namely a worst-case convergence rate of $O((1+\epsilon^2)/t)$ for entropic *tâtonnement* in homothetic Fisher markets.

THEOREM 4.2. Suppose $(\boldsymbol{u}, \boldsymbol{b})$ is homothetic Fisher market and $\varepsilon = \max_{\boldsymbol{p} \in \Delta_m, j, k \in [m]} |\varepsilon_{h_{ij}, p_k}(\boldsymbol{p}, 1)|$. Then, the following holds for entropic tâtonnement: for all $t \in \mathbb{N}$,

$$\varphi(\boldsymbol{p}^{(t)}) - \varphi(\boldsymbol{p}^*) \le \frac{\gamma \delta_{\mathrm{KL}}(\boldsymbol{p}^*, \boldsymbol{p}^0)}{t} , \qquad (8)$$

where $\gamma = \max_{j \in [m], t \in \mathbb{N}_+, c \in [0,1]} \left\{ q_j(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, \boldsymbol{b}) \right\} \left(6 + \frac{85\varepsilon}{12} + \frac{25\varepsilon^2}{72} \right) \delta_{\mathrm{KL}}(\boldsymbol{p}^{(t)} + \Delta \boldsymbol{p}, \boldsymbol{p}^{(t)}).$

Note, however, that for Theorem 4.2 to hold, the step size γ needs to be known in advance, so that it can be set at the start of the entropic *tâtonnement* process. In other words, we need to upper bound the aggregate demand for all goods all throughout tâtonnement. If the number of iterations *T* for which *tâtonnement* is to be run were known at the outset, then one could use a naive bound of $\max_{t \in \mathbb{N}} q_j^{(t)} \leq e^{T/5} \frac{\sum_{i \in [n]} b_i}{\min_{j \in [m]} p_j^0}$. This bound follows by induction, since for the first iteration the prices of the goods cannot change by more than $|\Delta p_j/p_j^{(t)}| \leq \frac{1}{4}$, and then, for all other iterations, the price for a good can decrease at most by a factor of $e^{-1/5}$ (Lemma A.2), ensuring that the learning rate is greater than the aggregate demand in the next iteration. This time-dependent upper bound could then be combined with the doubling trick to extend the convergence result to an unknown time horizon [Auer et al. 1995]. In addition, for markets with homothetic net complements, we provide the following bound, which holds even when the *tâtonnement* time horizon is not known. We note that if better upper bounds are found, they can be used to improve our convergence result by a constant factor.

LEMMA 4.3 (STEP SIZE FOR MARKETS WITH HOMOTHETIC NET COMPLEMENTS). If tâtonnement is run on a net complements Fisher market $(\boldsymbol{u}, \boldsymbol{b})$, then, for all $t \in \mathbb{N}_+$, prices satisfy $\left|\frac{\Delta p_j}{p_j^{(t)}}\right| \leq \frac{1}{4}$ if and only if $\gamma \geq \max\left\{d_{ij}^{(0)}, \max_{\boldsymbol{p} \in \Delta_m} \max_{k \in [m]: h_{ik}(\boldsymbol{p}, 1) > 0} \left\{\frac{h_{ij}(\boldsymbol{p}, 1)}{h_{ik}(\boldsymbol{p}, 1)}\right\}\right\}$.

5 CONCLUSION

We identified the maximum absolute value of the Hicksian price elasticity of demand as a sufficient parameter by which to analyze convergent and non-convergent behavior of *tâtonnement* in homothetic Fisher markets. We then showed that together with the KL divergence associated with a change in prices, we can use it to bound the percentage change in the expenditure of one unit of utility, assuming bounded price changes. This observation motivated us to consider analyzing the convergence of mirror descent with KL divergence on a recently proposed [Goktas et al. 2022] convex potential, making use of the expenditure function to characterize competitive equilibrium prices in homothetic Fisher markets. An important property of this convex potential is that its gradient is equal to the negative excess demand in the market, implying that mirror descent on it is equivalent to *tâtonnement*, an observation used to prove previous convergence results regarding tâtonnement in Fisher markets [Cheung et al. 2013]. Using the bound we derived on the change in the expenditure function as a function of the change in prices, we then showed that the potential function we considered is Bregman-smooth w.r.t. the KL divergence throughout a trajectory of tâtonnement. Combining this result with the sublinear convergence rate of mirror descent for Bregman-smooth functions [Birnbaum et al. 2011], we concluded that tâtonnement converges at a rate of $O((1+\varepsilon^2)/T)$, where ε is the maximum absolute value of the price elasticity of Hicksian demand across all buyers. Our result not only generalizes existing convergence results for CES and nested CES Fisher markets, but extends them beyond Fisher markets with concave utility functions. Our convergence rate covers the full spectrum of (nested) CES utilities, including Leontief and linear utilities, unifying previously existing disparate convergence and non-convergence results. In particular, for $\varepsilon = 0$, i.e., Leontief markets, we recover the best-known convergence rate of O(1/T)[Cheung et al. 2013], and as $\varepsilon \to \infty$, i.e., linear Fisher markets, we obtain the non-convergent behavior of *tâtonnement*.

Future work could investigate the space of homogeneous utility functions with negative crossprice elasticity of Hicksian demand to possibly derive faster convergence rates than those provided in this paper. Additionally, it remains to be seen if the bound we have provided in this paper is tight; the greatest lower bound known for the convergence of *tâtonnement* in homothetic Fisher markets is $O(1/t^2)$ for Leontief markets [Cheung et al. 2012], leaving space for improvement. Finally, Lemma 3.1 suggests that to extend convergence results for *tâtonnement* beyond homothetic domains, one might have to consider the Hicksian demand elasticity *w.r.t. utility level* rather than price.

ACKNOWLEDGMENTS

We would like to thank Richard Cole, Yun Kuen Cheung, and Yixin Tao for very carefully reading an earlier version of this paper and providing invaluable feedback. Their efforts helped us refine the results in the paper. Denizalp Goktas was supported by a JP Morgan AI Fellowship.

REFERENCES

- S. N. Afriat. 1971. Theory of Maxima and the Method of Lagrange. SIAM J. Appl. Math. 20, 3 (1971), 343-357. http://www.jstor.org/stable/2099955
- Gabriel Andrade, Rafael Frongillo, Sharadha Srinivasan, and Elliot Gorokhovsky. 2021. Graphical Economies with Resale. *Proceedings of the 22nd ACM Conference on Economics and Computation* (Jul 2021). https://doi.org/10.1145/3465456.3467628
- Kenneth Arrow. 1951a. An extension of the basic theorems of classical welfare economics. In *Proceedings of the second Berkeley symposium on mathematical statistics and probability.* The Regents of the University of California.
- Kenneth Arrow and Gerard Debreu. 1954. Existence of an equilibrium for a competitive economy. *Econometrica: Journal of the Econometric Society* (1954), 265–290.
- Kenneth J Arrow. 1951b. An extension of the basic theorems of classical welfare economics. In *Proceedings of the second* Berkeley symposium on mathematical statistics and probability, Vol. 2. University of California Press, 507–533.
- Kenneth Joseph Arrow, Frank Hahn, et al. 1971. General competitive analysis. (1971).
- Kenneth J. Arrow and Leonid Hurwicz. 1958. On the Stability of the Competitive Equilibrium, I. *Econometrica* 26, 4 (1958), 522–552. http://www.jstor.org/stable/1907515
- Kenneth J Arrow and Marc Nerlove. 1958. A note on expectations and stability. *Econometrica: Journal of the Econometric Society* (1958), 297–305.
- Peter Auer, Nicolo Cesa-Bianchi, Yoav Freund, and Robert E Schapire. 1995. Gambling in a rigged casino: The adversarial multi-armed bandit problem. In *Proceedings of IEEE 36th annual foundations of computer science*. IEEE, 322–331.
- Yves Balasko. 1975. Some results on uniqueness and on stability of equilibrium in general equilibrium theory. Journal of Mathematical Economics 2, 2 (1975), 95–118.
- Xiaohui Bei, Jugal Garg, and Martin Hoefer. 2015. Tatonnement for linear and gross substitutes markets. CoRR abs/1507.04925 (2015).
- Benjamin Birnbaum, Nikhil R. Devanur, and Lin Xiao. 2011. Distributed Algorithms via Gradient Descent for Fisher Markets. In Proceedings of the 12th ACM Conference on Electronic Commerce (San Jose, California, USA) (EC '11). Association for Computing Machinery, New York, NY, USA, 127–136. https://doi.org/10.1145/1993574.1993594
- Lawrence E. Blume. 2017. Duality. The New Palgrave Dictionary of Economics (2017), 1-7. https://doi.org/10.1057/978-1-349-95121-5_285-2
- KC Border. 2017. Euler's theorem for homogeneous functions. *Caltech Division of the Humanities and Social Sciences* 27 (2017), 16–34.
- Stephen Boyd, Stephen P Boyd, and Lieven Vandenberghe. 2004. Convex optimization. Cambridge university press.

William C Brainard, Herbert E Scarf, et al. 2000. How to compute equilibrium prices in 1891. Citeseer.

Simina Brânzei, Nikhil Devanur, and Yuval Rabani. 2021. Proportional dynamics in exchange economies. In Proceedings of the 22nd ACM Conference on Economics and Computation. 180–201.

Lev M Bregman. 1967. The relaxation method of finding the common point of convex sets and its application to the solution of problems in convex programming. USSR computational mathematics and mathematical physics 7, 3 (1967), 200–217.

Yun Kuen Cheung. 2014. Analyzing tatonnement dynamics in economic markets. Ph. D. Dissertation. New York University.

- Yun Kuen Cheung and Richard Cole. 2018. Amortized analysis of asynchronous price dynamics. arXiv preprint arXiv:1806.10952 (2018).
- Yun Kuen Cheung, Richard Cole, and Nikhil Devanur. 2013. Tatonnement beyond Gross Substitutes? Gradient Descent to the Rescue. In Proceedings of the Forty-Fifth Annual ACM Symposium on Theory of Computing (Palo Alto, California, USA) (STOC '13). Association for Computing Machinery, New York, NY, USA, 191–200. https://doi.org/10.1145/2488608.2488633
- Yun Kuen Cheung, Richard Cole, and Ashish Rastogi. 2012. Tatonnement in ongoing markets of complementary goods. In *Proceedings of the 13th ACM Conference on Electronic Commerce*. 337–354.
- Yun Kuen Cheung, Richard Cole, and Yixin Tao. 2018. Dynamics of distributed updating in Fisher markets. In Proceedings of the 2018 ACM Conference on Economics and Computation. 351–368.
- Yun Kuen Cheung, Martin Hoefer, and Paresh Nakhe. 2019. Tracing Equilibrium in Dynamic Markets via Distributed Adaptation. In Proceedings of the 18th International Conference on Autonomous Agents and MultiAgent Systems. 1225–1233.
- Bruno Codenotti, Benton McCune, and Kasturi Varadarajan. 2005. Market equilibrium via the excess demand function. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*. 74–83.
- Bruno Codenotti, Kasturi Varadarajan, Noam Nisan, Tim Roughgarden, Eva Tardos, and Vijay Vazirani. 2007. Algorithmic game theory. Cambridge university press, Chapter 5 and 6.
- Richard Cole and Lisa Fleischer. 2008. Fast-converging tatonnement algorithms for one-time and ongoing market problems. In Proceedings of the fortieth annual ACM symposium on Theory of computing. 315–324.
- Richard Cole, Lisa Fleischer, and Ashish Rastogi. 2010. Discrete Price Updates Yield Fast Convergence in Ongoing Markets with Finite Warehouses. arXiv:1012.2124 [cs.GT]

Richard Cole and Yixin Tao. 2019. Balancing the Robustness and Convergence of Tatonnement. arXiv:1908.00844 [cs.GT]

- Gerard Debreu. 1951. The Coefficient of Resource Utilization. *Econometrica* 19, 3 (1951), 273–292. http://www.jstor.org/stable/1906814
- N. R. Devanur, C. H. Papadimitriou, A. Saberi, and V. V. Vazirani. 2002. Market equilibrium via a primal-dual-type algorithm. In *The 43rd Annual IEEE Symposium on Foundations of Computer Science, 2002. Proceedings.* 389–395. https://doi.org/10.1109/SFCS.2002.1181963
- Nikhil R Devanur, Christos H Papadimitriou, Amin Saberi, and Vijay V Vazirani. 2008. Market equilibrium via a primal-dual algorithm for a convex program. *Journal of the ACM (JACM)* 55, 5 (2008), 1–18.
- Egbert Dierker. 1982. Chapter 17 Regular economies. Handbook of Mathematical Economics, Vol. 2. Elsevier, 795–830. https://doi.org/10.1016/S1573-4382(82)02012-8
- Edmund Eisenberg. 1961. Aggregation of utility functions. Management Science 7, 4 (1961), 337-350.
- Edmund Eisenberg and David Gale. 1959. Consensus of subjective probabilities: The pari-mutuel method. *The Annals of Mathematical Statistics* 30, 1 (1959), 165–168.
- Francisco Facchinei and Christian Kanzow. 2010. Generalized Nash equilibrium problems. Annals of Operations Research 175, 1 (2010), 177–211.
- Franklin M Fisher. 1975. The stability of general equilibrium: Results and problems. 1976), Essays in Economic Analysis: Proceedings of the Association of University Teachers of Economics Sheffield (1975), 3–29.
- Yuan Gao and Christian Kroer. 2020. First-order methods for large-scale market equilibrium computation. Advances in Neural Information Processing Systems 33 (2020), 21738–21750.
- Rahul Garg and Sanjiv Kapoor. 2004. Auction algorithms for market equilibrium. In Proceedings of the thirty-sixth annual ACM symposium on Theory of computing. 511–518.
- Benjamin J Gillen, Masayoshi Hirota, Ming Hsu, Charles R Plott, and Brian W Rogers. 2020. Divergence and convergence in Scarf cycle environments: experiments and predictability in the dynamics of general equilibrium systems. *Economic Theory* (2020), 1–52.
- Denizalp Goktas, Enrique Areyan Viqueira, and Amy Greenwald. 2022. A Consumer-Theoretic Characterization of Fisher Market Equilibria. In Web and Internet Economics: 17th International Conference, WINE 2021, Potsdam, Germany, December 14–17, 2021, Proceedings. Springer, 334–351.
- Noah Golowich, Sarath Pattathil, Constantinos Daskalakis, and Asuman E. Ozdaglar. 2020. Last Iterate is Slower than Averaged Iterate in Smooth Convex-Concave Saddle Point Problems. *CoRR* abs/2002.00057 (2020). arXiv:2002.00057 https://arxiv.org/abs/2002.00057
- Lawrence M Graves. 1927. Riemann integration and Taylor's theorem in general analysis. *Trans. Amer. Math. Soc.* 29, 1 (1927), 163–177.

Frank H Hahn. 1958. Gross substitutes and the dynamic stability of general equilibrium. *Econometrica (pre-1986)* 26, 1 (1958), 169.

P Jean-Jacques Herings. 1997. A globally and universally stable price adjustment process. Journal of Mathematical Economics 27, 2 (1997), 163–193.

Kamal Jain, Vijay V Vazirani, and Yinyu Ye. 2005. Market equilibria for homothetic, quasi-concave utilities and economies of scale in production. In SODA, Vol. 5. 63–71.

Shweta Jain, Narayanaswamy Balakrishnan, Yadati Narahari, Saiful A Hussain, and Nyuk Yoong Voo. 2013. Constrained tâtonnement for fast and incentive compatible distributed demand management in smart grids. In *Proceedings of the fourth international conference on Future energy systems*. 125–136.

Geoffrey Alexander Jehle. 2001. Advanced microeconomic theory. Pearson Education India.

James M Joyce. 2011. Kullback-leibler divergence. In International encyclopedia of statistical science. Springer, 720-722.

Sham M Kakade, Michael Kearns, and Luis E Ortiz. 2004. Graphical economics. In COLT. Springer, 17-32.

Nicholas Kaldor. 1934. A classificatory note on the determinateness of equilibrium. *The review of economic studies* 1, 2 (1934), 122–136.

Kazuya Kamiya. 1990. A globally stable price adjustment process. *Econometrica: Journal of the Econometric Society* (1990), 1481–1485.

Galina M Korpelevich. 1976. The extragradient method for finding saddle points and other problems. *Matecon* 12 (1976), 747–756.

Stefanos Leonardos, Barnabé Monnot, Daniël Reijsbergen, Efstratios Skoulakis, and Georgios Piliouras. 2021. Dynamical analysis of the eip-1559 ethereum fee market. In *Proceedings of the 3rd ACM Conference on Advances in Financial Technologies*. 114–126.

Jonathan Levin. 2004. Lecture notes on Consumer Theory.

Yulin Liu, Yuxuan Lu, Kartik Nayak, Fan Zhang, Luyao Zhang, and Yinhong Zhao. 2022. Empirical analysis of eip-1559: Transaction fees, waiting time, and consensus security. arXiv preprint arXiv:2201.05574 (2022).

Andreu Mas-Colell, Michael D. Whinston, and Jerry R. Green. 1995. Microeconomic Theory. Number 9780195102680 in OUP Catalogue. Oxford University Press. https://ideas.repec.org/b/oxp/obooks/9780195102680.html

Lionel W McKenzie. 1959. On the existence of general equilibrium for a competitive market. *Econometrica: journal of the Econometric Society* (1959), 54–71.

Lionel W McKenzie et al. 2005. Classical general equilibrium theory. MIT Press Books 1 (2005).

Paul Milgrom and Ilya Segal. 2002. Envelope theorems for arbitrary choice sets. Econometrica 70, 2 (2002), 583-601.

John F. Nash. 1950. Equilibrium points in <i>n</i>-person games. *Proceedings of the National Academy of Sciences* 36, 1 (1950), 48–49. https://doi.org/10.1073/pnas.36.1.48 arXiv:https://www.pnas.org/doi/pdf/10.1073/pnas.36.1.48

Hukukane Nikaidô and Hirofumi Uzawa. 1960. Stability and non-negativity in a Walrasian Tatonnement process. *International Economic Review* 1, 1 (1960), 50–59.

Christos H Papadimitriou and Mihalis Yannakakis. 2010. An impossibility theorem for price-adjustment mechanisms. *Proceedings of the National Academy of Sciences* 107, 5 (2010), 1854–1859.

I. F. Pearce and J. Wise. 1973. On the Uniqueness of Competitive Equilibrium: Part I, Unbounded Demand. *Econometrica* 41, 5 (1973), 817–828. http://www.jstor.org/stable/1913808

Jacob Ramskov and Jesper Munksgaard. 2001. Elasticities-a Theoretical Introduction. Balmorel Project (2001).

Daniël Reijsbergen, Shyam Sridhar, Barnabé Monnot, Stefanos Leonardos, Stratis Skoulakis, and Georgios Piliouras. 2021. Transaction fees on a honeymoon: Ethereum's eip-1559 one month later. In 2021 IEEE International Conference on Blockchain (Blockchain). IEEE, 196–204.

Herbert Scarf. 1960. Some Examples of Global Instability of the Competitive Equilibrium. *International Economic Review* 1, 3 (1960), 157–172. http://www.jstor.org/stable/2556215

Ronald William Shephard. 2015. Theory of cost and production functions. Princeton University Press.

Steve Smale. 1976. A convergent process of price adjustment and global Newton methods. Journal of Mathematical Economics 3, 2 (1976), 107–120.

Yoshihiro Tanaka. 2008. Nonsmooth optimization for production theory. Hokkaido University, Sapporo (2008).

Hirofumi Uzawa. 1960. Walras' tatonnement in the theory of exchange. The Review of Economic Studies 27, 3 (1960), 182–194.

Gerard van der Laan, AJJ Talman, et al. 1987. A convergent price adjustment process. *Economics Letters* 23, 2 (1987), 119–123.

Leon Walras. 1896. Elements de l'economie politique pure, ou, Theorie de la richesse sociale. F. Rouge.

Fang Wu and Li Zhang. 2007. Proportional Response Dynamics Leads to Market Equilibrium. In Proceedings of the Thirty-Ninth Annual ACM Symposium on Theory of Computing (San Diego, California, USA) (STOC '07). Association for Computing Machinery, New York, NY, USA, 354–363. https://doi.org/10.1145/1250790.1250844

Li Zhang. 2011. Proportional Response Dynamics in the Fisher Market. *Theor. Comput. Sci.* 412, 24 (May 2011), 2691–2698. https://doi.org/10.1016/j.tcs.2010.06.021

A OMMITED RESULTS AND PROOFS

We start by presenting the first lemma, which shows that the utility level elasticity of Hicksian demand is equal to 1 in homothetic Fisher markets.

LEMMA 3.1. For any Hicksian demand h_i associated with a homogeneous utility function u_i , for all $j, k \in [m], p \in \mathbb{R}^m_+, v_i \in \mathbb{R}_+$, it holds that $\varepsilon_{h_{ij},p_k}(p, v_i) = \varepsilon_{h_{ij},p_k}(p, 1) = 1$.

PROOF OF LEMMA 3.1. Recall from Goktas et al. [Goktas et al. 2022] that for homogeneous utility functions, the Hicksian demand is homogeneous in ν , i.e., for all $\lambda \ge 0$, $h_i(\mathbf{p}, \lambda \nu) = \lambda h_i(\mathbf{p}, \nu)$. Hence, we have:

$$\varepsilon_{h_{ij},p_k}(\boldsymbol{p},\boldsymbol{v}_i) = \partial_{p_k} h_{ij}(\boldsymbol{p},\boldsymbol{v}_i) \frac{p_k}{h_{ij}(\boldsymbol{p},\boldsymbol{v}_i)}$$
(9)

$$= v_i \partial_{p_k} h_{ij}(\boldsymbol{p}, 1) \frac{p_k}{v_i h_{ij}(\boldsymbol{p}, 1)}$$
(Homogeneity of Hicksian demand) (10)

$$=\frac{p_k}{h_{ij}(\boldsymbol{p},1)}\partial_{p_k}h_{ij}(\boldsymbol{p},1)$$
(11)

$$=\frac{p_k}{h_{ij}(\boldsymbol{p},1)}\partial_{p_k}h_{ij}(\boldsymbol{p},1)$$
(12)

$$=\varepsilon_{h_{ij},p_k}(\boldsymbol{p},1) \tag{13}$$

Additionally, looking back at Equation (12), since Hicksian demand is homogeneous of degree 1 for homogeneous utility function (see Lemma 5 of Goktas et al. [Goktas et al. 2022]), by Euler's theorem for homogeneous functions [Border 2017], we have: $\frac{p_k}{h_{ij}(p,1)} \partial_{p_k} h_{ij}(p,1) = \frac{h_{ij}(p,1)}{h_{ij}(p,1)} = 1$.

We recall Shephard's lemma which was used in the Equation (6):

LEMMA A.1 (SHEPHARD'S LEMMA [BLUME 2017; SHEPHARD 2015; TANAKA 2008]). Let $e_i(\mathbf{p}, v_i)$ be the expenditure function of buyer i and $\mathbf{h}_i(\mathbf{p}, v_i)$ be the Hicksian demand set of buyer i. The subdifferential $\partial_{\mathbf{p}} e_i(\mathbf{p}, v_i)$ is the Hicksian demand at prices \mathbf{p} and utility level v_i , i.e., $\partial_{\mathbf{p}} e_i(\mathbf{p}, v_i) = \mathbf{h}_i(\mathbf{p}, v_i)$.

We first prove that by setting γ to be 5 times the maximum demand for any good throughout the entropic tâtonnement process, we can bound the change in the prices of goods in each round. We will use the fact that the change in the price of each good is bounded as an assumption in most of the following results.

LEMMA A.2. Suppose that entropic tâtonnement process is run for all $t \in [T] \subseteq \mathbb{N}_+$ with $\gamma \geq 5 \max_{\substack{t \in [T] \\ j \in [m]}} \{1, q_j^{(t)}\}$ and let $\Delta \mathbf{p} = \mathbf{p}^{(t+1)} - \mathbf{p}^{(t)}$. then the following holds for all $t \in \mathbb{N}$:

$$e^{-\frac{1}{5}}p_j^{(t)} \le p_j^{(t+1)} \le e^{\frac{1}{5}}p_j^{(t)}$$
 and $\frac{|\Delta p_j|}{p_j^{(t)}} \le \frac{1}{4}$

LEMMA A.2. The price of of a good $j \in [m]$ can at most increase by a factor of $e^{\frac{1}{5}}$:

$$p_{j}^{(t+1)} = p_{j}^{(t)} \exp\left\{\frac{z_{j}(\boldsymbol{p}^{(t)})}{\gamma}\right\} = p_{j}^{(t)} \exp\left\{\frac{q_{j}^{(t)} - 1}{\gamma}\right\} \le p_{j}^{(t)} \exp\left\{\frac{q_{j}^{(t)}}{\gamma}\right\} \le p_{j}^{(t)} \exp\left\{\frac{q_{j}^{(t)}}{5\max_{\substack{t \in \mathbb{N} \\ j \in [m]}} \{1, q_{j}^{(t)}\}}\right\} \le p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in \mathbb{N} \\ t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in \mathbb{N} \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)}e^{\frac{1}{5}\sum_{\substack{t \in [m] \\ t \in [m]}} \{1, q_{j}^{(t)}\} = p_{j}^{(t)$$

 $\frac{1}{5}$

and decrease by a factor of $e^{-\frac{1}{5}}$:

$$p_{j}^{(t+1)} = p_{j}^{(t)} \exp\left\{\frac{z_{j}(\boldsymbol{p}^{(t)})}{\gamma}\right\} = p_{j}^{(t)} \exp\left\{\frac{q_{j}^{(t)} - 1}{\gamma}\right\} \ge p_{j}^{(t)} \exp\left\{\frac{-1}{\gamma}\right\} \ge p_{j}^{(t)} \exp\left\{\frac{-1}{5\max_{\substack{t \in \mathbb{N} \\ j \in [m]}} \{1, q_{j}^{(t)}\}}\right\} \ge p_{j}^{(t)}e^{-\frac{1}{\gamma}$$

Hence, we have $e^{-\frac{1}{5}}p_j^{(t)} \le p_j^{(t+1)} \le e^{\frac{1}{5}}p_j^{(t)}$. substracting $p_j^{(t)}$ from both sides and dividing by $p_j^{(t)}$, we obtain:

$$\frac{|\Delta p_j|}{p_j^{(t)}} = \frac{|p_j^{(t+1)} - p_j^{(t)}|}{p_j^{(t)}} \le e^{1/5} - 1 \le \frac{1}{4}$$

The following two results are due to Cheung et al. [Cheung et al. 2013]. We include their proofs for completeness. They allows us to relate the change in prices to the KL-divergence.

LEMMA A.3 (CHEUNG ET AL. [CHEUNG ET AL. 2013]). Fix $t \in \mathbb{N}_+$ and let $\Delta p = p^{(t+1)} - p^{(t)}$. Suppose that for all $j \in [m]$, $\frac{|\Delta p_j|}{p_i^{(t)}} \leq \frac{1}{4}$, then:

$$\frac{(\Delta p_j)^2}{p_j^{(t)}} \le \frac{9}{2} \delta_{\text{KL}}(p_j^{(t)} + \Delta p_j, p_j^{(t)})$$
(14)

LEMMA A.3. The bound $log(x) \ge x - x^2$ for $|x| \le \frac{1}{4}$ is used below:

$$\begin{split} \delta_{\mathrm{KL}}(p_{j}^{(t)} + \Delta p_{j}, p_{j}^{(t)}) &= (p_{j}^{(t)} + \Delta p_{j})(\log(p_{j}^{(t)} + \Delta p_{j})) - (p_{j}^{(t)} + \Delta p_{j} - p_{j}^{(t)}\log(p_{j}) + p_{j}^{(t)} - \log(p_{j}^{(t)})\Delta p_{j} \\ &= -\Delta p_{j} + (p_{j}^{(t)} + \Delta p_{j})\log\left(1 + \frac{\Delta p_{j}}{p_{j}^{(t)}}\right) \\ &\geq -\Delta p_{j} + (p_{j}^{(t)} + \Delta p_{j})\left(\frac{\Delta p_{j}}{p_{j}^{(t)}} - \frac{11}{18}\frac{(\Delta p_{j})^{2}}{(p_{j}^{(t)})^{2}}\right) \\ &\geq \frac{7}{18}\frac{(\Delta p_{j})^{2}}{p_{j}^{(t)}}\left(1 - \frac{11}{7}\frac{\Delta p_{j}}{p_{j}^{(t)}}\right) \\ &= \frac{7}{18}\frac{17}{28}\frac{(\Delta p_{j})^{2}}{p_{j}^{(t)}} \\ &\geq \frac{2}{9}\frac{(\Delta p_{j})^{2}}{p_{j}^{(t)}} \end{split}$$

LEMMA A.4. Fix $t \in \mathbb{N}_+$ and let $\Delta p = p^{(t+1)} - p^{(t)}$. Suppose that $\frac{|\Delta p_j|}{p_j} \leq \frac{1}{4}$, then for any $c \in (0, 1)$, and $A \in \mathbb{R}^{n \times m}$, and for all $j \in [m]$:

$$\frac{1}{b_i} \sum_{j \in [m]} \sum_{k \in [m]} a_{il} d_{ik} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) |\Delta p_j| |\Delta p_k| \le \frac{4}{3} \sum_{l \in [m]} \frac{a_{il}}{p_l^{(t)}} (\Delta p_l)^2$$

LEMMA A.4. First, note that since by our assumption the utilities are locally non-satiated, Walras' law is satisfied, i.e., we have $b_i = \sum_{k \in [m]} d_{ik}(\mathbf{p}^{(t)} + c\Delta \mathbf{p}, b_i)(p_k^{(t)} + c\Delta p_k);$

$$\begin{split} b_{l} \sum_{l \in [m]} \frac{a_{il}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \\ &= \sum_{l \in [m]} \frac{\left(\sum_{k \in [m]} d_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})(\boldsymbol{p}_{k}^{(t)} + c\Delta \boldsymbol{p}_{k})\right) d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})}{p_{l}^{(t)}} (\Delta p_{l})^{2} \\ &\geq \sum_{l \in [m]} \frac{\left(\sum_{k \in [m]} d_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})(\boldsymbol{p}_{k}^{(t)} - \frac{1}{4}p_{k}^{(t)})\right) a_{il}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \\ &= \sum_{l \in [m]} \frac{\left(\sum_{k \in [m]} d_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})(\frac{3}{4}p_{k}^{(t)})\right) a_{il}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \\ &= \frac{3}{4} \sum_{l \in [m]} \sum_{k \in [m]} a_{il} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \frac{p_{k}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \\ &= \frac{3}{4} \left[\sum_{l \in [m]} a_{il} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})(\Delta p_{l})^{2} + \sum_{l \in [m]} \sum_{k \neq l} a_{il} d_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \frac{p_{k}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \right] \\ &= \frac{3}{4} \left[\sum_{l \in [m]} a_{il} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i})(\Delta p_{l})^{2} + \sum_{k \in [m]} \sum_{k \neq l} a_{il} d_{ik}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \left(\frac{p_{k}^{(t)}}{p_{l}^{(t)}} |\Delta p_{l}|^{2} + \frac{p_{l}^{(t)}}{p_{k}^{(t)}} |\Delta p_{k}|^{2}\right) \right] \end{aligned}$$

Now, we apply the AM-GM inequality, i.e., for all $x, y \in \mathbb{R}_+$ since $\sqrt{xy} \le \frac{x+y}{2}$, we have:

$$b_{i} \sum_{l \in [m]} \frac{a_{il}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \geq \frac{3}{4} \sum_{l \in [m]} a_{il} d_{il} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) (\Delta p_{l})^{2} + \sum_{k < l} a_{il} d_{ik} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) (2|\Delta p_{l}||\Delta p_{k}||)$$
$$= \frac{3}{4} \sum_{j \in [m]} \sum_{k \in [m]} a_{il} d_{ik} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) |\Delta p_{j}| |\Delta p_{k}|$$

LEMMA A.5. [Cheung et al. 2013] For all $j \in [m]$:

$$\frac{1}{b_i} \sum_{j \in [m]} \sum_{k \in [m]} d_{ij}^{(t)} d_{ik}^{(t)} |\Delta p_j| |\Delta p_k| \le \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2$$

LEMMA A.5. First, note that by Walras' law we have $b_i = \sum_{k \in [m]} d_{ik}^{(t)} p_k^{(t)}$;

$$b_{i} \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} = \sum_{l \in [m]} \frac{\left(\sum_{k \in [m]} d_{ik}^{(t)} p_{k}^{(t)}\right) d_{il}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2}$$
$$= \sum_{l \in [m]} \sum_{k \in [m]} d_{il}^{(t)} d_{ik}^{(t)} \frac{p_{k}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2}$$

EC '23, July 9-12, 2023, London, United Kingdom

$$= \sum_{l \in [m]} (d_{il}^{(t)})^2 (\Delta p_l)^2 + \sum_{l \in [m]} \sum_{k \neq l} d_{il}^{(t)} d_{ik}^{(t)} \frac{p_k^{(t)}}{p_l^{(t)}} (\Delta p_l)^2$$

$$= \sum_{l \in [m]} (d_{il}^{(t)})^2 (\Delta p_l)^2 + \sum_{k \in [m]} \sum_{k \leq l} d_{ik}^{(t)} d_{il}^{(t)} \left(\frac{p_k^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{p_l^{(t)}}{p_k^{(t)}} (\Delta p_k)^2 \right)$$

Now, we apply the AM-GM inequality:

$$b_{i} \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} \ge \sum_{l \in [m]} (d_{il}^{(t)})^{2} (\Delta p_{l})^{2} + \sum_{k < l} d_{ik}^{(t)} d_{il}^{(t)} (2|\Delta p_{l}||\Delta p_{k}||)$$
$$= \sum_{j \in [m]} \sum_{k \in [m]} d_{ij}^{(t)} d_{ik}^{(t)} |\Delta p_{j}| |\Delta p_{k}|$$

An important result in microeconomics is the *law of demand* which states that when the price of a good increases, the Hicksian demand for that good decreases in a very general setting of utility functions [Levin 2004; Mas-Colell et al. 1995]. We state a weaker version of the law of demand which is re-formulated to fit the tâtonnement framework.

LEMMA A.6 (LAW OF DEMAND). [Levin 2004; Mas-Colell et al. 1995] Suppose that $\forall j \in [m], t \in \mathbb{N}, p_j^{(t)}, p_j^{(t+1)} \ge 0$ and u_i is continuous and concave. Then, $\sum_{j \in [m]} \Delta p_j \left(h_{ij}^{(t+1)} - h_{ij}^{(t)} \right) \le 0$.

A simple corollary of the law of demand which is used throughout the rest of this paper is that, during tâtonnement, the change in expenditure of the next time period is always less than or equal to the change in expenditure of the previous time period's.

COROLLARY A.7. Suppose that $\forall t \in \mathbb{N}, j \in [m], p_j^{(t)}, p_j^{(t+1)} \ge 0$ and u_i is continuous and concave then $\forall t \in \mathbb{N}, \sum_{j \in [m]} \Delta p_j h_{ij}^{(t+1)} \le \sum_{j \in [m]} \Delta p_j h_{ij}^{(t)}$.

The following lemma simply restates an essential fact about expenditure functions and Hicksian demand, namely that the Hicksian demand is the minimizer of the expenditure function.

LEMMA A.8. Suppose that $\forall j \in [m], p_j^{(t)}, t \in \mathbb{N}, h_{ij}^{(t)}, h_{ij}^{(t+1)} \ge 0$ and u_i is continuous and concave then $\sum_{j \in [m]} h_{ij}^{(t)} p_j^{(t)} \le \sum_{j \in [m]} h_{ij}^{(t+1)} p_j^{(t)}$.

LEMMA A.8. For the sake of contradiction, assume that $\sum_{j \in [m]} h_{ij}^{(t)} p_j^{(t)} > \sum_{j \in [m]} h_{ij}^{(t+1)} p_j^{(t)}$. By the definition of the Hicksian demand, we know that the bundle $h_i^{(t)}$ provides the buyer with one unit of utility. Recall that the expenditure at any price p is equal to the sum of the product of the Hicksian demands and prices, that is $e_i(p, 1) = \sum_{j \in [m]} h_{ij}(p, 1)p_j$. Hence, we have $e_i(p_j^{(t)}, 1) = \sum_{j \in [m]} h_{ij}^{(t)} p_j^{(t)} > \sum_{j \in [m]} h_{ij}^{(t+1)} p_j^{(t)} = e_i(p^{(t)}, 1)$, a contradiction.

We now introduce the following lemma which makes use of results on the behavior of Hicksian demand and expenditure functions in homothetic Fisher markets introduced by Goktas et al. [Goktas et al. 2022]. In conjunction with Corollary A.7 and Lemma A.8 are key in proving that Lemma 4.1 holds allowing us to establish convergence of tâtonnement in a general setting of utility functions. Additionally, the lemma relates the Marshallian demand of homogeneous utility functions to their Hicksian demand. Before we present the lemma, we recall the following identities [Mas-Colell et al. 1995]:

$$\forall b_i \in \mathbb{R}_+ \qquad e_i(\boldsymbol{p}, v_i(\boldsymbol{p}, b_i)) = b_i \qquad (15)$$

 $\forall v_i \in$

 $\forall b_i \in \mathbb{R}_+$

 $\forall v_i \in \mathbb{R}_+$

Denizalp Goktas, Jiayi Zhao, and Amy Greenwald

$$\mathbb{R}_{+} \qquad \qquad v_i(\boldsymbol{p}, e_i(\boldsymbol{p}, v_i)) = v_i \qquad (16)$$

$$\boldsymbol{h}_i(\boldsymbol{p}, \boldsymbol{v}_i(\boldsymbol{p}, b_i)) = \boldsymbol{d}_i(\boldsymbol{p}, b_i)$$
(17)

$$\boldsymbol{d}_i(\boldsymbol{p}, \boldsymbol{e}_i(\boldsymbol{p}, \boldsymbol{v}_i)) = \boldsymbol{h}_i(\boldsymbol{p}, \boldsymbol{v}_i)$$
(18)

LEMMA A.9. Suppose that u_i is continuous and homogeneous, then the following holds:

$$\forall j \in [m] \qquad \qquad d_{ij}(\boldsymbol{p}, b_i) = \frac{b_i h_{ij}(\boldsymbol{p}, 1)}{\sum_{j \in [m]} h_{ij}(\boldsymbol{p}, 1) p_j}$$

LEMMA A.9. We note that when utility function u_i is strictly concave, the Marshallian and Hicksian demand are unique making the following equalities well-defined.

$$\frac{b_i h_{ij}(\boldsymbol{p}, 1)}{\sum_{j \in [m]} h_{ij}(\boldsymbol{p}, 1) p_j} = \frac{b_i h_{ij}(\boldsymbol{p}, 1)}{e_i(\boldsymbol{p}, 1)}$$
(Definition of expenditure function)
$$= b_i v_i(\boldsymbol{p}, 1) h_{ij}(\boldsymbol{p}, 1)$$
(Corollary 1 of Goktas et al. [Goktas et al. 2022])
$$= v_i(\boldsymbol{p}, b_i) h_{ij}(\boldsymbol{p}, 1)$$
$$= h_{ij}(\boldsymbol{p}, v_i(\boldsymbol{p}, b_i))$$
$$= d_{ij}(\boldsymbol{p}, b_i)$$
(Marshallian Demand Identity Equation (17))

The following lemma proves that the relative change in expenditures at each iteration of tatonnement is bounded when the relative change in prices is bounded.

LEMMA A.10. Suppose that
$$\forall j \in [m], \frac{|\Delta p_j|}{p_j^{(t)}} \leq \frac{1}{4}$$
, then for any $t \in \mathbb{N}_+$ and $i \in [n]$:

$$\left| \frac{\left\langle \boldsymbol{h}_i(\boldsymbol{p}^{(t+1)}, 1), \boldsymbol{p}^{(t+1)} \right\rangle - \left\langle \boldsymbol{h}_i(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle}{\left\langle \boldsymbol{h}_i^{(t)}, \boldsymbol{p}^{(t)} \right\rangle} \right| \leq \frac{1}{4}$$
(19)

PROOF. Case 1: $\langle \boldsymbol{h}_i(\boldsymbol{p}^{(t+1)}, 1), \boldsymbol{p}^{(t+1)} \rangle \geq \langle \boldsymbol{h}_i(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \rangle$

$$\frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t+1)},1),\boldsymbol{p}^{(t+1)}\right\rangle - \left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1),\boldsymbol{p}^{(t)}\right\rangle}{\left\langle \boldsymbol{h}_{i}^{(t)},\boldsymbol{p}^{(t)}\right\rangle}$$
(20)

$$\leq \frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1), \boldsymbol{p}^{(t+1)} \right\rangle - \left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1), \boldsymbol{p}^{(t)} \right\rangle}{\left\langle \boldsymbol{h}_{i}^{(t)}, \boldsymbol{p}^{(t)} \right\rangle}$$
(Corollary A.7) (21)

$$=\frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1),\boldsymbol{p}^{(t+1)}\right\rangle}{\left\langle \boldsymbol{h}_{i}^{(t)},\boldsymbol{p}^{(t)}\right\rangle}-1$$
(22)

$$\leq \frac{5}{4} \frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1), \boldsymbol{p}^{(t)} \right\rangle}{\left\langle \boldsymbol{h}_{i}^{(t)}, \boldsymbol{p}^{(t)} \right\rangle} - 1$$
(23)

$$=\frac{1}{4} \tag{24}$$

where the penultimate line follows from the assumption that $\forall j \in [m], \frac{|\Delta p_j|}{p_j^{(t)}} \leq \frac{1}{4}$.

EC '23, July 9-12, 2023, London, United Kingdom

Case 2:
$$\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t+1)}, 1), \boldsymbol{p}^{(t+1)} \rangle \leq \langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \rangle$$

$$\frac{\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \rangle - \langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t+1)}, 1), \boldsymbol{p}^{(t+1)} \rangle}{\langle \boldsymbol{h}_{i}^{(t)}, \boldsymbol{p}^{(t)} \rangle} = 1 - \frac{\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t+1)}, 1), \boldsymbol{p}^{(t+1)} \rangle}{\langle \boldsymbol{h}_{i}^{(t)}, \boldsymbol{p}^{(t)} \rangle}$$
(25)

$$\leq 1 - \frac{3}{4} \frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t+1)}, 1), \boldsymbol{p}^{(t)} \right\rangle}{\left\langle \boldsymbol{h}_{i}^{(t)}, \boldsymbol{p}^{(t)} \right\rangle}$$
(26)

$$\leq 1 - \frac{3}{4} \frac{\left\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)}, 1), \boldsymbol{p}^{(t)} \right\rangle}{\left\langle \boldsymbol{h}_{i}^{(t)}, \boldsymbol{p}^{(t)} \right\rangle} \qquad \text{(Corollary A.7)} \quad (27)$$

$$=\frac{1}{4}$$
(28)

where the second line follows from the assumption that $\forall j \in [m], \frac{|\Delta p_j|}{p_j^{(t)}} \leq \frac{1}{4}$.

LEMMA A.11. Suppose that for all $j \in [m]$, $\frac{|\Delta p_j|}{p_j} \leq \frac{1}{4}$, then for some $c \in (0, 1)$ and $t \in \mathbb{N}_+$, we have:

$$\frac{1}{b_i} \left(\left\langle \boldsymbol{d}_i^{(t)}, \Delta \boldsymbol{p} \right\rangle + \frac{b_i}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^2 \boldsymbol{e}_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{\boldsymbol{e}_i(\boldsymbol{p}^{(t)}, 1)} \right)^2$$
(29)

$$\leq \left(1 + \frac{5\varepsilon}{9}\right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \left(\frac{25\varepsilon^2}{432}\right) \sum_{l \in [m]} \frac{d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i)}{p_l^{(t)}} (\Delta p_l)^2$$
(30)

Proof.

$$\frac{1}{b_i} \left(\left\langle \boldsymbol{d}_i^{(t)}, \Delta \boldsymbol{p} \right\rangle + \frac{b_i}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1) \Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2$$
(31)

$$=\frac{1}{b_{i}}\left[\left\langle \boldsymbol{d}_{i}^{(t)},\Delta\boldsymbol{p}\right\rangle^{2}+2\left\langle \boldsymbol{d}_{i}^{(t)},\Delta\boldsymbol{p}\right\rangle\left(\frac{b_{i}}{2}\frac{\left\langle \nabla_{\boldsymbol{p}}^{2}\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)},1)}\right)+\left(\frac{b_{i}}{2}\frac{\left\langle \nabla_{\boldsymbol{p}}^{2}\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)},1)}\right)^{2}\right]$$

$$(32)$$

$$\leq \frac{1}{b_{i}} \left[\left| \left\langle \boldsymbol{d}_{i}^{(t)}, \Delta \boldsymbol{p} \right\rangle^{2} \right| + 2 \left| \left\langle \boldsymbol{d}_{i}^{(t)}, \Delta \boldsymbol{p} \right\rangle \right| \left| \frac{b_{i}}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^{2} \boldsymbol{e}_{i}(\boldsymbol{p}^{(t)} + \boldsymbol{c}\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}, 1)} \right| + \left| \frac{b_{i}}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^{2} \boldsymbol{e}_{i}(\boldsymbol{p}^{(t)} + \boldsymbol{c}\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}, 1)} \right|^{2}$$

$$(33)$$

$$\leq \frac{1}{b_{i}} \left[\left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle^{2} + 2 \left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle \left| \frac{b_{i}}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^{2} \boldsymbol{e}_{i}(\boldsymbol{p}^{(t)} + \boldsymbol{c}\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}, 1)} \right| + \left| \frac{b_{i}}{2} \frac{\left\langle \nabla_{\boldsymbol{p}}^{2} \boldsymbol{e}_{i}(\boldsymbol{p}^{(t)} + \boldsymbol{c}\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}, 1)} \right|^{2}$$

$$(34)$$

where we denote $|\Delta \boldsymbol{p}| = (|\Delta p_1|, \dots, |\Delta p_m|).$

$$\leq \frac{1}{b_{i}} \left[\left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle^{2} + 2 \left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle \\ \left(\frac{5\varepsilon}{6} \sum_{j} \frac{(\Delta p_{j})^{2}}{p_{j}} d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \right) + \left(\frac{5\varepsilon}{6} \sum_{j} \frac{(\Delta p_{j})^{2}}{p_{j}} d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \right)^{2} \right] \quad (\text{Lemma 3.2}) \quad (35)$$
$$= \frac{1}{b_{i}} \left[\left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle^{2} + \frac{5\varepsilon}{3} \left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle \left(\sum_{j} \frac{(\Delta p_{j})^{2}}{p_{j}} d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \right) \\ + \frac{25\varepsilon^{2}}{36} \left(\sum_{j} \frac{(\Delta p_{j})^{2}}{p_{j}} d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \right)^{2} \right] \quad (36)$$

Since $\forall j \in [m], \frac{|\Delta p_j|}{p_j^{(t)}} \leq \frac{1}{4}$, we have:

$$\leq \frac{1}{b_{i}} \left[\left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle^{2} + \frac{5\varepsilon}{12} \left\langle \boldsymbol{d}_{i}^{(t)}, |\Delta \boldsymbol{p}| \right\rangle \left(\sum_{j} |\Delta p_{j}| d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1) \right) + \frac{25\varepsilon^{2}}{576} \left(\sum_{j} |\Delta p_{j}| d_{ij}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) \right)^{2} \right]$$

$$(37)$$

$$= \frac{1}{b_{i}} \sum_{j \in [m]} \sum_{k \in [m]} d_{ij}^{(t)} d_{ik}^{(t)} |\Delta p_{j}| |\Delta p_{k}| + \frac{1}{b_{i}} \frac{5\varepsilon}{12} \sum_{j} \sum_{k} d_{ik}^{(t)} d_{ij} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) |\Delta p_{k}| |\Delta p_{j}| \\ + \frac{1}{b_{i}} \frac{25\varepsilon^{2}}{576} \sum_{j} \sum_{k} d_{ij} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) d_{ik} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) |\Delta p_{k}| |\Delta p_{j}|$$

$$\leq \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_{l}^{(t)}} (\Delta p_{l})^{2} + \frac{1}{b_{i}} \frac{5\varepsilon}{12} \sum_{j} \sum_{k} d_{ik}^{(t)} d_{ij} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) |\Delta p_{k}| |\Delta p_{j}|$$

$$+ \frac{1}{b_{i}} \frac{25\varepsilon^{2}}{576} \sum_{j} \sum_{k} d_{ij} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) d_{ik} (\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_{i}) |\Delta p_{k}| |\Delta p_{j}|$$

$$(39)$$

where the last line was obtained by (Lemma A.5). Continuing, by Lemma A.4, we have:

$$\leq \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{5\varepsilon}{12} \frac{4}{3} \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{25\varepsilon^2}{576} \frac{4}{3} \sum_{l \in [m]} \frac{d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i)}{p_l^{(t)}} (\Delta p_l)^2$$
(40)

$$= \left(1 + \frac{5\varepsilon}{9}\right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \left(\frac{25\varepsilon^2}{432}\right) \sum_{l \in [m]} \frac{d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i)}{p_l^{(t)}} (\Delta p_l)^2$$
(41)

LEMMA A.12. Suppose that $\frac{|\Delta p_j|}{p_j^{(t)}} \leq \frac{1}{4}$, then

$$\begin{split} b_i \log \left(1 - \frac{e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1)}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(1 + \frac{e_i(\boldsymbol{p}^{(t+1)}, 1) - e_i(\boldsymbol{p}^{(t)}, 1)}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^{-1} \right) \\ \leq \left(\frac{4}{3} + \frac{20\varepsilon}{27} \right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \left(\frac{5\varepsilon}{6} + \frac{25\varepsilon^2}{324} \right) \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l^{(t)}} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) - \left\langle d_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \right\rangle \end{split}$$

LEMMA A.12. First, we note that $h_i^{(t)} \cdot p^{(t)} > 0$ because prices during our tâtonnement rule reach 0 only asymptotically and Hicksian demand for one unit of utility at prices $p^{(t)} > 0$ is strictly positive; and likewise, prices reach ∞ only asymptotically, which implies that Hicksian demand is always strictly positive. This fact will come handy, as we divide some expressions by $h_i^{(t)} \cdot p^{(t)}$.

Fix $t \in \mathbb{N}_+$ and $i \in [n]$. Since by our assumptions $\frac{|\Delta p_j|}{p_j^{(t)}} \le \frac{1}{4}$, by Lemma A.10, we have $0 \le \left|\frac{e_i(p^{(t+1)}, 1) - e_i(p^{(t)}, 1)}{e_i(p^{(t)}, 1)}\right| \le \frac{1}{4}$. We can then use the bound $1 - x(1+x)^{-1} \le 1 + \frac{4}{3}x^2 - x$, for $0 \le |x| \le \frac{1}{4}$, with $x = \frac{e_i(p^{(t+1)}, 1) - e_i(p^{(t)}, 1)}{e_i(p^{(t)}, 1)}$, to get:

$$b_{i} \log \left(1 - \frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)} \left(1 + \frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)^{-1}\right)$$

$$\leq b_{i} \log \left(1 + \frac{4}{3} \left(\frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)^{2} - \frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)$$

Let $a = \frac{4}{3} \left(\frac{e_i(p^{(t+1)}, 1) - e_i(p^{(t)}, 1)}{e_i(p^{(t)}, 1)} \right)^2 - \frac{e_i(p^{(t+1)}, 1) - e_i(p^{(t)}, 1)}{e_i(p^{(t)}, 1)}$. By Lemma A.10, we know that $0 + (-1/4) \le a \le \frac{1}{12} + \frac{1}{4} \Leftrightarrow -\frac{1}{4} \le a \le \frac{1}{3}$. We now use the bound $x \ge \log(1 + x)$ for x > -1, with x = a to get:

$$b_{i} \log \left(1 + \frac{4}{3} \left(\frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)^{2} - \frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)^{2} - \frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)^{2} - \frac{e_{i}(\boldsymbol{p}^{(t+1)}, 1) - e_{i}(\boldsymbol{p}^{(t)}, 1)}{e_{i}(\boldsymbol{p}^{(t)}, 1)}\right)$$

Using a first order Taylor expansion of $e_i(\mathbf{p}^{(t)} + \Delta \mathbf{p}, 1)$ around $\mathbf{p}^{(t)}$, by Taylor's theorem [Graves 1927], we have: $e_i(\mathbf{p}^{(t)} + \Delta \mathbf{p}, 1) = e_i(\mathbf{p}^{(t)}, 1) + \langle \nabla_{\mathbf{p}} e_i(\mathbf{p}^{(t)}, 1), \Delta \mathbf{p} \rangle + \frac{1}{2} \langle \nabla_{\mathbf{p}}^2 e_i(\mathbf{p}^{(t)} + c\Delta \mathbf{p}, 1)\Delta \mathbf{p}, \Delta \mathbf{p} \rangle$ for some $c \in (0, 1)$. Re-organizing terms around, we get $e_i(\mathbf{p}^{(t+1)}, 1) - e_i(\mathbf{p}^{(t)}, 1) = \langle \nabla_{\mathbf{p}} e_i(\mathbf{p}^{(t)}, 1), \Delta \mathbf{p} \rangle + \frac{1}{2} \langle \nabla_{\mathbf{p}}^2 e_i(\mathbf{p}^{(t)} + c\Delta \mathbf{p}, 1)\Delta \mathbf{p}, \Delta \mathbf{p} \rangle$, which gives us:

$$=b_{i}\left(\frac{4}{3}\left(\frac{\left\langle\nabla_{\boldsymbol{p}}\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)},1),\Delta\boldsymbol{p}\right\rangle+1/2\left\langle\nabla_{\boldsymbol{p}}^{2}\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)},1)}\right)^{2}-\frac{\left\langle\nabla_{\boldsymbol{p}}\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)},1),\Delta\boldsymbol{p}\right\rangle+1/2\left\langle\nabla_{\boldsymbol{p}}^{2}\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\right\rangle}{\boldsymbol{e}_{i}(\boldsymbol{p}^{(t)},1)}\right)$$

$$(42)$$

Continuing, by Shepherd's lemma [Shephard 2015], we have:

$$=\frac{4}{3}b_{i}\left(\frac{\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1),\Delta\boldsymbol{p}\rangle+\frac{1}{2}\left\langle\nabla_{\boldsymbol{p}}^{2}e_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\rangle\right\rangle}{e_{i}(\boldsymbol{p}^{(t)},1)}\right)^{2}-b_{i}\frac{\langle \boldsymbol{h}_{i}(\boldsymbol{p}^{(t)},1),\Delta\boldsymbol{p}\rangle+\frac{1}{2}\left\langle\nabla_{\boldsymbol{p}}^{2}e_{i}(\boldsymbol{p}^{(t)}+c\Delta\boldsymbol{p},1)\Delta\boldsymbol{p},\Delta\boldsymbol{p}\rangle}{e_{i}(\boldsymbol{p}^{(t)},1)}$$
(43)

Denizalp Goktas, Jiayi Zhao, and Amy Greenwald

$$= \frac{4}{3} \frac{1}{b_i} \left(\frac{b_i \langle \boldsymbol{h}_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} + \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)}$$

$$= \frac{4}{3} \frac{1}{b_i} \left(\langle \boldsymbol{d}_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \rangle + \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \left(\frac{44}{e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}} \rangle - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \left(\frac{44}{e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}} \rangle - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \left(\frac{44}{e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}} \rangle - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \left(\frac{44}{e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}} \rangle - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \left(\frac{44}{e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}} \rangle - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \frac{44}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(\frac{44}{e_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}} \rangle - \frac{b_i/2 \langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \rangle}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(\frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(\frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 + \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(\frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 + \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 - \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(\frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 + \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \left(\frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 + \frac{4}{e_i(\boldsymbol{p}^{(t)}, 1)} \right)^2 +$$

where the last line was obtained from Lemma A.9.

Using Lemma A.11, we have:

$$\leq \frac{4}{3} \left(\left(1 + \frac{5\varepsilon}{9} \right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{25\varepsilon^2}{432} \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l^{(t)}} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) \right) - \left(\frac{4}{q_i}(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \right) - \frac{b_{i/2} \left\langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{e_i(\boldsymbol{p}^{(t)}, 1)}$$

$$= \left(\frac{4}{3} + \frac{20\varepsilon}{27} \right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{25\varepsilon^2}{324} \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l^{(t)}} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) - \left(\frac{4}{q_i}(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \right) - \frac{b_{i/2} \left\langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{e_i(\boldsymbol{p}^{(t)}, 1)}$$

$$(46)$$

$$(46)$$

Finally, we note that $\nabla_{p}^{2} e_{i}$ is negative semi-definite, meaning that we have $\left\langle \nabla_{p}^{2} e_{i}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle \leq 0$, allowing us to re-express Equation (47) as follows:

$$= \left(\frac{4}{3} + \frac{20\varepsilon}{27}\right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{25\varepsilon^2}{324} \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l^{(t)}} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) - \left(\frac{d_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p}}{p_l^{(t)}}\right) + \left|\frac{b_{i/2} \left\langle \nabla_{\boldsymbol{p}}^2 e_i(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, 1)\Delta \boldsymbol{p}, \Delta \boldsymbol{p} \right\rangle}{e_i(\boldsymbol{p}^{(t)}, 1)}\right|$$
(48)

$$\leq \left(\frac{4}{3} + \frac{20\varepsilon}{27}\right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 + \frac{25\varepsilon^2}{324} \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l^{(t)}} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) - \left\langle d_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \right\rangle + \frac{5\varepsilon}{6} \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i)$$

$$\leq \left(\frac{4}{3} + \frac{20\varepsilon}{27}\right) \sum_{l \in [m]} \frac{d_{il}^{(t)}}{p_l^{(t)}} (\Delta p_l)^2 +$$

$$(49)$$

EC '23, July 9-12, 2023, London, United Kingdom

$$\left(\frac{5\varepsilon}{6} + \frac{25\varepsilon^2}{324}\right) \sum_{l \in [m]} \frac{(\Delta p_l)^2}{p_l^{(t)}} d_{il}(\boldsymbol{p}^{(t)} + c\Delta \boldsymbol{p}, b_i) - \left\langle \boldsymbol{d}_i(\boldsymbol{p}^{(t)}, 1), \Delta \boldsymbol{p} \right\rangle$$
(50)

where the penultimate line was obtained from Lemma 3.2.

LEMMA 4.3 (STEP SIZE FOR MARKETS WITH HOMOTHETIC NET COMPLEMENTS). If tâtonnement is run on a net complements Fisher market $(\boldsymbol{u}, \boldsymbol{b})$, then, for all $t \in \mathbb{N}_+$, prices satisfy $\left|\frac{\Delta p_j}{p_j^{(t)}}\right| \leq \frac{1}{4}$ if and only if $\gamma \geq \max\left\{d_{ij}^{(0)}, \max_{\boldsymbol{p}\in\Delta_m}\max_{k\in[m]:h_{ik}(\boldsymbol{p},1)>0}\left\{\frac{h_{ij}(\boldsymbol{p},1)}{h_{ik}(\boldsymbol{p},1)}\right\}\right\}$.

PROOF OF LEMMA 4.3. (\implies) Suppose that $d_{ij}^{(t)} \ge \max_{p \in \Delta_m} \max_{k \in [m]: h_{ik}(p,1) > 0} \left\{ \frac{h_{ij}(p,1)}{h_{ik}(p,1)} \right\}$. Then, for all $k \in [m]$:

$$d_{ik}^{(t)} = \frac{h_{ik}^{(t)}}{h_{ij}^{(t)}} d_{ij}^{(t)}$$
(51)

$$\geq \frac{h_{ik}^{(t)}}{h_{ij}^{(t)}} \max_{\boldsymbol{p} \in \Delta_m} \max_{l \in [m]: h_{il}(\boldsymbol{p}, 1) > 0} \left\{ \frac{h_{ij}(\boldsymbol{p}, 1)}{h_{il}(\boldsymbol{p}, 1)} \right\}$$
(52)

$$\geq \frac{h_{ik}^{(t)}}{h_{ij}^{(t)}} \frac{h_{ij}^{(t)}}{h_{ik}^{(t)}} = 1$$
(53)

This means that the price of good *j* will increase in the next time period, i.e., $\forall j \in [m], p_j^{(t+1)} \ge p_j^{(t)}$ which implies that $e_i(p^{(t+1)}, 1) \ge e_i(p^{(t)}, 1)$. Which gives us:

$$\frac{b_i}{e_i(\boldsymbol{p}^{(t+1)}, 1)} \le \frac{b_i}{e_i(\boldsymbol{p}^{(t)}, 1)}$$
(54)

$$\frac{b_i h_{ij}^{(t+1)}}{e_i(\boldsymbol{p}^{(t+1)}, 1)} \le \frac{b_i h_{ij}^{(t+1)}}{e_i(\boldsymbol{p}^{(t)}, 1)}$$
(55)

$$d_{ij}^{(t+1)} \le \frac{b_i h_{ij}^{(t+1)}}{e_i(\boldsymbol{p}^{(t)}, 1)}$$
(56)

$$d_{ij}^{(t+1)} \le \frac{b_i h_{ij} t]}{e_i(\boldsymbol{p}^{(t)}, 1)} = d_{ij}^{(t)}$$
(57)

where the last line follows from the net substitutes assumption, since all prices increase at time t + 1.

Now, instead suppose that $d_{ij}^{(t)} \leq \max_{p \in \Delta_m} \max_{k \in [m]: h_{ik}(p,1) > 0} \left\{ \frac{h_{ij}(p,1)}{h_{ik}(p,1)} \right\}$, then the demand for good *j* increases the most when the prices of all goods go down, which by our assumption that $\forall j \in [m], \frac{|\Delta p_j|}{p_j^{(t)}}$ gives us:

$$d_{ij}(\boldsymbol{p}^{(t+1)}, b_i) \le d_{ij}({}^{3}\!/_{4}\boldsymbol{p}^{(t)}, b_i)$$
(58)

$$\leq 4/3d_{ij}(\boldsymbol{p}^{(t)}, b_i) \tag{59}$$

$$\leq \max_{\boldsymbol{p} \in \Delta_m} \max_{k \in [m]: h_{ik}(\boldsymbol{p}, 1) > 0} \left\{ \frac{h_{ij}(\boldsymbol{p}, 1)}{h_{ik}(\boldsymbol{p}, 1)} \right\}$$
(60)

where the last line is obtained by the homogeneity of degree -1 of Marshallian demand in p for homothetic preferences [Goktas et al. 2022] and by our assumption. Then, by induction on t, we obtain:

$$d_{ij}^{(t)} \le \max\left\{d_{ij}^{0}, \max_{\boldsymbol{p}\in\Delta_{m}}\max_{k\in[m]:h_{ik}(\boldsymbol{p},1)>0}\left\{\frac{h_{ij}(\boldsymbol{p},1)}{h_{ik}(\boldsymbol{p},1)}\right\}\right\}$$
(61)

 (\rightleftharpoons)

The opposite direction follows by induction on the time steps of tâtonnement. On iteration 0, the condition holds trivially, then by the induction hypothesis can be proven by the same argument as above. $\hfill \Box$