QUANTITATIVE SPECTRAL STABILITY FOR AHARONOV-BOHM OPERATORS WITH MANY COALESCING POLES

VERONICA FELLI, BENEDETTA NORIS, ROBERTO OGNIBENE, AND GIOVANNI SICLARI

ABSTRACT. The behavior of simple eigenvalues of Aharonov-Bohm operators with many coalescing poles is discussed. In the case of half-integer circulation, a gauge transformation makes the problem equivalent to an eigenvalue problem for the Laplacian in a domain with straight cracks, laying along the moving directions of poles. For this problem, we obtain an asymptotic expansion for eigenvalues, in which the dominant term is related to the minimum of an energy functional associated with the configuration of poles and defined on a space of functions suitably jumping through the cracks.

Concerning configurations with an odd number of poles, an accurate blow-up analysis identifies the exact asymptotic behaviour of eigenvalues and the sign of the variation in some cases. An application to the special case of two poles is also discussed.

Keywords. Magnetic Schrödinger operators, Aharonov–Bohm potentials, asymptotics of eigenvalues, blow-up analysis.

MSC classification. 35J10, 35P20, 35J75,

1. INTRODUCTION

This paper deals with asymptotic expansions of the eigenvalue variation for Aharonov-Bohm operators with many coalescing poles with half-integer circulation, under homogeneous Dirichlet boundary conditions on a simply connected open bounded domain $\Omega \subset \mathbb{R}^2$. More precisely, we consider the case of any number k of poles moving along straight lines towards a collision point $P \in \Omega$, with distances from P vanishing with the same order. Without loss of generality, we assume that $P = 0 \in \Omega$, so that the moving poles can be written as multiples of k fixed points $\{a^j\}_{j=1,\dots,k}$ with the same multiplicative infinitesimal parameter $\varepsilon > 0$.

Since we are interested in the asymptotic behaviour of eigenvalues as $\varepsilon \to 0^+$, it is not restrictive to assume that there exists R < 1 such that

$$\{a^j\}_{j=1,\dots,k} \subset D_R(0) \subset \Omega,$$

where, for every r > 0 and $x \in \mathbb{R}^2$, we denote $D_r(x) := \{y \in \mathbb{R}^2 : |x - y| < r\}$. Henceforth, we denote $D_r(0)$ simply by D_r .

We assume that, among the k poles, there are k_1 poles that stand alone on their own straight line through the origin, while the remaining ones form k_2 pairs of poles staying on the same straight line but on different sides with respect to the origin. Hence

$$k = k_1 + 2k_2$$
 with $k_1, k_2 \in \mathbb{N}, \ (k_1, k_2) \neq (0, 0)$

and, for every j = 1, ..., k, there exist $r_j > 0$ and $\alpha^j \in (-\pi, \pi]$ such that $\alpha^j \neq \alpha^\ell$ if $j \neq \ell$ and

(1.1)
$$a^{j} = r_{j}(\cos(\alpha^{j}), \sin(\alpha^{j})),$$

where $\alpha^{j_1} \neq \alpha^{j_2} \pm \pi$ if $j_1 \neq j_2$ and $j_1, j_2 \in \{1, \ldots, k_1\}$, while $\alpha^j \in (-\pi, 0]$ and $\alpha^{j+k_2} = \alpha^j + \pi$ for every $j \in \{k_1 + 1, \ldots, k_1 + k_2\}$. For the sake of simplicity, we treat in detail configurations of the type described above, see Figure 1; in Section 8 we explain how our methods and results can be extended to more general configurations of poles.

For every $j = 1, \ldots, k$ and $\varepsilon \in (0, 1]$, we define

$$a^j_{\varepsilon} := \varepsilon a^j$$

Date: Revised version, December 7, 2023.



FIGURE 1. Configuration of poles $(k_1 + 1 \le j \le k_1 + k_2)$.

For every $b = (b_1, b_2) \in \mathbb{R}^2$, the Aharonov-Bohm vector potential with pole b and circulation $\rho \in \mathbb{R}$ is defined as

$$A_b^{\rho}(x_1, x_2) := \rho\left(\frac{-(x_2 - b_2)}{(x_1 - b_1)^2 + (x_2 - b_2)^2}, \frac{x_1 - b_1}{(x_1 - b_1)^2 + (x_2 - b_2)^2}\right), \quad (x_1, x_2) \in \mathbb{R}^2 \setminus \{b\}.$$

In this paper, we consider the case of half-integer circulations $\rho \in \frac{1}{2} + \mathbb{Z}$, which is of particular interest from the mathematical point of view due to applications to the problem of spectral minimal partitions, see [10, 24]. For $\rho = \frac{1}{2}$ we denote

(1.2)
$$A_b := A_b^{1/2}.$$

We are interested in the multi-singular vector potential

$$\mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)} := \sum_{j=1}^k A_{a_{\varepsilon}^j}^{n_j + \frac{1}{2}} = \sum_{j=1}^k (2n_j + 1) A_{a_{\varepsilon}^j}.$$

having at each pole a_{ε}^{j} half-integer circulation $n_{j} + \frac{1}{2}$ with $n_{j} \in \mathbb{Z}$, and in the corresponding eigenvalue problem

(1.3)
$$\begin{cases} \left(i\nabla + \mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)}\right)^2 u = \lambda u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega \end{cases}$$

where the magnetic Schrödinger operator $(i\nabla + \mathcal{A}_{\varepsilon}^{(n_1,n_2,...,n_k)})^2$ acts as

$$\left(i\nabla + \mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)}\right)^2 u := -\Delta u + 2i\,\mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)} \cdot \nabla u + \left|\mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)}\right|^2 u.$$

Since $n_j \in \mathbb{Z}$, $\mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)}$ is gauge equivalent to the vector potential

$$\mathcal{A}_{\varepsilon} := \sum_{j=1}^{k} (-1)^{j+1} A_{a_{\varepsilon}^{j}}.$$

Therefore the operators $(i\nabla + \mathcal{A}_{\varepsilon}^{(n_1, n_2, \dots, n_k)})^2$ and $(i\nabla + \mathcal{A}_{\varepsilon})^2$ are unitarily equivalent (see [19, Theorem 1.2] and [20, Proposition 2.2]), and consequently the spectrum of (1.3) coincides with that of

(1.4)
$$\begin{cases} (i\nabla + \mathcal{A}_{\varepsilon})^2 u = \lambda \, u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

Hence, to study the behaviour as $\varepsilon \to 0^+$ of the spectrum of (1.3), it is not restrictive to consider problem (1.4). We refer to (2.2) for the variational formulation of (1.4). From classical spectral theory, problem (1.4) has a diverging sequence of real positive eigenvalues $\{\lambda_{\varepsilon,n}\}_{n\in\mathbb{N}\setminus\{0\}}$; in the sequence $\{\lambda_{\varepsilon,n}\}_{n\in\mathbb{N}\setminus\{0\}}$ we repeat each eigenvalue according to its multiplicity. Moreover, the eigenspace associated to each eigenvalue has finite dimension. As $\varepsilon \to 0^+$, the following limit eigenvalue problem comes into play:

(1.5)
$$\begin{cases} \left(i\nabla + \frac{1+(-1)^{k+1}}{2}A_0\right)^2 u = \lambda u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega, \end{cases}$$

with A_0 defined as in (1.2) with b = 0. If k is odd, the operator in (1.5) is the Aharonov-Bohm operator with one pole in 0 and circulation $\frac{1}{2}$; as above, the classical Spectral Theorem applies and provides a diverging sequence of real positive eigenvalues $\{\lambda_{0,n}\}_{n \in \mathbb{N} \setminus \{0\}}$ with finite multiplicity. Furthermore, it is well-known that, in this case, eigenfunctions vanish in 0 with order $\frac{m}{2}$, for some odd $m \in \mathbb{N}$, and have exactly m nodal lines meeting at 0 and dividing the whole 2π -angle into m equal parts; see [15, Theorem 1.3, Section 7] and (3.14)–(3.15) for a description of the asymptotic behaviour at 0 of eigenfunctions of (1.5).

If k is even the nature of the limit eigenvalue problem undergoes a significant mutation. Indeed, for k even, the operator in (1.5) is the classical Dirichlet Laplacian and the eigenvalue problem (1.5) can be rewritten as

(1.6)
$$\begin{cases} -\Delta u = \lambda u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial \Omega. \end{cases}$$

We conclude that, for every $k \in \mathbb{N} \setminus \{0\}$, the spectrum of (1.5) is a diverging sequence $\{\lambda_{0,n}\}_{n \in \mathbb{N} \setminus \{0\}}$ of positive real eigenvalues.

We recall from [20, Theorem 1.2] that, whatever the number k of poles is,

the function $\varepsilon \mapsto \lambda_{\varepsilon,n}$ is continuous on [0,1],

so that, in particular,

(1.7)
$$\lim_{\varepsilon \to 0^+} \lambda_{\varepsilon,n} = \lambda_{0,n}$$

for every $n \in \mathbb{N} \setminus \{0\}$. The present paper aims at giving a sharp asymptotic expansion for the variation $\lambda_{\varepsilon,n} - \lambda_{0,n}$ of simple eigenvalues with respect to the moving configuration of poles.

In the case of one moving pole, [11] establishes a first relation between the rate of convergence (1.7) and the number of the nodal lines of the corresponding eigenfunction. Sharper asymptotic expansions for simple eigenvalues are obtained in [1], in the case of one pole moving along the tangent to a nodal line of the limit eigenfunction, and in [2], in the case of one pole moving along any direction. The case of one pole approaching the boundary is treated in [6] and [23]. The methods developed in [1], [6], and [23] are based on an Almgren type frequency formula, which provides local energy bounds for eigenfunctions. These are used to estimate the Rayleigh quotient, whose minimax levels characterize the eigenvalues, and to prove the convergence of a family of blown-up eigenfunctions to some non trivial limit profile. In particular, using the notation introduced above, in [1] it is proved that, for $k = k_1 = 1$ and $a_{\varepsilon}^1 = \varepsilon a^1 = \varepsilon r_1(\cos(\alpha^1), \sin(\alpha^1))$ moving along the tangent to one of the *m* nodal lines of the limit eigenfunction u_0 , if $\lambda_{0,n}$ is a simple, then

(1.8)
$$\lambda_{\varepsilon,n} - \lambda_{0,n} = 4 r_1^m (|\beta_1|^2 + |\beta_2|^2) \mathfrak{M} \varepsilon^m + o(\varepsilon^m) \quad \text{as } \varepsilon \to 0^+.$$

In (1.8) $(\beta_1, \beta_2) \neq (0, 0)$ is such that

$$\lim_{r \to 0^+} r^{-\frac{m}{2}} u_0(r \cos t, r \sin t) = \beta_1 e^{i\frac{t}{2}} \cos\left(\frac{m}{2}t\right) + \beta_2 e^{i\frac{t}{2}} \sin\left(\frac{m}{2}t\right),$$

see (3.14), and $\mathfrak{M} < 0$ is a negative constant depending only on m, which has the following variational characterization:

(1.9)
$$\mathfrak{M} = \min\left\{\frac{1}{2}\int_{\mathbb{R}^2_+} |\nabla u(x)|^2 \, dx - \frac{m}{2}\int_0^1 x_1^{\frac{m}{2}-1}u(x_1,0) \, dx_1 : u \in \mathcal{D}^{1,2}_s(\mathbb{R}^2_+)\right\},$$

where $s := \{(x_1, x_2) \in \mathbb{R}^2 : x_2 = 0 \text{ and } x_1 \geq 1\}$, $\mathbb{R}^2_+ = \{(x_1, x_2) \in \mathbb{R}^2 : x_2 > 0)\}$, and $\mathcal{D}^{1,2}_s(\mathbb{R}^2_+)$ is the completion of $C^{\infty}_c(\overline{\mathbb{R}^2_+} \setminus s)$ with respect to the norm $(\int_{\mathbb{R}^2_+} |\nabla u|^2 dx)^{1/2}$. For an explicit formula for \mathfrak{M} we refer to [5, Theorem 2.3]. The quantity appearing in (1.9) can be interpreted

as a weighted *torsional rigidity* of the segment along which the pole is moving. Concerning the classical notion of torsional rigidity of a set, the literature is vast; among many others, we cite the classical books [25, 17] for the basic definitions and some possible application in shape optimization and [27, 13, 12] for more recent investigations in the field. We also point out [8], where a notion of *thin torsional rigidity* is exploited in the study of spectral stability for some singularly perturbed problems.

In the case of one single pole, the study of Aharonov-Bohm eigenvalues benefits from the known regularity of the eigenvalue as a function of the pole position. Indeed, in [20] it is proved that, in the case of one moving pole, eigenvalues are analytic as functions of the pole, so that the eigenvalue variation admits a Taylor expansion. The sharp asymptotics on nodal lines (1.8) obtained in [1] is used in [2] to compute the leading term of such Taylor expansion, exploiting symmetry and periodicity properties of the Fourier coefficients of the blow-up profile with respect to the moving direction. In the case of many poles, the analyticity property is maintained as long as the poles are away from each other (see again [20]), but is lost in the case of a collision; indeed in [4] (and in [3] for symmetric domains) it is proved that, in the case of two poles colliding at a point outside the nodal set of the limit eigenfunction, the eigenvalue variation is asymptotic to the logarithm of the distance.

From the above discussion it therefore emerges that the case of multiple colliding poles presents additional significant difficulties. So far, up to our knowledge, in the literature only the case of two coalescing poles has been addressed with the aim of deriving precise asymptotic estimates in terms of the distance between the two poles. The paper [3] derives the asymptotic behaviour of eigenvalues of Aharonov–Bohm operators with two colliding poles moving on an axis of symmetry of the domain, which is assumed not to be tangent to any nodal line of the limit eigenfunction. The argument used in [3] is based on isospectrality with the Dirichlet Laplacian on the domain with a small segment removed, for which an asymptotic expansion of the eigenvalue variation is obtained by a capacity argument, in the spirit of [14]. The complementary case of two colliding poles, which move on an axis of symmetry coinciding with a nodal line of the limit eigenfunction, is treated in [5], exploiting an isospectrality result and a monotonicity formula in the spirit of [1]. The assumption of symmetry of the domain is removed in [4], in the case of two poles collapsing at an interior point out of nodal lines of the limit eigenfunction; this is possible thanks to an estimate of the diameter of the nodal set of magnetic eigenfunctions close to the collision point.

In the present paper we develop a new approach that provides asymptotic expansions of the eigenvalue variation in the most general case of any number of poles moving towards a collision point. We propose a method which combines the idea of torsional rigidity, naturally appearing in [1] (see also [6, Theorem 2.2]) to variationally characterize the coefficient of the leading term as in (1.9), with that of capacity, which [14] and [3] show to be the good small parameter in a spectral perturbation theory in domains with small holes.

Let us assume that there exists $n_0 \in \mathbb{N} \setminus \{0\}$ such that

(1.10)
$$\lambda_{0,n_0}$$
 is a simple eigenvalue of (1.5).

In view of (1.7), assumption (1.10) implies that also $\lambda_{\varepsilon,n_0}$ is simple as an eigenvalue of (1.4), provided ε is sufficiently small. Simplicity of the spectrum is a *generic* property for many differential operators. We refer e.g. to [26], where the author exhibits sufficient conditions for genericity of simplicity of the spectrum for various families of differential operators (including Aharonov-Bohm operators with a single pole). See also [7] for a focus on the particular case of Aharonov-Bohm operators.

The first step in our approach is to perform some gauge transformation, making the magnetic eigenvalue problem (1.4), and its corresponding limit one (1.5), equivalent to eigenvalue problems for the Laplacian in domains with straight cracks, laying along the moving directions of poles, see (2.10) and (2.14). Fixing a L^2 -normalized eigenfunction v_0 of the equivalent limit eigenvalue problem (2.14) associated to the eigenvalue λ_{0,n_0} , we prove in Theorem 2.1 the following asymptotic expansion:

(1.11)
$$\lambda_{\varepsilon,n_0} - \lambda_{0,n_0} = 2(\mathcal{E}_{\varepsilon} - L_{\varepsilon}(v_0)) + o(\|\nabla V_{\varepsilon}\|_{L^2(\Omega)}^2) \quad \text{as } \varepsilon \to 0^+,$$

where L_{ε} is the linear functional defined in (2.16), $\mathcal{E}_{\varepsilon}$ is the minimum of an energy functional associated with the configuration of poles and defined on a space of functions suitably jumping through the cracks, see (2.19), and V_{ε} is the potential attaining such a minimum. We observe that $\mathcal{E}_{\varepsilon}$ is a kind of intermediate quantity between torsional rigidity and capacity of the set obtained as the union of the segments connecting the poles to the origin. Indeed, the capacity of a set is defined by minimizing the L^2 -norm of the gradient among functions which are prescribed on the set; the torsional rigidity, instead, is constructed by minimizing an energy functional, which contains a linear term involving an integral on the set, without prescribing any condition. In the definition of $\mathcal{E}_{\varepsilon}$ given in (2.19), we minimize an energy functional over a family of functions which are only partially prescribed on the cracks, in the sense that we impose a jump condition on the functions across the segments, obtaining a jump of the normal derivatives as a consequent natural condition. The development of such an intermediate notion provides a unified approach, which does not require an a priori relation between the configuration of poles and the orientation of the nodal set of the limit eigenfunction. We mention that elliptic problems in cracked domains, with jumps of the unknown function and its normal derivative prescribed on the cracks, are studied in [22].

For k odd, a blow-up analysis allows us to identify the exact asymptotic behaviour of the quantities appearing in the right hand side of (1.11). In Theorem 2.2 we prove that $\lim_{\varepsilon \to 0^+} \varepsilon^{-m} \mathcal{E}_{\varepsilon} = \mathcal{E}$, where m is the vanishing order of v_0 at 0 and \mathcal{E} is the minimum of the energy functional defined in (2.28) over a space of suitably jumping functions, see (2.31). Thus we generalize (1.8) in the multipolar case, obtaining the following explicit expansion

(1.12)
$$\lambda_{\varepsilon,n_0} - \lambda_{0,n_0} = 2 \varepsilon^m \left(\mathcal{E} - L(\Psi_0) \right) + o(\varepsilon^m)$$

as $\varepsilon \to 0^+$, where L is the linear functional defined in (2.27) and Ψ_0 is the $\frac{m}{2}$ -homogeneous harmonic function introduced in (2.26). We note that the assumption that k is odd is crucial in the blow-up analysis, since it guarantees the validity of the Hardy-type inequality proved in Proposition 6.2, needed to characterize the functional space containing the limiting blow-up profile. In the particular case of all poles moving either along the tangents to nodal lines or along the bisectors between nodal lines of the limit eigenfunction, we can prove that the quantity $\mathcal{E} - L(\Psi_0)$, appearing as the coefficient of the leading term of the asymptotic expansion (1.12), does not vanish, see Proposition 2.3; this shows that m is exactly the vanishing order of the eigenvalue variation. On the other hand, the study of the continuity properties of the coefficients appearing in (1.12), see Theorem 6.8, allows us to prove the existence of configurations of poles for which $\mathcal{E} - L(\Psi_0) = 0$ and hence $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$ is an infinitesimal of higher order than m.

If k is even, a Hardy type inequality is no more available, and therefore the blow-up analysis meets the technical difficulty of identifying the limiting profile in an appropriate functional space. In spite of that, in the case of two poles colliding in a point of the nodal set of the limit eigenfunction and moving either along the tangents to its nodal lines or along its bisectors, in Theorems 2.6 and 2.7 we are able to derive the exact asymptotic behavior of $\mathcal{E}_{\varepsilon} - L_{\varepsilon}(v_0)$, and consequently of $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$ thanks to the use of elliptic coordinates; in this way we generalize the results of [3] and [5], which require an axial symmetry of the domain as a further hypothesis.

In the next section we state the main results of the paper, after having introduced the necessary notations.

2. Statement of the main results

To give a variational formulation of problem (1.4), we introduce the functional space $H^{1,\varepsilon}(\Omega, \mathbb{C})$, defined as the completion of

$$\{\phi \in H^1(\Omega, \mathbb{C}) \cap C^{\infty}(\Omega, \mathbb{C}) : \phi \equiv 0 \text{ in a neighbourhood of } a^j_{\varepsilon} \text{ for all } j = 1, \dots, k\}$$

with respect to the norm

(2.1)
$$\|w\|_{H^{1,\varepsilon}(\Omega,\mathbb{C})} := \left(\|w\|_{L^{2}(\Omega,\mathbb{C})}^{2} + \|\nabla w\|_{L^{2}(\Omega,\mathbb{C}^{2})}^{2} + \sum_{j=1}^{k} \left\|\frac{w}{|\cdot - a_{\varepsilon}^{j}|}\right\|_{L^{2}(\Omega,\mathbb{C})}^{2} \right)^{1/2}.$$

We observe that $H^{1,\varepsilon}(\Omega,\mathbb{C}) = \left\{ u \in H^1(\Omega,\mathbb{C}) : \frac{u}{|\cdot - a_{\varepsilon}^j|} \in L^2(\Omega,\mathbb{C}) \text{ for all } j = 1,\ldots,k \right\}.$ In [18] (see also [9] and [15, Lemma 3.1, Remark 3.2]), the following local magnetic Hardy-type

In [18] (see also [9] and [15, Lemma 3.1, Remark 3.2]), the following local magnetic Hardy-type inequality

$$\int_{D_r(b)} |i\nabla w + A_b^{\rho}w|^2 \, dx \ge \left(\min_{j\in\mathbb{Z}} |j-\rho|\right)^2 \int_{D_r(b)} \frac{|w(x)|^2}{|x-b|^2} \, dx$$

is proved for every $b \in \mathbb{R}^2$ and $w \in C_c^{\infty}(\overline{D_r(b)} \setminus \{b\}, \mathbb{C})$. It follows that the norm (2.1) is equivalent to the norm

$$\left(\left\|\left(i\nabla + \mathcal{A}_{\varepsilon}\right)u\right\|_{L^{2}(\Omega,\mathbb{C}^{2})}^{2} + \left\|u\right\|_{L^{2}(\Omega,\mathbb{C})}^{2}\right)^{1/2}.$$

To deal with homogeneous Dirichlet boundary conditions, we also introduce the space $H_0^{1,\varepsilon}(\Omega, \mathbb{C})$ defined as the closure of $C_c^{\infty}(\Omega \setminus \{a_{\varepsilon}^1, \ldots, a_{\varepsilon}^k\})$ in $H^{1,\varepsilon}(\Omega, \mathbb{C})$. The space $H_0^{1,\varepsilon}(\Omega, \mathbb{C})$ can be explicitly characterized as

$$H_0^{1,\varepsilon}(\Omega,\mathbb{C}) = \left\{ w \in H_0^1(\Omega,\mathbb{C}) : \frac{w}{|\cdot - a_{\varepsilon}^j|} \in L^2(\Omega,\mathbb{C}) \text{ for all } j = 1,\dots,k \right\}.$$

We say that $\lambda \in \mathbb{R}$ is an *eigenvalue* of (1.4) if there exists $u \in H_0^{1,\varepsilon}(\Omega, \mathbb{C}) \setminus \{0\}$ (called *eigenfunction*) such that

(2.2)
$$\int_{\Omega} (i\nabla + \mathcal{A}_{\varepsilon}) \, u \cdot \overline{(i\nabla + \mathcal{A}_{\varepsilon}) \, w} \, dx = \lambda \int_{\Omega} u \overline{w} \, dx \quad \text{for all } w \in H_0^{1,\varepsilon}(\Omega, \mathbb{C}).$$

We recall from the introduction that the eigenvalue problem (1.4) (and hence (2.2)) admits a diverging sequence of real positive eigenvalues

$$\lambda_{\varepsilon,1} \leq \lambda_{\varepsilon,2} \leq \lambda_{\varepsilon,3} \leq \cdots,$$

repeated in the enumeration according to their multiplicity.

In a similar way, the variational formulation of (1.5) in the case k odd (corresponding to a problem of type (1.4) with only one pole located at 0) can be be given in the functional space $\{w \in H_0^1(\Omega, \mathbb{C}) : \frac{w}{|x|} \in L^2(\Omega, \mathbb{C})\}$. In the case k even, instead, (1.5) takes the form of the classical eigenvalue problem for the Dirichlet Laplacian, whose variational formulation is well known. In both cases, (1.5) admits a diverging sequence of real positive eigenvalues

$$\lambda_{0,1} \leq \lambda_{0,2} \leq \lambda_{0,3} \leq \cdots,$$

repeated according to their multiplicity.

A suitable gauge transformation allows us to obtain equivalent formulations of (1.4) and (1.5) as eigenvalue problems for the Laplacian in domains with straight cracks. For every $\varepsilon \in [0, 1]$ we define

$$\begin{split} \Sigma^{j} &:= \{ ta^{j} : t \in \mathbb{R} \} \quad \text{for all } j = 1, \dots, k_{1} + k_{2}, \\ \Gamma^{j}_{\varepsilon} &:= \{ ta^{j} : t \in (-\infty, \varepsilon] \}, \quad S^{j}_{\varepsilon} := \{ ta^{j} : t \in [0, \varepsilon] \} \quad \text{for all } j = 1, \dots, k_{1}, \\ S^{j}_{\varepsilon} &:= \{ ta^{j} + (\varepsilon - t)a^{j+k_{2}} : t \in [0, \varepsilon] \} \quad \text{for all } j = k_{1} + 1, \dots, k_{1} + k_{2}, \\ \Gamma_{\varepsilon} &:= \left(\bigcup_{j=1}^{k_{1}} \Gamma^{j}_{\varepsilon} \right) \cup \left(\bigcup_{j=k_{1}+1}^{k_{1}+k_{2}} S^{j}_{\varepsilon} \right), \end{split}$$

see Figure 2. We note that, for every $j = 1, ..., k_1$, $\Gamma_0^j = \Gamma_{\varepsilon}^j \setminus S_{\varepsilon}^j$ is the straight half-line starting at 0 with slope $\alpha_j + \pi$. For every $\varepsilon \in [0, 1]$, we consider the functional space $\mathcal{H}_{\varepsilon}$ defined as the closure of

$$\left\{ w \in H^1(\Omega \setminus \Gamma_{\varepsilon}) = H^1(\Omega \setminus \Gamma_{\varepsilon}, \mathbb{R}) : w = 0 \text{ on a neighbourhood of } \partial \Omega \right\}$$

in $H^1(\Omega \setminus \Gamma_{\varepsilon})$ endowed with the norm $\|w\|_{H^1(\Omega \setminus \Gamma_{\varepsilon})} = \|\nabla w\|_{L^2(\Omega \setminus \Gamma_{\varepsilon})} + \|w\|_{L^2(\Omega)}$. From the Poincaré type inequality stated in Proposition 3.2, it follows that

$$\|w\|_{\mathcal{H}_{\varepsilon}} := \left(\int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^2 \, dx\right)^{1/2}$$



FIGURE 2. The sets Γ_{ε} , Γ_0 , S_{ε}^j $(1 \le j \le k_1 + k_2)$.

is a norm on $\mathcal{H}_{\varepsilon}$ equivalent to $\|w\|_{H^1(\Omega\setminus\Gamma_{\varepsilon})}$. The corresponding scalar product is denoted as $(\cdot, \cdot)_{\mathcal{H}_{\varepsilon}}$.

For every $j = 1, ..., k_1 + k_2$, with the notation $\nu^j := (-\sin(\alpha^j), \cos(\alpha^j))$ we consider the half-planes

$$\pi^j_+ := \{ x \in \mathbb{R}^2 : x \cdot \nu^j > 0 \} \text{ and } \pi^j_- := \{ x \in \mathbb{R}^2 : x \cdot \nu^j < 0 \}.$$

We observe that ν^j is the unit outer normal vector to π^j_- on $\partial \pi^j_-$. In view of classical trace results and embedding theorems for fractional Sobolev spaces in dimension 1, for every $j = 1, \ldots, k_1 + k_2$ and $p \in [2, +\infty)$ there exist continuous trace operators

(2.3)
$$\gamma^{j}_{+}: H^{1}(\pi^{j}_{+} \setminus \Gamma_{1}) \to L^{p}(\Sigma^{j}) \text{ and } \gamma^{j}_{-}: H^{1}(\pi^{j}_{-} \setminus \Gamma_{1}) \to L^{p}(\Sigma^{j}).$$

We also define the trace operators

(2.4)
$$T^{j}: H^{1}(\mathbb{R}^{2} \setminus \Gamma_{1}) \to L^{p}(\Sigma^{j}), \quad T^{j}(w) := \gamma^{j}_{+}(w|_{\pi^{j}_{+}}) + \gamma^{j}_{-}(w|_{\pi^{j}_{-}}),$$

for every $j = 1, \ldots, k_1 + k_2$ and $p \in [2, +\infty)$. For every $\varepsilon \in [0, 1]$, the restrictions to $\mathcal{H}_{\varepsilon}$ of the operators γ^j_+, γ^j_- and T^j are linear and continuous, since any element of $\mathcal{H}_{\varepsilon}$ can be trivially extended by 0 to an element of $H^1(\mathbb{R}^2 \setminus \Gamma_1)$; furthermore, due to the boundedness of Ω , such restrictions are continuous and compact from $\mathcal{H}_{\varepsilon}$ into $L^p(\Sigma^j \cap \Omega)$ for all $p \in [1, +\infty)$.

For every $\varepsilon \in (0, 1]$, we define the space

(2.5)
$$\widetilde{\mathcal{H}}_{\varepsilon} := \left\{ \begin{array}{ll} w \in \mathcal{H}_{\varepsilon} : & T^{j}(w) = 0 \text{ on } \Gamma_{\varepsilon}^{j} \text{ for all } j = 1, \dots, k_{1}, \\ & T^{j}(w) = 0 \text{ on } S_{\varepsilon}^{j} \text{ for all } j = k_{1} + 1, \dots, k_{1} + k_{2} \end{array} \right\},$$

and, for $\varepsilon = 0$,

(2.6)
$$\widetilde{\mathcal{H}}_0 := \left\{ w \in \mathcal{H}_0 : T^j(w) = 0 \text{ on } \Gamma_0^j \text{ for all } j = 1, \dots, k_1 \right\}.$$

In Section 3.3 we construct a function

(2.7)
$$\Theta_{\varepsilon} : \mathbb{R}^2 \setminus \{a_{\varepsilon}^j : j = 1, \dots, k\} \to \mathbb{R}$$

such that

(2.8)
$$\begin{cases} \Theta_{\varepsilon} \in C^{\infty}(\mathbb{R}^2 \setminus \Gamma_{\varepsilon}) \\ \nabla \Theta_{\varepsilon} \text{ can be extended to be in } C^{\infty}(\mathbb{R}^2 \setminus \{a_{\varepsilon}^j : j = 1, \dots, k\}) \text{ with } \nabla \Theta_{\varepsilon} = \mathcal{A}_{\varepsilon}, \end{cases}$$

see (3.5) for the definition of Θ_{ε} . The phase multiplication

(2.9)
$$u(x) \mapsto v(x) := e^{-i\Theta_{\varepsilon}(x)}u(x), \quad x \in \Omega \setminus \Gamma_{\varepsilon},$$

transforms any solution u to problem (1.4) into a solution v to

(2.10)
$$\begin{cases} -\Delta v = \lambda v, & \text{in } \Omega \setminus \Gamma_{\varepsilon}, \\ v = 0, & \text{on } \partial\Omega, \\ T^{j}(v) = 0, & \text{on } \Gamma_{\varepsilon}^{j} \text{ for all } j = 1, \dots, k_{1}, \\ T^{j}(\nabla v \cdot \nu^{j}) = 0, & \text{on } \Gamma_{\varepsilon}^{j} \text{ for all } j = 1, \dots, k_{1}, \\ T^{j}(v) = 0, & \text{on } S_{\varepsilon}^{j} \text{ for all } j = k_{1} + 1, \dots, k_{1} + k_{2}, \\ T^{j}(\nabla v \cdot \nu^{j}) = 0, & \text{on } S_{\varepsilon}^{j} \text{ for all } j = k_{1} + 1, \dots, k_{1} + k_{2}. \end{cases}$$

In (3.7) we also define a function

$$(2.11) \qquad \qquad \Theta_0: \mathbb{R}^2 \setminus \{0\} \to \mathbb{R}$$

satisfying

(2.12)
$$\begin{cases} \Theta_0 \in C^{\infty}(\mathbb{R}^2 \setminus \Gamma_0) \\ \nabla \Theta_0 \text{ can be extended to be in } C^{\infty}(\mathbb{R}^2 \setminus \{0\}) \text{ with } \nabla \Theta_0 = \frac{1 + (-1)^{k+1}}{2} A_0. \end{cases}$$

The gauge transformation

(2.13)
$$u(x) \mapsto v(x) := e^{-i\Theta_0(x)}u(x), \quad x \in \Omega \setminus \Gamma_0$$

shows that the limit eigenvalue problem (1.5) is equivalent to

(2.14)
$$\begin{cases} -\Delta v = \lambda v, & \text{in } \Omega \setminus \Gamma_0, \\ v = 0, & \text{on } \partial\Omega, \\ T^j(v) = 0, & \text{on } \Gamma_0^j \text{ for all } j = 1, \dots, k_1, \\ T^j(\nabla v \cdot \nu^j) = 0, & \text{on } \Gamma_0^j \text{ for all } j = 1, \dots, k_1, \end{cases}$$

in the sense that the two problems have the same eigenvalues and their eigenfunctions match each other via the phase multiplication (2.13), see Section 3.3 for details. Therefore, under assumption (1.10), λ_{0,n_0} is also a simple eigenvalue of (2.14). Let

(2.15) v_0 be an eigenfunction of (2.14) associated to λ_{0,n_0} such that $||v_0||_{L^2(\Omega)} = 1$;

it is not restrictive to assume that v_0 is real-valued, see Remark 3.5. Once v_0 is fixed as above, for every $\varepsilon \in (0, 1]$ we define

(2.16)
$$L_{\varepsilon}: \mathcal{H}_1 \to \mathbb{R}, \quad L_{\varepsilon}(w) := 2 \sum_{j=1}^{k_1+k_2} \int_{S_{\varepsilon}^j} \nabla v_0 \cdot \nu^j \gamma_+^j(w) \, dS$$

and

(2.17)
$$J_{\varepsilon} : \mathcal{H}_{\varepsilon} \to \mathbb{R}, \quad J_{\varepsilon}(w) := \frac{1}{2} \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^2 \, dx + L_{\varepsilon}(w)$$

As proved in Proposition 4.2, for every $\varepsilon \in (0,1]$ there exists a unique $V_{\varepsilon} \in \mathcal{H}_{\varepsilon}$ such that

(2.18)
$$V_{\varepsilon} - v_0 \in \widetilde{\mathcal{H}}_{\varepsilon}$$
 and $J_{\varepsilon}(V_{\varepsilon}) = \min\left\{J_{\varepsilon}(w) : w \in \mathcal{H}_{\varepsilon} \text{ and } w - v_0 \in \widetilde{\mathcal{H}}_{\varepsilon}\right\}.$

Our first main result is the following expansion of the eigenvalue variation $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$ in terms of

(2.19)
$$\mathcal{E}_{\varepsilon} = J_{\varepsilon}(V_{\varepsilon})$$

and $L_{\varepsilon}(v_0)$.

Theorem 2.1. Under assumption (1.10), let v_0 be as in (2.15). Then

(2.20)
$$\lambda_{\varepsilon,n_0} - \lambda_{0,n_0} = 2(\mathcal{E}_{\varepsilon} - L_{\varepsilon}(v_0)) + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2) \quad as \ \varepsilon \to 0^+,$$

where $\mathcal{E}_{\varepsilon}$ and V_{ε} are defined in (2.19) and (2.18), respectively.

2.1. The case k odd. For k odd, the asymptotic behaviour of $\mathcal{E}_{\varepsilon}$ as $\varepsilon \to 0^+$ can be quantified in terms of the vanishing order of v_0 at the collision point 0. Indeed, as detailed in Proposition 3.6, if k is odd, there exists $\beta \in \mathbb{R} \setminus \{0\}$ such that, as $\varepsilon \to 0^+$,

(2.21)
$$\varepsilon^{-\frac{m}{2}} v_0 \big(\varepsilon \cos t, \varepsilon \sin t \big) \to \beta f(t) \sin \big(\frac{m}{2} (t - \alpha_0) \big)$$

in $C^{1,\tau}([0,2\pi] \setminus \{\alpha^j + \pi\}_{j=1}^{k_1}, \mathbb{R})$ for all $\tau \in (0,1)$, where $m \in \mathbb{N}$ is odd and corresponds to the number of nodal lines of v_0 meeting at 0 (which equals the number of nodal lines of eigenfunctions of (1.5) associated to λ_{0,n_0}), $\alpha_0 \in [0, \frac{2\pi}{m})$ is the minimal slope of such nodal lines, and

(2.22)
$$f:[0,2\pi] \to \{-1,1\}, \quad f(t):=\prod_{j=1}^{\kappa_1} (-1)^{\chi_{[\alpha^j+\pi,2\pi)}(t)},$$

where

(2.23)
$$\chi_{[\alpha^j + \pi, 2\pi)}(t) := \begin{cases} 0, & \text{if } t \in [0, \alpha^j + \pi), \\ 1, & \text{if } t \in [\alpha^j + \pi, 2\pi). \end{cases}$$

From (2.21) we realize that the m nodal lines of v_0 which meet at 0 are tangent to the m straight half-lines

$$\mathcal{R}_j = \left\{ \left(\cos\left(\alpha_0 + j\frac{2\pi}{m}\right), \sin\left(\alpha_0 + j\frac{2\pi}{m}\right) \right) r : r \ge 0 \right\}, \quad j = 0, 1, \dots, m - 1,$$

which divide the whole 2π -angle into m equal sectors. We define the functional space

(2.24)
$$\widetilde{\mathcal{X}} := \left\{ \begin{array}{l} w \in L^1_{\text{loc}}(\mathbb{R}^2) : w \in H^1(D_r \setminus \Gamma_1) \text{ for all } r > 0, \\ \nabla w \in L^2(\mathbb{R}^2 \setminus \Gamma_1, \mathbb{R}^2), \ T^j(w) = 0 \text{ on } \Gamma_0^j \text{ for all } j = 1, \dots, k_1 \end{array} \right\},$$

and consider its closed subspace

(2.25)
$$\widetilde{\mathcal{H}} := \{ w \in \widetilde{\mathcal{X}} : T^j(w) = 0 \text{ on } S_1^j \text{ for any } j = 1, \dots, k_1 + k_2 \}.$$

Letting

(2.26)
$$\Psi_0(x) = \Psi_0(r\cos t, r\sin t) = \beta r^{\frac{m}{2}} f(t) \sin\left(\frac{m}{2}(t-\alpha_0)\right)$$

with f, m, β , and α_0 as in (2.21), we observe that the nodal set of Ψ_0 is given by $\bigcup_{j=0}^{m-1} \mathcal{R}_j$. We define

(2.27)
$$L: \widetilde{\mathcal{X}} \to \mathbb{R}, \quad L(w) := 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma^j_+(w) \, dS$$

and

(2.28)
$$J: \widetilde{\mathcal{X}} \to \mathbb{R}, \quad J(w) := \frac{1}{2} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx + L(w).$$

We observe that L(w) is well-defined also for any function $w \in H^1(D_1 \setminus \Gamma_1)$.

Let $\eta \in C_{c}^{\infty}(\mathbb{R}^{2})$ be a radial cut-off function such that

(2.29)
$$\begin{cases} 0 \le \eta(x) \le 1 \text{ for all } x \in \mathbb{R}^2, \\ \eta(x) = 1 \text{ if } x \in D_1, \quad \eta(x) = 0 \text{ if } x \in \mathbb{R}^2 \setminus D_2, \\ |\nabla \eta| \le 2 \text{ in } D_2 \setminus D_1. \end{cases}$$

As proved in Proposition 6.4, there exists a unique $\widetilde{V} \in \widetilde{\mathcal{X}}$ such that

(2.30)
$$\widetilde{V} - \eta \Psi_0 \in \widetilde{\mathcal{H}} \text{ and } J(\widetilde{V}) = \min\left\{J(w) : w \in \widetilde{\mathcal{X}} \text{ and } w - \eta \Psi_0 \in \widetilde{\mathcal{H}}\right\}.$$

Theorem 2.2. Let k be odd. Under assumption (1.10), let v_0 be as in (2.15). Then

(i) $\lim_{\varepsilon \to 0^+} \varepsilon^{-m} \mathcal{E}_{\varepsilon} = \mathcal{E}$, where m is the vanishing order of v_0 at 0 as in (2.21) and

(2.31)
$$\mathcal{E} = J(\widetilde{V}) = \min_{\eta \Psi_0 + \widetilde{\mathcal{H}}} J;$$

(ii) $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0} = 2 \varepsilon^m (\mathcal{E} - L(\Psi_0)) + o(\varepsilon^m) \text{ as } \varepsilon \to 0^+.$

The expansion proved in Theorem 2.2-(ii) identifies the sharp asymptotic behaviour of the eigenvalue variation $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$ if $\mathcal{E} - L(\Psi_0) \neq 0$; if instead $\mathcal{E} - L(\Psi_0) = 0$, Theorem 2.2-(ii) only provides the information that $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$ is an infinitesimal of higher order than m. It is therefore natural to ask whether there are configurations of poles $\{a^j\}$ for which the quantity $\mathcal{E} - L(\Psi_0)$ does or does not vanish. The following proposition gives an answer in this sense, also providing precise information on the sign of the eigenvalue variation in two remarkable cases: the case in which each pole moves along the tangent to a nodal line of the limit eigenfunction and the case in which each pole moves along the bisector between two nodal lines.

Proposition 2.3. Let $k = k_1 \leq m$ be odd and $k_2 = 0$. Under assumption (1.10), let v_0 be as in (2.15) and α_0 as in (2.21). For every $j \in \{1, \ldots, k_1\}$ let α^j be as in (1.1).

(i) If $\alpha^j \in \{\alpha_0 + \ell \frac{2\pi}{m} : \ell = 0, 1, 2, \dots, m-1\}$ for all $j \in \{1, \dots, k_1\}$, then

 $\mathcal{E} < 0$ and $L(\Psi_0) = 0;$

furthermore, $\lambda_{\varepsilon,n_0} < \lambda_{0,n_0}$ provided that $\varepsilon > 0$ is sufficiently small.

(ii) If $\alpha^j \in \{\alpha_0 + (1+2\ell)\frac{\pi}{m} : \ell = 0, 1, 2, \dots, m-1\}$ for all $j \in \{1, \dots, k_1\}$, then

 $\mathcal{E} > 0$ and $L(\Psi_0) = 0;$

furthermore, $\lambda_{\varepsilon,n_0} > \lambda_{0,n_0}$ provided that $\varepsilon > 0$ is sufficiently small.

(iii) There exists a choice of $\{\alpha^j : j = 1, ..., k\}$ such that $\mathcal{E} - L(\Psi_0) = 0$ and $\lambda_{\varepsilon, n_0} - \lambda_{0, n_0} = o(\varepsilon^m)$ as $\varepsilon \to 0^+$.

The proof of claim (iii) in Proposition 2.3 is based on a continuity argument. Indeed, the function $\mathcal{E} - L(\Psi_0)$ varies continuously under rotations of the configuration of poles, see Theorem 6.8. Hence (i) and (ii), together with Bolzano's Theorem, guarantee the existence of intermediate configurations for which $\mathcal{E} - L(\Psi_0)$ vanishes. The proof of claims (i) and (ii) highlights the fact that, analogously to $\mathcal{E}_{\varepsilon}$, also \mathcal{E} represents an intermediate notion between the capacity and the torsional rigidity of the set $\cup_{j=1}^{k_1} S_1^j$. Indeed, in case (i) it occurs that

$$\mathcal{E} = \min_{w \in \widetilde{\mathcal{H}}} \left\{ \frac{1}{2} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx + L(w) \right\} < 0,$$

see (6.41), i.e. \mathcal{E} is the minimum of a functional containing a (quadratic) energy term and a linear one, over a linear space: this makes it somehow behaving like a torsional rigidity of the set $\bigcup_{j=1}^{k_1} S_1^j$. On the other hand, in case (ii) we have the characterization

$$\mathcal{E} = \min\left\{\frac{1}{2}\int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx \colon w - \eta \Psi_0 \in \widetilde{\mathcal{H}}\right\} > 0,$$

see (6.42), which yields a notion resembling that of Ψ_0 -capacity of the set $\bigcup_{j=1}^{k_1} S_1^j$.

The proof of Theorem 2.2 is based on a blow-up analysis, which also provides the following result on the behavior of eigenfunctions, characterizing their blow-up profile and quantifying the convergence speed of the eigenfunctions of problem (1.4) towards the corresponding eigenfunction of the limit problem (1.5).

Theorem 2.4. Let k be odd and $n_0 \in \mathbb{N} \setminus \{0\}$ be such that (1.10) is satisfied. Let u_0 be an eigenfunction of (1.5) associated to λ_{0,n_0} such that $\int_{\Omega} |u_0|^2 dx = 1$. For every $\varepsilon \in (0,1]$, let $u_{\varepsilon} \in H_0^{1,\varepsilon}(\Omega, \mathbb{C})$ be the eigenfunction of (1.4) associated to the eigenvalue $\lambda_{n_0,\varepsilon}$ such that

(2.32)
$$\int_{\Omega} |u_{\varepsilon}|^2 dx = 1 \quad and \quad \int_{\Omega} e^{-i(\Theta_{\varepsilon} - \Theta_0)} u_{\varepsilon} \overline{u_0} dx \text{ is a positive real number,}$$

where Θ_{ε} and Θ_0 are as in (2.7)–(2.8) and (2.11)–(2.12), respectively. Then

(2.33)
$$\varepsilon^{-m/2} u_{\varepsilon}(\varepsilon \cdot) \to e^{i\Theta_1} (\Psi_0 - \widetilde{V}) \quad as \ \varepsilon \to 0^+$$

in
$$H^{1,1}(D_R,\mathbb{C})$$
 for all $R > 0$, where V and Ψ_0 are as in (2.30) and (2.26), respectively. Moreover,

(2.34)
$$\lim_{\varepsilon \to 0^+} \varepsilon^{-m} \int_{\mathbb{R}^2 \setminus \Gamma_1} \left| e^{-i(\Theta_{\varepsilon} - \Theta_0)} (i\nabla + \mathcal{A}_{\varepsilon}) u_{\varepsilon} - (i\nabla + A_0) u_0 \right|^2 dx = \|\nabla \widetilde{V}\|_{L^2(\mathbb{R}^2 \setminus \Gamma_1)}^2.$$

We observe that condition (2.32) allows us to identify, among all the eigenfunctions of (1.4) associated to the eigenvalue $\lambda_{n_0,\varepsilon}$ (that are multiples of a given one due to the simplicity of $\lambda_{n_0,\varepsilon}$), the one that converges to u_0 as $\varepsilon \to 0^+$.

2.2. The case of two opposite poles $(k_1 = 0, k_2 = 1)$. In the case of two opposite poles a_{ε}^1 , $a_{\varepsilon}^2 = -a_{\varepsilon}^1$ colliding to 0 from the two sides of the same straight line, we can rewrite the terms appearing in (2.20) in elliptic coordinates in the spirit of [3, Subsection 2.2], thus determining the dominant term in the asymptotic expansion. This allows us to generalize [5, Theorem 2.6, Theorem 2.8], see also [3, Theorem 1.16], removing any symmetry assumption on the domain Ω . Let us assume that

the n_0 -th eigenvalue λ_{n_0} of the Dirichlet Laplacian in Ω is simple.

We recall that, since k is even in this case, $\lambda_{n_0} = \lambda_{0,n_0}$ coincides with the n_0 -th eigenvalue of the limit problem (1.5). Let

(2.35)
$$u_0$$
 be an eigenfunction of (1.6) associated to $\lambda_{n_0} = \lambda_{0,n_0}$ such that $\int_{\Omega} u_0^2 dx = 1$.

If $u_0(0) \neq 0$, then, for any bounded simply connected domain Ω , a sharp expansion of the variation $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$ has already been obtained in [4, Theorem 1.2], see Remark 7.7. Hence we assume that $u_0(0) = 0$. Up to a suitable choice of the coordinate system, according to the notation introduced in (1.1), it is not restrictive to consider the case $\alpha^1 = 0$, $\alpha^2 = \pi$, so that, for some $r_1 \in (0, R)$, the configuration of the two opposite poles is given by

(2.36)
$$a_{\varepsilon}^{1} = r_{1}(\varepsilon, 0) \text{ and } a_{\varepsilon}^{2} = r_{1}(-\varepsilon, 0),$$

and

(2.37)
$$S_{\varepsilon} := S_{\varepsilon}^{1} = [-r_{1}\varepsilon, r_{1}\varepsilon] \times \{0\},$$

see Figure 3. Furthermore, since $u_0(0) = 0$, it is well known that there exists $m \in \mathbb{N} \setminus \{0\}$,



FIGURE 3. Two opposite poles colliding at 0 $(k_1 = 0, k_2 = 1)$.

 $\beta \in \mathbb{R} \setminus \{0\}$ and $\alpha_0 \in [0, \frac{\pi}{m})$ such that

(2.38) $r^{-m}u_0(r\cos t, r\sin t) \to \beta\sin(m(t-\alpha_0))$ in $C^{1,\tau}([0,2\pi],\mathbb{C})$ as $r \to 0^+$,

for any $\tau \in (0, 1)$. In particular, the 2m half-lines with slopes $\alpha_0 + j\frac{\pi}{m}$, $j = 0, \ldots, 2m - 1$, are tangent to the nodal lines of u_0 meeting at 0.

Remark 2.5. By standard regularity theory, u_0 is analytic in Ω . Let T_m be the Taylor polynomial of u_0 centered at 0 of order m, with $m \in \mathbb{N} \setminus \{0\}$ being as in (2.38). Then, in view of (2.38),

(2.39)
$$T_m(x_1, x_2) = \sum_{j=0}^m \frac{1}{(m-j)!j!} \frac{\partial^m u_0}{\partial x_1^{m-j} \partial x_2^j}(0) x_1^{m-j} x_2^j.$$

For every $t \in [0, 2\pi]$, we have

$$T_m(\cos t, \sin t) = \beta \sin(m(t - \alpha_0)),$$

(\nabla T_m)(\cos t, \sin t) \cdot (-\sin t, \cos t) = m\beta \cos(m(t - \alpha_0)).

Hence

$$\frac{1}{m!}\frac{\partial^m u_0}{\partial x_1^m}(0) = -\beta\sin(m\alpha_0) \quad \text{and} \quad \frac{1}{(m-1)!}\frac{\partial^m u_0}{\partial x_1^{m-1}\partial x_2^1}(0) = m\beta\cos(m\alpha_0),$$

so that, in particular,

(2.40)
$$\beta = \frac{(-1)^j}{m!} \frac{\partial^m u_0}{\partial x_1^{m-1} \partial x_2^1} (0) \text{ if } \alpha_0 = \frac{j\pi}{m} \text{ for some } j \in \{0, 1, \dots, 2m-1\},$$

and

(2.41)
$$\beta = \frac{(-1)^{j+1}}{m!} \frac{\partial^m u_0}{\partial x_1^m} (0) \quad \text{if } \alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m} \quad \text{for some } j \in \{0, 1, \dots, 2m-1\}.$$

If the segment S_{ε} is tangent to a nodal line of u_0 , i.e. if $\alpha_0 = \frac{j\pi}{m}$ for some $j \in \{0, 1, \dots, 2m-1\}$, we have the following result which generalizes [5, Theorem 2.8] dropping any symmetry assumption on Ω .

Theorem 2.6. Let λ_{n_0} be a simple eigenvalue of (1.6) and let u_0 be as in (2.35). Assume that $u_0(0) = 0$ and let $m \in \mathbb{N} \setminus \{0\}$ and α_0 be as in (2.38). Let $k_1 = 0$ and $k_2 = 1$ with the configuration of poles as in assumption (2.36). If $\alpha_0 = \frac{j\pi}{m}$ for some $j \in \{0, 1, \ldots, 2m - 1\}$, then

$$\lambda_{\varepsilon,n_0} - \lambda_{n_0} = -\frac{m\pi\beta^2 r_1^{2m}}{4^{m-1}} \binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}^2 \varepsilon^{2m} + o(\varepsilon^{2m}) \quad as \ \varepsilon \to 0^+,$$

with β as in (2.40).

On the other hand, if S_{ε} lays on the bisector of the angle between the tangents to nodal lines, i.e. if $\alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m}$ for some $j \in \{0, 1, \dots, 2m-1\}$, then we prove the following expansion.

Theorem 2.7. Let λ_{n_0} be a simple eigenvalue of (1.6) and let u_0 be as in (2.35). Assume that $u_0(0) = 0$ and let $m \in \mathbb{N} \setminus \{0\}$ and α_0 be as in (2.38). Let $k_1 = 0$ and $k_2 = 1$ with the configuration of poles as in assumption (2.36). If $\alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m}$ for some $j \in \{0, 1, \ldots, 2m - 1\}$, then

$$\lambda_{\varepsilon,n_0} - \lambda_{n_0} = \frac{m\pi\beta^2 r_1^{2m}}{4^{m-1}} {\binom{m-1}{\lfloor\frac{m-1}{2}\rfloor}}^2 \varepsilon^{2m} + o(\varepsilon^{2m}) \quad as \ \varepsilon \to 0^+,$$

with β as in (2.41).

We observe that Theorem 2.7 is a generalization of [5, Theorem 2.6] and [3, Theorem 1.16].

The rest of the paper is organized as follows. In Section 3 we collect some basic facts, such as the gauge invariance property of the problem, useful features of the functional spaces involved, and some known results that will be used in the rest of the paper. In Section 4 we provide some preliminary estimates on the quantities $\mathcal{E}_{\varepsilon}$ and L_{ε} that appear in formula (2.20); such estimates are used in Section 5, where the proof of Theorem 2.1 is completed. In Section 6 we perform a blow-up analysis of the potential V_{ε} appearing in (2.18), in the case k odd; this is the key ingredient in the proof of Theorems 2.2 and 2.4. In the same section we also complete the proof of Proposition 2.3. Finally, in Section 7 we consider the case of two poles colliding to 0 from opposite sides of the same straight line, thus proving Theorems 2.6 and 2.7.

3. Preliminaries

3.1. Scalar potential functions for A_b outside half-lines. The construction of the gauge transformation, which makes problems (1.4) and (1.5) equivalent to eigenvalue problems for the Laplacian in domains with straight cracks, is based on the remark that, since Aharonov-Bohm vector fields are irrotational, they are gradients of some scalar potential functions in simply connected domains, such as the complement of straight half-lines starting at the pole.

For every $b = (b_1, b_2) \in \mathbb{R}^2$, let $\theta_b : \mathbb{R}^2 \setminus \{b\} \to [0, 2\pi)$ be defined as

$$\theta_b(x_1, x_2) := \begin{cases} \arctan\left(\frac{x_2 - b_2}{x_1 - b_1}\right), & \text{if } x_1 > b_1, \ x_2 \ge b_2, \\ \frac{\pi}{2}, & \text{if } x_1 = b_1, \ x_2 > b_2, \\ \pi + \arctan\left(\frac{x_2 - b_2}{x_1 - b_1}\right), & \text{if } x_1 < b_1, \\ \frac{3}{2}\pi, & \text{if } x_1 = b_1, \ x_2 < b_2, \\ 2\pi + \arctan\left(\frac{x_2 - b_2}{x_1 - b_1}\right), & \text{if } x_1 > b_1, \ x_2 < b_2, \end{cases}$$

i.e.,

$$\theta_b(b + r(\cos t, \sin t)) = t$$
 for all $t \in [0, 2\pi)$ and $r > 0$.

We observe that $\theta_b \in C^{\infty}(\mathbb{R}^2 \setminus \{(x_1, b_2) : x_1 \ge b_1\})$ and $\nabla \theta_b$ can be extended to be in $C^{\infty}(\mathbb{R}^2 \setminus \{b\})$, with $\nabla(\frac{\theta_b}{2}) = A_b$ in $\mathbb{R}^2 \setminus \{b\}$. For every $b \in \mathbb{R}^2$, $\alpha \in \mathbb{R}$, and $x = (x_1, x_2) \in \mathbb{R}^2$, we define

(3.1)
$$R_{b,\alpha}(x) := \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} + M_{\alpha} \begin{bmatrix} x_1 - b_1 \\ x_2 - b_2 \end{bmatrix}$$

with

(3.2)
$$M_{\alpha} := \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix},$$

i.e., $R_{b,\alpha}$ is a rotation about b by an angle α . Let

(3.3)
$$\theta_{b,\alpha} := \theta_b \circ R_{b,\alpha},$$

so that $\theta_{b,\alpha}(b+r(\cos t,\sin t)) = \alpha + t$ for every r > 0 and $t \in [-\alpha, -\alpha + 2\pi)$. We observe that $\theta_{b,\alpha}$ is smooth in $\mathbb{R}^2 \setminus \{b+r(\cos \alpha, -\sin \alpha) : r \ge 0\}$ and $\nabla \theta_{b,\alpha}$ can be extended to be in $C^{\infty}(\mathbb{R}^2 \setminus \{b\})$, with $\nabla(\frac{\theta_{b,\alpha}}{2}) = A_b$, see Figure 4.



FIGURE 4. $\theta_{b,\alpha}$ is smooth in $\mathbb{R}^2 \setminus \{b + r(\cos \alpha, -\sin \alpha) : r \ge 0\}$.

3.2. Some remarks on functional spaces. In this subsection we describe some properties of the functional spaces $\mathcal{H}_{\varepsilon}$ introduced in Section 2.

Remark 3.1. The natural embedding $I : \mathcal{H}_{\varepsilon} \to L^2(\Omega)$ is compact. Indeed, we can cut Ω along the lines Σ^j for $j = 1 \dots, k_1 + k_2$, where Σ^j are defined in Section 2. Then we can use classical compact embedding results for each resulting subset, see for example [21, Theorem 12.30].

Arguing as in Remark 3.1, from the Poincaré inequality for functions vanishing on a portion of the boundary we can deduce the following Poincaré inequality in \mathcal{H}_1 , and hence in $\mathcal{H}_{\varepsilon}$ for any $\varepsilon \in [0, 1]$.

Proposition 3.2. There exists a constant $C_P > 0$ such that, for every $\varepsilon \in [0,1]$ and $w \in \mathcal{H}_{\varepsilon}$,

$$\int_{\Omega} w^2 dx \le C_P \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^2 \, dx.$$

Since $\Omega \setminus \Gamma_1 \subseteq \Omega \setminus \Gamma_{\varepsilon_1} \subseteq \Omega \setminus \Gamma_{\varepsilon_2}$, we have $\mathcal{H}_{\varepsilon_2} \subseteq \mathcal{H}_{\varepsilon_1} \subseteq \mathcal{H}_1$ for all $0 \leq \varepsilon_2 \leq \varepsilon_1 \leq 1$. Proposition 3.3 below establishes a Mosco-type convergence result for the spaces $\mathcal{H}_{\varepsilon}$ as $\varepsilon \to 0^+$.

Proposition 3.3. Let $\{\varepsilon_n\} \subset (0,1)$ be such that $\lim_{n\to\infty} \varepsilon_n = 0$. If $\{v_n\}_n \subset \mathcal{H}_1$ and $v \in \mathcal{H}_1$ are such that $v_n \in \mathcal{H}_{\varepsilon_n}$ for all $n \in \mathbb{N}$ and $v_n \rightharpoonup v$ in \mathcal{H}_1 as $n \rightarrow \infty$, then $v \in \mathcal{H}_0$.

Proof. For every $\varepsilon \in (0, 1]$, there exists $n(\varepsilon) \in \mathbb{N}$ such that $v_n \in \mathcal{H}_{\varepsilon}$ for all $n > n(\varepsilon)$. The weak convergence $v_n \rightharpoonup v$ in \mathcal{H}_1 then implies that $v \in \mathcal{H}_{\varepsilon}$ for all $\varepsilon \in (0, 1]$. It follows that there exists $\mathbf{f} \in L^2(\Omega, \mathbb{R}^N)$ such that $\nabla v = \mathbf{f}$ in $\mathcal{D}'(\Omega \setminus \Gamma_{\varepsilon})$ for all $\varepsilon \in (0, 1]$. Actually, $\nabla v = \mathbf{f}$ in $\mathcal{D}'(\Omega \setminus \Gamma_0)$, since, for every $\varphi \in C_c^{\infty}(\Omega \setminus \Gamma_0)$, supp $\varphi \subset \Omega \setminus \Gamma_{\varepsilon}$ for ε sufficiently small. Therefore, $v \in \mathcal{H}^1(\Omega \setminus \Gamma_0)$. From the fact that $v \in \mathcal{H}_1 \cap \mathcal{H}^1(\Omega \setminus \Gamma_0)$ it follows that $v \in \mathcal{H}_0$.

Since the singleton $\{0\}$ has null capacity in Ω , functions in \mathcal{H}_0 , respectively in \mathcal{H}_0 , can be approximated by functions vanishing in a neighbourhood of 0, as stated in Lemma 3.4.

Lemma 3.4.

- (i) The set $\mathcal{H}_{0,0} := \{ v \in \mathcal{H}_0 : v \equiv 0 \text{ in a neighbourhood of } 0 \}$ is dense in \mathcal{H}_0 .
- (ii) The set $\mathcal{H}_{0,0} := \{ v \in \mathcal{H}_0 : v \equiv 0 \text{ in a neighbourhood of } 0 \}$ is dense in \mathcal{H}_0 .

Proof. To prove (i) we first notice that, if $v \in \mathcal{H}_0$, then, defining v_n as

$$v_n(x) = \begin{cases} v(x), & \text{if } |v(x)| < n, \\ -n, & \text{if } v(x) < -n, \\ n, & \text{if } v(x) > n, \end{cases}$$

 $v_n \in \mathcal{H}_0 \cap L^{\infty}(\Omega)$ for all $n \in \mathbb{N}$ and $v_n \to v$ in \mathcal{H}_0 . Therefore it is enough to prove that $\mathcal{H}_{0,0} \cap L^{\infty}(\Omega)$ is dense in $\mathcal{H}_0 \cap L^{\infty}(\Omega)$. To this aim, let us fix some $v \in \mathcal{H}_0 \cap L^{\infty}(\Omega)$. For every $\varepsilon \in (0,1)$ we consider the cut-off function $\omega_{\varepsilon} \in W^{1,\infty}(\mathbb{R}^2)$ defined as

(3.4)
$$\omega_{\varepsilon}(x) := \begin{cases} 1, & \text{if } x \in D_{\varepsilon}, \\ \frac{2 \log |x| - \log \varepsilon}{\log \varepsilon}, & \text{if } x \in D_{\sqrt{\varepsilon}} \setminus D_{\varepsilon}, \\ 0, & \text{if } x \in \Omega \setminus D_{\sqrt{\varepsilon}}. \end{cases}$$

One may directly verify that $(1 - \omega_{\varepsilon})v \in \mathcal{H}_{0,0} \cap L^{\infty}(\Omega)$ for all $\varepsilon \in (0, 1)$ and $(1 - \omega_{\varepsilon})v \to v$ in \mathcal{H}_0 as $\varepsilon \to 0$. The proof of (i) is thereby complete. We can proceed in a similar way to obtain (ii). \Box

3.3. An equivalent eigenvalue problem by gauge transformation. For every $\varepsilon \in (0, 1]$, using the notation introduced in (3.3), we define

$$\theta_{\varepsilon}^{j} := \begin{cases} \theta_{a_{\varepsilon}^{j}, \pi-\alpha^{j}}, & \text{if } j = 1, \dots, k_{1} + k_{2}, \\ \theta_{a_{\varepsilon}^{j}, -\alpha^{j}}, & \text{if } j = k_{1} + k_{2} + 1, \dots, k_{1} + 2k_{2}, \end{cases}$$

with α^j as in (1.1), see Figure 5, and

(3.5)
$$\Theta_{\varepsilon} : \mathbb{R}^2 \setminus \{a^j_{\varepsilon} : j = 1, \dots, k\} \to \mathbb{R}, \quad \Theta_{\varepsilon} := \frac{1}{2} \sum_{j=1}^k (-1)^{j+1} \theta^j_{\varepsilon}.$$

We observe that Θ_{ε} verifies (2.8).

For any $\varepsilon \in (0,1]$, let $\lambda \in \mathbb{R}$ be an eigenvalue of problem (1.4) associated to the eigenfunction $u \in H_0^{1,\varepsilon}(\Omega, \mathbb{C}) \setminus \{0\}$. Then the function

$$v(x) := e^{-i\Theta_{\varepsilon}(x)}u(x), \quad x \in \Omega \setminus \Gamma_{\varepsilon},$$

belongs to $\mathcal{H}_{\varepsilon}$ and weakly solves (2.10), in the sense that $v \in \mathcal{H}_{\varepsilon}$ and

(3.6)
$$\int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla v \cdot \nabla w \, dx = \lambda \int_{\Omega} v w \, dx \quad \text{for all } w \in \widetilde{\mathcal{H}}_{\varepsilon},$$

where $\widetilde{\mathcal{H}}_{\varepsilon}$ is defined in (2.5). On the other hand, if $v \in \widetilde{\mathcal{H}}_{\varepsilon}$ solves (3.6), then $u = e^{i\Theta_{\varepsilon}}v$ solves (1.4). Therefore the eigenvalue problems (1.4) and (2.10) have the same eigenvalues and their eigenfunctions match each other via the phase $e^{-i\Theta_{\varepsilon}}$.



FIGURE 5. The angles θ_j^{ε} for $1 \leq j \leq k_1 + 2k_2$. The half-lines represent the singular set of the function θ_j^{ε} .

A similar gauge transformation can be made for solutions to (1.5). For every $j = 1, ..., k_1$, let α^j be as in (1.1) and

$$\theta_0^j := \theta_0 \circ R_{0,\pi-\alpha^j}.$$

We define

(3.7)
$$\Theta_0 : \mathbb{R}^2 \setminus \{0\} \to \mathbb{R}, \quad \Theta_0(x) = \frac{1}{2} \sum_{j=1}^{k_1} (-1)^{j+1} \theta_0^j(x)$$

If $k_1 = 0$ we just take $\Theta_0 \equiv 0$. We observe that Θ_0 satisfies (2.12). Furthermore, if $t \in [0, 2\pi)$,

(3.8)
$$\theta_0^j(\cos t, \sin t) = \begin{cases} t - \alpha^j + \pi, & \text{if } t \in [0, \alpha^j + \pi), \\ t - \alpha^j - \pi, & \text{if } t \in [\alpha^j + \pi, 2\pi), \end{cases}$$
$$= -\alpha^j + t + \pi(1 - 2\chi_{[\alpha^j + \pi, 2\pi)}),$$

where χ is defined in (2.23). We have that u is an eigenfunction of problem (1.5), associated to the eigenvalue λ , if and only if the function

(3.9)
$$v(x) := e^{-i\Theta_0(x)}u(x), \quad x \in \Omega \setminus \Gamma_0,$$

is a non-zero weak solution of (2.14) in the sense that $v \in \widetilde{\mathcal{H}}_0$ and

(3.10)
$$\int_{\Omega \setminus \Gamma_0} \nabla v \cdot \nabla w \, dx = \lambda \int_{\Omega} v w \, dx \quad \text{for all } w \in \widetilde{\mathcal{H}}_0,$$

where $\widetilde{\mathcal{H}}_0$ is defined in (2.6). We recall that, if k_1 is even, then, letting v as in (3.9), the function $ve^{i\Theta_0} = u$ is an eigenfunction of the Dirichlet Laplacian in Ω , hence it is smooth in Ω .

Remark 3.5. We may treat eigenfunctions of problems (2.10) and (2.14) as real-valued functions (thus justifying the choice to consider $\mathcal{H}_{\varepsilon}$ as a space of real functions). Indeed, since all the coefficients in (2.10) and (2.14) are real, both the real and the imaginary part of any eigenfunction are eigenfunctions, if not trivial. Hence, any eigenspace of (2.10) and (2.14) admits a basis made of reals eigenfunctions. See also [1, Subsection 2.3].

3.4. Asymptotics of solutions to the limit eigenvalue problem. Let $\{\alpha^j\}_{j=1}^{k_1}$ and $\chi_{[\alpha^j+\pi,2\pi)}$ be as in (1.1) and (2.23), respectively. Let f be the function defined in (2.22).

Proposition 3.6. Let k_1 be odd. If v is a non-trivial solution to (2.14), in the sense that $v \in \mathcal{H}_0$ satisfies (3.10), then there exist an odd number $m \in \mathbb{N}$, $\beta \in \mathbb{R} \setminus \{0\}$, and $\alpha_0 \in [0, \frac{2\pi}{m})$ such that

(3.11)
$$\varepsilon^{-\frac{m}{2}}v(\varepsilon\cos t, \varepsilon\sin t) \to \beta f(t)\sin\left(\frac{m}{2}(t-\alpha_0)\right)$$

in $C^{1,\tau}([0,2\pi] \setminus \{\alpha^j + \pi\}_{j=1}^{k_1}, \mathbb{R})$ as $\varepsilon \to 0^+$, for all $\tau \in (0,1)$. Moreover, there exists a constant C > 0 such that

$$(3.12) |v(x)| \le C|x|^{\frac{m}{2}} \quad and \quad |\nabla v(x)| \le C|x|^{\frac{m}{2}-1} \quad for \ all \ x \in \Omega \setminus \Gamma_0.$$

Furthermore, letting

 $\Psi(x) = \Psi(r\cos t, r\sin t) = \beta r^{\frac{m}{2}} f(t) \sin\left(\frac{m}{2}(t-\alpha_0)\right),$

with m, β , and α_0 as in (3.11) and f as in (2.22), we have that, as $\varepsilon \to 0^+$,

(3.13)
$$\varepsilon^{-\frac{m}{2}}v(\varepsilon \cdot) \to \Psi \quad in \ H^1(D_\rho \setminus \Gamma_0) \ for \ all \ \rho > 0.$$

Proof. As observed above, the function $u := e^{i\Theta_0} v$ is an eigenfunction of (1.5) with k odd, i.e.

$$\begin{cases} (i\nabla + A_0)^2 u = \lambda u, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

with A_0 defined in (1.2). From [15, Theorem 1.3, Section 7] it follows that there exist an odd $m \in \mathbb{N}$ and $\beta_1, \beta_2 \in \mathbb{C}$ such that $(\beta_1, \beta_2) \neq (0, 0)$ and, as $\varepsilon \to 0^+$,

(3.14)
$$\varepsilon^{-\frac{m}{2}}u(\varepsilon\cos t, \varepsilon\sin t) \to e^{\frac{i}{2}t}\left(\beta_1\cos\left(\frac{m}{2}t\right) + \beta_2\sin\left(\frac{m}{2}t\right)\right) \quad \text{in } C^{1,\tau}([0,2\pi],\mathbb{C})$$

and

$$(3.15) \quad \varepsilon^{1-\frac{m}{2}} \nabla u(\varepsilon \cos t, \varepsilon \sin t) \to \frac{m}{2} e^{\frac{i}{2}t} \left(\beta_1 \cos\left(\frac{m}{2}t\right) + \beta_2 \sin\left(\frac{m}{2}t\right) \right) \boldsymbol{\theta}(t) \\ \quad + \frac{d}{dt} \left(e^{\frac{i}{2}t} \left(\beta_1 \cos\left(\frac{m}{2}t\right) + \beta_2 \sin\left(\frac{m}{2}t\right) \right) \right) \boldsymbol{\tau}(t) \quad \text{in } C^{0,\tau}([0, 2\pi], \mathbb{C})$$

for all $\tau \in (0, 1)$, where $\theta(t) = (\cos t, \sin t)$ and $\tau(t) = (-\sin t, \cos t)$. Furthermore, by (3.8), for all $t \in [0, 2\pi]$ we have

(3.16)
$$\sum_{j=1}^{k_1} (-1)^{j+1} \theta_0^j(\varepsilon \cos t, \varepsilon \sin t) = t + \sum_{j=1}^{k_1} (-1)^j \alpha^j + \pi - 2\pi \sum_{j=1}^{k_1} (-1)^{j+1} \chi_{[\alpha^j + \pi, 2\pi)}(t).$$

From (3.14), the definition of u, and (3.16) it follows that

$$\varepsilon^{-\frac{m}{2}}v(\varepsilon\cos t,\varepsilon\sin t) \to f(t)e^{-\frac{i}{2}\left(\pi+\sum_{j=1}^{k_1}(-1)^j\alpha^j\right)}\left(\beta_1\cos\left(\frac{m}{2}t\right)+\beta_2\sin\left(\frac{m}{2}t\right)\right)$$

in $C^{1,\tau}([0,2\pi] \setminus \{\alpha^j + \pi\}_{j=1}^{k_1}, \mathbb{C})$ as $\varepsilon \to 0^+$, for all $\tau \in (0,1)$. Then, since v is real-valued (see Remark 3.5), we have proved that there exist $c_1, c_2 \in \mathbb{R}$ such that $(c_1, c_2) \neq (0,0)$ and

(3.17)
$$\varepsilon^{-\frac{m}{2}}v(\varepsilon\cos t,\varepsilon\sin t) \to f(t)\Big(c_1\cos\left(\frac{m}{2}t\right) + c_2\sin\left(\frac{m}{2}t\right)\Big).$$

Letting

$$\alpha_0 = \begin{cases} \frac{2}{m} \operatorname{arccot} \left(-\frac{c_2}{c_1} \right), & \text{if } c_1 \neq 0, \\ 0, & \text{if } c_1 = 0, \end{cases}$$

we can rewrite (3.17) as (3.11). Estimate (3.12) is a consequence of (3.14) and (3.15). Finally, to prove (3.13), we define

Finally, to prove
$$(3.15)$$
, we define

$$\tilde{u}_{\varepsilon}(x) := \varepsilon^{-\frac{m}{2}} u(\varepsilon x), \quad \Phi(x) = \Phi(r \cos t, r \sin t) = r^{\frac{m}{2}} e^{\frac{t}{2}t} \left(\beta_1 \cos\left(\frac{m}{2}t\right) + \beta_2 \sin\left(\frac{m}{2}t\right)\right).$$

We observe that (3.14), (3.15), and the Dominated Convergence Theorem imply that

$$abla ilde{u}_{arepsilon} o
abla \Phi \quad ext{and} \quad rac{ ilde{u}_{arepsilon}}{|x|} o rac{\Phi}{|x|} \quad ext{in } L^2(D_{
ho}) \quad ext{for all }
ho > 0,$$

which easily provides (3.13).

In the case k even, solutions to (2.14) are more regular.

Proposition 3.7. Let k_1 be even. If v is a non-trivial solution to (2.14), then there exist $m \in \mathbb{N}$, $\beta \in \mathbb{R} \setminus \{0\}$, and $\alpha_0 \in [0, \frac{\pi}{m})$ such that

(3.18)
$$\varepsilon^{-m}v(\varepsilon\cos t, \varepsilon\sin t) \to \beta f(t)\sin(m(t-\alpha_0))$$

in $C^{1,\tau}([0,2\pi] \setminus \{\pi + \alpha^j\}_{j=1}^{k_1}, \mathbb{R})$ as $\varepsilon \to 0^+$, for all $\tau \in (0,1)$. Moreover, there exists a constant C > 0 such that

(3.19)
$$|v(x)| \le C|x|^m \quad and \quad |\nabla v(x)| \le \begin{cases} C|x|^{m-1}, & \text{if } m \ge 1, \\ C, & \text{if } m = 0, \end{cases}$$

for all $x \in \Omega \setminus \Gamma_0$.

Proof. The function $u := e^{i\Theta_0}v$ is an eigenfunction of (1.5) with k even, i.e. u is an eigenfunction of the Dirichlet Laplacian. From (3.8) we deduce the analogue of (3.16) in the even case:

(3.20)
$$\sum_{j=1}^{k_1} (-1)^{j+1} \theta_0^j(\varepsilon \cos t, \varepsilon \sin t) = \sum_{j=1}^{k_1} (-1)^j \alpha^j - 2\pi \sum_{j=1}^{k_1} (-1)^{j+1} \chi_{[\alpha^j + \pi, 2\pi)}(t)$$

for all $t \in [0, 2\pi]$. Claims (3.18) and (3.19) follow from the fact that u is analytic, the definition of u and (3.20), observing that, since k_1 is even, $|\nabla u| = |\nabla v|$.

Remark 3.8. For the sake of simplicity, for any $w \in \mathcal{H}_0$ we simply write w instead of $\gamma^j_+(w)$ on S^j_{ε} , since $\gamma^j_+(w) = \gamma^j_-(w)$ on S^j_{ε} for any $j = 1, \ldots, k_1 + k_2$. We also simply write v_0 , ∇v_0 and $\nabla v_0 \cdot \nu^j$ when considering their traces on S^j_{ε} .

4. Definition and properties of $\mathcal{E}_{\varepsilon}$

For some $n_0 \in \mathbb{N} \setminus \{0\}$, let u_0 be an eigenfunction of (1.5) associated to the eigenvalue $\lambda_0 = \lambda_{0,n_0}$ and v_0 be as in (3.9), so that v_0 is a non-zero weak solution of (2.14) with $\lambda = \lambda_0$. By Remark 3.5 it is not restrictive to assume that v_0 is real-valued and $\|u_0\|_{L^2(\Omega,\mathbb{C})} = \|v_0\|_{L^2(\Omega)} = 1$.

Let L_{ε} be the functional introduced in (2.16). We observe that L_{ε} is well-defined; indeed, for every $j = 1, \ldots, k_1 + k_2$, we have $\nabla v_0 \in L^p(S^j_{\varepsilon})$ for all $p \in [1, 2)$ in view of (3.12) and (3.19), whereas $\gamma^j_+(w) \in L^q(S^j_{\varepsilon})$ for all $w \in \mathcal{H}_1$ and $q \in [2, +\infty)$ by (2.3). We provide below an estimate of the norm of L_{ε} in \mathcal{H}_1^* , where \mathcal{H}_1^* is the dual space of \mathcal{H}_1 .

Proposition 4.1. Let $m \in \mathbb{N}$ be as in Proposition 3.6 for $v = v_0$, if k is odd, or as in Proposition 3.7, if k is even. For every $\varepsilon \in (0, 1]$, the map L_{ε} defined in (2.16) belongs to \mathcal{H}_1^* and, as $\varepsilon \to 0^+$,

(4.1)
$$\|L_{\varepsilon}\|_{\mathcal{H}_{1}^{*}} = \begin{cases} O(\varepsilon^{\frac{m}{2}-1+\frac{1}{p}}), & \text{if } k \text{ is odd,} \\ O(\varepsilon^{\frac{1}{p}}), & \text{if } k \text{ is even and } m = 0, \\ O(\varepsilon^{m-1+\frac{1}{p}}), & \text{if } k \text{ is even and } m > 0, \end{cases}$$

for every $p \in (1,2)$. In particular, $L_{\varepsilon} \to 0$ in \mathcal{H}_1^* as $\varepsilon \to 0^+$.

Proof. If k is odd, for every $p \in (1, 2)$ and $w \in \mathcal{H}_1$, from the Hölder inequality, (2.3) and (3.12) it follows that, letting $p' = \frac{p}{p-1}$,

$$|L_{\varepsilon}(w)| \leq 2 \sum_{j=1}^{k_1+k_2} \|\nabla v_0\|_{L^p(S^j_{\varepsilon})} \|\gamma^j_+(w)\|_{L^{p'}(S^j_{\varepsilon})} \leq C\varepsilon^{\frac{m}{2}-1+\frac{1}{p}} \|w\|_{\mathcal{H}_1},$$

for some constant C > 0 independent of ε . If k is even, the proof is similar due to (3.19).

For every $\varepsilon \in (0, 1]$, we now consider the functional J_{ε} defined in (2.17) and, recalling the definition of $\widetilde{\mathcal{H}}_{\varepsilon}$ in (2.5), the minimization problem

(4.2)
$$\inf \left\{ J_{\varepsilon}(w) : w \in \mathcal{H}_{\varepsilon} \text{ and } w - v_0 \in \widetilde{\mathcal{H}}_{\varepsilon} \right\}.$$

Note that, since $v_0 \in \widetilde{\mathcal{H}}_0$, the condition $w - v_0 \in \widetilde{\mathcal{H}}_{\varepsilon}$ is equivalent to

(4.3)
$$T^{j}(w) = \begin{cases} 0, & \text{on } \Gamma_{0}^{j} \text{ for all } j = 1, \dots, k_{1}, \\ 2v_{0}, & \text{on } S_{\varepsilon}^{j} \text{ for all } j = 1, \dots, k_{1} + k_{2} \end{cases}$$

Proposition 4.2. The infimum in (4.2) is achieved by a unique $V_{\varepsilon} \in \mathcal{H}_{\varepsilon}$. Furthermore, V_{ε} weakly solves the problem

$$(4.4) \qquad \begin{cases} -\Delta V_{\varepsilon} = 0, & \text{in } \Omega \setminus \Gamma_{\varepsilon}, \\ V_{\varepsilon} = 0, & \text{on } \partial\Omega, \\ T^{j}(V_{\varepsilon} - v_{0}) = 0, & \text{on } \Gamma_{\varepsilon}^{j} \quad \text{for all } j = 1, \dots, k_{1}, \\ T^{j}(\nabla V_{\varepsilon} \cdot \nu^{j} - \nabla v_{0} \cdot \nu^{j}) = 0, & \text{on } \Gamma_{\varepsilon}^{j} \quad \text{for all } j = 1, \dots, k_{1}, \\ T^{j}(V_{\varepsilon} - v_{0}) = 0, & \text{on } S_{\varepsilon}^{j} \quad \text{for all } j = k_{1} + 1, \dots, k_{1} + k_{2}, \\ T^{j}(\nabla V_{\varepsilon} \cdot \nu^{j} - \nabla v_{0} \cdot \nu^{j}) = 0, & \text{on } S_{\varepsilon}^{j} \quad \text{for all } j = k_{1} + 1, \dots, k_{1} + k_{2}, \end{cases}$$

in the sense that $V_{\varepsilon} \in \mathcal{H}_{\varepsilon}, V_{\varepsilon} - v_0 \in \widetilde{\mathcal{H}}_{\varepsilon}$, and

(4.5)
$$\int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla V_{\varepsilon} \cdot \nabla w \, dx = -L_{\varepsilon}(w) \quad \text{for all } w \in \widetilde{\mathcal{H}}_{\varepsilon}$$

Proof. In view of (2.17), the continuity of the linear operator L_{ε} , and Proposition 3.2, we can easily verify that J_{ε} is continuous and coercive on the set $v_0 + \tilde{\mathcal{H}}_{\varepsilon} = \{w \in \mathcal{H}_{\varepsilon} : w - v_0 \in \tilde{\mathcal{H}}_{\varepsilon}\}$, which is closed and convex. Moreover, J_{ε} is convex. Therefore, the infimum in (4.2) is achieved by some V_{ε} , which weakly solves (4.4) in the sense of (4.5). If $V_{\varepsilon,1}, V_{\varepsilon,2} \in v_0 + \tilde{\mathcal{H}}_{\varepsilon}$ are weak solutions of (4.4), then $V_{\varepsilon,1} - V_{\varepsilon,2} \in \tilde{\mathcal{H}}_{\varepsilon}$ and

(4.6)
$$\int_{\Omega \setminus \Gamma_{\varepsilon}} (\nabla V_{\varepsilon,1} - \nabla V_{\varepsilon,2}) \cdot \nabla w \, dx = 0 \quad \text{for all } w \in \tilde{\mathcal{H}}_{\varepsilon}.$$

Testing (4.6) with $w = V_{\varepsilon,1} - V_{\varepsilon,2}$ we obtain $\nabla(V_{\varepsilon,1} - V_{\varepsilon,2}) = 0$ and hence, by Proposition 3.2, we conclude that $V_{\varepsilon,1} = V_{\varepsilon,2}$.

For every $\varepsilon \in (0, 1]$, let J_{ε} and V_{ε} be as (2.17) and Proposition 4.2, respectively. We consider the quantity $\mathcal{E}_{\varepsilon} := J_{\varepsilon}(V_{\varepsilon})$ as in (2.19). $\mathcal{E}_{\varepsilon}$ plays a significant role in the asymptotic expansion of the eigenvalue variation $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$, as the poles a_{ε}^j move towards the collision at 0.

To derive a first upper and lower bound for $\mathcal{E}_{\varepsilon}$, we consider, for every r > 0, the radial cut-off function $\eta_r \in C_c^{\infty}(\mathbb{R}^2)$ defined as

(4.7)
$$\eta_r(x) := \eta\left(\frac{x}{r}\right)$$

with η as in (2.29).

Proposition 4.3. Let $m \in \mathbb{N}$ be as in Proposition 3.6 for $v = v_0$, if k is odd, or as in Proposition 3.7, if k is even. Then there exists a constant $C_1 > 0$ such that, for all $\varepsilon \in (0, 1]$,

(4.8)
$$\mathcal{E}_{\varepsilon} \leq \begin{cases} C_1 \, \varepsilon^m, & \text{if } k \text{ is odd,} \\ C_1 \, \frac{1}{|\log \varepsilon|}, & \text{if } k \text{ is even and } m = 0, \\ C_1 \, \varepsilon^{2m}, & \text{if } k \text{ is even and } m > 0. \end{cases}$$

Moreover, for every $p \in (1,2)$ there exists $C_2 = C_2(p) > 0$ such that

(4.9)
$$\mathcal{E}_{\varepsilon} \geq \begin{cases} -C_2 \,\varepsilon^{m-2+\frac{2}{p}}, & \text{if } k \text{ is odd,} \\ -C_2 \,\varepsilon^{\frac{2}{p}}, & \text{if } k \text{ is even and } m = 0, \\ -C_2 \,\varepsilon^{2m-2+\frac{2}{p}}, & \text{if } k \text{ is even and } m > 0. \end{cases}$$

In particular, $\mathcal{E}_{\varepsilon} \to 0$ as $\varepsilon \to 0^+$.

Proof. If k is odd, let $\eta_{\varepsilon} \in C_{c}^{\infty}(\mathbb{R}^{2})$ be a cut-off function as in (4.7) with $r = \varepsilon$. From (2.16), (2.17), (4.2), (2.19), and (3.12) it follows that

$$\begin{aligned} J_{\varepsilon}(V_{\varepsilon}) &\leq J_{\varepsilon}(\eta_{\varepsilon}v_{0}) \leq \frac{1}{2} \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla(\eta_{\varepsilon}v_{0})|^{2} dx + 2 \sum_{j=1}^{k_{1}+k_{2}} \int_{S_{\varepsilon}^{j}} |\nabla v_{0}| |v_{0}| dS \\ &\leq \int_{(\Omega \cap D_{2\varepsilon}(0)) \setminus \Gamma_{\varepsilon}} |\nabla v_{0}|^{2} dx + \int_{\Omega \cap D_{2\varepsilon}(0)} |\nabla \eta_{\varepsilon}|^{2} v_{0}^{2} dx + 2 \sum_{j=1}^{k_{1}+k_{2}} \int_{S_{\varepsilon}^{j}} |\nabla v_{0}| |v_{0}| dS \leq C_{1} \varepsilon^{m} \end{aligned}$$

for some constant $C_1 > 0$ independent of ε . If k is even and $m \in \mathbb{N} \setminus \{0\}$, (4.8) can be proved arguing in a similar way and using (3.19) instead of (3.12).

If k is even and m = 0, for every $\varepsilon \in (0, 1]$ we consider the cut-off function $\omega_{\varepsilon} \in W^{1,\infty}(\mathbb{R}^2)$ defined in (3.4). We have $0 \le \omega_{\varepsilon} \le 1$ and, thanks to (2.16), (2.17), (4.2), (2.19), and (3.19) with m = 0,

$$J_{\varepsilon}(V_{\varepsilon}) \leq J_{\varepsilon}(\omega_{\varepsilon}v_{0}) \leq \frac{1}{2} \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla(\omega_{\varepsilon}v_{0})|^{2} dx + 2 \sum_{j=1}^{k_{1}+k_{2}} \int_{S_{\varepsilon}^{j}} |\nabla v_{0}| |v_{0}| dS$$
$$\leq \int_{(\Omega \cap D_{\sqrt{\varepsilon}}(0)) \setminus \Gamma_{\varepsilon}} |\nabla v_{0}|^{2} dx + \int_{\Omega \cap D_{\sqrt{\varepsilon}}(0)} |\nabla \omega_{\varepsilon}|^{2} v_{0}^{2} dx + 2 \sum_{j=1}^{k_{1}+k_{2}} \int_{S_{\varepsilon}^{j}} |\nabla v_{0}| |v_{0}| dS \leq C_{1} \frac{1}{|\log \varepsilon|}$$

for some constant $C_1 > 0$ independent of ε . Estimate (4.8) is thereby proved.

To prove (4.9), we observe that

$$\begin{aligned} \|V_{\varepsilon}\|_{\mathcal{H}_{1}}^{2} &= \|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2} = 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(V_{\varepsilon}) \leq 2\mathcal{E}_{\varepsilon} + 2|L_{\varepsilon}(V_{\varepsilon})| \\ &\leq 2\mathcal{E}_{\varepsilon} + 2\|L_{\varepsilon}\|_{\mathcal{H}_{1}^{*}}\|V_{\varepsilon}\|_{\mathcal{H}_{1}} \leq 2\mathcal{E}_{\varepsilon} + 2\|L_{\varepsilon}\|_{\mathcal{H}_{1}^{*}}^{2} + \frac{1}{2}\|V_{\varepsilon}\|_{\mathcal{H}_{1}}^{2}, \end{aligned}$$

and hence

(4.10)
$$\mathcal{E}_{\varepsilon} + \|L_{\varepsilon}\|_{\mathcal{H}_{1}^{*}}^{2} \geq \frac{1}{4} \|V_{\varepsilon}\|_{\mathcal{H}_{1}}^{2} \geq 0,$$

which, together with (4.1), implies (4.9).

Proposition 4.4. We have $V_{\varepsilon} \to 0$ as $\varepsilon \to 0^+$ strongly in \mathcal{H}_1 .

Proof. From Proposition 4.3 we have $\lim_{\varepsilon \to 0^+} \mathcal{E}_{\varepsilon} = 0$, whereas Proposition 4.1 implies that $\lim_{\varepsilon \to 0^+} \|L_{\varepsilon}\|_{\mathcal{H}^*_1} = 0$. The conclusion then follows from (4.10).

Proposition 4.5. We have $\mathcal{E}_{\varepsilon} = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} \right)$ as $\varepsilon \to 0^+$.

Proof. Proceeding similarly to the previous proof, we have

$$|\mathcal{E}_{\varepsilon}| \leq \frac{\|V_{\varepsilon}\|_{\mathcal{H}_{1}}^{2}}{2} + \|L_{\varepsilon}\|_{\mathcal{H}_{1}^{*}} \|V_{\varepsilon}\|_{\mathcal{H}_{1}}$$

and we can conclude thanks to (4.1) and Proposition 4.4.

Proposition 4.6. We have

$$\int_{\Omega} V_{\varepsilon}^2 \, dx = o\big(\left\| V_{\varepsilon} \right\|_{\mathcal{H}_{\varepsilon}}^2 \big) \quad as \ \varepsilon \to 0^+.$$

Proof. Let us assume by contradiction that there exist a positive constant C > 0 and a sequence $\{\varepsilon_n\}_{n \in \mathbb{N}} \subset (0, 1)$ such that $\lim_{n \to \infty} \varepsilon_n = 0$ and

(4.11)
$$\int_{\Omega} V_{\varepsilon_n}^2 dx \ge C \int_{\Omega \setminus \Gamma_{\varepsilon_n}} |\nabla V_{\varepsilon_n}|^2 dx \quad \text{for all } n \in \mathbb{N}.$$

For every $n \in \mathbb{N}$, we define $W_n := \frac{V_{\varepsilon_n}}{\|V_{\varepsilon_n}\|_{L^2(\Omega)}}$. Then $\|W_n\|_{L^2(\Omega)} = 1$ for every $n \in \mathbb{N}$ and $\{W_n\}_{n \in \mathbb{N}}$ is bounded in \mathcal{H}_1 thanks to (4.11). It follows that there exists $W \in \mathcal{H}_1$ such that $W_n \rightharpoonup W$ weakly

in \mathcal{H}_1 as $n \to \infty$, up to a subsequence. Since $W_n \in \mathcal{H}_{\varepsilon_n}$ for every *n*, from Proposition 3.3 we deduce that $W \in \mathcal{H}_0$, while Remark 3.1 ensures that

$$(4.12) ||W||_{L^2(\Omega)} = 1$$

Since $W_n - \|V_{\varepsilon_n}\|_{L^2(\Omega)}^{-1} v_0 \in \widetilde{\mathcal{H}}_{\varepsilon_n}$, we have $T^j(W_n) = 0$ on Γ_0^j for all $j = 1, \ldots, k_1$, see (4.3). By continuity of the trace operator (2.4), we deduce that $T^j(W) = 0$ on Γ_0^j for all $j = 1, \ldots, k_1$, hence $W \in \widetilde{\mathcal{H}}_0$.

Let $w \in \widetilde{\mathcal{H}}_{0,0}$, where $\widetilde{\mathcal{H}}_{0,0}$ is defined in Lemma 3.4. For *n* sufficiently large, $w \in \widetilde{\mathcal{H}}_{\varepsilon_n}$ and $L_{\varepsilon_n}(w) = 0$, hence we can test (4.5) with *w*, thus obtaining

$$\int_{\Omega \setminus \Gamma_1} \nabla W_n \cdot \nabla w \, dx = \int_{\Omega \setminus \Gamma_{\varepsilon_n}} \nabla W_n \cdot \nabla w \, dx = -L_{\varepsilon_n}(w) = 0.$$

Letting $n \to \infty$ in the above identity, we obtain $\int_{\Omega \setminus \Gamma_0} \nabla W \cdot \nabla w \, dx = 0$ for all $w \in \widetilde{\mathcal{H}}_{0,0}$ and hence, by the density of $\widetilde{\mathcal{H}}_{0,0}$ in $\widetilde{\mathcal{H}}_0$ established in Lemma 3.4,

(4.13)
$$\int_{\Omega \setminus \Gamma_0} \nabla W \cdot \nabla w \, dx = 0 \quad \text{for all } w \in \widetilde{\mathcal{H}}_0.$$

Choosing w = W in (4.13), we conclude that W = 0, thus contradicting (4.12).

5. Asymptotic expansion of the eigenvalue variation

For every $\varepsilon \in [0,1]$, we consider the bilinear form $q_{\varepsilon} : \widetilde{\mathcal{H}}_{\varepsilon} \times \widetilde{\mathcal{H}}_{\varepsilon} \to \mathbb{R}$ defined as

(5.1)
$$q_{\varepsilon}(w_1, w_2) := \int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla w_1 \cdot \nabla w_2 \, dx$$

where $\mathcal{H}_{\varepsilon}$ is as in (2.5). To simplify notation, we denote by q_{ε} both the bilinear form defined above and the associated quadratic form

$$q_{\varepsilon}(w) = \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^2 \, dx = \|w\|_{\mathcal{H}_{\varepsilon}}^2, \quad w \in \widetilde{\mathcal{H}}_{\varepsilon}.$$

The following preliminary result can be obtained in a standard way from the compactness properties pointed out in Remark 3.1 and abstract spectral theory, see for example [16, Theorems 6.16 and 6.21, Proposition 8.20].

Proposition 5.1. Let $\varepsilon \in [0,1]$ and $\mathcal{F}_{\varepsilon} : \widetilde{\mathcal{H}}_{\varepsilon} \to \widetilde{\mathcal{H}}_{\varepsilon}$ be the linear operator defined as

(5.2)
$$q_{\varepsilon}(\mathcal{F}_{\varepsilon}(w_1), w_2) = (w_1, w_2)_{L^2(\Omega)}$$

Then

- (i) $\mathcal{F}_{\varepsilon}$ is symmetric, non-negative and compact; in particular 0 belongs to its spectrum $\sigma(\mathcal{F}_{\varepsilon})$.
- (ii) $\sigma(\mathcal{F}_{\varepsilon}) \setminus \{0\} = \{\mu_{n,\varepsilon}\}_{n \in \mathbb{N} \setminus \{0\}}, \text{ where } \mu_{n,\varepsilon} := 1/\lambda_{\varepsilon,n} \text{ for every } n \in \mathbb{N} \setminus \{0\}.$
- (iii) For every $\mu \in \mathbb{R}$ and $w \in \widetilde{\mathcal{H}}_{\varepsilon}$,

$$\left(\operatorname{dist}(\mu, \sigma(\mathcal{F}_{\varepsilon}))\right)^2 \leq \frac{q_{\varepsilon}(\mathcal{F}_{\varepsilon}(w) - \mu w)}{q_{\varepsilon}(w)}$$

Letting $n_0 \in \mathbb{N} \setminus \{0\}$, v_0 and $\lambda_0 = \lambda_{0,n_0}$ be as in Section 4, to prove an asymptotic expansion of the eigenvalue variation we further assume that

(5.3)
$$\lambda_0$$
 is simple as an eigenvalue of (1.5),

and, consequently, as an eigenvalue of (2.14). Therefore, the continuity result of [20, Theorem 1.2], see (1.7), implies that also $\lambda_{\varepsilon,n_0}$ is simple for ε sufficiently small. From now on, we denote

$$\lambda_{\varepsilon} = \lambda_{\varepsilon, n_0}.$$

For ε small, let $v_{\varepsilon} \in \widetilde{\mathcal{H}}_{\varepsilon}$ be the unique eigenfunction of (2.10) associated to the eigenvalue $\lambda_{\varepsilon} = \lambda_{\varepsilon,n_0}$ satisfying

(5.4)
$$\int_{\Omega} v_{\varepsilon}^2 dx = 1 \quad \text{and} \quad \int_{\Omega} v_{\varepsilon} v_0 dx > 0.$$

We denote as Π_{ε} the projection onto the one-dimensional space spanned by v_{ε} , i.e.

(5.5)
$$\Pi_{\varepsilon} : L^{2}(\Omega) \to \widetilde{\mathcal{H}}_{\varepsilon},$$
$$w \mapsto (w, v_{\varepsilon})_{L^{2}(\Omega)} v_{\varepsilon}.$$

Theorem 2.1 is contained in the following result, the proof of which is inspired by [3, Appendix A].

Theorem 5.2. Under assumption (5.3), the following asymptotic expansion holds:

(5.6)
$$\lambda_{\varepsilon} - \lambda_0 = 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(v_0) + o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2\right) \quad as \ \varepsilon \to 0^+,$$

where V_{ε} is as Proposition 4.2. Furthermore,

(5.7)
$$\|v_0 - V_{\varepsilon} - \Pi_{\varepsilon} (v_0 - V_{\varepsilon})\|_{\mathcal{H}_{\varepsilon}} = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}\right) \quad as \ \varepsilon \to 0^+$$

(5.8)
$$\|v_0 - \Pi_{\varepsilon}(v_0 - V_{\varepsilon})\|_{L^2(\Omega)} = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}\right) \quad as \ \varepsilon \to 0^+$$

(5.9)
$$\|\Pi_{\varepsilon}(v_0 - V_{\varepsilon})\|_{L^2(\Omega)}^2 = 1 + o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}\right) \quad as \ \varepsilon \to 0^+$$

Proof. Let $\psi_{\varepsilon} := v_0 - V_{\varepsilon}$. We recall that we are assuming that v_0 is real-valued and $||v_0||_{L^2(\Omega)} = 1$. From (4.4) and (2.14) it follows that $\psi_{\varepsilon} \in \widetilde{\mathcal{H}}_{\varepsilon}$ is a weak solution of the problem

$$\begin{cases} -\Delta\psi_{\varepsilon} = \lambda_{0}v_{0}, & \text{in } \Omega \setminus \Gamma_{\varepsilon}, \\ \psi_{\varepsilon} = 0, & \text{on } \partial\Omega, \\ T^{j}(\psi_{\varepsilon}) = 0, & \text{on } \Gamma^{j}_{\varepsilon} \text{ for all } j = 1, \dots, k_{1}, \\ T^{j}(\nabla\psi_{\varepsilon} \cdot \nu^{j}) = 0, & \text{on } \Gamma^{j}_{\varepsilon} \text{ for all } j = 1, \dots, k_{1}, \\ T^{j}(\psi_{\varepsilon}) = 0, & \text{on } S^{j}_{\varepsilon} \text{ for all } j = k_{1} + 1, \dots, k_{1} + k_{2}, \\ T^{j}(\nabla\psi_{\varepsilon} \cdot \nu^{j}) = 0, & \text{on } S^{j}_{\varepsilon} \text{ for all } j = k_{1} + 1, \dots, k_{1} + k_{2}, \end{cases}$$

in the sense that, letting q_{ε} be as in (5.1),

(5.10)
$$q_{\varepsilon}(\psi_{\varepsilon}, w) = \lambda_0 (v_0, w)_{L^2(\Omega)} \quad \text{for all } w \in \mathcal{H}_{\varepsilon}.$$

Let v_{ε} be an eigenfunction of (2.10) associated to λ_{ε} chosen as in (5.4). Let Π_{ε} be the projection operator onto the one-dimensional space spanned by v_{ε} defined in (5.5). Moreover, we define

(5.11)
$$\hat{v}_{\varepsilon} := \frac{\Pi_{\varepsilon}(\psi_{\varepsilon})}{\|\Pi_{\varepsilon}(\psi_{\varepsilon})\|_{L^{2}(\Omega)}}$$

From (5.10) we deduce that

(5.12)
$$q_{\varepsilon}(\psi_{\varepsilon}, w) - \lambda_0 (\psi_{\varepsilon}, w)_{L^2(\Omega)} = \lambda_0 (V_{\varepsilon}, w)_{L^2(\Omega)} \text{ for all } w \in \mathcal{H}_{\varepsilon}.$$

Choosing $w = \hat{v}_{\varepsilon}$ in (5.12), by (3.6) and (5.11) we obtain

(5.13)
$$(\lambda_{\varepsilon} - \lambda_0) (\psi_{\varepsilon}, \hat{v}_{\varepsilon})_{L^2(\Omega)} = \lambda_0 (V_{\varepsilon}, v_0)_{L^2(\Omega)} + \lambda_0 (V_{\varepsilon}, \hat{v}_{\varepsilon} - v_0)_{L^2(\Omega)}.$$

We claim that

(5.14)
$$\lambda_0 \int_{\Omega} V_{\varepsilon} v_0 \, dx = 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(v_0).$$

Indeed, an integration by parts yields

(5.15)
$$\int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla v_0 \cdot \nabla V_{\varepsilon} \, dx - \lambda_0 \int_{\Omega} v_0 V_{\varepsilon} \, dx$$
$$= \sum_{j=1}^{k_1} \int_{\Gamma_0^j} \left(-\gamma_+^j (V_{\varepsilon}) \gamma_+^j (\nabla v_0 \cdot \nu^j) + \gamma_-^j (V_{\varepsilon}) \gamma_-^j (\nabla v_0 \cdot \nu^j) \right) dS$$
$$+ \sum_{j=1}^{k_1+k_2} \int_{S_{\varepsilon}^j} \left(-\gamma_+^j (V_{\varepsilon}) \nabla v_0 \cdot \nu^j + \gamma_-^j (V_{\varepsilon}) \nabla v_0 \cdot \nu^j \right) dS$$
$$= -2 \sum_{j=1}^{k_1+k_2} \int_{S_{\varepsilon}^j} \gamma_+^j (V_{\varepsilon}) \nabla v_0 \cdot \nu^j \, dS + 2 \sum_{j=1}^{k_1+k_2} \int_{S_{\varepsilon}^j} v_0 \nabla v_0 \cdot \nu^j \, dS,$$

thanks to (4.4). Testing (5.10) with $V_{\varepsilon} - v_0$ we obtain

$$\int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla (V_{\varepsilon} - v_0)|^2 = -\lambda_0 \int_{\Omega} v_0 (V_{\varepsilon} - v_0) \, dx,$$

and hence, in view of (3.10),

(5.16)
$$\int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla V_{\varepsilon} \cdot \nabla v_0 \, dx = \frac{1}{2} \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla V_{\varepsilon}|^2 \, dx + \frac{\lambda_0}{2} \int_{\Omega} v_0 V_{\varepsilon} \, dx.$$

Combining (2.16), (2.17), (2.19), (5.15) and (5.16), we derive (5.14).

From (5.13) and (5.14) we deduce that, for all $\varepsilon \in (0, 1)$,

(5.17)
$$(\lambda_{\varepsilon} - \lambda_0) (\psi_{\varepsilon}, \hat{v}_{\varepsilon})_{L^2(\Omega)} = 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(v_0) + \lambda_0 (V_{\varepsilon}, \hat{v}_{\varepsilon} - v_0)_{L^2(\Omega)}$$

Now we study the asymptotics, as $\varepsilon \to 0^+$, of each term in (5.17). For the sake of clarity, we divide the rest of the proof into several steps.

Step 1. We claim that

(5.18)
$$|\lambda_{\varepsilon} - \lambda_{0}| = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} \right) \quad \text{as } \varepsilon \to 0^{+}$$

Letting $\mu_0 := \lambda_0^{-1}$ and $\mu_{\varepsilon} := \lambda_{\varepsilon}^{-1}$, since λ_0 is simple and $\lambda_{\varepsilon} \to \lambda_0$ by (1.7), we have

(5.19)
$$|\lambda_{\varepsilon} - \lambda_{0}| = \lambda_{\varepsilon}\lambda_{0}|\mu_{\varepsilon} - \mu_{0}| \le 2\lambda_{0}^{2}\operatorname{dist}(\mu_{0}, \sigma(\mathcal{F}_{\varepsilon})) \le 2\lambda_{0}^{2}\left(\frac{q_{\varepsilon}(\mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon})}{q_{\varepsilon}(\psi_{\varepsilon})}\right)^{1/2},$$

where the last inequality is justified by Proposition 5.1. Since $||v_0||_{L^2(\Omega)} = 1$, Proposition 4.4 and the Cauchy-Schwarz inequality imply that

(5.20)
$$q_{\varepsilon}(\psi_{\varepsilon}) = \lambda_0 + \int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla V_{\varepsilon}|^2 \, dx - 2 \int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla V_{\varepsilon} \cdot \nabla v_0 \, dx = \lambda_0 + o(1).$$

Furthermore, in view of (5.2) and (5.10) tested with $\mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_0 \psi_{\varepsilon}$,

$$q_{\varepsilon}(\mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon}) = -(V_{\varepsilon}, \mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon})_{L^{2}(\Omega)} + (v_{0}, \mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon})_{L^{2}(\Omega)} - q_{\varepsilon}(\mu_{0}\psi_{\varepsilon}, \mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon}) = -(V_{\varepsilon}, \mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon})_{L^{2}(\Omega)}.$$

Hence, by Proposition 3.2, Proposition 4.6 and the Cauchy-Schwarz inequality we conclude that

(5.21)
$$(q_{\varepsilon}(\mathcal{F}_{\varepsilon}(\psi_{\varepsilon}) - \mu_{0}\psi_{\varepsilon}))^{1/2} = o(||V_{\varepsilon}||_{\mathcal{H}_{\varepsilon}}) \quad \text{as } \varepsilon \to 0^{+}.$$

Claim (5.18) is proved by combining (5.19), (5.20), and (5.21).

Step 2. We claim that

(5.22)
$$q_{\varepsilon}(\psi_{\varepsilon} - \Pi_{\varepsilon}\psi_{\varepsilon}) = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2} \right) \quad \text{as } \varepsilon \to 0^{+}.$$

Let

(5.23)
$$\chi_{\varepsilon} := \psi_{\varepsilon} - \Pi_{\varepsilon} \psi_{\varepsilon} \quad \text{and} \quad \xi_{\varepsilon} := \mathcal{F}_{\varepsilon}(\chi_{\varepsilon}) - \mu_{\varepsilon} \chi_{\varepsilon}.$$

By definition we have

$$\chi_{\varepsilon} \in N_{\varepsilon} := \{ w \in \mathcal{H}_{\varepsilon} : (w, v_{\varepsilon})_{L^{2}(\Omega)} = 0 \}$$

and, since v_{ε} is an eigenfunction of (2.10), from (5.2) it follows that $\mathcal{F}_{\varepsilon}(w) \in N_{\varepsilon}$ for all $w \in N_{\varepsilon}$. Hence the operator

$$\widetilde{\mathcal{F}}_{\varepsilon} := \mathcal{F}_{\varepsilon} \Big|_{N_{\varepsilon}} : N_{\varepsilon} \to N_{\varepsilon}$$

is well-defined. Furthermore, it is easy to verify that $\widetilde{\mathcal{F}}_{\varepsilon}$ satisfies properties (i)-(iii) of Proposition 5.1 and $\sigma(\widetilde{\mathcal{F}}_{\varepsilon}) = \sigma(\mathcal{F}_{\varepsilon}) \setminus \{\mu_{\varepsilon}\}$. In particular, there exists a constant K > 0, which does not depends on ε , such that $\left(\operatorname{dist}(\mu_{\varepsilon}, \sigma(\widetilde{\mathcal{F}}_{\varepsilon}))\right)^2 \geq K$. Then, by (5.23),

(5.24)
$$q_{\varepsilon}(\psi_{\varepsilon} - \Pi_{\varepsilon}\psi_{\varepsilon}) = q(\chi_{\varepsilon}) \leq \frac{1}{K} (\operatorname{dist}(\mu_{\varepsilon}, \sigma(\widetilde{\mathcal{F}}_{\varepsilon})))^2 q_{\varepsilon}(\chi_{\varepsilon}) \leq \frac{1}{K} q_{\varepsilon}(\widetilde{\mathcal{F}}_{\varepsilon}(\chi_{\varepsilon}) - \mu_{\varepsilon}\chi_{\varepsilon}) = \frac{1}{K} q_{\varepsilon}(\xi_{\varepsilon}).$$

To estimate $q_{\varepsilon}(\xi_{\varepsilon})$ we use (5.12) and (3.6) tested with ξ_{ε} , thus obtaining

(5.25)
$$q_{\varepsilon}(\chi_{\varepsilon},\xi_{\varepsilon}) - \lambda_{\varepsilon} (\chi_{\varepsilon},\xi_{\varepsilon})_{L^{2}(\Omega)} = \lambda_{0} (V_{\varepsilon},\xi_{\varepsilon})_{L^{2}(\Omega)} + (\lambda_{0} - \lambda_{\varepsilon}) (\psi_{\varepsilon},\xi_{\varepsilon})_{L^{2}(\Omega)}.$$

From (5.2) and (5.25) we deduce that

$$q_{\varepsilon}(\xi_{\varepsilon}) = q_{\varepsilon}(\mathcal{F}_{\varepsilon}(\chi_{\varepsilon}), \xi_{\varepsilon}) - \mu_{\varepsilon}q_{\varepsilon}(\chi_{\varepsilon}, \xi_{\varepsilon}) = -\mu_{\varepsilon}[q_{\varepsilon}(\chi_{\varepsilon}, \xi_{\varepsilon}) - \lambda_{\varepsilon}q_{\varepsilon}(\mathcal{F}_{\varepsilon}(\chi_{\varepsilon}), \xi_{\varepsilon})]$$
$$= -\frac{\lambda_{0}}{\lambda_{\varepsilon}}(V_{\varepsilon}, \xi_{\varepsilon})_{L^{2}(\Omega)} - \frac{(\lambda_{0} - \lambda_{\varepsilon})}{\lambda_{\varepsilon}}(\psi_{\varepsilon}, \xi_{\varepsilon})_{L^{2}(\Omega)}.$$

From the Cauchy-Schwarz inequality, Proposition 3.2, and (1.7) it follows that

(5.26)
$$(q_{\varepsilon}(\xi_{\varepsilon}))^{1/2} \le C \left(\|V_{\varepsilon}\|_{L^{2}(\Omega)} + |\lambda_{\varepsilon} - \lambda_{0}| \|\psi_{\varepsilon}\|_{L^{2}(\Omega)} \right)$$

for some constant C > 0 which does not depend on ε . Furthermore, (5.12) tested with ψ_{ε} , (5.20), Proposition 3.2, and Proposition 4.4 yield

$$\|\psi_{\varepsilon}\|_{L^{2}(\Omega)}^{2} - 1 = -(V_{\varepsilon}, \psi_{\varepsilon})_{L^{2}(\Omega)} + o(1) = o(1) \quad \text{as } \varepsilon \to 0^{+}.$$

Then (5.22) follows from Proposition 4.6, (5.18), (5.24), and (5.26). Estimate (5.7) is thereby proved.

Step 3. We claim that

(5.27)
$$\|v_0 - \hat{v}_{\varepsilon}\|_{L^2(\Omega)} = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}\right) \quad \text{as } \varepsilon \to 0^+.$$

By (5.11)

(5.28)
$$v_0 - \hat{v}_{\varepsilon} = v_0 - \frac{\Pi_{\varepsilon}\psi_{\varepsilon}}{\|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^2(\Omega)}} = \frac{1}{\|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^2(\Omega)}} \left(\left(\|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^2(\Omega)} - 1\right)v_0 + v_0 - \Pi_{\varepsilon}\psi_{\varepsilon} \right).$$

Furthermore, from the definition of ψ_{ε} , Proposition 4.6, Proposition 3.2 and (5.22) it follows that

(5.29)
$$\|v_0 - \Pi_{\varepsilon} \psi_{\varepsilon}\|_{L^2(\Omega)} \le \|v_0 - \psi_{\varepsilon}\|_{L^2(\Omega)} + \|\psi_{\varepsilon} - \Pi_{\varepsilon} \psi_{\varepsilon}\|_{L^2(\Omega)} = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}\right) \quad \text{as } \varepsilon \to 0^+$$

thus proving (5.8). Since $||v_0||_{L^2(\Omega)} = 1$, (5.29) and the Cauchy-Schwarz inequality imply that

(5.30)
$$\|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^{2}(\Omega)}^{2} = \|v_{0} - \Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^{2}(\Omega)}^{2} + \|v_{0}\|_{L^{2}(\Omega)}^{2} - 2(v_{0} - \Pi_{\varepsilon}\psi_{\varepsilon}, v_{0})_{L^{2}(\Omega)} = 1 + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}})$$

as $\varepsilon \to 0^{+}$, thus proving estimate (5.9). Combining (5.28), (5.29) and (5.30) we obtain (5.27).

Step 4. We claim that

(5.31)
$$(\psi_{\varepsilon}, \hat{v}_{\varepsilon})_{L^{2}(\Omega)} = 1 + o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} \right) \quad \text{as } \varepsilon \to 0^{+}$$

Indeed, by (5.11) we have

$$(\psi_{\varepsilon}, \hat{v}_{\varepsilon})_{L^{2}(\Omega)} = \frac{(\psi_{\varepsilon} - \Pi_{\varepsilon}\psi_{\varepsilon}, \Pi_{\varepsilon}\psi_{\varepsilon})_{L^{2}(\Omega)} + \|\Pi_{\varepsilon}\psi_{\varepsilon}\|^{2}_{L^{2}(\Omega)}}{\|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^{2}(\Omega)}}$$

Hence claim (5.31) follows from (5.22) and (5.30).

Putting together Proposition 4.6, (5.17), (5.27), and (5.31), we finally obtain

$$\lambda_{\varepsilon} - \lambda_{0} = (1 + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}})) \left(2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(v_{0}) + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2})\right)$$

$$= 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(v_{0}) + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2}) + \left(\mathcal{E}_{\varepsilon} - L_{\varepsilon}(v_{0})\right)o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}})$$

$$= 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(v_{0}) + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2}) \quad \text{as } \varepsilon \to 0^{+},$$

having used in the last estimate the fact that

$$\mathcal{E}_{\varepsilon} - L_{\varepsilon}(v_0) = o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}) \text{ as } \varepsilon \to 0^+,$$

due to (5.14) and Proposition 4.6. Expansion (5.6) is thereby proved.

6. Blow-up Analysis for k odd

In this section we assume that k, and consequently k_1 , are odd and we perform a blow-up analysis for the solution V_{ε} of problem (4.4). In order to characterize the functional space containing the limit profile, we first need a Hardy-type inequality, for the validity of which the assumption that k is odd is crucial.

6.1. A Hardy type inequality for functions jumping on an odd number of lines. Let $\hat{\mathcal{X}}$ and $\tilde{\mathcal{H}}$ be the functional spaces defined in (2.24) and (2.25), respectively. To prove a Hardy-type inequality in $\mathbb{R}^2 \setminus D_1$ for functions in $\tilde{\mathcal{X}}$, we first need the following Hardy inequality on annuli for functions jumping on an odd number of lines. For every r > 0, we define

$$\mathcal{X}_r := \{ w \in H^1((D_{2r} \setminus D_r) \setminus \Gamma_0) : T^j(w) = 0 \text{ on } \Gamma_0^j \text{ for all } j = 1, \dots, k_1 \}.$$

Lemma 6.1. Let k and k_1 be odd. There exists a constant $C_H > 0$ such that, for every r > 0 and $w \in \widetilde{\mathcal{X}}_r$,

(6.1)
$$r^{-2} \int_{D_{2r} \setminus D_r} w^2 \, dx \le C_H \int_{(D_{2r} \setminus D_r) \setminus \Gamma_0} |\nabla w|^2 \, dx$$

and

(6.2)
$$\int_{D_{2r}\setminus D_r} \frac{w^2}{|x|^2} dx \le C_H \int_{(D_{2r}\setminus D_r)\setminus \Gamma_0} |\nabla w|^2 dx.$$

Proof. Inequality (6.2) is a direct consequence of (6.1).

Let us first prove (6.1) for r = 1. We argue by contradiction and assume that there exists a sequence $\{w_n\}_{n \in \mathbb{N}} \subset \widetilde{\mathcal{X}}_1$ such that, for all $n \in \mathbb{N}$,

(6.3)
$$\int_{D_2 \setminus D_1} w_n^2 \, dx = 1 \quad \text{and} \quad \int_{(D_2 \setminus D_1) \setminus \Gamma_0} |\nabla w_n|^2 \, dx < \frac{1}{n}.$$

Hence $\{w_n\}_{n\in\mathbb{N}}$ is bounded in $\widetilde{\mathcal{X}}_1$ and, up to a subsequence, $w_n \rightharpoonup w$ weakly in $\widetilde{\mathcal{X}}_1$ for some $w \in \widetilde{\mathcal{X}}_1$. From (6.3) and weak lower semi-continuity of the L^2 -norm, we have $\nabla w \equiv 0$ in $(D_2 \setminus D_1) \setminus \Gamma_0$; furthermore, reasoning as in Remark 3.1, the natural embedding of $H^1((D_2 \setminus D_1) \setminus \Gamma_0)$) into $L^2(D_2 \setminus D_1)$ is compact, hence $\|w\|_{L^2(D_2 \setminus D_1)} = 1$. It follows that w is constant on each connected component of $(D_2 \setminus D_1) \setminus \Gamma_0$ and $w \neq 0$. Since $(D_2 \setminus D_1) \setminus \Gamma_0$ has k_1 connected components and k_1 is odd, a contradiction arises from the condition $T^j(w) = 0$, which is satisfied on Γ_0^j for all $j = 1, \ldots, k_1$.

For every r > 0 and $w \in \widetilde{\mathcal{X}}_r$, it is enough to write the proved inequality for the scaled function w(rx) to obtain (6.1).

We draw attention to the fact that the constant C_H in Lemma 6.1 does not depend on r. Hence, summing over annuli that fill $\mathbb{R}^2 \setminus D_1$, we obtain the following result.

Proposition 6.2. Let k and k_1 be odd. Let $C_H > 0$ be as in Lemma 6.1. Then, for every $w \in \mathcal{X}$,

(6.4)
$$\int_{\mathbb{R}^2 \setminus D_1} \frac{w^2}{|x|^2} dx \le C_H \int_{(\mathbb{R}^2 \setminus D_1) \setminus \Gamma_1} |\nabla w|^2 dx$$

Furthermore, there exists a constant $C'_H > 0$ such that, for all $w \in \widetilde{\mathcal{X}}$,

(6.5)
$$\int_{D_1} w^2 \, dx \le C'_H \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx.$$

Proof. If $w \in \widetilde{\mathcal{X}}$, then $w \in \widetilde{\mathcal{X}}_r$ for all r > 1. Hence, by (6.2),

$$\int_{\mathbb{R}^2 \setminus D_1} \frac{w^2}{|x|^2} dx = \sum_{h=0}^{\infty} \int_{D_{2^{h+1}} \setminus D_{2^h}} \frac{w^2}{|x|^2} dx$$
$$\leq C_H \sum_{h=0}^{\infty} \int_{(D_{2^{h+1}} \setminus D_{2^h}) \setminus \Gamma_0} |\nabla w|^2 dx = C_H \int_{(\mathbb{R}^2 \setminus D_1) \setminus \Gamma_1} |\nabla w|^2 dx$$

thus proving (6.4).

By integrating the identity $\operatorname{div}(u^2 x) = 2u\nabla u \cdot x + 2u^2$ on each subset of D_1 obtained by cutting along the lines Σ^{j} , $j = 1 \dots k_1 + k_2$, and using the Divergence Theorem, we can prove that, for all $w \in \mathcal{X}$,

$$\int_{D_1} w^2 \, dx \le \int_{\partial D_1} w^2 \, dS + \int_{D_1 \setminus \Gamma_1} |\nabla w|^2 \, dx$$

Then, by continuity of the trace operator from $H^1((D_2 \setminus D_1) \setminus \Gamma_0)$ into $L^2(\partial D_1)$ and (6.4), there exists a positive constant C > 0 such that

$$\begin{split} \int_{D_1} w^2 \, dx &\leq C \left(\int_{D_2 \setminus D_1} w^2 \, dx + \int_{(D_2 \setminus D_1) \setminus \Gamma_1} |\nabla w|^2 \, dx \right) + \int_{D_1 \setminus \Gamma_1} |\nabla w|^2 \, dx \\ &\leq 4C \int_{D_2 \setminus D_1} \frac{w^2}{|x|^2} \, dx + (C+1) \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx \leq (4CC_H + C + 1) \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx, \end{split}$$

this proving (6.5).

From Proposition 6.2 it follows that

(6.6)
$$\|w\|_{\widetilde{\mathcal{X}}} := \left(\int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx\right)^{1/2}$$

is a norm on $\widetilde{\mathcal{X}}$ and $\widetilde{\mathcal{X}}$ is a Hilbert space with respect to the corresponding scalar product. Proposition 6.2 also ensures that the restriction operator

(6.7)
$$\widetilde{\mathcal{X}} \to H^1(D_\rho \setminus \Gamma_1)$$

is continuous with respect to the norm defined in (6.6) for every $\rho > 0$. Hence, for every $p \in [1, +\infty)$, the trace operators

(6.8)
$$\gamma^j_+ : \widetilde{\mathcal{X}} \to L^p(S^j_1) \text{ and } \gamma^j_- : \widetilde{\mathcal{X}} \to L^p(S^j_1)$$

are well-defined and continuous with respect to the norm $\|\cdot\|_{\widetilde{\mathcal{X}}}$. In particular, since $\widetilde{\mathcal{H}} \subset \widetilde{\mathcal{X}}$,

(6.9)
$$\sup_{w \in \widetilde{\mathcal{H}} \setminus \{0\}} \frac{\|\gamma_{+}^{j}(w)\|_{L^{p}(S_{1}^{j})}^{2}}{\|w\|_{\widetilde{\mathcal{X}}}^{2}} < +\infty \quad \text{for every } p \in [1, +\infty) \text{ and } j = 1, \dots, k_{1} + k_{2}.$$

Using (6.4), we prove now that functions in $\widetilde{\mathcal{H}}$ can be approximated with functions with compact support. To this aim, we define

$$\mathcal{H}_c := \{ w \in \mathcal{H} : \text{ there exists } r > 0 \text{ such that } w \equiv 0 \text{ on } \mathbb{R}^2 \setminus D_r \}.$$

Proposition 6.3. $\widetilde{\mathcal{H}}_c$ is dense in $\widetilde{\mathcal{H}}$.

Proof. For every r > 1, let η_r be a cut-off function as in (4.7). If $w \in \mathcal{H}$, it is clear that $\{\eta_r w\}_{r>1} \subset \widetilde{\mathcal{H}}_c$; moreover, by (6.4) we have $\frac{w}{|x|} \in L^2(\mathbb{R}^2 \setminus D_1)$ and hence

$$\int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \eta_r|^2 w^2 \, dx \le 16 \int_{D_{2r} \setminus D_r} \frac{w^2}{|x|^2} \, dx \to 0^+ \quad \text{as } r \to \infty.$$

This implies that $\nabla(\eta_r w) \to \nabla w$ in $L^2(\mathbb{R}^2 \setminus \Gamma_1)$ and hence $\eta_r w \to w$ in $\widetilde{\mathcal{H}}$.

6.2. Limit profile for blown-up potentials. In this subsection, we introduce and characterize the function \tilde{V} appearing as limit profile in a blow-up analysis for the potentials V_{ε} .

Proposition 6.4. There exists a unique solution $\tilde{V} \in \tilde{\mathcal{X}}$ to the minimization problem (2.30). Furthermore, \tilde{V} satisfies

(6.10)
$$\begin{cases} V - \eta \Psi_0 \in \mathcal{H} \\ \int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V} \cdot \nabla w \, dx = -2 \sum_{j=1}^{k_1 + k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(w) \, dS \quad \text{for all } w \in \widetilde{\mathcal{H}}. \end{cases}$$

Proof. Since $\nabla \Psi_0 \in L^p(S_1^j)$ for all $p \in [1, 2)$, by continuity of the trace operators in (6.8) we have that the linear functional L defined in (2.27) is well-defined and continuous. Then the convex functional J defined in (2.28) is continuous and coercive on the closed and convex set $\eta \Psi_0 + \tilde{\mathcal{H}} = \{w \in \tilde{\mathcal{X}} : w - \eta \Psi_0 \in \tilde{\mathcal{H}}\}$. Therefore (2.30) admits a solution \tilde{V} , which satisfies (6.10).

If \widetilde{V}_1 and \widetilde{V}_2 are solutions of (6.10), then we may take the difference between (6.10) for \widetilde{V}_1 and (6.10) for \widetilde{V}_2 , both tested with $\widetilde{V}_1 - \widetilde{V}_2 \in \widetilde{\mathcal{H}}$, and conclude that $\widetilde{V}_1 = \widetilde{V}_2$ thanks to (6.4). Hence \widetilde{V} is the unique solution to (6.10).

6.3. An equivalent characterization of $\mathcal{E}_{\varepsilon}$. In this subsection, we obtain an equivalent characterization of the energy $\mathcal{E}_{\varepsilon}$ introduced in (2.19), which will be used to improve (4.9) and obtain an optimal estimate for $|\mathcal{E}_{\varepsilon}|$ in the case k odd.

Proposition 6.5. Let $\eta_{\varepsilon} \in C_{c}^{\infty}(\mathbb{R}^{2})$ be a cut-off function as in (4.7) with $r = \varepsilon$. Then, for every $\varepsilon \in (0, 1]$,

$$(6.11) \qquad \mathcal{E}_{\varepsilon} = -\frac{1}{2} \sup_{w \in \widetilde{\mathcal{H}}_{\varepsilon} \setminus \{0\}} \frac{\left(\int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla w \cdot \nabla(\eta_{\varepsilon} v_{0}) + L_{\varepsilon}(w)\right)^{2}}{\int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^{2} dx} + \frac{1}{2} \int_{\Omega \setminus \Gamma_{0}} |\nabla(\eta_{\varepsilon} v_{0})|^{2} dx + L_{\varepsilon}(v_{0}).$$

Proof. Since $\mathcal{E}_{\varepsilon}$ is the infimum in (4.2) and $\varphi - v_0 \in \widetilde{\mathcal{H}}_{\varepsilon}$ if and only if $\varphi - \eta_{\varepsilon} v_0 \in \widetilde{\mathcal{H}}_{\varepsilon}$, we have

(6.12)
$$\mathcal{E}_{\varepsilon} = \inf_{w \in \widetilde{\mathcal{H}}_{\varepsilon}} J_{\varepsilon}(w + \eta_{\varepsilon} v_0) = \inf_{w \in \widetilde{\mathcal{H}}_{\varepsilon} \setminus \{0\}} \left(\inf_{t \in [0, +\infty)} J_{\varepsilon}(tw + \eta_{\varepsilon} v_0) \right).$$

Moreover, by (2.17)

$$\begin{split} J_{\varepsilon}(tw+\eta_{\varepsilon}v_{0}) &= \frac{t^{2}}{2} \int_{\Omega\setminus\Gamma_{\varepsilon}} |\nabla w|^{2} \, dx + t \left(\int_{\Omega\setminus\Gamma_{\varepsilon}} \nabla w \cdot \nabla(\eta_{\varepsilon}v_{0}) \, dx + L_{\varepsilon}(w) \right) \\ &+ \frac{1}{2} \int_{\Omega\setminus\Gamma_{0}} |\nabla(\eta_{\varepsilon}v_{0})|^{2} \, dx + L_{\varepsilon}(v_{0}). \end{split}$$

Hence, for every $w \in \widetilde{\mathcal{H}}_{\varepsilon} \setminus \{0\}$,

$$\inf_{t\in[0,+\infty)} J_{\varepsilon}(tw+\eta_{\varepsilon}v_0) = -\frac{1}{2} \frac{\left(\int_{\Omega\setminus\Gamma_{\varepsilon}} \nabla w \cdot \nabla(\eta_{\varepsilon}v_0) \, dx + L_{\varepsilon}(w)\right)^2}{\int_{\Omega\setminus\Gamma_{\varepsilon}} |\nabla w|^2 \, dx} + \frac{1}{2} \int_{\Omega\setminus\Gamma_0} |\nabla(\eta_{\varepsilon}v_0)|^2 \, dx + L_{\varepsilon}(v_0),$$

which implies (6.11) in view of (6.12).

Proposition 6.6. Let k and k_1 be odd and $m \in \mathbb{N}$ be as in Proposition 3.6 for $v = v_0$. Then

$$\mathcal{E}_{\varepsilon} = O(\varepsilon^m) \quad as \ \varepsilon \to 0^+.$$

1~

Proof. From Proposition 6.5 and the Cauchy-Schwarz inequality it follows that

$$(6.13) \quad |\mathcal{E}_{\varepsilon}| \leq \frac{1}{2} \sup_{w \in \widetilde{\mathcal{H}}_{\varepsilon} \setminus \{0\}} \frac{\left(\int_{\Omega \setminus \Gamma_{\varepsilon}} \nabla w \cdot \nabla(\eta_{\varepsilon} v_{0}) + L_{\varepsilon}(w)\right)^{2}}{\int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^{2} dx} + \frac{1}{2} \int_{\Omega \setminus \Gamma_{0}} |\nabla(\eta_{\varepsilon} v_{0})|^{2} dx + |L_{\varepsilon}(v_{0})|$$
$$\leq \sup_{w \in \widetilde{\mathcal{H}}_{\varepsilon} \setminus \{0\}} \frac{|L_{\varepsilon}(w)|^{2}}{\int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^{2} dx} + \frac{3}{2} \int_{\Omega \setminus \Gamma_{0}} |\nabla(\eta_{\varepsilon} v_{0})|^{2} dx + |L_{\varepsilon}(v_{0})|.$$

From (3.12) and (2.16) it follows that

(6.14)
$$\int_{\Omega\setminus\Gamma_0} |\nabla(\eta_{\varepsilon}v_0)|^2 \, dx \le 2 \int_{D_{2\varepsilon}} |\nabla\eta_{\varepsilon}|^2 v_0^2 \, dx + 2 \int_{D_{2\varepsilon}\setminus\Gamma_0} |\nabla v_0|^2 \, dx = O(\varepsilon^m) \quad \text{as } \varepsilon \to 0^+$$

and

(6.15)
$$|L_{\varepsilon}(v_0)| = O(\varepsilon^m) \text{ as } \varepsilon \to 0^+.$$

By (2.16), the Hölder inequality, and (3.12), for every $p \in (1,2)$ and $p' = \frac{p}{p-1}$ we have

$$(6.16) \qquad \sup_{w\in\widetilde{\mathcal{H}}_{\varepsilon}\backslash\{0\}} \frac{|L_{\varepsilon}(w)|^{2}}{\int_{\Omega\backslash\Gamma_{\varepsilon}}|\nabla w|^{2}\,dx} \leq 4(k_{1}+k_{2})\sup_{w\in\widetilde{\mathcal{H}}_{\varepsilon}\backslash\{0\}} \frac{\sum_{j=1}^{k_{1}+k_{2}}\left(\int_{S_{\varepsilon}^{j}}|\nabla v_{0}||\gamma_{+}^{j}(w)|\,dS\right)^{2}}{\int_{\Omega\backslash\Gamma_{\varepsilon}}|\nabla w|^{2}\,dx}$$
$$\leq 4(k_{1}+k_{2})\sum_{j=1}^{k_{1}+k_{2}}\left(\int_{S_{\varepsilon}^{j}}|\nabla v_{0}|^{p}\,dS\right)^{2/p}\sup_{w\in\widetilde{\mathcal{H}}_{\varepsilon}\backslash\{0\}}\frac{\|\gamma_{+}^{j}(w)\|_{L^{p'}(S_{\varepsilon}^{j})}^{2}}{\int_{\Omega\backslash\Gamma_{\varepsilon}}|\nabla w|^{2}\,dx}$$
$$= O\left(\varepsilon^{m-2+\frac{2}{p}}\right)\sum_{j=1}^{k_{1}+k_{2}}\sup_{w\in\widetilde{\mathcal{H}}_{\varepsilon}\backslash\{0\}}\frac{\|\gamma_{+}^{j}(w)\|_{L^{p'}(S_{\varepsilon}^{j})}^{2}}{\int_{\Omega\backslash\Gamma_{\varepsilon}}|\nabla w|^{2}\,dx}.$$

A change of variables and (6.9) yield

$$\sup_{w\in\widetilde{\mathcal{H}}_{\varepsilon}\setminus\{0\}}\frac{\|\gamma^{j}_{+}(w)\|^{2}_{L^{p'}(S^{j}_{\varepsilon})}}{\int_{\Omega\setminus\Gamma_{\varepsilon}}|\nabla w|^{2}\,dx}\leq\varepsilon^{2/p'}\,\sup_{v\in\widetilde{\mathcal{H}}\setminus\{0\}}\frac{\|\gamma^{j}_{+}(v)\|^{2}_{L^{p'}(S^{j}_{1})}}{\|v\|^{2}_{\widetilde{\mathcal{X}}}}=O(\varepsilon^{2/p'})\quad\text{as }\varepsilon\to0^{+},$$

hence from (6.16) we deduce that

(6.17)
$$\sup_{w \in \widetilde{\mathcal{H}}_{\varepsilon} \setminus \{0\}} \frac{|L_{\varepsilon}(w)|^2}{\int_{\Omega \setminus \Gamma_{\varepsilon}} |\nabla w|^2 \, dx} = O(\varepsilon^m) \quad \text{as } \varepsilon \to 0^+.$$

The conclusion follows by combining estimates (6.13), (6.14), (6.15), and (6.17).

6.4. Blow-up analysis. Let k and k_1 be odd and $m \in \mathbb{N}$ be as in Proposition 3.6 for $v = v_0$. For every $\varepsilon \in (0, 1]$, letting V_{ε} be as Proposition 4.2, we define

(6.18)
$$\widetilde{V}_{\varepsilon}(x) := \varepsilon^{-\frac{m}{2}} V_{\varepsilon}(\varepsilon x) \quad \text{and} \quad \widetilde{V}_{0,\varepsilon}(x) := \varepsilon^{-\frac{m}{2}} v_0(\varepsilon x).$$

Extending trivially $\widetilde{V}_{\varepsilon}$ and $\widetilde{V}_{0,\varepsilon}$ in $\mathbb{R}^2 \setminus \Omega$, we have $\widetilde{V}_{\varepsilon}, \widetilde{V}_{0,\varepsilon} \in \widetilde{\mathcal{X}}$. Moreover

(6.19)
$$\widetilde{V}_{\varepsilon} - \widetilde{V}_{0,\varepsilon} \in \widetilde{\mathcal{H}}$$

and, by (4.5) and Proposition 6.3,

(6.20)
$$\int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V}_{\varepsilon} \cdot \nabla w \, dx = -2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \widetilde{V}_{0,\varepsilon} \cdot \nu^j \gamma^j_+(w) \, dS \quad \text{for all } w \in \widetilde{\mathcal{H}}.$$

Let Ψ_0 be as in (2.26). From (3.11) it follows that, for every $j = 1, \ldots, k_1 + k_2$,

(6.21)
$$\nabla \widetilde{V}_{0,\varepsilon}(x) \cdot \nu^j \to \nabla \Psi_0(x) \cdot \nu^j$$

as $\varepsilon \to 0^+$ for every $x \in S_1^j$, with (6.22)

 ∇T

$$= \begin{cases} \beta \frac{m}{2} |x|^{\frac{m}{2} - 1} f(\alpha^{j}) \cos\left(\frac{m}{2}(\alpha^{j} - \alpha_{0})\right), & \text{if } j = 1, \dots, k_{1}, \\ \beta \frac{m}{2} |x|^{\frac{m}{2} - 1} f(\alpha^{j}) \cos\left(\frac{m}{2}(\alpha^{j} - \alpha_{0})\right), & \text{if } x \in (S_{1}^{j})', \ j = k_{1} + 1, \dots, k_{1} + k_{2}, \\ -\beta \frac{m}{2} |x|^{\frac{m}{2} - 1} f(\alpha^{j} + \pi) \cos\left(\frac{m}{2}(\alpha^{j} + \pi - \alpha_{0})\right), & \text{if } x \in (S_{1}^{j})'', \ j = k_{1} + 1, \dots, k_{1} + k_{2}. \end{cases}$$

where, for every $j \in k_1, \ldots, k_1 + k_2$,

$$(S_1^j)' := \{ta^j : t \in [0,1]\}, \quad (S_1^j)'' := \{ta^{j+k_2} : t \in [0,1]\}.$$

On the other hand, (3.12) implies that

(6.23)
$$|\nabla \widetilde{V}_{0,\varepsilon}(x)| \le C|x|^{\frac{m}{2}-1} \quad \text{in } \mathbb{R}^2 \setminus \Gamma_0$$

From (6.21) and (6.23) we deduce that, for every $j = 1, ..., k_1 + k_2$ and $p \in [1, 2)$,

(6.24) $\nabla \Psi_0 \cdot \nu^j \in L^p(S_1^j) \text{ and } \nabla \widetilde{V}_{0,\varepsilon} \cdot \nu^j \to \nabla \Psi_0 \cdot \nu^j \text{ in } L^p(S_1^j) \text{ as } \varepsilon \to 0^+.$

Furthermore, by (3.13) we know that

(6.25)
$$\widetilde{V}_{0,\varepsilon} \to \Psi_0 \quad \text{in } H^1(D_\rho \setminus \Gamma_0) \text{ for all } \rho > 0$$

Proposition 6.7. Let k and k_1 be odd and $m \in \mathbb{N}$ be as in Proposition 3.6 for $v = v_0$. For every $\varepsilon \in (0, 1]$, let V_{ε} be as Proposition 4.2 and $\widetilde{V}_{\varepsilon}$ as in (6.18). Then

(6.26) $\widetilde{V}_{\varepsilon} \to \widetilde{V} \quad strongly \ in \ \widetilde{\mathcal{X}} \ as \ \varepsilon \to 0^+,$

where $\widetilde{V} \in \widetilde{\mathcal{X}}$ is the unique solution to the minimization problem (2.30) (and then to (6.10), see Proposition 6.4).

Proof. Taking into account (3.12), (2.19), and (6.18), a change of variables, (6.9), the Hölder inequality, and Proposition 6.6 imply that

$$(6.27) \quad \|\widetilde{V}_{\varepsilon}\|_{\widetilde{\mathcal{X}}}^{2} = \int_{\mathbb{R}^{2}\backslash\Gamma_{1}} |\nabla\widetilde{V}_{\varepsilon}|^{2} dx = \varepsilon^{-m} \|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2} = \varepsilon^{-m} (2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(V_{\varepsilon}))$$

$$\leq O(1) + 4\varepsilon^{-m} \sum_{j=1}^{k_{1}+k_{2}} \int_{S_{\varepsilon}^{j}} |\nabla v_{0}| |\gamma_{+}^{j}(V_{\varepsilon})| dS = O(1) + O(1) \sum_{j=1}^{k_{1}+k_{2}} \int_{S_{1}^{j}} |x|^{\frac{m}{2}-1} |\gamma_{+}^{j}(\widetilde{V}_{\varepsilon})| dS$$

$$= O(1) + O(1) \|\widetilde{V}_{\varepsilon}\|_{\widetilde{\mathcal{X}}}, \text{ as } \varepsilon \to 0^{+}.$$

Hence $\{\widetilde{V}_{\varepsilon}\}_{\varepsilon\in\{0,1]}$ is bounded in $\widetilde{\mathcal{X}}$. It follows that, for any sequence $\{\varepsilon_n\}_n$ such that $\varepsilon_n \to 0$ as $n \to \infty$, there exist a subsequence, still denoted by $\{\varepsilon_n\}_n$, and $V \in \widetilde{\mathcal{X}}$ such that $\widetilde{V}_{\varepsilon_n} \to V$ weakly in $\widetilde{\mathcal{X}}$ as $n \to \infty$. Therefore, from (6.20), (6.9), and (6.24) we deduce that V solves the variational equation in (6.10). Furthermore, by (6.19) we have $\widetilde{V}_{\varepsilon} - \eta \widetilde{V}_{0,\varepsilon} \in \widetilde{\mathcal{H}}$, hence (6.25) ensures that V satisfies the condition $V - \eta \Psi_0 \in \widetilde{\mathcal{H}}$. By the uniqueness part of Proposition 6.4 we conclude that $V = \widetilde{V}$.

Since $\widetilde{V} - \eta \Psi_0 \in \widetilde{\mathcal{H}}$, we may test (6.10) with $\widetilde{V} - \eta \Psi_0$, thus obtaining

(6.28)
$$\int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V}|^2 \, dx = \int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V} \cdot \nabla(\eta \Psi_0) \, dx - 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j (\widetilde{V} - \eta \Psi_0) \, dS.$$

On the other hand, testing (6.20) with $\widetilde{V}_{\varepsilon_n} - \eta \widetilde{V}_{0,\varepsilon_n} \in \widetilde{\mathcal{H}}$ we obtain

$$(6.29) \quad \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V}_{\varepsilon_n}|^2 \, dx = \int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V}_{\varepsilon_n} \cdot \nabla (\eta \widetilde{V}_{0,\varepsilon_n}) \, dx - 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \widetilde{V}_{0,\varepsilon_n} \cdot \nu^j \gamma_+^j (\widetilde{V}_{\varepsilon_n} - \eta \widetilde{V}_{0,\varepsilon_n}) \, dS.$$

In view of the weak convergence $\widetilde{V}_{\varepsilon_n} \to \widetilde{V}$ in $\widetilde{\mathcal{X}}$, (6.25), (6.24), and the continuity of the trace operators (6.8), the limit of the right hand side of (6.29) as $n \to \infty$ is equal to the right hand side

of (6.28), thus proving that $\widetilde{V}_{\varepsilon_n} \to \widetilde{V}$ strongly in $\widetilde{\mathcal{X}}$ as $n \to \infty$ by. Since \widetilde{V} is the unique solution of (6.10), (6.26) follows from the Urysohn Subsequence Principle.

In view of the blow-up analysis performed above, we are in position to prove Theorem 2.2.

Proof of Theorem 2.2. From (2.17), (2.19), (6.18), and a change of variables it follows that

(6.30)
$$\varepsilon^{-m} \mathcal{E}_{\varepsilon} = \frac{1}{2} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V_{\varepsilon}}|^2 \, dx + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \widetilde{V}_{0,\varepsilon} \cdot \nu^j \gamma_+^j(\widetilde{V_{\varepsilon}}) \, dS.$$

The convergences (6.26) and (6.24), together with the continuity of the trace operators in (6.8), allow us to pass to the limit in the right hand side of (6.30), thus yielding

(6.31)
$$\lim_{\varepsilon \to 0^+} \varepsilon^{-m} \mathcal{E}_{\varepsilon} = \frac{1}{2} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V}|^2 \, dx + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) = \mathcal{E}_{V_1}(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) = \mathcal{E}_{V_1}(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) = \mathcal{E}_{V_1}(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) = \mathcal{E}_{V_1}(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) = \mathcal{E}_{V_1}(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(\widetilde{V}) \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \, dS = J(\widetilde{V}) + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0$$

and proving claim (i). Furthermore, by (6.18), a change of variable, (6.24), and (6.25), we have

(6.32)
$$\varepsilon^{-m} L_{\varepsilon}(v_0) = 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \widetilde{V}_{0,\varepsilon} \cdot \nu^j \gamma^j_+(\widetilde{V}_{0,\varepsilon}) \, dS$$
$$= 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma^j_+(\Psi_0) \, dS + o(1) = L(\Psi_0) + o(1) \quad \text{as } \varepsilon \to 0^+.$$

Claim (ii) follows from (2.20), (6.31), (6.32), and estimate (6.27), which in particular ensures that $\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 = O(\varepsilon^m)$ as $\varepsilon \to 0^+$.

6.5. Continuity of $\mathcal{E} - L(\Psi_0)$ with respect to rotations of poles. In this subsection we prove the continuity of $\mathcal{E} - L(\Psi_0)$ with respect to rotations of the configuration of poles. We fix a configuration of poles $\{a^j\}$ as in (1.1). Then, for every $\zeta \in [-\pi, \pi)$, we define $\Psi_0^{(\zeta)}$, $L^{(\zeta)}(\Psi_0^{(\zeta)})$ and $\mathcal{E}^{(\zeta)}$ as in (2.26), (2.27), and (2.31), respectively, for a rotated configuration of poles $\{a^j_{\zeta}\}$, where a^j_{ζ} are defined as in (1.1) with angles $\alpha^j + \zeta$ instead of α^j , i.e.

$$a^j_{\zeta} = \mathcal{R}_{\zeta}(a^j),$$

being $\mathcal{R}_{\zeta} := R_{0,\zeta}$ with $R_{0,\zeta}$ as in (3.1), see Figure 6.



FIGURE 6. The rotated configuration $\{a_{\mathcal{L}}^j\}$.

In the next theorem we prove that the function $\zeta \mapsto \mathcal{E}^{(\zeta)} - L^{(\zeta)}(\Psi_0^{(\zeta)})$ is continuous. **Theorem 6.8.** The function $G : [-\pi, \pi) \to \mathbb{R}$, $G(\zeta) := \mathcal{E}^{(\zeta)} - L^{(\zeta)}(\Psi_0^{(\zeta)})$ is continuous. *Proof.* Through a rotation, the problem of continuity at any $\zeta \in [-\pi, \pi)$ can be reduced to the problem of continuity at $\zeta = 0$. Hence, it is enough to prove that $\lim_{\zeta \to 0} G(\zeta) = \mathcal{E} - L(\Psi_0)$.

We have

$$\Psi_0^{(\zeta)}(r\cos t, r\sin t) = f(t-\zeta)\phi_0(r\cos t, r\sin t),$$

where f is defined in (2.22) and

$$\phi_0(r\cos t, r\sin t) := \beta r^{\frac{m}{2}} \sin\left(\frac{m}{2}(t-\alpha_0)\right).$$

With a slight abuse of notation, henceforth we denote by f also the function $(r \cos t, r \sin t) \mapsto f(t)$ defined on $\mathbb{R}^2 \setminus \{0\}$.

A change of variables yields

$$\mathcal{E}^{(\zeta)} = \min\left\{I_{\zeta}(w) : w \in \widetilde{\mathcal{X}} \text{ and } w - \eta f(\phi_0 \circ \mathcal{R}_{\zeta}) \in \widetilde{\mathcal{H}}\right\},\$$

where $\eta \in C_c^{\infty}(\mathbb{R}^2)$ is a radial cut-off function as in (2.29) and

$$I_{\zeta}(w) = \frac{1}{2} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 \, dx + 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} f(\nabla \phi_0 \circ \mathcal{R}_{\zeta}) M_{\zeta} \cdot \nu^j \gamma_+^j(w) \, dS,$$

being M_{ζ} the matrix defined in (3.2). Moreover

$$L^{(\zeta)}(\Psi_0^{(\zeta)}) = 2\sum_{j=1}^{k_1+k_2} \int_{S_1^j} (\nabla \phi_0 \circ \mathcal{R}_\zeta) M_\zeta \cdot \nu^j(\phi_0 \circ \mathcal{R}_\zeta) \, dS$$

Since, in a neighbourhood of 0,

(6.33) $|\nabla \phi_0(\mathcal{R}_{\zeta}(x))| \le C|x|^{\frac{m}{2}-1} \text{ and } |\phi_0(\mathcal{R}_{\zeta}(x))| \le C|x|^{\frac{m}{2}}$

for some C > 0 independent of ζ , from the Dominated Convergence Theorem we deduce that

(6.34)
$$\lim_{\zeta \to 0} L^{(\zeta)}(\Psi_0^{(\zeta)}) = L(\Psi_0)$$

By Proposition 6.4, for every ζ there exists a unique $\widetilde{V}_{\zeta} \in \widetilde{\mathcal{X}}$ such that $\widetilde{V}_{\zeta} - \eta f(\phi_0 \circ \mathcal{R}_{\zeta}) \in \widetilde{\mathcal{H}}$ and $\mathcal{E}^{(\zeta)} = I_{\zeta}(\widetilde{V}_{\zeta})$; furthermore, \widetilde{V}_{ζ} satisfies

(6.35)
$$\int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V}_{\zeta} \cdot \nabla w \, dx = -2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} f(\nabla \phi_0 \circ \mathcal{R}_{\zeta}) M_{\zeta} \cdot \nu^j \gamma_+^j(w) \, dS \quad \text{for all } w \in \widetilde{\mathcal{H}}.$$

Choosing $w = \widetilde{V}_{\zeta} - \eta f(\phi_0 \circ \mathcal{R}_{\zeta})$ in (6.35) we obtain

$$(6.36) \qquad \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V}_{\zeta}|^2 \, dx = \int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V}_{\zeta} \cdot \nabla (\eta f(\phi_0 \circ \mathcal{R}_{\zeta})) \, dx - 2 \sum_{j=1}^{k_1 + k_2} \int_{S_1^j} f(\nabla \phi_0 \circ \mathcal{R}_{\zeta}) M_{\zeta} \cdot \nu^j \gamma^j_+ (\widetilde{V}_{\zeta}) \, dS + 2 \sum_{j=1}^{k_1 + k_2} \int_{S_1^j} f(\nabla \phi_0 \circ \mathcal{R}_{\zeta}) M_{\zeta} \cdot \nu^j \gamma^j_+ (f(\phi_0 \circ \mathcal{R}_{\zeta})) \, dS.$$

Using Young's inequality, estimate (6.33), and the continuity of the trace operators (6.8), from the above identity we deduce that

$$\|V_{\zeta}\|_{\widetilde{\mathcal{X}}} \leq C$$

for some C > 0 independent of ζ . It follows that every sequence $\zeta_n \to 0$ admits a subsequence $\{\zeta_{n_\ell}\}_\ell$ such that $\widetilde{V}_{\zeta_{n_\ell}} \to W$ weakly in $\widetilde{\mathcal{X}}$ as $\ell \to \infty$, for some $W \in \widetilde{\mathcal{X}}$. On account of (6.33) and (6.8), the Dominated Convergence Theorem yields

$$\int_{S_1^j} f(\nabla \phi_0 \circ \mathcal{R}_{\zeta_{n_\ell}}) M_{\zeta_{n_\ell}} \cdot \nu^j \gamma_+^j(w) \, dS \to \int_{S_1^j} f \nabla \phi_0 \cdot \nu^j \gamma_+^j(w) \, dS = \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(w) \, dS$$

30

as $\ell \to \infty$, for every $j = 1, \ldots k_1 + k_2$ and $w \in \widetilde{\mathcal{H}}$. By choosing $\zeta = \zeta_{n_\ell}$ in (6.35) and letting $\ell \to \infty$ we obtain that

(6.37)
$$\int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla W \cdot \nabla w \, dx = -2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j(w) \, dS \quad \text{for all } w \in \widetilde{\mathcal{H}}$$

Furthermore, since $\widetilde{V}_{\zeta} - \eta f(\phi_0 \circ \mathcal{R}_{\zeta}) \in \widetilde{\mathcal{H}}, \widetilde{\mathcal{H}}$ is a closed subspace of $\widetilde{\mathcal{X}}$, and $\eta f(\phi_0 \circ \mathcal{R}_{\zeta}) \to \eta \Psi_0$ as $\zeta \to 0$ in $\widetilde{\mathcal{X}}$ by the Dominated Convergence Theorem, we have

$$(6.38) W - \eta \Psi_0 \in \mathcal{H}.$$

From (6.37)-(6.38) and the uniqueness part of Proposition 6.4 we deduce that $W = \tilde{V}$. Having uniquely identified the weak limit independently of the subsequence, by the Urysohn subsequence principle we conclude that

(6.39)
$$\widetilde{V}_{\zeta} \rightharpoonup \widetilde{V} \quad \text{weakly in } \widetilde{\mathcal{X}} \quad \text{as } \zeta \to 0.$$

The weak convergence (6.39) allows us to pass to the limit as $\zeta \to 0$ on the right hand side of (6.36), thus proving that

(6.40)
$$\lim_{\zeta \to 0} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V}_{\zeta}|^2 \, dx = \int_{\mathbb{R}^2 \setminus \Gamma_1} \nabla \widetilde{V} \cdot \nabla(\eta \Psi_0) \, dx - 2 \sum_{j=1}^{k_1+k_2} \int_{S_1^j} \nabla \Psi_0 \cdot \nu^j \gamma_+^j (\widetilde{V} - \Psi_0) \, dS$$
$$= \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla \widetilde{V}|^2 \, dx,$$

the last equality being a consequence of (6.10) tested with $w = \tilde{V} - \eta \Psi_0$. From (6.40) it follows that $\lim_{\zeta \to 0} \mathcal{E}^{(\zeta)} = \lim_{\zeta \to 0} I_{\zeta}(\tilde{V}_{\zeta}) = J(\tilde{V}) = \mathcal{E}$, which, together with (6.34), yields the conclusion. \Box

When $k_2 = 0$ and the poles $\{a^j\}_{j=1,...,k_1}$ are on the tangents to nodal lines of v_0 (i.e. on the nodal set of Ψ_0), we have $\Psi_0 = 0$ on S_1^j for all $j = 1, ..., k_1$; on the other hand, if the poles are on the bisectors between nodal lines, then $\nabla \Psi_0 \cdot \nu^j = 0$ on S_1^j for all $j = 1, ..., k_1$. This leads to Proposition 2.3, which determines, in these particular cases, the sign of the dominant term in the asymptotic expansion obtained in Theorem 2.2, and, consequently, exploits the continuity result established in Theorem 6.8 to find configurations of poles for which the eigenvalue variation is an infinitesimal of higher order.

Proof of Proposition 2.3. (i) If $\alpha^j \in \{\alpha_0 + \ell \frac{2\pi}{m} : \ell = 0, 1, 2, \dots, m-1\}$ for all $j \in \{1, \dots, k_1\}$, then $\Psi_0 = 0$ on S_1^j for all $j \in \{1, \dots, k_1\}$, so that $L(\Psi_0) = 0$ and $\eta \Psi_0 + \widetilde{\mathcal{H}} = \widetilde{\mathcal{H}}$. It follows that

(6.41)
$$\mathcal{E} - L(\Psi_0) = \mathcal{E} = \min_{\widetilde{\mathcal{H}}} J.$$

Furthermore, $\nabla \Psi_0 \cdot \nu^j \neq 0$ on S_1^j for all $j \in \{1, \ldots, k_1\}$, see (6.22), hence $L \neq 0$ in $\widetilde{\mathcal{H}}$. Fixing some $w \in \widetilde{\mathcal{H}}$ such that $L(w) \neq 0$, we have then $J(tw) = \frac{t^2}{2} \int_{\mathbb{R}^2 \setminus \Gamma_1} |\nabla w|^2 dx + tL(w) < 0$ for some small t, thus implying that $\mathcal{E} = \min_{\widetilde{\mathcal{H}}} J < 0$. Once we have established that $\mathcal{E} - L(\Psi_0) = \mathcal{E} < 0$, from the asymptotic expansion of Theorem 2.2-(ii) we deduce that $\lambda_{\varepsilon,n_0} < \lambda_{0,n_0}$ for sufficiently small $\varepsilon > 0$.

(ii) If $\alpha^j \in \{\alpha_0 + (1+2\ell)\frac{\pi}{m} : \ell = 0, 1, 2, \dots, m-1\}$ for all $j \in \{1, \dots, k_1\}$, then $\nabla \Psi_0 \cdot \nu^j \equiv 0$ on S_1^j for all $j \in \{1, \dots, k_1\}$, see (6.22). It follows that $L \equiv 0$, and hence $J(w) = \frac{1}{2} ||w||_{\widetilde{\mathcal{X}}}^2$. Since, in this case, $\Psi_0 \neq 0$ on S_1^j for all $j \in \{1, \dots, k_1\}$, we have $w \neq 0$ for every $w \in \eta \Psi_0 + \widetilde{\mathcal{H}}$. Therefore

(6.42)
$$\mathcal{E} - L(\Psi_0) = \mathcal{E} = \min_{\eta \Psi_0 + \widetilde{\mathcal{H}}} J = \frac{1}{2} \min_{w \in \eta \Psi_0 + \widetilde{\mathcal{H}}} \|w\|_{\widetilde{\mathcal{X}}}^2 > 0.$$

From the asymptotic expansion of Theorem 2.2-(ii) we finally deduce that $\lambda_{\varepsilon,n_0} > \lambda_{0,n_0}$ for sufficiently small $\varepsilon > 0$.

(iii) Let us fix a configuration $\{a^j\}_{j=1}^k$ with $k = k_1 \leq m$ odd and $\alpha^j \in \{\alpha_0 + \ell \frac{2\pi}{m} : 0 \leq \ell \leq m-1\}$ for all $j \in \{1, \ldots, k_1\}$ as in (i). Then the rotated configuration $\{a_{\pi/m}^j\}$ is as in (ii). By (i)-(ii) we have G(0) < 0 and $G(\frac{\pi}{m}) > 0$. Since G is continuous by Theorem 6.8, Bolzano's Theorem ensures the existence of some $\zeta_0 \in (0, \frac{\pi}{m})$ such that $G(\zeta_0) = 0$, so that the angles $\{\alpha^j + \zeta_0 : j = 1, \ldots, k\}$ are as we are looking for. \Box



FIGURE 7. Nodal set of Ψ_0 and sign of $\mathcal{E} - L(\Psi_0)$ $(m = k = 3, \alpha_0 = 0)$.

Remark 6.9. Figure 7 and Figure 8 provide an example that helps to better visualize the result in Proposition 2.3. In Figure 7 we zoom in near a point (the origin) where the limit eigenfunction v_0 vanishes of order 3/2, namely (2.21) holds with m = 3. We consider the case $\alpha_0 = 0$. The function Ψ_0 as in (2.26) is the 3/2-homogeneous limit profile describing the local behavior of v_0 . In the image on the left, the black lines are the nodal lines of v_0 , which are tangent to the nodal lines of Ψ_0 (in green). The dotted lines denote the bisectors of the nodal lines of Ψ_0 . In the image on the right, we fix an admissible configuration of poles $\{a^j\}_{j=1,2,3}$ with k = 3 and $\alpha_j = 2\pi(j-1)/3$ for j = 1, 2, 3. From Proposition 2.3 we know that, if all the poles lie on the nodal set of Ψ_0 , then the coefficient $\mathcal{E} - L(\Psi_0)$ of the leading term in the asymptotic expansion stated in Theorem 2.2 is strictly negative. On the other hand, if all the poles lie on the bisectors of the nodal lines, then the coefficient $\mathcal{E} - L(\Psi_0)$ is strictly positive. In Figure 8 on the left, in the first picture (red



FIGURE 8. A visualization of Proposition 2.3.

32

arrows) we have our initial fixed configuration, which then provides a negative coefficient. In the second picture (blue arrows) we consider a rotation about the origin by an angle $\pi/m = \pi/3$: the rotated configuration ends up with all the poles lying on the bisectors, thus giving a positive coefficient $\mathcal{E} - L(\Psi_0)$. Furthermore, the continuity result in Theorem 6.8 ensures the existence of some $\zeta_0 \in (0, \pi/3)$ such that, if we rotate the initial configuration by an angle ζ_0 , we find a configuration of poles for which $\mathcal{E} - L(\Psi_0) = 0$: this is represented in the third picture on the left (yellow arrows). Finally, the right picture in Figure 8 presents the behavior of the perturbed eigenvalue in the three cases previously described. We point out that, when $\mathcal{E} - L(\Psi_0) = 0$ (yellow graph), it is currently not known what is the vanishing order of $\lambda_{\varepsilon,n_0} - \lambda_{0,n_0}$.

6.6. Blow-up and convergence rate for eigenfunctions. From the blow-up analysis for the potential V_{ε} performed in Subsection 6.4 and the energy estimate given in (5.7), we derive the following blow-up result for scaled eigenfunctions, together with a sharp estimate for their rate of convergence in the \mathcal{H}_1 -norm.

Proposition 6.10. Under assumption (5.3), let k be odd and v_0 be an eigenfunction of (2.14) associated to the eigenvalue $\lambda_0 = \lambda_{0,n_0}$ with $\|v_0\|_{L^2(\Omega)} = 1$. For $\varepsilon > 0$ small, let $\lambda_{\varepsilon} = \lambda_{\varepsilon,n_0}$ and v_{ε} be an eigenfunction of (2.10) associated to λ_{ε} and chosen as in (5.4). Let $m \in \mathbb{N}$ be given in Proposition 3.6 for $v = v_0$. Then

(6.43)
$$\varepsilon^{-\frac{m}{2}} v_{\varepsilon}(\varepsilon x) \to \Psi_0 - \widetilde{V} \quad as \ \varepsilon \to 0^+ \ in \ H^1(D_{\rho} \setminus \Gamma_1) \ for \ all \ \rho > 0,$$

where Ψ_0 is defined in (2.26) and \widetilde{V} is the unique solution to (6.10). Furthermore,

(6.44)
$$\lim_{\varepsilon \to 0^+} \varepsilon^{-\frac{m}{2}} \| v_{\varepsilon} - v_0 \|_{\mathcal{H}_1} = \| \widetilde{V} \|_{\widetilde{\mathcal{X}}}.$$

Proof. Using the same notation as in the proof of Theorem 5.2, let $\psi_{\varepsilon} = v_0 - V_{\varepsilon}$, where V_{ε} is defined as in Proposition 4.2. From (5.7) it follows that

$$\|\Pi_{\varepsilon}\psi_{\varepsilon} - \psi_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 = o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2\right) \quad \text{as } \varepsilon \to 0^+.$$

Therefore, defining

$$W_{\varepsilon}(x) := \varepsilon^{-\frac{m}{2}} (\Pi_{\varepsilon} \psi_{\varepsilon} - \psi_{\varepsilon})(\varepsilon x), \quad x \in \frac{1}{\varepsilon} \Omega,$$

and extending trivially W_{ε} in $\mathbb{R}^2 \setminus \frac{1}{\varepsilon}\Omega$, we have $W_{\varepsilon} \in \widetilde{\mathcal{H}}$ and, in view of Proposition 6.7,

$$\|W_{\varepsilon}\|_{\widetilde{\mathcal{X}}}^2 = \varepsilon^{-m} \|\Pi_{\varepsilon}\psi_{\varepsilon} - \psi_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 = \varepsilon^{-m} o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2\right) = \|\widetilde{V_{\varepsilon}}\|_{\widetilde{\mathcal{X}}}^2 o(1) = o(1)$$

as $\varepsilon \to 0^+$. By continuity of the restriction operator in (6.7) we deduce that

(6.45)
$$W_{\varepsilon} \to 0 \quad \text{as } \varepsilon \to 0^+ \text{ in } H^1(D_{\rho} \setminus \Gamma_1) \text{ for all } \rho > 0.$$

Let us define

(6.46)
$$U_{\varepsilon}(x) := \varepsilon^{-\frac{m}{2}} (\Pi_{\varepsilon} \psi_{\varepsilon})(\varepsilon x), \quad x \in \frac{1}{\varepsilon} \Omega,$$

and extend trivially U_{ε} in $\mathbb{R}^2 \setminus \frac{1}{\varepsilon} \Omega$. We have

$$U_{\varepsilon} = \widetilde{V}_{0,\varepsilon}(x) - \widetilde{V}_{\varepsilon} + W_{\varepsilon},$$

where $\widetilde{V}_{0,\varepsilon}$ and $\widetilde{V}_{\varepsilon}$ are defined in (6.18). Combining (6.25), (6.26), and (6.45), we conclude that

(6.47)
$$U_{\varepsilon} \to \Psi_0 - V$$
 as $\varepsilon \to 0^+$ in $H^1(D_{\rho} \setminus \Gamma_1)$ for all $\rho > 0$

From (5.8) it follows that

$$\int_{\Omega} v_0 \Pi_{\varepsilon} \psi_{\varepsilon} \, dx = 1 + o\left(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} \right) \quad \text{as } \varepsilon \to 0^+,$$

and hence, for $\varepsilon > 0$ small enough,

$$\int_{\Omega} \frac{\prod_{\varepsilon} \psi_{\varepsilon}}{\|\prod_{\varepsilon} \psi_{\varepsilon}\|_{L^{2}(\Omega)}} v_{0} \, dx > 0.$$

Since v_{ε} is the unique eigenfunction of (2.10) associated to λ_{ε} satisfying (5.4), we conclude that necessarily

(6.48)
$$v_{\varepsilon} = \frac{\prod_{\varepsilon} \psi_{\varepsilon}}{\|\prod_{\varepsilon} \psi_{\varepsilon}\|_{L^{2}(\Omega)}}$$

The convergence stated in (6.43) follows from (6.48), (6.46), (6.47), and (5.9). Moreover, (5.9) implies that

(6.49)
$$\|v_{\varepsilon} - \Pi_{\varepsilon}\psi_{\varepsilon}\|_{\mathcal{H}_{1}} = \frac{|1 - \|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^{2}(\Omega)}|}{\|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^{2}(\Omega)}} \|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{\mathcal{H}_{1}} = |1 - \|\Pi_{\varepsilon}\psi_{\varepsilon}\|_{L^{2}(\Omega)} |\|v_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} = o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}) \quad \text{as } \varepsilon \to 0^{+},$$

whereas (5.7) yields that

(6.50) $\|\Pi_{\varepsilon}\psi_{\varepsilon} - v_0\|_{\mathcal{H}_1}^2 = \|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 + \|\Pi_{\varepsilon}\psi_{\varepsilon} - \psi_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 - 2(V_{\varepsilon}, \Pi_{\varepsilon}\psi_{\varepsilon} - \psi_{\varepsilon})_{\mathcal{H}_{\varepsilon}} = \|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2)$ as $\varepsilon \to 0^+$. Combining (6.49) and (6.50) we deduce that

(6.51)
$$\|v_{\varepsilon} - v_0\|_{\mathcal{H}_1}^2 = \|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2 + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2) \quad \text{as } \varepsilon \to 0^+.$$

Letting $\widetilde{V}_{\varepsilon}$ be as in (6.18), from (6.51) and (6.26) we deduce that

$$\varepsilon^{-m} \|v_{\varepsilon} - v_0\|_{\mathcal{H}_1}^2 = \|\tilde{V}_{\varepsilon}\|_{\tilde{\mathcal{X}}}^2 (1 + o(1)) = \|\tilde{V}\|_{\tilde{\mathcal{X}}}^2 (1 + o(1)) \quad \text{as } \varepsilon \to 0^+,$$

thus proving (6.44).

Going back to the eigenfunctions of the original magnetic problem via the inverse of transformation (2.9), we deduce Theorem 2.4 from Proposition 6.10.

Proof of Theorem 2.4. If u_0 is an eigenfunction of (1.5) associated to the eigenvalue λ_{0,n_0} such that $\int_{\Omega} |u_0|^2 dx = 1$, and u_{ε} is the eigenfunction of (1.4) associated to $\lambda_{n_0,\varepsilon}$ satisfying (2.32), then $v_{\varepsilon} := e^{-i\Theta_{\varepsilon}}u_{\varepsilon}$ is an eigenfunction of (2.10) associated to $\lambda_{n_0,\varepsilon}$ and $v_0 := e^{-i\Theta_0}u_0$ is an eigenfunction of (2.14) associated to $\lambda_{n_0,0}$ such that condition (5.4) is satisfied. From Proposition 6.10 it follows that v_{ε} satisfies (6.43) and (6.44), in which we replace v_{ε} with $e^{-i\Theta_{\varepsilon}}u_{\varepsilon}$ and v_0 with $e^{-i\Theta_0}u_0$ to get exactly (2.33) and (2.34), taking into account that $\Theta_{\varepsilon}(\varepsilon x) = \Theta_1(x)$ for all $x \in \mathbb{R}^2 \setminus \{a^j : j = 1, \ldots, k\}$.

7. The case of two poles

The purpose of this section is to prove Theorems 2.6 and 2.7. We consider the case $k_1 = 0$ and $k_2 = 1$, with the configuration of poles as in assumption (2.36), being $r_1 \in (0, R)$ and $\varepsilon \in (0, 1]$. For the sake of simplicity, let us denote

$$T := T^1, \quad \gamma_+ := \gamma_+^1, \quad \gamma_- := \gamma_-^1, \text{ and } \nu := \nu^1 = (0, 1),$$

see (2.4). We first consider a linear functional $L_{\varepsilon,h,\Lambda}$ more general than the one introduced in (2.16), defined for a generic domain Λ and with the limit eigenfunction v_0 replaced by a generic function h; the corresponding minimal energy $\mathcal{E}_{\varepsilon,h,\Lambda}$ thus generalizes the energy $\mathcal{E}_{\varepsilon}$ defined in (2.19). For every simply connected open bounded domain $\Lambda \subset \mathbb{R}^2$ such that $D_R \subseteq \Lambda$ and every $h \in H_0^1(\Lambda) \cap C^{\infty}(\Lambda)$, let

$$L_{\varepsilon,h,\Lambda}: \mathcal{H}_{1,\Lambda} \to \mathbb{R}, \quad L_{\varepsilon,h,\Lambda}(w) := 2 \int_{S_{\varepsilon}} \frac{\partial h}{\partial x_2} \gamma_+(w) \, dS$$

and

$$J_{\varepsilon,h,\Lambda}: \mathcal{H}_{\varepsilon,\Lambda} \to \mathbb{R}, \quad J_{\varepsilon,h,\Lambda}(w) := \frac{1}{2} \int_{\Lambda \setminus S_{\varepsilon}} |\nabla w|^2 \, dx + L_{\varepsilon,h,\Lambda}(w),$$

where, for all $\varepsilon \in (0, 1]$, S_{ε} is defined in (2.37) and the functional space $\mathcal{H}_{\varepsilon,\Lambda}$ is the closure of

 $\left\{ w \in H^1(\Lambda \setminus S_{\varepsilon}) : w = 0 \text{ on a neighbourhood of } \partial \Lambda \right\}$

with respect to the norm $||w||_{H^1(\Omega \setminus S_{\varepsilon})}$. Then the minimization problem

(7.1)
$$\inf \left\{ J_{\varepsilon,h,\Lambda}(w) : w \in \mathcal{H}_{\varepsilon,\Lambda} \text{ and } w - h \in \widetilde{\mathcal{H}}_{\varepsilon,\Lambda} \right\}$$

with $\mathcal{H}_{\varepsilon,\Lambda} := \{ w \in \mathcal{H}_{\varepsilon,\Lambda} : T(w) = 0 \text{ on } S_{\varepsilon} \}$, is uniquely achieved, as stated in the following proposition. We omit the proof, being similar to the one of Proposition 4.2.

Proposition 7.1. The infimum in (7.1) is achieved by a unique $V_{\varepsilon,h,\Lambda} \in \mathcal{H}_{\varepsilon,\Lambda}$. Furthermore, $V_{\varepsilon,h,\Lambda}$ weakly solves the problem

(7.2)
$$\begin{cases} -\Delta V_{\varepsilon,h,\Lambda} = 0, & \text{in } \Lambda \setminus S_{\varepsilon} \\ V_{\varepsilon,h,\Lambda} = 0, & \text{on } \partial\Lambda, \\ T(V_{\varepsilon,h,\Lambda} - h) = 0, & \text{on } S_{\varepsilon}, \\ T\left(\frac{\partial V_{\varepsilon,h,\Lambda}}{\partial x_2} - \frac{\partial h}{\partial x_2}\right) = 0, & \text{on } S_{\varepsilon}, \end{cases}$$

in the sense that $V_{\varepsilon,h,\Lambda} \in \mathcal{H}_{\varepsilon,\Lambda}$, $V_{\varepsilon,h,\Lambda} - h \in \widetilde{\mathcal{H}}_{\varepsilon,\Lambda}$, and

(7.3)
$$\int_{\Lambda \setminus S_{\varepsilon}} \nabla V_{\varepsilon,h,\Lambda} \cdot \nabla w \, dx = -L_{\varepsilon,h,\Lambda}(w) \quad \text{for all } w \in \widetilde{\mathcal{H}}_{\varepsilon,\Lambda}$$

For every Λ , h as above and $\varepsilon \in (0, 1]$, let

(7.4)
$$\mathcal{E}_{\varepsilon,h,\Lambda} := J_{\varepsilon,h,\Lambda}(V_{\varepsilon,h,\Lambda}).$$

For every L > 0 and $\varepsilon > 0$, let $E_{\varepsilon}(L)$ be the ellipse defined as

$$E_{\varepsilon}(L) := \left\{ (x_1, x_2) \in \mathbb{R}^2 : \frac{x_1^2}{L^2 + r_1^2 \varepsilon^2} + \frac{x_2^2}{L^2} < 1 \right\}$$

We are going to compute $\mathcal{E}_{\varepsilon,P_m,E_{\varepsilon}(L)}$, where P_m is a homogeneous polynomial of degree $m \geq 1$. We shall later apply such estimate with P_m being the Taylor polynomial of u_0 centered at 0 of order m, with u_0 and m as in Section 2.2.

Proposition 7.2. Let $m \in \mathbb{N}$, $m \ge 1$, and let P_m be a homogeneous polynomial of degree m, i.e.

(7.5)
$$P_m(x_1, x_2) := \sum_{j=0}^m \ell_j x_1^{m-j} x_2^j$$

for some $\ell_0, \ell_1, \ldots, \ell_m \in \mathbb{R}$. Then, for every L > 0, we have

(7.6)
$$\int_{E_{\varepsilon}(L)\backslash S_{\varepsilon}} |\nabla V_{\varepsilon,P_m,E_{\varepsilon}(L)}|^2 dx = \pi (\varepsilon r_1)^{2m} \left(\ell_0^2 \sum_{j=1}^m j |c_j|^2 + \ell_1^2 \sum_{j=1}^m \frac{|d_j|^2}{j} \right) + o(\varepsilon^{2m})$$

as $\varepsilon \to 0^+$, where

(7.7)
$$c_j = \frac{1}{\pi} \int_0^{2\pi} (\cos \eta)^m \cos(j\eta) \, d\eta \quad \text{for every } j \in \mathbb{N},$$

(7.8)
$$d_j = \frac{1}{\pi} \int_0^{2\pi} (\cos \eta)^{m-1} \sin \eta \sin(j\eta) \, d\eta \quad \text{for every } j \in \mathbb{N} \setminus \{0\}.$$

Proof. We consider elliptic coordinates (ξ, η) defined as

(7.9)
$$\begin{cases} x_1 = \varepsilon r_1 \cosh(\xi) \cos(\eta), \\ x_2 = \varepsilon r_1 \sinh(\xi) \sin(\eta), \end{cases} \quad \xi \ge 0, \ \eta \in [0, 2\pi). \end{cases}$$

see e.g. [3, Section 2.2]. In this coordinates S_{ε} is described by the conditions

$$\xi = 0, \quad \eta \in [0, 2\pi),$$

whereas $E_{\varepsilon}(L)$ is described by

 $\xi \in [0, \xi_{\varepsilon}), \quad \eta \in [0, 2\pi),$

where ξ_{ε} is such that $r_1 \varepsilon \sinh(\xi_{\varepsilon}) = L$, that is

(7.10)
$$\xi_{\varepsilon} = \operatorname{arcsinh}\left(\frac{L}{r_{1}\varepsilon}\right) = \log\left(\frac{L}{r_{1}\varepsilon} + \sqrt{1 + \frac{L^{2}}{r_{1}^{2}\varepsilon^{2}}}\right).$$

In particular $\partial E_{\varepsilon}(L)$ is described by the conditions

$$\xi = \xi_{\varepsilon}, \quad \eta \in [0, 2\pi).$$

The map

$$F_{\varepsilon}: [0,\xi_{\varepsilon}) \times [0,2\pi) \to E_{\varepsilon}(L), \quad F_{\varepsilon}(\xi,\eta) = (x_1,x_2),$$

defined by (7.9), has a Jacobian matrix of the form

$$J_{F_{\varepsilon}}(\xi,\eta) = \varepsilon r_1 \sqrt{\cosh^2 \xi - \cos^2 \eta} \ O(\xi,\eta)$$

for some orthogonal matrix $O(\xi, \eta)$, and det $J_{F_{\varepsilon}}(\xi, \eta) = \varepsilon^2 r_1^2(\cosh^2 \xi - \cos^2 \eta)$. In particular F_{ε} is a conform mapping and $J_{F_{\varepsilon}}(\xi, \eta)$ is an invertible matrix if $(\xi, \eta) \neq (0, 0)$ and $(\xi, \eta) \neq (0, \pi)$.

Let $\widehat{V}_{\varepsilon,P_m,L} := V_{\varepsilon,P_m,E_{\varepsilon}(L)} \circ F_{\varepsilon}$, where $V_{\varepsilon,P_m,E_{\varepsilon}(L)}$ is the solution of (7.2) in the case $\Lambda = E_{\varepsilon}(L)$ and $h = P_m$. We observe that, since $F_{\varepsilon}(\xi,\eta) \in \mathbb{R}^2_+$ if $\eta \in (0,\pi)$ and $F_{\varepsilon}(\xi,\eta) \in \mathbb{R}^2_-$ if $\eta \in (\pi,2\pi)$,

$$\widehat{V}_{\varepsilon,P_m,L}(0,\eta) = \begin{cases} \gamma_+(V_{\varepsilon,P_m,E_\varepsilon(L)})(\varepsilon r_1 \cos \eta, 0), & \text{if } \eta \in (0,\pi), \\ \gamma_-(V_{\varepsilon,P_m,E_\varepsilon(L)})(\varepsilon r_1 \cos \eta, 0), & \text{if } \eta \in (\pi, 2\pi). \end{cases}$$

Furthermore,

$$\frac{\partial \widehat{V}_{\varepsilon,P_m,L}}{\partial \xi}(0,\eta) = \begin{cases} \varepsilon r_1(\sin\eta) \,\gamma_+ \left(\frac{\partial V_{\varepsilon,P_m,E_\varepsilon(L)}}{\partial x_2}\right) (\varepsilon r_1\cos\eta,0), & \text{if } \eta \in (0,\pi), \\ \varepsilon r_1(\sin\eta) \,\gamma_- \left(\frac{\partial V_{\varepsilon,P_m,E_\varepsilon(L)}}{\partial x_2}\right) (\varepsilon r_1\cos\eta,0), & \text{if } \eta \in (\pi,2\pi). \end{cases}$$

We also note that, for every $\eta \in [0, 2\pi)$,

$$P_m(F_{\varepsilon}(0,\eta)) = (\varepsilon r_1)^m \ell_0(\cos\eta)^m \quad \text{and} \quad \frac{\partial P_m}{\partial x_2}(F_{\varepsilon}(0,\eta)) = \ell_1(\varepsilon r_1)^{m-1}(\cos\eta)^{m-1}.$$

Therefore, $\hat{V}_{\varepsilon,P_m,L}$ solves the problem

$$(7.11) \begin{cases} -\Delta \widehat{V}_{\varepsilon,P_m,L} = 0, & \text{in } (0,\xi_{\varepsilon}) \times (0,2\pi), \\ \widehat{V}_{\varepsilon,P_m,L}(\xi_{\varepsilon},\eta) = 0, & \text{for all } \eta \in [0,2\pi), \\ \widehat{V}_{\varepsilon,P_m,L}(\xi,0) = \widehat{V}_{\varepsilon,P_m,L}(\xi,2\pi), & \text{for all } \xi \in (0,\xi_{\varepsilon}), \\ \widehat{V}_{\varepsilon,P_m,L}(0,\eta) + \widehat{V}_{\varepsilon,P_m,L}(0,2\pi-\eta) = 2\ell_0(\varepsilon r_1)^m (\cos \eta)^m, & \text{for all } \eta \in (0,\pi), \\ \frac{\partial \widehat{V}_{\varepsilon,P_m,L}}{\partial \xi}(0,\eta) - \frac{\partial \widehat{V}_{\varepsilon,P_m,L}}{\partial \xi}(0,2\pi-\eta) = 2\ell_1(\varepsilon r_1)^m (\cos \eta)^{m-1} \sin \eta, & \text{for all } \eta \in (0,\pi). \end{cases}$$

Let us consider the Fourier expansion of $(\varepsilon r_1)^{-m} \widehat{V}_{\varepsilon, P_m, L}$ with respect to the variable η

$$\frac{1}{(\varepsilon r_1)^m}\widehat{V}_{\varepsilon,P_m,L}(\xi,\eta) = \frac{a_{0,\varepsilon}(\xi)}{2} + \sum_{j=1}^{\infty} \Big(a_{j,\varepsilon}(\xi)\cos(j\eta) + b_{j,\varepsilon}(\xi)\sin(j\eta)\Big),$$

where

$$a_{j,\varepsilon}(\xi) := \frac{(\varepsilon r_1)^{-m}}{\pi} \int_0^{2\pi} \widehat{V}_{\varepsilon,P_m,L}(\xi,\eta) \cos(j\eta) \, d\eta \quad \text{for all } j \in \mathbb{N},$$
$$b_{j,\varepsilon}(\xi) := \frac{(\varepsilon r_1)^{-m}}{\pi} \int_0^{2\pi} \widehat{V}_{\varepsilon,P_m,L}(\xi,\eta) \sin(j\eta) \, d\eta \quad \text{for all } j \in \mathbb{N} \setminus \{0\}.$$

Since $\cos(2\pi - \eta) = \cos \eta$ for any $\eta \in (0, \pi)$, from (7.11) it follows that

$$a_{0,\varepsilon}(0) + 2\sum_{j=1}^{\infty} a_{j,\varepsilon}(0)\cos(j\eta) = 2\ell_0(\cos\eta)^m \quad \text{for all } \eta \in (0,2\pi),$$

hence $\{a_{j,\varepsilon}(0)\}_{j\in\mathbb{N}}$ are the Fourier coefficients of $\ell_0(\cos\eta)^m$ with respect to the orthonormal basis $\{\frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{\pi}}\cos(j\eta), \frac{1}{\sqrt{\pi}}\sin(j\eta)\}_{j\in\mathbb{N}\setminus\{0\}}$ of $L^2(0, 2\pi)$, i.e.

$$a_{j,\varepsilon}(0) = \ell_0 c_j$$
 for all $j \in \mathbb{N}$,

with c_j as in (7.7). In particular

(7.12)
$$a_{j,\varepsilon}(0) = \ell_0 c_j = 0 \quad \text{if } j > m.$$

On the other hand, the last condition in (7.11) reads as

$$\sum_{j=1}^{\infty} b'_{j,\varepsilon}(0) \sin(j\eta) = \ell_1(\cos\eta)^{m-1} \sin\eta \quad \text{for all } \eta \in (0, 2\pi).$$

It follows that $b'_{j,\varepsilon}(0)$ are independent of ε and

$$b'_{j,\varepsilon}(0) = \ell_1 d_j \quad \text{for all } j \in \mathbb{N} \setminus \{0\},$$

with d_j as in (7.8); hence

(7.13)
$$b'_{j,\varepsilon}(0) = \ell_1 d_j = 0 \text{ if } j > m.$$

From the equation in (7.11) it follows that

$$0 = \frac{1}{(\varepsilon r_1)^m} \Delta \widehat{V}_{\varepsilon, P_m, L}$$

= $\frac{a_{0,\varepsilon}''(\xi)}{2} + \sum_{j=1}^{\infty} \left((a_{j,\varepsilon}''(\xi) - j^2 a_{j,\varepsilon}(\xi)) \cos(j\eta) + (b_{j,\varepsilon}''(\xi) - j^2 b_{j,\varepsilon}(\xi)) \sin(j\eta) \right),$

hence

(7.14)
$$a_{0,\varepsilon}(\xi) = -\frac{a_{0,\varepsilon}(0)}{\xi_{\varepsilon}}\xi + a_{0,\varepsilon}(0) = -\frac{\ell_0 c_0}{\xi_{\varepsilon}}\xi + \ell_0 c_0 \quad \text{for all } \xi \in (0,\xi_{\varepsilon}),$$
$$a_{j,\varepsilon}(\xi) = \ell_0 c_j \left(\frac{e^{j\xi}}{1 - e^{2j\xi_{\varepsilon}}} + \frac{e^{-j\xi}}{1 - e^{-2j\xi_{\varepsilon}}}\right) \quad \text{for all } \xi \in (0,\xi_{\varepsilon}) \text{ and } j \in \mathbb{N} \setminus \{0\},$$
$$b_{j,\varepsilon}(\xi) = \frac{\ell_1 d_j}{j} \left(\frac{e^{j\xi}}{1 + e^{2j\xi_{\varepsilon}}} - \frac{e^{-j\xi}}{1 + e^{-2j\xi_{\varepsilon}}}\right) \quad \text{for all } \xi \in (0,\xi_{\varepsilon}) \text{ and } j \in \mathbb{N} \setminus \{0\},$$

with ξ_{ε} as in (7.10). Then, by (7.12) and (7.13), $a_{j,\varepsilon} \equiv b_{j,\varepsilon} \equiv 0$ for all j > m, so that

$$\frac{1}{(\varepsilon r_1)^m}\widehat{V}_{\varepsilon,P_m,L}(\xi,\eta) = \frac{a_{0,\varepsilon}(\xi)}{2} + \sum_{j=1}^m \Big(a_{j,\varepsilon}(\xi)\cos(j\eta) + b_{j,\varepsilon}(\xi)\sin(j\eta)\Big).$$

By a change of variables and the Parseval identity,

$$(7.15) \qquad \int_{E_{\varepsilon}(L)\setminus S_{\varepsilon}} |\nabla V_{\varepsilon,P_m,E_{\varepsilon}(L)}|^2 dx = \int_0^{\xi_{\varepsilon}} \int_0^{2\pi} |\nabla \widehat{V}_{\varepsilon,P_m,L}|^2 d\eta d\xi$$
$$= (\varepsilon r_1)^{2m} \frac{\pi}{2} \int_0^{\xi_{\varepsilon}} |a'_{0,\varepsilon}(\xi)|^2 d\xi$$
$$+ (\varepsilon r_1)^{2m} \pi \sum_{j=1}^m \int_0^{\xi_{\varepsilon}} \left(|a'_{j,\varepsilon}(\xi)|^2 + j^2 |b_{j,\varepsilon}(\xi)|^2 + |b'_{j,\varepsilon}(\xi)|^2 + j^2 |a_{j,\varepsilon}(\xi)|^2 \right) d\xi.$$

Let us compute each integral in the above formula. In view of (7.14) and (7.10), it is clear that

$$\int_0^{\xi_{\varepsilon}} |a_{0,\varepsilon}'(\xi)|^2 d\eta = \frac{\ell_0^2 c_0^2}{\xi_{\varepsilon}} = \frac{\ell_0^2 c_0^2}{|\log \varepsilon|} + O\left(\frac{1}{|\log \varepsilon|^2}\right) \quad \text{as } \varepsilon \to 0^+.$$

Furthermore, for every $j \in \mathbb{N} \setminus \{0\}$,

$$\begin{split} j^2 \int_0^{\xi_{\varepsilon}} |b_{j,\varepsilon}(\xi)|^2 d\xi &= \ell_1^2 d_j^2 \int_0^{\xi_{\varepsilon}} \left(\frac{e^{j\xi}}{1 + e^{2j\xi_{\varepsilon}}} - \frac{e^{-j\xi}}{1 + e^{-2j\xi_{\varepsilon}}} \right)^2 d\xi \\ &= \frac{\ell_1^2 d_j^2}{(1 + e^{2j\xi_{\varepsilon}})^2} \int_0^{\xi_{\varepsilon}} e^{2j\xi} d\xi + \frac{\ell_1^2 d_j^2}{(1 + e^{-2j\xi_{\varepsilon}})^2} \int_0^{\xi_{\varepsilon}} e^{-2j\xi} d\xi - \frac{2\ell_1^2 d_j^2 \xi_{\varepsilon}}{2 + e^{-2j\xi_{\varepsilon}} + e^{2j\xi_{\varepsilon}}} \\ &= \frac{\ell_1^2 d_j^2}{2j} \left(\frac{1}{(1 + e^{2j\xi_{\varepsilon}})^2} (e^{2j\xi_{\varepsilon}} - 1) + \frac{1}{(1 + e^{-2j\xi_{\varepsilon}})^2} (1 - e^{-2j\xi_{\varepsilon}}) \right) - \frac{2\ell_1^2 d_j^2 \xi_{\varepsilon}}{2 + e^{-2j\xi_{\varepsilon}} + e^{2j\xi_{\varepsilon}}} \\ &= \frac{\ell_1^2 d_j^2}{2j} (1 + o(1)) \quad \text{as } \varepsilon \to 0^+ \end{split}$$

and similarly

$$\int_0^{\xi_{\varepsilon}} |b'_{j,\varepsilon}(\xi)|^2 d\xi = \ell_1^2 d_j^2 \int_0^{\xi_{\varepsilon}} \left(\frac{e^{j\xi}}{1 + e^{2j\xi_{\varepsilon}}} + \frac{e^{-j\xi}}{1 + e^{-2j\xi_{\varepsilon}}} \right)^2 d\xi = \frac{\ell_1^2 d_j^2}{2j} (1 + o(1)) \quad \text{as } \varepsilon \to 0^+.$$

Finally, for every $j \in \mathbb{N} \setminus \{0\}$,

$$\int_{0}^{\xi_{\varepsilon}} |a_{j,\varepsilon}'(\xi)|^2 d\xi = j^2 \ell_0^2 c_j^2 \int_{0}^{\xi_{\varepsilon}} \left(\frac{e^{j\xi}}{1 - e^{2j\xi_{\varepsilon}}} - \frac{e^{-j\xi}}{1 - e^{-2j\xi_{\varepsilon}}}\right)^2 d\xi = \ell_0^2 c_j^2 \frac{j}{2} (1 + o(1)),$$

$$j^2 \int_{0}^{\xi_{\varepsilon}} |a_{j,\varepsilon}(\xi)|^2 d\xi = j^2 \ell_0^2 c_j^2 \int_{0}^{\xi_{\varepsilon}} \left(\frac{e^{j\xi}}{1 - e^{2j\xi_{\varepsilon}}} + \frac{e^{-j\xi}}{1 - e^{-2j\xi_{\varepsilon}}}\right)^2 d\xi = \ell_0^2 c_j^2 \frac{j}{2} (1 + o(1)),$$

as $\varepsilon \to 0^+$, as shown in the proof of [3, Lemma 2.3]. Replacing the above estimates in (7.15) we obtain (7.6).

Proposition 7.3. Let $m \in \mathbb{N} \setminus \{0\}$. For every $j \in \mathbb{N} \setminus \{0\}$, let c_j and d_j be as in (7.7) and (7.8), respectively. Then

(7.16)
$$\sum_{j=1}^{m} j |c_j|^2 = \frac{m}{4^{m-1}} {\binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}}^2,$$

(7.17)
$$\sum_{j=1}^{m} \frac{1}{j} |d_j|^2 = \frac{1}{m4^{m-1}} {\binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}}^2$$

Proof. For the proof of (7.16) we refer to [4, Proposition A.3]. To prove (7.17), we observe that, in view of (7.7),

$$(\cos \eta)^m = \frac{c_0}{2} + \sum_{j=1}^m c_j \cos(j\eta) \text{ for all } \eta \in [0, 2\pi].$$

Deriving the previous identity with respect to η , we obtain

$$(\cos \eta)^{m-1} \sin \eta = \frac{1}{m} \sum_{j=1}^{m} jc_j \sin(j\eta) = \sum_{j=1}^{m} d_j \sin(j\eta) \text{ for all } \eta \in [0, 2\pi],$$

in view of (7.8). It follows that $d_j = \frac{j}{m}c_j$ for all j = 1, ..., m, hence (7.17) follows from (7.16).

Remark 7.4. Let $m \in \mathbb{N}$, $m \geq 1$, and let P_m be a homogeneous polynomial of degree m as in (7.5). Let $\Lambda \subset \mathbb{R}^2$ be a simply connected open bounded domain such that $D_R \subseteq \Lambda$.

(i) If the coefficient ℓ_0 in (7.5) is zero, then $P_m \equiv 0$ on S_{ε} for all $\varepsilon \in (0, 1]$. Hence $V_{\varepsilon, P_m, \Lambda} \in \widetilde{\mathcal{H}}_{\varepsilon, \Lambda}$ and, in view of (7.3), $\int_{\Lambda \setminus S_{\varepsilon}} |\nabla V_{\varepsilon, P_m, \Lambda}|^2 dx = -L_{\varepsilon, P_m, \Lambda}(V_{\varepsilon, P_m, \Lambda})$, so that

$$\mathcal{E}_{\varepsilon,P_m,\Lambda} = -\frac{1}{2} \int_{\Lambda \setminus S_{\varepsilon}} |\nabla V_{\varepsilon,P_m,\Lambda}|^2 \, dx.$$

(ii) If the coefficient ℓ_1 in (7.5) is zero, then $\frac{\partial P_m}{\partial x_2} \equiv 0$ on S_{ε} for all $\varepsilon \in (0, 1]$. Hence $L_{\varepsilon, P_m, \Lambda} \equiv 0$ and

$$\mathcal{E}_{\varepsilon,P_m,\Lambda} = \frac{1}{2} \int_{\Lambda \setminus S_{\varepsilon}} |\nabla V_{\varepsilon,P_m,\Lambda}|^2 \, dx.$$

Proposition 7.5. Let $\Omega \subset \mathbb{R}^2$ be a simply connected open bounded domain with $0 \in D_R \subseteq \Omega$. For every $\varepsilon \in (0,1)$, let S_{ε} be defined in (2.37). Let P_m be a homogeneous polynomial of degree m as in (7.5) and $\mathcal{E}_{\varepsilon,P_m,\Omega}$ be defined in (7.4) with $\Lambda = \Omega$ and $h = P_m$. Then, letting ℓ_0 and ℓ_1 be as in (7.5), we have

(i) if $\ell_0 = 0$, then

$$\mathcal{E}_{\varepsilon,P_m,\Omega} = -\frac{\pi}{2} r_1^{2m} \ell_1^2 \varepsilon^{2m} \frac{1}{m4^{m-1}} \binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}^2 + o(\varepsilon^{2m}) \quad as \ \varepsilon \to 0^+;$$

(ii) if $\ell_1 = 0$, then

$$\mathcal{E}_{\varepsilon,P_m,\Omega} = \frac{\pi}{2} r_1^{2m} \ell_0^2 \varepsilon^{2m} \frac{m}{4^{m-1}} {\binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}}^2 + o(\varepsilon^{2m}) \quad as \ \varepsilon \to 0^+.$$

Proof. The set Ω is open and $0 \in \Omega$, hence there exist $L_1, L_2 > 0$ such that, for every $\varepsilon \in (0, 1]$, $S_{\varepsilon} \subset E_{\varepsilon}(L_1) \subset \Omega \subset E_{\varepsilon}(L_2)$ (e.g. we can choose any $0 < L_1 < \sqrt{R^2 - r_1^2}$ and $L_2 = \operatorname{diam} \Omega$). From (7.1), (7.4), and the space inclusions $\mathcal{H}_{\varepsilon,E_{\varepsilon}(L_1)} \subset \mathcal{H}_{\varepsilon,\Omega} \subset \mathcal{H}_{\varepsilon,E_{\varepsilon}(L_2)}, \widetilde{\mathcal{H}}_{\varepsilon,E_{\varepsilon}(L_1)} \subset \widetilde{\mathcal{H}}_{\varepsilon,\Omega} \subset \widetilde{\mathcal{H}}_{\varepsilon,E_{\varepsilon}(L_2)}$ obtained by trivial extension, we deduce that, for every $\varepsilon \in (0, 1]$,

(7.18)
$$\mathcal{E}_{\varepsilon,P_m,E_{\varepsilon}(L_2)} \leq \mathcal{E}_{\varepsilon,P_m,\Omega} \leq \mathcal{E}_{\varepsilon,P_m,E_{\varepsilon}(L_1)}$$

If $\ell_0 = 0$, from Remark 7.4, (7.6), and (7.17) it follows that, for i = 1, 2,

$$\begin{aligned} \mathcal{E}_{\varepsilon,P_m,E_{\varepsilon}(L_i)} &= -\frac{1}{2} \int_{E_{\varepsilon}(L_i) \setminus S_{\varepsilon}} |\nabla V_{\varepsilon,P_m,E_{\varepsilon}(L_i)}|^2 \, dx \\ &= -\frac{\pi}{2} (\varepsilon r_1)^{2m} \ell_1^2 \bigg(\sum_{j=1}^m \frac{|d_j|^2}{j} \bigg) + o(\varepsilon^{2m}) = -\frac{\pi}{2} r_1^{2m} \ell_1^2 \varepsilon^{2m} \frac{1}{m4^{m-1}} \binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}^2 + o(\varepsilon^{2m}) \end{aligned}$$

as $\varepsilon \to 0^+$, thus proving (i) in view of (7.18).

On the other hand, if $\ell_1 = 0$, then Remark 7.4, (7.6), and (7.16) imply that, for i = 1, 2,

$$\begin{aligned} \mathcal{E}_{\varepsilon,P_m,E_{\varepsilon}(L_i)} &= \frac{1}{2} \int_{E_{\varepsilon}(L_i) \setminus S_{\varepsilon}} |\nabla V_{\varepsilon,P_m,E_{\varepsilon}(L_i)}|^2 \, dx \\ &= \frac{\pi}{2} (\varepsilon r_1)^{2m} \ell_0^2 \bigg(\sum_{j=1}^m j |c_j|^2 \bigg) + o(\varepsilon^{2m}) = \frac{\pi}{2} r_1^{2m} \ell_0^2 \varepsilon^{2m} \frac{m}{4^{m-1}} {\binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}}^2 + o(\varepsilon^{2m}) \\ &\to 0^+, \text{ thus proving (ii) in view of (7.18).} \end{aligned}$$

as $\varepsilon \to 0^+$, thus proving (ii) in view of (7.18).

Let u_0 be as in (2.35) with $u_0(0) = 0$ and m, β, α_0 be as in (2.38). Let T_m be the Taylor polynomial of u_0 centered at 0 of order m written in (2.39). In particular T_m is of the form (7.5) with

$$\ell_j = \frac{1}{(m-j)!j!} \frac{\partial^m u_0}{\partial x_1^{m-j} \partial x_2^j}(0).$$

If $\alpha_0 = \frac{j\pi}{m}$ for some $j \in \{0, 1, \dots, 2m-1\}$, then, by Remark 2.5, $\ell_0 = T_m(1,0) = 0$

and

$$\ell_1 = \frac{\partial T_m}{\partial x_2}(1,0) = \nabla T_m(1,0) \cdot (0,1) = m\beta \cos(j\pi) = (-1)^j m\beta.$$

Hence, by Proposition 7.5, in this case we have

(7.19)
$$\mathcal{E}_{\varepsilon,T_m,\Omega} = -\frac{\pi}{2} r_1^{2m} \varepsilon^{2m} \frac{m\beta^2}{4^{m-1}} {\binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}}^2 + o(\varepsilon^{2m}) \quad \text{as } \varepsilon \to 0^+.$$

On the other hand, if $\alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m}$ for some $j \in \{0, 1, \dots, 2m-1\}$, then, by Remark 2.5,

$$\ell_0 = T_m(1,0) = -\beta \sin\left(\frac{\pi}{2} + j\pi\right) = (-1)^{j+1}\beta$$

and

$$\ell_1 = \frac{\partial T_m}{\partial x_2}(1,0) = m\beta \cos\left(\frac{\pi}{2} + j\pi\right) = 0$$

In this case, Proposition 7.5 then provides the expansion

(7.20)
$$\mathcal{E}_{\varepsilon,T_m,\Omega} = \frac{\pi}{2} r_1^{2m} \varepsilon^{2m} \frac{m\beta^2}{4^{m-1}} {\binom{m-1}{\lfloor \frac{m-1}{2} \rfloor}}^2 + o(\varepsilon^{2m}) \quad \text{as } \varepsilon \to 0^+.$$

Let $g := u_0 - T_m$. Since u_0 is smooth and T_m is its Taylor polynomial at 0 of order m, then

(7.21)
$$g(x) = O(|x|^{m+1})$$
 and $|\nabla g(x)| = O(|x|^m)$ as $x \to 0$.

Proposition 7.6. Let m and α_0 be as in (2.38). For every $\varepsilon \in (0, 1]$, let $V_{\varepsilon, T_m, \Omega}$ and $\mathcal{E}_{\varepsilon, T_m, \Omega}$ be as in (7.1) and (7.4), with $\Lambda = \Omega$ and $h = T_m$, and let $V_{\varepsilon} = V_{\varepsilon, u_0, \Omega}$ and $\mathcal{E}_{\varepsilon} = \mathcal{E}_{\varepsilon, u_0, \Omega}$ be as in (2.18) and (2.19), respectively. Then

(7.22)
$$\|V_{\varepsilon} - V_{\varepsilon, T_m, \Omega}\|_{\mathcal{H}_{\varepsilon}}^2 = O(\varepsilon^{2m+1}) = o(\varepsilon^{2m}) \quad as \ \varepsilon \to 0^+$$

and, if either $\alpha_0 = \frac{j\pi}{m}$ or $\alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m}$ for some $j \in \{0, 1, \dots, 2m-1\}$,

(7.23)
$$\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} = O(\varepsilon^m) \quad as \ \varepsilon \to 0^+,$$

(7.24)
$$\mathcal{E}_{\varepsilon} - \mathcal{E}_{\varepsilon, T_m, \Omega} = o(\varepsilon^{2m}) \quad as \ \varepsilon \to 0^+.$$

Proof. Let $W_{\varepsilon} := V_{\varepsilon} - V_{\varepsilon,T_m,\Omega}$. Then W_{ε} satisfies (7.3) with h := g. Let η_{ε} be as in (4.7). Testing (7.3) with $w = W_{\varepsilon} - \eta_{\varepsilon}g$, by Young's Inequality and (2.3) we obtain

$$\begin{split} \|W_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2} &= \int_{\Omega \setminus S_{\varepsilon}} \eta_{\varepsilon} \nabla W_{\varepsilon} \cdot \nabla g \, dx + \int_{\Omega \setminus S_{\varepsilon}} g \nabla W_{\varepsilon} \cdot \nabla \eta_{\varepsilon} \, dx - 2 \int_{S_{\varepsilon}} \frac{\partial g}{\partial x_{2}} \gamma^{+}(W_{\varepsilon}) \, dS + 2 \int_{S_{\varepsilon}} \frac{\partial g}{\partial x_{2}} g \, dS \\ &\leq \frac{1}{2} \|W_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2} + C \left(\int_{\Omega} \eta_{\varepsilon}^{2} |\nabla g|^{2} \, dx + \int_{\Omega} g^{2} |\nabla \eta_{\varepsilon}|^{2} \, dx + \int_{S_{\varepsilon}} \left| \frac{\partial g}{\partial x_{2}} \right|^{2} \, dS \right) + 2 \int_{S_{\varepsilon}} \left| \frac{\partial g}{\partial x_{2}} \right| |g| \, dS, \end{split}$$

for some positive constant C > 0. Hence (7.22) follows from (4.7) and (7.21). We have

(7.25)
$$\mathcal{E}_{\varepsilon} - \mathcal{E}_{\varepsilon,T_m,\Omega} = \frac{1}{2} \left(\left\| V_{\varepsilon} \right\|_{\mathcal{H}_{\varepsilon}}^2 - \left\| V_{\varepsilon,T_m,\Omega} \right\|_{\mathcal{H}_{\varepsilon}}^2 \right) + 2 \int_{S_{\varepsilon}} \left(\frac{\partial u_0}{\partial x_2} \gamma_+ (V_{\varepsilon}) - \frac{\partial T_m}{\partial x_2} \gamma_+ (V_{\varepsilon,T_m,\Omega}) \right) dS.$$

By Remark 7.4 and Proposition 7.5 we have that, if either $\alpha_0 = \frac{j\pi}{m}$ or $\alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m}$ for some $j \in \{0, 1, \ldots, 2m-1\}$, then $\|V_{\varepsilon, T_m, \Omega}\|_{\mathcal{H}_{\varepsilon}} = \sqrt{2|\mathcal{E}_{\varepsilon, T_m, \Omega}|} = O(\varepsilon^m)$ as $\varepsilon \to 0^+$. Then, (7.23) follows from (7.22). Using again (7.22) we conclude that

(7.26)
$$\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^{2} - \|V_{\varepsilon,T_{m},\Omega}\|_{\mathcal{H}_{\varepsilon}}^{2} = (V_{\varepsilon} - V_{\varepsilon,T_{m},\Omega}, V_{\varepsilon} + V_{\varepsilon,T_{m},\Omega})_{\mathcal{H}_{\varepsilon}} = o(\varepsilon^{2m}) \quad \text{as } \varepsilon \to 0^{+}.$$

Furthermore, fixing some p > 2 and letting $p' = \frac{p}{p-1}$, Hölder's inequality, (7.21), and the continuity of the trace operators (2.3) imply that

$$(7.27) \qquad \left| \int_{S_{\varepsilon}} \left(\frac{\partial u_{0}}{\partial x_{2}} \gamma_{+}(V_{\varepsilon}) - \frac{\partial T_{m}}{\partial x_{2}} \gamma_{+}(V_{\varepsilon,T_{m},\Omega}) \right) dS \right| \\ = \left| \int_{S_{\varepsilon}} \left(\frac{\partial g}{\partial x_{2}} \gamma_{+}(V_{\varepsilon}) + \frac{\partial T_{m}}{\partial x_{2}} (\gamma_{+}(V_{\varepsilon}) - \gamma_{+}(V_{\varepsilon,T_{m},\Omega})) \right) dS \right| \\ \leq \int_{S_{\varepsilon}} \left| \frac{\partial g}{\partial x_{2}} \right| |\gamma_{+}(V_{\varepsilon})| \, dS + \int_{S_{\varepsilon}} \left| \frac{\partial T_{m}}{\partial x_{2}} \right| |\gamma_{+}(W_{\varepsilon})| \, dS \\ \leq \operatorname{const} \left(\varepsilon^{m+\frac{1}{p'}} \left(\int_{S_{\varepsilon}} |\gamma_{+}(V_{\varepsilon})|^{p} \, dS \right)^{1/p} + \varepsilon^{m-1+\frac{1}{p'}} \left(\int_{S_{\varepsilon}} |\gamma_{+}(W_{\varepsilon})|^{p} \, dS \right)^{1/p} \right) \\ \leq \operatorname{const} \left(\varepsilon^{m+\frac{1}{p'}} \|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} + \varepsilon^{m-1+\frac{1}{p'}} \|W_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}} \right) \\ = O\left(\varepsilon^{2m+\frac{1}{p'}} \right) + O\left(\varepsilon^{2m-\frac{1}{2}+\frac{1}{p'}} \right) = o(\varepsilon^{2m}) \quad \text{as } \varepsilon \to 0^{+},$$

where we used estimates (7.23) and (7.22). Combining (7.25), (7.26), and (7.27) we finally obtain (7.24).

Proof of Theorems 2.6 and 2.7. Since we are considering only two opposite poles on the same line, we have $v_0 = e^{-i\Theta_0}u_0 = u_0$. Let $m \in \mathbb{N} \setminus \{0\}$ and $\alpha_0 \in [0, \frac{\pi}{m})$ be as in (2.38).

If $\alpha_0 = \frac{j\pi}{m}$ or $\alpha_0 = \frac{\pi}{2m} + \frac{j\pi}{m}$ for some $j \in \{0, 1, \dots, 2m-1\}$, then, by (2.38) (see Remark 2.5), $u_0(x) = T_m(x) + O(|x|^{m+1})$ and $\frac{\partial u_0}{\partial x_2}(x) = \frac{\partial T_m}{\partial x_2}(x) + O(|x|^m)$ as $x \to 0$, so that

(7.28)
$$L_{\varepsilon}(u_0) = 2 \int_{S_{\varepsilon}} \frac{\partial T_m}{\partial x_2} T_m \, dS + O(\varepsilon^{2m+1}) = O(\varepsilon^{2m+1}) \quad \text{as } \varepsilon \to 0^+$$

since, in this case, either $T_m|_{S_{\varepsilon}} \equiv 0$ or $\frac{\partial T_m}{\partial x_2}|_{S_{\varepsilon}} \equiv 0$. From Theorem 2.1, (7.28), (7.23), and (7.24), it follows that

(7.29)
$$\lambda_{\varepsilon,n_0} - \lambda_{0,n_0} = 2\mathcal{E}_{\varepsilon} - 2L_{\varepsilon}(u_0) + o(\|V_{\varepsilon}\|_{\mathcal{H}_{\varepsilon}}^2) = 2\mathcal{E}_{\varepsilon,T_m,\Omega} + o(\varepsilon^{2m}) \quad \text{as } \varepsilon \to 0^+.$$

Theorem 2.6 follows from (7.29) and (7.19), while Theorem 2.7 is a consequence of (7.29) and (7.20).

Remark 7.7. The case m = 0 has been omitted in the present section as, for $u_0(0) \neq 0$ the sharp expansion is already contained in [5] even without symmetry assumptions on the domain; however, the above argument could also apply in such a case, providing an alternative proof of the result of [5].

8. Dealing with more general configurations of poles

In this section, we give a hint on how our approach could be extended to treat other possible configurations of poles, which are not covered in the present paper for the sake of simplicity of exposition. By Theorem 2.1, the quantity that sharply measures the eigenvalue variation is $\mathcal{E}_{\varepsilon} - L_{\varepsilon}(v_0)$, where $\mathcal{E}_{\varepsilon}$ is as in (2.19), L_{ε} as in (2.16) and v_0 is the limit eigenfunction after a gauge transformation, thus solving (2.14). As explained in the introduction, $\mathcal{E}_{\varepsilon}$ is essentially an intermediate quantity between a capacity and a torsional rigidity, measuring the set $\bigcup_{i=1}^{k_1+k_2} S_{\varepsilon}^{j}$. For the success of our method it is important that the limit eigenfunction v_0 is regular on the sets S_{ε}^{j} , while the perturbed eigenfunction v_{ε} jumps on them, together with $\nabla v_{\varepsilon} \cdot \nu^{j}$. Our approach can be applied to all configurations of poles for which, after a gauge transformation as in Section 3.3, the origin belongs to the half-lines on which the perturbed eigenfunction v_{ε} jumps.

We provide below some examples. Since the gauge transformation for a configuration of poles is the composition of the gauge transformations of the families of poles lying on the same straight line, we now focus on a single set of k collinear poles. Hence, for sake of simplicity, we assume

$$\{a^j\}_{j=1,...,k} \subset D_R(0) \cap \{(x_1,0) \colon x_1 \in \mathbb{R}\} \subset \Omega.$$

More precisely, we assume that $k = n_1 + n_2$, where $n_1, n_2 \in \mathbb{N}$ denote, respectively, the number of poles which lie on the left and on the right side with respect to the origin (either n_1 or n_2 might be zero). Namely,

$$a^{j} = \begin{cases} (-\delta_{j}, 0), & \text{for } j = 1, \dots, n_{1}, \\ (\delta_{j}, 0), & \text{for } j = n_{1} + 1, \dots, n_{1} + n_{2}, \end{cases}$$

where $\delta_i > 0$ are such that

$$-\delta_1 < -\delta_2 < \dots < -\delta_{n_1} < 0 < \delta_{n_1+1} < \dots < \delta_{n_1+n_2}$$

For the above configuration, we consider problem (1.4). One of the following cases occurs:

- (i) n_1 and n_2 are both even;
- (ii) n_1 and n_2 are both odd;
- (iii) n_1 is odd and n_2 is even (or vice versa).

The procedure developed to prove our main result Theorem 2.1 can be reproduced in cases (i) and (ii), as well as in case (iii) if $n_2 = 0$.

Let us now briefly describe, in these cases, how problem (1.4) becomes after a tailored gauge transformation. Hereafter, we denote by $\Sigma := \mathbb{R} \times \{0\}$ the x_1 axis, by $T: H^1(\mathbb{R}^2 \setminus \Sigma) \to L^p(\Sigma)$ the jump trace operator defined as in (2.4) with Σ instead of Σ^j , and by $\nu := (0, 1)$.

Case (i): even number of poles evenly distributed, i.e. $n_1 = 2N$ and $n_2 = 2M$ for some $N, M \in \mathbb{N}$ (see Figure 9a). In this case, reasoning as in Section 3.3, it is possible to find a gauge transformation such that problem (1.4) is equivalent to

$$\begin{cases} -\Delta v = \lambda v, & \text{in } \Omega \setminus \bigcup_{j=1}^{N+M} S_{\varepsilon}^{j}, \\ v = 0, & \text{on } \partial \Omega, \\ T(v) = T(\nabla v \cdot \nu) = 0, & \text{on } \bigcup_{j=1}^{N+M} S_{\varepsilon}^{j}, \end{cases}$$

where

$$S^j_{\varepsilon} := \begin{cases} [-\varepsilon \delta_{2j-1}, -\varepsilon \delta_{2j}] \times \{0\}, & \text{if } j = 1, \dots, N, \\ [\varepsilon \delta_{2j-1}, \varepsilon \delta_{2j}] \times \{0\}, & \text{if } j = N+1, \dots, N+M. \end{cases}$$

Case (ii): even number of poles oddly distributed, i.e. $n_1 = 2N + 1$ and $n_2 = 2M + 1$ for some $N, M \in \mathbb{N}$ (see Figure 9b). Once again, reasoning as in Section 3.3, one can find a gauge transformation such that problem (1.4) is equivalent to

$$\begin{cases} -\Delta v = \lambda v, & \text{in } \Omega \setminus \bigcup_{j=1}^{N+M+1} S_{\varepsilon}^{j} \\ v = 0, & \text{on } \partial \Omega, \\ T(v) = T(\nabla v \cdot \nu) = 0, & \text{on } \bigcup_{j=1}^{N+M+1} S_{\varepsilon}^{j}, \end{cases}$$

where

$$S_{\varepsilon}^{j} := \begin{cases} [-\varepsilon \delta_{2j-1}, -\varepsilon \delta_{2j}] \times \{0\}, & \text{for } j = 1, \dots, N, \\ [-\varepsilon \delta_{2N+1}, \varepsilon \delta_{2N+2}] \times \{0\}, & \text{for } j = N+1, \\ [\varepsilon \delta_{2j-1}, \varepsilon \delta_{2j}] \times \{0\}, & \text{for } j = N+2, \dots, N+M+1. \end{cases}$$

(iii): odd number of poles all on the same side, i.e. $n_1 = 2N + 1$ and $n_2 = 0$ (see Figure 9c). In this case, problem (1.4) is equivalent to

$$\begin{cases} -\Delta v = \lambda v, & \text{in } \Omega \setminus \left[\Gamma_0 \cup \left(\bigcup_{j=1}^{N+1} S^j_{\varepsilon} \right) \right], \\ v = 0, & \text{on } \partial \Omega, \\ T(v) = T(\nabla v \cdot \nu) = 0, & \text{on } \Gamma_0, \\ T(v) = T(\nabla v \cdot \nu) = 0, & \text{on } \bigcup_{j=1}^{N+1} S^j_{\varepsilon}, \end{cases}$$

where

$$S_{\varepsilon}^{j} := \begin{cases} [-\varepsilon \delta_{2j-1}, -\varepsilon \delta_{2j}] \times \{0\}, & \text{for } j = 1, \dots, N, \\ [-\varepsilon \delta_{2N+1}, 0] \times \{0\}, & \text{for } j = N+1. \end{cases}$$

To conclude, the only case left open in the present work is case (iii) with $n_2 \neq 0$. This requires non-trivial technical adaptations and will be the object of future investigation.



FIGURE 9. The jumping set after gauge transformation in cases (i), (ii), and (iii).

Acknowledgments. B. Noris has been supported by the MUR grant Dipartimento di Eccellenza 2023-2027 and by the INdAM-GNAMPA Project 2022 "Studi asintotici in problemi parabolici ed ellittici". V. Felli, R. Ognibene, and G. Siclari are partially supported by the INDAM-GNAMPA Project 2022 "Questioni di esistenza e unicità per problemi non locali con potenziali di tipo Hardy". R. Ognibene is supported by the project ERC VAREG - Variational approach to the regularity of the free boundaries (grant agreement No. 853404).

References

- ABATANGELO, L., AND FELLI, V. Sharp asymptotic estimates for eigenvalues of Aharonov-Bohm operators with varying poles. Calc. Var. Partial Differential Equations 54, 4 (2015), 3857–3903.
- [2] ABATANGELO, L., AND FELLI, V. On the leading term of the eigenvalue variation for Aharonov-Bohm operators with a moving pole. SIAM J. Math. Anal. 48, 4 (2016), 2843–2868.
- [3] ABATANGELO, L., FELLI, V., HILLAIRET, L., AND LÉNA, C. Spectral stability under removal of small capacity sets and applications to Aharonov-Bohm operators. J. Spectr. Theory 9, 2 (2019), 379–427.
- [4] ABATANGELO, L., FELLI, V., AND LÉNA, C. On Aharonov-Bohm operators with two colliding poles. Adv. Nonlinear Stud. 17, 2 (2017), 283–296.
- [5] ABATANGELO, L., FELLI, V., AND LÉNA, C. Eigenvalue variation under moving mixed Dirichlet-Neumann boundary conditions and applications. ESAIM Control Optim. Calc. Var. 26 (2020), Paper No. 39, 47.
- [6] ABATANGELO, L., FELLI, V., NORIS, B., AND NYS, M. Sharp boundary behavior of eigenvalues for Aharonov-Bohm operators with varying poles. J. Funct. Anal. 273, 7 (2017), 2428–2487.
- [7] ABATANGELO, L., AND NYS, M. On multiple eigenvalues for Aharonov-Bohm operators in planar domains. Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods 169 (2018), 1–37.
- [8] ABATANGELO, L., AND OGNIBENE, R. Sharp behavior of dirichlet–laplacian eigenvalues for a class of singularly perturbed problems. *Preprint* (2023), arxiv.org/abs/2301.11729.
- [9] ALZIARY, B., FLECKINGER-PELLÉ, J., AND TAKÁČ, P. Eigenfunctions and Hardy inequalities for a magnetic Schrödinger operator in ℝ². Math. Methods Appl. Sci. 26, 13 (2003), 1093–1136.
- [10] BONNAILLIE-NOËL, V., HELFFER, B., AND HOFFMANN-OSTENHOF, T. Aharonov-Bohm Hamiltonians, isospectrality and minimal partitions. J. Phys. A 42, 18 (2009), 185203, 20.
- [11] BONNAILLIE-NOËL, V., NORIS, B., NYS, M., AND TERRACINI, S. On the eigenvalues of Aharonov-Bohm operators with varying poles. Anal. PDE 7, 6 (2014), 1365–1395.
- [12] BRASCO, L., DEL MAR GONZÁLEZ, M., AND ISPIZUA, M. A Steklov version of the torsional rigidity. Preprint (2022).
- [13] BRASCO, L., AND MAZZOLENI, D. On principal frequencies, volume and inradius in convex sets. NoDEA, Nonlinear Differ. Equ. Appl. 27, 2 (2020), 26. Id/No 12.
- [14] COURTOIS, G. Spectrum of manifolds with holes. J. Funct. Anal. 134, 1 (1995), 194–221.

- [15] FELLI, V., FERRERO, A., AND TERRACINI, S. Asymptotic behavior of solutions to Schrödinger equations near an isolated singularity of the electromagnetic potential. J. Eur. Math. Soc. (JEMS) 13, 1 (2011), 119–174.
- [16] HELFFER, B. Spectral theory and its applications, vol. 139 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2013.
- [17] HENROT, A., AND PIERRE, M. Shape variation and optimization. A geometrical analysis, vol. 28 of EMS Tracts Math. Zürich: European Mathematical Society (EMS), 2018.
- [18] LAPTEV, A., AND WEIDL, T. Hardy inequalities for magnetic Dirichlet forms. In Mathematical results in quantum mechanics (Prague, 1998), vol. 108 of Oper. Theory Adv. Appl. Birkhäuser, Basel, 1999, pp. 299– 305.
- [19] LEINFELDER, H. Gauge invariance of Schrödinger operators and related spectral properties. J. Operator Theory 9, 1 (1983), 163–179.
- [20] LÉNA, C. Eigenvalues variations for Aharonov-Bohm operators. J. Math. Phys. 56, 1 (2015), 011502, 18.
- [21] LEONI, G. A first course in Sobolev spaces, second ed., vol. 181 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2017.
- [22] MEDKOVÁ, D., AND KRUTITSKII, P. Neumann and Robin problems in a cracked domain with jump conditions on cracks. J. Math. Anal. Appl. 301, 1 (2005), 99–114.
- [23] NORIS, B., NYS, M., AND TERRACINI, S. On the Aharonov-Bohm operators with varying poles: the boundary behavior of eigenvalues. *Comm. Math. Phys.* 339, 3 (2015), 1101–1146.
- [24] NORIS, B., AND TERRACINI, S. Nodal sets of magnetic Schrödinger operators of Aharonov-Bohm type and energy minimizing partitions. *Indiana Univ. Math. J.* 59, 4 (2010), 1361–1403.
- [25] PÓLYA, G., AND SZEGÖ, G. Isoperimetric inequalities in mathematical physics, vol. 27 of Ann. Math. Stud. Princeton University Press, Princeton, NJ, 1951.
- [26] TEYTEL, M. How rare are multiple eigenvalues? Commun. Pure Appl. Math. 52, 8 (1999), 0917–0934.
- [27] VAN DEN BERG, M., BUTTAZZO, G., AND VELICHKOV, B. Optimization problems involving the first Dirichlet eigenvalue and the torsional rigidity. In New trends in shape optimization, vol. 166 of Internat. Ser. Numer. Math. Birkhäuser/Springer, Cham, 2015, pp. 19–41.

VERONICA FELLI AND GIOVANNI SICLARI DIPARTIMENTO DI MATEMATICA E APPLICAZIONI UNIVERSITÀ DEGLI STUDI DI MILANO-BICOCCA VIA COZZI 55, 20125 MILANO, ITALY Email address: veronica.felli@unimib.it, g.siclari2@campus.unimib.it

BENEDETTA NORIS DIPARTIMENTO DI MATEMATICA POLITECNICO DI MILANO PIAZZA LEONARDO DA VINCI 32, 20133 MILANO, ITALY *Email address*: benedetta.noris@polimi.it

ROBERTO OGNIBENE DIPARTIMENTO DI MATEMATICA UNIVERSITÀ DI PISA LARGO BRUNO PONTECORVO, 5, 56127 PISA, ITALY Email address: roberto.ognibene@dm.unipi.it