

# Bayesian Optimization of Expensive Nested Grey-Box Functions

Wenjie Xu · Yuning Jiang ·  
Bratislav Svetožarević · Colin N. Jones

the date of receipt and acceptance should be inserted later

**Abstract** We consider the problem of optimizing a *grey-box* objective function, i.e., nested function composed of both *black-box* and *white-box* functions. A general formulation for such grey-box problems is given, which covers the existing grey-box optimization formulations as special cases. We then design an *optimism*-driven algorithm to solve it. Under certain regularity assumptions, our algorithm achieves similar regret bound as that for the standard black-box Bayesian optimization algorithm, up to a constant multiplicative term depending on the Lipschitz constants of the functions considered. We further extend our method to the constrained case and discuss special cases. For the commonly used kernel functions, the regret bounds allow us to derive a convergence rate to the optimal solution. Experimental results show that our grey-box optimization method empirically improves the speed of finding the global optimal solution significantly, as compared to the standard black-box optimization algorithm.

## 1 Introduction

We consider the problem of optimizing a *grey-box* objective function, i.e., nested function composed of both *black-box* and *white-box* functions. Such problems arise from a wide variety of applications. For example, while we optimize an economic objective subject to some operational constraints in process

---

W. Xu, Y. Jiang, C. Jones  
Automatic Control Laboratory  
EPFL, Switzerland  
E-mail: wenjie.xu, yuning.jiang, colin.jones@epfl.ch

W. Xu and B. Svetožarević are also  
with Swiss Federal Laboratories  
for Materials Science and Technology (Empa)  
E-mail: bratislav.svetozarevic@empa.ch

control [10,27], the objective can be a bilinear function in known price variable and product fraction variables, where the latter are unknown black-box functions of the tunable operational parameters.

To solve such problems, one may ignore the grey-box structure and treat the nested function as a pure input-to-output black-box function. Bayesian optimization (BO) [11] has been shown to be an effective method to solve the purely black-box optimization problem [19]. However, the agnostic application of the black-box BO method to the grey-box problem has several limitations,

- Fails to utilize the intermediate observations and the known structure of how different *black-box* and *white-box* elements influence each other. However, it has been shown that such structural information can empirically boost the performance of Bayesian optimization [4].
- May violate the regularity assumptions necessary for convergence guarantee. For example, a common assumption for convergence guarantee of the black-box Bayesian optimization method is the bounded norm inside a reproducing kernel Hilbert space [20,27]. This assumption can be violated for the grey-box function with a discontinuous white-box element.

In this paper, we consider a general grey-box optimization problem and propose an *optimism*-driven algorithm to solve it. Specifically, our contributions are summarized as follows:

- A general formulation for grey-box optimization problems is given, which covers the existing grey-box optimization formulations [4,5,16] as special cases.
- An *optimism*-driven algorithm is designed to solve it. Under certain regularity assumptions, our algorithm achieves similar regret bound as that for the standard black-box Bayesian optimization algorithm, up to a constant multiplicative term depending on the Lipschitz constants of the functions considered. For the commonly used kernel functions, the regret bounds allow us to derive a convergence rate to the optimal solution.
- Our method is further extended to the constrained case and several special cases are discussed. Experimental results show that our grey-box optimization method empirically improves the speed of finding the global optimal solution significantly, as compared to the standard black-box optimization algorithm.

## 2 Related work

The problem we consider in this paper follows the research line of Bayesian optimization (BO), a framework of efficient global optimization of expensive-to-evaluate black-box function [30,17,14,26]. Standard Bayesian optimization [11] assumes that the expensive-to-evaluate functions are purely black-box. That is to say, the function is not known explicitly but only provides an oracle that can be queried to get the zero-order function evaluation. There are also

works [12, 13, 28, 25, 29] that extend the black-box Bayesian optimization to the constrained setting.

More recently, grey-box Bayesian optimization has been receiving more and more attention. For example, [15] and [24] use cheap approximations of the black-box functions to do multi-fidelity BO. [22, 7] propose a Bayesian optimization method for the unknown black-box function with integration structure. More generally, [4, 16] considers Bayesian optimization of a composite function, with an expensive-to-evaluate inner black-box function and cheap-to-evaluate outer known function. [4] further considers Bayesian optimization of a network of functions.

Another line of research considers causal Bayesian optimization [2, 1, 3, 21], where the relationship between intervention/input variables and target/output variables are modeled as a causal graph. However, this line of research results only considered the probabilistic modelling of the variables, but did not consider the known intermediate white-box functions.

Existing works on grey-box Bayesian optimization either only consider special structures such as addition, integration [22, 7], or one known function composed with black-box functions [4, 16], or can only provide very weak theoretical guarantee (such as the consistency of finding the global optimum). Our work can be regarded as a general extension of the research line for grey-box Bayesian optimization. Furthermore, we design an *optimism*-driven algorithm to solve the problem with a global convergence rate guarantee. To highlight the contributions of our work, we compare our method with the other state-of-the-art methods in Tab. 1.

**Table 1** The comparison of our method to existing state-of-the-art grey-box Bayesian optimization methods, where ‘hybrid’ means the grey-box function is composed of both unknown black-box functions and known white-box functions.

| Works                            | Regret Bound | Constraints Consideration | Multilevel Modelling | Function Type |
|----------------------------------|--------------|---------------------------|----------------------|---------------|
| Composite EI [4], etc.           | ✗            | ✗                         | ✗                    | hybrid        |
| BO of function network [5], etc. | ✗            | ✗                         | ✓                    | hybrid        |
| CUQB [16]                        | ✓            | ✓                         | ✗                    | hybrid        |
| Causal BO [2, 1]                 | ✗            | ✗                         | ✓                    | black-box     |
| Constrained Causal BO [3]        | ✗            | ✓                         | ✓                    | black-box     |
| Model-based Causal BO [21]       | ✓            | ✗                         | ✓                    | black-box     |
| Ours                             | ✓            | ✓                         | ✓                    | hybrid        |

### 3 Grey-box modelling

We use a hybrid factorable function to model the grey-box function. Sec. 3.1 will give the algebraic representation of the grey-box function, and Sec. 3.2 will

give its corresponding graph representation. Before introducing our grey-box modelling approach, we first clarify what we will mean by *black-box function* and *white-box function*. An expensive unknown function to which we can only query zero-order function evaluation is referred to as a *black-box function*. A cheap known function which has explicit mathematical expression is referred to as a *white-box function*.

### 3.1 Algebraic representation

We first introduce the algebraic representation of the grey-box function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . The modelling idea is introducing a set of intermediate variables  $z_i, i \in [m]$  and intermediate functions  $\phi_{i-1}, i \in [m]$ . Formally, the following definitions are needed.

**Definition 3.1 (Augmented State)**  $\forall x \in \mathbb{R}^n, z \in \mathbb{R}^m$ , we define the augmented state by,

$$s_0 = x = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}, s_1 = \begin{bmatrix} x_1 \\ \vdots \\ x_n \\ z_1 \end{bmatrix}, \dots, s_m = \begin{bmatrix} x_1 \\ \vdots \\ x_n \\ z_1 \\ \vdots \\ z_m \end{bmatrix}.$$

**Definition 3.2 (Augmented Elementary Function)** Augmented elementary function  $\Phi_i : \mathbb{R}^{n+i} \rightarrow \mathbb{R}^{n+i+1}$  is given as

$$\Phi_i(s_i) = \begin{bmatrix} x_1 \\ \vdots \\ x_n \\ z_1 \\ \vdots \\ z_i \\ \phi_i(x_1, \dots, x_n, z_1, \dots, z_i) \end{bmatrix}, s_{i+1} = \Phi_i(s_i),$$

where  $\phi_i : \mathbb{R}^{n+i} \rightarrow \mathbb{R}$  is either a known white-box function or an unknown black-box function. Thus, it can be seen that  $z_{i+1} = \phi_i(s_i), i \in \{0, \dots, m-1\}$ .

**Remark 3.1** Note that here  $\phi_i$  takes all the variables in  $\{x_1, x_2, \dots, x_n, z_1, \dots, z_m\}$  as input. In practice,  $\phi_i$  can be a function of a subset of variables  $\{x_1, \dots, x_n, z_1, \dots, z_i\}$ . All the following algorithms and analyses can be easily adapted to the practical setting of the subset input variables. For the simplicity of notation, we will focus on the setting that  $\phi_i$  takes the whole  $s_i$  as the input in the description of the algorithm and the theoretical analysis.


We can now define the *grey-box* objective function by

$$f(x) = e_{n+m}^T \Phi_{m-1} (\Phi_{m-2} (\dots \Phi_1 (\Phi_0(x)))) , \quad (1)$$

where  $e_{n+m} \in \mathbb{R}^{n+m}$  is the vector with the last element being 1 and all the other elements being 0. We use  $\mathcal{I}_{\mathcal{W}} \subset \{0, \dots, m-1\}$  ( $\mathcal{I}_{\mathcal{B}} \subset \{0, \dots, m-1\}$ , resp.) to denote the index set of the known white-box (unknown black-box, resp.) functions among  $\phi_i, i \in \{0, \dots, m-1\}$ . It holds that  $\mathcal{I}_{\mathcal{W}} \cup \mathcal{I}_{\mathcal{B}} = \{0, \dots, m-1\}$ .

### 3.2 Graph representation

The nested grey-box function can also be represented as an acyclic-directed graph  $\mathcal{G} := (\mathcal{V}, \mathcal{E})$ .  $\mathcal{V} = \mathcal{V}_x \cup \mathcal{V}_\phi$  is the vertices set, where  $\mathcal{V}_x$  corresponds to the input variables  $(x_i)_{i=1}^n$  and  $\mathcal{V}_\phi$  corresponds to the intermediate variables  $(z_i)_{i=1}^m$  that are outputs of the functions  $(\phi_i)_{i=0}^{m-1}$ .  $(v_i, v_j) \in \mathcal{E}$  if and only if the variable of  $v_j$  is either a black-box or white-box function that takes the variable of  $v_i$  as one of the inputs. For example, Fig. 1 gives the graph representation of a simple nested grey-box function  $f(x) = (\phi_1(x_1, x_2) - x_3)^2 + x_3^2$ , where  $\phi_1$  is a black-box function.



**Fig. 1** A simple example of the graph representation of a nested grey-box function.

### 3.3 Some examples

We now give examples to demonstrate that our grey-box modeling captures a rich class of functions.

1. Additive black-box functions with different input variables.

$$f(x) = \phi_1(x_1, x_2) + \phi_2(x_2, x_3) + \phi_3(x_1, x_3), \quad (2)$$

where  $\phi_i, i \in [3]$  are all black-box functions.

2. Black-box functions composed with white box functions.

$$f(x) = \phi_1(x)^2 + \phi_2(x)^2, \quad (3)$$

where  $\phi_1$  and  $\phi_2$  are black-box functions.

### 3. Hybrid chain of black-box/white-box functions

$$f(x) = \phi_3 (\phi_1(x)^2 + 3\phi_1(x) - 3), \quad (4)$$

where  $\phi_1$  and  $\phi_3$  are black-box functions.

## 4 Problem formulation

We consider the following unconstrained global optimization problem of grey-box functions.

$$\min_{x \in \mathcal{X}} f(x), \quad (5a)$$

$$\text{subject to } g_k(x) \leq 0, k \in [K], \quad (5b)$$

where  $\mathcal{X} \subset \mathbb{R}^n$  is a known compact subset of  $\mathbb{R}^n$ ,  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is a grey-box function as given in Eq. (1),  $g_k : \mathbb{R}^n \rightarrow \mathbb{R}$  is a constraint grey-box function similarly defined and  $K$  is the number of constraints.

Eq. (5) is a generic formulation of constrained grey-box optimization problem. This section focuses on the unconstrained case, where  $K = 0$  and we focus on  $\min_{x \in \mathcal{X}} f(x)$ . We will show later in Sec. 8 that our method can easily be extended to the constrained case by adopting a similar algorithmic idea as proposed in [25].

At step  $t$ , with the sample point  $x^t$ , there is a vector of augmented states  $\{z_{i+1}^t\}_{i \in \mathcal{I}_{\mathcal{B}}}$ , where  $z_{i+1}^t$  is the output of the black-box function  $\phi_i$  with input  $s_i^t$ . The relationship of  $(x^t, z^t, s^t)$  is as given in Def. 3.1 and Def. 3.2. Note that  $z_{i+1}^t$  may not be observable, or the observation is corrupted by noise in practice.

Similar to [25], we make some regularity assumptions on the grey-box optimization problem (5).

**Assumption 4.1** *Set  $\mathcal{X}$  is compact.*

**Assumption 4.2** *Functions  $\phi_i \in \mathcal{H}_{k_i}, i \in \mathcal{I}_{\mathcal{B}}$ , where  $k_i : \mathbb{R}^{n_i} \times \mathbb{R}^{n_i} \rightarrow \mathbb{R}, i \in \mathcal{I}_{\mathcal{B}}$  are symmetric, positive-semidefinite kernel functions,  $\mathcal{H}_{k_i}$  are the corresponding reproducing kernel Hilbert spaces (RKHSs, see [18]), and  $n_i$  is the dimension of the input variables for the function  $\phi_i$ . Furthermore, we assume  $\|\phi_i\|_{k_i} \leq B_i, i \in \mathcal{I}_{\mathcal{B}}$ , where  $\|\cdot\|_{k_i}$  is the norm induced by the inner product in the corresponding reproducing kernel Hilbert space.*

**Assumption 4.3**  $\forall s_i \in \mathbb{R}^{n+i}, i \in \mathcal{I}_{\mathcal{B}}, k_i(s_i, s_i) \leq 1$ .

**Assumption 4.4** *There exists a hyperbox*

$$\mathcal{C}_z := [-C_{z,1}, C_{z,1}] \times \cdots \times [-C_{z,m}, C_{z,m}]$$

with  $C_{z,i} > 0, i \in [m]$ , such that  $\forall x \in \mathcal{X}$ , the corresponding intermediate variables  $z \in \mathcal{C}_z$ . Furthermore, we choose  $C_{z,i+1} \geq B_i, i \in \mathcal{I}_{\mathcal{B}}$ .

To facilitate the following discussion, we use shorthand

$$\mathcal{S}_i = \mathcal{X} \times [-C_{z,1}, C_{z,1}] \times \cdots \times [-C_{z,i}, C_{z,i}].$$

**Assumption 4.5** *The functions  $\phi_i$ ,  $i \in \{0, \dots, m-1\}$  are Lipschitz continuous with respect to one norm over  $\mathcal{S}_i := \mathcal{X} \times [-C_{z,1}, C_{z,1}] \times \cdots \times [-C_{z,i}, C_{z,i}]$ . That is to say, there exist constants  $L_{\phi_i} > 0$ , such that  $\forall s_i^1, s_i^2 \in \mathcal{S}_i$ , we have,*

$$\|\phi_i(s_i^1) - \phi_i(s_i^2)\| \leq L_{\phi_i} \|s_i^1 - s_i^2\|_1, \quad (6)$$

**Assumption 4.6** *The black-box functions  $\phi_i$ ,  $\forall i \in \mathcal{I}_{\mathcal{B}}$  can be separately evaluated. That is to say we get access to an oracle to evaluate each black-box function  $\phi_i$ ,  $i \in \mathcal{I}_{\mathcal{B}}$ . Furthermore, each observation of the output for the black-box functions is corrupted by independent identically distributed  $\sigma$ -sub-Gaussian noise.<sup>1</sup>*

## 5 Learning of the black-box functions with Gaussian process

We use the independent Gaussian processes to learn the unknown black-box functions  $\phi_i$ ,  $i \in \mathcal{I}_{\mathcal{B}}$ . Similar to [8], we artificially introduce a Gaussian process  $\mathcal{GP}(0, k_i(\cdot, \cdot))$  for the surrogate modeling of the unknown black-box function  $\phi_i$ . We also adopt an *i.i.d* Gaussian zero-mean noise model with noise variance  $\lambda > 0$ .

**Remark 5.1** *Note that this Gaussian process model is only used to derive posterior mean functions, covariance functions, and maximum information gain for the purpose of algorithm design. We introduce the artificial Gaussian process only to make the algorithm more interpretable. It does not change our set-up that  $\phi_i, i \in \mathcal{I}_{\mathcal{B}}$  is a deterministic function and that the observational noise only needs to be *i.i.d*  $\sigma$ -sub-Gaussian. Indeed, we are considering the agnostic setting introduced in [20] and the noise  $\lambda$  here can also be interpreted as a regularizer that stabilizes the numerical computation.*

We use  $(\bar{s}_i^t, \tilde{z}_{i+1}^t), i \in \mathcal{I}_{\mathcal{B}}$  to denote the data we add to learn the black-box function  $\phi_i$  at step  $t$ , where  $\tilde{z}_{i+1}^t = \phi_i(\bar{s}_i^t) + \xi_i^t$ . We introduce the following functions for  $s_i, s'_i$  for the learning of the function  $\phi_i, i \in \mathcal{I}_{\mathcal{B}}$ ,

$$\mu_{i,t}(s_i) = k_i(\bar{s}_i^{1:t}, s_i)^\top (K_{i,t} + \lambda I)^{-1} \tilde{z}_{i+1}^{1:t}, \quad (7a)$$

$$k_{i,t}(s_i, s'_i) = k_i(s_i, s'_i) - k_i(\bar{s}_i^{1:t}, s_i)^\top (K_{i,t} + \lambda I)^{-1} k_i(\bar{s}_i^{1:t}, s'_i), \quad (7b)$$

$$\sigma_{i,t}^2(s_i) = k_{i,t}(s_i, s_i) \quad (7c)$$

with  $k_i(\bar{s}_i^{1:t}, s_i) = [k_i(\bar{s}_i^1, s_i), k_i(\bar{s}_i^2, s_i), \dots, k_i(\bar{s}_i^t, s_i)]^\top$ ,  $K_{i,t} = (k_i(\bar{s}_i^{\tau_1}, \bar{s}_i^{\tau_2}))_{\tau_1, \tau_2 \in [t]}$  and  $\tilde{z}_{i+1}^{1:t} = [\tilde{z}_{i+1}^1, \tilde{z}_{i+1}^2, \dots, \tilde{z}_{i+1}^t]^\top$ . We also introduce the maximum information gain for the function  $\phi_i$  as proposed in [20],

$$\gamma_{i,t} := \max_{S_i \subset \mathcal{S}_i; |S_i|=t} \frac{1}{2} \log |I + \lambda^{-1} K_{i,S_i}|, \quad (8)$$

<sup>1</sup> Note that the noise is only for observation. There is no process noise in the compositional chain of the grey-box function.

where  $K_{i,S_i} = (k_i(s_i, s'_i))_{s_i, s'_i \in S_i}$ .

**Lemma 5.1 (Theorem 2, [8])** *Let Assumptions 4.1, 4.2, 4.3, 4.4 and 4.6 hold. For any  $\delta \in (0, 1)$ , with probability at least  $1 - \delta/m$ , the following holds for all  $s_i \in \mathcal{S}_i, i \in \mathcal{I}_B$  and  $1 \leq t \leq T, T \in \mathbb{N}$ ,*

$$|\mu_{i,t-1}(s_i) - \phi_i(s_i)| \leq \left( B_i + \sigma \sqrt{2(\gamma_{i,t-1} + 1 + \ln(m/\delta))} \right) \sigma_{i,t-1}(s_i), \quad (9)$$

where  $\mu_{i,t-1}(s_i), \sigma_{i,t-1}^2(x)$  and  $\gamma_{i,t-1}$  are as given in Eq. (7) and Eq. (8), and  $\lambda$  is set to be  $1 + \frac{2}{T}$ . ( $\mu_{i,0}$  and  $\sigma_{i,0}$  are the prior mean and standard deviation, and  $\gamma_{i,0} = 0$ .)

**Remark 5.2** *Note that we replace the ‘ $\delta$ ’ in Theorem 2 of [8] by  $\delta/m$ , which will be useful to derive the confidence interval with  $1 - \delta$  probability in Lem. 5.2.*

To facilitate the following algorithm design and discussion, we introduce the lower confidence and upper confidence bound functions,

**Definition 5.1** For  $i \in \mathcal{I}_B, s_i \in \mathcal{S}_i$  and  $t \in [T]$ ,

$$l_{i,t}(s_i) := \max \{ \mu_{i,t-1}(s_i) - \beta_{i,t} \sigma_{i,t-1}(s_i), -B_i \} , \quad (10a)$$

$$u_{i,t}(s_i) := \min \{ \mu_{i,t-1}(s_i) + \beta_{i,t} \sigma_{i,t-1}(s_i), B_i \} \quad (10b)$$

with  $\beta_{i,t} = B_i + \sigma \sqrt{2(\gamma_{i,t-1} + 1 + \ln(m/\delta))}$ .

Lem. 5.2 then follows,

**Lemma 5.2 (Corollary 2.6, [25])** *Let Assumptions 4.1, 4.2, 4.3, 4.4 and 4.6 hold. With probability at least  $1 - \delta$ , the following holds for all  $s_i \in \mathcal{S}_i, i \in \mathcal{I}_B$  and  $1 \leq t \leq T$ ,*

$$\forall i \in \mathcal{I}_B, \phi_i(s_i) \in [l_{i,t}(s_i), u_{i,t}(s_i)] . \quad (11)$$

## 6 Algorithm

The Bayesian optimization algorithm of expensive nested grey-box functions is given as in Alg. 1. The idea of the Alg. 1 is that in each step, we solve an optimistic auxiliary problem of the original problem where we require the augmented states  $z_{i+1}, i \in \mathcal{I}_B$  need to be inside the confidence interval conditioned on the current observation history.<sup>2</sup>

**Remark 6.1** *We assume that an oracle can efficiently solve the auxiliary problem in line 2. In low dimensions, we can use pure grid search to solve it. In higher dimensional cases, we can apply a gradient-based optimization algorithm with different starting points.*

<sup>2</sup> Note that to observe these intermediate variables, we may suffer from observational noise, which may introduce noise perturbation to the subsequent Gaussian process input.



**Algorithm 1** Bayesian Optimization of Expensive Nested Grey Box Functions

- 
- 1: **for**  $t \in [T]$  **do**
  - 2: Find  $(x^t, \bar{z}^t)$  by solving
 
$$\min_{x \in \mathcal{X}, z \in \mathbb{R}^m} z_m \quad (12a)$$
 subject to  $l_{i,t}(s_i) \leq z_{i+1} \leq u_{i,t}(s_i), \forall i \in \mathcal{I}_{\mathcal{B}}, \quad (12b)$ 

$$z_{i+1} = \phi_i(s_i), \forall i \in \mathcal{I}_{\mathcal{W}}, \quad (12c)$$

$$s_i = [x_1, \dots, x_n, z_1, \dots, z_i]^\top, \forall i \in \{0\} \cup [m]. \quad (12d)$$
  - 3: Evaluate the grey-box function  $f$  at  $x^t$ , incur intermediate variables  $z^t$  and final objective  $f(x^t) = z_m^t$ .
  - 4: Separately get the noisy evaluation  $\bar{z}_{i+1}^t := \phi_i(\bar{s}_i^t) + \bar{\xi}_{i,t}$  of the black-box function  $\phi_i, i \in \mathcal{I}_{\mathcal{B}}$  at the point  $\bar{s}_i^t := [x_1^t, \dots, x_n^t, z_1^t, \dots, z_i^t]^\top$ .
  - 5: Update Gaussian process posterior mean and covariance for each black-box function with the new evaluations added.
  - 6: **end for**
- 

**7 Analysis**

We now analyze the performance of the algorithm. We use  $x^*$  to denote the optimal solution of the Prob. (5) and  $z^*$  to denote the corresponding intermediate variables. We are interested in bounding the cumulative regret,

$$R_T = \sum_{t=1}^T (f(x^t) - f(x^*)). \quad (13)$$

We first introduce a lemma to bound the discrepancy between the real output of a grey box function and the *possible* output of the grey-box function that is consistent with the posterior of Gaussian process with high probability.

**Lemma 7.1 (Output Discrepancy Bound)** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, and 4.6 hold. For any  $x^t \in \mathcal{X}$ , let  $z^t$  denote its corresponding augmented state when evaluating the grey-box function at  $x^t$ . Then for any  $z$  that satisfies,*

$$l_{i,t}(s_i) \leq z_{i+1} \leq u_{i,t}(s_i), \quad \forall i \in \mathcal{I}_{\mathcal{B}}, \quad (14)$$

$$z_{i+1} = \phi_i(s_i), \quad \forall i \in \mathcal{I}_{\mathcal{W}} \quad (15)$$

with  $s_i = [x_1^t, \dots, x_n^t, z_1, \dots, z_i]^\top, \forall i \in \{0\} \cup [m-1]$ . We have, with probability at least  $1 - \delta$ ,

$$|f(x^t) - z_m| = |z_m^t - z_m| \leq \sum_{i \in \mathcal{I}_{\mathcal{B}}} A_i \beta_{i,t} \sigma_{i,t-1}(s_i)$$

with  $A_i = 2 \left( \sum_{j=1}^{m-1-i} 2^j \prod_{i < k_1 < \dots < k_j = m-1} L_{\phi_{k_j}} \right)$ .

*Proof* We consider bounding  $|z_{i+1}^t - z_{i+1}|, i \in [m-1]$  recursively.

**Case 1:**  $\phi_i$  is a known white-box function, then,

$$|z_{i+1}^t - z_{i+1}| = |\phi_i(s_i^t) - \phi_i(s_i)| \leq L_{\phi_i} \|s_i^t - s_i\|_1 \leq 2L_{\phi_i} \sum_{j=1}^i |z_j^t - z_j|. \quad (16)$$

**Case 2:**  $\phi_i$  is an unknown black-box function, then,

$$|z_{i+1}^t - z_{i+1}| \quad (17a)$$

$$\leq |\phi_i(s_i^t) - l_{i,t}(s_i)| + |\phi_i(s_i^t) - u_{i,t}(s_i)| \quad (17b)$$

$$\leq |\phi_i(s_i^t) - \phi_i(s_i)| + |\phi_i(s_i) - l_{i,t}(s_i)| \\ + |\phi_i(s_i^t) - \phi_i(s_i)| + |\phi_i(s_i) - u_{i,t}(s_i)| \quad (17c)$$

$$\leq 2L_{\phi_i} \|s_i^t - s_i\| + 2\beta_{i,t}\sigma_{i,t-1}(s_i) \quad (17d)$$

$$= 2L_{\phi_i} \sum_{j=1}^i |z_j^t - z_j| + 2\beta_{i,t}\sigma_{i,t-1}(s_i). \quad (17e)$$

where the inequality (17b) follows by  $z_{i+1}^t = \phi_i(z_i^t)$  and  $z_{i+1} \in [l_{i,t}(s_i), u_{i,t}(s_i)]$ , the inequality (17c) follows by the triangle inequality, and the inequality (17d) follows by the Lipschitz assumption and that  $z_{i+1} \in [l_{i,t}(s_i), u_{i,t}(s_i)]$ . Solving the recursive inequalities above, we get

$$|z_m^t - z_m| \quad (18a)$$

$$\leq \sum_{i \in \mathcal{I}_{\mathcal{B}}} 2 \left( \sum_{j=1}^{m-1-i} 2^j \prod_{i < k_1 < \dots < k_j = m-1} L_{\phi_{k_j}} \right) \beta_{i,t}\sigma_{i,t-1}(s_i), \quad (18b)$$

which concludes the proof.  $\square$

We then have a lemma to bound the single-step instantaneous regret.

**Lemma 7.2** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, and 4.6 hold. There exists constants  $A_i, i \in \mathcal{I}_{\mathcal{B}}$ , such that with probability at least  $1 - \delta$ ,*

$$[f(x^t) - f(x^*)]^+ \leq \sum_{i \in \mathcal{I}_{\mathcal{B}}} A_i \beta_{i,t} \sigma_{i,t-1}(\bar{s}_i^t) \quad (19)$$

with  $A_i = 2 \left( \sum_{j=1}^{m-1-i} 2^j \prod_{i < k_1 < \dots < k_j = m-1} L_{\phi_{k_j}} \right)$ .

*Proof* With probability at least  $1 - \delta$ ,

$$[f(x^t) - f(x^*)]^+ \quad (20a)$$

$$= [z_m^t - \bar{z}_m^t + \bar{z}_m^t - f(x^*)]^+ \quad (20b)$$

$$\leq [z_m^t - \bar{z}_m^t]^+ + [\bar{z}_m^t - f(x^*)]^+ \quad (20c)$$

$$\leq [z_m^t - \bar{z}_m^t]^+ \quad (20d)$$

where the equality (20b) follows by the definition of  $z^t$ , the inequality (20d) follows by the optimality of  $(x^t, z^t)$  for the auxiliary problem in line 2 of the Alg. 1 and the feasibility of  $(x^*, z^*)$  for the same auxiliary problem. Applying the bound in Lemma 7.1 and setting  $A_i = 2 \left( \sum_{j=1}^{m-1-i} 2^j \prod_{i < k_1 < \dots < k_j = m-1} L_{\phi_{k_j}} \right)$  concludes the proof.  $\square$

We then restate a useful lemma to bound the cumulative posterior standard deviation along the sampling trajectory,

**Lemma 7.3 (Lemma 4, [9])** *If  $\bar{s}_i^1, \bar{s}_i^2, \dots, \bar{s}_i^T$  are the points sampled by Alg. 1 in line 4, then,*

$$\sum_{t=1}^T \sigma_{i,t-1}(\bar{s}_i^t) \leq \sqrt{4(T+2)\gamma_{i,T}}. \quad (21)$$

**Theorem 7.1** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, and 4.6 hold. Then, with probability at least  $1 - \delta$ , the sample points of Algorithm 1 satisfy,*

$$R_T \leq \sum_{i \in \mathcal{I}_B} 2A_i \beta_{i,T} \sqrt{(T+2)\gamma_{i,T}} = \mathcal{O}\left(\sum_{i \in \mathcal{I}_B} A_i \gamma_{i,T} \sqrt{T}\right), \quad (22)$$

where  $A_i$  is a Lipschitz constant dependent constant as given in Lem. 7.2.

*Proof* We have

$$R_T = \sum_{t=1}^T (f(x^t) - f(x^*)) \quad (23a)$$

$$\leq \sum_{t=1}^T \sum_{i \in \mathcal{I}_B} A_i \beta_{i,t} \sigma_{i,t-1}(\bar{s}_i^t) \quad (23b)$$

$$\leq \sum_{t=1}^T \sum_{i \in \mathcal{I}_B} A_i \beta_{i,T} \sigma_{i,t-1}(\bar{s}_i^t) \quad (23c)$$

$$= \sum_{i \in \mathcal{I}_B} A_i \beta_{i,T} \sum_{t=1}^T \sigma_{i,t-1}(\bar{s}_i^t) \quad (23d)$$

$$\leq \sum_{i \in \mathcal{I}_B} A_i \beta_{i,T} \sqrt{4(T+2)\gamma_{i,T}}, \quad (23e)$$

where the inequality (23b) follows by Lem. 7.2, the inequality (23c) follows by the monotonicity of  $\beta_{i,t}$ , and the inequality (23e) follows by Lem. 21. Plugging in the definition of  $\beta_{i,T}$  completes the proof.  $\square$

The best iterate convergence guarantee can be obtained, consequently given below.

**Theorem 7.2** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, and 4.6 hold. With probability at least  $1 - \delta$ , there exists  $\tilde{x}^T \in \{x^1, \dots, x^T\}$ , such that,*

$$f(\tilde{x}^T) - f(x^*) \leq \mathcal{O}\left(\frac{\sum_{i \in \mathcal{I}_B} A_i \gamma_{i,T}}{\sqrt{T}}\right).$$

*Proof* By Thm. 7.1, we get

$$R_T = \sum_{t=1}^T (f(x^t) - f(x^*)) \quad (24a)$$

$$\leq \mathcal{O} \left( \sum_{i \in \mathcal{I}_{\mathcal{B}}} A_i \gamma_{i,T} \sqrt{T} \right). \quad (24b)$$

Therefore,

$$\frac{R_T}{T} = \sum_{t=1}^T \frac{f(x^t) - f(x^*)}{T} \quad (25a)$$

$$\leq \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}} A_i \gamma_{i,T}}{\sqrt{T}} \right). \quad (25b)$$

Since  $f(x^t) - f(x^*) \geq 0$ , there exists  $\tilde{x}^T \in \{x^1, \dots, x^T\}$ , such that,

$$f(\tilde{x}^T) - f(x^*) \leq \frac{R_T}{T} \leq \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}} A_i \gamma_{i,T}}{\sqrt{T}} \right) \quad (26)$$

concludes the proof.  $\square$

If all the kernel functions corresponding to different black-box functions are of the same type, we can list the convergence rate as in the Tab. 2 by applying the bounds on maximum information gains from [20, 23].

**Table 2** Kernel-specific cumulative regret bounds and the convergence rate to the optimal solution. In the table,  $d = n + m$  and  $\nu$  represent the smoothness parameter of the Matérn kernel.

| Kernel                            | Maximum Information Gain  | Cumulative Regret  | Convergence Rate  |
|-----------------------------------|---|--|---|
| Linear                            | $\mathcal{O}(d \log T)$   | $\mathcal{O}(d \log T \sqrt{T})$   | $\mathcal{O}\left(\frac{d \log T}{\sqrt{T}}\right)$                                 |
| Squared Exponential               | $\mathcal{O}((\log T)^{d+1})$                                     | $\mathcal{O}((\log T)^{d+1} \sqrt{T})$                                   | $\mathcal{O}\left(\frac{(\log T)^{d+1}}{\sqrt{T}}\right)$                           |
| Matérn<br>( $\nu > \frac{d}{2}$ ) | $\mathcal{O}(T^{\frac{d}{2\nu+d}} \log^{\frac{2\nu}{2\nu+d}}(T))$ | $\mathcal{O}(T^{\frac{2\nu+3d}{4\nu+2d}} \log^{\frac{2\nu}{2\nu+d}}(T))$ | $\mathcal{O}\left(T^{-\frac{2\nu-d}{4\nu+2d}} \log^{\frac{2\nu}{2\nu+d}}(T)\right)$ |

## 8 Extension to the Constrained Case

We have given the results in the unconstrained case. Following the same idea as in [25], we can extend the Alg. 1 to the constrained case and derive bounds on the cumulative violations. We formulate our problem as in 5,

$$\min_{x \in \mathcal{X}} f(x), \quad (27a)$$

$$\text{subject to } g_k(x) \leq 0, k \in [K], \quad (27b)$$

where  $g_k, k \in [K]$  are also grey-box functions modeled in the same way as on the objective function  $f$ . We make the same regularity assumptions 4.2, 4.3, 4.4, 4.5 and 8.1 on  $g_k, k \in [K]$  as on the objective function  $f$ . To distinguish the grey-box modeling notations for different functions, we add a superscript of function. For example, we use  $z_i^{t,g_k}, i \in [m^{g_k}]$  to denote the  $i$ -th intermediate variable for the grey-box function  $g_k$  at step  $t$ . In this section, we first focus on the feasible case and introduce the feasibility assumption. We will later show an infeasibility detection scheme in Sec. 8.1.

**Assumption 8.1** *There exists  $\tilde{x} \in \mathcal{X}$ , such that  $g_k(\tilde{x}) \leq 0, \forall k \in [K]$ .*

To make the algorithmic description more compact, the following plausible set definition is introduced.

**Definition 8.1 (Plausible set of  $z$ )** Conditioned on the historical samples up to step  $t$ , the plausible set  $\mathcal{Z}_t^f(x)$  of  $z^f$  with input  $x$  is defined as the feasible set of the following feasibility problem,

$$\min_{z^f \in \mathbb{R}^{m^f}} 1 \quad (28a)$$

$$\text{subject to } l_{i,t}(s_i^f) \leq z_{i+1}^f \leq u_{i,t}(s_i^f), \forall i \in \mathcal{I}_{\mathcal{B}}^f, \quad (28b)$$

$$z_{i+1}^f = \phi_i^f(s_i^f), \forall i \in \mathcal{I}_{\mathcal{W}}^f, \quad (28c)$$

$$s_i^f = [x_1, \dots, x_n, z_1^f, \dots, z_i^f]^\top, \forall i \in \{0\} \cup [m^f - 1]. \quad (28d)$$

The plausible set corresponding to  $g_k$  can be defined in the same way. The

---

**Algorithm 2** Constrained Bayesian Optimization of Expensive Nested Grey Box Functions

---

- 1: **for**  $t \in [T]$  **do**
- 2: Find  $(x^t, \bar{z}^{t,f}, (\bar{z}^{t,g_k})_{k \in [K]})$  by solving

$$\min_{x \in \mathcal{X}, z^f \in \mathbb{R}^{m^f}, z^{g_k} \in \mathbb{R}^{m^{g_k}}, k \in [K]} z_{m^f}^f \quad (29a)$$

$$\text{subject to } z^f \in \mathcal{Z}_t^f(x), \quad (29b)$$

$$z^{g_k} \in \mathcal{Z}_t^{g_k}(x), \forall k \in [K] \quad (29c)$$

$$z_{m^{g_k}}^{g_k} \leq 0, \forall k \in [K] \quad (29d)$$

- 3: Evaluate the grey-box function  $f$  and  $g_k, k \in [K]$  at  $x^t$ , incur objective  $f(x^t)$ , and constraints  $g_k(x^t), k \in [K]$  with the input  $x^t$ .
  - 4: Separately evaluate the black-box function  $\phi_i^f, i \in \mathcal{I}_{\mathcal{B}}^f$  at the point  $\bar{s}_i^{t,f} := [x_1^t, \dots, x_n^t, \bar{z}_1^{t,f}, \dots, \bar{z}_i^{t,f}]^\top$ . Also separately evaluate the black-box function  $\phi_i^{g_k}, i \in \mathcal{I}_{\mathcal{B}}^{g_k}$  at the point  $\bar{s}_i^{t,g_k} := [x_1^t, \dots, x_n^t, \bar{z}_1^{t,g_k}, \dots, \bar{z}_i^{t,g_k}]^\top$ .
  - 5: Update Gaussian process posterior mean and covariance for each black-box function with the new evaluations added.
  - 6: **end for**
- 

observation 8.1 guarantees the feasibility of the auxiliary problem in Alg. 2.

**Observation 8.1** *Since  $x^*$  is a feasible point for the original constrained optimization problem, then with high probability,  $x^*$  with the corresponding  $z^{*,f}$  and  $z^{*,g^k}$ ,  $k \in [K]$  is a feasible solution for the auxiliary optimization problem in Alg. 2.*

For the constrained problem, violations need to be taken care of in addition to the cumulative regret [25]. We formally define cumulative violations up to step  $T$  as,

**Definition 8.2** Cumulative violation up to step  $T$  is defined as,

$$\mathcal{V}_{k,T} := \sum_{t=1}^T [g_k(x_t)]^+. \quad (30)$$

We then have a theorem to bound both the cumulative regret and the cumulative violations.

**Theorem 8.2** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, 4.6 and 8.1 hold. We have, with probability at least  $1 - \delta$ , the sample points of Algorithm 2 satisfy,*

$$\begin{aligned} R_T &\leq \sum_{t=1}^T [f(x^t) - f(x^*)]^+ \\ &\leq \sum_{i \in \mathcal{I}_B^f} 2A_i^f \beta_{i,T}^f \sqrt{(T+2)\gamma_{i,T}^f} = \mathcal{O} \left( \sum_{i \in \mathcal{I}_B^f} A_i^f \gamma_{i,T}^f \sqrt{T} \right), \end{aligned} \quad (31a)$$

$$\mathcal{V}_{k,T} \leq \sum_{i \in \mathcal{I}_B^{g^k}} 2A_i^{g^k} \beta_{i,T}^{g^k} \sqrt{(T+2)\gamma_{i,T}^{g^k}} \quad (31b)$$

$$= \mathcal{O} \left( \sum_{i \in \mathcal{I}_B^{g^k}} A_i^{g^k} \gamma_{i,T}^{g^k} \sqrt{T} \right), \forall k \in [K] \quad (31c)$$

where  $A_i^f$  and  $A_i^{g^k}$  are Lipschitz constant dependent constants.

*Proof* We have

$$R_T = \sum_{t=1}^T (f(x^t) - f(x^*)) \quad (32a)$$

$$\leq \sum_{t=1}^T [f(x^t) - f(x^*)]^+ \quad (32b)$$

$$\leq \sum_{t=1}^T \sum_{i \in \mathcal{I}_B^f} A_i^f \beta_{i,t}^f \sigma_{i,t-1}(\bar{s}_i^{t,f}) \quad (32c)$$

$$\leq \sum_{i \in \mathcal{I}_B^f} A_i^f \beta_{i,T}^f \sum_{t=1}^T \sigma_{i,t-1}(\bar{s}_i^{t,f}) \quad (32d)$$

$$\leq \sum_{i \in \mathcal{I}_B^f} A_i^f \beta_{i,T}^f \sqrt{4(T+2)\gamma_{i,T}^f}, \quad (32e)$$

where the inequality (32c) follows by the Lem. 7.1 since  $(x^*, z^{*,f}, (z^{*,g_k})_{k \in [K]})$  is a feasible solution for the constrained auxiliary optimization problem in line 2 of the Alg. 2.

Similarly, we have

$$\mathcal{V}_{k,T} = \sum_{t=1}^T [g_k(x_t)]^+ \quad (33a)$$

$$\leq \sum_{t=1}^T [g_k(x^t) - \bar{z}_{m^{g_k}}^{t,g_k}]^+ + [\bar{z}_{m^{g_k}}^{t,g_k}]^+ \quad (33b)$$

$$= \sum_{t=1}^T [g_k(x^t) - \bar{z}_{m^{g_k}}^{t,g_k}]^+ \quad (33c)$$

$$\leq \sum_{i \in \mathcal{I}_B^{g_k}} A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{4(T+2)\gamma_{i,T}^{g_k}}, \quad (33d)$$

where the equality (33c) follows by the feasibility of  $\bar{z}^{t,g_k}$  for the auxiliary constrained optimization problem and the inequality (33d) follows by the discrepancy bound in Lem. 7.1.  $\square$

Consequently, we can derive a convergence rate to the constrained global optimum.

**Theorem 8.3** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, 4.6 and 8.1 hold. We have, with probability at least  $1 - \delta$ , there exists  $\tilde{x}_T$  from the sample points of Algorithm 2 satisfying,*

$$f(\tilde{x}_T) - f(x^*) \leq \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_B^f} A_i^f \gamma_{i,T}^f + \sum_{k \in [K]} \sum_{i \in \mathcal{I}_B^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} \right), \quad (34a)$$

$$[g_k(\tilde{x}_T)]^+ \leq \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_B^f} A_i^f \gamma_{i,T}^f + \sum_{k \in [K]} \sum_{i \in \mathcal{I}_B^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} \right), \quad \forall k \in [K] \quad (34b)$$

where  $A_i^f$  and  $A_i^{g_k}$  are constants dependent on the Lipschitz constants.

*Proof* From theorem 8.2, we can derive,

$$\sum_{t=1}^T [f(x^t) - f(x^*)]^+ \leq \sum_{i \in \mathcal{I}_B^f} 2A_i^f \beta_{i,T}^f \sqrt{(T+2)\gamma_{i,T}^f}, \quad (35a)$$

$$\sum_{t=1}^T [g_k(x^t)]^+ \leq \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} 2A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{(T+2)\gamma_{i,T}^{g_k}}, \forall k \in [K]. \quad (35b)$$

Summing up the above inequalities and dividing by  $T$ , we get,

$$\frac{1}{T} \sum_{t=1}^T \left( [f(x^t) - f(x^*)]^+ + \sum_{k=1}^K [g_k(x^t)]^+ \right) \quad (36a)$$

$$\leq \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^f} 2A_i^f \beta_{i,T}^f \sqrt{(T+2)\gamma_{i,T}^f} + \sum_{k=1}^K \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} 2A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{(T+2)\gamma_{i,T}^{g_k}}}{T} \quad (36b)$$

$$= \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^f} A_i^f \gamma_{i,T}^f + \sum_{k \in [K]} \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} \right). \quad (36c)$$

Since  $[f(x^t) - f(x^*)]^+ + \sum_{k=1}^K [g_k(x^t)]^+ \geq 0$ , there exists  $\tilde{x}^T \in \{x^1, \dots, x^T\}$ , such that,

$$[f(\tilde{x}^T) - f(x^*)]^+ + \sum_{k=1}^K [g_k(\tilde{x}^T)]^+ \quad (37a)$$

$$\leq \frac{1}{T} \sum_{t=1}^T \left( [f(x^t) - f(x^*)]^+ + \sum_{k=1}^K [g_k(x^t)]^+ \right) \quad (37b)$$

$$= \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^f} A_i^f \gamma_{i,T}^f + \sum_{k \in [K]} \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} \right). \quad (37c)$$

The desired results then follow.  $\square$

## 8.1 Infeasibility Detection

The cumulative regret/violation bounds and convergence results are derived under the assumption that the problem is feasible. However, in practice, the grey-box problem may indeed be infeasible and therefore, an infeasibility detection scheme is highly desired. In the following, we show that if the original problem is infeasible, we will find that the auxiliary problem in line 2 of Alg. 2 infeasible within finite steps. That is to say, we are able to detect the infeasibility of the original problem through the detection of the infeasibility of the optimistic auxiliary problem.

For infeasibility detection, we try to avoid ‘false positives’, where a feasible problem is declared as infeasible, and ‘false negatives’, where an infeasible problem is not declared to be infeasible. The observation 8.1 shows that it is impossible to have ‘false positives’. The following theorem complements the observation 8.1 by showing that if the original problem is infeasible, our



algorithm is guaranteed to declare infeasibility within finite steps with high probability.

**Theorem 8.4** *Let assumptions 4.1, 4.2, 4.3, 4.4, 4.5, and 4.6 hold. Assume that Prob. (5) is infeasible, that is, there exists  $k \in [K]$ , such that*

$$\min_{x \in \mathcal{X}} g_k(x) = \epsilon > 0,$$

and  $\lim_{T \rightarrow \infty} \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} = 0$ . Given a desired confidence level  $\delta \in (0, 1)$ , the auxiliary problem in line 2 of the Alg. 2 will be found infeasible within a number of steps equal to

$$\bar{T} = \min_{T \in \mathbb{N}_+} \left\{ T \left| \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} \leq C\epsilon \right. \right\}, \quad (38)$$

with probability at least  $1 - \delta$ , where  $C$  is a constant independent of  $T$ .

*Proof* Suppose up to step  $T$ , the optimization problem in line 2 of Alg. 2 keeps feasible. We have,

$$\bar{z}_{m^{g_k}}^{t,g_k} \leq 0$$

and the corresponding real output  $z_{m^{g_k}}^{t,g_k}$  corresponding to input  $x_t$  satisfies

$$z_{m^{g_k}}^{t,g_k} = g_k(x_t) \geq \epsilon$$

with probability at least  $1 - \delta$ .

Accordingly, we have

$$\sum_{t=1}^T \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,t}^{g_k} \sigma_{i,t-1}(s_i) \geq \sum_{t=1}^T |g_k(x_t) - \bar{z}_{m^{g_k}}^{t,g_k}| \geq \sum_{t=1}^T \epsilon = T\epsilon,$$

where the first inequality follows by the output discrepancy bound in Lem. 7.1. Meanwhile,

$$\begin{aligned} \sum_{t=1}^T \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,t}^{g_k} \sigma_{i,t-1}(\bar{s}_i) &\leq \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,T}^{g_k} \sum_{t=1}^T \sigma_{i,t-1}(\bar{s}_i) \\ &\leq \sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{4(T+2)\gamma_{i,T}^{g_k}}, \end{aligned}$$

where the last inequality follows by Lem. 21. Therefore,

$$\sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{4(T+2)\gamma_{i,T}^{g_k}} \geq T\epsilon,$$

which implies  $\epsilon \leq \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{4(T+2)\gamma_{i,T}^{g_k}}}{T}$ . By Def. 5.1, we have

$$\frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \beta_{i,T}^{g_k} \sqrt{4(T+2)\gamma_{i,T}^{g_k}}}{T} = \mathcal{O} \left( \frac{\sum_{i \in \mathcal{I}_{\mathcal{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} \right).$$

Hence,  $\exists \tilde{C} > 0$ , such that,

$$\epsilon \leq \tilde{C} \frac{\sum_{i \in \mathcal{I}_{\mathbf{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}}. \quad (39)$$

That is

$$C\epsilon \leq \frac{\sum_{i \in \mathcal{I}_{\mathbf{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}}, \quad (40)$$

where  $C = \frac{1}{\tilde{C}}$ . However, since  $\lim_{T \rightarrow \infty} \frac{\sum_{i \in \mathcal{I}_{\mathbf{B}}^{g_k}} A_i^{g_k} \gamma_{i,T}^{g_k}}{\sqrt{T}} = 0$ , the inequality (40) will be violated when  $T$  is large enough. So the optimistic auxiliary problem will be found infeasible on or before the first time the above inequality is violated, which is  $\bar{T}$ .

*Remark 8.1* The proposed algorithm design and analysis in this paper focuses on the case where  $T$  is given. In cases where  $T$  is not given or not large enough to declare infeasibility, we can apply the ‘doubling trick’ [6]. The basic idea is that we can start running a round of steps with  $T = 1$  and double  $T$  after each round. As  $T$  grows larger and larger, it will finally be larger than  $\bar{T}$  and enough for infeasibility detection.

## 9 Discussion on Special Cases

### 9.1 Grey-box function with only one layer of black-box functions

We now restrict to the special case with only one layer of black-box functions. That is, the problem we consider can be formulated as,

$$\min_{x \in \mathcal{X}} F(x, \phi(x)), \quad (41a)$$

$$\text{subject to } G_k(x, \phi(x)) \leq 0, \quad \forall k \in [K], \quad (41b)$$

where  $\mathcal{X} \subset \mathbb{R}^n$  is a known compact subset of  $\mathbb{R}^n$ ,  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^{m-1}$ ,  $F : \mathbb{R}^n \times \mathbb{R}^{m-1} \rightarrow \mathbb{R}$  and  $G_k : \mathbb{R}^n \times \mathbb{R}^{m-1} \rightarrow \mathbb{R}$ ,  $k \in [K]$ . We assume that the functions  $F$  and  $G_k, k \in [K]$  are known and cheap to evaluate, and  $\phi$  is a potentially vector-valued black box function that is expensive to evaluate. When restricted to this special case, our algorithm 2 essentially recovers the algorithm proposed in [16].

## 10 Experiments

### 10.1 Linear Programming with Embedded Gaussian Process

Linear programming (LP) is a ubiquitous problem in engineering, operations research, and economics, etc. However, in many applications, LP problems

may involve some unknown black-box functions other than the known linear part. In this part, we consider a generic linear programming problem with unknown expensive black-box functions embedded as shown in (42).

$$\min_{x \in \mathcal{X}} \quad c_1^T x + c_2^T h(x), \quad (42a)$$

$$\text{subject to} \quad A_1 x + A_2 h(x) + b \leq 0. \quad (42b)$$

In this experiment, we consider  $x \in [-2, 2]^2 \subset \mathbb{R}^2$  and  $h : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ .

The corresponding auxiliary problem in the line 2 of the Alg. 2 is reduced to,

$$\min_{x \in \mathcal{X}} \min_{z \in [l_t(x), u_t(x)]} \quad c_1^T x + c_2^T z, \quad (43a)$$

$$\text{subject to} \quad A_1 x + A_2 z + b \leq 0. \quad (43b)$$

To compare the performance of different algorithms in terms of the speed of finding the global optimum, we also introduce the metric of constrained regret defined as,

$$\text{CR}_t := \min_{\tau \in [t]} [f(x_\tau) - f(x^*)]^+ + \|[g(x_\tau)]^+\|_1, \quad (44)$$


where  $x^*$  is the ground truth optimal solution. To evaluate the performance of different algorithms, we randomly sample  $c_1, c_2, b$  and the rows of  $A_1, A_2$  uniformly from the sphere  $\mathbb{S}^1$ . And we sample the function  $h$  from a Gaussian process with kernel function,

$$k(x, y) = \frac{1}{2} \exp\{-(x_1 - y_1)^2 - (x_2 - y_2)^2\}. \quad (45)$$


We compare our grey-box optimization method to the constrained black-box optimization method [25]. Fig. 2 to Fig. 5 show the evolutions of constrained regret, cumulative positive regret, cumulative violations for constraint 1 and constraint 2. It can be observed that our grey-box optimization method significantly outperforms the pure black-box optimization method in terms of all the above metrics.

## 11 Conclusion


In this paper, we have formulated a general *grey-box* optimization problem, i.e., with both objective and constraint functions composed of both *black-box* and *white-box* functions. Our formulation covers the existing grey-box optimization formulations as special cases. An *optimism*-driven algorithm is proposed to efficiently solve it. Under certain regularity assumptions, our algorithm achieves similar regret bound as that for the standard black-box Bayesian optimization algorithm, up to a constant term depending on the Lipschitz constants of the functions considered. The method is then extended to the constrained case and special cases are discussed. For the commonly used kernel functions,




**Fig. 2** The evolution of constrained regret with respect to sample step.



**Fig. 3** The evolution of cumulative positive regret with respect to sample step.



**Fig. 4** The evolution of the cumulative violation for the first constraint with respect to sample step.



**Fig. 5** The evolution of cumulative violation for the second constraint with respect to sample step.

kernel-specific convergence rate to the optimal solution is derived. Experimental results show that our grey-box optimization method empirically improves the speed of finding the global optimum solution significantly, as compared to the standard black-box optimization algorithm.

## References

1. Virginia Aglietti, Neil Dhir, Javier González, and Theodoros Damoulas. Dynamic causal Bayesian optimization. *Advances in Neural Information Processing Systems*, 34:10549–10560, 2021.
2. Virginia Aglietti, Xiaoyu Lu, Andrei Paleyes, and Javier González. Causal Bayesian optimization. In *International Conference on Artificial Intelligence and Statistics*, pages 3155–3164. PMLR, 2020.
3. Virginia Aglietti, Alan Malek, Ira Ktena, and Silvia Chiappa. Constrained causal Bayesian optimization. *arXiv preprint arXiv:2305.20011*, 2023.
4. Raul Astudillo and Peter Frazier. Bayesian optimization of composite functions. In *International Conference on Machine Learning*, pages 354–363. PMLR, 2019.
5. Raul Astudillo and Peter Frazier. Bayesian optimization of function networks. *Advances in neural information processing systems*, 34:14463–14475, 2021.
6. Lilian Besson and Emilie Kaufmann. What doubling tricks can and can’t do for multi-armed bandits. *arXiv preprint arXiv:1803.06971*, 2018.
7. Sait Cakmak, Raul Astudillo Marban, Peter Frazier, and Enlu Zhou. Bayesian optimization of risk measures. *Advances in Neural Information Processing Systems*, 33:20130–20141, 2020.
8. Sayak Ray Chowdhury and Aditya Gopalan. On kernelized multi-armed bandits. In *International Conference on Machine Learning*, pages 844–853. PMLR, 2017.
9. Sayak Ray Chowdhury and Aditya Gopalan. On kernelized multi-armed bandits. *arXiv preprint arXiv:1704.00445*, 2017.
10. Ehecatl Antonio del Rio Chanona, Panagiotis Petsagkourakis, Eric Bradford, JE Alves Graciano, and Benoît Chachuat. Real-time optimization meets Bayesian optimization and derivative-free optimization: A tale of modifier adaptation. *Comput. & Chem. Eng.*, 147:107249, 2021.
11. Peter I Frazier. A tutorial on Bayesian optimization. *arXiv preprint arXiv:1807.02811*, 2018.

12. Jacob R Gardner, Matt J Kusner, Zhixiang Eddie Xu, Kilian Q Weinberger, and John P Cunningham. Bayesian optimization with inequality constraints. In *ICML*, volume 2014, pages 937–945, 2014.
13. Michael A. Gelbart, Jasper Snoek, and Ryan P. Adams. Bayesian optimization with unknown constraints. In *Proc. of the 30th Conf. on Uncertainty in Artif. Intell.*, UAI’14, page 250–259, Arlington, Virginia, USA, 2014. AUAI Press.
14. Donald R Jones, Matthias Schonlau, and William J Welch. Efficient global optimization of expensive black-box functions. *Journal of Global Optimization*, 13(4):455–492, 1998.
15. Kirthevasan Kandasamy, Gautam Dasarathy, Jeff Schneider, and Barnabás Póczos. Multi-fidelity Bayesian optimisation with continuous approximations. In *International Conference on Machine Learning*, pages 1799–1808. PMLR, 2017.
16. Congwen Lu and Joel A Paulson. No-regret constrained Bayesian optimization of noisy and expensive hybrid models using differentiable quantile function approximations. *arXiv preprint arXiv:2305.03824*, 2023.
17. Jonas Moćkus. On Bayesian methods for seeking the extremum. In *Optimization Techniques IFIP Technical Conference: Novosibirsk, July 1–7, 1974*, pages 400–404. Springer, 1975.
18. Bernhard Schölkopf, Ralf Herbrich, and Alex J Smola. A generalized representer theorem. In *International Conference on Computational Learning Theory*, pages 416–426. Springer, 2001.
19. Jasper Snoek, Hugo Larochelle, and Ryan P Adams. Practical Bayesian optimization of machine learning algorithms. *Advances in Neural Inf. Process. Syst.*, 25, 2012.
20. Niranjan Srinivas, Andreas Krause, Sham M Kakade, and Matthias W Seeger. Information-theoretic regret bounds for Gaussian process optimization in the bandit setting. *IEEE Transactions on Information Theory*, 58(5):3250–3265, 2012.
21. Scott Sussex, Anastasiia Makarova, and Andreas Krause. Model-based causal Bayesian optimization. *arXiv preprint arXiv:2211.10257*, 2022.
22. Saul Toscano-Palmerin and Peter I Frazier. Bayesian optimization with expensive integrands. *arXiv preprint arXiv:1803.08661*, 2018.
23. Sattar Vakili, Kia Khezeli, and Victor Picheny. On information gain and regret bounds in Gaussian process bandits. In *International Conference on Artificial Intelligence and Statistics*, pages 82–90. PMLR, 2021.
24. Jian Wu, Saul Toscano-Palmerin, Peter I Frazier, and Andrew Gordon Wilson. Practical multi-fidelity Bayesian optimization for hyperparameter tuning. In *Uncertainty in Artificial Intelligence*, pages 788–798. PMLR, 2020.
25. Wenjie Xu, Yuning Jiang, and Colin N Jones. Constrained efficient global optimization of expensive black-box functions. *arXiv preprint arXiv:2211.00162*, 2022.
26. Wenjie Xu, Yuning Jiang, Emilio T Maddalena, and Colin N Jones. Lower bounds on the worst-case complexity of efficient global optimization. *arXiv preprint arXiv:2209.09655*, 2022.
27. Wenjie Xu, Yuning Jiang, Bratislav Svetozarevic, and Colin N Jones. CONFIG: Constrained efficient global optimization for closed-loop control system optimization with unmodeled constraints. *arXiv preprint arXiv:2211.11822*, 2022.
28. Wenjie Xu, Colin N Jones, Bratislav Svetozarevic, Christopher R Laughman, and Ankush Chakrabarty. VABO: Violation-Aware Bayesian Optimization for closed-loop control performance optimization with unmodeled constraints. *arXiv preprint arXiv:2110.07479*, 2021.
29. Wenjie Xu, Colin N Jones, Bratislav Svetozarevic, Christopher R Laughman, and Ankush Chakrabarty. Violation-aware contextual Bayesian optimization for controller performance optimization with unmodeled constraints. *arXiv preprint arXiv:2301.12099*, 2023.
30. AG Zhilinskias. Single-step Bayesian search method for an extremum of functions of a single variable. *Cybernetics*, 11(1):160–166, 1975.