RELATIONS AMONG HIGHER WHITEHEAD MAPS

JELENA GRBIĆ, GEORGE SIMMONS, MATTHEW STANIFORTH

9th June 2023

ABSTRACT. We define generalised higher Whitehead maps between polyhedral products. By investigating the interplay between the homotopy-theoretic properties of polyhedral products and the combinatorial properties of simplicial complexes, we describe new families of relations among these maps, while recovering and generalising known identities among Whitehead products.

Contents

1. Introduction	1
2. The higher Whitehead map	5
3. The nested higher Whitehead map	10
4. Folded higher Whitehead maps	13
5. Relations	18
5.1. Relations among higher Whitehead maps	21
6. Relations among folded higher Whitehead maps	24
6.1. Folds of identity complexes	24
6.2. Folding and substitution	29
7. Proof of main theorem	32
References	43

1. INTRODUCTION

In this paper we describe new relations among higher Whitehead maps in the group $[\Sigma X, W]$ of homotopy classes of maps from a suspension space ΣX to a topological space W. We obtain families of relations among *n*-ary higher Whitehead products for $n \ge 2$, recovering the Jacobi identity and Hardie's identity in homotopy groups as a special case.

Relations among Whitehead products in homotopy groups have been studied since the early 1950s. Given maps $f_i \in \pi_{d_i}(Y)$ for i = 1, 2 with $d_i \ge 1$, the Whitehead product $[f_1, f_2] \in \pi_{d_1+d_2-1}(Y)$ is the homotopy class of the map

$$[f_1, f_2]: S^{d_1-1} * S^{d_2-1} \xrightarrow{\rho} S^{d_1} \vee S^{d_2} \xrightarrow{f_1 \vee f_2} Y \vee Y \xrightarrow{\nabla} Y.$$

The Whitehead product equips the homotopy groups of Y with the structure of a graded quasi-Lie algebra. Motivated by the Jacobi identity for Lie algebras, Blakers

and Massey [5], and Samelson [12] independently conjectured the Jacobi identity as a relation among Whitehead products. Let $f_i \in \pi_{d_i}(Y)$ for i = 1, 2, 3, where $d_i \ge 2$. Then

(1)
$$(-1)^{d_1d_3}[f_1, [f_2, f_3]] + (-1)^{d_2d_3}[f_3, [f_1, f_2]] + (-1)^{d_1d_2}[f_2, [f_3, f_1]] = 0.$$

This relation was established by several authors in the mid 1950s, using a variety of methods. Notably, the proof of Massey and Uehara [13] is algebraic in nature, and is one of the first applications of the triple Massey product, while the proof of Nakaoka and Toda [10] uses a geometric approach considering relative homotopy theory.

Hardie [8] defined the exterior Whitehead product as an operation on homotopy groups generalising the Whitehead product. For $k \ge 2$, let $f_i \in \pi_{d_i}(Y_i)$ for $i = 1, \ldots, k$, where $d_i \ge 1$. The exterior Whitehead product $\{f_1, \ldots, f_k\} \in \pi_{d_1+\cdots+d_k-1}(FW(Y_1, \ldots, Y_k))$ is the homotopy class of the map

$$\{f_1,\ldots,f_k\}: \overset{k}{\underset{i=1}{\underbrace{\longrightarrow}}} S^{d_i-1} \xrightarrow{\rho} FW(S^{d_1},\ldots,S^{d_k}) \xrightarrow{FW(f_1,\ldots,f_k)} FW(Y_1,\ldots,Y_k)$$

where $FW(Y_1, \ldots, Y_k) = \bigcup_{i=1}^k Y_1 \times \ldots \times Y_{i-1} \times \ast \times Y_{i+1} \times \cdots \times Y_k$ is the fat wedge of the spaces Y_1, \ldots, Y_k and $FW(f_1, \ldots, f_k)$ is the map induced on the fat wedge by f_1, \ldots, f_k . For $i = 1, \ldots, k$, when $d_i \ge 2$, Hardie showed that exterior Whitehead products satisfy

(2)
$$\sum_{i=1}^{k} (-1)^{\eta(i)} \{ f_i, \{ f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_k \} \} = 0$$

in $\pi_{d_1+\dots+d_k-2}(Z)$, where $Z = \bigcup_{i=1}^k Y_i \vee FW(Y_1,\dots,Y_{i-1},Y_{i+1},\dots,Y_k)$.

Arkowitz [4] defined the generalised Whitehead product of non-spherical maps. Let $f_i \in [\Sigma X_i, Y]$ for i = 1, 2. The generalised Whitehead product $[f_1, f_2] \in [\Sigma X_1 \wedge X_2, Y]$ is the homotopy class of the composite

$$[f_1, f_2] \colon \Sigma X_1 \land X_2 \xrightarrow{\rho} \Sigma X_1 \lor \Sigma X_2 \xrightarrow{f_1 \lor f_2} Y \lor Y \xrightarrow{\nabla} Y.$$

Little is known about the existence of relations among non-spherical higher Whitehead maps. The only known relation is due to Arkowitz [4] who established the Jacobi identity (1) for generalised Whitehead products. This motivates us to ask whether analogues of Hardie's identity exist in the group of homotopy classes of maps $[\Sigma X, W]$ from a suspension ΣX to an arbitrary space W.

In this paper, we consider generalisations of relation (2) in the context of polyhedral products. A polyhedral product $(\underline{X}, \underline{A})^{\mathcal{K}}$ is a topological space built out of a simplicial complex \mathcal{K} on m vertices and an m-tuple of topological pairs $(\underline{X}, \underline{A})$, see Definition 2.1. It can be considered as a functor from topological pairs and simplicial complexes with respect to continuous maps of topological pairs and simplicial inclusions. Homotopy-theoretic properties of polyhedral products tend to be encoded by underlying combinatorial structures. By combining geometric and combinatorial

techniques, we generalise exterior Whitehead products to homotopy classes of maps between polyhedral products and study relations among them.

We begin by describing the exterior Whitehead product as the map of polyhedral products

$$\{f_1,\ldots,f_m\}: (\underline{D},\underline{S})^{\partial\Delta^{m-1}} \longrightarrow (\underline{Y},\underline{*})^{\partial\Delta^{m-1}}$$

induced by the maps of pairs $(D^{d_i}, S^{d_i-1}) \longrightarrow (Y_i, *)$ representing $f_i \in \pi_{d_i}(Y_i)$. Recognising that the boundary $\partial \Delta^{m-1}$ of a simplex governs the existence of the exterior Whitehead product, we define the higher Whitehead map $h_w(f_1, \ldots, f_m) \in [(\underline{CX}, \underline{X})^{\partial \Delta^{m-1}}, (\underline{Y}, *)^{\partial \Delta^{m-1}}]$ associated to maps of pairs $f_i: (CX_i, X_i) \longrightarrow (Y_i, *), i = 1, \ldots, m$, to be the map of polyhedral products

$$h_w(f_1,\ldots,f_m): (\underline{CX},\underline{X})^{\partial \Delta^{m-1}} \longrightarrow (\underline{Y},\underline{*})^{\partial \Delta^{m-1}}$$

induced by the maps f_i .

Higher Whitehead maps of spherical maps have been studied by Abramyan and Panov [2], and Zhuravleva [15], using algebraic techniques such as the Hurewicz map and Adams-Hilton models. In the non-spherical case, these algebraic techniques are not defined. We study higher Whitehead maps $h_w(f_1, \ldots, f_m)$ and relations between them geometrically by generalising the relative homotopy group approach of Nakaoka and Toda [10].

We generalise Hardie's identity to non-spherical maps. By using polyhedral products, we rewrite the maps in relation (2) in terms of the higher Whitehead map as

$$h_w(h_w(f_1,\ldots,f_{i-1},f_{i+1},\ldots,f_m),f_i):S^{d_1+\cdots+d_m-2}\longrightarrow (\underline{Y},\underline{*})^{\operatorname{sk}^{m-3}\Delta^{m-1}}$$

and say that the skeleton sk^{m-3} Δ^{m-1} is the simplicial complex which carries Hardie's identity. By considering non-spherical maps $f_i: \Sigma X_i \longrightarrow Y_i$, we study relations among higher Whitehead maps of the form

$$h_w(h_w(f_1,\ldots,f_{i-1},f_{i+1},\ldots,f_m),f_i):\Sigma^{m-2}X_1\wedge\cdots\wedge X_m\longrightarrow (\underline{Y},\underline{*})^{\mathrm{sk}^{m-3}\Delta^{m-1}}$$

We prove, as Theorem 5.6, a generalisation of relation (2) to these non-spherical higher Whitehead maps by analysing the homotopy-theoretic properties of the polyhedral product in terms of the underlying combinatorics, and developing the notion of relative higher Whitehead maps as an analogue to the relative Whitehead product.

Hardie's identity is the only known relation involving higher Whitehead maps in homotopy groups. The summands are exterior Whitehead products of a higher Whitehead map and an arbitrary map. We further generalise relation (2) to more general relations between higher Whitehead maps. We notice that the skeleton $\mathrm{sk}^{m-3} \Delta^{m-1}$ is the composition complex $\mathrm{sk}^{m-3} \Delta^{m-1}(\emptyset, \ldots, \emptyset)$, see Definition 3.2, and the vertex set $[m] = \{1, \ldots, m\}$ can be viewed as an *m*-partition of [m] into singletons. To obtain new relations, we consider an *m*-partition $\Pi = \{P_1, \ldots, P_m\}$ of a vertex set and replace the skeleton $\mathrm{sk}^{m-3} \Delta^{m-1}$ by changing $\emptyset = \partial \Delta^0$ to $\partial \Delta[P_i]$ in $\mathrm{sk}^{m-3} \Delta^{m-1}(\emptyset, \ldots, \emptyset)$. We define the identity complex \mathcal{K}_{Π} as the resulting complex $\mathrm{sk}^{m-3} \Delta^{m-1}(\partial \Delta[P_1], \ldots, \partial \Delta[P_m])$. The main result of this paper, given as Theorem 5.6, is that for every partition Π and associated identity complex \mathcal{K}_{Π} , there is a relation among higher Whitehead maps. Suppose that each X_i is a suspension space. Then

(3)
$$\sum_{i=1}^{\kappa} h_w^{\mathcal{K}_{\Pi}} \left(h_w \left(f_{j_1}, \dots, f_{j_{q_i}} \right), f_{i_1}, \dots, f_{i_{p_i}} \right) \circ \sigma_i = 0$$

in $[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}]$, where $P_i = \{i_1, \ldots, i_{p_i}\}$ and $\bigcup_{j=1}^m P_j \setminus P_i = \{j_1, \ldots, j_{q_i}\}$

 $\sigma_i: \Sigma^{m-2} X_1 \wedge \dots \wedge X_m \longrightarrow \Sigma^{p_i} (\Sigma^{q_i-2} X_{j_1} \wedge \dots \wedge X_{j_{q_i}}) \wedge X_{i_1} \wedge \dots \wedge X_{i_{p_i}}$

is the restriction of the coordinate permutation

$$CX_1 \times \cdots \times CX_m \longrightarrow CX_{j_1} \times \cdots \times CX_{j_{q_i}} \times CX_{i_1} \times \cdots \times CX_{i_{p_i}}.$$

All relations among higher Whitehead maps are governed by the shape of the identity complex \mathcal{K}_{Π} , and any simplicial complex \mathcal{K} that supports a relation has an identity complex at its core. We want to extend the family of simplicial complexes \mathcal{K} for which the polyhedral product $(\underline{Y}, \underline{*})^{\mathcal{K}}$ supports a relation. To do so, we propagate the structure of the identity complex \mathcal{K}_{Π} to more complex combinatorial objects by using the operation of simplicial substitution. For a simplicial complex \mathcal{K} on [m] and simplicial complexes $\mathcal{S}_1, \ldots, \mathcal{S}_m$, we denote the substitution complex of $\mathcal{S}_1, \ldots, \mathcal{S}_m$ into \mathcal{K} by $\mathcal{K}\langle \mathcal{S}_1, \ldots, \mathcal{S}_m \rangle$, see Definition 3.2. In Corollary 5.10, we propagate identity complexes to $\mathcal{K}_{\Pi}\langle \mathcal{S}_1, \ldots, \mathcal{S}_m \rangle$ which support new relations.

In homotopy groups, the Lie bracket given by the Whitehead product can have repeated factors; for example $[[f_1, f_2], f_1]$, in which f_1 is repeated. Up to now, a higher Whitehead map captured by a simplicial complex has all factors different as they correspond to different vertices of the simplicial complex. This motivates the question of whether higher Whitehead maps with repeated factors can also be encoded by a simplicial complex.

We introduce the combinatorial operation of folding. A fold of a simplicial complex is the image of the simplicial map induced by identifying two or more vertices. Starting with a relation with no repeated factors corresponding to the identity complex \mathcal{K}_{Π} , we fold \mathcal{K}_{Π} to a new simplicial complex which supports relations among higher Whitehead maps with repeated factors corresponding to the identified vertices under the fold.

We increase the family of simplicial complexes which detect relations among higher Whitehead maps with repeated factors by considering folds of $\mathcal{K}_{\Pi}\langle \mathcal{S}_1, \ldots, \mathcal{S}_m \rangle$ induced by folds of the simplicial complexes \mathcal{S}_i .

An L_{∞} -algebra, also known as a homotopy Lie algebra, is a higher generalisation of a Lie algebra, in which higher, multilinear brackets satisfy a strong homotopy Jacobi identity. In polyhedral products, we realise higher L_{∞} brackets as higher Whitehead maps, and using identity complexes and their folds, we realise the strong homotopy Jacobi identity. In the course of the paper we also study various homotopy-theoretical properties of higher Whitehead maps such as conditions under which they are null-homotopic. An application of folding simplicial complexes is the construction of 2-torsion elements in the homotopy groups of a wide range of polyhedral products.

2. The higher Whitehead map

Given maps $f_i \in \pi_{d_i}(Y_i)$ for $i = 1, ..., k, k \ge 2, d_i \ge 1$, Hardie [8] defined the exterior Whitehead product $\{f_1, \ldots, f_k\} \in \pi_{d_1+\cdots+d_k-1}(FW(Y_1, \ldots, Y_k))$ as the homotopy class of the composite

(4)
$$\{f_1, \dots, f_k\}: \overset{k}{\underset{i=1}{\longrightarrow}} S^{d_i-1} \xrightarrow{\rho} FW(S^{d_1}, \dots, S^{d_k}) \xrightarrow{FW(f_1, \dots, f_k)} FW(Y_1, \dots, Y_k)$$

where

(5)
$$\rho: \underset{i=1}{\overset{k}{\bigstar}} S^{d_i-1} = \bigcup_{i=1}^{k} D^{d_1} \times \dots \times S^{d_i-1} \times \dots \times D^{d_k} \to FW(S^{d_1}, \dots, S^{d_k})$$

is the restriction of the product of the quotient maps sending $\partial D^{d_i} = S^{d_i-1}$ to a point.

By expressing the exterior Whitehead product as a map of polyhedral products, we will use the underlying combinatorics of polyhedral products to study the exterior Whitehead product, and its generalisation to non-spherical maps $f_i \in [\Sigma X_i, Y_i]$. We start by defining polyhedral products.

Definition 2.1. Let \mathcal{K} be a simplicial complex on the vertex set [m], and let $(\underline{X}, \underline{A}) = \{(X_i, A_i)\}_{i=1}^m$ be an *m*-tuple of *CW*-pairs. The *polyhedral product* $(\underline{X}, \underline{A})^{\mathcal{K}}$ is defined as

$$(\underline{X},\underline{A})^{\mathcal{K}} = \bigcup_{\sigma \in \mathcal{K}} (\underline{X},\underline{A})^{\sigma} \subseteq \prod_{i=1}^{m} X_{i}, \quad \text{where } (\underline{X},\underline{A})^{\sigma} = \prod_{i=1}^{m} Y_{i}, \quad Y_{i} = \begin{cases} X_{i} & \text{for } i \in \sigma \\ A_{i} & \text{for } i \notin \sigma. \end{cases}$$

The polyhedral product is a covariant functor with respect to both inclusions of simplicial complexes, and continuous maps of *m*-tuples of *CW*-pairs. A simplicial inclusion $\mathcal{L} \longrightarrow \mathcal{K}$ induces an inclusion of polyhedral products

$$(\underline{X},\underline{A})^{\mathcal{L}} \longrightarrow (\underline{X},\underline{A})^{\mathcal{K}}$$

and maps of CW-pairs $(f_i, g_i): (X_i, A_i) \longrightarrow (Y_i, B_i)$ induce a map of polyhedral products

$$(\underline{f},\underline{g})^{\mathcal{K}}: (\underline{X},\underline{A})^{\mathcal{K}} \longrightarrow (\underline{Y},\underline{B})^{\mathcal{K}}.$$

In the case that $A_i = B_i = *$ for all $i \in [m]$, we suppress notation and denote this induced map by $(\underline{f})^{\mathcal{K}}$.

The fat-wedge and the join arise as special cases of polyhedral products. If $A_i = *$ for i = 1, ..., m, then

$$(\underline{X},\underline{*})^{\partial\Delta^{m-1}} = \bigcup_{i=1}^{m} (X_1 \times \cdots \times X_{i-1} \times \ast \times X_{i+1} \times \cdots \times X_m) = FW(X_1,\ldots,X_m)$$

and if $X_i = CA_i$ for $i = 1, \ldots, m$, then

$$(\underline{CA},\underline{A})^{\partial\Delta^{m-1}} = \bigcup_{i=1}^{m} CA_1 \times \cdots \times CA_{i-1} \times A_i \times CA_{i+1} \times \cdots \times CA_m = \overset{m}{\underset{i=1}{\overset{m}{\overset{m}{\overset{}}}} A_i.$$

Polyhedral products are the colimits of diagrams over the face category of a simplicial complex containing the empty set. Whenever all maps in such a diagram are cofibrant inclusions, the colimit is homotopy equivalent to the homotopy colimit (see [7, Proposition 8.1.1]), and therefore

(6)
$$(\underline{X},\underline{A})^{\mathcal{K}} \simeq \operatorname{hocolim}_{\sigma \in \mathcal{K}} (\underline{X},\underline{A})^{\sigma}.$$

This enables us to study the homotopy-theoretic properties of polyhedral products combinatorially.

Using the notion of polyhedral products, we rewrite the exterior Whitehead product. Let $(\underline{D}, \underline{S}) = \{(D^{d_i}, S^{d_i-1})\}_{i=1}^k$ and $(\underline{Y}, \underline{*}) = \{(Y_i, \ast)\}_{i=1}^k$ be k-tuples of pairs and for $i = 1, \ldots, k$, identify $f_i: S^{d_i} \longrightarrow Y_i$ with the map of pairs $(D^{d_i}, S^{d_i-1}) \longrightarrow (Y_i, \ast)$. Then the exterior Whitehead product (4) is the map of polyhedral products

(7)
$$\{f_1, \dots, f_k\}: (\underline{D}, \underline{S})^{\partial \Delta^{k-1}} \longrightarrow (\underline{Y}, \underline{*})^{\partial \Delta^{k-1}}$$

induced by the maps of pairs $(D^{d_i}, S^{d_i-1}) \longrightarrow (Y_i, *)$ for $i = 1, \ldots, k$. We generalise this construction to non-spherical maps $\Sigma X_i \longrightarrow Y_i$, which can be identified with maps of pairs $f_i: (CX_i, X_i) \longrightarrow (Y_i, *)$.

Definition 2.2. Let $m \ge 2$, and let $f_i: (CX_i, X_i) \longrightarrow (Y_i, *)$ be maps of pairs for $i = 1, \ldots, m$. Let $(\underline{CX}, \underline{X}) = \{(CX, X)\}_{i=1}^m$ and $(\underline{Y}, \underline{*}) = \{(Y_i, *)\}_{i=1}^m$ be *m*-tuples of pairs. The higher Whitehead map $h_w(f_1, \ldots, f_m)$ of the maps f_1, \ldots, f_m is the map of polyhedral products

(8)
$$h_w(f_1,\ldots,f_m): (\underline{CX},\underline{X})^{\partial \Delta^{m-1}} \longrightarrow (\underline{Y},\underline{*})^{\partial \Delta^{m-1}}$$

induced by the maps f_i .

Since the polyhedral product is a homotopy colimit (see (6)), the homotopy class of $h_w(f_1, \ldots, f_m)$ depends only on the homotopy classes of the maps f_1, \ldots, f_m .

We prove multilinearity, naturality and symmetry properties of the higher Whitehead map by expressing it as the composite

(9)
$$(\underline{CX},\underline{X})^{\partial\Delta^{m-1}} \xrightarrow{(\underline{\rho})^{\partial\Delta^{m-1}}} (\underline{\SigmaX},\underline{*})^{\partial\Delta^{m-1}} \xrightarrow{(\underline{f})^{\partial\Delta^{m-1}}} (\underline{Y},\underline{*})^{\partial\Delta^{m-1}}$$

Proposition 2.3. Let $f_i: (CX_i, X_i) \longrightarrow (Y_i, *)$ be maps of pairs for i = 1, ..., m, and for some $j \in \{1, ..., m\}$, let $f'_j: (CX_j, X_j) \longrightarrow (Y_j, *)$. If X_j is a suspension,

(10)
$$h_w(f_1, \ldots, f_j + f'_j, \ldots, f_m) = h_w(f_1, \ldots, f_j, \ldots, f_m) + h_w(f_1, \ldots, f'_j, \ldots, f_m)$$

Proof. Since X_j is a suspension, we decompose

$$X_j = \Sigma \widetilde{X}_j = C_+ \widetilde{X}_j \cup C_- \widetilde{X}_j$$

where $C_+\widetilde{X}_j$ and $C_-\widetilde{X}_j$ are two cones on \widetilde{X}_j . Let C_1X_j and C_2X_j be two cones on X_j and let $C_+^i\widetilde{X}_j \cup C_-^i\widetilde{X}_j$ be the copy of X_j inside C_iX_j for $i \in \{1, 2\}$. Define $D = C_1X_j \cup C_2X_j$ with the union taken by identifying $C_+^1\widetilde{X}_j = C_+^2\widetilde{X}_j$, and define $D' = C_-^1\widetilde{X}_j \cup C_-^2\widetilde{X}_j$ with the union taken by identifying the copies of \widetilde{X}_j . The sum $f_j + f'_j$ is represented by the map of pairs

$$(CX_j, X_j) \simeq (D, D') = (C_1 X_j \cup C_2 X_j, C_-^1 \widetilde{X}_j \cup C_-^2 \widetilde{X}_j) \longrightarrow (Y_j, *)$$

determined by f_j on C_1X_j and f'_j on C_2X_j . Then $h_w(f_1, \ldots, f_j + f'_j, \ldots, f_m)$ is the map of polyhedral products

$$(\underline{CX},\underline{X})^{\partial\Delta} \simeq (\underline{CX}',\underline{X}')^{\partial\Delta} \longrightarrow (\underline{Y},\underline{*})^{\partial\Delta}$$

induced by $f_1, ..., f_j + f'_j, ..., f_m$, where $(CX'_i, X'_i) = (CX_i, X_i)$ for $i \neq j$ and $(CX'_j, X'_j) = (D, D')$.

On the other hand, let $Z = CX_1 \times \cdots \times C_+^1 \widetilde{X}_j \times \cdots \times CX_m$. Then we decompose

$$(\underline{CX}',\underline{X}')^{\partial\Delta} = (\underline{CX},\underline{X})^{\partial\Delta} \cup_Z (\underline{CX},\underline{X})^{\partial\Delta} \simeq (\underline{CX},\underline{X})^{\partial\Delta} \vee (\underline{CX},\underline{X})^{\partial\Delta}$$

since Z is contractible. Therefore contracting Z gives a homotopy between $h_w(f_1, \ldots, f_j + f'_j, \ldots, f_m)$ and the sum $h_w(f_1, \ldots, f_j, \ldots, f_m) + h_w(f_1, \ldots, f'_j, \ldots, f_m)$.

The following result proves naturality.

Proposition 2.4. For i = 1, ..., m, let $f_i: (CX_i, X_i) \longrightarrow (Y_i, *)$ be maps of pairs, and let $g_i: Y_i \longrightarrow Z_i$ and $l_i: W_i \longrightarrow X_i$ be maps. Denote by $g: (\underline{Y}, \underline{*})^{\partial \Delta^{m-1}} \longrightarrow (\underline{Z}, \underline{*})^{\partial \Delta^{m-1}}$ the map of polyhedral products induced by the maps g_i . Then

(11)
$$h_w(g_1 \circ f_1, \dots, g_m \circ f_m) = g \circ h_w(f_1, \dots, f_m)$$

and

(12)
$$h_w(f_1 \circ \Sigma l_1, \dots, f_m \circ \Sigma l_m) = h_w(f_1, \dots, f_m) \circ \overset{m}{\underset{i=1}{\underbrace{\star}}} l_i$$

where $\Sigma l_i: (CW_i, W_i) \longrightarrow (CX_i, X_i)$ denotes the map of pairs induced by l_i .

Proof. Both follow by the functoriality of the polyhedral product with respect to continuous maps of pairs. \Box

To consider symmetry of higher Whitehead maps, let $\sigma: (1, \ldots, m) \mapsto (i_1, \ldots, i_m)$ denote a permutation. For a space Z, consider maps $\iota: FW(Y_1, \ldots, Y_m) \longrightarrow Z$ and $\iota': FW(Y_{i_1}, \ldots, Y_{i_m}) \longrightarrow Z$ such that $\iota'(Y_{\sigma(i)}) = \iota(Y_i)$ for $i = 1, \ldots, m$. Denote by

$$h_w^Z(f_1,\ldots,f_m) = \iota \circ h_w(f_1,\ldots,f_m) : (\underline{CX},\underline{X})^{\partial \Delta^{m-1}} \longrightarrow Z$$

and by

$$h_w^Z(f_{i_1},\ldots,f_{i_m}) = \iota' \circ h_w(\sigma(f_1,\ldots,f_m)) : (\underline{CX},\underline{X})_{\sigma}^{\partial \Delta^{m-1}} \longrightarrow Z$$

where $(\underline{CX}, \underline{X})_{\sigma}$ denotes the permuted tuple. The permutation σ induces the permutation on the products $\prod CX_i$ and $\prod Y_i$, which in turn induce the permutation on the join $X_1 * \cdots * X_m$ and on the fat wedge $FW(Y_1, \ldots, Y_m)$, respectively.

Proposition 2.5. For i = 1, ..., m, let $f_i: (CX_i, X_i) \longrightarrow (Y_i, *)$ be maps. Then

(13)
$$h_w^Z(f_1,\ldots,f_m) = h_w^Z(\sigma(f_1,\ldots,f_m)) \circ \sigma$$

and furthermore if $X_i = S^{p_i-1}$ for i = 1, ..., m, then

(14)
$$h_w^Z(f_1,\ldots,f_m) = \varepsilon(\sigma)h_w^Z(\sigma(f_1,\ldots,f_m))$$

where $\varepsilon(\sigma)$ is the Koszul sign of σ , the product of $(-1)^{p_i p_j}$ for every transposition (i, j) of σ .

Proof. Consider the following diagram

The square commutes since the higher Whitehead map is the restriction of the product map $\prod f_i$, and therefore acts coordinate-wise. The triangle commutes by the definition of the maps ι and ι' . Therefore the diagram commutes, and identity (13) follows. Identity (14) then follows from (13) since the map σ is a map $S^{p_1+\dots+p_m-1} \longrightarrow S^{p_1+\dots+p_m-1}$ of degree $\varepsilon(\sigma)$.

The higher Whitehead map is an element of the higher Whitehead product of Porter [11], which is defined as follows. Consider spaces X_1, \ldots, X_m, Z . Given maps $f_i: \Sigma X_i \longrightarrow Z$ for $i = 1, \ldots, m$, denote by

$$\omega(f_1,\ldots,f_m) = \{\phi: FW(\Sigma X_1,\ldots,\Sigma X_m) \longrightarrow Z \mid \phi|_{\Sigma X_i} \simeq f_i \text{ for } i = 1,\ldots,m\}$$

the set of extensions up to homotopy of $\bigvee f_i: \bigvee \Sigma X_i \longrightarrow Z$ to $FW(\Sigma X_1, \ldots, \Sigma X_m)$. The *k*-ary higher Whitehead product $[f_1, \ldots, f_m]$ of the maps f_i is the set of homotopy classes

(15)
$$[f_1, ..., f_m] = \{ [\phi \circ \rho] \mid \phi \in \omega(f_1, \ldots, f_m) \} \subseteq \begin{bmatrix} m \\ \bigstar X_i, Z \\ i=1 \end{bmatrix}.$$

Proposition 2.6. Denote by $i_j: Y_j \longrightarrow FW(Y_1, \ldots, Y_m)$ the inclusion of Y_j into the *j*th coordinate. Then

$$h_w(f_1,\ldots,f_m) \in [i_1 \circ f_1,\ldots,i_m \circ f_m] \subseteq \begin{bmatrix} m \\ \bigstar \\ i=1 \end{bmatrix} X_i, FW(Y_1,\ldots,Y_m) \end{bmatrix}.$$

Proof. The result follows from the fact that higher Whitehead map is the composite $(\underline{CX}, \underline{X})^{\partial \Delta^{m-1}} \xrightarrow{(\underline{\rho})^{\partial \Delta^{m-1}}} (\underline{\SigmaX}, \underline{*})^{\partial \Delta^{m-1}} \xrightarrow{(\underline{f})^{\partial \Delta^{m-1}}} (\underline{Y}, \underline{*})^{\partial \Delta^{m-1}}$, together with the observation that $(\underline{f})^{\partial \Delta^{m-1}} \in \omega(f_1, \ldots, f_m)$.

The higher Whitehead product is trivial when it contains a null-homotopic map. We now study when the higher Whitehead map is null-homotopic and therefore implies triviality of higher Whitehead products. By [11, Theorem 2.3], the top row of the diagram

is a homotopy cofibration. Therefore, $h_w(f_1, \ldots, f_m)$ is null-homotopic if and only if there exists an extension up to homotopy $\phi: (\underline{\Sigma X}, \underline{*})^{\Delta^{m-1}} \longrightarrow (\underline{Y}, \underline{*})^{\partial \Delta^{m-1}}$ of $(\underline{f})^{\partial \Delta^{m-1}}: (\underline{\Sigma X}, \underline{*})^{\partial \Delta^{m-1}} \longrightarrow (\underline{Y}, \underline{*})^{\partial \Delta^{m-1}}.$

Notice that if we consider instead the composite

$$\iota \circ h_w(f_1, \dots, f_m) \colon (\underline{CX}, \underline{X})^{\partial \Delta^{m-1}} \xrightarrow{h_w(f_1, \dots, f_m)} (\underline{Y}, \underline{*})^{\partial \Delta^{m-1}} \longrightarrow (\underline{Y}, \underline{*})^{\Delta^{m-1}}$$

then an extension ϕ always exists, given by the product $(\underline{f})^{\Delta^{m-1}}$, and the higher Whitehead map is trivial in the space $(\underline{Y}, \underline{*})^{\Delta^{m-1}}$ corresponding to the full simplex Δ^{m-1} . This observation motivates the question of whether there are other simplicial complexes \mathcal{K} such that the composite $\iota \circ h_w(f_1, \ldots, f_m) : (\underline{CX}, \underline{X})^{\partial \Delta} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}}$ is null-homotopic.

For a subset $J = \{j_1, \ldots, j_n\} \subseteq [m]$, denote by $\Delta[J]$ and $\partial \Delta[J]$ the simplex on the vertex set J and its boundary complex, respectively. For \mathcal{K} containing $\partial \Delta[J]$, define

(17)
$$h_w^{\mathcal{K}}(f_{j_1},\ldots,f_{j_n}): (\underline{CX},\underline{X})^{\partial\Delta[J]} \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}}$$

to be the composite

$$(\underline{CX},\underline{X})^{\partial\Delta[J]} \xrightarrow{h_w(f_{j_1},\dots,f_{j_n})} (\underline{Y},\underline{*})^{\partial\Delta[J]} \xrightarrow{\iota} (\underline{Y},\underline{*})^{\mathcal{K}}$$

where the map ι is induced by the inclusion $\iota: \partial \Delta[J] \longrightarrow \mathcal{K}$.

Lemma 2.7. Let \mathcal{K} be a simplicial complex which contains $\partial \Delta[J]$. Then the map

$$h_w^{\mathcal{K}}(f_{j_1},\ldots,f_{j_n}):(\underline{CX},\underline{X})^{\partial\Delta[J]}\longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}}$$

is null-homotopic if either f_{j_i} is null-homotopic for some $i \in \{1, \ldots, m\}$, or $\Delta[J] \subseteq \mathcal{K}$.

Proof. Consider diagram (16). If f_i is null-homotopic for some $i \in \{1, \ldots, m\}$, then ϕ can be chosen to be $f_1 \times \cdots \times f_{i-1} \times \cdots \times f_{i+1} \times \cdots \times f_m$. In this case, the diagram commutes up to homotopy. If $\Delta[J] \subseteq \mathcal{K}$, then ϕ can be chosen to be $f_1 \times \cdots \times f_m$. \Box

The previous lemma focuses on combinatorial criteria for the triviality of higher Whitehead maps, and it does not take into account trivialisation coming from topological properties of joins.

Example 2.8. Let $f_1: \Sigma M(\mathbb{Z}_2, 1) \longrightarrow Y_1$ and $f_2: \Sigma M(\mathbb{Z}_3, 1) \longrightarrow Y_2$ be non nullhomotopic, where M(G, n) denotes the Moore space with reduced homology G concentrated in degree n. The space $M(\mathbb{Z}_2, 1) \wedge M(\mathbb{Z}_3, 1)$ is simply-connected, and by the Künneth theorem has trivial homology in all positive degrees, and therefore is contractible. Thus the higher Whitehead map

$$h_w(f_1, f_2) \colon \Sigma M(\mathbb{Z}_2, 1) \land M(\mathbb{Z}_3, 1) \longrightarrow Y_1 \lor Y_2$$

is null-homotopic.

The above example is motivated by the observation that the higher Whitehead map $h_w(f_1, \ldots, f_m)$ is null-homotopic if and only if the join of the adjoint maps $\hat{f}_1 * \cdots * \hat{f}_m : \bigstar X_i \longrightarrow \bigstar \Omega Y_i$ is null-homotopic. To see this, in the following diagram

(18)
$$\begin{array}{cccc} \prod_{i=1}^{m} \Omega Y_{i} & \xrightarrow{*} & \Omega Y_{1} * \cdots * \Omega Y_{m} & \longrightarrow & FW(Y_{1}, \dots, Y_{m}) & \xrightarrow{\iota} & \prod_{i=1}^{m} Y_{i} \\ & & & & \uparrow \\ & & & & \uparrow \\ & & & & \hat{f}_{1} * \cdots * \hat{f}_{m} & & \uparrow \\ & & & & X_{1} * \cdots * X_{m} & \longrightarrow & \prod_{i=1}^{m} CX_{i} \end{array}$$

the top row is a homotopy fibration sequence. The composite $\iota \circ h_w(f_1, \ldots, f_m)$ is null-homotopic, and a lift to the homotopy fibre of ι is given by $\hat{f}_1 * \cdots * \hat{f}_m$. Therefore if $\hat{f}_1 * \cdots * \hat{f}_m$ is null-homotopic, so is $h_w(f_1, \ldots, f_m)$. On the other hand, if $h_w(f_1, \ldots, f_m)$ is null-homotopic, then there is a further lift $X_1 * \cdots * X_m \longrightarrow$ $\Omega Y_1 \times \cdots \times \Omega Y_m$ of the map $\hat{f}_1 * \cdots * \hat{f}_m$. Since $\Omega Y_1 \times \cdots \times \Omega Y_m \longrightarrow \Omega Y_1 * \cdots * \Omega Y_m$ is null-homotopic, it follows that $\hat{f}_1 * \cdots * \hat{f}_m$ is itself null-homotopic.

3. The nested higher Whitehead map

Higher Whitehead maps are defined as maps of polyhedral products over the boundary of a simplex. In this section, we identify a class of higher Whitehead maps with richer combinatorial structure, by supposing that the maps f_i themselves are higher Whitehead maps. Such maps appear, for example, in the Jacobi identity (1) and Hardie's identity (2).

Definition 3.1. A nested higher Whitehead map is a higher Whitehead map $h_w(f_1, \ldots, f_m)$ where for at least one $i \in \{1, \ldots, m\}$, the map f_i is itself a higher Whitehead map.

Given a nested higher Whitehead map $h_w(f_1, \ldots, f_m)$, if $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{\partial \Delta^{l_i-1}}$ is a higher Whitehead map, define $\partial \overline{\Delta}^{l_i-1} = \partial \Delta^{l_i-1}$, and otherwise let $\partial \overline{\Delta}^{l_i-1} = \bullet$, the simplex on one vertex. We describe the codomain of the nested higher Whitehead map $h_w(f_1, \ldots, f_m)$ as the polyhedral product

(19)
$$\left(\underline{(\underline{Y}_{i},\underline{*})}^{\partial \overline{\Delta}^{l_{i}-1}},\underline{*}\right)^{\partial \Delta^{m-1}}$$

To express the rich combinatorial structure arising within this polyhedral product, we introduce the polyhedral join product, defined by Vidaurre [14]. A simplicial pair $(\mathcal{S}, \mathcal{T})$ consists of simplicial complexes \mathcal{S} and \mathcal{T} both with vertex set [l] such that \mathcal{T} is a subcomplex of \mathcal{S} .

Definition 3.2. Let \mathcal{K} be a simplicial complex on [m], and let $(\mathcal{S}_i, \mathcal{T}_i)$ be a simplicial pair on $[l_i]$ for $i = 1, \ldots, m$. Let $(\underline{S}, \underline{\mathcal{T}}) = \{(\mathcal{S}_i, \mathcal{T}_i)\}_{i=1}^m$ be an *m*-tuple of simplicial pairs. The *polyhedral join product* is the simplicial complex on vertex set $[l_1] \sqcup \ldots \sqcup [l_m]$ defined by

Special cases of polyhedral join products have been studied. If $\mathcal{T}_i = \{\emptyset\}$ for all i, then we denote by

(20)
$$\mathcal{K}\langle \mathcal{S}_1, ..., \mathcal{S}_m \rangle = (\underline{\mathcal{S}}, \{\underline{\emptyset}\})^{*\mathcal{K}}$$

and call it the substitution of $S_1, ..., S_m$ into \mathcal{K} (see [2]). If $S_i = \Delta^{l_i - 1}$ for all *i*, then we denote by

(21)
$$\mathcal{K}(\mathcal{T}_1,\ldots,\mathcal{T}_m) = (\underline{\Delta}^{l_i-1},\underline{\mathcal{T}})^{*\mathcal{K}}$$

and call it the *composition* of \mathcal{K} with $\mathcal{T}_1, ..., \mathcal{T}_m$ (see [3]). In the case that $\mathcal{K} = \Delta^{m-1}$ or $\mathcal{K} = \partial \Delta^{m-1}$, we denote the corresponding substitution complexes by $\Delta \langle \mathcal{S}_1, \ldots, \mathcal{S}_m \rangle$ and $\partial \Delta \langle \mathcal{S}_1, \ldots, \mathcal{S}_m \rangle$, respectively, and adopt the similar abbreviation for composition complexes. We observe that for any simplicial complex $\mathcal{K}, \ \mathcal{K} \langle \bullet, \ldots, \bullet \rangle = \mathcal{K} = \mathcal{K}(\circ, \ldots, \circ)$, where \bullet denotes the simplicial complex consisting of a single vertex, and \circ denotes the empty complex on a single vertex.

In analogy with the polyhedral product, the polyhedral join product is covariantly functorial with respect to simplicial inclusions $\mathcal{L} \longrightarrow \mathcal{K}$, and simplicial maps of pairs of simplicial complexes.

Vidaurre [14, Theorem 2.9] showed that polyhedral products over polyhedral join products decompose in the following way

(22)
$$(\underline{X},\underline{A})^{(\underline{S},\underline{\mathcal{T}})^{*\mathcal{K}}} = \left(\underline{(X},\underline{A})^{\mathcal{S}_i},\underline{(X},\underline{A})^{\mathcal{T}_i}\right)^{\mathcal{K}}.$$

Using the combinatorics of the polyhedral join product, we analyse nested higher Whitehead maps.

By (22), we rewrite the codomain (19) of the nested higher Whitehead map as

$$\left(\underline{(\underline{Y}_{i},\underline{*})}^{\partial\overline{\Delta}^{l_{i}-1}},\underline{*}\right)^{\partial\Delta^{m-1}} = (\underline{Y},\underline{*})^{\left(\underline{\partial\overline{\Delta}}^{l_{i}-1},\underline{\emptyset}\right)^{\ast\partial\Delta^{m-1}}} = (\underline{Y},\underline{*})^{\partial\Delta^{m-1}\langle\partial\overline{\Delta}^{l_{1}-1},\dots,\partial\overline{\Delta}^{l_{m-1}}\rangle}$$

By Lemma 2.7, the composition of the nested higher Whitehead map with the inclusion

$$(\underline{Y},\underline{*})^{\partial\Delta^{m-1}\langle\partial\overline{\Delta}^{l_1-1},\ldots,\partial\overline{\Delta}^{l_m-1}\rangle} \longrightarrow (\underline{Y},\underline{*})^{\Delta^{m-1}\langle\partial\overline{\Delta}^{l_1-1},\ldots,\partial\overline{\Delta}^{l_m-1}\rangle} = \left(\underline{(\underline{Y}_i,\underline{*})}^{\partial\overline{\Delta}^{l_i-1}},\underline{*}\right)^{\Delta^{m-1}\langle\partial\overline{\Delta}^{l_1-1},\ldots,\partial\overline{\Delta}^{l_m-1}\rangle} = \left(\underline{(\underline{Y}_i,\underline{*})}^{\partial\overline{\Delta}^{l_1-1}},\ldots,\partial\overline{\Delta}^{l_m-1}\rangle$$

induced by the inclusion of simplicial complexes, is null-homotopic. This motivates the question of the description of subcomplexes \mathcal{K} of $\Delta^{m-1} \langle \partial \overline{\Delta}^{l_1-1}, \ldots, \partial \overline{\Delta}^{l_m-1} \rangle$ for which the map

(23)
$$h_w^{\mathcal{K}}(f_1,\ldots,f_m): (\underline{CX},\underline{X})^{\partial\Delta^{m-1}} \xrightarrow{h_w} (\underline{Y},\underline{*})^{\partial\Delta\langle\partial\overline{\Delta}^{l_1-1}},\ldots,\partial\overline{\Delta}^{l_m-1}\rangle \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}}$$

is null-homotopic. The following two propositions state topological and combinatorial conditions under which the higher Whitehead map is trivial.

Proposition 3.3. Let \mathcal{K} be a simplicial complex on $[l_1] \sqcup \cdots \sqcup [l_m]$ such that $\partial \Delta^{m-1} \langle \partial \overline{\Delta}^{l_1-1}, \ldots, \partial \overline{\Delta}^{l_m-1} \rangle \subseteq \mathcal{K}$. Suppose that for at least one $i \in \{1, \ldots, m\}$, $f_i = h_w(f_{i_1}, \ldots, f_{i_{l_i}}) \colon \Sigma X_i \longrightarrow (\underline{Y}_i, \underline{*})^{\partial \Delta^{l_i-1}}$ with $l_i \geq 2$. If one of the following is satisfied:

(i) the map
$$f_i: \Sigma X_i \longrightarrow Y_i$$
 is null-homotopic for some $i = 1, ..., m$;
(ii) $\Delta \langle \partial \overline{\Delta}^{l_1 - 1}, ..., \partial \overline{\Delta}^{l_m - 1} \rangle \subseteq \mathcal{K}$,

then $h_w^{\mathcal{K}}(f_1,\ldots,f_m)$ is null-homotopic.

Proof. The claim follows by Lemma 2.7, together with (22).

Proposition 3.4. Let \mathcal{K} be a simplicial complex on $[l_1] \sqcup \cdots \sqcup [l_m]$ such that $\partial \Delta^{m-1} \langle \partial \overline{\Delta}^{l_1-1}, \ldots, \partial \overline{\Delta}^{l_m-1} \rangle \subseteq \mathcal{K}$. Suppose that for at least one $i \in \{1, \ldots, m\}$, $f_i = h_w(f_{i_1}, \ldots, f_{i_{l_i}}) \colon \Sigma X_i \longrightarrow (\underline{Y}_i, \underline{*})^{\partial \Delta^{l_i-1}}$ with $l_i \geq 2$. If there exists $1 \leq i \leq m$ such that \mathcal{K} contains $\partial \Delta \langle \partial \overline{\Delta}^{l_1-1}, \ldots, \Delta^{l_i-1}, \ldots, \partial \overline{\Delta}^{l_m-1} \rangle$, where f_i is a higher Whitehead map, then $h_w^{\mathcal{K}}(f_1, \ldots, f_m)$ is null-homotopic.

Proof. Let
$$f'_i = h_w^{\Delta^{l_i-1}}(f_{i_1}, \dots, f_{i_{l_i}})$$
. Since $\partial \Delta^{l_i-1} \subseteq \Delta^{l_i-1}$,
 $h_w^{\mathcal{K}}(f_1, \dots, f_i, \dots, f_m) = h_w^{\mathcal{K}}(f_1, \dots, f'_i, \dots, f_m)$

The statement follows since the map $h_w^{\mathcal{K}}(f_1, \ldots, f'_i, \ldots, f_m)$ is null-homotopic by Lemma 2.7.

Whether the higher Whitehead map is null-homotopic is determined by both the combinatorics of \mathcal{K} and homotopy theoretic properties of the maps $f_i: \Sigma X_i \longrightarrow Y_i$, as shown in Example 2.8. In the following example, we study a family of nested higher Whitehead maps of inclusions $S^2 \longrightarrow DJ_{\mathcal{K}}$. Such maps were studied algebraically by Abramyan [1]. By considering a family of non null-homotopic maps, we construct non null-homotopic maps in trivial higher Whitehead products.

Example 3.5. Let $p \ge 2$, $q \ge 3$, and $S_1 = \partial \Delta^{p-1}$. Define the simplicial complex \mathcal{K} on the vertex set $[p+q-1] = \{1_1, \ldots, 1_p\} \sqcup \{2, \ldots, q\}$ by

$$\mathcal{K} = \partial \Delta^{q-1} \langle \mathcal{S}_1, p+1, \dots, p+q-1 \rangle \cup \Delta[1_1, \dots, 1_p]$$

= $\partial \Delta^{q-1} \langle \partial \Delta[1_1, \dots, 1_p], p+1, \dots, p+q-1 \rangle \cup \Delta[1_1, \dots, 1_p].$

Let $\mu_{1_i}: S_{1_i}^2 \longrightarrow \mathbb{C}P_{1_i}^\infty$ and $\mu_j: S_j^2 \longrightarrow \mathbb{C}P_j^\infty$ for $1 \leq i \leq p$ and $2 \leq j \leq q$ denote the inclusions of the bottom cell. The map

$$h_w^{\mathcal{K}}(h_w(\mu_{1_1},\ldots,\mu_{1_p}),\mu_2,\ldots,\mu_q):S^{2p+2q-5}\longrightarrow (\mathbb{C}P^{\infty},*)^{\partial\Delta^{q-1}\langle\partial\Delta^{q-1},\bullet,\ldots,\bullet\rangle}\longrightarrow (\mathbb{C}P^{\infty},*)^{\mathcal{K}}$$

is non null-homotopic, see [1, Proposition 7.2]. By Proposition 2.6, this map is an element of the higher Whitehead product $[\iota_1 \circ h_w(f_{1_1}, \ldots, f_{1_p}), \iota_2 \circ f_2, \ldots, \iota_q \circ f_q]$, which is trivial since $\iota_1 \circ h_w(f_{1_1}, \ldots, f_{1_p})$ is null-homotopic by Proposition 3.3. We therefore obtain higher Whitehead products with indeterminacy.

4. Folded higher Whitehead maps

The homotopy groups $\pi_*(X)$ of a space X are a central object of study in homotopy theory. The Whitehead product $[\cdot, \cdot]: \pi_m(X) \times \pi_n(X) \longrightarrow \pi_{m+n-1}(X)$ endows the graded module $\pi_*(X) = \bigoplus_{k \ge 1} \pi_k(X)$ with the structure of a graded quasi-Lie algebra. Under the isomorphism $\pi_{*+1}(X) \cong \pi_*(\Omega X)$, Whitehead products appear as commutators in the algebra $H_*(\Omega X; \mathbb{Z})$ under the Hurewicz map $\pi_*(\Omega X) \longrightarrow$ $H_*(\Omega X; \mathbb{Z})$. Rationalising, the algebra $H_*(\Omega X; \mathbb{Q})$ is the universal enveloping algebra of $\pi_*(\Omega X) \otimes \mathbb{Q}$, meaning that homotopy-theoretic information about X can be extracted from determining the primitive elements in a model for $H_*(\Omega X; \mathbb{Q})$.

For \mathcal{K} a simplicial complex, the homotopy Lie algebra $\pi_*(\Omega DJ_{\mathcal{K}}) \otimes \mathbb{Q}$ was studied by Zhuravleva [15]. Using an Adams–Hilton model of $H_*(\Omega DJ_{\mathcal{K}}, \mathbb{Q})$, she described the Hurewicz image of the adjoint of higher Whitehead maps.

For example, let $\mathcal{K} = \partial \Delta \langle \partial \Delta \langle 1, 2, 3 \rangle, 4, 5 \rangle$ and let $u_i: S^1 \longrightarrow \Omega \mathbb{C}P^{\infty}$ be the adjoint of the inclusion $\mu_i: S^2 \longrightarrow \mathbb{C}P^{\infty}$ of the bottom cell for $i = 1, \ldots, m$. The nested brackets $[[u_1, u_2, u_3], [u_1, u_4, u_5]]$ and $[[[u_1, u_2, u_3], u_4, u_5], u_1]$, the adjoints to $[h_w^{\mathcal{K}}(\mu_1, \mu_2, \mu_3), h_w^{\mathcal{K}}(\mu_1, \mu_4, \mu_5)]$ and $[h_w^{\mathcal{K}}(h_w(\mu_1, \mu_2, \mu_3), \mu_4, \mu_5), \mu_1^{\mathcal{K}}]$, respectively, are defined in $\pi_8(\Omega DJ_{\mathcal{K}}) \otimes \mathbb{Q}$.

We study these Whitehead products from the geometric perspective of the higher Whitehead map, and construct the maps considered by Zhuravleva integrally. Our construction realises 2-torsion elements of homotopy groups by giving a combinatorial interpretation of higher Whitehead maps with repeated factors.

Let \mathcal{K} be a simplicial complex realising a nested higher Whitehead map

$$h_w^{\mathcal{K}}(f_1,\ldots,f_m): (\underline{CX},\underline{X})^{\partial\Delta} \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}}$$

We study when the simplicial fold operation $\mathcal{K} \longrightarrow \overline{\mathcal{K}}$ identifying two or more vertices preserves the structure of a higher Whitehead map. Specifically, we ask when the simplicial fold $\mathcal{K} \longrightarrow \overline{\mathcal{K}}$ induces a map of polyhedral products $\nabla: (\underline{Y}, \underline{*})^{\mathcal{K}} \longrightarrow (\underline{Y}, \underline{*})^{\overline{\mathcal{K}}}$, in which case we will call the composite

$$\nabla h_w(f_1,\ldots,f_m): (\underline{CX},\underline{X})^{\partial\Delta} \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}} \longrightarrow (\underline{Y},\underline{*})^{\overline{\mathcal{K}}}$$

a folded higher Whitehead map. One situation where this is possible when the $Y_i = Y$, an associative *H*-space.

To build up to the definition of the folded higher Whitehead map, we begin by defining a fold of a simplicial complex.

Definition 4.1. A fold of a vertex set [m] consists of disjoint subsets $I, J \subset [m]$ and a surjective map $\psi: I \longrightarrow J$. Given a fold $\psi: I \longrightarrow J$ of [m], we denote $I_j = \psi^{-1}(j)$ for each $j \in J$, so that $I = \bigsqcup_{i \in J} I_j$.

Let \mathcal{K} be a simplicial complex on [m]. A fold $\psi: I \longrightarrow J$ of [m] extends to a map on \mathcal{K} by sending a simplex $(i_1, \ldots, i_k) \in \mathcal{K}$ to $(\overline{\psi}(i_1), \ldots, \overline{\psi}(i_k))$, where $\overline{\psi}(i) = \psi(i)$ if $i \in I$, and $\overline{\psi}(i) = i$, otherwise. We interchange freely between referring to $\psi: I \longrightarrow J$ as a fold of \mathcal{K} and of its vertex set [m]. We define the folded complex $\mathcal{K}_{\nabla(I,J)}$ of the fold of \mathcal{K} induced by ψ as the image $\overline{\psi}(\mathcal{K})$.

It follows from the definition that

(24) $\mathcal{K}_{\nabla(I,J)} = \{ \sigma \subseteq [m] \setminus I \mid \sigma \in \mathcal{K} \text{ or } (\sigma \setminus \{j\}) \sqcup \{i\} \in \mathcal{K} \text{ for some } j \in J, i \in I_j \}.$

When $J = \{j\}$ consists of one element we abbreviate $\mathcal{K}_{\nabla(I,\{j\})}$ to $\mathcal{K}_{\nabla(I,j)}$. If further $I = \{i\}$, then we abbreviate $\mathcal{K}_{\nabla(\{i\},j)} = \mathcal{K}_{\nabla(i,j)}$.

Example 4.2. Let \mathcal{K} be the simplicial complex shown on the left of Figure 1. Let $I = \{4\}, J = \{1\}$ and let $\psi: I \longrightarrow J$ be the map sending $\{4\}$ to $\{1\}$. Then $\mathcal{K}_{\nabla(I,J)}$ is the simplicial complex on vertices $\{1, 2, 3\}$ shown on the right of Figure 1.




FIGURE 1. The fold $\psi: \{4\} \longrightarrow \{1\}$ of a simplicial complex.

The following properties follow from Definition 4.1.

Proposition 4.3. Let \mathcal{K} be a simplicial complex on [m]. Then:

(i) if $i, j \in [m]$ with $i \neq j$, then $\mathcal{K}_{\nabla(i,j)} \cong \mathcal{K}_{\nabla(j,i)}$;

(ii) if $I = \{i_1, \ldots, i_n\} \subseteq [m]$, and $j \in [m]$ with $j \notin I$, then

$$\mathcal{K}_{\nabla(I,j)} = \mathcal{K}_{\nabla(i_1,j)\nabla(i_2,j)\dots\nabla(i_n,j)};$$

(iii) if $I_1, I_2 \subseteq [m]$ are such that $I_1 \cap I_2 = \emptyset$, and $j \notin I_1 \sqcup I_2$, then

$$\mathcal{K}_{\nabla(I_1,j)\nabla(I_2,j)} = \mathcal{K}_{\nabla(I_2,j)\nabla(I_1,j)}.$$

To describe the map of polyhedral products induced by a fold of simplicial complexes, let $\psi: I \longrightarrow J$ be a fold of \mathcal{K} . Suppose that Y_j is an associative H-space for each $j \in J$, and that $Y_i = Y_j$ for each $i \in I_j$. For $j \in J$, the H-multiplication map $\mu_j: Y_j \times Y_j \longrightarrow Y_j$ extends up to homotopy to a map

$$\prod_{i \in I_j} Y_i \times Y_j \longrightarrow Y_j$$

Therefore, the fold $\psi: I \longrightarrow J$ induces a map $\prod_{i \in [m]} Y_i \longrightarrow \prod_{i \in ([m]-I)} Y_i$ given by

(25)
$$\prod_{j\in J}\prod_{i\in I_j} (Y_i \times Y_j) \times \prod_{i\in ([m]-(I\sqcup J))} Y_i \longrightarrow \prod_{j\in J} Y_j \times \prod_{i\in ([m]-(I\sqcup J))} Y_i$$

The fold $\psi: I \longrightarrow J$ of \mathcal{K} induces the map of polyhedral products

(26)
$$\nabla_{(I,J)}: (\underline{Y}, \underline{*})^{\mathcal{K}} = \bigcup_{\sigma \in \mathcal{K}} \prod_{i \in \sigma} Y_i \longrightarrow \bigcup_{\sigma \in \mathcal{K}} \prod_{i \in \overline{\psi}(\sigma)} Y_i = (\underline{Y}, \underline{*})^{\mathcal{K}_{\nabla(I,J)}}$$

defined as the restriction of (25) to $(\underline{Y}, \underline{*})^{\mathcal{K}}$.

We call $(\underline{Y}, \underline{*})^{\mathcal{K}_{\nabla(I,J)}}$ the fold of $(\underline{Y}, \underline{*})^{\mathcal{K}}$ under the map $\psi: I \longrightarrow J$, and the map $\nabla_{(I,J)}$ the fold map on polyhedral products induced by the map $\psi: I \longrightarrow J$.

Since the multiplication $\mu: Y_j \times Y_j \longrightarrow Y_j$ extends the fold map $Y_j \vee Y_j \longrightarrow Y_j$ up to homotopy, the fold $\psi: I \longrightarrow J$ of \mathcal{K} also induces a map of polyhedral products whenever I_j is such that \mathcal{K}_{I_j} consists of disjoint vertices. In this case, the map (26) is defined without requiring Y_j to be an *H*-space.

We specialise to fold maps of polyhedral products related to higher Whitehead maps.

Definition 4.4. The folded higher Whitehead map is the composite

 $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m): (\underline{CX},\underline{X})^{\partial\Delta} \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}} \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}_{\nabla(I,J)}}.$

In analogy with the nested higher Whitehead map, the folded higher Whitehead map is an element of a particular higher Whitehead product.

Proposition 4.5. The folded higher Whitehead map is an element of the higher Whitehead product

$$\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m)\in [\nabla_{(I,J)}f_1^{\mathcal{K}},\ldots,\nabla_{(I,J)}f_m^{\mathcal{K}}]\subseteq [X_1*\cdots*X_m,(\underline{Y},\underline{*})^{\mathcal{K}_{\nabla(I,J)}}].$$

Proof. The folded higher Whitehead map $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m)$ is the composite along the top row of the diagram

Since for any *H*-space *Z* the multiplication $Z \times Z \longrightarrow Z$ extends the fold map $Z \vee Z \longrightarrow Z$, the right-hand square of the diagram commutes. It follows that the composite $(\underline{Y}, \underline{*})^{\partial \Delta} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_{\nabla(I,J)}}$ extends the map

$$\bigvee_{i\in[m]} \nabla_{(I,J)} f_i \colon \bigvee_{i\in[m]} \Sigma X_i \longrightarrow \bigvee_{i\in[m]} Y_i \longrightarrow \bigvee_{i\in[m]-I} Y_i \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}_{\nabla(I,J)}}$$

Therefore by definition of the higher Whitehead product, $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m) \in [\nabla_{(I,J)}f_1^{\mathcal{K}},\ldots,\nabla_{(I,J)}f_m^{\mathcal{K}}].$

Let


$$h_w^{\mathcal{K}}(f_1,\ldots,f_m)\colon (\underline{CX},\underline{X})^{\partial\Delta} \longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}}$$

be the nested higher Whitehead map, as in Definition 3.1. We study sufficient combinatorial conditions under which folded nested higher Whitehead maps are nullhomotopic. We begin with some preparatory results.

Lemma 4.6. Let S_1, \ldots, S_m be simplicial complexes on $[l_1], \ldots, [l_m]$, respectively. Let $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{S_i}$ be maps for $i = 1, \ldots, m$. Let $\mathcal{K}, \mathcal{K}'$ be simplicial complexes on $[l] = [l_1] \sqcup \cdots \sqcup [l_m]$ containing $\partial \Delta \langle S_1, \ldots, S_m \rangle$ and let $\psi: I \longrightarrow J$ be a fold of both \mathcal{K} and \mathcal{K}' such that $\mathcal{K}_{\nabla(I,J)} = \mathcal{K}'_{\nabla(I,J)}$. If Y_j is an associative H-space for each $j \in J$, and $Y_i = Y_j$ for each $i \in I_j$, then

$$\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m)=\nabla_{(I,J)}h_w^{\mathcal{K}'}(f_1,\ldots,f_m).$$

Proof. Since $\mathcal{K}_{\nabla(I,J)} = \mathcal{K}'_{\nabla(I,J)}$, there is a commutative diagram



Therefore, by functoriality of the polyhedral product with respect to simplicial inclusions, the following diagram commutes

where the composite around the top of the square is $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m)$ and the composite around the bottom of the square is $\nabla_{(I,J)}h_w^{\mathcal{K}'}(f_1,\ldots,f_m)$.

Whether the folded higher Whitehead map is null-homotopic is not necessarily determined by the complex \mathcal{K} since the fold of \mathcal{K} can factor through a different complex \mathcal{K}' which makes the map null-homotopic. To obtain a condition for when a folded higher Whitehead map $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m)$ is null-homotopic, we define a simplicial complex \mathcal{L}_{ψ} containing all complexes which fold onto $\mathcal{K}_{\nabla}(I,J)$ by $\psi: I \longrightarrow J$.

Definition 4.7. Let \mathcal{K} be a simplicial complex on [m], and let $\psi: I \longrightarrow J$ be a fold of \mathcal{K} for $I, J \subset [m]$. Denote by $\{k_1, \ldots, k_l\}$ the vertex set of the folded complex $\mathcal{K}_{\nabla(I,J)}$, and let $I_j = \psi^{-1}(j)$ for each $j \in J$. Define the simplicial complex \mathcal{L}_{ψ} on [m] by

$$\mathcal{L}_{\psi} = \mathcal{K}_{\nabla(I,J)} \langle \Delta[\{k_1\} \sqcup I_{k_1}], \dots, \Delta[\{k_l\} \sqcup I_{k_l}] \rangle$$

where $I_k = \emptyset$ if $k \notin J$.

Example 4.8. Consider the simplicial complex \mathcal{K} and the fold map $\psi: I \longrightarrow J$ from Example 4.2. The complex $\mathcal{K}_{\nabla(I,J)}$ is shown on the left of Figure 2, while the complex $\mathcal{L}_{\psi} = \mathcal{K}_{\nabla(I,J)} \langle \Delta[1,4], 2, 3 \rangle$ is shown on the right.

The complex \mathcal{L}_{ψ} is the largest simplicial complex which folds to $\mathcal{K}_{\nabla(I,J)}$ under $\psi: I \longrightarrow J$, and therefore in particular contains \mathcal{K} .

Lemma 4.9. Suppose that \mathcal{L}' is such that $\mathcal{L}'_{\nabla(I,J)} \subseteq \mathcal{K}_{\nabla(I,J)}$. Then $\mathcal{L}' \subseteq \mathcal{L}_{\psi}$.




FIGURE 2. Construction of the simplicial complex \mathcal{L}_{ψ} .

Proof. Let $J = \{j_1, \ldots, j_r\}$. For any $\sigma \in \mathcal{L}'$, write $\sigma = \sigma_{j_1} \sqcup \cdots \sqcup \sigma_{j_r} \sqcup \sigma'$, where $\sigma_{j_k} \subseteq I_{j_k}$ and $\sigma' \in [m] \setminus I$. Since $\mathcal{L}'_{\nabla(I,J)} \subseteq \mathcal{K}_{\nabla(I,J)}$, it follows that $\overline{\psi}(\sigma) = (j_1 \sqcup \cdots \sqcup j_r) \cup \sigma' \in \mathcal{K}_{\nabla(I,J)}$. By construction of L_{ψ} , $(I_{j_1} \sqcup \cdots \sqcup I_{j_r}) \cup (j_1 \sqcup \cdots \sqcup j_r) \cup \sigma' \in \mathcal{L}_{\psi}$. As $\sigma_{j_k} \subseteq I_{j_k}$, it follows that $\sigma \in \mathcal{L}_{\psi}$.

Finally, we state combinatorial conditions under which the folded higher Whitehead map is null-homotopic.

Proposition 4.10. Let S_1, \ldots, S_m be simplicial complexes on $[l_1], \ldots, [l_m]$, respectively. Let $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{S_i}$ be maps for $i = 1, \ldots, m$. Let \mathcal{K} be a simplicial complex on $[l_1] \sqcup \cdots \sqcup [l_m]$ containing $\partial \Delta \langle S_1, \ldots, S_m \rangle$ and let $\psi: I \longrightarrow J$ be a fold of \mathcal{K} .

The folded higher Whitehead map

$$\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1,\ldots,f_m):(\underline{CX},\underline{X})^{\partial\Delta}\longrightarrow (\underline{Y},\underline{*})^{\mathcal{K}_{\nabla(I,J)}}$$

is null-homotopic if one of the following holds:

- (i) $\mathcal{K}_{\nabla(I,J)}$ contains the folded complex $\mathcal{K}'_{\nabla(I,J)}$, where either $\mathcal{K}' = \partial \Delta \langle \mathcal{S}_1, \dots, \mathcal{S}'_i, \dots, \mathcal{S}_m \rangle$ for \mathcal{S}'_i a trivialising complex of f_i , or $\mathcal{K}' = \Delta \langle \mathcal{S}_1, \dots, \mathcal{S}_m \rangle$;
- (ii) the map

$$h_w^{\mathcal{L}_\psi}(f_1,\ldots,f_m):(\underline{CX},\underline{X})^{\partial\Delta}\longrightarrow (\underline{Y},\underline{*})^{\mathcal{L}_\psi}$$

is null-homotopic.

Proof. Suppose that (i) holds. Then $h_w^{\mathcal{K}'}(f_1, \ldots, f_m)$ is null-homotopic by Proposition 3.3. By Lemma 4.6, we have $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1, \ldots, f_m) = \nabla_{(I,J)}h_w^{\mathcal{K}'}(f_1, \ldots, f_m)$, and so the map $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1, \ldots, f_m)$ is null-homotopic. The same argument holds if instead (ii) holds, since \mathcal{L}_{ψ} folds to $\mathcal{K}_{\nabla(I,J)}$ by Lemma 4.9.

Example 4.11. Consider the simplicial complex \mathcal{K} and the fold map $\psi: I \longrightarrow J$ from Example 4.2. Let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps for $i = 1, \ldots, 4$. The nested higher Whitehead maps $h_w^{\mathcal{K}}(h_w(f_1, f_2, f_3), f_4)$ and $h_w^{\mathcal{K}}(h_w(f_1, f_4), f_2, f_3)$ are defined in $(\underline{Y}, \underline{*})^{\mathcal{K}}$. Therefore the folded higher Whitehead maps $\nabla_{(I,J)}h_w^{\mathcal{K}}(h_w(f_1, f_2, f_3), f_4)$ and $\nabla_{(I,J)}h_w^{\mathcal{K}}(h_w(f_1, f_4), f_2, f_3)$ are defined in $(\underline{Y}, \underline{*})^{\mathcal{K}_{\nabla(I,J)}}$. By Proposition 4.10, the folded higher Whitehead map $\nabla_{(I,J)}h_w^{\mathcal{K}}(h_w(f_1, f_4), f_2, f_3)$ is null-homotopic since $h_w^{\mathcal{L}_{\psi}}(h_w(h_w(f_1, f_4), f_2, f_3))$ is null-homotopic by Proposition 3.3.

By Proposition 4.5, for $\mathcal{L} = \mathcal{K}_{\nabla(I,J)}$, these folded higher Whitehead maps are elements of the higher Whitehead products $[h_w^{\mathcal{L}}(f_1, f_2, f_3), f_1^{\mathcal{L}}]$ and $[\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1, f_4), f_2^{\mathcal{L}}, f_3^{\mathcal{L}}]$,

respectively. Furthermore, by Proposition 4.5, the map $\nabla_{(I,J)}h_w^{\mathcal{K}}(f_1, f_4)$ is the composite of $[f_1, f_1]$ with the inclusion $Y_1 \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{L}}$.

5. Relations

We recall the relations among exterior Whitehead products (see (4)) due to Hardie [8]. Let $f_i: S^{q_i} \longrightarrow Y_i$ be maps for $i = 1, \ldots, m$, where $q_i \ge 2$, and let

$$Z = \bigcup_{i=1}^{m} Y_i \vee FW(Y_1, \dots, Y_{i-1}, Y_{i+1}, \dots, Y_m).$$

Then

(27)
$$\sum_{i=1}^{m} (-1)^{\eta(i)} [\theta_i f_i, \psi_i \{ f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_m \}] = 0 \in \pi_{q-2}(Z)$$

where $q = q_1 + \cdots + q_m - 1$, the maps $\theta_i: Y_i \longrightarrow Z$ and $\psi_i: FW(Y_1, \ldots, Y_{i-1}, Y_{i+1}, \ldots, Y_m) \longrightarrow Z$ are inclusions, and $\eta(i) = q_i(q_1 + \cdots + q_i) + 1$.

We rewrite relation (27) using polyhedral products in order to detect the combinatorial structure which captures it. Each summand $\{\theta_i f_i, \psi_i \{f_1, \ldots, f_{i-1}, f_{i+1}, \ldots, f_m\}\}$ is a map of polyhedral products with domain $(\underline{D}, \underline{S})^{\partial \Delta^1} = \sum^{m-2} \bigwedge_{i=1}^m S^{q_i-1}$ and codomain Z expressed as

$$Z = \bigcup_{i=1}^{m} Y_i \vee FW(Y_1, \dots, Y_{i-1}, Y_{i+1}, \dots, Y_m)$$

= { $(y_1, \dots, y_m) \in Y_1 \times \dots \times Y_m \mid y_i = y_j = * \text{ for } 1 \leq i < j \leq m$ }
= $(\underline{Y}, \underline{*})^{\text{sk}^{m-3} \Delta^{m-1}}$.

We analyse the combinatorial structure of the simplicial complex $\operatorname{sk}^{m-3} \Delta^{m-1}$. Each missing face $\{[m] \setminus \{i\} \mid i = 1, \ldots, m\}$ of $\operatorname{sk}^{m-3} \Delta^{m-1}$ gives rise to the higher Whitehead map $h_w(f_1, \ldots, f_{i-1}, f_{i+1}, \ldots, f_m)$ in $(\underline{Y}, \underline{*})^{\operatorname{sk}^{m-3} \Delta^{m-1}}$. Summing these, we rewrite relation (27) as

(28)
$$\sum_{i=1}^{m} (-1)^{\eta(i)} h_w^{\operatorname{sk}^{m-3} \Delta^{m-1}} (h_w(f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_m), f_i) = 0.$$

We replace the spherical maps f_i with maps $\Sigma X_i \longrightarrow Y_i$, where X_i is a suspension. To increase the relation complexity combinatorially, we propagate the missing face structure of $\operatorname{sk}^{m-3} \Delta^{m-1} = \operatorname{sk}^{k-3} \Delta^{k-1}(\emptyset, \ldots, \emptyset)$ by replacing $\emptyset = \partial \Delta^0$ with $\partial \Delta^n$ for various n > 0.

Given a vertex set [m], a k-partition Π is a collection of pairwise disjoint subsets $\{P_1, \ldots, P_k\}$ of [m] such that $\bigcup_{i=1}^k P_i = [m]$.

Definition 5.1. Let $\Pi = \{P_1, \ldots, P_k\}$ be a k-partition of [m], where $P_j = \{i_1, \ldots, i_{n_j}\}$ for $j = 1, \ldots, k$. The *identity complex* \mathcal{K}_{Π} associated to Π is defined by

$$\mathcal{K}_{\Pi} = \operatorname{sk}^{k-3} \Delta^{k-1}(\partial \Delta[P_1], \dots, \partial \Delta[P_k]).$$

The identity complex $\operatorname{sk}^{m-3} \Delta^{m-1}$ associated to Hardie's identity corresponds to the partition $\Pi = \{\{1\}, \ldots, \{m\}\}$ of [m].

The Alexander dual $\hat{\mathcal{K}}$ of a simplicial complex \mathcal{K} on [m] is the simplicial complex with simplices $\{[m] \setminus \sigma \mid \sigma \notin \mathcal{K}\}$. In particular, the maximal faces of $\hat{\mathcal{K}}$ are the complements in [m] of the minimal missing faces of \mathcal{K} . Since the minimal missing faces of sk^{m-3} Δ^{m-1} are $\{[m] \setminus \{i\} \mid i = 1, \ldots, m\}$, it is Alexander dual to the simplicial complex consisting of m disjoint vertices,

$$\operatorname{sk}^{m-3} \Delta^{m-1} = \widehat{\bullet_{[m]}}.$$

Identity complexes admit a similar description in terms of their Alexander duals, which we describe next.

Proposition 5.2. Let $\Pi = \{P_1, \ldots, P_k\}$ be a k-partition of [m]. Then

(29)
$$MF(\mathcal{K}_{\Pi}) = \{[m] \setminus P_i \mid i = 1, \dots, k\}$$

To prove Proposition 5.2, we first compute the minimal missing faces of the polyhedral join product.

Proposition 5.3. Let \mathcal{K} be a simplicial complex on [m], and let $(\mathcal{S}_1, \mathcal{T}_1), ..., (\mathcal{S}_m, \mathcal{T}_m)$ be simplicial pairs on vertex sets $[l_1], ..., [l_m]$, respectively. Then

$$MF\left((\underline{\mathcal{S}},\underline{\mathcal{T}})^{*\mathcal{K}}\right) = \{\tau \in MF(\mathcal{S}_i) \mid i \in \mathcal{K}\} \sqcup \left\{\bigsqcup_{i \in \kappa} \tau_i \mid \kappa \in MF(\mathcal{K}), \tau_i \in MF(\mathcal{T}_i), \tau_i \in \mathcal{S}_i\right\}$$

Proof. We first show that

$$\{J \in MF(\mathcal{S}_i) \mid i \in \mathcal{K}\} \sqcup \left\{ \bigsqcup_{i \in L} J_i \mid L \in MF(\mathcal{K}), J_i \in MF(\mathcal{T}_i), J_i \in \mathcal{S}_i \right\}$$
$$\subseteq MF\left((\underline{\mathcal{S}}, \underline{\mathcal{T}})^{*\mathcal{K}}\right).$$

For any $\{i\} \in \mathcal{K}, J \in MF(\mathcal{S}_i)$ implies that $J \in MF\left((\underline{\mathcal{S}}, \underline{\mathcal{T}})^{*\mathcal{K}}\right)$. Now consider $\bigsqcup_{i \in L} J_i$, where $L \in MF(\mathcal{K}), J_i \in MF(\mathcal{T}_i)$ and $J_i \in \mathcal{S}_i$ for all $i \in L$. This is a missing face of $(\underline{\mathcal{S}}, \underline{\mathcal{T}})^{*\mathcal{K}}$ by definition of the polyhedral join. Moreover, it is minimal since for any $i \in L$ and $s \in J_i$

$$\bigsqcup_{i \neq k \in L} J_k \sqcup (J_i - \{s\}) = \bigsqcup_{k \in \tau} J_k \sqcup \sigma_i \in (\underline{S}, \underline{\mathcal{T}})^{*\tau}$$

where $\tau \in \mathcal{K}$ since L is a minimal missing face, and $\sigma_i \in \mathcal{T}_i$ since J_i is a minimal missing face.

Now we show that

$$MF\left((\underline{\mathcal{S}},\underline{\mathcal{T}})^{*\mathcal{K}}\right) \subseteq \{J \in MF(\mathcal{S}_i) \mid i \in \mathcal{K}\}$$
$$\sqcup \left\{ \bigsqcup_{i \in L} J_i \mid L \in MF(\mathcal{K}), J_i \in MF(\mathcal{T}_i), J_i \in \mathcal{S}_i \right\}$$

Let $F \in MF((\underline{S},\underline{\mathcal{T}})^{*\mathcal{K}})$. We show that either $F \in \{J \in MF(\mathcal{S}_i) \mid i \in \mathcal{K}\}$ or $F \in \{\bigsqcup_{i \in L} F_i \mid L \in MF(\mathcal{K}), F_i \in MF(\mathcal{T}_i), F_i \in \mathcal{S}_i\}.$

Denote by $F_i = F|_{[l_i]}$ for each i = 1, ..., m. If $F_i \in MF(\mathcal{S}_i)$, then $F = F_i \in MF(\mathcal{K})$. Otherwise F would not be minimal, since $F_i \subseteq F$ is a missing face.

On the other hand, suppose that $F_i \in S_i$ for all *i*. Denote by $\sigma = \{i \in [m] \mid F \mid i \neq \emptyset\}$. Firstly, $\sigma \notin \mathcal{K}$, as otherwise $F \in \mathcal{K}$ since $F = \bigsqcup_{i \in \sigma} F_i$, where $F_i \in S_i$ for all *i*. For all $i \in \sigma$, F_i is a non-face of \mathcal{T}_i . Otherwise, by removing vertices from F_i , we obtain a smaller non-face of $(\underline{S}, \underline{\mathcal{T}})^{*\mathcal{K}}$. For such $i, F_i \in MF(\mathcal{T}_i)$. It follows that $\sigma \in MF(\mathcal{K})$, as otherwise we restrict to a minimal missing face $\tau \in MF(\mathcal{K})$ with $\tau \subseteq \sigma$, and $\bigsqcup_{i \in \tau} F_i$ is a missing face of $(\underline{S}, \underline{\mathcal{T}})^{*\mathcal{K}}$. Finally, for all i, F_i is a minimal missing face of \mathcal{T}_i . This follows by observing that otherwise for $\hat{F}_i \subsetneq F_i$ with $\hat{F}_i \in MF(\mathcal{S}_i)$, $(F - F_i) \sqcup \hat{F}_i \in (\underline{S}, \underline{\mathcal{T}})^{*\mathcal{K}}$, which is a contradiction. \Box

Proof of Proposition 5.2. By Proposition 5.3, the minimal missing faces of the composition complex $\mathcal{K}(\mathcal{T}_1, \ldots, \mathcal{T}_m)$ are

$$MF(\mathcal{K}(\mathcal{T}_1,\ldots,\mathcal{T}_m)) = \left\{ \bigsqcup_{i \in L} J_i \mid L \in MF(\mathcal{K}), J_i \in MF(\mathcal{T}_i) \right\}$$

so that

$$MF(\operatorname{sk}^{k-3}\Delta^{k-1}(\partial\Delta[I_1],\ldots,\partial\Delta[I_k])) = \{\{[m] \setminus \{i_1,\ldots,i_{n_i}\}\} \mid i=1,\ldots,k\}. \quad \Box$$

Since a simplicial complex is determined by the set of minimal missing faces, \mathcal{K}_{Π} is the simplicial complex on [m] given by

$$MF(\mathcal{K}_{\Pi}) = \{[m] \setminus P_i \mid i = 1, \dots, k\}.$$

It follows that the Alexander dual of \mathcal{K}_{Π} is given by

(30)
$$\widehat{\mathcal{K}_{\Pi}} = \bigsqcup_{i=1}^{k} \Delta[P_i]$$

We use this description of the Alexander dual to decompose \mathcal{K}_{Π} as the union of subcomplexes.

Proposition 5.4. Let $\Pi = \{P_1, \ldots, P_k\}$ be a k-partition of [m] and denote $P_i = \{i_1, \ldots, i_{p_i}\}$ and $[m] \setminus P_i = \{j_1, \ldots, j_{q_i}\}$. Then

(31)
$$\mathcal{K}_{\Pi} = \bigcup_{i=1}^{k} \partial \Delta \langle \partial \Delta[j_1, \dots, j_{q_i}], i_1, \dots, i_{p_i} \rangle.$$

Proof. Two finite simplicial complexes are isomorphic if and only if their Alexander duals are isomorphic. By (30), the Alexander dual of \mathcal{K}_{Π} has maximal faces $\bigsqcup_{i=1}^{k} \Delta[i_1, \ldots, i_{p_i}]$. On the other hand, if $\mathcal{K}^i = \partial \Delta \langle \partial \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle$, then by Proposition 5.3, the maximal faces $\widehat{\mathcal{K}^i}_{max}$ of $\widehat{\mathcal{K}^i}$ are given by

$$\{[m] \setminus \{j_1, \dots, j_{q_i}\} \sqcup \{[m] \setminus \{j, i_1, \dots, i_{p_i}\} \mid j \in Q_i\} = \{i_1, \dots, i_{p_i}\} \sqcup \{Q_i \setminus j \mid j \in Q_i\}$$

where $Q_i = [m] \setminus P_i = \{j_1, \dots, j_{q_i}\}$. Then $\widehat{\prod_{k=1}^{k} r_i} - \bigwedge^k \widehat{r_i} - \bigwedge^k (\widehat{r_i}) - \sum_{k=1}^{k} \widehat{r_i} - \sum_{k=1}^{k}$

$$\bigcup_{i=1}^{k} \mathcal{K}^{i} = \bigcap_{i=1}^{k} \widehat{\mathcal{K}^{i}} = \bigcap_{i=1}^{k} \left(\widehat{\mathcal{K}^{i}}_{max} \right) = \bigsqcup_{i=1}^{k} \Delta[i_{1}, \dots, i_{p_{i}}]$$

and the result follows.

We illustrate the three different decompositions of the complex \mathcal{K}_{Π} .

Example 5.5. Let $\Pi = \{\{1\}, \{2,3\}, \{4\}\}$ be a 3-partition of [4]. The complex $\mathcal{K}_{\Pi} = \mathrm{sk}^0 \Delta^2(\circ[1], \partial \Delta[2,3], \circ[4])$ is given by

$$\mathcal{K}_{\Pi} = \{1\} * \partial \Delta[2,3] \cup \partial \Delta[2,3] * \{4\} \cup \Delta[2,3].$$

This decomposition is shown in the following figure. The minimal missing faces of




FIGURE 3. Decomposition of the identity complex $\mathcal{K}_{\{1\},\{2,3\},\{4\}\}}$.

 \mathcal{K}_{Π} are $Q_1 = \{2, 3, 4\}$, $Q_2 = \{1, 4\}$, and $Q_3 = \{1, 2, 3\}$. Finally, since the complex \mathcal{K}_{Π} is also given by $\partial \Delta \langle \partial \Delta[1, 4], 2, 3 \rangle$, we obtain the decomposition

$$\mathcal{K}_{\Pi} = \partial \Delta \langle \partial \Delta[2,3,4], 1 \rangle \cup \partial \Delta \langle \partial \Delta[1,4], 2,3 \rangle \cup \partial \Delta \langle \partial \Delta[1,2,3], 4 \rangle.$$

5.1. Relations among higher Whitehead maps. Let $\Pi = \{P_1, \ldots, P_k\}$ be a *k*-partition of [m] and let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps for $i = 1, \ldots, m$. By Proposition 5.4, $\mathcal{K}_{\Pi} = \bigcup_{i=1}^k \partial \Delta \langle \partial \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle$. Therefore, the space $(\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}$ contains the codomains of the higher Whitehead maps

(32)
$$h_w \left(h_w \left(f_{j_1}, \dots, f_{j_{q_i}} \right), f_{i_1}, \dots, f_{i_{p_i}} \right) :$$
$$\Sigma^{p_i} \left(\Sigma^{q_i - 2} X_{j_1} \wedge \dots \wedge X_{j_{q_i}} \right) \wedge X_{i_1} \wedge \dots \wedge X_{i_{p_i}} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}^i}$$

for i = 1, ..., k.

We generalise the relation of Hardie (27) to relations among the higher Whitehead maps (32) in polyhedral products over the identity complex \mathcal{K}_{Π} .

Theorem 5.6. For i = 1, ..., m, let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps. Let $\Pi = \{P_1, ..., P_k\}$ be a k-partition of [m] for $k \ge 3$ and denote by $P_i = \{i_1, ..., i_{p_i}\}$ and $Q_i = [m] \setminus P_i = \{j_1, ..., j_{q_i}\}$. If X_i is a suspension for each i = 1, ..., m, then

(33)
$$\sum_{i=1}^{k} h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) \circ \sigma_{i} = 0$$

in $[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}]$, where

$$\sigma_i: \Sigma^{m-2} X_1 \wedge \cdots \wedge X_m \longrightarrow \Sigma^{p_i} (\Sigma^{q_i-2} X_{j_1} \wedge \cdots \wedge X_{j_{q_i}}) \wedge X_{i_1} \wedge \cdots \wedge X_{i_{p_i}}$$

is the restriction of the coordinate permutation

$$CX_1 \times \cdots \times CX_m \longrightarrow CX_{j_1} \times \cdots \times CX_{j_{q_i}} \times CX_{i_1} \times \cdots \times CX_{i_{p_i}}.$$

We delay the proof of Theorem 5.6 to Section 7. Now, we illustrate how Theorem 5.6 generalises Hardie's identity to higher Whitehead maps of non-spherical maps and obtain examples of new relations among higher Whitehead maps.

Example 5.7. For i = 1, ..., m, let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps such that X_i is a suspension.

(i) Let $\Pi = \{\{1\}, \{2\}, \{3\}\}$. Then $Q_1 = \{2, 3\}, Q_2 = \{1, 3\}$ and $Q_3 = \{1, 2\}$, and $\mathcal{K}_{\Pi} = \bullet_{[3]}$. We obtain the relation

$$h_w^{\mathcal{K}_{\Pi}}(h_w(f_2, f_3), f_1) \circ \sigma_1 + h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, f_3), f_2) \circ \sigma_2 + h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, f_2), f_3) \circ \sigma_3 = 0$$

in $[\Sigma X_1 \wedge X_2 \wedge X_3, Y_1 \vee Y_2 \vee Y_3]$. This recovers the generalised Jacobi identity, see [4].

(ii) Generalising the previous example, let $\Pi = \{\{1\}, \ldots, \{m\}\}$. This implies that $Q_i = \{1, \ldots, \hat{i}, \ldots, m\}$ for $i = 1, \ldots, m$ and $\mathcal{K}_{\Pi} = \operatorname{sk}^{m-3} \Delta^{m-1}$, giving rise to the relation

$$h_{w}^{\mathcal{K}_{\Pi}}\left(h_{w}\left(f_{2},\ldots,f_{m}\right),f_{1}\right)\circ\sigma_{1}+\cdots+h_{w}^{\mathcal{K}_{\Pi}}\left(h_{w}\left(f_{1},\ldots,f_{m-1}\right),f_{m}\right)\circ\sigma_{m}=0$$

in $\left[\Sigma^{m-2}X_{1}\wedge\cdots\wedge X_{m},\left(\underline{Y},\underline{*}\right)^{\mathrm{sk}^{m-3}}\Delta^{m-1}\right]$, generalising Hardie's identity (27) to non-spherical maps.

(iii) Define a 3-partition $\Pi = \{\{1\}, \{2,3\}, \{4\}\}\)$ of the vertex set [4]. In this case $Q_1 = \{2,3,4\}, Q_2 = \{1,4\}\)$ and $Q_3 = \{1,2,3\}\)$ and \mathcal{K}_{Π} is the simplicial complex shown in Example 5.5. Then

$$\begin{split} h_w^{\mathcal{K}_{\Pi}}(h_w(f_2, f_3, f_4), f_1) \circ \sigma_1 + h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, f_4), f_2, f_3) \circ \sigma_2 \\ &+ h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, f_2, f_3), f_4) \circ \sigma_3 = 0 \end{split}$$

in $[\Sigma^2 X_1 \wedge X_2 \wedge X_3 \wedge X_4, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}]$. In contrast to Hardie's identity (27), the summands of this relation include both 2-ary and 3-ary higher Whitehead maps, providing new forms of relations.

When the maps f_i are spherical, the permutation maps σ_i are degree maps. Recall the definition of the Koszul sign $\varepsilon(\sigma)$ of a permutation σ as given in Proposition 2.5.

Corollary 5.8. Let $f_i \in \pi_{r_i}(Y_i)$ for i = 1, ..., m. Let $\Pi = \{P_1, ..., P_k\}$ be a k-partition of [m] for $k \ge 3$ and denote $P_i = \{i_1, ..., i_{p_i}\}$ and $J_i = [m] \setminus P_i = \{j_1, ..., j_{q_i}\}$. If $r_i \ge 2$ for each i = 1, ..., m, then there is a relation

$$\sum_{i=1}^{k} \varepsilon(\sigma_i) h_w^{\mathcal{K}_{\Pi}} \left(h_w \left(f_{j_1}, \dots, f_{j_{q_i}} \right), f_{i_1}, \dots, f_{i_{p_i}} \right) = 0$$

22

in $\pi_{r_1+\dots+r_m-2}\left((\underline{Y},\underline{*})^{\mathcal{K}_{\Pi}}\right)$, where $\varepsilon(\sigma_i)$ is the Koszul sign of the permutation $\sigma_i:(1,\dots,m)\mapsto (j_1,\dots,j_{q_i},i_1,\dots,i_{p_i})$.

Proof. The map $\sigma_i: S^{r_1 + \dots + r_m - 2} \longrightarrow S^{r_1 + \dots + r_m - 2}$ from the statement of Theorem 5.6 has degree $\varepsilon(\sigma_i)$ and the result follows.

Example 5.9. In some cases, $\varepsilon(\rho)$ can be calculated explicitly. We revisit the relations from Example 5.7 when $f_i: S^{r_i} \longrightarrow Y_i$, with $r_i \ge 2$ for $i = 1, \ldots, m$.

(i) Let $\Pi = \{\{1\}, \ldots, \{m\}\}$. Denote by ι_i the inclusion $Y_i \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}$ and κ_i the inclusion $FW(Y_1, \ldots, Y_{i-1}, Y_{i+1}, \ldots, Y_m) \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}$. We recover relation (27) of Hardie,

$$\sum_{i=1}^{m} (-1)^{p_i(p_{i+1}+\dots+p_m)} [\theta_i h_w(f_1,\dots,f_{i-1},f_{i+1},\dots,f_m),\psi_i f_i] = 0$$

in $\pi_{p_1+\dots+p_m-2}\left((\underline{Y},\underline{*})^{\mathcal{K}_{\Pi}}\right)$. Moreover, if k=3, let $g_j = \iota_j \circ f_j$ for j=1,2,3. Multiplying by $(-1)^{p_1p_3}$ and applying Proposition 2.6, we recover the graded Jacobi identity for Whitehead products

$$(-1)^{p_1p_2}[[g_2,g_3],g_1] + (-1)^{p_2p_3}[[g_1,g_3],g_2] + (-1)^{p_1p_3}[[g_1,g_2],g_3] = 0.$$

(ii) for $\Pi = \{\{1\}, \{2, 3\}, \{4\}\}, \{4\}\}$, there is a relation

$$(-1)^{p_1(p_2+p_3)} h_w^{\mathcal{K}_{\Pi}}(h_w(f_2, f_3, f_4), f_1) + (-1)^{(p_2+p_3)p_4} h_w^{\mathcal{K}_{\Pi}}(h_w(f_4, f_1), f_2, f_3) + (-1)^{p_1p_4} h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, f_2, f_3), f_4) = 0$$

in $\pi_{p_1+\dots+p_4-2}\left((\underline{Y},\underline{*})^{\mathcal{K}_{\Pi}}\right)$.

As a further consequence of Theorem 5.6, we expand the family of simplicial complexes in which we can identify relations among higher Whitehead maps.

Corollary 5.10. For i = 1, ..., m, let S_i be a simplicial complex on $[l_i]$ and let $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{S_i}$ be maps. Let $\Pi = \{P_1, ..., P_k\}$ be a k-partition of [m] for $k \ge 3$ and denote by $P_i = \{i_1, ..., i_{p_i}\}$ and $Q_i = [m] \setminus P_i = \{j_1, ..., j_{q_i}\}$. Suppose that \mathcal{L} is a simplicial complex containing $\mathcal{K}_{\Pi} \langle S_1, ..., S_m \rangle$. Then if X_i is a suspension for each i = 1, ..., m,

$$\sum_{i=1}^{k} h_{w}^{\mathcal{L}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) \circ \sigma_{i} = 0$$

in $\left[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{\mathcal{L}}\right]$, where

$$\sigma_i: \Sigma^{m-2} X_1 \wedge \dots \wedge X_m \longrightarrow \Sigma^{p_i} (\Sigma^{q_i-2} X_{j_1} \wedge \dots \wedge X_{j_{q_i}}) \wedge X_{i_1} \wedge \dots \wedge X_{i_{p_i}}$$

is the restriction of the coordinate permutation

 $CX_1 \times \cdots \times CX_m \longrightarrow CX_{j_1} \times \cdots \times CX_{j_{q_i}} \times CX_{i_1} \times \cdots \times CX_{i_{p_i}}.$

Example 5.11. In some cases, the summands in relation (33) can be shown to be non-trivial. Suppose that for $i = 1, ..., m, f_i: S^2 \longrightarrow \mathbb{C}P^{\infty}$ is the inclusion of the bottom cell. Let Π be a k-partition of [m]. By Corollary 5.8

(34)
$$\sum_{i=1}^{k} \varepsilon(\rho) h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) = 0.$$

We show that none of the maps in (34) are null-homotopic. Abramyan and Panov showed in [2, Theorem 5.2] that the map $h_w^{\mathcal{K}_{\Pi}} \left(h_w \left(f_{j_1}, \ldots, f_{j_{q_i}} \right), f_{i_1}, \ldots, f_{i_{p_i}} \right)$ is null-homotopic if and only if $\Delta \langle \partial \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle \subseteq \mathcal{K}_{\Pi}$. The complex $\Delta \langle \partial \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle \subseteq \mathcal{K}_{\Pi}$ contains the simplex $Q_i = [m] \setminus P_i$. On the other hand, by the definition of the complex \mathcal{K}_{Π} associated to the partition Π , Q_i is a missing face of \mathcal{K}_{Π} . Therefore, $\Delta \langle \partial \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle$ is not a subcomplex of \mathcal{K}_{Π} . Hence each term in the relation

$$\sum_{i=1}^{\kappa} \varepsilon(\rho) h_w^{\mathcal{K}_{\Pi}} \left(h_w \left(f_{j_1}, \dots, f_{j_{q_i}} \right), f_{i_1}, \dots, f_{i_{p_i}} \right) = 0$$

is non-trivial.

6. Relations among folded higher Whitehead maps

Folded higher Whitehead maps provide a combinatorial framework to study elements of higher Whitehead products with repeated factors. By propagating the relations among higher Whitehead maps given in Theorem 5.6, we study relations among folded higher Whitehead maps.

Relations among folded higher Whitehead maps have been studied rationally by Zhuravleva [15]. Let $\mathcal{K} = \partial \Delta \langle \partial \Delta \langle 1, 2, 3 \rangle, 4, 5 \rangle$ and for $i = 1, \ldots, m$, let $u_i: S^1 \longrightarrow \Omega \mathbb{C}P^{\infty}$ be the adjoint of the inclusion $\mu_i: S^2 \longrightarrow \mathbb{C}P^{\infty}$ of the bottom cell. Zhuravleva showed that the relation

$$(35) \qquad \qquad [[u_1, u_2, u_3], [u_1, u_4, u_5]] + [[[u_1, u_2, u_3], u_4, u_5], u_1] = 0$$

holds in $\pi_8(\Omega DJ_{\mathcal{K}}) \otimes \mathbb{Q}$. We improve this result by showing that relation (35) holds integrally in $\pi_*(\Omega DJ_{\mathcal{K}})$.

6.1. Folds of identity complexes. For i = 1, ..., m, let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps. Let $\Pi = \{P_1, ..., P_k\}$ be a k-partition of [m], and denote by $P_i = \{i_1, ..., i_{p_i}\}$, and $[m] \setminus P_i = \{j_1, ..., j_{q_i}\}$. By Theorem 5.6,

(36)
$$\sum_{i=1}^{k} h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) \circ \sigma_{i} = 0$$

in $[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}]$. Let $\psi: I \longrightarrow J$ be a fold of \mathcal{K}_{Π} and suppose that Y_j is an associative *H*-space for each $j \in J$, and that $Y_i = Y_j$ for every $i \in I_j =$

 $\psi^{-1}(j)$. By composing each summand in (36) with the fold $\nabla_{(I,J)}: (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}} \longrightarrow (\underline{Y}, \underline{*})^{(\mathcal{K}_{\Pi})_{\nabla(I,J)}}$, we obtain the relation

(37)
$$\sum_{i=1}^{k} \nabla_{(I,J)} h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) \circ \sigma_{i} = 0$$

among folded higher Whitehead maps in $\left[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{(\mathcal{K}_{\Pi})_{\nabla(I,J)}}\right]$. We analyse the combinatorial properties of the folded complex $(\mathcal{K}_{\Pi})_{\nabla(I,J)}$ for various partitions Π and folds $\psi: I \longrightarrow J$, which govern the form of relation (37).

We determine, in terms of the partition Π and the fold $\psi: I \longrightarrow J$, which terms of relation (37) are null-homotopic after folding. By Proposition 4.10, the map $\nabla_{(I,J)}h_w^{\mathcal{K}_{\Pi}}\left(h_w\left(f_{j_1},\ldots,f_{j_{q_i}}\right),f_{i_1},\ldots,f_{i_{p_i}}\right)$ is null-homotopic if $h_w^{\mathcal{L}}\left(h_w\left(f_{j_1},\ldots,f_{j_{q_i}}\right),f_{i_1},\ldots,f_{i_{p_i}}\right)$ is null-homotopic, where

(38)
$$\mathcal{L} = (\mathcal{K}_{\Pi})_{\nabla(I,J)} \langle \Delta[\{k_1\} \sqcup I_{k_1}], \dots, \Delta[\{k_l\} \sqcup I_{k_l}] \rangle$$

where $I_k = \emptyset$ if $k \notin J$, and $\{k_1, \ldots, k_l\}$ is the vertex set of $(\mathcal{K}_{\Pi})_{\nabla(I,J)}$.

By first describing the folded identity complex $(\mathcal{K}_{\Pi})_{\nabla(I,J)}$ in terms of the partition Π and the fold $\psi: I \longrightarrow J$, we are able to express the complex \mathcal{L} associated to \mathcal{K}_{Π} in terms of Π and ψ . Applying Proposition 4.10, we identify the null-homotopic terms in relation (37).

Lemma 6.1. Let $\Pi = \{P_1, \ldots, P_k\}$ be a k-partition of [m]. Then the following hold: (i) if $I \cup J \subseteq P_l$ for some $l \in \{1, \ldots, k\}$, then

$$(\mathcal{K}_{\Pi})_{\nabla(I,J)} = \partial \Delta[[m] \setminus P_l] * \Delta[P_l \setminus I];$$

(ii) if |I| = |J| = 1 and $I \subseteq P_i$ and $J \subseteq P_j$ for $i \neq j$, then

$$(\mathcal{K}_{\Pi})_{\nabla(I,J)} = \partial \Delta[[m] \setminus I];$$

(iii) otherwise, $(\mathcal{K}_{\Pi})_{\nabla(I,J)} = \Delta[[m] \setminus I].$

Proof. We first prove (ii). Let $(\underline{\Delta}, \underline{\partial}\underline{\Delta})$ denote the tuple of pairs $((\Delta[P_1], \partial\Delta[P_1]), \dots, (\Delta[P_k], \partial\Delta[P_k]))$. By definition, $\mathcal{K}_{\Pi} = (\underline{\Delta}, \underline{\partial}\underline{\Delta})^{*sk^{k-3}\Delta[1,\dots,k]}$. For $i \neq j$, this polyhedral join decomposes into the union

$$\mathcal{K}_{\Pi} = \Delta[P_i] * \Delta[P_j] * \left((\underline{\Delta}, \underline{\partial}\underline{\Delta})^{*sk^{k-5}\Delta[[k] \setminus \{i,j\}]} \right)$$
$$\cup \left(\Delta[P_i] * \partial\Delta[P_j] \cup \partial\Delta[P_i] * \Delta[P_j] \right) * \left((\underline{\Delta}, \underline{\partial}\underline{\Delta})^{*\partial\Delta[[k] \setminus \{i,j\}]} \right)$$
$$\cup \partial\Delta[P_i] * \partial\Delta[P_j] * \left((\underline{\Delta}, \underline{\partial}\underline{\Delta})^{*\Delta[[k] \setminus \{i,j\}]} \right).$$

Suppose that $I = \{u\}$ and $J = \{v\}$ such that $u \in P_i$ and $v \in P_j$. Then

$$\nabla_{(u,v)}(\Delta[P_i] * \Delta[P_j]) = \Delta[P_i \sqcup P_j \setminus \{u\}]$$
$$\nabla_{(u,v)}(\Delta[P_i] * \partial \Delta[P_j]) = \Delta[P_i \sqcup P_j \setminus \{u\}]$$
$$\nabla_{(u,v)}(\partial \Delta[P_i] * \Delta[P_j]) = \Delta[P_i \sqcup P_j \setminus \{u\}]$$
$$\nabla_{(u,v)}(\partial \Delta[P_i] * \partial \Delta[P_j]) = \partial \Delta[P_i \sqcup P_j \setminus \{u\}]$$

Since $sk^{k-5}\Delta[[k] \setminus \{i, j\}]$ is a subcomplex of $\partial\Delta[[k] \setminus \{i, j\}]$, it follows that

$$\begin{aligned} (\mathcal{K}_{\Pi})_{\nabla(I,J)} &= \Delta[P_i \sqcup P_j \setminus \{u\}] * \left((\underline{\Delta}, \underline{\partial \Delta})^{*\partial \Delta[[k] \setminus \{i,j\}]} \right) \\ &\cup \partial \Delta[P_i \sqcup P_j \setminus \{u\}] * \left((\underline{\Delta}, \underline{\partial \Delta})^{*\Delta[[k] \setminus \{i,j\}]} \right) \\ &= ((\Delta[P_1], \partial \Delta[P_1]), \dots, (\Delta[P_i \sqcup P_j \setminus \{u\}], \partial \Delta[P_i \sqcup P_j \setminus \{u\}]), \dots \\ &\dots, (\Delta[P_k], \partial \Delta[P_k]))^{*\partial \Delta[[k] \setminus i]} \\ &= \partial \Delta[[m] \setminus \{u\}] \end{aligned}$$

proving (ii).

We now establish (i) and (iii). Assume that $u, v \in P_l$. The folding sends $\Delta[P_l] \mapsto \Delta[P_l \setminus \{u\}]$ and $\partial \Delta[P_l] \mapsto \Delta[P_l \setminus \{u\}]$. Then since

$$\mathcal{K}_{\Pi} = \Delta[P_l] * \left((\underline{\Delta}, \underline{\partial \Delta})^{*sk^{k-4}\Delta[[k] \setminus \{l\}]} \right) \cup \partial \Delta[P_l] * \left((\underline{\Delta}, \underline{\partial \Delta})^{*\partial \Delta[[k] \setminus \{l\}]} \right)$$

we obtain

$$(\mathcal{K}_{\Pi})_{\nabla(I,J)} = \Delta[P_l \setminus \{u\}] * \left((\underline{\Delta}, \underline{\partial \Delta})^{*\partial \Delta[[k] \setminus \{l\}]}\right) = \Delta[P_l \setminus \{u\}] * \partial \Delta[[m] \setminus P_l].$$

Next, for any I and J with $u \in I$ and $v \in J$, by Proposition 4.3,

$$(\mathcal{K}_{\Pi})_{\nabla(I,J)} = \left((\mathcal{K}_{\Pi})_{\nabla(u,v)} \right)_{\nabla(I \setminus \{u\}, J \setminus \{v\})} = \left(\Delta[P_l \setminus \{u\}] * \partial \Delta[[m] \setminus P_l] \right)_{\nabla(I \setminus \{u\}, J \setminus \{v\})}$$

Claim (i) then follows since for $I, J \subseteq P_l$, $(\Delta[P_l \setminus \{u\}] * \partial \Delta[[m] \setminus P_l])_{\nabla(I \setminus \{u\}, J \setminus \{v\})} = \Delta[P_l \setminus I] * \partial \Delta[[m] \setminus P_l].$

Alternatively, claim (iii) follows since for any fold not satisfying the hypotheses of (i) or (ii), there is $u' \in I$ with $u' \in P_{l'}$ for $l' \neq l$. Then $(\Delta[P_l \setminus \{u\}] * \partial \Delta[[m] \setminus P_l])_{\nabla(I \setminus \{u\}, J \setminus \{v\})} = \Delta[[m] \setminus I]$ since any fold of $\partial \Delta[[m] \setminus P_l]$ is a simplex. \Box

Lemma 6.2. Let \mathcal{L} denote the complex in (38). Then the following hold:

(i) if $I \cup J \subseteq P_l$ for some $l \in \{1, \ldots, k\}$, then

$$\mathcal{L} = \partial \Delta[[m] \setminus P_l] * \Delta[P_l];$$

- (ii) if |I| = |J| = 1 and $I \subseteq P_i$ and $J \subseteq P_j$ for $i \neq j$, then \mathcal{L} has minimal missing faces $[m] \setminus I$ and $[m] \setminus J$;
- (iii) otherwise, $\mathcal{L} = \Delta[[m]]$.

Proof. By Proposition 5.3,

$$MF(\mathcal{L}) = \left\{ \bigsqcup_{i \in \kappa} \tau_i \mid \kappa \in MF\left((\mathcal{K}_{\Pi})_{\nabla(I,J)} \right), \tau_i \in \Delta[\{k_i\} \sqcup I_{k_i}] \right\}.$$

Statement (i) then follows from Lemma 6.1 since the missing face of $\partial \Delta[[m] \setminus P_l] * \Delta[P_l \setminus I]$ is not supported by any vertex in J. Similarly, statements (ii) and (iii) also follow from Lemma 6.1.

Therefore the only folds of \mathcal{K}_{Π} that do not trivialise the map $\nabla_{(I,J)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_{j_1},\ldots,f_{j_{q_i}}),f_{i_1},\ldots,f_{i_{p_i}})$ are those identifying some i_s with some j_r , or vice-versa. The relations arising from folds of this form are described in the following theorem.

Theorem 6.3. Let $\Pi = \{P_1, \ldots, P_k\}$ be a k-partition of [m] and let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps for $i = 1, \ldots, m$, where each X_i is a suspension. Fix $u, v \in [k]$ with $u \neq v$ and write $P_u = \{i_1, \ldots, i_{p_i}\}$ and $P_v = \{j_1, \ldots, j_{p_j}\}$. Let $i \in P_u$ and $j \in P_v$ and let $\psi: \{i\} \longrightarrow \{j\}$ be a fold of \mathcal{K}_{Π} . If $Y_i = Y_j$ is an H-space, then there is a relation

$$\begin{aligned} \nabla_{(i,j)} h_w^{\mathcal{K}_{\Pi}}(h_w(f_{i'_1}, \dots, f_{i'_{q_i}}), f_{i_1}, \dots, f_{i_{p_i}}) \circ \sigma_i \\ &+ \nabla_{(i,j)} h_w^{\mathcal{K}_{\Pi}}(h_w(f_{j'_1}, \dots, f_{j'_{q_j}}), f_{j_1}, \dots, f_{j_{p_j}}) \circ \sigma_j = 0 \\ in \left[\Sigma^{m-2} X_1 \wedge \dots \wedge X_m, (\underline{Y}, \underline{*})^{\partial \Delta[1, \dots, i-1, i+1, \dots, m]} \right], where \ [m] \setminus P_u = \{i'_1, \dots, i'_{q_i}\} \ and \\ [m] \setminus P_v = \{j'_1, \dots, j'_{q_i}\}. \end{aligned}$$

Proof. We show that if either I or J contain more than one element, or if I and J are singletons both contained in one of $P_i = \{i_1, \ldots, i_{p_i}\}$ or $[m] \setminus P_i = \{j_1, \ldots, j_{q_i}\}$, then the folded higher Whitehead map

$$H_{i} = \nabla_{(I,J)} h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right)$$

is null-homotopic.

Suppose that at least one of I and J contains more than one element. Firstly, if $I \sqcup J$ is not contained in some P_r for $r \in \{1, \ldots, k\}$, then by Lemma 6.2(iii), $\mathcal{L} = \Delta[m]$. By Proposition 4.10, it follows that the map H_i is null-homotopic for each $i = 1, \ldots, k$.

Secondly, if $I \sqcup J \subseteq P_r$ for some $r = 1, \ldots, k$, then by Lemma 6.2(i), $\mathcal{L} = \partial \Delta[[m] \setminus P_r] * \Delta[P_r]$. Then H_r is null-homotopic, and furthermore for $i \neq r$, since $\partial \Delta \langle \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle \subseteq \mathcal{L}$, the map H_i is also null-homotopic for $i \neq r$.

Now suppose that |I| = |J| = 1, and that $I = \{u\} \in P_s$ and $J = \{v\} \in P_t$ for $s \neq t$. By Lemma 6.2(ii), \mathcal{L} has minimal missing faces $[m] \setminus \{u\}$ and $[m] \setminus \{v\}$. Therefore if $u = j_l$ and $v = j_k$, then $\partial \Delta \langle \Delta[j_1, \ldots, j_{q_i}], i_1, \ldots, i_{p_i} \rangle \subseteq \mathcal{L}$, and hence H_i is nullhomotopic. Observe that since $s \neq t$, it is not possible for $u = i_l$ and $v = i_k$ in this case.

The only remaining possibility is that |I| = |J| = 1 and that $I \subseteq P_u$ and $J \subseteq P_v$ for $u \neq v$, and so the theorem follows.

A special case of Theorem 6.3 gives an example of when a folded *n*-ary higher Whitehead map is identified with a folded 2-ary higher Whitehead map.

Corollary 6.4. Let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps for i = 1, ..., m, where each X_i is a suspension. Fix $i, j \in [m]$ with $i \neq j$ and let $\{i_1, \ldots, i_p\} \subseteq [m] \setminus \{i, j\}$ be any non-empty proper subset. If $Y_i = Y_j$ is an H-space, then there is a relation

$$\nabla_{(i,j)} h_w^{\mathcal{K}_{\Pi}}(h_w(f_{j_1}, \dots, f_{j_q}, f_i), f_{i_1}, \dots, f_{i_p}, f_j) \circ \sigma + \nabla_{(i,j)} h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_j, \dots, f_m), f_i) \circ \tau = 0$$

in $[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{\partial \Delta[1, \dots, i-1, i+1, \dots, m]}]$, where $[m] \setminus \{i, j, i_1, \dots, i_p\} = \{j_1, \dots, j_q\}$ and σ and τ are appropriate permutation maps.

Proof. The result follows by applying Theorem 6.3 to the 3-partition

$$\Pi = \{\{i\}, \{j, i_1, \dots, i_p\}, \{j_1, \dots, j_q\}\}\$$

of [m].

We demonstrate Corollary 6.4 by identifying new relations in homotopy groups.

Example 6.5. Let $\Pi = \{\{1\}, \{2, \ldots, m-1\}, \{m\}\}$ and let $f_i: S^{p_i} \longrightarrow Y_i, p_i \ge 2$, be maps for $i = 1, \ldots, m$ with $f_1 = f_m$ and $p_1 = p_m$ even. Consider the maps $h_w^{\mathcal{K}_{\Pi}}(h_w(f_1, \ldots, f_{m-1}), f_m)$ and $h_w^{\mathcal{K}_{\Pi}}(h_w(f_2, \ldots, f_m), f_1)$. Applying the fold $\psi: \{m\} \longrightarrow \{1\}$ of \mathcal{K}_{Π} and using Propositions 2.5 and 4.5,

$$\nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,\ldots,f_{m-1}),f_m) = \nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_2,\ldots,f_m),f_1).$$

On the other hand, by Corollary 6.4, we obtain

$$2\nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,\ldots,f_{m-1}),f_m) = \nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,\ldots,f_{m-1}),f_m) + \nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_2,\ldots,f_m),f_1) = 0$$

in $\pi_{p_1+\dots+p_m-2}((\underline{Y},\underline{*})^{\partial\Delta})$. Therefore our methods allow us to study, via higher Whitehead maps, elements of homotopy groups which cannot be seen with rational methods. We study when the map $\nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,\dots,f_{m-1}),f_m)$ is 2-torsion, that is, when it is not null-homotopic. In general, this depends on the properties of the space Y_i . To demonstrate this, let $p_i = 2$ for $i = 1, \dots, m$, so that $f_i: S^2 \longrightarrow Y_i$.

We first suppose that $Y_i = S^2$ and $f_i: S^2 \longrightarrow S^2$ are identity maps for $i = 1, \ldots, m$. While S^2 is not an *H*-space, we can still apply the fold ψ to the relation

$$h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,\ldots,f_{m-1}),f_m) + h_w^{\mathcal{K}_{\Pi}}(h_w(f_2,\ldots,f_m),f_1) \circ + h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,f_m),f_2,\ldots,f_{m-1}) = 0$$

to obtain that

to obtain that

$$2\nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,\ldots,f_{m-1}),f_m) + \nabla_{(m,1)}h_w^{\mathcal{K}_{\Pi}}(h_w(f_1,f_m),f_2,\ldots,f_{m-1}) = 0.$$

By naturality of the higher Whitehead map and Proposition 4.5, we have

$$\nabla_{(m,1)}h_w(h_w(f_1, f_m), f_2, \dots, f_{m-1}) = h_w([f_1, f_1], f_2, \dots, f_{m-1})$$

Since $[f_1, f_1] = 2\eta$, where $\eta \in \pi_3(S^2)$ is the Hopf map, we obtain from linearity of the higher Whitehead map that

(39)
$$\nabla_{(m,1)}h_w(h_w(f_1,\ldots,f_{m-1}),f_m) = -h_w(\eta,f_2,\ldots,f_{m-1}).$$

Therefore the folded higher Whitehead map $\nabla_{(m,1)}h_w(h_w(f_1,\ldots,f_{m-1}),f_m)$ is not null-homotopic since the higher Whitehead map $h_w(\eta, f_2,\ldots,f_{m-1})$ is not null-homotopic, see Example 2.8.

Now suppose that $Y_i = \mathbb{C}P^{\infty}$ and $f_i: S^2 \longrightarrow \mathbb{C}P^{\infty}$ is the inclusion of the bottom cell for i = 1, ..., m. Applying the map $(S^2, *)^{\partial \Delta} \longrightarrow (\mathbb{C}P^{\infty}, *)^{\partial \Delta}$ to each side of (39) trivialises the map $h_w(\eta, f_2, ..., f_{m-1})$, since $\eta = 0$ in $\pi_3(\mathbb{C}P^{\infty})$. Therefore by naturality of the higher Whitehead map, the folded map $\nabla_{(m,1)}h_w(h_w(f_1, ..., f_{m-1}), f_m)$ is trivial in $\pi_{2m-2}((\mathbb{C}P^{\infty}, *)^{\partial \Delta})$. Alternatively, if $Y_i = \Omega S^3$ and $f_i: S^2 \longrightarrow \Omega S^3$ is the suspension for $i = 1, \ldots, m$, then applying the map $(S^2, *)^{\partial \Delta} \longrightarrow (\Omega S^3, *)^{\partial \Delta}$ to (39) does not trivialise the map $h_w(\eta, f_2, \ldots, f_{m-1})$ since the composite $S^3 \xrightarrow{\eta} S^2 \longrightarrow \Omega \Sigma S^2$ is adjoint to $\Sigma \eta$, which generates $\pi_4(S^3)$. Therefore, the folded map $\nabla_{(m,1)}h_w(h_w(f_1, \ldots, f_{m-1}), f_m)$ is a 2-torsion element in $\pi_{2m-2}((\Omega S^3, *)^{\partial \Delta})$.

6.2. Folding and substitution. By introducing richer combinatorial structure into the identity complex \mathcal{K}_{Π} before folding, we obtain a broader family of relations. Suppose that $\mathcal{K}_1, \ldots, \mathcal{K}_m$ are simplicial complexes on $[l_1], \ldots, [l_m]$, respectively. Denote by $\mathcal{S}_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \ldots, \mathcal{K}_m \rangle$ the substitution complex on the vertex set [l] = $[l_1] \sqcup \cdots \sqcup [l_m]$ of $\mathcal{K}_1, \ldots, \mathcal{K}_m$ into the identity complex \mathcal{K}_{Π} . By analysing different partitions Π of [l] and folds $\psi: I \longrightarrow J$ of \mathcal{S}_{Π} , we obtain two further families of relations among folded higher Whitehead maps.

We first consider folds of S_{Π} which are induced by folds within the \mathcal{K}_i for $i = 1, \ldots, m$.

Lemma 6.6. Let \mathcal{L} be a simplicial complex on [m] and let $\mathcal{K}_1, \ldots, \mathcal{K}_m$ be simplicial complexes on $[l_1], \ldots, [l_m]$, respectively. Let $[l] = [l_1] \sqcup \cdots \sqcup [l_m]$ and suppose that $\psi: I \longrightarrow J$ is a fold of $\mathcal{L}\langle \mathcal{K}_1, \ldots, \mathcal{K}_m \rangle$ such that $\psi([l_i] \cap I) \subseteq [l_i]$ for each $i = 1, \ldots, m$. Then

$$\mathcal{L}\langle \mathcal{K}_1, \dots, \mathcal{K}_m \rangle_{\nabla(I,J)} = \mathcal{L}\langle (\mathcal{K}_1)_{\nabla(I,J)}, \dots, (\mathcal{K}_m)_{\nabla(I,J)} \rangle$$

where $(\mathcal{K}_i)_{\nabla(I,J)} = \mathcal{K}_i$ for $[l_i] \cap I = \emptyset$.

Proof. We extend the fold $\psi: I \longrightarrow J$ to a map $\overline{\psi}: [l] \longrightarrow [l]$ by setting $\overline{\psi}(i) = \psi(i)$ if $i \in I$ and $\overline{\psi}(i) = i$ otherwise. Then ψ and $\overline{\psi}$ induce the same fold of simplicial complexes. By definition,

$$\mathcal{L}\langle \mathcal{K}_1, \dots, \mathcal{K}_m \rangle = \left\{ \bigsqcup_{j \in \tau} \sigma_j \mid \sigma_j \in \mathcal{K}_j, \tau \in \mathcal{L} \right\}$$

and therefore

(40)
$$\mathcal{L}\langle \mathcal{K}_1, \dots, \mathcal{K}_m \rangle_{\nabla(I,J)} = \left\{ \bigsqcup_{j \in \tau} \overline{\psi}(\sigma_j) \mid \sigma_j \in \mathcal{K}_j, \tau \in \mathcal{L} \right\}.$$

Any simplex of $(\mathcal{K}_i)_{\nabla(I,J)}$ is either a simplex of \mathcal{K}_i , or $\psi(\sigma_i)$ for some $\sigma_i \in \mathcal{K}_i$. In either case, every simplex $(\mathcal{K}_i)_{\nabla(I,J)}$ can be written as $\overline{\psi}(\sigma_i)$ for some $\sigma_i \in \mathcal{K}_i$, and so the right-hand side of (40) is $\mathcal{L}\langle (\mathcal{K}_1)_{\nabla(I,J)}, \ldots, (\mathcal{K}_m)_{\nabla(I,J)} \rangle$.

Theorem 6.7. Let $\mathcal{K}_1, \ldots, \mathcal{K}_m$ be simplicial complexes on $[l_1], \ldots, [l_m]$, respectively, and let $[l] = [l_1] \sqcup \cdots \sqcup [l_m]$. Let $\Pi = \{P_1, \ldots, P_k\}$ be a k-partition of [m] and denote by $P_i = \{i_1, \ldots, i_{p_i}\}$ and $[m] \setminus P_i = \{j_1, \ldots, j_{q_i}\}$. Let $\mathcal{S}_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \ldots, \mathcal{K}_m \rangle$ and let $\psi: I \longrightarrow J$ be a fold of \mathcal{S}_{Π} such that $\psi([l_i] \cap I) \subseteq [l_i]$ for each $i = 1, \ldots, m$. Let $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_i}$ be nested higher Whitehead maps for $i = 1, \ldots, m$. If each X_i is a suspension, then

(41)
$$\sum_{i=1}^{k} h_{w}^{\mathcal{S}_{\Pi}} \left(h_{w} \left(\nabla_{(I,J)} f_{j_{1}}, \dots \nabla_{(I,J)} f_{j_{q_{i}}} \right), \nabla_{(I,J)} f_{i_{1}}, \dots, \nabla_{(I,J)} f_{i_{p_{i}}} \right) \circ \sigma_{i} = 0$$

 $in \left[\Sigma^{m-2} X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{(\mathcal{S}_{\Pi})_{\nabla(I,J)}} \right].$ Furthermore, all summands are null-homotopic if

$$\nabla_{(I,J)} f_j \colon \Sigma X_j \longrightarrow (\underline{Y}, \underline{*})^{(\mathcal{K}_j)_{\nabla(I,J)}}$$

is null-homotopic for some $j \in \{1, \ldots, m\}$.

Proof. By Lemma 6.6, $(\mathcal{S}_{\Pi})_{\nabla(I,J)} = \mathcal{K}_{\Pi} \langle (\mathcal{K}_1)_{\nabla(I,J)}, \dots, (\mathcal{K}_m)_{\nabla(I,J)} \rangle$. Therefore by Proposition 2.4,

$$\begin{aligned} \nabla_{(I,J)} h_w^{\mathcal{S}_{\Pi}} \left(h_w \left(f_{j_1}, \dots, f_{j_{q_i}} \right), f_{i_1}, \dots, f_{i_{p_i}} \right) \\ &= h_w^{\mathcal{S}_{\Pi}} \left(h_w \left(\nabla_{(I,J)} f_{j_1}, \dots, \nabla_{(I,J)} f_{j_{q_i}} \right), \nabla_{(I,J)} f_{i_1}, \dots, \nabla_{(I,J)} f_{i_{p_i}} \right) \end{aligned}$$

establishing the claimed relation. Moreover, if $\nabla_{(I,J)}f_j = 0$ for some $j \in \{1, \ldots, m\}$, every term in relation (41) is null-homotopic by Proposition 3.3.

The second type of folds of $S_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \ldots, \mathcal{K}_m \rangle$ we consider are those which fold \mathcal{K}_i onto \mathcal{K}_j for $i \neq j$.

Lemma 6.8. Let $\Pi = \{\{1\}, \{2, \ldots, m-1\}, \{m\}\}$ be a 3-partition of [m], and define $\mathcal{S}_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \ldots, \mathcal{K}_m \rangle$. Assume that \mathcal{K}_m is isomorphic to a full subcomplex of \mathcal{K}_1 , with the isomorphism given by a simplicial map $\psi: \mathcal{K}_m \longrightarrow \mathcal{K}_1$. Let $I = [l_m]$ and $J = \psi([l_m])$. Then

$$(\mathcal{S}_{\Pi})_{\nabla(I,J)} = \partial \Delta \langle \mathcal{K}_1, \dots, \mathcal{K}_{m-1} \rangle.$$

Proof. The subcomplexes \mathcal{K}_1 and \mathcal{K}_m of \mathcal{S}_{Π} are disjoint since there is no edge between the vertices $\{1\}$ and $\{m\}$ in \mathcal{K}_{Π} . Therefore, we rewrite \mathcal{S}_{Π} as

$$\mathcal{S}_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \dots, \mathcal{K}_m \rangle = \partial \Delta \langle \mathcal{K}_1 \sqcup \mathcal{K}_m, \mathcal{K}_2, \dots, \mathcal{K}_{m-1} \rangle.$$

The result follows by Lemma 6.6 since $(\mathcal{K}_1 \sqcup \mathcal{K}_m)_{\nabla(I,J)} = \mathcal{K}_1$.

For $i = 1, \ldots, m$, let $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_i}$ be nested higher Whitehead maps. If each X_i is a suspension, then there is a relation

(42)
$$\nabla_{(I,J)} h_w^{\mathcal{S}_{\Pi}} (h_w(f_2, \dots, f_m), f_1) \circ \sigma_1 + \nabla_{(I,J)} h_w^{\mathcal{S}_{\Pi}} (h_w(f_1, f_m), f_2, \dots, f_{m-1}) \circ \sigma_2$$

 $+ \nabla_{(I,J)} h_w^{\mathcal{S}_{\Pi}} (h_w(f_1, \dots, f_{m-1}), f_m) = 0$

in $\left[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{(\mathcal{S}_{\Pi})_{\nabla(I,J)}}\right]$. By Proposition 4.10(i), the second summand is null-homotopic if

$$(\partial \Delta \langle \Delta \langle \mathcal{K}_1, \mathcal{K}_m \rangle, \mathcal{K}_2, \dots, \mathcal{K}_{m-1} \rangle)_{\nabla(I,J)} = \partial \Delta \langle (\mathcal{K}_1 * \mathcal{K}_m)_{\nabla(I,J)}, \mathcal{K}_2, \dots, \mathcal{K}_{m-1} \rangle$$

is a subcomplex of $(\mathcal{S}_{\Pi})_{\nabla(I,J)} = \partial \Delta \langle \mathcal{K}_1, \ldots, \mathcal{K}_{m-1} \rangle$. Since $\mathcal{K}_1 \subseteq (\mathcal{K}_1 * \mathcal{K}_m)_{\nabla(I,J)}$, this condition is equivalent to $\mathcal{K}_1 = (\mathcal{K}_1 * \mathcal{K}_m)_{\nabla(I,J)}$. We therefore obtain the following.

Theorem 6.9. Let $\Pi = \{\{1\}, \{2, \ldots, m-1\}, \{m\}\}$ be a 3-partition of [m], and define $S_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \ldots, \mathcal{K}_m \rangle$. Assume that \mathcal{K}_m is isomorphic to a full subcomplex of \mathcal{K}_1 , with the isomorphism given by a simplicial map $\psi: \mathcal{K}_m \longrightarrow \mathcal{K}_1$. Let $I = [l_m]$ and $J = \psi([l_m])$. Let $f_i: \Sigma X_i \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_i}$ be nested higher Whitehead maps for $i = 1, \ldots, m$, with each X_i a suspension. Then

$$\nabla_{(I,J)} h_w^{S_{\Pi}} (h_w(f_2, \dots, f_m), f_1) \circ \sigma_1 + \nabla_{(I,J)} h_w^{S_{\Pi}} (h_w(f_1, f_m), f_2, \dots, f_{m-1}) \circ \sigma_2 + \nabla_{(I,J)} h_w^{S_{\Pi}} (h_w(f_1, \dots, f_{m-1}), f_m) = 0$$

 $in\left[\Sigma^{m-2}X_1\wedge\cdots\wedge X_m, (\underline{Y},\underline{*})^{(\mathcal{S}_{\Pi})_{\nabla(I,J)}}\right], \text{ where the second summand is trivial if } \mathcal{K}_1 = (\mathcal{K}_1 * \mathcal{K}_m)_{\nabla(I,J)}.$

As an application of Theorem 6.9, we return to relation (35), considered at the start of this section. We show that this relation holds integrally.

Example 6.10. Consider the 3-partition $\Pi = \{\{1\}, \{2,3\}, \{4\}\}\}$. Assume that $\mathcal{K}_1 = \partial \Delta[1_1, 1_2, 1_3]$ and $\mathcal{K}_i = \bullet$ for i = 2, 3, 4. Let

$$f_1 = h_w(f_{1_1}, f_{1_2}, f_{1_3}) \colon \Sigma^2 X_{1_1} \wedge X_{1_2} \wedge X_{1_3} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_1}$$

and $f_i: \Sigma X_i \longrightarrow Y_i$ for i = 2, 3, 4 be maps. Let $I = \{4\}, J = \{1_1\}$ and $\psi: I \longrightarrow J$ be the fold of $\mathcal{S}_{\Pi} = \mathcal{K}_{\Pi} \langle \mathcal{K}_1, \mathcal{K}_2, \mathcal{K}_3, \mathcal{K}_4 \rangle = \partial \Delta \langle \mathcal{K}_1 \sqcup \mathcal{K}_4, \mathcal{K}_2, \mathcal{K}_3 \rangle$ on $\{1_1, 1_2, 1_3, 2, 3, 4\}$ given by $\psi(4) = 1_1$. Suppose that X_i is a suspension for i = 2, 3, 4. Then since $(\mathcal{K}_1 * \mathcal{K}_4)_{\nabla(I,J)} = \Delta[1_1, 1_2, 1_3] \neq \mathcal{K}_1$, by Theorem 6.9 there is a relation (43)

$$\begin{split} \nabla_{(4,1_1)} h_w^{\mathcal{S}_{\Pi}}(h_w(f_2,f_3,f_4),h_w(f_{1_1},f_{1_2},f_{1_3})\circ\sigma_1 + \nabla_{(4,1_1)}h_w^{\mathcal{S}_{\Pi}}(h_w(h_w(f_{1_1},f_{1_2},f_{1_3}),f_4),f_2,f_3)\circ\sigma_2 \\ + \nabla_{(4,1_1)}h_w^{\mathcal{S}_{\Pi}}(h_w(h_w(f_{1_1},f_{1_2},f_{1_3}),f_2,f_3),f_4) = 0 \end{split}$$

in $\left[\Sigma^2 X_1 \wedge X_2 \wedge X_3 \wedge X_4, (\underline{Y}, \underline{*})^{(\mathcal{S}_{\Pi})_{\nabla(4,1_1)}}\right]$, where $(\mathcal{S}_{\Pi})_{\nabla(4,1_1)} = \partial \Delta \langle \partial \Delta [1_1, 1_2, 1_3], 2, 3 \rangle$. Specialising to the case that $X_i = S^1$ and $Y_i = \mathbb{C}P^{\infty}$ for all $i = 1, \ldots, m$, we

specialising to the case that $A_i = S^*$ and $I_i = \mathbb{C}I^*$ for all i = 1, ..., m, we recover integrally a relation detected in the rational homotopy groups of the Davis– Januszkiewicz space $DJ_{\mathcal{L}}$, where $\mathcal{L} = (\mathcal{S}_{\Pi})_{\nabla(4,1_1)}$. In particular, let each $f_j: S^2 \longrightarrow \mathbb{C}P^{\infty}$ be the inclusion of the bottom cell and let $u_j: S^1 \longrightarrow \Omega \mathbb{C}P^{\infty}$ be the adjoint of f_j . Zhuravleva [15] showed that

(44)
$$[[u_2, u_3, u_{1_1}], [u_{1_1}, u_{1_2}, u_{1_3}]] + [[[u_{1_1}, u_{1_2}, u_{1_3}], u_2, u_3], u_{1_1}] = 0$$

in $\pi_8(\Omega DJ_{\mathcal{L}}) \otimes \mathbb{Q}$, where $[\cdot, \cdot, \cdot]$ is the triple Lie bracket.

Integrally, by relation (43), the Whitehead products $[h_w^{\mathcal{L}}(f_2, f_3, f_{1_1}), h_w^{\mathcal{L}}(f_{1_1}, f_{1_2}, f_{1_3})]$ and $[h_w^{\mathcal{L}}(h_w(f_{1_1}, f_{1_2}, f_{1_3}), f_2, f_3), f_{1_1}^{\mathcal{L}}]$ differ up to sign by

(45)
$$\nabla_{(4,1_1)}h_w(h_w(f_{1_1}, f_{1_2}, f_{1_3}), f_4), f_2, f_3) \in [[h_w^{\mathcal{L}}(f_{1_1}, f_{1_2}, f_{1_3}), f_{1_1}^{\mathcal{L}}], f_2^{\mathcal{L}}, f_3^{\mathcal{L}}]$$

where $f_i^{\mathcal{L}}: \Sigma S^2 \xrightarrow{f_i} \mathbb{C}P_i^{\infty} \longrightarrow DJ_{\mathcal{L}}$. By Example 6.5, the folded map $\nabla_{(4,1_1)}h_w(h_w(f_{1_1}, f_{1_2}, f_{1_3}), f_4)$ is trivial, and therefore so is map (45). Hence relation (43) recovers relation (44) integrally.

Alternatively, the map (45) may not be null-homotopic for different maps f_i . By Example 6.5, if each $f_i: S^2 \longrightarrow \Omega S^3$ is adjoint to the suspension map on S^2 , then the folded map $\nabla_{(4,1_1)} h_w(h_w(f_{1_1}, f_{1_2}, f_{1_3}), f_4)$ is 2-torsion. It follows that map (45) is also 2-torsion. We therefore recover further integral relations which cannot be detected rationally.

7. Proof of main theorem

In this section we prove Theorem 5.6, which we now recall.

Theorem 5.6. For i = 1, ..., m, let $f_i: \Sigma X_i \longrightarrow Y_i$ be maps. Let $\Pi = \{P_1, ..., P_k\}$ be a k-partition of [m] for $k \ge 3$ and denote by $P_i = \{i_1, ..., i_{p_i}\}$ and $Q_i = [m] \setminus P_i = \{j_1, ..., j_{q_i}\}$. If X_i is a suspension for each i = 1, ..., m, then

(46)
$$\sum_{i=1}^{k} h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) \circ \sigma_{i} = 0$$

in $[\Sigma^{m-2}X_1 \wedge \cdots \wedge X_m, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}]$, where

$$\sigma_i: \Sigma^{m-2} X_1 \wedge \dots \wedge X_m \longrightarrow \Sigma^{p_i} (\Sigma^{q_i-2} X_{j_1} \wedge \dots \wedge X_{j_{q_i}}) \wedge X_{i_1} \wedge \dots \wedge X_{i_{p_i}}$$

is the restriction of the coordinate permutation

$$CX_1 \times \cdots \times CX_m \longrightarrow CX_{j_1} \times \cdots \times CX_{j_{q_i}} \times CX_{i_1} \times \cdots \times CX_{i_{p_i}}.$$

We establish Theorem 5.6 by generalising the proof of the Jacobi identity (1) and relation (27) due to Nakaoka–Toda [10] and Hardie [9], respectively. We extend their methods in two ways, first by using the combinatorial structure of \mathcal{K}_{Π} to detect the form of the nested higher Whitehead maps appearing in relation (46), and second by deriving the claimed relations for non-spherical maps $f_i: \Sigma X_i \longrightarrow Y_i$.

We begin by outlining the proof. To prove the Jacobi identity, Nakaoka-Toda studied relative Whitehead products, defined by Blakers–Massey [6], appearing in the long exact sequence of relative homotopy groups. We adapt their methods to higher Whitehead maps.

We will define relative higher Whitehead maps, which are a generalisation of the relative Whitehead products. In the long exact sequence in homotopy

$$(47) \qquad \cdots \longrightarrow [\Sigma W, X] \xrightarrow{j} [(CW, W), (X, A)] \xrightarrow{\partial} [W, A] \longrightarrow \cdots$$

the map j is the composite of the isomorphism $[\Sigma W, X] \longrightarrow [(CW, W), (X, *)]$ with the map induced by the inclusion $(X, *) \longrightarrow (X, A)$, and the map ∂ is the restriction sending $[f] \in [(CW, W), (X, A)]$ to $[f|_W] \in [W, A]$. We show that relative higher Whitehead maps have the property that

(48)
$$\partial h_w(g, g_1, \dots, g_n) = h_w(g|_{\Sigma X}, g_1, \dots, g_n)$$

where ∂ is the boundary map in (47), and g is a map of pairs.

Specialising to $W = \Sigma^{m-2} X_1 \wedge \cdots \wedge X_m$, $X = FW(Y_1, \ldots, Y_m)$, and $A = (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}$, we show that the higher Whitehead map $h_w(f_1, \ldots, f_m) \in [\Sigma W, X]$ when composed with the map j decomposes as

(49)
$$jh_w(f_1,\ldots,f_m) = \sum_{i=1}^k \varphi_i \sigma_i$$

where $\{\varphi_i\}$ is a family of relative higher Whitehead maps determined by the combinatorics of the partition Π , whose image under the boundary map in the long exact sequence (48) is

(50)
$$\partial \varphi_i = h_w^{\mathcal{K}_{\Pi}} \left(h_w \left(f_{j_1}, \dots, f_{j_{q_i}} \right), f_{i_1}, \dots, f_{i_{p_i}} \right)$$

The exactness of sequence (47) implies that

$$\sum_{i=1}^{k} h_{w}^{\mathcal{K}_{\Pi}} \left(h_{w} \left(f_{j_{1}}, \dots, f_{j_{q_{i}}} \right), f_{i_{1}}, \dots, f_{i_{p_{i}}} \right) \circ \sigma_{i} = \partial(j(h_{w}(f_{1}, \dots, f_{m}))) = 0$$

which proves Theorem 5.6.

We define the relative higher Whitehead map as a generalisation of the relative Whitehead product. For a sphere S^d , let D^d_+ and D^d_- denote the upper and lower hemispheres, respectively. Let $\alpha \in \pi_p(X, A)$ and $\beta \in \pi_q(A)$ be represented by $f: (D^p, S^{p-1}, D^{p-1}_+) \longrightarrow (X, A, *)$ and $g: (D^q, S^{q-1}) \longrightarrow (A, *)$, respectively. The relative Whitehead product $[\alpha, \beta] \in \pi_{p+q-1}(X, A)$ is the homotopy class of the map of pairs

$$(D^{p+q-1}, S^{p+q-2}) = (D^p \times S^{q-1} \cup D^{p-1}_+ \times D^q, D^{p-1}_- \times S^{q-1} \cup S^{p-2} \cup D^q)$$
$$\longrightarrow (X \lor A, A \lor A) \longrightarrow (X, A)$$

where the first map is induced by the product $f \times g$ and the second map is induced by the fold $\nabla: X \vee X \longrightarrow X$.

The relative Whitehead product has the properties of naturality and bilinearity, see [6], analogously with the Whitehead product. It also satisfies the following. Let $\partial: \pi_n(X, A) \longrightarrow \pi_{n-1}(A)$ denote the boundary operator in the long exact sequence

$$\cdots \longrightarrow \pi_n(X) \longrightarrow \pi_n(X, A) \xrightarrow{\partial} \pi_{n-1}(A) \longrightarrow \cdots$$

Then

(51)
$$\partial_{p+q-1}[\alpha,\beta] = -[\partial_p \alpha,\beta].$$

This is a key property used by Nakaoka–Toda [10] in the proof of the Jacobi identity. To analyse relations between higher Whitehead maps, we introduce the relative higher Whitehead map as an analogue of the relative Whitehead product. **Definition 7.1.** Let $g: (C\Sigma X, \Sigma X) \longrightarrow (Z, B)$ be a map of pairs, let $g_i: \Sigma X_i \longrightarrow Y_i$ be maps for $i = 1, \ldots, n$, and let $W = X * X_1 * \cdots * X_n$. The relative higher Whitehead map of g, g_1, \ldots, g_n is the composite

$$h_w(g, g_1, \dots, g_n): CW \simeq CX * X_1 * \dots * X_n$$
$$\xrightarrow{\rho} FW(\Sigma CX, \Sigma X_1, \dots, \Sigma X_n)$$
$$\xrightarrow{\simeq} FW(C\Sigma X, \Sigma X_1, \dots, \Sigma X_n)$$
$$\longrightarrow FW(Z, Y_1, \dots, Y_n)$$

where $\rho: CX * X_1 * \cdots * X_n \longrightarrow FW(\Sigma CX, \Sigma X_1, \ldots, \Sigma X_n)$ is the quotient map defined in (9), and the last map is induced by g, g_1, \ldots, g_n .

The restriction of $h_w(g, g_1, \ldots, g_n)$ to W,

$$W = X * X_1 * \dots * X_n$$

$$\stackrel{\rho}{\longrightarrow} FW(\Sigma X, \Sigma X_1, \dots, \Sigma X_n)$$

$$\longrightarrow (B, Y_1, \dots, Y_n)$$

is the higher Whitehead map $h_w(g|_{\Sigma X}, g_1, \ldots, g_n)$. The homotopy class of the relative higher Whitehead map $h_w(g, g_1, \ldots, g_n)$ is therefore an element of the relative homotopy group

 $[(CW, W), (FW(Z, Y_1, \ldots, Y_n), FW(B, Y_1, \ldots, Y_n))].$

The higher and relative higher Whitehead maps satisfy an analogous relation to that of Blakers–Massey (51).

Proposition 7.2. The relative higher Whitehead map $h_w(g, g_1, \ldots, g_n)$ satisfies

 $\partial h_w(g, g_1, \dots, g_n) = h_w(g|_{\Sigma X}, g_1, \dots, g_n)$

where

 $\partial: [(CW, W), (FW(Z, Y_1, \dots, Y_n), FW(B, Y_1, \dots, Y_n)] \longrightarrow [(W, FW(B, Y_1, \dots, Y_n)]$ is the boundary map in the exact sequence (47).

Proof. Since

$$\partial h_w(g, g_1, \dots, g_n) = h_w(g, g_1, \dots, g_n)|_W$$

the result follows immediately from the definition of $h_w(g, g_1, \ldots, g_n)$.

The following lemma enables us to identify the image of j with the sum of relative maps in the long exact sequence (47).

Lemma 7.3. Let W be a suspension space. For i = 1, ..., k, consider pairs (F_i, Z_i) with $F_i \subseteq \Sigma W$ satisfying the following conditions. For each i = 1, ..., k, Z_i is closed in F_i , pairwise intersections are such that $F_i \cap F_j = Z_i \cap Z_j$ for all $j \neq i$, and there exists a homotopy equivalence $e_i: \Sigma W \longrightarrow \Sigma W$ such that $e_i(F_i) = C_+W$, $e_i(Z_i) = W$, and $e_i((\Sigma W \setminus F_i) \cup Z_i) = C_-W$. Let (X, A) be a pair and suppose that $f: \Sigma W \longrightarrow X$ satisfies $f((\Sigma W \setminus \bigcup_i F_i) \cup (\bigcup_i Z_i)) \subseteq A$. Then

$$j(f) = \sum_{i=1}^{\kappa} f_i$$

where $f_i = f|_{F_i} \in [(F_i, Z_i), (X, A)] \cong [(CW, W), (X, A)]$ for each $i = 1, \ldots, k$ and $j: [\SigmaW, X] \longrightarrow [(CW, W), (X, A)]$ is the map in the long exact sequence (47).

To establish Lemma 7.3, we first establish the following lemma.

Lemma 7.4. Let $Z = C_1 \widetilde{Z} \cup C_2 \widetilde{Z}$ and write $\Sigma Z = C_+ Z \cup C_- Z$. Let (X, A) be a CW-pair. Let $g: \Sigma Z \longrightarrow X$ be a map such that $g(Z) \subseteq A \subseteq X$. Then $j(g) = g|_{C_+Z} + g|_{C_-Z}$.

Proof. The map j is the composite of the isomorphism $\psi: [\Sigma Z, X] \longrightarrow [(CZ, Z), (X, *)]$ with the map j' induced by the inclusion $(X, *) \longrightarrow (X, A)$. We describe a choice of map of pairs $\phi: (CZ, Z) \longrightarrow (\Sigma Z, *)$ which induces the isomorphism ψ .

Consider $CZ = C\Sigma \tilde{Z}$ and define

$$G_1 = \left\{ (s,t,z) \in C\Sigma\widetilde{Z} \mid 0 \leqslant t \leqslant \frac{1}{2} \right\}, \quad G_2 = \left\{ (s,t,z) \in C\Sigma\widetilde{Z} \mid \frac{1}{2} \leqslant t \leqslant 1 \right\}$$
$$G_3 = \left\{ (s,t,z) \in C\Sigma\widetilde{Z} \mid \frac{1}{2} \leqslant s \leqslant 1 \right\}, \quad G_4 = \left\{ (s,t,z) \in C\Sigma\widetilde{Z} \mid 0 \leqslant s \leqslant \frac{1}{2} \right\}.$$

Define the map $\phi: (CZ, Z) \longrightarrow (\Sigma Z, *)$ by $\phi(G_1) = C_+Z$, $\phi(G_2) = C_-Z$, $\phi(G_3) = \Sigma C_1 \widetilde{Z}$ and $\phi(G_4) = \Sigma C_2 \widetilde{Z}$. Then $\psi(g) = g\phi$.

In particular, ϕ sends $G_1 \cap G_2$ to $Z \subseteq \Sigma Z$, $G_1 \cap G_3$ to $C_+C_1\widetilde{Z}$ and $G_1 \cap G_4$ to $C_+C_2\widetilde{Z}$. Since $g(Z) \subseteq A$, then $\psi(g) = g\phi$ is a map such that $G_1 \cap G_2 \subseteq A$. Since $G_1 \cap G_2$ is contractible, there is a homotopy from $g\phi$ to a map sending $G_1 \cap G_2$ to the basepoint. Therefore $j(g) = j'(\psi g)$ is homotopic to the sum $g\phi|_{G_1} + g\phi|_{G_2} = g|_{C_+Z} + g|_{C_-Z}$.

We now prove Lemma 7.3.

Proof of Lemma 7.3. Write $\Sigma W = F_k \cup_{Z_k} ((\Sigma W \setminus F_k) \cup Z_k)$. Pre-composing with the homotopy equivalence e_k , the map f is homotopic to a map $f' = f \circ e_k \colon \Sigma W \longrightarrow X$, where $f'(W) \subseteq A$ since $f(Z_k) \subseteq A$. Then, by Lemma 7.4, $j(f) = j(f') = f'|_{C_-W} + f'|_{C_+W} = f|_{(\Sigma W \setminus F_k) \cup Z_k} + f|_{F_k}$. Since $f((\Sigma W \setminus \bigcup_i F_i) \cup (\bigcup_i Z_i)) \subseteq A$, collapsing $(\Sigma W \setminus \bigcup_i F_i) \cup (\bigcup_i Z_i)$ gives a homotopy of pairs between the map $f|_{(\Sigma W \setminus F_k) \cup Z_k} \colon (CW, W) \longrightarrow (X, A)$ and $\sum_{i=1}^{k-1} f|_{F_i}$.

We now specialise to the *j*-image of the higher Whitehead map $h_w(f_1, \ldots, f_m)$ and identify it as a sum of relative higher Whitehead maps.

Recall that for the partition $\Pi = \{P_1, \ldots, P_k\}$ of [m], we write $P_i = \{i_1, \ldots, i_{p_i}\}$ and $Q_i = [m] \setminus P_i = \{j_1, \ldots, j_{q_i}\}$. For $i = 1, \ldots, k$, the product $f_{j_1} \times \cdots \times f_{j_{q_i}}$ induces a map of pairs

$$\psi_i: (CX_{j_1} \times \dots \times CX_{j_{q_i}}, X_{j_1} \ast \dots \ast X_{j_{q_i}}) \longrightarrow \left((\underline{Y}, \underline{\ast})^{\Delta[j_1, \dots, j_{q_i}]}, (\underline{Y}, \underline{\ast})^{\partial \Delta[j_1, \dots, j_{q_i}]} \right)$$

such that the restriction $\psi_i|_{X_{j_1}*\cdots*X_{j_{q_i}}} = h_w(f_{j_1},\ldots,f_{j_{q_i}})$. The domain of ψ_i is homotopy equivalent to the pair $(C\Sigma W_{Q_i},\Sigma W_{Q_i})$, where $W_{Q_i} = \Sigma^{q_i-2}X_{j_1}\wedge\cdots\wedge X_{j_{q_i}}$. Therefore by Proposition 7.2, the relative higher Whitehead map

(52)
$$\varphi_i = h_w(\psi_i, f_{i_1}, \dots, f_{i_{p_i}}) \colon (CW_{Q_i} * X_{i_1} * \dots * X_{i_{p_i}}, W_{Q_i} * X_{i_1} * \dots * X_{i_{p_i}}) \\ \longrightarrow \left((\underline{Y}, \underline{*})^{\partial \Delta \langle \Delta[j_1, \dots, j_{q_i}], i_1, \dots, i_{p_i} \rangle}, (\underline{Y}, \underline{*})^{\partial \Delta \langle \partial \Delta[j_1, \dots, j_{q_i}], i_1, \dots, i_{p_i} \rangle} \right)$$

satisfies

(53)
$$\partial \varphi_i = h_w(h_w(f_{j_1}, \dots, f_{j_{q_i}}), f_{i_1}, \dots, f_{i_{p_i}})$$

for all $i = 1, \ldots, k$.

To apply Lemma 7.3, we need to check three conditions. We start by constructing pairs $(F_{Q_i}, Z_{Q_i}) \subseteq \Sigma W$ which are homotopy equivalent to the domains of the maps φ_i , and such that $F_{Q_i} \cap F_{Q_{i'}} = Z_{Q_i} \cap Z_{Q_{i'}}$ for $i \neq i'$.

Denote by (F'_{Q_i}, Z'_{Q_i}) the domain of the map φ_i , and observe that it decomposes as

$$(CW_{Q_{i}} * X_{i_{1}} * \dots * X_{i_{p_{i}}}, W_{Q_{i}} * X_{i_{1}} * \dots * X_{i_{p_{i}}})$$

= $(C'CW_{Q_{i}} \times (X_{i_{1}} * \dots * X_{i_{p_{i}}}) \cup CW_{Q_{i}} \times C(X_{i_{1}} * \dots * X_{i_{p_{i}}}),$
 $C'W_{Q_{i}} \times (X_{i_{1}} * \dots * X_{i_{p_{i}}}) \cup W_{Q_{i}} \times C(X_{i_{1}} * \dots * X_{i_{p_{i}}})).$

We construct a map from $C'CW_{Q_i}$ to a space G_{Q_i} which restricts to homotopy equivalences between CW_{Q_i} , $C'W_{Q_i}$ and W_{Q_i} and subspaces L_{Q_i} , M_{Q_i} , and N_{Q_i} of G_{Q_i} , respectively. This induces a homotopy equivalence of pairs from (F'_{Q_i}, Z'_{Q_i}) to (F_{Q_i}, Z_{Q_i}) such that $F_{Q_i} \cap F_{Q_{i'}} = Z_{Q_i} \cap Z_{Q_{i'}}$ for $i \neq i'$.

Decompositions of CX_i . Fix $i \in \{1, \ldots, m\}$. Since X_i is a suspension, we write $X_i = \Sigma \widetilde{X}_i = C_+ \widetilde{X}_i \cup C_- \widetilde{X}_i$. Consider $C\Sigma \widetilde{X}_i$ as

$$C\Sigma \widetilde{X}_i = \{(s,t,x) \in I \times I \times \widetilde{X}_i\}/(1,t,x) \sim (s,0,x) \sim (s,1,x) \sim (s,t,*)$$

so that $\Sigma \widetilde{X}_i$ is the subspace at s = 0, and $C_+ \widetilde{X}_i$, $C_- \widetilde{X}_i$ and \widetilde{X}_i are the subspaces of $\Sigma \widetilde{X}_i$ with $t \ge \frac{1}{2}$, $t \le \frac{1}{2}$, and $t = \frac{1}{2}$, respectively.

We define two decompositions $C\Sigma \widetilde{X}_i = D_i^- \cup D_i^+ = D_i^1 \cup D_i^2$, where

$$D_i^+ = \left\{ (s, (t, x)) \in C\Sigma \widetilde{X}_i \mid \frac{1}{2} \leqslant t \right\} \qquad D_i^- = \left\{ (s, (t, x)) \in C\Sigma \widetilde{X}_i \mid t \leqslant \frac{1}{2} \right\}$$
$$D_i^1 = \left\{ (s, (t, x)) \in C\Sigma \widetilde{X}_i \mid \frac{3}{4} - t \leqslant s \right\} \qquad D_i^2 = \left\{ (s, (t, x)) \in C\Sigma \widetilde{X}_i \mid s \leqslant \frac{3}{4} - t \right\}.$$

In order to create separation of subspaces in D_i^- and D_i^1 , we further decompose $D_i^1 = E_i^1 \cup C_i^1$ and $D_i^- = E_i^- \cup C_i^-$, where

$$E_i^- = \left\{ (s, (t, x)) \in D_i^- \mid t \leq \frac{1}{8} \right\} \qquad E_i^1 = \left\{ (s, (t, x)) \in D_i^1 \mid t \geq \frac{7}{8} \right\}$$
$$C_i^- = \left\{ (s, (t, x)) \in D_i^- \mid t \geq \frac{1}{8} \right\} \qquad C_i^1 = \left\{ (s, (t, x)) \in D_i^1 \mid t \leq \frac{7}{8} \right\}.$$

For $\delta \in \{-, 1\}$, we define maps $\alpha_i^{\delta} : C\Sigma \widetilde{X}_i \longrightarrow D_i^{\delta}$ and $\beta_i^{\delta} : D_i^{\delta} \longrightarrow E_i^{\delta}$ such that the restrictions of α_i^{δ} and $\beta_i^{\delta} \alpha_i^{\delta}$ to the subspaces $\Sigma \widetilde{X}_i, C_+ \widetilde{X}_i, C_- \widetilde{X}_i$, and \widetilde{X}_i are homotopy equivalences. We explicitly specify homotopy equivalent copies of $\Sigma \widetilde{X}_i, C_+ \widetilde{X}_i, C_- \widetilde{X}_i$, and \widetilde{X}_i inside D_i^{δ} and E_i^{δ} . Define

$$\alpha_i^-: C\Sigma \widetilde{X}_i \longrightarrow D_i^-, \qquad (s, (t, x)) \mapsto \begin{cases} (2s + 2t - 2st - 1, (\frac{1}{2}, x)) & \text{if } t \ge \frac{1}{2} \\ (s, (t, x)) & \text{otherwise} \end{cases}$$

and

$$\alpha_i^1 : C \Sigma \widetilde{X}_i \longrightarrow D_i^1, \qquad (s, (t, x)) \mapsto \begin{cases} (s, (\frac{3}{4} - s, x)) & \text{if } s \leqslant \frac{3}{4} - t, t \geqslant \frac{1}{2} \\ (1 - 2t + s, (2t - s - \frac{1}{4}, x)) & \text{if } s \leqslant t - \frac{1}{4}, t \leqslant \frac{1}{2} \\ (\frac{3}{4} - t, (t, x)) & \text{if } t - \frac{1}{4} \leqslant s \leqslant \frac{3}{4} - t \\ (s, (t, x)) & \text{if } s \geqslant \frac{3}{4} - t. \end{cases}$$

Then α_i^{δ} takes $\Sigma \widetilde{X}_i, C_- \widetilde{X}_i, C_+ \widetilde{X}_i$ and \widetilde{X}_i to homotopy equivalent spaces $B_i^{\delta}, (B_1)_i^{\delta}, (B_2)_i^{\delta}$ and $(B_1)_i^{\delta} \cap (B_2)_i^{\delta}$, respectively, where

$$(B_1)_i^- = \left\{ (s, (t, x)) \in D_i^- \mid t \leq \frac{1}{2}, s = 0 \right\} \quad (B_2)_i^- = \left\{ (s, (t, x)) \in D_i^- \mid t = \frac{1}{2} \right\}$$
$$(B_1)_i^1 = \left\{ (s, (t, x)) \in D_i^1 \mid s = \frac{3}{4} - t \right\} \qquad (B_2)_i^1 = \left\{ (s, (t, x)) \in D_i^1 \mid t \geq \frac{3}{4}, s = 0 \right\}$$

and $B_i^{\delta} = (B_1)_i^{\delta} \cup (B_2)_i^{\delta}$.

Next, define

$$\beta_i^-: D_i^- \longrightarrow E_i^-, \qquad (s, (t, x)) \mapsto \begin{cases} (s, (\frac{1}{8}, x)) & \text{if } t \ge \frac{1}{8} \\ (s, (t, x)) & \text{otherwise} \end{cases}$$

and

$$\beta_i^1 \colon D_i^1 \longrightarrow E_i^1, \qquad (s, (t, x)) \mapsto \begin{cases} (1 - 2t + 2st, (\frac{7}{8}, x)) & \text{if } t \leq \frac{1}{2} \\ (s, (\frac{7}{8}, x)) & \text{if } \frac{1}{2} \leq t \leq \frac{7}{8} \\ (s, (t, x)) & \text{otherwise.} \end{cases}$$

Then the images of $B_i^{\delta}, (B_1)_i^{\delta}, (B_2)_i^{\delta}$ and $(B_1)_i^{\delta} \cap (B_2)_i^{\delta}$ under β_i^{δ} are, respectively, the homotopy equivalent spaces $S_i^{\delta} \cup T_i^{\delta}, S_i^{\delta}, T_i^{\delta}$ and R_i^{δ} , where

$$\begin{split} R_i^- &= \left\{ (s,(t,x)) \in D_i^- \mid t = \frac{1}{8}, s = 0 \right\} \quad R_i^1 = \left\{ (s,(t,x)) \in D_i^1 \mid t = \frac{7}{8}, s = 0 \right\} \\ S_i^- &= \left\{ (s,(t,x)) \in D_i^- \mid t \leq \frac{1}{8}, s = 0 \right\} \quad S_i^1 = \left\{ (s,(t,x)) \in D_i^1 \mid t = \frac{7}{8} \right\} \\ T_i^- &= \left\{ (s,(t,x)) \in D_i^- \mid t = \frac{1}{8} \right\} \qquad T_i^1 = \left\{ (s,(t,x)) \in D_i^1 \mid t \geq \frac{7}{8}, s = 0 \right\}. \end{split}$$

Summarising, we have the following.

Lemma 7.5. There are homotopy equivalences of pairs

$$(C\Sigma\widetilde{X}_i, \Sigma\widetilde{X}_i) \xrightarrow{\alpha_i^{\delta}} (D_i^{\delta}, B_i^{\delta}) \xrightarrow{\beta_i^{\delta}} (E_i^{\delta}, S_i^{\delta} \cup T_i^{\delta}).$$

Subspaces of $CX_{j_1} \times \cdots \times CX_{j_{q_i}}$. Denote by

$$V = CX_1 \times \dots \times CX_m$$
$$V^* = \bigcup_{l=1}^m CX_1 \times \dots \times X_l \times \dots \times CX_m.$$

For $i = 1, \ldots, k$, we denote by

$$V_{Q_i} = CX_{j_1} \times \dots \times CX_{j_{q_i}}$$
$$V_{Q_i}^* = \bigcup_{l=1}^{q_i} CX_{j_1} \times \dots \times X_{j_l} \times \dots \times CX_{j_{q_i}}.$$

For each i = 1, ..., k, we define a pair $(G_{Q_i}, G_{Q_i}^*)$ homotopy equivalent to $(V_{Q_i}, V_{Q_i}^*)$ by applying coordinate-wise the homotopy equivalences of Lemma 7.5. We use matrix notation to encode the factors in the product. If $k \ge 3$ is odd, define the $k \times k$ matrix $H = [\eta(i, j)]$ by

$$\eta(i,j) = \begin{cases} * & \text{if } i = j \\ 1 & \text{if } i+j < k+1, \text{ or } i+j = k+1 \text{ and } i > \frac{k+1}{2} \\ - & \text{if } i+j > k+1, \text{ or } i+j = k+1 \text{ and } i < \frac{k+1}{2}. \end{cases}$$

If $k \ge 4$ is even, define the $k \times k$ matrix $H' = [\eta'(i, j)]$ by

$$\eta'(i,j) = \begin{cases} * & \text{if } i = j \\ 1 & \text{if } i+j < k+1, \text{ or } i+j = k+1 \text{ and } i \ge \frac{k}{2}+1 \\ - & \text{if } i+j > k+1, \text{ or } i+j = k+1 \text{ and } i \le \frac{k}{2} \end{cases}$$

and define the matrix $H = [\eta(i, j)]$ from H' by swapping the entries $\eta(\frac{k}{2}+1, \frac{k}{2}-1) = 1$ and $\eta(\frac{k}{2}+1, \frac{k}{2}) = -$. Figure 4 shows the matrix H, on the left for k = 7, and on the right for k = 8.

*	1	1	1	1	1	1
1	*	1	1	1	1	_
1	1	*	1	1	_	_
1	1	1	*	_	_	_
1	1	_	_	*	_	_
1	_	_	_	_	*	_
—	_	_	_	—	—	*
-		,		_		-
		(A) k =	= 7		

FIGURE 4. The matrix H for k = 7 and k = 8.

We define

$$G_{Q_i} = \prod_{n=1}^{q_i} D_{j_n}^{\eta(i,\pi(j_n))} = D_{j_1}^{\eta(i,\pi(j_1))} \times \cdots \times D_{j_{q_i}}^{\eta(i,\pi(j_{q_i}))}$$
$$G_{Q_i}^* = \bigcup_{n=1}^{q_i} D_{j_1}^{\eta(i,\pi(j_1))} \times \cdots \times B_{j_n}^{\eta(i,\pi(j_n))} \times \cdots \times D_{j_{q_i}}^{\eta(i,\pi(j_{q_i}))}$$

where if $j_i \in P_l$, then $\pi(j_i) = l$. The pair $(G_{Q_i}, G_{Q_i}^*)$ is the image of $(V_{Q_i}, V_{Q_i}^*)$ under the product of the maps of pairs $\alpha_{j_n}^{\eta(i,\pi(j_n))}: (C\Sigma \widetilde{X}_{j_n}, \Sigma \widetilde{X}_{j_n}) \longrightarrow (D_{j_n}^{\eta(i,\pi(j_n))}, B_{j_n}^{\eta(i,\pi(j_n))})$. We summarise this observation in the following lemma.

Lemma 7.6. For each i = 1, ..., k, there is a homotopy equivalence of pairs $(V_{Q_i}, V_{Q_i}^*) \longrightarrow (G_{Q_i}, G_{Q_i}^*)$.

The following property of our construction will later be used to establish properties of the pairwise intersections $F_{Q_i} \cap F_{Q_{i'}}$.

Lemma 7.7. Suppose that $1 \leq i < j \leq k$. Then there exists r, different to i, j, such that G_{Q_i} contains a factor D_r^1 and G_{Q_i} contains a factor D_r^- .

Proof. The statement follows from the definition of the matrix H since, given any two rows i and j, there exists r different from i, j such that $\eta(i, r) = 1$ and $\eta(j, r) = -$. Then by construction G_{Q_i} has a factor D_r^1 and G_{Q_j} has a factor D_r^- .

For $i = 1, \ldots, k$, let $W_{Q_i} = \Sigma^{q_i - 2} X_{j_1} \wedge \cdots \wedge X_{j_{q_i}}$. Observe that $G_{Q_i}^* \simeq X_{j_1} * \cdots * X_{j_{q_i}} \simeq \Sigma W_{Q_i}$. We decompose $G_{Q_i}^*$ as a union of two contractible spaces L_{Q_i} and M_{Q_i} such that $L_{Q_i} \simeq M_{Q_i} \simeq C W_{Q_i}$ and $L_{Q_i} \cap M_{Q_i} = N_{Q_i} \simeq W_{Q_i}$.

We define the spaces L_{Q_i} and M_{Q_i} to be the images of certain subspaces of V_i^* under the homotopy equivalences defined in Lemma 7.5. Denote by

$$U_{Q_i}^{-} = \bigcup_{n=1}^{q_i} C\Sigma \widetilde{X}_{j_1} \times \cdots \times C_{-} \widetilde{X}_{j_n} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i}}$$

and

$$U_{Q_i}^+ = \bigcup_{n=1}^{q_i} C\Sigma \widetilde{X}_{j_1} \times \cdots \times C_+ \widetilde{X}_{j_n} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i}}.$$

Both $U_{Q_i}^-$ and $U_{Q_i}^+$ are contractible, since they have the homotopy type of $C_-\widetilde{X}_{j_1} * \cdots * C_-\widetilde{X}_{j_{q_i}}$ and $C_+\widetilde{X}_{j_1} * \cdots * C_+\widetilde{X}_{j_{q_i}}$, respectively. Moreover,

$$U_{Q_i}^- \cup U_{Q_i}^+ = \bigcup_{n=1}^{q_i} C\Sigma \widetilde{X}_{j_1} \times \cdots \times X_{j_n} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i}} = V_i^*$$

and

$$\begin{aligned} U_{Q_i}^- \cap U_{Q_i}^+ &= \widetilde{X}_{j_1} \times C\Sigma \widetilde{X}_{j_2} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i}} \\ &\quad \cup C_- \widetilde{X}_{j_1} \times \left(C_+ \widetilde{X}_{j_2} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i}} \cup \cdots \cup C\Sigma \widetilde{X}_{j_2} \times \cdots \times C_+ \widetilde{X}_{j_{q_i}} \right) \\ &\quad \cup \cdots \cup C\Sigma \widetilde{X}_{j_1} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i-1}} \times \widetilde{X}_{j_{q_i}} \\ &\quad \left(C_+ \widetilde{X}_{j_1} \times \cdots \times C\Sigma \widetilde{X}_{j_{q_i-1}} \cup \cdots \cup C\Sigma \widetilde{X}_{j_1} \times \cdots \times C_+ \widetilde{X}_{j_{q_i-1}} \right) \times C_- \widetilde{X}_{j_{q_i}} \\ &\simeq \widetilde{X}_{j_1} * C_+ \widetilde{X}_{j_2} * \cdots * C_+ \widetilde{X}_{j_{q_i}} \cup \cdots \cup C_+ \widetilde{X}_{j_1} * C_+ \widetilde{X}_{j_2} * \cdots * \widetilde{X}_{j_{q_i}} \\ &\simeq \Sigma^{q_i - 1} \widetilde{X}_{j_1} * \cdots * \widetilde{X}_{j_{q_i}} \end{aligned}$$

where the final homotopy equivalence is established by observation that if B_i is a contractible space containing A_i for i = 1, ..., k, then there is a homotopy equivalence

$$\bigcup_{i=1}^{k} B_1 * \cdots * A_i * \cdots * B_k \simeq \Sigma^{k-1} A_1 * \cdots * A_k.$$

To see this, for k = 2, observe that

$$(A_1 * B_2) \cap (B_1 * A_2) = (CA_1 \times B_2 \cup A_1 \times CB_2) \cap (CB_1 \times A_2 \cup B_1 \times CA_2)$$

= $CA_1 \times A_2 \cup A_1 \times CA_2 = A_1 * A_2$

and therefore $(A_1 * B_2) \cup (B_1 * A_2) \simeq \Sigma A_1 * A_2$. The general case follows by induction.

By Lemma 7.5, the homotopy equivalence of pairs

$$\beta_{j_n}^{\eta(i,\pi(j_n))} \alpha_{j_n}^{\eta(i,\pi(j_n))} \colon (C\Sigma \widetilde{X}_{j_n}, \Sigma \widetilde{X}_{j_n}) \longrightarrow (E_{j_n}^{\eta(i,\pi(j_n))}, S_{j_n}^{\eta(i,\pi(j_n))} \cup T_{j_n}^{\eta(i,\pi(j_n))})$$

sends $C_{-}\widetilde{X}_{j_{n}} \longrightarrow S_{j_{n}}^{\eta(i,\pi(j_{n}))}, C_{+}\widetilde{X}_{j_{n}} \longrightarrow T_{j_{n}}^{\eta(i,\pi(j_{n}))}$ and $\widetilde{X}_{j_{n}} \longrightarrow R_{j_{n}}^{\eta(i,\pi(j_{n}))}$. Define $L_{Q_{i}}$ and $N_{Q_{i}}$ as the images of $U_{Q_{i}}^{-}$ and $U_{Q_{i}}^{-} \cap U_{Q_{i}}^{+}$ under the product of the maps $\beta_{j_{n}}^{\eta(i,\pi(j_{n}))}\alpha_{j_{n}}^{\eta(i,\pi(j_{n}))}$. Then by construction the pair $(L_{Q_{i}}, N_{Q_{i}})$ is homotopy equivalent to $(CW_{Q_{i}}, W_{Q_{i}})$. Define also $M_{Q_{i}}$ as the complement of $L_{Q_{i}}$ in $G_{Q_{i}}^{*} \simeq \Sigma W_{Q_{i}}$. Then the pair $(M_{Q_{i}}, N_{Q_{i}})$ is also homotopy equivalent to $(CW_{Q_{i}}, W_{Q_{i}})$.

40

Construction of the pairs (F_{Q_i}, Z_{Q_i}) . For i = 1, ..., k, we construct pairs (F_{Q_i}, Z_{Q_i}) homotopy equivalent to (CW, W), such that $F_{Q_i} \cap F_{Q_{i'}} = Z_{Q_i} \cap Z_{Q_{i'}}$, that is, the pairs (F_{Q_i}, Z_{Q_i}) satisfy the conditions of Lemma 7.3.

Let ρ_i be the permutation map defined by

$$\rho_i\left(\left(x_{j_1},\ldots,x_{j_{q_i}}\right),\left(x_{i_1},\ldots,x_{i_{p_i}}\right)\right)=(x_1,\ldots,x_m)$$

and define

$$F_{Q_i} = \rho_i \left(G_{Q_i} \times \left(\bigcup_{l=1}^{p_i} CX_{i_1} \times \cdots \times X_{i_l} \times \cdots \times CX_{i_{p_i}} \right) \cup L_{Q_i} \times \left(CX_{i_1} \times \cdots \times CX_{i_{p_i}} \right) \right)$$
$$Z_{Q_i} = \rho_i \left(M_{Q_i} \times \left(\bigcup_{l=1}^{p_i} CX_{i_1} \times \cdots \times X_{i_l} \times \cdots \times CX_{i_{p_i}} \right) \cup N_{Q_i} \times \left(CX_{i_1} \times \cdots \times CX_{i_{p_i}} \right) \right).$$

Since G_{Q_i} and L_{Q_i} have the homotopy type of $C\Sigma W_{Q_i}$ and $C_+W_{Q_i}$, respectively, F_{Q_i} is homotopy equivalent to

$$CW = \rho_i \left(CW_{Q_i} * X_{i_1} * \dots * X_{p_i} \right)$$
$$= C\Sigma W_{Q_i} \times \left(\bigcup_{l=1}^{p_i} CX_{i_1} \times \dots \times X_{i_l} \times \dots \times CX_{i_{p_i}} \right) \cup C_+ W_{Q_i} \times \left(CX_{i_1} \times \dots \times CX_{i_{p_i}} \right).$$

Moreover, the homotopy equivalences $C_-W_{Q_i} \simeq M_{Q_i}$ and $W_{Q_i} \simeq N_{Q_i}$ give a homotopy equivalence between Z_{Q_i} and

$$W = \rho_i \left(W_{Q_i} * X_{i_1} * \dots * X_{i_{p_i}} \right)$$
$$= C_- W_{Q_i} \times \left(\bigcup_{l=1}^{p_i} C X_{i_1} \times \dots \times X_{i_l} \times \dots \times C X_{i_{p_i}} \right) \cup W_{Q_i} \times \left(C X_{i_1} \times \dots \times C X_{i_{p_i}} \right).$$

We now study the intersection $F_{Q_i} \cap F_{Q_{i'}}$ for $i \neq i'$.

Lemma 7.8. For $i \neq i'$, we have $F_{Q_i} \cap F_{Q_{i'}} = Z_{Q_i} \cap Z_{Q_{i'}}$. Proof. We show that if i < i' then $F_{Q_i} \cap F_{Q_{i'}} \subseteq Z_{Q_i} \cap Z_{Q_{i'}}$. Let

$$J_{Q_i} = \rho_i \left(G_{Q_i} \times \bigcup_{l=1}^{p_i} CX_{i_1} \times \dots \times X_{i_l} \times \dots \times CX_{i_{p_i}} \right)$$
$$K_{Q_i} = \rho_i \left(L_{Q_i} \times \left(CX_{i_1} \times \dots \times CX_{i_{p_i}} \right) \right)$$

so that $F_{Q_i} = J_{Q_i} \cup K_{Q_i}$. Then

$$F_{Q_i} \cap F_{Q_{i'}} = (J_{Q_i} \cap J_{Q_{i'}}) \cup (J_{Q_i} \cap K_{Q_{i'}}) \cup (J_{Q_i} \cap K_{Q_{i'}}) \cup (K_{Q_i} \cap K_{Q_{i'}}).$$

First consider $J_{Q_i} \cap J_{Q_{i'}}$. Since for $l = 1, \ldots, m$, the intersection $D_l^{\delta} \cap X_l \subseteq B_k^{\delta}$, it follows that

$$J_{Q_i} \cap J_{Q_{i'}} \subseteq \rho_i \left(G_{Q_i}^* \times \bigcup_{l=1}^{p_i} CX_{i_1} \times \dots \times X_{i_l} \times \dots \times CX_{i_{p_i}} \right)$$
$$\cap \rho_{i'} \left(G_{Q_{i'}}^* \times \bigcup_{l=1}^{p_{i'}} CX_{i'_1} \times \dots \times X_{i'_l} \times \dots \times CX_{i'_{p_{i'}}} \right)$$

By Lemma 7.7, there is $r \neq i, i'$ such that G_{Q_i} contains a factor D_r^1 and $G_{Q_{i'}}$ contains a factor D_r^- . Moreover, L_{Q_i} contains a factor E_r^1 and $L_{Q_{i'}}$ contains a factor E_r^- . Since, by construction, $D_r^1 \cap E_r^- = \emptyset = D_r^- \cap E_r^1$, we obtain

$$(J_{Q_i} \cap J_{Q_{i'}}) \cap \rho_i \left(L_{Q_i} \times \bigcup_{l=1}^{p_i} CX_{i_1} \times \dots \times X_{i_l} \times \dots \times CX_{i_{p_i}} \right) = \emptyset$$

and

$$(J_{Q_i} \cap J_{Q_{i'}}) \cap \rho_{i'} \left(L_{Q_{i'}} \times \bigcup_{l=1}^{p_{i'}} CX_{i'_1} \times \cdots \times X_{i'_l} \times \cdots \times CX_{i'_{p_{i'}}} \right) = \emptyset.$$

Therefore, since $G_{Q_i}^* \setminus L_{Q_i} = M_{Q_i}$,

$$J_{Q_i} \cap J_{Q_{i'}} \subseteq \rho_i \left(M_{Q_i} \times \bigcup_{l=1}^{p_i} CX_{i_1} \times \cdots \times X_{i_l} \times \cdots \times CX_{i_{p_i}} \right)$$
$$\cap \rho_{i'} \left(M_{Q_{i'}} \times \bigcup_{l=1}^{p_{i'}} CX_{i'_1} \times \cdots \times X_{i'_l} \times \cdots \times CX_{i'_{p_{i'}}} \right) \subseteq Z_{Q_i} \cap Z_{Q_{i'}}.$$

Moreover, the conditions $D_r^1 \cap E_r^- = \emptyset = D_r^- \cap E_r^1$ imply that $J_{Q_i} \cap K_{Q'_i} = \emptyset = K_{Q_i} \cap J_{Q_{i'}}$. Finally, since $E_r^1 \cap E_r^- = \emptyset$, $K_{Q_i} \cap K_{Q_{i'}} = \emptyset$, the claim is established. \Box

We now prove Theorem 5.6.

Proof of Theorem 5.6. Since $X_l \subseteq D_l^+ \cup D_l^2$, we represent $f_l: \Sigma X_l \longrightarrow Y_l$ by a map of pairs $f_l: (C\Sigma \widetilde{X}_l, \Sigma \widetilde{X}_l) \longrightarrow (Y_l, *)$ such that $f_l(D_l^+ \cup D_l^2) = *$.

Let $\mathcal{L} = \partial \Delta[1, \ldots, m]$ and let $\mathcal{L}_i = \partial \Delta[j_1, \ldots, j_{q_i}]$ for $i = 1, \ldots, k$. Define the map $\psi: (V, V^*) \longrightarrow (\prod_{l=1}^m Y_l, (\underline{Y}, \underline{*})^{\mathcal{L}})$ by

$$\psi(x_1,\ldots,x_m) = (f_1(x_1),\ldots,f_m(x_m))$$

and for i = 1, ..., k, define maps $\psi_i: (V_{Q_i}, V_{Q_i}^*) \longrightarrow (\prod_{l=1}^{q_i} Y_{j_l}, (\underline{Y}, \underline{*})^{\mathcal{L}_i})$ by $\psi_i = \psi|_{V_{Q_i}}$. Since $f_l(D_l^+ \cup D_l^2) = *$ for $l = 1, ..., q_i$, the map ψ_i takes the complement of

Since $f_l(D_l \cup D_l) = *$ for $i = 1, ..., q_i$, the map ψ_i takes the complement of $\prod_{l=1}^{q_i} D_l^1 \cap D_l^-$ in V_{Q_i} to $(\underline{Y}, \underline{*})^{\partial \Delta[j_1, ..., j_{q_i}]}$. Equivalently, $\psi_i((V_{Q_i} \setminus G_{Q_i}) \cup G_{Q_i}^*) \subseteq (\underline{Y}, \underline{*})^{\mathcal{L}_i}$. Therefore, $\psi_i|_{G_{Q_i}}$ is homotopic as a map of pairs to ψ_i .

Now consider

$$\begin{split} \psi \rho_i |_{\rho_i^{-1} F_{Q_i}} &: G_{Q_i} \times \left(\bigcup_{l=1}^{p_i} CX_{i_1} \times \dots \times X_{i_l} \times \dots \times CX_{i_{p_i}} \right) \cup L_{Q_i} \times \left(CX_{i_1} \times \dots \times CX_{i_{p_i}} \right) \\ & \longrightarrow \psi_i (G_{Q_i}) \times \left(\bigcup_{l=1}^{p_i} \Sigma X_{i_1} \times \dots \times * \times \dots \times \Sigma X_{i_{p_i}} \right) \cup * \times \left(\Sigma X_{i_1} \times \dots \times \Sigma_{i_{p_i}} \right) \\ & \longrightarrow \prod_{l=1}^{q_i} Y_{j_l} \times (\underline{Y}, \underline{*})^{\partial \Delta[i_1, \dots, i_{p_i}]} \cup * \times \prod_{l=1}^{p_i} Y_{i_l} \\ &= FW \left(\prod_{l=1}^{q_i} Y_{j_l}, Y_{i_1}, \dots, Y_{i_{p_i}} \right) \\ &= (\underline{Y}, \underline{*})^{\partial \Delta \langle \Delta[j_1, \dots, j_{q_i}], i_1, \dots, i_{p_i} \rangle} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{L}} \end{split}$$

and the restriction

$$\begin{split} \psi \rho_i |_{\rho_i^{-1} Z_{Q_i}} &: M_{Q_i} \times \left(\bigcup_{l=1}^{p_i} C X_{i_1} \times \dots \times X_{i_l} \times \dots \times C X_{i_{p_i}} \right) \cup N_{Q_i} \times \left(C X_{i_1} \times \dots \times C X_{i_{p_i}} \right) \\ &\longrightarrow (\underline{Y}, \underline{*})^{\mathcal{L}_i} \times (\underline{Y}, \underline{*})^{\partial \Delta[i_1, \dots, i_{p_i}]} \cup \ast \times \prod_{l=1}^{p_i} Y_{i_l} \\ &= FW\left((\underline{Y}, \underline{*})^{\mathcal{L}_i}, Y_{i_1}, \dots, Y_{i_{p_i}} \right) \\ &= (\underline{Y}, \underline{*})^{\partial \Delta \langle \partial \Delta[j_1, \dots, j_{q_i}], i_1, \dots, i_{p_i} \rangle} \longrightarrow (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}. \end{split}$$

It follows that $\psi \rho_i|_{\rho_i^{-1}FQ_i} : (\rho_i^{-1}FQ_i, \rho_i^{-1}ZQ_i) \longrightarrow ((\underline{Y}, \underline{*})^{\mathcal{L}}, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}})$ is the relative higher Whitehead map $h_w(\psi_i|_{G_{Q_i}}, f_{i_1}, \dots, f_{i_{p_i}})$. Therefore, $\psi|_{F_{Q_i}}: (F_{Q_i}, Z_{Q_i}) \longrightarrow ((\underline{Y}, \underline{*})^{\mathcal{L}}, (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}})$ is the composite $h_w(\psi_i|_{G_{Q_i}}, f_{i_1}, \dots, f_{i_{p_i}}) \circ \rho_i^{-1}$. Finally, since $\psi_i|_{G_{Q_i}}$ is homotopic to ψ_i and $\rho_i^{-1} = \sigma_i$, the restriction $\psi|_{F_{Q_i}}$ is homotopic to $h_w(\psi_i, f_{i_1}, \dots, f_{i_{p_i}}) \circ \sigma_i.$ Let $W = \Sigma^{m-2} X_1 \wedge \dots \wedge X_m, X = (\underline{Y}, \underline{*})^{\mathcal{L}}$ and $A = (\underline{Y}, \underline{*})^{\mathcal{K}_{\Pi}}.$ Consider long exact

sequence (47)

$$\cdots \longrightarrow [\Sigma W, X] \xrightarrow{j} [(CW, W), (X, A)] \xrightarrow{\partial} [W, A] \longrightarrow \cdots$$

By Proposition 7.2 and Lemma 7.3,

$$\partial j(\psi|_{V^*}) \simeq \partial \left(\sum_{i=1}^k \psi|_{F_{Q_i}}\right)$$
$$\simeq \sum_{i=1}^k \partial h_w(\psi_i, f_{i_1}, \dots, f_{i_{p_i}}) \circ \sigma_i$$
$$= \sum_{i=1}^k h_w(\psi_i|_{V_{Q_i}^*}, f_{i_1}, \dots, f_{i_{p_i}}) \circ \sigma_i$$
$$= \sum_{i=1}^k h_w(h_w(f_{j_1}, \dots, f_{j_{q_i}}), f_{i_1}, \dots, f_{i_{p_i}}) \circ \sigma_i.$$

On the other hand, by exactness, ∂j is the zero homomorphism, completing the proof.

References

- [1] S. A. Abramyan. Iterated higher Whitehead products in the topology of moment-angle complexes. Sibirsk. Mat. Zh., 60(2):243-256, 2019.
- [2] S. A. Abramyan and T. E. Panov. Higher Whitehead products for moment-angle complexes and substitutions of simplicial complexes. Tr. Mat. Inst. Steklova, 305(Algebraicheskaya Topologiya Kombinatorika i Matematicheskaya Fizika):7-28, 2019.
- [3] A. Anton. Composition of simplicial complexes, polytopes and multigraded betti numbers, 2013.
- [4] M. Arkowitz. The generalized Whitehead product. Pacific Journal of Mathematics, 12(1):7 -23, 1962.

- [5] A. L. Blakers and W. S. Massey. The homotopy groups of a triad. iii. Annals of Mathematics, 58(3):409–417, 1953.
- [6] A. L. Blakers and W. S. Massey. Products in homotopy theory. Annals of Mathematics, 58(2):295–324, 1953.
- [7] V. M. Buchstaber and T. E. Panov. Toric topology, volume 204 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2015.
- [8] K. A. Hardie. Higher Whitehead products. The Quarterly Journal of Mathematics, 12(1):241– 249, 01 1961.
- [9] K. A. Hardie. A proof of the Nakaoka-Toda formula. Pacific Journal of Mathematics, 14(4):1249 - 1254, 1964.
- [10] M. Nakaoka and H. Toda. On Jacobi identity for Whitehead products. Journal of the Institute of Polytechnics, Osaka City University. Series A: Mathematics, 5(1):1 – 13, 1954.
- [11] G. J. Porter. Higher order Whitehead products. *Topology*, 3:123–135, 1965.
- [12] H. Samelson. A connection between the Whitehead and the Pontryagin product. American Journal of Mathematics, 75(4):744–752, 1953.
- [13] H. Uehara and W. S. Massey. The Jacobi identity for Whitehead products. 1957.
- [14] E. Vidaurre. On polyhedral product spaces over polyhedral joins. Homology, Homotopy and Applications, 20, 09 2017.
- [15] E. Zhuravleva. Adams-Hilton models and higher Whitehead brackets for polyhedral products. arXiv:2112.10013, 2021.