

ANISOTROPIC FLOWS OF FORCHHEIMER-TYPE IN POROUS MEDIA AND THEIR STEADY STATES

LUAN HOANG¹ AND THINH KIEU²

ABSTRACT. We study the anisotropic Forchheimer-typed flows for compressible fluids in porous media. The first half of the paper is devoted to understanding the nonlinear structure of the anisotropic momentum equations. Unlike the isotropic flows, the important monotonicity properties are not automatically satisfied in this case. Therefore, various sufficient conditions for them are derived and applied to the experimental data. In the second half of the paper, we prove the existence and uniqueness of the steady state flows subject to a nonhomogeneous Dirichlet boundary condition. It is also established that these steady states, in appropriate functional spaces, have local Hölder continuous dependence on the forcing function and the boundary data.

CONTENTS

1. Introduction	1
2. Preliminaries	4
2.1. Notation and inequalities	4
2.2. Functional spaces	5
3. Fundamental properties (I)	6
3.1. The two-dimensional case	8
3.2. The case of general dimensions	15
4. Fundamental properties (II)	17
5. Steady state flows	22
Forcing function.	23
Boundary data	23
5.1. Variational formulation on $V \times Z$	24
5.2. The regularized problem	26
5.3. Existence and uniqueness	30
5.4. Estimates and continuous dependence	30
5.5. Proof of Theorem 5.3	32
Appendix A.	32
References	34

1. INTRODUCTION

We study fluid flows in porous media. In this section, $p \in \mathbb{R}$ is the pressure, $\mathbf{v} \in \mathbb{R}^3$ is the velocity, $\mu > 0$ is the dynamic viscosity, $\rho \geq 0$ is the density, $k > 0$ is the permeability, and $\phi \in (0, 1)$ is the constant porosity.

Date: June 9, 2023.

2020 Mathematics Subject Classification. 76S05 , 76N10, 35F60.

Key words and phrases. porous media, anisotropic, Forchheimer, fluid flows, monotonicity, nonlinear partial differential equations, first-order system.

The standard momentum equation for fluid flows in porous media is the Darcy's law

$$-\nabla p = \frac{\mu}{k} \mathbf{v}.$$

This is used widely for flows with low Reynolds numbers. For higher Reynolds numbers, Forchheimer's equations are used, see [3, 11, 12, 19, 20]. The Forchheimer two-term law is

$$\frac{\mu}{k} \mathbf{v} + \frac{c_F \rho}{\sqrt{k}} |\mathbf{v}| \mathbf{v} = -\nabla p, \text{ where } c_F > 0. \quad (1.1)$$

The Forchheimer three-term law is

$$a_0 \mathbf{v} + a_1 |\mathbf{v}| \mathbf{v} + a_2 |\mathbf{v}|^2 \mathbf{v} = -\nabla p, \text{ where } a_0, a_1, a_2 > 0.$$

The Forchheimer power law is

$$\frac{\mu}{k} \mathbf{v} + c |\mathbf{v}|^{m-1} \mathbf{v} = -\nabla p, \text{ with } c > 0 \text{ and } m > 1.$$

The generalized Forchheimer equations are

$$\sum_{i=0}^m a_i |\mathbf{v}|^{\alpha_i} \mathbf{v} = -\nabla p, \quad (1.2)$$

where a_i 's are positive numbers and $0 = \alpha_0 < \alpha_1 < \dots < \alpha_m$. All of the above equations are written for isotropic porous media.

In their paper [1], Barak and Bear formulate many physical and mathematical models for fluid flows at high Reynolds numbers in anisotropic porous media. Their experiments confirm one of them, see equations (38)–(40) in [1]. We will take this as our starting point and consider the following equation

$$A_0 \mathbf{v} + |\mathbf{v}| A \mathbf{v} + \frac{1}{|\mathbf{v}|} B(\mathbf{v}, \mathbf{v}, \mathbf{v}) = -\nabla p, \quad (1.3)$$

where A_0 and A are square matrices, and $B(\cdot, \cdot, \cdot)$ is a trilinear mapping.

The first term in (1.3) is the anisotropic Darcy term. The second term in (1.3) is from equations (25), (27) and (30) in [1]. The third term in (1.3) is from equations (38)–(40) in [1].

Moreover, experiments reported in [1] yield a particular case for equation (1.3), namely,

$$\nu A_0 \mathbf{v} + \frac{1}{|\mathbf{v}|} \begin{pmatrix} (av_1^2 + bv_2^2)v_1 \\ (cv_1^2 + dv_2^2)v_2 \end{pmatrix} = -g \nabla p \text{ for } \mathbf{v} = (v_1, v_2) \in \mathbb{R}^2, \quad (1.4)$$

where ν is the kinetic viscosity, g is the gravity acceleration, $A_0 = \text{diag}[109, 220]$, and a, b, c, d are the following positive constants

$$a = 0.20, \quad b = 1.04, \quad c = 0.67, \quad d = 1.15. \quad (1.5)$$

When the dependence on the density is needed, we can use the dimension analysis, see [19], to modify equation (1.3). Similar to (1.1), it results in the following equation

$$A_0 \mathbf{v} + \rho |\mathbf{v}| A \mathbf{v} + \frac{\rho}{|\mathbf{v}|} B(\mathbf{v}, \mathbf{v}, \mathbf{v}) = -\nabla p. \quad (1.6)$$

The momentum equation (1.6) is coupled with the conservation of mass which is

$$\phi \rho_t + \nabla \cdot (\rho \mathbf{v}) = f, \quad (1.7)$$

where $f(x, t)$ is the source/sink term.

Multiplying equation (1.6) by ρ gives

$$A_0(\rho \mathbf{v}) + |\rho \mathbf{v}| A(\rho \mathbf{v}) \frac{1}{|\rho \mathbf{v}|} B(\rho \mathbf{v}, \rho \mathbf{v}, \rho \mathbf{v}) = -\rho \nabla p. \quad (1.8)$$

Denote the momentum by $\mathbf{m} = \rho \mathbf{v}$. Defining $F(\mathbf{v})$ to be the left-hand side of (1.3), we then obtain from (1.8) and (1.7) that

$$F(\mathbf{m}) = -\rho \nabla p, \quad \phi \rho_t + \nabla \cdot \mathbf{m} = f. \quad (1.9)$$

The case of isentropic flows. We have $p = \bar{c} \rho^\gamma$ with $\bar{c}, \gamma > 0$. Set

$$\tilde{p} = \frac{\bar{c}\gamma}{\gamma+1} \rho^{\gamma+1}, \quad \lambda = \frac{1}{\gamma+1}, \quad c = \left(\frac{\gamma+1}{\bar{c}\gamma} \right)^\lambda.$$

Then the system (1.9) becomes

$$F(\mathbf{m}) = -\nabla \tilde{p}, \quad c \phi (\tilde{p}^\lambda)_t + \nabla \cdot \mathbf{m} = f. \quad (1.10)$$

In particular, $\gamma = 1$ and $\lambda = 1/2$ for ideal fluids.

The case of slightly compressible flows. We have

$$\frac{1}{\rho} \frac{d\rho}{dp} = \kappa, \quad \text{where } \kappa \text{ is a positive constant.}$$

Observe that the first equation in (1.9) now is $F(\mathbf{m}) = -\kappa^{-1} \nabla \rho$. By setting

$$\tilde{p} = \rho/\kappa, \quad \lambda = 1 \text{ and } c = \kappa,$$

we obtain the system (1.10) again from (1.9)

Our next task is to analyze (1.10) which is a system of nonlinear partial differential equations (PDE). First, we study the nonlinear structure of the momentum equation (1.6) and its generalizations. The key properties that we investigate are the monotonicities for the function F , see Definition 3.2. Because of the counter examples 3.3 and 4.2, many sufficient conditions need be developed. Once the monotonicities are obtained, we can solve for $\mathbf{m} = -X(\nabla \tilde{p})$ from the first equation of (1.10) and substitute it into the second equation. It results in a second order scalar parabolic equation

$$c \phi \frac{\partial(\tilde{p}^\lambda)}{\partial t} + \nabla \cdot (X(\nabla \tilde{p})) = f.$$

This approach was widely exploited in our previous work, see [4, 5, 7, 8, 14–17] and references therein, and also [2, 9, 10, 13, 21] for the stationary problems and their numerical analysis. In the current work, we follow a more straightforward approach, namely, studying system (1.10) directly as a system of the first order PDE. As a demonstration, we will focus on the steady state flows with a nonhomogeneous Dirichlet boundary condition. We obtain the existence and uniqueness of these anisotropic flows for any order $s > 2$ in Assumption 5.1. They are more general than the counterpart in [18] for the isotropic Forcheimer flows corresponding to $s = 3$, as well as [10] for isotropic Forchheimer's power laws with the flux boundary condition. Moreover, we establish explicit estimates and continuous dependence results which are not obtained in [10, 18].

The paper is organized as follows. In Section 2, we present inequalities that will be used throughout the paper. Also, relevant functional spaces and their important properties are recalled. In Section 3, we study properties of the function G_B defined by (3.1) and (3.2). It is the n -dimensional version of the last term on the left-hand side of (1.3). Their upper and lower bounds, and Lipschitz estimates are obtained in Lemma 3.1. For our mathematical treatments, the monotonicities in Definition 3.2 are crucial. However, such monotonicities are not automatic under the general positivity condition for the coefficients, see Example 3.3. Therefore, the large part of this section is devoted to finding sufficient conditions for the monotonicities. Subsection 3.1 treats the two-dimensional case, while subsection 3.2 does the higher dimensional case. Our sufficient conditions turn out to be applicable to the real data obtained in [1], see Example 3.10. In Section 4, we consider other possible anisotropic flows in the form of the functions $F_{A,\alpha}$ and $\tilde{F}_{A,\alpha}$, see (4.2) and (4.3). These

represent different anisotropic versions of (1.2). While $\widetilde{F}_{A,\alpha}$ is $(\alpha + 2)$ -monotone under the coercivity condition (4.9), the function $F_{A,\alpha}$ may not be so, see Example 4.2. Therefore, many sufficient conditions for the monotonicities are derived in Theorems 4.3, 4.4 and 4.5. At the end of the section, many scenarios for the momentum equations are shown in Proposition 4.8. In Section 5, we study the steady state flows for a general system of the form (1.10) subject to a time-independent nonhomogeneous Dirichlet boundary condition, see Problem (5.1). Thanks to our understanding in Sections 3 and 4, natural and general conditions on the function F are imposed in Assumption 5.1. The variational formulation in suitable functional spaces for the weak solutions is established, see Definition 5.2. Theorem 5.3 contains the main results for system (5.1) including the existence, uniqueness of the weak solutions, together with their estimates and continuous dependence on the forcing function and boundary data. For its proof, we generalize the method used in [18]. In fact, a general functional equation is studied instead, see (5.20). Its regularized problem is introduced and investigated in subsection 5.2. Using this approximation, we obtain the existence and uniqueness in Theorem 5.12. Estimates of the solutions are obtained in Theorem 5.13 and the continuous dependence result is established in Theorem 5.14. Gathering the above general results gives a quick proof of Theorem 5.3 in subsection 5.5.

2. PRELIMINARIES

2.1. Notation and inequalities. For a vector $x \in \mathbb{R}^n$, its Euclidean norm is denoted by $|x|$.

Let $A = (a_{ij})_{1 \leq i, j \leq n}$ be an $n \times n$ matrices of real numbers. The Euclidean norm of A is $|A| = \left(\sum_{i, j=1}^n a_{ij}^2 \right)^{1/2}$. (We do not use $|A|$ to denote the determinant in this paper.) When A is considered as a linear operator, another norm is defined by

$$\|A\|_{\text{op}} = \max \left\{ \frac{|Ax|}{|x|} : x \in \mathbb{R}^n, x \neq 0 \right\} = \max\{|Ax| : x \in \mathbb{R}^n, |x| = 1\}.$$

It is well-known that

$$\|A\|_{\text{op}} \leq |A| \leq \bar{c}_0 \|A\|_{\text{op}}, \quad (2.1)$$

where $\bar{c}_0 = \bar{c}_0(n)$ is a positive constant independent of A .

For any $x \in \mathbb{R}^n$, we have

$$|Ax| \leq \|A\|_{\text{op}} |x| \leq |A| |x|, \quad (2.2)$$

and, in the case A is invertible,

$$|Ax| \geq \|A^{-1}\|_{\text{op}}^{-1} |x|. \quad (2.3)$$

The following are some commonly used consequences of Young's inequality. If $x, y \geq 0$, $\gamma \geq \beta \geq \alpha > 0$, $p, q > 1$ with $1/p + 1/q = 1$, and $\varepsilon > 0$, then

$$x^\alpha \leq 1 + x^\beta, \quad x^\beta \leq x^\alpha + x^\gamma, \quad xy \leq \varepsilon x^p + \varepsilon^{-q/p} y^q. \quad (2.4)$$

For $z \in \mathbb{R}$, denote $z^+ = \max\{0, z\}$. For $x, y \in \mathbb{R}^n$ and $p > 0$, one has

$$(|x| + |y|)^p \leq 2^{(p-1)^+} (|x|^p + |y|^p) = \begin{cases} |x|^p + |y|^p, & \text{for } p \in (0, 1], \\ 2^{p-1} (|x|^p + |y|^p), & \text{for } p > 1, \end{cases} \quad (2.5)$$

which consequently yields

$$(|x| + |y|)^p \leq 2^p (|x|^p + |y|^p) \quad \text{for } p > 0, \quad (2.6)$$

$$\left| |x|^p - |y|^p \right| \leq |x - y|^p \quad \text{for } p \in (0, 1]. \quad (2.7)$$

Regarding the last inequality, one has, in the case $p > 1$, that

$$\left| |x|^p - |y|^p \right| \leq 2^{(p-2)^+} (|x|^{p-1} + |y|^{p-1}) |x - y|. \quad (2.8)$$

Lemma 2.1. *If $p > 0$ and $x, y \in \mathbb{R}^n$, then*

$$||x|^p x - |y|^p y| \leq 2^{(p-1)^+} (|x|^p + |y|^p) |x - y|, \quad (2.9)$$

$$(|x|^p x - |y|^p y) \cdot (x - y) \geq \frac{1}{2} (|x|^p + |y|^p) |x - y|^2, \quad (2.10)$$

$$(|x|^p x - |y|^p y) \cdot (x - y) \geq \frac{1}{2^{1+(p-1)^+}} |x - y|^{p+2}. \quad (2.11)$$

If $p \in (-1, 0)$, then

$$||x|^p x - |y|^p y| \leq 2^{-p} |x - y|^{1+p} \quad \text{for all } x, y \in \mathbb{R}, \quad (2.12)$$

$$(|x|^p x - |y|^p y) \cdot (x - y) \geq (1 + p) (|x| + |y|)^p |x - y|^2 \quad \text{for all } x, y \in \mathbb{R}^n. \quad (2.13)$$

It is meant, naturally, in (2.12) and (2.13) that

$$|x|^p x, |y|^p y, (|x| + |y|)^p |x - y|^2 = 0 \text{ for } p \in (-1, 0) \text{ and } x = y = 0.$$

The inequalities (2.8)–(2.13) contain explicit constants and many of them are the best ones. For example, inequalities (2.10), (2.12), and particular cases of (2.8), respectively, (2.9), (2.11), when $p = 2$, respectively, $p = 1$, $p \geq 1$, have the best constants. Elementary proofs of these inequalities are provided in Appendix A below. For inequalities without explicit constants, see [9, 13].

We recall a simple yet very useful inequality from [6].

Lemma 2.2 ([6, from (2.29) to (2.31)]). *Let $x, y \in \mathbb{R}^n$ and $\alpha > 0$. Then*

$$\int_0^1 |tx + (1-t)y|^\alpha dt \geq \frac{|x - y|^\alpha}{2^{\alpha+1}(\alpha + 1)}. \quad (2.14)$$

2.2. Functional spaces. Next, we review the Sobolev spaces and trace theorems. *Hereafter, the spatial dimension $n \geq 2$ is fixed.*

Let Ω be an open, bounded subset of \mathbb{R}^n with the boundary $\partial\Omega$ of class C^1 .

For $1 \leq s < \infty$, let $L^s(\Omega)$ be the standard Lebesgue space of scalar functions and denote $\mathbf{L}^s(\Omega) = (L^s(\Omega))^n$. The notation $\|\cdot\|_{0,s}$ is used to denote both norms $\|\cdot\|_{L^s(\Omega)}$ and $\|\cdot\|_{\mathbf{L}^s(\Omega)}$.

For a nonnegative integer m , let $W^{m,p}(\Omega)$ be the standard Sobolev space with the norm

$$\|u\|_{m,s} = \left(\sum_{|\alpha| \leq m} \|D^\alpha u\|_{L^s(\Omega)}^s \right)^{\frac{1}{s}}.$$

For any normed space X , its dual space is denoted by X' and the product between X' and X is denoted by $\langle \cdot, \cdot \rangle_{X',X}$, i.e., $\langle y, x \rangle_{X',X} = y(x)$ for $y \in X'$ and $x \in X$.

Consider $1 < s < \infty$ now. The function $\gamma_0 : f \in C^\infty(\bar{\Omega}) \mapsto f|_{\partial\Omega}$ can be extended to a bounded linear mapping $\gamma_{0,s} : W^{1,s}(\Omega) \rightarrow L^s(\partial\Omega)$. The function $\gamma_{0,s}(f)$ is called the trace of f on $\partial\Omega$.

Define $X_s = W^{1-1/s,s}(\partial\Omega)$ to be the range of $\gamma_{0,s}$ equipped with the norm

$$\|f\|_{X_s} = \inf\{\|\varphi\|_{1,s} : \varphi \in W^{1,s}(\Omega), \gamma_{0,s}(\varphi) = f\}.$$

Define the space

$$\mathbf{W}_s(\text{div}, \Omega) = \{\mathbf{v} \in \mathbf{L}^s(\Omega) : \text{div } \mathbf{v} \in L^s(\Omega)\} \quad (2.15)$$

equipped with the norm

$$\|\mathbf{v}\|_{\mathbf{W}_s(\text{div}, \Omega)} = (\|\mathbf{v}\|_{0,s}^s + \|\nabla \cdot \mathbf{v}\|_{0,s}^s)^{1/s}. \quad (2.16)$$

Then $\mathbf{W}_s(\text{div}, \Omega)$ is a reflexive Banach space, see Lemma A.1 below.

Let $r > 1$ be the Hölder conjugate of s , i.e., $1/s + 1/r = 1$. Thanks to (2.5),

$$\|\mathbf{v}\|_{\mathbf{W}_s(\text{div}, \Omega)} \leq \|\mathbf{v}\|_{0,s} + \|\nabla \cdot \mathbf{v}\|_{0,s} \leq 2^{1/r} \|\mathbf{v}\|_{\mathbf{W}_s(\text{div}, \Omega)}. \quad (2.17)$$

Let $\vec{\nu}$ denote the outward normal vector to the boundary $\partial\Omega$. Then one can extend the normal trace $\gamma_n(\mathbf{v}) = \mathbf{v} \cdot \vec{\nu}$ for $\mathbf{v} \in (C^\infty(\bar{\Omega}))^n$ to a bounded, linear mapping $\gamma_{n,s}$ from $\mathbf{W}_s(\text{div}, \Omega)$ into X'_r . In particular, there is $\bar{c}_1 > 0$ such that

$$\|\gamma_{n,s}(\mathbf{v})\|_{X'_r} \leq \bar{c}_1 \|\mathbf{v}\|_{\mathbf{W}_s(\text{div}, \Omega)} \quad \text{for all } \mathbf{v} \in \mathbf{W}_s(\text{div}, \Omega), \quad (2.18)$$

and Green's formula

$$\int_{\Omega} \mathbf{v} \cdot \nabla q \, dx + \int_{\Omega} \nabla \cdot \mathbf{v} q \, dx = \langle \gamma_{n,s}(\mathbf{v}), \gamma_{0,r}(q) \rangle_{X'_r, X_r} \quad (2.19)$$

holds for every $\mathbf{v} \in \mathbf{W}_s(\text{div}, \Omega)$ and $q \in W^{1,r}(\Omega)$. In fact, the product $\langle \gamma_{n,s}(\mathbf{v}), \gamma_{0,r}(q) \rangle_{X'_r, X_r}$ in (2.19) is the extension of the boundary integral

$$\int_{\partial\Omega} (\mathbf{v} \cdot \vec{\nu}) q \, d\sigma \quad \text{for } \mathbf{v} \in (W^{1,s}(\Omega))^n, q \in W^{1,r}(\Omega).$$

Finally, we recall an important norm estimate, see [2, Inequality (4.2)] or [18, Lemma A.3].

Lemma 2.3. *Let $r, s \in (1, \infty)$ be Hölder conjugates of each other and let $V = \mathbf{W}_s(\text{div}, \Omega)$. Then there exists a constant $C_* > 0$ such that, for all $q \in L^r(\Omega)$, it holds*

$$\|q\|_{0,r} \leq C_* \sup_{\mathbf{v} \in V \setminus \{0\}} \frac{\int_{\Omega} (\text{div } \mathbf{v}) q \, dx}{\|\mathbf{v}\|_V}. \quad (2.20)$$

A direct proof of Lemma 2.3 is given in Appendix A.

3. FUNDAMENTAL PROPERTIES (I)

Let $B : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be defined by

$$B(u, v, w) = \text{diag}[u^T A_1 v, \dots, u^T A_n v] w \quad \text{for } u, v, w \in \mathbb{R}^n, \quad (3.1)$$

where each A_i , for $i = 1, 2, \dots, n$, is a nonzero $n \times n$ matrix. This form of B is a generalization of the corresponding term in (1.4).

Our focus in this section is the function $G_B : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by

$$G_B(u) = \begin{cases} 0 & \text{for } u = 0, \\ \frac{1}{|u|} B(u, u, u) & \text{for } u \in \mathbb{R}^n \setminus \{0\}. \end{cases} \quad (3.2)$$

The following are immediate properties of B and G_B .

Lemma 3.1. *The following statements hold true.*

- (i) B is a trilinear mapping, and hence, it is continuous.
- (ii) If A_i is symmetric for all $i = 1, 2, \dots, n$, then

$$B(u, v, w) = B(v, u, w) \quad \text{for any } u, v, w \in \mathbb{R}^n. \quad (3.3)$$

- (iii) If there is $C > 0$ such that, for all $i = 1, 2, \dots, n$,

$$u^T A_i u \geq C |u|^2 \quad \text{for any } u, v \in \mathbb{R}^n, \quad (3.4)$$

then one has

$$B(u, u, u) \cdot u \geq C |u|^4 \quad \text{and} \quad G_B(u) \cdot u \geq C |u|^3 \quad \text{for all } u \in \mathbb{R}^n. \quad (3.5)$$

- (iv) For any $u, v, w \in \mathbb{R}^n$,

$$|B(u, v, w)| \leq M_* |u| |v| |w|, \quad \text{where } M_* = \left(\sum_{i=1}^n |A_i|^2 \right)^{1/2}. \quad (3.6)$$

Consequently,

$$|G_B(u)| \leq M_* |u|^2 \quad \text{for all } u \in \mathbb{R}^n. \quad (3.7)$$

(v) The function G_B is continuous. More explicitly, one has, for any $u, v \in \mathbb{R}^n$,

$$|G_B(u) - G_B(v)| \leq 2M_*(|u| + |v|)|u - v|. \quad (3.8)$$

Proof. Parts (i)–(iii) are obvious. For part (iv), we have

$$|B(u, v, w)| \leq \left(\sum_{i=1}^n |u^T A_i v|^2 \right)^{1/2} |w| \leq \left(\sum_{i=1}^n |A_i|^2 \right)^{1/2} |u||v||w|,$$

which proves (3.6). It is clear that inequality (3.7) follows (3.6) with $u = v = w \neq 0$ and $G_B(0) = 0$.

We prove part (v) now. The fact that G_B is continuous comes from part (i) and (3.7). To prove (3.8), we first consider the case when the line segment connecting u and v does not contain the origin. For $t \in [0, 1]$, let $w(t) = tu + (1 - t)v$. Then $w(t) \neq 0$ for all $t \in [0, 1]$. Define $z = u - v$ and

$$h(t) = \frac{1}{|w(t)|} B(w(t), w(t), w(t)) \text{ for } t \in [0, 1].$$

Applying the Fundamental Theorem of Calculus to the function $h(t)$ on the interval $[0, 1]$, we obtain

$$|G_B(u) - G_B(v)| = |h(1) - h(0)| \leq \int_0^1 |h'(t)| dt. \quad (3.9)$$

Calculating $h'(t)$, we have

$$h'(t) = -\frac{1}{|w|^2} \frac{w \cdot z}{|w|} B(w, w, w) + \frac{1}{|w|} (B(z, w, w) + B(w, z, w) + B(w, w, z)).$$

Applying inequality (3.6) yields

$$|h'(t)| \leq 4M_* |w||z|. \quad (3.10)$$

Note that

$$\int_0^1 |w(t)| dt \leq \int_0^1 t|u| + (1 - t)|v| dt = \frac{1}{2}(|u| + |v|). \quad (3.11)$$

Combining (3.9), (3.10) and (3.11) yields (3.8).

Now, consider the case when the line segment connecting u and v contains the origin. Since the dimension n is greater than one, for $\varepsilon > 0$ sufficiently small, we can find $u^\varepsilon, v^\varepsilon \in \mathbb{R}^n \setminus \{0\}$ such that the line segment connecting u^ε and v^ε does not contain the origin, and $u^\varepsilon \rightarrow u, v^\varepsilon \rightarrow v$ as $\varepsilon \rightarrow 0$. Applying the inequality (3.8) to u^ε and v^ε , and then letting $\varepsilon \rightarrow 0$, we obtain (3.8) for u and v . \square

Although the coercivity of G_B in (3.5) is useful in later investigation of the corresponding systems of PDE, it is not enough. The following monotonicities will be needed.

Definition 3.2. Let F be a function from \mathbb{R}^n to \mathbb{R}^n .

(i) We say F is monotone if

$$(F(u) - F(v)) \cdot (u - v) \geq 0 \text{ for any } u, v \in \mathbb{R}^n.$$

(ii) For $\alpha > 0$, we say F is (power) α -monotone if there exists a constant $C > 0$ such that

$$(F(u) - F(v)) \cdot (u - v) \geq C|u - v|^\alpha \text{ for any } u, v \in \mathbb{R}^n.$$

Clearly, if F is α -monotone, then it is monotone. Our goal is to establish the monotonicities in Definition 3.2 for the function G_B .

3.1. The two-dimensional case. We consider the dimension $n = 2$ and a special case of B in (3.1). Specifically, given $a, b, c, d > 0$, we set

$$A_1 = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, \quad A_2 = \begin{pmatrix} c & 0 \\ 0 & d \end{pmatrix},$$

and write (3.1) explicitly as

$$B(u, v, w) = \begin{pmatrix} u^T A_1 v & 0 \\ 0 & u^T A_2 v \end{pmatrix} w = \begin{pmatrix} (au_1v_1 + bu_2v_2)w_1 \\ (cu_1v_1 + du_2v_2)w_2 \end{pmatrix}$$

for vectors $u = (u_1, u_2)$, $v = (v_1, v_2)$ and $w = (w_1, w_2)$ in \mathbb{R}^2 .

Then (3.4) is true with $c_0 = \min\{a, b, c, d\}$. Moreover, A_1 and A_2 are symmetric and, consequently, we have property (3.3).

For the sake of convenience in the calculations below, we denote $|v|_i = (v^T A_i v)^{1/2}$ for $i = 1, 2$. Then each $|\cdot|_i$ is a norm on \mathbb{R}^n .

Example 3.3 (Counter example for the monotonicities). We give an example to show that the monotonicities in Definition 3.2 are not automatically satisfied for any positive values of a, b, c, d .

- (a) Let $a = d = 1$, $b = c = 5$, $u_* = (2, 2)$ and $v_* = (3, 1)$. Then $|u_*| = 2\sqrt{2}$, $|v_*| = \sqrt{10}$, $B(u_*, u_*, u_*) = 48(1, 1)$, $B(v_*, v_*, v_*) = (42, 46)$ and $u_* - v_* = (-1, 1)$. We have

$$\left(\frac{1}{|u_*|} B(u_*, u_*, u_*) - \frac{1}{|v_*|} B(v_*, v_*, v_*) \right) \cdot (u_* - v_*) = \frac{-4}{\sqrt{10}} < 0.$$

Thus, G_B is not monotone.

- (b) Let A_0 be any 2×2 matrix. Set $u = tu_*$ and $v = tv_*$ for $t > 0$. Then

$$\left(A_0(u - v) + \frac{1}{|u|} B(u, u, u) - \frac{1}{|v|} B(v, v, v) \right) \cdot (u - v) = t^2(-1, 1) \cdot A_0(-1, 1) - \frac{4t^3}{\sqrt{10}}$$

which is negative for sufficiently large t . Therefore, the function $u \mapsto A_0 u + G_B(u)$ is not monotone.

Because of Example 3.3, it is not guaranteed that the experimental data (1.5) yields a corresponding monotone function G_B . Therefore, different sufficient conditions are needed.

Theorem 3.4. *The following statements hold true.*

- (i) *If*

$$\left(b + c - \frac{|w|_1^2 + |w|_2^2}{2} \right)^2 \leq 4ad \text{ for all } w \in \mathbb{R}^2 \text{ with } |w| = 1, \quad (3.12)$$

then G_B is monotone.

- (ii) *If the inequality in (3.12) is strict, then G_B is 3-monotone.*

Proof. We prove part (i) first. Assume (3.12). Let $u, v \in \mathbb{R}^2$. By the same perturbation arguments in the last part of the proof of Lemma 3.1, we can assume, without loss of generality, that the line segment connecting u and v does not contain the origin.

Let $z = u - v$, and, for $t \in [0, 1]$, $w(t) = tu + (1 - t)v$ and

$$h(t) = \frac{1}{|w(t)|} B(w(t), w(t), w(t)) \cdot z. \quad (3.13)$$

Note that $w(t) \neq 0$ for all $t \in [0, 1]$. We have

$$I \stackrel{\text{def}}{=} \left(\frac{1}{|u|} B(u, u, u) - \frac{1}{|v|} B(v, v, v) \right) \cdot (u - v) = h(1) - h(0) = \int_0^1 h'(t) dt. \quad (3.14)$$

We calculate the derivative

$$\begin{aligned} h'(t) &= -\frac{w \cdot z}{|w|^3} B(w, w, w) \cdot z + \frac{1}{|w|} (B(z, w, w) + B(w, z, w) + B(w, w, z)) \cdot z \\ &= \frac{1}{|w|} (J_1 + J_2 + J_3), \end{aligned}$$

where

$$J_1 = -\frac{w \cdot z}{|w|^2} B(w, w, w) \cdot z, \quad J_2 = (B(w, z, w) + B(z, w, w)) \cdot z, \quad J_3 = B(w, w, z) \cdot z.$$

More specifically,

$$\begin{aligned} J_1 &= -\frac{w \cdot z}{|w|^2} \left[(w^T A_1 w) w_1 z_1 + (w^T A_2 w) w_2 z_2 \right] \\ &= -\frac{1}{|w|^2} \left[|w|_1^2 w_1^2 z_1^2 + (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2 + |w|_2^2 w_2^2 z_2^2 \right]. \end{aligned} \quad (3.15)$$

By the symmetry (3.3),

$$\begin{aligned} J_2 &= 2B(w, z, w) \cdot z = 2 \left[(w^T A_1 z) w_1 z_1 + (w^T A_2 z) w_2 z_2 \right] \\ &= 2 \left[a w_1^2 z_1^2 + (b + c) w_1 w_2 z_1 z_2 + d w_2^2 z_2^2 \right]. \end{aligned} \quad (3.16)$$

The last term J_3 is

$$J_3 = (w^T A_1 w) z_1^2 + (w^T A_2 w) z_2^2 = |w|_1^2 z_1^2 + |w|_2^2 z_2^2. \quad (3.17)$$

Combining the terms J_1 and J_3 , we have

$$J_1 + J_3 = \frac{1}{|w|^2} \left[-|w|_1^2 w_1^2 z_1^2 - (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2 - |w|_2^2 w_2^2 z_2^2 + (|w|_1^2 z_1^2 + |w|_2^2 z_2^2) (w_1^2 + w_2^2) \right]$$

which yields

$$J_1 + J_3 = \frac{1}{|w|^2} \left[|w|_1^2 z_1^2 w_2^2 + |w|_2^2 z_2^2 w_1^2 - (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2 \right] \quad (3.18)$$

It follows (3.18) that

$$J_1 + J_3 \geq -\frac{1}{|w|^2} (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2.$$

Combining this with formula (3.16) of J_2 , we obtain

$$h'(t) \geq \frac{2}{|w|} \left\{ a w_1^2 z_1^2 + \left[(b + c) - \frac{|w|_1^2 + |w|_2^2}{2|w|^2} \right] w_1 w_2 z_1 z_2 + d w_2^2 z_2^2 \right\}. \quad (3.19)$$

Fix $t \in [0, 1]$. Let $\alpha(t) = w_1(t) z_1$ and $\beta(t) = w_2(t) z_2$. The expression between the parentheses on the right-hand side of (3.19) is $Q(\alpha(t), \beta(t))$, where Q is the following t -dependent quadratic function

$$Q(\alpha, \beta) = a\alpha^2 + \left[(b + c) - \frac{|w(t)|_1^2 + |w(t)|_2^2}{2|w(t)|^2} \right] \alpha\beta + d\beta^2 \text{ for } \alpha, \beta \in \mathbb{R}.$$

By (3.12), we have $Q(\alpha, \beta) \geq 0$ for all $\alpha, \beta \in \mathbb{R}$. Consequently, $h'(t) \geq 0$ for all $t \in [0, 1]$, which gives

$$\left(\frac{1}{|u|} B(u, u, u) - \frac{1}{|v|} B(v, v, v) \right) \cdot (u - v) \geq 0 \quad (3.20)$$

thanks to (3.14). Therefore, G_B is monotone.

We prove part (ii) now. Assume (3.12) with the strict inequality. Similar to the proof of part (i), we can assume that the line segment connecting u and v does not contain the origin. We refine

the estimates in part (i). Let $\theta, \theta' \in (0, 1)$ with $\theta + \theta' = 1$. Elementary manipulations starting with (3.15) and (3.17) give

$$\begin{aligned} J_1 + J_3 &= -\frac{1}{|w|^2} \left[(\theta + \theta') |w|_1^2 w_1^2 z_1^2 + (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2 + (\theta + \theta') |w|_2^2 w_2^2 z_2^2 \right] + |w|_1^2 z_1^2 + |w|_2^2 z_2^2 \\ &= \frac{1}{|w|^2} \left[-\theta (|w|_1^2 w_1^2 z_1^2 + |w|_2^2 w_2^2 z_2^2) - (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2 \right] + J_*, \end{aligned}$$

where

$$J_* = \left(1 - \theta' \frac{w_1^2}{|w|^2} \right) |w|_1^2 z_1^2 + \left(1 - \theta' \frac{w_2^2}{|w|^2} \right) |w|_2^2 z_2^2. \quad (3.21)$$

Adding this to the formula (3.16) of J_2 and combining the like-terms, we obtain

$$|w|h'(t) = J_1 + J_2 + J_3 = J_* + 2J_4, \quad (3.22)$$

where

$$J_4 = \left(a - \theta \frac{|w|_1^2}{2|w|^2} \right) w_1^2 z_1^2 + \left[(b + c) - \frac{|w|_1^2 + |w|_2^2}{2|w|^2} \right] w_1 w_2 z_1 z_2 + \left(d - \theta \frac{|w|_2^2}{2|w|^2} \right) w_2^2 z_2^2.$$

Letting $\bar{\alpha} = w_1 z_1$ and $\bar{\beta} = w_2 z_2$, we rewrite J_4 as

$$J_4 = \left(a - \theta \frac{|w|_1^2}{2|w|^2} \right) \bar{\alpha}^2 + \left[(b + c) - \frac{|w|_1^2 + |w|_2^2}{2|w|^2} \right] \bar{\alpha} \bar{\beta} + \left(d - \theta \frac{|w|_2^2}{2|w|^2} \right) \bar{\beta}^2. \quad (3.23)$$

The right-hand side of (3.23) is a quadratic function of $\bar{\alpha}$ and $\bar{\beta}$. Note that

$$C_1 \leq \frac{|w|_1^2}{|w|^2}, \frac{|w|_2^2}{|w|^2} \leq C_2 \text{ for all } w \in \mathbb{R}^2 \setminus \{0\}. \quad (3.24)$$

Thanks to the strict inequalities in (3.12), we can choose $\theta \in (0, 1)$ sufficiently small so that

$$a - C_2 \theta > 0 \text{ and } \max_{|w|=1} \left[b + c - \frac{|w|_1^2 + |w|_2^2}{2} \right]^2 \leq 4 \left(a - \frac{C_2 \theta}{2} \right) \left(d - \frac{C_2 \theta}{2} \right).$$

Therefore, for all $w \in \mathbb{R}^2 \setminus \{0\}$,

$$a - \theta \frac{|w|_1^2}{|w|^2} > 0 \quad (3.25)$$

and

$$\left[b + c - \frac{|w|_1^2 + |w|_2^2}{2|w|^2} \right]^2 \leq 4 \left(a - \theta \frac{|w|_1^2}{2|w|^2} \right) \left(d - \theta \frac{|w|_2^2}{2|w|^2} \right). \quad (3.26)$$

Then (3.25) and (3.26) imply that $J_4 \geq 0$ for all $t \in [0, 1]$. Therefore, $|w|h'(t) \geq J_*$.

Using the fact $w_1^2, w_2^2 \leq |w|^2$ in the formula (3.21) of J_* and then property (3.24), we have

$$J_* \geq (1 - \theta') (|w|_1^2 z_1^2 + |w|_2^2 z_2^2) \geq C_1 (1 - \theta') |w|^2 |z|^2 = c_* |w(t)|^2 |z|^2, \text{ where } c_* = C_1 \theta. \quad (3.27)$$

It follows that $h'(t) \geq c_* |w(t)| |z|^2$. Combining this estimate of $h'(t)$ with (3.14) and applying inequality (2.14) with $\alpha = 1$, we have

$$I \geq c_* |z|^2 \int_0^1 |w(t)| dt \geq \frac{c_*}{8} |z| \cdot |z|^2 = \frac{c_* |z|^3}{8}.$$

Thus, we obtain

$$\left(\frac{1}{|u|} B(u, u, u) - \frac{1}{|v|} B(v, v, v) \right) \cdot (u - v) \geq C |u - v|^3 \quad (3.28)$$

with $C = c_*/8$. Therefore, G_B is 3-monotone. \square

We find a consequence of Theorem 3.4 with simpler conditions which can be checked easily.

Corollary 3.5. *If*

$$\max\{a + c, b + d\} \leq 2(b + c + 2\sqrt{ad}) \quad (3.29)$$

and

$$2(b + c - 2\sqrt{ad}) \leq \min\{a + c, b + d\}, \quad (3.30)$$

then G_B is monotone.

If both inequalities (3.29) and (3.30) are strict, then G_B is 3-monotone.

Proof. Condition (3.12) is equivalent to

$$2(b + c - 2\sqrt{ad}) \leq |w|_1^2 + |w|_2^2 \leq 2(b + c + 2\sqrt{ad}) \quad \forall |w| = 1. \quad (3.31)$$

Note that

$$\min\{a + c, b + d\}|w|^2 \leq |w|_1^2 + |w|_2^2 \leq \max\{a + c, b + d\}|w|^2.$$

Then (3.29) and (3.30) are sufficient conditions for (3.31), and hence G_B is monotone thanks to Theorem 3.4.

The case of strict inequalities in (3.29) and (3.30) is similar. \square

Example 3.6. Define, for $a, b, c, d > 0$, the functions

$$\begin{aligned} g_1(a, b, c, d) &= 2(b + c + 2\sqrt{ad}) - \max\{a + c, b + d\}, \\ g_2(a, b, c, d) &= \min\{a + c, b + d\} - 2(b + c - 2\sqrt{ad}). \end{aligned}$$

(a) For any $a, b > 0$, taking $a = d$ and $b = c$, we have

$$g_1(a, b, b, a) = 3(a + b) > 0, \quad g_2(a, b, b, a) = a + b - 4(b - a) = 5a - 3b.$$

If $a = 3b/5$, then G_B is monotone. If $a > 3b/5$, then G_B is 3-monotone.

(b) In particular, when $a = b = c = d$, the function G_B is 3-monotone, which is well-known for the generalized Forchheimer equation (1.2).

(c) Suppose $a_* > 3b_*/5 > 0$. If (a, b, c, d) is sufficiently close to (a_*, b_*, b_*, a_*) , then, thanks to the continuity of g_1 and g_2 , one has $g_1(a, b, c, d) > 0$ and $g_2(a, b, c, d) > 0$ which imply G_B is 3-monotone.

(d) If the matrices A_1 and A_2 are close to a_*I_2 , for some $a_* > 0$, then, thanks to (c) with $a_* = b_*$, the function G_B is 3-monotone.

Example 3.7. With the values a, b, c, d in (1.5), we rewrite conditions (3.29) and (3.30) as

$$b + d \leq 2(b + c + 2\sqrt{ad}) \text{ and } 2(b + c - 2\sqrt{ad}) \leq a + c. \quad (3.32)$$

The values in (1.5) satisfy the first inequality in (3.32), but fail the second one. Thus, the monotonicity of G_B is not yet determined. Therefore, there is a need to derive different sufficient conditions for the monotonicities of G_B .

Below are other monotonicity conditions which resemble but neither contain nor are contained in those of Theorem 3.4.

Theorem 3.8. *The following statements hold true.*

(i) *If either*

$$\left[b + c - \frac{(|w|_1 + |w|_2)^2}{2} \right]^2 \leq 4ad \text{ for all } w \in \mathbb{R}^2 \text{ with } |w| = 1, \quad (3.33)$$

or

$$\left[b + c - \frac{(|w|_1 - |w|_2)^2}{2} \right]^2 \leq 4ad \text{ for all } w \in \mathbb{R}^2 \text{ with } |w| = 1, \quad (3.34)$$

then G_B is monotone.

(ii) *If either the inequality in (3.33) is strict, or the inequality in (3.34) is strict, then G_B is 3-monotone.*

Proof. We follow the proof of Theorem 3.4 up to (3.18).

Part (i). Applying Cauchy's inequality to the sum $|w|_1^2 z_1^2 w_2^2 + |w|_2^2 z_2^2 w_1^2$ in (3.18) gives

$$J_1 + J_3 \geq \frac{1}{|w|^2} \left[\pm 2|w|_1 z_1 w_2 |w|_2 z_2 w_1 - (|w|_1^2 + |w|_2^2) w_1 w_2 z_1 z_2 \right] = -\frac{1}{|w|^2} (|w|_1 \mp |w|_2)^2 w_1 w_2 z_1 z_2.$$

Combining with the formula (3.16) of J_2 , we obtain

$$h'(t) \geq \frac{2}{|w|} \left\{ a w_1^2 z_1^2 + \left[(b+c) - \frac{(|w|_1 \mp |w|_2)^2}{2|w|^2} \right] w_1 w_2 z_1 z_2 + d w_2^2 z_2^2 \right\}. \quad (3.35)$$

Fix $t \in [0, 1]$. Let $\alpha(t) = w_1(t)z_1$ and $\beta(t) = w_2(t)z_2$. The expression between the parentheses on the right-hand side of (3.35) is $Q_{\mp}(\alpha(t), \beta(t))$, where Q_{\mp} is the following t -dependent quadratic function

$$Q_{\mp}(\alpha, \beta) = a\alpha^2 + \left[(b+c) - \frac{(|w(t)|_1 \mp |w(t)|_2)^2}{2|w(t)|^2} \right] \alpha\beta + d\beta^2 \text{ for } \alpha, \beta \in \mathbb{R}.$$

Suppose (3.33) is true. Then

$$\left[b+c - \frac{(|w|_1 + |w|_2)^2}{2|w|^2} \right]^2 \leq 4ad \quad \forall w \in \mathbb{R}^2 \setminus \{0\},$$

which yields $Q_+(\alpha, \beta) \geq 0$ for all $\alpha, \beta \in \mathbb{R}$. Consequently, $h'(t) \geq 0$ for all $t \in [0, 1]$, which gives $I \geq 0$ thanks to (3.14) and proves (3.20).

In the case (3.34), the proof is similar by using Q_- in place of Q_+ .

The rest of the proof of part (i) is the same as that of Theorem 3.4(i).

Part (ii). We follow the proof of part (ii) of Theorem 3.4 up to (3.22). In the remainder of the proof, we use the plus sign for \pm in the case of the strict inequality in (3.33), and use the minus sign for \pm in the case of the strict inequality in (3.34)

Adding and subtracting $\pm 2 \frac{|w|_1 |w|_2}{|w|^2} w_1 w_2 z_1 z_2$ to the last sum of (3.22), we obtain

$$|w|h'(t) = J_* \pm 2 \frac{|w|_1 |w|_2}{|w|^2} w_1 w_2 z_1 z_2 + 2\tilde{J}_4, \quad (3.36)$$

where

$$\begin{aligned} \tilde{J}_4 &= J_4 - \left(\pm \frac{|w|_1 |w|_2}{|w|^2} w_1 w_2 z_1 z_2 \right) \\ &= \left(a - \theta \frac{|w|_1^2}{2|w|^2} \right) w_1^2 z_1^2 + \left[(b+c) - \frac{(|w|_1^2 + |w|_2^2 \pm 2|w|_1 |w|_2)}{2|w|^2} \right] w_1 w_2 z_1 z_2 + \left(d - \theta \frac{|w|_2^2}{2|w|^2} \right) w_2^2 z_2^2. \end{aligned}$$

Letting $\bar{\alpha} = w_1 z_1$ and $\bar{\beta} = w_2 z_2$, we rewrite \tilde{J}_4 as

$$\tilde{J}_4 = \left(a - \theta \frac{|w|_1^2}{2|w|^2} \right) \bar{\alpha}^2 + \left[(b+c) - \frac{(|w|_1 \pm |w|_2)^2}{2|w|^2} \right] \bar{\alpha}\bar{\beta} + \left(d - \theta \frac{|w|_2^2}{2|w|^2} \right) \bar{\beta}^2. \quad (3.37)$$

The right-hand side of (3.37) is a quadratic function of $\bar{\alpha}$ and $\bar{\beta}$. Then there is $\theta \in (0, 1)$ sufficiently small so that one has, for all $w \in \mathbb{R}^2 \setminus \{0\}$, (3.25) holds true and, similar to (3.26),

$$\left[b+c - \frac{(|w|_1 \pm |w|_2)^2}{2|w|^2} \right]^2 \leq 4 \left(a - \theta \frac{|w|_1^2}{2|w|^2} \right) \left(d - \theta \frac{|w|_2^2}{2|w|^2} \right). \quad (3.38)$$

Note that $\theta' = 1 - \theta$ is a fixed number in $(0, 1)$ now. Then (3.25) and (3.38) imply that $\tilde{J}_4 \geq 0$ for all $t \in [0, 1]$. Combining this fact with formulas (3.36) and (3.21) gives

$$\begin{aligned} |w|h'(t) &\geq J_* \pm 2 \frac{w_1 w_2}{|w|^2} (|w|_1 z_1)(|w|_2 z_2) \\ &= \left(1 - \theta' \frac{w_1^2}{|w|^2}\right) |w|_1^2 z_1^2 + \left(1 - \theta' \frac{w_2^2}{|w|^2}\right) |w|_2^2 z_2^2 \pm 2 \frac{w_1 w_2}{|w|^2} (|w|_1 z_1)(|w|_2 z_2). \end{aligned}$$

This implies

$$h'(t) \geq \frac{1}{|w|} \tilde{Q}(|w|_1 z_1, |w|_2 z_2), \quad (3.39)$$

where \tilde{Q} is now a new quadratic function defined for each t by

$$\tilde{Q}(\alpha, \beta) = \left(1 - \theta' \frac{w_1^2(t)}{|w(t)|^2}\right) \alpha^2 + \left(1 - \theta' \frac{w_2^2(t)}{|w(t)|^2}\right) \beta^2 \pm 2 \frac{w_1(t)w_2(t)}{|w(t)|^2} \alpha\beta \text{ for } \alpha, \beta \in \mathbb{R}.$$

We claim that

$$w_1^2 w_2^2 < (1 - \theta' w_1^2)(1 - \theta' w_2^2) \text{ for all } w \in \mathbb{R}^2 \text{ with } |w| = 1. \quad (3.40)$$

Indeed, (3.40) is equivalent to

$$0 < -w_1^2 w_2^2 + 1 - \theta'(w_1^2 + w_2^2) + \theta'^2 w_1^2 w_2^2 = 1 - \theta' - (1 - \theta'^2)w_1^2 w_2^2,$$

which, in turn, is equivalent to

$$w_1^2 w_2^2 < \frac{1}{1 + \theta'}. \quad (3.41)$$

Because

$$w_1^2 w_2^2 \leq \left(\frac{w_1^2 + w_2^2}{2}\right)^2 = \frac{1}{4}, \text{ while } \frac{1}{1 + \theta'} \geq \frac{1}{2},$$

one asserts that (3.41) is true, and, hence, so is the claim (3.40).

By (3.40) and the compactness of the unit sphere $\{w \in \mathbb{R}^2 : |w| = 1\}$, we have

$$1 > \max_{|w|=1} \frac{w_1^2 w_2^2}{(1 - \theta' w_1^2)(1 - \theta' w_2^2)} = (1 - \varepsilon)^2, \quad (3.42)$$

for some number $\varepsilon \in (0, 1)$. We split \tilde{Q} as

$$\tilde{Q}(\alpha, \beta) = \varepsilon \left(1 - \theta' \frac{w_1^2(t)}{|w(t)|^2}\right) \alpha^2 + \varepsilon \left(1 - \theta' \frac{w_2^2(t)}{|w(t)|^2}\right) \beta^2 + \hat{Q}(\alpha, \beta), \quad (3.43)$$

where, for each t ,

$$\hat{Q}(\alpha, \beta) = (1 - \varepsilon) \left(1 - \theta' \frac{w_1^2(t)}{|w(t)|^2}\right) \alpha^2 + (1 - \varepsilon) \left(1 - \theta' \frac{w_2^2(t)}{|w(t)|^2}\right) \beta^2 \pm 2 \frac{w_1(t)w_2(t)}{|w(t)|^2} \alpha\beta.$$

Because

$$(1 - \varepsilon) \left(1 - \theta' \frac{w_1^2}{|w|^2}\right) \geq (1 - \varepsilon)(1 - \theta') > 0,$$

and, thanks to (3.42),

$$\left(\frac{w_1}{|w|} \frac{w_2}{|w|}\right)^2 \leq (1 - \varepsilon)^2 \left(1 - \theta' \frac{w_1^2}{|w|^2}\right) \left(1 - \theta' \frac{w_2^2}{|w|^2}\right) \quad \forall w \in \mathbb{R}^2, w \neq 0,$$

we have $\hat{Q} \geq 0$. Together with (3.39) and (3.43), it implies $h'(t) \geq \varepsilon J_* / |w|$, where J_* is defined by (3.21). Thanks to the lower bound of J_* in (3.27), we derive $h'(t) \geq \varepsilon c_* |w(t)| |z|^2$. By using (3.14) and applying inequality (2.14) with $\alpha = 1$, we obtain (3.28) for $C = \varepsilon c_* / 8$ and complete proof of part (ii). \square

Note that condition (3.33) is equivalent to

$$2(b + c - 2\sqrt{ad}) \leq (|w|_1 + |w|_2)^2 \leq 2(b + c + 2\sqrt{ad}) \text{ for all } w \in \mathbb{R}^2 \text{ with } |w| = 1, \quad (3.44)$$

while condition (3.34) is equivalent to

$$2(b + c - 2\sqrt{ad}) \leq (|w|_1 - |w|_2)^2 \leq 2(b + c + 2\sqrt{ad}) \text{ for all } w \in \mathbb{R}^2 \text{ with } |w| = 1, \quad (3.45)$$

Based on Theorem 3.8, more specific conditions for the monotonicities can then be derived in the following.

Corollary 3.9. *If*

$$\max\{a + c, b + d\} + 2\sqrt{\max\{a, b\}}\sqrt{\max\{c, d\}} \leq 2(b + c + 2\sqrt{ad}) \quad (3.46)$$

and

$$2(b + c - 2\sqrt{ad}) \leq \min\{a + c, b + d\} + 2\sqrt{\min\{a, b\}}\sqrt{\min\{c, d\}}, \quad (3.47)$$

then G_B is monotone.

If both inequalities (3.46) and (3.47) are strict, then G_B is 3-monotone.

Proof. We use condition (3.44) for Theorem 3.8. Note, for any $w \in \mathbb{R}^2$, that

$$(|w|_1 + |w|_2)^2 = |w|_1^2 + |w|_2^2 + 2|w|_1|w|_2 \leq \max\{a + c, b + d\}|w|^2 + 2\sqrt{\max\{a, b\}}\sqrt{\max\{c, d\}}|w|^2,$$

and

$$(|w|_1 - |w|_2)^2 \geq \min\{a + c, b + d\}|w|^2 + 2\sqrt{\min\{a, b\}}\sqrt{\min\{c, d\}}|w|^2.$$

Hence, (3.46) is a sufficient condition for the second inequality in (3.44), and (3.47) is a sufficient condition for the first inequality in (3.44). Thus, the first statement follows part (i) of Proposition 3.8.

The arguments for the case (3.46) and (3.47) being strict inequalities are similar by using part (ii) of Theorem 3.8. \square

We are ready to check our mathematical criteria with the real data in [1].

Example 3.10. With the experimental values in (1.5) we have $a < c < b < d$. Then the strict inequalities in (3.46) and (3.47) become

$$b + d + 2\sqrt{bd} < 2(b + c + 2\sqrt{ad}) \quad (3.48)$$

and

$$2(b + c - 2\sqrt{ad}) < (a + c + 2\sqrt{ac}). \quad (3.49)$$

Simple calculations show that the values in (1.5) satisfy both (3.48) and (3.49). Therefore, G_B is 3-monotone. Moreover, thanks to the same continuity argument in part (b) of Example 3.6, G_B is still 3-monotone even when the measurements have small errors. This shows that our mathematical criteria work for real data.

Using condition (3.45) instead of (3.44), we obtain a counterpart of Corollary 3.9.

Corollary 3.11. *The following statements hold true.*

(i) *If*

$$\frac{\max\{a, d\}}{2} - 2\sqrt{ad} \leq b + c \leq 2\sqrt{ad}, \quad (3.50)$$

then G_B is monotone.

If

$$\frac{\max\{a, d\}}{2} - 2\sqrt{ad} < b + c < 2\sqrt{ad}, \quad (3.51)$$

then G_B is 3-monotone.

(ii) *Let $M = \max\{a, d\}$ and $m = \min\{a, d\}$.*

(a) If $M/m \leq 16$, then (3.50), respectively, (3.51), is equivalent to

$$b + c \leq 2\sqrt{ad}, \text{ respectively, } b + c < 2\sqrt{ad}.$$

(b) If $16 < M/m < 64$, then (3.50) and (3.51) stay the same.

(c) If $M/m = 64$, then (3.50) is equivalent to

$$b + c = 2\sqrt{ad},$$

while (3.51) is impossible.

(d) If $M/m > 64$, then (3.50) is impossible.

Proof. Observe, for all $w \in \mathbb{R}^2$, that

$$(|w|_1 - |w|_2)^2 \leq \max\{|w|_1^2, |w|_2^2\} = \max\{a, b, c, d\}|w|^2. \quad (3.52)$$

Part (i). Assume (3.50). By the virtue of part (ii) of Theorem 3.8, we can use (3.45) as a sufficient condition for G_B to be monotone. Thanks to (3.52), one sufficient condition for (3.45) is

$$b + c \leq 2\sqrt{ad}, \quad \max\{a, b, c, d\} \leq 2(b + c + 2\sqrt{ad}). \quad (3.53)$$

By dropping b and c on the left-hand side of the last inequality, we obtain the sufficient condition (3.50).

Now, assume (3.51). Then the second inequality in (3.50) implies the first inequality in (3.45) is strict. Also, the first inequality in (3.50) implies the second inequality in (3.53) is strict. This and (3.52) imply the second inequality in (3.45) is also strict. Thus, the inequality in (3.34) is strict. Therefore, according to part (ii) of Theorem 3.8, we have G_B is 3-monotone.

Part (ii). We rewrite the first group in (3.50) and (3.51) as

$$S_1 \stackrel{\text{def}}{=} \frac{\max\{a, d\}}{2} - 2\sqrt{ad} = \frac{M}{2} - 2\sqrt{Mm} = \frac{1}{2}\sqrt{Mm}(\sqrt{M/m} - 4).$$

If $M/m \leq 16$, then $S_1 \leq 0$. Therefore, the statements in part (a) are correct.

Comparing the last group with the first group in (3.50) and (3.51), we consider

$$S_2 \stackrel{\text{def}}{=} 2\sqrt{ad} - \left(\frac{\max\{a, d\}}{2} - 2\sqrt{ad} \right) = 4\sqrt{Mm} - \frac{M}{2} = \frac{1}{2}\sqrt{Mm}(8 - \sqrt{M/m}).$$

If $16 < M/m < 64$, then $S_1 > 0$ and $S_2 > 0$, which yield no changes to (3.50) and (3.51).

When $M/m = 64$, one has $S_1 > 0$ and $S_2 = 0$. Therefore, the statements in part (c) are correct.

If $M/m > 64$, then $S_2 < 0$, and, hence, inequality (3.50) is impossible. \square

Example 3.12. If $d = 36a$ and $6a < b + c < 12a$, then, by using (3.51), we have G_B is 3-monotone.

3.2. The case of general dimensions. We return to the general spatial dimension $n \geq 2$ with the trilinear mapping B defined by (3.1).

Theorem 3.13. *The following statements hold true.*

(i) *If there exists a number $\lambda > 0$ such that*

$$\|A_i - \lambda I_n\|_{\text{op}} \leq \frac{\lambda}{4} \quad \text{for all } i = 1, 2, \dots, n, \quad (3.54)$$

then G_B is monotone.

(ii) *If the inequality in (3.54) is strict for all $i = 1, \dots, n$, then G_B is 3-monotone.*

Proof. Part (i). Assume (3.54). Denote $A_* = \lambda I_n$ and, for $i = 1, 2, \dots, n$, $M_i = A_i - \lambda I_n$.

Let $u, v \in \mathbb{R}^n$. We can assume, without loss of generality, that the line segment connecting u and v does not contain the origin. Let $z, w(t)$ and $h(t)$ be as in (3.13), and I be as in (3.14). We calculate

$$\begin{aligned} h'(t) &= -\frac{w \cdot z}{|w|^3} B(w, w, w) \cdot z + \frac{1}{|w|} (B(z, w, w) + B(w, z, w) + B(w, w, z)) \cdot z \\ &= -\frac{w \cdot z}{|w|^3} \sum_{i=1}^n (w^T A_i w) w_i z_i + \frac{1}{|w|} \sum_{i=1}^n (z^T A_i w + w^T A_i z) w_i z_i + \frac{1}{|w|} \sum_{i=1}^n (w^T A_i w) z_i^2. \end{aligned}$$

For each i , we write $A_i = A_* + M_i$ and split the sums in $h'(t)$ accordingly to yield $h'(t) = H_1 + H_2$, where

$$\begin{aligned} H_1 &= -\frac{w \cdot z}{|w|^3} \sum_{i=1}^n (w^T A_* w) w_i z_i + \frac{1}{|w|} \sum_{i=1}^n (z^T A_* w + w^T A_* z) w_i z_i + \frac{1}{|w|} \sum_{i=1}^n (w^T A_* w) z_i^2 \\ H_2 &= -\frac{w \cdot z}{|w|^3} \sum_{i=1}^n (w^T M_i w) w_i z_i + \frac{1}{|w|} \sum_{i=1}^n (z^T M_i w + w^T M_i z) w_i z_i + \frac{1}{|w|} \sum_{i=1}^n (w^T M_i w) z_i^2. \end{aligned}$$

Regarding H_1 , one has

$$H_1 = -\frac{\lambda |w|^2 (w \cdot z)^2}{|w|^3} + \frac{2\lambda (w \cdot z)^2}{|w|} + \frac{1}{|w|} \lambda |w|^2 |z|^2 = \frac{\lambda (w \cdot z)^2}{|w|} + \lambda |w| |z|^2 \geq \lambda |w| |z|^2.$$

Set $\varepsilon_0 = \max\{\|M_i\|_{\text{op}} : 1 \leq i \leq n\}$. Applying the Cauchy–Schwarz inequality and the first inequality in (2.2) multiple times, we estimate H_2 by

$$\begin{aligned} |H_2| &\leq \frac{|w| \cdot |z|}{|w|^3} \sum_{i=1}^n \varepsilon_0 |w|^2 |w_i| |z_i| + \frac{1}{|w|} \sum_{i=1}^n 2\varepsilon_0 |w| |z| |w_i| |z_i| + \frac{1}{|w|} \sum_{i=1}^n \varepsilon_0 |w|^2 z_i^2 \\ &= \varepsilon_0 \left(3|z| \sum_{i=1}^n |w_i| |z_i| + |w| |z|^2 \right) \leq \varepsilon_0 (3|z| \cdot |w| |z| + |w| |z|^2) = 4\varepsilon_0 |w| |z|^2. \end{aligned}$$

Combining the above estimates of H_1 and H_2 gives

$$h'(t) \geq (\lambda - 4\varepsilon_0) |w| |z|^2.$$

Thanks to condition (3.54), we have $\varepsilon_0 \leq \lambda/4$, then $h'(t) \geq 0$ for all $t \in [0, 1]$. By this fact and (3.14), we obtain (3.20). Thus, G_B is monotone.

Part (ii). With strict inequalities in (3.54), we have $\lambda - 4\varepsilon_0 = c > 0$. Thus, $h'(t) \geq c|w| |z|^2$ for all $t \in [0, 1]$. Integrating this inequality from 0 to 1 and then applying inequality (2.14), we obtain (3.28) with $C = c/8$. Therefore, G_B is 3-monotone. \square

Remark 3.14. Compared with Theorem 3.8 in the two-dimension case, Theorem 3.13 does not require each matrix A_i to be diagonal. However, it must be a perturbation of the same multiple of the identity matrix for all i .

The next theorem contains very specific sufficient conditions for G_B 's monotonicities via justifying (3.54). Below, $\text{Tr}(\cdot)$ denotes the trace of square matrices.

Theorem 3.15. *If*

$$\frac{\sum_{i=1}^n \text{Tr}(A_i)}{\sqrt{\sum_{i=1}^n |A_i|^2}} \geq \sqrt{n^2 - \frac{1}{16}}, \quad (3.55)$$

then G_B is monotone.

If inequality (3.55) is strict, then G_B is 3-monotone.

Proof. Assume (3.55). Thanks to Theorem 3.13(i) and the relation (2.1), a sufficient condition for G_B being monotone is that there is $\lambda > 0$ such that

$$\sum_{i=1}^n |A_i - \lambda I_n|^2 \leq \lambda^2/16.$$

This can be rewritten as a quadratic inequality in λ as

$$\left(n^2 - \frac{1}{16}\right)\lambda^2 - 2\lambda \sum_{i=1}^n \text{Tr}(A_i) + \sum_{i=1}^n |A_i|^2 \leq 0. \quad (3.56)$$

By (3.55), we have

$$\sum_{i=1}^n \text{Tr}(A_i) > 0 \text{ and } \left(\sum_{i=1}^n \text{Tr}(A_i)\right)^2 - (n^2 - 1/16) \sum_{i=1}^n |A_i|^2 \geq 0.$$

Then the quadratic inequality (3.56) has a positive solution λ , which implies G_B is monotone.

If inequality (3.55) is strict, then there is a positive solution λ of the corresponding strict inequality in (3.56). By (2.1) and Theorem 3.13(ii), G_B is 3-monotone. \square

4. FUNDAMENTAL PROPERTIES (II)

Let A and K be $n \times n$ matrices, and α be a positive number. The functions of our interest are of the general form

$$F_{A,K,\alpha}(u) = |Ku|^\alpha Au \text{ for } u \in \mathbb{R}^n. \quad (4.1)$$

In particular, when $K = I_n$ in (4.1), we have the function

$$F_{A,\alpha}(u) = |u|^\alpha Au \text{ for } u \in \mathbb{R}^n. \quad (4.2)$$

We note that when $A = \text{diag}[a_1, a_2, \dots, a_n]$ and $\alpha = 1$,

$$F_{A,1}(u) = |u|^{-1} B(u, u, u) = G_B(u) \text{ for } u \neq 0,$$

with B defined as in (3.1) for $A_i = a_i I_n$. Thus, properties of $F_{A,1}$ in this case can be derived from those in Section 3. Nonetheless, we will focus on the general matrix A and power α in this section.

When A is symmetric, positive definite, and $K = A^{1/2}$ in (4.1), we have the functions

$$\tilde{F}_{A,\alpha}(u) = |A^{1/2}u|^\alpha Au \text{ for } u \in \mathbb{R}^n. \quad (4.3)$$

For our convenience, we also define

$$F_{A,0}(u) = \tilde{F}_{A,0}(u) = Au \text{ for } u \in \mathbb{R}^n.$$

The following are natural necessary conditions for the monotonicities for $F_{A,K,\alpha}$.

Proposition 4.1. *Let A, K be $n \times n$ matrices and $\alpha > 0$.*

(i) *If K is invertible and $F_{A,K,\alpha}$ is monotone, then*

$$u^T Au \geq 0 \text{ for all } u \in \mathbb{R}^n. \quad (4.4)$$

(ii) *If $F_{A,K,\alpha}$ is β -monotone for some $\beta > 0$, then A and K are invertible, $\beta = \alpha + 2$ and there is $C > 0$ such that*

$$u^T Au \geq C|u|^2 \text{ for all } u \in \mathbb{R}^n. \quad (4.5)$$

Proof. Part (i). Assume K is invertible and $F_{A,K,\alpha}$ is monotone. Then

$$(|Ku|^\alpha Au - |Kv|^\alpha Av) \cdot (u - v) \geq 0 \text{ for any } u, v \in \mathbb{R}^n. \quad (4.6)$$

By taking $v = 0$ in (4.6), we obtain (4.4).

Part (ii). Assume $F_{A,K,\alpha}$ is β -monotone. Then there is $C_0 > 0$ such that

$$(|Ku|^\alpha Au - |Kv|^\alpha Av) \cdot (u - v) \geq C_0|u - v|^\beta \text{ for any } u, v \in \mathbb{R}^n.$$

Taking $v = 0$ again yields

$$|Ku|^\alpha Au \cdot u \geq C_0|u|^\beta \text{ for any } u \in \mathbb{R}^n. \quad (4.7)$$

This implies A and K are invertible. It also follows (4.7) that

$$|K|^\alpha |A| |u|^{\alpha+2} \geq C_0 |u|^\beta \text{ for any } u \in \mathbb{R}^n. \quad (4.8)$$

Taking $u = tw$ for a fixed $w \neq 0$ in (4.8), and letting $t \rightarrow \infty$ and then $t \rightarrow 0^+$ yield $\beta = \alpha + 2$. As a consequence of this fact and (4.7),

$$|K|^\alpha |u|^\alpha Au \cdot u \geq C_0 |u|^{\alpha+2} \text{ for any } u \in \mathbb{R}^n,$$

which implies (4.5) with $C = C_0/|K|^\alpha$. \square

Example 4.2. In \mathbb{R}^2 , let

$$A = \begin{pmatrix} 5 & 1/5 \\ 1/5 & 1/100 \end{pmatrix}, \quad u = (3, 5), \text{ and } v = (4, 1).$$

Then A is symmetric and positive definite. However, $(|u|Au - |v|Av) \cdot (u - v) < 0$, which implies $F_{A,1}$ is not monotone.

Because of Example 4.2, we need to investigate the monotonicities in different situations.

Theorem 4.3. *Suppose there is a number $\kappa_0 > 0$ such that*

$$u^T Au \geq \kappa_0^2 |u|^2 \text{ for all } u \in \mathbb{R}^n. \quad (4.9)$$

- (i) *Then the function $F_{A,0} = \tilde{F}_{A,0}$ is 2-monotone.*
- (ii) *If A is symmetric and $\alpha > 0$, then the function $\tilde{F}_{A,\alpha}$ is $(\alpha + 2)$ -monotone.*
- (iii) *If $\alpha = \kappa_0^2/\|A\|_{\text{op}}$, then the function $F_{A,\alpha}$ is monotone.
If $\alpha < \kappa_0^2/\|A\|_{\text{op}}$, then the function $F_{A,\alpha}$ is $(\alpha + 2)$ -monotone.*

Proof. Part (i). Because $F_{A,0} = \tilde{F}_{A,0}$ is linear, the assumption (4.9) implies that it is 2-monotone.

Part (ii). Denote $F = \tilde{F}_{A,\alpha}$. In this case, we note, for any $u \in \mathbb{R}^n$, that

$$|A^{1/2}u| = (u \cdot Au)^{1/2} \geq \kappa_0 |u|,$$

and, hence,

$$F(u) \cdot u \geq |A^{1/2}u|^\alpha \kappa_0^2 |u|^2 \geq \kappa_0^{\alpha+2} |u|^{\alpha+2}. \quad (4.10)$$

We will prove that, for any $u, v \in \mathbb{R}^n$,

$$I \stackrel{\text{def}}{=} (F(u) - F(v)) \cdot (u - v) \geq \frac{\kappa_0^{\alpha+2}}{2^{\alpha+1}(\alpha+1)} |u - v|^{\alpha+2}. \quad (4.11)$$

Thanks to inequality (4.10), one obtains (4.11) when $u = 0$ or $v = 0$.

Consider $u, v \in \mathbb{R}^n \setminus \{0\}$ now. It suffices to establish (4.11) when the line segment connecting u and v does not contain the origin. Denote $z = u - v$, $w(t) = tu + (1-t)v$ for $t \in [0, 1]$, and

$$h(t) = F(w(t)) \cdot z = |A^{1/2}w(t)|^\alpha (A^{1/2}w(t) \cdot A^{1/2}z).$$

Then $w(t) \neq 0$ for all $t \in [0, 1]$, and

$$\begin{aligned} h'(t) &= |A^{1/2}w(t)|^\alpha |A^{1/2}z|^2 + \alpha |A^{1/2}w(t)|^{\alpha-2} (A^{1/2}w(t) \cdot A^{1/2}z)^2 \\ &\geq |A^{1/2}w(t)|^\alpha |A^{1/2}z|^2 \geq \kappa_0^{\alpha+2} |w(t)|^\alpha |z|^2. \end{aligned}$$

By The Fundamental Theorem of Calculus and inequality (2.14), we obtain

$$I = \int_0^1 h'(t) dt \geq \frac{\kappa_0^{\alpha+2}}{2^{\alpha+1}(\alpha+1)} |z|^{\alpha+2},$$

which proves (4.11). Therefore, F is $(\alpha+2)$ -monotone.

Part (iii). Denote $F = F_{A,\alpha}$. Note, for all $u \in \mathbb{R}^n$, that $F(u) \cdot u \geq \kappa_0^2 |u|^{\alpha+2}$.

Similar to part (ii), it suffices to consider $u, v \in \mathbb{R}^n \setminus \{0\}$ and the line segment connecting u and v does not contain the origin. With the same I, z and $w(t)$ as in part (ii), we define, for $t \in [0, 1]$, $h(t) = |w(t)|^\alpha Aw(t) \cdot z$. Then

$$\begin{aligned} h'(t) &= |w(t)|^\alpha (Az) \cdot z + \alpha |w(t)|^{\alpha-2} (w(t) \cdot z) Aw(t) \cdot z \\ &\geq \kappa_0^2 |w(t)|^\alpha |z|^2 - \alpha |w(t)|^{\alpha-2} \cdot |w(t)| |z| \cdot \|A\|_{\text{op}} |w(t)| |z| \\ &= (\kappa_0^2 - \alpha \|A\|_{\text{op}}) |w(t)|^\alpha |z|^2. \end{aligned}$$

If $\alpha = \kappa_0^2 / \|A\|_{\text{op}}$, then $h'(t) \geq 0$, which implies $I \geq 0$. Therefore, $F_{A,\alpha}$ is monotone.

If $\alpha < \kappa_0^2 / \|A\|_{\text{op}}$, Then $c \stackrel{\text{def}}{=} \kappa_0^2 - \alpha \|A\|_{\text{op}}$ is positive. By (2.14), we have

$$I = \int_0^1 h'(t) dt \geq \frac{c |z|^\alpha}{2^{\alpha+1}(\alpha+1)} |z|^2.$$

Thus, $F_{A,\alpha}$ is $(\alpha+2)$ -monotone. \square

Part (ii) of Theorem 4.3 means that the flow is close to (anisotropic) Darcy's flows ($\alpha = 0$). Also, if we assume additionally that A is symmetric, then we can take κ_0^2 to be the smallest eigenvalue λ_{\min} of A , while $\|A\|_{\text{op}}$ is then largest eigenvalue λ_{\max} of A . Thus, the quotient $\kappa_0^2 / \|A\|_{\text{op}}$ is simply $\lambda_{\min} / \lambda_{\max}$.

For large α , the following is a counterpart of Theorem 3.13.

Theorem 4.4. *Let $\alpha > 0$.*

(i) *If there exists $\lambda > 0$ such that*

$$\|A - \lambda I_n\|_{\text{op}} \leq \frac{\lambda}{1 + \alpha}, \quad (4.12)$$

then $F_{A,\alpha}$ is monotone.

(ii) *If inequality (4.12) is a strict inequality, then $F_{A,\alpha}$ is $(\alpha+2)$ -monotone.*

Proof. Part (i). Given $u, v \in \mathbb{R}^n$. Define $z = u - v$, and

$$w(t) = tu - (1-t)v, \quad h(t) = |w(t)|^\alpha Aw(t) \cdot z \text{ for } t \in [0, 1].$$

Again, without loss of generality, we assume $w(t) \neq 0$ for all $t \in [0, 1]$. Using the Fundamental Theorem of Calculus, we have

$$I \stackrel{\text{def}}{=} (|u|^\alpha Au - |v|^\alpha Av) \cdot (u - v) = h(1) - h(0) = \int_0^1 h'(t) dt.$$

We calculate

$$h'(t) = |w|^\alpha Az \cdot z + \alpha |w|^{\alpha-2} (w \cdot z) Aw \cdot z.$$

Denote $M = A - \lambda I_n$. Then, thanks to (4.12), $\|M\|_{\text{op}} \leq \lambda / (1 + \alpha)$.

With $A = \lambda I_n + M$, we have

$$\begin{aligned} h'(t) &= \lambda |w|^\alpha |z|^2 + |w|^\alpha Mz \cdot z + \lambda |w|^{\alpha-2} (w \cdot z)^2 + \alpha |w|^{\alpha-2} (w \cdot z) Mw \cdot z \\ &\geq (\lambda - (1 + \alpha) \|M\|_{\text{op}}) |w|^\alpha |z|^2. \end{aligned} \quad (4.13)$$

Thus, $h'(t) \geq 0$ which implies $I \geq 0$, that is, $F_{A,\alpha}$ is monotone.

Part (ii). In this case, we have $c \stackrel{\text{def}}{=} \lambda - (1 + \alpha) \|M\|_{\text{op}} > 0$. Combining this with (4.13) and then applying inequality (2.14) give

$$I \geq c |z|^2 \int_0^1 |w(t)| dt \geq \frac{c}{8} |z|^3.$$

Therefore, $F_{A,\alpha}$ is 3-monotone. \square

More specific criteria are next, which are counterparts of Theorem 3.15.

Theorem 4.5. *Let $\alpha > 0$. If*

$$\frac{\text{Tr}(A)}{|A|} \geq \sqrt{n - \frac{1}{(1 + \alpha)^2}}, \quad (4.14)$$

then $F_{A,\alpha}$ is monotone.

If inequality (4.14) is strict, then $F_{A,\alpha}$ is $(\alpha + 2)$ -monotone.

Proof. We solve for a solution $\lambda > 0$ of the inequality

$$|A - \lambda I_n|^2 \leq \lambda^2 / (1 + \alpha)^2$$

which is equivalent to

$$\left(n - \frac{1}{(1 + \alpha)^2} \right) \lambda^2 - 2\text{Tr}(A)\lambda + |A|^2 \leq 0. \quad (4.15)$$

Under condition (4.14), the quadratic inequality (4.15) has a solution $\lambda > 0$. Together with relation (2.1), we have (4.12), and, thanks to Theorem 4.4(i), $F_{A,\alpha}$ is monotone.

When the inequality (4.14) is strict, there is a solution $\lambda > 0$ of the strict inequality in (4.15). Thanks to (2.1) again, we have the strict inequality in (4.12), and, by Theorem 4.4(ii), $F_{A,\alpha}$ is $(\alpha + 2)$ -monotone. \square

Lemma 4.6. *Let $\alpha > 0$ and A, K be $n \times n$ matrices. If either $\alpha \geq 1$ or K is invertible, then there is a constant $C > 0$ such that, for any $u, v \in \mathbb{R}^n$,*

$$||Ku|^\alpha Au - |Kv|^\alpha Av| \leq C(|u|^\alpha + |v|^\alpha)|u - v|. \quad (4.16)$$

Proof. First, we observe that

$$\begin{aligned} ||Ku|^\alpha Au - |Kv|^\alpha Av| &\leq |Ku|^\alpha |A(u - v)| + ||Ku|^\alpha - |Kv|^\alpha| |Av| \\ &\leq \|K\|_{\text{op}}^\alpha \|A\|_{\text{op}} |u|^\alpha |u - v| + \|A\|_{\text{op}} ||Ku|^\alpha - |Kv|^\alpha| |v|. \end{aligned} \quad (4.17)$$

Case $\alpha = 1$. Estimating $||Ku| - |Kv|| \leq |Ku - Kv| \leq \|K\|_{\text{op}} |u - v|$ in (4.17), we obtain (4.16) with $C = \|K\|_{\text{op}} \|A\|_{\text{op}}$.

Case $\alpha > 1$. Using inequality (2.8), we estimate

$$||Ku|^\alpha - |Kv|^\alpha| \leq 2^{\alpha-1} (|Ku|^{\alpha-1} + |Kv|^{\alpha-1}) |Ku - Kv| \leq 2^{\alpha-1} \|K\|_{\text{op}}^\alpha (|u|^{\alpha-1} + |v|^{\alpha-1}) |u - v|.$$

Combining this with (4.17) yields

$$||Ku|^\alpha Au - |Kv|^\alpha Av| \leq 2^{\alpha-1} \|K\|_{\text{op}}^\alpha \|A\|_{\text{op}} (|u|^\alpha + |u|^{\alpha-1}|v| + |v|^\alpha) |u - v|.$$

By Young's inequality, we estimate $|u|^{\alpha-1}|v| \leq |u|^\alpha + |v|^\alpha$, and obtain (4.16) with constant $C = 2^\alpha \|K\|_{\text{op}} \|A\|_{\text{op}}$.

Case $\alpha \in (0, 1)$ and K is invertible. Without loss of generality, assume the line segment connecting u and v does not contain the origin. Let $z = u - v$ and define, for $t \in [0, 1]$,

$$w(t) = tu + (1 - t)v, \quad h(t) = |Kw(t)|^\alpha Aw(t) \cdot x.$$

By The Fundamental Theorem of Calculus,

$$I \stackrel{\text{def}}{=} |Ku|^\alpha Au - |Kv|^\alpha Av = h(1) - h(0) = \int_0^1 h'(t) dt.$$

With $w(t) \neq 0$ for all $t \in [0, 1]$, we have $Kw(t) \neq 0$, and

$$h'(t) = Kw(t)|^\alpha Az + \alpha |Kw(t)|^{\alpha-2} (Kw(t) \cdot Kz) Aw(t)$$

We estimate

$$|h'(t)| \leq \|K\|_{\text{op}}^\alpha \|A\|_{\text{op}} |w(t)|^\alpha |z| + \alpha \|K\|_{\text{op}} \|A\|_{\text{op}} |Kw(t)|^{\alpha-1} |w(t)| |z|.$$

With $\alpha - 1 < 0$, we use inequality (2.3) to have $|Kw(t)|^{\alpha-1} \leq \|K^{-1}\|_{\text{op}}^{1-\alpha} |w(t)|^{\alpha-1}$.

Thus, there is $C_0 > 0$ depending on A, K, α such that $|h'(t)| \leq C_0 |w(t)|^\alpha |z|$. Therefore,

$$\begin{aligned} |I| &\leq \int_0^1 |h'(t)| dt \leq C_0 |z| \int_0^1 |w(t)|^\alpha dt \\ &\leq C_0 |z| \int_0^1 t^\alpha |u|^\alpha + (1-t)^\alpha |v|^\alpha dt = \frac{C_0}{\alpha+1} (|u|^\alpha + |v|^\alpha) |z|, \end{aligned}$$

which proves (4.16). \square

Remark 4.7. In the case $\alpha \in (0, 1)$ and K is not invertible, we only have

$$||Ku|^\alpha Au - |Kv|^\alpha Av| \leq \|K\|_{\text{op}}^\alpha \|A\|_{\text{op}} (|u|^\alpha |u - v| + |v| |u - v|^\alpha). \quad (4.18)$$

This is weaker than (4.16) when $|u - v|$ is small. Inequality (4.18), in fact, is a consequence of (4.17) and the following estimate, thanks to inequality (2.7),

$$||Ku|^\alpha - |Kv|^\alpha| \leq |Ku - Kv|^\alpha \leq \|K\|_{\text{op}}^\alpha |u - v|^\alpha.$$

Consider an integer $N \geq 0$, real numbers $0 \leq \alpha_0 < \alpha_1 < \alpha_2 < \dots < \alpha_N$, and $n \times n$ matrices $\bar{A}_0, \bar{A}_1, \dots, \bar{A}_N$. For $0 \leq k \leq N$, let $F_k = F_{\bar{A}_k, \alpha_k}$ or $F_k = \tilde{F}_{\bar{A}_k, \alpha_k}$.

Let B be a trilinear mapping from $\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n$ to \mathbb{R}^n as in (3.1).

Define the mapping $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by

$$F(u) = G_B(u), \quad \text{or} \quad (4.19)$$

$$F(u) = \sum_{k=0}^N F_k(u), \quad \text{or} \quad (4.20)$$

$$F(u) = \sum_{k=0}^N F_k(u) + G_B(u). \quad (4.21)$$

Note that if $\alpha_0 = 0$ then we have the (anisotropic) Darcy term $\bar{A}_0 u$ in (4.20) and (4.21).

Proposition 4.8. *Let F be as in (4.19)–(4.21). Assume the followings.*

- (a) *In the case (4.19), the function G_B is 3-monotone.*
- (b) *In the case (4.20), each function F_k is monotone for $0 \leq k \leq N - 1$, and F_N is $(\alpha_N + 2)$ -monotone with $\alpha_N > 0$.*
- (c) *In the case (4.21), all functions F_k 's and G_B are monotone. In addition,*
 - (c.1) *if $\alpha_N < 1$, then G_B is 3-monotone,*
 - (c.2) *if $\alpha_N > 1$, then F_N is $(\alpha_N + 2)$ -monotone,*
 - (c.3) *if $\alpha_N = 1$, then F_N or G_B is 3-monotone.*

Let $\beta = 1$ in the case 4.19, $\beta = \alpha_N$ in the case (4.20), and $\beta = \max\{\alpha_N, 1\}$ in the case (4.21). Then there are positive constants c_1 and c_2 such that, for all $u, v \in \mathbb{R}^n$,

$$|F(u) - F(v)| \leq c_1(1 + |u|^\beta + |v|^\beta)|u - v|, \quad (4.22)$$

$$(F(u) - F(v)) \cdot (u - v) \geq c_2|u - v|^{\beta+2}. \quad (4.23)$$

Proof. In the case 4.21, inequalities (4.22) and (4.23) come from (3.8) and assumption (a).

Consider cases (4.20) and (4.21). In estimates below, C_1 and C_2 are positive constants. Let u, v be any vectors in \mathbb{R}^n . By Lemma 4.6 and Young's inequality, we have

$$\begin{aligned} \left| \sum_{k=0}^N (F_k(u) - F_k(v)) \right| &\leq C_1 \sum_{k=0}^N (|u|^{\alpha_k} + |v|^{\alpha_k})|u - v| \\ &\leq C_1(2N + 1)(1 + |u|^{\alpha_N} + |v|^{\alpha_N})|u - v|. \end{aligned} \quad (4.24)$$

If F_N is $(\alpha_N + 2)$ -monotone, then

$$\left(\sum_{k=0}^N F_k(u) - \sum_{k=0}^N F_k(v) \right) \cdot (u - v) \geq (F_N(u) - F_N(v)) \cdot (u - v) \geq C_2|u - v|^{\alpha_N+2}. \quad (4.25)$$

Therefore, (4.24) and (4.25) prove (4.22) and (4.23) in the case (4.20). In the case (4.20), combining (4.24) with (3.8) and the using Hölder's inequality, we obtain (4.22). Finally, (4.23) follows the fact G_B is 3-monotone for (c.1), or the fact (4.25) for (c.2), or one of these facts (or both) for (c.3). \square

5. STEADY STATE FLOWS

We study the steady states for system (1.10) with a nonhomogeneous Dirichlet boundary condition. Redenoting \mathbf{m} by \mathbf{u} , and \tilde{p} by p , we particularly consider

$$\begin{cases} F(\mathbf{u}) = -\nabla p & \text{in } \Omega, \\ \operatorname{div} \mathbf{u} = f & \text{in } \Omega, \\ p = \psi & \text{on } \partial\Omega, \end{cases} \quad (5.1)$$

where $f : \Omega \rightarrow \mathbb{R}$ and $\psi : \partial\Omega \rightarrow \mathbb{R}$ are given functions. The unknowns are $\mathbf{u} : \Omega \rightarrow \mathbb{R}^n$ and $p : \Omega \rightarrow \mathbb{R}$.

Although our original equation has F as in (1.2) and (1.3), we will study Problem (5.1) with a more general form of F . Motivated by Proposition 4.8, we impose the following conditions on F .

Assumption 5.1. *The function $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies $F(0) = 0$, and there exist constants $s > 2$ and $c_1, c_2 > 0$ such that, for all $x, y \in \mathbb{R}^n$,*

$$|F(x) - F(y)| \leq c_1(1 + |x|^{s-2} + |y|^{s-2})|x - y|, \quad (5.2)$$

$$(F(x) - F(y)) \cdot (x - y) \geq c_2|x - y|^s. \quad (5.3)$$

Taking $y = 0$ in (5.2) and (5.3), we obtain, for all $x \in \mathbb{R}^n$,

$$|F(x)| \leq c_1(|x| + |x|^{s-1}), \quad (5.4)$$

$$F(x) \cdot x \geq c_2|x|^s. \quad (5.5)$$

Note from (5.2) that F is locally Lipschitz continuous.

We find a variational formulation for Problem (5.1). By using testing functions $\mathbf{v} : \Omega \rightarrow \mathbb{R}^n$ for the first equation, and $q : \Omega \rightarrow \mathbb{R}$ for the second equation, integrating over Ω , using formal integration

by parts for the right-hand side of the first equations and utilizing the boundary condition in the third equation, we formally obtain

$$\begin{cases} \int_{\Omega} F(\mathbf{u}) \cdot \mathbf{v} \, dx - \int_{\Omega} p \operatorname{div} \mathbf{v} \, dx = - \int_{\partial\Omega} \psi \mathbf{v} \cdot \vec{\nu} \, d\sigma & \text{for all } \mathbf{v} \in C_c^\infty(\mathbb{R}^n, \mathbb{R}^n), \\ \int_{\Omega} q \operatorname{div} \mathbf{u} \, dx = \int_{\Omega} f q \, dx & \text{for all } q \in C_c^\infty(\mathbb{R}^n). \end{cases} \quad (5.6)$$

We need to find suitable functional spaces for solutions (\mathbf{u}, p) and testing functions \mathbf{v}, q . Throughout this section, r is the Hölder conjugate of s . Hence,

$$r = \frac{s}{s-1} \in (1, 2). \quad (5.7)$$

We define the spaces $V = \mathbf{W}_s(\operatorname{div}, \Omega)$ as in (2.15) and $Z = L^r(\Omega)$. Recall that their norms are $\|\cdot\|_V = \|\cdot\|_{\mathbf{W}_s(\operatorname{div}, \Omega)}$ as in (2.16) and $\|\cdot\|_Z = \|\cdot\|_{0,r}$.

For the first integral in (5.6), we define $a : \mathbf{L}^s(\Omega) \times \mathbf{L}^s(\Omega) \rightarrow \mathbb{R}$ by

$$a(\mathbf{u}, \mathbf{v}) = \int_{\Omega} F(\mathbf{u}(x)) \cdot \mathbf{v}(x) \, dx \text{ for } \mathbf{u}, \mathbf{v} \in \mathbf{L}^s(\Omega).$$

For the second and fourth integrals in (5.6), we define a bilinear form $b : V \times Z \rightarrow \mathbb{R}$ by

$$b(\mathbf{v}, q) = \int_{\Omega} (\operatorname{div} \mathbf{v}) q \, dx \text{ for } \mathbf{v} \in V, q \in Z.$$

Forcing function. For the fifth integral in (5.6), we assume $f \in L^s(\Omega)$ and define $\Phi_f : Z \rightarrow \mathbb{R}$ by

$$\Phi_f(q) = \int_{\Omega} f q \, dx \text{ for } q \in Z.$$

Then $\Phi_f \in Z'$ and

$$\|\Phi_f\|_{Z'} = \|f\|_{0,s}. \quad (5.8)$$

Boundary data. For the third integral in (5.6), we assume $\psi \in X_r$ and define $\mathcal{T}_\psi : V \rightarrow \mathbb{R}$ by

$$\mathcal{T}_\psi(\mathbf{v}) = \langle \gamma_{n,s}(\mathbf{v}), \psi \rangle_{X_r', X_r} \text{ for } \mathbf{v} \in V.$$

Thanks to the Green's formula (2.19), $\mathcal{T}_\psi(\mathbf{v})$ is the rigorous formulation for the boundary integral in (5.6). By (2.18), one has $\mathcal{T}_\psi \in V'$ and

$$\|\mathcal{T}_\psi\|_{V'} \leq \bar{c}_1 \|\psi\|_{X_r}. \quad (5.9)$$

Definition 5.2. Given $f \in L^s(\Omega)$ and $\psi \in X_r$, a weak solution of Problem (5.1) is a pair $(\mathbf{u}, p) \in V \times Z$ that satisfies

$$\begin{cases} a(\mathbf{u}, \mathbf{v}) - b(\mathbf{v}, p) = -\mathcal{T}_\psi(\mathbf{v}) & \text{for all } \mathbf{v} \in V, \\ b(\mathbf{u}, q) = \Phi_f(q) & \text{for all } q \in Z. \end{cases} \quad (5.10)$$

We obtain the existence, uniqueness, estimates and continuous dependence for weak solutions of Problem (5.1) in the next theorem.

Theorem 5.3. *The following statements hold true.*

- (i) For any $f \in L^s(\Omega)$ and $\psi \in X_r$, there exists a unique weak solution $(\mathbf{u}, p) \in V \times Z$ of Problem (5.1).
- (ii) There is $c_3 > 0$ such that if $f, \psi, (\mathbf{u}, p)$ are as in part (i) then

$$\|\mathbf{u}\|_{0,s} + \|\operatorname{div} \mathbf{u}\|_{0,s} + \|p\|_{0,r} \leq c_3 (\|f\|_{0,s}^{r-1} + \|f\|_{0,s}^{s-1} + \|\psi\|_{X_r}^{r-1} + \|\psi\|_{X_r}). \quad (5.11)$$

(iii) For $j = 1, 2$, assume $f_j, \psi_j, (\mathbf{u}_j, p_j)$ are as in part (i). Let

$$M_0 = 1 + \sum_{j=1}^2 (\|f_j\|_{0,s}^{s-2} + \|\psi_j\|_{X_r}^{2-r}).$$

Then

$$\begin{aligned} & \|\mathbf{u}_1 - \mathbf{u}_2\|_{0,s} + \|\operatorname{div}(\mathbf{u}_1 - \mathbf{u}_2)\|_{0,s} + \|p_1 - p_2\|_{0,r} \\ & \leq c_4 \left(\|f_1 - f_2\|_{0,s} + M_0^r \|f_1 - f_2\|_{0,s}^{r-1} + \|\psi_1 - \psi_2\|_{X_r} + M_0 \|\psi_1 - \psi_2\|_{X_r}^{r-1} \right). \end{aligned} \quad (5.12)$$

Consequently,

$$\begin{aligned} & \|\mathbf{u}_1 - \mathbf{u}_2\|_{0,s} + \|\operatorname{div}(\mathbf{u}_1 - \mathbf{u}_2)\|_{0,s} + \|p_1 - p_2\|_{0,r} \leq c_5 \left(1 + \sum_{i=1}^2 (\|f_i\|_{0,s} + \|\psi_i\|_{X_r}) \right)^{r(s-2)} \\ & \times \left(\|f_1 - f_2\|_{0,s}^{r-1} + \|f_1 - f_2\|_{0,s} + \|\psi_1 - \psi_2\|_{X_r}^{r-1} + \|\psi_1 - \psi_2\|_{X_r} \right). \end{aligned} \quad (5.13)$$

Above, c_4 and c_5 are positive constants independent of $f_j, \psi_j, \mathbf{u}_j, p_j$ for $j = 1, 2$.

The proof of Theorem 5.3 will be presented in subsection 5.5 below.

Remark 5.4. Part (iii) of Theorem 5.3 shows that the function that maps $(f, \psi) \in L^s(\Omega) \times X_r$ to the unique weak solution $(\mathbf{u}, p) \in \mathbf{W}_s(\operatorname{div}, \Omega) \times L^r(\Omega)$ of Problem (5.1) is locally Hölder continuous of order $(r - 1)$, which belongs to $(0, 1)$ thanks to property (5.7).

5.1. Variational formulation on $V \times Z$. Note that system (5.10) has variational formulation on V in its first equation and on Z in its second equation. We will convert (5.10) to a variational formulation on $V \times Z$.

Suppose $(\mathbf{u}, p) \in V \times Z$ is a solution of (5.10). Then adding two equations in (5.10) gives

$$\widehat{a}((\mathbf{u}, p), (\mathbf{v}, q)) = -\mathcal{T}_\psi(\mathbf{v}) + \Phi_f(q) \text{ for all } (\mathbf{v}, q) \in V \times Z. \quad (5.14)$$

Here, \widehat{a} is defined on $(V \times Z)^2$ by

$$\widehat{a}((\mathbf{u}, p), (\mathbf{v}, q)) = a(\mathbf{u}, \mathbf{v}) - b(\mathbf{v}, p) + b(\mathbf{u}, q) \text{ for } (\mathbf{u}, p), (\mathbf{v}, q) \in V \times Z.$$

Definition 5.5. Denote $\mathcal{W} = V \times Z$ with the norm $\|(\mathbf{u}, p)\|_{\mathcal{W}} = \|\mathbf{u}\|_V + \|p\|_Z$ for $\mathbf{u} \in V$ and $p \in Z$.

Because both V and Z are reflexive Banach spaces, so is \mathcal{W} .

For any $\mathbf{u}, \mathbf{v} \in \mathbf{L}^s(\Omega)$, one has from property (5.4) that

$$|a(\mathbf{u}, \mathbf{v})| \leq \int_{\Omega} |F(\mathbf{u}(x)) \cdot \mathbf{v}(x)| dx \leq c_1 \int_{\Omega} (|\mathbf{u}(x)| + |\mathbf{u}(x)|^{s-1}) |\mathbf{v}(x)| dx$$

Applying Hölder's inequality yields

$$|a(\mathbf{u}, \mathbf{v})| \leq c_1 \left(\|\mathbf{u}\|_{0,s} + \|\mathbf{u}\|_{0,s}^{s-1} \right) \|\mathbf{v}\|_{0,s}, \quad (5.15)$$

For any $\mathbf{v} \in V$ and $q \in Z$, applying Hölder's inequality gives

$$|b(\mathbf{v}, q)| \leq \|\operatorname{div} \mathbf{v}\|_{0,s} \|p\|_{0,r}. \quad (5.16)$$

For each $(\mathbf{u}, p) \in \mathcal{W}$, the mapping $(\mathbf{v}, q) \in \mathcal{W} \mapsto \widehat{a}((\mathbf{u}, p), (\mathbf{v}, q))$ is linear, and one has, following (5.15), (5.16) and (2.17),

$$\begin{aligned} |\widehat{a}((\mathbf{u}, p), (\mathbf{v}, q))| & \leq c_1 \left(\|\mathbf{u}\|_{0,s} + \|\mathbf{u}\|_{0,s}^{s-1} \right) \|\mathbf{v}\|_{0,s} + \|\operatorname{div} \mathbf{v}\|_{0,s} \|p\|_{0,r} + \|\operatorname{div} \mathbf{u}\|_{0,s} \|q\|_{0,r} \\ & \leq 2^{1/r} \max\{c_1, 1\} (\|\mathbf{u}\|_{0,s} + \|\mathbf{u}\|_{0,s}^{s-1} + \|p\|_{0,r} + \|\operatorname{div} \mathbf{u}\|_{0,s}) \|(\mathbf{v}, q)\|_{\mathcal{W}}. \end{aligned}$$

Therefore, one can define a mapping $\mathcal{A} : \mathcal{W} \rightarrow \mathcal{W}'$ by

$$\langle \mathcal{A}(\mathbf{u}, p), (\mathbf{v}, q) \rangle_{\mathcal{W}', \mathcal{W}} = \widehat{a}((\mathbf{u}, p), (\mathbf{v}, q)) \text{ for } (\mathbf{u}, p), (\mathbf{v}, q) \in \mathcal{W}.$$

For the right-hand side of (5.14), we define a linear functional $\mathcal{F}_{\psi,f}$ on \mathcal{W} by

$$\mathcal{F}_{\psi,f}(\mathbf{v}, q) = -\mathcal{T}_{\psi}(\mathbf{v}) + \Phi_f(q), \text{ for } (\mathbf{v}, q) \in \mathcal{W}. \quad (5.17)$$

For any $(\mathbf{v}, q) \in \mathcal{W}$, one has from (5.17) that

$$|\mathcal{F}_{\psi,f}(\mathbf{v}, q)| \leq \|\mathcal{T}_{\psi}\|_{V'} \|\mathbf{v}\|_V + \|\Phi_f\|_{Z'} \|q\|_Z \leq (\|\mathcal{T}_{\psi}\|_{V'} + \|\Phi_f\|_{Z'}) \|(\mathbf{v}, q)\|_{\mathcal{W}}.$$

Thus, $\mathcal{F}_{\psi,f} \in \mathcal{W}'$ and we can rewrite (5.14) as

$$\mathcal{A}(\mathbf{u}, p) = \mathcal{F}_{\psi,f}. \quad (5.18)$$

Now, suppose $(\mathbf{u}, p) \in \mathcal{W}$ is a solution of (5.18). By taking $q = 0$ and $\mathbf{v} \in V$ arbitrary in (5.14), we obtain the first equation of (5.10). By taking $\mathbf{v} = 0$ and $q \in Z$ arbitrary in (5.14), we obtain the second equation of (5.10). Therefore, (\mathbf{u}, p) is a solution of (5.10). We have proved that

$$\text{problems (5.10) and (5.18) are equivalent.} \quad (5.19)$$

Equation (5.18), in turn, is a special case of a more general class of equations

$$\mathcal{A}(\mathbf{u}, p) = \mathcal{F}, \quad \mathcal{F} \in \mathcal{W}'. \quad (5.20)$$

We will study (5.20) and apply the findings to (5.18).

Definition 5.6. For $\mathcal{F} \in \mathcal{W}'$, define $\pi_1 \mathcal{F} : V \rightarrow \mathbb{R}$ and $\pi_2 \mathcal{F} : Z \rightarrow \mathbb{R}$ by

$$(\pi_1 \mathcal{F})(\mathbf{v}) = \mathcal{F}(\mathbf{v}, 0) \text{ for } \mathbf{v} \in V \text{ and } (\pi_2 \mathcal{F})(q) = \mathcal{F}(0, q) \text{ for } q \in Z.$$

Then $\pi_1 : \mathcal{W}' \rightarrow V'$ and $\pi_2 : \mathcal{W}' \rightarrow Z'$ are linear, and

$$\|\pi_1 \mathcal{F}\|_{V'}, \|\pi_2 \mathcal{F}\|_{Z'} \leq \|\mathcal{F}\|_{\mathcal{W}'}. \quad (5.21)$$

One has, for all $(\mathbf{v}, q) \in \mathcal{W}$, $\mathcal{F}(\mathbf{v}, q) = (\pi_1 \mathcal{F})(\mathbf{v}) + (\pi_2 \mathcal{F})(q)$, which implies

$$|\mathcal{F}(\mathbf{v}, q)| \leq \|\pi_1 \mathcal{F}\|_{V'} \|\mathbf{v}\|_V + \|\pi_2 \mathcal{F}\|_{Z'} \|q\|_Z, \quad (5.22)$$

and, consequently, $\|\mathcal{F}\|_{\mathcal{W}'} \leq \|\pi_1 \mathcal{F}\|_{V'} + \|\pi_2 \mathcal{F}\|_{Z'}$.

Similar to (5.19), we have

Lemma 5.7. Equation (5.20) is equivalent to

$$\begin{cases} a(\mathbf{u}, \mathbf{v}) - b(\mathbf{v}, p) = (\pi_1 \mathcal{F})(\mathbf{v}) & \text{for all } \mathbf{v} \in V, \\ b(\mathbf{u}, q) = (\pi_2 \mathcal{F})(q) & \text{for all } q \in Z. \end{cases} \quad (5.23)$$

When $\mathcal{F} = \mathcal{F}_{\psi,f}$, we have

$$\pi_1(\mathcal{F}_{\psi,f}) = -\mathcal{T}_{\psi}, \quad \pi_2(\mathcal{F}_{\psi,f}) = \Phi_f, \quad (5.24)$$

and system (5.23) becomes (5.10).

In studying (5.20) and (5.23), the coercivity of $a(\cdot, \cdot)$ and \mathcal{A} plays an important role. For any $\mathbf{u} \in \mathbf{L}^s(\Omega)$, using property (5.5), we have

$$a(\mathbf{u}, \mathbf{u}) = \int_{\Omega} F(\mathbf{u}(x)) \cdot \mathbf{u}(x) dx \geq c_2 \int_{\Omega} |\mathbf{u}(x)|^s dx = c_2 \|\mathbf{u}\|_{0,s}^s. \quad (5.25)$$

Thus, $a(\cdot, \cdot)$ is coercive on $\mathbf{L}^s(\Omega)$. For \mathcal{A} to be coercive, it must satisfy

$$\lim_{\|(\mathbf{u}, p)\|_{\mathcal{W}} \rightarrow \infty} \frac{\langle \mathcal{A}(\mathbf{u}, p), (\mathbf{u}, p) \rangle_{\mathcal{W}', \mathcal{W}}}{\|(\mathbf{u}, p)\|_{\mathcal{W}}} = \infty. \quad (5.26)$$

However, for any $(\mathbf{u}, p) \in \mathcal{W}$,

$$\langle \mathcal{A}(\mathbf{u}, p), (\mathbf{u}, p) \rangle_{\mathcal{W}', \mathcal{W}} = \widehat{a}((\mathbf{u}, p), (\mathbf{u}, p)) = a(\mathbf{u}, \mathbf{u}). \quad (5.27)$$

Hence, (5.26) fails and \mathcal{A} is not coercive on \mathcal{W} . To overcome this issue, we need some regularized problem.

5.2. The regularized problem. For $\mathbf{u}, \mathbf{v} \in V$ and $p, q \in Z$, define

$$R_1(\mathbf{u}, \mathbf{v}) = \int_{\Omega} |\operatorname{div} \mathbf{u}|^{s-2} \operatorname{div} \mathbf{u} \cdot \operatorname{div} \mathbf{v} \, dx \quad \text{and} \quad R_2(p, q) = \int_{\Omega} |p|^{r-2} p \cdot q \, dx. \quad (5.28)$$

For $\varepsilon > 0$, we consider the following regularized problem of (5.20) and (5.23): Find $(\mathbf{u}, p) \in V \times Z$ such that

$$\begin{cases} a(\mathbf{u}, \mathbf{v}) + \varepsilon R_1(\mathbf{u}, \mathbf{v}) - b(\mathbf{v}, p) = (\pi_1 \mathcal{F})(\mathbf{v}) & \text{for all } \mathbf{v} \in V, \\ \varepsilon R_2(p, q) + b(\mathbf{u}, q) = (\pi_2 \mathcal{F})(q) & \text{for all } q \in Z. \end{cases} \quad (5.29)$$

Adding the equations in (5.29) yields

$$\widehat{a}_{\varepsilon}((\mathbf{u}, p), (\mathbf{v}, q)) = \mathcal{F}(\mathbf{v}, q) \quad \text{for all } (\mathbf{v}, q) \in V \times Z. \quad (5.30)$$

Here, $\widehat{a}_{\varepsilon}$ is defined on $(V \times Z)^2$ by

$$\widehat{a}_{\varepsilon}((\mathbf{u}, p), (\mathbf{v}, q)) = \widehat{a}((\mathbf{u}, \mathbf{v}), (\mathbf{v}, q)) + \varepsilon R_1(\mathbf{u}, \mathbf{v}) + \varepsilon R_2(p, q) \quad \text{for } (\mathbf{u}, p), (\mathbf{v}, q) \in V \times Z. \quad (5.31)$$

Note, for $\mathbf{u}, \mathbf{v} \in V$ and $p, q \in Z$, that

$$|R_1(\mathbf{u}, \mathbf{v})| \leq \|\operatorname{div} \mathbf{u}\|_{0,s}^{s-1} \|\operatorname{div} \mathbf{v}\|_{0,s}, \quad |R_2(p, q)| \leq \|p\|_{0,r}^{r-1} \|q\|_{0,r}, \quad (5.32)$$

and

$$\begin{aligned} |\widehat{a}_{\varepsilon}((\mathbf{u}, p), (\mathbf{v}, q))| &\leq \|\mathcal{A}(\mathbf{u}, p)\|_{\mathcal{W}'} \|(\mathbf{v}, q)\|_{\mathcal{W}} + \varepsilon \|\operatorname{div} \mathbf{u}\|_{0,s}^{s-1} \|\operatorname{div} \mathbf{v}\|_{0,s} + \varepsilon \|p\|_{0,r}^{r-1} \|q\|_{0,r} \\ &\leq (\|\mathcal{A}(\mathbf{u}, p)\|_{\mathcal{W}'} + \varepsilon \|(\mathbf{u}, p)\|_{\mathcal{W}}^{s-1} + \varepsilon \|(\mathbf{u}, p)\|_{\mathcal{W}}^{r-1}) \cdot 2 \|(\mathbf{v}, q)\|_{\mathcal{W}}. \end{aligned}$$

Then, similar to the definition of \mathcal{A} , there is a unique mapping $\mathcal{A}_{\varepsilon} : \mathcal{W} \rightarrow \mathcal{W}'$ so that

$$\langle \mathcal{A}_{\varepsilon}(\mathbf{u}, p), (\mathbf{v}, q) \rangle_{\mathcal{W}', \mathcal{W}} = \widehat{a}_{\varepsilon}((\mathbf{u}, p), (\mathbf{v}, q)) \quad \text{for all } (\mathbf{u}, p), (\mathbf{v}, q) \in \mathcal{W}.$$

Similar to Lemma 5.7, we have the following.

Lemma 5.8. *Problem (5.29) is equivalent to*

$$\mathcal{A}_{\varepsilon}(\mathbf{u}, p) = \mathcal{F}. \quad (5.33)$$

Together with (5.15) and (5.25), the following are the basic properties of $a(\cdot, \cdot)$.

Lemma 5.9. *For any $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbf{L}^s(\Omega)$, one has*

$$|a(\mathbf{u}, \mathbf{w}) - a(\mathbf{v}, \mathbf{w})| \leq c_6 \left(1 + \|\mathbf{u}\|_{0,s}^{s-2} + \|\mathbf{v}\|_{0,s}^{s-2}\right) \|\mathbf{u} - \mathbf{v}\|_{0,s} \|\mathbf{w}\|_{0,s}, \quad (5.34)$$

$$a(\mathbf{u}, \mathbf{u} - \mathbf{v}) - a(\mathbf{v}, \mathbf{u} - \mathbf{v}) \geq c_2 \|\mathbf{u} - \mathbf{v}\|_{0,s}^s, \quad (5.35)$$

where c_6 is a positive constant, and c_2 is from Assumption 5.1.

Proof. Using property (5.2), we have

$$|a(\mathbf{u}, \mathbf{w}) - a(\mathbf{v}, \mathbf{w})| \leq \int_{\Omega} |F(\mathbf{u}(x)) - F(\mathbf{v}(x))| |\mathbf{w}(x)| \, dx \leq c_1 \int_{\Omega} (1 + |\mathbf{u}|^{s-2} + |\mathbf{v}|^{s-2}) |\mathbf{u} - \mathbf{v}| |\mathbf{w}| \, dx.$$

Applying Hölder's inequality to three functions and powers $s/(s-2)$, s , s gives

$$|a(\mathbf{u}, \mathbf{w}) - a(\mathbf{v}, \mathbf{w})| \leq c_1 \left(\|1\|_{0,s}^{s-2} + \|\mathbf{u}\|_{0,s}^{s-2} + \|\mathbf{v}\|_{0,s}^{s-2} \right) \|\mathbf{u} - \mathbf{v}\|_{0,s} \|\mathbf{w}\|_{0,s}.$$

Thus, we obtain (5.34) with $c_6 = c_1 \max\{1, |\Omega|^{\frac{s-2}{s}}\}$. By (5.3), we have

$$a(\mathbf{u}, \mathbf{u} - \mathbf{v}) - a(\mathbf{v}, \mathbf{u} - \mathbf{v}) = \int_{\Omega} (F(\mathbf{u}) - F(\mathbf{v})) \cdot (\mathbf{u} - \mathbf{v}) \, dx \geq c_2 \int_{\Omega} |\mathbf{u} - \mathbf{v}|^s \, dx,$$

which proves (5.35). \square

Proposition 5.10. *For every $\varepsilon > 0$ and $\mathcal{F} \in \mathcal{W}'$, there exists a unique solution $(\mathbf{m}_{\varepsilon}, p_{\varepsilon}) \in V \times Z$ of the regularized problem (5.33).*

Proof. Below, we establish that \mathcal{A}_ε is continuous, coercive and strictly monotone.

Proof of the fact \mathcal{A}_ε is continuous. For any $(\mathbf{u}_1, p_1), (\mathbf{u}_2, p_2), (\mathbf{v}, q) \in V \times Z$, we have

$$\begin{aligned} J &\stackrel{\text{def}}{=} \langle \mathcal{A}_\varepsilon(\mathbf{u}_1, p_1) - \mathcal{A}_\varepsilon(\mathbf{u}_2, p_2), (\mathbf{v}, q) \rangle_{\mathcal{W}', \mathcal{W}} \\ &= J_1 + \varepsilon J_2 + \varepsilon J_3 - b(\mathbf{v}, p_1 - p_2) + b(\mathbf{u}_1 - \mathbf{u}_2, q), \end{aligned} \quad (5.36)$$

where

$$\begin{aligned} J_1 &= a(\mathbf{u}_1, \mathbf{v}) - a(\mathbf{u}_2, \mathbf{v}), \\ J_2 &= R_1(\mathbf{u}_1, \mathbf{v}) - R_1(\mathbf{u}_2, \mathbf{v}) = \int_{\Omega} (|\operatorname{div} \mathbf{u}_1|^{s-2} \operatorname{div} \mathbf{u}_1 - |\operatorname{div} \mathbf{u}_2|^{s-2} \operatorname{div} \mathbf{u}_2) \operatorname{div} \mathbf{v} \, dx, \\ J_3 &= R_2(p_1, q) - R_2(p_2, q) = \int_{\Omega} (|p_1|^{r-2} p_1 - |p_2|^{r-2} p_2) q \, dx. \end{aligned}$$

By (5.34),

$$\begin{aligned} |J_1| &\leq c_6 \left(1 + \|\mathbf{u}_1\|_{0,s}^{s-2} + \|\mathbf{u}_2\|_{0,s}^{s-2}\right) \|\mathbf{u}_1 - \mathbf{u}_2\|_{0,s} \|\mathbf{v}\|_{0,s} \\ &\leq c_6 \left(1 + \|\mathbf{u}_1\|_V^{s-2} + \|\mathbf{u}_2\|_V^{s-2}\right) \|\mathbf{u}_1 - \mathbf{u}_2\|_V \|\mathbf{v}\|_V. \end{aligned}$$

By (2.9) and Hölder's inequality we find that

$$\begin{aligned} |J_2| &\leq 2^{s-2} \int_{\Omega} (|\operatorname{div} \mathbf{u}_1|^{s-2} + |\operatorname{div} \mathbf{u}_2|^{s-2}) \cdot |\operatorname{div}(\mathbf{u}_1 - \mathbf{u}_2)| \cdot |\operatorname{div} \mathbf{v}| \, dx \\ &\leq 2^{s-2} (\|\operatorname{div} \mathbf{u}_1\|_{0,s}^{s-2} + \|\operatorname{div} \mathbf{u}_2\|_{0,s}^{s-2}) \|\operatorname{div}(\mathbf{u}_1 - \mathbf{u}_2)\|_{0,s} \|\operatorname{div} \mathbf{v}\|_{0,s} \\ &\leq 2^{s-2} (\|\mathbf{u}_1\|_V^{s-2} + \|\mathbf{u}_2\|_V^{s-2}) \|\mathbf{u}_1 - \mathbf{u}_2\|_V \|\mathbf{v}\|_V. \end{aligned}$$

By (2.12), using Hölder's inequality, and noting that $r - 2 \in (-1, 0)$, we have

$$\begin{aligned} |J_3| &\leq \int_{\Omega} \left| |p_1|^{r-2} p_1 - |p_2|^{r-2} p_2 \right| \cdot |q| \, dx \leq 2^{2-r} \int_{\Omega} |p_1 - p_2|^{r-1} |q| \, dx \leq C \|p_1 - p_2\|_{0,r}^{r-1} \|q\|_{0,r} \\ &\leq 2^{2-r} \|p_1 - p_2\|_Z^{r-1} \|q\|_Z. \end{aligned}$$

By (5.16),

$$\begin{aligned} |b(\mathbf{v}, p_1 - p_2)| + |b(\mathbf{u}_1 - \mathbf{u}_2, q)| &\leq \|\operatorname{div} \mathbf{v}\|_{0,s} \|p_1 - p_2\|_{0,r} + \|\operatorname{div}(\mathbf{u}_1 - \mathbf{u}_2)\|_{0,s} \|q\|_{0,r} \\ &\leq \|p_1 - p_2\|_Z \|\mathbf{v}\|_V + \|\mathbf{u}_1 - \mathbf{u}_2\|_V \|q\|_Z. \end{aligned}$$

From (5.36) and the above estimates, it follows that

$$\begin{aligned} |J| &\leq C_1 (1 + \varepsilon) \left(1 + \|\mathbf{u}_1\|_V^{s-2} + \|\mathbf{u}_2\|_V^{s-2}\right) \\ &\quad \times \left(\|\mathbf{u}_1 - \mathbf{u}_2\|_V + \|p_1 - p_2\|_Z + \|p_1 - p_2\|_Z^{r-1}\right) (\|\mathbf{v}\|_V + \|q\|_Z), \end{aligned}$$

where $C_1 = c_6 + 2^{s-2} + 2^{2-r} + 1$. This yields

$$\begin{aligned} \|\mathcal{A}_\varepsilon(\mathbf{u}_1, p_1) - \mathcal{A}_\varepsilon(\mathbf{u}_2, p_2)\|_{\mathcal{W}'} &\leq C_1 (1 + \varepsilon) \left(1 + \|\mathbf{u}_1\|_V^{s-2} + \|\mathbf{u}_2\|_V^{s-2}\right) \\ &\quad \times \left(\|\mathbf{u}_1 - \mathbf{u}_2\|_V + \|p_1 - p_2\|_Z + \|p_1 - p_2\|_Z^{r-1}\right). \end{aligned}$$

Thus, \mathcal{A}_ε is continuous.

Proof of the fact \mathcal{A}_ε is coercive. We need to prove (5.26) for \mathcal{A}_ε replacing \mathcal{A} . For any $(\mathbf{u}, p) \in \mathcal{W}$, we have from (5.25), (5.27), (5.28) and (5.31), that

$$\begin{aligned} \langle \mathcal{A}_\varepsilon(\mathbf{u}, p), (\mathbf{u}, p) \rangle_{\mathcal{W}', \mathcal{W}} &= \widehat{a}(\mathbf{u}, p), (\mathbf{u}, p) + \varepsilon R_1(\mathbf{u}, \mathbf{u}) + \varepsilon R_2(p, p) \geq c_2 \|\mathbf{u}\|_{0,s}^s + \varepsilon \|\operatorname{div} \mathbf{u}\|_{0,s}^s + \varepsilon \|p\|_{0,r}^r \\ &\geq \min\{c_2, \varepsilon\} (\|\mathbf{u}\|_V^s + \|p\|_Z^r). \end{aligned}$$

Note that $s > 2 > r > 1$. We consider $\|\mathbf{u}\|_V + \|p\|_Z \geq 2$. If $\|\mathbf{u}\|_V \geq 1$ then

$$\|\mathbf{u}\|_V^s + \|p\|_Z^r \geq \|\mathbf{u}\|_V^r + \|p\|_Z^r \geq 2^{1-r} (\|\mathbf{u}\|_V + \|p\|_Z)^r.$$

If $\|\mathbf{u}\|_V < 1$, then $\|p\|_Z > 1 > \|\mathbf{u}\|_V$, and

$$\|\mathbf{u}\|_V^s + \|p\|_Z^r \geq \|p\|_Z^r \geq \left(\frac{1}{2}\|p\|_Z + \frac{1}{2}\|\mathbf{u}\|_V\right)^r = 2^{-r}(\|\mathbf{u}\|_V + \|p\|_Z)^r.$$

In both cases, we find that

$$\frac{\langle \mathcal{A}_\varepsilon(\mathbf{u}, p), (\mathbf{u}, p) \rangle_{\mathcal{W}', \mathcal{W}}}{\|(\mathbf{u}, p)\|_{\mathcal{W}}} \geq 2^{-r} \min\{c_2, \varepsilon\} \|(\mathbf{u}, p)\|_{\mathcal{W}}^{r-1} \rightarrow \infty \text{ as } \|(\mathbf{u}, p)\|_{\mathcal{W}} \rightarrow \infty.$$

Therefore, \mathcal{A}_ε is coercive.

Proof of the fact \mathcal{A}_ε is strictly monotone. By ‘‘strictly monotone’’, we mean that

$$\langle \mathcal{A}_\varepsilon(\mathbf{u}, p) - \mathcal{A}_\varepsilon(\mathbf{v}, q), (\mathbf{u}, p) - (\mathbf{v}, q) \rangle_{\mathcal{W}', \mathcal{W}} > 0$$

for all $(\mathbf{u}, p), (\mathbf{v}, q) \in \mathcal{W}$ with $(\mathbf{u}, p) \neq (\mathbf{v}, q)$.

Let $(\mathbf{u}, p), (\mathbf{v}, q) \in \mathcal{W}$. We have

$$\begin{aligned} I &\stackrel{\text{def}}{=} \langle \mathcal{A}_\varepsilon(\mathbf{u}, p) - \mathcal{A}_\varepsilon(\mathbf{v}, q), (\mathbf{u} - \mathbf{v}, p - q) \rangle_{\mathcal{W}', \mathcal{W}} \\ &= a(\mathbf{u}, \mathbf{u} - \mathbf{v}) - a(\mathbf{v}, \mathbf{u} - \mathbf{v}) + \varepsilon(R_1(\mathbf{u}, \mathbf{u} - \mathbf{v}) - R_1(\mathbf{v}, \mathbf{u} - \mathbf{v})) + \varepsilon(R_2(p, p - q) - R_2(q, p - q)). \end{aligned}$$

By (5.35), we have

$$a(\mathbf{u}, \mathbf{u} - \mathbf{v}) - a(\mathbf{v}, \mathbf{u} - \mathbf{v}) \geq c_2 \|\mathbf{u} - \mathbf{v}\|_{0,s}^s.$$

By (2.11), we have

$$\begin{aligned} R_1(\mathbf{u}, \mathbf{u} - \mathbf{v}) - R_1(\mathbf{v}, \mathbf{u} - \mathbf{v}) &= \int_{\Omega} (|\operatorname{div} \mathbf{u}|^{s-2} \operatorname{div} \mathbf{u} - |\operatorname{div} \mathbf{v}|^{s-2} \operatorname{div} \mathbf{v}) \cdot (\operatorname{div} \mathbf{u} - \operatorname{div} \mathbf{v}) \, dx \\ &\geq 2^{1-s} \|\operatorname{div}(\mathbf{u} - \mathbf{v})\|_{0,s}^s. \end{aligned}$$

Utilizing inequality (2.13), we have

$$R_2(p, p - q) - R_2(q, p - q) = \int_{\Omega} (|p|^{r-2} p - |q|^{r-2} q) \cdot (p - q) \, dx \geq (r-1) \int_{\Omega} (|p| + |q|)^{r-2} |p - q|^2 \, dx.$$

Thus,

$$I \geq C_2 \left(\|\mathbf{u} - \mathbf{v}\|_{0,s}^s + \varepsilon \|\operatorname{div}(\mathbf{u} - \mathbf{v})\|_{0,s}^s + \varepsilon \int_{\Omega} (|p| + |q|)^{r-2} |p - q|^2 \, dx \right),$$

where $C_2 = \min\{c_2, 2^{1-s}, r-1\}$. This implies that I is positive whenever $(\mathbf{u}, p) \neq (\mathbf{v}, q)$. Therefore, \mathcal{A}_ε is strictly monotone.

By the Browder–Minty Theorem [22, Theorem 26.A], there exists a unique solution $(\mathbf{u}_\varepsilon, p_\varepsilon) \in \mathcal{W}$ of equation (5.33). \square

Next, we obtain a uniform upper bound for solutions $(\mathbf{u}_\varepsilon, p_\varepsilon)$ of (5.33) when ε is sufficiently small.

Lemma 5.11. *There exists $\varepsilon_0 > 0$ such that if $\varepsilon \in (0, \varepsilon_0)$, $\mathcal{F} \in \mathcal{W}'$, and $(\mathbf{u}_\varepsilon, p_\varepsilon)$ is the unique solution of (5.33), then*

$$\|(\mathbf{u}_\varepsilon, p_\varepsilon)\|_{\mathcal{W}} \leq \mathcal{C}, \tag{5.37}$$

where number $\mathcal{C} \geq 0$ is independent of ε and depends increasingly on $\|\mathcal{F}\|_{\mathcal{W}'}$.

Proof. Thanks to Lemma 5.8, $(\mathbf{u}_\varepsilon, p_\varepsilon)$ is the unique solution of the equivalent system (5.29). Denote $\beta = \|\pi_1 \mathcal{F}\|_{V'}$ and $\gamma = \|\pi_2 \mathcal{F}\|_{Z'}$. Let $\varepsilon > 0$. In calculations below, C denotes a positive constant independent of ε , β , γ that may have varied values from one line to another, while C_i , for $i = 0, 1, 2, 3$, denotes a positive constant independent of ε , β , γ with a fixed value.

Choosing $q = |\operatorname{div} \mathbf{u}_\varepsilon|^{s-2} \operatorname{div} \mathbf{u}_\varepsilon \in Z$ in the the second equation of (5.29) and using Hölder’s inequality, we find that

$$\|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}^s \leq \gamma \|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}^{s-1} + \varepsilon \|p_\varepsilon\|_{0,r}^{r-1} \|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}^{s-1}.$$

Consequently,

$$\|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s} \leq \gamma + \varepsilon \|p_\varepsilon\|_{0,r}^{r-1}. \quad (5.38)$$

Taking $(\mathbf{u}, p) = (\mathbf{v}, q) = (\mathbf{u}_\varepsilon, p_\varepsilon)$ in (5.30) and using (5.22), (2.17) give

$$a(\mathbf{u}_\varepsilon, \mathbf{u}_\varepsilon) + \varepsilon \|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}^s + \varepsilon \|p_\varepsilon\|_{0,r}^r = \mathcal{F}(\mathbf{u}_\varepsilon, p_\varepsilon) \leq \beta (\|\mathbf{u}_\varepsilon\|_{0,s} + \|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}) + \gamma \|p_\varepsilon\|_{0,r}. \quad (5.39)$$

Applying inequality (5.15) to the first term of (5.39), neglecting the next two terms, and utilizing the estimate (5.38) for the last divergence term, we obtain

$$c_2 \|\mathbf{u}_\varepsilon\|_{0,s}^s \leq \beta (\|\mathbf{u}_\varepsilon\|_{0,s} + \gamma + \varepsilon \|p_\varepsilon\|_{0,r}^{r-1}) + \gamma \|p_\varepsilon\|_{0,r}.$$

By Young's inequality, specifically, the last one in (2.4), one has $\beta \|\mathbf{u}_\varepsilon\|_{0,s} \leq (c_2/2) \|\mathbf{u}_\varepsilon\|_{0,s}^s + C\beta^r$. It follows that

$$\|\mathbf{u}_\varepsilon\|_{0,s}^s \leq C \left(\beta^r + \beta\gamma + \varepsilon\beta \|p_\varepsilon\|_{0,r}^{r-1} + \gamma \|p_\varepsilon\|_{0,r} \right). \quad (5.40)$$

Next, we use (2.20) to bound $\|p_\varepsilon\|_{0,r}$ and close the estimates (5.38) and (5.40). Using $q = p_\varepsilon$ in the first equation of (5.29) and the estimate of $\|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}$ in (5.38) above, we find that

$$\begin{aligned} b(\mathbf{v}, p_\varepsilon) &= a(\mathbf{u}_\varepsilon, \mathbf{v}) + \varepsilon R_1(\mathbf{u}_\varepsilon, \mathbf{v}) - (\pi_1 \mathcal{F}) \mathbf{v} \\ &\leq C (\|\mathbf{u}_\varepsilon\|_{0,s} + \|\mathbf{u}_\varepsilon\|_{0,s}^{s-1}) \|\mathbf{v}\|_{0,s} + \varepsilon \|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}^{s-1} \|\operatorname{div} \mathbf{v}\|_{0,s} + \beta \|\mathbf{v}\|_V \\ &\leq \left[C (\|\mathbf{u}_\varepsilon\|_{0,s} + \|\mathbf{u}_\varepsilon\|_{0,s}^{s-1}) + \varepsilon (\gamma + \varepsilon \|p_\varepsilon\|_{0,r}^{r-1})^{s-1} + \beta \right] \|\mathbf{v}\|_V. \end{aligned}$$

Applying (2.20) yields

$$\|p_\varepsilon\|_Z \leq C_* \sup_{\mathbf{v} \in V \setminus \{0\}} \frac{b(\mathbf{v}, p_\varepsilon)}{\|\mathbf{v}\|_V} \leq C_0 \left(\|\mathbf{u}_\varepsilon\|_{0,s} + \|\mathbf{u}_\varepsilon\|_{0,s}^{s-1} + \varepsilon \gamma^{s-1} + \varepsilon^s \|p_\varepsilon\|_Z + \beta \right). \quad (5.41)$$

Set $\varepsilon_0 = \min\{1, (2C_0)^{-1/s}\}$ and consider $\varepsilon \in (0, \varepsilon_0)$. We obtain from (5.41) that

$$\|p_\varepsilon\|_Z \leq 2C_0 (\|\mathbf{u}_\varepsilon\|_{0,s} + \|\mathbf{u}_\varepsilon\|_{0,s}^{s-1} + \varepsilon \gamma^{s-1} + \beta). \quad (5.42)$$

Consequently, noting that $(r-1)(s-1) = 1$, one has

$$\|p_\varepsilon\|_Z^{r-1} \leq C (\|\mathbf{u}_\varepsilon\|_{0,s}^{r-1} + \|\mathbf{u}_\varepsilon\|_{0,s} + \varepsilon^{r-1} \gamma + \beta^{r-1}). \quad (5.43)$$

Combining (5.42) and (5.43) with (5.40) leads to

$$\begin{aligned} \|\mathbf{u}_\varepsilon\|_{0,s}^s &\leq C\beta^r + C\beta\gamma + C\varepsilon\beta (\|\mathbf{u}_\varepsilon\|_{0,s}^{r-1} + \|\mathbf{u}_\varepsilon\|_{0,s} + \varepsilon^{r-1} \gamma + \beta^{r-1}) \\ &\quad + C\gamma (\|\mathbf{u}_\varepsilon\|_{0,s} + \|\mathbf{u}_\varepsilon\|_{0,s}^{s-1} + \varepsilon \gamma^{s-1} + \beta) \\ &\leq CJ + C\varepsilon\beta (\|\mathbf{u}_\varepsilon\|_{0,s}^{r-1} + \|\mathbf{u}_\varepsilon\|_{0,s}) + C\gamma (\|\mathbf{u}_\varepsilon\|_{0,s} + \|\mathbf{u}_\varepsilon\|_{0,s}^{s-1}), \end{aligned}$$

where $J = (1 + \varepsilon)\beta^r + (1 + \varepsilon^r)\beta\gamma + \varepsilon\gamma^s$.

Let $s_* = s/(s-r+1) = s(s-1)/(s(s-1)-1) \in (1, r)$. Then using Young's inequality we obtain

$$\|\mathbf{u}_\varepsilon\|_{0,s}^s \leq CJ + \frac{1}{2} \|\mathbf{u}_\varepsilon\|_{0,s}^s + C(\varepsilon^{s_*} \beta^{s_*} + \varepsilon^r \beta^r) + C(\gamma^r + \gamma^s).$$

This implies

$$\|\mathbf{u}_\varepsilon\|_{0,s} \leq C_1 d_1(\varepsilon, \beta, \gamma), \quad (5.44)$$

where

$$d_1(\varepsilon, \beta, \gamma) = \left[(1 + \varepsilon^r)\beta^r + \varepsilon^{s_*} \beta^{s_*} + (1 + \varepsilon^r)\beta\gamma + \gamma^r + (1 + \varepsilon)\gamma^s \right]^{1/s}.$$

Utilizing estimate (5.44) in (5.42) yields

$$\|p_\varepsilon\|_Z \leq C_2 d_2(\varepsilon, \beta, \gamma), \quad \text{where } d_2(\varepsilon, \beta, \gamma) = d_1(\varepsilon, \beta, \gamma) + d_1(\varepsilon, \beta, \gamma)^{s-1} + \varepsilon \gamma^{s-1} + \beta. \quad (5.45)$$

Combining (5.45) and (5.38), we have

$$\|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s} \leq C_3 d_3(\varepsilon, \beta, \gamma), \quad \text{where } d_3(\varepsilon, \beta, \gamma) = \gamma + \varepsilon d_2(\varepsilon, \beta, \gamma)^{r-1}. \quad (5.46)$$

Note that $d_i(\varepsilon, \beta, \gamma)$, for $i = 1, 2, 3$, are increasing in $\varepsilon, \beta, \gamma$, and, by (5.21), $\beta, \gamma \leq \|\mathcal{F}\|_{\mathcal{W}'}$. Summing up the estimates (5.44), (5.45) and (5.46) gives

$$\|(\mathbf{u}_\varepsilon, p_\varepsilon)\|_{\mathcal{W}} \leq \mathcal{C} \stackrel{\text{def}}{=} \sum_{i=1}^3 C_i d_i(1, \|\mathcal{F}\|_{\mathcal{W}'}, \|\mathcal{F}\|_{\mathcal{W}'}).$$

Thus, we obtain the desired estimate (5.37). \square

5.3. Existence and uniqueness.

Theorem 5.12. *For any $\mathcal{F} \in \mathcal{W}'$, equation (5.20) has a unique solution $(\mathbf{u}, p) \in \mathcal{W}$.*

Proof. Given $\mathcal{F} \in \mathcal{W}'$, let ε_0 and \mathcal{C} be as in Lemma 5.11. For $\varepsilon \in (0, \varepsilon_0)$, let $(\mathbf{u}_\varepsilon, p_\varepsilon)$ be the unique solution of the regularized problem (5.33). Note from (5.31), (5.30) and (5.32) that

$$\begin{aligned} & |\widehat{a}((\mathbf{u}_\varepsilon, p_\varepsilon), (\mathbf{v}, q)) - \mathcal{F}(\mathbf{v}, q)| = \varepsilon |R_1(\mathbf{u}_\varepsilon, \mathbf{v}) + R_2(p_\varepsilon, q)| \\ & \leq \varepsilon (\|\operatorname{div} \mathbf{u}_\varepsilon\|_{0,s}^{s-1} \|\operatorname{div} \mathbf{v}\|_{0,s} + \|p_\varepsilon\|_{0,r}^{r-1} \|q\|_{0,r}) \leq \varepsilon (\|\mathbf{u}_\varepsilon\|_V^{s-1} + \|p_\varepsilon\|_Z^{r-1}) \|(\mathbf{v}, q)\|_{\mathcal{W}}. \end{aligned}$$

Hence

$$\|\mathcal{A}(\mathbf{u}_\varepsilon, p_\varepsilon) - \mathcal{F}\|_{\mathcal{W}'} \leq \varepsilon (\|\mathbf{u}_\varepsilon\|_V^{s-1} + \|p_\varepsilon\|_Z^{r-1}) \leq 2\varepsilon(1 + \mathcal{C})^{s-1} \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (5.47)$$

Let $N > 1$ such that $1/N < \varepsilon_0$. By the virtue of Lemma 5.11, the sequence $((\mathbf{u}_{1/k}, p_{1/k}))_{k=N}^\infty$ is a bounded sequence in the reflexive Banach space \mathcal{W} . Then there exists a subsequence $((\mathbf{u}_{1/k_j}, p_{1/k_j}))_{j=1}^\infty$ that converges weakly in \mathcal{W} to an element $(\mathbf{u}, p) \in \mathcal{W}$. Combining this fact with the limit (5.47), and applying a general analysis result [22, part (c), p. 474], we obtain $\mathcal{A}(\mathbf{u}, p) = \mathcal{F}$. Therefore, (\mathbf{u}, p) is a solution of (5.20).

We prove the uniqueness next. Suppose (\mathbf{u}_1, p_1) and (\mathbf{u}_2, p_2) are two solutions of (5.20). Subtracting their equations and choosing the testing functions $(\mathbf{v}, q) = (\mathbf{u}_1 - \mathbf{u}_2, p_1 - p_2)$, we obtain

$$0 = \langle \mathcal{A}(\mathbf{u}_1, p_1) - \mathcal{A}(\mathbf{u}_2, p_2), (\mathbf{u}_1 - \mathbf{u}_2, p_1 - p_2) \rangle_{\mathcal{W}', \mathcal{W}} = a(\mathbf{u}_1, \mathbf{u}_1 - \mathbf{u}_2) - a(\mathbf{u}_2, \mathbf{u}_1 - \mathbf{u}_2).$$

Combining this with property (5.35) yields $0 \geq c_2 \|\mathbf{u}_1 - \mathbf{u}_2\|_{0,s}^s$, which implies $\mathbf{u}_1 = \mathbf{u}_2$.

For $i = 1, 2$, we have the variational equation $a(\mathbf{u}_i, \mathbf{v}) - b(\mathbf{v}, p_i) = (\pi_1 \mathcal{F})\mathbf{v}$ for all $\mathbf{v} \in V$. Subtracting these two equations implies $b(\mathbf{v}, p_1 - p_2) = a(\mathbf{u}_1, \mathbf{v}) - a(\mathbf{u}_2, \mathbf{v}) = 0$ for all $\mathbf{v} \in V$. By Lemma 2.3, we obtain $p_1 = p_2$. \square

5.4. Estimates and continuous dependence. Regarding the unique solutions of equation (5.20), we have the following estimates.

Theorem 5.13. *Assume $\mathcal{F} \in \mathcal{W}'$, $(\mathbf{u}, p) \in \mathcal{W}$, and $\mathcal{A}(\mathbf{u}, p) = \mathcal{F}$. Denote $\beta = \|\pi_1 \mathcal{F}\|_{V'}$ and $\gamma = \|\pi_2 \mathcal{F}\|_{Z'}$. Then*

$$\|\mathbf{u}\|_{0,s} \leq c_7(\beta^{r-1} + \gamma^{r-1} + \gamma), \quad \|\operatorname{div} \mathbf{u}\|_{0,s} \leq c_7\gamma, \quad \|p\|_Z \leq c_7(\gamma^{r-1} + \gamma^{s-1} + \beta^{r-1} + \beta), \quad (5.48)$$

and, hence,

$$\|(\mathbf{u}, p)\|_{\mathcal{W}} \leq c_8 (\gamma^{r-1} + \gamma^{s-1} + \beta^{r-1} + \beta). \quad (5.49)$$

Consequently,

$$\|(\mathbf{u}, p)\|_{\mathcal{W}} \leq c_9 (\|\mathcal{F}\|_{\mathcal{W}'}^{r-1} + \|\mathcal{F}\|_{\mathcal{W}'}^{s-1}). \quad (5.50)$$

Above, the positive constants c_7, c_8, c_9 are independent of $\mathcal{F}, \mathbf{u}, p$.

Proof. We repeat the calculations in Lemma 5.11 with $\varepsilon = 0$. It follows (5.44), (5.45) and (5.46) that

$$\|\mathbf{u}\|_{0,s} \leq C d_1, \quad \|p\|_Z \leq C d_2, \quad \|\operatorname{div} \mathbf{u}\|_{0,s} \leq C d_3,$$

where $d_1 = (\beta^r + \beta\gamma + \gamma^r + \gamma^s)^{1/s}$, $d_2 = d_1 + d_1^{s-1} + \beta$ and $d_3 = \gamma$.

Let C denote a generic positive constant as in the proof of Lemma 5.11. Using inequalities (2.6) and (2.4) yields

$$\begin{aligned} d_1 &\leq C(\beta^r + \gamma^r + \gamma^s)^{1/s} \leq C(\beta^{r-1} + \gamma^{r-1} + \gamma), \\ d_2 &\leq C(\beta^{r-1} + \gamma^{r-1} + \gamma) + C(\beta^{r-1} + \gamma^{r-1} + \gamma)^{s-1} + \beta \leq C(\beta^{r-1} + \beta + \gamma^{r-1} + \gamma^{s-1}). \end{aligned}$$

Therefore, we obtain the estimates in (5.48). Summing up the estimates in (5.48), noticing that $r-1 < 1 < s-1$ and applying inequality (2.4) yield (5.49).

Finally, inequality (5.50) follows (5.49), the fact $\beta, \gamma \leq \|\mathcal{F}\|_{\mathcal{W}'}$, which is due to (5.21), and, again, the use of (2.4). \square

It turns out that the solutions of (5.20) depend continuously on \mathcal{F} , as showed below.

Theorem 5.14. *For $j = 1, 2$, let $\mathcal{F}_j \in \mathcal{W}'$ and $(\mathbf{u}_j, p_j) \in \mathcal{W}$ satisfy $\mathcal{A}(\mathbf{u}_j, p_j) = \mathcal{F}_j$. Denote*

$$M = 1 + \sum_{j=1}^2 (\|\pi_1 \mathcal{F}_j\|_{V'}^{r-1} + \|\pi_2 \mathcal{F}_j\|_{Z'}), \quad \beta = \|\pi_1(\mathcal{F}_1 - \mathcal{F}_2)\|_{V'} \text{ and } \gamma = \|\pi_2(\mathcal{F}_1 - \mathcal{F}_2)\|_{Z'}. \quad (5.51)$$

Then

$$\|(\mathbf{u}_1, p_1) - (\mathbf{u}_2, p_2)\|_{\mathcal{W}} \leq c_{10}(\beta + M^{s-2}\beta^{r-1} + \gamma + M^{r(s-2)}\gamma^{r-1}). \quad (5.52)$$

Consequently,

$$\|(\mathbf{u}_1, p_1) - (\mathbf{u}_2, p_2)\|_{\mathcal{W}} \leq c_{11} (1 + \|\mathcal{F}_1\|_{\mathcal{W}'} + \|\mathcal{F}_2\|_{\mathcal{W}'})^{r(s-2)} (\|\mathcal{F}_1 - \mathcal{F}_2\|_{\mathcal{W}'}^{r-1} + \|\mathcal{F}_1 - \mathcal{F}_2\|_{\mathcal{W}'}). \quad (5.53)$$

Above positive constants c_{10} and c_{11} are independent of $\mathcal{F}_j, \mathbf{u}_j, p_j$ for $j = 1, 2$.

Proof. Below, C denotes a generic positive constant independent of $\mathcal{F}_j, \mathbf{u}_j, p_j$ for $j = 1, 2$.

Denote $\mathbf{u} = \mathbf{u}_1 - \mathbf{u}_2$, $p = p_1 - p_2$, and $\mathcal{F} = \mathcal{F}_1 - \mathcal{F}_2$. Then $\beta = \|\pi_1 \mathcal{F}\|_{V'}$, $\gamma = \|\pi_2 \mathcal{F}\|_{Z'}$ and, thanks to the first estimate in (5.48) for \mathbf{u}_1 and \mathbf{u}_2 ,

$$\|\mathbf{u}_1\|_{0,s} + \|\mathbf{u}_2\|_{0,s} \leq CM. \quad (5.54)$$

For $j = 1, 2$, one has

$$\begin{aligned} a(\mathbf{u}_j, \mathbf{v}) - b(\mathbf{v}, p_j) &= (\pi_1 \mathcal{F}_j)(\mathbf{v}) && \text{for all } \mathbf{v} \in V, \\ b(\mathbf{u}_j, q) &= (\pi_2 \mathcal{F}_j)(q) && \text{for all } q \in Z. \end{aligned}$$

It follows that

$$a(\mathbf{u}_1, \mathbf{v}) - a(\mathbf{u}_2, \mathbf{v}) - b(\mathbf{v}, p) = (\pi_1 \mathcal{F})(\mathbf{v}) \quad \text{for all } \mathbf{v} \in V, \quad (5.55)$$

$$b(\mathbf{u}, q) = (\pi_2 \mathcal{F})(q) \quad \text{for all } q \in Z. \quad (5.56)$$

Taking $q = |\operatorname{div} \mathbf{u}|^{s-2} \operatorname{div} \mathbf{u}$ in (5.56) gives $\|\operatorname{div} \mathbf{u}\|_{0,s}^s \leq \gamma \|\operatorname{div} \mathbf{u}\|_{0,s}^{s-1}$. Thus,

$$\|\operatorname{div} \mathbf{u}\|_{0,s} \leq \gamma. \quad (5.57)$$

For any $\mathbf{v} \in V$, we have from (5.55) and (5.34) that

$$b(\mathbf{v}, p) \leq |(\pi_1 \mathcal{F})\mathbf{v}| + |a(\mathbf{u}_1, \mathbf{v}) - a(\mathbf{u}_2, \mathbf{v})| \leq \beta \|\mathbf{v}\|_V + c_6 \left(1 + \|\mathbf{u}_1\|_{0,s}^{s-2} + \|\mathbf{u}_2\|_{0,s}^{s-2}\right) \|\mathbf{u}\|_{0,s} \|\mathbf{v}\|_V.$$

Hence, making use of Lemma 2.3 to estimate $\|p\|_Z$ and also (5.54) gives

$$\|p\|_Z \leq C(\beta + M^{s-2}\|\mathbf{u}\|_{0,s}). \quad (5.58)$$

Taking $\mathbf{v} = \mathbf{u}$ in (5.55), $q = p$ in (5.56), and adding the resulting equations yield

$$a(\mathbf{u}_1, \mathbf{u}) - a(\mathbf{u}_2, \mathbf{u}) = (\pi_1 \mathcal{F})\mathbf{u} + (\pi_2 \mathcal{F})p.$$

Applying inequality (5.35) to the left-hand side and inequality (5.22) to the right-hand side, we derive

$$\|\mathbf{u}\|_{0,s}^s \leq C(\beta \|\mathbf{u}\|_V + \gamma \|p\|_Z) \leq C(\beta(\|\mathbf{u}\|_{0,s} + \|\operatorname{div} \mathbf{u}\|_{0,s}) + \gamma \|p\|_Z). \quad (5.59)$$

Inserting (5.57) and (5.58) into (5.59) and using Young's inequality give

$$\|\mathbf{u}\|_{0,s}^s \leq C(\beta + M^{s-2}\gamma)\|\mathbf{u}\|_{0,s} + C\beta\gamma \leq \frac{1}{2}\|\mathbf{u}\|_{0,s}^s + C(\beta^r + M^{r(s-2)}\gamma^r + \beta\gamma).$$

Thus,

$$\|\mathbf{u}\|_{0,s} \leq C(\beta^{r-1} + M^{(r-1)(s-2)}\gamma^{r-1} + \beta^{1/s}\gamma^{1/s}). \quad (5.60)$$

Utilizing this estimate in (5.58) gives

$$\|p\|_Z \leq C(\beta + M^{r(s-2)}\gamma^{r-1} + M^{s-2}(\beta^{r-1} + \beta^{1/s}\gamma^{1/s})).$$

By Young's inequality again, $M^{s-2}\beta^{1/s}\gamma^{1/s} \leq \beta + M^{r(s-2)}\gamma^{r-1}$. We obtain

$$\|p\|_Z \leq C\beta + CM^{r(s-2)}\gamma^{r-1} + CM^{s-2}\beta^{r-1}. \quad (5.61)$$

Combining (5.57), (5.60), (5.61) gives

$$\|(\mathbf{u}, p)\|_{\mathcal{W}} \leq C\gamma + CM^{r(s-2)}\gamma^{r-1} + C\beta + CM^{s-2}\beta^{r-1} + C\beta^{1/s}\gamma^{1/s}. \quad (5.62)$$

By Young's inequality, $\beta^{1/s}\gamma^{1/s} \leq \beta + \gamma^{r-1}$. Hence, we obtain (5.52) from (5.62).

Finally, using the facts $\beta, \gamma \leq \|\mathcal{F}\|_{\mathcal{W}'}$ and $M \leq 3(1 + \|\mathcal{F}_1\|_{\mathcal{W}'} + \|\mathcal{F}_2\|_{\mathcal{W}'})$, we obtain (5.53) from (5.52). \square

5.5. Proof of Theorem 5.3.

Proof. Part (i). The statement follows Theorem 5.12 when applied to $\mathcal{F} = \mathcal{F}_{\psi, f}$.

Part (ii). We apply Theorem 5.13 to $\mathcal{F} = \mathcal{F}_{\psi, f}$. Note from (5.24), (5.8) and (5.9) that

$$\beta = \|\mathcal{T}_\psi\|_{V'} \leq \bar{c}_1\|\psi\|_{X^r} \text{ and } \gamma = \|\Phi_f\|_{Z'} = \|f\|_{0,s}.$$

Then estimate (5.11) follows (5.49).

Part (iii). We apply Theorem 5.14 to $\mathcal{F}_j = \mathcal{F}_{\psi_j, f_j}$ for $j = 1, 2$. Using properties (5.24) and (5.8), (5.9) we have M, β and γ defined in (5.51) satisfy $\beta \leq \bar{c}_1\|\psi_1 - \psi_2\|_{X^r}$, $\gamma = \|f_1 - f_2\|_{0,s}$, and, with the observation $(r-1)(s-2) = 2-r$,

$$M^{s-2} \leq C \left(1 + \sum_{j=1}^2 (\|\psi_j\|_{X^r}^{2-r} + \|f_j\|_{0,s}^{s-2}) \right) = CM_0,$$

for some positive constant C . Then estimate (5.12) follows (5.52). Recalling $2-r = (r-1)(s-2) < s-2$ and applying Young's inequality, we obtain (5.13) from (5.12). \square

APPENDIX A.

Lemma A.1. *For any $s \in (1, \infty)$, $\mathbf{W}_s(\text{div}, \Omega)$ is a reflexive Banach space.*

Proof. Firstly, one can verify that $\mathbf{W}_s(\text{div}, \Omega)$ is a Banach space. Same as in [18], we use the mapping $E\mathbf{v} = (\mathbf{v}, \text{div } \mathbf{v})$ for $\mathbf{v} \in \mathbf{W}_s(\text{div}, \Omega)$ to embed $\mathbf{W}_s(\text{div}, \Omega)$ into $(L^s(\Omega))^{n+1}$. Denote $\widetilde{\mathbf{W}}_s = E(\mathbf{W}_s(\text{div}, \Omega)) \subset (L^s(\Omega))^{n+1}$. Then the norm $\|E\mathbf{v}\|_{(L^s(\Omega))^{n+1}}$ in $\widetilde{\mathbf{W}}_s$ is equivalent to the norm $\|\mathbf{v}\|_V$. This way, we can identify $\mathbf{W}_s(\text{div}, \Omega)$ as $\widetilde{\mathbf{W}}_s$ and vice versa. As a consequence, $\widetilde{\mathbf{W}}_s$ is a closed subspace of $(L^s(\Omega))^{n+1}$, hence, it is a reflexive Banach space.

For dual and double dual spaces, we identify $F \in \mathbf{W}_s(\text{div}, \Omega)'$ as $\widetilde{F} = F \circ E^{-1} \in \widetilde{\mathbf{W}}_s'$, and identify $G \in \mathbf{W}_s(\text{div}, \Omega)''$ as $\widetilde{G} \in \widetilde{\mathbf{W}}_s''$ defined by $\widetilde{G}(\widetilde{F}) = G(\widetilde{F} \circ E)$ for any $\widetilde{F} \in \widetilde{\mathbf{W}}_s'$. Then $\widetilde{\mathbf{W}}_s$ being reflexive implies that $\mathbf{W}_s(\text{div}, \Omega)$ is a reflexive Banach space. \square

Proof of inequality (2.8). We call Scenario 1 the case when the origin is not on the line segment connecting x and y , and Scenario 2 otherwise. Consider Scenario 1 first. Let $\gamma(t) = tx + (1-t)y \neq 0$ for $t \in [0, 1]$ and $h_1(t) = |\gamma(t)|^p$. Then

$$\begin{aligned} ||x|^p - |y|^p| &= \left| \int_0^1 h_1'(t) dt \right| \leq \int_0^1 p|\gamma(t)|^{p-2} |\gamma(t) \cdot (x-y)| dt \leq p|x-y| \int_0^1 |\gamma(t)|^{p-1} dt \\ &\leq p|x-y| \int_0^1 2^{(p-2)^+} (t^{p-1}|x|^{p-1} + (1-t)^{p-1}|y|^{p-1}) dt \\ &= 2^{(p-2)^+} (|x|^{p-1} + |y|^{p-1})|x-y|, \end{aligned}$$

which yields (2.8). In Scenario 2, we have $y = -kx$ or $x = -ky$ for some number $k \in [0, \infty)$. Without loss of generality, assume the former situation. Then we have

$$||x|^p - |y|^p| = |x|^p|1 - k^p| \leq |x|^p(1 + k^p) \leq |x|^p(1 + k^{p-1})(1 + k) = (|x|^{p-1} + |y|^{p-1})|x-y|,$$

which shows (2.8) again. \square

Proof of Lemma 2.1. Consider $p > 0$ and $x, y \in \mathbb{R}^n$. We use the same named scenarios and the function $\gamma(t)$ as in the proof of inequality (2.8).

Proof of inequality (2.9). Consider Scenario 1 and define $h_2(t) = |\gamma(t)|^p \gamma(t)$ for $t \in [0, 1]$. Then

$$\begin{aligned} ||x|^p x - |y|^p y| &= \left| \int_0^1 h_2'(t) dt \right| = \left| \int_0^1 |\gamma(t)|^p (x-y) + p|\gamma(t)|^{p-2} (\gamma(t) \cdot (x-y)) \gamma(t) dt \right| \\ &\leq (1+p)|x-y| \int_0^1 |\gamma(t)|^p dt \leq (1+p)|x-y| \int_0^1 2^{(p-1)^+} (t^p|x|^p + (1-t)^p|y|^p) dt \\ &= 2^{(p-1)^+} (|x|^p + |y|^p)|x-y|. \end{aligned}$$

This proves (2.9). In Scenario 2, we can assume $y = -kx$ for some $k \geq 0$. We have

$$||x|^p x - |y|^p y| = |x|^{p+1}(1 + k^{p+1}) \leq |x|^{p+1}(1 + k^p)(1 + k) = (|x|^p + |y|^p)|x-y|.$$

Hence, we obtain (2.9).

Proofs of inequalities (2.10) and (2.11). Letting $z = x - y$, we have

$$\begin{aligned} (|x|^p x - |y|^p y) \cdot (x-y) &= \left(|x|^p \left(\frac{x+y}{2} + \frac{z}{2} \right) - |y|^p \left(\frac{x+y}{2} - \frac{z}{2} \right) \right) \cdot z \\ &= (|x|^p - |y|^p) \frac{x+y}{2} \cdot z + \frac{1}{2} (|x|^p + |y|^p) |z|^2 \\ &= \frac{1}{2} (|x|^p - |y|^p) (|x|^2 - |y|^2) + \frac{1}{2} (|x|^p + |y|^p) |z|^2. \end{aligned}$$

Since $(|x|^p - |y|^p)(|x|^2 - |y|^2) \geq 0$, we obtain (2.10). Using $(|x| + |y|)^p \geq 2^{-(p-1)^+} |x-y|^p$, we then deduce (2.11) from (2.10).

Now, consider $p \in (-1, 0)$.

Proof of inequality (2.12). Let $x, y \in \mathbb{R}$. The inequality obviously holds true when $x = 0$ or $y = 0$. Also, by switching the roles of x and y , we can assume $x > 0$ and $y \neq 0$.

If $y > 0$, then $||x|^p x - |y|^p y| = |x|^{1+p} - |y|^{1+p}$. Noting that $1+p \in (0, 1)$, we apply inequality (2.7) to have

$$||x|^p x - |y|^p y| \leq |x-y|^{1+p} \leq 2^{-p} |x-y|^{1+p}.$$

If $y < 0$, then $||x|^p x - |y|^p y| = |x|^{1+p} + |y|^{1+p}$. Applying Hölder's inequality to the dot product of two vectors $(|x|^{1+p}, |y|^{1+p})$ and $(1, 1)$ with powers $1/(1+p)$ and $-1/p$, we obtain

$$||x|^p x - |y|^p y| \leq (|x| + |y|)^{1+p} \cdot 2^{-p} = 2^{-p} |x-y|^{1+p},$$

which yields (2.12).

Proof of inequality (2.13). Let $x, y \in \mathbb{R}^n$. Consider Scenario 1 and define the function

$$h_3(t) = |\gamma(t)|^p \gamma(t) \cdot (x - y) \text{ for } t \in [0, 1].$$

Then

$$\begin{aligned} (|x|^p x - |y|^p y) \cdot (x - y) &= \int_0^1 h_3'(t) dt = \int_0^1 |\gamma(t)|^p |x - y|^2 + p |\gamma(t)|^{p-2} |\gamma(t) \cdot (x - y)|^2 dt \\ &\geq (1 + p) |x - y|^2 \int_0^1 |\gamma(t)|^p dt. \end{aligned}$$

Note that $-p \in (0, 1)$, hence $|\gamma(t)|^{-p} \leq (|x| + |y|)^{-p}$. Therefore, we obtain (2.13).

In Scenario 2, we can assume $y = -kx$ for some $k \geq 0$. We have

$$(|x|^p x - |y|^p y) \cdot (x - y) = |x|^{2+p} (1 + k^{1+p}) (1 + k).$$

Since $0 < 1 + p < 1$, we have from (2.6) that $1 + k^{1+p} \geq (1 + k)^{1+p}$. Hence,

$$(|x|^p x - |y|^p y) \cdot (x - y) \geq |x|^{2+p} (1 + k)^{2+p} = |x - y|^2 (|x| + |y|)^p,$$

which proves (2.13) again. \square

Proof of Lemma 2.3. Let $q \in L^r(\Omega)$. If $q = 0$ then (2.20) holds true. Consider $q \neq 0$. Denote by $W_0^{1,r}(\Omega)$ the space of functions in $W_0^{1,r}(\Omega)$ having zero trace on the boundary. Note that $|q|^{r-2} q \in L^s(\Omega)$. By the Browder–Minty Theorem, there exists a unique solution $w \in W_0^{1,s}(\Omega)$ of the problem

$$\int_{\Omega} |\nabla w|^{r-2} \nabla w \cdot \nabla v dx = \int_{\Omega} |q|^{r-2} q v dx \quad \text{for all } v \in W_0^{1,r}(\Omega). \quad (\text{A.1})$$

Choosing $v = w$ in (A.1) and applying the Hölder and Poincaré inequalities give

$$\|\nabla w\|_{0,r}^r = \int_{\Omega} |\nabla w|^r dx = \int_{\Omega} |q|^{r-2} q w dx \leq \|q\|_{0,r}^{r-1} \|w\|_{0,r} \leq C \|q\|_{0,r}^{r-1} \|\nabla w\|_{0,r},$$

where C is a positive constant. Hereafter, C denotes a generic positive constant. It follows that

$$\|\nabla w\|_{0,r} \leq C \|q\|_{0,r}.$$

Set $\mathbf{u} = -|\nabla w|^{r-2} \nabla w$. Then $\mathbf{u} \in \mathbf{L}^s(\Omega)$ and, by (A.1), $\operatorname{div} \mathbf{u} = |q|^{r-2} q \in L^s(\Omega)$. Therefore, $\mathbf{u} \in V \setminus \{0\}$. Observe that

$$\|\mathbf{u}\|_V^s = \|\mathbf{u}\|_{0,s}^s + \|\operatorname{div} \mathbf{u}\|_{0,s}^s = \|\nabla w\|_{0,r}^r + \|q\|_{0,r}^r \leq C \|q\|_{0,r}^r.$$

Thus, $\|\mathbf{u}\|_V \leq C \|q\|_{0,r}^{r-1}$. We then have

$$\int_{\Omega} (\operatorname{div} \mathbf{u}) q dx = \int_{\Omega} |q|^r dx = \|q\|_{0,r}^r = \|q\|_{0,r} \|q\|_{0,r}^{r-1} \geq C \|q\|_{0,r} \|\mathbf{u}\|_V.$$

Consequently, we obtain inequality (2.20). \square

REFERENCES

- [1] A. Z. Barak and J. Bear. Flow at high reynolds numbers through anisotropic porous media. *Advances in Water Resources*, 4(2):54–66, 1981.
- [2] J. Baranger and K. Najib. Numerical analysis of quasi-Newtonian flow obeying the power law or the carreau flow. *Numerische Mathematik*, 58(1):35–49, 1990.
- [3] J. Bear. *Dynamics of Fluids in Porous Media*. Dover Publications, 1988. Reprint of the American Elsevier Publishing Company, Inc., New York, 1972 edition.
- [4] E. Celik and L. Hoang. Generalized Forchheimer flows in heterogeneous porous media. *Nonlinearity*, 29(3):1124–1155, 2016.
- [5] E. Celik and L. Hoang. Maximum estimates for generalized Forchheimer flows in heterogeneous porous media. *J. Differential Equations*, 262(3):2158–2195, 2017.

- [6] E. Celik, L. Hoang, A. Ibragimov, and T. Kieu. Fluid flows of mixed regimes in porous media. *J. Math. Phys.*, 58(2):023102, 30 pp, 2017.
- [7] E. Celik, L. Hoang, and T. Kieu. Doubly nonlinear parabolic equations for a general class of Forchheimer gas flows in porous media. *Nonlinearity*, 31(8):3617–3650, 2018.
- [8] E. Celik, L. Hoang, and T. Kieu. Studying a doubly nonlinear model of slightly compressible Forchheimer flows in rotating porous media. *Turkish Journal of Mathematics*, 47(3):949–987, 2023.
- [9] S. S. Chow. Finite element error estimates for non-linear elliptic equations of monotone type. *Numerische Mathematik*, 54(4):373–393, 1989.
- [10] P. Fabrie. Regularity of the solution of Darcy-Forchheimer’s equation. *Nonlinear Anal.*, 13(9):1025–1049, 1989.
- [11] P. Forchheimer. Wasserbewegung durch Boden. *Zeit. Ver. Deut. Ing.*, 45:1781–1788, 1901.
- [12] P. Forchheimer. *Hydraulik*. Number Leipzig, Berlin, B. G. Teubner. 1930. 3rd edition.
- [13] R. Glowinski and A. Marroco. Sur l’approximation, par éléments finis d’ordre un, et la résolution, par pénalisation-dualité d’une classe de problèmes de Dirichlet non linéaires. *Revue française d’automatique, informatique, recherche opérationnelle. Analyse numérique*, 9(R2):41–76, 1975.
- [14] L. Hoang and A. Ibragimov. Qualitative study of generalized Forchheimer flows with the flux boundary condition. *Adv. Differential Equations*, 17(5–6):511–556, 2012.
- [15] L. Hoang, A. Ibragimov, T. Kieu, and Z. Sobol. Stability of solutions to generalized Forchheimer equations of any degree. *J. Math. Sci. (N.Y.)*, 210(4, Problems in mathematical analysis. No. 81 (Russian)):476–544, 2015.
- [16] L. Hoang and T. Kieu. Global estimates for generalized Forchheimer flows of slightly compressible fluids. *J. Anal. Math.*, 137(1):1–55, 2019.
- [17] L. T. Hoang, T. T. Kieu, and T. V. Phan. Properties of generalized Forchheimer flows in porous media. *J. Math. Sci.*, 202(2):259–332, 2014.
- [18] P. Knabner and G. Summ. Solvability of the mixed formulation for Darcy-Forchheimer flow in porous media. pages 1–26, 2016. preprint. DOI:10.48550/arXiv.1608.08829.
- [19] M. Muskat. *The flow of homogeneous fluids through porous media*. McGraw-Hill Book Company, inc., 1937.
- [20] D. A. Nield and A. Bejan. *Convection in porous media*. Springer-Verlag, New York, fourth edition, 2013.
- [21] Sandri, D. Sur l’approximation numérique des écoulements quasi-newtoniens dont la viscosité suit la loi puissance ou la loi de carreau. *ESAIM: M2AN*, 27(2):131–155, 1993.
- [22] E. Zeidler. *Nonlinear functional analysis and its applications. II/B*. Springer-Verlag, New York, 1990. Nonlinear monotone operators, Translated from the German by the author and Leo F. Boron.

¹DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, 1108 MEMORIAL CIRCLE, LUBBOCK, TX 79409–1042, U. S. A.

Email address: luan.hoang@ttu.edu

²DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH GEORGIA, GAINESVILLE CAMPUS, 3820 MUNDY MILL RD., OAKWOOD, GA 30566, U. S. A.

Email address: tinh.kieu@ung.edu