Fourier Transform of Anisotropic Hardy Spaces Associated with Ball Quasi-Banach Function Spaces and Its Applications to Hardy–Littlewood Inequalities

Chaoan Li, Xianjie Yan and Dachun Yang *

Abstract Let *A* be a general expansive matrix and *X* be a ball quasi-Banach function space on \mathbb{R}^n , whose certain power (namely its convexification) supports a Fefferman–Stein vectorvalued maximal inequality and the associate space of whose other power supports the boundedness of the powered Hardy–Littlewood maximal operator. Let $H_X^A(\mathbb{R}^n)$ be the anisotropic Hardy space associated with *A* and *X*. The authors first prove that the Fourier transform of $f \in H_X^A(\mathbb{R}^n)$ coincides with a continuous function *F* on \mathbb{R}^n in the sense of tempered distributions. Moreover, the authors obtain a pointwise inequality that the function *F* is less than the product of the anisotropic Hardy space norm of *f* and a step function with respect to the transpose matrix of the expansive matrix *A*. Applying this, the authors further induce a higher order convergence for the function *F* at the origin and give a variant of the Hardy–Littlewood inequality in $H_X^A(\mathbb{R}^n)$. All these results have a wide range of applications. Particularly, the authors apply these results, respectively, to classical (variable and mixed-norm) Lebesgue spaces, Morrey spaces, Lorentz spaces, Orlicz spaces, Orlicz-slice spaces, and local generalized Herz spaces and, even on the last five function spaces, the obtained results are completely new.

1 Introduction

In 1972, Fefferman and Stein [24] introduced a famous problem, that is, what is the characterization of the Fourier transform \widehat{f} of a distribution f from the classical Hardy space $H^p(\mathbb{R}^n)$. Recall that, in 1974, Coifman [19] characterized \hat{f} via the entire function of exponential type for $n = 1$, where $f \in H^p(\mathbb{R})$ with $p \in (0, 1]$. Since then, many researchers investigated the characterization of f with the distribution f from Hardy spaces with $n \geq 2$; see, for instance, [7, 22, 25, 56]. In particular, Taibleson and Weiss [56] proved that, for any given $p \in (0, 1]$, the Fourier transform of $f \in H^p(\mathbb{R}^n)$ coincides with a continuous function *F* in the sense of tempered distributions and

²⁰²⁰ *Mathematics Subject Classification*. Primary 42B10; Secondary 42B30, 42B35, 46E30.

Key words and phrases. expansive matrix, ball quasi-Banach function space, anisotropic Hardy space, Fourier transform, Hardy–Littlewood inequality.

This project is partially supported by the National Key Research and Development Program of China (Grant No. 2020YFA0712900), the National Natural Science Foundation of China (Grant Nos. 11971058 and 12071197), China Postdoctoral Science Foundation (Grant No. 2022M721024), and the Open Project Program of Key Laboratory of Mathematics and Complex System of Beijing Normal University (Grant No. K202304).

^{*}Corresponding author, E-mail: dcyang@bnu.edu.cn/July 13, 2023/Final version.

there exists a positive constant *C*, independent of *f* and *F*, such that, for any $x \in \mathbb{R}^n$,

(1.1)
$$
|F(x)| \leq C ||f||_{H^p(\mathbb{R}^n)} |x|^{n(\frac{1}{p}-1)}.
$$

This further implies the following generalization of the Hardy–Littlewood inequality that

(1.2)
$$
\left[\int_{\mathbb{R}^n} |x|^{n(p-2)} |F(x)|^p dx\right]^{1/p} \leq C \|f\|_{H^p(\mathbb{R}^n)},
$$

where *C* is a positive constant independent of *f* and *F* (see [55, p. 128]).

Recently, Sawano et al. [53] originally introduced the ball quasi-Banach function space *X* and the associated Hardy space $H_X(\mathbb{R}^n)$. In their article [53], by assuming that the Hardy–Littlewood maximal operator satisfies a Fefferman–Stein vector-valued inequality on certain power (namely its convexification) of *X* and the powered Hardy–Littlewood maximal operator is bounded on the associate space of certain power of *X*, Sawano et al. established various maximal function characterizations and several other characterizations of $H_X(\mathbb{R}^n)$, respectively, in terms of atoms, molecules, and Lusin area functions. Indeed, the real-variable theory of the Hardy space $H_X(\mathbb{R}^n)$ associated with the ball quasi-Banach function space *X* provides an unified framework of various types of Hardy spaces, which includes many important Hardy spaces that have been studied before, such as classical Hardy spaces, mixed-norm Hardy spaces, variable Hardy spaces, and Orlicz– Hardy spaces. For more recent developments on this topic, we refer the reader to [13, 28, 53, 57, 62, 66]. Based on the recent rapid developments of the theory of the Hardy space $H_X(\mathbb{R}^n)$ and the aforementioned works on the characterization of the Fourier transform of the classical Hardy space $H^p(\mathbb{R}^n)$ or their generalization, very recently, Huang et al. [36] showed that both (1.1) and (1.2) hold true in $H_X(\mathbb{R}^n)$.

On the other hand, in 2003, motivated by the application of discrete groups of dilations in wavelet theory, Bownik [4] introduced and investigated the anisotropic Hardy space H_A^p $_{A}^{p}(\mathbb{R}^{n})$ with $p \in (0, \infty)$, where *A* is a general expansive matrix on \mathbb{R}^n , which includes both the classical Hardy space and the parabolic Hardy space of Calder on and Torchinsky [12] as special cases. Since then, various variants of classical Hardy spaces over anisotropic Euclidean spaces (see, for instance, [5, 6, 15, 16, 17, 35, 38, 43, 44, 47]) or, more generally, over spaces of homogeneous type in the sense of Coifman and Weiss [20, 21] (see, for instance, [18, 26, 27, 29, 30]) have been introduced and their real-variable theories have been well developed. In 2013, Bownik and Wang [7] generalized inequalities (1.1) and (1.2) to the anisotropic Hardy space H_A^p $_{A}^{p}(\mathbb{R}^{n})$ with the known characterization of \overline{H}_A^p $_A^p(\mathbb{R}^n)$. Furthermore, Liu [44] pointed out that (1.1) and (1.2) also apply to the setting of the variable anisotropic Hardy space $H_A^{p(\cdot)}(\mathbb{R}^n)$. Later, Liu et al. [46] proved that (1.1) and (1.2) hold true for the anisotropic mixed-norm Hardy space $H_A^{\vec{p}}$ $A^{\vec{p}}(\mathbb{R}^n)$. Recall that the anisotropic Hardy space $H_X^A(\mathbb{R}^n)$ associated with both *A* and *X* was first introduced and studied by Wang et al. [59], in which Wang et al. characterized $H_X^A(\mathbb{R}^n)$ in terms of maximal functions, atoms, finite atoms, and molecules. Moreover, the variable anisotropic Hardy space $H_A^{p(\cdot)}(\mathbb{R}^n)$ and the anisotropic mixednorm Hardy space $H_A^{\vec{p}}$ $\overrightarrow{P}_A(\mathbb{R}^n)$ are the special cases of $H_X^A(\mathbb{R}^n)$ with $X := L^{p(\cdot)}(\mathbb{R}^n)$ or $X := L^{\overrightarrow{p}}(\mathbb{R}^n)$, respectively. Based on these results, it is natural to ask whether (1.1) and (1.2) also hold true for $H_X^A(\mathbb{R}^n)$. The goal of this article is to give a positive answer to this question.

Let *A* be a dilation and *X* a ball quasi-Banach function space on \mathbb{R}^n . In this article, under the assumptions that the Hardy–Littlewood maximal operator satisfies some Fefferman–Stein vectorvalued inequality on certain power of *X*, the powered Hardy–Littlewood maximal operator is bounded on the associate space of certain power of *X*, and the *X*-quasi-norm of characteristic functions of anisotropic balls has a lower bound, we get rid of the dependence on the concavity of $\|\cdot\|_X$. With these mild assumptions and two uniform pointwise estimates we show that the Fourier transform \widehat{f} of $f \in H_X^A(\mathbb{R}^n)$ coincides with a continuous function F on \mathbb{R}^n in the sense of tempered distributions and prove that an inequality similar to (1.1) also holds true for any $f \in H_X^A(\mathbb{R}^n)$. Furthermore, applying this and a technical inequality about the value of the Fourier transform of atoms, we further conclude a higher order convergence of the continuous function *F* at the origin and then show that an inequality similar to (1.2) holds true for $H_X^A(\mathbb{R}^n)$, which is a variant of the Hardy–Littlewood inequality in $H_X^A(\mathbb{R}^n)$. It is remarkable that the results obtained in this article have a wide range of generality because ball quasi-Banach function spaces include lots of important function spaces. In particular, when these results are applied to classical Lebesgue spaces, variable Lebesgue spaces, and mixed-norm Lebesgue spaces, the obtained conclusions coincide with the known ones; morever, when these results are applied, respectively, to Morrey spaces, Lorentz spaces, Orlicz spaces, Orlicz-slice spaces, and local generalized Herz spaces, the obtained conclusions are completely new. More applications of these results to new-found function spaces are quite possible.

The remainder of this article is organized as follows.

In Sect. 2, we first present definitions of expansive matrices, ball quasi-Banach function spaces, and anisotropic Hardy spaces associated with both *A* and *X*; see Definitions 2.1, 2.4, and 2.10 below.

The aim of Sect. 3 is to prove the main result (see Theorem 3.1 below), that is, the Fourier transform \widehat{f} of $f \in H_X^A(\mathbb{R}^n)$ coincides with a continuous function *F* in the sense of tempered distributions. In order to achieve this, we apply Lemmas 3.5 (some subtle estimates on derivatives of the Fourier transform of the dilation of atoms) and 3.7 (some exquisite relations between the Euclidean norm and the step homogeneous quasi-norm ρ under consideration) to establish a uniform pointwise estimate for atoms (see Lemma 3.6 below). Then Theorem 3.1 is proved by this and some real-variable characterizations from [59], especially its atomic decompositions. Morever, we apply these results, respectively, to classical Lebesgue spaces, variable Lebesgue spaces, and mixed-norm Lebesgue spaces and show that the obtained conclusions coincide with the known ones [see Remark 3.2(i)-(iv) below]. At the same time, we obtain a pointwise inequality of the continuous function *F*, which suggests that the anisotropic mixed-norm atoms must possess vanishing moments in some sense [see Remark 3.2(v) below].

Applying the Fourier transform, in Sect. 4, we present some further applications of Theorem 3.1. First, we prove that the above function F has a higher order convergence at the origin (see Theorem 4.1 below). Second, we show that the term

$$
|F(\cdot)|\min\left\{[\rho_*(\cdot)]^{1-\frac{1}{p_-}-\frac{1}{q_0}+(d+1)\frac{\ln\lambda_-}{\ln b}}, [\rho_*(\cdot)]^{1-\frac{2}{q_0}+(d+1)\frac{\ln\lambda_-}{\ln b}}\right\}
$$

is $L^{q_0}(\mathbb{R}^n)$ -integrable and a positive constant multiple of the anisotropic Hardy space norm of f can uniformly controll this integral. Thus, we extend the Hardy–Littlewood inequality to the setting of anisotropic Hardy spaces associated with ball quasi-Banach function spaces (see Theorem 4.2 below). Further, we apply these results, respectively, to classical Lebesgue spaces, variable Lebesgue spaces, and mixed-norm Lebesgue spaces and prove that the obtained conclusions coincide with the known ones (see Remark 4.3 below).

As applications, in Sect. 5, we apply Theorems 3.1, 4.1, and 4.2, via verfiying all the necessary assumptions, to five concrete examples of ball quasi-Banach function spaces, namely Morrey spaces (see Subsection 5.1 below), Lorentz spaces (see Subsection 5.2 below), Orlicz spaces (see Subsection 5.3 below), Orlicz-slice spaces (see Subsection 5.4 below), and local Herz–Hardy spaces (see Subsection 5.5 below). In particular, we show that the anisotropic local Herz space is a quasi-banach function space (see Theorem 5.16 below). Through both a boundedness criterion of sublinear operators on anisotropic local generalized Herz spaces and its simple corollary (see Lemma 5.18 and Corollary 5.20 below), we prove that the anisotropic local generalized Herz space supports a Fefferman–Stein vector-valued maximal inequality and the associate space of whose other power supports the boundedness of the powered Hardy–Littlewood maximal operator (see Lemma 5.17 and Theorem 5.21 below).

At the end of this section, we make some conventions on notation. Let $\mathbb{N} := \{1, 2, ...\}$, $\mathbb{Z}_+ :=$ $\mathbb{N} \cup \{0\}$, $\mathbb{Z}_{+}^{n} := (\mathbb{Z}_{+})^{n}$, and 0 be the *origin* of \mathbb{R}^{n} . For any multi-index $\alpha := (\alpha_{1}, \dots, \alpha_{n}) \in \mathbb{Z}_{+}^{n}$ and any $x := (x_1, \ldots, x_n) \in \mathbb{R}^n$, let $|\alpha| := \alpha_1 + \cdots + \alpha_n$, $\partial^{\alpha} := (\frac{\partial}{\partial x_1}, \ldots, x_n)$ $\frac{\partial}{\partial x_1}$ ^{ou}₁ ... ($\frac{\partial}{\partial x_1}$ $\frac{\partial}{\partial x_n}$ ^{2*a*}, and $x^\alpha := x_1^{\alpha_1}$ $\frac{\alpha_1}{1} \cdots \frac{\alpha_n}{n}$. We denote by *C* a *positive constant* which is independent of the main parameters involved, but may vary from line to line. We use $C_{(\alpha,\dots)}$ to denote a positive constant depending on the indicated parameters α , The symbol $f \leq g$ means $f \leq Cg$. If $f \leq g$ and $g \leq f$, we then write $f \sim g$. If $f \leq Cg$ and $g = h$ or $g \leq h$, we then write $f \leq g = h$ or $f \leq g \leq h$. For any $q \in [1, \infty]$, we denote by *q'* its *conjugate index*, that is, $1/q + 1/q' = 1$. For any $x \in \mathbb{R}^n$, we denote by |*x*| the *n*-dimensional *Euclidean metric* of *x*. If *E* is a subset of \mathbb{R}^n , we denote by $\mathbf{1}_E$ its *characteristic function* and by E^C the set $\mathbb{R}^n \setminus E$. For any $r \in (0, \infty)$ and $x \in \mathbb{R}^n$, we denote by $B(x, r)$ the ball centered at *x* with the radius *r*, that is, $B(x, r) := \{y \in \mathbb{R}^n : |x - y| < r\}$. For any ball *B*, we use x_B to denote its center and r_B its radius and we denote by λB for any $\lambda \in (0, \infty)$ the ball concentric with *B* having the radius λr_B . We also use $\epsilon \to 0^+$ to denote $\epsilon \in (0, \infty)$ and $\epsilon \to 0$. Let *X* and *Y* be two normed vector spaces, respectively, with the norm $\| \cdot \|_X$ and the norm $\| \cdot \|_Y$; then we use $X \hookrightarrow Y$ to denote $X \subset Y$ and there exists a positive constant *C* such that, for any $f \in X$, $||f||_Y \leq C||f||_X$. At last, when we prove a theorem or the like, we always use the same symbols in the wanted proved theorem or the like.

2 Preliminaries

In this section, we first recall some symbols and concepts on dilations (see, for instance, [4, 53]) as well as ball quasi-Banach function spaces (see, for instance, [53, 57, 58, 62, 65]). We begin by recalling the concept of expansive matrices from [4].

Definition 2.1. A real $n \times n$ matrix *A* is called an *expansive matrix* (shortly, a *dilation*) if

$$
\min_{\lambda \in \sigma(A)} |\lambda| > 1,
$$

here and hereafter, σ(*A*) denotes the *set of all eigenvalues of A*.

Fourier Transform of Anisotropic Hardy Spaces 5 5 5 5 5 6 1 3 April 1988 6 6 1 3 April 19

Let *A* be a dilation and

$$
(2.1) \t\t b := |\det A|,
$$

where det *A* denotes the determinant of *A*. Then it follows from [4, p. 6, (2.7)] that $b \in (1, \infty)$. By the fact that there exists an open and symmetry ellipsoid Δ , with $|\Delta| = 1$, and an $r \in (1, \infty)$ such that $\Delta \subset r\Delta \subset A\Delta$ (see [4, p. 5, Lemma 2.2]), we find that, for any $k \in \mathbb{Z}$,

$$
(2.2) \t\t B_k := A^k \Delta
$$

is open, $B_k \subset rB_k \subset B_{k+1}$, and $|B_k| = b^k$. For any $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}$, an ellipsoid $x + B_k$ is called a *dilated ball*. In what follows, we always let B be the set of all such dilated balls, that is,

(2.3)
$$
\mathcal{B} := \{x + B_k : x \in \mathbb{R}^n, k \in \mathbb{Z}\}\
$$

and let

$$
\tau := \inf \left\{ l \in \mathbb{Z} : r^l \geq 2 \right\}.
$$

Let $\lambda_-, \lambda_+ \in (0, \infty)$ satisfy that

$$
1 < \lambda_- < \min\{|\lambda| : \lambda \in \sigma(A)\} \le \max\{|\lambda| : \lambda \in \sigma(A)\} < \lambda_+.
$$

We point out that, if A is diagonalizable over \mathbb{R} , then we may let

 $\lambda_- := \min\{|\lambda| : \lambda \in \sigma(A)\}\$ and $\lambda_+ := \max\{|\lambda| : \lambda \in \sigma(A)\}.$

Otherwise, we may choose them sufficiently close to these equalities in accordance with what we need in our arguments.

The following definition of the homogeneous quasi-norm is just [4, p. 6, Definition 2.3].

Definition 2.2. A *homogeneous quasi-norm*, associated with a dilation *A*, is a measurable mapping $\varrho: \mathbb{R}^n \to [0, \infty)$ such that

- (i) $\varrho(x) = 0 \Longleftrightarrow x = 0$, where 0 denotes the origin of \mathbb{R}^n ;
- (ii) $\varrho(Ax) = b\varrho(x)$ for any $x \in \mathbb{R}^n$;
- (iii) there exists an $A_0 \in [1, \infty)$ such that, for any $x, y \in \mathbb{R}^n$,

$$
\varrho(x+y) \le A_0 \left[\varrho(x) + \varrho(y) \right].
$$

In the standard Euclidean space case, let $A := 2 I_{n \times n}$ and, for any $x \in \mathbb{R}^n$, $\varrho(x) := |x|^n$. Then ϱ is an example of homogeneous quasi-norms associated with *A* on \mathbb{R}^n . Here and thereafter, $I_{n \times n}$ always denotes the $n \times n$ *unit matrix* and $|\cdot|$ the *Euclidean norm* in \mathbb{R}^n .

For a fixed dilation *A*, by [4, p. 6, Lemma 2.4], we introduce the following quasi-norm which is used throughout this article.

Definition 2.3. Define the *step homogeneous quasi-norm* ρ on \mathbb{R}^n , associated with the dilation A, by setting

$$
\rho(x) := \begin{cases} b^k & \text{if } x \in B_{k+1} \setminus B_k, \\ 0 & \text{if } x = \mathbf{0}, \end{cases}
$$

where *b* is the same as in (2.1) and, for any $k \in \mathbb{Z}$, B_k the same as in (2.2).

Then (\mathbb{R}^n, ρ, dx) is a space of homogeneous type in the sense of Coifman and Weiss [20], where *dx* denotes the *n*-dimensional Lebesgue measure. For more studies on the real-variable theory of function spaces over spaces of homogeneous type, we refer the reader to [8, 9, 10, 11, 39, 40, 41, 60, 61, 63].

Throughout this article, we always let *A* be a dilation in Definition 2.1, *b* the same as in (2.1), ρ the step homogeneous quasi-norm in Definition 2.3, \mathcal{B} the set of all dilated balls in (2.3), $\mathcal{M}(\mathbb{R}^n)$ the *set of all measurable functions* on \mathbb{R}^n and, for any $k \in \mathbb{Z}$, B_k the same as in (2.2). Now, we recall the definition of ball quasi-norm Banach function spaces (see [53]).

Definition 2.4. A quasi-normed linear space $X \subset \mathcal{M}(\mathbb{R}^n)$, equipped with a quasi-norm $\|\cdot\|$ which makes sense for the whole $\mathcal{M}(\mathbb{R}^n)$, is called a *ball quasi-Banach function space* if it satisfies

- (i) for any $f \in \mathcal{M}(\mathbb{R}^n)$, $||f||_X = 0$ implies that $f = 0$ almost everywhere;
- (ii) for any $f, g \in \mathcal{M}(\mathbb{R}^n)$, $|g| \le |f|$ almost everywhere implies that $||g||_X \le ||f||_X$;
- (iii) for any $\{f_m\}_{m\in\mathbb{N}} \subset \mathcal{M}(\mathbb{R}^n)$ and $f \in \mathcal{M}(\mathbb{R}^n)$, $0 \le f_m \uparrow f$ as $m \to \infty$ almost everywhere implies that $||f_m||_X \uparrow ||f||_X$ as $m \to \infty$;
- (iv) $\mathbf{1}_B \in X$ for any dilated ball $B \in \mathcal{B}$.

Moreover, a ball quasi-Banach function space *X* is called a *ball Banach function space* if it satisfies:

- (v) for any $f, g \in X$, $||f + g||_X \le ||f||_X + ||g||_X$;
- (vi) for any given dilated ball *B* ∈ *B*, there exists a positive constant $C_{(B)}$ such that, for any $f \in X$, $f \in X$,

$$
\int_B |f(x)| dx \le C_{(B)} ||f||_X.
$$

Now, we recall the concept of the *p*-convexification of ball quasi-Banach function spaces, which is a part of [53, Definition 2.6].

Definition 2.5. Let *X* be a ball quasi-Banach function space and $p \in (0, \infty)$. The *p-convexification* X^p of *X* is defined by setting

$$
X^p := \{ f \in \mathcal{M}(\mathbb{R}^n) : |f|^p \in X \}
$$

equipped with the *quasi-norm* $||f||_{X^p} := |||f|^p||_X^{1/p}$ $\int_X^{1/p}$ for any $f \in X^p$.

The associate space *X* ′ of any given ball Banach function space *X* is defined as follows; see [3, Chapter 1, Section 2] or [53, p. 9].

Definition 2.6. For any given ball Banach function space *X*, its *associate space* (also called the *Köthe dual space*) X' is defined by setting

$$
X':=\left\{f\in \mathscr{M}(\mathbb{R}^n):\; \|f\|_{X'}:=\sup_{g\in X,\; \|g\|_X=1}\|fg\|_{L^1(\mathbb{R}^n)}<\infty\right\},
$$

where $\|\cdot\|_{X'}$ is called the *associate norm* of $\|\cdot\|_{X}$.

Now, we recall the concept of the Hardy–Littlewood maximal operator. In what follows, for any given $p \in (0, \infty]$ and any given subset $E \subset \mathbb{R}^n$, let $L^p_{loc}(E)$ denote the *set of all p-order locally integrable functions* on *E*. Recall that the *Hardy–Littlewood maximal operator* M(*f*) of $f \in L^1_{loc}(\mathbb{R}^n)$ is defined by setting, for any $x \in \mathbb{R}^n$,

(2.5)
$$
\mathcal{M}(f)(x) := \sup_{k \in \mathbb{Z}} \sup_{y \in x + B_k} \frac{1}{|B_k|} \int_{y + B_k} |f(z)| dz = \sup_{x \in B \in \mathcal{B}} \frac{1}{|B|} \int_B |f(z)| dz,
$$

where $\mathcal B$ is the same as in (2.3) and the supremum in the second equality is taken over all the balls $B \in \mathcal{B}$. For any given $\alpha \in (0, \infty)$, the *powered Hardy–Littlewood maximal operator* $\mathcal{M}^{(\alpha)}$ is defined by setting, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$
\mathcal{M}^{(\alpha)}(f)(x) := {\mathcal{M}(|f|^{\alpha})(x)}^{\frac{1}{\alpha}}.
$$

Throughout this article, we also need the following fundamental assumptions about the boundedness of M on the convexification of the given ball quasi-Banach function space and the boundedness of certain powered of M on the associate space of its convexification.

Assumption 2.7. *Let X be a ball quasi-Banach function space. Assume that there exists a p*[−] ∈ (0, ∞) *such that, for any p* ∈ (0, *p*−) *and u* ∈ (1, ∞)*, there exists a positive constant C, depending on both p and r, such that, for any* $\{f_k\}_{k=1}^{\infty} \subset \mathcal{M}(\mathbb{R}^n)$,

$$
\left\| \left\{ \sum_{k=1}^{\infty} \left[\mathcal{M}(f_k) \right]^u \right\}^{\frac{1}{u}} \right\|_{X^{\frac{1}{p}}} \leq C \left\| \left\{ \sum_{k=1}^{\infty} |f_k|^u \right\}^{\frac{1}{u}} \right\|_{X^{\frac{1}{p}}}.
$$

In what follows, for any given $p_$ ∈ (0, ∞), let

$$
\underline{p} := \min\{p_-, 1\}.
$$

Assumption 2.8. *Let p*[−] ∈ (0, ∞) *and X be a ball quasi-Banach function space. Assume that there exists a* $\theta_0 \in (0, \underline{p})$, with <u>*p*</u> *in* (2.6)*, and a* $p_0 \in (\theta_0, \infty)$ *such that* X^{1/θ_0} *is a ball Banach function space and, for any* $f \in (X^{1/\theta_0})'$,

$$
\left\|{\cal M}^{((p_0/\theta_0)')}(f)\right\|_{(X^{1/\theta_0})'}\le C\|f\|_{(X^{1/\theta_0})'},
$$

where C is a positive constant, independent of f, and $\frac{1}{p_0/\theta_0} + \frac{1}{(p_0/\theta_0)'} = 1$.

Next, recall that a *Schwartz function* is a function $\varphi \in C^{\infty}(\mathbb{R}^n)$ satisfying that, for any $k \in \mathbb{Z}_+$ and any multi-index $\alpha \in \mathbb{Z}_+^n$,

(2.7)
$$
\|\varphi\|_{\alpha,k} := \sup_{x \in \mathbb{R}^n} [\rho(x)]^k |\partial^{\alpha} \varphi(x)| < \infty.
$$

Denote by $S(\mathbb{R}^n)$ the *set of all Schwartz functions*, equipped with the topology determined by $\{\|\cdot\|_{\alpha,k}\}_{\alpha\in\mathbb{Z}_{+}^{n},k\in\mathbb{Z}_{+}}$. Then $\mathcal{S}'(\mathbb{R}^{n})$ is defined to be the *dual space* of $\mathcal{S}(\mathbb{R}^{n})$, equipped with the weak-* topology. For any $N \in \mathbb{Z}_+$, let

$$
\mathcal{S}_N(\mathbb{R}^n) := \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n) : \ \|\varphi\|_{\alpha,k} \leq 1, |\alpha| \leq N, k \leq N \right\},\
$$

equivalently,

$$
\varphi \in \mathcal{S}_N(\mathbb{R}^n) \iff ||\varphi||_{\mathcal{S}_N(\mathbb{R}^n)} := \sup_{|\alpha| \le N} \sup_{x \in \mathbb{R}^n} \max\{1, [\rho(x)]^N\} |\partial^{\alpha} \varphi(x)| \le 1.
$$

Now, we recall the definitions of the anisotropic non-tangential maximal function and the anisotropic non-tangential grand maximal function from [59, Definition 2.15]. In what follows, for any $\varphi \in \mathcal{S}(\mathbb{R}^n)$ and $k \in \mathbb{Z}$, let $\varphi_k(\cdot) := b^{-k} \varphi(A^{-k} \cdot)$.

Definition 2.9. Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$ and $f \in \mathcal{S}'(\mathbb{R}^n)$. The *anisotropic non-tangential maximal function* $M_{\varphi}(f)$, with respect to φ , is defined by setting, for any $x \in \mathbb{R}^n$,

$$
M_{\varphi}(f)(x) := \sup_{k \in \mathbb{Z}, y \in x + B_k} |f * \varphi_k(y)|.
$$

Moreover, for any given $N \in \mathbb{N}$, the *anisotropic non-tangential grand maximal function* $M_N(f)$ is defined by setting, for any $x \in \mathbb{R}^n$,

(2.8)
$$
M_N(f)(x) := \sup_{\varphi \in S_N(\mathbb{R}^n)} M_{\varphi}(f)(x).
$$

We present the definition of $H_X^A(\mathbb{R}^n)$ from [59] as follows. In what follows, for any $\alpha \in \mathbb{R}$, we denote by the *symbol* $\lfloor \alpha \rfloor$ the largest integer not greater than α .

Definition 2.10. Let *A* be a dilation and *X* a ball quasi-Banach function space satisfying both Assumption 2.7 with $p_$ ∈ (0, ∞) and Assumption 2.8 with the same $p_$, θ_0 ∈ (0, p), and p_0 ∈ (θ_0, ∞) , where *p* is the same as in (2.6). Assume that

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{\theta_0} - 1 \right) \frac{\ln b}{\ln(\lambda_-)} \right| + 2, \infty \right).
$$

The *anisotropic Hardy space* $H^A_{X,N}(\mathbb{R}^n)$, associated with both *A* and *X*, is defined by setting

$$
H_{X,N}^A(\mathbb{R}^n) := \{ f \in \mathcal{S}'(\mathbb{R}^n) : ||M_N(f)||_X < \infty \},\
$$

where $M_N(f)$ is the same as in (2.8). Moreover, for any $f \in H^A_{X,N}(\mathbb{R}^n)$, let

$$
||f||_{H^A_{X,N}(\mathbb{R}^n)} := ||M_N(f)||_X.
$$

Let *A* be a dilation and *X* the same as in Definition 2.10. In the remainder of this article, we always let

$$
N_{X,A} := \left\lfloor \left(\frac{1}{\theta_0} - 1\right) \frac{\ln b}{\ln(\lambda_-)} \right\rfloor + 2.
$$

- **Remark 2.11.** (i) As was mentioned in [59, Remark 2.17(i)], the space $H^A_{X,N}(\mathbb{R}^n)$ is independent of the choice of *N* as long as $N \in \mathbb{N} \cap [N_{X, A}, \infty)$. Thus, in what follows, when $N \in \mathbb{N} \cap [N_{X,A}, \infty)$, we always write $H^A_{X,N}(\mathbb{R}^n)$ simply by $H^A_X(\mathbb{R}^n)$.
	- (ii) If $A := 2I_{n \times n}$, then $H_X^A(\mathbb{R}^n)$ coincides with $H_X(\mathbb{R}^n)$ which was introduced by Sawano et al. in [53].

3 Fourier Transforms of *H A* $^A_X(\mathbb{R}^n)$

Let *A* be a dilation, *X* a ball quasi-Banach function space satisfying some mild assumptions, and $f \in H_X^A(\mathbb{R}^n)$. In this section, we aim to study the Fourier transform of *f*. Recall that, for any $\varphi \in \mathcal{S}(\mathbb{R}^n)$, its *Fourier transform*, denoted by $\mathscr{F}(\varphi)$ or $\widehat{\varphi}$, is defined by setting, for any $\xi \in \mathbb{R}^n$,

$$
\mathscr{F}(\varphi)(\xi)=\widehat{\varphi}(\xi):=\int_{\mathbb{R}^n}\varphi(x)e^{-2\pi\iota x\cdot\xi}\,dx,
$$

here and thereafter, $\iota := \sqrt{-1}$ and, for any $x := (x_1, \ldots, x_n), \xi := (\xi_1, \ldots, \xi_n) \in \mathbb{R}^n, x \cdot \xi := \sum_{i=1}^n x_i \xi_i$. For any $f \in S'(\mathbb{R}^n)$, \widehat{f} is defined by setting, for any $\varphi \in S(\mathbb{R}^n)$, $\langle \widehat{f}, \varphi \rangle := \langle f, \widehat{\varphi} \rangle$; also, for any *f* ∈ $S(\mathbb{R}^n)$ [resp. $S'(\mathbb{R}^n)$], f^{\vee} denotes its *inverse Fourier transform* which is defined by setting, for any $\xi \in \mathbb{R}^n$, $f^{\vee}(\xi) := \widehat{f}(-\xi)$ [resp., for any $\varphi \in \mathcal{S}(\mathbb{R}^n)$, $\langle f^{\vee}, \varphi \rangle := \langle f, \varphi^{\vee} \rangle$].

Now, we present the main result of this section as follows.

Theorem 3.1. *Let A be a dilation and X a ball quasi-Banach function space satisfying both Assumption 2.7 with p*_− ∈ (0, ∞) *and Assumption 2.8 with the same p*_−*,* θ_0 ∈ (0, *p*)*, and p*₀ ∈ (θ_0 , ∞)*, where p is the same as in* (2.6)*. Further assume that there exists a* $q_0 \in [\theta_0, 1]$ *such that:*

(i) *for any non-negative measurable functions* $\{f_k\}_{k=1}^{\infty}$,

(3.1)
$$
\sum_{k=1}^{\infty} ||f_k||_{X^{\frac{1}{q_0}}} \lesssim \left\| \sum_{k=1}^{\infty} f_k \right\|_{X^{\frac{1}{q_0}}},
$$

where the implicit positive constant is independent of $\{f_k\}_{k=1}^{\infty}$;

(ii) *for any* $B \in \mathcal{B}$ *with* \mathcal{B} *in* (2.3)*,*

(3.2)
$$
\|1_B\|_X \gtrsim \min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{\theta_0}}\right\},\,
$$

where the implicit positive constant is independent of B.

Then, for any $f \in H_X^A(\mathbb{R}^n)$ *, there exists a continuous function F on* \mathbb{R}^n such that

$$
\widehat{f} = F \quad in \quad \mathcal{S}'(\mathbb{R}^n)
$$

and there exists a positive constant C, depending only on A and X, such that, for any $x \in \mathbb{R}^n$,

(3.4)
$$
|F(x)| \leq C||f||_{H_X^A(\mathbb{R}^n)} \max\left\{[\rho_*(x)]^{\frac{1}{q_0}-1}, [\rho_*(x)]^{\frac{1}{\theta_0}-1}\right\},
$$

here and thereafter, $ρ_*$ *is defined as in Definition 2.3 with A replaced by its transposed matrix A[∗].*

Remark 3.2. (i) If $A := 2I_{n \times n}$, then Theorem 3.1 was obtained in [36, Theorem 2.1].

(ii) For any given measurable set $E \subset \mathbb{R}^n$ and any given $p \in (0, \infty)$, the *Lebesgue space* $L^p(E)$ is defined by setting,

(3.5)
$$
L^{p}(E) := \left\{ f \text{ is measurable on } E: ||f||_{L^{p}(E)} := \left[\int_{E} |f(x)|^{p} dx \right]^{1/p} < \infty \right\}.
$$

Let *A* be a dilation, $p \in (0, 1)$, and

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{p} - 1 \right) \frac{\ln b}{\ln(\lambda_-)} \right| + 2, \infty \right).
$$

Then, by [61, Remarks 2.7(i) and 4.21(i)], we conclude that $L^p(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10 with $X := L^p(\mathbb{R}^n)$, $p_- \in (0, p]$, $\theta_0 \in (0, p_-)$, and $p_0 \in (p, \infty)$. Moreover, choose $q_0 \in (p, 1]$. Then it follows from (3.5) that, for any non-negative measurable functions $\{f_k\}_{k=1}^{\infty}$ and any $B \in \mathcal{B}$,

$$
\sum_{k=1}^{\infty} \left\| f_k \right\|_{L^{\frac{p}{q_0}}(\mathbb{R}^n)} \le \left\| \sum_{k=1}^{\infty} f_k \right\|_{L^{\frac{p}{q_0}}(\mathbb{R}^n)}
$$

and

$$
\|{\bf 1}_B\|_{L^p(\mathbb{R}^n)}=|B|^{\frac{1}{p}}>\min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{q_0}}\right\}.
$$

Thus, $L^p(\mathbb{R}^n)$ satisfies all the assumptions of Theorem 3.1 with $X := L^p(\mathbb{R}^n)$. In this case, Theorem 3.1 was obtained in [7, Theorem 1].

(iii) Recall that $P(\mathbb{R}^n)$ is defined to be the *set of all the measurable functions* $p(\cdot)$ on \mathbb{R}^n satisfying

$$
0 < \widetilde{p_-} := \operatorname*{ess\,inf}_{x \in \mathbb{R}^n} p(x) \le \operatorname*{ess\,sup}_{x \in \mathbb{R}^n} p(x) =: \widetilde{p_+} < \infty.
$$

For any $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$, the *variable Lebesgue space* $L^{p(\cdot)}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$
\int_{\mathbb{R}^n} |f(x)|^{p(x)} dx < \infty,
$$

equipped with the *quasi-norm* $||f||_{L^{p(\cdot)}(\mathbb{R}^n)}$ defined by setting

$$
(3.6) \t\t\t\t||f||_{L^{p(\cdot)}(\mathbb{R}^n)} := \inf \left\{\lambda \in (0, \infty): \int_{\mathbb{R}^n} \left[\frac{|f(x)|}{\lambda}\right]^{p(x)} dx \le 1\right\}.
$$

Denote by $C^{\log}(\mathbb{R}^n)$ the set of all the functions $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfying the *globally log*-*Hölder continuous condition*, that is, there exist $C_{\log}(p)$, $C_{\infty} \in (0, \infty)$ and $p_{\infty} \in \mathbb{R}$ such that, for any $x, y \in \mathbb{R}^n$,

$$
|p(x) - p(y)| \le \frac{C_{\log}(p)}{\log (e + 1/|x - y|)}
$$

and

$$
|p(x) - p_{\infty}| \leq \frac{C_{\infty}}{\log(e + |x|)}.
$$

Let *A* be a dilation and $p(\cdot) \in C^{\log}(\mathbb{R}^n)$ satisfy $0 < \widetilde{p_-} \le \widetilde{p_+} < 1$ and

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{\widetilde{p_{-}}} - 1 \right) \frac{\ln b}{\ln(\lambda_{-})} \right| + 2, \infty \right).
$$

Then, by [61, Remarks 2.7(iv) and 4.21(v)], we conclude that $L^{p(\cdot)}(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10 with $X := L^{p(\cdot)}(\mathbb{R}^n)$, $p_-\in (0,\widetilde{p_-}], \theta_0 \in (0,p_-)$, and $p_0 \in (0,\widetilde{p_-}],$ $(\widetilde{p_+}, \infty)$. Moreover, choose $q_0 \in (\widetilde{p_+}, 1]$. On the one hand, from [64, Remark 2.1(iv)], we deduce that, for any non-negative measurable functions $\{f_k\}_{k=1}^{\infty}$,

$$
\sum_{k=1}^\infty\|f_k\|_{L^{\frac{p(\cdot)}{q_0}}(\mathbb{R}^n)}\leq \left\|\sum_{k=1}^\infty f_k\right\|_{L^{\frac{p(\cdot)}{q_0}}(\mathbb{R}^n)}.
$$

On the other hand, by (3.6), we find that, for any $B \in \mathcal{B}$,

$$
\|{\bf 1}_B\|_{L^{p(\cdot)}({\mathbb R}^n)}\gtrsim \min\left\{|B|^{\frac{1}{\widetilde{p_+}}},|B|^{\frac{1}{\widetilde{p_-}}}\right\}> \min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{\theta_0}}\right\}.
$$

Thus, $L^{p(\cdot)}(\mathbb{R}^n)$ satisfies all the assumptions of Theorem 3.1 with $X := L^{p(\cdot)}(\mathbb{R}^n)$. In this case, Theorem 3.1 was obtained in [45, Theorem 1].

(iv) Let $\vec{p} := (p_1, \ldots, p_n) \in (0, \infty]^n$. Recall that the *mixed-norm Lebesgue space* $L^{\vec{p}}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$
(3.7) \qquad ||f||_{L^{\vec{p}}(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}} \cdots \left[\int_{\mathbb{R}} \left\{ \int_{\mathbb{R}} |f(x_1, \ldots, x_n)|^{p_1} dx_1 \right\}^{\frac{p_2}{p_1}} dx_2 \right\}^{\frac{p_3}{p_2}} \cdots dx_n \right\}^{\frac{1}{p_n}}
$$

is finite with the usual modifications made when $p_i = \infty$ for some $i \in \{1, ..., n\}$. Let \vec{p} ∈ (0, 1)^{*n*}, $\widehat{p_{-}}$:= min{*p*₁, ..., *p_n*}, $\widehat{p_{+}}$:= max{*p*₁, ..., *p_n*}, and

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{\widehat{p_{-}}} - 1 \right) \frac{\ln b}{\ln(\lambda_{-})} \right| + 2, \infty \right).
$$

Then, by both [65, p. 2047] and [65, Lemmas 7.22 and 7.26], we conclude that $L^{\vec{p}}(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10 with $X := L^{\vec{p}}(\mathbb{R}^n)$, $p_{-} \in (0, \widehat{p_{-}}]$, $\theta_0 \in (0, p_{-})$, and $p_0 \in (\widehat{p_+}, \infty)$. Moreover, choose $q_0 \in (\widehat{p_+}, 1]$. From (3.7) and [46, (9)], we deduce that, for any non-negative measurable functions $\{f_k\}_{k=1}^{\infty}$ and any $B \in \mathcal{B}$,

$$
\sum_{k=1}^{\infty} \left\|f_k\right\|_{L^{\frac{\vec{p}}{q_0}}(\mathbb{R}^n)} \leq C \left\|\sum_{k=1}^{\infty} f_k\right\|_{L^{\frac{\vec{p}}{q_0}}(\mathbb{R}^n)}
$$

and

$$
\|1_B\|_{L^{\vec{p}}(\mathbb{R}^n)} \gtrsim \min\left\{|B|^{\frac{1}{\vec{p}+}},|B|^{\frac{1}{\vec{p}^-}}\right\} = \min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{\theta_0}}\right\}.
$$

Thus, $L^{\vec{p}}(\mathbb{R}^n)$ satisfies all the assumptions of Theorem 3.1 with $X := L^{\vec{p}}(\mathbb{R}^n)$. In this case, Theorem 3.1 was obtained in [46, Theorem 3.1].

(v) As was mentioned in [36, Remark 2.1(ii)], (3.4) implies that the function $f \in H_X^A(\mathbb{R}^n) \cap$ $L^1(\mathbb{R}^n)$ has a vanishing moment. This illustrates the necessity of the vanishing moment of atoms in some sense.

To prove Theorem 3.1, we need more preparations. Let *A* be a dilation. Recall that the dilation operator *D_A* is defined by setting, for any $f \in \mathcal{M}(\mathbb{R}^n)$,

$$
D_A(f)(\cdot) := f(A\cdot).
$$

Then, by an elementary calculation (see also [7, (3.1)]), we find that, for any $k \in \mathbb{Z}$, $f \in L^1(\mathbb{R}^n)$, and $x \in \mathbb{R}^n$,

(3.8)
$$
\widehat{f}(x) = b^k \left(D_{A^*}^k \left(\mathcal{F} \left(D_A^k f \right) \right) \right)(x).
$$

Next, we recall the definitions of anisotropic (X, q, d) -atoms and anisotropic atomic Hardy spaces $H_{X,\text{atom}}^{A,q,d}(\mathbb{R}^n)$ which were first introduced in [59, Definitions 4.1 and 4.2].

Definition 3.3. Let A, X, θ_0 , and p_0 be the same as in Definition 2.10. Further assume that $q \in$ (max $\{p_0, 1\}$, ∞) and

(3.9)
$$
d \in \left[\left| \left(\frac{1}{\theta_0} - 1 \right) \frac{\ln b}{\ln(\lambda_-)} \right|, \infty \right) \cap \mathbb{Z}_+.
$$

- (i) An *anisotropic* (X, q, d) *-atom a* is a measurable function on \mathbb{R}^n satisfying that
	- (i)₁ supp (*a*) := {*x* ∈ ℝ^{*n*} : *a*(*x*) ≠ 0} ⊂ *B*, where *B* ∈ *B* and *B* is the same as in (2.3);
	- $(i)_2$ $||a||_{L^q(\mathbb{R}^n)} \leq |B|^{\frac{1}{q}}||1_B||_X^{-1};$
	- (i)₃ $\int_{\mathbb{R}^n} a(x) x^{\gamma} dx = 0$ for any $\gamma \in \mathbb{Z}_+^n$ with $|\gamma| \leq d$, here and thereafter, for any $x :=$ $(x_1, \ldots, x_n) \in \mathbb{R}^n$ and $\gamma := {\gamma_1, \ldots, \gamma_n} \in \mathbb{Z}_+^n$, $|\gamma| := \gamma_1 + \cdots + \gamma_n$ and $x^\gamma := x_1^{\gamma_1}$ $\frac{\gamma_1}{1} \cdots \frac{\gamma_n}{n}$.

Fourier Transform of Anisotropic Hardy Spaces 13 and 14 and 15 and 16 and 16 and 1

(ii) The *anisotropic atomic Hardy space* $H_{X,\text{atom}}^{A,q,d}(\mathbb{R}^n)$ is defined to be the set of all the $f \in \mathcal{S}'(\mathbb{R}^n)$ satisfying that there exists a sequence $\{\lambda_j\}_{j\in\mathbb{N}} \subset \mathbb{C}$ and a sequence $\{a_j\}_{j\in\mathbb{N}}$ of (X, q, d) -atoms supported, respectively, in $\{B^j\}_{j\in\mathbb{N}} \subset \mathcal{B}$ such that

$$
f = \sum_{j \in \mathbb{N}} \lambda_j a_j
$$

in $S'(\mathbb{R}^n)$ and that

$$
\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{|\lambda_j| \mathbf{1}_{B^j}}{\|\mathbf{1}_{B^j}\|_X} \right]^{\theta_0} \right\}^{\frac{1}{\theta_0}} \right\|_X < \infty.
$$

Moreover, for any $f \in H_{X, \text{atom}}^{A,q,d}(\mathbb{R}^n)$, let

$$
\|f\|_{H^{A,q,d}_{X,\text{atom}}(\mathbb{R}^n)}:=\inf\left\|\left\{\sum_{j\in\mathbb{N}}\left[\frac{|\lambda_j|\mathbf{1}_{B^j}}{\|\mathbf{1}_{B^j}\|_X}\right]^{\theta_0}\right\}^{\frac{1}{\theta_0}}\right\|_X,
$$

where the infimum is taken over all the decompositions of *f* as above.

The following atomic characterization of $H_X^A(\mathbb{R}^n)$, which was established in [59, Theorem 4.3], is needed in the proof of Theorem 3.1.

Lemma 3.4. Let A, X, q, and d be the same as in Definition 3.3. Then $H_X^A(\mathbb{R}^n) = H_{X,\text{atom}}^{A,q,d}(\mathbb{R}^n)$ with *equivalent quasi-norms.*

By an argument similar to that used in proof of [7, Lemma 4], we immediately obtain the following conclusioan.

Lemma 3.5. *Let A, X, q, and d be the same as in Definition 3.3. Assume that a is an anisotropic* (X, q, d) -atom supported in $x_0 + B_{i_0}$ with some $x_0 \in \mathbb{R}^n$ and $i_0 \in \mathbb{Z}$. Then there exists a positive *constant C such that, for any* $\alpha \in \mathbb{Z}_+^n$ *with* $|\alpha| \leq d$ *and for any* $x \in \mathbb{R}^n$ *,*

$$
(3.10) \qquad \left|\partial^{\alpha}\left(\mathscr{F}\left(D_A^{i_0}a\right)\right)(x)\right| \leq C \left\|\mathbf{1}_{B_{i_0}}\right\|_{X}^{-1} \min\left\{1, |x|^{d-|\alpha|+1}\right\},\
$$

where C is also independent of a.

Proof. Without loss of generality, we may assume that *a* is supported in B_{i_0} . Thus, supp $(D_A^{i_0})$ *A a*) ⊂ B_0 . On the one hand, by [23, (1.20)], Definition 3.3(i)₃, the Taylor remainder theorem, the Hölder inequality, and Definition 3.3(i)₂, we conclude that, for any $\alpha \in \mathbb{Z}_{+}^{n}$ with $|\alpha| \leq d$ and for any $x \in \mathbb{R}^n$,

(3.11)
$$
\left| \partial^{\alpha} \left(\mathcal{F} \left(D_A^{i_0} a \right) \right) (x) \right| = \left| \int_{B_0} (-2\pi i \xi)^{\alpha} \left(D_A^{i_0} a \right) (\xi) e^{-2\pi i x \cdot \xi} d\xi \right|
$$

$$
= \left| \int_{B_0} (-2\pi i \xi)^{\alpha} \left(D_A^{i_0} a \right) (\xi) \left[e^{-2\pi i x \cdot \xi} - T(\xi) \right] d\xi \right|
$$

$$
\begin{split} &\lesssim \int_{B_0} |\xi|^{|\alpha|} \left| a\left(A^{i_0}\xi\right) \right| |x|^{d-|\alpha|+1} |\xi|^{d-|\alpha|+1} \, d\xi \\ &\lesssim |x|^{d-|\alpha|+1} b^{-i_0} \int_{B_{i_0}} |a\left(\xi\right) | \, d\xi \\ &\leq |x|^{d-|\alpha|+1} \left\| \mathbf{1}_{B_{i_0}} \right\|_{X}^{-1}, \end{split}
$$

where $T(\xi)$ is the $(d - |\alpha|)$ th-order Taylor polynomial of the function $\xi \to e^{-2\pi ix\cdot\xi}$ at the origin. On the other hand, from [23, (1.20)], the Hölder inequality, and Definition $3.3(i)_2$, we deduce that, for any $\alpha \in \mathbb{Z}_+^n$ with $|\alpha| \le d$ and for any $x \in \mathbb{R}^n$,

$$
\left| \partial^{\alpha} \left(\mathcal{F} \left(D_A^{i_0} a \right) \right) (x) \right| = \left| \int_{B_0} (-2\pi \iota \xi)^{\alpha} \left(D_A^{i_0} a \right) (\xi) e^{-2\pi \iota x \cdot \xi} d\xi \right|
$$

$$
\lesssim \int_{B_0} |\xi|^{\alpha} |a \left(A^{i_0} \xi \right) | d\xi \lesssim b^{-i_0} \int_{B_{i_0}} |a(\xi)| d\xi
$$

$$
\leq \left\| \mathbf{1}_{B_{i_0}} \right\|_{X}^{-1},
$$

which, combined with (3.11) , further implies (3.10) and hence completes the proof of Lemma $3.5.$

Applying Lemma 3.5, we obtain the following uniform estimate for anisotropic (X, q, d) -atoms, which plays a key role in the proof of Theorem 3.1.

Lemma 3.6. *Let A, X, q, d, and* θ_0 *be the same as in Definition 3.3. Further assume that X satisfies* (3.2) *with* $q_0 \in [\theta_0, 1]$ *. Then there exists a positive constant C such that, for any anisotropic* (X, q, d) -atom a and for any $x \in \mathbb{R}^n$,

(3.12)
$$
|\widehat{a}(x)| \leq C \max \left\{ [\rho_*(x)]^{\frac{1}{q_0}-1}, [\rho_*(x)]^{\frac{1}{q_0}-1} \right\},
$$

where $ρ_*$ *is the same as in Theorem 3.1.*

The proof of Lemma 3.6 needs the following inequalities which are just [4, p. 11, Lemma 3.2]. **Lemma 3.7.** Let A be a dilation. Then there exists a positive constant C such that, for any $x \in \mathbb{R}^n$,

$$
\frac{1}{C}[\rho(x)]^{\ln(\lambda_-)/\ln b} \le |x| \le C[\rho(x)]^{\ln(\lambda_+)/\ln b} \text{ when } \rho(x) \in (1, \infty)
$$

and

$$
\frac{1}{C}[\rho(x)]^{\ln(\lambda_+)/\ln b} \le |x| \le C[\rho(x)]^{\ln(\lambda_-)/\ln b} \text{ when } \rho(x) \in [0, 1].
$$

Now, we give the proof of Lemma 3.6.

Proof of Lemma 3.6. Let *a* be an anisotropic (X, q, d) -atom supported in $x_0 + B_{i_0}$ with some $x_0 \in \mathbb{R}^n$ and $i_0 \in \mathbb{Z}$. Without loss of generality, we may assume $x_0 = 0$. By (3.8), Lemma 3.5 with *n* times

 $\alpha = \overbrace{0, \ldots, 0}^{n \text{ times}}$ $\widehat{0, \ldots, 0}$, and (3.2), we conclude that, for any $x \in \mathbb{R}^n$,

(3.13)
$$
\left| \widehat{a}(x) \right| = \left| b^{i_0} \left(D_{A^*}^{i_0} \left(\mathcal{F} \left(D_A^{i_0} a \right) \right) \right) (x) \right| = \left| b^{i_0} \mathcal{F} \left(D_A^{i_0} a \right) \left((A^*)^{i_0} x \right) \right|
$$

Fourier Transform of Anisotropic Hardy Spaces 15 and 16 and 1

$$
\leq b^{i_0} \left\| \mathbf{1}_{B_{i_0}} \right\|_X^{-1} \min \left\{ 1, \left| (A^*)^{i_0} x \right|^{d+1} \right\} \leq b^{i_0} \max \left\{ b^{-\frac{i_0}{q_0}}, b^{-\frac{i_0}{\theta_0}} \right\} \min \left\{ 1, \left| (A^*)^{i_0} x \right|^{d+1} \right\}.
$$

Next, we prove (3.12) by considering two cases: $\rho_*(x) \le b^{-i_0}$ and $\rho_*(x) > b^{-i_0}$.

Case 1) $\rho_*(x) \le b^{-i_0}$. In this case, note that

(3.14)
$$
\rho_*\left((A^*)^{i_0}x\right) = b^{i_0}\rho_*(x) \le 1.
$$

Moreover, by (3.9), we find that

$$
1 - \frac{1}{q_0} + (d+1)\frac{\ln(\lambda_-)}{\ln b} \ge 1 - \frac{1}{\theta_0} + (d+1)\frac{\ln(\lambda_-)}{\ln b} > 0.
$$

From this, (3.13), (3.14), and Lemma 3.7, we infer that, for any $x \in \mathbb{R}^n$ satisfying $\rho_*(x) \le b^{-i_0}$,

$$
(3.15) \qquad |\widehat{a}(x)| \lesssim b^{i_0} \max \left\{ b^{-\frac{i_0}{q_0}}, b^{-\frac{i_0}{\theta_0}} \right\} \left[\rho_* \left((A^*)^{i_0} x \right) \right]^{(d+1)\frac{\ln(2-)}{\ln b}} = \max \left\{ b^{i_0[1-\frac{1}{q_0}+(d+1)\frac{\ln(2-)}{\ln b}]}, b^{i_0[1-\frac{1}{q_0}+(d+1)\frac{\ln(2-)}{\ln b}]} \right\} \left[\rho_*(x) \right]^{(d+1)\frac{\ln(2-)}{\ln b}} = \max \left\{ \left[\rho_*(x) \right]^{\frac{1}{q_0}-1}, \left[\rho_*(x) \right]^{\frac{1}{\theta_0}-1} \right\}.
$$

This shows (3.12) in Case 1).

Case 2) $\rho_*(x) > b^{-i_0}$. In this case, note that

$$
\rho_*\left((A^*)^{i_0}x\right) = b^{i_0}\rho_*(x) > 1.
$$

Using this, (3.13), Lemma 3.7, and the fact that

$$
\frac{1}{\theta_0} - 1 \ge \frac{1}{q_0} - 1 \ge 0,
$$

we conclude that, for any $x \in \mathbb{R}^n$ satisfying $\rho_*(x) > b^{-i_0}$,

$$
|\widehat{a}(x)| \le b^{i_0} \max \left\{ b^{-\frac{i_0}{q_0}}, b^{-\frac{i_0}{\theta_0}} \right\} = \max \left\{ b^{-i_0(\frac{1}{q_0}-1)}, b^{-i_0(\frac{1}{\theta_0}-1)} \right\}
$$

$$
\le \max \left\{ \left[\rho_*(x) \right]^{\frac{1}{q_0}-1}, \left[\rho_*(x) \right]^{\frac{1}{\theta_0}-1} \right\},
$$

which, combined with (3.15), then completes the proof of (3.12) and hence Lemma 3.6. \Box

The following inequality is basic and used throughout this article.

Lemma 3.8. *Let* $\{a_j\}_{j \in \mathbb{N}}$ ⊂ [0, ∞)*. If* $\alpha \in (0, 1]$ *, then*

$$
\left(\sum_{j\in\mathbb{N}}a_j\right)^\alpha\leq \sum_{j\in\mathbb{N}}a_j^\alpha.
$$

.

The following conclusion is also used in the proof of Theorem 3.1.

Lemma 3.9. Let A, X, and θ_0 be the same as in Definition 3.3. Further assume that X satisfies (3.1) *with* $q_0 \in [\theta_0, 1]$ *. Then there exists a positive constant C such that, for any* $\{\lambda_i\}_{i\in\mathbb{N}} \subset \mathbb{C}$ *and* {*B* (*i*) }*i*∈^N ⊂ B*,*

$$
\sum_{i\in\mathbb{N}}|\lambda_i|\leq C\left\|\left\{\sum_{i\in\mathbb{N}}\left[\frac{|\lambda_i|\mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X}\right]^{ \theta_0}\right\}_{\theta}\right\|_X.
$$

Proof. Indeed, by Lemma 3.8, Definition 2.5, (3.1), and Definition 2.4(ii), we find that, for any $\{\lambda_i\}_{i\in\mathbb{N}}\subset\mathbb{C}$ and $\{B^{(i)}\}_{i\in\mathbb{N}}\subset\mathcal{B}$,

$$
\sum_{i=1}^{\infty} |\lambda_i| \le \left(\sum_{i=1}^{\infty} |\lambda_i|^{q_0} \right)^{\frac{1}{q_0}} = \left\{ \sum_{i=1}^{\infty} \left\| \frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X} \right\|_{X}^{q_0} \right\}^{\frac{1}{q_0}}
$$
\n
$$
= \left\{ \sum_{i=1}^{\infty} \left\| \frac{|\lambda_i|^{q_0} \mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X^{q_0}} \right\|_{X^{\frac{1}{q_0}}} \right\}^{\frac{1}{q_0}} \lesssim \left\| \sum_{i=1}^{\infty} \left[\frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X} \right]^{q_0} \right\|_{X^{\frac{1}{q_0}}}
$$
\n
$$
= \left\| \left\{ \sum_{i=1}^{\infty} \left[\frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X} \right]^{q_0} \right\}^{\frac{1}{q_0}} \right\|_{X} \le \left\| \left\{ \sum_{i=1}^{\infty} \left[\frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X} \right]^{q_0} \right\}^{\frac{1}{q_0}}
$$

This finishes the proof of Lemma 3.9.

Next, we show Theorem 3.1.

Proof of Theorem 3.1. Let *q* and *d* be the same as in Definition 3.3. Without loss of generality, we may assume that $||f||_{H_X^A(\mathbb{R}^n)} > 0$. Then, by Lemma 3.4 and Definition 3.3(ii), we find that there exists a sequence $\{\lambda_i\}_{i\in\mathbb{N}} \subset \mathbb{C}$ and a sequence $\{a_i\}_{i\in\mathbb{N}}$ of anisotropic (X, q, d) -atoms supported, respectively, in ${B^{(i)}}_{i \in \mathbb{N}} \subset \mathcal{B}$ such that

(3.16)
$$
f = \sum_{i \in \mathbb{N}} \lambda_i a_i \text{ in } \mathcal{S}'(\mathbb{R}^n)
$$

and

(3.17)
$$
||f||_{H_X^A(\mathbb{R}^n)} \sim \left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{||\mathbf{1}_{B^{(i)}}||_X} \right]^{ \theta_0} \right\}^{1/\theta_0} \right\|_X.
$$

First, we try to find the desired function *F*. Note that a function $g \in L^1(\mathbb{R}^n)$ implies that \widehat{g} is well defined in \mathbb{R}^n (see, for instance, [23, (1.11)]), so does $\widehat{a_i}$ for any $i \in \mathbb{N}$. Moreover, from Lemmas 3.6 and 3.9 and from (3.17), it follows that, for any $x \in \mathbb{R}^n$,

$$
(3.18) \qquad \sum_{i \in \mathbb{N}} |\lambda_i| |\widehat{a_i}(x)| \lesssim \sum_{i \in \mathbb{N}} |\lambda_i| \max \left\{ [\rho_*(x)]^{\frac{1}{q_0} - 1}, [\rho_*(x)]^{\frac{1}{\theta_0} - 1} \right\} \\ \lesssim ||f||_{H_X^A(\mathbb{R}^n)} \max \left\{ [\rho_*(x)]^{\frac{1}{q_0} - 1}, [\rho_*(x)]^{\frac{1}{\theta_0} - 1} \right\} < \infty.
$$

Fourier Transform of Anisotropic Hardy Spaces 17 and 17

Therefore, the function

$$
F(\cdot) := \sum_{i \in \mathbb{N}} \lambda_i \widehat{a_i}(\cdot)
$$

is well defined pointwisely on \mathbb{R}^n and, for any $x \in \mathbb{R}^n$,

$$
|F(x)| \lesssim ||f||_{H_X^A(\mathbb{R}^n)} \max \left\{ [\rho_*(x)]^{\frac{1}{q_0}-1}, [\rho_*(x)]^{\frac{1}{\theta_0}-1} \right\},\,
$$

which completes the proof of (3.4) .

Second, we show the continuity of F on \mathbb{R}^n . If we can prove that F is continuous on any compact subset of \mathbb{R}^n , then the continuity of *F* on \mathbb{R}^n is obvious. Let $E \subset \mathbb{R}^n$ be any given compact set. Then there exists a positive constant *K*, depending only on *A* and *E*, such that $\rho_*(x) \leq K$ holds true for any $x \in E$. By this and (3.18), we conclude that, for any $x \in E$,

$$
\sum_{i\in\mathbb{N}}|\lambda_i||\widehat{a_i}(x)|\lesssim \max\left\{K^{\frac{1}{q_0}-1},\ K^{\frac{1}{\theta_0}-1}\right\}||f||_{H_X^A(\mathbb{R}^n)}<\infty.
$$

Thus, the summation $\sum_{i\in\mathbb{N}} \lambda_i \widehat{a_i}(\cdot)$ converges uniformly on *E*. This, together with the fact that $\widehat{a_i}$ is continuous for any $i \in \mathbb{N}$, further implies that *F* is also continuous on *E* and hence on \mathbb{R}^n .

Finally, we show (3.3). By (3.16) and the continuity of the Fourier transform in $\mathcal{S}'(\mathbb{R}^n)$ (see, for instance, [23, Theorem 1.17]), we obtain

$$
\widehat{f} = \sum_{i \in \mathbb{N}} \lambda_i \widehat{a_i} \text{ in } \mathcal{S}'(\mathbb{R}^n).
$$

Thus, to prove (3.3), we only need to show that

(3.20)
$$
F = \sum_{i \in \mathbb{N}} \lambda_i \widehat{a_i} \text{ in } \mathcal{S}'(\mathbb{R}^n).
$$

Indeed, from Lemma 3.6 and the definition of Schwartz functions [see (2.7)], we deduce that, for any $i \in \mathbb{N}$ and $\varphi \in \mathcal{S}(\mathbb{R}^n)$,

$$
\begin{split}\n|\langle \widehat{a_i}, \varphi \rangle| &= \left| \int_{\mathbb{R}^n} \widehat{a_i}(x) \varphi(x) \, dx \right| \\
&\leq \sum_{k=1}^{\infty} \int_{(A^*)^{k+1} B_0^* \setminus (A^*)^k B_0^*} \max \left\{ [\rho_*(x)]^{\frac{1}{q_0} - 1}, [\rho_*(x)]^{\frac{1}{q_0} - 1} \right\} |\varphi(x)| \, dx \\
&+ \|\varphi\|_{L^1(\mathbb{R}^n)} \\
&\lesssim \sum_{k=1}^{\infty} b^{k+1} b^{k(\frac{1}{q_0} - 1)} b^{-k(\lfloor \frac{1}{q_0} \rfloor + 2)} + \|\varphi\|_{L^1(\mathbb{R}^n)} \\
&\leq \sum_{k=1}^{\infty} b^{-k} + \|\varphi\|_{L^1(\mathbb{R}^n)},\n\end{split}
$$

where B_0^* is the unit dilated ball with respect to A^* . This further implies that there exists a positive constant *C* such that $|\langle \hat{a}_i, \varphi \rangle| \le C$ holds true uniformly for any $i \in \mathbb{N}$. Combining this and (3.17), we have

$$
\lim_{I\to\infty}\sum_{i=I+1}^{\infty}|\lambda_i||\langle\widehat{a_i},\varphi\rangle|\lesssim \lim_{I\to\infty}\sum_{i=I+1}^{\infty}|\lambda_i|=0.
$$

Therefore, for any $\varphi \in \mathcal{S}(\mathbb{R}^n)$,

$$
\langle F, \varphi \rangle = \lim_{I \to \infty} \left\langle \sum_{i=1}^{I} \lambda_i \widehat{a_i}, \varphi \right\rangle.
$$

This finishes the proof of (3.20) and hence Theorem 3.1. \square

4 Hardy–Littlewood Inequalities on H_X^A $^A_X(\mathbb{R}^n)$

In this section, as applications of Theorem 3.1, we first prove that the function *F* given in Theorem 3.1 has a higher order convergence at the origin (see Theorem 4.1 below). Then we extend the Hardy–Littlewood inequality to the setting of anisotropic Hardy spaces associated with ball quasi-Banach function spaces (see Theorem 4.2 below). In what follows, $\varepsilon \to 0^+$ means that there exists an $\alpha_0 \in (0, \infty)$ such that $\varepsilon \in (0, \alpha_0)$ and $\varepsilon \to 0$.

Theorem 4.1. *Let A, X, q*₀*, and* ρ_* *be the same as in Theorem 3.1. Then, for any* $f \in H_X^A(\mathbb{R}^n)$ *, there exists a continuous function* F *on* \mathbb{R}^n *such that* $\widehat{f} = F$ *in* $S'(\mathbb{R}^n)$ *and*

(4.1)
$$
\lim_{|x| \to 0^+} \frac{F(x)}{[\rho_*(x)]^{\frac{1}{\theta_0}-1}} = 0.
$$

Proof. Let $f \in H_X^A(\mathbb{R}^n)$ and *q* and *d* be the same as in Definition 3.3. Then, by Lemma 3.4 and Definition 3.3(ii), we find that there exists a sequence $\{\lambda_i\}_{i\in\mathbb{N}} \subset \mathbb{C}$ and a sequence $\{a_i\}_{i\in\mathbb{N}}$ of anisotropic (X, q, d) -atoms supported, respectively, in ${B^{(i)}}_{i \in \mathbb{N}} \subset \mathcal{B}$ such that

$$
f = \sum_{i \in \mathbb{N}} \lambda_i a_i \text{ in } \mathcal{S}'(\mathbb{R}^n)
$$

and

(4.2)
$$
||f||_{H_X^A(\mathbb{R}^n)} \sim \left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{||\mathbf{1}_{B^{(i)}}||_X} \right]^{\theta_0} \right\}_{0}^{\frac{1}{\theta_0}} \right\|_X.
$$

Moreover, from the proof of Theorem 3.1, it follows that, for any $x \in \mathbb{R}^n$,

(4.3)
$$
F(x) = \sum_{i \in \mathbb{N}} \lambda_i \widehat{a_i}(x)
$$

is continuous and satisfies that $\hat{f} = F$ in $\mathcal{S}'(\mathbb{R}^n)$. Thus, to show the present theorem, we only need to prove that (4.1) holds true for *F* in (4.3) . On the one hand, by an argument similar to that used

in the proof of (3.15), we conclude that, for any anisotropic (X, q, d) -atom *a* supported in $x_0 + B_{k_0}$ with some $x_0 \in \mathbb{R}^n$ and $k_0 \in \mathbb{Z}$ and for any $x \in \mathbb{R}^n$ satisfying $\rho_*(x) \leq b^{-k_0}$,

$$
\left|\widehat{a}(x)\right|\lesssim \max\left\{b^{k_0[1-\frac{1}{q_0}+(d+1)\frac{\ln(1_-)}{\ln b}]},\; b^{k_0[1-\frac{1}{\theta_0}+(d+1)\frac{\ln(1_-)}{\ln b}]}\right\}\left[\rho_*(x)\right]^{(d+1)\frac{\ln(1_-)}{\ln b}},
$$

which, together with (3.9) and Lemma 3.7, further implies that

$$
(d+1)\frac{\ln(\lambda_-)}{\ln b} > \frac{1}{\theta_0} - 1
$$

and

(4.4)
$$
\lim_{|x| \to 0^+} \frac{|\widehat{a}(x)|}{[\rho_*(x)]^{\frac{1}{\theta_0}-1}} = 0.
$$

On the other hand, from (4.3), Lemmas 3.6, 3.7, and 3.9, and (4.2), we deduce that, for any $x \in \mathbb{R}^n$ satisfying $|x|$ < 1,

$$
(4.5) \qquad \qquad \frac{|F(x)|}{[\rho_*(x)]^{\frac{1}{\theta_0}-1}} \leq \sum_{i\in\mathbb{N}} |\lambda_i| \frac{|\widehat{a_i}(x)|}{[\rho_*(x)]^{\frac{1}{\theta_0}-1}} \lesssim \sum_{i\in\mathbb{N}} |\lambda_i| \lesssim ||f||_{H_X^A(\mathbb{R}^n)} < \infty.
$$

Using this, the dominated convergence theorem, and (4.4), we find that

$$
\lim_{|x| \to 0^+} \frac{F(x)}{\left[\rho_*(x)\right]^{\frac{1}{q_0}-1}} = 0,
$$

which completes the proof of Theorem 4.1.

As another application of Theorem 3.1, we extend the Hardy–Littlewood inequality to the setting of anisotropic Hardy spaces associated with ball quasi-Banach function spaces as follows.

Theorem 4.2. *Let A, X,* θ_0 *, and* q_0 *be the same as in Theorem 3.1. Then, for any* $f \in H_X^A(\mathbb{R}^n)$ *, there exists a continuous function* F *on* \mathbb{R}^n *such that* $\widehat{f} = F$ *in* $S'(\mathbb{R}^n)$ *and*

$$
(4.6) \qquad \left[\int_{\mathbb{R}^n} |F(x)|^{q_0} \min \left\{ \left[\rho_*(x) \right]^{q_0 - \frac{q_0}{\theta_0} - 1}, \left[\rho_*(x) \right]^{q_0 - 2} \right\} dx \right]^{\frac{1}{q_0}} \le C \|f\|_{H^A_X(\mathbb{R}^n)},
$$

where C is a positive constant depending only on A and X.

Proof. Let p_0 and *d* be the same as in Definition 3.3, $q \in (\max\{p_0, 2\}, \infty]$, and $f \in H_X^A(\mathbb{R}^n)$. Then, by Lemma 3.4 and Definition 3.3, we find that there exists a sequence $\{\lambda_i\}_{i\in\mathbb{N}}\subset\mathbb{C}$ and a sequence { a_i }_{*i*∈N} of (*X*, *q*, *d*)-atoms supported, respectively, in { B ^(*i*)}_{*i*∈N} ⊂ B such that

$$
f = \sum_{i \in \mathbb{N}} \lambda_i a_i \text{ in } \mathcal{S}'(\mathbb{R}^n)
$$

and

$$
(4.7) \qquad \qquad \left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{|\lambda_i| \mathbf{1}_{B^{(i)}}}{\|\mathbf{1}_{B^{(i)}}\|_X} \right]^{\theta_0} \right\}^{1/\theta_0} \right\|_X \sim \|f\|_{H_X^A(\mathbb{R}^n)} < \infty.
$$

.

By Theorem 3.1, we find that, to prove the present theorem, it suffices to show that (4.6) holds true for *F* in (3.19). For this purpose, we first prove that there exists a positive constant *M* such that, for any (X, q, d) -atom a , it holds true that

$$
(4.8) \qquad \left(\int_{\mathbb{R}^n} \left[\left| \widehat{a}(x) \right| \min \left\{ \left[\rho_*(x) \right]^{1 - \frac{1}{\theta_0} - \frac{1}{q_0}}, \left[\rho_*(x) \right]^{1 - \frac{2}{q_0}} \right\} \right]^{q_0} dx \right)^{\frac{1}{q_0}} \le M.
$$

Without loss of generality, we may assume that *a* is supported in $x_0 + B_{i_0}$ with some $x_0 \in \mathbb{R}^n$ and $i_0 \in \mathbb{Z}$. Then it is easy to conclude that

$$
(4.9) \qquad \left\{ \int_{\mathbb{R}^n} \left[\left| \widehat{a}(x) \right| \min \left\{ \left[\rho_*(x) \right]^{1 - \frac{1}{\theta_0} - \frac{1}{q_0}}, \left[\rho_*(x) \right]^{1 - \frac{2}{q_0}} \right\} \right]^{q_0} dx \right\}^{\frac{1}{q_0}} \leq \left\{ \int_{(A^*)^{-i_0 + 1} B_0^*} \left[\left| \widehat{a}(x) \right| \min \left\{ \left[\rho_*(x) \right]^{1 - \frac{1}{\theta_0} - \frac{1}{q_0}}, \left[\rho_*(x) \right]^{1 - \frac{2}{q_0}} \right\} \right]^{q_0} dx \right\}^{\frac{1}{q_0}} \newline + \left\{ \int_{((A^*)^{-i_0 + 1} B_0^*)} \cdots dx \right\}^{\frac{1}{q_0}} \newline =: I_1 + I_2,
$$

where B_0^* is the unit dilated ball with respect to A^* . Let θ be a fixed positive constant such that

$$
1 - \frac{1}{q_0} + (d+1)\frac{\ln(\lambda_-)}{\ln b} - \theta \ge 1 - \frac{1}{\theta_0} + (d+1)\frac{\ln(\lambda_-)}{\ln b} - \theta > 0.
$$

Using this and (3.15), we find that

$$
(4.10) \tI_{1} \leq b^{i_{0}[1+(d+1)\frac{\ln \lambda}{\ln b}]} \max\left\{b^{-\frac{i_{0}}{\theta_{0}}}, b^{-\frac{i_{0}}{\theta_{0}}}\right\}
$$

$$
\times \left\{\int_{(A^{*})^{-i_{0}+1}B_{0}^{*}} \left[\min\left\{\left[\rho_{*}(x)\right]^{1-\frac{1}{\theta_{0}}-\frac{1}{\eta_{0}}+(d+1)\frac{\ln \lambda}{\ln b}}\right\}\right]^{q_{0}}
$$

$$
\leq b^{i_{0}[1+(d+1)\frac{\ln \lambda}{\ln b}]} \max\left\{b^{-\frac{i_{0}}{\theta_{0}}}, b^{-\frac{i_{0}}{\theta_{0}}}\right\}
$$

$$
\times \min\left\{b^{-i_{0}[1-\frac{1}{\theta_{0}}+(d+1)\frac{\ln \lambda}{\ln b}-\theta]}, b^{-i_{0}[1-\frac{1}{\theta_{0}}+(d+1)\frac{\ln \lambda}{\ln b}-\theta]}\right\}
$$

$$
\times \left\{\int_{(A^{*})^{-i_{0}+1}B_{0}^{*}} \left[\rho_{*}(x)\right]^{q_{0}-1} dx\right\}^{\frac{1}{q_{0}}}
$$

$$
= b^{i_{0}\theta}\left[\sum_{k \in \mathbb{Z} \backslash \mathbb{N}} b^{-i_{0}+k}(b-1)b^{(-i_{0}+k)(q_{0}-1)}\right]^{\frac{1}{q_{0}}} = \left(\frac{b-1}{1-b^{-\theta_{0}}}\right)^{\frac{1}{q_{0}}}
$$

For I₂, by the Hölder inequality, the Plancherel theorem (see [23, Theorem 1.18]), $q_0 \in [\theta_0, 1]$, Definition $3.3(i)_2$, and (3.2) , we obtain

$$
I_{2} \leq \left\{ \int_{((A^{*})^{-i_{0}+1}B_{0}^{*})} \left[\overline{a}(x)\right]^{2} dx \right\}^{\frac{1}{2}}
$$
\n
$$
\times \left\{ \int_{((A^{*})^{-i_{0}+1}B_{0}^{*})} \left[\min \left\{ \left[\rho_{*}(x)\right]^{1-\frac{1}{p_{-}}-\frac{1}{q_{0}}}, \left[\rho_{*}(x)\right]^{1-\frac{2}{q_{0}}} \right\} \right]^{\frac{2q_{0}}{2-q_{0}}} dx \right\}^{\frac{2-q_{0}}{2q_{0}}}
$$
\n
$$
\leq ||a||_{L^{2}(\mathbb{R}^{n})} \left\{ \sum_{k \in \mathbb{N}} b^{-i_{0}+k} (b-1) \left[\min \left\{ b^{(-i_{0}+k)(1-\frac{1}{p_{-}}-\frac{1}{q_{0}})}, b^{(-i_{0}+k)(1-\frac{2}{q_{0}})} \right\} \right]^{\frac{2q_{0}}{2-q_{0}}}} \right\}^{\frac{2-q_{0}}{2q_{0}}}
$$
\n
$$
\leq ||a||_{L^{2}(\mathbb{R}^{n})} \left\{ b^{-i_{0}} \left[\min \left\{ b^{-i_{0}(1-\frac{1}{p_{-}}-\frac{1}{q_{0}})}, b^{-i_{0}(1-\frac{2}{q_{0}})} \right\} \right]^{\frac{2q_{0}}{2-q_{0}}}} \right\}^{\frac{2-q_{0}}{2q_{0}}}
$$
\n
$$
\leq \max \left\{ b^{i_{0}(\frac{1}{2}-\frac{1}{p_{-}})}, b^{i_{0}(\frac{1}{2}-\frac{1}{q_{0}})} \right\} \min \left\{ b^{-i_{0}(\frac{1}{2}-\frac{1}{p_{-}})}, b^{-i_{0}(\frac{1}{2}-\frac{1}{q_{0}})} \right\}
$$
\n= 1,

which, together with (4.9) and (4.10) , further implies (4.8) .

Next, we prove (4.6). From Lemma 3.8, Definition 2.5, (3.1), Definition 2.4(ii), an argument similar to that used in the proof of Lemma 3.9, and (4.7), we deduce that

$$
\left(\sum_{i=1}^{\infty}|\lambda_i|^{q_0}\right)^{\frac{1}{q_0}} \lesssim ||f||_{H_X^A(\mathbb{R}^n)}.
$$

By this, (3.19) , $q_0 \in [\theta_0, 1]$, Lemma 3.8, the Fatou lemma, and (4.8) , we conclude that

$$
\left[\int_{\mathbb{R}^n} |F(x)|^{q_0} \min\left\{ [\rho_*(x)]^{q_0 - \frac{q_0}{\theta_0} - 1}, [\rho_*(x)]^{q_0 - 2} \right\} dx \right]^{\frac{1}{q_0}} \n\leq \left\{ \sum_{i \in \mathbb{N}} |\lambda_i|^{q_0} \int_{\mathbb{R}^n} \left[|\widehat{a_i}(x)| \min\left\{ [\rho_*(x)]^{1 - \frac{1}{\theta_0} - \frac{1}{q_0}}, [\rho_*(x)]^{1 - \frac{2}{q_0}} \right\} \right]^{q_0} dx \right\}^{\frac{1}{q_0}} \n\leq M \left(\sum_{i \in \mathbb{N}} |\lambda_i|^{q_0} \right)^{\frac{1}{q_0}} \lesssim ||f||_{H^A_X(\mathbb{R}^n)}.
$$

This finishes the proof of Theorem 4.2.

- **Remark 4.3.** (i) If $A := 2I_{n \times n}$, then Theorems 4.1 and 4.2 were obtained, respectively, in [36, Theorems 2.2 and 2.3].
	- (ii) Let *A* be a dilation and $p \in (0, 1)$. Then, by Remark 3.2(ii), we find that $L^p(\mathbb{R}^n)$ satisfies all the assumptions of Theorems 4.1 and 4.2 with $X := L^p(\mathbb{R}^n)$. In this case, Theorems 4.1 and 4.2 were obtained, respectively, in [7, Corollaries 6 and 8].

- (iii) Let *A* be a dilation and $p() \in C^{\log}(\mathbb{R}^n)$ satisfy $0 < \widetilde{p_-} \le \widetilde{p_+} < 1$. Then, by Remark 3.2(iii), we conclude that $L^{p(\cdot)}(\mathbb{R}^n)$ satisfies all the assumptions of Theorems 4.1 and 4.2 with $X := L^{p(\cdot)}(\mathbb{R}^n)$. In this case, Theorems 4.1 and 4.2 were obtained, respectively, in [45, Theorems 2 and 3].
- (iv) Let $\vec{p} \in (0, 1)^n$. Then, by Remark 3.2(iv), we conclude that $L^{\vec{p}}(\mathbb{R}^n)$ satisfies all the assumptions of Theorems 4.1 and 4.2 with $X := L^{\vec{p}}(\mathbb{R}^n)$. In this case, Theorems 4.1 and 4.2 were obtained, respectively, in [46, Theorems 4.1 and 4.3].

5 Several Applications

In this section, we apply Theorems 3.1, 4.1, and 4.2 to five concrete examples of ball quasi-Banach function spaces, namely Morrey spaces (see Subsection 5.1 below), Lorentz spaces (see Subsection 5.2 below), Orlicz spaces (see Subsection 5.3 below), Orlicz-slice spaces (see Subsection 5.4 below), and local generalized Herz–Hardy spaces (see Subsection 5.5 below).

5.1 Morrey Spaces

Recall that the classical Morrey space $M_q^p(\mathbb{R}^n)$ with $0 < q \leq p < \infty$, originally introduced by Morrey [50] in 1938, plays a key role in harmonic analysis and partial differential equations. Since then, various variants of Morrey spaces over different underlying spaces have been investigated and developed (see, for instance, [14, 31, 32, 33, 34, 54]).

Definition 5.1. Let $0 < q \le p < \infty$. The *anisotropic Morrey space* $M_{q,A}^p(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$
||f||_{M_{q, A}^p(\mathbb{R}^n)} := \sup_{B \in \mathcal{B}} |B|^{1/p-1/q} ||f||_{L^q(B)} < \infty,
$$

where \mathcal{B} is the same as in (2.3).

If $0 < q \le p < 1$, then, obviously, M_{q}^{p} $_{q, A}^p(\mathbb{R}^n)$ is a ball quasi-Banach function space. From these and [59, Remark 8.4], we deduce that M_a^p $_{q, A}^p(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10 with $X := M_q^p$ *p*_{*q*,*A*}(\mathbb{R}^n), *p*_− ∈ (0, *q*], θ_0 ∈ (0, *p*_−), and p_0 ∈ (*p*, ∞). Moreover, choose a q_0 ∈ (*p*, 1]. Then, from Definition 5.1, we infer that, for any non-negative measurable functions ${f_k}_{k=1}^{\infty}$ and any $B \in \mathcal{B}$,

$$
\sum_{k=1}^{\infty}\|f_k\|_{M_{q/q_0,A}^{p/q_0}(\mathbb{R}^n)}\leq \left\|\sum_{k=1}^{\infty}f_k\right\|_{M_{q/q_0,A}^{p/q_0}(\mathbb{R}^n)}
$$

and

$$
\|{\bf 1}_B\|_{{M^p_{q,A}(\mathbb{R}^n)}}\geq |B|^{\frac{1}{p}}>\min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{q_0}}\right\}.
$$

Therefore, all the assumptions of Theorems 3.1, 4.1, and 4.2 are satisfied with $X := M_a^p$ $_{q,A}^p(\mathbb{R}^n)$. Applying Theorems 3.1, 4.1, and 4.2, we obtain the following conclusion.

Theorem 5.2. *If* $0 < q \le p < 1$, then Theorems 3.1, 4.1, and 4.2 still hold true with X replaced by $M_{q,A}^p(\mathbb{R}^n)$.

Remark 5.3. We point out that Theorem 5.2 even when $A := 2I_{n \times n}$ is completely new.

Fourier Transform of Anisotropic Hardy Spaces 23 23

5.2 Lorentz Spaces

Let $p \in (0, \infty)$ and $q \in (0, \infty)$. Recall that the *Lorentz space L^{p,q}*(\mathbb{R}^n) is defined to be the set of all the measurable functions f on \mathbb{R}^n with the following finite *quasi-norm*

(5.1)
$$
||f||_{L^{p,q}(\mathbb{R}^n)} := \begin{cases} \left[\frac{q}{p} \int_0^{\infty} \left\{t^{\frac{1}{p}} f^*(t)\right\}^q \frac{dt}{t}\right]^{\frac{1}{q}} & \text{if } q \in (0, \infty),\\ \sup_{t \in (0, \infty)} \left[t^{\frac{1}{p}} f^*(t)\right] & \text{if } q = \infty \end{cases}
$$

with the usual modification made when $p = \infty$, where f^* denotes the *non-increasing rearrangement* of *f*, that is, for any $t \in (0, \infty)$,

$$
f^*(t) := \inf \left\{ \alpha \in (0, \infty) : d_f(\alpha) \le t \right\}
$$

with $d_f(\alpha) := |\{x \in \mathbb{R}^n : |f(x)| > \alpha\}|$ for any $\alpha \in (0, \infty)$.

Let $p \in (0, 1)$, $q \in (0, 1)$, and

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{p} - 1 \right) \frac{\ln b}{\ln(\lambda_-)} \right| + 2, \infty \right).
$$

Then, by [61, Remarks 2.7(ii) and 4.21(ii)], we conclude that $L^{p,q}(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10 with $X := L^{p,q}(\mathbb{R}^n)$, $p_- \in (0, \min\{p, q\}]$, $\theta_0 \in (0, p_-)$, and $p_0 \in (\max\{p, q\}, \infty)$. Moreover, choose a $q_0 \in (\max\{p, q\}, 1]$. From (5.1), we further deduce that, for any non-negative measurable functions $\{f_k\}_{k=1}^{\infty}$ and any $B \in \mathcal{B}$,

$$
\sum_{k=1}^{\infty} ||f_k||_{L^{p/q_0,q/q_0}(\mathbb{R}^n)} \leq \left\|\sum_{k=1}^{\infty} f_k\right\|_{L^{p/q_0,q/q_0}(\mathbb{R}^n)}
$$

and

$$
\|\mathbf{1}_B\|_{L^{p,q}(\mathbb{R}^n)}=\left\{\frac{q}{p}\int_0^{|B|}t^{\frac{q}{p}-1}dt\right\}^{\frac{1}{q}}=|B|^{\frac{1}{p}}\geq\min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{q_0}}\right\}.
$$

Therefore, all the assumptions of Theorems 3.1, 4.1, and 4.2 are satisfied with $X := L^{p,q}(\mathbb{R}^n)$. Applying Theorems 3.1, 4.1, and 4.2, we obtain the following conclusion.

Theorem 5.4. *If* $p \in (0, 1)$ *and* $q \in (0, 1)$ *, then Theorems 3.1, 4.1, and 4.2 still hold true with* X *replaced by* $L^{p,q}(\mathbb{R}^n)$ *.*

Remark 5.5. We point out that Theorem 5.4 even when $A := 2I_{n \times n}$ is completely new.

5.3 Orlicz Spaces

Recall that a function $\Phi : [0, \infty) \to [0, \infty)$ is called an *Orlicz function* if it is non-decreasing, $\Phi(0) = 0$, $\Phi(t) > 0$ for any $t \in (0, \infty)$, and $\lim_{t \to \infty} \Phi(t) = \infty$. The function Φ is said to be of *upper* (resp. *lower*) *type* p for some $p \in [0, \infty)$ if there exists a positive constant *C* such that, for any $s \in [1, \infty)$ (resp. $s \in [0, 1]$) and $t \in [0, \infty)$, $\Phi(st) \leq C s^p \Phi(t)$. The *Orlicz space* $L^{\Phi}(\mathbb{R}^n)$ is defined to be the set of all the measurable functions f on \mathbb{R}^n such that

$$
||f||_{L^{\Phi}(\mathbb{R}^n)} := \inf \left\{ \lambda \in (0, \infty) : \int_{\mathbb{R}^n} \Phi\left(\frac{|f(x)|}{\lambda}\right) dx \le 1 \right\} < \infty.
$$

Let Φ be an Orlicz function with lower type p_{Φ}^- and upper type p_{Φ}^+ satisfying $0 < p_{\Phi}^- \le p_{\Phi}^+ < 1$ and let

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{p_{\Phi}^-} - 1 \right) \frac{\ln b}{\ln(\lambda_-)} \right| + 2, \infty \right).
$$

By [61, Remarks 2.7(iii) and 4.21(iv)], we conclude that $L^{\Phi}(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10 with *X* := $L^{\Phi}(\mathbb{R}^n)$, $p_-\in(0, p_0^-]$, $\theta_0 \in (0, p_0^-)$, and $p_0 \in (p_0^+, \infty)$. Moreover, choose a $q_0 \in (p_0^+, 1]$. Then, from [67, Remark 5.3] and [36, (25)], we deduce that, for any non-negative measurable functions $\{f_k\}_{k=1}^{\infty}$ and any $B \in \mathcal{B}$,

$$
\sum_{k=1}^{\infty} ||f_k||_{L^{\Phi}(\mathbb{R}^n)^{\frac{1}{q_0}}} \le \left\| \sum_{k=1}^{\infty} f_k \right\|_{L^{\Phi}(\mathbb{R}^n)^{\frac{1}{q_0}}}
$$

and

$$
\|1_B\|_{L^{\Phi}(\mathbb{R}^n)} \gtrsim \min\left\{|B|^{\frac{1}{p_{\Phi}^-}},|B|^{\frac{1}{p_{\Phi}^+}}\right\} \ge \min\left\{|B|^{\frac{1}{q_0}},|B|^{\frac{1}{q_0}}\right\}.
$$

Therefore, all the assumptions of Theorems 3.1, 4.1, and 4.2 are satisfied with $X := L^{\Phi}(\mathbb{R}^n)$. Applying Theorems 3.1, 4.1, and 4.2, we obtain the following conclusion.

Theorem 5.6. Let Φ *be an Orlicz function with lower type* p_{Φ}^- *and upper type* p_{Φ}^+ *satisfying* 0 < $p_{\Phi}^- \leq p_{\Phi}^+ < 1$ *. Then Theorems 3.1, 4.1, and 4.2 still hold true with X replaced by* $\tilde{L}^{\Phi}(\mathbb{R}^n)$ *.*

Remark 5.7. We point out that Theorem 5.6 even when $A := 2I_{n \times n}$ is completely new.

5.4 Orlicz-Slice Spaces

Recently, Zhang et al. [67] originally introduced the Orlicz-slice space on \mathbb{R}^n , which generalizes both the slice space in [2] and the Wiener-amalgam space in [1]. They also introduced the Orlicz-slice (local) Hardy spaces and developed a complete real-variable theory of these function spaces in [66, 67]. Next, we recall the definition of anisotropic Orlicz-slice spaces.

Definition 5.8. Let $\ell \in \mathbb{Z}$, $q \in (0, \infty)$, and Φ be an Orlicz function. The *anisotropic Orlicz-slice space* (*E q* ${}_{\Phi}^{q}$)_{ℓ ,*A*} (\mathbb{R}^{n}) is defined to be the set of all the measurable functions *f* on \mathbb{R}^{n} such that

$$
||f||_{(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)} := \left\{ \int_{\mathbb{R}^n} \left[\frac{||f \mathbf{1}_{x+B_\ell}||_{L^\Phi(\mathbb{R}^n)}}{||\mathbf{1}_{x+B_\ell}||_{L^\Phi(\mathbb{R}^n)}} \right]^q dx \right\}^{\frac{1}{q}} < \infty,
$$

where B_{ℓ} is the same as in (2.2) with *k* replaced by ℓ .

Fourier Transform of Anisotropic Hardy Spaces 25 and 25 and 25 and 25 and 25 and 26 and 26 anisotropic Hardy Spaces 25 and 25 and 2012 and 20

Let $\ell \in \mathbb{Z}$, $q \in (0, 1)$, Φ be an Orlicz function with positive lower type p_{Φ}^- and positive upper type p_{Φ}^+ satisfying $0 < p_{\Phi}^- \le p_{\Phi}^+ < 1$, and

$$
N \in \mathbb{N} \cap \left[\left| \left(\frac{1}{\min\{p_{\Phi}^-, q\}} - 1 \right) \frac{\ln b}{\ln(\lambda_-)} \right| + 2, \infty \right).
$$

Then, by [59, Remark 8.14], we conclude that (E_q^q) ${}_{\Phi}^{q}$ $)$ _{ℓ} \mathcal{A} (\mathbb{R}^{n}) satisfies all the assumptions of Definition 2.10 with $X := (E_d^q)$ θ_{Φ}^{q}) $\ell_{A}(\mathbb{R}^{n})$, $p_{-} \in (0, \min\{p_{\Phi}^{-}, q\}], \theta_{0} \in (0, p_{-})$, and $p_{0} \in (\max\{p_{\Phi}^{+}, q\}, \infty)$. Moreover, choose a $q_0 = 1$. On the one hand, from [67, Lemma 5.4], we infer that, for any non-negative measurable functions $\{f_k\}_{k=1}^{\infty}$,

$$
\sum_{k=1}^\infty\|f_k\|_{[(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)]^{\frac{1}{q_0}}}\le \left\|\sum_{k=1}^\infty f_k\right\|_{[(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)]^{\frac{1}{q_0}}}
$$

.

On the other hand, we have, for any $B \in \mathcal{B}$,

(5.2)
$$
||1_B||_{(E^q_{\Phi})_{\ell,A}(\mathbb{R}^n)} \gtrsim \min \{ |B|, |B|^{\frac{1}{\theta_0}} \}.
$$

Indeed, for any $B \in \mathcal{B}$ with $|B| \geq |B_{\ell}|$,

(5.3)
$$
||\mathbf{1}_B||_{(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)} = \left\{ \int_{\mathbb{R}^n} \left[\frac{||\mathbf{1}_B \mathbf{1}_{x+B_\ell}||_{L^\Phi(\mathbb{R}^n)}}{||\mathbf{1}_{x+B_\ell}||_{L^\Phi(\mathbb{R}^n)}} \right]^q dx \right\}^{\frac{1}{q}}
$$

$$
\geq \left(\int_B 1 dx \right)^{\frac{1}{q}} = |B|^{\frac{1}{q}}.
$$

On the other hand, for any $x_0 \in \mathbb{R}^n$, $k \in \mathbb{Z}$ with $|B_k| \leq |B_\ell|$, $x \in B(x_0, \lambda_\ell^{\ell})$, and $\eta \in (0, p_0^{-})$, by [61, Remark 4.21(iv)], we conclude that $L^{\Phi}(\mathbb{R}^n)$ satisfies Assumption 2.7 with $X := L^{\Phi}(\mathbb{R}^n)$, $u := 1/\eta$, and $p := \eta$. Thus, we obtain

$$
||\mathbf{1}_{x_0+B_k}||_{L^{\Phi}(\mathbb{R}^n)}^{\eta} \gtrsim ||[\mathcal{M}(\mathbf{1}_{x_0+B_k})]^{1/\eta}||_{L^{\Phi}(\mathbb{R}^n)}^{\eta}.
$$

For any $y \in x + B_\ell$, we have

(5.5)
$$
\mathcal{M}(\mathbf{1}_{x_0+B_k})(y) \geq \frac{1}{|B_\ell|} \int_{x+B_\ell} \mathbf{1}_{x_0+B_k}(z) dz \gtrsim \frac{|B_k|}{|B_\ell|}.
$$

Combining (5.4) and (5.5), we conclude that

$$
\|{\bf 1}_{x_0+B_k}\|_{L^{\Phi}(\mathbb{R}^n)}^{\eta} \gtrsim \left\| \left[\frac{|B_k|}{|B_{\ell}|}({\bf 1}_{x+B_{\ell}})\right]^{1/\eta} \right\|_{L^{\Phi}(\mathbb{R}^n)}^{\eta} \gtrsim |B_k| \, \|{\bf 1}_{x+B_{\ell}}\|_{L^{\Phi}(\mathbb{R}^n)}^{\eta}.
$$

Therefore, we obtain

(5.6)
$$
||\mathbf{1}_{x_0+B_k}||_{(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)} = \left\{ \int_{\mathbb{R}^n} \left[\frac{||\mathbf{1}_{x_0+B_k} \mathbf{1}_{x+B_\ell}||_{L^\Phi(\mathbb{R}^n)}}{||\mathbf{1}_{x+B_\ell}||_{L^\Phi(\mathbb{R}^n)}} \right]^q dx \right\}^{\frac{1}{q}}
$$

$$
\geq \left\{ \int_{B(x_0,\lambda_-^{\ell})} \left[\frac{\|{\bf 1}_{x_0+B_k}\|_{L^{\Phi}(\mathbb{R}^n)}}{\|{\bf 1}_{x+B_\ell}\|_{L^{\Phi}(\mathbb{R}^n)}} \right]^q dx \right\}^{\frac{1}{q}}.
$$

$$
\geq |B_k|^{1/\eta} \left\{ \int_{B(x_0,\lambda_-^{\ell})} 1 dx \right\}^{\frac{1}{q}} \sim |B_k|^{1/\eta}.
$$

By (5.3) and (5.6), we find that, for any $B \in \mathcal{B}$,

$$
||\mathbf{1}_B||_{(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)} \gtrsim |B|^{1/q} \gtrsim |B| \text{ if } |B| \ge |B_\ell|
$$

and

$$
||1_B||_{(E^q_\Phi)_{\ell,A}(\mathbb{R}^n)} \gtrsim |B|^{1/p_\Phi^-} \gtrsim |B|^{1/\theta_0} \text{ if } |B| \leq |B_\ell|.
$$

This finishes the proof of (5.2).

Therefore, all the assumptions of Theorems 3.1, 4.1, and 4.2 are satisfied with *X* := (E_q^q) ${}_{\Phi}^{q}$ $)_{\ell,A}$ (\mathbb{R}^{n}). Applying Theorems 3.1, 4.1, and 4.2, we obtain the following conclusion.

Theorem 5.9. *Let* $\ell \in \mathbb{Z}$, $q \in (0, 1)$, and Φ *be an Orlicz function with lower type* p_{Φ}^- *and upper type* p_{Φ}^+ satisfying $0 \le p_{\Phi}^- \le p_{\Phi}^+ \le 1$. Then Theorems 3.1, 4.1, and 4.2 still hold true with X *replaced by* (E_d^q) $_{\Phi}^{q}$ $)_{\ell,A}$ (\mathbb{R}^{n}).

Remark 5.10. We point out that Theorem 5.9 even when $A := 2I_{n \times n}$ is completely new.

5.5 Local Generalized Herz Spaces

In what follows, we always let $\mathbb{R}_+ := (0, \infty)$. A nonnegative function ω on \mathbb{R}_+ is said to be *almost increasing* (resp. *almost decreasing*) on \mathbb{R}_+ if there exists a constant $C \in [1, \infty)$ such that, for any $s, t \in \mathbb{R}_+$ satisfying $s \le t$ (resp. $s \ge t$),

$$
\omega(s) \leq C \omega(t)
$$

(see, for instance, [37, 42]). Now, we recall the concept of the function class $M(\mathbb{R}_+)$ as follows (see, for instance, [37, 51]).

Definition 5.11. The *function class* $M(\mathbb{R}_+)$ is defined to be the set of all the positive functions ω on \mathbb{R}_+ such that, for any $0 < \delta_1 < \delta_2 < \infty$,

$$
0 < \inf_{t \in (\delta_1, \delta_2)} \omega(t) \leq \sup_{t \in (\delta_1, \delta_2)} \omega(t) < \infty
$$

and there exist four constants $\alpha_0, \beta_0, \alpha_\infty, \beta_\infty \in \mathbb{R}$ such that

- (i) for any $t \in (0, 1]$, $\omega(t)t^{-\alpha_0}$ is almost increasing and $\omega(t)t^{-\beta_0}$ is almost decreasing;
- (ii) for any $t \in [1, \infty)$, $\omega(t)t^{-\alpha_{\infty}}$ is almost increasing and $\omega(t)t^{-\beta_{\infty}}$ is almost decreasing.

Next, we introduce anisotropic local generalized Herz spaces as follows.

Fourier Transform of Anisotropic Hardy Spaces 27 and 27

Definition 5.12. Let *A* be a dilation, $p, q \in (0, \infty]$, and $\omega \in M(\mathbb{R}_+)$. The *anisotropic local gener*alized Herz space $\dot{\mathcal{K}}^{p,q}_{A,\omega}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$ is defined to be the set of all $f \in L^p_{loc}(\mathbb{R}^n \setminus \{0\})$ such that

$$
\|f\|_{\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n)} := \left\{\sum_{k\in\mathbb{Z}}\left[\omega\left(b^k\right)\right]^q\left\|f\mathbf{1}_{B_k\setminus B_{k-1}}\right\|_{L^p(\mathbb{R}^n)}^q\right\}^{\frac{1}{q}} < \infty
$$

with the usual modification made when $q = \infty$, where *b* is the same as in (2.1) and, for any $k \in \mathbb{Z}$, B_k the same as in (2.2).

Now, we introduce the following Matuszewska–Orlicz indices (see, for instance, [48, 49]).

Definition 5.13. Let ω be a positive function on \mathbb{R}_+ . Then the *Matuszewska–Orlicz indices* $m_0(\omega)$, $M_0(\omega)$, $m_\infty(\omega)$, and $M_\infty(\omega)$ of ω are defined, respectively, by setting, for any $h \in (0, \infty)$,

$$
m_0(\omega) := \sup_{t \in (0,1)} \frac{\ln(\overline{\lim}_{h \to 0^+} \frac{\omega(ht)}{\omega(h)})}{\ln t}, \ M_0(\omega) := \inf_{t \in (0,1)} \frac{\ln(\frac{\lim}{\lim}_{h \to 0^+} \frac{\omega(ht)}{\omega(h)})}{\ln t},
$$

$$
m_{\infty}(\omega) := \sup_{t \in (1,\infty)} \frac{\ln(\frac{\lim}{\lim}_{h \to \infty} \frac{\omega(ht)}{\omega(h)})}{\ln t}, \ \text{and } M_{\infty}(\omega) := \inf_{t \in (1,\infty)} \frac{\ln(\overline{\lim}_{h \to \infty} \frac{\omega(ht)}{\omega(h)})}{\ln t}.
$$

The following property about the function class $M(\mathbb{R}_+)$ and the Matuszewska–Orlicz indices can be found in $[42,$ Lemma 1.1.6] (see also $[52, (6.4), (6.5),$ and $(6.14)]$). In what follows, for any $\omega \in M(\mathbb{R}_+)$ and $t \in \mathbb{R}_+$, $\omega^s(t) := [\omega(t)]^s$.

Lemma 5.14. *Let* $\omega \in M(\mathbb{R}_+)$ *. Then, for any given* $t \in (0, \infty)$ *, it holds true that* $1/\omega, \omega^t \in M(\mathbb{R}_+)$ *and*

- (i) $m_0(1/\omega) = -M_0(\omega)$ and $M_0(1/\omega) = -m_0(\omega)$;
- (ii) $m_{\infty}(1/\omega) = -M_{\infty}(\omega)$ and $M_{\infty}(1/\omega) = -m_{\infty}(\omega)$;
- (iii) $m_0(\omega^t) = tm_0(\omega)$ *and* $M_0(\omega^t) = tM_0(\omega)$;
- (iv) $m_{\infty}(\omega^t) = tm_{\infty}(\omega)$ and $M_{\infty}(\omega^t) = tM_{\infty}(\omega)$.

Remark 5.15. Let $\omega \in M(\mathbb{R}_+)$. Then, by [42, Remark 1.1.5(ii)], we conclude that

$$
-\infty < m_0(\omega) \le M_0(\omega) < \infty
$$

and

$$
-\infty < m_{\infty}(\omega) \leq M_{\infty}(\omega) < \infty.
$$

The following theorem shows that anisotropic local generalized Herz spaces $\dot{K}^{p,q}_{A,\omega}$ $A^{p,q}_{A,\omega}(\mathbb{R}^n)$ are ball quasi-Banach function spaces under some additional assumptions on the exponent $\omega \in M(\mathbb{R}_+)$; see [42, Theorem 1.2.42] for the standard Euclidean space case.

Theorem 5.16. *Let A be a dilation, p, q* \in (0, ∞)*, and* $\omega \in M(\mathbb{R}_+)$ *with* $m_0(\omega) \in (-\frac{1}{p}, \infty)$ *. Then* the anisotropic local generalized Herz space $\dot{\mathcal{K}}^{p,q}_{A,\omega}$ $\mathcal{L}_{A,\omega}^{p,q}(\mathbb{R}^n)$ *is a ball quasi-Banach function space.*

Proof. Indeed, by an argument similar to that used in the proof of [42, Theorem 1.2.38], we find that $\dot{K}^{p,q}_{A,\omega}$ $P_{A,\omega}^{p,q}(\mathbb{R}^n)$ is a quasi-Banach space satisfying (i), (ii), and (iii) of Definition 2.4. Next, we prove that $\dot{K}^{p,q}_{A,\omega}$ $P^{q}(R^n)$ satisfies Definition 2.4(iv). To this end, let $x_0 \in \mathbb{R}^n$, $k_0 \in \mathbb{Z}$, and B_{k_0} be the same as in (2.2) . Then

$$
(5.7) \t\t || \mathbf{1}_{x_0 + B_{k_0}} ||_{\dot{K}^{p,q}_{A,\omega}(\mathbb{R}^n)} = \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^k \right) \right]^q \left\| \mathbf{1}_{x_0 + B_{k_0}} \mathbf{1}_{B_k \setminus B_{k-1}} \right\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}} \n\leq \left\{ \sum_{k \in \mathbb{Z} \setminus \mathbb{N}} \left[\omega \left(b^k \right) \right]^q \left\| \mathbf{1}_{x_0 + B_{k_0}} \mathbf{1}_{B_k \setminus B_{k-1}} \right\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}} + \left\{ \sum_{k \in \mathbb{N}} \cdots \right\}^{\frac{1}{q}}
$$
\n= : I + II.

We first deal with I. From [42, Lemma 1.1.12], we deduce that, for any $k \in \mathbb{Z}\backslash\mathbb{N}$,

$$
\omega\left(b^k\right) \lesssim b^{k[m_0(\omega)-\varepsilon]},
$$

where $\varepsilon \in (0, m_0(\omega) + \frac{1}{p})$ is a fixed positive constant. This, combined with (5.7), further implies that

$$
(5.8) \t I \lesssim \left\{ \sum_{k \in \mathbb{Z} \backslash \mathbb{N}} b^{kq[m_0(\omega)-\varepsilon]} \| 1_{B_k \backslash B_{k-1}} \|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

$$
\lesssim \left\{ \sum_{k \in \mathbb{Z} \backslash \mathbb{N}} b^{kq[m_0(\omega)+\frac{1}{p}-\varepsilon]} \right\}^{\frac{1}{q}} < \infty.
$$

Next, we deal with II. To this end, we first claim that, for any $k \in \mathbb{N} \cap (\tau + |k_0| + |[\log_b \rho(x_0)]| + 3, \infty)$ with τ in (2.4),

$$
(5.9) \qquad \qquad (x_0 + B_{k_0}) \cap (B_k \backslash B_{k-1}) = \emptyset.
$$

Indeed, by the inequality in line 26 of [4, p. 7], we conclude that, for any $y \in x_0 + B_{k_0}$,

$$
\rho(y) \le b^{\tau} \left[\rho(x_0 - y) + \rho(x_0) \right] < b^{\tau} \left[b^{k_0 - 1} + b^{\lfloor \log_b \rho(x_0) \rfloor + 1} \right] \\
 \le b^{\tau + |k_0| + |\lfloor \log_b \rho(x_0) \rfloor + 1} < b^{k - 2},
$$

which further implies *y* ∈ *B*_{*k*−1} and hence $x_0 + B_{k_0} \subset B_{k-1}$. Thus, (5.9) holds true. From (5.7) and (5.9), we deduce that

$$
\mathrm{II} = \left\{\sum_{k \in \mathbb{N} \cap [1,\tau + |k_0| + |\mathrm{llog}_b \rho(x_0)|] + 3]} \left[\omega \left(b^k\right)\right]^q \left\| \mathbf{1}_{x_0 + B_{k_0}} \mathbf{1}_{B_k \setminus B_{k-1}} \right\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}} < \infty.
$$

This, together with (5.7) and (5.8) , then finishes the proof of Theorem 5.16.

The following lemma gives the Fefferman–Stein vector-valued inequality on anisotropic local generalized Herz spaces; see [42, Theorem 1.6.1] for the standard Euclidean space case.

Lemma 5.17. *Let A be a dilation, p, r* \in (1, ∞)*, q* \in (0, ∞)*, and* $\omega \in M(\mathbb{R}_+)$ *satisfy*

$$
-\frac{1}{p}
$$

and

$$
-\frac{1}{p} < m_{\infty}(\omega) \le M_{\infty}(\omega) < \frac{1}{p'},
$$

where $\frac{1}{p} + \frac{1}{p'} = 1$ *. Then there exists a positive constant C such that, for any* $\{f_j\}_{j\in\mathbb{N}} \subset L^1_{loc}(\mathbb{R}^n)$ *,*

$$
\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(f_j) \right]^r \right\}^{\frac{1}{r}} \right\|_{\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n)} \leq C \left\| \left(\sum_{j \in \mathbb{N}} |f_j|^r \right)^{\frac{1}{r}} \right\|_{\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n)}
$$

.

To prove this lemma, we need more preparations. Recall that an operator T defined on $\mathcal{M}(\mathbb{R}^n)$ is called a *sublinear operator* if, for any $f, g \in \mathcal{M}(\mathbb{R}^n)$ and $\lambda \in \mathbb{C}$,

$$
|T(f+g)|\leq |T(f)|+|T(g)|
$$

and

$$
|T(\lambda f)| = |\lambda||T(f)|.
$$

Moreover, for any normed linear space *X* and any operator *T* on *X*, the *operator norm* $||T||_{X\to X}$ of *T* is defined by setting

$$
||T||_{X\to X} := \sup_{\{x\in X:\;||x||_X=1\}} ||T(x)||_X.
$$

The following lemma is a boundedness criterion of sublinear operators on anisotropic local generalized Herz spaces; see [42, Theorem 1.5.1] for the standard Euclidean space case.

Lemma 5.18. *Let A be a dilation, p* ∈ (1, ∞)*, q* ∈ (0, ∞)*, and* $ω ∈ M(\mathbb{R}_+)$ *satisfy*

$$
-\frac{1}{p} < m_0(\omega) \le M_0(\omega) < \frac{1}{p'}
$$

and

$$
-\frac{1}{p} < m_{\infty}(\omega) \le M_{\infty}(\omega) < \frac{1}{p'},
$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. Assume that T is a sublinear operator satisfying that T is bounded on $L^p(\mathbb{R}^n)$ *and that there exists a positive constant* C_0 *such that, for any* $f \in \dot{K}^{p,q}_{A,\omega}$ $\lim_{A,\omega} \mathbb{R}^n$ *and* $x \notin \overline{\text{supp}(f)} :=$ ${x \in \mathbb{R}^n : f(x) \neq 0},$

(5.10)
$$
|T(f)(x)| \leq C_0 \int_{\mathbb{R}^n} \frac{|f(y)|}{\rho(x - y)} dy,
$$

where ρ *is the same as in Definition 2.3. Then there exists a positive constant C, independent of T*, such that, for any $f \in \mathcal{K}_{A,\omega}^{p,q}$ $A^{p,q}_{A,\omega}(\mathbb{R}^n)$,

$$
(5.11) \t\t ||T(f)||_{\dot{K}^{p,q}_{A,\omega}(\mathbb{R}^n)} \leq C \left[C_0 + ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)}\right] ||f||_{\dot{K}^{p,q}_{A,\omega}(\mathbb{R}^n)}.
$$

Proof. Let $f \in \mathcal{K}_{A,\omega}^{p,q}$ $\chi_{A,\omega}^{p,q}(\mathbb{R}^n)$. For any given $k \in \mathbb{Z}$, let

$$
f_{k,1} := f \mathbf{1}_{B_{k-\tau-1}}, \ f_{k,2} := f \mathbf{1}_{B_{k+\tau+2} \setminus B_{k-\tau-1}}, \text{ and } f_{k,3} := f \mathbf{1}_{[B_{k+\tau+2}]}[c,
$$

where τ is the same as in (2.4). Obviously, for any $k \in \mathbb{Z}$, it holds true that

$$
f = f_{k,1} + f_{k,2} + f_{k,3}.
$$

From this, Definition 5.12, the sublinearity of *T*, and the Minkowski inequality, we deduce that

(5.12)
$$
||T(f)||_{\dot{K}_{A,\omega}^{p,q}(\mathbb{R}^{n})} = \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^{k} \right) \right]^{q} ||T(f)1_{B_{k} \setminus B_{k-1}}||_{L^{p}(\mathbb{R}^{n})}^{q} \right\}^{\frac{1}{q}} \n\leq \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^{k} \right) \right]^{q} ||T(f_{k,1})1_{B_{k} \setminus B_{k-1}}||_{L^{p}(\mathbb{R}^{n})}^{q} \right\}^{\frac{1}{q}} \n+ \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^{k} \right) \right]^{q} ||T(f_{k,2})1_{B_{k} \setminus B_{k-1}}||_{L^{p}(\mathbb{R}^{n})}^{q} \right\}^{\frac{1}{q}} \n+ \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^{k} \right) \right]^{q} ||T(f_{k,3})1_{B_{k} \setminus B_{k-1}}||_{L^{p}(\mathbb{R}^{n})}^{q} \right\}^{\frac{1}{q}} \n=:\left[1_{1} + I_{2} + I_{3},\n\right\}
$$

where the implicit positive constant is independent of both T and f . Next, we deal with I_1 , I_2 , and I³ successively. To this end, let

(5.13)
$$
\varepsilon \in \left(0, \min\left\{\min\{m_0(\omega), m_\infty(\omega)\} + \frac{1}{p}, \frac{1}{p'} - \max\{M_0(\omega), M_\infty(\omega)\}\right\}\right)
$$

be a fixed positive constant. Then, by [42, Lemma 1.5.2], we find that, for any $0 < s < t < \infty$,

(5.14)
$$
\frac{\omega(s)}{\omega(t)} \lesssim \left(\frac{s}{t}\right)^{\min\{m_0(\omega), m_\infty(\omega)\}-\varepsilon}
$$

and

(5.15)
$$
\frac{\omega(t)}{\omega(s)} \lesssim \left(\frac{t}{s}\right)^{\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon},
$$

where the implicit positive constant is independent of both *T* and *f* .

We first deal with I₁. From the inequality in line 26 of [4, p. 7], we infer that, for any $k, i \in \mathbb{Z}$ satisfying $i \in (-\infty, k - \tau - 1]$ and for any $x, y \in \mathbb{R}^n$ satisfying $x \in B_k \setminus B_{k-1}$ and $y \in B_i \setminus B_{i-1}$,

$$
\rho(x - y) \ge b^{-\tau} \rho(x) - \rho(y) = b^{k - \tau - 1} - b^{i - 1}
$$

$$
\ge b^{k - \tau - 1} - b^{k - \tau - 2} = (b^{-\tau - 1} - b^{-\tau - 2})b^k.
$$

By this, (5.10), and the Hölder inequality, we conclude that, for any $k \in \mathbb{Z}$ and $x \in B_k \setminus B_{k-1}$,

$$
(5.16) \t\t |T(f_{k,1})(x)| \leq C_0 \int_{\mathbb{R}^n} \frac{|f_{k,1}(y)|}{\rho(x-y)} dy = C_0 \sum_{i=-\infty}^{k-\tau-1} \int_{B_i \setminus B_{i-1}} \frac{|f(y)|}{\rho(x-y)} dy
$$

$$
\leq C_0 \sum_{i=-\infty}^{k-\tau-1} b^{-k+\frac{i}{p'}} ||f\mathbf{1}_{B_i \setminus B_{i-1}}||_{L^p(\mathbb{R}^n)},
$$

where the implicit positive constant is independent of both T and f . Moreover, from (5.15), it follows that, for any $k, i \in \mathbb{Z}$ satisfying $i \in (-\infty, k - \tau - 1]$,

$$
\frac{\omega(b^k)}{\omega(b^i)} \lesssim b^{(k-i)[\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon]},
$$

which, combined with (5.16), further implies that, for any $k \in \mathbb{Z}$,

$$
(5.17) \qquad \omega\left(b^{k}\right) \| T(f_{k,1}) \mathbf{1}_{B_{k} \setminus B_{k-1}} \|_{L^{p}(\mathbb{R}^{n})}
$$
\n
$$
\leq C_{0} \omega\left(b^{k}\right) \sum_{i=-\infty}^{k-\tau-1} b^{-k+\frac{i}{p'}} \| f \mathbf{1}_{B_{i} \setminus B_{i-1}} \|_{L^{p}(\mathbb{R}^{n})} \| \mathbf{1}_{B_{k} \setminus B_{k-1}} \|_{L^{p}(\mathbb{R}^{n})}
$$
\n
$$
\leq C_{0} \sum_{i=-\infty}^{k-\tau-1} b^{\frac{1}{p'}(i-k)} \frac{\omega(b^{k})}{\omega(b^{i})} \omega\left(b^{i}\right) \| f \mathbf{1}_{B_{i} \setminus B_{i-1}} \|_{L^{p}(\mathbb{R}^{n})}
$$
\n
$$
\leq C_{0} \sum_{i=-\infty}^{k-\tau-1} b^{(k-i)[\max\{M_{0}(\omega), M_{\infty}(\omega)\}+\varepsilon-\frac{1}{p'}]} \omega\left(b^{i}\right) \| f \mathbf{1}_{B_{i} \setminus B_{i-1}} \|_{L^{p}(\mathbb{R}^{n})},
$$

where the implicit positive constants are independent of both *T* and *f*. Next, we show the desired estimate of I_1 by considering the following two cases on q .

Case 1) *q* ∈ (0, 1]. In this case, by (5.12), (5.17), Lemma 3.8, and (5.13), we conclude that

$$
(5.18) \tI1 \leq C_0 \left\{ \sum_{k \in \mathbb{Z}} \sum_{i=-\infty}^{k-\tau-1} b^{q(k-i)[\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon-\frac{1}{p'}]}\left[\omega(b^i)\right]^q \left\|f\mathbf{1}_{B_i\setminus B_{i-1}}\right\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

\n
$$
= C_0 \left\{ \sum_{i\in \mathbb{Z}} \sum_{k=i+\tau+1}^{\infty} b^{q(k-i)[\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon-\frac{1}{p'}]}\left[\omega(b^i)\right]^q \left\|f\mathbf{1}_{B_i\setminus B_{i-1}}\right\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

\n
$$
\sim C_0 \left\{ \sum_{i\in \mathbb{Z}} \left[\omega(b^i)\right]^q \left\|f\mathbf{1}_{B_i\setminus B_{i-1}}\right\|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

\n
$$
\leq [C_0 + ||T||_{L^p(\mathbb{R}^n)\to L^p(\mathbb{R}^n)}] ||f||_{\dot{K}^{p,q}_{A,\omega}(\mathbb{R}^n)},
$$

where the implicit positive constants are independent of both *T* and *f* .

Case 2) $q \in (1, \infty]$. In this case, from (5.17), the Hölder inequality, and (5.13), we deduce that, for any $k \in \mathbb{Z}$,

$$
(5.19) \t\t \omega(b^k)\|T(f_{k,1})\mathbf{1}_{B_k\setminus B_{k-1}}\|_{L^p(\mathbb{R}^n)}
$$

$$
\leq C_0 \left\{ \sum_{i=-\infty}^{k-\tau-1} b^{\frac{(k-i)q'}{2} [\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon-\frac{1}{p'}]}\right\}^{\frac{1}{q'}} \times \left\{ \sum_{i=-\infty}^{k-\tau-1} b^{\frac{(k-i)q}{2} [\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon-\frac{1}{p'}]}\left[\omega(b^i)\right]^q \left\|f\mathbf{1}_{B_i\setminus B_{i-1}}\right\|_{L^p(\mathbb{R}^n)}^q\right\}^{\frac{1}{q}} \times C_0 \left\{ \sum_{i=-\infty}^{k-\tau-1} b^{\frac{(k-i)q}{2} [\max\{M_0(\omega), M_\infty(\omega)\}+\varepsilon-\frac{1}{p'}]}\left[\omega(b^i)\right]^q \left\|f\mathbf{1}_{B_i\setminus B_{i-1}}\right\|_{L^p(\mathbb{R}^n)}^q\right\}^{\frac{1}{q}},
$$

which, together with (5.12) and (5.13) , further implies that

$$
(5.20) \tI1 \leq C_0 \left\{ \sum_{k \in \mathbb{Z}} \sum_{i=-\infty}^{k-\tau-1} b^{\frac{(k-i)q}{2} [\max\{M_0(\omega), M_\infty(\omega)\} + \varepsilon - \frac{1}{p'}] } [\omega(b^i)]^q \| f \mathbf{1}_{B_i \setminus B_{i-1}} \|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

\n
$$
= C_0 \left\{ \sum_{i \in \mathbb{Z}} \sum_{k=i+\tau+1}^{\infty} b^{\frac{(k-i)q}{2} [\max\{M_0(\omega), M_\infty(\omega)\} + \varepsilon - \frac{1}{p'}] } [\omega(b^i)]^q \| f \mathbf{1}_{B_i \setminus B_{i-1}} \|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

\n
$$
\sim C_0 \left\{ \sum_{i \in \mathbb{Z}} [\omega(b^i)]^q \| f \mathbf{1}_{B_i \setminus B_{i-1}} \|_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

\n
$$
\leq [C_0 + ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)}] \|f\|_{\dot{K}_{A,\omega}^{p,q}(\mathbb{R}^n)},
$$

where the implicit positive constants are independent of both *T* and *f* .

Second, we deal with I_2 . By the boundedness of *T* on $L^p(\mathbb{R}^n)$ and the Minkowski inequality, we find that, for any $k \in \mathbb{Z}$,

$$
(5.21) \t\t ||T(f_{k,2})\mathbf{1}_{B_k\setminus B_{k-1}}||_{L^p(\mathbb{R}^n)} \leq ||T(f_{k,2})||_{L^p(\mathbb{R}^n)} \leq ||T||_{L^p(\mathbb{R}^n)\to L^p(\mathbb{R}^n)} ||f_{k,2}||_{L^p(\mathbb{R}^n)}
$$

$$
\leq ||T||_{L^p(\mathbb{R}^n)\to L^p(\mathbb{R}^n)} \sum_{i=-\tau}^{\tau+2} ||f\mathbf{1}_{B_{k+i}\setminus B_{k+i-1}}||_{L^p(\mathbb{R}^n)}.
$$

Moreover, from [42, Lemma 1.1.3], it follows that, for any $k \in \mathbb{Z}$ and $i \in [-\tau, \tau + 2] \cap \mathbb{Z}$,

$$
\omega\left(b^k\right)\sim \omega\left(b^{k+i}\right),\,
$$

which, combined with (5.12) and (5.21) , further implies that

$$
(5.22) \qquad I_2 \leq ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)} \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^k \right) \right]^q \sum_{i=-\tau}^{\tau+2} ||f \mathbf{1}_{B_{k+i} \setminus B_{k+i-1}}||_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}} \n\leq ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)} \sum_{i=-\tau}^{\tau+2} \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^{k+i} \right) \right]^q ||f \mathbf{1}_{B_{k+i} \setminus B_{k+i-1}}||_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}} \n\sim ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)} \left\{ \sum_{k \in \mathbb{Z}} \left[\omega \left(b^k \right) \right]^q ||f \mathbf{1}_{B_k \setminus B_{k-1}}||_{L^p(\mathbb{R}^n)}^q \right\}^{\frac{1}{q}}
$$

Fourier Transform of Anisotropic Hardy Spaces 33

$$
\leq [C_0 + ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)}] ||f||_{\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n)}.
$$

Finally, we deal with I_3 . By the inequality in line 26 of $[4, p.7]$, we conclude that, for any $k, i \in \mathbb{Z}$ satisfying $i \in [k + \tau + 3, \infty)$ and for any $x, y \in \mathbb{R}^n$ satisfying $x \in B_k \setminus B_{k-1}$ and $y \in B_i \setminus B_{i-1}$,

$$
\rho(x - y) \ge b^{-\tau} \rho(y) - \rho(x) = b^{i - \tau - 1} - b^{k - 1}
$$

$$
\ge b^{i - \tau - 1} - b^{i - \tau - 4} = (b^{-\tau - 1} - b^{-\tau - 4})b^{i}.
$$

From this, (5.10), and the Hölder inequality, we infer that, for any $k \in \mathbb{Z}$ and $x \in B_k \setminus B_{k-1}$,

$$
(5.23) \t\t |T(f_{k,3})(x)| \leq C_0 \int_{\mathbb{R}^n} \frac{|f_{k,3}(y)|}{\rho(x-y)} dy = C_0 \sum_{i=k+\tau+3}^{\infty} \int_{B_i \setminus B_{i-1}} \frac{|f(y)|}{\rho(x-y)} dy
$$

$$
\lesssim C_0 \sum_{i=k+\tau+3}^{\infty} b^{-\frac{i}{p}} ||f\mathbf{1}_{B_i \setminus B_{i-1}}||_{L^p(\mathbb{R}^n)},
$$

where the implicit positive constant is independent of both T and f . Moreover, by (5.14), we conclude that, for any $k, i \in \mathbb{Z}$ satisfying $i \in [k + \tau + 3, \infty)$,

$$
\frac{\omega(b^k)}{\omega(b^i)} \lesssim b^{(k-i)[\min\{m_0(\omega), m_\infty(\omega)\}-\varepsilon]},
$$

which, combined with (5.23), further implies that, for any $k \in \mathbb{Z}$,

$$
(5.24) \qquad \omega\left(b^{k}\right) \| T(f_{k,3}) \mathbf{1}_{B_{k} \setminus B_{k-1}} \|_{L^{p}(\mathbb{R}^{n})}
$$
\n
$$
\leq C_{0} \omega\left(b^{k}\right) \sum_{i=k+\tau+3}^{\infty} b^{-\frac{i}{p}} \| f \mathbf{1}_{B_{i} \setminus B_{i-1}} \|_{L^{p}(\mathbb{R}^{n})} \| \mathbf{1}_{B_{k} \setminus B_{k-1}} \|_{L^{p}(\mathbb{R}^{n})}
$$
\n
$$
\leq C_{0} \sum_{i=k+\tau+3}^{\infty} b^{\frac{1}{p}(k-i)} \frac{\omega(b^{k})}{\omega(b^{i})} \omega\left(b^{i}\right) \| f \mathbf{1}_{B_{i} \setminus B_{i-1}} \|_{L^{p}(\mathbb{R}^{n})}
$$
\n
$$
\leq C_{0} \sum_{i=k+\tau+3}^{\infty} b^{(k-i)[\min\{m_{0}(\omega), m_{\infty}(\omega)\} - \varepsilon + \frac{1}{p}]} \omega\left(b^{i}\right) \| f \mathbf{1}_{B_{i} \setminus B_{i-1}} \|_{L^{p}(\mathbb{R}^{n})},
$$

where the implicit positive constants are independent of both T and f . From (5.24), (5.13), and an argument similar to that used in the estimations of (5.18) and (5.20), we deduce that

$$
I_3 \lesssim [C_0 + ||T||_{L^p(\mathbb{R}^n) \to L^p(\mathbb{R}^n)}] ||f||_{\dot{K}^{p,q}_{A,\omega}(\mathbb{R}^n)},
$$

where the implicit positive constant is independent of both *T* and *f*. This, together with (5.12), (5.18) , (5.20) , and (5.22) , further implies (5.11) and hence finishes the proof of Lemma 5.18. \Box

The following conclusion is a simple corollary of both (2.5) and Lemma 5.18.

Corollary 5.19. *Let A, p, q, and* ω *be the same as in Lemma 5.18 and* M *the Hardy–Littlewood maximal operator in* (2.5). Then there exists a positive constant C such that, for any $f \in \dot{\mathcal{K}}_{A,\omega}^{p,q}$ $A^{p,q}_{A,\omega}(\mathbb{R}^n)$,

$$
\|\mathcal{M}(f)\|_{\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n)} \leq C \|f\|_{\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n)}.
$$

Now, we show Lemma 5.17.

Proof of Lemma 5.17. Let $\{f_j\}_{j\in\mathbb{N}} \subset L^1_{loc}(\mathbb{R}^n)$ be any given sequence and, for any $r \in (1, \infty]$, $g \in \mathcal{M}(\mathbb{R}^n)$, and $x \in \mathbb{R}^n$, define

$$
A(g)(x):=\left\{\sum_{j\in\mathbb{N}}\left[\mathcal{M}(g\xi_j)(x)\right]^r\right\}^{\frac{1}{r}},
$$

where, for any $j \in \mathbb{N}$ and $y \in \mathbb{R}^n$,

$$
\xi_j(y):=\frac{f_j(y)}{[\sum_{j\in\mathbb{N}}|f_j(y)|^r]^{1/r}}\text{ if }\left[\sum_{j\in\mathbb{N}}\left|f_j(y)\right|^r\right]^{1/r}\neq 0
$$

and $\xi_j(y) := 0$ otherwise. Obviously, for any $\lambda \in \mathbb{C}$ and $g \in \mathcal{M}(\mathbb{R}^n)$,

$$
A(\lambda g) = |\lambda| A(g).
$$

Moreover, by the Minkowski inequality, we find that, for any $g_1, g_2 \in \mathcal{M}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$
A(g_1 + g_2)(x) = \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M} \left([g_1 + g_2] \xi_j \right)(x) \right]^r \right\}^{\frac{1}{r}}
$$

\n
$$
\leq \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M} (g_1 \xi_j)(x) + \mathcal{M} (g_2 \xi_j)(x) \right]^r \right\}^{\frac{1}{r}}
$$

\n
$$
\leq \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M} (g_1 \xi_j)(x) \right]^r \right\}^{\frac{1}{r}} + \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M} (g_2 \xi_j)(x) \right]^r \right\}^{\frac{1}{r}}
$$

\n
$$
= A(g_1)(x) + A(g_2)(x).
$$

Thus, *A* is sublinear. Next, we prove that *A* satisfies (5.10). Indeed, notice that, for any $y \in \mathbb{R}^n$,

$$
(5.25) \qquad \qquad \left[\sum_{j\in\mathbb{N}}\left|\xi_j(y)\right|^r\right]^{1/r}\leq 1,
$$

which, combined with (2.5) and the Minkowski integral inequality, further implies that, for any $x \notin \overline{\text{supp}(g)}$,

$$
(5.26) \qquad |A(g)(x)| = \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(g\xi_j)(x) \right]^r \right\}^{\frac{1}{r}} \lesssim \left\{ \sum_{j \in \mathbb{N}} \left[\int_{\mathbb{R}^n} \frac{|g(y)\xi_j(y)|}{\rho(x-y)} dy \right]^r \right\}^{\frac{1}{r}}
$$

$$
\leq \int_{\mathbb{R}^n} \left\{ \sum_{j \in \mathbb{N}} \left[\frac{|g(y)\xi_j(y)|}{\rho(x-y)} \right]^r \right\}^{\frac{1}{r}} dy \leq \int_{\mathbb{R}^n} \frac{|g(y)|}{\rho(x-y)} dy,
$$

where the implicit positive constant is independent of $\{f_i\}_{i\in\mathbb{N}}$. Thus, *A* satisfies (5.10). In addition, from the Fefferman–Stein vector-valued inequality of \tilde{M} on the Lebesgue space $L^r(\mathbb{R}^n)$ (see, for instance, [4, p. 104, Theorem 6.2]) and (5.25), we infer that, for any $g \in \mathcal{M}(\mathbb{R}^n)$,

(5.27)
$$
||A(g)||_{L^p(\mathbb{R}^n)} = \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(g\xi_j) \right]^r \right\}^{\frac{1}{r}} \right\|_{L^p(\mathbb{R}^n)} \leq ||g||_{L^p(\mathbb{R}^n)},
$$

where the implicit positive constant is independent of $\{f_i\}_{i\in\mathbb{N}}$. Thus, *A* satisfies all the assumptions of Lemma 5.18. This, together with the fact that the implicit positive constants in both (5.26) and (5.27) are independent of $\{f_j\}_{j\in\mathbb{N}}$, further implies that *A* is bounded on $\mathcal{K}_{A,\omega}^{p,q}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$. Letting $g := (\sum_{j \in \mathbb{N}} |f_j|^r)^{1/r}$, by Lemma 5.18, we conclude that

$$
\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(f_j) \right]^r \right\}^{\frac{1}{r}} \right\|_{\dot{\mathcal{K}}^{p,q}_{A,\omega}(\mathbb{R}^n)} = \|A(g)\|_{\dot{\mathcal{K}}^{p,q}_{A,\omega}(\mathbb{R}^n)} \leq \|g\|_{\dot{\mathcal{K}}^{p,q}_{A,\omega}(\mathbb{R}^n)} = \left\| \left(\sum_{j \in \mathbb{N}} \left| f_j \right|^r \right)^{1/r} \right\|_{\dot{\mathcal{K}}^{p,q}_{A,\omega}(\mathbb{R}^n)},
$$

which completes the proof of Lemma 5.17.

By Lemma 5.17 and an argument similar to that used in the proof of [42, Lemma 4.3.10], we obtain the following conclusion.

Theorem 5.20. *Let A be a dilation, p, q* \in (0, ∞)*, and* $\omega \in M(\mathbb{R}_+)$ *satisfy* $m_0(\omega) \in (-\frac{1}{p}, \infty)$ *and* $m_{\infty}(\omega) \in (-\frac{1}{p}, \infty)$ *. Then, for any given r* ∈ (1, ∞) *and*

(5.28)
$$
u \in \left(0, \min\left\{p, \frac{1}{\max\{M_0(\omega), M_\infty(\omega)\} + 1/p}\right\}\right),\,
$$

there exists a positive constant C such that, for any $\{f_j\}_{j\in\mathbb{N}} \subset L^1_{\text{loc}}(\mathbb{R}^n)$,

$$
\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(f_j) \right]^r \right\}^{\frac{1}{r}} \right\|_{\left[\mathcal{K}^{p,q}_{A,\omega}(\mathbb{R}^n) \right]^{1/u}} \leq C \left\| \left\{ \sum_{j \in \mathbb{N}} |f_j|^r \right\}^{\frac{1}{r}} \right\|_{\left[\mathcal{K}^{p,q}_{A,\omega}(\mathbb{R}^n) \right]^{1/u}}.
$$

Proof. On the one hand, from [42, Lemma 1.3.1], we deduce that

(5.29)
$$
\left[\dot{K}_{A,\omega}^{p,q}(\mathbb{R}^n)\right]^{1/u} = \dot{K}_{A,\omega^u}^{p/u,q/u}(\mathbb{R}^n).
$$

On the other hand, applying Lemma 5.14, $m_0(\omega) > -\frac{1}{p}$, $m_\infty(\omega) > -\frac{1}{p}$, and (5.28), we conclude that

$$
\min\{m_0(\omega^u), m_\infty(\omega^u)\} = u \min\{m_0(\omega), m_\infty(\omega)\} > -\frac{1}{(p/u)}
$$

and

$$
\max\{M_0(\omega^u), M_\infty(\omega^u)\} = u \max\{M_0(\omega), M_\infty(\omega)\} < u\left(\frac{1}{u} - \frac{1}{p}\right) = \frac{1}{(p/u)'}.
$$

These, combined with (5.29) and Lemma 5.17, further implies that

$$
\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(f_j) \right]^r \right\}^{\frac{1}{r}} \right\|_{\left[\mathcal{K}^{p,q}_{A,\omega}(\mathbb{R}^n) \right]^{1/n}} \n= \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\mathcal{M}(f_j) \right]^r \right\}^{\frac{1}{r}} \right\|_{\mathcal{K}^{p,u,q/u}_{A,\omega^{u}}} \lesssim \left\| \left\{ \sum_{j \in \mathbb{N}} |f_j|^r \right\}^{\frac{1}{r}} \right\|_{\mathcal{K}^{p/u,q/u}_{A,\omega^{u}}} \n= \left\| \left\{ \sum_{j \in \mathbb{N}} |f_j|^r \right\}^{\frac{1}{r}} \right\|_{\left[\mathcal{K}^{p,q}_{A,\omega}(\mathbb{R}^n) \right]^{1/u}},
$$

which completes the proof of Theorem 5.20.

The following conclusion shows the boundedness of certain powered Hardy–Littlewood maximal operator on the associate space of certain power of $\dot{K}^{p,q}_{A,\omega}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$; see [42, Lemma 1.8.6] for the standard Euclidean space case.

Theorem 5.21. *Let A be a dilation, p, q* \in (0, ∞)*, and* $\omega \in M(\mathbb{R}_+)$ *satisfy* $m_0(\omega) \in (-\frac{1}{p}, \infty)$ *and* $m_{\infty}(\omega) \in \left(-\frac{1}{p}\right)$ $\frac{1}{p}$, ∞). *Then, for any given*

$$
s \in \left(0, \min\left\{p, q, \frac{1}{\max\{M_0(\omega), M_\infty(\omega)\} + 1/p}\right\}\right)
$$

and

$$
t \in \left(\max\left\{p, \frac{1}{\min\{m_0(\omega), m_\infty(\omega)\} + 1/p}\right\}, \infty\right],
$$

the Herz space $[\dot{\mathcal{K}}_{A,\omega}^{p,q}]$ *A*,ω (R *n*)]1/*^s is a ball Banach function space and there exists a positive constant C* such that, for any $f \in L^1_{loc}(\mathbb{R}^n)$,

$$
(5.30) \t\t ||\mathcal{M}^{((t/s)')}(f)||_{([\dot{K}_{A,\omega}^{p,q}(\mathbb{R}^n)]^{1/s})'} \leq C \, ||f||_{([\dot{K}_{A,\omega}^{p,q}(\mathbb{R}^n)]^{1/s})'}.
$$

To prove this theorem, we need more preparations. By Theorem 5.16 and an argument similar to that used in the proof of [42, Theorem 1.2.46], we obtain the following conclusion; we omit the details.

Fourier Transform of Anisotropic Hardy Spaces 37 37

Lemma 5.22. *Let A be a dilation, p, q* \in [1, ∞]*, and* $\omega \in M(\mathbb{R}_+)$ *satisfy*

$$
-\frac{1}{p} < m_0(\omega) \le M_0(\omega) < \frac{1}{p'},
$$

where $\frac{1}{p} + \frac{1}{p'}$ $\frac{1}{p'}$ = 1*. Then the anisotropic local generalized Herz space* $\dot{K}^{p,q}_{A,\omega}$ *A*,ω (R *n*) *is a ball Banach function space.*

Proof of Theorem 5.21. Obviously,

$$
s < \frac{1}{\max\{M_0(\omega), M_\infty(\omega)\} + 1/p} \le \frac{1}{M_0(\omega) + 1/p},
$$

which, combined with Lemma 5.14, further implies that

(5.31)
$$
M_0(\omega^s) = sM_0(\omega) < s\left(\frac{1}{s} - \frac{1}{p}\right) = \frac{1}{(p/s)'}
$$

and

(5.32)
$$
m_0(\omega^s) = sm_0(\omega) > -\frac{s}{p} = -\frac{1}{p/s}.
$$

From these, an argument similar to that used in the proof of [42, Lemma 1.3.1], the assumptions that $p/s > 1$ and $q/s > 1$, and Lemma 5.22, we infer that

(5.33)
$$
\left[\mathbf{K}_{A,\omega}^{p,q}(\mathbb{R}^n)\right]^{1/s} = \mathbf{K}_{A,\omega^s}^{p/s,q/s}(\mathbb{R}^n)
$$

is a ball Banach function space.

Next, we prove (5.30). On the one hand, by (5.33) and arguments similar to those used in the proofs of [42, Lemma 1.3.1 and Theorem 1.7.9], we conclude that

$$
(5.34) \qquad \qquad \left[\left(\left[\dot{\mathcal{K}}_{A,\omega}^{p,q}(\mathbb{R}^n) \right]^{1/s} \right)' \right]^{1/(t/s)'} = \left[\dot{\mathcal{K}}_{A,1/\omega^s}^{(p/s)',(q/s)',(q/s)'}(\mathbb{R}^n) \right]^{1/(t/s)'} \\qquad \qquad = \dot{\mathcal{K}}_{A,\omega^{-s(t/s)'} }^{(p/s)'/(t/s)',(q/s)'/(t/s)'}(\mathbb{R}^n).
$$

On the other hand, from Lemma 5.14 and the assumptions that

$$
s < \frac{1}{\max\{M_0(\omega), M_\infty(\omega)\} + 1/p}
$$

and

$$
t > \frac{1}{\min\{m_0(\omega), m_\infty(\omega)\} + 1/p},
$$

we deduce that

$$
\min\left\{m_0\left(\omega^{-s(t/s)'}\right), m_{\infty}\left(\omega^{-s(t/s)'}\right)\right\} = -s(t/s)'\max\{M_0(\omega), M_{\infty}(\omega)\}\
$$

$$
> -s(t/s)'\left(\frac{1}{s} - \frac{1}{p}\right) = -\frac{1}{(p/s)'/(t/s)'}
$$

and

$$
\max\left\{M_0\left(\omega^{-s(t/s)'}\right), M_\infty\left(\omega^{-s(t/s)'}\right)\right\} = -s(t/s)'\min\{m_0\left(\omega\right), m_\infty\left(\omega\right)\}
$$

$$
< -s(t/s)'\left(\frac{1}{t} - \frac{1}{p}\right) = 1 - \frac{1}{(p/s)'/(t/s)'}
$$

$$
= \frac{1}{[(p/s)'/(t/s)']'}.
$$

These, together with (5.34) and Corollary 5.19, further imply that

$$
\| \mathcal{M}^{((t/s)')}(f) \|_{([\vec{K}_{A,\omega}^{p,q}(\mathbb{R}^n)]^{1/s})'} = \| \mathcal{M}(|f|^{(t/s)'}) \|_{\vec{K}_{A,\omega^{-s(t/s)'}(t/s)'(q/s)'/(t/s)'(\mathbb{R}^n))}^{(\mathbb{R}^n)}
$$

$$
\leq \| |f|^{(t/s)'} \|_{\vec{K}_{A,\omega^{-s(t/s)'}}^{(p/s)/(t/s)',(q/s)'/(t/s)'(\mathbb{R}^n))}}
$$

$$
= \| f \|_{([\vec{K}_{A,\omega}^{p,q}(\mathbb{R}^n)]^{1/s})'} .
$$

This finishes the proof of (5.30) and hence Theorem 5.21.

Remark 5.23. Let $p, q \in (0, \infty)$ and $\omega \in M(\mathbb{R}_+)$ satisfy $m_0(\omega) \in (-\frac{1}{p}, \infty)$ and $m_\infty(\omega) \in (-\frac{1}{p}, \infty)$. Choose

$$
p_{-} := \min \left\{ p, q, \frac{1}{\max \{M_0(\omega), M_{\infty}(\omega)\} + 1/p} \right\}, \ \theta_0 \in (0, \underline{p}),
$$

and

$$
p_0 \in \left(\max\left\{1, p, \frac{1}{\min\{m_0(\omega), m_\infty(\omega)\} + 1/p}\right\}, \infty\right),\,
$$

where p is the same as in (2.6) . From these and Theorems 5.16, 5.20, and 5.21, we deduce that $\dot{\mathcal{K}}_{A,\omega}^{p,q}$ $P_{A,\omega}^{p,q}(\mathbb{R}^n)$ satisfies both Assumptions 2.7 and 2.8. Further assume that

$$
N\in\mathbb{N}\cap\left[\left|\left(\frac{1}{\min\{p,q,\frac{1}{\max\{M_0(\omega),M_\infty(\omega)\}+1/p}\}}-1\right)\frac{\ln b}{\ln(\lambda_-)}\right|+2,\infty\right).
$$

Then $\dot{\mathcal{K}}^{p,q}_{A,\omega}$ $H_{A,\omega}^{p,q}(\mathbb{R}^n)$ satisfies all the assumptions of Definition 2.10. Let $H\dot{\mathcal{K}}_{A,\omega}^{p,q}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$ be the anisotropic local generalized Herz–Hardy space defined in Definition 2.10 with $X := \mathcal{K}_{A,\omega}^{p,q}$ $L_{A,\omega}^{p,q}(\mathbb{R}^n)$. Then the main results of [59] and [38], for example, the maximal function characterizations of $H\dot{\mathcal{K}}_{A,\omega}^{p,q}$ $\chi_{A,\omega}^{p,q}(\mathbb{R}^n)$ (see [59, Theorem 3.9]), the (finite) atomic characterizations of $H\dot{\mathcal{K}}_{A,\omega}^{p,q}$ $A^{p,q}_{A,\omega}(\mathbb{R}^n)$ (see [59, Theorems 4.3 and 5.4]), the molecular characterization of $H\dot{\mathcal{K}}_{A,\omega}^{p,q}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$ (see [59, Theorem 6.3]), the boundedness of Calderón–Zygmund operators on $H\dot{K}^{p,q}_{A,\omega}$ $P_{A,\omega}^{p,q}(\mathbb{R}^n)$ (see [59, Theorems 7.10 and 7.11]), the Littlewood– Paley function characterizations of $H\widetilde{K}_{A,\omega}^{p,q}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$ (see [38, Theorems 5.4, 5.5, and 5.6]), and the dual theorem of $H\dot{\mathcal{K}}_{A,\omega}^{p,q}$ $\mathcal{A}_{A,\omega}^{p,q}(\mathbb{R}^n)$ (see [38, Theorem 3.25]) still hold true.

Let $p, q \in (0, 1)$ and $\omega \in M(\mathbb{R}_+)$ satisfy $m_0(\omega) \in (-\frac{1}{p}, \infty)$ and $m_\infty(\omega) \in (-\frac{1}{p}, \infty)$. Choose $a q_0 = 1$. By the above discussion, an argument similar to that used in the proof of [42, Lemma 4.9.4], and [42, (4.135)], we find that all the assumptions of Theorems 3.1, 4.1, and 4.2 are satisfied with $X := \dot{K}^{p,q}_{A,\omega}$ $A_{A,\omega}^{p,q}(\mathbb{R}^n)$. Applying Theorems 3.1, 4.1, and 4.2, we obtain the following result.

Fourier Transform of Anisotropic Hardy Spaces 39 and the United States 39 and 39

Theorem 5.24. *Let p*, $q \in (0, 1)$ *and* $\omega \in M(\mathbb{R}_+)$ *satisfy* $m_0(\omega) \in (-\frac{1}{p}, \infty)$ *and* $m_\infty(\omega) \in (-\frac{1}{p}, \infty)$ *. Then Theorems 3.1, 4.1, and 4.2 still hold true with X replaced by* $\dot{K}^{p,q}_{A,\omega}$ $P^{p,q}_{A,\omega}(\mathbb{R}^n)$.

Remark 5.25. We point out that Theorem 5.24 is completely new. When $A := 2I_{n \times n}$, the corresponding results of Theorem 5.24 can be found in [42, Theorem 4.9.1].

References

- [1] Z. V. P. Ablé and J. Feuto, Atomic decomposition of Hardy-amalgam spaces, J. Math. Anal. Appl. 455 (2017), 1899–1936.
- [2] P. Auscher and M. Mourgoglou, Representation and uniqueness for boundary value elliptic problems via first order systems, Rev. Mat. Iberoam. 35 (2019), 241–315.
- [3] C. Bennett and R. Sharpley, Interpolation of Operators, Pure Appl. Math. 129, Academic Press, Boston, MA, 1988.
- [4] M. Bownik, Anisotropic Hardy spaces and wavelets, Mem. Amer. Math. Soc. 164 (2003), no. 781, vi+122 pp.
- [5] M. Bownik, B. Li, D. Yang and Y. Zhou, Weighted anisotropic Hardy spaces and their applications in boundedness of sublinear operators, Indiana Univ. Math. J. 57 (2008), 3065–3100.
- [6] M. Bownik, B. Li, D. Yang and Y. Zhou, Weighted anisotropic product Hardy spaces and boundedness of sublinear operators, Math. Nachr. 283 (2010), 392–442.
- [7] M. Bownik and L.-A. D. Wang, Fourier transform of anisotropic Hardy spaces, Proc. Amer. Math. Soc. 141 (2013), 2299–2308.
- [8] T. A. Bui, P. D'Ancona and F. Nicola, Sharp L^p estimates for Schrödinger groups on spaces of homogeneous type, Rev. Mat. Iberoam. 36 (2020) 455–484.
- [9] T. A. Bui, X.-T. Duong and F. K. Ly, Maximal function characterizations for new local Hardy-type spaces on spaces of homogeneous type, Trans. Amer. Math. Soc. 370 (2018), 7229–7292.
- [10] T. A. Bui, X.-T. Duong and F. K. Ly, Maximal function characterizations for Hardy spaces on spaces of homogeneous type with finite measure and applications, J. Funct. Anal. 278 (2020), 108423, 55 pp.
- [11] T. A. Bui and J. Li, Orlicz–Hardy spaces associated to operators satisfying bounded *H*[∞] functional calculus and Davies–Gaffney estimates, J. Math. Anal. Appl. 373 (2011), 485– 501.
- [12] A.-P. Calder on and A. Torchinsky, Parabolic maximal functions associated with a distribution, Adv. Math. 16 (1975), 1–64.
- [13] D.-C. Chang, S. Wang, D. Yang and Y. Zhang, Littlewood–Paley characterizations of Hardytype spaces associated with ball quasi-Banach function spaces, Complex Anal. Oper. Theory 14 (2020), Paper No. 40, 33 pp.
- [14] J. Chou, X. Li, Y. Tong and H. Lin, Generalized weighted Morrey spaces on RD-spaces, Rocky Mountain J. Math. 50 (2020), 1277–1293.
- [15] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Anisotropic mixed-norm Hardy spaces, J. Geom. Anal. 27 (2017), 2758–2787.
- [16] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Molecular decomposition of anisotropic homogeneous mixed-norm spaces with applications to the boundedness of operators, Appl. Comput. Harmon. Anal. 47 (2019), 447–480.
- [17] G. Cleanthous, A. G. Georgiadis and M. Nielsen, Fourier multipliers on anisotropic mixednorm spaces of distributions, Math. Scand. 124 (2019), 289–304.
- [18] G. Cleanthous, A. G. Georgiadis and E. Porcu, Oracle inequalities and upper bounds for kernel density estimators on manifolds and more general metric spaces, J. Nonparametr. Stat. 34 (2022), 734–757.
- [19] R. R. Coifman, Characterization of Fourier transforms of Hardy spaces, Proc. Nat. Acad. Sci. U.S.A. 71 (1974), 4133–4134.
- [20] R. R. Coifman and G. Weiss, Analyse Harmonique Non-commutative sur Certains Espaces Homogènes. (French) Étude de Certaines Intégrales Singulières, Lecture Notes in Math. 242, Springer-Verlag, Berlin–New York, 1971.
- [21] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, Bull. Amer. Math. Soc. 83 (1977), 569–645.
- [22] L. Colzani, Fourier transform of distributions in Hardy spaces, Boll. Un. Mat. Ital. A (6) 1 (1982), 403–410.
- [23] J. Duoandikoetxea, Fourier Analysis, Graduate Studies in Mathematics 29, American Mathematical Society, Providence, RI, 2001.
- [24] C. Fefferman and E. M. Stein, H^p spaces of several variables, Acta Math. 129 (1972), 137– 193.
- [25] J. García-Cuerva and V. I. Kolyada, Rearrangement estimates for Fourier transforms in L^p and H^p in terms of moduli of continuity, Math. Nachr. 228 (2001), 123-144.
- [26] A. G. Georgiadis, G. Kyriazis and P. Petrushev, Product Besov and Triebel–Lizorkin spaces with application to nonlinear approximation, Constr. Approx. 53 (2021), 39–83.
- [27] A. G. Georgiadis and M. Nielsen, Spectral multipliers on spaces of distributions associated with non-negative self-adjoint operators, J. Approx. Theory 234 (2018), 1–19.
- [28] N. Hatano, R. Kawasumi and Y. Sawano, Sparse non-smooth atomic decomposition of quasi-Banach lattices, J. Fourier Anal. Appl. 28 (2022), Paper No. 61, 21 pp.
- [29] Z. He, Y. Han, J. Li, L. Liu, D. Yang and W. Yuan, A complete real-variable theory of Hardy spaces on spaces of homogeneous type, J. Fourier Anal. Appl. 25 (2019), 2197–2267.
- [30] Z. He, L. Liu, D. Yang and W. Yuan, New Calder on reproducing formulae with exponential decay on spaces of homogeneous type, Sci. China Math. 62 (2019), 283–350.
- [31] K.-P. Ho, Atomic decomposition of Hardy–Morrey spaces with variable exponents, Ann. Acad. Sci. Fenn. Math. 40 (2015), 31–62.
- [32] K.-P. Ho, Hardy's inequality on Hardy–Morrey spaces with variable exponents, Mediterr. J. Math. 14 (2017), Paper No. 79, 19 pp.
- [33] K.-P. Ho, Fractional integral operators on Morrey spaces built on rearrangement-invariant quasi-Banach function spaces, Positivity 27 (2023), Paper No. 26, 14 pp.
- [34] K.-P. Ho, Grand Morrey spaces and grand Hardy–Morrey spaces on Euclidean space, J. Geom. Anal. 33 (2023), Paper No. 180, 23 pp.
- [35] L. Huang, D.-C. Chang and D. Yang, Fourier transform of anisotropic mixed-norm Hardy spaces, Front. Math. China. 16 (2021), 119–139.
- [36] L. Huang, D.-C. Chang and D. Yang, Fourier transform of Hardy spaces associated with ball quasi-Banach function spaces, Appl. Anal. 101 (2022), 3825–3840.
- [37] V. Kokilashvili, A. Meskhi, H. Rafeiro and S. Samko, Integral Operators in Non-Standard Function Spaces, Vol. 1, Variable Exponent Lebesgue and Amalgam Spaces, Operator Theory: Advances and Applications 248, Birkhüser/Springer, [Cham], 2016, xx+567 pp.
- [38] C. Li, X. Yan and D. Yang, Anisotropic ball Campanato-type function spaces and their applications, Anal. Math. Phys. 13 (2023), Paper No. 50, 71 pp.
- [39] J. Li, Atomic decomposition of weighted Triebel–Lizorkin spaces on spaces of homogeneous type, J. Aust. Math. Soc. 89 (2010), 255–275.
- [40] J. Li, L. Song and C. Tan, Various characterizations of product Hardy space, Proc. Amer. Math. Soc. 139 (2011), 4385–4400.
- [41] J. Li and L. A. Ward, Singular integrals on Carleson measure spaces CMO^p on product spaces of homogeneous type, Proc. Amer. Math. Soc. 141 (2013), 2767–2782.
- [42] Y. Li, D. Yang and L. Huang, Real-Variable Theory of Hardy Spaces associated with Generalized Herz Spaces of Rafeiro and Samko, Lecture Notes in Mathematics 2320, Springer, Cham, 2022.
- [43] Y. Liang, J. Huang and D. Yang, New real-variable characterizations of Musielak–Orlicz Hardy spaces, J. Math. Anal. Appl. 395 (2012), 413–428.
- [44] J. Liu, Molecular characterizations of variable anisotropic Hardy spaces with applications to boundedness of Calder *ón*–Zygmund operators, Banach J. Math. Anal. 15 (2021), 1–24.
- [45] J. Liu, Fourier transform of variable anisotropic Hardy spaces with applications to Hardy– Littlewood inequalities, Math. Inequal. Appl. 25 (2022), 447–465.
- [46] J. Liu, Y. Lu and M. Zhang, Fourier transform of anisotropic mixed-norm Hardy spaces with applications to Hardy–Littlewood inequalities, J. Korean Math. Soc. 59 (2022), 927–944.
- [47] J. Liu, D. Yang and M. Zhang, Sharp bilinear decomposition for products of both anisotropic Hardy spaces and their dual spaces with its applications to endpoint boundedness of commutators, Sci. China Math. (2023), https://doi.org/10.1007/s11425-023-2153-y.
- [48] W. Matuszewska and W. Orlicz, On certain properties of φ -functions, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 8 (1960), 439-443.
- [49] W. Matuszewska and W. Orlicz, On some classes of functions with regard to their orders of growth, Studia Math. 26 (1965), 11–24.
- [50] C. B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, Trans. Amer. Math. Soc. 43 (1938), 126–166.
- [51] H. Rafeiro and S. Samko, Herz spaces meet Morrey type spaces and complementary Morrey type spaces, J. Fourier Anal. Appl. 26 (2020), Paper No. 74, 14 pp.
- [52] N. Samko, Weighted Hardy operators in the local generalized vanishing Morrey spaces, Positivity 17 (2013), 683–706.
- [53] Y. Sawano, K.-P. Ho, D. Yang and S. Yang, Hardy spaces for ball quasi-Banach function spaces, Dissertationes Math. 525 (2017), 1–102.
- [54] Y. Sawano and H. Tanaka, Predual spaces of Morrey spaces with non-doubling measures, Tokyo J. Math. 32 (2009), 471–486.
- [55] E. M. Stein, Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals, Princeton Mathematical Series 43, Monographs in Harmonic Analysis III, Princeton University Press, Princeton, NJ, 1993.
- [56] M. H. Taibleson and G. Weiss, The molecular characterization of certain Hardy spaces, in: Representation theorems for Hardy spaces, pp. 67–149, Astérisque, 77, Soc. Math. France, Paris, 1980.
- [57] F. Wang, D. Yang and S. Yang, Applications of Hardy spaces associated with ball quasi-Banach function spaces, Results Math. 75 (2020), Paper No. 26, 58 pp.
- [58] S. Wang, D. Yang, W. Yuan and Y. Zhang, Weak Hardy-type spaces associated with ball quasi-Banach function spaces II: Littlewood–Paley characterizations and real interpolation, J. Geom. Anal. 31 (2021), 631–696.
- [59] Z. Wang, X. Yan and D. Yang, Anisotropic Hardy spaces associated with ball Quasi-Banach function spaces and their applications, Kyoto J. Math. (to appear).
- [60] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Littlewood–Paley characterizations with applications to boundedness of Calder on–Zygmund operators, Acta Math. Sin. (Engl. Ser.) 38 (2022), 1133–1184.
- [61] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Characterizations of maximal functions, decompositions, and dual spaces, Math. Nachr. 296 (2023), 3056–3116.
- [62] X. Yan, D. Yang and W. Yuan, Intrinsic square function characterizations of Hardy spaces associated with ball quasi-Banach function spaces, Front. Math. China 15 (2020), 769–806.
- [63] X. Yan and D. Yang, New molecular characterization of Musielak–Orlicz Hardy spaces on spaces of homogeneous type and its applications, Chinese Ann. Math. Ser. B (to appear).
- [64] X. Yan, D. Yang, W. Yuan and C. Zhuo, Variable weak Hardy spaces and their applications, J. Funct. Anal. 271 (2016), 2822–2887.
- [65] Y. Zhang, S. Wang, D. Yang and W. Yuan, Weak Hardy-type spaces associated with ball quasi-Banach function spaces I: Decompositions with applications to boundedness of Calderón–Zygmund operators, Sci. China Math. 64 (2021), 2007–2064.
- [66] Y. Zhang, D. Yang and W. Yuan, Real-variable characterizations of local Orlicz-slice Hardy spaces with application to bilinear decompositions, Commun. Contemp. Math. 24 (2022), Paper No. 2150004, 35 pp.
- [67] Y. Zhang, D. Yang, W. Yuan and S. Wang, Real-variable characterizations of Orlicz-slice Hardy spaces, Anal. Appl. (Singap.) 17 (2019), 597–664.

Chaoan Li and Dachun Yang (Corresponding author)

Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, The People's Republic of China

E-mails: cali@mail.bnu.edu.cn (C. Li) dcyang@bnu.edu.cn (D. Yang)

Xianjie Yan

Institute of Contemporary Mathematics, School of Mathematics and Statistics, Henan University, Kaifeng 475004, The People's Republic of China

E-mail: xianjieyan@henu.edu.cn