

An Accelerated Stochastic ADMM for Nonconvex and Nonsmooth Finite-Sum Optimization

Yuxuan Zeng, Zhiguo Wang, Jianchao Bai, and Xiaojing Shen

July 4, 2023

Abstract

The nonconvex and nonsmooth finite-sum optimization problem with linear constraint has attracted much attention in the fields of artificial intelligence, computer, and mathematics, due to its wide applications in machine learning and the lack of efficient algorithms with convincing convergence theories. A popular approach to solve it is the stochastic Alternating Direction Method of Multipliers (ADMM), but most stochastic ADMM-type methods focus on convex models. In addition, the variance reduction (VR) and acceleration techniques are useful tools in the development of stochastic methods due to their simplicity and practicability in providing acceleration characteristics of various machine learning models. However, it remains unclear whether accelerated SVRG-ADMM algorithm (ASVRG-ADMM), which extends SVRG-ADMM by incorporating momentum techniques, exhibits a comparable acceleration characteristic or convergence rate in the nonconvex setting. To fill this gap, we consider a general nonconvex nonsmooth optimization problem and study the convergence of ASVRG-ADMM. By utilizing a well-defined potential energy function, we establish its sublinear convergence rate $O(1/T)$, where T denotes the iteration number. Furthermore, under the additional Kurdyka-Lojasiewicz (KL) property which is less stringent than the frequently used conditions for showcasing linear convergence rates, such as strong convexity, we show that the ASVRG-ADMM sequence almost surely has a finite length and converges to a stationary solution with a linear convergence rate. Several experiments on solving the graph-guided fused lasso problem and regularized logistic regression problem validate that the proposed ASVRG-ADMM performs better than the state-of-the-art methods.

Keywords— Nonconvex and nonsmooth optimization, acceleration technique, stochastic ADMM, convergence.

I. INTRODUCTION

In recent years, machine learning has been studied and applied extensively in system identification [1], and automatic control [2]. In this paper, we consider a class of the nonconvex nonsmooth finite-sum

Yuxuan Zeng, Zhiguo Wang (corresponding author), and Xiaojing Shen are with College of Mathematics, Sichuan University, 610064, Chengdu, China (e-mail: 2020222010085@stu.scu.edu.cn, wangzhiguo@scu.edu.cn, shenxj@scu.edu.cn).

Jianchao Bai (corresponding author) is with Research & Development Institute of Northwestern Polytechnical University in Shenzhen, Shenzhen 518057, China; School of Mathematics and Statistics, Northwestern Polytechnical University, Xi'an 710129, China (e-mail: jianchaobai@nwpu.edu.cn).

optimization problems with a linear constraint in machine learning, as follows:

$$\min_{\mathbf{x} \in \mathbb{R}^{d_1}, \mathbf{y} \in \mathbb{R}^{d_2}} f(\mathbf{x}) + g(\mathbf{y}), \text{ s.t. } A\mathbf{x} + B\mathbf{y} = \mathbf{c}, \quad (1)$$

where $f(\mathbf{x})$ consists of the summation of n components, i.e., $f(\mathbf{x}) := \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{x})$ and $f_i: \mathbb{R}^{d_1} \rightarrow \mathbb{R}$ is assumed smooth, but can be nonconvex; $g(\mathbf{y})$ is convex but possibly nonsmooth; the constraint is used for encoding the structure pattern of model parameters, where $A \in \mathbb{R}^{d \times d_1}$, $B \in \mathbb{R}^{d \times d_2}$, $\mathbf{c} \in \mathbb{R}^d$. To motivate our work, we first briefly discuss one common setting related to problem (1).

Let us consider a binary classification task. Specifically, given a set of training samples (a_i, b_i) , $i = 1, \dots, n$, where a_i is the input data and its corresponding label is $b_i \in \{1, -1\}$. Then we use the famous graph-guided fused Lasso model [3] to learn the parameter, which can be represented as

$$\min_{\mathbf{x}} \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{x}) + \lambda_1 \|A\mathbf{x}\|_1, \quad (2)$$

where the sigmoid loss function f_i is as defined by $f_i(\mathbf{x}) = \frac{1}{1 + \exp(b_i a_i^T \mathbf{x})}$, which is a nonconvex function. A is a given matrix decoded the sparsity pattern of the graph, which is obtained by sparse inverse covariance matrix estimation [4]. In order to solve (2), we can introduce an additional primal variable \mathbf{y} , and reformulate the problem (2) as a special case of (1)

$$\min_{\mathbf{x}, \mathbf{y}} f(\mathbf{x}) + g(\mathbf{y}) \quad \text{s.t. } A\mathbf{x} - \mathbf{y} = 0,$$

where $f(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{x})$ and $g(\mathbf{y}) = \lambda_1 \|\mathbf{y}\|_1$.

The problem (1) also attracts attention and arises in many other fields such as statistical learning [5], computer vision [6] and 3D CT image reconstruction [7]. In the following text, we focus on the composite finite-sum equality-constrained optimization problem (1) and check the efficiency of this paper from both algorithmic and theoretical perspectives.

A. Related work

One of the popular methods to solve the general optimization (1) is ADMM algorithm [8]–[10]. Specifically, at the t -th iteration, the typical sequential update steps are

$$\begin{cases} \mathbf{y}_{t+1} = \arg \min_{\mathbf{y}} \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}, \boldsymbol{\lambda}_t), \\ \mathbf{x}_{t+1} = \arg \min_{\mathbf{x}} \mathcal{L}_\rho(\mathbf{x}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t), \\ \boldsymbol{\lambda}_{t+1} = \boldsymbol{\lambda}_t - \rho (A\mathbf{x}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}), \end{cases} \quad (3)$$

where $\boldsymbol{\lambda}$ is the dual variable and

$$\mathcal{L}_\rho(\mathbf{x}, \mathbf{y}, \boldsymbol{\lambda}) = f(\mathbf{x}) + g(\mathbf{y}) - \langle \boldsymbol{\lambda}, A\mathbf{x} + B\mathbf{y} - \mathbf{c} \rangle + \frac{\rho}{2} \|A\mathbf{x} + B\mathbf{y} - \mathbf{c}\|^2 \quad (4)$$

is an augmented Lagrangian (AL) function and $\rho > 0$ is a penalty parameter. Many ADMM-type algorithms, such as proximal ADMM [9], [11], [12], inexact ADMM [8], [13], linearized/relaxed ADMM [14], [15], and consensus ADMM [16], [17] have been proposed to solve the subproblem in (3) efficiently. Recently, a proximal AL method with an exponential averaging scheme was proposed by [18] to handle nonconvex optimizations.

Large-scale optimization problems (1) typically involve a large sum of n component functions, making it infeasible for deterministic ADMMs to compute the full gradient on all training samples at each iteration. Stochastic gradient descent (SGD) has a much lower per-iteration complexity than deterministic methods by utilizing only one sample's gradient at each iteration. Stochastic ADMM (SADMM), proposed in [19], combines SGD and ADMM, and the update step of variable \mathbf{x}_{t+1} in SADMM is approximated as follows:

$$\mathbf{x}_{t+1} = \arg \min_{\mathbf{x}} \mathbf{x}^T \nabla f_{i_t}(\mathbf{x}_t) + \frac{1}{2\eta_t} \|\mathbf{x} - \mathbf{x}_t\|_Q^2 + \frac{\rho}{2} \|A\mathbf{x} + B\mathbf{y}_{t+1} - \mathbf{c} - \frac{\boldsymbol{\lambda}_t}{\rho}\|^2, \quad (5)$$

where we select the random variable i_t uniformly at random from $[n] := \{1, \dots, n\}$, $\eta_t \propto 1/\sqrt{t}$ is the step-size, and $\|\mathbf{x}\|_Q^2 = \mathbf{x}^T Q \mathbf{x}$ with given positive semi-definite matrix Q .

However, when the objective function is convex, stochastic ADMMs have been proved to own the worst-case $O(1/\sqrt{T})$ convergence rate, which is lower than the deterministic ADMMs with $O(1/T)$ convergence rate. The gap in convergence caused by the high variances of stochastic gradients can be tackled by combining stochastic variance reduction gradient (SVRG) methods [20] with ADMM, resulting in the SVRG-ADMM in [21]. Unlike SADMM, variance-reduced ADMM aims to find an unbiased gradient estimator whose variance will vanish as the algorithm converges. This encourages the selection of a larger step-size, such as a constant step-size η , to improve the convergence rate. Specifically, we compute the gradient $\hat{\nabla} f(\mathbf{x}_t^{s+1})$ for each iteration as follows:

$$\hat{\nabla} f(\mathbf{x}_t^{s+1}) = \nabla f_{i_t}(\mathbf{x}_t^{s+1}) - \nabla f_{i_t}(\tilde{\mathbf{x}}^s) + \nabla f(\tilde{\mathbf{x}}^s), \quad (6)$$

where i_t is defined in the equation (5), and $\hat{\nabla} f(\mathbf{x}_t^{s+1})$ is an unbiased estimator of the gradient $\nabla f(\mathbf{x}_t^{s+1})$, i.e., $\mathbb{E}[\hat{\nabla} f(\mathbf{x}_t^{s+1})] = \nabla f(\mathbf{x}_t^{s+1})$.

The Nesterov acceleration technique [22] enjoys a fast convergence rate and has been successfully applied to train many large-scale machine learning models, including deep network models. Directly extending the Nesterov acceleration technique to the stochastic settings may aggravate the inaccuracy of the stochastic gradient and harm the convergence performance. [13] proposed ASVRG-ADMM, an accelerated SVRG-ADMM method by introducing a new Katyusha momentum acceleration trick [23] into SVRG-ADMM. Table I presents a comparison of several stochastic ADMM algorithms. All algorithms

TABLE I: Comparison of convergence rates of some stochastic ADMM algorithms.

Algorithms	Strongly convex	Convex	Nonconvex	Momentum	KL property
SAG-ADMM [21]	\	\	$\mathcal{O}(1/T)$	×	×
SDCA-ADMM [20]	Linear Rate	\	\	×	×
SVRG-ADMM [21]	\	\	$\mathcal{O}(1/T)$	×	×
SPIDER-ADMM [26]	\	\	$\mathcal{O}(1/T)$	×	×
ASVRG-ADMM [13]	Linear Rate	$\mathcal{O}(1/T^2)$	\	✓	×
AS-ADMM [8]	Linear Rate	$\mathcal{O}(1/T)$	\	✓	×
ASVRG-ADMM (ours)	\	\	$\mathcal{O}(1/T)$ & linear rate	✓	✓

in the table incorporate the VR technique, while only the last three employ momentum acceleration techniques.

Although ASVRG-ADMM has been successful in convex problems, its behavior in nonconvex nonsmooth problems remains largely unknown. Many important applications, such as signal/image processing [24], computer vision [6], and video surveillance [25], involve nonconvex objective functions that are beyond the scope of the theoretical conditions under which ASVRG-ADMM [13] has been proven to converge. Therefore, there exists a gap in the theoretical convergence analysis between extant ASVRG-ADMM in the convex scenario and ASVRG-ADMM in the nonconvex scenario.

To fill this gap, this paper addresses two crucial problems:

- Q1. Whether the ASVRG-ADMM algorithm can converge to the stationary point for the common nonconvex nonsmooth finite-sum optimization?
- Q2. If the algorithm converges, can we establish its linear convergence rate such as R-linear¹ convergence?

B. Contributions

In this paper, we develop ASVRG-ADMM for the nonconvex and nonsmooth problems (1). ASVRG-ADMM is inspired by the momentum [8], [13], [23] and the variance reduction [20], [21] technique.

¹

The sequence $\{x_n\}_{n \in \mathbb{N}}$ converges to x^* R-linearly if

$$\|x_n - x^*\| \leq y_n,$$

for all n , and the sequence $\{y_n\}_{n \in \mathbb{N}}$ converges to zero Q-linearly. Then one can always find a suitable sequence $\{y_n\}_{n \in \mathbb{N}}$ as $y_n = c * \xi^n$, such that

$$\|x_n - x^*\| \leq c * \xi^n,$$

for all n , where $0 < \xi < 1$.

In addition, it has several new features. First, compared with [13], we consider the nonconvex objective function rather than the convex function, which poses a major obstacle to establishing the convergence of our algorithm without the sufficiently decreasing property of the AL function. Second, compared with SVRG-ADMM in [21], our proposed algorithm employs the Katyusha momentum technique to improve the accuracy of the stochastic gradient and facilitate faster convergence in practice. In summary, our main contributions include four folds as follows:

- (C1) We propose a novel accelerated stochastic ADMM (ASVRG-ADMM) for the nonconvex and nonsmooth problem (1), which enjoys the advantages of both the momentum acceleration technique [13], [23] and the variance reduction technique [20], [21].
- (C2) By introducing a potential energy function related to the AL function of (1), we establish the sublinear convergence rate $O(1/T)$, then we answer **Q1**. While ASVRG-ADMM has the same convergence rate as SVRG-ADMM [21], theoretically, we can achieve faster decay of the potential energy function than SVRG-ADMM by selecting the optimal momentum parameter. Furthermore, our algorithm achieves better convergence in numerical experiments.
- (C3) We establish the almost sure R-linear convergence rate of ASVRG-ADMM by utilizing the extra KL property to answer **Q2**, as listed in Table I and demonstrated in Theorem 2. Answering **Q2** under the nonconvex and nonsmooth setting is not a trivial task due to the inclusion of both the momentum term and variance reduction technique. To the best of our knowledge, this is the first almost sure linear convergence result for the accelerated stochastic ADMM algorithm with variance reduction technique.

Various numerical experiments demonstrate the effectiveness of our ASVRG-ADMM, as it exhibits both lower variance and faster convergence compared to several advanced stochastic ADMM-type methods [19], [21].

Synopsis: The remainder of this paper is organized as follows. Section II introduces the notations and mild assumptions. We present the framework of ASVRG-ADMM in Section III. We study its convergence and extend the theoretical analysis of the almost sure linear convergence rate under the extra KL property in Section IV. Section V presents some numerical results to demonstrate the effectiveness and efficiency of the method. Finally, we conclude the paper with discussions.

Notation: The symbol $\|\cdot\|$ denotes the Euclidean norm of a vector (or the spectral norm of a matrix), and $\|\cdot\|_1$ is the ℓ_1 -norm, i.e., $\|\mathbf{x}\|_1 = \sum_i \|x_i\|$. We use \mathbb{R} , \mathbb{R}^d , and $\mathbb{R}^{d \times d_1}$ to denote the sets of real numbers, d dimensional real column vectors, and $d \times d_1$ real matrices, respectively. Let (Ω, \mathcal{F}, P) be the probability space, $\mathbf{x} : \Omega \rightarrow \mathbb{R}^d$ be a random variable, $\mathbb{E}[\mathbf{x}]$ denote the expectation, $\mathbb{E}[\mathbf{x} \mid \mathcal{F}_t^s]$

denote the conditional expectation of \mathbf{x} given sub- σ -algebra \mathcal{F}_t^s , and employ the abbreviations ‘‘a.s.’’ for ‘‘almost surely’’. Let I denote the identity matrix and $\mathbf{0}$ denote the zero matrix. We denote by $\nabla f(\mathbf{x})$ the gradient of $f(\mathbf{x})$ if it is differentiable, or ∂f any of the subgradients of the function $f(\cdot)$. Let σ_{min}^A and σ_{max}^A be the smallest and largest eigenvalues of matrix AA^T , respectively; and let ϕ_{min} and ϕ_{max} denote the smallest and largest eigenvalues of positive matrix Q , respectively. For a nonempty closed set \mathcal{C} , $dist(\mathbf{x}, \mathcal{C}) = \inf_{\mathbf{y} \in \mathcal{C}} \|\mathbf{x} - \mathbf{y}\|$ denotes the distance from \mathbf{x} to set \mathcal{C} .

II. PRELIMINARIES

A. Basic Assumptions and Definitions

Before establishing the convergence analysis, we clarify the following proper assumptions.

Assumption 1 For $\forall i \in \{1, 2, \dots, n\}$, each function f_i possesses a Lipschitz continuous gradient with the constant $L_i > 0$, that is

$$\|\nabla f_i(\mathbf{x}_1) - \nabla f_i(\mathbf{x}_2)\| \leq L \|\mathbf{x}_1 - \mathbf{x}_2\|, \quad \forall \mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^{d_1},$$

where $L = \max_i L_i$.

Assumption 2 $f(\mathbf{x})$ and $g(\mathbf{y})$ are lower bounded.

Assumption 3 (feasibility) $(\text{Im}(B) \cup \mathbf{c}) \subseteq \text{Im}(A)$, where $\text{Im}(\cdot)$ returns the image of a matrix.

Assumption 4 (Lipschitz sub-minimization paths) For any fixed \mathbf{x} , $\text{argmin}_{\mathbf{y}} \{f(\mathbf{x}) + g(\mathbf{y}) : B\mathbf{y} = \mathbf{u}\}$ has a unique minimizer. In addition, $H : \text{Im}(B) \rightarrow \mathbb{R}^{d_2}$ defined by $H(\mathbf{u}) \triangleq \text{argmin}_{\mathbf{y}} \{f(\mathbf{x}) + g(\mathbf{y}) : B\mathbf{y} = \mathbf{u}\}$ is a Lipschitz continuous map.

Assumption 1 is a standard smoothness assumption. Note that Assumptions 1-2 are satisfied for the case of $f(\mathbf{x}) = \frac{1}{1+\exp(\mathbf{x})}$, $g(\mathbf{y}) = \lambda_1 \|\mathbf{y}\|_1$. Assumptions 3-4 are proposed in [27], which weaken the full column rank assumption typically imposed on matrices A and B . Moreover, Assumptions 3-4 are the mild assumptions commonly used to ensure that the dual variable λ is controlled by the primal variable, see Lemma 1.

Next, we introduce the critical point and KL property used in this paper.

Definition 1 The point $(\mathcal{X}^*, \mathcal{Y}^*)$ is denoted as the feasible solution set of the problem (1). When there exist a optimal point $(\mathbf{x}^*, \mathbf{y}^*)$ and a dual variable $\boldsymbol{\lambda}^*$ for the problem (4), the KKT conditions are satisfied as follows:

$$\begin{cases} A^T \boldsymbol{\lambda}^* = \nabla f(\mathbf{x}^*), \\ B^T \boldsymbol{\lambda}^* \in \partial g(\mathbf{y}^*), \\ A\mathbf{x}^* + B\mathbf{y}^* - \mathbf{c} = 0. \end{cases}$$

We also say that $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ satisfy the KKT conditions is exactly a critical point of the AL function, denoted as $\text{crit } \mathcal{L}_\rho$.

Definition 2 (KL property [28]) A proper lower semicontinuous function $F : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to possess the KL property at $\mathbf{x}^* \in \text{dom } \partial F$ if there exist a constant $\eta \in (0, \infty]$, a neighborhood U of \mathbf{x}^* and a continuous concave function $\varphi : [0, \eta) \rightarrow \mathbb{R}_+$, satisfying the following requirements:

- $\varphi(0) = 0$, $\varphi \in C^1((0, \eta))$, and $\nabla \varphi(s) > 0$ for all $s \in (0, \eta)$;
- for all $\mathbf{x} \in U \cap [F(\mathbf{x}^*) < F < F(\mathbf{x}^*) + \eta]$, the KL inequality holds:

$$\nabla \varphi(F(\mathbf{x}) - F(\mathbf{x}^*)) \text{dist}(0, \partial F(\mathbf{x})) \geq 1. \quad (7)$$

Many effective functions satisfy the KL property, including but not limited to $\|\mathbf{x}\|_1$, $\|\mathbf{A}\mathbf{x} - \mathbf{b}\|^2$, log-exp, and the logistic loss function $\psi(t) = \log(1 + e^{-t})$. For more examples, readers can refer to [28], [29], [30], [31], and [32](see Page 919). Therefore, the potential energy function defined in (15) can easily satisfy the KL property.

Remark 1 When function F in Definition 2 is differentiable, then the KL property becomes the PL property [33], which is much weaker than the strongly convex property, and widely used to show the global convergence in the field of deep learning. However, the PL property only deals with smooth differentiable functions. In this paper, the considered problem (1) not only has the smooth term but also has the nonsmooth term, thus we use the KL property to show the linear convergence.

III. DEVELOPMENTS OF ASVRG-ADMM

We propose ASVRG-ADMM, which incorporates momentum and variance reduction techniques to ensure fast convergence of the nonsmooth and nonconvex problem (1). The algorithm, presented in Algorithm 1, consists of S epochs, each containing m iterations with m typically chosen to be $O(n)$.

Algorithm 1: ASVRG-ADMM for solving (1)

Input: parameter $m, T, S = \lceil T/m \rceil, \eta, \gamma, \rho, 0 \leq \theta \leq 1$, and initial values of $\tilde{\mathbf{x}}^0 = \mathbf{x}_m^0, \mathbf{y}_m^0$,

$$\mathbf{z}_m^0 = \mathbf{x}_m^0 \text{ and } \boldsymbol{\lambda}_m^0.$$

Output: Iterate \mathbf{x} and \mathbf{y} chosen from $\{(\mathbf{x}_m^S, \mathbf{y}_m^S)\}$.

for $s = 0, 1, \dots, S - 1$ **do**

$$\mathbf{x}_0^{s+1} = \mathbf{x}_m^s, \mathbf{y}_0^{s+1} = \mathbf{y}_m^s \text{ and } \boldsymbol{\lambda}_0^{s+1} = \boldsymbol{\lambda}_m^s;$$

$$\nabla f(\tilde{\mathbf{x}}^s) = \frac{1}{n} \sum_{i=1}^n \nabla f_i(\tilde{\mathbf{x}}^s);$$

for $t = 0, 1, \dots, m - 1$ **do**

Uniformly and randomly choose a random variable i_t from $\{1, \dots, n\}$, and compute

$$\hat{\nabla} f(\mathbf{x}_t^{s+1}) \text{ in (6) ;}$$

$$\mathbf{y}_{t+1}^{s+1} = \arg \min_{\mathbf{y}} \mathcal{L}_\rho(\mathbf{x}_t^{s+1}, \mathbf{y}, \boldsymbol{\lambda}_t^{s+1}); \quad (8)$$

$$\mathbf{z}_{t+1}^{s+1} = \mathbf{z}_t^{s+1} - \frac{\eta}{\gamma\theta} \left[\hat{\nabla} f(\mathbf{x}_t^{s+1}) + \rho A^T \left(A\mathbf{z}_t^{s+1} + B\mathbf{y}_{t+1}^{s+1} - \mathbf{c} - \frac{\boldsymbol{\lambda}_t^{s+1}}{\rho} \right) \right]; \quad (9)$$

$$\mathbf{x}_{t+1}^{s+1} = \theta \mathbf{z}_{t+1}^{s+1} + (1 - \theta) \tilde{\mathbf{x}}^s; \quad (10)$$

$$\boldsymbol{\lambda}_{t+1}^{s+1} = \boldsymbol{\lambda}_t^{s+1} - \rho(A\mathbf{z}_{t+1}^{s+1} + B\mathbf{y}_{t+1}^{s+1} - \mathbf{c}); \quad (11)$$

end

$$\tilde{\mathbf{x}}^{s+1} = \mathbf{x}_m^{s+1}.$$

end

A. Update of Primal Variable \mathbf{y}

The first step is to update the primal variable \mathbf{y} in (8) by solving

$$\begin{aligned} \mathbf{y}_{t+1}^{s+1} &= \arg \min_{\mathbf{y}} \mathcal{L}_\rho(\mathbf{x}_t^{s+1}, \mathbf{y}, \boldsymbol{\lambda}_t^{s+1}) \\ &= \arg \min_{\mathbf{y}} g(\mathbf{y}) + \frac{\rho}{2} \left\| A\mathbf{x}_t^{s+1} + B\mathbf{y} - \mathbf{c} - \frac{\boldsymbol{\lambda}_t^{s+1}}{\rho} \right\|^2. \end{aligned} \quad (12)$$

If $g(\mathbf{y}) = \lambda_1 \|\mathbf{y}\|_1$, one can observe that the optimization (12) has closed form solution with

$$\mathbf{y}_{t+1}^{s+1} = \max \left(\mathbf{0}, A\mathbf{x}_t^{s+1} - \frac{\boldsymbol{\lambda}_t^{s+1}}{\rho} - \frac{\lambda_1}{\rho} \right) - \max \left(\mathbf{0}, -A\mathbf{x}_t^{s+1} + \frac{\boldsymbol{\lambda}_t^{s+1}}{\rho} - \frac{\lambda_1}{\rho} \right),$$

where λ_1 is the regularization parameter.

B. Update of Auxiliary Variable \mathbf{z}

Inspired by Katyusha [13], [23] acceleration tricks, the second step is adopting an auxiliary variable \mathbf{z} before updating the primal variable \mathbf{x} . A surrogate function is first introduced to approximate the objective function in (3) and the update rule of \mathbf{z} is formulated as follows:

$$\mathbf{z}_{t+1}^{s+1} = \arg \min_{\mathbf{z}} (\mathbf{z} - \mathbf{z}_t^{s+1})^T \hat{\nabla} f(\mathbf{x}_t^{s+1}) + \frac{\theta}{2\eta} \|\mathbf{z} - \mathbf{z}_t^{s+1}\|_Q^2 + \frac{\rho}{2} \|\mathbf{A}\mathbf{z} + \mathbf{B}\mathbf{y}_{t+1}^{s+1} - \mathbf{c} - \frac{\boldsymbol{\lambda}_t^{s+1}}{\rho}\|^2, \quad (13)$$

where $\eta > 0$ is the learning rate or step-size, and $\hat{\nabla} f(\mathbf{x}_t^{s+1})$ is the stochastic variance reduced gradient estimator defined in (6).

In general, the direct solution of (13) can be challenging or even infeasible since it requires the inversion of $\frac{\theta}{\eta}Q + \rho A^T A$. To alleviate this computational burden, the inexact Uzawa method can be employed to linearize the last term (13). Specifically, we can select $Q = \gamma I - \frac{\eta\rho}{\theta}A^T A$ with $\gamma \geq 1 + \frac{\eta\rho\|A^T A\|_2}{\theta}$ to ensure that $Q \succ I$, which results in the simpler closed-form solution of (13) as (9).

C. Momentum Accelerated Update Rule for \mathbf{x}

The next step is our momentum accelerated update rule for the primal variable \mathbf{x} , as given below:

$$\mathbf{x}_{t+1}^{s+1} = \theta \mathbf{z}_{t+1}^{s+1} + (1 - \theta) \tilde{\mathbf{x}}^s = \tilde{\mathbf{x}}^s + \theta(\mathbf{z}_{t+1}^{s+1} - \tilde{\mathbf{x}}^s), \quad (14)$$

where θ is a momentum parameter, and $\theta(\mathbf{z}_{t+1}^{s+1} - \tilde{\mathbf{x}}^s)$ is a momentum term, which uses the final output iterate from the previous epoch (updated every m iterations in the inner loop), i.e., $\tilde{\mathbf{x}}^s$, to speed up our algorithm. In (14), we leverage the Katyusha momentum technique [23], which uses a convex combination of the latest \mathbf{z}_{t+1}^{s+1} and snapshot $\tilde{\mathbf{x}}^s$, whereas the Nesterov-type momentum technique uses a nonconvex extrapolation of the two latest iterates. Compared with Nesterov momentum, Katyusha momentum ensures that the iterates do not deviate too far from $\tilde{\mathbf{x}}^s$, which ensures accurate gradient estimation.

D. Update of Dual Variable $\boldsymbol{\lambda}$

The final step is to update the dual variable by

$$\boldsymbol{\lambda}_{t+1}^{s+1} = \boldsymbol{\lambda}_t^{s+1} - \rho(\mathbf{A}\mathbf{z}_{t+1}^{s+1} + \mathbf{B}\mathbf{y}_{t+1}^{s+1} - \mathbf{c}).$$

Note that we use auxiliary variable \mathbf{z}_{t+1}^{s+1} rather than the primal variable \mathbf{x}_{t+1}^{s+1} [19] in the dual update scheme, which helps us to establish the convergence of the proposed algorithm under nonconvex setting.

Before ending the section, let's make a few remarks about ASVRG-ADMM.

Remark 2 Compared with SVRG-ADMM [21], our ASVRG-ADMM incorporates a new acceleration step (14) and its sublinear and linear convergence have been established under weaker Assumption 3-

4. Additionally, if we set θ in the update step of \mathbf{x}_{t+1}^{s+1} to 1, it results in the algorithm degenerating into SVRG-ADMM [21], where $\mathbf{x}_{t+1}^{s+1} = \mathbf{z}_{t+1}^{s+1}$, which can be evidenced by the overlapping descent curves of ASVRG-ADMM1 ($\theta = 1$) and SVRG-ADMM in Figure 3 (left).

IV. CONVERGENCE ANALYSIS

This section establishes the sublinear and linear convergence of our proposed ASVRG-ADMM for nonconvex and nonsmooth problems. In the literature, the AL function [27] is frequently used as the potential energy function with sufficiently decreasing property (see [27]). However, in the nonconvex setting, our Algorithm 1 deviates from this approach. To ensure sufficient decrease and convergence, we present the following lemmas and theorems with a practical sequence of potential energy functions $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$ defined as follows:

$$\Psi_t^s := \mathcal{L}_\rho(\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s) + \beta_5 \|\mathbf{x}_t^s - \mathbf{x}_{t-1}^s\|^2 + h_t^s (\|\mathbf{x}_t^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \|\mathbf{x}_{t-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2), \quad (15)$$

where the positive sequence $\{(h_t^s)_{t=1}^m\}_{s=1}^S$ and the parameter $\beta_5 > 0$ will be decided in the following Lemma 2.

Let $\{(\mathbf{x}_t^s)_{t=1}^m\}_{s=1}^S$ be a stochastic process adapted to the filtration $\{(\mathcal{F}_t^s)_{t=1}^m\}_{s=1}^S$, where $\mathcal{F}_t^s = \sigma(\mathbf{x}_1^1, \mathbf{x}_2^1, \dots, \mathbf{x}_t^s)$, $\sigma(\cdot)$ is the σ -field, and we abbreviate $\mathbb{E}[\cdot \mid \mathcal{F}_t^s]$ as $\mathbb{E}_t^s[\cdot]$, see more details in [7], [30]. Additionally, we define Ψ^* as $\min_{s,t} \Psi_t^s$. Now, we first provide an outline of the proof in this section.

- Firstly, we propose a new potential energy function, which differs from the one in [21], and prove the monotonicity of this function.
- Next, we leverage the monotonicity property of the potential energy function to establish the convergence rate of the newly defined sequence in (24).
- Lastly, we leverage the KL property along with finite-length property in Lemma 3 to achieve a almost sure linear convergence result for ASVRG-ADMM.

Our convergence analysis will utilize a bound on the term $\mathbb{E}_t^{s+1} \|\boldsymbol{\lambda}_{t+1}^{s+1} - \boldsymbol{\lambda}_t^{s+1}\|^2$, which is given by Lemma 1.

Lemma 1 (Upper bound of $\mathbb{E}_t^{s+1} \|\boldsymbol{\lambda}_{t+1}^{s+1} - \boldsymbol{\lambda}_t^{s+1}\|^2$) *Given Assumptions 1-3, for $\{(\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s)_{t=1}^m\}_{s=1}^S$ generated by the Algorithm 1, we can have the following result*

$$\begin{aligned} \mathbb{E}_t^{s+1} [\|\boldsymbol{\lambda}_{t+1}^{s+1} - \boldsymbol{\lambda}_t^{s+1}\|^2] &\leq \frac{5L^2}{\sigma_{\min}^A} [\|\mathbf{x}_t^{s+1} - \tilde{\mathbf{x}}^s\|^2] + \frac{5L^2}{\sigma_{\min}^A} \|\mathbf{x}_{t-1}^{s+1} - \tilde{\mathbf{x}}^s\|^2 + \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \mathbb{E}_t^{s+1} [\|\mathbf{x}_{t+1}^{s+1} - \mathbf{x}_t^{s+1}\|^2] \\ &\quad + \frac{5(L^2 \eta^2 + \phi_{\max}^2)}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_t^{s+1} - \mathbf{x}_{t-1}^{s+1}\|^2. \end{aligned} \quad (16)$$

Proof 1 See Appendix A.

Lemma 2 For the sequence $\{(\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s)_{t=1}^m\}_{s=1}^S$ generated by Algorithm 1, assuming that Assumptions 1-3 hold, and the positive sequences $\{(h_t^s)_{t=1}^m\}_{s=1}^S$ and $\{(\Gamma_t^s)_{t=1}^m\}_{s=1}^S$ exist, we choose positive parameters $\eta, \theta, l_1, l_2, l_3, \alpha_1$, and ρ such that

$$\Gamma_t^s > 0, \forall t \in [m], \forall s \in [S],$$

then the sequence $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$ is sufficiently decreasing:

$$\mathbb{E}_t^{s+1}[\Psi_{t+1}^{s+1}] \leq \Psi_t^{s+1} - \Gamma_t^{s+1} \mathbb{E}_t^{s+1}[\|\mathbf{x}_{t+1}^{s+1} - \mathbf{x}_t^{s+1}\|^2] - [h_{t+1}^{s+1} + (1 + \alpha_1)(h_{t+1}^{s+1} + \beta_1)] \|\mathbf{x}_{t-1}^{s+1} - \tilde{\mathbf{x}}^s\|^2. \quad (17)$$

Proof 2 The specific expression of the tuple $(\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6)$ and the sequences $\{(h_t^s)_{t=1}^m\}_{s=1}^S$, $\{(\Gamma_t^s)_{t=1}^m\}_{s=1}^S$ are provided in Appendix B.

Remark 3 As we see, there are some crucial parameters in Lemma 2. In this remark, we further clarify how to choose these parameters θ, η, ρ . One can observe that there exists a parameter Γ_t^s in the right side of the inequality (17). If Γ_t^s promotes larger, then the sequence $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$ decreases faster.

- Firstly, we rewritten Γ_t^s defined in (S.5) as follows:

$$\Gamma_t^s = F(\theta) + H(\eta) - \frac{L}{2} - \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2}{\sigma_{min}^A} - h_{t+1}^s \left(1 + \frac{1}{\alpha_1}\right), \quad (18)$$

where

$$F(\theta) = \frac{\rho \sigma_{min}^A (l_2 - (1 - \theta))}{2\theta^2 l_2} - \left(\frac{\rho + l_1}{2}\right) \left(\frac{1 - \theta}{\theta}\right)^2 \sigma_{max}^A \left(1 + \frac{1}{\alpha_1}\right), \quad (19)$$

$$H(\eta) = \frac{\phi_{min}}{\eta} - \frac{10\phi_{max}^2}{\sigma_{min}^A \eta^2} \left(\frac{1}{\rho} + \frac{1}{2l_1}\right), \quad (20)$$

- Secondly, in order to obtain the maximum value of Γ_t^s , then the optimal parameters θ^* and η^* can be obtained

$$\theta^* = \frac{2l_2(\rho + l_1)(1 + \alpha_1)\sigma_{max}^A - 2\alpha_1\rho\sigma_{min}^A(l_2 + 1)}{2l_2(\rho + l_1)(1 + \alpha_1)\sigma_{max}^A + \alpha_1\rho\sigma_{min}^A(l_2 + 1)} \quad (21)$$

$$\frac{1}{\eta^*} = \frac{\phi_{min}\sigma_{min}^A}{20\phi_{max}^2 \left(\frac{1}{\rho} + \frac{1}{2l_1}\right)}, \quad (22)$$

- Finally, from (21) and (22), we can see that the optimal θ^* and η^* become larger, the parameter ρ should be adjusted to smaller.

Remark 4 Now we show that there also exist a lower bound for the dual step size ρ . Taking the optimal value of η^* in (22) into Γ_t^s in (18), we study the requirement of ρ with the inequality $\Gamma_t^s > 0$, which is simplified as follows:

$$0 < \frac{\phi_{\min}^2 \sigma_{\min}^A}{40\phi_{\max}^2 \left(\frac{1}{\rho} + \frac{1}{2l_1}\right)} - \frac{L}{2} + \frac{\rho\sigma_{\min}^A(l_2 - (1 - \theta))}{2\theta^2 l_2} - \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2}{\sigma_{\min}^A} - (h_{t+1}^s + \beta_1)\left(1 + \frac{1}{\alpha_1}\right).$$

Considering the setting of $\sigma_{\min}^A \rho > 1$, the parameter ρ should satisfy the following inequality

$$\begin{aligned} & 10L^2 + (h_{t+1}^s + \beta_1) \left(1 + \frac{1}{\alpha_1}\right) \\ & < \frac{\phi_{\min}^2 \sigma_{\min}^A \theta^2 l_2 + 40\sigma_{\min}^A [l_2 - (1 - \theta)] \phi_{\max}^2}{80\phi_{\max}^2 \theta^2 l_2} \rho, \end{aligned}$$

which indicates

$$\rho > \frac{80\phi_{\max}^2 \theta^2 l_2 \left[10L^2 + (h_{t+1}^s + \beta_1) \left(1 + \frac{1}{\alpha_1}\right)\right]}{\phi_{\min}^2 \sigma_{\min}^A \theta^2 l_2 + 40\sigma_{\min}^A [l_2 - (1 - \theta)] \phi_{\max}^2}. \quad (23)$$

Based on the above important descending Lemma 2, we next analyze the convergence and iteration complexity of the ASVRG-ADMM with the aid of a simple notation:

$$R_t^s := \mathbb{E}\|\mathbf{x}_t^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \mathbb{E}\|\mathbf{x}_{t-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \mathbb{E}\|\mathbf{x}_{t+1}^s - \mathbf{x}_t^s\|^2 + \mathbb{E}\|\mathbf{x}_t^s - \mathbf{x}_{t-1}^s\|^2. \quad (24)$$

We also denote

$$(\hat{t}, \hat{s}) = \arg \min_{1 \leq t \leq m, 1 \leq s \leq S} R_t^s.$$

Theorem 1 Assuming that Assumptions 1-3 hold, let $\{(\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s)_{t=1}^m\}_{s=1}^S$ be the sequence generated by Algorithm 1, using the identical notations as in Lemma 2. For $\tau = \min\{\gamma, \omega\} > 0$, where $\gamma = \min_{t,s} \Gamma_t^s$ and $\omega = \frac{5L^2}{\sigma_{\min}^A \rho}$, the following can be derived:

$$R_{\hat{t}}^{\hat{s}} = \min_{s,t} R_t^s \leq \frac{1}{\tau T} \mathbb{E}(\Psi_1^1 - \Psi^*), \quad (25)$$

where Ψ^* denotes a lower bound of the sequence $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$.

Proof 3 See Appendix C.

Theorem 1 shows that Algorithm 1 converges with the worst-case $\mathcal{O}(1/T)$ convergence rate, where $T = mS$.

Lemma 3 (The Finite Length Property) Let the sequence $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ be generated by Algorithm 1 under Assumptions 1-4 and Ψ_t has the KL property at the stationary point $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ with KL

exponent $\tilde{\mu} \in [0; 1)$ (see the Definition 2). Suppose that f and g are semi-algebraic functions satisfying the KL inequality, then $\{\mathbf{w}_t\}$ has a finite length, that is,

$$\sum_{t=0}^{+\infty} \|\mathbf{w}_{t+1} - \mathbf{w}_t\| < +\infty \quad a.s.$$

Moreover, $\{\mathbf{w}_t\}$ converges a.s. to a critical point of $\mathcal{L}_\rho(\cdot)$.

Proof 4 See Appendix F.

Now, combined with the KL property defined in Definition 2 (see [28]), we can obtain the linear convergence rate of ASVRG-ADMM.

Theorem 2 (Linear Convergence) For notational simplicity, we also omit the upper script s . The sequence $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ is generated by Algorithm 1, and it converges to $\{\mathbf{w}^* = (\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)\}$. Additionally, we define $\varphi(s) = \tilde{c}s^{1-\tilde{\mu}}$, $\tilde{\mu} \in [0, 1)$, $\tilde{c} > 0$. Assume that Ψ_t defined in (15) has the KL property at the stationary point $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ with an exponent $\tilde{\mu}$. Then, we draw the following estimations under Assumptions 1-4:

- i) If $\tilde{\mu} = 0$, then the sequence $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ converges a.s. in a finite number of steps.
- ii) If $\tilde{\mu} \in (0, \frac{1}{2}]$, then there exists $\hat{c} > 0$ and $\xi_3 \in [0, 1)$, such that

$$\|(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) - (\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)\| \leq \hat{c}\xi_3^t \quad a.s.$$

- iii) If $\tilde{\mu} \in (\frac{1}{2}, 1)$, then there exists $\hat{c} > 0$, such that

$$\|(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) - (\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)\| \leq \hat{c}t^{(\tilde{\mu}-1)/(2\tilde{\mu}-1)} \quad a.s.$$

Proof 5 We also give a sketch of the proof and refer to Appendix G for more details. This proof is structured by two main steps which contain:

- First, we define a new sequence and prove it converges a.s. to zero Q -linearly.
- Second, we conclude that the sequence $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ is bounded by the newly defined sequence which converges a.s. Q -linearly, thus the sequence $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ converges a.s. to $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ R -linearly.

Remark 5 We prove in Theorem 2 that our ASVRG-ADMM almost surely achieves R -linear convergence at the rate of ξ_3 for nonconvex nonsmooth optimization, in accordance with the definition of R -linear convergence presented in Section I. We construct a potential energy function Ψ_t^s in (15) that satisfies the KL property to ensure convergence when $\tilde{\mu} \in (0, 1/2]$. This is the first demonstration of linear convergence for a stochastic ADMM incorporating momentum and variance reduction techniques.




Fig. 1: Comparison of the SADMM, SVRG-ADMM, and ASVRG-ADMM algorithms for the nonconvex quadratic problem in the synthetic setting. The objective values (left) and variances of gradient (right).

V. NUMERICAL EXPERIMENTS

In this section, we examine the numerical performance of the proposed ASVRG-ADMM algorithm and present comparison results with the existing methods. We conducted several different real dataset experiments to validate our theoretical results and show the high efficiency of the implementation of our proposed ASVRG-ADMM algorithmic framework. We compare ASVRG-ADMM with the following state-of-the-art methods for nonconvex problems: SADMM in [19], SVRG-ADMM in [21], and SAG-ADMM in [21]. In the following, all algorithms were implemented in MATLAB, and the experiments were performed on a PC with an Intel i7-12700F CPU and 16GB memory.

A. Synthetic Data For the Nonconvex Quadratic Problem

Our setup for the synthetic nonconvex quadratic problem is as follows. We first generate the symmetric matrix A_i from the standard $d_1 \times d_1$ Gaussian distribution. Next, the matrix $A = I$ or $A = [G; I]$, where G is obtained from a sparse inverse covariance estimation given in [3]. Then we solve the following nonconvex quadratic problem:

$$\frac{1}{n} \sum_{i=1}^n \frac{\mathbf{x}^T A_i \mathbf{x}}{2} + \lambda_1 \|\mathbf{y}\|_1, \text{ s.t. } \mathbf{y} = A\mathbf{x},$$

where $\lambda_1 = 10^{-4}$ is a positive regularization parameter. Finally, all experimental results are averaged over 300 Monte Carlo experiments.

From Fig.1, we can see the training loss (i.e., the training objective value minus the best value) and the variance of SADMM [19], SVRG-ADMM [21], and ASVRG-ADMM for the nonconvex quadratic problem. All the experimental results show that our ASVRG-ADMM method with smaller variance

converges consistently much faster than both SADMM and SVRG-ADMM, which empirically verifies our theoretical results of the effectiveness of the momentum technique resulting in faster convergence.

B. Graph-Guided Fused Lasso




Fig. 2: Comparison of different stochastic ADMM methods for graph-guided fused lasso problems on the two data sets: ijcn1 (left) and w8a (right).

We first evaluate the proposed method for solving the graph-guided fused Lasso problem (2). We set $\eta = 2$, $\rho = 6$, $\lambda_1 = 10^{-5}$ and $Q = I$, similar to [21]. Two cases are taken into account in Algorithm 1, particularly: a time-varying stepsize parameter $\eta_t = 2\sqrt{t+2}$ for the SADMM; a fixed parameter $\eta = 2$ for the SADMM-F. The parameter $m = n$ has been set for the SVRG-ADMM algorithm. At last, the result of all experiments is averaged over 30 Monte Carlos experiments.

We use real datasets ijcn1, a9a, and w8a downloaded from LIBSVM to test our proposed method. The training error of all the methods is shown in Fig.2. The results indicate that compared to algorithms without variance reduction techniques, e.g., SADMM and SADMM-F, the variance reduced stochastic ADMM algorithms (including both SVRG-ADMM and ASVRG-ADMM) converge substantially more quickly. Notably, in terms of the experimental convergence performance, ASVRG-ADMM consistently outperforms all comparison algorithms under all datasets, which is consistent with our theoretical analysis.

We compare various values of momentum θ in Figure 3, namely the larger value $\theta = 0.9$ (ASVRG-ADMM2), the smaller but empirically superior value $\theta = 0.19$ (ASVRG-ADMM3), and the Nesterov-type momentum parameter [13], [22] (ASVRG-ADMM-Nes). The results indicate that smaller values of θ lead to improved experimental outcomes while employing the Nesterov-type decaying momentum parameter results in optimal performance. Furthermore, we conduct additional comparisons using different values




Fig. 3: Comparison of ASVRG-ADMM algorithm for different θ (left) and ρ (right) on Graph-Guided Fused Lasso using synthetic data.

of ρ as shown in Figure 3. The value of $\rho = 60$ in ASVRG-ADMM1 exceeded the experimentally optimal value of $\rho = 6$ in ASVRG-ADMM2. To satisfy the lower bound inequality (23), an adaptive approach known as ASVRG-ADMM-Adap [34] can be employed. This approach gradually increases ρ by setting $\rho_{t+1} = \kappa * \rho_t$, where $\kappa > 1$, to ensure the validity of (23). Smaller values of ρ yield improved experimental performance, with the adaptive approach achieving the best results.

C. Regularized Logistic Regression

In this subsection, we evaluate the test data performance of ASVRG-ADMM for solving the non-convex Regularized Logistic Regression (RLR) problem

$$\min_{\mathbf{x}} \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{x}) + \lambda_1 \|\mathbf{x}\|_1,$$

where $f_i(\mathbf{x})$ is defined the same as in (2), and the parameter $\lambda_1 = 2 * 10^{-5}$. In addition, we randomly select 60% of the data to train the models and the rest to test the performance.

Fig.4 shows the experimental results of datasets w8a (left) and a9a (right) respectively. Fig.4 indicates that both the algorithm SADMM and SADMM-F consistently converge more slowly than the others and the variances of these two algorithms are larger than the others in all settings, which empirically verifies that variance reduced stochastic ADMM algorithms with smaller variance can perform better. We observe that SVRG-ADMM and ASVRG-ADMM consistently outperform the others. Moreover, ASVRG-ADMM performs much better than the other methods in all settings, which again verifies the effectiveness of our ASVRG-ADMM method with the accelerated momentum technique.




Fig. 4: Comparison of different stochastic ADMM methods for the accuracy of logistic regression problems.

VI. CONCLUSION AND FUTURE WORK

In this paper, we integrated both momentum acceleration and the variance reduction trick into stochastic ADMM and proposed an efficient accelerated stochastic variance reduced ADMM for solving the common nonconvex nonsmooth problems. Furthermore, we theoretically proved that ASVRG-ADMM obtains a similar convergence to SVRG-ADMM [21] for nonconvex problems. Especially under the KL property, the convergence rate of ASVRG-ADMM can be improved to the almost sure linear convergence rate. Compared to those of the state-of-the-art stochastic ADMM methods such as SVRG-ADMM, massive experimental results validate the numerical advantages of our proposed algorithm. For future work, there are some interesting directions such as the research of applying the momentum acceleration trick to other algorithms like SAG-ADMM and SAGA-ADMM [21], and the research of the theoretical improvement of the convergence rate of the proposed method. In addition, it is also interesting to extend ASVRG-ADMM and theoretical results to the distributed optimization and nonsmooth and nonconvex optimization.

REFERENCES

- [1] G. Pillonetto, F. Dinuzzo, T. Chen, G. De Nicolao, and L. Ljung, “Kernel methods in system identification, machine learning and function estimation: A survey,” *Automatica*, vol. 50, no. 3, pp. 657–682, 2014.
- [2] L. Li, H. Luo, S. X. Ding, Y. Yang, and K. Peng, “Performance-based fault detection and fault-tolerant control for automatic control systems,” *Automatica*, vol. 99, pp. 308–316, 2019.
- [3] S. Kim, K.-A. Sohn, and E. P. Xing, “A multivariate regression approach to association analysis of a quantitative trait network,” *Bioinformatics*, vol. 25, no. 12, pp. i204–i212, 2009.
- [4] J. Friedman, H. Trevor, and R. Tibshirani, “Sparse inverse covariance estimation with the graphical lasso,” *Biostatistics*, vol. 9, no. 3, pp. 432–441, 2008.

- [5] J. Bai, H. Zhang, and J. Li, “A parameterized proximal point algorithm for separable convex optimization,” *Optimization Letters*, vol. 12, no. 7, pp. 1589–1608, 2018.
- [6] N. P. Papanikolopoulos and P. K. Khosla, “Adaptive robotic visual tracking: Theory and experiments,” *IEEE Transactions on Automatic Control*, vol. 38, no. 3, pp. 429–445, 1993.
- [7] F. Bian, J. Liang, and X. Zhang, “A stochastic alternating direction method of multipliers for non-smooth and non-convex optimization,” *Inverse Problems*, vol. 37, p. 075009, 2021.
- [8] J. Bai, W. Hager, and H. Zhang, “An inexact accelerated stochastic ADMM for separable convex optimization,” *Computational Optimization and Applications*, vol. 81, no. 2, pp. 479–518, 2022.
- [9] Y. Yang, Q. Jia, and e. a. Xu, Zhanbo, “Proximal ADMM for nonconvex and nonsmooth optimization,” *Automatica*, vol. 146, p. 110551, 2022.
- [10] M. Chao, D. Han, and X. Cai, “Convergence of the peaceman–rachford splitting method for a class of nonconvex programs,” *Numerical Mathematics: Theory, Methods and Applications*, vol. 14, no. 2, pp. 438–460, 2021.
- [11] K. Guo, D. Han, and T. Wu, “Convergence of ADMM for optimization problems with nonseparable nonconvex objective and linear constraints,” *Pacific Journal of Optimization*, vol. 14, pp. 489–506, 2018.
- [12] Z. Wu and M. Li, “General inertial proximal gradient method for a class of nonconvex nonsmooth optimization problems,” *Computational Optimization and Applications*, vol. 73, pp. 129–158, 2019.
- [13] Y. Liu, F. Shang, H. Liu, L. Kong, L. Jiao, and Z. Lin, “Accelerated variance reduction stochastic ADMM for large-scale machine learning,” *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol. 43, no. 12, pp. 4242–4255, 2020.
- [14] J. Bai, D. Han, H. Sun, and H. Zhang, “Convergence on a symmetric accelerated stochastic ADMM with larger stepsizes,” *CSIAM Transactions on Applied Mathematics*, vol. 31, no. 3, pp. 448–479, 2022.
- [15] M. Tao, “Convergence study of indefinite proximal ADMM with a relaxation factor,” *Computational Optimization and Applications*, vol. 77, no. 1, pp. 91–123, 2020.
- [16] M. Hong, Z.-Q. Luo, and M. Razaviyayn, “Convergence analysis of alternating direction method of multipliers for a family of nonconvex problems,” *SIAM Journal on Optimization*, vol. 26, no. 1, pp. 337–364, 2016.
- [17] Z. Wang, J. Zhang, T.-H. Chang, J. Li, and Z.-Q. Luo, “Distributed stochastic consensus optimization with momentum for nonconvex nonsmooth problems,” *IEEE Transactions on Signal Processing*, vol. 69, pp. 4486–4501, 2021.
- [18] J. Zhang and Z.-Q. Luo, “A global dual error bound and its application to the analysis of linearly constrained nonconvex optimization,” *SIAM Journal on Optimization*, vol. 32, no. 3, pp. 2319–2346, 2022.
- [19] H. Ouyang, N. He, L. Tran, and A. Gray, “Stochastic alternating direction method of multipliers,” in *International Conference on Machine Learning*. PMLR, 2013, pp. 80–88.
- [20] T. Suzuki, “Stochastic dual coordinate ascent with alternating direction method of multipliers,” in *International Conference on Machine Learning*. PMLR, 2014, pp. 736–744.
- [21] F. Huang, S. Chen, and Z. Lu, “Stochastic alternating direction method of multipliers with variance reduction for nonconvex optimization,” *arXiv preprint arXiv:1610.02758*, 2016.
- [22] Y. Nesterov, *Introductory Lectures on Convex Optimization: A Basic Course*. Springer Science & Business Media, 2013, vol. 87.
- [23] Zeyuan Allen-Zhu, “Katyusha: The first direct acceleration of stochastic gradient methods,” *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, 2017.
- [24] Z. Ge, X. Zhang, and Z. Wu, “A fast proximal iteratively reweighted nuclear norm algorithm for nonconvex low-rank matrix minimization problems,” *Applied Numerical Mathematics*, vol. 179, pp. 66–86, 2022.

- [25] B. Gao and F. Ma, “Alternating direction method of multipliers for a class of nonconvex and nonsmooth problems with applications to background/foreground,” *SIAM Journal on Imaging Sciences*, vol. 10, no. 1, pp. 74–110, 2017.
- [26] F. Huang, S. Chen, and H. Huang, “Faster stochastic alternating direction method of multipliers for nonconvex optimization,” in *International Conference on Machine Learning*. PMLR, 2019, pp. 2839–2848.
- [27] Y. Wang, W. Yin, and J. Zeng, “Global convergence of ADMM in nonconvex nonsmooth optimization,” *Journal of Scientific Computing*, vol. 78, no. 1, pp. 29–63, 2019.
- [28] H. Attouch, J. Bolte, P. Redont, and A. Soubeyran, “Proximal alternating minimization and projection methods for nonconvex problems: An approach based on the Kurdyka-Lojasiewicz inequality,” *Mathematics of Operations Research*, vol. 35, no. 2, pp. 438–457, 2010.
- [29] J. Bolte, S. Sabach, and M. Teboulle, “Proximal alternating linearized minimization for nonconvex and nonsmooth problems,” *Mathematical Programming*, vol. 146, no. 1, pp. 459–494, 2014.
- [30] A. Milzarek and J. Qiu, “Convergence of a normal map-based prox-sgd method under the kl inequality,” *arXiv preprint arXiv:2305.05828*, 2023.
- [31] E. Chouzenoux, J.-B. Fest, and A. Repetti, “A kurdyka-lojasiewicz property for stochastic optimization algorithms in a non-convex setting,” 2023.
- [32] M. Yashtini, “Convergence and rate analysis of a proximal linearized admm for nonconvex nonsmooth optimization,” *Journal of Global Optimization*, vol. 84, no. 4, pp. 913–939, 2022.
- [33] X. Yi, S. Zhang, T. Yang, and K. H. Johansson, “Zeroth-order algorithms for stochastic distributed nonconvex optimization,” *Automatica*, vol. 142, p. 110353, 2022.
- [34] B. He, H. Yang, and S. Wang, “Alternating direction method with self-adaptive penalty parameters for monotone variational inequalities,” *Journal of Optimization Theory and applications*, vol. 106, pp. 337–356, 2000.
- [35] F. Huang and S. Chen, “Mini-batch stochastic admm for nonconvex nonsmooth optimization,” *arXiv preprint arXiv:1802.03284*, 2018.

APPENDIX A

PROOF OF THE LEMMA 1

Proof: For notational simplicity, we denote the stochastic gradient $\Delta_t^{s+1} := \hat{\nabla} f(\mathbf{x}_t^{s+1}) - \nabla f(\mathbf{x}_t^{s+1})$, where $\hat{\nabla} f(\mathbf{x}_t^{s+1}) = \nabla f_{i_t}(\mathbf{x}_t^{s+1}) - \nabla f_{i_t}(\tilde{\mathbf{x}}^s) + \nabla f(\tilde{\mathbf{x}}^s)$. We also denote the variance of stochastic gradient $\hat{\nabla} f(\mathbf{x}_t^{s+1})$ as $\mathbb{E}_t[\|\Delta_t^{s+1}\|^2]$, and omit the label s as $\|\Delta_t^{s+1}\|^2 := \|\Delta_t\|^2$, $\mathbf{x}_t^{s+1} := \mathbf{x}_t$, $\mathbf{y}_t^{s+1} := \mathbf{y}_t$, $\boldsymbol{\lambda}_t^{s+1} := \boldsymbol{\lambda}_t$, $\tilde{\mathbf{x}}^s := \tilde{\mathbf{x}}$, and the conditional expectation operator $\mathbb{E}_t^s := \mathbb{E}_t$. By the optimal condition of \mathbf{z} -update (9) in Algorithm 1, we have

$$\begin{aligned} 0 &= \hat{\nabla} f(\mathbf{x}_t) + \rho A^\top (A\mathbf{z}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}) - A^\top \boldsymbol{\lambda}_t + \frac{\theta}{\eta} Q(\mathbf{z}_{t+1} - \mathbf{z}_t) \\ &= \hat{\nabla} f(\mathbf{x}_t) - A^\top \boldsymbol{\lambda}_{t+1} + \frac{\theta}{\eta} Q(\mathbf{z}_{t+1} - \mathbf{z}_t), \end{aligned}$$

where the second equality follows from the update of $\boldsymbol{\lambda}$ (11) in Algorithm 1. Thus, we have $A^\top \boldsymbol{\lambda}_{t+1} = \hat{\nabla} f(\mathbf{x}_t) + \frac{\theta}{\eta} Q(\mathbf{z}_{t+1} - \mathbf{z}_t)$. Recall that $\mathbf{x}_{t+1} = \theta \mathbf{z}_{t+1} + (1-\theta)\tilde{\mathbf{x}}$, then it yields $\mathbf{x}_{t+1} - \mathbf{x}_t = \theta(\mathbf{z}_{t+1} - \mathbf{z}_t)$.

$$A^\top \boldsymbol{\lambda}_{t+1} = \hat{\nabla} f(\mathbf{x}_t) - \frac{1}{\eta} Q(\mathbf{x}_t - \mathbf{x}_{t+1}). \quad (\text{S.1})$$

Combining (S.1) and the properties of $\|\cdot\|$

$$\begin{aligned} &\|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\|^2 \\ &\leq (\sigma_{\min}^A)^{-1} \|A^\top \boldsymbol{\lambda}_{t+1} - A^\top \boldsymbol{\lambda}_t\|^2 \\ &= (\sigma_{\min}^A)^{-1} \left\| \hat{\nabla} f(\mathbf{x}_t) - \hat{\nabla} f(\mathbf{x}_{t-1}) - \frac{1}{\eta} Q(\mathbf{x}_t - \mathbf{x}_{t+1}) - \frac{1}{\eta} Q(\mathbf{x}_t - \mathbf{x}_{t-1}) \right\|^2 \\ &= (\sigma_{\min}^A)^{-1} \left\| \hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_t) + \nabla f(\mathbf{x}_t) - \nabla f(\mathbf{x}_{t-1}) + \nabla f(\mathbf{x}_{t-1}) - \hat{\nabla} f(\mathbf{x}_{t-1}) + \frac{1}{\eta} Q(\mathbf{x}_{t+1} - \mathbf{x}_t) \right. \\ &\quad \left. - \frac{1}{\eta} Q(\mathbf{x}_t - \mathbf{x}_{t-1}) \right\|^2 \\ &\stackrel{(i)}{\leq} \frac{5}{\sigma_{\min}^A} \|\Delta_t\|^2 + \frac{5}{\sigma_{\min}^A} \|\Delta_{t-1}\|^2 + \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2 + \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\ &\quad + \frac{5}{\sigma_{\min}^A} \|\nabla f(\mathbf{x}_t) - \nabla f(\mathbf{x}_{t-1})\|^2, \end{aligned} \quad (\text{S.2})$$

and applying the conditional expectation operator \mathbb{E}_t , we obtain

$$\begin{aligned} &\mathbb{E}_t \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\|^2 \\ &\leq \frac{5}{\sigma_{\min}^A} \|\Delta_t\|^2 + \frac{5}{\sigma_{\min}^A} \|\Delta_{t-1}\|^2 + \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2 + \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\ &\quad + \frac{5}{\sigma_{\min}^A} \|\nabla f(\mathbf{x}_t) - \nabla f(\mathbf{x}_{t-1})\|^2, \end{aligned}$$

where the first inequality is based on Assumption 3 and the update rule of dual variable given by $\boldsymbol{\lambda}_t^{s+1} - \boldsymbol{\lambda}_{t+1}^{s+1} = \rho(A\mathbf{z}_{t+1}^{s+1} + B\mathbf{y}_{t+1}^{s+1} - c) \in \text{Im}(A)$, the inequality (i) holds by the Cauchy-Schwartz inequality, ϕ_{\max} denotes the largest eigenvalue of positive matrix Q , and $\|Q(x - \mathbf{y})\|^2 \leq \phi_{\max}^2 \|x - \mathbf{y}\|^2$.

We recall the [21, Lemma 1] as follows:

$$\mathbb{E}_t[\|\Delta_t^{s+1}\|^2] \leq L^2 \|\mathbf{x}_t^{s+1} - \tilde{\mathbf{x}}^s\|^2. \quad (\text{S.3})$$

By inserting (S.3) and Assumption 1 to the above inequality (S.2), we can estimate the upper bound of $\mathbb{E}_t[\|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\|^2]$ in (16). This completes the proof of Lemma 1. \blacksquare

Remark 6 Huang and Chen [35] employed a general unbiased stochastic gradient estimator with bounded variance:

$$\mathbb{E} \left[\|G(\mathbf{x}, \xi_{\mathcal{I}}) - \nabla f(\mathbf{x})\|^2 \right] \leq \sigma^2/M,$$

where $G(\mathbf{x}, \xi_{\mathcal{I}}) = \frac{1}{M} \sum_{i \in \mathcal{I}} G(\mathbf{x}, \xi_i)$ represents the stochastic gradient estimator, M denotes the mini-batch size, and $\xi_{\mathcal{I}} = \{\xi_1, \xi_2, \dots, \xi_M\}$ represents a set of i.i.d. random variables.

Compared to the stochastic gradient estimator with bounded variance in [35], the variance-reduced stochastic gradient defined in Equation (6) exhibits a decrease in variance with an increasing iteration number, as shown in Equation (S.3). In contrast, the variance of the general stochastic gradient does not decrease. Our paper leverages the property of variance reduction in the gradient to achieve a linear convergence rate superior to sublinear using the KL property in subsequent analysis.

APPENDIX B

PROOF OF LEMMA 2

We first define the positive sequences $\{(h_t^s)_{t=1}^m\}_{s=1}^S$, $\{(\Gamma_t^s)_{t=1}^m\}_{s=1}^S$ and the tuple $(\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6)$ to be used in constructing the potential energy function and proving the sufficient descent inequality of the potential energy function:

$$h_t^s = \begin{cases} (2 + \alpha_1)h_{t+1}^s + [\beta_3 + (1 + \alpha_1)\beta_1], & 1 \leq t \leq m - 1, \\ \frac{5L^2}{\sigma_{\min}^A} \left(\frac{2}{\rho} + \frac{1}{2l_1} \right), & t = m, \end{cases} \quad (\text{S.4})$$

$$\Gamma_t^s = \begin{cases} \beta_2 - \beta_5 - (h_{t+1}^s + \beta_1) \left(1 + \frac{1}{\alpha_1} \right), & 1 \leq t \leq m - 1, \\ \beta_6 - \beta_5 - h_1^s, & t = m. \end{cases} \quad (\text{S.5})$$

$$\begin{aligned}
\beta_1 &= \left(\frac{\rho + l_1}{2}\right) \left(\frac{1 - \theta}{\theta}\right)^2 \sigma_{max}^A, \\
\beta_2 &= \frac{\phi_{\min}}{\eta} - \frac{L}{2} - \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) + \frac{\rho \sigma_{\min}^A (l_2 - (1 - \theta))}{2\theta^2 l_2}, \\
\beta_3 &= -\frac{\rho \sigma_{\min}^A}{2} \left[\left(\frac{1 - \theta}{\theta}\right)^2 - \frac{(1 - \theta)l_2}{\theta^2} \right] + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2}{\sigma_{\min}^A}, \\
\beta_4 &= \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2}{\sigma_{\min}^A}, \\
\beta_5 &= \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2 \eta^2 + 5\phi_{\max}^2}{\sigma_{\min}^A \eta^2}, \\
\beta_6 &= \frac{\phi_{\min}}{\eta} - \frac{L}{2} - \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} + \frac{\sigma_{\min}^A \rho}{2\theta^2} - \sigma_{max}^A \left(\frac{\rho + l_1}{2}\right) \left(\frac{1 - \theta}{\theta}\right)^2,
\end{aligned} \tag{S.6}$$

Proof: This proof is structured by two main parts which contain:

- First, we will prove that $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$ is sufficiently and monotonically decreasing over $t \in \{1, 2, \dots, m\}$ in each iteration $s \in \{1, 2, \dots, S\}$.
- Second, we will prove that $\Psi_m^s \geq \mathbb{E}_0 \Psi_1^{s+1}$ for any $s \in \{1, 2, \dots, S\}$.

For notational simplicity, we omit the superscript s in the first part, i.e., let $\mathbf{x}_t^{s+1} = \mathbf{x}_t$, $\mathbf{y}_t^{s+1} = \mathbf{y}_t$, $\boldsymbol{\lambda}_t^{s+1} = \boldsymbol{\lambda}_t$, $\tilde{\mathbf{x}}^s = \tilde{\mathbf{x}}$. By the \mathbf{y} -update (8) in Algorithm 1, we have

$$\mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t) \leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t). \tag{S.7}$$

The optimal condition of \mathbf{z} -update (9) in Algorithm 1 implies

$$\begin{aligned}
0 &= (\mathbf{x}_t - \mathbf{x}_{t+1})^T \left[\hat{\nabla} f(\mathbf{x}_t) - A^T \boldsymbol{\lambda}_t + \rho A^T (A \mathbf{z}_{t+1} + B \mathbf{y}_{t+1} - \mathbf{c}) + \frac{\theta}{\eta} Q(\mathbf{z}_{t+1} - \mathbf{z}_t) \right] \\
&= (\mathbf{x}_t - \mathbf{x}_{t+1})^T \left[\hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_t) + \nabla f(\mathbf{x}_t) - A^T \boldsymbol{\lambda}_t \right] + \rho A^T (A \mathbf{z}_{t+1} + B \mathbf{y}_{t+1} - \mathbf{c}) \\
&\quad - (\mathbf{x}_t - \mathbf{x}_{t+1})^T \frac{1}{\eta} Q(\mathbf{x}_t - \mathbf{x}_{t+1}) \\
&\leq f(\mathbf{x}_t) - f(\mathbf{x}_{t+1}) + (\mathbf{x}_t - \mathbf{x}_{t+1})^T (\hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_t)) + \frac{L}{2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 - \boldsymbol{\lambda}_t^T (A \mathbf{x}_t - A \mathbf{x}_{t+1}) \\
&\quad - \frac{1}{\eta} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|_Q^2 + \rho (A \mathbf{x}_t - A \mathbf{x}_{t+1})^T (A \mathbf{z}_{t+1} + B \mathbf{y}_{t+1} - \mathbf{c}),
\end{aligned} \tag{S.8}$$

where the inequality holds by the Assumption 1. Inserting the equality $(a - b)^T (c - d) = \frac{1}{2} (\|a - d\|^2 - \|b - d\|^2 + \|b - c\|^2 - \|c - a\|^2)$ on the term $\rho (A \mathbf{x}_t - A \mathbf{x}_{t+1})^T (A \mathbf{z}_{t+1} + B \mathbf{y}_{t+1} - \mathbf{c})$ into the above

inequality (S.8), we have the following trivial inequality

$$\begin{aligned}
0 &\leq f(\mathbf{x}_t) - f(\mathbf{x}_{t+1}) + (\mathbf{x}_t - \mathbf{x}_{t+1})^T (\hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_t)) + \frac{L}{2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 - \frac{1}{\eta} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|_Q^2 \\
&\quad - \boldsymbol{\lambda}_t^T (A\mathbf{x}_t + B\mathbf{y}_{t+1} - \mathbf{c}) + \boldsymbol{\lambda}_t^T (A\mathbf{x}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}) \\
&\quad + \frac{\rho}{2} \left(\|A\mathbf{x}_t + B\mathbf{y}_{t+1} - \mathbf{c}\|^2 + \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \|A\mathbf{x}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}\|^2 - \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \right) \\
&= \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t) - \mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t) + (\mathbf{x}_t - \mathbf{x}_{t+1})^T (\hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_t)) + \frac{L}{2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\
&\quad + \frac{\rho}{2} \left(\|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \right) - \frac{1}{\eta} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|_Q^2. \tag{S.9}
\end{aligned}$$

By inserting $-\phi_{\min} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \geq -\|\mathbf{x}_{t+1} - \mathbf{x}_t\|_Q^2$ into the above inequality (S.9), we have

$$\begin{aligned}
0 &\leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t) - \mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t) + (\mathbf{x}_t - \mathbf{x}_{t+1})^T (\hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_t)) \\
&\quad + \frac{\rho}{2} \left(\|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \right) + \frac{L}{2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 - \frac{\phi_{\min}}{\eta} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2. \tag{S.10}
\end{aligned}$$

Then, applying conditioned expectation \mathbb{E}_t on information i_t to (S.10), and using $\mathbb{E}_t[\hat{\nabla} f(\mathbf{x}_t)] = \nabla f(\mathbf{x}_t)$, we have

$$\begin{aligned}
\mathbb{E}_t[\mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t)] &\leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t) + \left(\frac{L}{2} - \frac{\phi_{\min}}{\eta} \right) \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\
&\quad + \frac{\rho}{2} \left(\mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \right). \tag{S.11}
\end{aligned}$$

By the $\boldsymbol{\lambda}$ -update (11) in Algorithm 1, and applying conditioned expectation \mathbb{E}_t on information i_t again, we have

$$\begin{aligned}
&\mathbb{E}_t[\mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_{t+1}) - \mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_t)] \\
&= \frac{1}{\rho} \mathbb{E}_t \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\|^2 + \mathbb{E}_t \langle \boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}, A\mathbf{x}_{t+1} - A\mathbf{z}_{t+1} \rangle. \tag{S.12}
\end{aligned}$$

Combine (S.7), (S.11) and (S.12) to get

$$\begin{aligned}
\mathbb{E}_t[\mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_{t+1})] &\leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) + \left(\frac{L}{2} - \frac{\phi_{\min}}{\eta} \right) \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\
&\quad + \frac{\rho}{2} \left(\mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \right) \\
&\quad + \frac{1}{\rho} \mathbb{E}_t \|\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}\|^2 + \mathbb{E}_t \langle \boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}, A\mathbf{x}_{t+1} - A\mathbf{z}_{t+1} \rangle. \tag{S.13}
\end{aligned}$$

Apply the Cauchy-Schwartz inequality to the term $\langle \boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}, A\mathbf{x}_{t+1} - A\mathbf{z}_{t+1} \rangle$ to drive

$$\langle \boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}, A\mathbf{x}_{t+1} - A\mathbf{z}_{t+1} \rangle \leq \frac{1}{2l_1} \|\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}\|^2 + \frac{l_1}{2} \|A\mathbf{x}_{t+1} - A\mathbf{z}_{t+1}\|^2, \forall l_1 > 0.$$

Combining the above inequality with (S.13), we have

$$\begin{aligned}
& \mathbb{E}_t[\mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_{t+1})] \\
& \leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) + \left(\frac{L}{2} - \frac{\phi_{\min}}{\eta}\right) \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \frac{\rho}{2} \mathbb{E}_t \left(\|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \right) \\
& \quad + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \mathbb{E}_t \|\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}\|^2 + \frac{l_1}{2} \mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 \\
& \stackrel{(i)}{\leq} \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) + \frac{\rho + l_1}{2} \mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \left[\frac{\phi_{\min}}{\eta} - \frac{L}{2} - \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \right] \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\
& \quad - \frac{\rho}{2} \mathbb{E}_t \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \left(\frac{5L^2}{\sigma_{\min}^A} \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 + \frac{5L^2}{\sigma_{\min}^A} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \right) \\
& \quad + \frac{5L^2\eta^2 + 5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2, \tag{S.14}
\end{aligned}$$

where the inequality (i) holds by Lemma 1.

Notice by the update of \mathbf{x}_{t+1} in Algorithm 1 that $\mathbf{z}_{t+1} - \mathbf{x}_t = \frac{1}{\theta}(\mathbf{x}_{t+1} - \mathbf{x}_t) + \frac{1-\theta}{\theta}(\tilde{\mathbf{x}} - \mathbf{x}_t)$ and $\mathbf{z}_{t+1} - \mathbf{x}_{t+1} = \frac{1-\theta}{\theta}(\mathbf{x}_{t+1} - \tilde{\mathbf{x}})$. Substitute these two equations into $\|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2$ and $\|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2$ to have

$$\begin{aligned}
& \frac{\rho + l_1}{2} \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \frac{\rho}{2} \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \\
& \leq \frac{\rho + l_1}{2} \sigma_{\max}^A \|\mathbf{z}_{t+1} - \mathbf{x}_{t+1}\|^2 - \frac{\rho}{2} \sigma_{\min}^A \|\mathbf{z}_{t+1} - \mathbf{x}_t\|^2 \\
& = -\frac{\rho \sigma_{\min}^A}{2} \left[\frac{1}{\theta^2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \left(\frac{1-\theta}{\theta}\right)^2 \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \right] - \frac{\rho \sigma_{\min}^A}{\theta} \left(1 - \frac{1}{\theta}\right) \langle \mathbf{x}_{t+1} - \mathbf{x}_t, \mathbf{x}_t - \tilde{\mathbf{x}} \rangle \\
& \quad + \frac{\rho + l_1}{2} \sigma_{\max}^A \left(\frac{1-\theta}{\theta}\right)^2 \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2, \tag{S.15}
\end{aligned}$$

then we can drive

$$\begin{aligned}
& \frac{\rho + l_1}{2} \|A\mathbf{z}_{t+1} - A\mathbf{x}_{t+1}\|^2 - \frac{\rho}{2} \|A\mathbf{z}_{t+1} - A\mathbf{x}_t\|^2 \\
& \stackrel{(i)}{\leq} \left(\frac{\rho + l_1}{2}\right) \left(\frac{1-\theta}{\theta}\right)^2 \sigma_{\max}^A \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2 - \frac{\rho \sigma_{\min}^A}{2} \left[\frac{1}{\theta^2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \left(\frac{1-\theta}{\theta}\right)^2 \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \right] \\
& \quad + \frac{\rho \sigma_{\min}^A}{2} \frac{(1-\theta)}{\theta^2 l_2} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \frac{\rho \sigma_{\min}^A}{2} \frac{(1-\theta) l_2}{\theta^2} \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2, \tag{S.16}
\end{aligned}$$

where the first equality holds by the equality $\mathbf{x}_{t+1} = \theta \mathbf{z}_{t+1} + (1-\theta)\tilde{\mathbf{x}}$ and the inequality (i) holds by using the Cauchy-Schwartz on the term $\langle \mathbf{x}_{t+1} - \mathbf{x}_t, \mathbf{x}_t - \tilde{\mathbf{x}} \rangle$.

Thus, by the inequalities (S.13), (S.14) and (S.16), we have

$$\begin{aligned}
& \mathbb{E}_t[\mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_{t+1})] \\
& \leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) - \left(\frac{\phi_{\min}}{\eta} - \frac{L}{2} - \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \left(\frac{1}{\rho} + \frac{1}{2l_1} \right) + \frac{\rho \sigma_{\min}^A (l_2 - (1-\theta))}{2\theta^2 l_2} \right) \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\
& \quad + \left[\left(\frac{1}{\rho} + \frac{1}{2l_1} \right) \frac{5L^2}{\sigma_{\min}^A} - \frac{\rho \sigma_{\min}^A}{2} \left(\left(\frac{1-\theta}{\theta} \right)^2 - \frac{(1-\theta)l_2}{\theta^2} \right) \right] \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \\
& \quad + \left(\frac{1}{\rho} + \frac{1}{2l_1} \right) \frac{5L^2}{\sigma_{\min}^A} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 + \left(\frac{1}{\rho} + \frac{1}{2l_1} \right) \frac{5L^2 \eta^2 + 5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2 \\
& \quad + \left(\frac{\rho + l_1}{2} \right) \left(\frac{1-\theta}{\theta} \right)^2 \sigma_{\max}^A \mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2. \tag{S.17}
\end{aligned}$$

The definitions of $\beta_1, \beta_2, \beta_3, \beta_4$, and β_5 given in Lemma 2, (S.17) can be rewritten as

$$\begin{aligned}
& \mathbb{E}_t[\mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_{t+1})] \\
& \leq \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) - \beta_2 \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \beta_3 \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 + \beta_4 \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \\
& \quad + \beta_5 \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2 + \beta_1 \mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2.
\end{aligned}$$

For notational simplicity, we let $\mathcal{L}_\rho(t) := \mathcal{L}_\rho(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)$, $\mathcal{L}_\rho(t+1) := \mathbb{E}_t \mathcal{L}_\rho(\mathbf{x}_{t+1}, \mathbf{y}_{t+1}, \boldsymbol{\lambda}_{t+1})$ then we have

$$\begin{aligned}
& \mathcal{L}_\rho(t+1) + \beta_5 \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + h \left(\mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2 + \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \right) \\
& \leq \mathcal{L}_\rho(t) - (\beta_2 - \beta_5) \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + (h + \beta_1) \mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2 + (h + \beta_3) \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \\
& \quad + \beta_4 \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 + \beta_5 \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2. \tag{S.18}
\end{aligned}$$

By the Cauchy-Schwartz inequality again, for any $\alpha_1 > 0$ we have

$$(h + \beta_1) \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2 \leq (h + \beta_1) \left\{ \left(1 + \frac{1}{\alpha_1} \right) \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + (1 + \alpha_1) \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \right\}.$$

Plugging it into (S.18) is to derive

$$\begin{aligned}
& \mathcal{L}_\rho(t+1) + \beta_5 \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + h \left(\mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|^2 + \|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 \right) \\
& \leq \mathcal{L}_\rho(t) + \beta_5 \|\mathbf{x}_t - \mathbf{x}_{t-1}\|^2 - \left[\beta_2 - \beta_5 - (h + \beta_1) \left(1 + \frac{1}{\alpha_1} \right) \right] \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 \\
& \quad - [h + \beta_3 - \beta_4 + (1 + \alpha_1) (h + \beta_1)] \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \\
& \quad + [h + \beta_3 + (1 + \alpha_1) (h + \beta_1)] \left(\|\mathbf{x}_t - \tilde{\mathbf{x}}\|^2 + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \right). \tag{S.19}
\end{aligned}$$

Using the definition of $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$, $h = h_{t+1}^s$ and inserting both (S.4) and (S.5) into (S.19), we have

$$\mathbb{E}_t \Psi_{t+1}^{s+1} \leq \Psi_t^{s+1} - [h_{t+1}^{s+1} + \beta_3 - \beta_4 + (1 + \alpha_1)(h_{t+1}^{s+1} + \beta_1)] \|\mathbf{x}_{t-1}^{s+1} - \tilde{\mathbf{x}}^s\|^2 - \Gamma_t^{s+1} \mathbb{E}_t \|\mathbf{x}_{t+1}^{s+1} - \mathbf{x}_t^{s+1}\|^2, \quad (\text{S.20})$$

for any $s \in \{0, 1, \dots, S-1\}$. Since $\Gamma_t^s > 0$, $\forall t \in \{1, 2, \dots, m\}$, we have proved the first part.

In the following, we will prove the second part. We begin by estimating the upper bound of $\mathbb{E}_0 \|\boldsymbol{\lambda}_0^{s+1} - \boldsymbol{\lambda}_1^{s+1}\|^2$.

Since $\boldsymbol{\lambda}_0^{s+1} = \boldsymbol{\lambda}_m^s$, $\mathbf{y}_0^{s+1} = \mathbf{y}_m^s$ and $\mathbf{x}_0^{s+1} = \mathbf{x}_m^s = \tilde{\mathbf{x}}^s$, we have

$$\begin{aligned} \mathbb{E}_0 \|\boldsymbol{\lambda}_0^{s+1} - \boldsymbol{\lambda}_1^{s+1}\|^2 &= \mathbb{E}_0 \|\boldsymbol{\lambda}_m^s - \boldsymbol{\lambda}_1^{s+1}\|^2 \\ &\leq (\sigma_{\min}^A)^{-1} \mathbb{E}_0 \|A^T \boldsymbol{\lambda}_m^s - A^T \boldsymbol{\lambda}_1^{s+1}\|^2 \\ &\stackrel{(i)}{=} (\sigma_{\min}^A)^{-1} \mathbb{E}_0 \|\hat{\nabla} f(\mathbf{x}_{m-1}^s) - \hat{\nabla} f(\mathbf{x}_0^{s+1}) - \frac{1}{\eta} Q(\mathbf{x}_{m-1}^s - \mathbf{x}_m^s) - \frac{1}{\eta} Q(\mathbf{x}_0^{s+1} - \mathbf{x}_1^{s+1})\|^2 \\ &\stackrel{(ii)}{=} (\sigma_{\min}^A)^{-1} \mathbb{E}_0 \|\hat{\nabla} f(\mathbf{x}_{m-1}^s) - \nabla f(\mathbf{x}_{m-1}^s) + \nabla f(\mathbf{x}_{m-1}^s) - \nabla f(\mathbf{x}_m^s) - \frac{1}{\eta} Q(\mathbf{x}_{m-1}^s - \mathbf{x}_m^s) \\ &\quad - \frac{1}{\eta} Q(\mathbf{x}_0^{s+1} - \mathbf{x}_1^{s+1})\|^2 \\ &\stackrel{(iii)}{\leq} \frac{5L^2}{\sigma_{\min}^A} \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \frac{5L^2\eta^2 + 5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_{m-1}^s - \mathbf{x}_m^s\|^2 + \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \mathbb{E}_0 \|\mathbf{x}_0^{s+1} - \mathbf{x}_1^{s+1}\|^2, \quad (\text{S.21}) \end{aligned}$$

where the equality (i) holds by the equality (S.1), the inequality (iii) holds by Assumption 1 and the (S.3), and the equality (ii) holds by the following result:

$$\begin{aligned} \hat{\nabla} f(\mathbf{x}_0^{s+1}) &= \nabla f_{i_t}(\mathbf{x}_0^{s+1}) - \nabla f_{i_t}(\tilde{\mathbf{x}}^s) + \nabla f(\tilde{\mathbf{x}}^s) \\ &= \nabla f_{i_t}(\mathbf{x}_m^s) - \nabla f_{i_t}(\mathbf{x}_m^s) + \nabla f(\mathbf{x}_m^s) \\ &= \nabla f(\mathbf{x}_m^s). \end{aligned}$$

By (S.7), we have

$$\begin{aligned} \mathcal{L}_\rho(\mathbf{x}_0^{s+1}, \mathbf{y}_1^{s+1}, \boldsymbol{\lambda}_0^{s+1}) &\leq \mathcal{L}_\rho(\mathbf{x}_0^{s+1}, \mathbf{y}_0^{s+1}, \boldsymbol{\lambda}_0^{s+1}) \\ &= \mathcal{L}_\rho(\mathbf{x}_m^s, \mathbf{y}_m^s, \boldsymbol{\lambda}_m^s). \quad (\text{S.22}) \end{aligned}$$

Similarly, using (S.11), we have

$$\begin{aligned} \mathbb{E}_0 [\mathcal{L}_\rho(\mathbf{x}_1^{s+1}, \mathbf{y}_1^{s+1}, \boldsymbol{\lambda}_0^{s+1})] &\leq \mathcal{L}_\rho(\mathbf{x}_0^{s+1}, \mathbf{y}_1^{s+1}, \boldsymbol{\lambda}_0^{s+1}) + \left(\frac{L}{2} - \frac{\phi_{\min}}{\eta}\right) \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 \\ &\quad + \frac{\rho}{2} \left(\mathbb{E}_0 \|A\mathbf{z}_1^{s+1} - A\mathbf{x}_1^{s+1}\|^2 - \mathbb{E}_0 \|A\mathbf{z}_1^{s+1} - A\mathbf{x}_0^{s+1}\|^2 \right). \quad (\text{S.23}) \end{aligned}$$

For the update of λ (11):

$$\begin{aligned} & \mathbb{E}_0[\mathcal{L}_\rho(\mathbf{x}_1^{s+1}, \mathbf{y}_1^{s+1}, \lambda_1^{s+1})] \\ & \leq \mathbb{E}_0[\mathcal{L}_\rho(\mathbf{x}_1^{s+1}, \mathbf{y}_1^{s+1}, \lambda_0^{s+1})] + \frac{l_1}{2} \mathbb{E}_0 \|A\mathbf{z}_1^{s+1} - A\mathbf{x}_1^{s+1}\|^2 + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \mathbb{E}_0 \|\lambda_1^{s+1} - \lambda_0^{s+1}\|^2, \end{aligned} \quad (\text{S.24})$$

where the inequality holds by the Cauchy-Schwartz inequality just like in (S.13). Combine (S.22), (S.23) with (S.24) and use similar tricks, we have

$$\begin{aligned} & \mathbb{E}_0[\mathcal{L}_\rho(\mathbf{x}_1^{s+1}, \mathbf{y}_1^{s+1}, \lambda_1^{s+1})] \\ & \leq \mathbb{E}_0 \left[\mathcal{L}_\rho(\mathbf{x}_0^{s+1}, \mathbf{y}_0^{s+1}, \lambda_0^{s+1}) + \left(\frac{L}{2} - \frac{\phi_{\min}}{\eta}\right) \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \|\lambda_0^{s+1} - \lambda_1^{s+1}\|^2 \right. \\ & \quad \left. + \frac{\rho + l_1}{2} \|A\mathbf{z}_1^{s+1} - A\mathbf{x}_1^{s+1}\|^2 - \frac{\rho}{2} \|A\mathbf{z}_1^{s+1} - A\mathbf{x}_0^{s+1}\|^2 \right] \\ & \stackrel{(i)}{\leq} - \left[\frac{\phi_{\min}}{\eta} - \frac{L}{2} - \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} + \frac{\sigma_{\min}^A \rho}{2\theta^2} - \left(\frac{\rho + l_1}{2}\right) \left(\frac{1-\theta}{\theta}\right)^2 \sigma_{\max}^A \right] \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 \\ & \quad + \mathcal{L}_\rho(\mathbf{x}_0^{s+1}, \mathbf{y}_0^{s+1}, \lambda_0^{s+1}) + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2}{\sigma_{\min}^A} \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 \\ & \quad + \left(\frac{1}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2 \eta^2 + 5\phi_{\max}^2}{\sigma_{\min}^A \eta^2} \|\mathbf{x}_{m-1}^s - \mathbf{x}_m^s\|^2 \\ & \leq \mathcal{L}_\rho(\mathbf{x}_0^{s+1}, \mathbf{y}_0^{s+1}, \lambda_0^{s+1}) - \beta_6 \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + \beta_4 \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \beta_5 \|\mathbf{x}_{m-1}^s - \mathbf{x}_m^s\|^2, \end{aligned} \quad (\text{S.25})$$

where β_6 is given in Lemma 2, the equality (i) holds by using the (S.21), the equality $\mathbf{x}_{t+1}^{s+1} = \theta \mathbf{z}_{t+1}^{s+1} + (1-\theta)\tilde{\mathbf{x}}^s$ and the following Cauchy-Schwartz inequality for the last two terms of the first inequality:

$$\frac{1-\theta}{\theta^2} \langle \mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}, \tilde{\mathbf{x}}^s - \mathbf{x}_0^{s+1} \rangle \leq \frac{1}{2\theta^2} \left(\|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + (1-\theta)^2 \|\tilde{\mathbf{x}}^s - \mathbf{x}_0^{s+1}\|^2 \right).$$

Also, for notational simplicity, let $\mathcal{L}_\rho(0) := \mathcal{L}_\rho(\mathbf{x}_0, \mathbf{y}_0, \lambda_0)$, $\mathcal{L}_\rho(1) := \mathbb{E}_0 \mathcal{L}_\rho(\mathbf{x}_1, \mathbf{y}_1, \lambda_1)$, and sum

$$\beta_5 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + h_1^{s+1} \left[\mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \tilde{\mathbf{x}}^s\|^2 + \|\mathbf{x}_0^{s+1} - \tilde{\mathbf{x}}^s\|^2 \right]$$

to both sides of the (S.25) then we have

$$\begin{aligned} & \mathcal{L}_\rho(1) + \beta_5 \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + h_1^{s+1} \left[\mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \tilde{\mathbf{x}}^s\|^2 + \|\mathbf{x}_0^{s+1} - \tilde{\mathbf{x}}^s\|^2 \right] \\ & \leq \mathcal{L}_\rho(0) - \beta_6 \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + \beta_4 \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \beta_5 \|\mathbf{x}_{m-1}^s - \mathbf{x}_m^s\|^2 + \beta_5 \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 \\ & \quad + h_1^{s+1} \left[\mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \tilde{\mathbf{x}}^s\|^2 + \|\mathbf{x}_0^{s+1} - \tilde{\mathbf{x}}^s\|^2 \right] \\ & = \mathcal{L}_\rho(0) - (\beta_6 - \beta_5 - h_1^{s+1}) \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 + h_1^{s+1} \left[\|\mathbf{x}_m^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 \right] \\ & \quad + \beta_5 \|\mathbf{x}_{m-1}^s - \mathbf{x}_m^s\|^2 - (h_1^{s+1} - \beta_4) \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 - h_1^{s+1} \|\mathbf{x}_m^s - \tilde{\mathbf{x}}^{s-1}\|^2. \end{aligned}$$

By the notation $h_1^{s+1} = \left(\frac{2}{\rho} + \frac{1}{2l_1}\right) \frac{5L^2}{\sigma_{\min}^A}$ in (S.4) and the definition of the sequence $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$, we have

$$\mathbb{E}_0 \Psi_1^{s+1} \leq \Psi_m^s - \Gamma_m^s \mathbb{E}_0 \|\mathbf{x}_1^{s+1} - \mathbf{x}_0^{s+1}\|^2 - \frac{5L^2}{\sigma_{\min}^A \rho} \|\mathbf{x}_{m-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2. \quad (\text{S.26})$$

Since $\Gamma_m^s > 0$, $\forall s \geq 1$, we can obtain the above result of the second part.

Thus, we prove the above conclusion. \blacksquare

APPENDIX C

PROOF OF THEOREM 1

Proof: Using the above proofs, inequalities (S.20) and (S.26), we have

$$\mathbb{E}_t^{s+1}[\Psi_{t+1}^{s+1}] \leq \Psi_t^{s+1} - \Gamma_t^{s+1} \mathbb{E}_t^{s+1}[\|\mathbf{x}_{t+1}^{s+1} - \mathbf{x}_t^{s+1}\|^2] - [h_{t+1}^{s+1} + (1 + \alpha_1)(h_{t+1}^{s+1} + \beta_1)] \|\mathbf{x}_{t-1}^{s+1} - \tilde{\mathbf{x}}^s\|^2. \quad (\text{S.27})$$

The above inequality requires $\beta_3 - \beta_4 \geq 0$ which can be ensured if we take $l_2 \geq 1 - \theta$, and

$$\mathbb{E}_0^{s+1}[\Psi_1^{s+1}] \leq \Psi_m^s - \Gamma_m^s \mathbb{E}_0^{s+1}[\|\mathbf{x}_0^{s+1} - \mathbf{x}_1^{s+1}\|^2] - \frac{5L^2}{\sigma_{\min}^A \rho} \|\mathbf{x}_m^s - \tilde{\mathbf{x}}^{s-1}\|^2. \quad (\text{S.28})$$

for any $s \in \{1, 2, \dots, S\}$ and $t \in \{1, 2, \dots, m\}$. To establish the convergence of the sequence defined in equation (24), we calculate the above conditional expectations in expressions (S.27) and (S.28). By leveraging the property $\mathbb{E}[\mathbb{E}[\cdot | \mathcal{F}_t^s]] = \mathbb{E}[\cdot]$, we then evaluate the full expectation of (S.27) and (S.28). Additionally, we sum up the resulting expressions for (S.27) and (S.28) over the ranges $t = 1, 2, \dots, m$ and $s = 1, 2, \dots, S$ to obtain

$$\mathbb{E} \Psi_m^S - \mathbb{E} \Psi_1^1 \leq -\gamma \sum_{s=1}^S \sum_{t=1}^m \mathbb{E} \|\mathbf{x}_t^s - \mathbf{x}_{t-1}^s\|^2 - \omega \sum_{s=1}^S \sum_{t=1}^m \mathbb{E} \|\mathbf{x}_{t-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2, \quad (\text{S.29})$$

where the parameter $\gamma = \min_{s,t} \Gamma_t^s$, and

$$\begin{aligned} \omega &= \min_{s,t} \{ [h_{t+1}^{s+1} + (1 + \alpha_1)(h_{t+1}^{s+1} + \beta_1)], \frac{5L^2}{\sigma_{\min}^A \rho} \} \\ &= \frac{5L^2}{\sigma_{\min}^A \rho}. \end{aligned}$$

From Assumption 2, there exists a low bound Ψ^* of the sequence $\{\Psi_t^s\}$, i.e., $\Psi_t^s \geq \Psi^*$. Using the definition of R_t^s , we have

$$R_t^{\hat{s}} = \min_{s,t} R_t^s \leq \frac{1}{\tau T} \mathbb{E}(\Psi_1^1 - \Psi^*), \quad (\text{S.30})$$

where $\tau = \min(\gamma, \omega)$ and $T = mS$. This completes the whole proof. \blacksquare

APPENDIX D

PROOF OF THE PROPERTY OF $\|\mathbf{w}_t^s - \mathbf{w}_{t+1}^s\|^2$

In this section, we establish the linear convergence rate of our ASVRG-ADMM under the so-called KL condition. We first draw the following Lemma of the property of $\|\mathbf{w}_{t+1}^s - \mathbf{w}_t^s\|^2$, where $\mathbf{w} = (\mathbf{x}, \mathbf{y}, \boldsymbol{\lambda})$ represents the sequence generated by our algorithm.

Lemma 4 *Let $\{\mathbf{w}_t^s = (\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s)\}$ be the sequence generated by ADMM Algorithm 1 under Assumptions 1-4. Then*

$$\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{w}_{t+1}^s - \mathbf{w}_t^s\|^2 < +\infty.$$

Proof: From the inequality (S.29), it follows that

$$\begin{aligned} \tau \sum_{s=1}^S \sum_{t=1}^m R_t^s &\leq \mathbb{E} \Psi_1^1 - \mathbb{E} \Psi_m^s \leq +\infty, \\ \sum_{s=1}^S \sum_{t=1}^m R_t^s &\leq +\infty, \end{aligned}$$

where $\tau = \min(\gamma, \omega)$ with γ and ω mentioned in the proof of Lemma 1. By the definitions of R_t^s in (24), we can have that

$$\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{x}_{t+1}^s - \mathbf{x}_t^s\|^2 < +\infty$$

It follows from Lemma 1 that $\boldsymbol{\lambda}$ can be bounded by R_t^s . Thus,

$$\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 < +\infty.$$

Next, we give an upper bound of

$$\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{y}_{t+1}^s - \mathbf{y}_t^s\|^2$$

based on the following equations

$$\begin{cases} \boldsymbol{\lambda}_{t+1}^s = \boldsymbol{\lambda}_t^s - \rho (A\mathbf{z}_{t+1}^s + B\mathbf{y}_{t+1}^s - c), \\ \boldsymbol{\lambda}_t^s = \boldsymbol{\lambda}_{t-1}^s - \rho (A\mathbf{z}_t^s + B\mathbf{y}_t^s - c) \\ \boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s = (\boldsymbol{\lambda}_t^s - \boldsymbol{\lambda}_{t-1}^s) + \rho (A\mathbf{z}_t^s - A\mathbf{z}_{t+1}^s) + \rho (B\mathbf{y}_t^s - B\mathbf{y}_{t+1}^s). \end{cases}$$

Together with $\mathbf{x}_{t+1} = \theta \mathbf{z}_{t+1} + (1 - \theta) \tilde{\mathbf{x}}$, simple algebra shows that

$$\begin{aligned} &\mathbb{E} \|\rho (B\mathbf{y}_t^s - B\mathbf{y}_{t+1}^s)\|^2 \\ &\leq \mathbb{E} \|(\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s) - (\boldsymbol{\lambda}_t^s - \boldsymbol{\lambda}_{t-1}^s) - \rho (A\mathbf{z}_t^s - A\mathbf{z}_{t+1}^s)\|^2 \\ &\leq 3\mathbb{E} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 + 3\mathbb{E} \|\boldsymbol{\lambda}_t^s - \boldsymbol{\lambda}_{t-1}^s\|^2 + 3\rho^2 \frac{\sigma_{\max}^A}{\theta^2} \mathbb{E} \|\mathbf{x}_{t+1}^s - \mathbf{x}_t^s\|^2 \end{aligned} \quad (\text{S.31})$$

Then we can use (S.31) drive

$$\mathbb{E} \|\mathbf{y}_t^s - \mathbf{y}_{t+1}^s\|^2 \leq \frac{3\bar{M}^2}{\rho^2} \mathbb{E} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 + \frac{3\bar{M}^2}{\rho^2} \mathbb{E} \|\boldsymbol{\lambda}_t^s - \boldsymbol{\lambda}_{t-1}^s\|^2 + \frac{3\bar{M}^2 \sigma_{\max}^A}{\theta^2} \mathbb{E} \|\mathbf{x}_{t+1}^s - \mathbf{x}_t^s\|^2,$$

where the inequality is derived from [27, Lemma 1] with the Assumption 4.

We set $\zeta_{11} = \frac{3\bar{M}^2}{\rho^2}$ and $\zeta_{12} = \frac{3\bar{M}^2 \sigma_{\max}^A}{\theta^2}$, such that

$$\begin{aligned} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{y}_t^s - \mathbf{y}_{t+1}^s\|^2 &\leq \zeta_{11} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 + \zeta_{11} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t-1}^s - \boldsymbol{\lambda}_t^s\|^2 \\ &\quad + \zeta_{12} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{x}_t^s - \mathbf{x}_{t+1}^s\|^2. \end{aligned} \quad (\text{S.32})$$

Recall that R_t^s defined in (24) is

$$R_t^s := \mathbb{E} [\|\mathbf{x}_t^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \|\mathbf{x}_{t-1}^s - \tilde{\mathbf{x}}^{s-1}\|^2 + \|\mathbf{x}_{t+1}^s - \mathbf{x}_t^s\|^2 + \|\mathbf{x}_t^s - \mathbf{x}_{t-1}^s\|^2].$$

By Lemma 1, the first term

$$\zeta_{11} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 + \zeta_{11} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t-1}^s - \boldsymbol{\lambda}_t^s\|^2$$

can be bounded as follows:

$$\begin{aligned} &\zeta_{11} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 + \zeta_{11} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t-1}^s - \boldsymbol{\lambda}_t^s\|^2 \\ &\leq \zeta_{13} R_t^s, \end{aligned} \quad (\text{S.33})$$

where $\zeta_{13} = 2\zeta_{11} \max \left\{ \frac{5L^2}{\sigma_{\min}^A}, \frac{5\phi_{\max}^2}{\sigma_{\min}^A \eta^2}, \frac{5(\eta^2 L^2 + \phi_{\max}^2)}{\sigma_{\min}^A \eta^2} \right\}$. The second term $\zeta_{12} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{x}_t^s - \mathbf{x}_{t+1}^s\|^2$ in (S.32) can be bounded by

$$\zeta_{12} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{x}_t^s - \mathbf{x}_{t+1}^s\|^2 \leq \zeta_{12} R_t^s. \quad (\text{S.34})$$

Every term on the right-hand side of (S.32) can be bounded by R_t^s . Thus, there exists $\zeta_{14} > 0$ such that the upper bound of all above terms on the right-hand side can be limited by

$$\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{y}_t^s - \mathbf{y}_{t+1}^s\|^2 \leq \zeta_{14} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} R_t^s < +\infty, \quad (\text{S.35})$$

where $\zeta_{14} = \zeta_{12} + \zeta_{13}$. As a result, we obtain

$$\begin{aligned} &\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{y}_t^s - \mathbf{y}_{t+1}^s\|^2 < +\infty, \\ &\sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \mathbb{E} \|\mathbf{w}_t^s - \mathbf{w}_{t+1}^s\|^2 < +\infty. \end{aligned}$$

This completes the whole proof. ■

APPENDIX E
PROOF OF LEMMA 5

Now, based on Lemma 4 we can demonstrate the upper bound of $\mathbb{E} \|\partial\mathcal{L}_\rho(\mathbf{w}_{t+1})\|$ which is important for the linear convergence of ASVRG-ADMM.

Lemma 5 *Let $\{\mathbf{w}_t^s = (\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s)\}$ be the sequence generated by Algorithm 1 under Assumptions 1-4. For notational simplicity, we omit the upper script s with setting $(\mathbf{x}_j, \mathbf{y}_j, \boldsymbol{\lambda}_j) := (\mathbf{x}_t^s, \mathbf{y}_t^s, \boldsymbol{\lambda}_t^s)$, where $j = s * m + t$. Then, there exists $\xi_1 > 0$ such that*

$$\begin{aligned} & \mathbb{E}_t \|\partial\mathcal{L}_\rho(\mathbf{w}_{t+1})\| \\ & \leq \xi_1 (\mathbb{E}_t[\|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|] + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \mathbf{x}_{t-1}\| + \mathbb{E}_t[\|\mathbf{x}_{t+1} - \mathbf{x}_t\|]). \end{aligned}$$

Lemma 5 shows that $\|\partial\mathcal{L}_\rho(\mathbf{w}_{t+1})\|$ in the Definition 2 deducing the linear convergence with the KL property, is upper bounded by some primal/iterative residuals. Based on this Lemma, we will show that the sequence $\{\mathbf{w}_t^s\}$ converges to a critical point of the problem (1).

Proof: From the definition of $\mathcal{L}_\rho(\cdot)$ in (4), it follows

$$\begin{aligned} \frac{\partial\mathcal{L}_\rho(\mathbf{w}_{t+1})}{\partial\mathbf{x}} &= \nabla f(\mathbf{x}_{t+1}) - A^\top \boldsymbol{\lambda}_{t+1} + \rho A^\top (A\mathbf{x}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}), \\ \frac{\partial\mathcal{L}_\rho(\mathbf{w}_{t+1})}{\partial\mathbf{y}} &= \partial g(\mathbf{y}_{t+1}) - B^\top \boldsymbol{\lambda}_{t+1} + \rho B^\top (A\mathbf{x}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}), \\ \frac{\partial\mathcal{L}_\rho(\mathbf{w}_{t+1})}{\partial\boldsymbol{\lambda}} &= - (A\mathbf{x}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}). \end{aligned}$$

Recalling the first-order optimality conditions of the subproblems in Algorithm1 together with the update of $\boldsymbol{\lambda}_{t+1}$, we have

$$\begin{aligned} \mathbf{x} : \quad & \nabla \hat{f}(\mathbf{x}_t) = A^\top \boldsymbol{\lambda}_{t+1} - \frac{\theta}{\eta} Q(\mathbf{z}_{t+1} - \mathbf{z}_t), \\ \mathbf{y} : \quad & B^\top \boldsymbol{\lambda}_{t+1} - \rho B^\top A(\mathbf{x}_t - \mathbf{z}_{t+1}) \in \partial g(\mathbf{y}_{t+1}), \\ \boldsymbol{\lambda} : \quad & \boldsymbol{\lambda}_{t+1} = \boldsymbol{\lambda}_t - \rho (A\mathbf{z}_{t+1} + B\mathbf{y}_{t+1} - \mathbf{c}). \end{aligned} \tag{S.36}$$

Invoking the above optimality conditions of the Algorithm1 yields

$$\frac{\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t}{\rho} + A \frac{1-\theta}{\theta} (\mathbf{x}_{t+1} - \tilde{\mathbf{x}}) \in \partial_{\boldsymbol{\lambda}} \mathcal{L}_\rho(\mathbf{w}_{t+1}), \tag{S.37}$$

$$B^\top (\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}) - \rho B^\top A(\mathbf{z}_t - \mathbf{x}_{t+1}) \in \partial_{\mathbf{y}} \mathcal{L}_\rho(\mathbf{w}_{t+1}), \tag{S.38}$$

$$\begin{aligned} & \nabla f(\mathbf{x}_{t+1}) - \nabla \hat{f}(\mathbf{x}_t) + \frac{1}{\eta} Q(\mathbf{x}_t - \mathbf{x}_{t+1}) + A^\top (\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}) \\ & + \rho \frac{1-\theta}{\theta} A^\top A(\tilde{\mathbf{x}} - \mathbf{x}_{t+1}) \in \partial_{\mathbf{x}} \mathcal{L}_\rho(\mathbf{w}_{t+1}). \end{aligned} \tag{S.39}$$

Thus, we can obtain

$$\begin{aligned}
& \mathbb{E}_t \text{dist}(0, \partial \mathcal{L}_\rho(\mathbf{w}_{t+1})) \\
&= \mathbb{E}_t \|\partial_\lambda \mathcal{L}_\rho(\mathbf{w}_{t+1})\| + \mathbb{E}_t \|\partial_y \mathcal{L}_\rho(\mathbf{w}_{t+1})\| + \mathbb{E}_t \|\partial_x \mathcal{L}_\rho(\mathbf{w}_{t+1})\| \\
&\leq \mathbb{E}_t \left[\left\| \frac{\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t}{\rho} \right\| + \left\| A \frac{1-\theta}{\theta} (\mathbf{x}_{t+1} - \tilde{\mathbf{x}}) \right\| + \left\| B^\top (\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}) \right\| + \left\| \rho B^\top A (\mathbf{z}_t - \mathbf{x}_{t+1}) \right\| \right. \\
&\quad \left. + \left\| \nabla f(\mathbf{x}_{t+1}) - \nabla f(\mathbf{x}_t) \right\| + \left\| \rho \frac{1-\theta}{\theta} A^\top A (\tilde{\mathbf{x}} - \mathbf{x}_{t+1}) \right\| + \left\| \frac{1}{\eta} Q (\mathbf{x}_t - \mathbf{x}_{t+1}) \right\| + \left\| A^\top (\boldsymbol{\lambda}_t - \boldsymbol{\lambda}_{t+1}) \right\| \right] \\
&\leq \mathbb{E}_t \left[\zeta_{21} \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\| + \zeta_{22} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \zeta_{23} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \left\| \hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_{t+1}) \right\| \right] \\
&\leq \zeta_{24} \mathbb{E}_t \left(\|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\| + \left\| \hat{\nabla} f(\mathbf{x}_t) - \nabla f(\mathbf{x}_{t+1}) \right\| \right) \\
&\stackrel{(i)}{\leq} \zeta_{24} \left(\mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E}_t \left\| \hat{\nabla} f(\mathbf{x}_{t+1}) - \nabla f(\mathbf{x}_t) \right\| \right) \\
&\quad + \zeta_{24} \zeta_{25} \mathbb{E}_t \|\mathbf{x}_{t-1} - \mathbf{x}_t\| + \zeta_{24} \zeta_{25} \left(\|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\| \right), \tag{S.40}
\end{aligned}$$

where σ_{max}^B is the largest positive eigenvalue of $B^\top B$ (or equivalently the smallest positive eigenvalue of BB^\top), ϕ_{max} is the largest positive eigenvalue of the matrix Q , and the inequality (i) is due to Lemma 1 and the inequality $\sqrt{a^2 + b^2 + c^2 + d^2} \leq a + b + c + d$, for any $a, b, c, d \geq 0$.

We can further obtain

$$\begin{aligned}
& \mathbb{E}_t [\text{dist}(0, \partial \mathcal{L}_\rho(\mathbf{w}^{t+1}))] \\
&\stackrel{(i)}{\leq} \zeta_{24} L [\|\mathbf{x}_t - \tilde{\mathbf{x}}\|] + \zeta_{24} L \mathbb{E}_t [\|\mathbf{x}_{t+1} - \mathbf{x}_t\|] + \zeta_{24} \mathbb{E}_t \left(\|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t+1} - \mathbf{x}_t\| \right. \\
&\quad \left. + \zeta_{25} (\|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \|\mathbf{x}_{t-1} - \mathbf{x}_t\|) \right) \\
&\leq (\zeta_{24} L + \zeta_{24} + \zeta_{24} \zeta_{25}) [\|\mathbf{x}_t - \tilde{\mathbf{x}}\|] + \zeta_{24} \mathbb{E}_t [\|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\|] + \zeta_{24} \zeta_{25} [\|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|] \\
&\quad + (\zeta_{24} L + \zeta_{24} + \zeta_{24} \zeta_{25}) \mathbb{E}_t [\|\mathbf{x}_{t+1} - \mathbf{x}_t\|] + \zeta_{24} \zeta_{25} [\|\mathbf{x}_t - \mathbf{x}_{t-1}\|], \\
&\leq \xi_1 (\mathbb{E}_t \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \mathbf{x}_{t-1}\|),
\end{aligned}$$

where the parameters ζ_{21}, ζ_{22} and ζ_{23} are

$$\left\{ \begin{array}{l} \zeta_{21} = \frac{1}{\rho} + \sqrt{\sigma_{max}^B} + \sqrt{\sigma_{max}^A}, \\ \zeta_{22} = \rho \frac{1-\theta}{\theta} \sigma_{max}^A, \\ \zeta_{23} = \frac{\phi_{max}}{\eta} + \rho \sqrt{\sigma_{max}^B} \sigma_{max}^A, \\ \zeta_{24} = \max \{ \zeta_{21}, \zeta_{22}, \zeta_{23}, 1 \}, \\ \zeta_{25} = \max \left\{ \sqrt{\frac{5L^2}{\sigma_{min}^A}}, \sqrt{\frac{5\phi_{max}^2}{\sigma_{min}^A \eta^2}}, \sqrt{\frac{5(\eta^2 L^2 + \phi_{max}^2)}{\sigma_{min}^A \eta^2}} \right\}, \\ \xi_1 = \zeta_{24} L + \zeta_{24} + \zeta_{24} \zeta_{25}, \end{array} \right. \tag{S.41}$$

where the inequality (i) is due to the triangle inequality $\|a + b\| \leq \|a\| + \|b\|$, Assumption 1, the inequality $(\mathbb{E}\|\mathbf{x}\|)^2 \leq \mathbb{E}(\|\mathbf{x}\|^2)$ and the inequality (S.3).

This completes the proof. \blacksquare

APPENDIX F

PROOF OF LEMMA 3

The convergence properties of the stochastic sequence $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ under the Kurdyka-Lojasiewicz (KL) inequality condition will be investigated in the following sections. It is important to acknowledge that the implementation of the KL technique displays slight differences between stochastic and deterministic algorithms. For further details, readers are encouraged to refer to [30], [31].

Before proving the key Lemma 3, we first prove the Lemma 6.

Lemma 6 *Let $\{\mathbf{w}_t = (\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ (for notational simplicity, we omit the label s) be the stochastic sequence generated by ADMM procedure. Let $S(\mathbf{w}_0)$ denote the set of its limit points. With Definition 1, then we have*

- i) $S(\mathbf{w}_0)$ is a.s. a nonempty compact set, and $\text{dist}(\mathbf{w}_t, S(\mathbf{w}_0))$ converges a.s. to 0;
- ii) $S(\mathbf{w}_0) \subset \text{crit } \mathcal{L}_\rho$ a.s.;
- iii) $\mathcal{L}_\rho(\cdot)$ is a.s. finite and constant on $S(\mathbf{w}_0)$, equal to $\inf_{t \in \mathbb{N}} \mathcal{L}_\rho(\mathbf{w}_t) = \lim_{t \rightarrow +\infty} \mathcal{L}_\rho(\mathbf{w}_t)$ a.s.

Proof: We prove the results item by item.

i) By applying the descent inequalities (S.27) and (S.28) along with the supermartingale convergence theorem, we can establish that

$$\left\{ \begin{array}{l} \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \|\mathbf{x}_{t+1}^s - \mathbf{x}_t^s\|^2 < +\infty \quad \text{a.s.}, \\ \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \|\mathbf{y}_{t+1}^s - \mathbf{y}_t^s\|^2 < +\infty \quad \text{a.s.}, \\ \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \|\boldsymbol{\lambda}_{t+1}^s - \boldsymbol{\lambda}_t^s\|^2 < +\infty \quad \text{a.s.}, \\ \sum_{s=1}^{+\infty} \sum_{t=1}^{+\infty} \|\mathbf{w}_{t+1}^s - \mathbf{w}_t^s\|^2 < +\infty \quad \text{a.s.}, \end{array} \right. \quad (\text{S.42})$$

then we can further have $\|\mathbf{w}_{t+1} - \mathbf{w}_t\| \rightarrow 0$ a.s.

Consequently, for any sequence satisfying $\|\mathbf{w}_{t+1} - \mathbf{w}_t\| \rightarrow 0$ a.s., claim i) holds. And we refer to [31, proposition 2.3], [7, Lemma A.14] for more details.

ii) Let $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*) \in S(\mathbf{w}_0)$, then there exists a subsequence $\{(\mathbf{x}_{t_j}, \mathbf{y}_{t_j}, \boldsymbol{\lambda}_{t_j})\}$ of $\{(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ converging a.s. to $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$. Note that (S.42) implies

$$\|\mathbf{w}_{t+1} - \mathbf{w}_t\| \rightarrow 0 \quad \text{a.s.}, \quad (\text{S.43})$$

which means \mathbf{w}_{t+1} converges a.s. to \mathbf{w}_t and $\{(\mathbf{x}_{t_j+1}, \mathbf{y}_{t_j+1}, \boldsymbol{\lambda}_{t_j+1})\}$ also converges a.s. to $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$. It is important to highlight that the convergence of random variables in an almost sure manner is observed in $\mathbf{w}_{t+1} \rightarrow \mathbf{w}_t$ a.s. As a result, any newly derived conclusions based on this convergence will also hold with an almost sure guarantee. As a result, many of the conclusions presented in the subsequent chapters differ from those in the previous chapters but can be considered almost surely valid. Since \mathbf{y}_{t_j+1} is a minimizer of $\mathcal{L}_\rho(\mathbf{x}_{t_j}, \mathbf{y}, \boldsymbol{\lambda}_{t_j})$ for the variable \mathbf{y} , it holds that

$$\mathcal{L}_\rho(\mathbf{x}_{t_j}, \mathbf{y}_{t_j+1}, \boldsymbol{\lambda}_{t_j}) \leq \mathcal{L}_\rho(\mathbf{x}_{t_j}, \mathbf{y}^*, \boldsymbol{\lambda}_{t_j}). \quad (\text{S.44})$$

Then, it follows from Equations (S.43), (S.44) and the continuity of $\mathcal{L}_\rho(\cdot)$ with respect to \mathbf{x} and $\boldsymbol{\lambda}$ that

$$\begin{aligned} & \limsup_{j \rightarrow +\infty} \mathcal{L}_\rho(\mathbf{x}_{t_j}, \mathbf{y}_{t_j+1}, \boldsymbol{\lambda}_{t_j}) \\ &= \limsup_{j \rightarrow +\infty} \mathcal{L}_\rho(\mathbf{x}_{t_j+1}, \mathbf{y}_{t_j+1}, \boldsymbol{\lambda}_{t_j+1}) \leq \mathcal{L}_\rho(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*) \text{ a.s.} \end{aligned} \quad (\text{S.45})$$

On the other hand, by the lower semicontinuity of $\mathcal{L}_\rho(\cdot)$, we know

$$\liminf_{j \rightarrow +\infty} \mathcal{L}_\rho(\mathbf{x}_{t_j+1}, \mathbf{y}_{t_j+1}, \boldsymbol{\lambda}_{t_j+1}) \geq \mathcal{L}_\rho(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*) \text{ a.s.} \quad (\text{S.46})$$

The above two relations (S.45) and (S.46) show that $\lim_{j \rightarrow +\infty} g(\mathbf{y}_{t_j+1}) = g(\mathbf{y}^*)$ a.s. Because of the continuity of ∇f and the closeness of ∂g , taking limits in Equation (S.36) along the subsequence $\{(\mathbf{x}_{t_j+1}, \mathbf{y}_{t_j+1}, \boldsymbol{\lambda}_{t_j+1})\}$ and using Equation (S.43) again, we almost surely have

$$\begin{aligned} A^T \boldsymbol{\lambda}^* &= \partial f(\mathbf{x}^*), \\ B^T \boldsymbol{\lambda}^* &\in \nabla g(\mathbf{y}^*), \\ A\mathbf{x}^* + B\mathbf{y}^* - \mathbf{c} &= 0. \end{aligned}$$

Then, $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ is a.s. a critical point of the problem (1), hence $w^* \in \text{crit } \mathcal{L}_\beta$ a.s.

iii) For any point $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*) \in S(w^0)$, there exists a subsequence $\{(\mathbf{x}_{t_j}, \mathbf{y}_{t_j}, \boldsymbol{\lambda}_{t_j})\}$ of $\{(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ converging to $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$. Combining Equations (S.45) and (S.46), we obtain

$$\lim_{j \rightarrow +\infty} \mathcal{L}_\rho(\mathbf{x}_{t_j}, \mathbf{y}_{t_j}, \boldsymbol{\lambda}_{t_j}) = \mathcal{L}_\rho(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*) \quad \text{a.s.}$$

Therefore, $\mathcal{L}_\rho(\cdot)$ is a.s. constant on $S(w^0)$. Moreover, $\inf_{t \in N} \mathcal{L}_\rho(\mathbf{w}_t) = \lim_{t \rightarrow +\infty} \mathcal{L}_\rho(\mathbf{w}_t)$ a.s. ■

With the established conclusions, we are now prepared to give the proof of Lemma 3.

Proof: From the proof of Lemma 6, it follows that $\mathcal{L}_\rho(\mathbf{w}_t) \xrightarrow{\text{a.s.}} \mathcal{L}_\rho(\mathbf{w}^*)$ for all $\mathbf{w}^* \in S(w^0)$.

We consider two cases.

(i) We first consider this case: there exists a finite positive discrete random variable t_0 for which $\Psi_{t_0} = \mathcal{L}_\rho(\mathbf{w}^*)$ a.s.

Taking full expectation operator to the inequality (S.27), we can have

$$\begin{aligned} \mathbb{E}\Psi_{t+1} &\stackrel{(i)}{\leq} \mathbb{E}\Psi_t - \Gamma_t \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 - w \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2, \\ \Gamma_t \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + w \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 &\leq \mathbb{E}\Psi_t - \mathbb{E}\Psi_{t+1} \leq \mathbb{E}\Psi_{t_0} - \mathbb{E}\mathcal{L}_\rho(\mathbf{w}^*) = 0 \quad \text{a.s.}, \end{aligned} \quad (\text{S.47})$$

where the parameters Γ_t, w are defined in (S.29). Then we can drive

$$\begin{aligned} \Gamma_t \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + w \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 &= 0, \\ \mathbf{x}_{t+1} = \mathbf{x}_t, \mathbf{x}_{t-1} = \tilde{\mathbf{x}} &\quad \text{a.s.}, \\ \boldsymbol{\lambda}_{t+1} = \boldsymbol{\lambda}_t, \mathbf{y}_{t+1} = \mathbf{y}_t &\quad \text{a.s.} \end{aligned} \quad (\text{S.48})$$

Thus, for any $t \geq t_0$, we have $\mathbf{x}_{t+1} = \mathbf{x}_t$, $\mathbf{x}_{t-1} = \tilde{\mathbf{x}}$, $\boldsymbol{\lambda}_{t+1} = \boldsymbol{\lambda}_t$ and $\mathbf{y}_{t+1} = \mathbf{y}_t$ a.s. Then the assertion holds.

(ii) We define $\Pi = \liminf_{t \rightarrow +\infty} \{\omega \in \Omega \mid \Psi_t = \Psi_t(\omega) > \mathcal{L}_\rho(\mathbf{w}^*(\omega))\}$. We assume $\mathbb{P}(\Pi) = 1$ and $\Psi_t > \mathcal{L}_\rho(\mathbf{w}^*)$ for all t over the set Π . Since $\text{dist}(\mathbf{w}_t, S(\mathbf{w}^0)) \xrightarrow{\text{a.s.}} 0$, it follows that for all $\epsilon > 0$, there exists a finite positive discrete random variable $k_1 > 0$, such that for any $t > k_1$, $\text{dist}(\mathbf{w}_t, S(\mathbf{w}^0)) < \epsilon$ a.s. Again since $\Psi_t \xrightarrow{\text{a.s.}} \mathcal{L}_\rho(\mathbf{w}^*)$, it follows for all $\eta > 0$, there exists a finite positive discrete random variable $k_2 > 0$, such that for any $t > k_2$, $\Psi_t < \mathcal{L}_\rho(\mathbf{w}^*) + \eta$ a.s. Consequently, for all $\epsilon, \eta > 0$, when $t > \tilde{k} = \max\{k_1, k_2\}$, we almost surely have $\text{dist}(\mathbf{w}_t, S(\mathbf{w}^0)) < \epsilon$, $\mathcal{L}_\rho(\mathbf{w}^*) < \Psi_t < \mathcal{L}_\rho(\mathbf{w}^*) + \eta$. Since $S(\mathbf{w}^0)$ is a.s. a nonempty compact set and $\mathcal{L}_\rho(\cdot)$ is a.s. constant on $S(\mathbf{w}^0)$, applying the definition of KL property with $\Omega = S(\mathbf{w}^0)$, we deduce that for any $t > \tilde{k}$

$$\nabla\varphi(\Psi_t - \mathcal{L}_\rho(\mathbf{w}^*)) \text{dist}(0, \partial\Psi_t) \geq 1 \quad \text{a.s.}$$

We denote the $\Psi^* = \mathcal{L}_\rho(\mathbf{w}^*)$, then the above inequality becomes

$$\nabla\varphi(\Psi_t - \Psi^*) \text{dist}(0, \partial\Psi_t) \geq 1 \quad \text{a.s.} \quad (\text{S.49})$$

From the concavity of φ , we get that

$$\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t\Psi_{t+1} - \Psi^*) \geq \nabla\varphi(\Psi_t - \Psi^*)(\Psi_t - \mathbb{E}_t\Psi_{t+1}) \quad \text{a.s.}, \quad (\text{S.50})$$

and it amount to

$$\begin{aligned} \Psi_t - \mathbb{E}_t\Psi_{t+1} &\leq \frac{\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t\Psi_{t+1} - \Psi^*)}{\nabla\varphi(\Psi_t - \Psi^*)} \\ &\stackrel{(i)}{\leq} [\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t\Psi_{t+1} - \Psi^*)] \times \text{dist}(0, \partial\Psi_t) \quad \text{a.s.}, \end{aligned}$$

where the inequality (i) holds by the inequality (S.49), and we set

$$\Pi_{p,q} := \varphi(\Psi_p - \Psi^*) - \varphi(\Psi_q - \Psi^*). \quad (\text{S.51})$$

Moreover, recalling the equations (S.27) and (S.28), we have

$$\Psi_t - \mathbb{E}_t \Psi_{t+1} \geq \tau \left(\mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \right), \quad (\text{S.52})$$

where $\tau = \min(\gamma, \omega)$ with γ and ω being given by Theorem 1.

From the definition of the sequence $\{(\Psi_t^s)_{t=1}^m\}_{s=1}^S$, we obtain

$$\begin{aligned} \text{dist}(0, \partial\Psi_t) &= \text{dist}(0, (\partial\mathcal{L}_\rho(\mathbf{w}_t) + 2h_t(\mathbf{x}_t - \tilde{\mathbf{x}}) + 2\beta_5(\mathbf{x}_t - \mathbf{x}_{t-1}))) \\ &\leq \text{dist}(0, \partial\mathcal{L}_\rho(\mathbf{w}_t)) + 2h_t \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + 2\beta_5 \|\mathbf{x}_t - \mathbf{x}_{t-1}\|. \end{aligned} \quad (\text{S.53})$$

Combine the (S.50)-(S.53) we have

$$\begin{aligned} &\tau \left(\mathbb{E}_t \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \right) \\ &\leq \left[\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t \Psi_{t+1} - \Psi^*) \right] \left[\text{dist}(0, \partial\mathcal{L}_\rho(\mathbf{w}_t)) + 2h_t \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + 2\beta_5 \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right] \quad \text{a.s.} \end{aligned} \quad (\text{S.54})$$

Taking the full expectation and inserting Lemma 5 into the above inequality, we see that

$$\begin{aligned} &\mathbb{E}[\text{dist}(0, \partial\Psi_t)] + 2h_t \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + 2\beta_5 \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \\ &\leq \xi_{max} \left[\mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right], \end{aligned} \quad (\text{S.55})$$

where $\xi_{max} = \xi_1 + 2h_t + 2\beta_5$ and the inequality (S.55) becomes

$$\begin{aligned} \mathbb{E}[\text{dist}(0, \partial\Psi_t)] &\leq \xi_{max} \left[\mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| \right. \\ &\quad \left. + \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right]. \end{aligned} \quad (\text{S.56})$$

Now, combine (S.54) and (S.56) and take the full expectation on (S.54) to obtain

$$\begin{aligned} &\tau \left(\mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|^2 \right) \\ &\leq \xi_{max} \mathbb{E} \left[\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t \Psi_{t+1} - \Psi^*) \right] \mathbb{E} \left[\|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{t+1} - \mathbf{x}_t\| \right. \\ &\quad \left. + \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right] \quad \text{a.s.} \end{aligned}$$

It follows from the inequality $\sqrt{a^2 + b^2} \geq \frac{\sqrt{2}}{2}(a + b)$ that

$$\begin{aligned}
& 4(\mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|) \\
& \leq 4\sqrt{\xi_{max}} \sqrt{\frac{2}{\tau}} \left[\mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right]^{\frac{1}{2}} \\
& \quad \times \left\{ \mathbb{E} [\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t \Psi_{t+1} - \Psi^*)] \right\}^{\frac{1}{2}} \quad \text{a.s.}, \tag{S.57}
\end{aligned}$$

which by the Cauchy-Schwartz inequality $2\sqrt{ab} \leq a + b$ further gives

$$\begin{aligned}
& 4(\mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\|) \\
& \leq \left(2\sqrt{\xi_{max}} \sqrt{\frac{2}{\tau}} \right)^2 \mathbb{E} [\varphi(\Psi_t - \Psi^*) - \varphi(\mathbb{E}_t \Psi_{t+1} - \Psi^*)] + \left[\mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| \right. \\
& \quad \left. + \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right], \\
& \stackrel{(i)}{\leq} \left(2\sqrt{\xi_{max}} \sqrt{\frac{2}{\tau}} \right)^2 \mathbb{E} [\varphi(\Psi_t - \Psi^*) - \varphi(\Psi_{t+1} - \Psi^*)] + \left[\mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| \right. \\
& \quad \left. + \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right], \\
& = \left(2\sqrt{\xi_{max}} \sqrt{\frac{2}{\tau}} \right)^2 \mathbb{E}(\Pi_{t,t+1}) + \left[\mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{x}_t - \tilde{\mathbf{x}}\| \right. \\
& \quad \left. + \mathbb{E} \|\mathbf{x}_t - \mathbf{x}_{t-1}\| \right] \quad \text{a.s.} \tag{S.58}
\end{aligned}$$

In the above analysis, we use the abbreviation of $\mathbb{E}[\cdot | \mathcal{F}_t]$ as $\mathbb{E}_t[\cdot]$, and apply the conditional Jensen's inequality to concave function φ as

$$\begin{aligned}
& \mathbb{E} [\varphi(\mathbb{E}_t \Psi_{t+1} - \Psi^*)] = \mathbb{E} [\varphi(\mathbb{E} \Psi_{t+1} - \Psi^* | \mathcal{F}_t)] \\
& \geq \mathbb{E} [\mathbb{E} [\varphi(\Psi_{t+1} - \Psi^*) | \mathcal{F}_t]] = \mathbb{E} [\varphi(\Psi_{t+1} - \Psi^*)].
\end{aligned}$$

Summing up (S.58) from $t = \tilde{t} + 1, \dots, m$ yields

$$\begin{aligned}
& 2\mathbb{E} \|\mathbf{x}_{\tilde{t}} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{\tilde{t}+1} - \tilde{\mathbf{x}}\| + 2 \sum_{t=\tilde{t}+1}^m \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \sum_{t=\tilde{t}+1}^m \mathbb{E} \|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{m+1} - \mathbf{x}_m\| \\
& \leq 2\mathbb{E} \|\mathbf{x}_m - \tilde{\mathbf{x}}\| + \mathbb{E} \|\mathbf{x}_{m+1} - \tilde{\mathbf{x}}\| + (2\sqrt{\xi_{max}} \sqrt{\frac{2}{\tau}})^2 \times \mathbb{E} [\varphi(\psi_{\tilde{t}+1} - \psi^*) - \varphi(\psi_{m+1} - \psi^*)] \\
& \quad + \mathbb{E} \|\mathbf{x}_{\tilde{t}+1} - \mathbf{x}_{\tilde{t}}\| \\
& < +\infty \quad \text{a.s.} \tag{S.59}
\end{aligned}$$

Let m be $+\infty$, by the equation (S.48), it yields

$$\sum_{t=\tilde{t}+1}^{+\infty} \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| < +\infty \quad \text{a.s.} \quad \text{and} \quad \sum_{t=\tilde{t}+1}^{+\infty} \mathbb{E} \|\mathbf{x}_{t+1} - \tilde{\mathbf{x}}\| < +\infty \quad \text{a.s.}$$

Similarly we can obtain that

$$\sum_{t=\tilde{t}+1}^{+\infty} \mathbb{E} \|\mathbf{y}_{t+1} - \mathbf{y}_t\| < +\infty \quad \text{a.s.} \quad \text{and} \quad \sum_{t=\tilde{t}+1}^{+\infty} \mathbb{E} \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\| < +\infty \quad \text{a.s.}$$

Hence, $\sum_{k=1}^{+\infty} \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| < +\infty$ a.s. Besides, we note that

$$\begin{aligned} & \mathbb{E} \|\mathbf{w}_{t+1} - \mathbf{w}_t\| \\ &= \mathbb{E} \left(\|\mathbf{x}_{t+1} - \mathbf{x}_t\|^2 + \|\mathbf{y}_{t+1} - \mathbf{y}_t\|^2 + \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\|^2 \right)^{1/2} \\ &\leq \mathbb{E} \|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \mathbb{E} \|\mathbf{y}_{t+1} - \mathbf{y}_t\| + \mathbb{E} \|\boldsymbol{\lambda}_{t+1} - \boldsymbol{\lambda}_t\| \quad \text{a.s.} \end{aligned}$$

Consequently,

$$\begin{aligned} & \sum_{t=0}^{+\infty} \mathbb{E} \|\mathbf{w}_{t+1} - \mathbf{w}_t\| < +\infty \quad \text{a.s.}, \\ & \sum_{t=0}^{+\infty} \|\mathbf{w}_{t+1} - \mathbf{w}_t\| < +\infty \quad \text{a.s.} \end{aligned} \tag{S.60}$$

This completes the whole proof. ■

Remark 7 *In Lemma 1, we have presented the framework of convergence analysis and obtained*

$$\sum_{t=0}^{+\infty} \|\mathbf{w}_{t+1} - \mathbf{w}_t\|^2 < +\infty \quad \text{a.s.}$$

However, it is unclear whether their results can be extended to

$$\sum_{t=0}^{+\infty} \|\mathbf{w}_{t+1} - \mathbf{w}_t\| < +\infty \quad \text{a.s.}$$

Lemma 3 gives some sufficient conditions to guarantee the finite length property. Combining Lemma 4 and Lemma 3, we can draw the gradient can be bounded by the iteration points.

G PROOF OF THEOREM 2

Now, under the KL property defined in Definition 2 (see [28]), we make full use of the decreasing property of the potential energy function Ψ_t^s and the boundedness of $\text{dist}(0, \partial\Psi_t)$ (see Lemma 5) to prove Theorem 2.

Proof: First, we prove a newly defined sequence N_t in (S.73) converges a.s. to zero Q-linearly. By the KL property at $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ we have

$$\nabla\varphi(\Psi_{\tilde{t}+1} - \Psi^*) \text{dist}(0, \partial\Psi_{\tilde{t}+1}) \geq 1 \quad \text{a.s.} \tag{S.61}$$

Using the definition of $\varphi(s) = \tilde{c}s^{1-\tilde{\mu}}, \tilde{\mu} \in [0, 1), \tilde{c} > 0$ in Theorem 2, we can insert $\nabla\varphi(s) = \tilde{c}(1-\tilde{\mu})s^{-\tilde{\mu}}$ into (S.61) to deduce

$$\begin{aligned} \tilde{c}(1-\tilde{\mu})(\Psi_{\tilde{t}+1} - \Psi^*)^{-\tilde{\mu}} \text{dist}(0, \partial\Psi_{\tilde{t}+1}) &\geq 1 \quad \text{a.s.}, \\ (\Psi_{\tilde{t}+1} - \Psi^*)^{\tilde{\mu}} &\leq \tilde{c}(1-\tilde{\mu}) \text{dist}(0, \partial\psi_{\tilde{t}} + 1) \quad \text{a.s.} \end{aligned} \quad (\text{S.62})$$

Using the expression for $\varphi(s) = \tilde{c}s^{1-\tilde{\mu}}$, and the equation (S.55) again to obtain

$$\begin{aligned} &\mathbb{E}\varphi(\Psi_{\tilde{t}+1} - \Psi^*) \\ &\leq \gamma \mathbb{E} \left[\|\mathbf{x}_{\tilde{t}+1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{\tilde{t}-1} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{\tilde{t}+1} - \mathbf{x}_{\tilde{t}}\| + \|\mathbf{x}_{\tilde{t}} - \tilde{\mathbf{x}}\| + \|\mathbf{x}_{\tilde{t}} - \mathbf{x}_{\tilde{t}-1}\| \right]^{\frac{1-\tilde{\mu}}{\tilde{\mu}}} \quad \text{a.s.}, \end{aligned} \quad (\text{S.63})$$

where $\gamma = \tilde{c}[\tilde{c}(1-\tilde{\mu})\xi_{\max}]^{\frac{1-\tilde{\mu}}{\tilde{\mu}}}$.

Next, we focus on the case where $\tilde{\mu} \in (0, \frac{1}{2}]$ and $\frac{1-\tilde{\mu}}{\tilde{\mu}} \geq 1$. When $\tilde{t} \rightarrow +\infty$, the number under the $\frac{1-\tilde{\mu}}{\tilde{\mu}}$ root is infinitely small. And other cases can be proved similarly, which has been studied in [29]. In this case, it follows from Equation (S.63) that

$$\begin{aligned} \mathbb{E}\varphi(\Psi_{\tilde{t}+1} - \Psi^*) &\leq \gamma \left[\mathbb{E}\|\mathbf{x}_{\tilde{t}+1} - \tilde{\mathbf{x}}\| + \mathbb{E}\|\mathbf{x}_{\tilde{t}-1} - \tilde{\mathbf{x}}\| + \mathbb{E}\|\mathbf{x}_{\tilde{t}+1} - \mathbf{x}_{\tilde{t}}\| + \mathbb{E}\|\mathbf{x}_{\tilde{t}} - \tilde{\mathbf{x}}\| \right. \\ &\quad \left. + \mathbb{E}\|\mathbf{x}_{\tilde{t}} - \mathbf{x}_{\tilde{t}-1}\| \right] \quad \text{a.s.} \end{aligned} \quad (\text{S.64})$$

Setting m in the equation (S.59) to be $+\infty$ with $\lim_{m \rightarrow +\infty} \mathbb{E}\|\mathbf{x}_m - \tilde{\mathbf{x}}\| = 0, \lim_{m \rightarrow +\infty} \mathbb{E}\|\mathbf{x}_{m+1} - \mathbf{x}_m\| = 0$ a.s., then we can see that

$$\begin{aligned} &2 \sum_{t=\tilde{t}+1}^{+\infty} \mathbb{E}\|\mathbf{x}_{t+1} - \mathbf{x}_t\| + \sum_{t=\tilde{t}+1}^{+\infty} \mathbb{E}\|\mathbf{x}_{t-1} - \tilde{\mathbf{x}}\| + \mathbb{E}\|\mathbf{x}_{\tilde{t}+1} - \tilde{\mathbf{x}}\| + 2\mathbb{E}\|\mathbf{x}_{\tilde{t}} - \tilde{\mathbf{x}}\| \\ &\leq \mathbb{E}\|\mathbf{x}_{\tilde{t}+1} - \mathbf{x}_{\tilde{t}}\| + (2\sqrt{\xi_{\max}}\sqrt{\frac{2}{\tau}})^2 \mathbb{E}\varphi(\Psi_{\tilde{t}+1} - \Psi^*) < +\infty \quad \text{a.s.} \end{aligned} \quad (\text{S.65})$$

Now set

$$\begin{aligned} \Delta_{\tilde{t}}^1 &:= \sum_{i=\tilde{t}}^{+\infty} \mathbb{E}\|\mathbf{x}_{i+1} - \mathbf{x}_i\|, \\ \Delta_{\tilde{t}}^2 &:= \sum_{i=\tilde{t}}^{+\infty} \mathbb{E}\|\mathbf{x}_i - \tilde{\mathbf{x}}\|, \end{aligned} \quad (\text{S.66})$$

and $L_\varphi = (2\sqrt{\xi_{\max}}\sqrt{\frac{2}{\tau}})^2$, it follows from Equations (S.64) and (S.65) that

$$\begin{aligned} &2\Delta_{\tilde{t}+1}^1 + \Delta_{\tilde{t}}^2 + 2\Delta_{\tilde{t}}^2 - 2\Delta_{\tilde{t}+1}^2 + \Delta_{\tilde{t}+1}^2 - \Delta_{\tilde{t}+2}^2 \\ &\leq \Delta_{\tilde{t}}^1 - \Delta_{\tilde{t}+1}^1 + L_\varphi \gamma [\Delta_{\tilde{t}+1}^2 - \Delta_{\tilde{t}+2}^2 + \Delta_{\tilde{t}}^1 - \Delta_{\tilde{t}+1}^1 + \Delta_{\tilde{t}}^2 - \Delta_{\tilde{t}+1}^2 + \Delta_{\tilde{t}-1}^1 - \Delta_{\tilde{t}}^1 + \Delta_{\tilde{t}-1}^2 - \Delta_{\tilde{t}}^2] \quad \text{a.s.} \end{aligned}$$

To simplify this expression, let's replace the label \tilde{t} with t and set

$$a_t := \Delta_{\tilde{t}}^1, \quad b_t := \Delta_{\tilde{t}}^2, \quad (\text{S.67})$$

the above inequality can be rewritten as the following form

$$0 \leq -(L_\varphi\gamma + 3)a_{t+1} + a_t + L_\varphi\gamma a_{t-1} + L_\varphi\gamma b_{t-1} - 3b_t + b_{t+1} + (1 - L_\varphi\gamma)b_{t+2} \quad \text{a.s.} \quad (\text{S.68})$$

Let us introduce some constants A, B, C, D, E, F , and G satisfied the following conditions, which is the key trick to driving the linear convergence rate

$$\left\{ \begin{array}{l} A = L_\varphi\gamma, \\ B - AC = 1, \\ -BC = -(L_\varphi\gamma + 3), \\ B = 1 + L_\varphi\gamma \times C, \\ D = L_\varphi\gamma, \\ E - GD = -3, \\ F - GE = 1, \\ -GF = 1 - L_\varphi\gamma. \end{array} \right. \quad (\text{S.69})$$

Then (S.68) can be rewritten as

$$0 \leq -BCa_{t+1} + (B - AC)a_t + Aa_{t-1} + Db_{t-1} + (E - GD)b_t + (F - GE)b_{t+1} - GFb_{t+2} \quad \text{a.s.} \quad (\text{S.70})$$

We can rearrange the terms in the above inequality (S.70) to obtain

$$C(Aa_t + Ba_{t+1}) + G(Db_t + Eb_{t+1} + Fb_{t+2}) \leq (Aa_{t-1} + Ba_t) + (Db_{t-1} + Eb_t + Fb_{t+1}) \quad \text{a.s.} \quad (\text{S.71})$$

Let $I = \min\{C, G\} > 1$, from (S.71), we have

$$I[(Aa_t + Ba_{t+1}) + (Db_t + Eb_{t+1} + Fb_{t+2})] \leq Aa_{t-1} + Ba_t + (Db_{t-1} + Eb_t + Fb_{t+1}) \quad \text{a.s.} \quad (\text{S.72})$$

Denote

$$N_t := Aa_{t-1} + Ba_t + (Db_{t-1} + Eb_t + Fb_{t+1}), \quad (\text{S.73})$$

and insert the symbol N_t and N_{t-1} into the inequality (S.72) we get the desired result

$$\frac{N_{t+1}}{N_t} \leq \frac{1}{I} \quad \text{a.s.} \quad (\text{S.74})$$

Then we prove the existence of the parameter. First, we obtain the part of the solution in the equation

(S.69):

$$\begin{cases} L_\varphi\gamma C^2 + C - (L_\varphi\gamma + 3) = 0, \\ B = L_\varphi\gamma C + 1, \\ C = \frac{-1 + \sqrt{1 + 4L_\varphi\gamma(3 + L_\varphi\gamma)}}{2L_\varphi\gamma} > 1, 0 < \frac{1}{C} < 1. \end{cases}$$

Also, we have the rest part of the solution in the equation (S.69):

$$\begin{cases} D = L_\varphi\gamma, \\ E - GD = -3, \\ F - GE = 1, \\ 1 - L_\varphi\gamma = -GF. \end{cases} \quad (\text{S.75})$$

Using the equation (S.75), we have

$$\begin{cases} E = L_\varphi\gamma G - 3, \\ F = GE + 1, \\ L_\varphi\gamma G^3 - 3G^2 + G + (1 - L_\varphi\gamma) = 0. \end{cases}$$

Set $H(G) = L_\varphi\gamma G^3 - 3G^2 + G + (1 - L_\varphi\gamma)$, $H(1) < 0, H(+\infty) > 0$. Thus we can choose the root $G > 1$ of $H(G)$. Since the parameters G and C defined in (S.69) are both greater than 1, we can obtain that I satisfies this condition $I = \min\{C, G\} > 1$. The existence of these parameters defined in (S.69) and I are proved.

Second, the above shows that the sequence $\{N_t\}$ converges a.s. to zero Q-linearly². As a result, we can use the triangle inequality with the notations in (S.66) and (S.67) to yield:

$$\begin{aligned} A\mathbb{E} \|\mathbf{x}^* - \mathbf{x}_t\| &= A\mathbb{E} \left\| \sum_{i=t}^{+\infty} (\mathbf{x}_{i+1} - \mathbf{x}_i) \right\|, \\ &\leq A \sum_{i=t}^{+\infty} \mathbb{E} \|\mathbf{x}_{i+1} - \mathbf{x}_i\|, \\ &\leq N_t \quad \text{a.s.}, \end{aligned} \quad (\text{S.76})$$

and it is sufficient to say the sequence $\{\mathbf{x}_t\}$ converges a.s. to \mathbf{x}^* R-linearly from the definition of R-linear convergence in Section I. Furthermore, from the formulas (S.32) and (16) we can see the sequences $\{\mathbf{y}_t\}$

²For the sequence $\{x_n\}_{n \in \mathbb{N}}$ with $\lim_{n \rightarrow \infty} x_n = x^*$, if

$$\lim_{n \rightarrow \infty} \frac{\|x_{n+1} - x^*\|}{\|x_n - x^*\|} \leq \gamma,$$

where $0 < \gamma < 1$, then the sequence $\{x_n\}_{n \in \mathbb{N}}$ is said to converge to x^* Q-linearly, and the constant γ is called the rate of (linear) convergence.

and $\{\boldsymbol{\lambda}_t\}$ can be controlled by the sequence $\{\mathbf{x}_t\}$ and converge a.s. to \mathbf{y}^* and $\boldsymbol{\lambda}^*$ R-linearly, respectively. Combing the R-linear convergences of the sequences $\{\mathbf{x}_t\}$, $\{\mathbf{y}_t\}$, and $\{\boldsymbol{\lambda}_t\}$ and the definition of R-linear convergence in Section I, we can derive the desired result: the sequence $\{(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t)\}$ converges a.s. to $(\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)$ R-linearly with a existing sequence $\{\hat{c}\xi_3^t\}$ as follows:

$$\mathbb{E} \|(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) - (\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)\| \leq \hat{c}\xi_3^t \quad \text{a.s.}$$

and

$$\|(\mathbf{x}_t, \mathbf{y}_t, \boldsymbol{\lambda}_t) - (\mathbf{x}^*, \mathbf{y}^*, \boldsymbol{\lambda}^*)\| \leq \hat{c}\xi_3^t \quad \text{a.s.}$$

This completes the whole proof. ■