A dichotomy theorem for Γ -switchable *H*-colouring on *m*-edge coloured graphs

Richard Brewster^{*}, Arnott Kidner[†], Gary MacGillivray[‡]

Abstract

Let G be a graph in which each edge is assigned one of the colours $1, 2, \ldots, m$, and let Γ be a subgroup of S_m . The operation of switching at a vertex x of G with respect to an element π of Γ permutes the colours of the edges incident with x according to π . We investigate the complexity of whether there exists a sequence of switches that transforms a given m-edge coloured graph G so that it has a colour-preserving homomorphism to a fixed m-edge coloured graph H and give a dichotomy theorem in the case that Γ acts transitively.

1 Introduction

A homomorphism of a graph G to a graph H is a mapping of the vertices of G to the vertices of H that preserves edges. If the vertices of H are regarded as colours, then a homomorphism f of G to H is an assignment of these colours to the vertices of G so that adjacent vertices are assigned different colours but there are restrictions on the pairs of colours that can appear on the ends of edges. Natural questions about colourings extend to homomorphisms and their generalizations (see [11] and its references). For example, the result that k-COLOURING is polynomial-time solvable for $k \leq 2$ and NP-complete for $k \geq 3$ generalizes to the Hell-Nešetřil Homomorphism Dichotomy Theorem: HOM-H, the problem of deciding whether a given (input) graph G has a homomorphism to a fixed simple (target) graph H is polynomial-time solvable if H is bipartite, and NP-complete if H is not bipartite.

If we move from graphs to relational systems, a set of vertices together with a collection of relations over some fixed signature, then a homomorphism is similarly defined as a vertex mapping which preserves all relations. For a fixed relational system **H**, the problem from the graph setting naturally generalizes to HOM-**H**. The settling of the Constraint Satisfaction Problem (CSP)

^{*}Department of Mathematics and Statistics, Thompson Rivers University, Kamloops, BC, Canada

[†]Department of Mathematics and Statistics, Memorial University of Newfoundland, St, John's, NL, Canada

[‡]Department of Mathematics and Statistics, University of Victoria, Victoria, BC, Canada

dichotomy conjecture [8] by Bulatov [6] and Zhuk [19] gives that HOM-**H** is either polynomial-time solvable or NP-complete for any fixed **H**. The description of the dichotomy is through symmetries and algebra. For many families of targets, there is a combinatorial (structural) description of the dichotomy for the family. Such a dichotomy for the family of all undirected simple graphs is given above. In this paper we give a structural description of the dichotomy for the family of *m*-edge coloured graphs with a switching operation (defined below).

The general setting for our work is as follows. Let G be a graph. We obtain a mixed graph from, G by choosing a subset $A \subseteq E(G)$ and an orientation for each element of A, so the resulting structure has both edges and arcs. We restrict attention to the case where G is a simple graph, thus obtaining what could (and should) be called a *simple* mixed graph but drop the qualifier "simple" because the context is clear throughout the manuscript. Mixed graphs were introduced by Nešetřil and Raspaud [17] in an attempt to unify the homomorphism results for oriented graphs and 2-edge-coloured graphs. For fixed integers m, n > 1, an (m,n)-mixed graph is a mixed graph G where each edge has been assigned a colour from $\{1, 2, \ldots, m\}$ and each arc has been assigned a colour from $\{m +$ $1, m+2, \ldots, m+n$. The operation of *switching* at a vertex of an (m, n)-mixed graph permutes the colours of edges, the colours of arcs, and the direction of arcs incident with the vertex with respect to some element of a group of permutations of the colours and directions. Homomorphisms of (m, n)-mixed graphs with a switching operation are studied in [4, 12], see also [18]. We give a structural description of a homomorphism dichotomy in the case of coloured edges only. The rationale for restricting our attention to coloured edges is discussed below.

In [9], it is shown that for each general Constraint Satisfaction Problem (CSP), there is a directed acyclic graph (DAG) H such that the CSP and HOM-H are polynomially equivalent. Thus a structural description of the dichotomy for HOM-H when restricted to DAGs is likely complicated. In [2] the authors modify this proof to show for each CSP there is a 2-edge coloured graph H with a polynomially equivalent HOM-H problem. Since complete structural description of the homomorphism dichotomy for arc-coloured digraphs would imply such a description for DAGS, it too is likely to be complicated.

By contrast, the introduction of a switching operation seems to lead to simpler dichotomy descriptions. For oriented graphs this has been observed in [14] and for 2-edge coloured graphs, i.e., *signed graphs*, a simple dichotomy description is given in [2, 5].

For a fixed non-negative integer m, an *m*-edge coloured graph is an ordered pair $G = (G_u, \sigma)$ where G_u is a graph and σ is a function that assigns each edge of G_u a colour from $\{1, 2, ..., m\}$. We call G_u the underlying graph of G. As noted above, we restrict attention to the case where G_u is simple. We say G is a cycle if the underlying graph G_u is a cycle, and similarly for paths, trees, etc..

Given *m*-edge coloured graphs *G* and *H*, a homomorphism of *G* to *H* is a function $f: V(G) \to V(H)$ such that if uv is an edge of colour *i* in *G*, then f(u)f(v) is an edge of colour *i* in *H*. We sometimes use the notation $f: G \to H$, or $G \to H$ to denote a homomorphism *f* of *G* to *H*, or when the name of the function is not required,

Two *m*-edge coloured graphs H_1 and H_2 are homomorphically equivalent if there exist homomorphisms $H_1 \to H_2$ and $H_2 \to H_1$. Homomorphic equivalence is an equivalence relation on the set of *m*-edge coloured graphs. If H_1 and H_2 are homomorphically equivalent, then there is a homomorphism of a *m*edge coloured graph G to H_1 if and only if there is a homomorphism of Gto H_2 . After noting that every bipartite graph is homomorphically equivalent to K_1 or K_2 , the Hell-Nešetřil Homomorphism Dichotomy Theorem can be restated in the language of homomorphic equivalence as: If the fixed graph His homomorphically equivalent to a bipartite graph, then HOM-H is solvable in polynomial time. If H is not homomorphically equivalent to a bipartite graph, then HOM-H is NP-complete.

Let $\Gamma \subseteq S_m$ be a permutation group acting on $\{1, 2, \ldots, m\}$. Let G be a m-edge coloured graph. Given a permutation $\pi \in \Gamma$ and a vertex $v \in V(G)$, the m-edge coloured graph $G^{(v,\pi)}$ is obtained by applying π to each edge incident with v as follows. If vu is an edge of colour i in G, then vu is an edge of colour $\pi(i)$ in $G^{(v,\pi)}$. We call this operation switching at v with respect to π . Given a sequence $\Sigma = (v_1, \pi_1), (v_2, \pi_2), \ldots, (v_k, \pi_k)$, switching with respect to Σ is defined as

$$G^{\Sigma} = (G^{(v_1, \pi_1)})^{(v_2, \pi_2), \dots, (v_k, \pi_k)}.$$

Whenever G' can be obtained from G through a sequence of switches, i.e., $G' = G^{\Sigma}$, we say G and G' are Γ -equivalent, denoted $G \cong_{\Gamma} G'$. (Note that the underlying graph of G and G' is the same.)

For ease of terminology, we call a group Γ a *m*-switching group if $\Gamma \subseteq S_m$ and Γ acts transitively on $\{1, 2, \ldots, m\}$.

We say G admits a Γ -homomorphism to H, and write $G \to_{\Gamma} H$, if $G' \to H$ for some G' which is Γ -equivalent to G. If $G \to_{\Gamma} H$ and H' is Γ -equivalent to H, then it is easy to verify that $G \to_{\Gamma} H'$ (apply switches on H to pre-images in G). A similar argument is used to prove the following proposition, which has previously appeared in several contexts.

Proposition 1.1 ([2,3,13,15]). Let H_1, H_2 and H_3 be m-edge coloured graphs, and let Γ be a m-switching group. If f is a Γ -homomorphism of H_1 to H_2 and g is a Γ -homomorphism of H_2 to H_3 , then $g \circ f$ is a Γ -homomorphism of H_1 to H_3 .

The concept of Γ -homomorphic equivalence, defined analogously as above, will allow the main results of this paper to be presented in a way that shows them to be natural generalizations of the Hell-Nešetřil Homomorphism Dichotomy Theorem.

It is not immediately clear whether testing the existence of a Γ -homomorphism to a fixed target is a CSP as each CSP is equivalent to testing the existence of a homomorphism to a fixed target H. That is, a Γ -homomorphism is a two step process: switch G to some G', then map $G' \to H$. The following construction has appeared in many contexts [2,3,12,13,15]. Given a *m*-edge coloured graph H and a fixed *Abelian* permutation group Γ acting (perhaps not transitively) on $\{1, 2, \ldots, m\}$, the Γ -switch graph, $sw_{\Gamma}(H)$, has vertices $V(H) \times \Gamma$ and edges defined as follows. If $v_j v_k$ is an edge of colour i in H, then $(v_j, \pi_r)(v_k, \pi_s)$ is an edge of colour $\pi_r \pi_s(i)$ in $sw_{\Gamma}(H)$. As a convention we shall use e_{Γ} to denote the identity of Γ . In particular, the vertices of the form (v, e_{Γ}) induce a copy of H in $sw_{\Gamma}(H)$. The following proposition shows that the existence of a Γ -homomorphism $G \to_{\Gamma} H$ is equivalent to the existence of a homomorphism $G \to sw_{\Gamma}(H)$ (without switching).

Proposition 1.2 ([2,3,12,13,15]). Let G and H be m-coloured graphs and $\Gamma \subseteq S_m$ be an Abelian group acting on $\{1, 2, ..., m\}$. The following are equivalent.

(a) $G \to_{\Gamma} H$, (b) $G \to sw_{\Gamma}(H)$, (c) $sw_{\Gamma}(G) \to sw_{\Gamma}(H)$.

The focus of this paper is the following decision problem. Let H be a fixed *m*-edge coloured graph and Γ be a fixed permutation group acting on $\{1, 2, \ldots, m\}$.

 Γ -HOM-HInstance: a *m*-edge coloured graph G. Question: Is there a Γ -homomorphism of G to H?

For Abelian groups, Proposition 1.2 gives an equivalence between the above problem and the HOM- $sw_{\Gamma}(H)$ problem. Since $sw_{\Gamma}(H)$ is a fixed target, by the CSP dichotomy, each Γ -HOM-H problem is polynomial-time solvable or NPcomplete. In Section 2 we provide a structural description of the dichotomy for Γ -HOM-H in the case that Γ is Abelian.

For non-Abelian groups it is neither clear that Γ -HOM-H is in NP nor that it is a CSP. Our second contribution, in Section 3, is to show for each m-switching group Γ there is an integer m' and an Abelian m'-switching group Γ' and an edge-coloured graph H' such that Γ -HOM-H and Γ' -HOM-H' are polynomially equivalent. Combining these two results gives us the main result of the paper which is a structural description of the dichotomy for Γ -HOM-H for all fixed simple m-edge coloured graphs H and all m-switching groups Γ .

We conclude with a final remark on restricting to m-edge coloured graphs. As noted above, the description of the dichotomy for HOM-H is likely complicated when restricted to DAGs or 2-edge coloured graphs (without switching). If we allow both edges and arcs, together with a switching operation that does not interchange edges and arcs, the same proof as in [9], can be used to show that Γ -HOM-H captures the full complexity of all CSPs. Thus we shall restrict our attention to m-edge coloured graphs.

2 A dichotomy for Abelian Γ

In this section we establish a dichotomy theorem for Γ -HOM-H when the m-switching group Γ is Abelian. As cycles play an important roll in this dichotomy we first determine important facts about switching cycles with Abelian groups.

We say a graph or subgraph (usually a cycle or spanning tree) is monochromatic of colour i if all of its edges are of colour i. In the case that a cycle has all edges but one of colour i and one edge of colour $j \neq i$, we say that it is nearly monochromatic of colours (i, j). When no confusion arises, we may omit the colours and say the graph or subgraph is monochromatic or nearlymonochromatic. These concepts generalize the situation for signed graphs to m-edge coloured graphs (see [16] and the references within).

Proposition 2.1. Suppose Γ is an Abelian *m*-switching group and let *C* be a *m*-edge coloured cycle. Then exactly one of the following statements holds:

- (i) C has even length and for any colour i, C can be switched using Γ to be monochromatic of colour i;
- (ii) C has even length and for any colour i, C can be switched using Γ to be nearly monochromatic of colours (i, j) where j depends on i; or
- (iii) C has odd length and for some colour j, C can be switched using Γ to be monochromatic of colour j.

Proof. Let the cycle C have vertices $v_0, v_1, \ldots, v_{n-1}$ in the natural order. Since Γ acts transitively, for any colour i, we can switch at $v_1, v_2, \ldots, v_{n-1}$ so that all edges with the possible exception of v_0v_{n-1} are of colour i. Hence every cycle can be switched to be monochromatic or nearly monochromatic using Γ . We therefore assume C is monochromatic of colour i or nearly monochromatic of colours (i, j), where j depends on i.

Suppose that C is of even length. We now show that statements (i) and (ii) are mutually exclusive. Let C be a nearly monochromatic even cycle. Suppose to the contrary that it is possible to switch C to be monochromatic. By statement (i) we may assume C can be switched to be monochromatic of colour i. Since Γ is Abelian, we can assume that at most one switch is applied at each vertex. Let (π_s, v_s) be the switch applied to v_s for $0 \le s \le n-1$. For each edge $v_s v_{s+1}$, $0 \le s \le n-2$, we have $\pi_{s+1}\pi_s(i) = i$, or $\pi_{s+1} = \pi_s^{-1}$. This implies π_0 is applied at all vertices with an even subscript and π_0^{-1} at all vertices with an odd subscript. However this implies the edge $v_0 v_{n-1}$ is switched from colour j to colour i by $\pi_0 \pi_0^{-1}$, a contradiction.

Since *i* is arbitrary, it follows that (i) and (ii) hold. Note that if *C* is monochromatic and Γ acts transitively, *C* can be transformed to be monochromatic of any colour *j* by switching at every other vertex around cycle.

Finally, suppose C is an odd cycle. If C is monochromatic, then we are done. Otherwise, v_0v_{n-1} is colour $j \neq i$. Let $\pi \in \Gamma$ satisfy $\pi(i) = j$. Switching at vertices $v_1, v_3, \ldots, v_{n-2}$ with π switches C to be monochromatic of colour j.

Since statements (i), (ii) and (iii) are mutually exclusive, the result follows.

We denote by K_2^i the monochromatic *m*-edge coloured graph of colour *i* whose underlying graph is K_2 . In [4,12] a polynomial algorithm for Γ -HOM- K_2^i is given. In the Abelian case it is described as follows. If the given graph is not bipartite, then we have a no-instance. Otherwise, let *G* be a connected,

bipartite *m*-edge coloured graph. Fix a spanning tree *T* of *G* and switch so that *T* is monochromatic of colour *i* (this is always possible). Next we look at the cotree edges. If each of them is of colour *i*, then *G* is a yes-instance; otherwise some co-tree edge is of colour $j \neq i$ and so *G* contains a cycle that is Γ -equivalent to a nearly monochromatic cycle of colour (i, j). By Proposition 2.1(ii), the cycle cannot be switched to be monochromatic of colour *i* and hence neither can *G*. Thus *G* is a no-instance.

Alternatively, we can leverage the switch graph using Proposition 1.2. Since Γ is Abelian and acts transitively it is regular. Thus in $sw_{\Gamma}(K_2^i)$ each vertex is incident with exactly one edge of each colour. Thus one can test $G \rightarrow_{\Gamma} K_2^i$ by first mapping one vertex v of G to some vertex (w, e_{Γ}) in $sw_{\Gamma}(K_2^i)$. Each neighbour of v in G must then map to the unique neighbour of (w, e_{Γ}) joined by an edge of the same colour. Continuing, we either construct a homomorphism or obtain an contradiction. This approach avoids the need to switch. If the answer is "no" then one needs to trace backwards to find a certificate that no homomorphism exists.

We conclude this section with a dichotomy theorem for Γ -HOM-H when the m-switching group Γ is Abelian. The proof uses the indicator construction [11] and a theorem of Barto, Kozig and Niven [1]. These are described next.

Let A be a m-edge coloured graph with specified vertices a and b. Given a m-edge-coloured graph H, the *indicator construction with respect to* (A, a, b)produces a digraph H^* with vertex set $V(H^*) = V(H)$ and an arc from x to y whenever there is a homomorphism of A to H that maps a to x and b to y.

Lemma 2.2 ([11]). HOM- H^* polynomially transforms to HOM-H.

A digraph D is called *smooth* if it has no vertex of in-degree 0 and no vertex of out-degree 0. A dichotomy theorem for homomorphisms to smooth digraphs is known:

Theorem 2.3 ([1]). Let H be a smooth digraph, and let C be a shortest directed cycle in H. If H is homomorphically equivalent to C, then HOM-H is solvable in polynomial time; otherwise HOM-H is NP-complete.

Corollary 2.4. Let H be a smooth digraph with no loops. If some component of H contains directed cycles whose lengths are relatively prime, then HOM-H is NP-complete.

The polynomial cases of the dichotomy theorem below rely on the following theorem. It was the first non-trivial result which applies to all groups that act transitively on the edge colours.

Theorem 2.5 ([4,12]). Let Γ be a *m*-switching group and $i \in \{1, 2, ..., m\}$. Then HOM- K_2^i is polynomial-time solvable.

Theorem 2.6. Let Γ be an Abelian m-switching group. For a fixed m-edge coloured graph H, if H is Γ -homomorphically equivalent to a monochromatic bipartite m-edge coloured graph, then Γ -HOM-H is polynomial-time solvable. If H is not Γ -homomorphically equivalent to a monochromatic m-edge coloured graph, then Γ -HOM-H is NP-complete.

Proof. a *m*-edge coloured graph *G* has a Γ-homomorphism to a monochromatic bipartite graph if and only if it has a homomorphism to a monochromatic K_1 or a monochromatic K_2 . The case for K_1 is easy to test. There is a Γ-homomorphism to a monochromatic K_2 if and only if *G* is bipartite and can be switched to be monochromatic. In this case Γ-HOM-*H* is polynomial-time solvable by Theorem 2.5.

Now suppose H is Γ -homomorphically equivalent to neither a monochromatic K_1 nor a monochromatic K_2 . Thus either H is bipartite and cannot be switched to be monochromatic, or H is not bipartite.

Suppose H is bipartite and cannot be switched to be monochromatic. Choose a spanning tree T of H and apply a sequence of switches Σ so that in H^{Σ} every edge of T has colour i. Since G cannot be switched to be monochromatic, there must be some co-tree edge of colour $j \neq i$. Let C be a nearly monochromatic cycle in H^{Σ} on vertices $v_0, v_1, \ldots, v_{2k-1}$ with all edges in T except the co-tree edge v_0v_{2k-1} , which has colour $j \neq i$.

For ease of notation denote $sw_{\Gamma}(H)$ by SH. Let A be the 2-edge coloured graph consisting of a path of length 2 with end points a and b with the edge incident with a of colour i and the edge incident with b of colour j. Let SH^* be the result of applying the indicator construction with respect to (A, a, b) to SH. By Lemma 2.2, we need to show the HOM- SH^* problem is NP-complete. This is accomplished by proving SH^* is smooth, and some component has directed cycles of relatively prime lengths.

Claim 2.7. The digraph SH^* is smooth.

Consider a non-trivial component of H. Then each vertex (u, π) in SH is incident with an edge $(u, \pi)(w, \pi)$ of some colour, say c. Since Γ acts transitively, there is $\tau \in \Gamma$ such that $\tau(c) = i$, or equivalently the edge $(u, \pi)(w, \tau\pi)$ is of colour i. Similarly the vertex $(w, \tau\pi)$ is incident with an edge of colour j, say $(w, \tau\pi)(v, \sigma)$. Hence there exists a homomorphism from A to SH where a is mapped to (u, π) and b is mapped to (v, σ) . We conclude that in SH^* , the vertex (u, π) is the tail of an arc. Using similar reasoning, the vertex (u, π) is also the head of an arc in SH^* . This proves the claim.

Claim 2.8. The digraph SH* has directed cycles of relatively prime lengths.

By hypothesis, SH contains the nearly monochromatic cycle

$$(v_0, e_{\Gamma}), (v_1, e_{\Gamma}), \dots, (v_{2n-1}, e_{\Gamma}), (v_0, e_{\Gamma})$$

with all edges of colour *i* except the edge $(v_0, e_{\Gamma})(v_{2n-1}, e_{\Gamma})$ which is of colour *j*.

Since Γ acts transitively, there exists $\pi \in \Gamma$ such that $\pi(i) = j$. Let the order of π be d, that is, $\pi^d = e_{\Gamma}$. Observe that the edge $(v_0, \pi^s)(v_1, \pi^t)$ will be of colour i when $s+t \equiv 0 \pmod{d}$ and will be of colour j when $s+t \equiv 1 \pmod{d}$. In $sw_{\Gamma}(H)$ the cycle

$$C_1 = (v_0, e_{\Gamma}), (v_1, e_{\Gamma}), (v_0, \pi), (v_1, \pi^{d-1}), (v_0, \pi^2), (v_1, \pi^{d-2}), \\ \dots, (v_0, \pi^{\ell}), (v_1, \pi^{d-\ell}), \dots, (v_0, \pi^{d-1}), (v_1, \pi), (v_0, e_{\Gamma})$$

is a cycle of length 2d whose edges alternate colour i and colour j. An example is shown in Figure 1.



Figure 1: A blue-red alternating cycle of length 12 in $sw_{\Gamma}(H)$ where π has order 6.

Let P_t be the path given by

$$P_t = (v_t, e_{\Gamma}), (v_{t+1}, \pi), (v_t, \pi^{d-1}), \dots, (v_{t+1}, \pi^{\ell}), (v_t, \pi^{d-\ell}), \\ \dots, (v_{t+1}, \pi^{d-1}), (v_t, \pi), (v_{t+1}, e_{\Gamma}), (v_{t+2}, e_{\Gamma})$$

Then the following cycle is of length 2d(k-1) + 2.

$$C_2 = (v_0, e_{\Gamma}), P_1, P_3, \dots, P_{2k-3}, (v_{2k-1}, e_{\Gamma}), (v_0, e_{\Gamma})$$

It is clear by construction that C_2 is a cycle of length 2d(k-1) + 2 whose edges alternate colours *i* and *j*. An example is shown in Figure 2.



Figure 2: A red-blue alternating cycle of length 12(k-1) + 2 in $sw_{\Gamma}(H)$ where π has order 6.

The cycles C_1 and C_2 show that SH^* has a component containing directed cycles of relatively prime lengths d and d(k-1) + 1 (on the vertices of C_1 and C_2 at even distance along the cycle from (v_0, e_{Γ})), which completes the proof of the claim.

As SH does not contain any cycles of length 1 or 2, we have SH^* is loopfree. Thus HOM- SH^* is NP-complete by Corollary 2.4, and hence HOM-SH is NP-complete.

Finally, suppose H is not bipartite. Thus it contains an odd cycle C. By Proposition 2.1(iii), H is Γ -switch equivalent to a graph with a monochromatic odd cycle. Thus, $sw_{\Gamma}(H)$ contains a monochromatic odd cycle, say of colour i. By [10], the HOM- $sw_{\Gamma}(H)$ problem is NP-complete by restricting the inputs to colour i. By Proposition 1.2, Γ -HOM-H is NP-complete.

This completes the proof of the theorem.

3 Group Abelianization

We now turn our attention to non-Abelian m-switching groups. Some special cases have been considered in earler work [7].

Let H be a *m*-edge coloured graph and let Γ be a *m*-switching group. In this section we show there is a set of m' colours, a m'-edge coloured graph H'with the same underlying graph as H, and an Abelian m'-switching group Γ' such that Γ' -HOM-H' and Γ -HOM-H are polynomially equivalent.

We recall some group theory terminology. For $\pi, \phi \in \Gamma$, the element $[\pi, \phi] = \pi \phi \pi^{-1} \phi^{-1}$ is a *commutator* of Γ . The *commutator subgroup of* Γ , denoted $[\Gamma, \Gamma]$, is the subgroup generated by the commutators of Γ . We now show how commutators can be used to generate a sequence of switches that allow the colour of a single edge to be changed.

Lemma 3.1. Let H be a m-edge coloured graph and Γ be a m-switching group. For any $\tau \in [\Gamma, \Gamma]$ and any edge uw in H of colour i, there is a sequence Σ of switches such that in H^{Σ} the edge uw has colour $\tau(i)$ and all other edges in Hand H^{Σ} have the same colour.

Proof. Since $\tau \in [\Gamma, \Gamma]$, we have $\tau = [\pi_1, \phi_1] \cdots [\pi_k, \phi_k]$. Let

$$H^{\Sigma} = H^{(w,\phi_k^{-1})(u,\pi_k^{-1})(w,\phi_k)(u,\pi_k)\cdots(w,\phi_1^{-1})(u,\pi_1^{-1})(w,\phi_1)(u,\pi_1)}.$$

The switches at u are the product $\pi_k^{-1}\pi_k\cdots\pi_1^{-1}\pi_1$ and at w are $\phi_k^{-1}\phi_k\cdots\phi_1^{-1}\phi_1$. Thus each edge other that uw has the same colour in H and H^{Σ} . The edge uw, on other the hand, has colour $\tau(i)$ as required.

It is well known that $[\Gamma, \Gamma]$ is a smallest normal subgroup of Γ such that the quotient group $\Gamma/[\Gamma, \Gamma]$ is Abelian. The orbits of $[\Gamma, \Gamma]$ partition $\{1, 2, \ldots, m\}$ as $D = \{\Delta(1), \ldots, \Delta(m)\}$, where $j \in \Delta(i)$ if there is $\tau \in [\Gamma, \Gamma]$ such that $\tau(i) = j$. Note that $\Delta(i) = \Delta(j)$ if and only if $j \in \Delta(i)$. The partition D is a block system for Γ . Let m' be the number of distinct blocks in D. In the sequel we will regard these as edge colours.

Given a *m*-edge coloured graph G, we define the Abelianization of G with respect to Γ , denoted G^{Γ} , to be the *m'*-edge coloured graph obtained from G by replacing each edge of colour i with an edge of colour $\Delta(i)$.

The quotient group $\Gamma/[\Gamma, \Gamma]$ acts transitively on the set of m' colours. It may not be a m'-switching group because it may have several permutations that act identically on the block system. To see this, consider $\Gamma = S_m$. Then $[\Gamma, \Gamma] = A_m$ and $\Gamma/[\Gamma, \Gamma] = S_2$. But in this case m' = 1 (since A_m acts transitively), hence the quotient group is not a m'-switching group as it isn't a subgroup of $S_{m'}$. However, both elements of the quotient group act identically on the block system.

Let Γ^{ab} be the subgroup of $S_{m'}$ which is isomorphic to the subgroup of $\Gamma/[\Gamma, \Gamma]$ arising from defining two elements of $\Gamma/[\Gamma, \Gamma]$ to be equivalent if they act identically on the block system D. Then Γ^{ab} is a m'-switching group. Since $\Gamma/[\Gamma, \Gamma]$ is Abelian, so is Γ^{ab} . We call Γ^{ab} the Abelianization of Γ .

We now define switching on G^{Γ} with respect to Γ^{ab} . Recall that each $\pi' \in \Gamma^{ab}$ corresponds to an equivalence class of cosets of the form $\pi[\Gamma, \Gamma]$. To switch G^{Γ} at u with respect to π' we apply $\pi \in \Gamma$ to each of the edges incident with u as follows: an edge uv of colour $\Delta(i)$ in G^{Γ} switches to colour $\Delta(\pi(i))$. We claim that such switching is well defined in that for any element in the coset $\pi[\Gamma, \Gamma]$ and any colour $j \in \Delta(i)$, the resulting colour of uv in G^{Γ} is the same. To see this, given $j \in \Delta(i)$ there is $\tau \in [\Gamma, \Gamma]$ such that $\tau(i) = j$. Then $\pi(j) = \pi(\tau(i))$. Since $[\Gamma, \Gamma]$ is a normal subgroup, $\pi\tau = \tau'\pi$ for some $\tau' \in [\Gamma, \Gamma]$. Thus $\pi(j) = \tau'(\pi(i))$ giving $\pi(j) \in \Delta(\pi(i))$ as required. Similarly, we can argue $\phi(i) \in \Delta(\pi(i))$ for any $\phi \in \pi[\Gamma, \Gamma]$ (or any other coset of $[\Gamma, \Gamma]$ with the same action on D). This proves the claim. As a result, if $\pi' = \pi[\Gamma, \Gamma]$, we can write $\pi'(\Delta(i)) = \Delta(\pi(i))$.

We note two special cases of interest. If Γ is Abelian, then $[\Gamma, \Gamma] = \{e_{\Gamma}\}$ and each block is a singleton, i.e., $\Delta(i) = \{i\}$. Hence, $\Gamma = \Gamma^{ab}$ and the action of the two groups is the same. Conversely, if $[\Gamma, \Gamma]$ acts transitively on $\{1, 2, \ldots, m\}$, then the block system consists of a single block, i.e. $\Delta(i) = \{1, 2, \ldots, m\}$ for all *i*. In this case for a given *G*, the *m*-edge coloured graph G^{Γ} has a single edge colour (and is invariant under switching by Γ^{ab}). By Lemma 3.1 it is possible to switch each edge of *G* individually to any colour of $\{1, 2, \ldots, m\}$ implying all edge colourings of *G* are Γ -equivalent. See [7] for examples.

Theorem 3.2. Let G and H be m-edge coloured graphs and Γ be a m-switching group. Then $G \cong_{\Gamma} H$ if and only if $G^{\Gamma} \cong_{\Gamma^{ab}} H^{\Gamma}$.

Proof. Suppose $H = G^{\Sigma}$. By induction it suffices to consider the case $\Sigma = \{(u,\pi)\}$. Consider the edges incident with u. Suppose one such edge uv has colour i in G and colour $j = \pi(i)$ in H. In G^{Γ} the edge uv has colour $\Delta(i)$ and it has the colour $\Delta(j)$ in H^{Γ} . Letting $\pi' = \pi[\Gamma, \Gamma]$, we have $\pi'(\Delta(i)) = \Delta(\pi(i)) = \Delta(j)$. This holds for all edges incident with u and thus $(G^{\Gamma})^{(u,\pi')} = H^{\Gamma}$.

Now suppose $G^{\Gamma} \cong_{\Gamma^{ab}} H^{\Gamma}$. As above it suffices to consider the case $H^{\Gamma} = (G^{\Gamma})^{(u,\pi')}$ for some $\pi' \in \Gamma^{ab}$. Again we consider the edges incident with u and focus on an arbitrary edge uv. Suppose in G^{Γ} the edge uv is colour $\Delta(s)$ and colour $\Delta(t)$ in H^{Γ} where $\pi'(\Delta(s)) = \Delta(t)$. By construction, in G the edge uv is some colour $i \in \Delta(s)$ and in H some colour $j \in \Delta(t)$.

Suppose π' is the coset $\pi[\Gamma, \Gamma]$. Since the switching operation is well defined, we have $\pi(i) = j'$ for some $j' \in \Delta(t)$. Thus, in $G^{(u,\pi)}$, the edge uv has colour j'. Since $j, j' \in \Delta(t)$, there is $\tau \in [\Gamma, \Gamma]$ such that $\tau(j') = j$. Using Lemma 3.1, there is a sequence Σ that changes uv from colour j' to j and leaves all other edges unchanged. That is, in $G^{(u,\pi)\Sigma}$ the edge uv has colour j. After doing this for each edge incident at u, we see $G \cong_{\Gamma} H$.

Corollary 3.3. Let G and H be m-edge coloured graphs and Γ be a m-switching group. Then $G \to_{\Gamma} H$ if and only if $G^{\Gamma} \to_{\Gamma^{ab}} H^{\Gamma}$.

Proof. Let G and H be *m*-edge coloured graphs with Abelianizations G^{Γ} and H^{Γ} respectively. Suppose $f: G \to_{\Gamma} H$. By definition, $f: G^{\Sigma} \to H$ for some sequence of switches Σ . Consider an edge uv of colour i in G, and colour i' in G^{Σ} . Then by Theorem 3.2 there is a sequence Σ' of switches on G^{Γ} so that edge uv switches from $\Delta(i)$ to $\Delta(i')$. In H, the edge f(u)f(v) has colour i' and thus in H^{Γ} it has colour $\Delta(i')$. Hence, the mapping $f: (G^{\Gamma})^{\Sigma'} \to H^{\Gamma}$ is a homomorphism as required.

Similarly, suppose $f: G^{\Gamma} \to_{\Gamma^{ab}} H^{\Gamma}$, so that $f: (G^{\Gamma})^{\Sigma} \to H^{\Gamma}$. Let uv have colour $\Delta(s)$ in G^{Γ} and colour $\Delta(t)$ in $(G^{\Gamma})^{\Sigma}$. In G the edge uv has colour $i \in \Delta(s)$. After applying the corresponding switches from Σ (by selecting a representative from each coset) to G to obtain $G^{\Sigma'}$, the edge uw has colour jfor some $j \in \Delta(t)$. In H^{Γ} , the edge f(u)f(v) must have colour $\Delta(t)$ and hence f(u)f(v) in H has some colour $j' \in \Delta(t)$. Thus, there is $\tau \in [\Gamma, \Gamma]$ such that $\tau(j) = j'$. Hence, by Lemma 3.1, we can switch $G^{\Sigma'}$ so that uv has colour j'. Processing each edge in this way shows there is a sequence of switches Σ'' such that $G \cong_{\Gamma} G^{\Sigma''}$ and $f: G^{\Sigma''} \to H$ is a homomorphism as required. \Box

It is not immediately clear that the number of switches required to switch a *m*-edge coloured graph G to a Γ -equivalent *m*-edge coloured graph G^{Σ} is polynomial in the size of G. If Γ is Abelian, then there is a sequence that switches each vertex at most once. As Γ^{ab} is Abelian, we have the following corollary.

Corollary 3.4. For any fixed m-switching group Γ , Γ -Hom-H is in NP.

Proof. As Γ is fixed, the groups Γ^{ab} and $[\Gamma, \Gamma]$ are of fixed sized. We can certify G is a yes-instance of Γ -HOM-H by providing the switching sequence Σ and the function $f: V(G) \to V(H)$. By Corollary 3.3, we know the switches of G consist of switches from Γ^{ab} and switches from $[\Gamma, \Gamma]$. As Γ^{ab} is Abelian, each vertex needs to be switched at most once by an element of Γ^{ab} , and as $[\Gamma, \Gamma]$ is fixed, the sequence used to change the colour of a single edge (as in Lemma 3.1) is of bounded sized. Thus, the switching sequence Σ has length O(|E(G)|). \Box

We now state the full dichotomy theorem for Γ -HOM-H. A fundamental step in the polynomial-time case is testing if a given G can be switched to be monochromatic of some colour i.

As Γ^{ab} is Abelian, this can be done as described previously.

Theorem 3.5. Let Γ be a *m*-switching group. For a fixed *m*-edge coloured graph *H*, if *H* is Γ -homomorphically equivalent to (a monochromatic) bipartite graph, then Γ -HOM-*H* is polynomial-time solvable. If *H* is not Γ -homomorphically equivalent to a monochromatic bipartite graph, then Γ -HOM-*H* is NP-complete.

Proof. Note that H is Γ-homomoprically equivalent to a monochromatic bipartite graph if and only if it is Γ-homomophically equivalent to a monochromatic K_1 or K_2 . The case for K_1 is trivial. Consider the case where H is Γ-homomophically equivalent to a monochromatic K_2 . Without loss of generality, $H = K_2^i$. Then $H^{\Gamma} = K_2^{\Delta(i)}$. By Corollary 3.3, $G \to_{\Gamma} H$ if and only if $G^{\Gamma} \to_{\Gamma^{ab}} H^{\Gamma}$, which can be tested in polynomial time by Theorem 2.5.

Conversely if H is not Γ -homomorphically equivalent to a monochromatic bipartite graph, then H^{Γ} contains an odd cycle, or is bipartite but with a cycle that switches to be nearly monochromatic of some colours $(\Delta(i), \Delta(j))$. Thus, Γ^{ab} -HOM- H^{Γ} is NP-complete. Thus Γ -HOM-H is NP-complete by Corollary 3.3.

The following corollary gives the structural characterization of the dichotomy.

Corollary 3.6. Let Γ be a m-switching group. For a fixed m-edge coloured graph H, if

- (i) H is bipartite; and
- (ii) can be switched to be monochromatic,

then Γ -Hom-H is polynomial-time solvable. Otherwise, if

- (iii) H contains an odd cycle, or
- (iv) H^{Γ} contains an (even) length cycle that can be switched (by Γ^{ab}) to be nearly monochromatic of colours $(\Delta(k), \Delta(l))$,

then Γ -HOM-H is NP-complete.

The importance of nearly monochromatic cycles was identified in [12] where the agreeance class of i (later renamed the substitution class of i in [4]) is defined to be the set of all colours j for which a nearly monochromatic cycle of colours (i, j) can be switched to be monochromatic of colour i. For even cycles these classes turn out to be exactly the block system $D = \{\Delta(1), \Delta(2), \ldots, \Delta(m)\}$ as defined by the action of the commutator subgroup.

References

- Libor Barto, Marcin Kozik, and Todd Niven. The CSP dichotomy holds for digraphs with no sources and no sinks (a positive answer to a conjecture of Bang-Jensen and Hell). SIAM J. Comput., 38(5):1782–1802, 2008/09.
- [2] Richard C. Brewster, Florent Foucaud, Pavol Hell, and Reza Naserasr. The complexity of signed graph and edge-coloured graph homomorphisms. *Discrete Mathematics*, 340(2):223–235, Feb 2017.
- [3] Richard C. Brewster and Timothy Graves. Edge-switching homomorphisms of edge-coloured graphs. *Discrete Mathematics*, 309(18):5540–5546, 2009.

- [4] Richard C. Brewster, Arnott Kidner, and Gary MacGillivray. The 2colouring problem for (m, n)-mixed graphs with switching is polynomial, November 2022.
- [5] Richard C. Brewster and Mark Siggers. A complexity dichotomy for signed H-colouring. *Discrete Math.*, 341(10):2768–2773, 2018.
- [6] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), pages 319–330, 2017.
- [7] Christopher Duffy, Gary MacGillivray, and Ben Tremblay. Switching medge-coloured graphs with non-Abelian groups, Manuscript, 2023.
- [8] Tomás Feder and Moshe Y. Vardi. Monotone monadic SNP and constraint satisfaction. In *Proceedings of the Twenty-Fifth Annual ACM Symposium* on Theory of Computing, STOC '93, page 612–622, New York, NY, USA, 1993. Association for Computing Machinery.
- [9] Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic snp and constraint satisfaction: A study through datalog and group theory. SIAM Journal on Computing, 28(1):57–104, 1998.
- [10] Pavol Hell and Jaroslav Nešetřil. On the complexity of H-coloring. Journal of Combinatorial Theory, Series B, 48(1):92–110, 1990.
- [11] Pavol Hell and Jaroslav Nešetřil. Graphs and Homomorphisms. Oxford Univ. Press, 2008.
- [12] Arnott Kidner. Gamma-switchable 2-colourings of (m, n)-mixed graphs. Master's thesis, University of Victoria, Victoria, Canada, 2021. https://dspace.library.uvic.ca/handle/1828/13337.
- [13] William F. Klostermeyer. Pushing vertices and orienting edges. Ars Combinatorial, 51:65–75, 1999.
- [14] William F. Klostermeyer and Gary MacGillivray. Homomorphisms and oriented colorings of equivalence classes of oriented graphs. *Discrete Mathematics*, 274(1):161–172, 2004.
- [15] Etienne Leclerc, Gary MacGillivray, and J. Maria Warren. Switching (m,n)mixed graphs with respect to Abelian groups, Manuscript, 2021.
- [16] Reza Naserasr, Éric Sopena, and Thomas Zaslavsky. Homomorphisms of signed graphs: an update. *European J. Combin.*, 91:Paper No. 103222, 20, 2021.
- [17] Jaroslav Nešetřil and André Raspaud. Colored homomorphisms of colored mixed graphs. Journal of Combinatorial Theory, Series B, 80(1):147–155, 2000.

- [18] Sagnik Sen, Éric Sopena, and S. Taruni. Homomorphisms of (n,m)-graphs with respect to generalised switch, 2022.
- [19] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. J. ACM, 67(5), August 2020.