DYNAMICAL PROPERTIES FOR COMPOSITION OPERATORS ON $H^2(\mathbb{C}_+)$

CARLOS F. ÁLVAREZ AND JAVIER HENRÍQUEZ-AMADOR

ABSTRACT. Expansivity, Li-Yorke chaos and shadowing are popular and wellstudied notions of dynamical systems. Several simple and useful characterizations of these notions within the setting of linear dynamics were obtained recently. We explore these three dynamical properties for composition operators $C_{\phi}f = f \circ \phi$ induced by affine self-maps ϕ of the right half-plane \mathbb{C}_+ on the Hardy-Hilbert space $H^2(\mathbb{C}_+)$.

CONTENTS

1. Introduction	1
2. Preliminaries	2
2.1. Notions of linear dynamics	3
2.2. Hardy space $H^2(\mathbb{C}_+)$	5
3. Expansivity of C_{ϕ}	6
3.1. Expansivity in a positive sense of C_{ϕ}	6
3.2. Forward and backward expansivity for C_{ϕ}	8
4. Shadowing for C_{ϕ}	9
5. Li-Yorke chaos for C_{ϕ}	11
Acknowledgments	11
References	11

1. INTRODUCTION

Linear Dynamics field focuses on the study of dynamical systems that involve continuous linear operators on infinite-dimensional Banach (or Fréchet) spaces. Within this area, important concepts such as *chaos, expansivity* and *shadowing* have emerged. Exploring these dynamical properties has become a common interest among researchers in the fields of functional analysis and dynamical systems. For a more detailed understanding, readers are encouraged to refer to the books [1, 11] and the papers [3, 4], which provide additional references. To delve into the dynamics of linear operators on Hilbert spaces, readers can consult [10, 16, 17].

In the early 1970's Eisenberg and Hedlund studied relationships between expansivity and spectrum of operators on Banach spaces [6]. Starting the 2000's, Mazur investigated expansivity and shadowing for the class of normal operators on Hilbert spaces [16]. Recently, the notions of expansivity and shadowing has been explored for continuous linear operators on Banach spaces [4]. The *Li-Yorke chaos* notion was introduced by Li and Yorke in [12] for interval maps. This version of chaos

Date: October 23, 2023.

²⁰²⁰ Mathematics Subject Classification. Primary 47A16, 47B33; Secondary 37D45. Key words and phrases. Expansivity; Shadowing; Li-Yorke chaos; Composition operators.

CF. A. was supported by ANID Proyecto FONDECYT Postdoctorado 3230048, Chile.

has also been studied within context of linear dynamics, for instance, as exemplified [2, 3] and related references. It is worth mentioning that Godefroy and Shapiro introduced chaos for linear operators on Fréchet spaces [9], adopting Devaney's definition of chaos.

Composition operators are a class of linear operators very studied in complex and functional analysis. There have been several works exploring composition operators within the framework of linear dynamics. For instance, Li-Yorke chaos and expansivity has been characterized for composition operators $T_{\phi} : f \mapsto f \circ \phi$ acting on L^p -spaces with $1 \leq p < \infty$, see [5] and [13], respectively. Other kind of dynamical properties for weighted composition operators acting on the smooth function space were investigated in [18].

Let Y be a Banach space of holomorphic functions on an open set $U \subset \mathbb{C}$ and $\phi: U \to U$ is a holomorphic map. Then, the *composition operator with symbol* ϕ is defined by $C_{\phi}f = f \circ \phi$, for any $f \in Y$.

In this paper, we study expansivity, shadowing and Li-Yorke chaos for the operators C_{ϕ} defined on the Hardy Hilbert space of the open right half-plane $H^2(\mathbb{C}_+)$, this operators has been studied in [7, 14, 15, 20]. The only linear fractional self-maps of \mathbb{C}_+ that induce continuous composition operators on $H^2(\mathbb{C}_+)$ are the *affine maps*, that is, $\phi(w) = aw + b$ with a > 0 and $Re(b) \ge 0$.

This work was inspired by the results of Noor and Severiano in [20], since the class of composition operators under investigation is not hypercyclic. Moreover, the notions of expansivity and shadowing has not been explored in this context. We provide examples of operators exhibiting expansivity and the shadowing property which are not hyperbolic, but they are not Li-Yorke chaotic. These examples show the richness of linear dynamics and how this differs from finite nonlinear dimensional dynamics.

Let us now describe the organization of the article. In section 2 we fix the notations and recall the definitions and a few results which will be important in our work. In section 3, we characterize some notions of expansivity and show that in certain cases, there exist some equivalences between these notions. In section 4, we provide conditions to guarantee or not the shadowing property and we give a characterization of the positive shadowing property. Finally, in section 5, we show that composition operators on the Hardy space of the open right half-plane can not be Li-Yorke chaotic.

2. Preliminaries

In this section, we fix some terminology and recall some definitions and results about spectral theory, linear dynamics and Hardy space on the open right half-plane \mathbb{C}_+ . Here, \mathbb{D} and \mathbb{T} denote the open unit disc and the unit circle in the complex plane \mathbb{C} , respectively. X is a complex Banach space, the set $S_X = \{x \in X : ||x|| = 1\}$ is the *unit sphere* of X and (M, d) is a compact metric space. In addition, by an operator on X we mean a continuous linear map $T : X \to X$.

Let $T: X \to X$ be an operator. The spectrum $\sigma(T)$ of T is the set

 $\sigma(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not invertible continuous operator}\}.$

Recall that $\sigma(T) \subset \mathbb{C}$ is always a non-empty compact set. The *point spectrum*, i.e. the set of eigenvalues of T, is denoted by $\sigma_p(T)$. The *approximative point spectrum* of T is defined by

$$\sigma_A(T) = \{\lambda \in \mathbb{C} : \text{there exists sequence } (h_n) \subset S_X \text{ such that } \| (T - \lambda I) h_n \| \xrightarrow{n \to \infty} 0 \}.$$

Obviously, $\sigma_p(T) \subset \sigma_A(T)$. The following inclusions always hold

$$\partial \sigma(T) \subset \sigma_A(T) \subset \sigma(T)$$

where $\partial \sigma(T)$ denotes the boundary of $\sigma(T)$. Moreover, if T is a self-adjoint operator on a complex Hilbert space H and $x \in H$, then there is a unique positive Radon measure μ on $\sigma(T)$ such that

$$\langle f(T)x,x\rangle = \int_{\sigma(T)} f(t)d\mu(t) \text{ for all } f \in C(\sigma(T)).$$

So, $\mu(\sigma(T)) = ||x||^2$. The measure μ is called the *spectral measure* associated to T and x.

2.1. Notions of linear dynamics. We start this section introducing a definition of expansivity for homeomorphisms defined on compact metric spaces. A homeomorphism $f: M \to M$ is expansive if there is C > 0 such that if $x, y \in M$ and $d(f^n(x), f^n(y)) \leq C$ for all $n \in \mathbb{Z}$ then x = y. Expansivity is a well known property of dynamical systems. For continuous operators, we provide a definition that contain several versions of expansivity.

Definition 1. Let $T: X \to X$ be an invertible operator on X. We say that

- (1) T is expansive if for every $z \in S_X$ there exists $n \in \mathbb{Z}$ such that $||T^n z|| \ge 2$.
- (2) T is positively expansive if for every $z \in S_X$ there exists $n \in \mathbb{N}$ such that $||T^n z|| \ge 2$.
- (3) T is uniformly expansive if there exists $n \in \mathbb{N}$ such that

$$z \in S_X \Longrightarrow ||T^n z|| \ge 2 \text{ or } ||T^{-n} z|| \ge 2.$$

(4) T is uniformly positively expansive if there exists $n \in \mathbb{N}$ such that

$$z \in S_X \Longrightarrow ||T^n z|| \ge 2.$$

The following result characterizes the expansivity for operators defined on Hilbert spaces.

Theorem 1 ([4]). Let T be a continuous operator on a Hilbert space H. The following assertions hold:

- (1) T is positively expansive if and only if $\sup_{n \in \mathbb{N}} ||T^n(x)|| = +\infty$, for every nonzero $x \in H$
- (2) T is uniformly positively expansive if and only if $\lim_{n \to +\infty} ||T^n(x)|| = +\infty$ uni-
- formly on S_H . If, in addition, T is an invertible normal operator then
- (3) T is expansive if and only if $\sigma_p(T^*T) \cap \mathbb{T} = \emptyset$.

Now, we introduce the notion of shadowing for homeomorphisms over compact metric spaces. Given a homeomorphism $h: M \to M$, recall that $(x_n)_{n \in \mathbb{Z}}$ is a δ -pseudotrajectory of h ($\delta > 0$) if $d(h(x_n), x_{n+1}) \leq \delta$ for all $n \in \mathbb{Z}$. h has the shadowing property if for every $\epsilon > 0$ there is $\delta > 0$ such that every δ -pseudotrajectory $(x_n)_{n \in \mathbb{Z}}$ of h is ϵ -shadowed by a real trajectory of h, that is, there is $x \in M$ such that $d(x_n, h^n(x)) < \epsilon$ for all $n \in \mathbb{Z}$. The definition of shadowing has a simplified formulation in the setting of linear dynamics. Here, we use this formulation.

Definition 2. Let $T: X \to X$ be an operator on X. A sequence $(x_n)_{n \in \mathbb{Z}} \subset X$ is called a δ -pseudotrajectory of T, where $\delta > 0$, if

$$||Tx_n - x_{n+1}|| \leq \delta$$
, for all $n \in \mathbb{Z}$.

Definition 3. Let $T: X \to X$ be an invertible operator on X. Then T is said to have the *shadowing property* if for every $\epsilon > 0$ there exists $\delta > 0$ such that every δ -pseudotrajectory $(x_n)_{n \in \mathbb{Z}}$ of T is ϵ -shadowed by a real trajectory of T, that is, there exists $x \in X$ such that

$$||T^n x - x_n|| \le \epsilon$$
, for all $n \in \mathbb{Z}$.

When T is an operator not necessarily invertible, we can define the notion of positive shadowing property for T. In such case, we replace the set \mathbb{Z} by \mathbb{N} in the above definition.

Theorem 2. Let $T: X \to X$ be an invertible operator on X. Suppose that $X = M \oplus N$, where M and N are closed subspaces of X with $T(M) \subset M$ and $T^{-1}(N) \subset N$. If $\sigma(T|M) \subset \mathbb{D}$ and $\sigma(T^{-1}|N) \subset \mathbb{D}$, then T has the shadowing property.

An operator $T: X \to X$ is said to be *hyperbolic* if $\sigma(T) \cap \mathbb{T} = \emptyset$. As consequence of Theorem 2, we have the following results:

Corollary 1. Every invertible hyperbolic operator $T : X \to X$ has the shadowing property.

Corollary 2. If T is an invertible normal operator on a Hilbert space H, then T has the shadowing property if and only if T is hyperbolic.

Finally, we introduce the notion of Li-Yorke chaos for continuous maps over compact metric spaces. The self-map $g: M \to M$ is said to be *Li-Yorke chaotic* if there exists an uncountable set $S \subset M$ (called a *scrambled set* for g) such that each pair $(x, y) \in S \times S$ with $x \neq y$, is a Li-Yorke pair for g, in the sense that

$$\liminf_{n \to \infty} d(g^n x, g^n y) = 0 \text{ and } \limsup_{n \to \infty} d(g^n x, g^n y) > 0.$$

The following result is well-known with respect to the recent advances of linear dynamics.

Theorem 3 ([3]). Let T be an operator on X. The following assertions are equivalent:

(1) T is Li-Yorke chaotic;

(2) T admits a semi-irregular vector, that is, a vector $y \in X$ such that

$$\liminf_{n \to \infty} \|T^n y\| = 0 \text{ and } \limsup_{n \to \infty} \|T^n y\| > 0;$$

(3) T admits an irregular vector, that is, a vector $z \in X$ such that

$$\liminf_{n \to \infty} \|T^n z\| = 0 \text{ and } \limsup_{n \to \infty} \|T^n z\| = \infty.$$

An important consequence of the previous result is the following corollary.

Corollary 3 ([3]). Let T be a Li-Yorke chaotic operator. The following assertions hold:

(1) $\sigma(T) \cap \mathbb{T} \neq \emptyset$. (2) T is not normal. (3) T is not compact. 2.2. Hardy space $H^2(\mathbb{C}_+)$. Now, we consider the Hardy space $H^2(\mathbb{C}_+)$ of the open right half-plane \mathbb{C}_+ . This is a Hilbert space of analytic functions on \mathbb{C}_+ for which the norm

$$||f||_{2}^{2} = \sup_{0 < x < \infty} \frac{1}{\pi} \int_{-\infty}^{\infty} |f(x+iy)|^{2} dy$$

is finite. For each $\beta \in \mathbb{C}_+$, let k_β denote the *reproducing kernel* for $H^2(\mathbb{C}_+)$ at β , that is,

$$k_{\beta}(w) = \frac{1}{w + \overline{\beta}}.$$

These kernels satisfy the fundamental relation $\langle f, k_{\beta} \rangle = f(\beta)$ for all $f \in H^2(\mathbb{C}_+)$. If $\phi : \mathbb{C}_+ \to \mathbb{C}_+$ is a holomorphic map and C_{ϕ} is the composition operator with symbol ϕ , then we have that $C_{\phi}^* k_{\beta} = k_{\phi(\beta)}$, for every $\beta \in \mathbb{C}_+$.

Elliott and Jury [7, Theorem 3.1] proved that a holomorphic map $\phi : \mathbb{C}_+ \to \mathbb{C}_+$ induces a continuous operator C_{ϕ} on $H^2(\mathbb{C}_+)$ if and only if $\phi(\infty) = \infty$ and if the non-tangential limit

$$\phi'(\infty) := \lim_{w \to \infty} \frac{w}{\phi(w)} \tag{1}$$

exists and is finite. Moreover, the norm of C_{ϕ} on $H^2(\mathbb{C}_+)$ is given by $||C_{\phi}|| = \sqrt{\phi'(\infty)}$. Matache [15] proved that the only linear fractional self-maps of \mathbb{C}_+ that induce continuous composition operators on $H^2(\mathbb{C}_+)$ are the *affine maps*

$$\phi(w) = aw + b,\tag{2}$$

where a > 0 and $Re(b) \ge 0$. Such a map ϕ is said to be of *parabolic-type* if a = 1 and is a *parabolic automorphism* if additionally Re(b) = 0. These symbols are useful to characterize normal composition operators C_{ϕ} .

Theorem 4 ([20]). Let $\phi(w) = aw + b$ with a > 0 and $Re(b) \ge 0$. Then

- (1) C_{ϕ} is normal if and only if a = 1 or Re(b) = 0,
- (2) C_{ϕ} is self-adjoint if and only if a = 1 and $b \ge 0$,
- (3) C_{φ} is unitary if and only if a = 1 and Re(b) = 0.

On the other hand, ϕ is of hyperbolic-type if $a \neq 1$ and is a hyperbolic automorphism if additionally Re(b) = 0. A hyperbolic non-automorphisms, these are precisely the symbols $\phi(w) = aw + b$ with $a \in (0, 1) \cup (1, \infty)$ and Re(b) > 0.

For simplicity, we call as hyperbolic of type I to every map $\phi(w) = aw + b$ with $a \in (0, 1)$ and $Re(b) \ge 0$. Similarly, we call as hyperbolic of type II to every map $\phi(w) = aw + b$ with $a \in (1, \infty)$ and $Re(b) \ge 0$.

We finish this section recalling known results about the spectrum of the operator C_{ϕ} on $H^2(\mathbb{C}_+)$ (see Theorems 1.1 and 1.2 of [19]).

Theorem 5 ([19]). Let $\phi(w) = w + b$ be a parabolic self-map of \mathbb{C}_+ with $Re(b) \ge 0$. Then the spectrum of C_{ϕ} acting on $H^2(\mathbb{C}_+)$ is

- (1) $\sigma(C_{\phi}) = \mathbb{T}$, when Re(b) = 0,
- (2) $\sigma(C_{\phi}) = \{e^{bt} : t \in [0, \infty)\} \cup \{0\}, when b \in \mathbb{C}_+.$

Theorem 6 ([19]). Let $\phi(w) = aw + b$ be a hyperbolic self-map of \mathbb{C}_+ with $Re(b) \ge 0$ and $b \ne 0$. Then the spectrum of C_{ϕ} acting on $H^2(\mathbb{C}_+)$ is

(1) $\sigma(C_{\phi}) = \{\lambda \in \mathbb{C} : |\lambda| = a^{-1/2}\}, \text{ when } Re(b) = 0,$ (2) $\sigma(C_{\phi}) = \{\lambda \in \mathbb{C} : |\lambda| \le a^{-1/2}\}, \text{ when } b \in \mathbb{C}_+.$

3. Expansivity of C_{ϕ}

In this section, we fully characterize the different notions of expansivity for the family of all composition operators on $H^2(\mathbb{C}_+)$ induced by a certain class of affine maps.

3.1. Expansivity in a positive sense of C_{ϕ} . To explore the (uniformly) positive expansivity of C_{ϕ} on $H^2(\mathbb{C}_+)$, we restrict ourselves to a particular class of affine symbols ϕ given by:

$$\phi(z) = az + b, \ z \in \mathbb{C}_+, \ a \in (0, 1) \text{ and } Re(b) \ge 0$$

i.e. ϕ is hyperbolic of type I. It is well known that hyperbolic automorphic maps induce invertible and normal composition operators on $H^2(\mathbb{C}_+)$ (see, [15, 20]), the following proposition shows that, in addition, they also induce positively expansive composition operators on $H^2(\mathbb{C}_+)$.

Proposition 1. Let $\phi(w) = aw + b$ be a self-map of \mathbb{C}_+ . If ϕ is hyperbolic of type *I*, then C_{ϕ} is positively expansive on $H^2(\mathbb{C}_+)$.

Proof. Suppose that ϕ is hyperbolic of type I and let $f \in H^2(\mathbb{C}_+)$ nonzero. Since $Re(b) \geq 0$ by the Blaschke zeros condition (see [8, Pag.53]), for f we can assume without loss of generality that $f\left(1+\frac{b}{1-a}\right) \neq 0$ so we can take $0 < \delta < \left|f\left(1+\frac{b}{1-a}\right)\right|$. Then, by using $\lim_{n \to +\infty} a^n = 0$ there exists $n_1 \in \mathbb{N}$ such that

$$\left| f\left(1 + \frac{1 - a^{n_1}}{1 - a}b\right) \right| > \left| f\left(1 + \frac{b}{1 - a}\right) \right| - \delta.$$
(3)

Moreover, for a > 0 and $w \in \mathbb{C}_+$ by using induction on ϕ^n iterations it is easily obtained that

$$\phi^{[n]}(w) = \begin{cases} w + nb & \text{if } a = 1\\ a^n w + \frac{(1 - a^n)}{1 - a}b & \text{if } a \neq 1 \end{cases}$$
(4)

Besides, by fundamental relation of reproductive nucleus with $\beta := a^{-n} \in \mathbb{C}_+$ follows that

$$\langle C^n_{\phi}f, k_{a^{-n}} \rangle = \left(C^n_{\phi}f \right) \left(a^{-n} \right) = f \left(\phi^n(a^{-n}) \right),$$

and $||k_{a^{-n}}|| = \frac{\sqrt{2}}{2}a^{n/2}$ for $n \in \mathbb{N}$. Applying Cauchy-Schwarz inequality, we have

$$\left\|C_{\phi}^{n}f\right\| \ \left\|k_{a^{-n}}\right\| \ge \left|f(\phi^{n}(a^{-n}))\right| \quad n \in \mathbb{N},$$

hence, by equation 4 with $w := a^{-n}$ the following estimate for the C_{ϕ} -orbit of f holds

$$\left\|C_{\phi}^{n}f\right\| \geq \frac{2}{\sqrt{2}a^{n/2}} \left\|f\left(1+\frac{1-a^{n}}{1-a}b\right)\right\|, \quad n \in \mathbb{N}.$$

Finally, since $a \in (0, 1)$ is clear that $\lim_{n \to +\infty} \frac{1}{a^{n/2}} = +\infty$ so we can fix $n_2 \in \mathbb{N}$ such that

$$\frac{1}{a^{n_2/2}} > \frac{\sqrt{2}}{\left|f\left(1 + \frac{b}{1-a}\right)\right| - \delta} \tag{5}$$

Consequently, by combining 3 and 5 we obtain $||C_{\phi}^{n}f|| > 2$ for $n := \max\{n_1, n_2\}$. \Box

Remark 1. Let $\delta > 0$, $a \in (0, 1)$ and $Re(b) \ge 0$. Applying fundamental relation of reproductive nucleus in $H^2(\mathbb{C}_+)$ we show that $S_{H^2(\mathbb{C}_+)}$ is locally bounded in $H^2(\mathbb{C}_+)$. In fact, for $z \in \mathbb{C}_+$ choose $0 < \xi < Re(z)$ and consider $B(z,\xi)$, so if $w \in B(z,\xi)$ then $0 < Re(z) - \xi < Re(w)$. Consequently,

$$|f(w)| \le \frac{1}{\sqrt{2Re(w)}} \le \frac{1}{\sqrt{2(Re(z) - \xi)}}$$

for any $f \in S_{H^2(\mathbb{C}_+)}$ and all $w \in B(z,\xi)$. By using Arzela-Ascoli Theorem we can choose $n_0 \in \mathbb{N}$ such that

$$\left| f\left(1 + \frac{1 - a^{n_0}}{1 - a}b\right) - f\left(1 + \frac{b}{1 - a}\right) \right| < \delta, \text{ for any } f \in S_{H^2(\mathbb{C}_+)}.$$

The rest of the reasoning is identical to the proof of Proposition 1 but it should be noted that n does not depend on any $f \in S_{H^2(\mathbb{C}_+)}$. Consequently, we obtain the following result:

Proposition 2. Let $\phi(w) = aw + b$ be a self-map of \mathbb{C}_+ . If ϕ is hyperbolic of type I then C_{ϕ} is uniformly positively expansive on $H^2(\mathbb{C}_+)$.

In the following result we show that all affine maps ϕ hyperbolic of type I ($a \in (0, 1)$ and $Re(b) \geq 0$) provides a necessary condition for an affine (not necessarily invertible) composition operator C_{ϕ} to be (uniformly) positively expansive on $H^2(\mathbb{C}_+)$.

Proposition 3. Let $\phi(w) = aw + b$ be a self-map of \mathbb{C}_+ satisfies one of following assertions:

- (1) ϕ is of parabolic type or the identity map;
- (2) ϕ is hyperbolic of type II.

Then C_{ϕ} is not (uniformly) positively expansive on $H^2(\mathbb{C}_+)$.

Proof. Let us show that if $a \geq 1$ then $H^2(\mathbb{C}_+)$ contains elements whose orbits are bounded. In fact, recall that $C_{\phi}^* = a^{-1}C_{\varphi}$ where $\varphi(w) = a^{-1}w + a^{-1}\overline{b}$ for any $w \in \mathbb{C}_+$, so a simple calculation shows that

$$\phi^{[n]}(\varphi^{[n]}(w)) = \begin{cases} w + 2nRe(b) & \text{if } a = 1\\ \\ w + 2(1-a)^{-1}(1-a^n)Re(b) & \text{if } a \neq 1 \end{cases}$$

Consequently,

$$\begin{split} \left\| C_{\phi}^{n} k_{w} \right\|^{2} &= \left(C_{\phi}^{n} k_{w}, C_{\phi}^{n} k_{w} \right) \\ &= \left(\left(C_{\phi}^{n} \right)^{*} C_{\phi}^{n} k_{w}, k_{w} \right) \\ &= a^{-n} \left(C_{\varphi}^{n} C_{\phi}^{n} k_{w}, k_{w} \right) \\ &= a^{-n} k_{w} \left(\phi^{n} \left(\varphi^{n} (w) \right) \right) \\ &= \begin{cases} \frac{1}{2Re(w) + 2nRe(b)} & \text{if } a = 1 \\ \frac{1}{2a^{n}Re(w) + 2(1-a)^{-1}a^{n}(1-a^{n})Re(b)} & \text{if } a \neq 1 \end{cases}$$

i.e.,

$$\left\|C_{\phi}^{n}k_{w}\right\| = \begin{cases} \frac{1}{\sqrt{2Re(w) + 2nRe(b)}} & \text{if } a = 1\\ \frac{1}{\sqrt{2a^{n}Re(w) + 2(1-a)^{-1}a^{n}(1-a^{n})Re(b)}} & \text{if } a > 1 \end{cases}$$
(6)

so the C_{ϕ} -orbit of k_w is bounded by $1/\sqrt{2Re(w)}$ when $a \in [1, +\infty)$. Thus, we have that C_{ϕ} is not positively expansive on $H^2(\mathbb{C}_+)$ by Theorem 1 part (1). Now, let us show that C_{ϕ} is not uniformly positively expansive on $H^2(\mathbb{C}_+)$. In fact, suppose that ϕ is *parabolic-type*, that is, a = 1. Notice that $k_w \in S_{H^2(\mathbb{C}_+)}$ if and only if $Re(w) = \frac{1}{2}$ so take $w_n := \frac{1}{2} + ni$, then by using 6 we obtain

$$\lim_{n \to +\infty} \left\| C_{\phi}^{n} k_{w_{n}} \right\| = \begin{cases} 0, & Re(b) > 0\\ 1, & Re(b) = 0 \end{cases}$$

In any case, this implies C_{ϕ} is not uniformly positively expansive on $H^2(\mathbb{C}_+)$ by Theorem 1 part (2). Besides, if ϕ is hyperbolic of type II then for all $n \in \mathbb{N}$ follows that

$$f \in S_{H^2(\mathbb{C}_+)}$$
 implies $||C_{\phi}^n f|| \le ||C_{\phi}^n|| ||f|| = \frac{1}{\sqrt{a^n}} < 1$,

which shows that C_{ϕ} is not uniformly positively expansive on $H^2(\mathbb{C}_+)$.

Combining Propositions 1, 2 and 3, we obtain the following result:

Theorem 7. Let $\phi(w) = aw + b$ be a self-map of \mathbb{C}_+ . Then, the following statements are equivalent:

- (1) C_{ϕ} is positively expansive on $H^2(\mathbb{C}_+)$;
- (2) ϕ is hyperbolic of type I;
- (3) C_{ϕ} is uniformly positively expansive on $H^2(\mathbb{C}_+)$.

3.2. Forward and backward expansivity for C_{ϕ} . We characterize the expansivity for invertible composition operators on $H^2(\mathbb{C}_+)$.

Theorem 8. Let $\phi(w) = aw + ib$ with $b \in \mathbb{R}$. Then, the following statements are equivalent:

- (1) C_{ϕ} is expansive on $H^2(\mathbb{C}_+)$;
- (2) C_{ϕ} is uniformly expansive on $H^2(\mathbb{C}_+)$;
- (3) ϕ is hyperbolic-type.

Proof. (1) \iff (3). Since Re(ib) = 0 then by Theorem 4 and [15, Theorem 2.4] is known that C_{ϕ} is normal and invertible. Let's show that $\frac{1}{a} \in \sigma_p(C_{\phi}^*C_{\phi})$. Indeed, for $w \in \mathbb{C}_+$ we obtain that

$$(C_{\phi}^*C_{\phi}) k_w(z) = a^{-1}C_{(\phi \circ \varphi)}k_w(z) = a^{-1}k_w((\phi \circ \varphi)(z)) = \frac{1}{az + a\overline{w}} = \frac{1}{a} k_w(z),$$

by using $C_{\phi}^* = a^{-1}C_{\varphi}$ with $\varphi(w) = a^{-1}w + a^{-1}\overline{ib}$ (see [20, Proposition 1]) and $\phi(\varphi(z)) = z$, so $(C_{\phi}^*C_{\phi}) k_w = \frac{1}{a}k_w$. Hence, for $\lambda \in \sigma_p(C_{\phi}^*C_{\phi})$ we get

$$\lambda f(z) = \left(C_{\phi}^* C_{\phi}\right) f(z) = a^{-1} f(\phi(\varphi(z))) = a^{-1} f(z)$$

i.e., $\lambda f(z) = a^{-1}f(z)$ for some $f \neq 0$, so $\lambda = \frac{1}{a}$ and consequently $\sigma_p(C_{\phi}^*C_{\phi}) = \{\frac{1}{a}\}$ and by Theorem 1 part (3) C_{ϕ} is expansive if and only if $a \neq 1$. This proves that (1) and (3) are equivalent.

(2) \iff (3). Let ϕ is hyperbolic-type; then, when $a \in (1, +\infty)$ we get $||C_{\phi}|| = \frac{1}{\sqrt{a}} < 1$ so $|\lambda| < 1$ for all $\lambda \in \sigma(C_{\phi})$, i.e., $\sigma(C_{\phi}) \cap \mathbb{T} = \emptyset$. Besides, if $a \in (0, 1)$ then by [15, Theorem 2.12] and [19, Theorem 4.1] is known that

$$\sigma(C_{\phi}) = \left\{ \lambda \in \mathbb{C} : |\lambda| = \frac{1}{\sqrt{a}} \right\}$$

since $a \in (0,1)$ this implies $\sigma(C_{\phi}) \cap \mathbb{T} = \emptyset$. In any case, since $\sigma_A(C_{\phi}) \subset \sigma(C_{\phi})$ we obtain $\sigma_A(C_{\phi}) \cap \mathbb{T} = \emptyset$, i.e., C_{ϕ} is uniformly expansive. Reciprocally, if ϕ is *parabolic-type* then by taking $k_{w_n} \in S_{H^2(\mathbb{C}_+)}$ with $w_n := \frac{1}{2} + in$ is easily obtained

$$||C_{\phi}^{n}k_{w_{n}}|| = ||C_{\phi}^{-n}k_{w_{n}}|| = 1, \ n \in \mathbb{N}$$

so C_{ϕ} is not uniformly expansive on $H^2(\mathbb{C}_+)$.

4. Shadowing for C_{ϕ}

In this section, we characterize positive shadowing for continuous composition operators on $H^2(\mathbb{C}_+)$ induced for a certain class of affine maps.

Proposition 4. Let $\phi : \mathbb{C}_+ \to \mathbb{C}_+$ be a holomorphic map with $\phi(\eta) = \eta$ for some $\eta \in \mathbb{C}_+$ such that C_{ϕ} is bounded on $H^2(\mathbb{C}_+)$. Then, C_{ϕ} does not have the positive shadowing property on $H^2(\mathbb{C}_+)$.

Proof. Let $\delta > 0$ and consider $\{f_n^{\delta}\}_{n \in \mathbb{N}} \subset H^2(\mathbb{C}_+)$ given by

$$f_n^{\delta} := \frac{\delta}{\|C_{\phi}f\|} \sum_{k=0}^{n-1} C_{\phi}^{n-k} f, \quad n = \in \mathbb{N},$$

where $f \in H^2(\mathbb{C}_+)$ is chosen such that $f(\eta) \neq 0$. Then, $(f_n^{\delta})_{n \in \mathbb{N}}$ is a δ -pseudotrajectory of C_{ϕ} for all $\delta > 0$. In fact, by standard arguments we obtain

$$\begin{aligned} \|C_{\phi}f_{n}^{\delta} - f_{n+1}^{\delta}\| &= \frac{\delta}{\|C_{\phi}f\|} \left\| \left[\sum_{k=0}^{n-1} C_{\phi}^{n-k+1}f - \sum_{k=0}^{n} C_{\phi}^{n-k+1}f \right] \right\| \\ &= \frac{\delta}{\|C_{\phi}f\|} \left\| \left[\sum_{k=0}^{n-1} C_{\phi}^{n-k+1}f - \sum_{k=0}^{n-1} C_{\phi}^{n-k+1}f - C_{\phi}f \right] \right\| \\ &= \frac{\delta}{\|C_{\phi}f\|} \|C_{\phi}f\| \\ &= \delta \end{aligned}$$

Now, notice that

$$f_n^{\delta}(\eta) = \frac{n\delta}{\|C_{\phi}f\|} f(\eta), \quad n \in \mathbb{N}.$$

Consequently, for any $g \in H^2(\mathbb{C}_+)$ applying fundamental relation of reproductive nucleus with $\beta := \eta \in \mathbb{C}_+$ we get

$$\|C_{\phi}^{n}g - f_{n}^{\delta}\| \geq \sqrt{2Re(\eta)} |g(\phi^{n}(\eta)) - f_{n}^{\delta}(\eta)| \geq \sqrt{2Re(\eta)} \left|\frac{n\delta}{\|C_{\phi}f\|}|f(\eta)| - |g(\eta)|\right|$$

that is,

$$\|C_{\phi}^{n}g - f_{n}^{\delta}\| \ge \left|\frac{n\delta}{\|C_{\phi}f\|}|f(\eta)| - |g(\eta)|\right|, \quad n \in \mathbb{N}.$$

When $n \to +\infty$, we obtain $||C^n_{\phi}g - f^{\delta}_n|| \to +\infty$, which shows that the $(f^{\delta}_n)_{n \in \mathbb{N}}$ cannot be ϵ -shadowed for any $\epsilon > 0$. Therefore, C_{ϕ} does not have the positive shadowing property on $H^2(\mathbb{C}_+)$.

The main goal of this section is to classify the symbols that induce a composition operator satisfying some type of shadowing on $H^2(\mathbb{C}_+)$. Proposition 4 states that existence of fixed points for the symbol implies non-existence of positive shadowing property.

Theorem 9. Let $\phi(w) = aw + b$ be a self-map of \mathbb{C}_+ with a > 0 and $Re(b) \ge 0$. The following assertions hold:

- S.1 If ϕ is a hyperbolic automorphism or a hyperbolic non-automorphism of type II, then C_{ϕ} has the positive shadowing property on $H^2(\mathbb{C}_+)$.
- S.2 If ϕ is parabolic-type or a hyperbolic non-automorphism of type I, then C_{ϕ} does not have the positive shadowing property on $H^2(\mathbb{C}_+)$.

Proof. S.1 Suppose ϕ is a hyperbolic automorphism, that is, $a \in (0, 1) \cup (1, +\infty)$ and Re(b) = 0. Here, C_{ϕ} is an invertible normal operator on $H^2(\mathbb{C}_+)$. From Theorem 6 follows that $\sigma(C_{\phi}) = \{z : |z| = \frac{1}{\sqrt{a}}\}$. When $a \in (0, 1)$, Corollary 1 guarantees shadowing property for C_{ϕ} . On the other hand, when $a \in (1, +\infty)$, we have that $\sigma(C_{\phi}) \subset \mathbb{D}$. Then, there exist $p \in (0, 1)$ and $K \ge 1$ such that $\|C_{\phi}^n\| \le Kp^n$ for all $n \in \mathbb{N} \cup \{0\}$. We argue as in [4, Theorem 13], for $\epsilon > 0$, we choose $\delta = \frac{(1-p)\epsilon}{2K} > 0$. Let $(f_n)_{n \in \mathbb{N} \cup \{0\}}$ be a δ -pseudotrajectory of C_{ϕ} and define the sequence $g_n = f_n - C_{\phi}f_{n-1}$. By induction,

$$f_n = C_{\phi}^n f_0 + \sum_{i=1}^n C_{\phi}^{n-i} g_i \text{ for all } n \in \mathbb{N}.$$

Hence,

$$||f_n - C^n_{\phi} f_0|| \le \sum_{i=1}^n ||C^{n-i}_{\phi}|| ||g_i||$$

Since $||g_n|| < \delta$ for all $n \in \mathbb{N}$, it follows that

$$\|f_n - C^n_\phi f_0\| \le \sum_{i=1}^n \delta K p^{n-i} < \frac{K\delta}{1-p} = \frac{\epsilon}{2}$$

for $n \in \mathbb{N}$. Therefore, $(f_n)_{n \in \mathbb{N} \cup \{0\}}$ is ϵ -shadowed by $(C_{\phi}^n f_0)_{n \in \mathbb{N} \cup \{0\}}$ and C_{ϕ} has the positive shadowing property. Finally, if ϕ is a hyperbolic non-automorphism of type II, that is, $a \in (1, +\infty)$ and Re(b) > 0. From Theorem 6 follows that $\sigma(C_{\phi}) = \{z : |z| \leq \frac{1}{\sqrt{a}}\}$. Then, $\sigma(C_{\phi}) \subset \mathbb{D}$ and we proceed as in the case of hyperbolic automorphism with $a \in (1, \infty)$.

S.2 If ϕ is a parabolic map, then a = 1 and Re(b) = 0 or a = 1 and Re(b) > 0. When a = 1 and Re(b) = 0, then C_{ϕ} is a unitary operator. In particular, C_{ϕ} is a normal and invertible operator. From Theorem 5 follows that C_{ϕ} is not hyperbolic and it does not have the shadowing property by Corollary 2. Now, if a = 1 and Re(b) > 0, then C_{ϕ} is a normal operator (not invertible). By Theorem 5 the spectrum $\sigma(C_{\phi})$ intersects T. Now, we argue as in [4, Theorem 30], assuming that C_{ϕ} has positive shadowing property. We argue as in the proof of [4, Theorem 27] to obtain a function $f \in H^2(\mathbb{C}_+)$ such that

$$\frac{n\delta}{3} - 1 < \|C_{\phi}^n f\| < n\delta + 1 \quad \text{for all } n \in \mathbb{N} \cup \{0\}, \tag{7}$$

where $\delta > 0$ is the constant in the definition of shadowing associated to $\epsilon = 1$. Let $S_{\phi} = C_{\phi}^* C_{\phi}$ and let μ be the spectral measure associated to S_{ϕ} and f. Then, $\mu(\sigma(S_{\phi}) \cap (1, +\infty)) = 0$, because

$$0 \le \int_{\sigma(S_{\phi})} t^n d\mu(t) = \langle S_{\phi}^n f, f \rangle = \|C_{\phi}^n f\|^2 \text{ for all } n \in \mathbb{N} \cup \{0\}$$

Hence, using Hölder inequality we obtain that $||C_{\phi}^{n}f|| \leq (\mu(\sigma(S_{\phi})))^{\frac{1}{2}} = ||f||$ for all $n \in \mathbb{N} \cup \{0\}$, which contradicts (7). Consequently, C_{ϕ} does not have the positive

shadowing property. Finally, if ϕ is a hyperbolic non-automorphism of type I, then $a \in (0, 1), Re(b) > 0$ and $\frac{b}{1-a}$ is a fixed point of ϕ . Thus, Proposition 4 guarantees that C_{ϕ} does not have positive shadowing property on $H^2(\mathbb{C}_+)$.

To combining the assertions S.1 and S.2 of the previous theorem, we have the following result that characterizes the positive shadowing property.

Corollary 4. Let $\phi(w) = aw + b$ be a self-map of \mathbb{C}_+ with a > 0 and $Re(b) \ge 0$. Then, C_{ϕ} has positive shadowing property if and only if ϕ is a hyperbolic automorphism or a hyperbolic non-automorphism of type II.

5. LI-YORKE CHAOS FOR C_{ϕ}

Finally we prove that $H^2(\mathbb{C}_+)$ does not support any Li-Yorke chaotic composition operator with linear fractional symbol.

Theorem 10. C_{ϕ} is not Li-Yorke chaotic on $H^2(\mathbb{C}_+)$ for any affine symbol ϕ .

Proof. For $w \in \mathbb{C}_+$, let $\phi(w) = aw + b$ with a > 0 and $Re(b) \ge 0$. If ϕ is to be parabolic-type or Re(b) = 0, then C_{ϕ} is normal by Theorem 4, so C_{ϕ} is not Li-Yorke by Corollary 3. The same conclusion follows for the case $b \ge 0$, because C_{ϕ} is self-adjoint. Now, suppose that ϕ is to be a hyperbolic non-automorphism, that is,

$$\phi(w) = aw + b$$
, with $a \in (0, 1) \cup (1, +\infty)$ and $Re(b) > 0$.

By using 1 we get $||C_{\phi}|| = \frac{1}{\sqrt{a}}$. Hence, for $a \in (1, +\infty)$ we obtain $||C_{\phi}|| < 1$ and this implies $|\lambda| < 1$ for all $\lambda \in \sigma(C_{\phi})$ so that $\sigma(C_{\phi}) \cap \mathbb{T} = \emptyset$. By Corollary 3, C_{ϕ} is not Li-Yorke chaotic . Besides, for the case $a \in (0, 1)$; suppose that $f \in H^2(\mathbb{C}_+)$ nonzero then there exists $N \in \mathbb{N}$ such that $||C_{\phi}^N f|| > 2$ (see proof of Proposition 1) this implies that $\lim \inf ||C_{\phi}^n f|| \ge 2$. Therefore, f cannot be an irregular vector for C_{ϕ} in $H^2(\mathbb{C}_+)$ (see also, [4, Theorem A]).

Acknowledgments

The authors thank to the anonymous referees for their valuable comments to improve this article. CF.A. was supported by ANID Chile and BASEX-PD/2023-02 project of University of Sinú, Cartagena.

References

- [1] BAYART, F., AND MATHERON, E. Dynamics of linear operators, vol. 179 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2009.
- [2] BERMÚDEZ, T., BONILLA, A., MARTÍNEZ-GIMÉNEZ, F., AND PERIS, A. Li-Yorke and distributionally chaotic operators. J. Math. Anal. Appl. 373, 1 (2011), 83–93.
- [3] BERNARDES, JR., N. C., BONILLA, A., MÜLLER, V., AND PERIS, A. Li-Yorke chaos in linear dynamics. *Ergodic Theory Dynam. Systems* 35, 6 (2015), 1723–1745.
- [4] BERNARDES, JR., N. C., CIRILO, P. R., DARJI, U. B., MESSAOUDI, A., AND PUJALS, E. R. Expansivity and shadowing in linear dynamics. J. Math. Anal. Appl. 461, 1 (2018), 796–816.
- [5] BERNARDES, JR., N. C., DARJI, U. B., AND PIRES, B. Li-Yorke chaos for composition operators on L^p-spaces. Monatsh. Math. 191, 1 (2020), 13–35.
- [6] EISENBERG, M., AND HEDLUND, J. H. Expansive automorphisms of Banach spaces. Pacific J. Math. 34 (1970), 647–656.
- [7] ELLIOTT, S., AND JURY, M. T. Composition operators on Hardy spaces of a half-plane. Bull. Lond. Math. Soc. 44, 3 (2012), 489–495.
- [8] GARNETT, J. B. Bounded analytic functions, first ed., vol. 236 of Graduate Texts in Mathematics. Springer, New York, 2007.

- [9] GODEFROY, G., AND SHAPIRO, J. H. Operators with dense, invariant, cyclic vector manifolds. J. Funct. Anal. 98, 2 (1991), 229–269.
- [10] GRIVAUX, S., MATHERON, E., AND MENET, Q. Linear dynamical systems on Hilbert spaces: typical properties and explicit examples. *Mem. Amer. Math. Soc.* 269, 1315 (2021), v+147.
- [11] GROSSE-ERDMANN, K.-G., AND PERIS MANGUILLOT, A. *Linear chaos*. Universitext. Springer, London, 2011.
- [12] LI, T. Y., AND YORKE, J. A. Period three implies chaos. Amer. Math. Monthly 82, 10 (1975), 985–992.
- [13] MAIURIELLO, M. Expansivity and strong structural stability for composition operators on L^p spaces. Banach J. Math. Anal. 16, 4 (2022), Paper No. 51, 20.
- [14] MATACHE, V. Composition operators on Hardy spaces of a half-plane. Proc. Amer. Math. Soc. 127, 5 (1999), 1483–1491.
- [15] MATACHE, V. Invertible and normal composition operators on the Hilbert Hardy space of a half-plane. Concr. Oper. 3, 1 (2016), 77–84.
- [16] MAZUR, M. Hyperbolicity, expansivity and shadowing for the class of normal operators. Funct. Differ. Equ. 7, 1-2 (2000), 147–156 (2001).
- [17] PRĂJITURĂ, G. T. Irregular vectors of Hilbert space operators. J. Math. Anal. Appl. 354, 2 (2009), 689–697.
- [18] PRZESTACKI, A. Dynamical properties of weighted composition operators on the space of smooth functions. J. Math. Anal. Appl. 445, 1 (2017), 1097–1113.
- [19] SCHRODERUS, R. Spectra of linear fractional composition operators on the Hardy and weighted Bergman spaces of the half-plane. J. Math. Anal. Appl. 447, 2 (2017), 817–833.
- [20] WALEED NOOR, S., AND SEVERIANO, O. R. Complex symmetry and cyclicity of composition operators on H²(C₊). Proc. Amer. Math. Soc. 148, 6 (2020), 2469–2476.

Área de Ciencias Básicas Exactas, Universidad del Sinú Seccional Cartagena, Av. El Bosque, Transversal 54 № 30-72, Cartagena de Indias 130001, Colombia

Current address: Instituto de Matemáticas, Pontificia Universidad Católica de Valparaíso, Blanco Viel 596, Cerro Barón, Valparaíso, Chile

Email address: carlosfalvarez@unisinu.edu.co; carlos.alvarez.e@pucv.cl

Departamento de Ciencias Naturales y Exactas, Universidad de la Costa, Calle 58 $\mathrm{N}^\circ55\text{-}56,$ Barranquilla, Colombia

Email address: jhenriqu13@cuc.edu.co