Edson de Faria and Sebastian van Strien

Before going into Dennis Sullivan's work on dynamics, we would like to share some of our reminiscences on the remarkable way in which he influenced a huge number of mathematicians, including the two of us. Both while at IHES and at CUNY, Dennis had an office which came with an anteroom. Our impression is that he would spend most of his time in this anteroom, talking about mathematics with whoever he had invited or whoever was around. Quite often while listening to somebody, he would end up giving a new spin or a new interpretation to what they had been saying. Similarly, he would explain what he was working on, trying out new ideas, and also often explaining results of others. Spending time with him was always an incredible experience.

In this spirit, Dennis explained much of his work on renormalisation to Welington de Melo. In turn, Welington would then try to explain what he had heard and learned to SvS. When he could not convince SvS of some argument, Welington would go back to Dennis and this process would repeat again, sometimes many times. This is how the final chapter of the book *One dimensional dynamics* of SvS with Welington de Melo came into being, see [30]. This chapter contains a full exposition of Dennis' remarkable renormalisation theory, arguably the only place in which it was published with full details.

At the Graduate Center of CUNY, Dennis coordinated the *Einstein Chair Seminar*, more informally known as the Sullivan seminar, bringing in speakers from all over the world. The seminar ran once a week with talks in the afternoon, but the invited speakers usually came in the morning, and intense mathematical discussions ensued through lunch, and oftentimes even after the speaker's talk, following a short break for tea. During the talks, Dennis often asked questions not necessarily to know the answer himself, but because he knew that somebody else in the audience would

E. de Faria

Instituto de Matematica e Estatistica, Universidade de Sao Paulo, e-mail: edson@ime.usp.br S. van Strien

Department of Mathematics, Imperial College London, e-mail: s.van-strien@imperial.ac.uk

find the answer helpful. In this way, Dennis took on the role of introducing two people to each other. His presence in the audience would usually make a talk much more accessible and interesting. His questions would often clarify connections that would have remained implicit otherwise.

When Dennis invented or learned about a new mathematical idea, he would push this idea to the limit. For him it was very important to understand what this idea would give, and equally important to find out what the limitations of this idea might be. Moreover, whenever possible, he liked to associate names to arguments such as the *dollar argument, smallest interval argument* or the *non-coiling argument* in order to synthesise a complex proof into its core ideas.

Often he mentioned that to understand a proof properly, you should treat it like a three dimensional object. You should not only look at it from one side, but from all sides. So in this sense, in his view, a proof was about mathematical understanding rather than about 'killing' a theorem.

Indeed, Dennis' interest in a result might not necessarily be in the power of the statement per se, but in the tools that are used in the proof of this result. Once he understands the tools and ideas, then he probably can recover the statement of the results by himself.

So let us turn to the field of dynamical systems. Dennis Sullivan always had a keen interest in the field of dynamical systems, and already in the 1970's published several high impact papers in this area, many of them remarkably short. Let us first highlight a few of his papers on smooth dynamical systems, and then his ground-breaking papers on Kleinian groups and holomorphic dynamics.

1 Smooth dynamics

Although his early papers on dynamics treat a diverse range of problems, they all have an overarching theme: what smoothness (or other) structures are compatible with a particular dynamical setting.

1.1 From topology to dynamics

In the mid-seventies, Dennis started to become more and more interested in dynamics, transitioning from the pure study of structures on manifolds to the study of dynamical objects such as flows and, more generally, foliations on manifolds. One of his first striking results in this direction is the following.

Theorem 1 (Counterexample to the periodic orbit conjecture, see [100], [98]) There exists a flow on a compact five-dimensional manifold all of whose orbits are periodic, and yet the lenghts of such orbits are not bounded.

This theorem¹ was announced in [100], and a more detailed argument given in [98]. Dennis' topological-geometric construction in the latter paper yields a flow on a smooth 5-dimensional manifold M which is Lipschitz, but he states that he sees no reason why the example could not be made smooth. In an addendum at the end of the paper, Dennis briefly explains an idea due to Kuiper that results in an example in which the flow is C^{∞} . He also explains a beautiful construction due to Thurston which yields a real-analytic flow with the desired properties on the manifold $M = \mathbb{R}^5/(\Gamma \times \mathbb{Z} \times \mathbb{Z})$, where Γ is the group

$$\Gamma = \left\{ \begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix} : x, y, z \in \mathbb{Z} \right\} .$$

In other words, Γ is the so-called *discrete Heisenberg group*.

1.2 Rigidity in smooth dynamics

The following paper, joint with Shub, dates back to 1985.

Theorem 2 (Expanding maps of the circle, see [94]) *Two* C^r , $r \ge 2$, *expanding maps of the circle which are absolutely continuously conjugate are* C^r *conjugate.*

The proof of this result is short and starts by invoking well-known results to reduce the problem to the setting where both maps preserve the Lebesgue measure. Using the assumption that the maps are C^2 and expanding makes it possible to consider an iterate of the maps taking a small interval to big scale with bounded distortion. This then implies that the assumption that the conjugacy *h* is absolutely continuous gives that *h* is, in fact, Lipschitz. Going on from there, one obtains that *h* is C^r .

In some broader sense the main idea of this paper was the starting point for quite a lot of later research. Indeed, the pullback argument that Dennis introduced in the field of holomorphic dynamics is somewhat similar in spirit. Quite a few other papers followed on from this work. For example,

- if the multipliers of corresponding periodic points of two topologically conjugate unimodal interval are equal, then the conjugacy is smooth, see [75, 64] (there are corresponding results in higher dimensions);
- if a conjugacy between two interval maps is smooth at some point, then it is smooth everywhere, see [2];
- there are quite a few very interesting related results for group actions of circle maps, see for example [31];

¹ This theorem is also discussed in McMullen's beautiful talk on Dennis' work at MSRI in the spring of 2022, see https://vimeo.com/702914316?embedded=true&source= vimeo_logo&owner=106107493

• when studying the smoothness of a conjugacy between two maps on defined on Cantor sets, Sullivan introduced the notion of a scaling function on a Cantor set, see [107] and also [89].

Another paper, joint with Norton, on rigidity in dynamics is concerned with Denjoy examples of C^1 torus diffeomorphisms $T^2 \rightarrow T^2$ which are isotopic to the identity. These are maps f which are topologically semi-conjugate to a minimal translation on the torus, i.e. hf = Rh, where h is a continuous map of T^2 onto itself, homotopic to the identity, such that the set of $x \in T^2$ for which the cardinality of $h^{-1}(x)$ is greater than 1 is nonempty and countable. Then the interior of any $h^{-1}(x)$, if nonempty, is a wandering domain for f.

Theorem 3 (Smoothness and wandering domains for torus maps, see [86]) Let f be a torus map of Denjoy type, and let $\Gamma \neq T^2$ be its minimal set. Then (i) if f preserves a measurable, essentially bounded conformal structure on Γ , then the maps f^n , restricted to the prime end boundaries of the wandering domains, cannot be uniformly quasiconformal; and (ii) if f preserves a C^2 conformal structure on Γ , then f cannot be C^3 .

One consequence of this theorem is that one cannot have a C^3 Denjoy diffeomorphism so that the iterates of some disc are all disjoint. Of course this theorem is a partial higher dimensional analogue of Denjoy's famous theorem showing that a C^2 diffeomorphism of the circle without periodic orbits cannot have wandering intervals, and therefore must be topologically conjugate to an irrational rotation. Again quite a few papers followed on from this work, for example:

- it is not possible to have C¹ toral diffeomorphism with wandering round discs, see [82];
- very recently it was shown that there exist smooth and even real analytic diffeomorphisms of Denjoy type on the torus with a wandering topological disc, see [82, 115]; see also [4].
- wandering topological discs were also established for smooth two dimensional diffeomorphisms, see [56], and even for polynomial maps in higher dimensions [3, 12].

Another theorem Dennis proved, jointly with Gambaudo and Tresser, is the following (informally stated).

Theorem 4 (Smoothness and linking number of periodic orbits for diffeomorphism of the disc, see [41]) Let f be a C^1 diffeomorphism of the disc with periodic orbits p_n , $n \ge 0$, so that for each $n \ge 0$ the periodic orbit p_{n+1} 'cycles as a satelite' around p_n . Then the average linking number between p_{n+1} and p_n must converge as $n \to \infty$.

One of the inspirations for this paper was a question by Smale, who asked whether it was possible to construct a smooth diffeomorphism on the disc with infinitely many hyperbolic periodic saddles, but without periodic sinks or sources (or

4

neutral points). In [10, 40] such examples were constructed in the C^1 respectively C^2 category. It was subsequently shown that one can construct smooth and even real analytic diffeomorphisms with these properties, see [42] and also [22, 67], namely with a Feigenbaum-Coullet-Tresser Cantor attractor. The constructions in those papers build on the renormalisation theory developed for interval maps. The braid type of the periodic orbits is quite different from those in [10, 40], and as far as we know it is not yet known whether there are smooth diffeomorphisms which are topologically conjugate to the ones constructed in those papers (in which it is conjectured that one cannot construct C^3 diffeomorphisms topologically conjugate to their examples).

What the theorems discussed in this section have in common is that they are about invariant structures, and that the full theory in this direction has not yet been completed. For this reason having the additional conformal structure was quite appealing to Dennis. Another reason to start working on holomorphic dynamics in the 1980's was of course that he saw a compelling analogue with the theory on Kleinian groups that he had been working on previously – see section 2 below.

1.3 Further results

Another research direction we would like to explicitly mention is Dennis' work on currents, see [99, 92]. In his beautiful survey talk at the Abel Prize lectures at the University of Oslo on Dennis' work, Étienne Ghys singles out this work (this talk is available on YouTube²).

2 Dynamics and ergodic theory of Kleinian groups

Dennis's interest in the geometric, dynamical and ergodic properties of discrete groups of hyperbolic isometries dates back to the mid to late 1970s. His work in this area was motivated in part by Mostow's rigidity theorem from two decades earlier, and in part by the work of Ahlfors on Kleinian groups, especially his famous finiteness theorem from 1965. Dennis was also greatly influenced by Thurston's work on geometric structures over 3-manifolds. It was Lipman Bers who first told Dennis about the so-called *Ahlfors conjecture*, according to which the limit set of a Kleinian group acting in hyperbolic 3-space either has zero Lebesgue measure in the sphere at infinity, or else it is equal to the entire sphere. This is now a theorem, thanks to the work of several mathematicians – see for instance [72] and references therein.

Let us present a brief account of the contributions of Dennis to this beautiful subject. Before we do that, we need to recall a few preliminary notions. For general

² See https://www.youtube.com/watch?v=reC5-XUeH_4.

background on the geometry of discrete groups and hyperbolic geometry, especially in dimension 3, we recommend [8], [77] and [72]. For a systematic exposition of the work of Dennis (and Patterson) on the ergodic theory of discrete groups, see [85].

The Moebius group

Consider the one-point compactification $\widehat{\mathbb{R}}^n = \mathbb{R}^n \cup \{\infty\}$ of Euclidean *n*-space. The Moebius group in dimension *n* is the group $MG(\mathbb{R}^n)$ consisting of all transformations $T: \widehat{\mathbb{R}}^n \to \widehat{\mathbb{R}}^n$ which arise as all possible compositions of (linear) conformal transformations of the form $x \mapsto Ax + b$, where *A* is a scalar multiple of an orthogonal matrix and $b \in \mathbb{R}^n$, with the inversion $J: \widehat{\mathbb{R}}^n \to \widehat{\mathbb{R}}^n$ given by $J(x) = x/|x|^2$ for $x \neq 0, J(0) = \infty$ and $J(\infty) = 0$. The elements of $MG(\mathbb{R}^n)$ are called *Moebius transformations*.

Hyperbolic space

Let us denote by \mathbb{H}^n hyperbolic *n*-space, which we view as the open unit ball $B^n = \{x : |x| < 1\} \subset \mathbb{R}^n$ endowed with the *hyperbolic metric* (also called *Poincaré metric*) given by

$$ds = \frac{2|dx|}{1 - |x|^2}$$

The *ideal boundary* or *sphere at infinity* of hyperbolic *n*-space is by definition the sphere $S^{n-1} = \partial B^n$, endowed with the standard conformal structure inherited from \mathbb{R}^n . It is customary to denote the sphere at infinity by S_{∞} . The group $\mathrm{Isom}^+(\mathbb{H}^n)$ of orientation-preserving isometries of this metric consists precisely of all *Moebius transformations* that preserve the unit ball, *i.e.*, those $T \in MG(\mathbb{R}^n)$ such that $T(B^n) = B^n$. Every $T \in \mathrm{Isom}^+(\mathbb{H}^n)$ acts on the sphere at infinity as a *conformal automorphism*. The elements of $\mathrm{Isom}^+(\mathbb{H}^n)$ are classified according to their action on S_{∞} as follows. If $T \in \mathrm{Isom}^+(\mathbb{H}^n)$ has exactly one fixed point in S_{∞} , then *T* is said to be a *parabolic* transformation. If it has exactly two fixed points in S_{∞} , then it is called a *loxodromic* transformation. All other elements of $\mathrm{Isom}^+(\mathbb{H}^n)$ are said to be *elliptic*.

Kleinian groups

A (generalized) *Kleinian group* is a discrete subgroup $\Gamma \subset \text{Isom}^+(\mathbb{H}^n)$. Discreteness means in particular that the orbit $\Gamma(x) = \{\gamma x : \gamma \in \Gamma\}$ of any point $x \in B^n$ can only accumulate on the sphere at infinity. The set $\Lambda(\Gamma) \subseteq S_{\infty}$ of all such accumulation points is the *limit set* of Γ (see figure 1). Its complement $\Omega(\Gamma) = S_{\infty} \setminus \Lambda(\Gamma)$ is the *domain of discontinuity* or *ordinary set* of Γ . Clearly, both $\Lambda(\Gamma)$ and $\Omega(\Gamma)$ are completely invariant under the action of Γ . When n = 3, the ordinary set $\Omega(\Gamma)$ is

precisely the *domain of normality* of Γ , that is to say, the set of all points $z \in S_{\infty} \equiv \widehat{\mathbb{C}}$ having a neighborhood $V_z \subset \widehat{\mathbb{C}}$ such that $\{\gamma|_{V_z} : \gamma \in \Gamma\}$ is a normal family in the sense of Montel (thus, $\Omega(\Gamma)$ is the analogue of the Fatou set for a rational map, and the limit set $\Lambda(\Gamma)$ is the analogue of the Julia set – see §4). A Kleinian group is said to be *non-elementary* if its limit set consists of more than two points.




Fig. 1 The limit set of a Kleinian group sitting in the sphere at infinity is oftentimes a fractal object. [Credit: This picture was generated using C. McMullen's program "lim", available at https://people.math.harvard.edu/ ctm/programs/.]

There are various ways under which the Γ -orbit of a point $x \in B^n$ can accumulate on a point of $\Lambda(\Gamma)$ in the sphere at infinity. The two most important are a *conical* approach and a *horospherical* approach. Let us be more precise.

Definition 5 Let $\Gamma \subset \text{Isom}^+(\mathbb{H}^n)$ be a Kleinian group and let $\xi \in \Lambda(\Gamma)$.

(i) We say that ξ is a *conical limit point* of Γ if for each x ∈ Bⁿ there exists a sequence {γ_n} ⊂ Γ such that the ratio

$$\frac{|\xi - \gamma_n(x)|}{1 - |\gamma_n(x)|}$$

remains bounded as $n \to \infty$.

(ii) We say that ξ is a *horospherical limit point* of Γ if for each $x \in B^n$ there exists a sequence $\{\gamma_n\} \subset \Gamma$ such that the ratio

$$\frac{|\xi - \gamma_n(x)|^2}{1 - |\gamma_n(x)|}$$

goes to zero as $n \to \infty$.

It is an exercise to show that if ξ is the fixed point of a loxodromic element of Γ , then ξ is a conical limit point of Γ . The set of all conical limit points of Γ is called the *conical limit set*, and it is denoted $\Lambda_c(\Gamma)$. The set of all horospherical limit points of Γ is called the *horospherical limit set*, and it is denoted $\Lambda_h(\Gamma)$. Clearly, these are both Γ -invariant.

An important class of Kleinian groups is the class consisting of so-called *convex co-compact groups*. Given a (non-elementary) Kleinian group Γ , let $\Lambda = \Lambda(\Gamma)$ be its limit set, and consider the convex hull $C(\Lambda)$ of Λ inside hyperbolic space. Then $C(\Lambda)$ is invariant under Γ , and we say that Γ is convex co-compact if the quotient $C(\Lambda)/\Gamma$ is compact. It is not difficult to see that if Γ is convex co-compact, then every element of Λ is a conical limit point – in other words, $\Lambda_c(\Gamma) = \Lambda(\Gamma)$ in this case.

Hyperbolic manifolds

The quotient space $M_{\Gamma} = \mathbb{H}^n/\Gamma$ of hyperbolic *n*-space by a Kleinian group Γ is what one calls an *orbifold* (after Thurston). Such quotient is always a manifold when n = 2, 3, but it may fail to be one when n > 3. However, if Γ acts *freely and properly discontinuously* on \mathbb{H}^n , then M_{Γ} is indeed a manifold. Such manifolds are called *hyperbolic*. The natural quotient projection $\mathbb{H}^n \to M_{\Gamma}$ is a proper covering map, and therefore the hyperbolic metric of \mathbb{H}^n descends to M_{Γ} . Thus, \mathbb{H}^n is the universal covering space of M_{Γ} , and the fundamental groups Γ_1, Γ_2 are *conjugate* subgroups of Isom⁺(\mathbb{H}^n), *i.e.*, if there exists $\gamma \in \text{Isom}^+(\mathbb{H}^n)$ such that $\Gamma_1 = \gamma^{-1}\Gamma_2\gamma$, then the corresponding orbifolds M_{Γ_i} , i = 1, 2, are *isometric*, and conversely.

Quasi-conformal homeomorphisms

A quasiconformal homeomorphism $h: S_{\infty} \to S_{\infty}$ is a homeomorphism which is differentiable Lebesgue almost-everywhere, and whose derivative at each point of differentiability maps round spheres onto ellipsoids whose ratios between the largest axis and the smallest axis yield a measurable function on the sphere that is essentially bounded. The essential norm of this function is called the quasi-conformal distortion of *h*, denoted *K_h*. It turns out that if *K_h* = 1 then *h* is in fact *conformal*. Every quasi-conformal homeomorphism determines a measurable field of ellipsoids, also known as a *measurable conformal structure* on *S*_∞. In dimension two, such measurable conformal structures can be integrated to recover *h* up to post-composition by a conformal map – a famous result known as the measurable Riemann mapping theorem – but no such theorem exists in higher dimensions.

2.1 Sullivan's rigidity theorem

It is a truly remarkable theorem due to G. Mostow [84] that complete finite-volume hyperbolic *n*-manifolds are determined up to isometry by their fundamental groups when $n \ge 3$. This is the famous *Mostow rigidity theorem*, which can be formally stated as follows.

Theorem 6 (Mostow Rigidity) *Let* M *and* N *be two complete, finite-volume hyperbolic n-manifolds with* $n \ge 3$ *, and let* $\theta : \pi_1(M) \to \pi_1(N)$ *be an isomorphism between their fundamental groups. Then there exists an isometry* $f : M \to N$ *between both manifolds such that the induced isomorphism* $f_* : \pi_1(M) \to \pi_1(N)$ *agrees with* θ .

In fact, this theorem was proved by Mostow for *closed* manifolds, *i.e.*, compact manifolds without boundary. It was then extended to finite volume manifolds by Marden [71] in dimension n = 3, and by Prasad [90] in all dimensions $n \ge 3$.

The way Mostow proved his theorem was by first showing that the manifolds M,N are *pseudo-isometric* in the following sense. A continuous, surjective map $\phi : M \to N$ between two hyperbolic manifolds is a *pseudo-isometry* if (a) it induces an isomorphism between the fundamental groups of both manifolds and moreover (b) there exist constants K > 1 and $\delta > 0$ such that

$$\frac{1}{K} \le \frac{d_N(\phi(x), \phi(y))}{d_M(x, y)} \le K$$

for each pair of points $x, y \in M$ such that $d_M(x, y) \ge \delta$. Here d_M, d_N denote the hyperbolic distances in M and N, respectively. Condition (b) is saying that a pseudoisometry distorts hyperbolic distances between points by a bounded amount, provided these points are sufficiently far apart.

If one lifts a given pseudo-isometry between M and N to their universal covering space, one gets a pseudo-isometry of hyperbolic n-space. Mostow proved in [84]

that every pseudo-isometry of \mathbb{H}^n extends continuously to the sphere at infinity as a *quasiconformal homeomorphism* $h: S_{\infty} \to S_{\infty}$. The key step in the proof is to show by means of an ergodic argument that if Γ is a finite-volume Kleinian group then there is only one measurable conformal structure on S_{∞} which is Γ -invariant. This implies that the quasiconformal homeomorphism h is actually conformal, which in turn means that the two associated Kleinian groups $\Gamma_M \simeq \pi_1(M), \Gamma_N \simeq \pi_1(N)$ are conjugate subgroups of $\mathrm{Isom}^+(\mathbb{H}^n)$, and therefore the hyperbolic manifolds M and N must be isometric.

Here is another way of stating Mostow's theorem. Following [102], we say that a hyperbolic manifold M is *Mostow-rigid* if any pseudo-isometry between M and another hyperbolic manifold N is homotopic to an isometry. Recast in this language, Mostow's theorem states that every complete, finite-volume hyperbolic M is Mostow-rigid.

Dennis proved in [102], in his own words, a *maximal extension* of Mostow's theorem. In order to state his theorem, let us introduce some notation. Given a hyperbolic manifold M, a point $p \in M$ and r > 0, let $V_M(p,r)$ denote the hyperbolic volume of the set $\{x \in M : d_M(p,x) < r\}$. For example, for the hyperbolic ball of radius r in hyperbolic n-space, we have

$$V_{\mathbb{H}^n}(0,r) = \omega_n \int_0^{\tanh(r/2)} \frac{2^n |x|^{n-1} d|x|}{(1-|x|^2)^n} = \omega_n \int_0^r \sinh^{n-1} t \, dt \sim \operatorname{const} \cdot e^{(n-1)r} \, .$$

Here, ω_n denotes the euclidean area of $S_{\infty} = S^{n-1}$.

Lemma 7 Let $M = \mathbb{H}^n / \Gamma$ be a hyperbolic *n*-manifold. Then the following assertions are equivalent.

(i) The ratio $V_M(p,r)/V_{\mathbb{H}^n}(0,r)$ goes to zero as $r \to \infty$.

(ii) The Kleinian group Γ acts conservatively on S_{∞} .

(iii) The horospherical limit set of Γ has full Lebesgue measure on S_{∞} .

A fourth assertion equivalent to the above three is that the fundamental domain of Γ in \mathbb{H}^n has zero Lebesgue measure on S_{∞} . We now have all the necessary elements to state the Sullivan rigidity theorem.

Theorem 8 (Sullivan Rigidity) Let M be a complete hyperbolic n-manifold, and suppose that M satisfies any of the equivalent conditions of Lemma 7. Then M is Mostow rigid.

Dennis deduces this theorem from the following result, also due to him.

Theorem 9 Let $\Gamma \subset \text{Isom}^+(\mathbb{H}^n)$ be a Kleinian group, and consider its action on the sphere at infinity. Suppose v is a measurable conformal structure (i.e., a measurable field of ellipsoids) which is Lebesgue almost everywhere invariant under Γ . Then v agrees a.e. with the standard conformal structure of S_{∞} on the limit set Λ_{Γ} .

In the case of hyperbolic 3-manifolds, *i.e.*, when n = 3, a major consequence of this theorem is obtained by combining it with the Ahlfors finiteness theorem. Recall

that a Riemann surface *X* is said to be of *finite type* if *X* is obtained from a compact Riemann surface by removing from it a finite set of points.

Theorem 10 (Ahlfors Finiteness Theorem) Let $\Gamma \subset PSL(2, \mathbb{C})$ be a finitely generated Kleinian group. Then $\Omega(\Gamma)/\Gamma$ is a finite union of Riemann surfaces of finite type.

In particular, the *Teichmüller space* Teich $(\Omega(\Gamma)/\Gamma)$ is finite-dimensional. Hence we have the following result.

Corollary 11 Let $\Gamma \subset PSL(2,\mathbb{C})$ be a finitely generated Kleinian group. Then the space of quasi-conformal deformations of Γ is parametrized by $\text{Teich}(\Omega(\Gamma)/\Gamma)$, and is therefore finite-dimensional.

In particular, if Γ has a dense orbit on the sphere S^2 , then the space of quasiconformal deformations of Γ reduces to a point, *i.e.*, Γ is quasi-conformally rigid.

2.2 Conformal densities and Patterson-Sullivan measures

Let Γ be a non-elementary Kleinian group acting on $\mathbb{H}^n \cup S_{\infty}$, and as before let $\Lambda(\Gamma) \subseteq S_{\infty}$ be its limit set. Also as before, let $\Lambda_c(\Gamma) \subseteq \Lambda(\Gamma)$ be its conical limit set. Generalizing work of Patterson for Fuchsian groups [88], Dennis was able to construct, in [101], an invariant measure for the geodesic flow on the unit tangent bundle of the hyperbolic manifold \mathbb{H}^n/Γ . This measure comes from a *conformal density* μ on the sphere at infinity, and the geodesic flow is either ergodic or dissipative, depending on whether μ assigns positive or zero measure to $\Lambda_c(\Gamma)$, respectively. We proceed to a brief description of the construction. Details can be found either in the original paper by Dennis, or in the book by Nicholls [85].

Conformal densities

Let us start by clarifying what is meant by conformal density. Let *M* be a smooth manifold, let \mathscr{R} be a non-empty collection of Riemannian metrics on *M*, and let $\alpha > 0$. Following [101, p. 421], we define a *conformal density of dimension* α , or α -conformal density, on *M* (relative to \mathscr{R}) to be a function that assigns to each element $g \in \mathscr{R}$ a positive, finite Borel measure μ_g on *M* in such a way that, whenever g_1 and g_2 are in the same conformal class (*i.e.*, whenever $g_1 = \varphi g_2$ for some positive function φ), then μ_{g_1} and μ_{g_2} are in the same measure class, and the Radon-Nikodym derivative $d\mu_{g_1}/d\mu_{g_2}$ satisfies

$$\frac{d\mu_{g_1}}{d\mu_{g_2}} = \left(\frac{g_1}{g_2}\right)^{\alpha} \,.$$

We are not interested in conformal densities in such vast generality, but rather in the following specific context. We take $M = S_{\infty}$, and let $\mathscr{R} = \{g_x : x \in B^n\}$, where

 g_0 is the standard (euclidian) Riemannian metric on the sphere S_{∞} , and for each $x \in B^n$ the Riemannian metric g_x is obtained by transporting g_0 via any hyperbolic isometry mapping 0 to x. These metrics are all conformally equivalent. Thus, in the present context, we can think of a α -conformal density on the sphere at infinity as an assignment $x \mapsto \mu_x$ from points on hyperbolic space to positive measures on S_{∞} , all in the same measure class. Shortening the notation to $\mu_x = \mu_{g_x}$, we deduce after a simple calculation that

$$\frac{d\mu_{x_1}}{d\mu_{x_2}}(\xi) = \left(\frac{P(x_1,\xi)}{P(x_2,\xi)}\right)^{\alpha},\tag{1}$$

for each pair of points $x_1, x_2 \in B^n$ and all $\xi \in S^{n-1} \equiv S_{\infty}$, where

$$P(x,\xi) = \frac{1 - |x|^2}{|x - \xi|^2}$$

is the well-known Poisson kernel.

Given a non-elementary Kleinian group Γ , we are interested in conformal densities of the type just described *that are entirely supported in the limit set of* Γ *and that are* Γ *-invariant.* More precisely, we want to know whether there exists an α conformal density $\mu = {\mu_x : x \in B^n}$ such that

- (1) For each *x*, the measure μ_x has support in the limit set $\Lambda(\Gamma)$.
- (2) For each pair of points $x_1, x_2 \in B^n$, the measures μ_{x_1}, μ_{x_2} are mutually absolutely continuous, and the Radon-Nikodym derivative $d\mu_{x_1}/d\mu_{x_2}$ satisfies (1).
- (3) For all $x \in B^n$ and each $\gamma \in \Gamma$, we have $\gamma_* \mu_x = \mu_{\gamma x}$.

Given a Γ -invariant conformal density of dimension α in the sense just described, each of its associated measures μ_x is an α -conformal measure in the sense that

$$\mu_x(\gamma(E)) = \int_E |\gamma'_x(\xi)|^{\alpha} d\mu_x(\xi)$$

for each Borel set $E \subset S_{\infty}$ and each $\gamma \in \Gamma$ (*cf.* the discussion on conformal measures for rational maps in §3.2). The question as to whether such *Patterson-Sullivan measures* exist is examined below.

Patterson-Sullivan measures: construction

The *Poincaré series* of the (non-elementary) Kleinian group Γ is defined as

$$g_s(x,y) = \sum_{\gamma \in \Gamma} e^{-sd(x,\gamma y)} , \qquad (2)$$

where $x, y \in \mathbb{H}^n$, *d* is the hyperbolic metric on \mathbb{H}^n , and s > 0 is a real parameter. Whether the series (2) converges or not for a given value of *s* is independent of which points *x*, *y* one chooses. In order to state this more precisely, define the *critical*

exponent of Γ to be the number

$$\delta(\Gamma) = \inf\{s > 0 : g_s(0,0) < \infty\} ,$$

which turns out to be strictly positive when Γ is non-elementary³. Then it is a fact that the series (2) converges for all $s > \delta(\Gamma)$, and diverges for all $0 < s < \delta(\Gamma)$. It is also not difficult to prove that $\delta(\Gamma) \le n - 1$. What happens when $s = \delta(\Gamma)$? Obviously, only one of two things:

- (i) if $g_{\delta(\Gamma)}(0,0) < \infty$, we say that Γ is a group of *convergence type*;
- (ii) if $g_{\delta(\Gamma)}(0,0) = \infty$, we say that Γ is a group of *divergence type*.

For the construction to follow, let us fix a point $y \in B^n$ once and for all (for example, we could take y = 0). For each $x \in B^n$ and each $s > \delta(\Gamma)$, one constructs a positive Borel measure $\mu_{x,s}$ on the closure of B^n as follows. The rough idea is to place a Dirac mass at each point of the Γ -orbit of y, with weights that depend on the hyperbolic distance between each such point and x in a suitable way. Let us be more precise.

When the group Γ is of *divergence* type, one simply defines⁴

$$\mu_{x,s} = \frac{1}{g_s(y,y)} \sum_{\gamma \in \Gamma} e^{-sd(x,\gamma y)} \delta_{\gamma x}$$

With this definition, one can consider the weak limits of such measures when $s \searrow \delta(\Gamma)$. Since $g_s(y, y) \to \infty$ as $s \to \delta(\Gamma)$, the point masses are swept off to the sphere at infinity, and any weak limit will be a measure supported on the sphere (actually on the limit set). The existence of limits is guaranteed by a classical result in real analysis (namely, Helly's theorem).

However, when the group is of *convergence* type, the above will not work, because $g_s(y,y)$ remains bounded as $s \to \delta(\Gamma)$, and whatever limiting measure we get will still have an atom at each point in the Γ -orbit of y (recall that the goal is to obtain measures supported on the limit set of Γ). To circumvent this problem, Dennis borrows an idea due to Patterson [88] (in the Fuchsian case, n = 2) and introduces a *mollifier*, called, not surprisingly, the *Patterson auxiliary function*. This is a continuous non-decreasing function $h : \mathbb{R}^+ \to \mathbb{R}^+$ having the following properties:

- (1) For each $\varepsilon > 0$ there exists $r_0 > 0$ such that $h(tr) \le t^{\varepsilon} h(r)$ for all $r > r_0$ and all t > 1.
- (2) The series

$$\sum_{\in \Gamma} e^{-sd(x,\gamma y)} h(e^{d(x,\gamma y)})$$

converges for $s > \delta(\Gamma)$ and diverges for $s \leq \delta(\Gamma)$.

Using this function, one defines the modified Poincaré series

Ŷ

³ This was first proved by Beardon [7].

⁴ We denote by δ_z the Dirac probability measure concentrated at $z \in B^n$.

Edson de Faria and Sebastian van Strien

$$g_s^*(x,y) = \sum_{\gamma \in \Gamma} e^{-sd(x,\gamma y)} h(e^{d(x,\gamma y)})$$

This now diverges when $s \searrow \delta(\Gamma)$. Thus, for each $x \in B^n$ and each $s > \delta(\Gamma)$ we may now consider the positive Borel measure $\mu_{x,s}$ defined by

$$\mu_{x,s} = rac{1}{g_s^*(y,y)} \sum_{\gamma \in \Gamma} e^{-sd(x,\gamma y)} h(e^{d(x,\gamma y)}) \delta_{\gamma x} \ .$$

As before, the resulting weak limits as $s \searrow \delta(\Gamma)$ are positive Borel measures on the sphere at infinity, and their supports are contained in the limit set $\Lambda(\Gamma)$.

In either case, we have for each $x \in B^n$ a non-empty closed subset $M_x(\Gamma) \subset \mathcal{M}^+(S_\infty)$ of the space of all positive Borel measures on the sphere at infinity (endowed with the topology of weak convergence of measures). It is possible to prove that the $M_x(\Gamma)$'s are all homeomorphic (see for instance [85, Th. 3.4.1]).

We are now ready to summarize some of the main results obtained by Dennis in [101]. That paper is very rich, and we can hardly do any justice to it in such a short exposition.

The first theorem generalizes results obtained by Patterson and Bowen in the Fuchsian case.

Theorem 12 (Patterson-Sullivan Measures and Hausdorff Dimension, see [101]) Let $\Gamma \subset \text{Isom}^+(\mathbb{H}^n)$ be a non-elementary Kleinian group, and let $\delta = \delta(\Gamma)$ be its critical exponent.

- (i) There exists a δ -conformal density $\mu = {\mu_x : x \in B^n}$ on the sphere at infinity which is Γ -invariant and satisfies $\mu_x \in M_x(\Gamma)$ for each x.
- (ii) We have $\dim_H(\Lambda_c(\Gamma)) \leq \delta$, i.e., the conical limit set of Γ has Hausdorff dimension less than or equal to its critical exponent
- (iii) If Γ is convex co-compact, then the δ -conformal density in (i) is unique up to a scalar multiple, and for each euclidian ball $B(\xi,r)$ centered at a point $\xi \in \Lambda_c(\Gamma)$, and each $x \in B^n$, we have $\mu_x(B(\xi,r) \cap \Lambda_c(\Gamma)) \approx r^{\delta}$. In particular, $\dim_H(\Lambda_c(\Gamma)) = \delta$, i.e., the Hausdorff dimension of the conical limit set is equal to δ in this case.

The last item in the above statement is a very elegant result which generalizes an equally elegant result for Fuchsian groups due to Bowen [9].

Another striking result obtained by Dennis in [101] states that the *total-mass* function of a Γ -invariant conformal density in dimension $\delta(\Gamma)$ is an eigenfunction of the hyperbolic Laplacian. Let us state this result a bit more precisely, explaining the meaning of these terms. The hyperbolic Laplacian in $\mathbb{H}^n \equiv B^n$ (written in generalized polar coordinates) is the second-order differential operator

$$arDelta_h \,=\, rac{(1-r^2)^2}{4} \left[arDelta + rac{2(n-2)r}{1-r^2} rac{\partial}{\partial r}
ight] \,,$$

where Δ is the standard (euclidian) Laplacian – see for instance [1, p. 56]. If $\mu = {\mu_x : x \in B^n}$ is a Γ -invariant conformal density in dimension $\delta(\Gamma)$, its *total-mass*

function is the function $\varphi : \mathbb{H}^n \to \mathbb{R}$ given by

$$\varphi(x) = \int_{\partial B^n} d\mu_x(\xi) = \int_{S_\infty} d\mu_x(\xi) \; .$$

Theorem 13 *The total-mass function* φ *is an eigenfunction of the hyperbolic Laplacian,* i.e., *it satisfies*

$$\Delta_h \varphi = \lambda(\Gamma) \varphi$$
,

with eigenvalue $\lambda(\Gamma) = \delta(\Gamma)(1 + \delta(\Gamma) - n)$.

This statement is perhaps made more plausible if one takes into account that, for each α , we have

$$\Delta_h(P(x,\xi)^{\alpha}) = \alpha(\alpha - n + 1)P(x,\xi)^{\alpha}$$

a fact that can easily be checked by direct calculation.

Finally, as Dennis explains in [101], a Γ -invariant conformal density μ gives rise to a measure *m* on the unit tangent bundle of the quotient hyperbolic manifold \mathbb{H}^n/Γ which is *invariant under the geodesic flow*. In addition, the normalized probability measures $\varphi(x)^{-1}\mu_x$ can be used to generate a Markovian stochastic process on \mathbb{H}^n/Γ akin to Brownian motion. The beautiful synthesis obtained by Dennis in [101] relates the recurrent properties of this Markovian process with the ergodic properties of the geodesic flow on the quotient manifold, and can be informally stated as follows.

Theorem 14 (Ergodic Measures for the Geodesic Flow, see [101]) Let $\Gamma \subset \text{Isom}^+(\mathbb{H}^n)$ be a non-elementary Kleinian group, and let $\delta = \delta(\Gamma)$ be its critical exponent. Also, let μ be a Γ -invariant δ -conformal density. Consider the following assertions:

- (1) The conical limit set $\Lambda_c(\Gamma)$ has positive μ -measure.
- (2) The action of Γ on $S_{\infty} \times S_{\infty}$ minus the diagonal is ergodic with respect to $\mu \times \mu$.
- (3) The geodesic flow on the unit tangent bundle to \mathbb{H}^n/Γ is ergodic with respect to m_{μ} .
- (4) The group Γ is of divergence type.
- (5) The Markov process on \mathbb{H}^n/Γ is recurrent.

Then (1), (2) and (3) are equivalent, and they imply (4). If in addition $2\delta > n - 1$, then (4) implies (5), and (5) implies all the others (i.e., all five assertions are equivalent in this case).

2.3 Further results

Dennis has written a number of other very interesting papers on the geometry and dynamics of Kleinian groups. For example, in [105] he extended some of the above results from the convex co-compact case to the case of *geometrically finite groups* – one of the main consequences being the fact that the Hausdorff dimension of the

limit set of a geometrically finite group is equal to the critical exponent of the group. In [103], he investigated the excursion of geodesics on hyperbolic surfaces, relating their behaviour near cusps with classical results in Diophantine approximations.

3 Holomorphic dynamics

3.1 The reemergence of holomorphic dynamics in Paris in the 1980's

Right from the start the city of Paris has played a key role in the study of holomorphic dynamical systems. In the 1920's Julia and Fatou developed many results on the theory of iterations of rational maps $f: \overline{\mathbb{C}} \to \overline{\mathbb{C}}$. They introduced what is now called the *Fatou set*, the set of points in $\overline{\mathbb{C}}$ which have a neighbourhood N on which the iterates $f^n|N, n \in \mathbb{N}$, form a normal family, i.e. are equicontinuous. Similarly, the Julia set J(f) is defined as the complement of F(f). At the time the main tool Julia and Fatou had at their disposal was the Montel theorem, which states that a family of maps $f^n|N, n \in \mathbb{N}$ defined on an open set $N \subset \overline{\mathbb{C}}$ is normal if it has the property that there are three points in $\overline{\mathbb{C}}$ which are omitted in $\cup_n f^n(N)$. Among the many results they showed is that the Julia set is the closure of the set of repelling periodic points. They also developed a theory on the local dynamics near periodic points.

In the early 1980's there was a huge revival of this theory in Paris, with main drivers being Dennis Sullivan, Adrien Douady, Hamal Hubbard and Michael Herman. One of the main reasons for this resurgence was that it became increasingly clear that there were *new powerful tools* available, namely the *Measurable Riemann Mapping Theorem* (MRMT) and the notion of *quasiconformal maps*. These would make it possible to complete and go much beyond the theory initiated by Julia and Fatou in the 1920's.

There are quite a few equivalent definitions of the notion of a *quasiconformal* map, and all reflect that such maps are generalisations of conformal maps. One of these definitions is that $h: U \to V$ is a quasiconformal map if it is an orientation preserving homeomorphism between two domains U, V on \mathbb{C} so that the Beltrami equation

$$\frac{\partial h}{\partial \bar{z}} = \mu(z) \frac{\partial h}{\partial z}$$

makes sense and so that μ is Lebesgue measurable essentially bounded, *i.e.*, satisfies $\|\mu\|_{\infty} < 1$. This means that at each point $z \in U$ at which *h* is differentiable, the derivative Dh(z) maps circles to ellipses with uniformly bounded eccentricity.

The MRMT implies that each such μ is associated to a quasiconformal map h_{μ} and, crucially, that h_{μ} depends analytically on μ .

Dennis was amongst the first to realise the power of the MRMT in the field of holomorphic dynamics, partly because he had previously used it very successfully in the study of Kleinian groups. Parallel to Douady and Hubbard's seminal Orsay notes

[32, 33, 34, 35], which are a tour the force through the entire subject of holomorphic dynamics, Dennis proved the following remarkable theorem:

Theorem 15 (No-wandering-domains Theorem, see [106]) *Let f be a rational map on the Riemann sphere. Then each component of the Fatou set is eventually periodic.*

More precisely, for each component U of F(f) there exists $m \ge 0$ so that $V = f^m(U)$ is periodic, i.e. there exists $p \ge 1$ so that $f^p(V) = V$. Moreover, each periodic component V of F(f) can be classified. Indeed, it contains one of the following:

- 1. a periodic point of eigenvalue $\lambda = 0$ (called *superattractive*),
- 2. a periodic point of eigenvalue $0 < |\lambda| < 1$ (called *attractive*), or
- 3. ∂V contains a periodic point whose multiplier is a root of unity (*rational indifferent*),
- 4. f^p is analytically conjugate to an irrational rotation on V and either
 - a. V is a simply-connected Siegel disc or
 - b. a doubly-connected *Herman ring*.

To prove the first part of the statement one needs to show that if f is a rational map, then no component of its Fatou set is a wandering domain, i.e. a domain so that all its iterates are pairwise disjoint. In a nutshell Dennis' proof of this theorem goes as follows: suppose by contradiction that f has a wandering domain W. Then this makes it possible to construct an infinite dimensional space of deformations of f, contradicting that the space of rational maps of a given degree is finite dimensional.

That the space of rational maps is finite dimensional is crucial: shortly after the preprint version of [106] appeared, Baker constructed entire functions $f: \mathbb{C} \to \mathbb{C}$ which do have wandering domains. Sullivan's no wandering theorem has also been extended to the setting of entire maps (and similar spaces) with a finite number of singular values, see for example [37] and [43]. A very elegant proof of the no wandering domains theorem due to McMullen – which circumvents the use of the MRMT and uses an infinitesimal deformations argument more in line with Ahlfors' original proof of his finiteness theorem – can be found in [78, p. 90].

Unfortunately, as there is no corresponding MRMT in the real one-dimensional case, the analogous theory in the real one-dimensional case requires a careful combinatorial analysis together with an understanding of the non-linearity of the map. For circle diffeomorphisms this goes back to Denjoy in the 1930's and from this paper Dennis learned the *smallest interval argument*: Assume that *W* is a maximal wandering interval, i.e. that *W* is not contained in a larger wandering interval. Then for each $n \ge 3$ consider the smallest, say $f^i(W)$, amongst the collection of disjoint intervals $W, \ldots, f^n(W)$. Then $f^i(W)$ has neighbours on each side which are larger (or empty space in the case of an interval map). So $f^i(W)$ is well-inside the convex hull W'_i of the two neighbours. Using the way W'_i is chosen, the interval W'_i can be pulled back to an interval $W'_0 \supset W$ so that the pullbacks W'_0, \ldots, W'_i are essentially disjoint. This disjointness and the fact that f is a C^2 diffeomorphism implies that W is δ -well-inside W'_0 where δ does not depend on n. Using the maximality of W

this gives a contradiction. That $f^i(W)$ is well-inside the convex hull W'_i is often called *Koebe space* and is a property that is often used both in real and holomorphic dynamics.

The terminology *Koebe space* comes from the Koebe Lemma in complex analysis, which states that for each $\delta > 0$ there exists K > 0 so that when $U_0 \subset U$ are topological discs and $\phi : U \to \mathbb{C}$ is a univalent map and the modulus of $U \setminus U_0$ is at least δ then

$$rac{D\phi(z)|}{D\phi(z')|} \leq K ext{ for all } z, z' \in U_0.$$

The power of this Lemma is that *K* does not depend on ϕ . It turns out that in the real case there is an analogous result: for each $\delta > 0$ there exists K > 0 so that when $I_0 \subset I$ are intervals and $g: I \to \mathbb{R}$ is a diffeomorphism so that $Sg \ge 0$ then

$$\frac{|Dg(z)|}{|Dg(z')|} \le K \text{ for all } z, z' \in I_0.$$

In applications, g is usually the inverse of a diffeomorphic branch of an interval map f^n where f is assumed to have negative Schwarzian Sf < 0. The reason this is useful is that the Schwarzian property has the property that Sf < 0 implies $Sf^n < 0$ and that $Sf^n < 0$ implies $Sf^{-n} > 0$ (on diffeomorphic branches). Furthermore, if f is a real polynomial with only real critical points then Sf < 0. This observation that the negative Schwarzian could be used to bound the number of periodic attractors of an interval map was first made by Singer, see [95] but also appeared at around the same time in for example Herman's work [48]. A version of the Koebe Lemma in this setting was proved for the first time in [114]. That the Schwarzian derivative is related the distortion of cross-ratio was already known by E. Cartan in the 1930's, see the discussion in [30, Sections IV.1 and IV.2]. As the use of the above distortion estimate is so widespread, one often refers to the *Koebe Principle* and the assumption on the domains $U_0 \subset U$ (resp. $I_0 \subset I$) as *Koebe space*.

For interval maps and critical circle maps, the presence of critical points implies that one cannot control the non-linearity of the map and its iterates. Instead, it turns out that it is enough (i) to consider the *cross-ratio distortion* of a triple of adjacent intervals under iterates, (ii) assume that the map has some local symmetry around the critical points (e.g. the maps are non-flat at the critical points) and (iii) a more elaborate combinatorial analysis of orbits of wandering intervals. This was done by Guckenheimer, Yoccoz, Lyubich, Block, de Melo, van Strien, Martens in various generalities. For a history and a full analogue of Theorem 15 see [76, 30]. Probably the most elegant way of proving absence of wandering intervals in this setting can be found in [113]. Interestingly, it was Dennis who emphasised and insisted that the right smoothness class for (i) is $C^{1+Zygmund}$, whereas the earlier results required that the map was C^3 and even assumed that the map has negative Schwarzian. See [108], [109] or [30] for the definition of the classes $C^{1+Zygmund}$ and $C^{1+zygmund}$. The analogue of Sullivan's no-wandering Theorem 15 is:

Theorem 16 (See [76, 30]) Assume that f is an interval map which is $C^{1+Zygmund}$ and has non-flat critical points. Then f has no wandering interval. Moreover, if f

is a $C^{2+zygmund}$ map with non-flat critical points, then there exist $\kappa > 1$ and $n_0 \in \mathbb{N}$ such that

$$|Df^n(p)| > \kappa$$

for every periodic point p of f of period $n \ge n_0$.

Interestingly, it is not clear to what extent local symmetry around a critical point is crucial. Indeed, consider a map of the form

$$f(x) = \begin{cases} x^{\alpha} + c & \text{for } x > 0\\ x^{\beta} + c & \text{for } x < 0 \end{cases}$$

with $\alpha \neq \beta, \alpha, \beta > 1$ and *c* real. It is not known whether such a map can have wandering intervals. For $\alpha = \beta$ the answer is no, due to the previous theorem. For the case that $\alpha \neq \beta$ very little is known, except for the case that $\alpha = 1 < \beta$ and *f* has Feigenbaum-Coullet-Tresser dynamics, see [61] (and the proof of absence of wandering intervals in that case follows a rather curious approach).

The real bounds that go into the proof of the absence of wandering intervals for real maps, certainly inspired Dennis' proofs of complex bounds which are crucial in his renormalisation theory.

As mentioned, a crucial ingredient in the proof of real bounds is Schwarzian derivative, or more generally the notion of cross-ratio. A special cross-ratio inequality was used by Yoccoz to show that smooth circle homeomorphisms with a unique non-flat critical point cannot have wandering intervals, see [118]. More general cross-ratio inequalities were then used in [29] for the interval case and subsequently in [112] for circle endomorphisms, see [30, Sections IV.1 and IV.2] for a discussion of the connection between cross-ratio and Schwarzian derivative. In particular, the cross-ratio distortion arguments (i) suggest the relevance of the Poincaré metric on $(\mathbb{C} \setminus \mathbb{R}) \cup J$, which is the complex analogue of the cross-ratio on a real interval J. Indeed, let $D_r(J)$ be the set of points consisting of the set of points with distance to J of at most r with respect to the Poincaré metric on $(\mathbb{C} \setminus \mathbb{R}) \cup J$. This set is often called a Poincaré disc, and is bounded by two arcs of the circles through a, b. Using the Schwartz inclusion lemma, it then follows that if f is (for example) a real polynomial so that $f: J' \to J$ is a diffeomorphism and so that all critical values of f lie in $\mathbb{R} \setminus J$, then the component of $f^{-1}(D_r(J))$ intersecting J' is contained in $D_r(J')$. This turned out to be a key ingredient to the proof of his theorem on complex bounds for renormalisable maps, see Theorem 30.

Naturally, Dennis did ask himself whether there are analogues of his no wandering domain theorem in the higher dimensional case in the smooth category. A partial answer to this question is given by Theorem 3 for toral diffeomorphisms of Denjoy type.

3.2 Conformal measures for rational maps

Soon after developing his no wandering theorem, Dennis introduced the notion of *conformal* δ *-measure* for a rational map f. This is a Borel probability measure m so that

$$m(fA) = \int_A |Df|^\delta \, dm,$$

for every Borel measurable set $A \subset \mathbb{C}$ and where it is assumed that $\delta \ge 0$, see [104]. Dennis then showed that one can also construct conformal measures on the Julia set analogous to what he had done before in the setting of Kleinian groups, extending earlier work by Patterson:

Theorem 17 (Existence of conformal measures for rational maps, see [104]) For every rational map there exists a conformal measure. In the hyperbolic case, the exponent δ is positive and is equal to the Hausdorff dimension of the Julia set of the rational map.

The paper [104] is also shows that for Dennis the theory of dynamical systems is unified: topological, smooth and ergodic aspects are all connected. Moreover, in his view the theory of real and complex one-dimensional systems together with the theory of Kleinian groups all should be viewed as highly interwoven.

3.3 The λ -Lemma

One of Dennis' most used and cited papers on holomorphic dynamics is one in which he, and his coauthors Mañé and Sad, proved that most maps are *stable*. The main technical tool in that paper is the celebrated:

Theorem 18 (λ -Lemma, see [70]) Let A be a subset of $\overline{\mathbb{C}}$, \mathbb{D} the open unit disc and $i_{\lambda} : A \to \overline{\mathbb{C}}$ a family of maps so that

1. for each $z \in A$, $\mathbb{D} \ni \lambda \mapsto i_{\lambda}(z)$ is analytic; 2. $A \ni z \mapsto i_{\lambda}(z)$ is injective for each $\lambda \in \mathbb{D}$; 3. $i_0 = id$.

Then every $i_{\lambda} : A \to \overline{\mathbb{C}}$ has a quasiconformal extension to a continuous map $i_{\lambda} : \overline{A} \to \overline{\mathbb{C}}$, which for fixed λ is a topological embedding, and so that $\mathbb{D} \ni \lambda \mapsto i_{\lambda}(z)$ is analytic for each fixed $z \in \overline{A}$.

The proof of the λ -Lemma is surprisingly simple, and is based on the Schwarz's lemma which states that any analytic map $\xi : \mathbb{D} \to \mathbb{C} \setminus \{0, 1, \infty\}$ is contracting w.r.t. the Poincaré metric on these sets. Now consider the cross-ratio distortion of any distinct four points in *A*. Using that the cross-ratio distortion omits the values $0, 1, \infty$ and this version of the Schwarz lemma, one obtains the above λ -Lemma.

In addition to the MRMT and the theory of quasiconformal homoeomorphisms, the λ -Lemma has become one of the most widely used tools in the field of holomorphic dynamical systems.⁵

Later, jointly with Thurston, Dennis improved this λ -Lemma to show that one can extend $i_{\lambda} : A \to \overline{\mathbb{C}}$ to a qc map $i_{\lambda} : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$, provided we restrict λ to a suitable disc $\mathbb{D}_0 \subseteq \mathbb{D}$ (here the choice of \mathbb{D}_0 is universal), see [111].

3.4 Density of stable maps

The initial motivation for the λ -Lemma, and the main purpose of the paper [70], was to prove that stable maps are dense (within the space of rational maps). As a first step towards proving this, the class of *J*-stable maps is considered. Here *f* is called *J*-stable if for each *g* near *f* there exists a homeomorphism *h* of J(f) to J(g) so that $h \circ f = g \circ h$ on J(f) and so that J(g) depends continuously on *g* in the sense of Hausdorff distance between closed sets. *f* is called *structurally stable* if the conjugacy holds on $\overline{\mathbb{C}}$.

Consider a family $f_w(z)$ or rational maps depending on $w \in W$, where W is a connected complex submanifold of \mathbb{C}^{2d+1} , and so that $(w,z) \to f_w(z)$ is analytic in w, z. Let $H(f) \subset W$ be the set of $w \in W$ for which there exists a neigbourhood V with $w \in V \subset W$ so that each periodic point p_w of f_w depends analytically on $w \in V$ and so that their multiplier satisfies either $\lambda(w) \neq 1$ for all $w \in V$ or $\lambda(w) \equiv constant$ for all $w \in V$.

Theorem 19 (*J*-stability, see [70]) H(f) is open and dense in W. Moreover, f_w is *J*-stable if and only if $w \in W$ and the conjugating homeomorphism h_w can be taken to be analytic in w and quasiconformal in z.

Analogous to the set H(f), the authors introduce the set $C(f) \subset W$ of points w for which there is a neighbourhood V with $w \in V \subset W$ so that each critical point $c_i(w)$ of f_w depends analytically on $w \in V$ and so that any critical relation relation $f_w^n(c_i(w)) = f_w^m(c_j(w))$ holds either for all w in V or for none.

Theorem 20 (Structurally stable maps are dense, see [70]) C(f) *is an open and dense subset of* H(f)*. Moreover, if* $w \in C(f)$ *then* f *is structurally stable.*

In the late 1960's Smale suggested that a similar result should hold for general smooth dynamical systems, but this turned out to be false (due to examples by Newhouse and others).

⁵ Independently, Lyubich proved an analogous result, see [69].

3.5 Towards the Fatou conjecture: absence of line fields

A map *f* is said to be *Axiom A* (or *hyperbolic*) if there exist $\rho > 1$ and C > 0 so that $|(f^k)'(z)| > C\rho^k$ for all $k \ge 0$ and $z \in J(f)$. It is not hard to see that *f* is Axiom A if and only if the periodic components of F(f) are superattractive or attractive and if the orbit of every critical point of *f* is eventually contained in one of these components.

Fatou already stated the following:

Conjecture 21 (Fatou Conjecture) Each rational map can be approximated by an Axiom A rational map of the same degree.

No doubt Dennis tried to prove this conjecture but to this day nobody has succeeded in doing so. One of the main appealing properties of Axiom A maps is that they are stable, provided they satisfy some mild additional conditions, and that their dynamics is very well-understood. For example, for such a map Lebesque almost every initial point converges under iterates to a periodic attractor.

Amongst many other results in [81], Dennis, together with McMullen, shows that the above conjecture can be reduced to proving absence of *measurable invariant line fields* supported on Julia sets. Here a measurable line field on a forward invariant subset $K \subset J(f)$ of positive Lebesgue measure is a measurable function $z \mapsto \mu(z)$ on K. Here one can think of $\mu(z)$ as a line through z, and invariance means that $\mu(f(z)) = Df_z\mu(z)$.

Theorem 22 (A conditional proof of the Fatou conjecture, see [81]) Assume that any rational map which supports a measurable invariant line field on its Julia set is a Lattès map. Then the above Fatou Conjecture holds.

3.6 Monotonicity of entropy and the pullback argument

Another problem which was extensively studied in the early 1980's was whether the topological entropy of the family $f_a: [0,1] \rightarrow [0,1]$, $a \in [0,4]$ defined by $f_a(x) = ax(1-x)$ is a monotone function in a. This problem was solved by several people independently and using different methods. For a history of this problem see [62]. Dennis' approach was particularly important because it became a key ingredient in the proof of density of hyperbolicity within interval maps, see Theorem 31 below.

Monotonicity follows immediately from the following:

Theorem 23 (Monotonicity of entropy) *Consider the family* $f_a: [0,1] \rightarrow [0,1], a \in [0,4]$ *defined by* $f_a(x) = ax(1-x)$. *Then no periodic orbit disappears as a increases. If* $f_a, f_{a'}$ *are two such maps which are topologically conjugate and whose critical points are eventually periodic, then* a = a'.

This theorem is non-trivial. Indeed, it is not known whether within the family $x \mapsto x^d + c$ with d > 1 fixed but not necessarily an integer, bifurcations are monotone

in *c*. Partial results, and monotonicity for other non-trivial families of intervals maps, are given in [62]. One also has monotonicity within families of real polynomials of higher degree, namely each set of parameter for which the topological entropy is constant is connected. For the case of real cubic critical polynomials, see [83] and for the general case of real polynomials with all critical points real, see [11] and also [57].

The rigidity statement in the second part of Theorem 23 follows immediately from Thurston's famous theorem, see [36]. The approach proposed by Dennis uses the *pullback argument*. This argument is formalised in the following theorem and also applies to the setting of polynomial-like maps discussed below. Let P(f) be the closure of the forward iterates of critical points of f.

Theorem 24 (Pullback argument) Let h_0 be a quasiconformal homeomorphism so that

- 1. $h_0(P(f_a)) = P(f_{a'}),$
- 2. there exists a qc map h_1 for which $f_{a'} \circ h_1 = h_0 \circ f_a$ so that $h_1 = h_0$ on $P(f_a)$ and which is homotopic to h_0 rel. $P(f_a)$ and
- 3. h_0 is a conformal conjugacy between f_a and $f_{a'}$ near ∞ (and near their periodic attractor if they exist).

Then there exists a qc homeomorphism h so that $f_{a'} \circ h = h \circ f_a$. If h is conformal near all periodic attractors (if they exist) and if f_a does not carry an invariant line field on $J(f_a)$ then h is conformal.

Proof By the Homotopy Lifting Theorem, there exists a sequence of homeomorphism h_n so that $f_{a'} \circ h_{n+1} = h_n \circ f_a$ which are homotopic to h_0 rel. $P(f_a)$. Since f_a and $f_{a'}$ are conformal, h_{n+1} will have the same qc dilatation as h_n . Moreover, h_{n+1} agrees with h_n on a set F_n so that $F_{n+1} \supset F_n$ so that $\cup F_n$ is dense in \overline{C} . Since the space of qc maps is compact, h_n converges to a qc map h. If the Julia set of f_a has zero Lebesgue measure (or, even more generally, does not carry invariant line fields) then h is conformal.

One can deduce Theorem 23 quite easily from the pullback argument, using the *open-closed argument*. Indeed, assume by contradiction that the conclusion of Theorem 23 is wrong. Then there exists two topologically conjugate post-critically finite real quadratic maps f_a and $f_{a'}$ with $a \neq a'$. Choose [a, a'] 'maximal' i.e., so that there exists no real $a'' \notin [a, a']$ for which $f_{a''}$ is a real quadratic map which again is topologically conjugate to f_a . The pullback argument implies that f_a and $f_{a'}$ are in fact quasiconformally conjugate. Let h be a qc-conjugacy so that $f_{a'} \circ h = h \circ f_a$ and let μ be its Beltrami coefficient. Then the MRMT gives a (normalised) family of qc maps h_t whose Beltrami coefficient is $t\mu$ for $|t| \leq 1 + \varepsilon$, provided $\varepsilon > 0$ is small. A simple calculation then shows that each of the maps $g_t := h_t \circ f_a \circ h_t^{-1}$ is again conformal. This is because h_t sends the ellipse field determined by $t\mu$ to a field of circles, and the invariance of the Beltrami coefficient implies that this ellipse field is preserved by Df. Moreover, g_t depends analytically on t. That g_t is in fact is quadratic follows from the degree of the map (and a suitable normalisation of h_t). It follows that there

exists $\varepsilon > 0$ so that for $s \in (a - \varepsilon, a' + \varepsilon)$ each f_s is quasiconformally conjugate to f_a . But this contradicts the maximality of the choice of [a, a'].

This proof is very interesting because it makes it possible to reduce density of hyperbolicity with real quadratic maps to quasisymmetric rigidity in the unicritical setting, see Section 3.11. In fact, even if in the context of map with several critical points, quasisymmetric rigidity can be used to derive density of hyperbolicity, see [60, 58, 91].

Another reason that makes Dennis' proof of Theorem 23 so interesting is that it also applies to the following setting, introduced by Douady and Hubbard [34].

Definition 25 Assume that U and V are simply connected domains in \mathbb{C} . Then a holomorphic map $F: U \to V$ is called *quadratic-like* if the closure of U is contained in V and if there exists a unique critical point c of F such that F restricted to $U \setminus \{c\}$ is a covering map of degree two onto $V \setminus \{F(c)\}$. The subset

$$K(F) = \{ z \in U : F^n(z) \in U \text{ for all } n \ge 0 \}$$

is called the *filled Julia set* of F.

Any quadratic map is quadratic-like (just take U to be some very large disc). Moreover, by the so-called Straightening Theorem, see [34], any quadratic-like map is quasiconformally conjugate to a quadratic map.

One step in the renormalisation theory developed by Dennis (and also in subsequent developments) is to show that certain iterates of a given map have quadraticlike restrictions $f^n: U_n \to V_n$ with the additional property that the modulus $mod(V_n \setminus U_n)$ is bounded from below uniformly in *n*. Such bounds are called *a*-priori bounds or complex bounds.

3.7 Renormalisation theory for interval maps

Consider the family $f_a(x) = ax(1-x)$. A simple computer simulation shows that this family of maps undergoes a period doubling bifurcation from period 2^n to period 2^{n+1} at some parameter a_n . That these parameters a_n are in fact unique (and increasing) can be deduced from a result similar to Theorem 23.

One of the reasons that iterations of interval maps attracted so much attention from the late 1970's was the observation by Feigenbaum and independently by Coullet and Tresser of metric *universality* within a wide class of such families. Namely, it turns out that the parameters a_n converge to some limit value a_{∞} at a particular rate $\delta > 1$:

$$\frac{a_{n-1} - a_{n-2}}{a_n - a_{n-1}} \to \delta = 4.669201...$$

Remarkably, if one takes some other family such as $f_a(x) = a \sin(\pi x)$ then the corresponding rate is the same! Moreover, the map $f_{a_{\infty}}$ has an invariant Cantor set and the scaling structure of this Cantor set also displays metric universality.

Feigenbaum, Coullet and Tresser already suggested a mechanism which would be responsible for this universality. The key idea is renormalisation. Indeed, there exists a (real) parameter interval $[u_1, v_1] \ni a_{\infty}$ so that for each $a \in [u_1, v_1]$ there exists an interval $J_a^1 \ni 1/2$ so that $f_a^2(J_a^1) \subset J_a^1$ and so that $f_a(J_a^1)$ and J_a^1 have disjoint interiors. Such maps are called 2-*renormalisable*. Note that 1/2 is the critical point of f_a . It turns out that there exists an interval $[u_2, v_2] \subset [u_1, v_1]$ so that for each $a \in [u_2, v_2]$ the map $f_a^2|J_a^1$ (rescaled) is again 2-renormalisable. In other words, there exists an interval J_a^2 with $1/2 \in J_a^2 \subset J_a^1$ so that $f_a^4(J_a^2) \subset J_a^2$ and so that $J_a^2, \ldots, f_a^3(J_a^2)$ have disjoint interiors. Moreover, $J_a^2, f_a^2(J_a^2) \subset J_a^1$ and $f(J_a^2), f_a^3(J_a^2) \subset f(J_a^1)$. So these maps are twice 2-renormalisable. Continuing like this, for each k there exists an interval $[u_k, v_k]$ so that for each $a \in [u_k, v_k]$ the map f_a is 2^k -renormalisable. For $a_{\infty} \in \cap[u_k, v_k]$ the set

$$\Lambda_{a_{\infty}} := \bigcap_{k \ge 0} \left(J_a^j \cup \dots \cup f_a^{2^k - 1}(J_a^k) \right)$$

is a Cantor set.

To formalise this one can define, near the limit map $f_{a_{\infty}}$, the renormalisation operator

$$R(f) = f^2 | J$$
 rescaled

where *J* is the maximal interval of renormalisation of period two, i.e. the maximal interval so that $f^2(J) \subset J$ and so that f(J) and *J* have disjoint interiors. This operator is well-defined for all maps which are at least once 2-renormalisable.

More generally one has the following

Definition 26 A interval map *f* is *renormalisable* if there exist p > 0 and an interval *J* around a critical point of *f* so that $J, \ldots, f^{p-1}(J)$ have disjoint interiors and so that $f^p(J) \subset J$. The operator $R(f) = f^p | J$ rescaled is then called the renormalisation operator. See figure 2.

If *f* is unimodal, then $f^p|J$ is again a unimodal map and it makes sense to require that *p* is minimal and *J* is maximal with the above properties. If R(f) is again renormalisable, then we say that *f* is twice renormalisable. Similarly, *f* is called *infinitely renormalisable* if this process can be repeated infinitely often. If the corresponding integers p_1, p_2, \ldots are all bounded by some number $P < \infty$ then *f* is called infinitely renormalisable of bounded type.

It turns out that the renormalisation conjectures of Feigenbaum and Coullet & Tresser of metric universality follow from (i) the existence of a fixed point ψ of the operator R, (ii) that the spectrum of the operator $DR\psi$ lies off the unit circle and (iii) that $DR\psi$ has a unique expanding eigenvalue. The universal parameter scaling constant δ is equal to this expanding eigenvalue. The universal dynamical scaling structure of $\Lambda_{a_{\infty}}$ follows from the largest contracting eigenvalue of DR.

Edson de Faria and Sebastian van Strien




Fig. 2 Renormalising a unimodal map. Here, *J* is the red interval, p = 4, and $\Re(f)$ is simply R(f) rescaled by the affine map that takes the blue interval onto the red interval.

The universality from the Feigenbaum-Coullet-Tresser conjectures then follows from the fact that for any family which crosses the stable manifold of *R* transversally, the parameter scalings δ and the dynamical scaling can be obtained from the spectrum of $DR(\psi)$.

The existence of the fixed point ψ , and an analysis of the spectrum of the linear map *DR* were established by Lanford before Dennis started working on the renormalisation conjectures. This was done in part using careful rigorous computer estimates. However, there were three limitations to Lanford's results.

Firstly, Lanford's proof did not establish which maps are contained in the stable manifold of *R*. In other words, what remained unclear whether any 2^{∞} -infinitely renormalisable unimodal maps (with a quadratic critical point) would be in the stable manifold of of the period doubling operator. Secondly, Lanford's proof did not establish a conceptual proof of why this result was true. Finally, Lanford's proof also did not cover a more general situation of maps which are infinitely renormalisable of bounded type, but only of constant type $p_1 = p_2 = \dots$

The huge result which Dennis managed to obtain is the following:

Theorem 27 (Renormalisation for unimodal interval maps, see [109] and also [30]) There exists a Cantor set \mathbb{K}_p of infinitely renormalisable maps of bounded type $\leq p$, of real analytic unimodal maps, which form an invariant subset for the corresponding renormalisation operator. The renormalisation operator acts on \mathbb{K}_p as a full shift on finitely many symbols. Each real analytic unimodal map with a

quadratic critical point which is infinitely renormalisable map and of bounded type is in the stable manifold of the corresponding renormalisation operator.

After this, other proofs of the renormalisation conjectures appeared. McMullen [79, 80] and Avila and Lyubich [5] gave easier proofs that the stable manifold of the renormalisation operator contains all the relevant infinitely renormalisable maps using 'towers' respectively based on a Schwarz Lemma. These proofs additionally give that under renormalisation infinitely renormalisable maps of bounded type converge with an exponential rate to the above Cantor set of infinitely renormalisable maps of bounded type. Moreover, Lyubich [66] gave a conceptual proof showing that the renormalisation operator has a unique expanding eigenvalue.

The above proofs require that the maps are real analytic. In the C^r context, these theorems also go through, see [27]. An alternative approach to extend the renormalisation theory for real analytic maps to smooth maps is via asymptotically holomorphic maps, see [17].

In the multimodal setting, the renormalisation picture is not complete yet. In this setting, the conjecture could be

- Topologically conjugate mappings converge exponentially quickly under renormalisation.
- 2. The stable manifolds of renormalisation are smooth.
- 3. The transverse directions to the stable manifolds are exponentially expanded by renormalisation.

Part 2 of this conjecture has been proved in [20]. For the case of bounded combinatorics, see [97].

3.8 Real and complex bounds

The first step towards proving Theorem 27 is to show that the space of renormalisable maps is compact. To do this, Dennis established apriori bounds, first in the real and then in the complex setting:

Theorem 28 (Real Bounds) Let f be a real analytic map which is infinitely renormalisable and of bounded type. Then the C^2 norm of the maps $\mathbb{R}^n f$ is uniformly bounded.

The first part of this theorem shows that $R^n f$ is a composition of a quadratic map and a map g whose non-linearity is bounded from above. The main ingredient he used for this is the (real) Koebe Principle and the *smallest interval argument* discussed above. Indeed, let J be the first renormalisation interval and that $f^p(J) \subset J$ with $p \ge 1$ minimal. Then the intervals $J, \ldots, f^{p-1}(J)$ are disjoint and one among them, let us say $f^i(J)$, is the smallest. This means that, unless $f^i(J)$ is one of the two extreme intervals in this collection, the interval $f^i(J)$ is contained in an interval $T = [f^l(J), f^r(J)]$ which has the property that both components of $T \setminus f^i(J)$ are not small compared to $f^i(J)$. This simple idea can be used to obtain *Koebe space*, namely that the map $f^{p-1}: f(J) \to f^p(J) \subset J$ extends to a diffeomorphism with range $T' \supset J$ so that J is well-inside J. Using the real Koebe Principle the map $f^{p-1}: f(J) \to f^p(J)$ then has bounded non-linearity.

Extending this argument, and using the pullback argument from above, one then obtains one of the key steps:

Theorem 29 (Quasisymmetric Rigidity in the Renormalisable Case) Let f,g be two infinitely renormalisable real-analytic maps of bounded type and with quadratic critical points. Then if f,g are topologically conjugate they are in fact quasisymmetrically conjugate.

To obtain Theorem 27 Dennis needed to extend the real maps $f^p: J \rightarrow J$ to quadratic-like maps and to obtain compactness with the space of such maps:

Theorem 30 (Complex Bounds) Let f be an infinitely renormalisable real-analytic unimodal map of bounded type $p_i \leq p$ for all $i \geq 0$ and with quadratic critical point. Then, for every n sufficiently large, $R^n f$ extends to a quadratic-like map $F: U \rightarrow V$ so that the modulus of $V \setminus U$ is bounded by some number $\rho > 0$ which does not depend on f but only the upper bound p.

Moreover, any limit of the sequence $\{R^n f\}$ has a complex analytic extension which is in the so-called Epstein class.

Real bounds were proved in a much more general context, see (in increasing generality) [74, 93, 113]. Complex bounds for real unicritical maps were proved in [63, 68] and for multicritical maps in (in increasing generality) in [96, 21]. Complex bounds do not hold for general (non-real) quadratic maps: there are examples of infinitely renormalisable quadratic maps for which no modulus bounds as in the above theorem hold. On the other hand, for non-renormalisable polynomials maps (with only hyperbolic periodic points) one does have complex bounds, see [60] and [18]. An important ingredient in the latter developments is the quasi-additive lemma by Kahn and Lyubich [54]. This lemma was also used to treat some maps which are infinitely renormalisable, see [52, 53]. It is not known how to extend complex bounds to the case of general rational maps, as in general it is not clear how to construct an initial puzzle partition.

3.9 Riemann surface laminations and the non-coiling lemma

To complete the proof of Theorem 27, Dennis introduced a new tool, namely his *non-coiling principle* and his *almost geodesic principle*. To explain this, consider a qc conjugacy *H* between F_0 and F_1 . Its Beltrami coefficient $\mu_H = \bar{\partial}H/\partial H$ its invariant under F_0 . It follows that the family of qc maps H_t associated to $\mu_t = t\mu_H$ (coming from the MRMT) defines a family of quadratic like maps $F_t = H_t \circ F_0 \circ H_t^{-1}$ connecting F_0 to F_1 . This is called a *Beltrami path* between F_0 and F_1 . Dennis' *almost geodesic principle* shows that the Beltrami path corresponding to an *almost*

extremal vector does not coil: if the tangent Beltrami vector is almost extremal then the Beltrami path remains almost a geodesic for a long (but a priori fixed) time.

To show that the renormalisation operator is (weakly) contracting he then argued as follows. From the complex bounds discussed in the previous subsection, he obtained that there exists a compact space \mathcal{K} , so that if we take an arc connecting two conjugate maps which are infinitely renormalisable and of bounded type, then after *n* renormalisations this arc is mapped in \mathcal{K} . Here *n* depends on the choice of the chosen maps, but \mathcal{K} does not. Now take an almost geodesic path between two maps F_0, F_1 . Extend this path to a geodesic path between two maps \tilde{F}_0, \tilde{F}_1 which are extremely far apart. Now apply renormalisation. For *n* sufficiently large, the renormalisations $R^n(\tilde{F}_0), R^n(\tilde{F}_1)$ are in \mathcal{K} and so not far apart. But then, by the *almost geodesic principle*, the renormalisations $R^n(F_0), R^n(F_1)$ are extremely close.

To make all this work, Dennis had to consider germs of quadratic-like maps. For this reason he considered inverse limits of the quadratic-like maps $F_i: U_i \to V_i$ which led to the study of *Riemann surface laminations*. This is not the place to go into a full description of the beautiful theory of such objects, but let us at least say a word or two. Roughly speaking, a Riemann surface lamination (RSL) is a space akin to a foliated space in which the chart domains are homeomorphic to $D \times T$, where $D \subset \mathbb{C}$, is a disk and the transversal $T \subset \mathbb{R}$ is typically a Cantor set (or an interval), and the chart transitions are holomorphic along the horizontal leaves. In the present context, the main example is the following. Let $F: U \to V$ be a quadratic-like map, let $K(F) \subseteq \mathbb{C}$ be its filled-in Julia set, which we assume to contain the critical point of F (so that it is connected), set $W = V \setminus K(F)$, and consider the inverse system of holomorphic covering maps:

$$\cdots \to F^{-n-1}W \to F^{-n}W \to \cdots \to F^{-1}W \to W$$
.

The inverse limit space $\mathscr{L}(F)$ of this system is a fibration over W, the fiber above each $x \in W$ being a Cantor set (the binary Cantor set at the end of the tree giving the full backward orbit of x). From this it follows that $\mathscr{L}(F)$ is an RSL in a natural way.

The inverse limit map $F_{\infty} : \mathscr{L}(F) \to \mathscr{L}(F)$ is invertible and acts properly discontinuously on $\mathscr{L}(F)$, and the quotient $X_F = \mathscr{L}(F)/F_{\infty}$ is a *compact* RSL. In [109], Dennis defined a deformation space or *Teichmüller space* of X_F (and more general RSLs) in such a way that every Beltrami path between two quadratic-like maps F_0 and F_1 as above can be lifted to a Beltrami path between the corresponding laminations X_{F_0} and X_{F_1} , and all deformations are encoded in this fashion. What makes this possible is the fact that the Julia set of an infinitely renormalisable quadraticlike map with complex bounds does not carry any non-trivial quasi-conformal deformations. Dennis then proved the non-coiling principle and the almost-geodesic principle at the level of laminations, transporting the resulting contraction of the Teichmüller distance downstairs, at the level of maps.

We will not go further into Dennis proof of this tour de force (a full description can be found in [30, Chapter VI], in addition to the papers [109] and [110]). After all, as mentioned, this last step was improved in subsequent proofs which show

that the invariant Cantor set of *R* attracts other maps with an *exponential rate*, see McMullen [79, 80] and Avila and Lyubich [5]. Still, the reader who wants to learn more about RSLs should consult the elegant survey written by Ghys in [15].

3.10 Renormalisation theory for circle maps

While working on the renormalisation problem for unimodal maps, Dennis was well aware that similar experimental discoveries to those made by Feigenbaum and Coullet-Tresser had been made by physicists concerning circle homeomorphisms having a single non-flat critical point (of power-law type). The topological classification of such maps had been accomplished by Yoccoz in [118].

Dennis also knew that, in the circle context, Lanford had formulated a renormalisation conjecture akin to the one for unimodal maps, using the language of commuting pairs. Dennis then suggested to EdF, as a thesis problem, to adapt his holomorphic ideas to the case of such critical circle maps. The key step was to find an analogue of quadratic-like maps in the context of critical circle maps. This was accomplished in [23] (see also [24]) with the notion of *holomorphic commuting pair* (inspired in part by a computer picture drawn by H. Epstein), alongside a proof of complex bounds (as well as a pull-back argument) for such objects, assuming the rotation number of the underlying critical circle map to be of bounded type, and also that the circle maps belonged to a special class of maps known as *Epstein class*. The necessary real bounds had already been established by Herman (unpublished manuscript, but see [49]) and Swiatek [112]. The bounded type assumption was removed by Yampolsky [116], still assuming the Epstein property. The latter was finally removed by EdF and Welington de Melo in [26].

For unicritical circle maps, the fact that, under suitable full-measure conditions on the rotation number, exponential convergence of renormalisations leads to $C^{1+\alpha}$ rigidity was established in [25] (counterexamples to this ansatz for rotation numbers in a special zero-measure class were constructed in the same paper). The analogous conditional statement obtained replacing $C^{1+\alpha}$ by C^1 holds under no restriction on the rotation number (other than being irrational), as shown by [55]. The exponential convergence of renormalisations for *real-analytic* unicritical circle maps with bounded type rotation number was proved in [26]. Using the concept of parabolic renormalisation, Yampolsky [117] was able to remove the bounded type hypothesis, and in fact proved that the renormalisation operator attractor is globally hyperbolic in the analytic context. In the larger space of C^3 unicritical circle maps, exponential convergence towards the attractor was first proved by Guarino in his thesis under de Melo – see [46] – assuming rotation numbers of bounded type only. The bounded type hypothesis was later removed in [47], at the cost of assuming the maps to be C^4 .

In recent years, considerable work has been done to extend these rigidity, universality and renormalisation convergence results to multicritical circle maps – see for instance [39] and references therein. An important step towards this goal is to first

establish the *quasisymmetric rigidity* of such maps – this was accomplished in [38] for multicritical circle maps with critical points having arbitrary (real) power-law criticalities. For this and much more about multicritical circle maps, see [28].

3.11 The Fatou conjecture in the real setting

As mentioned, Fatou conjectured that each rational map can be approximated by an Axiom A rational map of the same degree. This question is still wide open, even in the quadratic case. However, in the setting of real maps the corresponding result has been answered completely:

Theorem 31 (The Real Fatou Conjecture) *Each real polynomial can can be approximated by a real Axiom A polynomial of the same degree.*

In the setting of real quadratic polynomials, the real Fatou conjecture was proved independently by Lyubich [65] and Graczyk & Swiátek [44, 45]. The setting of real polynomials of higher degree d > 2 was solved (using entirely different tools) by Kozloski, Shen and van Strien, see [59] and [58]. In the non-real non-renormalisable case see also [60] and [18]. An important ingredient in the latter developments is the quasi-additive lemma by Kahn and Lyubich [54]. This lemma was also used to treat some maps which are infinitely renormalisable, see [52, 53].

Although the proofs of these results are not due to Dennis, he played an important role in them. Indeed, his work suggested that to prove density of hyperbolicity that it would be enough to prove the following

Theorem 32 (Quasisymmetric Rigidity in one-dimensions) If f,g are topologically conjugate real polynomials with only real critical points, and all their critical points are quadratic, then these maps are quasisymmetrically conjugate.

Dennis' renormalisation theory for infinitely renormalisable unimodal (unicritical) maps of bounded type, relied on this result (in this setting). In the quadratic case, the above theorem is due to [65] and Graczyk & Swiátek [44, 45]. Their proof relied on the property that, in this setting, the moduli of certain annuli tends to infinity. This growth of moduli is a deep and subtle result, but this does not hold for unimodal maps with a degenerate critical point nor for mulimodal maps with nondegenerate critical points. So for the general case a different approach is needed. The approach by Kozlosvki, Shen and van Strien in [59] uses the enhanced nest, which is a particular choice of a sequence of puzzle pieces that turn out to have Koebe space. An introductory survey on this technique can be found in [18]. The most general quasisymmetric result is contained in joint work of Clark and van Strien, see [19].

Kleinian Groups	Iterated Analytic Maps
Kleinian group Γ	Holomorphic map f
Γ finitely generated	f rational map
Γ is Fuchsian	f is a Blaschke product
Domain of discontinuity $\Omega(\Gamma)$	Fatou set $\mathscr{F}(f)$
$\begin{array}{c} \text{Limit set } \Lambda(\Gamma) \\ \Lambda(\Gamma) \neq \emptyset \end{array}$	Julia set $J(f)$ $J(f) \neq \emptyset$
$\Omega(\Gamma)$ has either 0, 1, 2 or infinitely many components	$\mathscr{F}(f)$ has either 0, 1, 2 or infinitely many components
Either $\Lambda(\Gamma) = \widehat{\mathbb{C}}$ or $\Lambda(\Gamma)$ has empty interior	Either $J(f) = \widehat{\mathbb{C}}$ or $J(f)$ has empty interior
Ahlfors finiteness theorem	Sullivan's no-wandering-domains theorem
Bers area theorem	Shishikura's bound on the number of non-repelling periodic cycles
Mostow's rigidity theorem	Thurston's uniqueness theorem on post-critically finite rational maps
Patterson-Sullivan measures on $\Lambda(\Gamma)$	Sullivan's conformal measures on $J(f)$
The quotient manifold \mathbb{H}/Γ	Lyubich-Minsky lamination
Geometrically finite groups with no cusps are dense	Are hyperbolic rational maps dense?

 Table 1
 Some entries in Sullivan's dictionary

3.12 Sullivan's quasisymmetry rigidity programme

Even though there is no analogue of the MRMT in the real setting, one of the insights of Dennis was that quasisymmetric rigidity should still be a very powerful tool in addressing questions about the topological structure of conjugacy classes of interval maps. For example, whether such a conjugacy class is a connected manifold. This insight turned out to be justified. Indeed, let $\mathscr{A}^{\underline{v}}$ be the space of real analytic maps with precisely v real critical points $c_1 < \cdots < c_v$ of order ℓ_1, \ldots, ℓ_v . The following theorem was shown by Clark and van Strien [20].

Theorem 33 Let $f \in \mathscr{A}^{\underline{v}}$. Then the space \mathscr{T}_f of real analytic maps in $\mathscr{A}^{\underline{v}}$ which are topologically conjugate to f forms an analytic manifold. This manifold is connected and simply connected.

This theorem extends results of Avila-Lyubich-de Melo [6] for the quasi-quadratic unimodal case and of Clark [16] for the more general unimodal case. Their methods fail in the case where there are several critical points. For this reason, the notion of *pruned Julia set* is introduced in [20]. This set is a version of the Julia set (but pruned) but depends on where one 'prunes'. A pruned Julia set can be defined for each real analytic map f. The real analytic map f, together with its pruned Julia set, define a real analytic *external map* of the circle *with discontinuities*. Using this external map, one can construct a *a pruned polynomial-like* complex extension of the real analytic map. Finally, from all this one is able to show that topological conjugacy classes are connected (something which was not even known in the general unimodal setting). Even more, this space is contractible and forms an analytic manifold.

4 Sullivan's dictionary

Dennis' wide-range view of Mathematics allows him to draw fruitful analogies between different theories, leading to several conjectures on either side. A case in point is what is now known as the *Sullivan dictionary* between the theory of Kleinian groups (in dimension n = 3) on one side and the theory of iterated holomorphic maps on the other side. A sample of entries in this dictionary is shown in Table 1. Note that the last entry in the table has a question mark: that is none other than the famous Fatou Conjecture, widely regarded as the main classical open problem about the dynamics of rational maps.

Not all meaningful analogies, however, deserve to be in the dictionary. For instance, a famous conjecture by Ahlfors in the 1960's stated that the limit set of a finitely generated Kleinian group is either the entire sphere or else has zero Lebesgue measure. As we mentioned in the beginning of section 2, this is now a theorem, thanks to the combined efforts of several mathematicians. The final piece of the puzzle was laid down by Canary [14], building primarily on previous works by Thurston and Bonahon – see for instance [72] for a description of the whole story, and references therein. The corresponding statement for iterated holomorphic maps – to wit, that the Julia set of a rational map is either the entire sphere or else has zero Lebesgue measure – was thought for a long time to be true, until X. Buff and A. Chéritat [13] found an example of a quadratic polynomial whose Julia set has positive measure.

As Dennis himself has explained to us, he never put the Ahlfors conjecture in the dictionary because – after working on the problem for about 13 months in the late seventies and exhausting all available ergodic arguments that would have solved it he came to the conclusion that no one would be able to prove it using what was currently known about finitely generated Kleinian groups. After proving his finiteness theorem, what Ahlfors really wanted to know was under which conditions the limit set of a Kleinian group could support non-trivial quasiconformal deformations. He asked the question about the Lebesgue measure of the limit set in the finitely generated case because, if the measure indeed turned out to be zero in that case, there would be no such deformations. Thus, Dennis realized that the real question was not the measure zero question, but rather to describe, if any, the quasiconformal deformations on the limit set. He was able to prove a very general result that states that, given a Kleinian group Γ and any Γ -invariant subset $E \subset \Lambda_{\Gamma}$ of its limit set, there are quasiconformal deformations of Γ supported in E if and only if there are *positive measure wandering sets* inside E, and when this happens, the space of such nontrivial deformations is infinite dimensional. In particular, since for finitely generated groups the space of deformations is a-priori known to be finite-dimensional, that are no quasiconformal deformations supported on Λ_{Γ} when Γ is finitely generated. This holds even if the limit set happens to be the whole sphere. This absence of invariant line fields supported in the limit set is stated in Dennis' Theorem 9. After more than 40 years, the corresponding statement for rational maps⁶ remains an open problem. And it is a fundamental problem: indeed it is possible to prove that if the statement for rational maps is true, then so is the Fatou conjecture. Thus, the question of absence of invariant line fields certainly deserves its place as an entry in the Sullivan dictionary (albeit being conspicuously absent from Table 1).

Over the years, the Sullivan dictionary has continued to inspire new results. A recent example is provided by the work of Hee Oh. Working on the Kleinian side of the dictionary, in collaboration with Margulis and Mohammadi [73], she examined closed geodesics and holonomies for hyperbolic 3-manifolds. She was then asked by Dennis himself about an analogue of her results for rational maps. This resulted in her paper [87] in collaboration with Winter, in which they establish estimates on the number of primitive periodic orbits of a hyperbolic rational map.

⁶ To wit, that a rational map is either a Lattès example or else carries no invariant line fields in its Julia set.

5 Final words

Dennis Sullivan's major contributions to the field of Dynamical Systems, some of which we attempted to describe here, constitute but one facet of his extraordinary work as a mathematician. There are several other facets. Thus, for his fundamental work in Topology, especially regarding the study of geometric and/or algebraic structures on manifolds, see the article by Shmuel Weinberger in the present volume. In more recent years, Dennis has essentially founded, in collaboration with M. Chas, the sub-field of Topology now known as String Topology. We have heard it said elsewhere that *Dennis Sullivan is a mathematician who has re-invented himself several times*, and that seems to us a very accurate statement.

We have not included anything about Dennis' recent work on fluid dynamics. Nor have we mentioned any work on dynamics that Dennis co-wrote with some of his students and/or post-docs, such as the work with Jiang and Morita [51], his work with Hu [50] or his work with Pinto [89], nor with many other collaborators from the dynamical systems community.

In closing, it is important to add that Dennis has always been extremely generous when sharing his ideas with other researchers, as well as in guiding young mathematicians. According to the Math Genealogy Project, he has had so far 40 students, and a total of 155 descendants. Among his students, those who have written a thesis in dynamics under his supervision include Andre de Carvalho, Adam Epstein, Jun Hu, Yunping Jiang, Curt McMullen, Waldemar Paluba, Guiai Peng, Meiyu Su, as well as one of us (EdF). But many more mathematicians, young and old, although not formally his students, have been directly influenced by him. Through his insightful lectures, and his inquisitive quest not merely for results, but for *understanding* Mathematics, Dennis has inspired and will continue to inspire us all.

Acknowledgements We would like to thank Curt McMullen, Leon Staresinic, Edson Vargas for their useful comments, and especially Dennis Sullivan for explaining to us the origins of his dictionary.

References

- 1. Ahlfors, L. V. (1981). *Möbius transformations in several dimensions*. Ordway Professorship Lectures in Mathematics. University of Minnesota, School of Mathematics, Minneapolis, Minn.
- Alves, J. F., Pinheiro, V., and Pinto, A. A. (2014). Explosion of smoothness for conjugacies between multimodal maps. J. Lond. Math. Soc. (2), 89(1):255–274.
- Astorg, M., Buff, X., Dujardin, R., Peters, H., and Raissy, J. (2016). A two-dimensional polynomial mapping with a wandering Fatou component. *Ann. of Math.* (2), 184(1):263–313.
- 4. Avila, A., Fayad, B., Le Calvez, P., Xu, D., and Zhang, Z. (2020). On mixing diffeomorphisms of the disc. *Invent. Math.*, 220(3):673–714.
- Avila, A. and Lyubich, M. (2011). The full renormalization horseshoe for unimodal maps of higher degree: exponential contraction along hybrid classes. *Publ. Math. Inst. Hautes Études Sci.*, (114):171–223.

 Avila, A., Lyubich, M., and de Melo, W. (2003). Regular or stochastic dynamics in real analytic families of unimodal maps. *Invent. Math.*, 154(3):451–550.

- 8. Beardon, A. F. (1983). *The geometry of discrete groups*, volume 91 of *Graduate Texts in Mathematics*. Springer-Verlag, New York.
- Bowen, R. (1979). Hausdorff dimension of quasicircles. Inst. Hautes Études Sci. Publ. Math., (50):11–25.
- 10. Bowen, R. and Franks, J. (1976). The periodic points of maps of the disk and the interval. *Topology*, 15(4):337–342.
- 11. Bruin, H. and van Strien, S. (2015). Monotonicity of entropy for real multimodal maps. J. Amer. Math. Soc., 28(1):1–61.
- 12. Buff, X. (2018). Wandering Fatou component for polynomials. *Ann. Fac. Sci. Toulouse Math.* (6), 27(2):445–475.
- 13. Buff, X. and Chéritat, A. (2012). Quadratic Julia sets with positive area. Ann. of Math. (2), 176(2):673–746.
- 14. Canary, R. D. (1993). Ends of hyperbolic 3-manifolds. J. Amer. Math. Soc., 6(1):1-35.
- 15. Cerveau, D., Ghys, E., Sibony, N., and Yoccoz, J.-C. (2003). Complex dynamics and geometry, volume 10 of SMF/AMS Texts and Monographs. American Mathematical Society, Providence, RI; Société Mathématique de France, Paris. With the collaboration of Marguerite Flexor, Papers from the Meeting "State of the Art of the Research of the Société Mathématique de France" held at the École Normale Supérieure de Lyon, Lyon, January 1997, Translated from the French by Leslie Kay.
- 16. Clark, T. (2014). Regular or stochastic dynamics in families of higher-degree unimodal maps. *Ergodic Theory Dynam. Systems*, 34(5):1538–1566.
- Clark, T., de Faria, E., and Van Strien, S. (2022a). Asymptotically holomorphic methods for infinitely renormalizable c^r unimodal maps. *Ergodic Theory and Dynamical Systems*, page 1–49.
- Clark, T., Drach, K., Kozlovski, O., and van Strien, S. (2022b). The dynamics of complex box mappings. Arnold Math. J., 8(2):319–410.
- 19. Clark, T. and van Strien, S. (2018). Quasisymmetric rigidity in one-dimensional dynamics.
- Clark, T. and van Strien, S. (2023). Conjugacy classes of real analytic one-dimensional maps are analytic connected manifolds.
- 21. Clark, T., van Strien, S., and Trejo, S. (2017). Complex bounds for real maps. *Comm. Math. Phys.*, 355(3):1001–1119.
- De Carvalho, A., Lyubich, M., and Martens, M. (2005). Renormalization in the Hénon family. I. Universality but non-rigidity. J. Stat. Phys., 121(5-6):611–669.
- 23. de Faria, E. (1992). *Proof of universality for critical circle mappings*. ProQuest LLC, Ann Arbor, MI. Thesis (Ph.D.)–City University of New York.
- 24. de Faria, E. (1999). Asymptotic rigidity of scaling ratios for critical circle mappings. *Ergodic Theory Dynam. Systems*, 19(4):995–1035.
- 25. de Faria, E. and de Melo, W. (1999). Rigidity of critical circle mappings. I. J. Eur. Math. Soc. (JEMS), 1(4):339–392.
- 26. de Faria, E. and de Melo, W. (2000). Rigidity of critical circle mappings. II. J. Amer. Math. Soc., 13(2):343–370.
- 27. de Faria, E., de Melo, W., and Pinto, A. (2006). Global hyperbolicity of renormalization for *C^r* unimodal mappings. *Ann. of Math.* (2), 164(3):731–824.
- 28. de Faria, E. and Guarino, P. (2022). *Dynamics of circle mappings*. 33 ^o Colóquio Brasileiro de Matemática. Instituto Nacional de Matemática Pura e Aplicada (IMPA), Rio de Janeiro.
- 29. de Melo, W. and van Strien, S. (1989). A structure theorem in one-dimensional dynamics. *Ann. of Math.* (2), 129(3):519–546.
- de Melo, W. and van Strien, S. (1993). One-dimensional dynamics, volume 25 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin.

36

^{7.} Beardon, A. F. (1968). The exponent of convergence of Poincaré series. *Proc. London Math. Soc.* (3), 18:461–483.

- Deroin, B., Kleptsyn, V., and Navas, A. (2007). Sur la dynamique unidimensionnelle en régularité intermédiaire. Acta Math., 199(2):199–262.
- 32. Douady, A. and Hubbard, J. H. (1982). Itération des polynômes quadratiques complexes. *C. R. Acad. Sci. Paris Sér. I Math.*, 294(3):123–126.
- 33. Douady, A. and Hubbard, J. H. (1984). Étude dynamique des polynômes complexes. Partie I, volume 84 of Publications Mathématiques d'Orsay [Mathematical Publications of Orsay]. Université de Paris-Sud, Département de Mathématiques, Orsay.
- 34. Douady, A. and Hubbard, J. H. (1985a). Étude dynamique des polynômes complexes. Partie II, volume 85 of Publications Mathématiques d'Orsay [Mathematical Publications of Orsay]. Université de Paris-Sud, Département de Mathématiques, Orsay. With the collaboration of P. Lavaurs, Tan Lei and P. Sentenac.
- 35. Douady, A. and Hubbard, J. H. (1985b). On the dynamics of polynomial-like mappings. *Ann. Sci. École Norm. Sup.* (4), 18(2):287–343.
- Douady, A. and Hubbard, J. H. (1993). A proof of Thurston's topological characterization of rational functions. *Acta Math.*, 171(2):263–297.
- Erëmenko, A. E. and Lyubich, M. Y. (1992). Dynamical properties of some classes of entire functions. *Ann. Inst. Fourier (Grenoble)*, 42(4):989–1020.
- Estevez, G. and de Faria, E. (2018). Real bounds and quasisymmetric rigidity of multicritical circle maps. *Trans. Amer. Math. Soc.*, 370(8):5583–5616.
- Estevez, G., Smania, D., and Yampolsky, M. (2022). Renormalization of analytic multicritical circle maps with bounded type rotation numbers. *Bull. Braz. Math. Soc.* (N.S.), 53(3):1053–1071.
- Franks, J. and Young, L. S. (1981). A C² Kupka-Smale diffeomorphism of the disk with no sources or sinks. In *Dynamical systems and turbulence, Warwick 1980 (Coventry, 1979/1980)*, volume 898 of *Lecture Notes in Math.*, pages 90–98. Springer, Berlin-New York.
- 41. Gambaudo, J.-M., Sullivan, D., and Tresser, C. (1994). Infinite cascades of braids and smooth dynamical systems. *Topology*, 33(1):85–94.
- 42. Gambaudo, J.-M., van Strien, S., and Tresser, C. (1989). Hénon-like maps with strange attractors: there exist C[∞] Kupka-Smale diffeomorphisms on S² with neither sinks nor sources. *Nonlinearity*, 2(2):287–304.
- Goldberg, L. R. and Keen, L. (1986). A finiteness theorem for a dynamical class of entire functions. *Ergodic Theory Dynam. Systems*, 6(2):183–192.
- 44. Graczyk, J. and Świątek, G. (1997). Generic hyperbolicity in the logistic family. *Ann. of Math.* (2), 146(1):1–52.
- 45. Graczyk, J. and Świątek, G. (1998). *The real Fatou conjecture*, volume 144 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ.
- 46. Guarino, P. and de Melo, W. (2017). Rigidity of smooth critical circle maps. J. Eur. Math. Soc. (JEMS), 19(6):1729–1783.
- 47. Guarino, P., Martens, M., and de Melo, W. (2018). Rigidity of critical circle maps. *Duke Math. J.*, 167(11):2125–2188.
- Herman, M.-R. (1979). Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Inst. Hautes Études Sci. Publ. Math.*, (49):5–233.
- 49. Herman, M. R. (2018). Notes inachevées—sélectionnées par Jean-Christophe Yoccoz, volume 16 of Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Société Mathématique de France, Paris.
- Hu, J. and Sullivan, D. P. (1997). Topological conjugacy of circle diffeomorphisms. *Ergodic Theory Dynam. Systems*, 17(1):173–186.
- Jiang, Y. P., Morita, T., and Sullivan, D. (1992). Expanding direction of the period doubling operator. *Comm. Math. Phys.*, 144(3):509–520.
- 52. Kahn, J. and Lyubich, M. (2008). A priori bounds for some infinitely renormalizable quadratics. II. Decorations. Ann. Sci. Éc. Norm. Supér. (4), 41(1):57–84.
- Kahn, J. and Lyubich, M. (2009a). A priori bounds for some infinitely renormalizable quadratics. III. Molecules. In *Complex dynamics*, pages 229–254. A K Peters, Wellesley, MA.
- 54. Kahn, J. and Lyubich, M. (2009b). The quasi-additivity law in conformal geometry. Ann. of Math. (2), 169(2):561–593.

- 55. Khanin, K. and Teplinsky, A. (2007). Robust rigidity for circle diffeomorphisms with singularities. *Invent. Math.*, 169(1):193–218.
- Kiriki, S., Nakano, Y., and Soma, T. (2017). Non-trivial wandering domains for heterodimensional cycles. *Nonlinearity*, 30(8):3255–3270.
- 57. Kozlovski, O. (2019). On the structure of isentropes of real polynomials. J. Lond. Math. Soc. (2), 100(1):159–182.
- Kozlovski, O., Shen, W., and van Strien, S. (2007a). Density of hyperbolicity in dimension one. Ann. of Math. (2), 166(1):145–182.
- 59. Kozlovski, O., Shen, W., and van Strien, S. (2007b). Rigidity for real polynomials. Ann. of Math. (2), 165(3):749-841.
- Kozlovski, O. and van Strien, S. (2009). Local connectivity and quasi-conformal rigidity of non-renormalizable polynomials. *Proc. Lond. Math. Soc.* (3), 99(2):275–296.
- Kozlovski, O. and van Strien, S. (2020). Asymmetric unimodal maps with non-universal period-doubling scaling laws. *Comm. Math. Phys.*, 379(1):103–143.
- Levin, G., Shen, W., and van Strien, S. (2020). Positive transversality via transfer operators and holomorphic motions with applications to monotonicity for interval maps. *Nonlinearity*, 33(8):3970–4012.
- 63. Levin, G. and van Strien, S. (1998). Local connectivity of the Julia set of real polynomials. *Ann. of Math.* (2), 147(3):471–541.
- 64. Li, S. and Shen, W. (2006). Smooth conjugacy between S-unimodal maps. *Nonlinearity*, 19(7):1629–1634.
- 65. Lyubich, M. (1997). Dynamics of quadratic polynomials. I, II. Acta Math., 178(2):185–247, 247–297.
- Lyubich, M. (1999). Feigenbaum-Coullet-Tresser universality and Milnor's hairiness conjecture. Ann. of Math. (2), 149(2):319–420.
- 67. Lyubich, M. and Martens, M. (2011). Renormalization in the Hénon family, II: the heteroclinic web. *Invent. Math.*, 186(1):115–189.
- 68. Lyubich, M. and Yampolsky, M. (1997). Dynamics of quadratic polynomials: complex bounds for real maps. *Ann. Inst. Fourier (Grenoble)*, 47(4):1219–1255.
- 69. Lyubich, M. Y. (1983). Some typical properties of the dynamics of rational mappings. *Uspekhi Mat. Nauk*, 38(5(233)):197–198.
- Mañé, R., Sad, P., and Sullivan, D. (1983). On the dynamics of rational maps. Ann. Sci. École Norm. Sup. (4), 16(2):193–217.
- 71. Marden, A. (1974). The geometry of finitely generated kleinian groups. Ann. of Math. (2), 99:383-462.
- 72. Marden, A. (2007). *Outer circles*. Cambridge University Press, Cambridge. An introduction to hyperbolic 3-manifolds.
- Margulis, G., Mohammadi, A., and Oh, H. (2014). Closed geodesics and holonomies for Kleinian manifolds. *Geom. Funct. Anal.*, 24(5):1608–1636.
- 74. Martens, M. (1994). Distortion results and invariant Cantor sets of unimodal maps. *Ergodic Theory Dynam. Systems*, 14(2):331–349.
- 75. Martens, M. and de Melo, W. (1999). The multipliers of periodic points in one-dimensional dynamics. *Nonlinearity*, 12(2):217–227.
- Martens, M., de Melo, W., and van Strien, S. (1992). Julia-Fatou-Sullivan theory for real one-dimensional dynamics. *Acta Math.*, 168(3-4):273–318.
- 77. Maskit, B. (1988). Kleinian groups, volume 287 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin.
- 78. McMullen, C. (2020). Riemann surfaces, dynamics and geometry. Harvard University.
- 79. McMullen, C. T. (1994). *Complex dynamics and renormalization*, volume 135 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ.
- 80. McMullen, C. T. (1996). *Renormalization and 3-manifolds which fiber over the circle*, volume 142 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ.
- McMullen, C. T. and Sullivan, D. P. (1998). Quasiconformal homeomorphisms and dynamics. III. The Teichmüller space of a holomorphic dynamical system. *Adv. Math.*, 135(2):351–395.

38

Merenkov, S. (2019). No round wandering domains for C¹-diffeomorphisms of tori. *Ergodic Theory Dynam. Systems*, 39(11):3127–3135.

Milnor, J. and Tresser, C. (2000). On entropy and monotonicity for real cubic maps. *Comm. Math. Phys.*, 209(1):123–178. With an appendix by Adrien Douady and Pierrette Sentenac.

- Mostow, G. D. (1968). Quasi-conformal mappings in n-space and the rigidity of hyperbolic space forms. *Inst. Hautes Études Sci. Publ. Math.*, (34):53–104.
- 85. Nicholls, P. J. (1989). The ergodic theory of discrete groups, volume 143 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge.
- Norton, A. and Sullivan, D. (1996). Wandering domains and invariant conformal structures for mappings of the 2-torus. *Ann. Acad. Sci. Fenn. Math.*, 21(1):51–68.
- 87. Oh, H. and Winter, D. (2017). Prime number theorems and holonomies for hyperbolic rational maps. *Invent. Math.*, 208(2):401–440.
- 88. Patterson, S. J. (1976). The limit set of a Fuchsian group. Acta Math., 136(3-4):241-273.
- 89. Pinto, A. A. and Sullivan, D. (2006). The circle and the solenoid. *Discrete Contin. Dyn. Syst.*, 16(2):463–504.
- 90. Prasad, G. (1973). Strong rigidity of Q-rank 1 lattices. Invent. Math., 21:255-286.
- Rempe-Gillen, L. and van Strien, S. (2015). Density of hyperbolicity for classes of real transcendental entire functions and circle maps. *Duke Math. J.*, 164(6):1079–1137.
- 92. Ruelle, D. and Sullivan, D. (1975). Currents, flows and diffeomorphisms. *Topology*, 14(4):319–327.
- 93. Shen, W. (2003). Bounds for one-dimensional maps without inflection critical points. *J. Math. Sci. Univ. Tokyo*, 10(1):41–88.
- Shub, M. and Sullivan, D. (1985). Expanding endomorphisms of the circle revisited. *Ergodic Theory Dynam. Systems*, 5(2):285–289.
- 95. Singer, D. (1978). Stable orbits and bifurcation of maps of the interval. *SIAM J. Appl. Math.*, 35(2):260–267.
- 96. Smania, D. (2001). Complex bounds for multimodal maps: bounded combinatorics. *Nonlinearity*, 14(5):1311–1330.
- 97. Smania, D. (2020). Solenoidal attractors with bounded combinatorics are shy. *Ann. of Math.* (2), 191(1):1–79.
- 98. Sullivan, D. (1976a). A counterexample to the periodic orbit conjecture. *Inst. Hautes Études Sci. Publ. Math.*, (46):5–14.
- Sullivan, D. (1976b). Cycles for the dynamical study of foliated manifolds and complex manifolds. *Invent. Math.*, 36:225–255.
- 100. Sullivan, D. (1976c). A new flow. Bull. Amer. Math. Soc., 82(2):331-332.
- 101. Sullivan, D. (1979). The density at infinity of a discrete group of hyperbolic motions. *Inst. Hautes Études Sci. Publ. Math.*, (50):171–202.
- 102. Sullivan, D. (1981). On the ergodic theory at infinity of an arbitrary discrete group of hyperbolic motions. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, volume 97 of *Ann. of Math. Stud.*, pages 465–496. Princeton Univ. Press, Princeton, N.J.
- 103. Sullivan, D. (1982). Disjoint spheres, approximation by imaginary quadratic numbers, and the logarithm law for geodesics. *Acta Math.*, 149(3-4):215–237.
- 104. Sullivan, D. (1983). Conformal dynamical systems. In *Geometric dynamics (Rio de Janeiro, 1981)*, volume 1007 of *Lecture Notes in Math.*, pages 725–752. Springer, Berlin.
- 105. Sullivan, D. (1984). Entropy, Hausdorff measures old and new, and limit sets of geometrically finite Kleinian groups. *Acta Math.*, 153(3-4):259–277.
- 106. Sullivan, D. (1985). Quasiconformal homeomorphisms and dynamics. I. Solution of the Fatou-Julia problem on wandering domains. *Ann. of Math.* (2), 122(3):401–418.
- 107. Sullivan, D. (1988). Differentiable structures on fractal-like sets, determined by intrinsic scaling functions on dual Cantor sets. In *The mathematical heritage of Hermann Weyl (Durham, NC, 1987)*, volume 48 of *Proc. Sympos. Pure Math.*, pages 15–23. Amer. Math. Soc., Providence, RI.

- Sullivan, D. (1991). Renormalization, Zygmund smoothness and the Epstein class. In *Chaos, order, and patterns (Lake Como, 1990)*, volume 280 of *NATO Adv. Sci. Inst. Ser. B: Phys.*, pages 25–34. Plenum, New York.
- Sullivan, D. (1992). Bounds, quadratic differentials, and renormalization conjectures. In American Mathematical Society centennial publications, Vol. II (Providence, RI, 1988), pages 417–466. Amer. Math. Soc., Providence, RI.
- 110. Sullivan, D. (1993). Linking the universalities of Milnor-Thurston, Feigenbaum and Ahlfors-Bers. In *Topological methods in modern mathematics (Stony Brook, NY, 1991)*, pages 543–564. Publish or Perish, Houston, TX.
- 111. Sullivan, D. P. and Thurston, W. P. (1986). Extending holomorphic motions. *Acta Math.*, 157(3-4):243–257.
- 112. Świątek, G. (1988). Rational rotation numbers for maps of the circle. *Comm. Math. Phys.*, 119(1):109–128.
- 113. van Strien, S. and Vargas, E. (2004). Real bounds, ergodicity and negative Schwarzian for multimodal maps. J. Amer. Math. Soc., 17(4):749–782.
- 114. van Strien, S. J. (1981). On the bifurcations creating horseshoes. In *Dynamical systems and turbulence, Warwick 1980 (Coventry, 1979/1980)*, volume 898 of *Lecture Notes in Math.*, pages 316–351. Springer, Berlin-New York.
- 115. Wang, J. and Yang, H. (2021). A question of Norton-Sullivan in the analytic case. *Int. Math. Res. Not. IMRN*, (22):17201–17219.
- 116. Yampolsky, M. (1999). Complex bounds for renormalization of critical circle maps. *Ergodic Theory Dynam. Systems*, 19(1):227–257.
- 117. Yampolsky, M. (2002). Hyperbolicity of renormalization of critical circle maps. *Publ. Math. Inst. Hautes Études Sci.*, (96):1–41 (2003).
- 118. Yoccoz, J.-C. (1984). Il n'y a pas de contre-exemple de Denjoy analytique. C. R. Acad. Sci. Paris Sér. I Math., 298(7):141–144.