

Centrally Essential Factor Rings and Subdirect Indecomposability

Oleg Lyubimtsev¹, Askar Tuganbaev²

Abstract. Let R be a ring and let $J(R)$, $C(R)$ be its Jacobson radical and center, correspondingly. If R is a centrally essential ring and the factor ring $R/J(R)$ is commutative, then any minimal right ideal is contained in the center $C(R)$. A right Artinian (or right Noetherian subdirectly indecomposable) centrally essential ring is a right and left Artinian local ring. We describe centrally essential Noetherian subdirectly indecomposable rings and centrally essential rings with subdirectly indecomposable center. We give examples of non-commutative subdirectly indecomposable, centrally essential rings.

Key words. centrally essential ring, subdirectly indecomposable ring.

The work of Oleg Lyubimtsev is supported by Ministry of Education and Science of the Russian Federation, project FSWR-2023-0034. The study of Askar Tuganbaev is supported by grant of Russian Science Foundation (=RSF), project 22-11-00052, <https://rscf.ru/en/project/22-11-00052>.

MSC2020 database 16D25, 16R99

1 Introduction

We consider only associative unital non-zero rings. A ring R is said to be centrally essential if either R is commutative or for every non-central element a of R , there exist non-zero central elements x, y with $ax = y$. It is clear that a ring R with center $C(R)$ is centrally essential if and only if the module $R_{C(R)}$ is an essential extension of the module $C(R)_{C(R)}$. In addition, any commutative ring is centrally essential.

Remark 1.1. There exists a non-commutative ring R such that all factor rings of R are centrally essential (see Example 3.3).

Remark 1.2. In [7, Proposition 2.8], it is proved that a right uniserial, right Artinian, centrally essential ring is left Artinian.

¹Nizhny Novgorod State University, Nizhny Novgorod, Russia; email: oleg_lyubimcev@mail.ru .

²National Research University MPEI, Moscow, Russia; Lomonosov Moscow State University, Moscow, Russia; tuganbaev@gmail.com .

Remark 1.3. There exist right and left subdirectly indecomposable rings R which are not centrally essential. For example, let V be a vector space over a field \mathbb{F} of characteristic 0 or $p \neq 2$ with basis $\{e_1, e_2\}$, and let $\Lambda(V)$ be the exterior algebra of the space V , i.e., $\Lambda(V)$ is an algebra with operation \wedge , generators e_1, e_2 and defining relations $e_i \wedge e_j + e_j \wedge e_i = 0$ for all $i, j = 1, 2$. Then $R = \Lambda(V)$ is a non-commutative 4-dimensional \mathbb{F} -algebra with basis $\{1, e_1, e_2, e_1 \wedge e_2\}$ and the center $C(R)$ of R is $\{q_0 \cdot 1 + q_1 \cdot e_1 \wedge e_2 \mid q_i \in \mathbb{F}\}$. It is easy to see that R is a right and left subdirectly indecomposable ring with subdirectly indecomposable center. In addition, the ring R is not centrally essential; see [5, Theorem 1].

In connection to Remarks 1.1, 1.2 and 1.3, we prove Theorems 1.4, 1.5 and 1.6 which are the main results of this paper.

Theorem 1.4. Let R be a centrally essential ring.

1. If the factor ring $R/J(R)$ is commutative, then any minimal right ideal S of the ring R is contained in the center $C(R)$. In particular, all minimal right ideals of the ring R are ideals and its right socle is contained in the center of the ring R .
2. If R is a right Artinian ring (or right Noetherian subdirectly indecomposable) ring, then R is a right and left Artinian local ring.
3. If the center of the ring R is subdirectly indecomposable, then the ring R is right and left subdirectly indecomposable.

Theorem 1.5. If R is a right Noetherian ring such that all factor rings of R are centrally essential, then R is a subdirect product of local, right and left Artinian rings.

Theorem 1.6. Let R be a right Noetherian ring with center $C(R) = C$. We set $\overline{C} = J(R) \setminus J(C)$.

1. If the ring R is subdirectly indecomposable, then the ring R is centrally essential if and only if $\text{Ann}_{\overline{C}} J(C) = \text{Ann}_{\overline{C}} J(R) = 0$.
2. If the center C of the ring R is subdirectly indecomposable with core H , then the ring R is centrally essential if and only if $rC \cap H \neq 0$ for any non-zero element $r \in R$.

We give some necessary definitions and notation.

For a ring R , the Jacobson radical of R is denoted by $J(R)$. A module is said to be **subdirectly indecomposable** if it has the least non-zero submodule. A ring R is said to be **right subdirectly indecomposable** (or *RSI-ring* [2]) if the intersection H_r of all its non-zero right ideals is non-zero. The **left core**

H_ℓ of the ring R is defined similarly. A ring R is said to be **subdirectly indecomposable** with core H if the intersection H of all its non-zero ideals is non-zero. A ring R is said to be **centrally subdirectly indecomposable** if the center of R is subdirectly indecomposable.

For a module M , the socle $Soc M$ is the sum of all simple submodules in M ; if M does not contain simple submodules, then $Soc M = 0$, by definition. A module M is said to be **uniserial** if any two submodules of M are comparable with respect to inclusion.

For a ring R , the Lie ring $G(R)$ of R consists of elements of the ring R with multiplication $[x, y] = xy - yx$ for any $x, y \in R$. An element $r \in R$ is said to be **right regular** (resp., **left regular**) if $rx \neq 0$ (resp., $xr \neq 0$) for every non-zero element $x \in R$. Right and left regular elements are called **regular**. A non-zero element $r \in R$ is $C(R)$ -torsion free if $rc \neq 0$ for every non-zero element $c \in C(R)$. If S and T are two subsets of the ring R , then $Ann_S T = \{s \in S \mid sT = 0\}$. If \mathbb{F} is a field and V is a vector \mathbb{F} -space with basis e_1, \dots, e_n , then the exterior algebra $\Lambda(V)$ is an algebra with operation \wedge , generators e_1, \dots, e_n , defining relations $e_i \wedge e_j + e_j \wedge e_i = 0$ for all $i, j = 1, \dots, n$, and the basis

$$\{1, e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_k} \mid k = 1, 2, \dots, n; 1 \leq i_1 < i_2 < \dots < i_k \leq n\}.$$

2 Proof of Main Results

Proposition 2.1. For a centrally essential ring R , the following conditions are equivalent.

- 1) The ring R is right subdirectly indecomposable.
- 2) The ring R is left subdirectly indecomposable.
- 3) The ring R is subdirectly indecomposable.

In addition, the cores H_r , H_ℓ and H coincide in each of cases **1)–3)**.

Proof. Let the ring R be right subdirectly indecomposable and let $\{J_t\}_{t \in T}$ be the set of all non-zero left ideals of the ring R . Since the ring R is centrally essential, each of the left ideals J_t contains a non-zero central element c_t . Then $0 \neq \bigcap_{t \in T} c_t R \subseteq \bigcap_{t \in T} J_t$. Therefore, **1)** implies **2)**.

It is clear that **2)** implies **3)**. Since each of the ideals $c_t R$ is two-sided, it follows from **3)** that **1)** is true.

It is clear that $H_r = hR$ for any $0 \neq h \in H_r$. Since R is a centrally essential

ring, we have that $0 \neq hc = d$ for some central elements c and d . Then $H_r = dR$ is a two-sided ideal. Therefore, $H_r = H$. The relation $H_\ell = H$ can be proved similarly. \square

We remark that the analogue of Proposition 2.1 is not true for rings which are not centrally essential. For example, the ring of all $n \times n$ ($n \geq 2$) matrices over a division ring is subdirectly indecomposable but it is not right or left subdirectly indecomposable. In addition, there exists a right subdirectly indecomposable ring which is not left subdirectly indecomposable; see [2, Example 4.5].

It follows from Proposition 2.1 that it is sufficient to restrict ourself to centrally essential, subdirectly indecomposable rings in the study of right (or left) subdirectly indecomposable, centrally essential rings.

Proposition 2.2. If R is a centrally essential ring and the factor ring $R/J(R)$ is commutative, then any minimal right ideal S of the ring R is contained in $C(R)$. In particular, all minimal right ideals of the ring R are ideals and $\text{Soc } R_R$ is contained in the center of the ring R .

Proof. The simple R -module S_R is a cyclic module with non-zero generator $s \in S$. We set $K = S \cap C(R)$. By assumption, there exist non-zero central elements $x, y \in C(R)$ such that $0 \neq sx = y \in S$. Then $S = yR$ and $y \in K \neq 0$. Since $J(R)$ is the intersection of the annihilators of all simple right R -modules, we have that $SJ(R) = 0$. It follows from commutativity of the ring $R/J(R)$ that $ab - ba \in J(R)$ for any elements $a, b \in R$. In addition, $k(ab - ba) = 0$ for any $k \in K$. Since $k \in C(R)$, we have that $(ka)b = kba = b(ka)$. Therefore, $ka \in C(R)$. Further, it follows from $ka \in S$ that K is a right ideal of the ring R . Since S is a minimal right ideal, we have $S = K \subseteq C(R)$. \square

Proposition 2.3. Let R be a centrally essential ring.

1. An element $r \in R$ is regular if and only if r is $C(R)$ -torsion free.
- 2, [7, Lemma 2.2 (2.2.1)]. In the ring R , every one-sided zero-divisor is a two-sided zero-divisor.

Proof. 1. It is clear that if r is regular, then it is $C(R)$ -torsion free. Conversely, let $rs = 0$ for some non-zero element $s \in R$. Then $s \notin C(R)$ and there exist $c, d \in C(R)$ such that $0 \neq sc = d$. Since r is $C(R)$ -torsion free, we have that $rd \neq 0$ and $rsc \neq 0$. This is a contradiction.

2. If $rs = 0$ for some $0 \neq s \in R$, then 1 implies that $rd = dr = 0$ for some $0 \neq d \in C(R)$. \square

Proposition 2.4. Let R be a right Noetherian, centrally essential, subdi-

rectly indecomposable ring with core H .

1. R is a right Artinian local ring.
2. The factor ring $R/J(R)$ is commutative.
3. The core H is contained in the center $C(R)$.

Proof. 1. It follows from Proposition 2.3(2) that any right regular element is left regular. Then the assertion follows from [2, Theorem 2.1].

2. The assertion follows from 1 and [6, Proposition 2.4].

3. By 2, the ring $R/J(R)$ is commutative. Now we use Proposition 2.2. \square

Proposition 2.5. Let R be a centrally essential ring.

1. If the ring R is right Artinian, then R is left Artinian.
2. If R is a right Noetherian, subdirectly indecomposable ring, then R is a right and left Artinian ring.
3. If the ring R is local and $rJ(C) = 0$ for $r \in R$, then $r \in J(C)$.
4. If the center of the ring R is subdirectly indecomposable, then R is right and left subdirectly indecomposable.

Proof. 1. It follows from [6, Proposition 2.4] that the ring $R/J(R)$ is commutative. Then $[x, y] \in J(R)$ for any $x, y \in G(R)$. In addition, the radical $J(R)$ is nilpotent with some nilpotence index n . Then

$$[x_1, y_1] \cdot \dots \cdot [x_n, y_n] = 0$$

for all elements $[x_i, y_i] \in G(R)$, $i = 1, \dots, n$. In this case, the Lie ring $G(R)$ is nilpotent. Then the result follows from [1, Theorem 2.2].

2. It follows from Proposition 2.4(1) that the ring R is right Artinian. By 1, the ring R is left Artinian.

3. We set $C = C(R)$. We assume that $rJ(C) = 0$. Since R is a centrally essential ring, we have $0 \neq rc = d \in J(C)$. It follows from the relation $J(C) = C \cap J(R)$ that the element c is invertible. Then $r = c^{-1}d \in J(C)$.

4. Let $M = \{I_k\}_{k \in K}$ be the set of all non-zero right ideals of the ring R . Since R is a centrally essential ring, each of the ideals I_k contains a non-zero central element c_k . We consider ideals $c_k C(R)$ of the center $C(R)$. Since the center is subdirectly indecomposable, we have $0 \neq \bigcap_{k \in K} c_k C(R) \subseteq \bigcap_{k \in K} I_k$. \square

2.6. The proof of Theorem 1.4.

1. The assertion follows from Proposition 2.2.
2. The assertion follows from Proposition 2.4(1) and Proposition 2.5(1,2).

3. The assertion follows from Proposition 2.5(4). \square

2.7. The proof of Theorem 1.5.

Any ring is a subdirect product of subdirectly indecomposable rings. Therefore, the right Noetherian ring R is a subdirect product of right Noetherian, subdirectly indecomposable, centrally essential rings R_i , $i \in I$. It follows from Propositions 2.4(1) and 2.5(2) that all these rings are right and left Artinian local rings. \square

2.8. The proof of Theorem 1.6. We recall that R is a right Noetherian ring with center C and $\overline{C} = J(R) \setminus J(C)$.

1. If R is a centrally essential ring, then it follows from Propositions 2.4(1) and 2.5(3) that $\text{Ann}_{\overline{C}}J(C) = 0$. Then $\text{Ann}_{\overline{C}}J(R) = 0$.

Conversely, let $0 \neq r \notin C$ and $rJ(C) \neq 0$. By Proposition 2.4(1), the ring R is local; therefore, $r \in \overline{C}$ or r is invertible. Then R is right Artinian and we can choose central elements c_1, \dots, c_{k-1} such that $rc_1 \dots c_{k-1} \neq 0$, but $rc_1 \dots c_{k-1}c = 0$ for all $c \in J(C)$. We set $c' = c_1 \dots c_{k-1}$. If $0 \neq rc' \notin J(C)$, then $rc' \in \text{Ann}_{\overline{C}}J(C) = 0$. This is a contradiction. Consequently, rc' is a non-zero central element and R is a centrally essential ring.

2. Let R be a centrally essential ring and $r \notin H$. There exist central elements c and d such that $0 \neq rc = d$. If $d \in H$, then it is nothing to prove. Let $d \notin H$. Since R is a centrally subdirectly indecomposable ring, we have $dC \cap H \neq 0$, i.e., $0 \neq dc' = h$ for some $c' \in C$ and $h \in H$. Then

$$0 \neq dc' = (rc)c' = r(cc') = rc'' \in rC,$$

whence we obtain that $0 \neq rc'' \in rC \cap H$.

Conversely, for any $0 \neq r \in R$, there exists an element $c \in C$ such that $0 \neq rc = h \in H \subseteq C$, i.e., R is a centrally essential ring. \square

3 Examples and Remarks

A ring is said to be **invariant** if all its one-sided ideals are two-sided ideals.

Example 3.1. We give an example of an Artinian, centrally essential, right and left subdirectly indecomposable, invariant ring R such that R is not a centrally subdirectly indecomposable ring and all proper centrally essential factor rings of R are commutative. However,, there exists a proper factor ring of the ring R which is not subdirectly indecomposable or centrally essential. A ring R is a non-commutative 8-dimensional algebra over an arbitrary field

\mathbb{F} of characteristic 0 or $p \neq 2$. If we take $\mathbb{F} = \mathbb{Z}/p\mathbb{Z}$ for prime integer $p > 2$, then the ring R is finite.

Proof. Let $R = \Lambda(V)$ be the exterior algebra with generators e_1, e_2, e_3 . Then R is an 8-dimensional \mathbb{F} -algebra with basis

$$\{1, e_1, e_2, e_3, e_1 \wedge e_2, e_2 \wedge e_3, e_1 \wedge e_3, e_1 \wedge e_2 \wedge e_3\}$$

and R is a non-commutative centrally essential ring (see details in [5, Example 1]) with center

$$C(R) = \{q_0 \cdot 1 + q_1 \cdot e_1 \wedge e_2 + q_2 \cdot e_2 \wedge e_3 + q_3 \cdot e_1 \wedge e_3 + q_4 \cdot e_1 \wedge e_2 \wedge e_3, q_i \in \mathbb{F}\}.$$

However, $I_1 \cap I_2 = \{0\}$ for central ideals $I_1 = \langle e_1 \wedge e_2 \rangle_{\mathbb{F}}$ and $I_2 = \langle e_1 \wedge e_2 \wedge e_3 \rangle_{\mathbb{F}}$, i.e., the center of the ring R is not a subdirectly indecomposable ring. We remark that R is an invariant ring and the ideal $H = I_2$ is contained in the intersection of any ideals of R . Consequently, R is an Artinian, centrally essential, right and left subdirectly indecomposable ring which is not centrally subdirectly indecomposable.

We prove that all proper centrally essential factor rings of R are commutative. We consider the factor ring $\overline{R} = R/I$, where I is a non-zero ideal in R . If $e_i \notin I$ and $\overline{e}_i \notin C(\overline{R})$ for some i , then $\overline{e}_i J(C(\overline{R})) = \overline{0}$, since I contains the core H and $e_i \wedge (e_i \wedge e_j) = 0$. Then the ring \overline{R} is not centrally essential, by Proposition 2.5(3). In addition, if, for example, $I = \langle \overline{e}_1 \wedge \overline{e}_3, \overline{e}_1 \wedge \overline{e}_2 \wedge \overline{e}_3 \rangle_{\mathbb{F}}$, then the ring \overline{R} is not subdirectly indecomposable. Indeed, principal ideals generated by elements $\overline{e}_1 \wedge \overline{e}_2$ and $\overline{e}_2 \wedge \overline{e}_3$ have the zero intersection. \square

Example 3.2. Let Q_8 be the quaternion group, i.e., the group with two generators a, b and defining relations $a^4 = 1, a^2 = b^2$ and $aba^{-1} = b^{-1}$; e.g., see [3, Section 4.4]. We have

$$Q_8 = \{e, a, a^2, b, ab, a^3, a^2b, a^3b\},$$

with center $C(Q_8) = \{e, a^2\}$. We consider the group algebra $R = \mathbb{Z}_2 Q_8$ with center

$$C(R) = \{\lambda_0 + \lambda_1 a^2 + \lambda_2 (a + a^3) + \lambda_3 (b + a^3 b) + \lambda_4 (ab + a^3 b), \lambda_i \in \mathbb{Z}_2\};$$

e.g., see [9, Part 2]. Then R is a ring with subdirectly indecomposable center and core $H = \{0, \widehat{Q}_8\}$, where $\widehat{Q}_8 = \sum_{g \in Q_8} g$. In addition, R is a centrally essential ring; see, [4]. It follows from Theorem 1.6(2) that for every $0 \neq x \in R$, there exists an element $c \in C(R)$ such that $xc = \widehat{Q}_8$. Then the factor ring $\overline{R} = R/H$ with basis $\{\overline{e}, \overline{a}, \overline{a}^2, \overline{a}^3, \overline{b}, \overline{ab}, \overline{a^2 b}\}$ and with center

$$C(\overline{R}) = \{\alpha_0 \overline{e} + \alpha_1 \overline{a}^2 + \alpha_2 (\overline{a} + \overline{a}^3) + \alpha_3 (\overline{b} + \overline{a^2 b}) + \alpha_4 (\overline{e} + \overline{a} + \overline{a}^2 + \overline{a}^3 + \overline{b} + \overline{a^2 b})\}$$

is not a centrally essential ring. This property can be directly verified. For example, let $x \in R$ and $xa \neq ax$ but $\bar{x}\bar{a} = \bar{a}\bar{x}$ in \bar{R} . Then $xa = ax + \widehat{Q}_8$, whence $\sum x_i a = \sum ax_i + \widehat{Q}_8$. Since only basis elements b, ab, a^2b and a^3b do not commute with the element a , we obtain the contradictory relation

$$\alpha_1 b + \alpha_2 ab + \alpha_3 a^2 b + \alpha_4 a^3 b = \widehat{Q}_8, \quad \alpha_i \in \mathbb{Z}_2.$$

The case $xb \neq bx$ is considered similarly, but $\bar{x}\bar{b} = \bar{b}\bar{x}$. When we multiply the element $\bar{r} = (\bar{e} + \bar{a} + \bar{b} + \bar{ab}) \notin C(\bar{R})$ by central basis elements, we obtain either \bar{r} (under multiplication by \bar{a}^2) or $\bar{0}$. Consequently, \bar{R} is not a centrally essential ring. Similarly, we can verify that all centrally essential proper factor rings of the ring R are commutative.

Example 3.3. We give an example of a non-commutative ring R with subdirectly indecomposable center such that all factor rings of R are centrally essential. Let $F = \mathbb{Q}(x, y)$ be the field of rational functions. We consider two partial derivations $d_1 = \frac{\partial}{\partial x}$ and $d_2 = \frac{\partial}{\partial x}$. Then the ring $R = T(F, F)$ of matrices

$$\left\{ \left(\begin{array}{ccc} f & d_1(f) & g \\ 0 & f & d_2(f) \\ 0 & 0 & f \end{array} \right) \mid f, g \in F \right\}$$

is a centrally essential ring; see [6]. In addition, R is a ring with subdirectly indecomposable center and with core

$$H = \left\{ \left(\begin{array}{ccc} 0 & 0 & g \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right) \mid g \in F \right\}.$$

Since $R/H \cong \mathbb{Q}(x, y)$ and H is the the least ideal in R , the ring R does not have non-zero proper ideals not equal to H .

Remark 3.4. In [5, Proposition 3.3], it is proved that if R is a centrally essential local ring, then the ring $R/J(R)$ is commutative (in particular, $R/J(R)$ is centrally essential). This is not true in general case; see [8, Theorem 1.5].

Remark 3.5. If R is not centrally essential, then the assertion of Proposition 2.4(1) is not true: there exist right Noetherian, right subdirectly indecomposable rings which are not right Artinian; see [2, Example 2.2].

Remark 3.6. If R is a centrally essential, right and left subdirectly indecomposable ring with core H , then either $H^2 = 0$ or R is a field. Indeed, if $H^2 = H$, then R is a division ring; see [2, Proposition 1.3]. Since R is a centrally essential ring, it follows from [4, Proposition 3.3] that R is a field.

Open question 3.7. Is it true that there exists a centrally essential ring which has a proper non-commutative centrally essential factor ring?

Open question 3.8. Is it true that a centrally essential subdirectly indecomposable (or centrally subdirectly indecomposable) ring is invariant?

References

- [1] Bjork J.E. Conditions which imply that subrings of Artinian rings are Artinian // J. Reine Angew. Math. – 1971. – Vol. 247. – P 123–138.
- [2] Deshpande M.G. Structure of right subdirectly irreducible ring I // J. Algebra. – 1971. – Vol. 17. – P. 317–325.
- [3] M. Hall, The Theory of Groups, Macmillan, New York, 1959.
- [4] Markov V.T., Tuganbaev A.A. Centrally essential group algebras // J. Algebra. – 2018. – Vol. 512. – P. 109–118.
- [5] Markov V.T., Tuganbaev A.A. Centrally essential rings // Discrete Math. Appl. – 2019. – Vol. 29, no. 3. – P. 189–194.
- [6] Markov V.T., Tuganbaev A.A. Rings essential over their centers // Comm. Algebra. – 2019. – Vol. 47, no. 4. – P. 1642–1649.
- [7] Markov V.T., Tuganbaev A.A. Uniserial Noetherian Centrally Essential Rings // Comm. Algebra. – 2020. – Vol. 48, no. 1. – P. 149–153.
- [8] Markov, V.T., Tuganbaev, A.A. Constructions of Centrally Essential Rings // Communications in Algebra. – 2020. – Vol. 48. – № 1. – P. 198–203.
- [9] Passman D.S. The Algebraic Structure of Group Rings// John Wiley and Sons, New York, 1977.