

APPLICATIONS OF P -FUNCTIONS TO FULLY NONLINEAR ELLIPTIC EQUATIONS: GRADIENT ESTIMATES AND RIGIDITY RESULTS

DIMITRIOS GAZOULIS

ABSTRACT. We introduce the notion of P -functions for fully nonlinear equations and establish some general criterion for obtaining such quantities for this class of equations. Some applications are gradient bounds, De Giorgi-type properties of entire solutions and rigidity results. Furthermore, we prove Harnack-type inequalities and local pointwise estimates for the gradient of solutions to fully nonlinear elliptic equations. In addition, we consider such quantities for higher order nonlinear equations and for equations of order greater than two we obtain Liouville-type theorems and pointwise estimates for the Laplacian.

1. INTRODUCTION

In this work we introduce the notion of P -function for fully nonlinear partial differential equations or differential inequalities and we incorporate the “ P -function technique” in a general setting. This abstract setting allow us to obtain many applications by only determining an example of a P -function. Some of these applications are gradient bounds for entire solutions, Harnack -type inequalities for the gradient of solutions and rigidity results.

The structure of this paper is as follows. In section 2 we define the notion of P -functions and study such quantities for fully nonlinear elliptic equations. We provide two general criteria for obtaining P -functions for this class of equations. Different examples of such quantities may give various types of gradient bounds for solutions of a particular equation. For instance, in section 3, we obtain a gradient bound for entire solutions of the Allen-Cahn equation that differs from the Modica inequality and in Appendix A we provide examples of gradient bounds for quasi-linear equations that are different from the one in [4]. So, these criteria allow us to obtain many types of gradient bounds by determining a particular P -function and we provide some examples.

Additionally, in section 3, we prove an abstract pointwise estimate, i.e. Theorem 3.1, for a class of P -functions that are associated to any given fully nonlinear equation. This pointwise estimate is in fact a gradient bound for entire solutions in a wide variety of fully nonlinear elliptic equations and some examples of such gradient bounds are given. One such bound, generalizes the one in [4] for a class of quasi-linear equations. Other consequences are also illustrated in section 4 where we establish a Liouville-type theorem and a De Giorgi-type property for entire solutions. One of the most important applications, is an abstract rigidity result for entire solutions of fully nonlinear elliptic equations i.e. Theorem 4.1. This result recovers as particular cases the classical results of J. Serrin in [19].

In section 5 we establish a Harnack-type inequality and local pointwise estimates for the gradient of solutions to quasi-linear equations. These estimates can be extended to fully nonlinear elliptic equations, such as for the Monge-Ampère equation.

Moreover, in section 6, we study such quantities for nonlinear equations of order greater than two together with some applications. For example, we establish an a priori bound for the Laplacian and pointwise estimates through the mean value properties for higher order equations. Also, some Liouville-type properties can be established for nonlinear equations of order greater than two. In this setting, we believe that one can obtain many other types of bounds for any order of derivatives, assuming a $C^{k,\alpha}$ a priori estimate and that we have an appropriate P -function related to the respective equation.

We will now briefly discuss some of the most important contributions on the “ P -function technique” and its applications. Perhaps the most well-known example is $P(u, x) = \frac{1}{2}|\nabla u|^2 - W(u)$ that is related to the Allen-Cahn equation

$$\Delta u = W'(u) \quad , \quad u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R} \quad (1.1)$$

and Modica in [16] proved the well-known gradient bound

$$\frac{1}{2}|\nabla u|^2 \leq W(u) \quad (1.2)$$

for every bounded entire solution of (1.1).

Later, Caffarelli et al in [4] generalized this gradient bound for a class of variational quasi-linear equations and proved Liouville-type and De Giorgi-type properties for a particular choice of P -function related to the equation $\operatorname{div}(\Phi'(|\nabla u|^2)\nabla u) = F'(u)$. This bound was extended for anisotropic partial differential equations and other general types of equations in [5, 8, 9]. The gradient bound (1.2) also holds in unbounded domains with nonnegative mean curvature as proved in [10].

Furthermore, P -functions had been already studied by Sperb in [20], Payne and Philippin in [17, 18] who studied other types of quasilinear equations for the form $\operatorname{div}(A(u, |\nabla u|^2)\nabla u) = B(u, |\nabla u|^2)$, which are not necessarily Euler-Lagrange equations of an elliptic integrand. They derived maximum principles for some appropriate P -functions. Due to the greater generality, however, the relevant P and the conditions under which satisfies an elliptic differential inequality are rather implicitly given while in [4, 6] are given explicitly.

There are many other applications of P -functions that can be found in [20], among others, such as lower bounds for eigenvalue problems. One additional important application is in [1], where they showed that the monotonicity assumption $u_{x_n} > 0$, that is also stated in the De Giorgi’s conjecture, does in fact imply the local minimality of u . Such implication is by no means trivial and it is based on the construction of a so-called *calibration* associated to the energy functional. Such notion is intimately connected to the theory of null-Lagrangians, see [13], chapter 1 and chapter 4, section 2.4. In Theorem 4.4 in [1], they carry out the construction of the appropriate calibration for general integrands of the calculus of variations and such construction relies explicitly on the P -function.

Last but not least, there are applications such as gradient bounds similar to (1.2) and Liouville-type properties for vector equations. To be more precise, in Theorem 3.5 in [21], is a gradient bound is proved for the Ginzburg–Landau system of equations.

2. P -FUNCTIONS FOR FULLY NONLINEAR ELLIPTIC EQUATIONS

We begin by defining the notion of P -function

Definition 2.1. *Let $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^d$ be a smooth solution or subsolution of*

$$F(x, u, \nabla u, \dots, \nabla^m u) = 0 \quad (2.1)$$

where F is a continuous function.

We say that $P = P(x, u, \nabla u, \dots, \nabla^{m-1} u)$ is a P -function of (2.1) if there exists an elliptic operator L and a non negative function $\mu = \mu(x) \geq 0$

$$L = - \sum_{i,j=1}^n a_{ij} \partial_{x_i x_j} + \sum_{i=1}^n b_i \partial_{x_i} + c \quad , \quad \text{with } c \geq 0 \quad (2.2)$$

such that $\mu L P \leq 0$, in Ω .

an immediate corollary is that any P -function related to an equation or to a differential inequality attains its maximum at the boundary $\partial\Omega$ or at a point $x \in \Omega$ such that $\mu(x) = 0$.

We initially state as a direct consequence a strong maximum principle that holds in general (see Theorem 2.2 in [4] or Theorem 4.7 in [6]).

Theorem 2.2. *Let u be a smooth solution or subsolution of*

$$F(x, u, \nabla u, \dots, \nabla^m u) = 0 \quad , \quad u : \Omega \rightarrow \mathbb{R}^d \quad (2.3)$$

where Ω is a connected, bounded subset of \mathbb{R}^n

such that $\inf_{\overline{\Omega}} g(\nabla^k u) > 0$ for some $g : \mathbb{R}^{n^k \times d} \rightarrow [0, +\infty)$, $k \in \{1, \dots, m-1\}$ and suppose that $P = P(x, u, \nabla u, \dots, \nabla^{m-1} u)$ is a P -function of (2.3) with $\mu = \mu(g(\nabla^k u))$, $\mu(t) > 0$, $\forall t > 0$.

If there exists $x_0 \in \Omega$ such that

$$P(x_0, u(x_0), \dots, \nabla^{m-1} u(x_0)) = \sup_{\Omega} P(x, u, \dots, \nabla^{m-1} u) \quad (2.4)$$

then $P(x, u, \nabla u, \dots, \nabla^{m-1} u)$ is constant in Ω .

Proof. The proof is an immediate consequence of the strong maximum principle since $\mu(g(\nabla^k u)) > 0$ in Ω . □

The most common choice of g in Theorem 2.2 above is the Euclidean norm. For example, if $k = 1$, $g(\nabla u) = |\nabla u|$. If $\mu > 0$, $\forall t \geq 0$, then the assumption $\inf_{\overline{\Omega}} g(\nabla^k u) > 0$ is dismissed.

Remark 2.3. *The constancy of P -functions with a particular form hides geometric information on the level sets $\{x \in \mathbb{R}^n \mid u(x) = t\}$ of the solution u , such as the property of being surfaces of zero mean curvature (see Proposition 4.11 in [6]).*

We now focus on fully nonlinear elliptic equations. Let $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth solution of

$$F(x, u, \nabla u, \nabla^2 u) = 0 \quad (2.5)$$

where $F : \Omega \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is a continuous function and satisfies the ellipticity condition

$$\lambda |\xi|^2 \leq \sum_{i,j} F_{a_{ij}}(x, u, \nabla u, \nabla^2 u) \xi_i \xi_j \leq \Lambda |\xi|^2, \quad \forall x \in \Omega, \forall \xi \in \mathbb{R}^n \quad (2.6)$$

Here we use the notation $F = F(x, s, q, A)$, $s \in \mathbb{R}$, $q \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$ and $F_{a_{ij}} = \frac{\partial F}{\partial a_{ij}}$.

Some important examples of fully nonlinear elliptic equations are

(1) Monge-Ampère's equation

$$\det(\nabla^2 u) = f(x)$$

for strictly convex solutions u and $f > 0$.

(2) Equation of Prescribed Gauss curvature

$$\det(\nabla^2 u) = K(x)(1 + |\nabla u|^2)^{\frac{n+2}{2}}$$

$K(x)$ is the Gauss curvature of the graph u at $(x, u(x))$. Again, this equation is elliptic for strictly convex solutions u .

(3) Quasi-Linear equations of the form

$$\sum_{i,j} a_{ij}(\nabla u) u_{x_i x_j} = F(x, u, \nabla u)$$

where a_{ij} satisfy the ellipticity condition $\lambda |\xi|^2 \leq \sum_{i,j} a_{ij} \xi_i \xi_j \leq \Lambda |\xi|^2$. The p -Laplace equation, the Allen-Cahn equation and the minimal surface equation belong in this class of equations. Such equations are thoroughly studied in [4, 6, 9] among others.

There are many other important examples of fully nonlinear elliptic equations, such as Pucci's equation, Bellman equation, Isaacs equation (see [2]).

We now provide two general criteria for obtaining P -functions.

Lemma 2.4. *Let u be a smooth solution of (2.5) and F satisfies (2.6). Consider the quantity*

$$I := (\lambda g''(u) - 2F_u)|\nabla u|^2 + (g'(u)\nabla_q F - 2\nabla_x F)\nabla u + g'(u) \sum_{i,j} F_{a_{ij}} u_{x_i x_j} \quad (2.7)$$

and assume that $I \geq 0$ for some function $g : \mathbb{R} \rightarrow \mathbb{R}$.

Then $P(u, \nabla u) = |\nabla u|^2 + g(u)$ is a P -function of (2.5).

Proof. Assume that $I \geq 0$ for some $g : \mathbb{R} \rightarrow \mathbb{R}$ and let $P(u, |\nabla u|^2) = |\nabla u|^2 + g(u)$ (i.e. $P(s, t) = t + g(s)$).

We have

$$\begin{aligned} P_{x_i} &= 2 \sum_k u_{x_k} u_{x_k x_i} + g'(u) u_{x_i} \\ \Rightarrow P_{x_i x_j} &= 2 \sum_k (u_{x_k x_j} u_{x_k x_i} + u_{x_k} u_{x_k x_i x_j}) + g''(u) u_{x_i} u_{x_j} + g'(u) u_{x_i x_j} \\ \Rightarrow \sum_{i,j} d_{ij} P_{x_i x_j} &= 2 \sum_{i,j,k} (d_{ij} u_{x_k x_j} u_{x_k x_i} + d_{ij} u_{x_k} u_{x_k x_i x_j}) + g''(u) \sum_{i,j} d_{ij} u_{x_i} u_{x_j} \\ &\quad + g'(u) \sum_{i,j} d_{ij} u_{x_i x_j} \\ &\geq 2\lambda |Hes u|^2 + 2 \sum_{i,j,k} d_{ij} u_{x_k} u_{x_k x_i x_j} + \lambda g''(u) |\nabla u|^2 + g'(u) \sum_{i,j} d_{ij} u_{x_i x_j} \end{aligned} \quad (2.8)$$

where $d_{ij} = F_{a_{ij}} = \frac{\partial F}{\partial a_{ij}}$.

Differentiating (2.5) over x_k , and then multiplying by u_k , we obtain

$$\begin{aligned} F_{x_k} + F_u u_{x_k} + \sum_m F_{q_m} u_{x_m x_k} + \sum_{m,l} d_{ml} u_{x_m x_l x_k} &= 0 \\ \Rightarrow \sum_{m,l,k} d_{ml} u_k u_{x_m x_l x_k} &= -\nabla_x F \nabla u - F_u |\nabla u|^2 - \frac{1}{2} \nabla_q F \nabla_x P + \frac{1}{2} g'(u) \nabla_q F \nabla u \end{aligned} \quad (2.9)$$

Therefore the last equation of (2.8) becomes

$$\begin{aligned} \sum_{i,j} d_{ij} P_{x_i x_j} + \nabla_q F \nabla_x P &\geq 2\lambda |Hes u|^2 + g'(u) \nabla_q F \nabla u + (\lambda g''(u) - 2F_u) |\nabla u|^2 \\ &\quad - 2\nabla u \nabla_x F + g'(u) \sum_{i,j} F_{a_{ij}} u_{x_i x_j} \geq 0 \end{aligned} \quad (2.10)$$

□

Lemma 2.5. *Let u be a smooth solution of (2.5) and F satisfies (2.6). Consider the quantity*

$$\begin{aligned} J := & (\lambda B''(u) - 2A'(|\nabla u|^2)F_u)|\nabla u|^2 + (B'(u)\nabla_q F - 2A'(|\nabla u|^2)\nabla_x F)\nabla u \\ & + B'(u) \sum_{i,j} F_{a_{ij}} u_{x_i x_j} + \frac{\lambda(B'(u))^2}{2A'(|\nabla u|^2)} \end{aligned} \quad (2.11)$$

and assume that $J \geq 0$ for some functions $A, B : \mathbb{R} \rightarrow \mathbb{R}$ with $A' > 0$, $\forall t > 0$ and $A'' \geq 0$. Then $P(u, \nabla u) = A(|\nabla u|^2) + B(u)$ is a P -function of (2.5).

Proof. We argue as in Lemma 2.4 and obtain

$$\begin{aligned} \sum_{i,j} d_{ij} P_{x_i x_j} &= A'' \left(\sum_{i,j} d_{ij} [|\nabla u|^2]_{x_i} [|\nabla u|^2]_{x_j} \right) + 2A' \left(\sum_{i,j,k} d_{ij} [u_{x_k x_j} u_{x_k x_i} + u_{x_k} u_{x_k x_i x_j}] \right) \\ &\quad + \sum_{i,j} d_{ij} (B''(u) u_{x_i} u_{x_j} + B'(u) u_{x_i x_j}) \\ &\geq 2A' \lambda |Hes u|^2 + 2A' \sum_{i,j,k} d_{ij} u_{x_k} u_{x_k x_i x_j} + \lambda B'' |\nabla u|^2 + B' \sum_{i,j} d_{ij} u_{x_i x_j} \end{aligned} \quad (2.12)$$

by (2.6) and since $A'' \geq 0$, where $d_{ij} = \frac{\partial F}{\partial a_{ij}}$.

We also calculate from the first equation of (2.8) and the Cauchy-Schwarz inequality

$$\begin{aligned} \sum_i (P_{x_i} - B' u_{x_i})^2 &\leq 4(A')^2 |\nabla u|^2 |Hes u|^2 \\ \Rightarrow 2A' |Hes u|^2 &\geq \frac{1}{2|\nabla u|^2 A'} (|\nabla P|^2 - 2B' \nabla u \nabla P + (B')^2 |\nabla u|^2) \end{aligned} \quad (2.13)$$

In addition, similarly to (2.9) we have

$$\sum_{m,l,k} d_{ml} u_k u_{x_m x_l x_k} = -\nabla_x F \nabla u - F_u |\nabla u|^2 - \frac{1}{2A'} (\nabla_q F \nabla_x P - B' \nabla u \nabla_q F) \quad (2.14)$$

We plug (2.13) and (2.14) into (2.12) and thus

$$\begin{aligned} \sum_{i,j} d_{ij} P_{x_i x_j} &\geq \frac{\lambda}{2|\nabla u|^2 A'} (|\nabla P|^2 - 2B' \nabla u \nabla P + (B')^2 |\nabla u|^2) + \lambda B'' |\nabla u|^2 \\ &\quad + B' \sum_{i,j} d_{ij} u_{x_i x_j} + 2A' (-\nabla_x F \nabla u - F_u |\nabla u|^2 - \frac{1}{2A'} (\nabla_q F \nabla_x P - B' \nabla u \nabla_q F)) \end{aligned} \quad (2.15)$$

which gives

$$\sum_{i,j} d_{ij} P_{x_i x_j} + \left(\frac{\lambda B'(u)}{A'(|\nabla u|^2) |\nabla u|^2} \nabla u + \nabla_q F \right) \nabla_x P \geq J \quad (2.16)$$

and we conclude. \square

A direct consequence of Lemma 2.5 is the following.

Corollary 2.6. *Let $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth solution of*

$$\Delta u = f(u) \quad (2.17)$$

and let $P(s, t) = A(t) + B(s)$ such that $A' > 0$ for $t > 0$, $A'' \geq 0$ and assume that

$$t^2 B''(s) + B'(s) f(s) + \frac{(B'(s))^2}{2A'(t^2)} + 2A'(t^2) t^2 f'(s) \geq 0 \quad (2.18)$$

Then $P = P(u, |\nabla u|^2)$ is a P -function of (2.17).

2.1. Examples of P -functions.

(1) The well known P -function of (2.17) is

$$P(u, |\nabla u|^2) = \frac{|\nabla u|^2}{2} - F(u) \quad (2.19)$$

where $F'(u) = f(u)$

(see [16] or Chapter 5 in [20]).

It is easy to see that (2.19) satisfies (2.18) in Corollary 2.6.

(2) Another general example of P -function of (2.17) is

$$P(u, |\nabla u|^2) = \frac{|\nabla u|^4}{2} + 2 \int_0^u \left(\int_0^y \sqrt{f(z) f'(z)} dz \right)^2 dy \quad , \quad \text{if } f(t) f'(t) \geq 0, \forall t \in \mathbb{R} \quad (2.20)$$

$$P(u, |\nabla u|^2) = \frac{|\nabla u|^4}{2} - 2 \int_0^u \left(\int_0^y \sqrt{-f(z) f'(z)} dz \right)^2 dy \quad , \quad \text{if } f(t) f'(t) \leq 0$$

and satisfies condition (2.18) of Corollary 2.6.

Note that the above example is not in the form $P = g(u) |\nabla u|^2 + h(u)$ that we see in [20] as general form for P related to equation (2.17).

(3) Let u be a solution of

$$F(|\nabla u|^2 - cu, \nabla^2 u) = 0 \quad (2.21)$$

where F satisfies the ellipticity condition (2.6) and assume $\sum_{i,j} F_{a_{ij}} u_{x_i x_j} \leq \frac{\lambda c}{2}$, $c > 0$.

Then $P = P(u, |\nabla u|^2) = |\nabla u|^2 - cu$ is a P -function of (2.21) since P satisfies condition (2.11) of Lemma 2.5.

(4) The following example is in [18] (see Theorem 1).

Let u be a solution of

$$\operatorname{div}(\Phi'(|\nabla u|^2)\nabla u) = \rho(|\nabla u|^2)F'(u) \quad (2.22)$$

with $\Phi'(t), \rho(t) > 0$ and $\Phi'(t) + 2t\Phi''(t) > 0$, $\forall t \geq 0$.

Consider the function

$$P(s, t) = \int_0^t \frac{\Phi'(y) + 2y\Phi''(y)}{\rho(y)} dy - 2F(s) \quad (2.23)$$

Then $P = P(u, |\nabla u|^2)$ is a P -function of (2.22).

Note that for $\rho \equiv 1$, we have the one studied in [4].

3. GRADIENT BOUNDS AND PROPERTIES OF ENTIRE SOLUTIONS OF FULLY NONLINEAR EQUATIONS

In this section we will see that utilizing the techniques of [4, 9], we can obtain gradient bounds for solutions of equations of the form (2.5). To be more precise, for any explicit example of P -function with a specific property, we obtain a particular gradient bound.

Some of the regularity assumptions in this work can be relaxed for some classes of equations. In the study of Quasi-linear equations for example, we can only assume that $u \in W_{loc}^{1,p}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$, as in assumption (i) in Theorem 1.6 in [4] and utilize regularity results in [22] afterwards. For fully nonlinear elliptic equations also, we can relax the regularity of solutions and then utilize regularity results from [2, 23]. However, our main goal is not the optimal regularity assumptions since we state the results in an abstract form. Therefore, we will assume that the solutions are smooth and satisfy an analog of assumption (ii) in Theorem 1.6 in [4].

Assumption.

$$u \in C^2(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n), \nabla u \in C_{loc}^\alpha(\mathbb{R}^n; \mathbb{R}^n) \text{ for some } \alpha \in (0, 1) \quad (3.1)$$

and there exists $C = C(\|u\|_{L^\infty(\mathbb{R}^n)}) > 0$ such that $|\nabla u(x)| \leq C$, for any $x \in \mathbb{R}^n$

The next theorem provides an a priori pointwise estimate for solutions of (2.5). In contrast to the gradient bounds in [4, 6], the theorem below holds for any P -function that satisfies $P(u, 0) \leq 0$. When P is of the form $P = P(u, \nabla u)$ we use the notation $P(u, 0)$ instead of $P(u, 0, \dots, 0)$ and also we sometimes write $P = P(u; x)$ for simplicity.

Theorem 3.1. *Let u be an entire solution of*

$$F(x, u, \nabla u, \nabla^2 u) = 0 \quad (3.2)$$

that satisfy assumption (3.1). If $P = P(u, \nabla u)$ is a P -function of (3.2), with $\mu = \mu(|\nabla u|)$, $\mu(t) > 0, \forall t > 0$, such that $P(s, 0) \leq 0$,

Then

$$P(u(x), \nabla u(x)) \leq 0, \forall x \in \mathbb{R}^n \quad (3.3)$$

Proof. Let u be a solution of (3.2) that satisfies assumption (3.1) and consider the family of all translations of u ,

$$\mathcal{F} = \{v : \mathbb{R}^n \rightarrow \mathbb{R} \mid \exists z \in \mathbb{R}^n \text{ such that } v(x) = u(x+z) \forall x \in \mathbb{R}^n\} \quad (3.4)$$

\mathcal{F} is non empty since $u \in \mathcal{F}$.

Let P be a P -function of (3.2), with $\mu = \mu(|\nabla u|)$, $\mu(t) > 0, \forall t > 0$, such that $P(u, 0) \leq 0$. For simplicity, we denote $P = P(u; x)$ instead of $P = P(u(x), \nabla u(x))$.

Consider now

$$P_0 = \sup\{P(v; x) \mid v \in \mathcal{F}, x \in \mathbb{R}^n\} \quad (3.5)$$

We will prove that

$$P_0 \leq 0 \quad (3.6)$$

and from this we conclude.

We argue by contradiction, so we suppose that

$$P_0 > 0 \quad (3.7)$$

Then, by (3.5) there exist $(w_k)_{k \in \mathbb{N}}$ in \mathcal{F} and $(x_k)_{k \in \mathbb{N}}$ in \mathbb{R}^n such that

$$\lim_{k \rightarrow +\infty} P(w_k; x_k) = P_0 \quad (3.8)$$

Let $v_k(x) = w(x + x_k)$. Also, by definition we have that $v_k \in \mathcal{F}$ and $P(v_k; 0) = P(w_k; x_k)$, so that (3.8) can be rewritten as

$$\lim_{k \rightarrow +\infty} P(v_k; 0) = P_0 \quad (3.9)$$

Since $v_k \in C_{loc}^{1,\alpha}(\mathbb{R}^n)$, by the Ascoli-Arzelà theorem together with a diagonal argument, we can extract from $(v_k)_{k \in \mathbb{N}}$ a subsequence, denoted by $(v_k^{(k)})_{k \in \mathbb{N}}$ that converges with its first-order derivatives, uniformly on compact subsets of \mathbb{R}^n . Denote by \tilde{v} the limit function.

By the assumption (3.1) we have that

$$\mathcal{F} \text{ is relatively compact in } C_{loc}^{1,\alpha}(\mathbb{R}^n) \quad (3.10)$$

Thus $\tilde{v} \in \mathcal{F}$ and

$$P(\tilde{v}; 0) = \lim_{k \rightarrow +\infty} P(v_k^{(k)}; 0) = P_0 \quad (3.11)$$

by (3.9).

Consider now the set

$$U = \{x \in \mathbb{R}^n \mid P(\tilde{v}; x) = P_0\} \quad (3.12)$$

from the continuity of P on \mathbb{R}^n , U is closed and non empty since $0 \in U$. We will prove that U is also open. Let $x_0 \in U$, we observe that $|\nabla \tilde{v}(x_0)| \neq 0$, otherwise we would have

$$P_0 = P(\tilde{v}; x_0) = P(\tilde{v}(x_0), \nabla \tilde{v}(x_0)) = P(\tilde{v}(x_0), 0) \leq 0$$

against the assumption that $P_0 > 0$.

By continuity, there exists $\delta > 0$ such that

$$\inf_{\overline{B}_\delta(x_0)} |\nabla \tilde{v}| > 0 \text{ and thus } \inf_{\overline{B}_\delta(x_0)} \mu(|\nabla \tilde{v}|) > 0 \quad (3.13)$$

and by Theorem 2.2 we conclude that

$$P(\tilde{v}; x) \equiv P_0 \text{ in } B_\delta(x_0) \quad (3.14)$$

So U is open and it follows that $U = \mathbb{R}^n$ by connectedness.

On the other hand, since \tilde{v} is bounded it holds that there exists a sequence $(y_l)_{l \in \mathbb{N}}$ in \mathbb{R}^n such that

$$\lim_{l \rightarrow +\infty} \nabla \tilde{v}(y_l) = 0 \quad (3.15)$$

By the boundedness of \tilde{v} we also have $\tilde{v}(y_l) = \tilde{v}_l \rightarrow v_0$ up to a subsequence that we still denote as y_l , and so we obtain

$$0 < P_0 = \lim_{l \rightarrow \infty} P(\tilde{v}(y_l), \nabla \tilde{v}(y_l)) = P(v_0, 0)$$

which contradicts the assumption $P(s, 0) \leq 0$. Therefore $P_0 \leq 0$ and we conclude. \square

A direct consequence is a general gradient bound for fully nonlinear equations.

Corollary 3.2. *Let u be an entire solution of*

$$F(x, u, \nabla u, \nabla^2 u) = 0 \quad (3.16)$$

that satisfy assumption (3.1) and F satisfies the ellipticity condition (2.6). Consider $P = P(u, \nabla u) = A(|\nabla u|^2) + B(u)$ is a P -function from Lemma 2.5 such that $B \leq 0$.

Then

$$\begin{aligned} |\nabla u|^2 &\leq \Psi(u) \\ \text{where } \Psi(s) &= A^{-1}(-B(s)) \end{aligned} \quad (3.17)$$

Similarly, if P is obtained from Lemma 2.4 and $g \leq 0$, we have $|\nabla u|^2 \leq G(u)$, where $G = -g \geq 0$.

Remark 3.3. *Note that the condition $P(u, 0) \leq 0$ can be removed. Consider for example $P = P(u, \nabla u)$ be a P -function of (3.2) such that the condition $P(u, 0) \leq 0$ is not satisfied and set $\tilde{P}(u, \nabla u) = P(u, \nabla u) - \sup_{\mathbb{R}^n} P(u, 0)$. Then \tilde{P} is also a P -function of (3.2) and satisfies $\tilde{P}(u, 0) \leq 0$. The only difference is that the gradient bound in this case takes the form $P(u(x), \nabla u(x)) \leq \sup_{\mathbb{R}^n} P(u(x), 0)$.*

3.1. Gradient Bounds for entire solutions by the Examples of subsection 2.1.

Next, we observe that in Example (2) above, for solutions of (2.17),

$$P(u, |\nabla u|^2) = \frac{|\nabla u|^4}{2} - 2 \int_0^u \left(\int_0^y \sqrt{-f(z)f'(z)} dz \right)^2 dy \quad , \quad \text{if } f(t)f'(t) \leq 0 \quad (3.18)$$

satisfies $P(u, 0) \leq 0$.

Therefore, we have

Corollary 3.4. *Let u be a smooth and bounded entire solution to*

$$\Delta u = f(u) \quad (3.19)$$

where $f \in C^{1,\alpha}(\mathbb{R}^n)$ and $f(t)f'(t) \leq 0$.

Then

$$\frac{|\nabla u|^4}{4} \leq \int_0^u \left(\int_0^y \sqrt{-f(z)f'(z)} dz \right)^2 dy \quad (3.20)$$

Proof. By elliptic regularity theory (see [14]) we have that $u \in C^{2,\alpha}(\mathbb{R}^n)$ and that $|\nabla u|$ is bounded in \mathbb{R}^n . It suffices to prove that P defined in (3.18) is a P -function of (3.19) and then the conclusion is direct application of Theorem 3.1.

We have that P satisfies (2.18), $P(s, t) = \frac{t^2}{2} - 2 \int_0^s \left(\int_0^y \sqrt{-f(z)f'(z)} dz \right)^2 dy = \frac{t^2}{2} + q(s)$, so $P_t = t > 0$ for $t > 0$, $P_{tt} \geq 0$ and $\mu = P_t(u, |\nabla u|^2) |\nabla u|^2 = \frac{1}{2} |\nabla u|^4$.

Finally,

$$t^2 P_{ss}(s, t^2) + P_s(s, t^2)f(s) + 2P_t(s, t^2)t^2 f'(s) = 2t^4 f'(s) + t^2 q''(s) + q'(s)f(s) \geq 0$$

since the above polynomial has zero discriminant. \square

In addition, we have the following gradient bound for Example (3)

Corollary 3.5. *Let u be a non negative entire solution of*

$$F(|\nabla u|^2 - cu, \nabla^2 u) = 0 \tag{3.21}$$

that satisfy (3.1), where F satisfies the ellipticity condition (2.6) and assume

$$\sum_{i,j} F_{a_{ij}} u_{x_i x_j} \leq \frac{\lambda c}{2} \text{ for some } c > 0.$$

Then

$$|\nabla u|^2 \leq cu \tag{3.22}$$

Proof. We have that the function

$$P(s, t) = t - cs$$

satisfy the condition (2.11) and also, $P(u, 0) = -cu \leq 0$ since u is non negative by assumption. Therefore we conclude by the Theorem 3.1. \square

Another application of interest of Theorem 3.1, utilizing the Example (4), is the following gradient bound that is a quite more general form of the gradient bound in [4].

Corollary 3.6. *Let u be an entire solution of*

$$\operatorname{div}(\Phi'(|\nabla u|^2)\nabla u) = \rho(|\nabla u|^2)F'(u) \text{ , } F \geq 0 \tag{3.23}$$

that satisfy assumption (3.1), with $\Phi'(t), \rho(t) > 0$ and $\Phi'(t) + 2t\Phi''(t) > 0$, $\forall t \geq 0$.

Then

$$|\nabla u|^2 \leq \Psi(u) \text{ , where } \Psi(u) = Q^{-1}(2F(u)) \tag{3.24}$$

$$\text{and } Q(t) = \int_0^t \frac{\Phi'(y) + 2y\Phi''(y)}{\rho(y)} dy$$

Proof. By Theorem 1 in [18], we have that $P(u, |\nabla u|^2) = Q(|\nabla u|^2) - 2F(u)$ is a P -function of (3.23) with $\mu(t) > 0$, $\forall t \geq 0$ and satisfies $P(u, 0) \leq 0$ since $F \geq 0$. Thus we apply Theorem 3.1 and we conclude. \square

4. RIGIDITY RESULTS AND PROPERTIES OF ENTIRE SOLUTIONS OF FULLY NONLINEAR EQUATIONS

In this section we will see that if an equation admits a P -function of the form $P = |\nabla u|^2$, then the solutions that satisfy assumption (3.1) are constant. Special cases of this result are some of the well-known results of J. Serrin in [19].

In particular we have

Theorem 4.1. *Let u be an entire solution of*

$$F(u, \nabla u, \nabla^2 u) = 0 \quad (4.1)$$

that satisfy assumption (3.1) where F satisfies the condition (2.6) and assume $F_u \leq 0$.

Then u is a constant

Proof. The proof of Theorem 4.1 is a consequence of Theorem 3.1 and Lemma 2.4 by considering $P = |\nabla u|^2$. \square

Remark 4.2. (1) *If $F = F(x, u, \nabla u, \nabla^2 u)$ and assume in addition that $\nabla_x F \cdot \nabla u \leq 0$, then the conclusion of Theorem 4.1 still holds by Theorem 3.1 and Lemma 2.4. Note also that Theorem 1, Theorem 6 and Theorem 8 in [19] are recovered.*

(2) *We note that, in the special case where equation (4.1) takes the form*

$$\operatorname{div}(\Phi'(|\nabla u|^2)\nabla u) = f'(u) \quad (4.2)$$

then the condition $F_u \leq 0$ reads $f''(u) \geq 0$ which implies stability of the solutions, i.e. the second variation of the associated energy functional $J(u) = \int (\frac{1}{2}\Phi(|\nabla u|^2) + f(u))dx$ is non negative (see also Theorem 4.5 in [4]).

Another consequence of Theorem 3.1 is the following Liouville-type result

Theorem 4.3. *Let u be an entire solution of (3.2) that satisfies assumption (3.1) and P is a P -function from 2.5 that satisfies $P(u, 0) \leq 0$. If there exists $x_0 \in \mathbb{R}^n$ such that $B(u(x_0)) = 0$, then $u \equiv \text{const.}$ in \mathbb{R}^n .*

Proof. We argue as in the proof of Theorem 1.8 in [4] with slight modifications. For the convenience of the reader we provide the details.

Suppose that $B(u(x_0)) = 0$, let $u_0 = u(x_0)$ and consider the set

$$V = \{x \in \mathbb{R}^n \mid u(x) = u_0\} \quad (4.3)$$

V is a closed set and by the assumption, non empty. Let $x_1 \in V$ and consider the function $\phi(t) = u(x_1 + t\omega) - u_0$, where $|\omega| = 1$ is arbitrarily fixed. We have $|\phi'(t)| = |\nabla u(x_1 + t\omega)|$. By the gradient bound in Corollary 3.2 we have,

$$|\nabla u|^2 \leq \Psi(u) \quad , \quad \text{where } \Psi(s) = A^{-1}(-B(s)) \quad (4.4)$$

Since $\Psi \in C^2(\mathbb{R})$ and $\Psi(u_0) = 0$, we have $\Psi(u) = O(|u - u_0|^2)$, as $|u - u_0| \rightarrow 0$. So, we conclude from (4.4) that $|\phi'(t)| \leq C|\phi(t)|$ for t small enough. Since $\phi(0) = 0$, we must have $\phi \equiv 0$ on $[-\delta, \delta]$, for some $\delta > 0$. Thus V is open, which gives that $V = \mathbb{R}^n$. \square

Also, we have a De Giorgi type property for solutions that attain the equality at a point in the gradient bound obtained in Corollary 3.2.

Theorem 4.4. *Let u be an entire solution of*

$$F(u, \nabla u, \nabla^2 u) = 0 \quad (4.5)$$

that satisfy assumption (3.1) and let $P = P(u, |\nabla u|^2)$ be a P -function of (4.5) obtained in Lemma 2.5 that satisfies $P(u, 0) \leq 0$. If there exists $x_0 \in \mathbb{R}^n$ such that

$$P(u(x_0), |\nabla u(x_0)|^2) = 0 \quad (4.6)$$

then there exists a function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\begin{aligned} \text{either } u(x) &= g(a \cdot x + b) \quad , \quad a \in \mathbb{R}^n \text{ with } |a| = 1, \quad , \quad b \in \mathbb{R} \\ \text{or } u(x) &= g(|x - z_0| + c) \quad , \quad z_0 \in \mathbb{R}^n \text{ and } c \in \mathbb{R} \end{aligned} \quad (4.7)$$

Proof. By Corollary 3.2, we have that $P(u, |\nabla u|^2) \leq 0$.

We begin by considering the set

$$\mathcal{A} = \{x \in \mathbb{R}^n : P(u, |\nabla u|^2) = 0\} \quad (4.8)$$

\mathcal{A} is closed and by the assumption $\mathcal{A} \neq \emptyset$. We are going to prove that \mathcal{A} is open.

Let $x_1 \in \mathcal{A}$, if $\nabla u(x_1) = 0$, we obtain by the form $P(s, t) = A(t) + B(s)$ that $P(u(x_1), 0) = -B(u(x_1)) = 0$. By Theorem 4.3, we conclude that $u \equiv u(x_1)$ and $\nabla u \equiv 0$ and hence $P \equiv 0$.

On the other hand, if $\nabla u(x_1) \neq 0$, we have $\inf_{\overline{B}_{\delta_1}(x_1)} |\nabla u| > 0$ for some $\delta_1 > 0$ and by Theorem 2.2 we conclude that $P(u, |\nabla u|^2) \equiv 0$ in $B_{\delta_1}(x_1)$ and therefore \mathcal{A} is open.

By connectedness, we have that $\mathcal{A} = \mathbb{R}^n$, that is,

$$P(u, |\nabla u|^2) \equiv 0 \quad , \quad \forall x \in \mathbb{R}^n \quad (4.9)$$

and $P_t = A'(t) > 0$, thus

$$|\nabla u|^2 = \Psi(u) \quad , \quad \text{in } \mathbb{R}^n \quad , \quad \text{where } \Psi(u) = A^{-1}(-B(u)) \quad (4.10)$$

Now, if there exists $x_2 \in \mathbb{R}^n$ such that $\Phi(u(x_2)) = 0$, so $|\nabla u(x_2)| = 0$, again by Theorem 4.3 we have that $u \equiv u(x_2)$.

If, on the other hand $\Psi(u(x)) > 0$, $\forall x \in \mathbb{R}^n$, we set

$$v = G(u) \quad , \quad \text{where } G'(s) = \frac{1}{\Psi(s)} \quad (4.11)$$

$$\text{and } |\nabla v|^2 = 1 \quad \text{in } \mathbb{R}^n$$

Therefore, by the result in [3], we have that

$$\begin{aligned} \text{either } v(x) &= a \cdot x + b \quad , \quad a \in \mathbb{R}^n \quad \text{with } |a| = 1 \quad \text{and } b \in \mathbb{R} \\ \text{or } v(x) &= |x - z_0| + c \quad , \quad z_0 \in \mathbb{R}^n \quad \text{and } c \in \mathbb{R} \end{aligned} \quad (4.12)$$

So we conclude that

$$\begin{aligned} \text{either } u(x) &= g(a \cdot x + b) \quad , \quad a \in \mathbb{R}^n \quad \text{with } |a| = 1 \quad , \quad b \in \mathbb{R} \quad \text{where } g(s) = G^{-1}(s) \\ \text{or } u(x) &= g(|x - z_0| + c) \quad , \quad z_0 \in \mathbb{R}^n \quad \text{and } c \in \mathbb{R} \end{aligned} \quad (4.13)$$

□

Remark 4.5. Note that if $u : \Omega \rightarrow \mathbb{R}$ where Ω is an open and connected domain in \mathbb{R}^n and $P = P(u, |\nabla u|^2) = A(|\nabla u|^2) + B(u)$ with $A' > 0$ that attains its maximum at a point then u will be a solution of the Eikonal equation $|\nabla u|^2 = \Psi(u)$. If in addition $u_{x_n} > 0$ and consider $F_i = \frac{u_{x_i}}{u_{x_n}}$, by Proposition 2.1 in [11], the function $F = (F_1, \dots, F_{n-1})$ will satisfy the Isobaric Euler equation.

5. A HARNACK-TYPE INEQUALITY AND LOCAL ESTIMATES FOR THE GRADIENT

5.1. Estimates for Quasi-Linear equations. We will establish a Harnack inequality and local estimates for the gradient of solutions to Quasi-linear equations in a domain $\Omega \subset \mathbb{R}^n$. Properties of entire solutions for such equations have been studied in [4, 6, 9] among others. These estimates can be extended for some fully nonlinear equations such as for the Monge-Ampère equation.

Let $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth solution of

$$\operatorname{div}(\Phi'(|\nabla u|^2)\nabla u) = f'(u) \quad (5.1)$$

or equivalently,

$$\sum_{i,j} a_{ij}(\nabla u) u_{x_i x_j} = f'(u) \quad (5.2)$$

$$\text{where } a_{ij}(\sigma) = 2\Phi''(|\sigma|^2)\sigma_i\sigma_j + \Phi'(|\sigma|^2)\delta_{ij}$$

and we assume that a_{ij} satisfy the ellipticity condition

$$\theta|\xi|^2 \leq \sum_{i,j} a_{ij}\xi_i\xi_j \leq \Theta|\xi|^2 \quad (5.3)$$

Some important examples are

(1) The p -Laplacian

$$\operatorname{div}(|\nabla u|^{p-2}\nabla u) = f'(u) \quad , \quad \text{for } p > 1 \quad (5.4)$$

(2) The Allen-Cahn equation (for $p = 2$)

$$\Delta u = f'(u) \quad (5.5)$$

(3) The minimal surface equation

$$\operatorname{div}\left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}}\right) = f'(u) \quad (5.6)$$

Assume also that $a_{ij} \in L^\infty(\Omega)$. First, we have Local Boundedness on the gradient of solutions.

Theorem 5.1. *Let $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth solution of (5.1) and assume (5.3). Suppose $f''(u)|\nabla u|^2 \in L^q(\Omega)$ for some $q > \frac{n}{2}$.*

Then for any $B_R \subset \Omega$, any $0 < r < R$ and any $p \geq 1$

$$\sup_{B_r} |\nabla u|^2 \leq C \left[\frac{\|\nabla u\|_{L^{2p}(B_R)}^2}{(R-r)^{n/p}} + R^{2-n/q} \|f''(u)|\nabla u|^2\|_{L^q(B_R)} \right] \quad (5.7)$$

where $C = C(n, \theta, \Theta, p, q)$ is a positive constant.

In addition, if $f'' \geq 0$, the assumption on $f''(u)|\nabla u|^2$ is dismissed and we have

$$\sup_{B_r} |\nabla u|^2 \leq \frac{C}{(R-r)^{n/p}} \|\nabla u\|_{L^{2p}(B_R)}^2 \quad (5.8)$$

Proof. Arguing as in Lemma 2.4 with $g = 0$, we have

$$\sum_{i,j} a_{ij} P_{x_i x_j} \geq 2\theta |\operatorname{Hes} u|^2 + 2 \sum_{i,j,k} a_{ij} u_{x_k} u_{x_k x_i x_j} \quad (5.9)$$

Differentiating (5.2) over x_k , multiplying with u_{x_k} and summing over k ,

$$\sum_{i,j,k} (a_{ij} u_{x_i x_j})_{x_k} u_{x_k} = f''(u) |\nabla u|^2 \quad (5.10)$$

Now since $\frac{\partial a_{ij}}{\partial x_k} = \frac{\partial a_{ki}}{\partial x_j}$ we have

$$\begin{aligned} & \sum_{i,j,k} (a_{ij} u_{x_k x_i})_{x_j} u_{x_k} = f''(u) |\nabla u|^2 \\ \Rightarrow & \sum_{i,j,k} a_{ij} u_{x_k x_i x_j} u_{x_k} = f''(u) |\nabla u|^2 - \sum_{i,j,k} (a_{ij})_{x_j} u_{x_k x_i} u_{x_k} \end{aligned} \quad (5.11)$$

and thus (5.9) becomes

$$\sum_{i,j} (a_{ij} P_{x_i})_{x_j} \geq 2\theta |Hes u|^2 + 2f''(u) |\nabla u|^2 \quad (5.12)$$

Finally, by the local boundedness for elliptic equations we conclude (see for example Theorem 4.14 in [15]).

In the case where $f'' \geq 0$, we have that $-\sum_{i,j} (a_{ij} P_{x_i})_{x_j} \leq 0$ and similarly we have (5.8). \square

Additionally, a Harnack-type inequality is established with similar arguments.

Theorem 5.2. *Let $u : \Omega \rightarrow \mathbb{R}$ be a smooth solution of (5.1) and assume (5.3). Suppose $|Hes u|^2, f''(u) |\nabla u|^2 \in L^q(\Omega)$ for some $q > \frac{n}{2}$. Then for any $B_R \subset \Omega$ there holds that for any $0 < p < \frac{n}{n-2}$ and any $0 < \eta < \tau < 1$*

$$C \left(\frac{1}{R^n} \int_{B_{\tau R}} |\nabla u|^{2p} \right)^{\frac{1}{p}} \leq \inf_{B_{\eta R}} |\nabla u|^2 + R^{2-\frac{n}{p}} \| |Hes u|^2 \|_{L^{2q}(B_R)}^2 + \| f''(u) |\nabla u|^2 \|_{L^q(B_R)} \quad (5.13)$$

where $C = C(n, p, q, \theta, \Theta, \eta, \tau)$.

Moreover, if $f'' \leq 0$, the assumption on $f''(u) |\nabla u|^2$ is dismissed and we have

$$C \left(\frac{1}{R^n} \int_{B_{\tau R}} |\nabla u|^{2p} \right)^{\frac{1}{p}} \leq \inf_{B_{\eta R}} |\nabla u|^2 + R^{2-\frac{n}{p}} \| |Hes u|^2 \|_{L^{2q}(B_R)}^2 \quad (5.14)$$

Proof. As in the proof of Theorem 5.1, together with the ellipticity of a_{ij} there holds

$$\sum_{i,j} (a_{ij} P_{x_i})_{x_j} \leq 2\Theta |Hes u|^2 + 2f''(u) |\nabla u|^2 \quad (5.15)$$

and by the Harnack inequality applied in $P = |\nabla u|^2$ we conclude (see for example Theorem 4.15 in [15]).

If $f'' \leq 0$, (5.15) becomes

$$-\sum_{i,j} (a_{ij} P_{x_i})_{x_j} \geq -2\Theta |Hes u|^2 \quad (5.16)$$

and similarly we conclude. \square

5.2. Estimates for Fully-Nonlinear Elliptic equations. The local estimates and the Harnack-type inequality can be extended to fully nonlinear elliptic equations. By Lemma 2.4 with $g = 0$ or by Lemma 2.5 with $B = 0$, we can utilize the elliptic inequality (2.10) or (2.16) respectively and assuming that $F_{a_{ij}} \in L^\infty(\Omega)$, we can apply Local Estimates for subsolutions to general elliptic operators to the P -function and obtain similar local estimates for the gradient of solutions to fully nonlinear equations.

So, we consider solutions of the equation

$$F(x, u, \nabla u, \nabla^2 u) = 0 \quad (5.17)$$

where $F : \Omega \times \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^{n \times n} \rightarrow \mathbb{R}$ is a continuous function and satisfies the ellipticity condition (2.6). We denote $F = F(x, s, q, A)$, $F_{q_i} = \frac{\partial F}{\partial q_i}$, $F_{a_{ij}} = \frac{\partial F}{\partial a_{ij}}$.

In this subsection we will assume the bound

$$|F_{a_{ij}}|, |F_{q_i}| \leq M \quad (5.18)$$

We first establish a local pointwise estimate for the gradient of solutions.

Theorem 5.3. *Let $u : \Omega \rightarrow \mathbb{R}$ be a smooth solution of (5.17) and assume $\nabla_x F \cdot \nabla u \leq 0$ and $F_u |\nabla u|^2 \in L^n(\Omega)$.*

Then for any $B_{2R} \subset \Omega$ and any $p \geq 1$

$$\sup_{B_R} |\nabla u|^2 \leq C \left[\frac{\|\nabla u\|_{L^{2p}(B_R)}^2}{|B_{2R}|^{1/p}} + \frac{R}{\lambda} \|F_u(u) |\nabla u|^2\|_{L^n(B_{2R})} \right] \quad (5.19)$$

where $C = C(n, p, \lambda, \Lambda, MR^2)$ is a positive constant.

In addition, if $F_u \leq 0$, the assumption on $F_u(u) |\nabla u|^2$ is dismissed and we have

$$\sup_{B_R} |\nabla u|^2 \leq \frac{C}{|B_{2R}|^{1/p}} \|\nabla u\|_{L^{2p}(B_{2R})}^2 \quad (5.20)$$

Proof. Set $P = |\nabla u|^2$, by Lemma 2.4 and the assumption $\nabla_x F \cdot \nabla u \leq 0$ we have

$$\sum_{i,j} d_{ij} P_{x_i x_j} + \nabla_q F \nabla_x P \geq -2F_u |\nabla u|^2 \quad (5.21)$$

where $d_{ij} = F_{a_{ij}}$. That is

$$L P \leq 2F_u |\nabla u|^2 \quad (5.22)$$

Thus, by classical local pointwise estimates for subsolutions of elliptic equations (see for example Theorem 9.20 in [14]) we obtain (5.19).

In the case where $F_u \leq 0$ we have $L P \leq 0$ and similarly we conclude. \square

As in the previous subsection, we obtain the following Harnack estimate

Theorem 5.4. *Let $u : \Omega \rightarrow \mathbb{R}$ be a smooth solution of (5.17). Assume $\nabla_x F \cdot \nabla u \geq 0$, $|Hes u|^2, F_u(u)|\nabla u|^2 \in L^n(\Omega)$.*

Then for any $B_{2R} \subset \Omega$ there holds that for any $p \geq 1$,

$$\left(\frac{1}{|B_R|} \int_{B_R} |\nabla u|^{2p}\right)^{\frac{1}{p}} \leq C(\inf_{B_R} |\nabla u|^2 + \frac{R}{\lambda} \|Hes u\|_{L^{2n}(B_{2R})}^2 + \frac{R}{\lambda} \|F_u(u)|\nabla u|^2\|_{L^n(B_{2R})}) \quad (5.23)$$

where $C = C(n, p, \lambda, \Lambda, MR^2)$.

Moreover, if $F_u \geq 0$, the assumption on $F_u(u)|\nabla u|^2$ is dismissed and we have

$$\left(\frac{1}{|B_R|} \int_{B_R} |\nabla u|^{2p}\right)^{\frac{1}{p}} \leq C(\inf_{B_R} |\nabla u|^2 + \frac{R}{\lambda} \|Hes u\|_{L^{2n}(B_{2R})}^2) \quad (5.24)$$

Proof. Set $P = |\nabla u|^2$. We argue as in the proof of Lemma 2.4 and by the assumption $\nabla_x F \cdot \nabla u \geq 0$ to obtain

$$\sum_{i,j} d_{ij} P_{x_i x_j} + \nabla_q F \nabla_x P \leq -2F_u |\nabla u|^2 + 2\Lambda |Hes u|^2 \quad (5.25)$$

So, by the Harnack inequality for supersolutions of elliptic equations (see Theorem 9.22 in [14] for instance), we obtain (5.23).

If in addition $F_u \geq 0$, we have $LP \geq -2\Lambda |Hes u|^2$, where $L = -\sum_{i,j} d_{ij} \partial_{x_i x_j} - \sum_i F_{q_i} \partial_{x_i}$ and we conclude. \square

Last but not least, we have the following estimates for the Monge-Ampère equation

Corollary 5.5. *Let $u : \Omega \rightarrow \mathbb{R}$ be a smooth and convex solution of*

$$\det(\nabla^2 u) = f(u, \nabla u) \quad (5.26)$$

where $f > 0$ and assume $|f_{q_i}|, |adj^T(\nabla^2 u)_{ij}| \leq M$.

(1) *If $f_u \geq 0$, then for any $B_{2R} \subset \Omega$ and any $p \geq 1$ we have*

$$\sup_{B_R} |\nabla u|^2 \leq \frac{C}{R^{n/p}} \|\nabla u\|_{L^{2p}(B_{2R})}^2 \quad (5.27)$$

(2) *If $f_u \leq 0$ and $|Hes u|^2 \in L^n(\Omega)$, then for any $B_{2R} \subset \Omega$ and any $p \geq 1$ we have*

$$\left(\frac{1}{|B_R|} \int_{B_R} |\nabla u|^{2p}\right)^{\frac{1}{p}} \leq C(\inf_{B_R} |\nabla u|^2 + \frac{R}{\lambda} \|Hes u\|_{L^{2n}(B_{2R})}^2) \quad (5.28)$$

where $C = C(n, p, \lambda, \Lambda, MR^2)$.

Proof. The proof (1) is a consequence of Theorem 5.3 and the proof of (2) is a consequence of Theorem 5.4 and since $\frac{\partial F}{\partial a_{ij}}(A) = \text{adj}^T(A)_{ij}$, for $F(A) = \det(A)$ by Jacobi's formula. \square

Note: If $\Omega = \mathbb{R}^n$, the fact that $|\nabla u|^2$ is a P -function of (5.26) in view of Theorem 4.1 states that there is no solution of (5.26) that satisfies the assumption (3.1). Indeed, if we assume that u is an entire solution of (5.26) that satisfies (3.1), then by Theorem 4.1 we have that u is constant in \mathbb{R}^n , which contradicts the fact that it's Hessian has positive determinant. We can also see this as follows, if $|\nabla u|$ is bounded in \mathbb{R}^n , then ∇u can not be a global diffeomorphism and thus $\det(\nabla^2 u)$ can not be strictly positive in \mathbb{R}^n .

6. HIGHER ORDER NONLINEAR EQUATIONS

In this last section, we will provide examples of P -functions for higher order nonlinear equations and their applications. In particular, an analogous version of Theorems 3.1 and 4.1, allow us to obtain properties and pointwise estimates of entire solutions even in this case. Moreover, we establish local pointwise estimates for nonlinear equations of order greater than two, through the mean value properties of the P -functions or with analogous arguments to that of section 5, applied in higher order equations. This method can be applied to many other classes of higher order nonlinear equations.

We begin by stating the analogous Theorem 3.1 for equations of general order.

Assumption

$$\begin{aligned} u \in C^m(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n) \text{ , } \nabla^{m-1}u \in C_{loc}^\alpha(\mathbb{R}^n) \text{ for some } \alpha \in (0, 1) \\ \text{and there exists } C > 0 \text{ such that } |\nabla^l u| \leq C \text{ , } l = 1, \dots, m-1. \end{aligned} \quad (6.1)$$

Theorem 6.1. *Let u be an entire solution or subsolution of*

$$F(x, u, \nabla u, \dots, \nabla^m u) = 0 \quad (6.2)$$

that satisfies assumption (6.1) and let $P = P(u, \dots, \nabla^{m-1}u) = P(u; x)$ be a P -function of (6.2) such that one of the following holds:

- (i) $\mu = \mu(g(\nabla^k u))$ for some $g : \mathbb{R}^{n^k} \rightarrow \mathbb{R}$, $g(z) > 0$, $\forall z \neq 0$, $g((0, \dots, 0)) = 0$, $\mu(t) > 0$, $\forall t > 0$ and $P(u; x) \leq 0$, when $\nabla^k u = (0, \dots, 0)$, $k \in \{1, \dots, m-1\}$,
- (ii) $\mu = \mu(g(\nabla^k u))$ for some $g : \mathbb{R}^{n^k} \rightarrow \mathbb{R}$, $g(z) > 0$, $\forall z \neq 0$, $g((0, \dots, 0)) = 0$, $\mu(t) > 0$, $\forall t > 0$, $P(u; x) \leq 0$, when $\nabla^l u = (0, \dots, 0)$, $k \neq l$, $k, l \in \{1, \dots, m-1\}$ and $g(\nabla^k u) > 0$, $\forall x \in \mathbb{R}^n$.

Then $P(u, \dots, \nabla^{m-1}u) \leq 0 \forall x \in \mathbb{R}^n$.

Proof. The proof is similar to that of Theorem 3.1 with minor modifications. \square

We now provide the generalization of Theorem 4.1 in the higher order case.

Theorem 6.2. *Let u be an entire solution of*

$$F(x, u, \nabla u, \dots, \nabla^m u) = 0 \quad (6.3)$$

and let $P = P(u, \dots, \nabla^{m-1}u) = P(u; x)$ be a P -function of (6.2) such that $\mu = \mu(g(\nabla^k u))$ for some $g : \mathbb{R}^{n^k} \rightarrow \mathbb{R}$, $g(z) > 0, \forall z \neq 0$, $g((0, \dots, 0)) = 0$, $\mu(t) > 0, \forall t > 0$ and

$$\begin{aligned} P &= H(\nabla^k u) \quad , \quad \text{where } H : \mathbb{R}^{n^k} \rightarrow [0, +\infty) \\ \text{and } \{H = 0\} &= \{0 \in \mathbb{R}^{n^k}\} \quad , \quad k \in \{1, \dots, m-1\} \end{aligned} \quad (6.4)$$

Then $\nabla^{k-1}u$ is a constant.

Proof. The proof is direct consequence of Theorem 6.1 since $P = H(\nabla^k u) = 0$ when $\nabla^k u$ vanish which gives $P \equiv 0$ in \mathbb{R}^n . \square

6.1. Local and Global Pointwise estimates. The arguments of section 5, can be applied for higher order nonlinear equations. In this case, we extract local and global estimates for higher order of derivatives of u , such as for the Laplacian.

Proposition 6.3. *Let $u : \Omega \rightarrow \mathbb{R}$ be a smooth and convex subsolution of*

$$\Delta^2 u - F(x, u, \nabla u, \nabla^2 u, \nabla^3 u) = 0 \quad , \quad \text{with } F \geq 0 \quad (6.5)$$

Then for any $B_R \subset \Omega$, any $0 < r < R$ and any $p \geq 1$,

$$\sup_{B_r} (\Delta u)^2 \leq \frac{C}{(R-r)^{n/p}} \|\Delta u\|_{L^{2p}(B_R)}^2 \quad (6.6)$$

where $C = C(n, p)$.

Proof.

$$\begin{aligned} P_{x_i} &= 2\Delta u \Delta u_{x_i} \\ P_{x_i x_i} &= 2(\Delta u_{x_i})^2 + 2\Delta u \Delta u_{x_i x_i} \\ \Rightarrow \Delta P &= 2|\nabla \Delta u|^2 + 2\Delta u \Delta^2 u \geq 2|\nabla \Delta u|^2 + F \Delta u \geq 0 \end{aligned} \quad (6.7)$$

Therefore, by classical estimates for subsolutions of elliptic equations he conclude (see for example Theorem 4.14 in [15]). \square

Furthermore, we give some examples of P -functions of the form $P = P(u, |\nabla u|, \Delta u)$ related to fourth order nonlinear equations together with applications.

Proposition 6.4. *Let u be a smooth solution of*

$$a(\Delta u)[|\nabla u|^2 \Delta^2 u - \Delta u(\nabla u \cdot \nabla \Delta u)] = b(u)|\nabla u|^4 \quad (6.8)$$

where $a, b : \mathbb{R} \rightarrow \mathbb{R}$ and $a > 0$, $a' \geq 0$

and set $P(s, t) = A(t) - B(s)$ such that $A' = a$ and $B'' = b$.

Then $P = P(u, \Delta u) = A(\Delta u) - B(u)$ is a P -function of (6.8).

In addition, if u satisfies (6.1) with $m = 4$, $B(u) \geq 0$ and $u_{x_n} > 0$, then

$$\Delta u \leq \Gamma(u) \quad \forall x \in \mathbb{R}^n, \quad \text{where } \Gamma(u) = A^{-1}(B(u)). \quad (6.9)$$

Proof. We have

$$P_{x_i} = P_s u_{x_i} + P_t \Delta u_{x_i} \quad (6.10)$$

and so,

$$\begin{aligned} \Delta u(\nabla P \cdot \nabla u) &= P_s |\nabla u|^2 \Delta u + P_t \Delta u \sum_{i=1}^n u_{x_i} \Delta u_{x_i} \\ \Leftrightarrow -B'(u)|\nabla u|^2 \Delta u &= \Delta u(\nabla P \cdot \nabla u) - A'(\Delta u) \Delta u \sum_{i=1}^n u_{x_i} \Delta u_{x_i} \end{aligned} \quad (6.11)$$

on the other hand we have

$$\begin{aligned} P_{x_i x_i} &= P_{ss} u_{x_i}^2 + 2P_{st} u_{x_i} \Delta u_{x_i} + P_{tt} (\Delta u_{x_i})^2 + P_s u_{x_i x_i} + P_t \Delta u_{x_i x_i} \\ \Rightarrow \Delta P &= (-B''(u))|\nabla u|^2 + A''(\Delta u) \sum_{i=1}^n (\Delta u_{x_i})^2 - B'(u) \Delta u + A'(\Delta u) \Delta^2 u \end{aligned} \quad (6.12)$$

and by (6.11) and the assumptions of A and B , (6.12) becomes

$$|\nabla u|^2 \Delta P - \Delta u(\nabla P \cdot \nabla u) \geq a(\Delta u)[|\nabla u|^2 \Delta^2 u - \Delta u(\nabla u \cdot \nabla \Delta u)] - b(u)|\nabla u|^4 = 0 \quad (6.13)$$

For the bound of the Laplacian, we have $P(u, 0) = -B(u) \leq 0$ and $\mu = |\nabla u|^2 > 0 \quad \forall x \in \mathbb{R}^n$ since $u_{x_n} > 0$, so the assumption (i) in Theorem 6.1 is satisfied and we conclude. \square

Proposition 6.5. *Let u be a smooth solution of*

$$|\text{Hes } u|^2 = F(u, |\nabla u|^2, \Delta u) + \frac{u}{2} \Delta^2 u \quad (6.14)$$

where $F : \mathbb{R}^3 \rightarrow \mathbb{R}$ is such that $F(s, t, w) \geq \frac{1}{2} w^2$.

Then $P = P(u, |\nabla u|^2, \Delta u) = |\nabla u|^2 - u \Delta u$ is a P -function of (6.16).

In addition, if u is non negative, convex solution of (6.16) that satisfies assumption (6.1), then

$$|\nabla u|^2 \leq u \Delta u \quad , \quad \forall x \in \mathbb{R}^n. \quad (6.15)$$

Proof. We have that

$$P_{x_i} = 2 \sum_{j=1}^n u_{x_j} u_{x_j x_i} - u_{x_i} \Delta u - u \Delta u_{x_i}$$

and

$$\Delta P = 2|Hes u|^2 + 2\nabla u \nabla \Delta u - (\Delta u)^2 - 2\nabla u \nabla \Delta u - u \Delta^2 u$$

so by (6.16),

$$\Delta P = 2F(u, |\nabla u|^2, \Delta u) - (\Delta u)^2 \geq 0$$

For the gradient bound we see that $P(u, 0, \Delta u) = -u \Delta u \leq 0$ since u is non negative and convex, so the assumption (i) of Theorem 6.1 is satisfied and we conclude. \square

As a result, we have the following pointwise estimate

Corollary 6.6. *Let $u : B_2 \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth solution of*

$$|Hes u|^2 = F(u, |\nabla u|^2, \Delta u) + \frac{u}{2} \Delta^2 u \quad (6.16)$$

$$\text{where } F : \mathbb{R}^3 \rightarrow \mathbb{R} \text{ is such that } F(s, t, w) \geq \frac{1}{2} w^2.$$

Then

$$\begin{aligned} |\nabla u(x)|^2 - u(x) \Delta u(x) &\leq C(\|u\|_{H^1(B_2)} + \|\Delta u\|_{L^2(B_2)}), \\ \forall x \in B_1 = \{y \in \mathbb{R}^n : |y| < 1\}, \text{ and } C \text{ depends only on } n. \end{aligned} \quad (6.17)$$

Proof. By Proposition 6.5, we have that $P = |\nabla u|^2 - u \Delta u = P(u; x)$ is subharmonic. Therefore we have

$$P(u; x) \leq \frac{1}{|B(x, r)|} \int_{B(x, r)} P(u; y) dy, \quad \forall B(x, r) \subset B_2 \quad (6.18)$$

Also, $P \leq |\nabla u|^2 + \frac{1}{2}(u^2 + (\Delta u)^2)$.

So,

$$\int_{B(x, r)} P(u; y) dy \leq \|u\|_{H^1(B_2)} + \|\Delta u\|_{L^2(B_2)}, \quad \forall B(x, r) \subset B_2 \quad (6.19)$$

Thus, for any $x \in B_1$ (since $B(x, 1) \subset B_2$), we have

$$P(u; x) \leq \frac{1}{|B_1|} (\|u\|_{H^1(B_2)} + \|\Delta u\|_{L^2(B_2)}) \quad (6.20)$$

\square

Remark 6.7. *Note that if $F(u, |\nabla u|^2, \Delta u) = \frac{1}{2}(\Delta u)^2$, we have a reduction of order result, that is, if u is a smooth and bounded entire solution of*

$$2|Hes u|^2 = (\Delta u)^2 + u \Delta^2 u \quad (6.21)$$

such that $\nabla u, \Delta u \in L^\infty(\mathbb{R}^n)$, then u satisfies $u\Delta u = |\nabla u|^2 + c$ for some $c \in \mathbb{R}$. We can see this from the proof of Proposition 6.5, where $P = |\nabla u|^2 - u\Delta u$ will be harmonic for this particular equation. Also, $|P| \leq M$ for some $M = M(\|u\|_{L^\infty(\mathbb{R}^n)}, \|\nabla u\|_{L^\infty(\mathbb{R}^n)}, \|\Delta u\|_{L^\infty(\mathbb{R}^n)}) > 0$ and thus $P \equiv \text{constant}$.

A special case of interest of Corollary 6.6 is the following

Corollary 6.8. *Let $u : B_2 \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth solution of*

$$\det(\nabla^2 u) = G(u, |\nabla u|^2) - \frac{u}{4}\Delta^2 u \quad (6.22)$$

where $G : \mathbb{R}^2 \rightarrow [0, +\infty)$

Then

$$|\nabla u(x)|^2 - u(x)\Delta u(x) \leq C(\|u\|_{H^1(B_2)} + \|\Delta u\|_{L^2(B_2)}) \quad (6.23)$$

Proof. We write $\det(\nabla^2 u) = \frac{1}{2}((\Delta u)^2 - |Hess u|^2)$ since u is defined a domain in the plane and then the proof is a consequence of Corollary 6.6 for $F = G(u, |\nabla u|^2) + (\Delta u)^2$, $G \geq 0$. \square

6.2. A Liouville theorem and a De Giorgi-type property. A direct consequence of Theorem 6.2 is the following

Corollary 6.9. *Let u be a convex entire subsolution of*

$$\sum_{i,j} a_{ij}(x, u, \nabla u, \nabla^2 u, \nabla^3 u) \Delta u_{x_i x_j} - F(x, u, \nabla u, \nabla^2 u, \nabla^3 u) = 0 \quad (6.24)$$

that satisfies assumption (6.1) with $m = 4$ and assume a_{ij} satisfy the ellipticity condition (2.6) and $F \geq 0$.

Then u is constant.

Proof. Consider $P = P(u, \nabla u, \Delta u) = (\Delta u)^2$, so as in the proof of Proposition 6.3 we calculate

$$\begin{aligned} P_{x_i x_j} &= 2\Delta u_{x_i} \Delta u_{x_j} + 2\Delta u \Delta u_{x_i x_j} \\ \Rightarrow \sum_{i,j} a_{ij} P_{x_i x_j} &\geq 2\lambda |\nabla \Delta u|^2 + F \Delta u \geq 0 \end{aligned} \quad (6.25)$$

and $P(u, \nabla u, 0) = 0$ with $\mu = 1$, so by Theorem 6.2 we obtain $\Delta u \equiv 0$ in \mathbb{R}^n and u is bounded by (6.1), so u is constant. \square

Finally, we have the following De Giorgi-type property

Proposition 6.10. *Let $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a smooth and bounded solution of*

$$F(u, \nabla u, \nabla^2 u, \nabla^3 u, \nabla^4 u) = 0 \quad (6.26)$$

such that $u_y > 0$ and assume $P = P(u, \Delta u)$ is a P -function of (6.26), such that $P_t > 0$ ($P = P(s, t)$) with $\mu = \mu(|\nabla u|)$, $\mu(t) > 0$, $\forall t > 0$.

If there exists $x_0 \in \mathbb{R}^2$ such that

$$P(u(x_0), \Delta u(x_0)) = \sup_{\mathbb{R}^n} P(u, \Delta u) < +\infty \quad (6.27)$$

then there exists a function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$u(x) = g(ax + by) \quad , \quad \text{for } a, b \in \mathbb{R} \quad (6.28)$$

Proof. Arguing as in the proof of Theorem 4.4 we obtain that

$$P(u, \Delta u) \equiv c_0 \quad , \quad \text{where } c_0 = \sup_{\mathbb{R}^n} P(u, \Delta u) \quad (6.29)$$

since $P_t > 0$ we have

$$\Delta u = f(u) \quad , \quad \text{for some } f : \mathbb{R} \rightarrow \mathbb{R} \quad (6.30)$$

and u is bounded entire solution of (6.30) such that $u_y > 0$.

Therefore, by Theorem 1.1 in [12], we conclude that

$$u(x) = g(ax + by) \quad , \quad \text{for some } g : \mathbb{R} \rightarrow \mathbb{R} \quad (6.31)$$

□

Acknowledgments: I would like to thank my advisor professors N. Alikakos and C. Makridakis for their support. Also, I would like to thank professors N. Alikakos and A. Farina for their useful suggestions that lead to various improvements.

APPENDIX A. SOME ADDITIONAL EXAMPLES OF P -FUNCTIONS AND THEIR GRADIENT BOUNDS

(1) The following example can be found in [17]. Let u be a solution of

$$\Delta u = u(k|\nabla u|^2 + \lambda e^{-cu^2})$$

$$\text{and let } P(s, t) = \begin{cases} te^{-ks^2} + \frac{\lambda}{k+c} e^{-s^2(k+c)} & , \quad k \neq -c \\ te^{cs^2} - \lambda s^2 & , \quad k = -c \end{cases} \quad (A.1)$$

Then $P = P(u, |\nabla u|^2)$ is a P -function of (A.1).

Corollary A.1. *Let u be an entire solution of*

$$\Delta u = u(k|\nabla u|^2 + \lambda e^{-cu^2}) \quad (\text{A.2})$$

that satisfy (3.1).

Then

$$|\nabla u|^2 \leq \begin{cases} -\frac{\lambda}{k+c}e^{-cu^2} & , \text{ if } \lambda(k+c) < 0 \\ \lambda u^2 e^{-cu^2} & , \text{ if } k = -c \text{ and } \lambda \geq 0 \end{cases} \quad (\text{A.3})$$

Proof. By [17], we have that

$$P(s, t) = \begin{cases} te^{-ks^2} + \frac{\lambda}{k+c}e^{-s^2(k+c)} & , k \neq -c \\ te^{cs^2} - \lambda s^2 & , k = -c \end{cases} \quad (\text{A.4})$$

is a P -function of (A.2) with $\mu(t) > 0$, $\forall t \geq 0$ and $P(s, 0) \leq 0$ in both cases since either $\lambda(k+c) < 0$ or $k = -c$ and $\lambda \geq 0$. Therefore by Theorem 3.1 we conclude that $P(u, |\nabla u|^2) \leq 0 \forall x \in \mathbb{R}^n$ and we obtain the gradient bound (A.3). \square

Remark A.2. *For $\lambda = 0$, Corollary A.1 says that $|\nabla u| \equiv 0$ and thus u is a constant. That is a Liouville-type result and can also be obtained either as an application of Theorem 4.1 or by Liouville's theorem by setting $v = g(u)$, where $g(z) = \int_0^y e^{-kz^2} dz$ and then $\Delta v = 0$ and v is bounded since u is bounded.*

(2) Consider the equation

$$\operatorname{div}(|\nabla u|^2 \nabla u) = f'(u) \quad (\text{A.5})$$

where f is such that $f \geq 0$ and $f'' \leq 0$.

This is a quasi-linear equation of the form (5.1) with $\Phi(t) = t^2$. In view of Theorem 4.1, the assumption $f'' \leq 0$ is reasonable since the entire solutions would be constant in the case where $f'' \geq 0$.

By Lemma 2.5 we have that

$$P(u, |\nabla u|^2) = |\nabla u|^2 - \frac{6}{\lambda} f(u) \quad (\text{A.6})$$

is a P -function of (A.5), where λ is the ellipticity constant.

Indeed, we calculate J in (2.11) with $A(t) = t$, $B(s) = -\frac{6}{\lambda} f(u)$

$$\begin{aligned} J &= (\lambda B''(u) + 2f''(u))|\nabla u|^2 + B'(u) \sum_{i,j,k} \frac{\partial a_{ij}}{\partial \sigma_k} (\nabla u) u_{x_i x_j} u_{x_k} + B'(u) \sum_{i,j} a_{ij} (\nabla u) u_{x_i x_j} + \frac{\lambda}{2} (B'(u))^2 \\ &\geq -4f''(u)|\nabla u|^2 - \frac{6}{\lambda} f'(u) \sum_{i,j,k} \frac{\partial a_{ij}}{\partial \sigma_k} (\nabla u) u_{x_i x_j} u_{x_k} - \frac{6}{\lambda} (f'(u))^2 + \frac{18}{\lambda} (f'(u))^2 \end{aligned}$$

where $a_{ij} = 4\sigma_i \sigma_j + 2|\sigma|^2 \delta_{ij}$ as defined in (5.2).

Therefore we have

$$\begin{aligned} \frac{\partial a_{ij}}{\partial \sigma_k} &= 4\sigma_i \delta_{jk} + 4\sigma_j \delta_{ik} + 4\sigma_k \delta_{ij} \\ \Rightarrow \sum_{i,j,k} \frac{\partial a_{ij}}{\partial \sigma_k} (\nabla u)_{x_k} u_{x_i x_j} &= 2f'(u) \end{aligned}$$

So,

$$J \geq -4f''(u)|\nabla u|^2 \geq 0$$

since $f'' \leq 0$.

In addition, $P(u, 0) = -\frac{6}{\lambda}f(u) \leq 0$, since f is non negative. Thus, by Theorem 3.1 we obtain the bound

Corollary A.3. *Let u be an entire solution of*

$$\operatorname{div}(|\nabla u|^2 \nabla u) = f'(u) \tag{A.7}$$

that satisfy (3.1), where f is such that $f \geq 0$ and $f'' \leq 0$.

Then

$$|\nabla u|^2 \leq \frac{6}{\lambda}f(u) \tag{A.8}$$

REFERENCES

- [1] Alberti, G., Ambrosio, L., Cabre, X. *On a Long-Standing Conjecture of E. De Giorgi Symmetry in 3D for General Nonlinearities and a Local Minimality Property*, Acta Applicandae Mathematicae, 65: 9-33 (2001).
- [2] Caffarelli, L., Cabré, X. *Fully Nonlinear Elliptic Equations*, American Mathematical Society, Vol 43 (1995).
- [3] Caffarelli, L., Crandall, M. *Distance Functions and Almost Global Solutions of Eikonal Equations*, Com. Partial Differential Equations, 35, no 3, 391-414 (2010).
- [4] Caffarelli, L., Garofalo, N., Segala, F. *A Gradient Bound for Entire Solutions of Quasi-Linear Equations and Its Consequences*, Communications on Pure and Applied Mathematics, Vol. XLVII, 1457-1473 (1994).
- [5] Cavaterra, C., Dipierro, S., Farina, A., Gao, Z., Valdinoci, E. *Pointwise gradient bounds for entire solutions of elliptic equations with non-standard growth conditions and general nonlinearities*, Journal of Differential Equations, 270, pp. 435-475 (2021).
- [6] Danielli, D., Garofalo, N. *Properties of entire solutions of non-uniformly elliptic equations arising in geometry and in phase transitions*, Calculus of Variations, 15, 451-491 (2002).
- [7] Dupaigne, L., *Stable solutions of Elliptic Partial Differential Equations*, Chapman and Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, 143 (2011).
- [8] Farina, A., Valdinoci, E. *Gradient bounds for anisotropic partial differential equations*, Calculus of Variations, 49, 923-936 (2013).
- [9] Farina, A., Valdinoci, E. *Pointwise estimates and rigidity results for entire solutions of nonlinear elliptic pde's*, ESAIM: Control, Optimisation and Calculus of Variations, Volume 19 no. 2, pp. 616-627 (2013).

- [10] Farina, A., Valdinoci, E. *A pointwise gradient estimate in possibly unbounded domains with nonnegative mean curvature*, Advances in Mathematics Volume 225, Issue 5, Pages 2808-2827 (2010).
- [11] Gazoulis, D. *A Relation of the Allen-Cahn equations and the Euler equations and applications of the Equipartition*, arXiv:2203.13960 (2022).
- [12] Ghoussoub, N., Gui, C. *On a conjecture of De Giorgi and some related problems*, Math. Ann. 311, 481-491 (1998).
- [13] Giaquinta, M., Hildebrandt, S. *Calculus of Variations I*, Berlin: Springer-Verlag, Heidelberg (1996).
- [14] Gilbarg, D., Trudinger, N., S. *Elliptic Partial Differential Equations of Second Order*, Berlin: Springer-Verlag, reprint of 1998 edition (2001).
- [15] Han, Q., Lin, F. *Elliptic Partial Differential Equations*, AMS, Courant Institute of Mathematical Sciences, Second Ed. (2011).
- [16] Modica, L. *A Gradient Bound and a Liouville Theorem for Nonlinear Poisson Equations*, Communications on Pure and Applied Mathematics, Vol. XXXVIII 679-684 (1985).
- [17] Payne, L., E., Philippin, G., A. *On Maximum Principles for a Class of Nonlinear Second-Order Elliptic Equations*, Journal of Differential Equations, 37, 39-48 (1980).
- [18] Payne, L., E., Philippin, G., A. *Some Maximum Principles for Nonlinear Elliptic Equations in divergence form with applications to capillary surfaces and to surfaces of constant mean curvature*, Nonlinear Analysis, Theory, Methods & Applications, Vol. 3, No. 2, pp. 193-211 (1979).
- [19] Serrin, J. *Entire solutions of nonlinear Poisson equations*, Proc. London Math. Soc. **24**, 348-366 (1972).
- [20] Sperb, R., P. *Maximum Principles and their Applications*, Zurich, Switzerland, Academic Press (1981).
- [21] Smyrnelis, P. *Gradient estimates for semilinear elliptic systems and other related results*, Proceedings of the Royal Society of Edinburgh, 145A, pp. 1313-1330 (2015).
- [22] Tolksdorf, P. *Regularity for a More General Class of Quasilinear Elliptic Equations*, Journal of Differential Equations, 51, 126-150 (1984).
- [23] Trudinger, N. S. *Fully nonlinear, uniformly elliptic equations under natural structure conditions*, Trans. Amer. Math. Soc. 278, 751-769 (1983).

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF CRETE, 70013 HERAKLION, GREECE, AND INSTITUTE OF APPLIED AND COMPUTATIONAL MATHEMATICS, FORTH, 700 13 HERAKLION, CRETE, GREECE

Email address: dgazoulis@math.uoa.gr