

Cost-reduction implicit exponential Runge–Kutta methods for highly oscillatory systems

Xianfa Hu^a, Wansheng Wang^a, Bin Wang^b, Yonglei Fang^{c,*}

^a*Department of Mathematics, Shanghai Normal University, Shanghai 200234, P.R.China*

^b*School of Mathematics and Statistics, Xi'an Jiaotong University, Xi'an 710049, Shannxi, P.R.China*

^c*School of Mathematics and Statistics, Zaozhuang University, Zaozhuang 277160, P.R. China*

Abstract

In this paper, two novel classes of implicit exponential Runge–Kutta (ERK) methods are studied for solving highly oscillatory systems. Firstly, we analyze the symplectic conditions for two kinds of exponential integrators and obtain the symplectic method. In order to effectively solve highly oscillatory problems, we try to design the highly accurate implicit ERK integrators. By comparing the Taylor series of numerical solution with exact solution, it can be verified that the order conditions of two new kinds of exponential methods are identical to classical Runge–Kutta (RK) methods, which implies that using the coefficients of RK methods, some highly accurate numerical methods are directly formulated. Furthermore, we also investigate the linear stability regions for these exponential methods. Finally, numerical results not only display the long time energy preservation of the symplectic method, but also illustrate the accuracy and efficiency of these formulated methods in comparison with standard ERK methods.

Keywords: Implicit exponential Runge–Kutta methods, symplectic conditions, order conditions, linear stability analysis, highly oscillatory systems

Mathematics Subject Classification (2000): 65L05, 65L06

1. Introduction

As is known that the classical Runge–Kutta methods have a wide range of applications in scientific computing. Especially, considering implicit numerical methods with symplecticity or symmetry, we applied the numerical methods to highly oscillatory problems or Hamiltonian systems, which not only reduce the restriction on stepsize, but also preserve the long time energy. Symplectic algorithms for Hamiltonian systems appeared in 1980s, and the earliest significant contribution to this field were due to Feng Kang (see [12, 13]). It is also worth noting the earlier important work on symplectic integration by J. M. Sanz-Serna, who first found and analysed symplectic Runge-Kutta schemes for Hamiltonian systems (see [29]). Symplectic exponential Runge–Kutta methods for solving Hamiltonian systems were proposed by Mei et al. [28]. Symplectic exponential Runge–Kutta methods show better performance than standard symplectic Runge–Kutta schemes. However those symplectic exponential Runge–Kutta methods are expensive because the coefficients of these exponential integrators are strongly dependent on the evaluations of matrix exponentials. Consequently, we try to design two novel classes of implicit exponential Runge–Kutta methods, which can reduce the computational cost to some extent, we focus our attention on symplectic or highly accurate exponential Runge–Kutta methods.

*Corresponding author

Email addresses: zzxyhxf@163.com (Xianfa Hu), w.s.wang@163.com (Wansheng Wang), wangbinmaths@xjtu.edu.cn (Bin Wang), ylfangmath@163.com (Yonglei Fang)

In this work, we consider the initial problems of first-order differential equations

$$\begin{cases} y'(t) + My(t) = f(y(t)), & t \in [t_0, t_{\text{end}}], \\ y(t_0) = y_0, \end{cases} \quad (1)$$

where the matrix $M \in \mathbb{R}^{m \times m}$ is symmetric positive definite or skew-Hermitian with eigenvalues of large modulus. Problems of the form (1) arise frequently in a variety of applied science such as quantum mechanics, flexible mechanics, and electrodynamics. It's true that some highly oscillatory problems (see, e.g. [20, 32]), Schrödinger equations (see, e.g. [3, 6, 35]) and KdV equations (see, e.g. [36]) can be converted into (1) with appropriate spatial discretization. The exact solution of problem (1) is obtained by the Volterra integral formula

$$y(t_0 + h) = e^{-hM}y(t_0) + h \int_0^1 e^{-(1-\tau)hM} f(y(t_0 + h\tau))d\tau. \quad (2)$$

As we known, it's a challenge for effectively solving the problem (1) once it has the stiffness matrix M . In formula (2), the e^{-hM} is usually the matrix exponential function, and exponential integrators can exactly integrate the linear equation $y'(t) + My(t) = 0$, which indicates exponential integrators have unique advantages for solving stiff or highly oscillatory problems than non-exponential integrators. Exponential Runge–Kutta methods have the relatively complete theory and wide applications [11, 16, 17, 18, 19, 20, 22, 23, 24, 25, 28, 33]. It is worth mentioning that extended Runge-Kutta-Nyström (ERKN) methods [9, 10, 31, 37, 38, 39, 40], as the exponential integrators, which are formulated for effectively solving second-order oscillatory systems.

On the other hands, (1) frequently possesses some important geometrical or physical properties. When $f(y) = J^{-1}\nabla U(y)$ and $-M = J^{-1}Q$, with the skew-symmetric

$$J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix},$$

where $U(y)$ is a smooth potential function, Q is a symmetric matrix and I is the identity matrix, the problem (1) can be converted into a Hamiltonian system. It is well known that the symplectic methods not only preserve the symplecticity of the initial systems, but also possess the long time energy preservation and stability [12, 15]. Owing to this, our study starts by the research of symplectic methods and further explores the highly accurate and effective exponential algorithms.

It is noted that the study of standard exponential Runge-Kutta (ERK) methods is based on the stiff-order conditions [17, 18, 19, 20]. However, as claimed by Berland et al. in [2], the stiff-order conditions are relatively strict. In this paper, our study is related to the classical order (nonstiff). We have presented that two new kinds of explicit ERK methods up to order four, which reduce to classical Runge-Kutta (RK) methods once $M \rightarrow \mathbf{0}$ and their coefficients are independent of matrix-valued exponentials [21, 35]. In what follows, we will study the implicit ERK methods.

The paper is organized as follows. In Section 2, we investigate the symplectic conditions for the simplified version of ERK (SVERK) and modified version of ERK (MVERK) methods respectively, and verify the existence of the symplectic method. In Section 3, we derive the order conditions of implicit SVERK and MVERK methods, further obtain the highly accurate implicit ERK algorithms. In Section 4, the linear stability regions of explicit and implicit SVERK and MVERK methods are analyzed. In Section 5, numerical experiments illustrate the structure-preserving property of the symplectic method, and show the comparable accuracy and efficiency of these implicit ERK methods. We make a conclusion for this paper in last section.

2. The symplectic conditions for two new classes of ERK methods

In our previous work, we have formulated the modified and simplified versions of explicit ERK methods for solving stiff or highly oscillatory problems, and shown the convergence of these two new explicit exponential methods. Also, we have pointed out that the internal stages and update of the simplified version of ERK methods preserve some properties of matrix exponentials, and the modified version of ERK methods inherit the internal stages and modify the update of classical RK methods, but their coefficients are independent of matrix exponentials.

Definition 2.1. ([35]) An s -stage SVERK method for the numerical integration (1) is defined as

$$\begin{cases} Y_i = e^{-c_i h M} y_0 + h \sum_{j=1}^s a_{ij} f(Y_j), \quad i = 1, \dots, s, \\ y_1 = e^{-h M} y_0 + h \sum_{i=1}^s b_i f(Y_i) + w_s(z), \end{cases} \quad (3)$$

where a_{ij}, b_i are real constants for $i, j = 1, \dots, s$, $Y_i \approx y(t_0 + c_i h)$ for $i = 1, \dots, s$, $w_s(z)$ depends on h, M , and $w_s(z) \rightarrow 0$ when $M \rightarrow \mathbf{0}$.

Definition 2.2. ([35]) An s -stage MVERK method for the numerical integration (1) is defined as

$$\begin{cases} \bar{Y}_i = y_0 + h \sum_{j=1}^s \bar{a}_{ij} (-M \bar{Y}_j + f(\bar{Y}_j)), \quad i = 1, \dots, s, \\ \bar{y}_1 = e^{-h M} y_0 + h \sum_{i=1}^s \bar{b}_i f(\bar{Y}_i) + \bar{w}_s(z), \end{cases} \quad (4)$$

where \bar{a}_{ij}, \bar{b}_i are real constants for $i, j = 1, \dots, s$, $\bar{Y}_i \approx y(t_0 + \bar{c}_i h)$ for $i = 1, \dots, s$, $\bar{w}_s(z)$ is related to h and M , and $\bar{w}_s(z) \rightarrow 0$ once $M \rightarrow \mathbf{0}$.

In fact, the $w_s(z)$ and $\bar{w}_s(z)$ also depend on the term $f(\cdot)$ and initial value y_0 once we consider the order of the SVERK and MVERK methods which satisfies $p \geq 1$. However, the $w_s(z)$ and $\bar{w}_s(z)$ are independent of matrix-valued exponentials. We have presented that the same order of these two new exponential methods share the same $w_s(z)$ or $\bar{w}_s(z)$, and $w_s(z)$ is different from $\bar{w}_s(z)$ when $p \geq 3$ in [21, 35]. The SVERK method (3) and MVERK method (4) can exactly integrate the first-order homogeneous linear system

$$y'(t) = -M y(t), \quad y(0) = y_0, \quad (5)$$

with the exact solution

$$y(t) = e^{-tM} y_0.$$

The SVERK method (3) can be displayed by the following Butcher Tableau

$$\begin{array}{c|c|c} c & \mathbf{e}^{-c h M} & A \\ \hline e^{-h M} & w_s(z) & b^\top \end{array} = \begin{array}{c|c|c} c_1 & e^{-c_1 h M} & a_{11} \quad \cdots \quad a_{1s} \\ \vdots & \vdots & \vdots \quad \vdots \quad \vdots \\ c_s & e^{-c_s h M} & a_{s1} \quad \cdots \quad a_{ss} \\ \hline e^{-h M} & w_s(z) & b_1 \quad \cdots \quad b_s \end{array} \quad (6)$$

where $c_i = \sum_{j=1}^s a_{ij}$ for $i = 1, \dots, s$. Similarly, the MVERK method (4) can be expressed in the Butcher tableau

$$\begin{array}{c|c|c} \bar{c} & \mathbf{I} & \bar{A} \\ \hline e^{-h M} & \bar{w}_s(z) & \bar{b}^\top \end{array} = \begin{array}{c|c|c} \bar{c}_1 & I & \bar{a}_{11} \quad \cdots \quad \bar{a}_{1s} \\ \vdots & \vdots & \vdots \quad \vdots \quad \vdots \\ \bar{c}_s & I & \bar{a}_{s1} \quad \cdots \quad \bar{a}_{ss} \\ \hline e^{-h M} & \bar{w}_s(z) & \bar{b}_1 \quad \cdots \quad \bar{b}_s \end{array} \quad (7)$$

with $\bar{c}_i = \sum_{j=1}^s \bar{a}_{ij}$ for $i = 1, \dots, s$.

It is true that (1) becomes a Hamiltonian system when $f(y) = J^{-1} \nabla U(y)$ and $M = -J^{-1} Q$, with $U(y)$ is a smooth potential function and Q is a symmetric matrix. Thus, we consider the following Hamiltonian system

$$\begin{cases} y'(t) - J^{-1} Q y(t) = J^{-1} \nabla U(y), \quad t \in [t_0, t_{\text{end}}], \\ y(t_0) = y_0. \end{cases} \quad (8)$$

Under the assumptions of $w_s(z) = 0$ and $\bar{w}_s(z) = 0$, we will analyze the symplectic conditions for SVERK and MVERK methods at first, then we obtain the symplectic SVERK method. Since $w_s(z) \neq 0$ and $\bar{w}_s(z) \neq 0$, the symplectic conditions of the SVERK and MVERK methods can be discussed in the same way.

Theorem 2.3. *If the coefficients of an s -stage SVERK method with $w_s(z) = 0$, which satisfy the following conditions:*

$$\begin{cases} e^{-(1-c_k)hM} G_k^{-1} = G_k^{-1} (e^{-(1-c_k)hM})^\top, & k = 1, \dots, s, \\ b_k b_l = b_k a_{kl} J e^{-(1-c_k)hM} J^{-1} + b_l a_{lk} (e^{-(1-c_k)hM})^\top, & k, l = 1, \dots, s \end{cases} \quad (9)$$

where $G_k^{-1} = \nabla^2 U(y_n + h \sum_{l=1}^s a_{kl} \xi_l)$ and $\xi_k = f(y_n + h \sum_{l=1}^s a_{kl} \xi_l)$, then the SVERK method is symplectic.

Proof. A numerical method is said to be symplectic if the numerical solution y_{n+1} satisfies $(\frac{\partial y_{n+1}}{\partial y_0})^\top J (\frac{\partial y_{n+1}}{\partial y_0}) = (\frac{\partial y_n}{\partial y_0})^\top J (\frac{\partial y_n}{\partial y_0})$. Under the assumption $w_s(z) = 0$, we rewrite the SVERK method as

$$\begin{cases} \xi_k = f(e^{-c_k h M} y_n + h \sum_{l=1}^s a_{kl} \xi_l), & i = 1, \dots, s, \\ y_{n+1} = e^{-hM} y_n + h \sum_{k=1}^s b_k \xi_k. \end{cases} \quad (10)$$

Letting $\Xi_k = \frac{\partial \xi_k}{\partial y_0}$, $\Psi_{n+1} = \frac{\partial y_{n+1}}{\partial y_0}$ and $G_k = \nabla^2 U(e^{-c_k h M} y_n + h \sum_{l=1}^s a_{kl} \xi_l)$, $k = 1, \dots, s$. Furthermore, assuming the symmetric matrix G_1, \dots, G_s is nonsingular. Applying (10) to a Hamiltonian system (8), and the derivative of y_{n+1} with respect to y_0 is obtained by

$$\Psi_{n+1} = e^{-hM} \Psi_n + h \sum_{k=1}^s b_k \Xi_k. \quad (11)$$

It is easy to see that

$$\begin{aligned} \Psi_{n+1}^\top J \Psi_{n+1} &= \left(e^{-hM} \Psi_n + h \sum_{k=1}^s b_k \Xi_k \right)^\top J \left(e^{-hM} \Psi_n + h \sum_{k=1}^s b_k \Xi_k \right) \\ &= (e^{-hM} \Psi_n)^\top J (e^{-hM} \Psi_n) + h \sum_{k=1}^s b_k (e^{-hM} \Psi_n)^\top J \Xi_k + h \sum_{k=1}^s b_k \Xi_k^\top J e^{-hM} \Psi_n + h^2 \sum_{k=1}^s \sum_{l=1}^s b_k b_l \Xi_k^\top J \Xi_l. \end{aligned} \quad (12)$$

From the representation of (10), we give that

$$\Xi_k = J^{-1} G_k \left(e^{-c_k h M} \Psi_n + h \sum_{l=1}^s a_{kl} \Xi_l \right), \quad k = 1, \dots, s, \quad (13)$$

thus

$$\Psi_n = e^{c_k h M} G_k^{-1} J \Xi_k - h \sum_{l=1}^s a_{kl} e^{c_k h M} \Xi_l, \quad k = 1, \dots, s. \quad (14)$$

According to (14), we have

$$\begin{aligned} h \sum_{k=1}^s b_k (e^{-hM} \Psi_n)^\top J \Xi_k &= h \sum_{l=1}^s b_l \left(e^{c_l h M} G_l^{-1} J \Xi_l - h \sum_{k=1}^s a_{lk} e^{c_l h M} \Xi_k \right)^\top (e^{-hM})^\top J \Xi_l \\ &= h \sum_{k=1}^s b_k \Xi_k^\top J^\top G_k^{-1} (e^{-(1-c_k)hM})^\top J \Xi_k - h^2 \sum_{k=1}^s \sum_{l=1}^s b_l a_{lk} \Xi_k^\top (e^{-(1-c_k)hM})^\top J \Xi_l, \end{aligned} \quad (15)$$

and

$$\begin{aligned}
h \sum_{k=1}^s b_k \Xi_k^\top J e^{-hM} \Psi_n &= h \sum_{k=1}^s b_k \Xi_k^\top J e^{-hM} \left(e^{c_k h M} G_k^{-1} J \Xi_k - h \sum_{l=1}^s a_{kl} e^{c_k h M} \Xi_l \right) \\
&= h \sum_{k=1}^s b_k \Xi_k^\top J e^{-(1-c_k)hM} G_k^{-1} J \Xi_k - h^2 \sum_{k=1}^s \sum_{l=1}^s b_k a_{kl} \Xi_k^\top J e^{-(1-c_k)hM} \Xi_l.
\end{aligned} \tag{16}$$

Inserting (15) and (16) into (12) yields

$$\begin{aligned}
\Psi_{n+1}^\top J \Psi_{n+1} &= \Psi_n^\top J \Psi_n + h \sum_{k=1}^s b_k \Xi_k^\top \left(J^\top G_k^{-1} (e^{-(1-c_k)hM})^\top + J e^{-(1-c_k)hM} G_k^{-1} \right) J \Xi_k \\
&\quad - h^2 \sum_{k=1}^s \sum_{l=1}^s b_l a_{lk} \Xi_k^\top (e^{-(1-c_k)hM})^\top J \Xi_l - h^2 \sum_{k=1}^s \sum_{l=1}^s b_k a_{kl} \Xi_k^\top J e^{-(1-c_k)hM} \Xi_l \\
&\quad + h^2 \sum_{k=1}^s \sum_{l=1}^s b_k b_l \Xi_k^\top J \Xi_l \\
&= \Psi_n^\top J \Psi_n + h \sum_{k=1}^s b_k \Xi_k^\top J \left(e^{-(1-c_k)hM} G_k^{-1} - G_k^{-1} (e^{-(1-c_k)hM})^\top \right) J \Xi_k \\
&\quad + h^2 \sum_{k=1}^s \sum_{l=1}^s \Xi_k^\top (b_k b_l - b_k a_{kl} J e^{-(1-c_k)hM} J^{-1} - b_l a_{lk} (e^{-(1-c_k)hM})^\top) J \Xi_l.
\end{aligned} \tag{17}$$

As the coefficients of the method satisfy the conditions (9), the direct calculation gives

$$\Psi_{n+1} J \Psi_{n+1} = \Psi_n J \Psi_n.$$

Therefore the SVERK method with the coefficients satisfying (9) is symplectic. The proof is complete. \square

We remark that the symplectic conditions of SVERK methods with $w_s(z) = 0$ reduce to the symplectic conditions of classical RK methods once $M \rightarrow \mathbf{0}$, and it's obvious that taking $s = 1$ and $c_1 = 1$, we give the first-order symplectic SVERK method

$$y_1 = e^{-hM} y_0 + h f(y_1). \tag{18}$$

The symplectic conditions of (3) with $w_s(z) \neq 0$ can be analyzed by the same way, unfortunately, there is no existing the symplectic method when we consider the SVERK method with order $p \geq 2$.

The next theorem will present the symplectic conditions of the MVERK methods with $\bar{w}_s(z) = 0$.

Theorem 2.4. *If the coefficients of an s -stage MVERK method with $\bar{w}_s(z) = 0$, which satisfy the following conditions:*

$$\begin{cases} D_i^\top J e^{-hM} + (e^{-hM})^\top J D_i = 0, \quad i = 1, \dots, s \\ \bar{b}_i \bar{b}_j = \bar{b}_i \bar{a}_{ij} J e^{-hM} J^{-1} + \bar{b}_j \bar{a}_{ji} e^{-hM}, \quad i, j = 1, \dots, s, \\ \bar{b}_j \bar{a}_{ji} M^\top (e^{-hM})^\top J D_j + \bar{b}_i \bar{a}_{ij} D_i^\top J e^{-hM} M = 0, \quad i, j = 1, \dots, s, \end{cases} \tag{19}$$

where $D_i = \frac{\partial f(Y_i)}{\partial y_0}$, then the MVERK method is symplectic.

Proof. If the numerical method is symplectic, the numerical solution \bar{y}_{n+1} satisfies

$$\left(\frac{\partial \bar{y}_{n+1}}{\partial y_0} \right)^\top J \left(\frac{\partial \bar{y}_{n+1}}{\partial y_0} \right) = \left(\frac{\partial \bar{y}_n}{\partial y_0} \right)^\top J \left(\frac{\partial \bar{y}_n}{\partial y_0} \right).$$

Setting $D_i = \frac{\partial f(Y_i)}{\partial y_0}$, $X_i = \frac{\partial \bar{Y}_i}{\partial y_0}$ and $\Psi_{n+1} = \frac{\partial \bar{y}_{n+1}}{y_0}$. When we apply the MVERK method (4) with $\bar{w}_s(z) = 0$ to a Hamiltonian system (8), the derivatives of this scheme (4) with respect to y_0 are

$$\begin{cases} X_i = \frac{\partial \bar{Y}_i}{\partial y_0} = \Psi_n + h \sum_{j=1}^s \bar{a}_{ij}(-MX_j + D_j X_j), \\ \Psi_{n+1} = e^{-hM} \Psi_n + h \sum_{i=1}^s \bar{b}_i D_i X_i. \end{cases} \quad (20)$$

Then, we have

$$\begin{aligned} \Psi_{n+1}^\top J \Psi_{n+1} &= \left(e^{-hM} \Psi_n + h \sum_{i=1}^s \bar{b}_i D_i X_i \right)^\top J \left(e^{-hM} \Psi_n + h \sum_{i=1}^s \bar{b}_i D_i X_i \right) \\ &= (e^{-hM} \Psi_n)^\top J (e^{-hM} \Psi_n) + h \sum_{i=1}^s \bar{b}_i (e^{-hM} \Psi_n)^\top J D_i X_i + h \sum_{i=1}^s \bar{b}_i (D_i X_i)^\top J e^{-hM} \Psi_n + h^2 \sum_{i=1}^s \sum_{j=1}^s \bar{b}_i \bar{b}_j (D_i X_i)^\top J D_j X_j. \end{aligned} \quad (21)$$

Using the first formula of (20), we obtain

$$X_i^\top (e^{-hM})^\top J D_i X_i = \Psi_n^\top (e^{-hM})^\top J D_i X_i + h \sum_{j=1}^s \bar{a}_{ij}(-MX_j + D_j X_j)^\top (e^{-hM})^\top J D_i X_i, \quad (22)$$

$$(D_i X_i)^\top J e^{-hM} X_i = (D_i X_i)^\top J e^{-hM} \Psi_n + h \sum_{j=1}^s \bar{a}_{ij} (D_i X_i)^\top J e^{-hM} (-MX_j + D_j X_j),$$

thus

$$\Psi_n^\top (e^{-hM})^\top J D_i X_i = X_i^\top (e^{-hM})^\top J D_i X_i - h \sum_{j=1}^s \bar{a}_{ij}(-MX_j + D_j X_j)^\top (e^{-hM})^\top J D_i X_i, \quad (23)$$

$$(D_i X_i)^\top J e^{-hM} \Psi_n = (D_i X_i)^\top J e^{-hM} X_i - h \sum_{j=1}^s \bar{a}_{ij} (D_i X_i)^\top J e^{-hM} (-MX_j + D_j X_j).$$

Inserting (23) into (21) leads to

$$\begin{aligned} \Psi_{n+1}^\top J \Psi_{n+1} &= \Psi_n^\top J \Psi_n + h \sum_{i=1}^s \bar{b}_i X_i^\top (e^{-hM})^\top J D_i X_i - h^2 \sum_{i=1}^s \sum_{j=1}^s \bar{b}_j \bar{a}_{ji} (-MX_i + D_i X_i)^\top (e^{-hM})^\top J D_j X_j \\ &\quad + h \sum_{i=1}^s \bar{b}_i (D_i X_i)^\top J e^{hM} X_i - h^2 \sum_{i=1}^s \sum_{j=1}^s \bar{b}_i \bar{a}_{ij} (D_i X_i)^\top J S (-MX_j + D_j X_j) \\ &\quad + h^2 \sum_{i=1}^s \sum_{j=1}^s \bar{b}_i \bar{b}_j (D_i X_i)^\top J (D_j X_j) \\ &= \Psi_n^\top J \Psi_n + h \sum_{i=1}^s \bar{b}_i X_i^\top (D_i^\top J e^{hM} + (e^{hM})^\top J D_i) X_i \\ &\quad + h^2 \sum_{i=1}^s \sum_{j=1}^s (D_i X_i)^\top (\bar{b}_i \bar{b}_j - \bar{b}_j \bar{a}_{ij} (e^{-hM})^\top - \bar{b}_i \bar{a}_{ij} J S J^{-1}) J D_j X_j \end{aligned}$$

$$+h^2 \sum_{i=1}^s \sum_{j=1}^s X_i^\top (\bar{b}_j \bar{a}_{ji} M^\top (e^{-hM})^\top J D_j + \bar{b}_i \bar{a}_{ij} D_i^\top J e^{-hM} M) X_j.$$

As the coefficients of the MVERK method satisfy conditions (19), it can be easily verified that

$$\Psi_{n+1}^\top J \Psi_{n+1} = \Psi_n^\top J \Psi_n,$$

so the MVERK method is symplectic. The proof is complete. \square

Remark 2.5. *The Theorem 2.4 presents the symplectic conditions of MVERK methods with $w_s(z) = 0$. However, when b_i ($i = 1, \dots, s$) are constants, the second formula of (19) can never be satisfied. The symplectic conditions of MVERK methods with $w_s(z) \neq 0$ are obtained by the same way, unfortunately, we have the same result that some formulas of symplectic conditions for MVERK methods with $w_s(z) \neq 0$ can never be satisfied.*

Corollary 2.6. *There does not exist any symplectic MVERK methods.*

3. Highly accurate implicit ERK methods

In the previous section, we have analyzed the symplectic conditions for the SVERK and MVERK methods. However, the symplectic SVERK method only has order one, and there is no existing symplectic MVERK methods. In practice, we need some highly accurate and effective numerical methods, thus the high-order implicit SVERK (IMSVERK) and implicit MVERK (IMMVERK) methods are studied respectively in this section.

We consider the one-stage IMSVERK method with $w_2(z) = -\frac{h^2 M f(y_0)}{2!}$

$$\begin{cases} Y_1 = e^{-c_1 h M} y_0 + h a_{11} f(Y_1), \\ y_1 = e^{-h M} y_0 + h b_1 f(Y_1) - \frac{h^2 M f(y_0)}{2!}, \end{cases} \quad (24)$$

and the one-stage IMMVERK method with $\bar{w}_2(z) = -\frac{h^2 M f(y_0)}{2!}$

$$\begin{cases} \bar{Y}_1 = y_0 + h \bar{a}_{11} (-M \bar{Y}_1 + f(\bar{Y}_1)), \\ \bar{y}_1 = e^{-h M} y_0 + h \bar{b}_1 f(\bar{Y}_1) - \frac{h^2 M f(y_0)}{2!}. \end{cases} \quad (25)$$

It is noted that our study is based on the classical order (nonstiff). A numerical method is said to be of order p if the Taylor series of numerical solution y_1 or \bar{y}_1 and exact solution $y(t_0 + h)$ coincides up to h^p about y_0 . To simplify the calculation, we denote $g(t_0) = -M y(t_0) + f(y(t_0))$, the Taylor series for exact solution $y(t_0 + h)$ is given by

$$\begin{aligned} y(t_0 + h) &= y(t_0) + h y'(t_0) + \frac{h^2}{2!} y''(t_0) + \frac{h^3}{3!} y'''(t_0) + \frac{h^4}{4!} y^{(4)}(t_0) + \mathcal{O}(h^5) + \dots \\ &= y(t_0) + h g(t_0) + \frac{h^2}{2!} (-M + f'_y(y_0)) g(t_0) + \frac{h^3}{3!} (M^2 g(t_0) + (-M + f'_y(y(t_0))) f''_y(y(t_0)) g(t_0) - f'_y(y(t_0)) M g(t_0) \\ &\quad + f''_{yy}(y(t_0))(g(t_0), g(t_0))) + \frac{h^4}{4!} \left(-M^3 g(t_0) + M^2 f'_y(y(t_0)) g(t_0) - M f'_y(y(t_0)) (-M + f'_y(y(t_0))) g(t_0) \right. \\ &\quad \left. - M f''_{yy}(y(t_0))(g(t_0), g(t_0)) + f'''_{yyy}(y(t_0))(g(t_0), g(t_0), g(t_0)) + 3 f''_{yy}(y(t_0))((-M + f'_y(y(t_0))) g(t_0), g(t_0)) \right. \\ &\quad \left. + f'_y(y(t_0))(-M + f'_y(y(t_0)))(-M + f'_y(y(t_0))) g(t_0) + f'_y(y(t_0)) f''_{yy}(y(t_0))(g(t_0), g(t_0)) \right) + \mathcal{O}(h^5) + \dots \end{aligned}$$

Under $y_0 = y(t_0)$, the Taylor series for numerical solution y_1 and \bar{y}_1 are

$$\begin{aligned}
y_1 &= (I - hM + \frac{h^2 M^2}{2!} + \mathcal{O}(h^3))y_0 + hb_1 f(y_0 - c_1 hM y_0 + a_{11} h f(y_0) + \mathcal{O}(h^2)) - \frac{h^2 M f(y_0)}{2!} \\
&= y_0 - hM y_0 + \frac{h^2 M^2}{2!} y_0 + hb_1 f(y_0) + h^2 b_1 c_1 f'_y(y_0) (-M y_0 + f(y_0)) - \frac{h^2 M f(y_0)}{2!} + \mathcal{O}(h^3) \\
&= y_0 - hM y_0 + hb_1 f(y_0) + \frac{h^2 M^2}{2!} y_0 + h^2 b_1 c_1 f'_y(y_0) (-M y_0 + f(y_0)) - \frac{h^2 M f(y_0)}{2!} + \mathcal{O}(h^3),
\end{aligned} \tag{26}$$

and

$$\begin{aligned}
\bar{y}_1 &= (I - hM + \frac{h^2 M^2}{2!} + \mathcal{O}(h^3))y_0 + h\bar{b}_1 [f(y_0) + h\bar{a}_{11} f'_y(y_0) (-M y_0 + f(y_0) + \mathcal{O}(h))] - \frac{h^2 M f(y_0)}{2!} \\
&= y_0 - hM y_0 + h\bar{b}_1 f(y_0) + \frac{h^2 M^2}{2!} y_0 + h^2 \bar{b}_1 \bar{a}_{11} f'_y(y_0) (-M y_0 + f(y_0)) - \frac{h^2 M f(y_0)}{2!} + \mathcal{O}(h^3).
\end{aligned} \tag{27}$$

Compared with the Taylor series of exact solution $y(t_0 + h)$, if we consider the second-order implicit ERK method with one stage, we then have $b_1 = \bar{b}_1 = 1$ and $a_{11} = \bar{a}_{11} = \frac{1}{2}$. Therefore, the unique second-order IMSVERK method with one stage is given by

$$\begin{cases} Y_1 = e^{-\frac{1}{2}hM} y_0 + \frac{h}{2} f(Y_1), \\ y_1 = e^{-hM} y_0 + h f(Y_1) - \frac{h^2 M f(y_0)}{2!}, \end{cases} \tag{28}$$

which can be denoted by the Butcher tableau

$$\begin{array}{c|c|c} \frac{1}{2} & e^{-\frac{1}{2}hM} & \frac{1}{2} \\ \hline e^{-hM} & w_2(z) & 1 \end{array} \tag{29}$$

and the unique second-order IMMVERK method with one stage is shown as

$$\begin{cases} \bar{Y}_1 = y_0 + \frac{h}{2} (-M \bar{Y}_1 + f(\bar{Y}_1)), \\ y_1 = e^{-hM} y_0 + h f(\bar{Y}_1) - \frac{h^2 M f(y_0)}{2!}, \end{cases} \tag{30}$$

which also can be indicated by the Butcher tableau

$$\begin{array}{c|c|c} \frac{1}{2} & I & \frac{1}{2} \\ \hline e^{-hM} & \bar{w}_2(z) & 1 \end{array} \tag{31}$$

Hochbruck and Ostermann [19] have presented that using the collocation code $c_1 = \frac{1}{2}$, then the second-order (stiff) ERK method with one stage can be denoted by

$$\begin{array}{c|c} \frac{1}{2} & \frac{1}{2} \varphi_1(-\frac{1}{2}hM) \\ \hline & \varphi_1(-hM) \end{array} \tag{32}$$

with

$$\varphi_{ij}(-hM) = \varphi_i(-c_j hM) = \int_0^1 e^{-(1-\tau)c_j hM} \frac{\tau^{i-1}}{(i-1)!} d\tau. \tag{33}$$

Now, we consider the fourth-order IMSVERK and IMMVERK methods with two stages. The order conditions of the fourth-order implicit RK method with two stages are

$$\begin{aligned}
\bullet & \quad b_1 + b_2 & & = 1, \\
\downarrow & \quad b_1c_1 + b_2c_2 & & = \frac{1}{2}, \\
\forall & \quad b_1c_1^2 + b_2c_2^2 & & = \frac{1}{3}, \\
\downarrow & \quad b_1(a_{11}c_1 + a_{12}c_2) + b_2(a_{21}c_1 + a_{22}c_2) & & = \frac{1}{6}, \\
\forall & \quad b_1c_1^3 + b_2c_2^3 & & = \frac{1}{4}, \\
\downarrow & \quad b_1c_1(a_{11}c_1 + a_{12}c_2) + b_2c_2(a_{21}c_1 + a_{22}c_2) & & = \frac{1}{8}, \\
\forall & \quad b_1(a_{11}c_1^2 + a_{12}c_2^2) + b_2(a_{21}c_1^2 + a_{22}c_2^2) & & = \frac{1}{12}, \\
\downarrow & \quad (b_1a_{11} + b_2a_{21})(a_{11}c_1 + a_{12}c_2) + (b_1a_{12} + b_2a_{22})(a_{21}c_1 + a_{22}c_2) & & = \frac{1}{24}.
\end{aligned} \tag{34}$$

Under the assumptions $c_1 = a_{11} + a_{12}$ and $c_2 = a_{21} + a_{22}$, there exists a unique solution [15, 14]. The following theorem will state that the order conditions of fourth-order IMSVERK methods with two stages are identical to (34).

Theorem 3.1. *Assume the coefficients of the two-stage IMSVERK method with $w_4(z)$*

$$\begin{cases} Y_1 = e^{-c_1 h M} y_0 + h [a_{11} f(Y_1) + a_{12} f(Y_2)], \\ Y_2 = e^{-c_2 h M} y_0 + h [a_{21} f(Y_1) + a_{22} f(Y_2)], \\ y_1 = e^{-h M} y_0 + h (b_1 f(Y_1) + b_2 f(Y_2)) + w_4(z), \end{cases} \tag{35}$$

where

$$\begin{aligned}
w_4(z) = & -\frac{h^2}{2!} M f(y_0) + \frac{h^3}{3!} \left((M - f'_y(y_0)) M f(y_0) - M f'_y(y_0) g(y_0) \right) + \frac{h^4}{4!} \left((-M + f'_y(y_0)) M^2 f(y_0) + M^2 f'_y(y_0) g(y_0) \right. \\
& - M f''_{yy}(y_0) (g(y_0), g(y_0)) - M f'_y(y_0) (-M + f'_y(y_0)) g(y_0) - f'_y(y_0) M f'_y(y_0) g(y_0) - f'_y(y_0) f'_y(y_0) M f(y_0) \\
& \left. + 3 f''_{yy}(y_0) (-M f(y_0), g(y_0)) \right),
\end{aligned}$$

which satisfying (34), then the IMSVERK method has order four.

Proof. The Taylor series expansion about y_0 of y_1 agrees with $y(t_0 + h)$ up to the term h^p , the numerical

method is said to be of order p . Then, the Taylor series expansion for numerical solution y_1 is shown by

$$\begin{aligned}
y_1 &= e^{-hM}y_0 + hb_1f\left((I - c_1hM + \frac{(c_1hM)^2}{2!} - \frac{(c_1hM)^3}{3!} + O(h^4))y_0 + ha_{11}f(Y_1) + ha_{12}f(Y_2)\right) + hb_2 \\
&\cdot f\left((I - c_2hM + \frac{(c_2hM)^2}{2!} - \frac{(c_2hM)^3}{3!} + O(h^4))y_0 + ha_{21}f(Y_1) + ha_{22}f(Y_2)\right) + w_4(z) + O(h^5) \\
&= (I - hM + \frac{h^2M^2}{2} - \frac{h^3M^3}{3!} + \frac{h^4M^4}{4!} + O(h^5))y_0 + h(b_1 + b_2)f(y_0) + h^2b_1f'_y(y_0)(-c_1My_0 + a_{11}f(Y_1) \\
&+ a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0 - \frac{h^2(c_1M)^3}{3!}y_0 + O(h^3)) + h^2b_2f'_y(y_0)(-c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2) \\
&+ \frac{h(c_2M)^2}{2}y_0 - \frac{h^2(c_2M)^3}{3!}y_0 + O(h^3)) + \frac{h^3}{2}b_1f''_{yy}(y_0)(-c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0, \\
&-c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0) + \frac{h^3}{2}b_2f''_{yy}(y_0)(-c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2) \\
&+ \frac{h(c_2M)^2}{2}y_0, -c_2My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_2M)^2}{2}y_0) + \frac{h^4}{3!}b_1f'''_{yyy}(y_0)(-c_1My_0 + a_{11}f(Y_1) \\
&+ a_{12}f(Y_2), -c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2), -c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2)) + \frac{h^4}{3!}b_2f'''_{yyy}(y_0)(-c_2My_0 \\
&+ a_{21}f(Y_1) + a_{22}f(Y_2), -c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2), -c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2)) + w_4(z) + O(h^5).
\end{aligned} \tag{36}$$

Under the appropriate assumptions $c_i = \sum_{j=1}^s a_{ij}$ for $i, j = 1, \dots, s$, we ignore the term $O(h^5)$ and obtain

$$\begin{aligned}
y_1 &= e^{-hM}y_0 + h(b_1 + b_2)f(y_0) + h^2b_1f'_y(y_0)(-c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0 - \frac{h^2(c_1M)^3}{3!}y_0) \\
&+ h^2b_2f'_y(y_0)(-c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2) + \frac{h(c_2M)^2}{2}y_0 - \frac{h^2(c_2M)^3}{3!}y_0) \\
&+ \frac{h^3}{2}b_1f''_{yy}(y_0)(-c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0, -c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0) \\
&+ \frac{h^3}{2}b_2f''_{yy}(y_0)(-c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2) + \frac{h(c_2M)^2}{2}y_0, -c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2) + \frac{h(c_2M)^2}{2}y_0) \\
&+ \frac{h^4}{3!}b_1f'''_{yyy}(y_0)(c_1(-My_0 + f(y_0)), c_1(-My_0 + f(y_0)), c_1(-My_0 + f(y_0))) \\
&+ \frac{h^4}{3!}b_2f'''_{yyy}(y_0)(c_2(-My_0 + f(y_0)), c_2(-My_0 + f(y_0)), c_2(-My_0 + f(y_0))) + w_4(z) + O(h^5).
\end{aligned} \tag{37}$$

Inserting Y_1 and Y_2 of (35) into $f(Y_1)$ and $f(Y_2)$ yields to

$$\begin{aligned}
a_{11}f(Y_1) &= a_{11}f(y_0 - c_1hMy_0 + ha_{11}f(Y_1) + ha_{12}f(Y_2) + \frac{(c_1hM)^2}{2!}y_0) \\
&= a_{11}f(y_0) + hf'_y(y_0)(-c_1My_0 + a_{11}f(y_0) + ha_{11}f'_y(y_0)(-c_1My_0 + a_{11}f(y_0) + a_{12}f(y_0)) + a_{12}f(y_0)) \\
&\quad + ha_{12}f'_y(y_0)(-c_2My_0 + a_{21}f(y_0) + a_{22}f(y_0)) + \frac{h(c_1M)^2}{2}y_0 + \frac{h^2}{2}a_{11}c_1^2f''_{yy}(y_0)(g(y_0), g(y_0)) \\
&= a_{11}f(y_0) + ha_{11}c_1f'_y(y_0)g(y_0) + h^2(a_{11}^2c_1 + a_{11}a_{12}c_2)f'_y(y_0)f'_y(y_0)g(y_0) + \frac{h^2}{2}a_{11}c_1^2f'_y(y_0)M^2y_0 \\
&\quad + \frac{h^2}{2}a_{11}c_1^2f''_{yy}(y_0)(g(y_0), g(y_0)),
\end{aligned}$$

and

$$\begin{aligned}
a_{12}f(Y_2) &= a_{12}f(y_0) + ha_{12}c_2f'_y(y_0)g(y_0) + h^2(a_{12}a_{21}c_1 + a_{12}a_{22}c_2)f'_y(y_0)f'_y(y_0)g(y_0) + \frac{h^2}{2}a_{12}c_2^2f'_y(y_0)M^2y_0 \\
&\quad + \frac{h^2}{2}a_{12}c_2^2f''_{yy}(y_0)(g(y_0), g(y_0)).
\end{aligned}$$

Hence, we have

$$\begin{aligned}
h^2b_1f'(y_0)(-c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0 - \frac{h^2(c_1M)^3}{3!}y_0) &= h^2b_1c_1f'_y(y_0)g(y_0) + \frac{h^3}{2}b_1c_1^2f'_y(y_0) \\
\cdot M^2y_0 - \frac{h^4}{3!}b_1c_1^3f'_y(y_0)M^3y_0 + h^3b_1(a_{11}c_1 + a_{12}c_2)f'_y(y_0)f'_y(y_0)g(y_0) &+ h^4b_1(a_{11}^2c_1 + a_{11}a_{12}c_2 + a_{12}a_{21}c_1 + a_{12} \\
\cdot a_{22}c_2)f'_y(y_0)f'_y(y_0)f'_y(y_0)g(y_0) + \frac{h^4}{2}b_1(a_{11}c_1^2 + a_{12}c_2^2)(f'_y(y_0)f'_y(y_0)M^2y_0 &+ f'_y(y_0)f''_{yy}(y_0)(g(y_0), g(y_0))),
\end{aligned}$$

and

$$\begin{aligned}
h^2b_2f'(y_0)(-c_2My_0 + a_{21}f(Y_1) + a_{22}f(Y_2) + \frac{h(c_2M)^2}{2}y_0 - \frac{h^2(c_2M)^3}{3!}y_0) &= h^2b_2c_2f'_y(y_0)g(y_0) + \frac{h^3}{2}b_2c_2^2f'_y(y_0) \\
\cdot M^2y_0 - \frac{h^4}{3!}b_2c_2^3f'_y(y_0)M^3y_0 + h^3b_2(a_{21}c_1 + a_{22}c_2)f'_y(y_0)f'_y(y_0)g(y_0) &+ h^4b_2(a_{22}^2c_2 + a_{22}a_{21}c_1 + a_{21}a_{12}c_2 + a_{21} \\
\cdot a_{11}c_1)f'_y(y_0)f'_y(y_0)f'_y(y_0)g(y_0) + \frac{h^4}{2}b_2(a_{21}c_1^2 + a_{22}c_2^2)(f'_y(y_0)f'_y(y_0)M^2y_0 &+ f'_y(y_0)f''_{yy}(y_0)(g(y_0), g(y_0))).
\end{aligned}$$

Similarly, we consider the term $\mathcal{O}(h^3)$ and get

$$\begin{aligned}
\frac{h^3}{2}b_1f''_{yy}(y_0)(-c_1My_0 + a_{11}f(Y_1) + a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0, -c_1My_0 + a_{11}f(Y_1) &+ a_{12}f(Y_2) + \frac{h(c_1M)^2}{2}y_0) \\
= \frac{h^3}{2}b_1f''_{yy}(y_0)(c_1g(y_0) + h(a_{11}c_1 + a_{12}c_2)f'_y(y_0)g(y_0) + \frac{h(c_1M)^2}{2}y_0, &c_1g(y_0) + h(a_{11}c_1 + a_{12}c_2)f'_y(y_0)g(y_0) \\
+ \frac{h(c_1M)^2}{2}y_0)f'_y(y_0) & \\
= \frac{h^3}{2}b_1c_1^2f''_{yy}(y_0)(g(y_0), g(y_0)) + \frac{h^4}{2}b_1c_1^3f''_{yy}(y_0)(M^2y_0, g(y_0)) &+ h^4b_1(a_{11}c_1 + a_{12}c_2)c_1f''_{yy}(y_0)(f'_y(y_0)g(y_0), g(y_0)),
\end{aligned}$$

and

$$\begin{aligned} & \frac{h^3}{2} b_2 f''_{yy}(y_0) \left(-c_2 M y_0 + a_{21} f(Y_1) + a_{22} f(Y_2) + \frac{h(c_2 M)^2}{2} y_0, -c_2 M y_0 + a_{21} f(Y_1) + a_{22} f(Y_2) + \frac{h(c_2 M)^2}{2} y_0 \right) \\ &= \frac{h^3}{2} b_2 c_2^2 f''_{yy}(y_0)(g(y_0), g(y_0)) + \frac{h^4}{2} b_2 c_2^3 f''_{yy}(y_0)(M^2 y_0, g(y_0)) + h^4 b_2 (a_{21} c_1 + a_{22} c_2) c_2 f''_{yy}(y_0)(f'_y(y_0)g(y_0), g(y_0)). \end{aligned}$$

Summing up, the Taylor series for numerical solution y_1 is

$$\begin{aligned} y_1 = & y_0 - hM y_0 + h(b_1 + b_2)f(y_0) - \frac{h^2 M g(y_0)}{2!} + h^2(b_1 c_1 + b_2 c_2) f'_y(y_0)g(y_0) - \frac{h^3}{3!} f'_y(y_0) M f(y_0) \\ & + \frac{h^3}{3!} M(M - f'_y(y_0))g(y_0) + \frac{h^3}{2} (b_1 c_1^2 + b_2 c_2^2) (f'_y(y_0) M^2 y_0 + f''_{yy}(y_0)(g(y_0), g(y_0))) \\ & + h^3 (b_1(a_{11} c_1 + a_{12} c_2) + b_2(a_{21} c_1 + a_{22} c_2)) f'_y(y_0) f'_y(y_0) g(y_0) \\ & + \frac{h^4}{4!} \left(-M^3 g(y_0) + M^2 f'_y(y_0)g(y_0) - M f'_y(y_0)(-M + f'_y(y_0))g(y_0) - M f''_{yy}(y_0)(g(y_0), g(y_0)) \right. \\ & \left. - f'_y(y_0) M f'_y(y_0)g(y_0) - f'_y(y_0) f'_y(y_0) M f(y_0) + f'_y(y_0) M^2 f(y_0) + 3 f''_{yy}(y_0)(-M f(y_0), g(y_0)) \right) \\ & + h^4 (b_1 c_1^3 + b_2 c_2^3) \left(\frac{1}{3!} f'''_{yyy}(y_0)(g(y_0), g(y_0), g(y_0)) + \frac{1}{2} f''_{yy}(y_0)(M^2 y_0, g(y_0)) - \frac{1}{3!} f'_y(y_0) M^3 y_0 \right) \\ & + h^4 (b_1 c_1 (a_{11} c_1 + a_{12} c_2) + b_2 c_2 (a_{21} c_1 + a_{22} c_2)) f''_{yy}(y_0)(f'_y(y_0)g(y_0), g(y_0)) \\ & + h^4 (b_1 (a_{11}^2 c_1 + (a_{11} + a_{22}) a_{12} c_2 + a_{12} a_{21} c_1) + b_2 (a_{22}^2 c_2 + (a_{11} + a_{22}) a_{21} c_1 + a_{21} a_{12} c_2)) (f'_y(y_0))^3 g(y_0) \\ & + \frac{h^4}{2} (b_1 (a_{11} c_1^2 + a_{12} c_2^2) + b_2 (a_{21} c_1^2 + a_{22} c_2^2)) f'_y(y_0)(f''_{yy}(y_0)(g(y_0), g(y_0)) + f'_y(y_0) f'_y(y_0) M^2 y_0) + \mathcal{O}(h^5). \end{aligned} \tag{38}$$

Compared with the Taylor series for exact solution $y(t_0 + h)$, we illustrate that the IMSVERK method with coefficients satisfying the order conditions (34), has order four. The proof of the theorem is complete. \square

The Theorem 3.1 indicates that there exists the unique fourth-order IMSVERK method with two stages:

$$\left\{ \begin{aligned} Y_1 &= e^{-\frac{3-\sqrt{3}}{6} hM} y_0 + h \left[\frac{1}{4} f(Y_1) + \frac{3-2\sqrt{3}}{12} f(Y_2) \right], \\ Y_2 &= e^{-\frac{3+\sqrt{3}}{6} hM} y_0 + h \left[\frac{3+2\sqrt{3}}{12} f(Y_1) + \frac{1}{4} f(Y_2) \right], \\ y_1 &= e^{-hM} y_0 + \frac{h}{2} (f(Y_1) + f(Y_2)) - \frac{h^2}{2!} M f(y_0) + \frac{h^3}{3!} \left((M - f'_y(y_0)) M f(y_0) - M f'_y(y_0) g(y_0) \right. \\ & \left. + \frac{h^4}{4!} \left((-M + f'_y(y_0)) M^2 f(y_0) + M^2 f'_y(y_0) g(y_0) - M f''_{yy}(y_0)(g(y_0), g(y_0)) - M f'_y(y_0)(-M + f'_y(y_0))g(y_0) \right. \right. \\ & \left. \left. - f'_y(y_0) M f'_y(y_0) g(y_0) - f'_y(y_0) f'_y(y_0) M f(y_0) + 3 f''_{yy}(y_0)(-M f(y_0), g(y_0)) \right) \right). \end{aligned} \right. \tag{39}$$

The method (39) can be displayed by the Butcher tableau

$$\begin{array}{c|cc|cc}
 \frac{1}{2} - \frac{\sqrt{3}}{6} & e^{-\frac{3-\sqrt{3}}{6}hM} & \frac{1}{4} & \frac{3-2\sqrt{3}}{12} \\
 \frac{1}{2} + \frac{\sqrt{3}}{6} & e^{-\frac{3+\sqrt{3}}{6}hM} & \frac{3+2\sqrt{3}}{12} & \frac{1}{4} \\
 \hline
 e^{-hM} & w_4(z) & \frac{1}{2} & \frac{1}{2}
 \end{array} \quad (40)$$

We have presented the order conditions for the fourth-order IMSVERK method with $w_s(z)$ reduce to (34), the order conditions for the fourth-order IMSVERK method with $\bar{w}_s(z)$ are exactly identical to (34) as well. The following theorem shows this point.

Theorem 3.2. *Supposed that the coefficients of the two-stage IMMVERK method with $\bar{w}_4(z)$*

$$\begin{cases}
 \bar{Y}_1 = y_0 + h[\bar{a}_{11}(-M\bar{Y}_1 + f(\bar{Y}_1)) + \bar{a}_{12}(-M\bar{Y}_2 + f(\bar{Y}_2))], \\
 \bar{Y}_2 = y_0 + h[\bar{a}_{21}(-M\bar{Y}_1 + f(\bar{Y}_1)) + \bar{a}_{22}(-M\bar{Y}_2 + f(\bar{Y}_2))], \\
 \bar{y}_1 = e^{-hM}y_0 + h(\bar{b}_1f(\bar{Y}_1) + \bar{b}_2f(\bar{Y}_2)) + \bar{w}_4(z),
 \end{cases} \quad (41)$$

where

$$\begin{aligned}
 \bar{w}_4(z) = & -\frac{h^2}{2!}Mf(y_0) + \frac{h^3}{3!}(M^2f(y_0) - Mf'_y(y_0)g(y_0)) + \frac{h^4}{4!}(-M^3f(y_0) + M^2f'_y(y_0)g(y_0) - Mf''_{yy}(y_0)(g(y_0), g(y_0)) \\
 & - Mf'_y(y_0)(-M + f'_y(y_0))g(y_0)),
 \end{aligned}$$

which satisfy (34), then the IMMVERK method has order four.

Proof. Under the assumptions $\bar{c}_1 = \bar{a}_{11} + \bar{a}_{12}$ and $\bar{c}_2 = \bar{a}_{21} + \bar{a}_{22}$, by comparing the Taylor series of numerical solution \bar{y}_1 with exact solution $y(t_0 + h)$, we can verify the conclusion. Therefore, we omit the details. \square

The Theorem 3.2 reveals that order conditions of fourth-order IMMVERK methods with two stages, which are exactly identical to (34). Thus, we obtain the unique fourth-order IMMVERK method with two stages

$$\begin{cases}
 Y_1 = y_0 + h\left[\frac{1}{4}(-MY_1 + f(Y_1)) + \frac{3-2\sqrt{3}}{12}(-MY_2 + f(Y_2))\right], \\
 Y_2 = y_0 + h\left[\frac{3+2\sqrt{3}}{12}(-MY_1 + f(Y_1)) + \frac{1}{4}(-MY_2 + f(Y_2))\right], \\
 y_1 = e^{-hM}y_0 + \frac{h}{2}(f(Y_1) + f(Y_2)) - \frac{h^2}{2!}Mf(y_0) + \frac{h^3}{3!}(M^2f(y_0) - Mf'_y(y_0)g(y_0)) \\
 \quad + \frac{h^4}{4!}\left(-M^3f(y_0) + M^2f'_y(y_0)g(y_0) - Mf''_{yy}(y_0)(g(y_0), g(y_0)) - Mf'_y(y_0)(-M + f'_y(y_0))g(y_0)\right),
 \end{cases} \quad (42)$$

in here $g(y_0) = -My_0 + f(y_0)$, which can be denoted by the Butcher tableau

$$\begin{array}{c|cc|cc}
 \frac{1}{2} - \frac{\sqrt{3}}{6} & I & \frac{1}{4} & \frac{3-2\sqrt{3}}{12} \\
 \frac{1}{2} + \frac{\sqrt{3}}{6} & I & \frac{3+2\sqrt{3}}{12} & \frac{1}{4} \\
 \hline
 e^{-hM} & \bar{w}_4(z) & \frac{1}{2} & \frac{1}{2}
 \end{array} \quad (43)$$

In view of (39) and (42), two methods reduce to the classical fourth-order implicit RK method with two stages by Hammer and Hollingsworth (see, e.g. [14]) once $M \rightarrow \mathbf{0}$. It should be noted that (39) and (42) use the Jacobian matrix and Hessian matrix of $f(y)$ with respect to y at each step, however, as we known that the

idea for stiff problems is no by means new (see, e.g. [5, 8]). Using the Gaussian points $c_1 = \frac{1}{2} - \frac{\sqrt{3}}{6}$ and $c_2 = \frac{1}{2} + \frac{\sqrt{3}}{6}$, the ERK method of collocation type with two stages was formulated by Hochbruck et al. [19], which can be indicated by the Butcher tableau

$$\begin{array}{c|cc} \frac{1}{2} - \frac{\sqrt{3}}{6} & \frac{\sqrt{3}}{6}\varphi_1(-c_1hM) - \sqrt{3}c_1^2\varphi_2(-c_1hM) & -\sqrt{3}c_1^2\varphi_1(-c_1hM) + \sqrt{3}c_1^2\varphi_2(-c_1hM) \\ \frac{1}{2} + \frac{\sqrt{3}}{6} & \sqrt{3}c_2^2\varphi_1(-c_2hM) - \sqrt{3}c_2^2\varphi_2(-c_2hM) & -\frac{\sqrt{3}}{6}\varphi_1(-c_2hM) + \sqrt{3}c_2^2\varphi_2(-c_2hM) \\ \hline & \sqrt{3}c_2\varphi_1(-hM) - \sqrt{3}\varphi_2(-hM) & -\sqrt{3}c_1\varphi_1(-hM) + \sqrt{3}\varphi_2(-hM) \end{array} \quad (44)$$

Differently from the classical order (nonstiff), the order of (44) is based on the stiff conditions. It is clear that the coefficients of (44) are matrix exponentials, which means their implementation depends on evaluations of matrix exponentials. When we consider the variable stepsize technique, we have to recalculate the coefficients of these exponential integrators. By contrast, the coefficients of SVERK and MVERK methods are the real constants, which can reduce to the computation of matrix exponentials to some extent. On the other hands, compared with standard RK methods, SVERK and MVERK methods can reduce the limit on stepsize for solving (1), because these ERK methods reserve some properties of matrix exponentials.

4. Linear stability

In what follows we investigate the linear stability properties of explicit and implicit MVERK and SVERK methods. For classical RK methods, the linear stability analysis of RK methods is related to the Dahlquist equation [15]

$$y' = \lambda y, \quad \lambda \in \mathbb{R}.$$

When we consider the exponential integrators, as stated in [4], the stability properties of a exponential method is shown by applying the method to the partitioned Dahlquist equation

$$y' = i\lambda_1 y + i\lambda_2 y, \quad y(t_0) = y_0, \quad \lambda_1, \lambda_2 \in \mathbb{R}. \quad (45)$$

In fact, we solve (45) by a partitioned exponential integrator, treat the $i\lambda_1$ exponentially and the $i\lambda_2$ explicitly, then we have the explicit scalar form

$$y_{n+1} = R(ik_1, ik_2)y_n, \quad k_1 = h\lambda_1, \quad k_2 = h\lambda_2. \quad (46)$$

Definition 4.1. *The set S of the stability function $R(ik_1, ik_2)$, which satisfies*

$$S = \{(k_1, k_2) \in \mathbb{R}^2 : |R(ik_1, ik_2)| \leq 1\},$$

then the set S is called the stability region of a exponential method.

Applying the implicit MVERK method (30) and SVERK method (28) to (45), the stability regions of implicit second-order ERK method are respectively depicted in Fig. 1 (b) and (c), and we also plot the stability region of the explicit second-order MVERK method with two stages (see, e.g. [35]) in Fig. 1 (a). For fourth-order ERK methods, we select the IMMVERK method (42), IMSVERK method (39) and the explicit SVERK method with four stages (see, e.g. [21]) to make a comparison, the stability regions of these methods are shown in Fig. 2. The Figs. 1 and 2 reveal that the implicit exponential methods possess the comparable stability regions than explicit exponential methods.

5. Numerical Experiments

In this section, we respectively apply the symplectic method (18) and some highly accurate IMSVERK and IMMVERK methods to some highly oscillatory problems. In order to show the accuracy and efficiency of these new methods, we select the standard implicit ERK methods to make comparisons. Since these methods are implicit, the iteration is required, and we use the fixed-point iteration. In all numerical experiments, we take the Euclidean norm for global errors (GE) and denote the global error of Hamiltonian by GEH. The numerical results will demonstrate the long time energy preservation of (18) and the efficiency of these highly accurate IMSVERK and IMMVERK methods. The following numerical methods are chosen for comparison:




Figure 1: (a): Stability region for second-order explicit MVERK (EXMVERK21) method with two stages. (b) Stability region for second-order implicit MVERK (IMMVERK12) method. (c): Stability region for second-order implicit SVERK (IMSVERK12) method.




Figure 2: (a): Stability region for fourth-order explicit SVERK (EXSVERK41) method with four stages. (b) Stability region for fourth-order implicit MVERK (IMMVERK24) method. (c): Stability region for fourth-order implicit SVERK (IMSVERK24) method.

- First-order methods:
 - IMSVERK1: the 1-stage implicit SVERK method (18) of order one presented in this paper;
 - EEuler: the 1-stage explicit exponential Euler method of order one proposed in [18];
 - IMEEuler: the 1-stage implicit exponential Euler method of order one proposed in [19].
- Second-order methods:
 - IMMVERK12: the 1-stage implicit MVERK method (30) of order two presented in this paper;
 - IMSVERK12: the 1-stage implicit SVERK method (28) of order two presented in this paper;
 - IMERK12: the 1-stage implicit ERK method (32) of order two proposed in [19].
- Fourth-order methods:
 - IMMVERK24: the 2-stage implicit MVERK method (42) of order four presented in this paper;
 - IMSVERK24: the 2-stage implicit SVERK method (39) of order four presented in this paper;
 - IMERK24: the 2-stage implicit ERK method (44) proposed in [19].

Problem 1. The Hénon-Heiles Model used to describe the stellar motion (see, e.g. [15]), which has the following identical form

$$\begin{pmatrix} x_1 \\ x_2 \\ y_1 \\ y_2 \end{pmatrix}' + \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ -2x_1x_2 \\ -x_1^2 + x_2^2 \end{pmatrix}.$$

The Hamiltonian function of this system is

$$H(x, y) = \frac{1}{2}(y_1^2 + y_2^2) + \frac{1}{2}(x_1^2 + x_2^2) + x_1^2 x_2 - \frac{1}{3}x_2^3.$$

We select the initial values as

$$(x_1(0), x_2(0), y_1(0), y_2(0))^T = \left(\sqrt{\frac{11}{96}}, 0, 0, \frac{1}{4}\right)^T.$$

At first, Fig. 3 (a) presents that the problem is solved on the interval $[0, 10]$ with stepsizes $h = 1/2^k$, $k = 3, \dots, 7$ for IMSVERK1s1, EEuler, IMEEuler, then we integrate this problem over the interval $[0, 100]$ and the stepsize is chosen as $h = \frac{1}{40}$, the relative errors $RGEH = \frac{GEH}{H_0}$ of Hamiltonian energy for IMSVERK1s1, EEuler and IMEEuler are presented by the Fig. 3 (b) and Fig. 4. Finally, we integrate this system over the interval $[0, 10]$ with stepsizes $h = 1/2^k$ for $k = 3, \dots, 7$, Figs. 5 and 6 display the global errors (GE) against the stepsizes and the CPU time (seconds) for IMMVERK12, IMSVERK12, IMERK12, IMMVERK24, IMSVERK24, IMERK24.




Figure 3: Results for Problem 1. (a): The log-log plots of global errors (GE) against h . (b): the energy preservation for method IMSVERK1s1.




Figure 4: Results for Problem 1. : the energy preservation for methods EEuler (a) and IMEEuler (b).

Problem 2. Consider the Duffing equation [28]

$$\begin{cases} \ddot{q} + \omega^2 q = k^2(2q^3 - q), \\ q(0) = 0, \dot{q}(0) = \omega, \end{cases} \quad (47)$$




Figure 5: Results for Problem 1. (a): The log-log plots of global errors (GE) against h . (b): The log-log plots of global errors against the CPU time.




Figure 6: Results for Problem 1. (a): The log-log plots of global errors (GE) against h . (b): The log-log plots of global errors against the CPU time.

where $0 \leq k < \omega$.

Let $p = \dot{q}$, $z = (p, q)^T$. We rewrite the Duffing equation as

$$\begin{pmatrix} p \\ q \end{pmatrix}' + \begin{pmatrix} 0 & \omega^2 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} k^2(2q^3 - q) \\ 0 \end{pmatrix}.$$

It is a Hamiltonian system with the Hamiltonian

$$H(p, q) = \frac{1}{2}p^2 + \frac{1}{2}\omega^2 q^2 + \frac{k^2}{2}(q^2 - q^4).$$

The exact solution of this problem is

$$q(t) = \text{sn}(\omega t; k/\omega),$$

where sn is the Jacobian elliptic function.

Let $\omega = 10$, $k = 0.01$. This problem is solved on the interval $[0, 100]$ with the stepsize $h = \frac{1}{40}$, the Fig. 7 (b) and Fig. 8 show the energy preservation behaviour for IMSVERK1s1, EEuler and IMEEuler. We also integrate the system over the interval $[0, 10]$ with stepsizes $h = 1/2^k$, $k = 4, \dots, 8$ for IMSVERK1s1, EEuler, IMEEuler, IMMVERK12, IMSVERK12, IMERK12, IMMVERK24, IMSVERK24, IMERK24, which are shown in Figs 7 (a), 9 and 10.

Problem 3. Consider the sine-Gorden equation with periodic boundary conditions [28]

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2} - \sin(u), & -1 < x < 1, t > 0, \\ u(-1, t) = u(1, t). \end{cases}$$




Figure 7: Results for Problem 2. (a): The log-log plots of global errors (GE) against h . (b): the energy preservation for method IMSVERK1s1.




Figure 8: Results for Problem 2. : the energy preservation for methods EEuler (a) and IMEEuler (b).




Figure 9: Results for Problem 2. (a): The log-log plots of global errors (GE) against h . (b): The log-log plots of global errors against the CPU time.

Discretising the spatial derivative ∂_{xx} by the second-order symmetric differences, which leads to the following Hamiltonian system

$$\frac{d}{dt} \begin{pmatrix} U' \\ U \end{pmatrix} + \begin{pmatrix} \mathbf{0} & M \\ -I & \mathbf{0} \end{pmatrix} \begin{pmatrix} U' \\ U \end{pmatrix} = \begin{pmatrix} -\sin(U) \\ \mathbf{0} \end{pmatrix}, \quad t \in [0, t_{\text{end}}],$$




Figure 10: Results for Problem 2. (a): The log-log plots of global errors (GE) against h . (b): The log-log plots of global errors against the CPU time.

whose Hamiltonian is shown by

$$H(U', U) = \frac{1}{2} U'^T U' + \frac{1}{2} U^T M U^T - (\cos u_1 + \cdots + u_N).$$

In here, $U(t) = (u_1(t), \dots, u_N(t))^T$ with $u_i(t) \approx u(x_i, t)$ for $i = 1, \dots, N$, with $\Delta x = 2/N$ and $x_i = -1 + i\Delta x$, $F(t, U) = -\sin(u) = -(\sin(u_1), \dots, \sin(u_N))^T$, and

$$M = \frac{1}{\Delta x^2} \begin{pmatrix} 2 & -1 & & -1 \\ -1 & 2 & -1 & \\ & \ddots & \ddots & \ddots \\ -1 & & -1 & 2 & -1 \\ & & & -1 & 2 \end{pmatrix}.$$

In this test, we choose the initial value conditions

$$U(0) = (\pi)_{i=1}^N, \quad U'(0) = \sqrt{N} \left(0.01 + \sin\left(\frac{2\pi i}{N}\right) \right)_{i=1}^N$$

with $N = 32$, and solve the problem on the interval $[0, 1]$ with stepsizes $h = 1/2^k$, $k = 5, \dots, 9$. The global errors GE against the stepsizes and the CPU time (seconds) for IMSVERK1s1, EEuler, IMEEuler, IMMVERK12, IMSVERK12, IMERK12, IMMVERK24, IMSVERK24, IMERK24, which are respectively presented in Figs 11 (a), 13 and 14. The Fig. 11 (b) and Fig. 12 display the energy preservation behaviour of this problem integrated on the interval $[0, 100]$ with stepsize $h = 1/100$ for IMSVERK1s1, EEuler, IMEEuler.

6. Conclusion

It is known that exponential Runge–Kutta methods have the unique advantage to solve highly oscillatory systems, however, the implementation of ERK depends on the evaluations of matrix exponentials. Actually, the coefficients of these exponential integrators are related to the matrix exponentials. In order to reduce computational cost, two new classes of explicit ERK integrators are formulated in recent papers [21, 35]. As a sequel to our work, implicit ERK methods are studied in this paper. First of all, we analyzed the symplectic conditions and obtained the numerical method with symplecticity, which exhibited the structure-preserving property for solving highly oscillatory problems or Hamiltonian systems. But the symplectic method only had order one, we needed some practical and effective numerical methods. Highly accurate ERK methods were presented, and the order conditions of these ERK methods were exactly identical to standard RK methods. Furthermore, we presented the linear stability regions for explicit and implicit ERK methods. Finally, in




Figure 11: Results for Problem 3. (a): The log-log plots of global errors (GE) against h . (b): the energy preservation for method IMSVERK1s1.




Figure 12: Results for Problem 3. : the energy preservation for methods EEuler (a) and IMEEuler (b).




Figure 13: Results for Problem 3. (a): The log-log plots of global errors (GE) against h . (b): The log-log plots of global errors against the CPU time.

numerical experiments, we took IMSVERK1s1 and highly accurate methods (IMMVERK12, IMMVERK24, IMSVERK12, IMSVERK24) by comparison with the standard ERK methods, the numerical results not only shown the good energy preservation behaviour for IMSVERK1s1, but also demonstrated the comparable accuracy and efficiency for IMMVERK12, IMMVERK24, IMSVERK12, IMSVERK24, when applied to the Hénon-Heiles Model, the Duffing equation and the sine-Gordon equation.

Exponential integrators show the better performance than non-exponential integrators. High accuracy and




Figure 14: Results for Problem 3. (a): The log-log plots of global errors (GE) against h . (b): The log-log plots of global errors against the CPU time.

structure preservation for exponential integrators can be further investigated.

References

References

- [1] S. M. Allen, J. W. Cahn, A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening, *Acta Metall.* 27 (1979) 1085-1095.
- [2] H. Berland, B. Owren, B. Skaflestad, B-series and order conditions for exponential integrators, *SIAM J. Numer. Anal.* 43 (2005) 1715-1727.
- [3] L. Brugnano, C. Zhang, D. Li, A class of energy-conserving Hamiltonian boundary value methods for nonlinear Schrödinger equation with wave operator, *Commun. Nonlinear Sci. Numer. Simul.* 60 (2018) 33-49.
- [4] T. Buvoli, M. L. Minion, On the stability of exponential integrators for non-diffusive equations, *J. Comput. Appl. Math.* 409 (2022) 114126.
- [5] J. R. Cash, Second derivative extended backward differentiation formulas for the numerical integration of stiff systems, *SIAM J. Numer. Anal.* 18 (1981) 21-36.
- [6] E. Celledoni, D. Cohen, B. Owren, Symmetric exponential integrators with an application to the cubic Schrödinger equation, *Found. Comput. Math.* 8 (2008) 303-317.
- [7] J. Chen, M. Qin, Multisymplectic Fourier pseudospectral method for the nonlinear Schrödinger equation, *Electron. Trans. Numer. Anal.* 12 (2001) 193-204.
- [8] W. H. Enright, Second derivative multistep methods for stiff ordinary differential equations, *SIAM J. Numer. Anal.* 11 (1974) 321-331.
- [9] Y. Fang, Q. Ming, Embedded pair of extended Runge-Kutta-Nystrom type methods for perturbed oscillators, *Appl. Math. Model.* 34 (2010) 2665-2675.
- [10] Y. Fang, Q. Ming, X. Wu, Extended RKN-type methods with minimal dispersion error for perturbed oscillators, *Comput. Phys. Commun.* 181 (2010) 639-650.
- [11] Y. Fang, X. Hu, J. Li, Explicit pseudo two-step exponential Runge-Kutta methods for the numerical integration of first-order differential equations, *Numer. Algor.* 86 (2021) 1143-1163.
- [12] K. Feng, On difference schemes and symplectic geometry. *Proceedings of the 5th International Symposium on Differential Geometry & Differential Equations*. Beijing: Science Press, 1985: 42-58.
- [13] K. Feng, Difference schemes for Hamiltonian formalism and symplectic geometry, *J. Comp. Math.* 4 (1986) 279-289.
- [14] P. C. Hammer, J. W. Hollingsworth, Trapezoidal methods of approximating solutions of differential equations, *MTAC.* 9 (1955) 92-96.
- [15] E. Hairer, C. Lubich, G. Wanner, *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*, 2nd edn. Springer, Berlin (2006).
- [16] M. Hochbruck, C. Lubich, On Krylov subspace approximations to the matrix exponential operator, *SIAM J. Numer. Anal.* 34 (1997) 1911-1925.
- [17] M. Hochbruck, C. Lubich, H. Selhofer, Exponential integrators for large systems of differential equations, *SIAM J. Sci. Comput.* 19 (1998) 1552-1574.
- [18] M. Hochbruck, A. Ostermann, Explicit exponential Runge-Kutta methods for semilinear parabolic problems, *SIAM J. Numer. Anal.* 43 (2005) 1069-1090.
- [19] M. Hochbruck, A. Ostermann, Exponential Runge-Kutta methods for parabolic problems, *Appl. Numer. Math.* 53 (2005) 323-339.
- [20] M. Hochbruck, A. Ostermann, Exponential integrators, *Acta Numer.* 19 (2010) 209-286.
- [21] X. Hu, Y. Fang, B. Wang, Two new families of fourth-order explicit exponential Runge-Kutta methods with four stages for stiff or highly oscillatory systems, *arXiv: 2210.12407*. 2022.
- [22] S. Krogstad, Generalized integrating factor methods for stiff PDEs, *J. Comput. Phys.* 203 (2005) 72-88.

- [23] J. D. Lawson, Generalized Runge-Kutta processes for stable systems with large Lipschitz constants, *SIAM J. Numer. Anal.* 4 (1967) 372-380.
- [24] J. D. Lambert, S. T. Sigurdsson, Multistep methods with variable matrix coefficients, *SIAM J. Numer. Anal.* 9 (1972) 715-733.
- [25] Y. Li, X. Wu, Exponential integrators preserving first integrals or Lyapunov functions for conservative or dissipative systems, *SIAM J. Sci. Comput.* 38 (2016) A1876-A1895.
- [26] R. I. McLachlan, G. R. W. Quispel, N. Robidoux, A unified approach to Hamiltonian systems, Poisson systems, gradient systems, and systems with Lyapunov functions or first integrals, *Phys. Rev. Lett.* 81 (1998) 2399-2411.
- [27] L. Mei, L. Huang, X. Wu, Energy-preserving continuous-stage exponential Runge-Kutta integrators for efficiently solving Hamiltonian systems, *SIAM J. Sci. Comput.* 44 (2022) A1092-A1115.
- [28] L. Mei, X. Wu, Symplectic exponential Runge-Kutta methods for solving nonlinear Hamiltonian systems, *J. Comput. Phys.* 338 (2017) 567-584.
- [29] J. M. Sanz-Serna, Runge-Kutta schemes for Hamiltonian systems, *BIT Numer. Math.* 28 (1988) 877-883.
- [30] L. N. Trefethen, *Spectral methods in MATLAB*, SIAM, Philadelphia, 2000.
- [31] B. Wang, X. Wu, Explicit multi frequency symmetric extended RKN integrators for solving multi-frequency and multidimensional oscillatory reversible systems, *Calcolo* 52 (2015) 207-231.
- [32] B. Wang, X. Wu, F. Meng, Y. Fang, Exponential Fourier collocation methods for solving first-order differential equations, *J. Comput. Math.* 35 (2017) 711-736.
- [33] B. Wang, X. Wu, Exponential collocation methods for conservative or dissipative systems, *J. Comput. Appl. Math.* 360 (2019) 99-116.
- [34] B. Wang, X. Wu, *Geometric Integrators for Differential Equations with Highly Oscillatory Solutions*, Springer Nature Singapore Pte Ltd. 2021.
- [35] B. Wang, X. Hu, X. Wu, Two new classes of exponential Runge-Kutta integrators for efficiently solving stiff systems or highly oscillatory problems, *arXiv: 2210.00685*. 2022.
- [36] M. Wang, D. Li, C. Zhang, Y. Tang, Long time behavior of solutions of gKdV equations, *J. Math. Anal. Appl.* 390 (2012) 136-150.
- [37] X. Wu, X. You, W. Shi, B. Wang, ERKN integrators for systems of oscillatory second-order differential equations, *Comput. Phys. Commun.* 181 (2010) 1873-1887.
- [38] X. Wu, X. You, B. Wang, *Structure-Preserving Algorithms for Oscillatory Differential Equations*, Springer-Verlag, Berlin, Heidelberg, 2013.
- [39] X. Wu, K. Liu, W. Shi, *Structure-Preserving Algorithms for Oscillatory Differential Equations II*, Springer-Verlag, Berlin, Heidelberg, 2015.
- [40] X. Wu, B. Wang, *Recent Developments in Structure-Preserving Algorithms for Oscillatory Differential Equations*, Springer-Verlag, Singapore, 2018.