

Spanning subdivisions in dense digraphs

Hyunwoo Lee*

June 13, 2023

Abstract

We prove that an n -vertex digraph D with minimum semi-degree at least $(\frac{1}{2} + \varepsilon)n$ and $n = \Omega_\varepsilon(m)$ contains a subdivision of all m -arc digraphs. This is the best possible and settles a conjecture raised by Pavez-Singé [11] in a stronger form.

1. Introduction

Embedding a spanning sparse (di)graph into a dense (di)graph has been extensively studied over decades. Finding a sufficient condition that ensures the existence of a spanning structure is one of the central topics in this theme. Arguably, the most natural condition is the minimum degree condition. For example, a classical result of Dirac asserts that every n -vertex graph G with minimum degree at least $\frac{n}{2}$ contains a Hamiltonian cycle.

Inspired by Dirac's theorem, there were various problems and results regarding the minimum degree threshold for the existence of certain spanning structures. One direction is a perfect tiling problem. For graphs G and H , we say G has a perfect H -tiling if G contains a collection of vertex disjoint copies of H that covers every vertex of G . For a complete graph K_r on r vertices, Hajnal and Szemerédi [4] proved that every n -vertex graph G with minimum degree at least $\frac{r-1}{r}n$, where n divisible by r , contains a perfect K_r -tiling. For a general graph H , Kühn and Osthus [8] determines the exact minimum degree threshold for the existence of perfect H -tiling up to an additive constant.

Since then, many variants of the Kühn-Osthus theorem on graph tiling has been considered (for instance, see [1, 5, 6, 10]). Recently, the author [9] considered a new variant of the tiling problem, subdivision tilings. For graphs G and H we say that G has a perfect H -subdivision tiling if G contains a collection of vertex disjoint copies of subdivisions of H that covers all vertices of G . Here, we say that H' is a subdivision of H if H' is obtained by replacing each edge of H with vertex-disjoint path. The asymptotic behavior of the tight minimum degree threshold for the existence of perfect H -subdivision tiling was determined for all graph H in [9].

We note that indeed, Dirac's theorem determined the minimum degree threshold for the existence of spanning K_3 -subdivision. On the other hand, for a graph H , [9] determined the minimum degree threshold for the existence of spanning structure that each can be partitioned into vertex disjoint copies of subdivisions of H . Comparing these two results, it is very natural to ask the minimum degree threshold for the existence of a spanning subdivision of general graphs H . Very recently, Pavez-Singé [11] proved that for every $\varepsilon > 0$, there is a constant C such that every graph G on $n \geq Ck^2$ vertices with minimum degree at least $(\frac{1}{2} + \varepsilon)n$ contains a spanning subdivision of K_k for any positive integer k . More generally,

*Department of Mathematical Sciences, KAIST, South Korea and Extremal Combinatorics and Probability Group (ECOPRO), Institute for Basic Science (IBS). E-mail: hyunwoo.lee@kaist.ac.kr. Supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government(MSIT) No. RS-2023-00210430, and the Institute for Basic Science (IBS-R029-C4).

they showed that if H is an k -vertex d -regular graph with $\log k \leq d \leq k$, then the same result holds for a spanning subdivision of H as long as $n \geq Cdk$. For general graphs, they propose the following natural conjecture.

Conjecture 1.1 ([11]). *For every $\varepsilon > 0$, there is a constant C such that the following holds. For every graph G on $n \geq Cm$ vertices, if the minimum degree of G is at least $(\frac{1}{2} + \varepsilon)n$, then G contains a spanning subdivision of any m -edge graph H with no isolated vertices.*

In this article, we prove Conjecture 1.1 in a stronger form considering digraphs. Let H be a digraph. We say a digraph H' is a subdivision of H if H' is obtained by replacing each arc of H with a vertex-disjoint directed path with the same consistent orientation. For a digraph D and $v \in D$, the semi-degree of v in D is the minimum of the out-degree of v and in-degree of v in D .

Theorem 1.2. *Let $\varepsilon > 0$ be a positive real number. Then there is a constant $C > 0$ such that the following holds. Let D be a digraph on $n \geq Cm$ vertices with minimum degree at least $(\frac{1}{2} + \varepsilon)n$. Then D contains a spanning subdivision of any m -arc digraph H with no isolated vertices.*

By replacing edges of G with two arcs in both directions, it is straightforward to check that Theorem 1.2 implies Conjecture 1.1.

The minimum semi-degree condition in Theorem 1.2 is asymptotically best possible. Let H be an m -arc digraph on k vertices without isolated vertices. We note that $k \leq 2m$. Let A and B be disjoint sets with $|A| = \lfloor \frac{n}{2} \rfloor - (m + k)$ and $|B| = n - |A|$. Let D be a digraph on the vertex set $A \cup B$ such that $A(D) = \{\overrightarrow{uv}, \overrightarrow{vu} : (u, v) \in A \times B\}$. Then the minimum semi-degree of D is at least the size of A , that is $\lfloor \frac{n}{2} \rfloor - (m + k)$. We now claim that D does not contain a spanning subdivision of H . Assume D has a spanning subdivision of H . Then at least $n - k$ vertices of D covered by at most m pairwise vertex disjoint directed paths. Since the underlying graph of D is a bipartite graph on the bipartition (A, B) , the inequality $||A| - |B|| \leq m + k$ must hold, a contradiction. Thus, D does not contain a spanning subdivision of H .

Since a directed Hamiltonian cycle is a spanning subdivision of a directed 2-cycle, we can consider Theorem 1.3 as the asymptotic generalization of the following classical digraph analog to Dirac's theorem.

Theorem 1.3 (Ghouila-Houri [3]). *Let D be an n -vertex digraph with minimum semi-degree at least $\frac{n}{2}$. Then D contains a directed Hamiltonian cycle.*

The proof of the main theorem uses the absorption method introduced by Rödl, Ruciński, and Szemerédi [12] and implicitly used in [2, 7]. The absorption method is an extremely useful tool for finding spanning structures. In order to get more information about the absorption method, we recommend seeing the survey [13].

2. Preliminaries

2.1. Notations

We write $[n] = \{1, 2, \dots, n\}$. For a statement regarding some parameters $\beta, \alpha_1, \dots, \alpha_t$, we say that it holds when $0 < \beta \ll \alpha_1, \dots, \alpha_t < 1$ if there exists a function f such that for every $0 < \beta \leq f(\alpha_1, \dots, \alpha_t)$ the statement holds. In this article, we will not compute these functions explicitly.

We use standard notations in digraph theory. For a digraph D , we let $V(D)$ be the vertex set of D and $A(D)$ be the arc set of D . For a vertex v in a directed graph D , we denote by $N_D^+(v)$ and $N_D^-(v)$ the out-neighborhood and in-neighborhood of v , respectively. We write $d_D^+(v)$ and $d_D^-(v)$ as the size of the

out-neighborhood and in-neighborhood of v and we call them out-degree and in-degree of v , respectively. We denote by $\delta^0(D)$ the minimum semi-degree, that is $\min\{\delta^+(D), \delta^-(D)\}$.

For a vertex subset X and Y , we let $A_D[X, Y]$ be the set of arcs starting from X and ending at Y . We note that X and Y do not need to be disjoint. For a pair of vertices $(u, v) \in V(D) \times V(D)$, we define $N_D(u, v) := \{w \in V(D) : \overrightarrow{uw}, \overrightarrow{wv} \in A(D)\}$. We denote by $d_D(u, v)$ the number $|N_D(u, v)|$.

For a digraph D and vertex subsets $X, Y \subseteq V(D)$, let $D - X + Y$ be a digraph that is an induced digraph of D on the vertex subset $(V(D) \setminus X) \cup Y$.

Let X be a set and t be a positive integer. We write X^t be the t -fold cartesian product of X , that is $X \times \cdots \times X$. We say two element $(y_1, \dots, y_t), (z_1, \dots, z_t)$ of X^t are tuple-disjoint if $\{y_1, \dots, y_t\} \cap \{z_1, \dots, z_t\} = \emptyset$.

3. (n, t, d) -tuple systems

Definition 3.1. Let n and t be two positive integers and let d be a real number lies on $(0, 1)$. For two sets X and Y where $|X| \leq n^2$ and $|Y| = n$, we say a collection of pairs $\mathcal{P} \subseteq X \times Y^t$ is an (n, t, d) -tuple system on (X, Y) if \mathcal{P} satisfies the following: For all element x in X , the size of the set $\{Z \subseteq Y : (x, Z) \in \mathcal{P}\}$ is at least dn^t .

The following lemma is the key lemma for constructing a connector and absorber via the (n, t, d) -tuple systems.

Lemma 3.2. *We fix the parameters below.*

$$0 < \frac{1}{n}, \beta \ll d, \alpha, \frac{1}{t} \leq 1$$

where n and t are integers.

Let \mathcal{P} be an (n, t, d) -tuple system on the pair of sets (X, Y) . Then there is a collection of subsets $\mathcal{Y} \subseteq Y^t$ satisfies the followings:

- Elements in \mathcal{Y} are pairwise tuple-disjoint,
- The size of \mathcal{Y} are at most $\frac{\alpha}{t}n$,
- For every $x \in X$, the number of elements Z in \mathcal{Y} where $(x, Z) \in \mathcal{P}$ is at least βn .

Indeed, the similar notion with (n, t, d) -tuple systems and Lemma 3.2 were already considered in [9] in the notion of (n, t, d) -systems. The proof of Lemma 3.2 is almost the same as the proof of Lemma 4.9 in [9]. As the proof of Lemma 3.2 is almost identical to the proof of Lemma 4.9 in [9] up to straightforward modification, we omit the proof here.

We want to remark that the notion of (n, t, d) -systems, (n, t, d) -tuple systems, and Lemma 3.2 provide systematic frameworks for the absorption method that appeared in [12].

4. Proof of Theorem 1.2

In this section, we will prove Theorem 1.2.

Lemma 4.1. *We fix the parameters as follows.*

$$0 < \frac{1}{n} \ll \beta \ll \alpha \ll \varepsilon < 1.$$

Let D an n -vertex digraph with $\delta^0(D) \geq (\frac{1}{2} + \varepsilon)n$. Then there is a set $R \subseteq V(D)$ with size $|R| \leq \alpha n$ such that the following holds. For any distinct vertices $u, v \in V(D)$, we have $|N_D(u, v) \cap R| \geq \beta n$.

Proof. Let $X = V(D) \times V(D) \setminus \{(v, v) : v \in V(D)\}$ and let $Y = V(D)$. Let $\mathcal{P} = \{((u, v), w) \in X \times Y : w \in N_D(u, v)\}$. By the pigeonhole principle, every distinct pair of vertices (u, v) of $V(G)$, we have $d_D(u, v) \geq 2\varepsilon n$. This implies that \mathcal{P} is a $(n, 1, 2\varepsilon)$ -system on (X, Y) . By Lemma 3.2, there is $\mathcal{Y} \subseteq Y$ with $|\mathcal{Y}| \leq \alpha n$ and for every $(u, v) \in X$, we have $|\{w\} \in \mathcal{Y} : ((u, v), w) \in \mathcal{P}| \geq \beta n$. This implies that $R = \mathcal{Y}$ is the desired set. \square

Lemma 4.2. *We fix the positive parameters as follows.*

$$0 < \frac{1}{n} \ll \beta \ll \alpha \ll \varepsilon < 1.$$

Then the following holds. Let D be an n -vertex graph with $\delta^0(D) \geq (\frac{1}{2} + \varepsilon)n$. Then there is a directed path A in D satisfying $|V(A)| \leq \alpha n$ such that for every $U \subseteq V(G) \setminus A$ with $|U| \leq \beta n$, there is a directed path P in D on the vertex set $V(A) \cup U$ that has the same endpoints with the path A .

Proof. For a vertex $u \in V(D)$, we say a tuple $(v, w) \in V(D) \times V(D)$ is a *good tuple* for u if $v \neq w$ and $\overrightarrow{vu}, \overrightarrow{uw}, \overrightarrow{vw} \in A(D)$. We claim that every vertex of D has many good tuples.

Claim 1. *For every $u \in V(D)$, there are at least $4\varepsilon^2 n^2$ good tuples for u .*

Proof. Let U be a subsets of $V(D)$ with size at least $(\frac{1}{2} + \varepsilon)n$ and U' be a subset of U with size at least $2\varepsilon n$. Then for every $u' \in U'$, we have $|N_D^+(u') \cap U| \geq 2\varepsilon n$. Thus, the inequality $|A_D[U', U]| = \sum_{u' \in U'} |N_D^+(u') \cap U| \geq 4\varepsilon^2 n^2$ holds. Take $U = N^+(u)$ and $U' = U \cap N^-(u)$. Then we have $|U| \geq (\frac{1}{2} + \varepsilon)n$ and $|U'| \geq 2\varepsilon n$. Then for every $\overrightarrow{vw} \in A_D[U', U]$, the tuple (v, w) is good tuple for u . Since $|A_D[U', U]| \geq 4\varepsilon^2 n^2$, the claim holds. \blacksquare

Let $X = V(D)$ and $Y = V(D)$. Let $\mathcal{P} = \{(u, (v, w)) \in X \times Y^2 : (v, w) \text{ is a good tuple for } u\}$. Then by Claim 1, \mathcal{P} is a $(n, 2, 4\varepsilon^2)$ -system on (X, Y) . Thus, by Lemma 3.2, there is a set $\mathcal{Y} \subseteq V(D)^2$ with size at most $\frac{\alpha}{3}$ such that elements in \mathcal{Y} are pairwise tuple-disjoint and for every $u \in V(D)$, we have $|\{(v, w) \in \mathcal{Y} : (u, (v, w)) \in \mathcal{P}\}| \geq \beta n$. Let $\mathcal{Y} = \{(v_1, w_1), \dots, (v_\ell, w_\ell)\}$. Since we have $d_D(x, y) \geq 2\varepsilon n \geq \alpha n \geq 3|\mathcal{Y}|$ for every pair of distinct vertices (x, y) , we can greedily choose distinct vertices $x_1, \dots, x_{\ell-1}$ which are disjoint from $\{v_1, \dots, v_\ell, w_1, \dots, w_\ell\}$ such that $\overrightarrow{w_i x_i}, \overrightarrow{x_i v_{i+1}} \in A(D)$ for each $i \in [\ell - 1]$. This implies that there is a directed path $A = v_1 w_1 x_1 v_2 w_2 x_2 \dots x_{\ell-1} v_\ell w_\ell$ in D .

We now show that A is a desired directed path. Note that $|V(A)| \leq 3|\mathcal{Y}| \leq \alpha n$. Let $U \subseteq V(D) \setminus V(A)$ be a vertex set with size at most βn . By our choice of \mathcal{Y} , for each $u \in U$, there are at least βn pairs of (v_i, w_i) such that $\overrightarrow{v_i u}, \overrightarrow{u w_i} \in A(D)$. Thus, we can greedily absorb all the vertices of U by replacing the arc $\overrightarrow{v_i w_i}$ with $\overrightarrow{v_i u}, \overrightarrow{u w_i}$ and still maintain the same endpoints v_1 and w_ℓ . This proves the lemma. \square

We now prove Theorem 1.2. We fix the parameters as follows.

$$0 < \frac{1}{C} \ll \gamma \ll \beta \ll \alpha \ll \varepsilon < 1.$$

Let D be an n -vertex digraph with $\delta^0(D) \geq (\frac{1}{2} + \varepsilon)n$ where $n \geq Cm$. Let H be an m -arc digraph without isolated vertices. By Lemma 4.2, there is a path A in D that satisfy the Lemma 4.2 with parameters α and β . Let w_1, w_2 be the endpoints of the path A . Then we have $|V(A)| \leq \alpha n$ and the following holds. For every $U \subseteq V(D) \setminus V(A)$ of size at most βn , there is a directed path P in D which starts at w_1 and ends at w_2 where $V(P) = V(A) \cup U$.

Let $D_1 := D - V(A) + \{w_1, w_2\}$. Then we have $\delta^0(D_1) \geq (\frac{1}{2} + \frac{\varepsilon}{2})n$. By Lemma 4.1, there is a set $R \subseteq V(D_1)$ that satisfies the conclusions of Lemma 4.1 with parameters $\frac{\alpha}{2}, \beta, 2\gamma$. If $R \cap \{w_1, w_2\} \neq \emptyset$, then we simply remove w_1 and w_2 from R . Then R is disjoint from $\{w_1, w_2\}$ and we have $|R| \leq \beta n$ and

for any two distinct vertices x, y in $V(D_1)$, we have $|N_D(x, y) \cap R| \geq \gamma n$. Let $D_2 := D - (V(A) \cup R)$. Then we have $\delta^0(D_2) \geq (\frac{1}{2} + \frac{\varepsilon}{4})n$.

Let $V(H) = \{x_1, \dots, x_k\}$. We arbitrary choose vertices v_1, \dots, v_k in D_2 . Let $D_3 := D_2 - \{v_1, \dots, v_k\}$. Note that $k \leq 2m \leq \frac{\varepsilon}{4}Cm \leq \frac{\varepsilon}{4}n$. Thus, we have $\delta^0(D_3) \geq \frac{n}{2}$. By Theorem 1.3, the digraph D_3 has a directed Hamiltonian path starts from u_1 and ends at u_2 . Without loss of generality, we may assume that $\overrightarrow{x_1x_2} \in A(H)$. By our choice of R , there are three distinct vertices $z_1, z_2, z_3 \in R$ such that $\overrightarrow{v_1z_1}, \overrightarrow{z_1u_1}, \overrightarrow{u_2z_2}, \overrightarrow{z_2w_1}, \overrightarrow{w_2z_3}, \overrightarrow{z_3v_2} \in A(D)$. This implies that there is a directed path Q starts from v_1 and ends at v_2 such that Q contains the directed path A , all vertices of D_3 , and exactly three elements of R .

Let $A(H) \setminus \{\overrightarrow{x_1x_2}\} = \{e_1, \dots, e_{m-1}\}$. Since $\gamma n \geq \gamma Cm \geq m+2$, for each $i \in [m-1]$ where $e_i = \overrightarrow{x_ax_b}$, we can greedily choose distinct vertices $z_{e_i} \in R$ which is disjoint from $\{z_1, z_2, z_3\}$ such that $\overrightarrow{v_az_i}, \overrightarrow{z_iv_b} \in E(G)$. Thus, we can obtain a subdivision of H that contains the directed path A , and the remaining vertices are contained in R . Since $|R| \leq \beta n$, by our choice of A , we can absorb the remaining vertices, hence we obtain a spanning subdivision of H in D . This completes the proof.

References

- [1] József Balogh, Lina Li, and Andrew Treglown. “Tilings in vertex ordered graphs”. In: *Journal of Combinatorial Theory, Series B* 155 (2022), pp. 171–201.
- [2] Paul Erdős, András Gyárfás, and László Pyber. “Vertex coverings by monochromatic cycles and trees”. In: *Journal of Combinatorial Theory, Series B* 51.1 (1991), pp. 90–95.
- [3] Alain GHOUILAHOURLI. “Une condition suffisante d’existence d’un circuit hamiltonien”. In: *COMPTES RENDUS HEBDOMADAIRES DES SEANCES DE L’ACADEMIE DES SCIENCES* 251.4 (1960), pp. 495–497.
- [4] András Hajnal and Endre Szemerédi. “Proof of a conjecture of P. Erdős”. In: *Combinatorial theory and its applications 2* (1970), pp. 601–623.
- [5] Jie Han, Patrick Morris, and Andrew Treglown. “Tilings in randomly perturbed graphs: Bridging the gap between Hajnal-Szemerédi and Johansson-Kahn-Vu”. In: *Random Structures & Algorithms* 58.3 (2021), pp. 480–516.
- [6] Joseph Hyde and Andrew Treglown. “A degree sequence version of the Kuhn-Osthus tiling theorem”. In: *arXiv preprint arXiv:1909.12670* (2019).
- [7] Michael Krivelevich. “Triangle factors in random graphs”. In: *Combinatorics, Probability and Computing* 6.3 (1997), pp. 337–347.
- [8] Daniela Kühn and Deryk Osthus. “The minimum degree threshold for perfect graph packings”. In: *Combinatorica* 29.1 (2009), pp. 65–107.
- [9] Hyunwoo Lee. “On perfect subdivision tilings”. In: *arXiv preprint arXiv:2302.09393* (2023).
- [10] Allan Lo and Klas Markström. “ F -Factors in Hypergraphs Via Absorption”. In: *Graphs and Combinatorics* 31.3 (2015), pp. 679–712.
- [11] Matías Pavez-Signé. “Spanning subdivisions in Dirac graphs”. In: *arXiv preprint arXiv:2306.03994* (2023).
- [12] Vojtěch Rödl, Andrzej Ruciński, and Endre Szemerédi. “A Dirac-type theorem for 3-uniform hypergraphs”. In: *Combinatorics, Probability and Computing* 15.1-2 (2006), pp. 229–251.
- [13] Yi Zhao. “Recent advances on Dirac-type problems for hypergraphs”. In: *Recent trends in combinatorics* (2016), pp. 145–165.