Order-One Convergence of the Backward Euler Method for Random Periodic Solutions of Semilinear SDEs [†]

Yujia Guo^a , Xiaojie Wang^{*a}, and Yue Wu^b

^a School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha,

Hunan, P. R. China

^b Department of Mathematics and Statistics, University of Strathclyde, Glasgow G1 1XH, UK

December 12, 2023

Abstract

In this paper, we revisit the backward Euler method for numerical approximations of random periodic solutions of semilinear SDEs with additive noise. Improved L^p -estimates of the random periodic solutions of the considered SDEs are obtained under a more relaxed condition compared to literature. The backward Euler scheme is proved to converge with an order one in the mean square sense, which also improves the existing order-half convergence. Numerical examples are presented to verify our theoretical analysis.

AMS subject classification: 37H99, 60H10, 60H35, 65C30.

Keywords: Backward Euler method, Random periodic solution, Stochastic differential equations, Additive noise, Pull-back.

1 Introduction

Many phenomena in the real world exhibit both periodic and random nature, for instance, the daily temperature, energy consumption, and airline passenger volumes. To understand the long-time behaviour of these laws of random motion, where the underlying dynamics are usually modeled via stochastic differential equations (SDEs), it is crucial to study their random periodic solutions [6, 18].

Though the periodic solution has been a central concept in the field of deterministic dynamical systems since Poincaré's seminal work [11], its stochastic counterpart has not been properly defined or studied until the last decade. Zhao and Zheng [18] first formulated the definition of the pathwise random periodic solutions for C^1 -cocycles of random dynamical systems and Feng, Zhao and Zhou [6] further developed the definition for semiflows. This pioneering study boosts a series of work, including the study of anticipating random solutions of SDEs with multiplicative linear

[†]This work was supported by Natural Science Foundation of China (12071488, 12371417, 11971488) and Natural Science Foundation of Hunan Province (2020JJ2040).

E-mail addresses: y.j.guo@csu.edu.cn, x.j.wang7@csu.edu.cn, yue.wu@strath.ac.uk.

noise [3], the existence of random solutions generated by non-autonomous SPDEs with additive noise [4], periodic measures and ergodicity [5], etc. Note that almost all the sequel works are based on the definition of pathwise random periodic solutions for semiflows of random dynamical systems.

As the random periodic solution is not constructed explicitly, it is useful to study its numerical approximation. This is an infinite time horizon problem. The relevant research on the numerics of random periodic solutions for SDEs has made rapid progress recently [1,8–10,12,14,15]. The first paper [2] to approximate the random period trajectory considered an Euler-Maruyama method and a modified Milstein method for a dissipative system with a global Lipschitz condition. Wei and Chen [16] later generalised the Euler-Maruyama method to the stochastic theta method by showing that the approximation converges to the exact one at an order 1/4. Wu [17] studied the existence and uniqueness of the random periodic solution for an additive SDE with a one-sided Lipschitz condition and gave the analysis of the order-half convergence of its numerical approximation via the backward Euler method.

In this paper, we mainly consider the strong convergence rate of the backward Euler method for the random periodic solution of semilinear SDE with a one-sided Lipschitz condition. Our contribution is two-folded:

- Improved L^p -estimates of the random periodic solutions of the considered SDEs can be guaranteed under a more relaxed condition compared to [17];
- The order of convergence of the backward Euler method used to approximate random periodic solutions can be lifted from half in [17] to one for SDEs with an additive noise.

The outline of the paper is as follows. In section 2, we present some standard notation and assumptions that will be employed in our proofs, and give the existence and uniqueness of the random periodic solution. In Section 3we consider the L^p -estimates of the random periodic solutions under a relaxed condition compared to [17]. Section 4 is devoted to the error analysis of order-one convergence of the backward Euler method, where the supporting evidence is shown via numerical experiments in Section 5.

2 Random Periodic Solutions of SDEs

Let us recall the definition of the random periodic solution for stochastic semi-flows given in [6]. Let X be a separable Banach space. Denote by $(\Omega, \mathcal{F}, \mathbb{P}, (\theta_s)_{s \in \mathbb{R}})$ a metric dynamical system and $\theta_s : \Omega \to \Omega$ is assumed to be a measurably invertible for all $s \in \mathbb{R}$. Denote $\Delta := \{(t, s) \in \mathbb{R}^2, s \leq t\}$. Consider a stochastic periodic semi-flow $u : \Delta \times \Omega \times X \to X$ of period τ , which satisfies the following standard condition

$$u(t, r, \omega) = u(t, s, \omega) \circ u(s, r, \omega) \quad a.e. \ \omega \in \Omega,$$

$$(2.1)$$

for all $r \leq s \leq t, r, s, t \in \mathbb{R}$.

Definition 2.1. Given $\tau > 0$. A random periodic solution of period τ of a semi-flow $u : \Delta \times \Omega \times X \to X$ is an \mathcal{F} -measurable map $Y : \mathbb{R} \times \Omega \to X$ such that

$$u(t-s, s, Y(s, \omega), \omega) = Y(t, \omega) \text{ and } Y(t+\tau, \omega) = Y(t, \theta_{\tau}\omega)$$
(2.2)

for any $(t,s) \in \Delta$ and a.e. $\omega \in \Omega$.

Throughout this paper the following notation is frequently used. First, we have $d \in \mathbb{N}$. Denote $[d] := \{1, ..., d\}$ and let the letter C to denote a generic positive constant independent of the time and temporal stepsize of the numerical scheme we will introduce later. Let $|\cdot|, ||\cdot||$ and $\langle \cdot, \cdot \rangle$ be the absolute value of a scalar, the Euclidean norm and the inner product of vectors in \mathbb{R}^d , respectively. By A^T we denote the transpose of vector or matrix. Given a matrix A, we use $||A|| := \sqrt{trace(A^T A)}$ to denote the trace norm of A. On a canonical probability space $(\Omega, \mathcal{F}, \mathbb{P})$, we use \mathbb{E} to denote the expectation and $L^p(\Omega; \mathbb{R}^d)$ to denote the family of \mathbb{R}^d -valued variables with the norm defined by $\|\xi\|_{L^p(\Omega; \mathbb{R}^d)} = (\mathbb{E}[||\xi||^p])^{\frac{1}{p}} < \infty$. For a integrable random variable X on $(\Omega, \mathcal{F}, \mathbb{P})$ and for \mathcal{G} a σ -algebra such that $\mathcal{G} \subset \mathcal{F}$, we use $\mathbb{E}[X|\mathcal{G}]$ to denote the conditional expectation. Let $W : \mathbb{R} \times \Omega \to \mathbb{R}^d$ be a standard two-sided Wiener process on $(\Omega, \mathcal{F}, \mathbb{P})$. The filtration is defined as $\mathcal{F}_s^t := \sigma\{W_u - W_v : s \leq v \leq u \leq t\}$ and $\mathcal{F}^t = \mathcal{F}_{-\infty}^t = \bigvee_{s \leq t} \mathcal{F}_s^t$. Denote the standard \mathbb{P} -preserving ergodic Wiener shift by $\theta : \mathbb{R} \times \Omega \to \Omega$, which has the property $\theta_t(\omega)(s) := W_{t+s} - W_t$ for any $s, t \in \mathbb{R}$.

Given a standard two-sided Wiener process W on a metric dynamical system $(\Omega, \mathcal{F}, \mathbb{P}, (\theta_s)_{s \in \mathbb{R}})$, where θ is a \mathbb{P} -preserving ergodic Wiener shift, we revisit the long time behaviour of the following stochastic differential equation with an additive noise:

$$\begin{cases} dX_t^{t_0} = \left(-\Lambda X_t^{t_0} + f(t, X_t^{t_0}) \right) dt + g(t) \, dW(t), \quad t \in (t_0, T], \\ X_{t_0}^{t_0} = \xi, \end{cases}$$
(2.3)

where $f : \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}^d$, $g : \mathbb{R} \to \mathbb{R}$ and Λ is $d \times d$ matrix, and reconsider its numerical treatment. Note that we use $X_{t_1}^{t_0}$ to emphasise a process X evaluated at t_1 which starts from t_0 . The random initial value ξ is assumed to be \mathcal{F}^{t_0} -measurable. Further conditions on the initial value, the drift and diffusion coefficients are collected below.

Assumption 2.2. Suppose the following conditions are satisfied.

(i) Λ is self-adjoint and positive definite operator and there exists a non-decreasing sequence $(\lambda_i)_{i \in [d]} \subset \mathbb{R}$ of positive real numbers and an orthonormal basis $(e_i)_{i \in [d]}$, such that

$$\Lambda e_i = \lambda_i e_i, \tag{2.4}$$

for every $i \in [d]$.

(ii) The drift coefficient functions f is continuous and $f(t,x) = f(t+\tau,x)$. There exists a constant $0 < c_f < \lambda_1$ such that for any $x, y \in \mathbb{R}^d$ and $t \in [0, \tau)$

$$\langle x - y, f(t, x) - f(t, y) \rangle \leq c_f ||x - y||^2, \langle x, f(t, x) \rangle \leq c_f (1 + ||x||^2).$$

$$(2.5)$$

(iii) The diffusion coefficient functions g is continuous and $g(t) = g(t + \tau)$ and there exists a constant $c_q > 0$ such that $\sup_{t \in [0,\tau)} ||g(t)|| \le c_q$ and

$$\|g(t_1) - g(t_2)\| \le c_g |t_2 - t_1|, \quad \forall t_1, t_2 \in [0, \tau).$$
(2.6)

(iv) There exists a constant $C^* > 0$ such that $\mathbb{E}[\|\xi\|^2] \leq C^*$.

Assumption 2.3. There exists a constant $\tilde{c_f}$ such that

$$\left\| f(t,x) - \frac{\langle f(t,x), x \rangle}{\|x\|^2} x \right\| \le \tilde{c}_f (1 + \|x\|^2),$$
(2.7)

for $x \in \mathbb{R}^d, t \in [0, \tau)$.

Following a similar argument as in [13, Proposition 7.1], the SDE (2.3) admits a unique global semiflow under Assumption 2.2 and Assumption 2.3. Note that by the variation of constant formula, the solution of (2.3) is written as

$$X_t^{t_0}(\xi) = e^{-\Lambda(t-t_0)}\xi + \int_{t_0}^t e^{-\Lambda(t-s)}f(s, X_s^{t_0})\,\mathrm{d}s + \int_{t_0}^t e^{-\Lambda(t-s)}g(s)\,\mathrm{d}W_s.$$
 (2.8)

These assumptions can ensure the existence and uniqueness of the random periodic solution, which has been proposed in [17, Theorem 7].

Theorem 2.4. Let Assumptions 2.2 and Assumptions 2.3 be hold, then there exists a unique random periodic solution $X_t^*(\cdot) \in L^2(\Omega)$ with the form

$$X_t^* = \int_{-\infty}^t e^{-\Lambda(t-s)} f(s, X_s^*) \, ds + \int_{-\infty}^t e^{-\Lambda(t-s)} g(s) \, dW_s \tag{2.9}$$

such that X^* is a limit of the pull-back $X_t^{-k\tau}(\xi)$ of (2.3) when $k \to \infty$, ie,

$$\lim_{k \to \infty} \mathbb{E} \left[\|X_t^{-k\tau}(\xi) - X_t^*\|^2 \right] = 0.$$
(2.10)

3 Improved L^p -estimates of the random periodic solutions

In this section, our objective is to demonstrate the time-uniform moment boundedness of the random periodic solutions. While the work of Wu (2022) established the uniform boundedness for the *p*-th moment of the SDE solution as referenced in [17], our contribution lies in surpassing the limitation imposed on the drift functions (see Remark 3.3 for details). To accomplish this objective, we begin by deriving a generalized lemma, building upon the principles outlined in [7, Lemma 8.1].

Lemma 3.1. Let $m : [a, \infty) \to [0, \infty), \psi : [a, \infty) \to [0, \infty)$ be nonnegative continuous functions for $a \in \mathbb{R}$. If there exists a positive constant δ such that

$$m(t) - m(s) \le -\delta \int_s^t m(u) \, du + \int_s^t \psi(u) \, du \tag{3.1}$$

for any $a \leq s < t < \infty$, then

$$m(t) \le m(a) + \int_{a}^{t} e^{-\delta(t-u)} \psi(u) \, du.$$
 (3.2)

Proof of Lemma 3.1. Denote $m_1(t) := m(a) + \int_a^t e^{-\delta(t-u)} \psi(u) \, du$. It is easy to derive that

$$m'_{1}(t) = -\delta \int_{a}^{t} e^{-\delta(t-u)} \psi(u) \, \mathrm{d}u + \psi(t)$$

= $-\delta(m_{1}(t) - m(a)) + \psi(t).$ (3.3)

So we get

$$m_1(t) - m_1(s) = -\delta \int_s^t (m_1(u) - m(a)) \,\mathrm{d}u + \int_s^t \psi(u) \,\mathrm{d}u.$$
(3.4)

Now set $m_2(t) = m(t) - m_1(t)$. Due to (3.1), (3.4) and noting $m(a) \ge 0$, we have

$$m_{2}(t) - m_{2}(s) = m(t) - m(s) - (m_{1}(t) - m_{1}(s))$$

$$\leq -\delta \int_{s}^{t} (m(u) - m_{1}(u) + m(a)) du$$

$$\leq -\delta \int_{s}^{t} (m(u) - m_{1}(u)) du$$

$$= -\delta \int_{s}^{t} m_{2}(u) du.$$
(3.5)

To prove (3.2), it suffices to show that $m_2(t) \leq 0$ for any $t \in \mathbb{R}$. If $m_2(t) > 0$ for some t, then $m_2(a) = 0$ implies that there exists an interval $[s_1, t_1] \subset [a, \infty]$, $m_2(t_1) > m_2(s_1)$ and $m_2 > 0$ on $[s_1, t_1]$, which contradict (3.5).

We now give the time-uniform moment boundedness of the exact solution of SDE (2.3).

Theorem 3.2. Let Assumptions 2.2 hold. Let $X_t^{-k\tau}$ be the solution of SDE (2.3) with the initial value $X_{-k\tau}^{-k\tau} = \xi$ obeying $\mathbb{E}[\|\xi\|^{2p}] < \infty$ for any $p \in [1, \infty)$. Then there exists a positive constant C such that

$$\sup_{k\in\mathbb{N}}\sup_{t\geq -k\tau}\mathbb{E}\left[\left\|X_{t}^{-k\tau}\right\|^{2p}\right]\leq C\left(1+\mathbb{E}\left[\left\|\xi\right\|^{2p}\right]\right).$$
(3.6)

Proof of Theorem 3.2. Using the Itô formula gives

$$(1 + ||X_t^{-k\tau}||^2)^p = (1 + ||\xi||^2)^p + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, -\Lambda X_s^{-k\tau} \rangle \,\mathrm{d}s + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, f(s, X_s^{-k\tau}) \rangle \,\mathrm{d}s + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, g(s) \,\mathrm{d}W_s \rangle$$

$$+ p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} ||g(s)||^2 \,\mathrm{d}s + 2p(p-1) \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-2} ||(X_s^{-k\tau})^T g(s)||^2 \,\mathrm{d}s.$$

$$(3.7)$$

This directly leads to

$$(1 + ||X_t^{-k\tau}||^2)^p \leq (1 + ||\xi||^2)^p + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, -\Lambda X_s^{-k\tau} \rangle \,\mathrm{d}s + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, f(s, X_s^{-k\tau}) \rangle \,\mathrm{d}s + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, g(s) \,\mathrm{d}W_s \rangle + p(2p-1) \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} ||g(s)||^2 \,\mathrm{d}s.$$
 (3.8)

Combining this with (2.4) and (2.5) we can get

$$(1 + \|X_t^{-k\tau}\|^2)^p \leq (1 + \|\xi\|^2)^p + 2p \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^{p-1} (-\lambda_1 (1 + \|X_s^{-k\tau}\|^2)) ds + 2pc_f \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^p ds + 2p \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^{p-1} \langle X_s^{-k\tau}, g(s) dW_s \rangle + p((2p-1)c_g^2 + 2\lambda_1) \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^{p-1} ds \leq (1 + \|\xi\|^2)^p - 2p(\lambda_1 - c_f) \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^p ds + 2p \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^{p-1} \langle X_s^{-k\tau}, g(s) dW_s \rangle + p((2p-1)c_g^2 + 2\lambda_1) \int_{-k\tau}^t (1 + \|X_s^{-k\tau}\|^2)^{p-1} ds,$$

$$(3.9)$$

where

$$p((2p-1)c_g^2 + 2\lambda_1) (1 + ||X_s^{-k\tau}||^2)^{p-1} = p(\lambda_1 - c_f) (1 + ||X_s^{-k\tau}||^2)^{p-1} \times \frac{(2p-1)c_g^2 + 2\lambda_1}{\lambda_1 - c_f}$$

$$\leq p(\lambda_1 - c_f) \times \frac{p-1}{p} (1 + ||X_s^{-k\tau}||^2)^{p-1} + p(\lambda_1 - c_f) \times \frac{1}{p} \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(\lambda_1 - c_f)^p}$$

$$= (p-1)(\lambda_1 - c_f) (1 + ||X_s^{-k\tau}||^2)^p + \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(\lambda_1 - c_f)^{p-1}}$$
(3.10)

by the Young inequality $a^{p-1}b \leq \frac{p-1}{p}a^p + \frac{1}{p}b^p$. Therefore

$$(1 + ||X_t^{-k\tau}||^2)^p \leq (1 + ||\xi||^2)^p - (p+1)(\lambda_1 - c_f) \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^p \,\mathrm{d}s + \int_{-k\tau}^t \frac{((2p-1)c_g^2 + 2\lambda_1))^p}{(\lambda_1 - c_f)^{p-1}} \,\mathrm{d}s + 2p \int_{-k\tau}^t (1 + ||X_s^{-k\tau}||^2)^{p-1} \langle X_s^{-k\tau}, g(s) \,\mathrm{d}W_s \rangle.$$

$$(3.11)$$

For every interval $n \geq 1$, define the stopping time

$$\tau_n = \inf\{s \in [-k\tau, t] : \|X_s^{-k\tau}\| \ge n\}.$$
(3.12)

Clearly, $\tau_n \uparrow t$ a.s.. Moreover, it follows from (3.11) and the property of the Itô integral that

$$\mathbb{E}[(1 + \|X_{t \wedge \tau_n}^{-k\tau}\|^2)^p] \leq \mathbb{E}[(1 + \|\xi\|^2)^p] - (p+1)(\lambda_1 - c_f)\mathbb{E}\Big[\int_{-k\tau}^{t \wedge \tau_n} (1 + \|X_s^{-k\tau}\|^2)^p \,\mathrm{d}s\Big] \\
+ \mathbb{E}\Big[\int_{-k\tau}^{t \wedge \tau_n} \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(\lambda_1 - c_f)^{p-1}} \,\mathrm{d}s\Big].$$
(3.13)

Letting $n \to \infty$ and by the Fatou's lemma, we have

$$\mathbb{E}[(1 + \|X_t^{-k\tau}\|^2)^p] \leq \mathbb{E}[(1 + \|\xi\|^2)^p] - (p+1)(\lambda_1 - c_f) \int_{-k\tau}^t \mathbb{E}\Big[(1 + \|X_s^{-k\tau}\|^2)^p\Big] ds + \int_{-k\tau}^t \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(\lambda_1 - c_f)^{p-1}} ds.$$
(3.14)

Now applying Lemma 3.1 to (3.14) with $\delta = (p+1)(\lambda_1 - c_f) > 0$ and $\psi = \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(\lambda_1 - c_f)^{p-1}}$ gives

$$\mathbb{E}[(1 + \|X_t^{-k\tau}\|^2)^p] \leq \mathbb{E}[(1 + \|\xi\|^2)^p] + \int_{-k\tau}^t e^{-(p+1)(\lambda_1 - c_f)(t-u)} \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(\lambda_1 - c_f)^{p-1}} \, \mathrm{d}u \\
\leq \mathbb{E}[(1 + \|\xi\|^2)^p] + \frac{((2p-1)c_g^2 + 2\lambda_1)^p}{(p+1)(\lambda_1 - c_f)^p} (1 - e^{-(p+1)(\lambda_1 - c_f)(t+k\tau)}) \\
\leq C(1 + \mathbb{E}[\|\xi\|^{2p}]),$$
(3.15)

which completes the proof.

Remark 3.3. The above time-uniform moment boundedness was obtained under more relaxed conditions, compared to [17, Proposition 14], where the author additionally required $(c_f + \frac{(p-1)c_g}{2})(2 + p + 2^{p+1}) < p\lambda_1$ for some positive number $p \ge 4\gamma - 2$ (cf. [17, Assumption 13]). Instead, here we just require $0 < c_f < \lambda_1$, which significantly relaxes the aforementioned condition.

4 The order-one convergence of the backward Euler method

This section is devoted to the error analysis of the strong convergence rate of the backward Euler approximation to SDE (2.3), where we lift the order of convergence to one from half as obtained in [17]. Take an equidistant partition $\mathcal{T}^h := \{jh, j \in \mathbb{Z}\}$, such that $h \in (0, 1)$. Note that \mathcal{T}^h stretch along the real line because we are dealing with an infinite time horizon problem. The backward Euler method applied to SDE (2.3) takes the following form:

$$\tilde{X}_{-k\tau+jh}^{-k\tau} = \tilde{X}_{-k\tau+(j-1)h}^{-k\tau} - \Lambda h \tilde{X}_{-k\tau+jh}^{-k\tau} + h f \left(jh, \tilde{X}_{-k\tau+jh}^{-k\tau} \right) + g((j-1)h) \Delta W_{-k\tau+(j-1)h}$$
(4.1)

for all $j \in \mathbb{N}$, where $\Delta W_{-k\tau+(j-1)h} := W_{-k\tau+jh} - W_{-k\tau+(j-1)h}$, and the initial value $\tilde{X}_{-k\tau}^{-k\tau} = \xi$. Because of the periodicity of f and g, we have that $f(-k\tau + jh, \tilde{X}_{-k\tau+jh}^{-k\tau}) = f(jh, \tilde{X}_{-k\tau+jh}^{-k\tau})$, $g(-k\tau + jh) = g(jh)$.

Before proceeding to the assumptions and the main proof of the error analysis, we present some existing and essential results regarding (4.1) from literature. The uniform bounds for the second moment of the numerical approximation have been established in [17, Lemma 17] as follows.

Proposition 4.1. Let Assumptions 2.2 be satisfied. Then there exists a constant $\tilde{C} > 0$ such that

$$\sup_{k,j\in\mathbb{N}} \mathbb{E}[\|\tilde{X}_{-k\tau+jh}^{-k\tau}\|^2] \le \tilde{C},\tag{4.2}$$

where $\{\tilde{X}^{-k\tau}_{-k\tau+jh}\}_{k,j\in\mathbb{N}}$ is given by (4.1).

The existence and uniqueness of random periodic solutions associated to the backward Euler method has been guaranteed in [17].

Theorem 4.2. Let Assumptions 2.2 be satisfied. For $h \in (0,1)$, the time domain is divided as \mathcal{T}^h . The backward Euler method (4.1) admits a random period solution $\tilde{X}^* \in L^2(\Omega)$ such that

$$\lim_{k \to \infty} \mathbb{E} \left[\| \tilde{X}_{-k\tau+jh}^{-k\tau}(\xi) - \tilde{X}^* \|^2 \right] = 0.$$
(4.3)

In order to establish a strong convergence rate as high as one of the backward Euler method, we need the following conditions on the drift as well as the initial condition besides Assumption 2.2.

Assumption 4.3. Assume the drift coefficient functions $f : \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}^d$ are continuously differentiable, and there exists a constant $\gamma \in [1, \infty)$ such that

$$\left\| \left(\frac{\partial f}{\partial x}(t,x) - \frac{\partial f}{\partial x}(t,\bar{x}) \right) y \right\| \le C(1 + \|x\| + \|\bar{x}\|)^{\gamma-2} \|x - \bar{x}\| \|y\|, \quad \forall x, \bar{x}, y \in \mathbb{R}^d,$$

$$(4.4)$$

$$||f(t,x) - f(s,x)|| \le C(1 + ||x||^{\gamma})|t - s|, \quad \forall x \in \mathbb{R}^d, s, t \in [0,\tau),$$
(4.5)

and $\|\xi\|_{L^{\max\{4\gamma, 8\gamma-8\}}(\Omega; \mathbb{R}^d)} < \infty.$

Note that the condition (4.4) implies

$$\left\|\frac{\partial f}{\partial x}(t,x)y\right\| \le C(1+\|x\|)^{\gamma-1}\|y\|, \quad \forall x,y \in \mathbb{R}^d,$$
(4.6)

which in turn implies

$$\|f(t,x) - f(t,\bar{x})\| \le C(1+\|x\|+\|\bar{x}\|)^{\gamma-1}\|x-\bar{x}\|, \quad \forall x,\bar{x} \in \mathbb{R}^d,$$
(4.7)

$$\|f(t,x)\| \le C(1+\|x\|)^{\gamma}, \quad \forall x \in \mathbb{R}^d.$$

$$(4.8)$$

We now present two lemmas with estimates which play important role in proving the order of convergence of the backward Euler method in Theorem 4.6. The first one is the Hölder continuity of the exact solution of (2.3) with respect to the norm in $L^p(\Omega; \mathbb{R}^d)$ and the second one is about the regularity of f.

Lemma 4.4. Let Assumptions 2.2 and 4.3 be hold. Then there exists a positive constant C which depends on γ , d, Λ , f, g only, such that

$$\|X_{t_1}^{-k\tau} - X_{t_2}^{-k\tau}\|_{L^p(\Omega;\mathbb{R}^d)} \le C \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \ge -k\tau} \|X_t^{-k\tau}\|_{L^{p\gamma}(\Omega;\mathbb{R}^d)}^{\gamma}\Big) |t_2 - t_1| + C|t_2 - t_1|^{\frac{1}{2}}, \tag{4.9}$$

for all $t_1, t_2 \ge -k\tau$ and $p \in [2, \infty)$, where $X_t^{-k\tau}$ denotes the exact solution to the SDE (2.3).

Proof of Lemma 4.4. Without loss of generality we set $t_1 \leq t_2$ and get

$$\|X_{t_{1}}^{-k\tau} - X_{t_{2}}^{-k\tau}\|_{L^{p}(\Omega;\mathbb{R}^{d})} = \left\|\int_{t_{1}}^{t_{2}} \left(-\Lambda X_{r}^{k\tau} + f(r, X_{t}^{-k\tau}) \,\mathrm{d}r + \int_{t_{1}}^{t_{2}} g(r) \,\mathrm{d}W_{r}\right\|_{L^{p}(\Omega;\mathbb{R}^{d})}$$

$$\leq \left\|\int_{t_{1}}^{t_{2}} \left(\Lambda X_{r}^{-k\tau} + f(r, X_{t}^{-k\tau})\right) \,\mathrm{d}r\right\|_{L^{p}(\Omega;\mathbb{R}^{d})}$$

$$+ \left\|\int_{t_{1}}^{t_{2}} g(r) \,\mathrm{d}W_{r}\right\|_{L^{p}(\Omega;\mathbb{R}^{d})}.$$
(4.10)

For the first term, we can get the following estimate using the Hölder inequality, (3.6) and (4.8)

$$\begin{aligned} \left\| \int_{t_1}^{t_2} \left(-\Lambda X_r^{-k\tau} + f(r, X_t^{-k\tau}) \right) \mathrm{d}r \right\|_{L^p(\Omega; \mathbb{R}^d)} \\ & \leq \int_{t_1}^{t_2} \|\Lambda X_r^{-k\tau}\|_{L^p(\Omega; \mathbb{R}^d)} \mathrm{d}r + \int_{t_1}^{t_2} \|f(r, X_t^{-k\tau})\|_{L^p(\Omega; \mathbb{R}^d)} \mathrm{d}r \\ & \leq C \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \ge -k\tau} \|X_t^{-k\tau}\|_{L^{p\gamma}(\Omega; \mathbb{R}^d)}^{\gamma} \Big) |t_2 - t_1|. \end{aligned}$$
(4.11)

Applying the Burkholder-Davis-Gundy inequality to the last term of (4.10) gives

$$\left\|\int_{t_1}^{t_2} g(r) \,\mathrm{d}W_r\right\|_{L^p(\Omega;\mathbb{R}^d)} \le C \left(\int_{t_1}^{t_2} \|g(r)\|_{L^p(\Omega;\mathbb{R}^d)}^2 \,\mathrm{d}r\right)^{\frac{1}{2}} \le C|t_2 - t_1|^{\frac{1}{2}}.$$
(4.12)

This completes proof.

Lemma 4.5. Let Assumptions 2.2 and 4.3 be satisfied. Consider the exact solution $X_t^{-k\tau}$ of SDE (2.3) over $[-k\tau, T]$ for arbitrary $k \in \mathbb{N}$ and $T \geq -k\tau$. Then there exists a positive constant C which depends on γ, d, Λ, f, g only, such that for all $t_1, t_2 \in [-k\tau, T]$ and $s \in [t_1, t_2]$, it holds that

$$\begin{aligned} \left\| f(s, X_s^{-k\tau}) - f(t_2, X_{t_2}^{-k\tau}) \right\|_{L^2(\Omega; \mathbb{R}^d)} &\leq C \left(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \| X_t^{-k\tau} \|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{\gamma-1} \right) |t_2 - t_1|^{\frac{1}{2}} \\ &+ C \left(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \| X_t^{-k\tau} \|_{L^{4\gamma-2}(\Omega; \mathbb{R}^d)}^{2\gamma-1} \right) |t_2 - t_1|. \end{aligned}$$

$$(4.13)$$

Proof of Lemma 4.5. For all $s, t_2 \in [-k\tau, T]$, it follows from (4.7) and (4.8) that

$$\begin{aligned} \left\| f(s, X_s^{-k\tau}) - f(t_2, X_{t_2}^{-k\tau}) \right\| &= \left\| f(s, X_s^{-k\tau}) - f(s, X_{t_2}^{-k\tau}) + f(s, X_{t_2}^{-k\tau}) - f(t_2, X_{t_2}^{-k\tau}) \right\| \\ &\leq \left\| f(s, X_s^{-k\tau}) - f(s, X_{t_2}^{-k\tau}) \right\| + \left\| f(s, X_{t_2}^{-k\tau}) - f(t_2, X_{t_2}^{-k\tau}) \right\| \\ &\leq C(1 + \left\| X_s^{-k\tau} \right\| + \left\| X_{t_2}^{-k\tau} \right\|)^{\gamma - 1} \left\| X_{t_2}^{-k\tau} - X_s^{-k\tau} \right\| \\ &+ C(1 + \left\| X_{t_2}^{-k\tau} \right\|^{\gamma}) |t_2 - s|. \end{aligned}$$
(4.14)

Taking the expectation on both sides and using the Hölder inequality $\|f^{\gamma-1}g\|_{L^2(\Omega;\mathbb{R}^d)} \leq \|f\|_{L^{2\rho_1(\gamma-1)}(\Omega;\mathbb{R}^d)}^{\gamma-1}$

 $+ \|g\|_{L^{2\rho_2}(\Omega;\mathbb{R}^d)} \left(\frac{1}{2\rho_1} + \frac{1}{2\rho_2} = 1\right)$ with exponents $\rho_1 =: \frac{2\gamma - 1}{\gamma - 1}$ and $\rho_2 =: \frac{2\gamma - 1}{\gamma}$ yield that

$$\begin{split} \left\| f(s, X_{s}^{-k\tau}) - f(t_{2}, X_{t_{2}}^{-k\tau}) \right\|_{L^{2}(\Omega; \mathbb{R}^{d})} \\ &\leq C \left\| (1 + \|X_{s}^{-k\tau}\| + \|X_{t_{2}}^{-k\tau}\|)^{\gamma-1} \|X_{t_{2}}^{-k\tau} - X_{s}^{-k\tau}\| \right\|_{L^{2}(\Omega; \mathbb{R})} \\ &+ C \left\| (1 + \|X_{t_{2}}^{-k\tau}\|^{\gamma})|t_{2} - s| \right\|_{L^{2}(\Omega; \mathbb{R})} \\ &\leq C \left(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \|X_{t}^{-k\tau}\|_{L^{2\rho_{1}(\gamma-1)}(\Omega; \mathbb{R}^{d})}^{\gamma-1} \right) \|X_{t_{2}}^{-k\tau} - X_{s}^{-k\tau}\|_{L^{2\rho_{2}}(\Omega; \mathbb{R}^{d})} \\ &+ C \left(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \|X_{t}^{-k\tau}\|_{L^{2\gamma}(\Omega; \mathbb{R}^{d})}^{\gamma} \right) |t_{2} - t_{1}|. \end{split}$$

$$(4.15)$$

Moreover, through Lemma 4.4 with $p = 2\rho_2$ we have that

$$\begin{aligned} \|X_{s}^{-k\tau} - X_{t_{2}}^{-k\tau}\|_{L^{2\rho_{2}}(\Omega;\mathbb{R}^{d})} &\leq C \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \|X_{t}^{-k\tau}\|_{L^{2\gamma\rho_{2}}(\Omega;\mathbb{R}^{d})}^{\gamma} \Big) |t_{2} - s| + C|t_{2} - s|^{\frac{1}{2}} \\ &\leq C \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \|X_{t}^{-k\tau}\|_{L^{4\gamma-2}(\Omega;\mathbb{R}^{d})}^{\gamma} \Big) |t_{2} - t_{1}| + C|t_{2} - t_{1}|^{\frac{1}{2}}. \end{aligned}$$
(4.16)

Note that $2\rho_1(\gamma - 1) = 4\gamma - 2$. Altogether, it follows that for $s \in [t_1, t_2]$

$$\begin{split} \left\| f(s, X_{s}^{-k\tau}) - f(t_{2}, X_{t_{2}}^{-k\tau}) \right\|_{L^{2}(\Omega; \mathbb{R}^{d})} \\ &\leq C \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \| X_{t}^{-k\tau} \|_{L^{4\gamma-2}(\Omega; \mathbb{R}^{d})}^{\gamma-1} \Big) |t_{2} - t_{1}|^{\frac{1}{2}} \\ &+ C \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \left(\| X_{t}^{-k\tau} \|_{L^{4\gamma-2}(\Omega; \mathbb{R}^{d})}^{2\gamma-1} \right) |t_{2} - t_{1}|. \end{split}$$

$$(4.17)$$

This thus finishes the proof of the lemma.

We are now ready to give the main result of this section that reveals the order-one convergence of the backward Euler scheme to the SDE (2.3) in the long run.

Theorem 4.6. Under Assumptions 2.2 and 4.3. For $h \in (0,1)$, the time domain is divided as \mathcal{T}^h . If $X_{-k\tau+jh}^{-k\tau}$ and $\tilde{X}_{-k\tau+jh}^{-k\tau}$ are the exact and the numerical solutions given by (2.3) and (4.1), respectively, then there exists a constant C that depends on the γ, Λ, f, g and d such that

$$\sup_{k,j} \|X_{-k\tau+jh}^{-k\tau} - \tilde{X}_{-k\tau+jh}^{-k\tau}\|_{L^2(\Omega;\mathbb{R}^d)} \le Ch.$$
(4.18)

Proof of Theorem 4.6. First note that

$$\begin{aligned} X_{-k\tau+jh}^{-k\tau} &= X_{-k\tau+(j-1)h}^{-k\tau} - \int_{-k\tau+(j-1)h}^{-k\tau+jh} \Lambda X_s^{-k\tau} \,\mathrm{d}s \\ &+ \int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_s^{-k\tau}) \,\mathrm{d}s + \int_{-k\tau+(j-1)h}^{-k\tau+jh} g(s) \,\mathrm{d}W_s \\ &= X_{-k\tau+(j-1)h}^{-k\tau} - \Lambda h X_{-k\tau+jh}^{-k\tau} + h f(jh, X_{-k\tau+jh}^{-k\tau}) \\ &+ g((j-1)h) \Delta W_{-k\tau+(j-1)h} + \mathcal{R}_j, \end{aligned}$$
(4.19)

where

$$\mathcal{R}_{j} := -\int_{-k\tau+(j-1)h}^{-k\tau+jh} \Lambda(X_{s}^{-k\tau} - X_{-k\tau+jh}^{-k\tau}) \,\mathrm{d}s + \int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{s}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \,\mathrm{d}s + \int_{-k\tau+(j-1)h}^{-k\tau+jh} g(s) - g((j-1)h) \,\mathrm{d}W_{s}.$$
(4.20)

Subtracting (4.1) from this yields

$$X_{-k\tau+jh}^{-k\tau} - \tilde{X}_{-k\tau+jh}^{-k\tau} = X_{-k\tau+(j-1)h}^{-k\tau} - \tilde{X}_{-k\tau+(j-1)h}^{-k\tau} - \Lambda h(X_{-k\tau+jh}^{-k\tau} - \tilde{X}_{-k\tau+jh}^{-k\tau}) + h(f(jh, X_{-k\tau+jh}^{-k\tau}) - f(jh, \tilde{X}_{-k\tau+jh}^{-k\tau})) + g((j-1)h)\Delta W_{-k\tau+(j-1)h} - g((j-1)h)\Delta W_{-k\tau+(j-1)h} + \mathcal{R}_j.$$
(4.21)

For brevity, we denote

$$D_j := X_{-k\tau+jh}^{-k\tau} - \tilde{X}_{-k\tau+jh}^{-k\tau}, \quad \Delta f_j := f(jh, X_{-k\tau+jh}^{-k\tau}) - f(jh, \tilde{X}_{-k\tau+jh}^{-k\tau}).$$
(4.22)

We emphasize that D_j and Δf_j are $\mathcal{F}_{jh-k\tau}$ -measurable. Using (4.22), (4.21) can be now rewritten as

$$D_j = D_{j-1} - \Lambda h D_j + h \Delta f_j + \mathcal{R}_j.$$
(4.23)

This leads to

$$||D_j + \Lambda h D_j - h \Delta f_j||^2 = ||D_{j-1} + \mathcal{R}_j||^2.$$
(4.24)

Using (2.4) and (2.5) gives

$$\begin{split} \|D_{j} + \Lambda h D_{j} - h \Delta f_{j}\|^{2} \\ &= \|D_{j}\|^{2} + 2h \langle D_{j}, \Lambda D_{j} \rangle - 2h \langle D_{j}, \Delta f_{j} \rangle + h^{2} \|\Lambda D_{j} - \Delta f_{j}\|^{2} \\ &\geq \|D_{j}\|^{2} + 2\lambda_{1}h \|D_{j}\|^{2} - 2c_{f}h \|D_{j}\|^{2} \\ &= \left(1 + 2(\lambda_{1} - c_{f})h\right) \|D_{j}\|^{2}. \end{split}$$

$$(4.25)$$

Meanwhile,

$$||D_{j-1} + \mathcal{R}_j||^2 = ||D_{j-1}||^2 + 2\langle D_{j-1}, \mathcal{R}_j \rangle + ||\mathcal{R}_j||^2.$$
(4.26)

As a result,

$$(1 + 2(\lambda_1 - c_f)h) \|D_j\|^2 \le \|D_{j-1}\|^2 + 2\langle D_{j-1}, \mathcal{R}_j \rangle + \|\mathcal{R}_j\|^2.$$
(4.27)

Denoting $v := \lambda_1 - c_f$ and taking expectation yield,

$$(1+2vh)\mathbb{E}[\|D_{j}\|^{2}] \leq \mathbb{E}[\|D_{j-1}\|^{2}] + 2\mathbb{E}[\langle D_{j-1}, \mathcal{R}_{j}\rangle] + \mathbb{E}[\|\mathcal{R}_{j}\|^{2}].$$
(4.28)

Recalling D_{j-1} is $\mathcal{F}_{(j-1)h-k\tau}$ -measurable, we deduce

$$\mathbb{E}[\langle D_{j-1}, \mathcal{R}_j \rangle] = \mathbb{E}\big[\mathbb{E}[\langle D_{j-1}, \mathcal{R}_j \rangle | \mathcal{F}_{(j-1)h-k\tau}]\big] = \mathbb{E}\big[\langle D_{j-1}, \mathbb{E}[\mathcal{R}_j | \mathcal{F}_{(j-1)h-k\tau}]\rangle\big].$$
(4.29)

Further, noting v > 0 by (2.5), and using the Cauchy-Schwartz inequality $2ab \le vha^2 + \frac{1}{vh}b^2$ for arbitrary positive h, we obtain

$$(1+2vh)\mathbb{E}[\|D_{j}\|^{2}] \leq \mathbb{E}[\|D_{j-1}\|^{2}] + 2\mathbb{E}[\langle \sqrt{vh}D_{j-1}, \frac{1}{\sqrt{vh}}\mathbb{E}[\mathcal{R}_{j}|\mathcal{F}_{(j-1)h-k\tau}]\rangle] + \mathbb{E}[\|\mathcal{R}_{j}\|^{2}] \\ \leq (1+vh)\mathbb{E}[\|D_{j-1}\|^{2}] + \mathbb{E}[\|\mathcal{R}_{j}\|^{2}] + \frac{1}{vh}\mathbb{E}[\|\mathbb{E}[\mathcal{R}_{j}|\mathcal{F}_{(j-1)h-k\tau}]\|^{2}].$$

$$(4.30)$$

Hence,

$$\mathbb{E}[\|D_{j}\|^{2}] \leq \frac{1+vh}{1+2vh} \mathbb{E}[\|D_{j-1}\|^{2}] + \frac{1}{1+2vh} \mathbb{E}[\|\mathcal{R}_{j}\|^{2}] + \frac{1}{(1+2vh)vh} \mathbb{E}[\|\mathbb{E}[\mathcal{R}_{j}|\mathcal{F}_{(j-1)h-k\tau}]\|^{2}] \\ = (1 - \frac{v}{1+2vh}h) \mathbb{E}[\|D_{j-1}\|^{2}] + \frac{1}{1+2vh} \mathbb{E}[\|\mathcal{R}_{j}\|^{2}] + \frac{1}{(1+2vh)vh} \mathbb{E}[\|\mathbb{E}[\mathcal{R}_{j}|\mathcal{F}_{(j-1)h-k\tau}]\|^{2}].$$
(4.31)

Therefore, we only need to estimate two error terms $\mathbb{E}[||\mathcal{R}_j||^2]$ and $\mathbb{E}[||\mathbb{E}[\mathcal{R}_j|\mathcal{F}_{(j-1)h-k\tau}]||^2]$. Recalling the definition of \mathcal{R}_j given by (4.20) and using an triangle inequality yield

$$\begin{aligned} \|\mathcal{R}_{j}\|_{L^{2}(\Omega;\mathbb{R}^{d})} &\leq \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} \Lambda(X_{s}^{-k\tau} - X_{-k\tau+jh}^{-k\tau}) \,\mathrm{d}s \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\ &+ \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{s}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \,\mathrm{d}s \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\ &+ \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} g(s) - g((j-1)h) \,\mathrm{d}W_{s} \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\ &=: \mathbb{I}_{1} + \mathbb{I}_{2} + \mathbb{I}_{3}. \end{aligned}$$
(4.32)

For the term \mathbb{I}_1 , following the Hölder inequality and (4.9) shows

$$\mathbb{I}_{1} \leq \int_{-k\tau+(j-1)h}^{-k\tau+jh} \|\Lambda(X_{s}^{-k\tau} - X_{-k\tau+jh}^{-k\tau})\|_{L^{2}(\Omega;\mathbb{R}^{d})} \,\mathrm{d}s \\
\leq Ch^{\frac{3}{2}} \Big(1 + \sup_{k\in\mathbb{N}} \sup_{t\geq -k\tau} \|X_{t}^{-k\tau}\|_{L^{2\gamma}(\Omega;\mathbb{R}^{d})}^{\gamma}\Big).$$
(4.33)

With the help of the Hölder inequality and (4.14), one can obtain

$$\mathbb{I}_{2} \leq \int_{-k\tau+(j-1)h}^{-k\tau+jh} \left\| f(s, X_{s}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \right\|_{L^{2}(\Omega; \mathbb{R}^{d})} \mathrm{d}s \\
\leq Ch^{\frac{3}{2}} \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \| X_{t}^{-k\tau} \|_{L^{4\gamma-2}(\Omega; \mathbb{R}^{d})}^{2\gamma-1} \Big).$$
(4.34)

In view of the Itô isomery and by Assumption 2.2 (iii)

$$\mathbb{I}_{3} = \left(\int_{-k\tau+(j-1)h}^{-k\tau+jh} \|g(s) - g((j-1)h)\|^{2} \,\mathrm{d}s\right)^{\frac{1}{2}} \le Ch^{\frac{3}{2}}.$$
(4.35)

Putting all the above estimates together we derive from (4.32) that

$$\|\mathcal{R}_{j}\|_{L^{2}(\Omega;\mathbb{R}^{d})} \leq Ch^{\frac{3}{2}} \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \geq -k\tau} \|X_{t}^{-k\tau}\|_{L^{4\gamma-2}(\Omega;\mathbb{R}^{d})}^{2\gamma-1} \Big).$$
(4.36)

Note that $\mathbb{E}\left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} g(s) - g((j-1)h) \, \mathrm{d}W_s | \mathcal{F}_{(j-1)h-k\tau}\right] = 0$. Thus

$$\begin{split} \|\mathbb{E}[\mathcal{R}_{j}|\mathcal{F}_{(j-1)h-k\tau}]\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\ &\leq \left\|\mathbb{E}\Big[\int_{-k\tau+(j-1)h}^{-k\tau+jh} -\Lambda(X_{s}^{-k\tau}-X_{-k\tau+jh}^{-k\tau})\,\mathrm{d}s\Big|\mathcal{F}_{(j-1)h-k\tau}\Big]\right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\ &+ \left\|\mathbb{E}\Big[\int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s,X_{s}^{-k\tau}) - f(jh,X_{-k\tau+jh}^{-k\tau})\,\mathrm{d}s\Big|\mathcal{F}_{(j-1)h-k\tau}\Big]\right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\ &=: \mathbb{I}_{4} + \mathbb{I}_{5}. \end{split}$$
(4.37)

In order to estimate \mathbb{I}_4 , we first show that

$$\mathbb{E}\left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_r \, \mathrm{d}s \left| \mathcal{F}_{(j-1)h-k\tau} \right] \\
= \int_{-k\tau+(j-1)h}^{-k\tau+jh} \mathbb{E}\left[\int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_r \left| \mathcal{F}_{(j-1)h-k\tau} \right] \, \mathrm{d}s \\
= \int_{-k\tau+(j-1)h}^{-k\tau+jh} \mathbb{E}\left[\mathbb{E}\left[\int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_r \left| \mathcal{F}_s \right] \left| \mathcal{F}_{(j-1)h-k\tau} \right] \, \mathrm{d}s \\
= 0.$$
(4.38)

As a result, we derive from (4.19) that

$$\mathbb{E}\left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} -\Lambda(X_s^{-k\tau} - X_{-k\tau+jh}^{-k\tau}) \,\mathrm{d}s \middle| \mathcal{F}_{(j-1)h-k\tau}\right] \\
= \mathbb{E}\left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} (-\Lambda^2 X_r^{-k\tau} + \Lambda f(r, X_r^{-k\tau})) \,\mathrm{d}r \,\mathrm{d}s\right].$$
(4.39)

Based on the Jensen inequality and the Hölder inequality, according to (3.6) and (4.8), one can infer

$$\mathbb{I}_{4} = \left\| \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} \left(-\Lambda^{2}X_{r}^{-k\tau} + \Lambda f(r, X_{r}^{-k\tau}) \right) dr ds \Big| \mathcal{F}_{(j-1)h-k\tau} \right] \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
\leq \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} \left(-\Lambda^{2}X_{r}^{-k\tau} + \Lambda f(r, X_{r}^{-k\tau}) \right) dr ds \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
\leq \int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} \left\| -\Lambda^{2}X_{r}^{-k\tau} + \Lambda f(r, X_{r}^{-k\tau}) \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} dr ds \\
\leq \int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} \left\| \Lambda^{2}X_{r}^{-k\tau} \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} dr ds \\
+ \int_{-k\tau+(j-1)h}^{-k\tau+jh} \int_{s}^{-k\tau+jh} \left\| \Lambda f(r, X_{r}^{-k\tau}) \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} dr ds \\
\leq Ch^{2} \left(1 + \left(\| X_{t}^{-k\tau} \|_{L^{2\gamma}(\Omega;\mathbb{R}^{d})}^{\gamma} \right) \right).$$
(4.40)

With regard to \mathbb{I}_5 , we first rewrite it as follows

$$\mathbb{I}_{5} = \left\| \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{s}^{-k\tau}) - f(j, X_{-k\tau+jh}^{-k\tau}) \, \mathrm{d}s \middle| \mathcal{F}_{(j-1)h-k\tau} \right] \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
= \left\| \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{s}^{-k\tau}) - f(s, X_{-k\tau+jh}^{-k\tau}) \, \mathrm{d}s \middle| \mathcal{F}_{(j-1)h-k\tau} \right] \\
+ \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{-k\tau+jh}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \, \mathrm{d}s \middle| \mathcal{F}_{(j-1)h-k\tau} \right] \right\|_{L^{2}(\Omega;\mathbb{R}^{d})}.$$
(4.41)

Because of the existence of the first derivative of f with respect to the spatial variable, for $s \in [-k\tau + (j-1)h, -k\tau + jh]$,

$$\begin{split} f(s, X_{-k\tau+jh}^{-k\tau+jh}) &- f(s, X_{s}^{-k\tau}) \\ &= \int_{0}^{1} \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \, \mathrm{d}\zeta \\ &= \int_{0}^{1} \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) \int_{s}^{-k\tau+jh} \big(-\Lambda X_{r}^{-k\tau} + f(r, X_{r}^{-k\tau}) \big) \, \mathrm{d}r \, \mathrm{d}\zeta \\ &+ \int_{0}^{1} \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) \int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_{r} \, \mathrm{d}\zeta \\ &= \underbrace{\int_{0}^{1} \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) \int_{s}^{-k\tau+jh} \big(-\Lambda X_{r}^{-k\tau} + f(r, X_{r}^{-k\tau}) \big) \, \mathrm{d}r \, \mathrm{d}\zeta \\ &= \underbrace{\int_{0}^{1} \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) \int_{s}^{-k\tau+jh} \big(-\Lambda X_{r}^{-k\tau} + f(r, X_{r}^{-k\tau}) \big) \, \mathrm{d}r \, \mathrm{d}\zeta \\ &= \underbrace{\int_{0}^{1} \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) - \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + f(r, X_{r}^{-k\tau}) \big) \, \mathrm{d}r \, \mathrm{d}\zeta \\ &+ \underbrace{\int_{0}^{1} \Big(\frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \big) - \frac{\partial f}{\partial x} \big(s, X_{s}^{-k\tau} \big) \int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_{r} \, \mathrm{d}\zeta \\ &= \underbrace{:\Phi_{2}} \end{aligned}$$

Following a similar argument as (4.38), we have that

$$\mathbb{E}\Big[\int_{-k\tau+(j-1)h}^{-k\tau+jh}\int_{0}^{1}\frac{\partial f}{\partial x}(s,X_{s}^{-k\tau})\int_{s}^{-k\tau+jh}g(r)\,\mathrm{d}W_{r}\,\mathrm{d}\zeta\,\mathrm{d}s\Big|\mathcal{F}_{(j-1)h-k\tau}\Big]=0.$$
(4.43)

Using the Jensen inequality and the Hölder inequality gives

$$\mathbb{I}_{5} \leq \left\| \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} \Phi_{1} \Big| \mathcal{F}_{(j-1)h-k\tau} \right] \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} + \left\| \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} \Phi_{2} \Big| \mathcal{F}_{(j-1)h-k\tau} \right] \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
+ \left\| \mathbb{E} \left[\int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{-k\tau+jh}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \, \mathrm{d}s \Big| \mathcal{F}_{(j-1)h-k\tau} \right] \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
\leq \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} \Phi_{1} \, \mathrm{d}s \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} + \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} \Phi_{2} \, \mathrm{d}s \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
+ \left\| \int_{-k\tau+(j-1)h}^{-k\tau+jh} f(s, X_{-k\tau+jh}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \, \mathrm{d}s \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \\
\leq \int_{-k\tau+(j-1)h}^{-k\tau+jh} \left\| \Phi_{1} \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \, \mathrm{d}s + \int_{-k\tau+(j-1)h}^{-k\tau+jh} \left\| \Phi_{2} \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \, \mathrm{d}s \\
+ \int_{-k\tau+(j-1)h}^{-k\tau+jh} \left\| f(s, X_{-k\tau+jh}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau}) \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \, \mathrm{d}s.$$

In the following, we cope with the last three terms separately. According to the Hölder inequality and (3.6), (4.6) and (4.8), we can get

$$\begin{split} \|\Phi_{1}\|_{L^{2}(\Omega;\mathbb{R}^{d})} &\leq \int_{0}^{1} \int_{s}^{-k\tau+jh} \left\|\frac{\partial f}{\partial x} \left(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau})\right) \right\|_{\times} \left(-\Lambda X_{r}^{-k\tau} + f(r, X_{r}^{-k\tau})\right) \|_{L^{2}(\Omega;\mathbb{R}^{d})} \,\mathrm{d}r \,\mathrm{d}\zeta \\ &\leq \int_{0}^{1} \int_{s}^{-k\tau+jh} \left\|C \left(1 + \left\|\zeta X_{-k\tau+jh}^{-k\tau} + (\zeta - 1)X_{s}^{-k\tau}\right\|\right)^{\gamma-1} \right. \tag{4.45} \\ &\qquad \times \left(1 + \left\|X_{r}^{-k\tau}\right\|\right)^{\gamma} \right\|_{L^{2}(\Omega;\mathbb{R})} \,\mathrm{d}r \,\mathrm{d}\zeta \\ &\leq Ch \left(1 + \sup_{k\in\mathbb{N}} \sup_{t\geq -k\tau} \left\|X_{t}^{-k\tau}\right\|_{L^{4\gamma-2}(\Omega;\mathbb{R}^{d})}^{2\gamma-1}\right). \end{split}$$

For the second term $\|\Phi_2\|_{L^2(\Omega;\mathbb{R}^d)}$, we can get the following estimate through the Hölder inequality,

Burkholder-Davis-Gundy inequality, (4.4) and (4.9),

$$\begin{split} \|\Phi_{2}\|_{L^{2}(\Omega;\mathbb{R}^{d})} &\leq \int_{0}^{1} \left\| \left(\frac{\partial f}{\partial x} \left(s, X_{s}^{-k\tau} + \zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau}) \right) - \frac{\partial f}{\partial x} \left(s, X_{s}^{-k\tau} \right) \right) \\ &\qquad \times \int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_{r} \right\|_{L^{2}(\Omega;\mathbb{R}^{d})} \, \mathrm{d}\zeta \\ &\leq C \int_{0}^{1} \left\| \left(1 + \|\zeta X_{-k\tau+jh}^{-k\tau} + (\zeta - 1)X_{s}^{-k\tau}\| + \|X_{s}^{-k\tau}\| \right)^{\gamma-2} \\ &\qquad \times \|\zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau})\| \right\|_{L^{4}(\Omega;\mathbb{R})} \left\| \int_{s}^{-k\tau+jh} g(r) \, \mathrm{d}W_{r} \right\|_{L^{4}(\Omega;\mathbb{R}^{d})} \, \mathrm{d}\zeta \qquad (4.46) \\ &\leq Ch^{\frac{1}{2}} \int_{0}^{1} \left\| \left(1 + \|\zeta X_{-k\tau+jh}^{-k\tau} + (\zeta - 1)X_{s}^{-k\tau}\| + \|X_{s}^{-k\tau}\| \right)^{\gamma-2} \\ &\qquad \times \|\zeta (X_{-k\tau+jh}^{-k\tau} - X_{s}^{-k\tau})\| \right\|_{L^{4}(\Omega;\mathbb{R})} \, \mathrm{d}\zeta \\ &\leq Ch \left(1 + \sup_{k\in\mathbb{N}} \sup_{t\geq -k\tau} \left(\|X_{t}^{-k\tau}\|_{L^{4\gamma}(\Omega;\mathbb{R}^{d})} \vee \|X_{t}^{-k\tau}\|_{L^{8\gamma-8}(\Omega;\mathbb{R}^{d})} \right) \right). \end{split}$$

Using (4.5), we can easily get

$$\|f(s, X_{-k\tau+jh}^{-k\tau}) - f(jh, X_{-k\tau+jh}^{-k\tau})\|_{L^{2}(\Omega; \mathbb{R}^{d})}$$

$$\leq Ch \Big(1 + \sup_{k \in \mathbb{N}} \sup_{t \ge -k\tau} \|X_{t}^{-k\tau}\|_{L^{2\gamma}(\Omega; \mathbb{R}^{d})}^{\gamma}\Big).$$

$$(4.47)$$

Inserting (4.45)-(4.47) into (4.44) implies

$$\mathbb{I}_{5} \leq Ch^{2} \Big(\Big(\|X_{t}^{-k\tau}\|_{L^{4\gamma}(\Omega;\mathbb{R}^{d})}^{\gamma} \vee \|X_{t}^{-k\tau}\|_{L^{8\gamma-8}(\Omega;\mathbb{R}^{d})}^{2\gamma-2} \Big) \Big).$$

$$(4.48)$$

Therefore, from (4.37) it immediately follows that

$$\|\mathbb{E}[\mathcal{R}_j|\mathcal{F}_{(j-1)h-k\tau}]\|_{L^2(\Omega;\mathbb{R}^d)} \le Ch^2 \Big(\big(\|X_t^{-k\tau}\|_{L^{4\gamma}(\Omega;\mathbb{R}^d)}^{\gamma} \vee \|X_t^{-k\tau}\|_{L^{8\gamma-8}(\Omega;\mathbb{R}^d)}^{2\gamma-2} \big) \Big).$$
(4.49)

Considering (4.31), define $\hat{v} := \frac{v}{1+2vh}$, we get

$$\mathbb{E}[\|D_{j}\|^{2}] \leq (1 - \hat{v}h)\mathbb{E}[\|D_{j-1}\|^{2}] + \frac{1}{1+2vh}Ch^{3}$$

$$\leq (1 - \hat{v}h)^{j}\mathbb{E}[\|D_{0}\|^{2}] + \sum_{i=0}^{j-1}(1 - \hat{v}h)^{i}Ch^{3}$$

$$= (1 - \hat{v}h)^{j}\mathbb{E}[\|D_{0}\|^{2}] + \frac{1 - (1 - \hat{v}h)^{j}}{\hat{v}h}Ch^{3}$$
(4.50)

By observing $D_0 = 0$,

$$\mathbb{E}[\|D_j\|^2] \le Ch^2. \tag{4.51}$$

 \Box

Then the assertion follows.

Corollary 4.7. Under Assumptions 2.2, let X_t^* be the random periodic solution of SDE (2.3) and \tilde{X}_t^* be the random periodic solution of the backward Euler numerical approximation. Then there exists a constant C that depends on q, Λ, f, g and d such that

$$\sup_{t \in \mathcal{T}^h} \mathbb{E}([\|X_t^* - \tilde{X}_t^*\|^2])^{1/2} \le Ch,$$
(4.52)

and \tilde{X}_t^* satisfies the random periodicity property.

Proof of Corollary 4.7. Due to

$$\|X_t^* - \tilde{X}_t^*\|^2 \le \limsup_k \left[\|X_t^* - X_t^{-k\tau}\|^2 + \|X_t^{-k\tau} - \tilde{X}_t^{-k\tau}\|^2 + \|\tilde{X}_t^{-k\tau} - \tilde{X}_t^*\|^2 \right],$$
(4.53)

thus the conclusion can be obtained by Theorem 2.4, Theorem 4.2 and Theorem 4.6.

Corollary 4.7 implies that the order-one convergence can be achieved.

5 Numerical experiments

In this Section, we present two numerical experiments to support the theoretical findings. To evaluate the effectiveness of the proposed numerical method under different scenarios, we vary the drift coefficients while maintaining a constant diffusion coefficient, ie, g(t) = 1.

5.1 Example 1

In the first example, we test the performance of the backward Euler method (4.1) on a linear drift coefficient as follows:

$$dX_t^{t_0} = -\pi X_t^{t_0} dt + \sin(2\pi t) dt + dW_t.$$
(5.1)

To fit into the general form of SDE in (2.3), we have that in this case

$$\Lambda = \pi$$
, and $f(t, X_t^{t_0}) = \sin(2\pi t)$. (5.2)

It is easy to verify that the associated period is 1 and Assumptions 2.2 are fulfilled with $\lambda_1 = \pi$, $c_f = 1$ and $c_g = 1$. According to Theorem 2.4, SDE (5.1) admits a unique random periodic solution. By Theorem 4.2, its backward Euler simulation also has a random periodic path.

We will first verify the periodicity by examining the dynamics of two processes under the same realisation ω : $\tilde{X}_t^{-5}(\omega, 0.3)$ over $-5 \leq t \leq 13$ and $\tilde{X}_t^{-5}(\theta_{-1}\omega, 0.3)$ over $-5 \leq t \leq 14$, where 0.3 is the initial condition of both processes. Because of Theorem 4.2, it is expected that $\tilde{X}_t^{-5}(\omega, 0.3) \approx \tilde{X}_t^*(\omega)$ and $\tilde{X}_t^{-5}(\theta_{-1}\omega, 0.3) \approx \tilde{X}_t^*(\theta_{-1}\omega)$ after a sufficiently long time, and we may then observe that $\tilde{X}_{t-1}^{-5}(\omega, 0.3) \approx \tilde{X}_t^{-5}(\theta_{-1}\omega, 0.3)$ due to the fact $\tilde{X}_{t-1}^*(\omega) = \tilde{X}_t^*(\theta_{-1}\omega)$ in Definition 2.1.

Figure 1 illustrates the evolution of processes before t = 0, where two trajectories separate from the initial time point and resemble each other with an increasing time gap. For example, it can be observed that $\tilde{X}_{-4.35}^{-5}(\omega, 0.3) \approx \tilde{X}_{-4.15}^{-5}(\theta_{-1}\omega, 0.3), \ \tilde{X}_{-2.5}^{-5}(\omega, 0.3) \approx \tilde{X}_{-1.9}^{-5}(\theta_{-1}\omega, 0.3)$ and



Figure 1: Simulations of the processes $\{\tilde{X}_t^{-5}(\omega, 0.3), -5 \le t \le -1\}$ and $\{\tilde{X}_t^{-5}(\theta_{-1}\omega, 0.3), -5 \le t \le 0\}$.



Figure 2: Simulations of the processes $\{\tilde{X}_t^{-5}(\omega, 0.3), 10 \le t \le 13\}$ and $\{\tilde{X}_t^{-5}(\theta_{-1}\omega, 0.3), 11 \le t \le 14\}$.



Figure 3: The mean-square error plot of the backward Euler method (4.1) for simulating the solution of (5.1).

 $\tilde{X}_{-1}^{-5}(\omega, 0.3) \approx \tilde{X}_{0}^{-5}(\theta_{-1}\omega, 0.3)$. Figure 2 demonstrates both processes resemble each other with a stable time gap 1, that is, $\tilde{X}_{t-1}^{-5}(\omega, 0.3) \approx \tilde{X}_{t}^{-5}(\theta_{-1}\omega, 0.3)$ over $11 \le t \le 14$.

Next, to test the strong convergence rate of backward Euler method, we simulate the solution of (5.1) over $t \in [-5, 15]$ with 5000 samples. The reference solution is obtained via the same numerical method at a fine stepsize $h_{exact} = 2^{-15} \times 20$. We plot in Figure 3 mean-square approximation errors D_h against six different stepsizes $h = 2^{-i} \times 20$, i = 7, 8, ..., 12 on a log-log scale.

From Figure 3, one can clearly observe that the mean-square error decreases at a slope close to 1, consistent with Theorem 4.6. Suppose that the approximation error D_h obeys a power law relation $D_h = Ch^{\kappa}$ for $C, \kappa > 0$, so that $\log D_h = \log C + \kappa \log h$. Then we do a least squares power law fit for κ and get the value 0.9836 for the rate κ with residual of 0.0697. Again, this confirms the expected convergence rate in Theorem 4.6.

5.2 Example 2

In the second example, we test the performance of the backward Euler method (4.1) on a cubic drift coefficient as follows:

$$dX_t^{t_0} = -2\pi X_t^{t_0} dt + (X_t^{t_0} - (X_t^{t_0})^3 + \cos(\pi t)) dt + dW_t.$$
(5.3)

Similarly, we run a similar experiment to verify the periodicity as described in Section 5.1 and present the identical patterns of $\tilde{X}_{t-2}^{-6}(\omega, 0.5)$ over $10 \le t \le 14$ and $\tilde{X}_t^{-6}(\theta_{-2}\omega, 0.5)$ over $12 \le t \le 16$ under the same realisation ω in Figure 4.



Figure 4: Simulations of the processes $\{\tilde{X}_t^{-6}(\omega, 0.5), 10 \le t \le 14\}$ and $\{\tilde{X}_t^{-6}(\theta_{-2}\omega, 0.5), 12 \le t \le 16\}$.

We also test the performance of the backward Euler method, in terms of mean-square error, in simulating SDE (5.3) over [-6, 14]. Figure 5 confirms an order-one convergence, again consistent with Theorem 4.6. A least squares fit produces a rate 0.9495 with residual of 0.1153 for (5.3). Hence, numerical result is consistent with strong order of convergence equal to one, as already revealed in 4.6.

References

- Kevin Burrage, PM Burrage, and Tianhai Tian. Numerical methods for strong solutions of stochastic differential equations: an overview. Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences, 460(2041):373-402, 2004.
- [2] Chunrong Feng, Yu Liu, and Huaizhong Zhao. Numerical approximation of random periodic solutions of stochastic differential equations. Zeitschrift f
 ür angewandte Mathematik und Physik, 68(5):119, 2017.
- [3] Chunrong Feng, Yue Wu, and Huaizhong Zhao. Anticipating random periodic solutions—I. SDEs with multiplicative linear noise. *Journal of Functional Analysis*, 271(2):365–417, 2016.
- [4] Chunrong Feng and Huaizhong Zhao. Random periodic solutions of SPDEs via integral equations and Wiener–Sobolev compact embedding. *Journal of Functional Analysis*, 262(10):4377– 4422, 2012.



Figure 5: The mean-square error plot of the backward Euler method (4.1) for simulating the solution of (5.3).

- [5] Chunrong Feng and Huaizhong Zhao. Random periodic processes, periodic measures and ergodicity. *Journal of Differential Equations*, 269(9):7382–7428, 2020.
- [6] Chunrong Feng, Huaizhong Zhao, and Bo Zhou. Pathwise random periodic solutions of stochastic differential equations. *Journal of Differential Equations*, 251(1):119–149, 2011.
- [7] Kiyosi Itô and Makiko Nisio. On stationary solutions of a stochastic differential equation. J. Math. Kyoto Univ, 4(1):1–75, 1964.
- [8] Grigori Noyhovich Mil'shtein. Approximate integration of stochastic differential equations. Theory of Probability & Its Applications, 19(3):557–562, 1975.
- [9] Bernt Øksendal and Bernt Øksendal. Stochastic Differential Equations. Springer, 2003.
- [10] Eckhard Platen. An introduction to numerical methods for stochastic differential equations. Acta numerica, 8:197–246, 1999.
- [11] Henri Poincaré. Mémoire sur les courbes définies par une équation différentielle. Journal de mathématiques pures et appliquées, 7:375–422, 1881.
- [12] W Rüemelin. Numerical treatment of stochastic differential equations. SIAM Journal on Numerical Analysis, 19(3):604–613, 1982.
- [13] Michael Scheutzow and Susanne Schulze. Strong completeness and semi-flows for stochastic differential equations with monotone drift. Journal of Mathematical Analysis and Applications, 446(2):1555–1570, 2017.

- [14] Xiaojie Wang. Mean-square convergence rates of implicit Milstein type methods for SDEs with non-Lipschitz coefficients. Advances in Computational Mathematics, 49(3): 37, 2023.
- [15] Xiaojie Wang, Jiayi Wu, and Bozhang Dong. Mean-square convergence rates of stochastic theta methods for SDEs under a coupled monotonicity condition. *BIT Numerical Mathematics*, 60(3):759–790, 2020.
- [16] Rong Wei and Chuan-zhong Chen. Numerical approximation of stochastic theta method for random periodic solution of stochastic differential equations. Acta Mathematicae Applicatae Sinica, English Series, 36(3):689–701, 2020.
- [17] Yue Wu. Backward Euler-Maruyama method for the random periodic solution of a stochastic differential equation with a monotone drift. *Journal of Theoretical Probability*, pages 1–18, 2022.
- [18] Huaizhong Zhao and Zuohuan Zheng. Random periodic solutions of random dynamical systems. Journal of Differential equations, 246(5):2020–2038, 2009.