A CHARACTERIZATION OF POSITROIDS, WITH APPLICATIONS TO AMALGAMS AND EXCLUDED MINORS

JOSEPH E. BONIN

ABSTRACT. A matroid of rank r on n elements is a positroid if it has a representation by an r by n matrix over \mathbb{R} , each r by r submatrix of which has nonnegative determinant. Earlier characterizations of connected positroids and results about direct sums of positroids involve connected flats and non-crossing partitions. We prove another characterization of positroids of a similar flavor and give some applications of the characterization. We show that if M and N are positroids and the intersection of their ground sets is an independent set and a set of clones in both M and N, then the free amalgam of M and N is a positroid, and we prove a second result of that type. Also, we identify several multi-parameter infinite families of excluded minors for the class of positroids.

1. INTRODUCTION

A *positroid* is a matroid M on a set E(M), say of rank r and with |E(M)| = n, that has a representation by an r by n matrix over \mathbb{R} , each r by r submatrix of which has nonnegative determinant. Since Blum [3] (as base-orderable matroids) and Postnikov [23] introduced positroids, many papers have developed this topic, spurred by the important connections with other branches of mathematics that led to the introduction of positroids as well as the wealth of lenses through which they can be viewed (e.g., Grassmann necklaces, reduced plabic graphs, decorated permutations).

Positroids are often defined to have a fixed linear order on the ground set, as is inherent in the columns of a matrix (and so E(M) is often taken to be $[n] = \{1, 2, ..., n\}$). With our focus on matroids per se, we find it useful to separate the matroid and the linear order, so in this paper, a positroid is a matroid M for which there is a linear order on the ground set E(M), which we call a *positroid order* for M, so that the elements, when put in that order, correspond to the columns of a matrix representation of the type described above.

Blum [3] and Ardila, Rincón, and Williams [2] treat some of the fundamental properties of the class of positroids. In particular, they show that the class of positroids is closed under the operations of minors, duals, and direct sums. Thus, a matroid is a positroid if and only if the restrictions to its connected components are positroids. This raises the issue of determining how to combine the positroid orders on the connected components to get a positroid order on the direct sum. Ardila, Rincón, and Williams settled this question using the well-studied notion of a *non-crossing partition*, that is, a partition $\{X_1, X_2, \ldots, X_k\}$ of a linearly ordered set E for which, for any two blocks X_i and X_j , there is a subset A of E such that $X_i \subseteq A, X_j \subseteq E - A$, and either A or E - A is an interval in the linear order. They proved the following result [2, Theorem 7.6].

Theorem 1.1. Let $X_1, X_2, ..., X_k$ be the connected components of a matroid M. For any positroid order for M, the partition $\{X_i : i \in [k]\}$ of E(M) is non-crossing. Conversely,

Date: June 18, 2024.

Key words and phrases. positroid, base-sortable matroid, amalgam, excluded minor.

JOSEPH E. BONIN

any linear order on E(M) for which the induced order on each X_i is a positroid order for $M|X_i$ and the partition $\{X_i : i \in [k]\}$ of E(M) is non-crossing is a positroid order for M.

That result along with the following characterization of positroid orders for connected matroids [2, Proposition 5.6] give a two-part characterization of positroid orders: Theorem 1.2 characterizes the orders on the connected components while Theorem 1.1 characterizes how the orders on those components are assembled globally.

Theorem 1.2. Let M be a connected matroid with $r(M) \ge 2$. A linear order \le on E(M) is a positroid order for M if and only if, for every flat F such that both M|F and M/F are connected, at least one of F and E(M) - F is an interval in \le .

The following similar characterization of positroid orders for connected matroids was given by Rincón, Vinzant, and Yu [24, Proposition 6.5], and likewise can be paired with Theorem 1.1 to characterize positroids.

Theorem 1.3. Let M be a rank-r, connected matroid. Fix an integer k with $1 < k \le r$. A linear order on E(M) is a positroid order for M if and only if, for each flag

$$\emptyset = F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \cdots \subsetneq F_{k-1} \subsetneq F_k = E(M)$$

of flats of M for which each minor $M|F_i/F_{i-1}$, for $i \in [k]$, is connected, the set of their ground sets, $\{F_i - F_{i-1} : i \in [k]\}$, is a non-crossing partition of E(M).

In Theorem 3.2, we give a similar characterization of positroids, namely, the following result, which, along with its corollaries, the rest of the paper exploits.

Let M be a matroid with no loops. A linear order \leq on E(M) is a positroid order if and only if, for each connected flat F of M, if K is a connected component of the contraction M/F, then there is a subset I of E(M) such that $F \subseteq E(M) - I$, $K \subseteq I$, and either I or E(M) - I is an interval in \leq .

This result applies to all loopless matroids, whether connected or not, thereby making it easier to use. There is much interest in characterizations of positroid orders; for instance, see Agarwala, Delaney, Yeats [1, Remark 5.27], which relates Theorem 5.17 in that paper to Theorem 3.2 in this paper. In Section 3 we also treat some corollaries of Theorem 3.2 and, in Theorem 3.5, relate it to several earlier characterizations of positroids.

We develop more extensive applications of Theorem 3.2 in Sections 4 and 5, the first of which concerns free amalgams. The free amalgam of matroids M and N, which exists only for certain pairs (M, N), is the freest matroid on $E(M) \cup E(N)$ that has M and N as restrictions. A subset X of E(M) is a set of clones if each permutation of E(M) that fixes each element of E(M) - X is an automorphism of M. We show that if M and N are positroids and $E(M) \cap E(N)$ is an independent set of clones in both M and N, then the free amalgam of M and N exists and is a positroid (Theorem 4.17). We also show that the same conclusion holds under conditions on the independent set $E(M) \cap E(N)$ that are in one way weaker and in another stronger (Theorem 4.21). A contraction of the free amalgam of M and N need not be the free amalgam of the corresponding contractions of M and N; because of this, we introduce and work with a more general way to glue matroids together, one that has useful contraction properties. Defining and developing the basic properties of this construction, which we call bonding, occupies much of Section 4. Proving all relevant properties of bonding makes the paper more self-contained; no knowledge of free amalgams is required. In Section 5, we identify many excluded minors for the class of positroids, including multi-parameter infinite families of excluded minors. We do not identify all excluded minors for the class of positroids; for some excluded minors not given here, see Park [22].

In Section 2 we review the background that is needed for these results. To make the paper accessible to a wide audience, we give complete background on, for instance, the orders used when treating Grassmann necklaces, as well as sketches of some of the less widely known topics in matroid theory. We assume familiarity with basic matroid theory, as set out in [20], and we follow the matroid notation used in [20].

2. BACKGROUND

We use \mathbb{Z} for the set of integers, \mathbb{N} for $\{n \in \mathbb{Z} : n > 0\}$, and [n] for $\{1, 2, \dots, n\}$.

2.1. Some orders related to a linear order. For a linear order \leq on a set E, say given by $e_1 < e_2 < \cdots < e_n$, its *i-shift* \leq_i is the cyclic shift in which e_i is least, that is,

 $e_i <_i e_{i+1} <_i \dots <_i e_n <_i e_1 <_i \dots <_i e_{i-1}.$

A cyclic interval of \leq is an interval in some *i*-shift of \leq . Elements *e* and *f* of *E* are cyclically consecutive in \leq if $\{e, f\}$ is a cyclic interval of \leq . In an ordered set (P, \leq) , an *ideal* is a subset *I* of *P* for which if $x \in I$ and $y \in P$ with $y \leq x$, then $y \in I$; also, $F \subseteq P$ is a *filter* if whenever $x \in F$ and $z \in P$ with $x \leq z$, then $z \in F$. For a linear order, an ideal is an interval that contains the least element; a filter is an interval that contains the greatest element. Note that for $A \subseteq E$, the following statements are equivalent to saying that *A* is a cyclic interval: E - A is a cyclic interval; *A* or E - A is an interval; *A* is an ideal of some shift \leq_i (so e_i is the least element of *A* using \leq_i); *A* is a filter of some shift \leq_i (so e_{i-1} is the greatest element of *A* using \leq_i).

In some examples we will need to show that no linear order on E has all sets in some set of sets being cyclic intervals; for instance, $\{1,3,5\}$, $\{2,3,4\}$, $\{4,5,6\}$, and $\{1,2,6\}$ cannot all be cyclic intervals in any linear order on [6]. For this, the following observation and lemma are useful. Given two cyclic intervals A and B in a linear order on E, if $A \cap B \neq \emptyset$, then either $A \cup B = E$ or $A \cap B$ is a cyclic interval.

Lemma 2.1. Let A, B, C, and D be cyclic intervals in a linear order on E with $A \not\subseteq B$, $A \cup B \neq E$, $A \cap B \neq \emptyset$, $C \not\subseteq B$, $B \cup C \neq E$, $B \cap C \neq \emptyset$, and $A \cap B \cap C = \emptyset$. If D is disjoint from $A \cap B$ and $B \cap C$ but not from $B - (A \cup C)$, then $D \subseteq B - (A \cup C)$.

Proof. By replacing the order by a cyclic shift if needed, we may assume that B is an interval. The result follows by observing that, by the hypotheses, one of $A \cap B$ and $B \cap C$ is an ideal in the order that is induced on B, and the other is a filter, so only one of the two cyclic intervals whose union is $E - (B \cap (A \cup C))$ can contain elements of D.

For a rank-r matroid M, if the matrix representation A shows that \leq is a positroid order for M, then the matrix A' that shifts the first column of A to be last and multiplies that column by $(-1)^{r-1}$ shows that \leq_2 is also a positroid order for M since one can shift the first column of an r by r matrix to the last place with r-1 column transpositions. Thus, \leq is a positroid order for a matroid M if and only if \leq_i is a positroid order for M.

Given a linear order \leq on an *n*-element set *E* and an integer $k \in [n]$, we consider two associated orders on the set of *k*-element subsets of *E*: for *k*-subsets *X* and *Y* of *E* where the elements of *X* are, in order, $x_1 < x_2 < \cdots < x_k$, and likewise $y_1 < y_2 < \cdots < y_k$ for *Y*, we have

- the Gale order: $X \leq_G Y$ if $x_i \leq y_i$ for all $i \in [k]$, and
- the *lexicographic order*: $X \leq_L Y$ if X = Y or $x_i < y_i$ for the least *i* with $x_i \neq y_i$.

While \leq_L is a linear order, \leq_G is not except for certain n and k. Also, \leq_L is an extension of \leq_G , that is, if $X \leq_G Y$, then $X \leq_L Y$.

By the next lemma, as one would expect, replacing an element in a set by a smaller element yields a smaller set in the Gale order.

Lemma 2.2. Let \leq be a linear order on E. Fix a subset X of E and elements $a \in X$ and $b \in E - X$. If b < a, then $(X - a) \cup b <_G X$. If a < b, then $X <_G (X - a) \cup b$.

Proof. Assume that b < a. To get $(X - a) \cup b <_G X$, compare the elements of X, in order, (the first line below) to those of $(X - a) \cup b$ (the second line):

$$x_1 < \dots < x_{j-1} < x_j < x_{j+1} \cdots < x_{i-1} < a < x_{i+1} < \dots < x_k$$

 $x_1 < \dots < x_{j-1} < b < x_j < \dots < x_{i-2} < x_{i-1} < x_{i+1} < \dots < x_k$

where *j* is the least integer with $b < x_j$. The second statement follows from the first. \Box

The next lemma is from Gale [16].

Lemma 2.3. Let M be a matroid. Let \leq be a linear order on E(M). If B is the least basis of M in the lexicographic order, then, in the Gale order, $B \leq_G B'$ for all bases B' of M.

Proof. We induct on the distance between bases B and B' in the lexicographic order. If B' = B, then $B \leq_G B'$, so we may assume that $B \neq B'$ and, inductively, that $B \leq_G A$ for all bases A for which $A <_L B'$. Fix $a \in B' - B$. By the symmetric exchange property, for some $b \in B - B'$, both $(B - b) \cup a$ and $(B' - a) \cup b$ are bases of M. If a < b, then Lemma 2.2 would give $(B - b) \cup a <_G B$, so $(B - b) \cup a <_L B$, contrary to B being the least basis in the lexicographic order. Thus, b < a, so Lemma 2.2 gives $(B' - a) \cup b <_G B'$. By the inductive assumption, $B \leq_G (B' - a) \cup b$, so $B <_G B'$.

The least basis of a matroid M in the Gale order induced by a linear order is the *Gale* basis for that order. (In some sources, the Gale basis is taken to be the greatest basis.)

2.2. Transversal and nested matroids. Let $\mathcal{A} = (A_i : i \in [r])$ be an indexed family of subsets of a set E. A partial transversal of \mathcal{A} is a subset X of E for which there is an injection $\phi : X \to [r]$ with $e \in A_{\phi(e)}$ for all $e \in X$. Edmonds and Fulkerson [15] showed that the partial transversals of \mathcal{A} are the independent sets of a matroid on E. This matroid, denoted $M[\mathcal{A}]$, is a transversal matroid, and \mathcal{A} is a presentation of it. In [10], Brylawski showed how to view transversal matroids geometrically: transversal matroids of rank r with no loops are precisely the matroids that have geometric representations on an r-vertex simplex in which, for all $k \in [r]$, each circuit of rank k spans a k-vertex face of the simplex (e.g., a 3-element circuit must span an edge of the simplex).

Given a linear order \leq on E and an r-subset I of E, for each $i \in I$, let F_i be the filter $\{j : i \leq j\}$, and let $\mathcal{A} = (F_i : i \in I)$. Let $N(I, \leq)$ be the transversal matroid $M[\mathcal{A}]$. Note that if i < k, then $F_k \subsetneq F_i$, so the presentation \mathcal{A} of $N(I, \leq)$ is a chain of sets under inclusion. Transversal matroids that have presentations that are chains of sets are called *nested matroids*. (They are also called shifted matroids and Schubert matroids, among other names; see [6, Section 4].) Note that I is the Gale basis of $N(I, \leq)$, and an r-subset J of E is a basis of $N(I, \leq)$ if and only if $I \leq_G J$.

2.3. Grassmann necklaces and base-sortable matroids. For a matroid M and linear order $e_1 < e_2 < \cdots < e_n$ on E(M), if I_i is the Gale basis of M using the *i*-shift \leq_i of \leq , then the *n*-tuple (I_1, I_2, \ldots, I_n) of bases has the properties

• if $e_i \in I_i$, then $I_{i+1} = (I_i - e_i) \cup e_j$ for some $j \in [n]$ (j can be i), and

4

• if $e_i \notin I_i$, then $I_{i+1} = I_i$,

where $I_{n+1} = I_1$. An *n*-tuple of *r*-element subsets of a linearly ordered *n*-element set with these properties is a *Grassmann necklace of type* (n, r). The next result combines a result of Postnikov [23] (the first part) with one of Oh [19] (the converse). In this result and the rest of this paper, $\mathcal{B}(M)$ denotes the set of bases of a matroid M.

Theorem 2.4. Let \leq be a linear order on E. If (I_1, I_2, \ldots, I_n) is a Grassmann necklace of type (n, r) on E, then

(2.1)
$$\bigcap_{i \in [n]} \mathcal{B}(N(I_i, \leq_i))$$

is the set of bases of a rank-r positroid on E and \leq is a positroid order for this matroid. Conversely, if \leq is a positroid order for a matroid M, then $\mathcal{B}(M)$ is the intersection (2.1) where I_i is the Gale basis of M for \leq_i .

Positroids first appeared, under the name of base-sortable matroids, in Blum [3]. Let \leq be a linear order on the ground set E(M) of a matroid M of rank r. Given bases B and B' of M, list the elements of the multiset union $B \cup B'$ in order, taking multiplicities into account, as $x_1 \leq x_2 \leq \cdots \leq x_{2r}$. Let $B_e = \{x_{2i} : i \in [r]\}$, the set of elements in the even positions in the list, and $B_o = \{x_{2i-1} : i \in [r]\}$, the set of elements in the odd positions. Set $\operatorname{sort}_{\leq}(B, B') = \{B_e, B_o\}$. Thus, $\operatorname{sort}_{\leq}(B, B') = \operatorname{sort}_{\leq}(B', B)$. Under the order dual \leq^d of \leq (i.e., $e \leq^d f$ if and only if $f \leq e$), the elements in the even positions are switched with those in the odd positions, so $\operatorname{sort}_{\leq d}(B, B') = \operatorname{sort}_{\leq}(B, B')$. Likewise, for the *i*-shift, $\operatorname{sort}_{\leq_i}(B, B') = \operatorname{sort}_{\leq}(B, B')$.

Definition 2.5. A matroid M is base-sortable if there is a linear order on E(M) for which, for every pair B, B' of bases of M, the sets B_e and B_o in sort(B, B') are bases of M. Such a linear order on E(M) is called a base-sorting order for M.

Base-sortable matroids are also often called *sort-closed matroids*. As Blum noted, the observations above about *i*-shifts and the dual order show that \leq is a base-sorting order for M if and only if \leq_i is a base-sorting order for M, and likewise for the dual order.

Lam and Postnikov [18, Corollary 9.4] proved that positroids and base-sortable matroids are the same. (Key results are in [17, Section 4.2].)

Theorem 2.6. Let M be a matroid. A linear order on E(M) is a positroid order if and only if it is a base-sorting order.

2.4. **Connectivity.** For a matroid M, let $e, f \in E(M)$ be related precisely when e = f or there is a circuit C of M with $\{e, f\} \subseteq C$. This relation is an equivalence relation and the equivalence classes are the *connected components* of M (see [20, Proposition 4.1.2]). A matroid M is *connected* if it has just one connected component; equivalently, for each pair of distinct elements $e, f \in E(M)$, there is a circuit C of M with $\{e, f\} \subseteq C$. A *separator* of M is a union of connected components of M. The *trivial separators* are \emptyset and E(M). For a partition $\{X_1, X_2, \ldots, X_k\}$ of E(M) with k > 1, we have

$$(2.2) M = (M|X_1) \oplus (M|X_2) \oplus \dots \oplus (M|X_k)$$

if and only if M and the direct sum have the same circuits, and that holds if and only if each set X_i is a nontrivial separator of M. Since taking direct sums and the dual commute, it follows that a matroid and its dual have the same connected components, so M is connected if and only if for each pair of distinct elements $e, f \in E(M)$, there is a cocircuit C^* of Mwith $\{e, f\} \subseteq C^*$. **Lemma 2.7.** For a matroid M and a partition $\{X_1, X_2, \ldots, X_k\}$ of E(M), Equation (2.2) holds if and only if $r(M) = r(X_1) + r(X_2) + \cdots + r(X_k)$.

Proof. Equation (2.2) clearly gives $r(M) = r(X_1) + \cdots + r(X_k)$. For the converse, it suffices to prove that if the rank equality holds and C is a circuit of M, then $C \subseteq X_i$ for some $i \in [k]$. Assume that this is false. Set $Y_i = X_i \cap C$ for $i \in [k]$. By our assumption, each set Y_i is independent. Let B_i be a basis of $M|X_i$ with $Y_i \subseteq B_i$. This gives the contradiction that $B_1 \cup \cdots \cup B_k$ spans M, has size r(M), but contains the circuit C. \Box

We say that a set X in a matroid M is *connected* if the restriction M|X is connected.

Lemma 2.8. Let F be a nonempty connected flat of a matroid M. If A is a nonempty connected flat of the contraction M/F, then either A or $A \cup F$ is a connected flat of M.

Proof. Now $A \cup F$ is a flat of M since A is a flat of M/F. Let $M|(A \cup F)$ be the direct sum $M_1 \oplus M_2 \oplus \cdots \oplus M_k$ where each M_i is connected. Since F is connected in M, we may assume that $F \subseteq E(M_1)$. If $F = E(M_1)$, then $M/F|A = M_2 \oplus \cdots \oplus M_k$, so, since A is a connected flat of M/F, we must have k = 2, and therefore A is a connected flat of M/F, then, since $M/F|A = (M_1/F) \oplus M_2 \oplus \cdots \oplus M_k$ and A is a connected flat of M/F, we must have k = 1, so the flat $A \cup F$ of M is connected.

The simple properties in the next lemma will come up at several points.

Lemma 2.9. All circuits of a restriction of a matroid M are circuits of M. Dually, all cocircuits of a contraction of M are cocircuits of M. In particular, coloops of contractions of M are coloops of M.

Recall that the closure operator cl of a matroid M can be cast in terms of circuits as follows: for $X \subseteq E(M)$, we have $cl(X) = X \cup \{e : Y \cup e \text{ is a circuit for some } Y \subseteq X\}$.

Lemma 2.10. Let M be a matroid with no loops.

- (1) If C is a circuit of M, then cl(C) is connected.
- (2) If M has a spanning circuit C, i.e., r(C) = r(M), then M is connected.
- (3) If some $e \in E(M)$ is not a coloop and is in no proper connected nonsingleton flat of M, then M is connected, as is M/F for any flat F of M with $e \notin F$.

Proof. Part (1) follows from two items recalled above, the equivalence relation, applied to $M | \operatorname{cl}(C)$, and the formulation of $\operatorname{cl}(C)$. Part (2) follows from that since $\operatorname{cl}(C) = E(M)$. For part (3), since e is not a coloop, e is in at least one circuit, say C, and C spans M since $\operatorname{cl}(C)$ cannot be a proper flat, so M is connected by part (2). For the second assertion, it suffices to show that the hypotheses of part (3) hold for M/F. Now e is not a coloop of M/F by Lemma 2.9. Let the connected components of M | F be F_1, F_2, \ldots, F_t . No proper connected nonsingleton flat of M/F_1 contains e since otherwise e would be in a proper connected nonsingleton flat of M by Lemma 2.8. The same argument, using M/F_1 , then shows that e is in no proper connected nonsingleton flat of $M/(F_1 \cup F_2)$. Iterating this show that, as needed, e is in no proper connected nonsingleton flat of M/F. \Box

2.5. Cyclic flats, clones, and paving matroids. Some of the excluded minors for the class of positroids that we present in Section 5 will be defined by giving their cyclic flats and the ranks of those flats, so we briefly review this perspective on matroids. Given a matroid M, a subset A of E(M) is cyclic if the restriction M|A has no coloops. Equivalently, A is cyclic if it is a (possibly empty) union of circuits. A cyclic flat is a flat that is cyclic. Let $\mathcal{Z}(M)$ denote the set of cyclic flats of M. Note that $\mathcal{Z}(M)$ is a lattice: the join of cyclic

6



FIGURE 1. A graph whose cycle matroid is a positroid.

flats A and B is $cl(A \cup B)$, and their meet is $(A \cap B) - C$ where C is the set of coloops of $M|(A \cap B)$. All connected flats F with $|F| \ge 2$ are cyclic flats, but not conversely.

There is no connection between the terms *cyclic interval* and *cyclic flat*, both of which are well established in the literature. For instance, using the usual order, in the cycle matroid of the graph shown in Figure 1, which is a positroid, the set $\{3, 6, 9\}$ is a cyclic flat that is not a cyclic interval, while $\{6, 7\}$ is a cyclic interval that is a flat, but it is not a cyclic flat.

For any set X in a matroid M, if F is the closure of the union of the circuits of M|X, then F is a cyclic flat of M and r(X) = r(F) + |X - F| since the elements of X - F are coloops of M|X. If $A \subseteq E(M)$, then $r(X) \leq r(X \cap A) + |X - A| \leq r(A) + |X - A|$. Thus, $r(X) = \min\{r(F) + |X - F| : F \in \mathcal{Z}(M)\}$. So M is determined by its cyclic flats and their ranks, that is, by the set $\{(A, r(A)) : A \in \mathcal{Z}(M)\}$. The next result, from [25, 7], provides an axiom scheme for matroids from this perspective.

Theorem 2.11. For a set Z of subsets of a set E and a function $r : Z \to \mathbb{Z}$, there is a matroid M on E with Z(M) = Z and $r_M(X) = r(X)$ for all $X \in Z$ if and only if

- (Z0) (\mathcal{Z}, \subseteq) is a lattice,
- (Z1) $r(0_{\mathcal{Z}}) = 0$, where $0_{\mathcal{Z}}$ is the least set in \mathcal{Z} ,

(Z2) 0 < r(Y) - r(X) < |Y - X| for all $X, Y \in \mathbb{Z}$ with $X \subsetneq Y$, and

(Z3) for all sets X, Y in Z (or, equivalently, just incomparable sets in Z),

(2.3)
$$r(X \lor Y) + r(X \land Y) + |(X \cap Y) - (X \land Y)| \le r(X) + r(Y).$$

A set A is cyclic in a matroid M if and only if E(M) - A is a flat of the dual matroid, M^{*}, since A being a union of circuits of M is equivalent to E(M) - A being an intersection of hyperplanes (cocircuit complements) of M^{*}. Thus, X is a cyclic flat of M if and only if E(M) - X is a cyclic flat of M^{*}, and so $\mathcal{Z}(M^*)$, the lattice of cyclic flats of M^{*}, is isomorphic to the order dual of $\mathcal{Z}(M)$. In particular, just as $cl_M(\emptyset)$, the set of loops of M, is the least cyclic flat of M, the set $E(M) - cl_{M^*}(\emptyset)$ is the greatest cyclic flat of M. The first assertion in the lemma below is easy to see directly, and the second follows by duality.

Lemma 2.12. Let X be a cyclic flat of M. The lattice $\mathcal{Z}(M|X)$ of cyclic flats of the restriction M|X is the interval $[cl(\emptyset), X]$ in $\mathcal{Z}(M)$. The lattice $\mathcal{Z}(M/X)$ of cyclic flats of the contraction M/X is $\{F - X : F \in \mathcal{Z}(M) \text{ and } X \subseteq F\}$, which is isomorphic to the upper interval $[X, E(M) - cl_{M^*}(\emptyset)]$ of cyclic flats that contain X in $\mathcal{Z}(M)$.

From the remarks before Theorem 2.11, it follows that the automorphisms of M are the permutations of E(M) that map each cyclic flat of M to a cyclic flat of the same rank. Elements e and f in a matroid M are *clones of* M if the transposition (e, f), which maps e to f, and f to e, and fixes all other elements of E(M), is an automorphism of M. Since M and M^* have the same automorphisms, two elements are clones in a matroid if and only if

JOSEPH E. BONIN

they are clones in its dual. A subset X of E(M) is a set of clones of M if $e, f \in X$ are clones whenever $e \neq f$. The next lemma holds since any permutation of E(M) that fixes each element that is not in a set X of clones is a composition of transpositions of clones.

Lemma 2.13. For any matroid M and set X of clones of M, any permutation ϕ of E(M) for which $\phi(e) = e$ for all $e \in E(M) - X$ is an automorphism of M,

Note that e and f are clones in M if and only if they are in the same cyclic flats of M, and, if M has no loops, that is equivalent to e and f being in the same connected flats Fof M with $|F| \ge 2$ since such flats, which are cyclic, are the connected components of all cyclic flats. Thus, if X is a set of clones and F is a connected flat with $|F| \ge 2$, then either $X \cap F = \emptyset$ or $X \subseteq F$. For any $Y \subseteq E(M)$, the closure in M of any cyclic flat of M|Y is a cyclic flat of M, so clones in M that are in Y are clones in M|Y. By duality, the same holds for M/Y. Thus, if X is a set of clones of M, then X is a set of clones in any minor M' of M for which $X \subseteq E(M')$. The characterization of clones in terms of cyclic flats makes it evident that the relation on E(M) in which $e, f \in E(M)$ are related when e = f or e and f are clones is an equivalence relation; the equivalence classes are the *clonal classes* of M.

A matroid M of rank r is a paving matroid if each subset X of E(M) with |X| < ris independent. Thus, each set of size at most r-2 is a flat, so the only flats of a paving matroid M that may be dependent are E(M) and the hyperplanes. A paving matroid that has no dependent hyperplanes is a uniform matroid: in the *uniform matroid* $U_{r,n}$ on an *n*-element set E, the bases are all r-subsets of E. The cyclic flats of a connected paving matroid M are \emptyset , E(M), and (if any) the dependent hyperplanes. A paving matroid is sparse paving if all of its dependent hyperplanes are circuits (that is, circuit-hyperplanes). For example, the rank-3 matroids W_3 and W^3 shown in Figure 2 and discussed in the next subsection are sparse paving matroids; in both, $\{1, 2, 6\}$, $\{2, 3, 4\}$, and $\{4, 5, 6\}$ are circuit-hyperplanes, while $\{1, 3, 5\}$ is a circuit-hyperplane of \mathcal{W}_3 but not of \mathcal{W}^3 . It follows from Theorem 2.11 that a connected paving matroid M on E of rank $r \ge 2$ that is not uniform can be specified by giving subsets H_1, H_2, \ldots, H_t (the dependent hyperplanes) of E for which (a) $r \leq |H_i| \leq |E| - 2$, for all $i \in [t]$, and (b) $|H_i \cap H_j| \leq r - 2$, for all distinct $i, j \in [t]$; given such sets, letting \mathcal{Z} be $\{\emptyset, E, H_1, H_2, \ldots, H_t\}$ and setting $r(\emptyset) = 0, r(H_i) = r - 1$ for all $i \in [t]$, and r(E) = r defines a paving matroid (condition (a) gives property (Z2) and condition (b) gives the only nontrivial cases of property (Z3)).

2.6. Lattice path and multi-path matroids. Two special classes of positroids are the class of lattice path matroids [5] and the larger class of multi-path matroids [8]. A matroid M is a *lattice path matroid* if and only if it is transversal and there is a linear order on E(M) and a presentation \mathcal{A} of M that is an antichain of intervals in E(M), that is, each set in \mathcal{A} is an interval in E(M) and no set in \mathcal{A} is a subset of another set in \mathcal{A} . A multi-path matroid is a transversal matroid M that has a presentation by an antichain of cyclic intervals in some linear order on E(M).

An important matroid that is a multi-path matroid but not a lattice path matroid is the *n*-whirl, W^n , for $n \ge 3$. This rank-*n* matroid is formed from the cycle matroid W_n of an *n*-spoke wheel (an *n*-vertex cycle with one more vertex, the hub, that is adjacent to all other vertices) by relaxing a circuit-hyperplane. (Only when $n = 3 \operatorname{does} W_n$ have more than one circuit-hyperplane.) The sets $\{1, 2, 3\}, \{3, 4, 5\}, \ldots, \{2n - 1, 2n, 1\}$ give a presentation of W^n by cyclic intervals in the ground set [2n] under the usual linear order. See Figure 2.

By [8, Theorem 6.3], restricting a multi-path matroid to a proper flat gives a lattice path matroid. The paragraph after that result notes that, with that result, some properties of



FIGURE 2. The 3-spoke wheel, its cycle matroid W_3 , and the 3-whirl W^3 . The 3-spoke wheel is the complete graph K_4 , so W_3 is $M(K_4)$.

lattice path matroids automatically extend to multi-path matroids. For this paper, the most relevant such property is [6, Theorem 3.11]: any connected flat in a lattice path matroid M is an interval in the linear order on E(M). Thus, any connected flat in a multi-path matroid M is a cyclic interval in the linear order on E(M).

2.7. Amalgams and parallel connections. For matroids M and N, any matroid K on $E(M) \cup E(N)$ for which K|E(M) = M and K|E(N) = N is an *amalgam* of M and N. The equality M|T = N|T, where T is $E(M) \cap E(N)$, is a necessary (but not sufficient) condition for amalgams to exist. As Figure 3 shows, when M and N have an amalgam, there may be many amalgams; in that example, a different rank-4 amalgam results if $\{d, f, g, i\}$ is a circuit-hyperplane, and another rank-3 amalgam has, instead of the line $\{a, e, f, i, h\}$, the lines $\{a, e, f\}$ and $\{a, i, h\}$. If K is the freest amalgam of M and N (i.e., any set that is independent in any amalgams of M and N may exist without the free amalgam of M and N existing. The next result is one implication in [20, Proposition 11.4.3].

Theorem 2.14. Let K be an amalgam of M and N, and set $T = E(M) \cap E(N)$. If, for each flat F of K, we have

$$r(F) = r(F \cap E(M)) + r(F \cap E(N)) - r(F \cap T),$$

then K is the free amalgam of M and N.

Note that [20, Proposition 11.4.3] is about the proper amalgam, for which we refer to [20, Section 11.4] for the definition and a complete account. The proper amalgam, when it exists, is the free amalgam, but the free amalgam can exist without the proper amalgam existing. Two general conditions under which the proper amalgam is known to exist are (a) when T is a modular flat of either M or N, and (b) when the common restriction, M|T = N|T, is a modular matroid. Condition (b) applies to the amalgams that we consider here since, in what we treat, T will be independent in both M and N.

When T is the singleton $\{p\}$ and $r_M(p) = r_N(p)$, the free amalgam of M and N is the *parallel connection*, P(M, N), of M and N at the base point p. (When $r_M(p) = r_N(p)$, parallel connection generalizes the operation of gluing two graphs together along an edge. This operation is often defined by giving the set of circuits, as in [20, Proposition 7.1.4]. By [20, Exercise 7.1.1] and Theorem 2.14, when $r_M(p) = r_N(p)$, the parallel connection, as defined in [20], is the free amalgam, and thinking about parallel connection as the free amalgam is more useful for this paper, so we take that as the definition.) If p is a loop of just one of M and N, then although there are no amalgams of M and N, the parallel



FIGURE 3. Two rank-3 whirls, their free amalgam (which has rank 4), and a rank-3 amalgam.

connection is still defined: if p is a loop of M but not of N, then P(M, N) is defined to be $M \oplus (N/p)$; if p is a loop of just N, then P(M, N) is defined to be $(M/p) \oplus N$.

2.8. **Principal extension, series extension, quotients, and truncation.** A matroid N is an *extension* of a matroid M if M is a restriction of N. Single-element extensions have been studied extensively, starting with Crapo [11] (see also [20, Section 7.2]). We focus on principal extensions. Given a matroid M, a set $X \subseteq E(M)$, and an element $e \notin E(M)$, define $r' : 2^{E(M) \cup e} \to \mathbb{Z}$ by, for all $Y \subseteq E(M)$, setting $r'(Y) = r_M(Y)$ and

$$r'(Y \cup e) = \begin{cases} r_M(Y), & \text{if } X \subseteq \operatorname{cl}_M(Y), \\ r_M(Y) + 1, & \text{otherwise.} \end{cases}$$

It is routine to show that r' is the rank function of a matroid on $E(M) \cup e$. This matroid, which is denoted $M +_X e$, is the *principal extension* of M in which e is *added freely* to X. The *free extension* of M by e is $M +_{E(M)} e$; we shorten $M +_{E(M)} e$ to M + e. If $r_M(f) = 1$, then $M +_f e$ is a *parallel extension* of M. The parallel extension $M +_f e$ is the parallel connection of M and the rank-1 uniform matroid on $\{e, f\}$.

Note that e is in the closure of Y in $M +_X e$ if and only if either $e \in Y$ or $X \subseteq \operatorname{cl}_M(Y)$. Also, $M +_X e = M +_{\operatorname{cl}(X)} e$. It is easy to check that if $e, f \notin E(M)$ and X and Y are subsets of E(M), then $(M +_X e) +_Y f = (M +_Y f) +_X e$, so we do not need to specify an order for multiple principal extensions. If M has no loops, then the flat $\operatorname{cl}_M(X) \cup e$ of $M +_X e$ is connected by Lemma 2.10; in particular, M + e is connected. If M has no loops and $f \in E(M)$, then there is a minimum nonsingleton connected flat of M that contains f if and only if M is a principal extension $(M \setminus f) +_X f$ of $M \setminus f$ for some $X \subseteq E(M \setminus f)$.

Series extension (or, more properly, series coextension) is dual to parallel extension: $M \times_f e = ((M^*) +_f e)^*$. Thus, $M \times_f e$ is defined when f is not a coloop of M; also, $r(M \times_f e) = r(M) + 1$. The cocircuits of the parallel extension $M +_f e$ are those of M, except that those that contain f are augmented by e, so in the series extension $M \times_f e$, the circuits are those of M except that those that contain f are augmented by e.

For a matroid M, a matroid Q with E(Q) = E(M) is a *quotient* of M if, for some extension N of M with r(N) = r(M), we have Q = N/A where A = E(N) - E(M). For example, if M is the rank-4 matroid in Figure 3 and $X = \{a, e, f, i, h\}$, the quotient $(M +_X j)/j$ is the right-most matroid shown. If $A = \{e_1, e_2, \ldots, e_k\}$ with $k \le r(M)$ and $N = (((M + e_1) + e_2) \cdots) + e_k$, then Q = N/A is the *truncation* of M to rank r(M) - k, and $r_Q(X) = \min\{r_M(X), r(M) - k\}$ for all $X \subseteq E(Q)$. Observe that each basis of a rank-r matroid M is a spanning circuit of its truncation to rank r - 1, so, if M has no loops, its truncations to positive ranks less than r are connected by Lemma 2.10. Theorem 3.2 below, the proof of which uses Theorem 2.4, characterizes positroids with no loops by characterizing positroid orders. The equivalence of statements (1) and (2) is known (Theorem 2.6, due to Lam and Postnikov), but we include (2) since it fits naturally into our proof of the equivalence of (1) and (3). Statement (4), which uses the following definition, recasts (3) in a way that will prove to be useful.

Definition 3.1. For a matroid M that has no loops, a linear order on E(M) has the cyclic interval property if, for each proper connected flat F of M with $|F| \ge 2$ and each connected component K of the contraction M/F with $|K| \ge 2$, the set K is a subset of a cyclic interval that is disjoint from F.

We require $|F| \ge 2$ and $|K| \ge 2$ so that we can focus on the substantial cases when applying Theorem 3.2; the condition trivially holds if either inequality fails. We assume that M has no loops since otherwise the condition on connected flats would hold vacuously. A matroid is a positroid if and only if the matroid obtained by deleting all of its loops is a positroid, and the placement of loops has no impact on whether a linear order is a positroid order, so the assumption of no loops does not limit the applicability of Theorem 3.2. In this result, the matroid is not assumed to be connected.

Theorem 3.2. Let M be a matroid that has no loops and let \leq be a linear order on E(M). The following statements are equivalent:

- (1) \leq is a positroid order for M,
- (2) \leq is a base-sorting order for M,

(3) if F is a flat of M for which $2 \le |F| \le |E(M)| - 2$ and M|F is connected, then

(3.1)
$$M/F = (M/F|U_1) \oplus (M/F|U_2) \oplus \cdots \oplus (M/F|U_h)$$

where U_1, U_2, \ldots, U_h are the maximal cyclic intervals that are disjoint from F, and

(4) \leq has the cyclic interval property.

Proof. It is easy to see that statements (3) and (4) are equivalent. Below we show that statements (1)–(3) are equivalent. Let n = |E(M)| and r = r(M).

Assume that statement (1) holds. By Theorem 2.4, the set of bases of M is

$$\bigcap_{i\in[n]} \mathcal{B}\big(N(I_i,\leq_i)\big)$$

where I_i is the Gale basis of M in the *i*-shift \leq_i of \leq . Let B and B' be bases of M, and let $\operatorname{sort}_{\leq}(B, B') = \{B_e, B_o\}$. To show that B_e and B_o are bases of M, we show that $B_e, B_o \in \mathcal{B}(N(I_i, \leq_i))$ for all $i \in [n]$. Now $B, B' \in \mathcal{B}(N(I_i, \leq_i))$ since both B and B'are bases of M, so, for each $j \in [r]$, each of B and B' has at most j - 1 elements that are less than the *j*th element of I_i using \leq_i ; thus, the elements in positions 2j - 1 and 2j when we list the elements in the multiset union $B \cup B'$ in order, using \leq_i , are both at least the *j*th element of I_i . The elements in positions 2j - 1 and 2j are the *j*th elements of B_o and B_e , so $B_e, B_o \in \mathcal{B}(N(I_i, \leq_i))$, so statement (2) holds.

Now assume that statement (2) holds. Let F be a connected flat F of M for which $2 \leq |F| \leq |E(M)| - 2$. Let U_1, U_2, \ldots, U_h be the maximal cyclic intervals that are disjoint from F. By replacing \leq by a shift \leq_j if needed, we may assume that the elements of E(M) are, in order,

$$f_1^1, f_2^1, \dots, f_{p_1}^1, u_1^1, u_2^1, \dots, u_{q_1}^1, \dots, f_1^h, f_2^h, \dots, f_{p_h}^h, u_1^h, u_2^h, \dots, u_{q_h}^h,$$

JOSEPH E. BONIN

where all $p_i, q_i \in \mathbb{N}$ and $f_j^i \in F$, and $U_i = \{u_1^i, u_2^i, \dots, u_{q_i}^i\}$. Set $F_i = \{f_1^i, f_2^i, \dots, f_{p_i}^i\}$ for $i \in [h]$. Statement (3) holds trivially if h = 1, so assume that h > 1.

Since M|F is connected, there is a circuit $C \subseteq F$ with $\{f_{p_1}^1, f_1^2\} \subseteq C$. Let B be a basis of M|F with $C - f_{p_1}^1 \subseteq B$. Let B_1 and B_2 be bases of M with $B \subseteq B_1 \cap B_2$. Let t be $|B_1 \cap U_1| + |B_2 \cap U_1|$. We claim that t is even, so $|B_1 \cap U_1|$ and $|B_2 \cap U_1|$ have the same parity. To see this, let the elements in the multiset union $B_1 \cup B_2$, listed in order, be

$$a_1 = a_1 < a_2 = a_2 < \dots < a_i = a_i < z_1 \le z_2 \le \dots \le z_t < f_1^2 = f_1^2 < \dots$$

where $\{a_1, a_2, \ldots, a_i\} \subseteq F_1 - f_{p_1}^1$ and $z_j \in U_1$ for $j \in [t]$. Since $\{f_{p_1}^1, f_1^2\} \subseteq C$ and $C - f_{p_1}^1 \subseteq B$, the set $B'_1 = (B_1 - f_1^2) \cup f_{p_1}^1$ is a basis of M. The elements in the multiset union $B'_1 \cup B_2$, listed in order, are

$$a_1 = a_1 < a_2 = a_2 < \dots < a_i = a_i < f_{p_1}^1 < z_1 \le z_2 \le \dots \le z_t < f_1^2 < \dots$$

Both lists have 2r(F) elements of F, so since the sets in $\operatorname{sort}_{\leq}(B_1, B_2)$ and $\operatorname{sort}_{\leq}(B'_1, B_2)$ are bases, in each list, r(F) elements of F are in even positions and the other r(F) elements of F are in odd positions. The only difference in the positions of elements in F in the two lists is that the first copy of f_1^2 in the first list changes to $f_{p_1}^1$ and moves t places earlier. Therefore t must be even, and so $|B_1 \cap U_1|$ and $|B_2 \cap U_1|$ have the same parity.

We next draw a sharper conclusion: $|B_1 \cap U_1| = |B_2 \cap U_1|$ for any bases B_1 and B_2 of M with $B \subseteq B_1 \cap B_2$. This holds since we can get B_2 from B_1 by a sequence of singleelement exchanges, and no such exchange involves elements of F (since $B \subseteq B_1 \cap B_2$), so, by the parity result above, each exchange either exchanges an element in U_1 for an element in U_1 , or an element in $E(M) - (F \cup U_1)$ for an element in $E(M) - (F \cup U_1)$.

For any two bases B and B' of M|F, and any subset I of E(M) - F, the set $B \cup I$ is a basis of M if and only if $B' \cup I$ is a basis of M. This gives the stronger conclusion that for all bases B_1 and B_2 of M that contain bases of M|F, we have $|B_1 \cap U_1| = |B_2 \cap U_1|$.

The same argument applied to cyclic shifts shows that the same result holds for each of U_2, \ldots, U_h . For each $j \in [h]$, let $s_j = r(F \cup U_j) - r(F)$, so $s_j = r(M/F|U_j)$. We can extend a basis B of M|F to a basis of $M|(F \cup U_j)$, and then extend that to a basis of M, so what we just showed implies that for each basis B_1 of M with $B \subseteq B_1$, we have $|B_1 \cap U_j| = s_j$ for each $j \in [h]$. Thus, $r(M) = r(F) + s_1 + s_2 + \cdots + s_h$, and so

(3.2)
$$r(M/F) = r(M/F|U_1) + r(M/F|U_2) + \dots + r(M/F|U_h),$$

and so the direct sum decomposition in statement (3) follows by Lemma 2.7.

Now assume that statement (3) holds. Recall that n = |E(M)| and r = r(M). Let I_i be the Gale basis of M using \leq_i . By Theorem 2.4, the matroid M' for which

$$\mathcal{B}(M') = \bigcap_{i \in [n]} \mathcal{B}\big(N(I_i, \leq_i)\big)$$

is a positroid. Since I_i is the Gale basis of M using \leq_i , we have $\mathcal{B}(M) \subseteq \mathcal{B}(N(I_i, \leq_i))$ by Lemma 2.3, so $\mathcal{B}(M) \subseteq \mathcal{B}(M')$. Showing equality in that inclusion will give M = M', and so M is a positroid and \leq is a positroid order, thus proving statement (1). To show that equality, it suffices to show that no r-subset of E(M) that contains a (necessarily non-spanning) circuit of M is in $\mathcal{B}(M')$. Let C be a non-spanning circuit of M, let Fbe the connected flat $cl_M(C)$ of M, and let $C \subseteq D \subseteq E(M)$ where |D| = r. Let U_1, U_2, \ldots, U_h be the maximal cyclic intervals that are disjoint from F, and let s_j be $r(M/F|U_j)$. Equation (3.1) gives equation (3.2), so $r(M) = r_M(F) + s_1 + s_2 + \cdots + s_h$. Since $|D \cap F| \geq |C| > r_M(F)$, we have $|D \cap U_j| < s_j$ for at least one set U_j . Pick the i for which U_j is a filter in the i-order \leq_i . Equation (3.1) gives $M/F|U_j = M/(E(M)-U_j)$, so

12

the Gale basis I_i must contain s_j elements of U_j . Thus, each basis of $N(I_i, \leq_i)$ contains at least s_j elements in the filter U_j of the *i*-order \leq_i . Since $|D \cap U_j| < s_j$, we have $D \notin \mathcal{B}(N(I_i, \leq_i))$, so $D \notin \mathcal{B}(M')$, as needed.

The corollary below is obtained from condition (3) in Theorem 3.2 by duality and the following results: F is a flat of M if and only if E(M) - F is a cyclic set of M^* ; the positroid orders for M are the same as the positroid orders for M^* .

Corollary 3.3. Let M be a matroid that has no coloops. A linear order \leq on E(M) is a positroid order for M if and only if, for all cyclic sets A of M for which M/A is connected and $2 \leq |A| \leq |E(M)| - 2$, we have $M|A = (M|V_1) \oplus (M|V_2) \oplus \cdots \oplus (M|V_h)$ where V_1, V_2, \ldots, V_h are the maximal cyclic intervals that are subsets of A.

The next corollary is useful both for limiting the search for potential positroid orders and for deducing that certain matroids are not positroids. This result is part of [2, Proposition 5.6] by Ardila, Rincón, and Williams, as cast in Theorem 1.2 above.

Corollary 3.4. Let F be a flat in a matroid M that has no loops. If both M|F and M/F are connected, then F is a cyclic interval in any positroid order for M.

Proof. This follows since the integer h in condition (3) in Theorem 3.2 must be 1.

For example, the cycle matroid $M(K_4)$ is not a positroid since the four 3-point lines are connected, as are the corresponding contractions, but, by Lemma 2.1, there is no linear order on the ground set in which each of those lines is a cyclic interval. (The example before Lemma 2.1 uses the 3-point lines of $M(K_4)$ as labeled in Figure 2.) This idea is taken much further in Examples 4–8 of Section 5.

Example 1. Let M be the cycle matroid of the graph in Figure 1, which has rank five and is the parallel connection of four copies of the uniform matroid $U_{2,3}$, one on each of the sets $\{3, 6, 9\}$, $\{1, 2, 3\}$, $\{4, 5, 6\}$, and $\{7, 8, 9\}$. By Theorem 3.2, M is a positroid since the cyclic interval property clearly holds for each proper connected flat with at least two elements that is a cyclic interval (so, $\{1, 2, 3\}$, $\{4, 5, 6\}$, and $\{7, 8, 9\}$), and all other connected flats F with $|F| \ge 2$ contain $\{3, 6, 9\}$, so each connected component of M/F is one of the intervals $\{1, 2\}$, $\{4, 5\}$, and $\{7, 8\}$.

To illustrate a point that is used at the end of the proof of Theorem 3.2, we explain how each 5-element subset X of E(M) that contains a non-spanning circuit of M fails to be a basis of some matroid $N(I_i, \leq_i)$. If $\{1, 2, 3\} \subseteq X$, then X is not a basis of $N(I_1, \leq_1)$ since $I_1 = \{1, 2, 4, 5, 7\}$ but at most two elements $j \in X$ satisfy $4 \leq_1 j$. The cases of $\{4, 5, 6\} \subseteq X$ (use I_4) and $\{7, 8, 9\} \subseteq X$ (use I_7) are similar. If X contains a 4- or 5circuit, then X is disjoint from $\{1, 2\}, \{4, 5\}$, or $\{7, 8\}$. If $X \cap \{1, 2\} = \emptyset$, then X is not a basis of $N(I_3, \leq_3)$ since the fifth element of X is less than that of $I_3 = \{3, 4, 5, 7, 1\}$ in \leq_3 . The cases of $X \cap \{4, 5\} = \emptyset$ (use I_6) and $X \cap \{7, 8\} = \emptyset$ (use I_9) are similar.

Figure 4 shows the rank-4 truncation of M. Applying Corollary 3.4 to the connected flat $\{3, 6, 9\}$ shows that the usual order is not a positroid order for the truncation of M to rank 3 or 4. Indeed, these truncations are excluded minors for the class of positroids (see Section 5). Thus, the class of positroids is not closed under truncation.

The proof of the next result shows that (1) the cyclic interval property, (2) a condition that Blum conjectured to characterize base-sorting orders [3, Conjecture 4.9], and (3) a characterization of positroid orders via forbidden induced orders on certain 4-element rank-2 minors can easily be derived from each other. The third characterization, while widely known among researchers in the field, seems to never have appeared in the literature.



FIGURE 4. A rank-4 excluded minor for the class of positroids that is the truncation of a rank-5 positroid.

Theorem 3.5. Let M be a matroid that has no loops and let \leq be a linear order on E(M). The following statements are equivalent:

- (1) \leq has the cyclic interval property;
- (2) for every minor N of M that has neither loops nor coloops, each circuit-hyperplane of N is a cyclic interval in the linear order that \leq induces on E(N);
- (3) for every minor N of M for which E(N) is the disjoint union of a 2-element circuit C and a 2-element cocircuit, the circuit C is a cyclic interval in the linear order that ≤ induces on E(N).

Proof. Assume that statement (1) holds, so \leq is a positroid order by Theorem 3.2. Let N be a minor of M that has neither loops nor coloops and let H be a circuit-hyperplane of N. By Corollary 3.4 applied to the positroid N and its connected flat H, the set H is a cyclic interval in the linear order that \leq induces on E(N). Thus, statement (2) holds, as does (3), which is a special case of (2).

Now assume that statement (1) fails. We claim that statement (3), and hence (2), also fail. Let F be a connected flat of M and let K be a connected component of M/F that is not contained in any cyclic interval of E(M) that is disjoint from F. Thus, there are elements $a, b \in F$ and $e, f \in K$ for which $\{a, b\}$ is not a cyclic interval in the induced order on $\{a, b, e, f\}$. Since F is connected, some circuit C of M|F has $a, b \in C$. Since M/F|K is connected, some cocircuit C^* of M/F has $e, f \in C^*$, and C^* is a cocircuit of M by Lemma 2.9. Extend C-a to a basis B of $M-C^*$. Set $N = M/(B-b)|\{a, b, e, f\}$. Thus, N has rank two, $\{a, b\}$ is a circuit of N, and $\{e, f\}$ is a cocircuit of N. Since $\{a, b\}$ is not a cyclic interval in the induced order on $\{a, b, e, f\}$, statement (3) fails, as claimed. \Box

Statement (3) in the result above is an excluded-minor characterization of positroids if one considers the linear order to be part of the positroid. In contrast, in this paper, we distinguish between the matroid and positroid orders for it, so the excluded minors that we consider in Section 5 are for the class of matroids for which positroid orders exist.

The cyclic interval property clearly holds under the hypotheses of the next corollary, which are less restrictive than the hypotheses of [3, Proposition 4.5].

Corollary 3.6. Let M be a matroid with no loops. If there is a linear order on E(M) in which each connected flat of M is a cyclic interval, then M is a positroid and the linear order is a positroid order.

In particular, any matroid with no loops in which the proper nonsingleton connected flats are pairwise disjoint is a positroid. Thus, every rank-2 matroid is a positroid.

If a matroid M satisfies the hypothesis of Corollary 3.6, then so do its truncations since all proper connected flats of a truncation are proper connected flats of M. The same does not apply to quotients other than truncations. Also, the same is not true of Theorem 3.2. Indeed, the class of positroids is not closed under truncations, as Example 1 shows.

Oh [19] proved that lattice path matroids are positroids. From Corollary 3.6 and the remarks in Section 2.6, noting our observation on loops at the beginning of this section, it is easy to see that the same is true of the larger class of multi-path matroids (this also follows from Blum [3, Theorem 5.2]), and likewise for truncations of lattice path matroids (which need not be lattice path), and truncations of multi-path matroids (which need not be multi-path). As an aside, we note that the class of lattice path matroids is closed under direct sums, but that of multi-path matroids is not; also, truncations of direct sums of multi-path matroids.

We next show that the class of positroids is closed under circuit-hyperplane relaxation and a more general relaxation operation that we now discuss. Assume that a cyclic flat X of a matroid M is neither $cl_M(\emptyset)$ nor $E(M) - cl_{M^*}(\emptyset)$, that the only cyclic flat that properly contains X is $E(M) - cl_{M^*}(\emptyset)$, and that the only cyclic flat that X properly contains is $cl_M(\emptyset)$. (Thus, X could be a circuit-hyperplane of a connected matroid of rank at least two, but X is not limited to such sets.) Observe that properties (Z0)–(Z3) in Theorem 2.11 hold for $\mathcal{Z} = \mathcal{Z}(M) - \{X\}$ and $r : \mathcal{Z} \to \mathbb{Z}$ where $r(A) = r_M(A)$ for each $A \in \mathcal{Z}$. By that result, this reduced cyclic flat and rank data defines another matroid, a generalized relaxation of M, in which X is independent.

Corollary 3.7. Assume that X is a proper, nonempty cyclic flat of a matroid M that has no loops and no coloops, and that no other proper nonempty cyclic flat of M either contains X or is contained in X. Let M' be the matroid for which $\mathcal{Z}(M') = \mathcal{Z}(M) - \{X\}$ and $r_{M'}(A) = r_M(A)$ for all $A \in \mathcal{Z}(M')$. If M is a positroid, then so is M', and any positroid order for M is a positroid order for M'. Thus, any circuit-hyperplane relaxation of a connected positroid of rank at least two is a positroid.

Proof. It follows from Lemma 2.12 that for any proper, nonempty cyclic flat F of M', we have M|F = M'|F and M/F = M'/F. Thus, the proper connected flats F of M' with $|F| \ge 2$ (all of which are cyclic flats of M') are those of M other than X. For a positroid order for M, the cyclic interval property holds for M, so the cyclic interval property also holds for M', so the linear order is a positroid order for M'.

The same result and proof apply to the more general relaxation M' of M where $\mathcal{Z}(M')$ is $\mathcal{Z}(M) - \mathcal{S}$ for some $\mathcal{S} \subseteq \mathcal{Z}(M) - \{\emptyset, E(M)\}$ where $\mathcal{S} \cup \{\emptyset, E(M)\}$ is both a filter and an ideal of $\mathcal{Z}(M)$.

Since the class of positroids is closed under contraction but not under truncation, it is not closed under free extension. (It is therefore also not closed under the operation of free product, of which free extension is a special case; see [12, 13, 20].) The following corollary characterizes when free extensions of positroids are positroids.

Corollary 3.8. Let M be a matroid with no loops, and fix $e \notin E(M)$. The free extension M + e is a positroid if and only if there is a linear order on E(M) in which each proper connected flat of M is an interval.

Proof. First assume that M + e is a positroid, and let \leq be a positroid order for M + e. By replacing \leq by a cyclic shift if needed, we may assume that e is the greatest element of E(M + e). Since e is neither a coloop of M + e nor in any proper connected nonsingleton flat of M + e, by Lemma 2.10 any contraction (M + e)/F by a flat F with $e \notin F$ is connected. Let F be a proper connected flat of M. Then F is a proper connected flat of M + e, and (M + e)/F is connected, so F is a cyclic interval in \leq by Corollary 3.4. Therefore F is an interval in \leq since $e \notin F$. Thus, in the positroid order that \leq induces on E(M), every proper connected flat of M is an interval.

The converse follows easily from Corollary 3.6 since the proper connected nonsingleton flats of M are precisely those of M + e, so if there is a linear order on E(M) in which each proper connected flat of M is an interval, then the same holds for M + e by extending that positroid order for M to make e the greatest element in the linear order.

Corollary 3.9. If the ground sets of M_1, M_2, \ldots, M_t are pairwise disjoint and the free extension $M_i + e$, for each $i \in [t]$, is a positroid, then any iterated free extension of any truncation of $M_1 \oplus M_2 \oplus \cdots \oplus M_t$ is a positroid.

Proof. Let $N = M_1 \oplus M_2 \oplus \cdots \oplus M_t$. For each $i \in [t]$, since $M_i + e$ is a positroid, there is a linear order $e_{i,1}, e_{i,2}, \ldots, e_{i,n_i}$ on $E(M_i)$ in which each connected flat of M_i is an interval. Concatenating these linear orders as

$$e_{1,1}, e_{1,2}, \ldots, e_{1,n_1}, e_{2,1}, e_{2,2}, \ldots, e_{2,n_2}, \ldots, e_{t,1}, e_{t,2}, \ldots, e_{t,n_t}$$

gives a linear order on E(N) in which each connected flat of N is an interval. By Corollary 3.8, adding elements freely to N yields a positroid, and contracting some of them gives an iterated free extension of a truncation of N.

We next identify some cases in which principal extensions of positroids are positroids. The assumption in this result amounts to M being a principal extension of $M \setminus f$.

Corollary 3.10. Let M be a positroid with no loops. Fix $e \notin E(M)$. Assume that for some nonsingleton connected flat A of M and $f \in E(M)$, for each nonsingleton connected flat F of M, we have $f \in F$ if and only if $A \subseteq F$. Then the principal extension $M +_A e$ of M is a positroid.

Proof. By construction, e and f are clones in $M +_A e$. Let \leq be a positroid order for M. We may assume that f is the greatest element of E(M) under \leq . Define \leq' on $E(M) \cup e$ by appending e as the new greatest element, so E(M) is an ideal in the extension \leq' of \leq . The connected flats of $M +_A e$ with at least two elements are of two types: (i) connected flats F of M with $|F| \geq 2$ and $f \notin F$, and (ii) $F \cup e$ where F is a connected flat of Mwith $|F| \geq 2$ and $f \in F$. For a connected flat F of $M +_A e$ of type (i), the connected components of $(M +_A e)/F$ are those of M/F, but with the one that contains f augmented by e. Also, for a connected flat $F \cup e$ of $M +_A e$ of type (ii), the connected components of $(M +_A e)/(F \cup e)$ are those of M/F. With those observations, it is easy to see that \leq' has the cyclic interval property.

There are infinite antichains of positroids in the minor quasi-order (where $N \leq M$ if N is isomorphic to a minor of M), so the class of positroids is not well-quasi-ordered. Some examples of infinite antichains of positroids are: the excluded minors for the class of nested matroids (the truncation, to rank n, of the direct sum of two n-circuits, for $n \geq 2$; see [21]); apart from $M(K_4)$, the excluded minors for the class of lattice path matroids (see [4]); for all $n \geq 3$, the truncation of the rank-n whirl W^n to rank 3.

The next result will be useful in Section 4.

Corollary 3.11. Let $X_1, X_2, ..., X_d$ be pairwise disjoint sets of clones in a positroid M for which, in some positroid order for M, some $x \in X_1$ and $y \in X_2$ are cyclically consecutive. Then there is a positroid order for M in which each X_i , for $i \in [d]$, is a cyclic

interval, as is $X_1 \cup X_2$. In particular, there is a positroid order for M in which each clonal class is a cyclic interval.

Proof. Assume that the following linear order, \leq , on E(M), written as a list from the minimum to the maximum element, is a positroid order for M:

 $x_1^1, \dots, x_{t_1}^1, a_1^1, \dots, a_{s_1}^1, x_1^2, \dots, x_{t_2}^2, a_1^2, \dots, a_{s_2}^2, \dots, x_1^k, \dots, x_{t_k}^k, a_1^k, \dots, a_{s_k}^k,$

where all t_i , s_i are in \mathbb{N} , each x_i^j is in X_1 , and each a_i^j is in $E(M) - X_1$. Also, we can take the elements x and y in the hypothesis to be $x = x_1^1$ and $y = a_{s_k}^k$. (By using a cyclic shift and the order dual as needed, any other pair of cyclically consecutive elements $x \in X_1$ and $y \in X_2$ can be brought into the analogous position.) Consider the linear order \leq_1 that, written as a list, is given by

$$x_1^1, \dots, x_{t_1}^1, x_1^2, \dots, x_{t_2}^2, \dots, x_1^k, \dots, x_{t_k}^k, a_1^1, \dots, a_{s_1}^1, a_1^2, \dots, a_{s_2}^2, \dots, a_1^k, \dots, a_{s_k}^k,$$

so \leq and \leq_1 induce the same linear orders on X_1 and on $E(M) - X_1$; also, X_1 is an ideal of \leq_1 . If k = 1, then \leq_1 is the positroid order \leq for M. We next show that \leq_1 is a positroid order for M even if k > 1.

Let F be a connected flat of M with $|F| \ge 2$ and let K be a connected component of M/F with $|K| \ge 2$. By Theorem 3.2, there is a cyclic interval I for \le so that $K \subseteq I$ and $I \cap F = \emptyset$. We must show that $K \subseteq I'$ and $I' \cap F = \emptyset$ for some cyclic interval I' for \le_1 . If $X_1 \subseteq F$, then since F and the cyclic interval I are disjoint, $I \subseteq \{a_1^i, a_2^i, \ldots, a_{s_i}^i\}$ for some $i \in [k]$, and so I is a cyclic interval for \le_1 and we can take I' = I. Now assume that $X_1 \not\subseteq F$. Then $X_1 \cap F = \emptyset$ since X_1 is a set of clones of M. Since X_1 is also a set of clones of M/F, either $X_1 \cap K = \emptyset$ or $X_1 \subseteq K$. First assume that $X_1 \cap K = \emptyset$. If $\{x_1^1, \ldots, x_{t_1}^1\} \cap I = \emptyset$, then $I' = I - X_1$ is a cyclic interval for \le_1 and satisfies $K \subseteq I'$ and $I' \cap F = \emptyset$. If $\{x_1^1, \ldots, x_{t_1}^1\} \cap I \neq \emptyset$. Now assume that $X_1 \subseteq K$. Then $X_1 \subseteq I$, so the cyclic interval E(M) - I for \le , which contains F, is $\{a_p^h, a_{p+1}^h, \ldots, a_q^h\}$ for some h, p, and q. Since $\{a_p^h, a_{p+1}^h, \ldots, a_q^h\}$ is also a cyclic interval for \le_1 , so is I, so we can take I' = I.

Starting with the dual of \leq_1 , apply the argument above to X_2 in place of X_1 , keeping $x = x_1^1$ and $y = a_{s_k}^k$ cyclically consecutive, giving \leq_2 . Since the restrictions of the orders \leq_1 and \leq_2 to $E(M) - X_2$ are the same, X_1 is a cyclic interval in \leq_2 , as are X_2 and $X_1 \cup X_2$. To complete the proof, apply the same modification of the linear order, one step at a time, for each of X_3, \ldots, X_d .

We close this section with a corollary of Theorem 3.2 that we will use in Section 5.

Corollary 3.12. Let M be a matroid that has exactly three proper, nonempty cyclic flats, Z_1, Z_2 , and Z_3 . If $E(M) = Z_1 \cup Z_2 \cup Z_3$ and $Z_1 \cap Z_2 \cap Z_3 = \emptyset$, then M is a positroid.

Proof. Any proper connected flat with at least two elements is cyclic and so must be Z_1 , Z_2 , or Z_3 , so, by Corollary 3.6, it suffices to find a linear order on E(M) in which Z_1 , Z_2 , and Z_3 are cyclic intervals. Consider the following six subsets of E(M), some of which may be empty: $Z_1 \cap Z_3$, $Z_1 - (Z_2 \cup Z_3)$, $Z_1 \cap Z_2$, $Z_2 - (Z_1 \cup Z_3)$, $Z_2 \cap Z_3$, and $Z_3 - (Z_1 \cup Z_2)$; for each set, pick a linear order on the elements, and then concatenate these six linear orders in the order given for these sets (so, for example, $Z_1 \cap Z_3$ is an ideal in this linear order). The result is a positroid order for M.

Similarly, any matroid that has at most four cyclic flats (and so at most one pair of incomparable cyclic flats) is a positroid.



FIGURE 5. Two rank-2 matroids M and N, the matroid H, and the bonding $B_T(M, N)$, which is $H/\{q_1, q_2\}\setminus\{s_1, s_2\}$.

4. BONDING AND ITS APPLICATION TO AMALGAMS OF POSITROIDS

In this section, we prove Theorem 4.17: if M and N are positroids and $E(M) \cap E(N)$ is an independent set of clones in M and in N, then the free amalgam of M and N is a positroid. We also prove a second result of this type (Theorem 4.21).

A difficulty that arises is that a contraction of an amalgam of M and N need not be an amalgam of the corresponding contractions of M and N. To deal with this, we introduce a more general way to glue matroids M and N together along $T = E(M) \cap E(N)$ that coincides with the free amalgam when T is independent in both M and N, but that has crucial contraction properties that come from the greater generality of the construction.

This section has two parts. In the first, we define the new construction, which we call bonding, and develop the properties of bonding that we need for our work on positroids, and then we treat those applications. In the second, we prove other properties of bonding that may help the future development of this topic.

4.1. Bonding and its application to positroids. The notation that we introduce next is used throughout this section. Let M and N be matroids for which $E(M) \cap E(N)$ is nonempty; let T denote this intersection. We form the *bonding of* M and N at T, denoted $B_T(M, N)$, via the following steps.

- Say that T is $\{t_1, t_2, \ldots, t_k\}$. Fix sets $S = \{s_1, s_2, \ldots, s_k\}$ and $Q = \{q_1, q_2, \ldots, q_k\}$ that are disjoint from each other and from E(M) and E(N).
- Form N' from N by, for each $i \in [k]$, relabeling t_i as s_i . Thus, $E(M) \cap E(N') = \emptyset$.
- Extend the direct sum $M \oplus N'$ by, for each integer $i \in [k]$, adding q_i freely to the flat $\operatorname{cl}_{M \oplus N'}(\{t_i, s_i\})$, giving the matroid H on $E(M) \cup E(N') \cup Q$.
- The bonding $B_T(M, N)$ is the minor $H/Q \setminus S$ of H.

Thus, $E(B_T(M, N)) = E(M) \cup E(N)$. Note that $B_T(M, N) = B_T(N, M)$. Also, the point q_i is added freely to the line $cl_{M \oplus N'}(\{t_i, s_i\})$ of $M \oplus N'$ if t_i is a loop of neither M nor N, while it is parallel to s_i if t_i is a loop of M only, and parallel to t_i if t_i is a loop of N only, and q_i is a loop if t_i is a loop of both M and N.

Figure 5 gives a simple example that will show the necessity of certain hypotheses in some results below. Figure 6 gives an example that is more representative of what we will see: if T is independent in both M and N, then $B_T(M, N)$ is the free amalgam of M and N; also, if, in addition, T is a set of clones, and M and N are positroids, then $B_T(M, N)$ is a positroid. That example shows that bonding two transversal matroids (in fact, multipath matroids) need not yield a transversal matroid. If M and N are the rank-2 uniform



FIGURE 6. Two rank-3 positroids M and N, and the bonding (their free amalgam) $B_T(M, N)$. Here, $T = \{1, 2\}$ is an independent set of clones in M and N. The bonding is a rank-4 positroid.

matroids on [q + 1] and [2q] - [q - 1], respectively, so $T = \{q, q + 1\}$, then $B_T(M, N)$ is the rank-2 uniform matroid on [2q]. Thus, bonding need not preserve representability over a fixed finite field. However, bonding preserves membership in classes of matroids that are closed under direct sum, deletion, contraction, and principal extension, such as the class of matroids that are representable over a given infinite field.

The way that bonding interacts with direct sums, which we treat in the next lemma, is crucial in our work.

Lemma 4.1. Let M be $M_1 \oplus M_2 \oplus \cdots \oplus M_s$ and N be $N_1 \oplus N_2 \oplus \cdots \oplus N_t$. Let $h \leq \min(s,t)$ and let $\{T_1, T_2, \ldots, T_h\}$ be a partition of T. If, for each $i \in [h]$, we have $T_i = E(M_i) \cap E(N_i)$, then the bonding $B_T(M,N)$ is the direct sum of the bondings $B_{T_i}(M_i, N_i)$, for $i \in [h]$, along with M_{h+1}, \ldots, M_s and N_{h+1}, \ldots, N_t . Also, for i with $h < i \leq s$, $E(M_i)$ is a connected component of M if and only if $E(M_i)$ is a connected component of $B_T(M, N)$, and likewise for $E(N_j)$ with $h < j \leq t$.

Proof. This follows from the construction since the matroid H that is used to construct $B_T(M, N)$ is the direct sum of its counterparts in the construction of $B_{T_i}(M_i, N_i)$, for $i \in [h]$, with the other direct sum factors M_{h+1}, \ldots, M_s and N_{h+1}, \ldots, N_t .

We next relate the flats of $B_T(M, N)$ to those of H.

Lemma 4.2. Let F be a flat of $B_T(M, N)$, and set $S_F = \{s_i : t_i \in F \cap T\}$ and $Q_F = \{q_i : t_i \in F \cap T\}$. Then $F \cup Q \cup S_F$ is a flat of H, each $q_j \in Q - Q_F$ is a coloop of $H|(F \cup Q \cup S_F)$, and $B_T(M, N)|F = H|(F \cup Q_F \cup S_F)/Q_F \setminus S_F$.

Proof. Since F is a flat of $B_T(M, N)$, which is $H \setminus S/Q$, the set $F \cup Q$ is a flat of $H \setminus S$. Its closure $cl_H(F \cup Q)$ in H is $F \cup Q \cup S_F$ since any two of t_i , s_i , and q_i have the third in their closure. If $q_j \in Q - Q_F$, then q_j is a coloop of $H|(F \cup Q \cup S_F)$ since q_j was added freely to the flat spanned by t_j and s_j , which are not in the flat $F \cup Q \cup S_F$. Thus,

$$B_T(M,N)|F = H|(F \cup Q \cup S_F)/Q \setminus S_F = H|(F \cup Q_F \cup S_F)/Q_F \setminus S_F. \square$$

The next lemma allows us to deduce that certain flats of $B_T(M, N)$ are disconnected. In this lemma and throughout this section, for $X \subseteq E(M) \cup E(N)$, we shorten $X \cap E(M)$ to X_M , and likewise set $X_N = X \cap E(N)$. **Lemma 4.3.** Let F be a flat of $B_T(M, N)$. If there are separators X of $M|F_M$ and Y of $N|F_N$, not both trivial, for which $X \cap T = Y \cap T$, then $X \cup Y$ is a nontrivial separator of $B_T(M, N)|F$, so $B_T(M, N)|F$ is disconnected.

Proof. We have $B_T(M, N)|F = H|(F \cup Q_F \cup S_F)/Q_F \setminus S_F$ where S_F and Q_F are as in Lemma 4.2. By symmetry, we may assume that the separator X of $M|F_M$ is neither \emptyset nor F_M . Set $T' = X \cap T = Y \cap T$, and let $S_{T'} = \{s_i : t_i \in T'\}$ and $Q_{T'} = \{q_i : t_i \in T'\}$. Thus, $H|(F \cup Q_F \cup S_F)$ is the direct sum

$$(H|(X \cup Y \cup Q_{T'} \cup S_{T'})) \oplus (H|((F - (X \cup Y)) \cup (Q_F - Q_{T'}) \cup (S_F - S_{T'}))),$$

and neither $X \cup Y$ nor $F - (X \cup Y)$ is empty. Therefore $X \cup Y$ is a nontrivial separator of $H|(F \cup Q_F \cup S_F)/Q_F \setminus S_F$, that is, of $B_T(M, N)|F$.

We next show that bonding and restriction commute when restricting to supersets of T.

Lemma 4.4. If $T \subseteq X \subseteq E(M) \cup E(N)$, then $B_T(M, N)|X = B_T(M|X_M, N|X_N)$.

Proof. Now $B_T(M, N)|X = H/Q \setminus S|X = H|(X \cup Q \cup S)/Q \setminus S$ since deletion and contraction commute. With that, the lemma follows by showing that $H|(X \cup Q \cup S)$ plays the role of H in the construction of $B_T(M|X_M, N|X_N)$. That holds since (a) $H|(X \cup S)$ is $(M|X_M) \oplus (N'|X_{N'})$ where $X_{N'} = (X_N - T) \cup S$ and (b) for each $t_i \in T$, the point q_i is added freely to $cl_{M \oplus N'}(\{t_i, s_i\})$ in $M \oplus N'$ to get H, so q_i is added freely to $cl_{H|(X \cup S)}(\{t_i, s_i\})$ in $H|(X \cup S)$.

We next treat the counterpart for contraction.

Lemma 4.5. If $T \subseteq X \subseteq E(M) \cup E(N)$, then $B_T(M, N)/X = (M/X_M) \oplus (N/X_N)$. *Proof.* We show the special case $B_T(M, N)/T = (M/T) \oplus (N/T)$, from which the

general result follows. Observe that $cl_H(S \cup T) = cl_H(S \cup Q) = cl_H(Q \cup T)$. Thus,

$$B_T(M, N)/T = H/(Q \cup T) \setminus S$$

= $H/(S \cup T) \setminus Q$
= $H \setminus Q/(S \cup T)$
= $(M/T) \oplus (N'/S)$
= $(M/T) \oplus (N/T).$

We next treat a key contraction property for bonding that shows the traction we gain by using bonding. When T is independent in both M and N, as we will see, $B_T(M, N)$ is the free amalgam of M and N, but the bonding $B_T(M/X, N/Y)$ in the next lemma need not be an amalgam of M/X and N/Y; indeed, we may have $M/X|T \neq N/Y|T$.

Lemma 4.6. If $X \subseteq E(M) - T$ and $Y \subseteq E(N) - T$, then

 $B_T(M, N)/(X \cup Y) = B_T(M/X, N/Y).$

Proof. This holds since deletion and contraction commute:

$$B_T(M,N)/(X \cup Y) = H/Q \setminus S/(X \cup Y)$$
$$= H/(X \cup Y)/Q \setminus S$$
$$= B_T(M/X, N/Y)$$

since, to construct $B_T(M/X, N/Y)$, the matroid $H/(X \cup Y)$ plays the role of H.

The next lemma relates $B_T(M, N)/P$ and $B_{T-P}(M/P, N/P)$ when $\emptyset \neq P \subsetneq T$.

Lemma 4.7. If $\emptyset \neq P \subsetneq T$, then $B_T(M, N)/P = B_{T-P}(M/P, N/P)$.

Proof. It suffices to prove that $B_T(M, N)/t_1 = B_{T-t_1}(M/t_1, N/t_1)$ for any $t_1 \in T$, so we focus on that. Since deletion and contraction commute,

$$B_T(M,N)/t_1 = H/Q \backslash S/t_1 = H/q_1 \backslash s_1/t_1/(Q-q_1) \backslash (S-s_1).$$

Thus, it suffices to show that $H/q_1 \setminus s_1/t_1$ is the counterpart of H in the construction of $B_{T-t_1}(M/t_1, N/t_1)$, that is, it is the direct sum of M/t_1 and N'/s_1 (the copy of N/t_1 where $T - t_1$ is replaced by $S - s_1$), to which, for i with $2 \le i \le k$, the element q_i is added freely to the flat $cl_{(M/t_1)\oplus(N'/s_1)}(\{t_i, s_i\})$. For a set $X \subseteq E(H) - \{q_1, s_1, t_1, q_i\}$ where $2 \le i \le k$, the following statements are equivalent:

- $q_i \in \operatorname{cl}_{H/q_1 \setminus s_1/t_1}(X)$,
- $q_i \in \operatorname{cl}_H(X \cup \{q_1, t_1\}),$
- $t_i, s_i \in cl_H(X \cup \{q_1, t_1\}),$
- $t_i, s_i \in \operatorname{cl}_{H/q_1 \setminus s_1/t_1}(X).$

Thus, q_i is added freely to the flat $cl_{(M/t_1)\oplus (N'/s_1)}(\{t_i, s_i\})$ of $(M/t_1) \oplus (N'/s_1)$. Since $r_H(\{q_1, t_1\}) = r_H(\{s_1, t_1\})$, we have

$$r(H/q_1 \setminus s_1/t_1) = r(H) - r_H(\{q_1, t_1\})$$

= $r(M) + r(N') - r_H(\{s_1, t_1\})$
= $r(M/t_1) + r(N'/s_1).$

If $r_M(t_1) = 1 = r_N(t_1)$, then q_1 is in the span of neither E(M) nor E(N') in H, so M and N' are restrictions of H/q_1 , and so, since t_1 and s_1 are parallel in H/q_1 , both M/t_1 and N'/s_1 are restrictions of $H/q_1 \setminus s_1/t_1$. If $r_M(t_1) = 0$ and $r_N(t_1) = 1$, then q_1 and s_1 are parallel in H, so $H/q_1 \setminus s_1 = H \setminus q_1/s_1$, and so, with t_1 being a loop, N'/s_1 is a restriction of $H/q_1 \setminus s_1/t_1$. The other two options for $r_M(t_1)$ and $r_N(t_1)$ similarly imply that M/t_1 and N'/s_1 are restrictions of $H/q_1 \setminus s_1/t_1$. So, to complete the proof, observe that with the rank equality above, we get $H/q_1 \setminus s_1/t_1 \setminus (Q-q_1) = (M/t_1) \oplus (N'/s_1)$.

By the next lemma, $B_T(M, N)$ is an amalgam of M and N if T is independent in both M and N. Figures 5 (with T dependent) and 6 (with T independent) illustrate the lemma.

Lemma 4.8. If T is independent in M, then the restriction $B_T(M, N)|E(N)$ is N. If T is dependent in M and no element of T is a loop of N, then $B_T(M, N)|E(N)$ is a proper quotient of N.

Proof. Let T_0 be a basis of M|T. Set $Q_0 = \{q_i : t_i \in T_0\}$. For all j, the sets $\{s_j, q_j\}$ and $\{s_j, t_j\}$ have the same closure in H, so the four sets

$$E(N') \cup Q$$
, $E(N') \cup T$, $E(N') \cup T_0$, and $E(N') \cup Q_0$

have the same closure in H. The rank of that closure is

$$r_H(E(N') \cup Q) = r(N) + r_M(T) = r(N) + |T_0| = r(N) + |Q_0|.$$

Thus, Q_0 is independent in H and $H/Q_0|E(N') = N'$.

If T is independent in M, then $T_0 = T$, so $Q_0 = Q$, and so H/Q|E(N') is N'. The first assertion follows since t_i and s_i are either parallel in H/Q or, if t_i is a loop of N, then t_i and q_i are parallel in H, so t_i is a loop of H/Q.

Now assume that T is dependent in M and contains no loops of N. Since $|T_0| < |T|$, we have $Q - Q_0 \neq \emptyset$, so $H|(E(N') \cup Q)/Q_0$ is a rank-r(N) extension of N'. If $q_j \in Q - Q_0$, then $s_j \notin cl_H(Q_0)$, for otherwise there would be a circuit C so that $s_j \in C \subseteq Q_0 \cup s_j$, which would give $q_i \in cl_H(E(N') \cup (Q_0 - q_i))$ for any $q_i \in C$, contrary to the equality

JOSEPH E. BONIN



FIGURE 7. Two connected rank-3 matroids whose bonding at the circuit $T = \{1, 2\}$ is disconnected; $B_T(M, N)$ has six connected components, namely, $\{a, b\}, \{c, d\}, \{e, f\}, \{g, h\}, \{1\}$, and $\{2\}$.

 $r(E(N') \cup Q_0) = r(N') + |Q_0|$; so $q_j \notin cl_H(Q_0)$ since q_j was added freely to the flat spanned by s_j and t_j . Thus, $cl_H(Q_0) \cap (Q - Q_0) = \emptyset$, so $Q - Q_0$ has positive rank in the extension $H|(E(N') \cup Q)/Q_0$ of N'. To get $B_T(M, N)|E(N)$ from that extension of N', besides the relabeling relating N and N', we contract $Q - Q_0$, so $B_T(M, N)|E(N)$ is a proper quotient of N.

Thus, in general, $B_T(M, N)$ is an amalgam of quotients of M and N.

Corollary 4.9. Assume that the set T is independent in both M and N. If F is a flat of $B_T(M, N)$, then F_M is a flat of M and F_N is a flat of N. If both M and N are connected, then so is $B_T(M, N)$.

Proof. Lemma 4.8 gives the result about flats. For distinct $e, f \in E(M) \cup E(N)$, if $T \cap \{e, f\} \neq \emptyset$, then e and f are in a circuit of either M or N; otherwise each of e and f is in a circuit of either M or N with some $t \in T$. Such circuits are circuits of $B_T(M, N)$, so e and f are in the same component of $B_T(M, N)$. Thus, $B_T(M, N)$ is connected. \Box

If T is dependent in both M and N, then $B_T(M, N)$ can be disconnected even if both M and N are connected; see Figure 7 for an example. Also, $B_T(M, N)$ may be connected even if both M and N are disconnected and T is independent in both: this occurs, for instance, when M is the direct sum of a 4-circuit on $\{1, 2, a, b\}$ and a coloop 3, while N is the direct sum of a 4-circuit on $\{2, 3, c, d\}$ and a coloop 1. Thus, when F is a connected flat of $B_T(M, N)$, without additional hypotheses (as in Lemmas 4.16 and 4.20 below), we cannot deduce that either F_M is connected in M or F_N is connected in N.

The next result gives the ranks of the flats of $B_T(M, N)$.

Lemma 4.10. Assume that T is independent in M. The rank r(F) of a flat F of $B_T(M, N)$ is $r_M(F_M) + r_N(F_N) - |F \cap T|$.

Proof. Lemma 4.2 gives $B_T(M, N)|F = H|(F \cup Q_F \cup S_F)/Q_F \setminus S_F$ where Q_F and S_F are defined in that lemma. Since $cl_H(F \cup Q_F) = cl_H(F \cup S_F)$, we have

$$r(F) = r(B_T(M, N)|F) = r_H(F \cup S_F) - r_H(Q_F).$$

By how *H* is constructed, $H|(F \cup S_F)$ is the direct sum of $M|F_M$ and the restriction of N' that is a relabeling of $N|F_N$. As shown in the proof of Lemma 4.8, having *T* independent in *M* implies that *Q* is independent in *H*. Thus, $r(F) = r_M(F_M) + r_N(F_N) - |Q_F|$. The equality in the lemma follows since $|Q_F| = |T \cap F|$.

This lemma and Theorem 2.14 have the following corollary.

Corollary 4.11. If T is independent in both M and N, then $B_T(M, N)$ is the free amalgam of M and N.

The terminology of free amalgams is well established in the literature, so we will use that name when T is independent in both M and N, but, noting Lemma 4.6 and the remarks before it, we cannot work solely with free amalgams; we sometimes need the more general setting of bonding.

The next result follows from Corollary 4.11 when $r_M(p) = 1 = r_N(p)$. The other cases are easy to verify directly from the definitions of $B_p(M, N)$ and P(M, N).

Corollary 4.12. If $E(M) \cap E(N) = \{p\}$, then $B_p(M, N)$ is the parallel connection P(M, N) of M and N at the base point p.

We are most interested in connected flats. The next corollary narrows the collection of flats that we must consider.

Corollary 4.13. Assume that T is independent in both M and N. If F is a flat of $B_T(M, N)$ with $F \cap T = \emptyset$ and neither F_M nor F_N is empty, then F is disconnected.

The bonding $B_T(M, N)$ has additional useful properties when T is a set of clones in both M and N. We first show that clones in both M and N are clones in $B_T(M, N)$.

Lemma 4.14. If $t_i, t_j \in T$ are clones in both M and N, then they are clones in $B_T(M, N)$. Thus, if T is a set of clones in both M and N, then T is a set of clones in $B_T(M, N)$.

Proof. By symmetry, it suffices to show that if F is a cyclic flat of $B_T(M, N)$ and $t_i \in F$, then $t_j \in F$. Set $S_F = \{s_h : t_h \in F \cap T\}$. By Lemma 4.2, the set $F \cup Q \cup S_F$ is a flat of H. At least one of t_i and s_i is not a coloop of $H|(F \cup S_F)$, for otherwise t_i would be a coloop of both $M|F_M$ and $N|F_N$, so t_i would be a coloop of $B_T(M, N)|F$ by Lemma 4.3, contrary to the flat F being cyclic. If t_i is not a coloop of $H|(F \cup S_F)$, then its clone t_j in M must be in the flat $F \cup Q \cup S_F$, so $t_j \in F$; otherwise s_i is not a coloop of $H|(F \cup S_F)$, so its clone s_j must be in the flat $F \cup Q \cup S_F$, and so $t_j \in F$.

Since two elements are clones in M if and only if they are in the same nonsingleton connected flats of M, we get the next result.

Lemma 4.15. If T is a set of clones in both M and N, and if F is a connected flat of $B_T(M, N)$ with $|F| \ge 2$, then either $T \subseteq F$ or $F \cap T = \emptyset$.

We have seen that, when T is independent in M and N, if F is a connected flat of $B_T(M, N)$ and $F \cap T = \emptyset$, then F is a connected flat of one of M or N. The next lemma treats the case of $T \subsetneq F$, assuming that T is a set of clones. One can give a shorter proof of this lemma in the case of a connected flat by applying Lemma 4.3 (and that would suffice for our applications of the next lemma), but the next lemma treats connected sets, not just connected flats. Note also that the lemma does not assume that T is independent in M and N, so M and N might not be restrictions of $B_T(M, N)$.

Lemma 4.16. Assume that T is a set of clones in M and in N. Let X be a connected set in $B_T(M, N)$ with $T \subsetneq X$. If $X_M \neq T$, then $M|X_M$ is connected, and likewise for $N|X_N$ if $X_N \neq T$.

Proof. By symmetry, it suffices to prove the result for $M|X_M$ when $X_M \neq T$. Lemma 4.4 gives $B_T(M,N)|X = B_T(M|X_M,N|X_N)$. Since T is a set of clones in M, either (i) $|T| \geq 2$ and T contains only loops or only coloops of $M|X_M$, or (ii) $T \subseteq K$ for some connected component K of $M|X_M$. Since $T \neq X_M$, if option (i) held, then $M|X_M$ would have more than |T| connected components, so Lemmas 4.4 and 4.1 would give the contradiction that $B_T(M,N)|X$, which is connected, is a direct sum of at least two

matroids. So option (ii) holds. By Lemmas 4.4 and 4.1, if $M|X_M$ had any connected components besides K, then $B_T(M, N)|X$ would be a direct sum of at least two matroids, but $B_T(M, N)|X$ is connected. Thus, $M|X_M$ is connected.

We turn to the applications to positroids.

Theorem 4.17. Let M and N be positroids with no loops, and let $E(M) \cap E(N)$ be T. If T is nonempty, T is independent in both M and N, and T is a set of clones in both M and N, then the free amalgam $B_T(M, N)$ is a positroid.

Proof. Let $T = \{t_1, t_2, \ldots, t_k\}$. Observe that if the elements of T are coloops of M, then $B_T(M, N) = (M \setminus T) \oplus N$, in which case the result follows since the class of positroids is closed under deletion and direct sum. Thus, by symmetry and the assumption that T is a set of clones of both M and N, we may assume that no element of T is a coloop of either M or N. By Lemma 2.13 and Corollary 3.11, some positroid order for M and some positroid order for N have the same restriction to T and have T being an interval. Let

$$e_1 <_M e_2 <_M \cdots <_M e_m <_M t_1 <_M t_2 <_M \cdots <_M t_k$$

and

$$t_1 <_N t_2 <_N \cdots <_N t_k <_N f_1 <_N f_2 <_N \cdots <_N f_r$$

be such positroid orders for M and N, respectively. We claim that the linear order

 $e_1 < e_2 < \dots < e_m < t_1 < t_2 < \dots < t_k < f_1 < f_2 < \dots < f_n$

of $E(B_T(M, N))$ satisfies the cyclic interval property, so $B_T(M, N)$ is a positroid. Note that each of \leq_M and \leq_N is a restriction of \leq to an interval. The following observations about cyclic intervals in these orders are used below.

- (I₁) Assume that $F \subseteq E(M)$ and $F \cap T = \emptyset$, and that I is a cyclic interval in \leq_M with $I \cap F = \emptyset$. If $I \cap T = \emptyset$, then I is also a cyclic interval in \leq ; if $I \cap T \neq \emptyset$, then $I \cup E(N)$ is a cyclic interval in \leq and it is disjoint from F. The same statement holds with the roles of M and N reversed.
- (I₂) If $T \subseteq F \subseteq E(M) \cup E(N)$, then any cyclic interval in \leq_M that is disjoint from F is a cyclic interval in \leq , as is any cyclic interval in \leq_N that is disjoint from F.

Let F be a connected flat of $B_T(M, N)$ with $|F| \ge 2$. By the hypotheses about clones and Lemma 4.15, either $F \cap T = \emptyset$ or $T \subsetneq F$; we address each case below.

First assume that $F \cap T = \emptyset$. By Corollaries 4.9 and 4.13, the set F is a connected flat of either M or N, say of M. No element of T is a coloop of M, so no element of T is a coloop of M/F by Lemma 2.9, and so all elements of T, which are clones in M/F, are in the same connected component, say X, of M/F. Lemma 4.6 gives $B_T(M, N)/F = B_T(M/F, N)$. (Unless T is independent in M/F, there is no amalgam of M/F and N, but their bonding is defined. While M/F is a restriction of $B_T(M/F, N)$, only when T is independent in M/F will N be a restriction of $B_T(M/F, N)$.) Let K be a connected component of $B_T(M, N)/F$ with $|K| \ge 2$. Since T is a set of clones of $B_T(M, N)$ and so of $B_T(M, N)/F$, either $T \subseteq K$ or $T \cap K = \emptyset$. Now either

(a) $K \subseteq E(N)$,

or, by applying Lemma 4.1 to $B_T(M/F, N)$, one of the following statements holds:

- (b) K is a connected component of M/F for which $K \cap T = \emptyset$, or
- (c) K is a connected component of the bonding $B_T(M/F|X, N|Y)$ for some set Y with $T \subseteq Y \subseteq E(N)$.

The cyclic interval property holds in case (a) since E(N) is a cyclic interval in \leq . In case (b), there is a cyclic interval I in \leq_M that contains K and has $I \cap F = \emptyset$, so observation (I₁) above shows that the cyclic interval property holds. In case (c), for any cyclic interval I in \leq_M that contains X and has $I \cap F = \emptyset$, by observation (I₁), the cyclic interval $I \cup E(N)$ in \leq shows that the cyclic interval property holds.

Now assume that $T \subsetneq F$. By Lemma 4.5, we have $B_T(M, N)/T = (M/T) \oplus (N/T)$. If $F \subseteq E(M)$, then F is a connected flat of M and $B_T(M, N)/F = (M/F) \oplus (N/T)$, so each connected component K of $B_T(M, N)/F$ is a connected component of either M/For N/T; thus, K is contained either in a cyclic interval in \leq_M that is disjoint from F, or in the interval E(N) - T; such cyclic intervals are cyclic intervals in \leq , so the cyclic interval property holds. The same idea applies if $F \subseteq E(N)$. Now assume that $F \not\subseteq E(M)$ and $F \not\subseteq E(N)$. By Corollary 4.9 and Lemma 4.16, the sets F_M and F_N are connected flats of M and N, respectively. Also, $B_T(M, N)/F = (M/F_M) \oplus (N/F_N)$. Thus, each connected component of $B_T(M, N)/F$ is a connected component of one of M/F_M and N/F_N , and the cyclic interval property holds since it holds in M and N.

With the assumptions above, while $B_T(M, N)$ is a positroid, the matroid H that we use to construct $B_T(M, N)$ need not be a positroid. For instance, H is not a positroid when M and N are both the uniform matroid $U_{3,4}$ and |T| = 3.

Corollary 4.12 and Theorem 4.17 give one case of the next result on parallel connection; the other case, when p is a loop of one or both of M and N, holds since contractions and direct sums of positroids are positroids. Recall that duals of positroids are positroids, and series connections are dual to parallel connections: $S(M, N) = (P(M^*, N^*))^*$. Blum [3, Corollary 4.2] treated the special cases of parallel and series extension.

Corollary 4.18. If M and N are positroids and $|E(M) \cap E(N)| = 1$, then the parallel connection P(M, N) and the series connection S(M, N) are positroids.

A series-parallel network is a graph that can be obtained from a graph with a single edge (perhaps a loop) by repeatedly applying parallel extension and series extension. Duffin [14] proved that a 2-connected graph is a series-parallel network if and only if it has no K_4 -minor. Binary matroids with no $M(K_4)$ minor are graphic since the excluded minors for graphic matroids, other than the excluded minor for binary matroids, have $M(K_4)$ as a minor. (See Tutte [26] for the excluded minors.) So a matroid is the cycle matroid of a series-parallel network if and only if it is connected, binary, and has no $M(K_4)$ -minor. The next corollary, which is also [3, Theorem 5.1], follows since the class of positroids is closed under parallel extension and series extension, and $M(K_4)$ is not a positroid.

Corollary 4.19. A binary matroid is a positroid if and only if the restriction to each of its connected components is the cycle matroid of a series-parallel network.

In particular, cycle matroids of series-parallel networks are positroids.

In the proof of Theorem 4.17, to get a linear order < that extends both $<_M$ and $<_N$, we can start with E(M) - T as an interval followed by E(N) - T, and then put T either between them (as we did) or after E(N) - T. Having two options suggests that a similar argument may work if T is the union of two sets of clones; we may be able to put one set of clones between E(M) - T and E(N) - T, and the other after E(N) - T. Theorem 4.21 below, which we illustrate in Figure 8, treats that. The conditions on T in Theorem 4.21 are stronger than those in Theorem 4.17 in that $M | cl_M(T)$ and $N | cl_N(T)$ are required to be connected, but weaker in that, instead of requiring T to be a set of clones, T is partitioned into two sets of clones, P and T - P, where all nonsingleton connected flats of either M or N that contain at least one element of T - P contain all of T.



FIGURE 8. The free amalgam $B_T(M, N)$ of the rank-3 positroids M and N, where $T = \{5, 10\}$. The free amalgam is a positroid, but 5 and 10 are not clones. The set P in the hypothesis of Theorem 4.21 is $\{5\}$.

Many positroids satisfy the hypotheses of Theorem 4.21. For instance, start with an n-spoke wheel, delete at most n - 2 spokes (see Figure 9), take the cycle matroid, and relax the circuit-hyperplane of rim edges. Fix two consecutive spokes that remain, let P contain one of those spokes, and let T - P be the set of rim edges between those spokes (e.g., T can be the set of the thick edges in Figure 9). The set T is independent, its closure is connected (it is the circuit consisting of the two consecutive spokes and the rim edges between them), and all nonsingleton connected flats that contain one of those rim edges contain that circuit (relaxing the circuit-hyperplane is crucial for that property). To get examples for which both |P| and |T - P| are arbitrarily large, start with the matroids just constructed and apply the operation of t-expansion, from [9], which, given a matroid M, produces a matroid M^t with similar geometric structure, but of larger rank and size. To get M^t , where $t \in \mathbb{N}$ and $t \ge 2$, first extend M to M' by, for each $e \in E(M)$, adding t - 1 elements parallel to e if r(e) = 1, or as loops if r(e) = 0; then M^t is the matroid union of t copies of M'. By [9, Theorem 3.15] (another corollary of Theorem 3.2), a matroid M is a positroid if and only if its t-expansion M^t is a positroid.

We will need the following variation on Lemma 4.16.

Lemma 4.20. Assume that T is independent in both M and N, that $\emptyset \neq P \subsetneq T$, and that P is a set of clones in both M and N. Let F be a connected flat of $B_T(M, N)$ with $F \cap T = P$. If $F \subseteq E(M)$, then the flat F of M is connected, and likewise if $F \subseteq E(N)$. If $F \not\subseteq E(M)$ and $F \not\subseteq E(N)$, then the flat F_M of M is connected, as is the flat F_N of N.

Proof. The first assertions are immediate since M and N are restrictions of $B_T(M, N)$ by Lemma 4.8. Now assume that $F \not\subseteq E(M)$ and $F \not\subseteq E(N)$, and, contrary to what we must show, that $M|F_M$ is disconnected. The elements of P, which are clones in M, either are all coloops of $M|F_M$ or are all in the same connected component of $M|F_M$. Now $F \not\subseteq E(N)$, so in either case we have $M|F_M = (M|X) \oplus (M|(F_M - X))$ for some set X where $P \subseteq X \subsetneq F_M$. Lemma 4.3 gives the contradiction that $B_T(M, N)|F$ is disconnected (as Y in that lemma, use F_N). Thus, $M|F_M$ is connected. \Box

Theorem 4.21. Let M and N be positroids with no loops and let $E(M) \cap E(N)$ be T. Assume that



FIGURE 9. A wheel with some spokes deleted. The thick edges are the rim edges between two consecutive spokes along with one of those spokes.

- (1) T is independent in both M and N,
- (2) $M | \operatorname{cl}_M(T)$ and $N | \operatorname{cl}_N(T)$ are connected, and
- (3) there is a nonempty proper subset P of T for which(3a) P is a set of clones in both M and N, and
 - (3b) if *F* is a nonsingleton connected flat of either *M* or *N* with $F \cap (T P) \neq \emptyset$, then $T \subseteq F$.

If some positroid order for M has some element in P and some element in T - P cyclically consecutive, and some positroid order for N has some element in P and some element in T - P cyclically consecutive, then the free amalgam $B_T(M, N)$ is a positroid.

Proof. Let $P = \{p_1, p_2, \dots, p_h\}$ and $T - P = \{t_1, t_2, \dots, t_k\}$. Assumption (3b) implies that T - P is a set of clones in both M and N. By that observation, assumption (3a), Lemma 2.13, and the hypothesis in the last sentence of the theorem, p_h and t_1 are cyclically consecutive in some positroid order for M, as are t_k and p_1 in some positroid order for N. With P and T - P being sets of clones in both M and N, Lemma 2.13, Corollary 3.11, and the observation that the order dual of a positroid order is a positroid order, we can assume that the following linear orders are positroid orders for M and N, respectively:

 $e_1 <_M e_2 <_M \dots <_M e_m <_M p_1 <_M p_2 <_M \dots <_M p_h <_M t_1 <_M t_2 <_M \dots <_M t_k$ and

 $t_1 <_N t_2 <_N \dots <_N t_k <_N p_1 <_N p_2 <_N \dots <_N p_h <_N f_1 <_N f_2 <_N \dots <_N f_n.$

The linear order

 $e_1 < e_2 < \cdots < e_m < p_1 < p_2 < \cdots < p_h < f_1 < f_2 < \cdots < f_n < t_1 < t_2 < \cdots < t_k$ has both \leq_M and a shift of \leq_N as the induced orders on E(M) and E(N), respectively. We claim that the linear order \leq on $E(B_T(M, N))$ satisfies the cyclic interval property, so $B_T(M, N)$ is a positroid. Note that observations (I₁) and (I₂) from the proof of Theorem 4.17 apply to the linear order < just defined on $E(B_T(M, N))$.

Let F be a connected flat of $B_T(M, N)$ with $|F| \ge 2$. If F contained some but not all elements of T - P, then, by assumption (3b), each $t_i \in F \cap (T - P)$ would be a coloop of $M|F_M$ and of $N|F_N$, so setting $X = Y = \{t_i\}$ in Lemma 4.3 would give the contradiction that $B_T(M, N)|F$ is disconnected. Thus, by the hypotheses, there are three options: $F \cap T = \emptyset$, $T \subsetneq F$, or $F \cap T = P$.

The cyclic interval property holds when $T \subsetneq F$ by the same argument as in the proof of Theorem 4.17 once we make the following observations. Since F contains the connected flats $\operatorname{cl}_M(T)$ of M and $\operatorname{cl}_N(T)$ of N, there are connected components X of $M|F_M$ and

Y of $N|F_N$ that contain T. Since $B_T(M, N)|F$, which is $B_T(F_M, F_N)$, is connected, Lemma 4.1 implies that $F_M = X$ and $F_N = Y$, so F_M is a connected flat of M, as is F_N in N. We turn to the other two cases.

Assume that $F \cap T = \emptyset$. As before, F is a connected flat of either M or N, say of M. Thus, $B_T(M, N)/F = B_T(M/F, N)$. It follows from Lemma 2.8 that assumption (3b) also holds for M/F. Also, since $cl_M(T)$ is connected, no element of T is a coloop of M, so no element of T is a coloop of M/F. Thus, all elements of T are in the same connected component, say X, of M/F. It follows that any connected component K of $B_T(M, N)/F$ with $|K| \ge 2$ falls under one of cases (a)–(c) in the proof of Theorem 4.17. The cyclic interval property holds in each case by the argument given in that proof.

Now assume that $F \cap T = P$. We first treat the case with $F \not\subseteq E(M)$ and $F \not\subseteq E(N)$, in which case, by Lemma 4.20, the restrictions $M|F_M$ and $N|F_N$ are connected. By applying Lemma 4.7 and then Lemma 4.6, we have

$$B_T(M,N)/F = (B_{T-P}(M/P,N/P))/(F-T)$$

= $B_{T-P}(M/F_M,N/F_N).$

Since no element of T is a coloop of M, no element of T - P is a coloop of M/F_M . Also, T - P is a set of clones of M and so of M/F_M . Therefore, some connected component X of M/F_M has $T - P \subseteq X$. Similarly, some connected component Y of N/F_N has $T - P \subseteq Y$. Thus, any connected component of $B_{T-P}(M, N)/F$ is either

- (a) a connected component of M/F_M or of N/F_N that is disjoint from T P, or
- (b) a connected component of the bonding $B_{T-P}(M/F_M|X, N/F_N|Y)$.

For case (a), say K is a connected component of M/F_M ; then any cyclic interval in \leq_M that contains K and is disjoint from F_M is a cyclic interval in \leq , so the cyclic interval property holds. For a connected component K that falls under case (b), note that the union of a cyclic interval I in \leq_M for which $I \cap T = T - P$ and a cyclic interval J in \leq_N for which $J \cap T = T - P$ is a cyclic interval in \leq . Choosing such I and J for which $X \subseteq I$, $Y \subseteq J$, and $F \cap (I \cup J) = \emptyset$ shows that the cyclic interval property holds in case (b).

Finally, to complete the case with $F \cap T = P$, we can assume, by symmetry, that $F \subseteq E(M)$, so F is a connected flat of M. Now $B_T(M, N)/P = B_{T-P}(M/P, N/P)$ by Lemma 4.7, so

$$B_T(M, N)/F = B_{T-P}(M/F, N/P).$$

As above, all elements of T - P are in the same connected component of M/F, say X. Thus, any connected component K of $B_T(M, N)/F$ falls under one of three cases:

- (a) $K \subseteq E(N) P$,
- (b) K is a connected component of M/F with $K \cap (T P) = \emptyset$, or
- (c) K is a connected component of $B_{T-P}(M/F|X, N/P|Y)$ for some set Y for which $T P \subseteq Y \subseteq E(N) P$.

The cyclic interval property holds in case (a) since E(N) - P is a cyclic interval in \leq . The argument in case (b) is exactly as for case (a) in the previous paragraph. In case (c), X is a subset of a cyclic interval I in \leq_M with $I \cap F = \emptyset$ and so $I \cap T = T - P$. Therefore $I \cup (E(N) - P)$ is a cyclic interval in \leq , and it shows that the cyclic interval property holds in this case, thereby completing the proof of the theorem.

4.2. **Further properties of bonding.** It may be possible to prove other results in the spirit of Theorems 4.17 and 4.21. In this section, we treat some results about bonding that may be useful for the further development of this operation and its applications.

The connected components of bondings play crucial roles above. Even if M and N are connected, $B_T(M, N)$ might be disconnected; see Figure 7. The next lemma relates the connected components of M and N to those of $B_T(M, N)$ under certain hypotheses.

Lemma 4.22. Let K be a connected component of M. If $T \subseteq K$ and T is independent in N, then K is contained in a connected component of $B_T(M, N)$, say X, and $X_M = K$.

Proof. We have $B_T(M, N)|E(M) = M$ by Lemma 4.8 since T is independent in N. Thus, any two elements of K are contained in a circuit of $B_T(M, N)$, and so are in the same connected component, say X, of $B_T(M, N)$. Lemma 4.1 gives $X_M = K$.

The next lemma treats a partial converse of the first assertion in Corollary 4.9. Recall that (X, Y) is a *modular pair* in a matroid M if $r(X) + r(Y) = r(X \cup Y) + r(X \cap Y)$. When Y is independent, this equality can be rewritten as $r(X \cup Y) = r(X) + |Y - X|$.

Lemma 4.23. Assume that T is independent in both M and N. Fix $F \subseteq E(M) \cup E(N)$ where F_M is a flat of M and F_N is a flat of N. If (F_M, T) is a modular pair in M and (F_N, T) is a modular pair in N, then F is a flat of $B_T(M, N)$. In particular, if $|T-F| \leq 1$, then F is a flat of $B_T(M, N)$.

Proof. Recall that q_i , for each $i \in [k]$, is added freely to the line $\operatorname{cl}_{M \oplus N'}(\{t_i, s_i\})$ of $M \oplus N'$ to form H. We may assume that $F \cap T$ is $T_h = \{t_1, t_2, \ldots, t_h\}$, where h = 0 and $T_0 = \emptyset$ if $F \cap T = \emptyset$. Set $S_h = \{s_1, s_2, \ldots, s_h\}$ and $Q_h = \{q_1, q_2, \ldots, q_h\}$, which are empty if h = 0. Thus, $F \cup S_h \cup Q_h$ is a flat of H. Let $F_{N'} = (F \cap E(N')) \cup S_h$. By the two assumed modular pairs,

$$r_H(F \cup T \cup S) = r_M(F_M \cup T) + r_{N'}(F_{N'} \cup S)$$

= $r_M(F_M) + |T - T_h| + r_{N'}(F_{N'}) + |S - S_h|$
= $r_H(F \cup S_h) + 2(k - h).$

We claim that $F \cup S_h \cup Q$ is a flat of H. Assume instead that $F \cup S_h \cup Q$ is not a flat of H, so for some $e \notin F \cup S_h \cup Q$, there is a circuit C with $e \in C \subseteq F \cup S_h \cup Q \cup e$. Let $I = \{i : q_i \in C \cap (Q - Q_h)\}$ and $J = [k] - ([h] \cup I)$. Now $I \neq \emptyset$ since $F \cup S_h \cup Q_h$ is a flat of H. If $i \in I$, then $\{t_i, s_i\} \subseteq cl_H(C)$, so

$$cl_H(F \cup S_h \cup C \cup \{t_j, s_j : j \in J\}) = cl_H(F \cup T \cup S).$$

The union of $\{q_i : i \in I\} \cup \{t_j, s_j : j \in J\}$ and a basis of $F \cup S_h$ spans $F \cup T \cup S$, so $r_H(F \cup T \cup S) \leq r_H(F \cup S_h) + |I| + 2|J|$. However, |I| + 2|J| < 2(k - h), contrary to what we just showed. Thus, $F \cup S_h \cup Q$ is a flat of H, and so F is a flat of $B_T(M, N)$. \Box

The set $F = \{4, 6, 9, 11\}$ in the example in Figure 6 shows that the hypothesis about modular pairs in Lemma 4.23 is needed. In the proofs of the corollaries below, we will use two observations: (1) if (X, Y) is a modular pair in M, then so is $(cl_M(X), Y)$, and (2) if, in addition, Y is independent, then $X \cap Y = cl_M(X) \cap Y$.

Corollary 4.24. Assume that T is independent in both M and N, and $X \subseteq E(M) \cup E(N)$. If (X_M, T) is a modular pair in M and (X_N, T) is a modular pair in N, then the closure $\operatorname{cl}_B(X)$ of X in $B_T(M, N)$ is $\operatorname{cl}_M(X_M) \cup \operatorname{cl}_N(X_N)$.

Proof. We shorten $B_T(M, N)$ to B. Now $\operatorname{cl}_M(X_M) \subseteq \operatorname{cl}_B(X)$ since B|E(M) is M. Likewise, $\operatorname{cl}_N(X_N) \subseteq \operatorname{cl}_B(X)$. Thus, $\operatorname{cl}_M(X_M) \cup \operatorname{cl}_N(X_N) \subseteq \operatorname{cl}_B(X)$. Since (X_M, T) is a modular pair in M, so is $(\operatorname{cl}_M(X_M), T)$, and $\operatorname{cl}_M(X_M) \cap T = X \cap T$. Likewise, $(\operatorname{cl}_N(X_N), T)$ is a modular pair in N and $\operatorname{cl}_N(X_N) \cap T = X \cap T$. Thus, by Lemma 4.23,



FIGURE 10. The rank-4 truncation of the positroid in Example 1 and two of its contractions.

the union $\operatorname{cl}_M(X_M) \cup \operatorname{cl}_N(X_N)$ is a flat of B. Since this flat of B contains X, we have $\operatorname{cl}_B(X) \subseteq \operatorname{cl}_M(X_M) \cup \operatorname{cl}_N(X_N)$. With the inclusion proven above, equality follows. \Box

Corollary 4.25. Assume that T is independent in both M and N, and $X \subseteq E(M) \cup E(N)$. If (X_M, T) is a modular pair in M and (X_N, T) is a modular pair in N, then the rank $r_B(X)$ of X in $B_T(M, N)$ is $r_M(X_M) + r_N(X_N) - |X \cap T|$.

Proof. We again shorten $B_T(M, N)$ to B. Corollary 4.24, Lemma 4.10, and the equality $(cl_M(X_M) \cup cl_N(X_N)) \cap T = X \cap T$ give

$$r_B(X) = r_B(\operatorname{cl}_B(X))$$

= $r_B(\operatorname{cl}_M(X_M) \cup \operatorname{cl}_N(X_N))$
= $r_M(\operatorname{cl}_M(X_M)) + r_N(\operatorname{cl}_N(X_N)) - |X \cap T|$
= $r_M(X_M) + r_N(X_N) - |X \cap T|.$

5. Some excluded minors for the class of positroids

As [3, 2] show, the class of positroids is minor-closed, that is, every minor of a positroid is a positroid. Thus, the class of positroids is characterized by its excluded minors, which are the matroids that are not positroids, but all of their proper minors are positroids. Blum [3, Corollary 4.12] found the excluded minors for the class of positroids of rank at most three. In this section, we use Theorem 3.2 and its corollaries to identify infinitely many excluded minors for the class of positroids, focusing on excluded minors of rank greater than three. We do not identify all excluded minors for the class of positroids; indeed, more are given in Park [22]. We start with the truncation to rank 4 of the positroid in Example 1. (The truncation to rank 3 is treated in [3].)

Example 2. Let M be the rank-4 truncation of the positroid in Example 1, labeled as in Figure 10. Let $A = \{a, s, t\}, B = \{b, p, q\}, C = \{c, u, v\}, and X = \{a, b, c\}$. The proper connected flats F of M with $|F| \ge 2$ are those four sets along with $A \cup X, B \cup X$, and $C \cup X$. For each such F, the contraction M/F is connected, so by Corollary 3.4, if M were a positroid, then there would be a linear order on E(M) in which each of those flats is a cyclic interval. By Lemma 2.1, there is no such linear order, so M is not a positroid.

The automorphism group of M is transitive on $\{s, t, p, q, u, v\}$ and on $\{a, b, c\}$, so in order to show that all proper minors of M are positroids, it suffices to show that $M \setminus u$, M/u, $M \setminus c$, and M/c are positroids. We show that Corollary 3.6 applies to these minors.



FIGURE 11. Two rank-3 positroids M and N, and their free amalgam $B_T(M, N)$, which is a rank-4 excluded minor for the class of positroids.

- The proper connected flats *F* of *M**u* with |*F*| ≥ 2 are *A*, *B*, *X*, *A* ∪ *X*, and *B* ∪ *X*, which are intervals in the linear order *s* < *t* < *a* < *c* < *b* < *p* < *q* < *v*.
- The proper connected flats F of M\c with |F| ≥ 2 are A, B, and (C ∪ X) c, which are intervals in the linear order s < t < a < u < v < b < p < q.
- The proper connected flats *F* of *M*/*u* with |*F*| ≥ 2 are *A*, *B*, *C* − *u*, and *X* ∪ *v*, which are intervals in the linear order *s* < *t* < *a* < *c* < *v* < *b* < *p* < *q*.
- The proper connected flats F of M/c with |F| ≥ 2 are A ∪ b, B ∪ a, C c, and X c, which are intervals in the linear order s < t < a < b < p < q < u < v.

Thus, M is an excluded minor for the class of positroids.

Part of the interest in the next example is to show that the free amalgam of positroids, when the hypotheses of neither Theorem 4.17 nor Theorem 4.21 hold, need not yield a positroid; indeed, it may produce an excluded minor for the class of positroids.

Example 3. Consider the free amalgam $B = B_T(M, N)$ of the matroids M and N shown in Figure 11. There is no positroid order for B since, if there were, then by Corollary 3.4, the sets $\{a, b, c\}$, $\{c, d, e\}$, $\{e, g, h\}$, and $\{h, i, a\}$ would have to be cyclic intervals in this linear order, but that is impossible since f would have to be in one of those cyclic intervals. Thus, while M and N are positroids, their free amalgam is not. To show that B is an excluded minor for the class of positroids, by the symmetry of B, it suffices to show that $B \setminus x$ and B/x are positroids for each $x \in \{a, b, c, f\}$.

Each contraction B/x with $x \in \{a, b, c, f\}$ is a parallel extension of a matroid that either has at most two proper nonempty cyclic flats or satisfies the hypothesis of Corollary 3.12, so B/x is a positroid by Corollary 4.18. The deletion $B \setminus f$ is the whirl W^4 , which is a multi-path matroid, and $B \setminus b$ is a parallel connection of W^3 and $U_{2,3}$, and so is a positroid. The deletion $B \setminus a$ is a positroid since its four proper connected flats F with $|F| \ge 2$, namely, $\{c, d, e\}$, $\{e, g, h\}$, $\{f, b, c, d, e\}$, $\{e, g, h, i, f\}$, are cyclic intervals in the linear order b < c < d < e < g < h < i < f. Lastly, for $B \setminus c$, the proper connected flats $\{h, i, a\}$, $\{e, g, h\}$, $\{a, b, d, f, e\}$, and $\{f, e, g, h, i, a\}$ are cyclic intervals in the linear order a < b < d < f < e < g < h < i, and for $F = \{a, f, e\}$, the only proper connected flat with $|F| \ge 2$ that is not a cyclic interval, the two connected components of $(B \setminus c)/F$, both of rank 1, are $\{b, d\}$ and $\{g, h, i\}$, which, as needed, are cyclic intervals. Examples 4–12 give infinite families of excluded minors for the class of positroids, most with multiple parameters. Example 4 includes $M(K_4)$ and Example 9 includes free extensions of rank-3 truncations of whirls.

Example 4. Let \mathcal{L} be $\{\{1,3,5\}, \{2,3,4\}, \{4,5,6\}, \{1,2,6\}\}$, the set of 3-point lines of the cycle matroid $M(K_4)$ as labeled in Figure 2. Let X_1, X_2, \ldots, X_6 be pairwise disjoint nonempty sets that satisfy the following conditions. Let $x_i = |X_i|$. The set X_2 is arbitrary. Pick X_1, X_3, X_4 , and X_6 so that $x_3 + x_4 = x_1 + x_6$. Let r be $x_2 + x_3 + x_4$, which is also $x_1 + x_2 + x_6$. We may assume that $x_1 + x_3 \leq x_4 + x_6$. Choose X_5 so that $x_1 + x_3 + x_5 \leq r \leq x_4 + x_5 + x_6$. Let $E = X_1 \cup \cdots \cup X_6$, and let \mathcal{Z} consist of \emptyset , E, and the four sets $X_i \cup X_j \cup X_k$ for $\{i, j, k\} \in \mathcal{L}$, and let the ranks of these sets be as given in the table below.

Ø	$X_1 \cup X_3 \cup X_5$	$X_2 \cup X_3 \cup X_4$	$X_4 \cup X_5 \cup X_6$	$X_1 \cup X_2 \cup X_6$	E
0	$x_1 + x_3 + x_5 - 1$	r-1	r-1	r-1	r

Properties (Z0) and (Z1) in Theorem 2.11 clearly hold. The only case of property (Z2) that is not immediate is when $X = X_1 \cup X_3 \cup X_5$ and Y = E, in which case we need r(Y) - r(X) < |Y - X|, that is, $r - (x_1 + x_3 + x_5 - 1) < x_2 + x_4 + x_6$, that is, r + 1 < |E|, which is clear. Thus, property (Z2) holds. Each instance of the inequality in property (Z3) for two incomparable cyclic flats other than $X_1 \cup X_3 \cup X_5$ is an inequality of the form $r + x_i \le 2(r - 1)$, i.e., $x_i \le r - 2$, which holds since the sum of x_i and two other positive integers is at most r. The inequality required for property (Z3) in each case that involves $X_1 \cup X_3 \cup X_5$ has the form $r + x_i \le r - 1 + x_1 + x_3 + x_5 - 1$ with $i \in \{1, 3, 5\}$, i.e., $x_i \le x_1 + x_3 + x_5 - 2$, which holds since $x_1, x_3, x_5 \in \mathbb{N}$. Thus, the cyclic flats and ranks given above define a matroid M on E.

We now show that M is an excluded minor for the class of positroids. By Lemma 2.12, the restriction to and contraction by $X_i \cup X_j \cup X_k$, for each $\{i, j, k\} \in \mathcal{L}$, is a uniform matroid of positive nullity, and so is connected. If M were a positroid, then each of these connected flats would be a cyclic interval in any positroid order for M; that is impossible by Lemma 2.1, so M is not a positroid. Each cyclic flat $X_i \cup X_j \cup X_k$ for $\{i, j, k\} \in \mathcal{L} - \{\{4, 5, 6\}\}$ is a circuit $(X_4 \cup X_5 \cup X_6 \text{ might be a circuit})$ and each cyclic flat $X_i \cup X_j \cup X_k$ for $\{i, j, k\} \in \mathcal{L} - \{\{4, 5, 6\}\}$ is a circuit $(X_4 \cup X_5 \cup X_6 \text{ might be a circuit})$ and each cyclic flat $X_i \cup X_j \cup X_k$ for $\{i, j, k\} \in \mathcal{L} - \{\{1, 3, 5\}\}$ is a hyperplane $(X_1 \cup X_3 \cup X_5 \text{ might be a hyperplane})$. Thus, each element e is in either one or two cyclic flats that are circuits, and not in either one or two cyclic flats that are hyperplanes. So $M \setminus e$ either has only four cyclic flats or the hypotheses of Corollary 3.12 apply, and likewise for M/e, so these minors of M are positroids. Thus, M is an excluded minor for the class of positroids.

In Example 4, the proper, nonempty cyclic flat (if any) that is not a circuit and the one (if any) that is not a hyperplane are not equal. In the next example, they are the same.

Example 5. Fix $a, b, c, s, r \in \mathbb{N}$ for which (i) $\max(a, b, c) < s < a + b + c$, (ii) r has the same parity as a + b + c, and (iii) $\max(s, a + b - c, a + c - b, b + c - a) < r$. Let \mathcal{L} be as in Example 4. Pick pairwise disjoint sets X_1, X_2, \ldots, X_6 for which $|X_4| = a, |X_5| = c$, $|X_6| = b$, and

$$|X_1| = \frac{r+a-b-c}{2}, \qquad |X_2| = \frac{r-a-b+c}{2} \qquad |X_3| = \frac{r-a+b-c}{2}.$$

Thus, $|X_4 \cup X_5 \cup X_6| = a + b + c$ and $|X_i \cup X_j \cup X_k| = r$, for $\{i, j, k\} \in \mathcal{L} - \{\{4, 5, 6\}\}$. Let $E = X_1 \cup \cdots \cup X_6$, and let \mathcal{Z} consist of \emptyset , E, and the four sets $X_i \cup X_j \cup X_k$ for $\{i, j, k\} \in \mathcal{L}$, and let the ranks of these sets be as given in the table below.

Ø	$X_1 \cup X_3 \cup X_5$	$X_2 \cup X_3 \cup X_4$	$X_1\cup X_2\cup X_6$	$X_4 \cup X_5 \cup X_6$	E
0	r-1	r-1	r-1	s	r

32

Properties (Z0) and (Z1) in Theorem 2.11 clearly hold. The only case of property (Z2) that is not transparent is when $X = X_4 \cup X_5 \cup X_6$ and Y = E, for which we must show that r(Y) - r(X) < |Y - X|, that is, $r - s < |X_1 \cup X_2 \cup X_3|$. That inequality simplifies to a + b + c < r + 2s, which holds since a + b + c < 3s < r + 2s by conditions (i) and (iii). Thus, property (Z2) holds. Each instance of the inequality in property (Z3) for two incomparable cyclic flats other than $X_4 \cup X_5 \cup X_6$ is an inequality of the form $r + |X_i| \le 2(r - 1)$, i.e., $|X_i| \le r - 2$, which holds since $|X_i|$ and two other positive integers add to r. The inequality required for property (Z3) in each case that involves $X_4 \cup X_5 \cup X_6$ has the form $r + |X_i| \le r - 1 + s$ with $i \in \{4, 5, 6\}$, i.e., $|X_i| < s$, which holds by the inequality max(a, b, c) < s in condition (i). Thus, the cyclic flats and their ranks define a matroid M on E. The same argument as used in the last example shows that M is an excluded minor for the class of positroids.

We next give three more infinite families of excluded minors for the class of positroids, along with their duals, that are different variations on the idea in Example 4. We omit the proofs that these are excluded minors since no new ideas are required.

Example 6. Fix $a, b, c, k \in \mathbb{N}$ and let X_1, \ldots, X_6 be pairwise disjoint sets with $|X_3| = a$, $|X_6| = a + k, |X_4| = b, |X_1| = b + k, |X_5| = c$, and $|X_2| = c + k$. Fix an element p that is in none of X_1, \ldots, X_6 , and let $E = X_1 \cup \cdots \cup X_6 \cup p$. Let the paving matroid M of rank a+b+c+k+1 on E have as its dependent hyperplanes the three circuit-hyperplanes $X_1 \cup X_3 \cup X_5 \cup p, X_2 \cup X_3 \cup X_4 \cup p$, and $X_4 \cup X_5 \cup X_6 \cup p$, along with $X_1 \cup X_2 \cup X_6$, which has nullity 2k. It is easy to check that M is an excluded minor for the class of positroids. In the dual, M^* , of M, which has rank a+b+c+2k, the sets $X_2 \cup X_4 \cup X_6$, $X_1 \cup X_5 \cup X_6$, and $X_1 \cup X_2 \cup X_3$ are circuit-hyperplanes, while $X_3 \cup X_4 \cup X_5 \cup p$ is a circuit but not a hyperplane.

Example 7. Fix $a, b, c \in \mathbb{N}$ and let the sets X_1, \ldots, X_6 be pairwise disjoint and satisfy $|X_3| = |X_6| = a$, $|X_1| = b + 1$, $|X_4| = b$, and $|X_2| = |X_5| = c$. Fix elements p and q that are in none of X_1, \ldots, X_6 , and let $E = X_1 \cup \cdots \cup X_6 \cup \{p, q\}$. The sets

 $X_1 \cup X_3 \cup X_5 \cup p, \ X_2 \cup X_3 \cup X_4 \cup \{p,q\}, \ X_4 \cup X_5 \cup X_6 \cup \{p,q\}, \ X_1 \cup X_2 \cup X_6 \cup q$

are the circuit-hyperplanes of a sparse paving matroid M of rank a + b + c + 2 on E. It is easy to check that M is an excluded minor for the class of positroids. The sparse paving matroid M^* has rank a + b + c + 1 and its circuit-hyperplanes are $X_2 \cup X_4 \cup X_6 \cup q$, $X_1 \cup X_5 \cup X_6, X_1 \cup X_2 \cup X_3$, and $X_3 \cup X_4 \cup X_5 \cup p$.

Example 8. Fix $a, b, c \in \mathbb{N}$ and let the sets X_1, \ldots, X_6 be pairwise disjoint and satisfy $|X_3| = |X_6| = a$, $|X_1| = |X_4| = b$, and $|X_2| = |X_5| = c$. Fix four elements p, q, s, t that are in none of X_1, \ldots, X_6 , and let $E = X_1 \cup \cdots \cup X_6 \cup \{p, q, s, t\}$. The sets

$X_1 \cup X_3 \cup X_5 \cup \{q, s, t\},$	$X_2 \cup X_3 \cup X_4 \cup \{p, s, t\}$
$X_4 \cup X_5 \cup X_6 \cup \{p, q, t\},$	$X_1 \cup X_2 \cup X_6 \cup \{p, q, s\}$

are the circuit-hyperplanes of a sparse-paving matroid M on E of rank a + b + c + 3. It is easy to check that M is an excluded minor for the class of positroids. The sparse-paving matroid M^* has the circuit-hyperplanes $X_2 \cup X_4 \cup X_6 \cup p$, $X_1 \cup X_5 \cup X_6 \cup q$, $X_1 \cup X_2 \cup X_3 \cup s$, and $X_3 \cup X_4 \cup X_5 \cup t$, and its rank is a + b + c + 1.

Let L be the lattice of flats of $M(K_4)$ and let M be a loopless matroid whose lattice of cyclic flats is isomorphic to L. Note that if the cyclic flats of a matroid N are either (i) just \emptyset and E(N), or (ii) just \emptyset , E(N), and three other sets, none of which contains either of the other two, then N is connected since its lattice of cyclic flats is not a direct



FIGURE 12. The matroids in Example 10 for ranks 3 and 4, with the 4-circuits in the rank-4 matroid drawn in the faces of a triangular prism.

product of two other lattices. By Lemma 2.12, for any cyclic flat F_1 of M that corresponds to a point of $M(K_4)$ and any cyclic flat F_2 of M that corresponds to a 3-point line of $M(K_4)$, condition (i) holds for $\mathcal{Z}(M|F_1)$ and $\mathcal{Z}(M/F_2)$ while condition (ii) holds for $\mathcal{Z}(M/F_1)$ and $\mathcal{Z}(M|F_2)$, so all such minors are connected. As above, it follows that M is not a positroid. Thus, while every lattice is isomorphic to the lattice of cyclic flats of some matroid [7], the same is not true for positroids.

The rank-*n* whirl, W^n , is a multi-path matroid, so it and its rank-3 truncation are positroids, but the free extension of the rank-3 truncation of W^n is an excluded minor for the class of positroids. The next example treats these excluded minors and many more.

Example 9. Fix $n, r, x_1, x_2, \ldots, x_{2n} \in \mathbb{N}$, with $n \ge 3$, that satisfy the properties below, where we interpret indices modulo 2n and set $m_i = x_{2i-2} + x_{2i-1} + x_{2i}$ for all $i \in [n]$:

- (i) $3 \le m_i \le r$ for all $i \in [n]$,
- (ii) if n = 3, then all m_i are r, while if n > 3, then $m_i = r = m_j$ for at least two elements i and j of [n] that are not cyclically consecutive,

(iii) if
$$i, j \in [n]$$
 with $j \neq i$ and $j \neq i+1$, then $r < x_{2i} + x_{2j} + \sum_{k=i+1}^{j} x_{2k-1}$.

(To be clear, the sum in property (iii) is over the odd terms in a cyclic interval in [2n].) Let e_1, e_2, \ldots, e_{2n} be the elements of the *n*-whirl \mathcal{W}^n , labeled so that, for each $i \in [n]$, the set $\{e_{2i-2}, e_{2i-1}, e_{2i}\}$ is a 3-circuit, where $e_0 = e_{2n}$. For each $i \in [2n]$, apply series extension $x_i - 1$ times to e_i , so the 3-circuit $\{e_{2i-2}, e_{2i-1}, e_{2i}\}$ of \mathcal{W}^n yields an m_i -circuit C_i . Let M' denote the resulting matroid. Truncate the free extension M' + f to rank r to get a matroid M. We claim that M is an excluded minor for the class of positroids.

The circuits of the *n*-whirl W^n are the symmetric differences of consecutive 3-circuits, so the circuits of M' are the symmetric differences of the form $C_i \triangle C_{i+1} \triangle \cdots \triangle C_j$, where we interpret the subscripts modulo n. Thus, by property (iii), the only circuits of M of rank less than r are the circuits C_i for $i \in [n]$, so they are the only proper connected flats F of M with $|F| \ge 2$. Also, M/C_i is connected by part (3) of Lemma 2.10. So if M were a positroid, each circuit C_i would have to be an interval in any positroid order for M, but, since f is in no circuit C_i and $C_i \cap C_{i+1} \neq \emptyset$ for all $i \in [n]$, there is no positroid order for M, so M is not a positroid.

For each $e \in E(M)$, at least one circuit C_i of M fails to be a circuit of M/e by property (ii), so a now-routine use of Corollary 3.6 implies that M/e is a positroid. The same holds for $M \setminus e$ if $e \neq f$. Adapting a positroid order for a whirl to $M \setminus f$ shows that it too is a positroid. Thus, M is an excluded minor for the class of positroids. The next infinite family of excluded minors for the class of positroids can be seen as a variation on a 3-whirl, but with two *r*-circuits and a 3-circuit, and instead of taking a free extension, another 3-circuit is added.

Example 10. Fix an integer $r \ge 3$, take the parallel connection of two *r*-circuits, *A* and *B*, with base point *e*, truncate to rank *r*, fix $a_1, a_2 \in A - e$ and $b_1, b_2 \in B - e$, and then, using principal extension, add a point p_1 to the line $\{a_1, b_1\}$ and a point p_2 to the line $\{a_2, b_2\}$. The resulting matroid *M* has rank *r* and its proper, nonempty connected flats, each of which is a circuit, are *A*, *B*, $\{a_1, b_1, p_1\}$, and $\{a_2, b_2, p_2\}$. (See Figure 12 for the cases r = 3 and r = 4.) Contracting any of the proper, nonempty connected flats yields a connected matroid, and, by Lemma 2.1, no linear order on E(M) has each of *A*, *B*, $\{a_1, b_1, p_1\}$, and $\{a_2, b_2, p_2\}$ being a cyclic interval, so *M* is not a positroid. It is routine to check that all proper minors of *M* are positroids.

We close with two more infinite families of excluded minors for the class of positroids. With the ideas above, verifying that these are excluded minors is routine. Also, applying duality yields more excluded minors.

Example 11. For integers $n \ge k \ge 3$, take the parallel connection of two *n*-circuits and a *k*-circuit (or three *n*-circuits if n = k) at a common base point, and then truncate to rank *n*. The matroid obtained is an excluded minor for the class of positroids.

Example 12. For integers $n \ge k \ge 3$, take two *n*-circuits and a *k*-circuit (or three *n*-circuits if n = k) and a 3-point line; for each of the first three circuits, take its parallel connection with the line, using three different base points on the line, and then truncate to rank *n*. The resulting matroid is an excluded minor for the class of positroids.

ACKNOWLEDGMENTS

The author thanks Kolja Knauer for pointing out the work of Blum and the result of Lam and Postnikov that base-sortable matroids are the same as positroids. The author thanks Felipe Rincón for pointing him to Theorems 1.2 and 1.3 and recasting those results, which were originally stated in terms of matroid polytopes, in a form closer to what is given here, and for a useful discussion about Blum's conjecture that led to Theorem 3.5. The author also thanks both referees for their careful reading and helpful comments that should result in making this paper accessible to a wider audience.

REFERENCES

- S. Agarwala, C. Delaney, K. Yeats, Rado matroids and a graphical calculus for boundaries of Wilson loop diagrams, preprint, https://arxiv.org/abs/2401.05592 (2024).
- [2] F. Ardila, F. Rincón, and L. Williams, Positroids and non-crossing partitions, *Trans. Amer. Math. Soc.*, 368 (2016) 337–363, https://doi.org/10.1090/tran/6331.
- [3] S. Blum, Base-sortable matroids and Koszulness of semigroup rings, *European J. Combin.* (2001) 937–951, https://doi.org/10.1006/eujc.2001.0516.
- [4] J. Bonin, Lattice path matroids: the excluded minors, J. Combin. Theory Ser. B 100 (2010) 585–599, https://doi.org/10.1016/j.jctb.2010.05.001.
- [5] J. Bonin, A. de Mier, and M. Noy, Lattice path matroids: enumerative aspects and Tutte polynomials, J. Combin. Theory Ser. A, 104 (2003) 63–94, https://doi.org/10.1016/S0097-3165(03)00122-5.
- [6] J. Bonin and A. de Mier, Lattice path matroids: structural properties, *European J. Combin.*, 27 (2006) 701–738, https://doi.org/10.1016/j.ejc.2005.01.008.
- [7] J. Bonin and A. de Mier, The lattice of cyclic flats of a matroid, Ann. Comb. 12 (2008) 155–170, https://doi.org/10.1007/s00026-008-0344-3.
- [8] J. Bonin and O. Giménez, Multi-path matroids, Combin. Probab. Comput., 16 (2007) 193–217, https://doi.org/10.1017/S0963548306007942.

JOSEPH E. BONIN

- [9] J. Bonin and K. Long, Connectivity gaps among matroids with the same enumerative invariants, Adv. in Appl. Math. 154 (2024) Paper No. 102648, https://doi.org/10.1016/j.aam.2023.102648.
- [10] T.H. Brylawski, An affine representation for transversal geometries, *Studies in Appl. Math.* 54 (1975) 143–160, https://doi.org/10.1002/sapm1975542143.
- [11] H.H. Crapo, Single-element extensions of matroids, J. Res. Nat. Bur. Standards Sect. B 69B (1965) 55–65, https://doi.org/10.6028/jres.069b.003.
- [12] H. Crapo and W. Schmitt, The free product of matroids, *European J. Combin.*, 26 (2005) 1060–1065, https://doi.org/10.1016/j.ejc.2004.05.005.
- [13] H. Crapo and W. Schmitt, A unique factorization theorem for matroids, J. Combin. Theory Ser. A, 112, (2005) 222–249, https://doi.org/10.1016/j.jcta.2005.02.004.
- [14] R.J. Duffin, Topology of series-parallel networks, J. Math. Anal. Appl., 10 (1965) 303–318, https://doi.org/10.1016/0022-247X(65)90125-3.
- [15] J. Edmonds and D. R. Fulkerson, Transversals and matroid partition, J. Res. Nat. Bur. Standards Sect. B 69B (1965) 147–153, https://doi.org/10.6028/jres.069b.016.
- [16] D. Gale, Optimal assignments in an ordered set: An application of matroid theory, J. Combinatorial Theory, 4 (1968) 176–180, https://doi.org/10.1016/S0021-9800(68)80039-0.
- [17] T. Lam and A. Postnikov, Alcoved polytopes. I, Discrete Comput. Geom. 38 (2007) 453–478, https://doi.org/10.1007/s00454-006-1294-3.
- [18] T. Lam and A. Postnikov, Polypositroids, Forum Math. Sigma 12 (2024) Paper No. e42, 67, https://doi.org/10.1017/fms.2024.11.
- [19] S. Oh, Positroids and Schubert matroids, J. Combin. Theory Ser. A, 118 (2011) 2426–2435, https://doi.org/10.1016/j.jcta.2011.06.006.
- [20] J. Oxley, Matroid Theory, second edition (Oxford University Press, Oxford, 2011), https://doi.org/10.1093/acprof:oso/9780198566946.001.0001.
- [21] J. Oxley, K. Prendergast, and D. Row, Matroids whose ground sets are domains of functions, J. Austral. Math. Soc. Ser. A 32 (1982) 380–387, https://doi.org/10.1017/S1446788700024939.
- [22] H. Park, Excluded minors of interval positroids that are paving matroids, preprint https://arxiv.org/abs/2312.03525 (2023).
- [23] A. Postnikov, Total positivity, Grassmannians, and networks, https://arxiv.org/abs/math/0609764 (2006).
- [24] F. Rincón, C. Vinzant, and J. Yu, Positively hyperbolic varieties, tropicalization, and positroids, Adv. Math. 383 (2021) Paper No. 107677, 35, https://doi.org/10.1016/j.aim.2021.107677.
- [25] J.A. Sims, Some Problems in Matroid Theory, (Ph.D. Dissertation, Linacre College, Oxford University, Oxford, 1980).
- [26] W.T. Tutte, Matroids and graphs, Trans. Amer. Math. Soc., 90 (1959) 527–552, https://doi.org/10.1090/S0002-9947-1959-0101527-3.

DEPARTMENT OF MATHEMATICS, THE GEORGE WASHINGTON UNIVERSITY, WASHINGTON, D.C. 20052, USA

Email address: jbonin@gwu.edu

36